Wasserstein Bounds for generative diffusion models with Gaussian tail targets

Xixian Wang[∗] , Zhongjian Wang†

December 17, 2024

Abstract

We present an estimate of the Wasserstein distance between the data distribution and the generation of score-based generative models, assuming an ϵ -accurate approximation of the score and a Gaussian-type tail behavior of the data distribution. The complexity bound in dimension is $O(\sqrt{d})$, with a logarithmic constant. Such Gaussian tail assumption applies to the distribution of a compact support target with early stopping technique and the Bayesian posterior with a bounded observation operator. Corresponding convergence and complexity bounds are derived.

The crux of the analysis lies in the Lipchitz bound of the score, which is related to the Hessian estimate of a viscous Hamilton-Jacobi equation (vHJ). This latter is demonstrated by employing a dimension independent kernel estimate. Consequently, our complexity bound scales linearly (up to a logarithmic constant) with the square root of the trace of the covariance operator, which relates to the invariant distribution of forward process. Our analysis also extends to the probabilistic flow ODE, as the sampling process.

Keywords: Diffusion model, Lipschitz estimate, Score function, viscous Hamilton-Jacobi equation, convergence

1 Introduction

Diffusion models (DM) are among the state-of-the-art tools in the new GenAI era. As a generative model, the diffusion model first links its target distribution to some distribution easy to sample via a diffusive process (forward). The generative process (backward) then involves "reversing" the diffusion to enable sampling of the target distribution from a distribution that is easily sampled. A well-known mathematical model that encapsulates this approach is the score-based stochastic differential equation (SDE) model[\[25\]](#page-11-0), where the forward and backward processes are represented by two SDEs that share the same marginal distribution [\[1\]](#page-10-0). In most cases, the forward process is assumed to be an Ornstein-Uhlenbeck (OU) process. The backward SDE incorporates the gradient of the logarithmic density (score) of the forward process. When the explicit form of the score is unknown, the score is then estimated by a neural network from discrete samples of the target distribution.

A major direction for the theoretical study of DMs is the convergence of the approximated backward process with limited assumptions of data. When the Lipschitz bound of the score is available (regular target), the backward process is generally well-defined until $t = 0$, and convergence results are related to the Lipschitz bound. Otherwise, when the bound is unavailable (singular target), the early stopping technique is introduced, and convergence results are related to the stopping time. Various analytical approaches (reviewed in the related work part) have been adapted for these two types of assumptions.

In this work, we aim to present a general error analysis that applies to both regular and singular target distributions and generalizes to an infinite dimensional setting. The analysis is based on an dimensionless and uniform-in-space Lipschitz bound of the score, derived from a partial differential equation (PDE) analysis approach. Our bounds are under the Wasserstein metric and are linear (with a logarithmic constant) in the square root of variance of Brownian motion in the forward process (Theorem [3.3\)](#page-5-0). In the finite dimensional case with white noise as invariant measure, the variance is linear in the dimension and hence the complexity is $\mathcal{O}(\sqrt{d})$. For the infinite dimensional case, a Gaussian random field is taken as the base measure and the variance is linear with the trace of the covariance operator.

[∗]Division of Mathematical Sciences, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore. xixian001@e.ntu.edu.sg

[†]Division of Mathematical Sciences, School of Physical and Mathematical Sciences, Nanyang Technological University,Singapore. zhongjian.wang@ntu.edu.sg (Corresponding)

Table 1: Summary of previous bounds and our results for score based diffusion models in d dimensions. Bounds expressed in terms of the number of steps (N) required to guarantee an error of at most ϵ_0 in the stated metric. For ease of presentation, we assume \overrightarrow{P}_0 has finite second moments M_2 and is normalized so that $Cov(\vec{P}_0) = I_d$, and the forward process is an OU with standard Brownian motion. We ignore the dependence on Lipschitz constants of the score at $t = 0$.

TARGET P_0	METRIC	COMPLEXITY	RESULT
SUPP $B_R(0)$	$\mathcal{W}_1(\vec{P}_0, \overleftarrow{Q}_{T-\delta})$	$\log N = \mathcal{O}(\frac{R^2(d+R^4)^2(\log R)^2}{\epsilon_2^2})$	[10]: THM. $1 + \text{COR. } 2$
SUPP $B_R(0)$	$\mathcal{W}_2(\vec{P}_0, \vec{Q}_{T-\delta})$	$\log N = \mathcal{O}(\frac{R^2 d^2}{\epsilon^4})$	THIS WORK: COR. 3.13
SUPP $B_R(0)$	$\mathcal{W}_2(\vec{P}_\delta, \overset{\leftarrow}{Q}_{T-\delta})$	$N = \mathcal{O}(\frac{\sqrt{d}}{\epsilon_0} \log \frac{d}{\epsilon_0^2} \exp(\frac{1}{\delta} + 3\frac{R^2}{\delta^2})\frac{R^4}{\delta^4})$	THIS WORK: COR. 3.12
$\mathbb{E} X_0 ^2 < \infty$	$\text{KL}(\overrightarrow{P}_{\delta} \overleftarrow{Q}_{T-\delta})$	$N = \mathcal{O}(\frac{d \log^2 \frac{1}{\delta}}{\epsilon^2})$	$ 3 $ Cor 1
∇ log \overrightarrow{P}_0 <i>L</i> -LIP	$KL(\vec{P}_0 \vec{Q}_T)$	$N = \mathcal{O}(\frac{d^2}{\epsilon_0})$	[6] THM 5
$\mathcal{F}(\vec{P}_0 \mathcal{N}(0,I_d)) \lesssim d$ (*)	$KL(\vec{P}_0 \vec{Q}_T)$	$N = \mathcal{O}(\frac{d}{\epsilon_0^2} \log \frac{d}{\epsilon_0})$	$[7]$ Thm 1
P_0 LOG-CONCAVE	$\mathcal{W}_2(\overrightarrow{P}_0,\overleftarrow{Q}_T)$	$N = \mathcal{O}(\frac{d}{\epsilon_0^2} \log \frac{d}{\epsilon_0})$	[13] TABLE 2
G TAIL, ASS.2	$\mathcal{W}_2(\vec{P}_0, \overleftarrow{Q}_T)$	$N = \mathcal{O}(\frac{\sqrt{d}}{\epsilon_0} \log \frac{d}{\epsilon_0^2})$	THIS WORK: COR. 3.4

^{*} Here $\mathcal F$ denotes relative Fisher information. One can show our Gaussian tail Assumption [2](#page-3-0) \Rightarrow $\mathcal{F}(\vec{P}_0|\mathcal{N}(0,I_d)) \lesssim d$, while in standard Gaussian case, they are equivalent.

It is worth noting that the results of this work adopt the Wasserstein-2 distance as the metric of error (instead of Kullback–Leibler (KL) divergence) due to its flexibility. More precisely, the reasons are two-fold: (1) In practical applications of diffusion models for the generation of structured data (image, text, video, protein...), the target distributions mostly find their support in a compact sub-manifold, see further discussion of the manifold hypothesis in [\[27,](#page-11-2) [2\]](#page-10-5). Consequently, under this hypothesis, standard KL divergence cannot be consistently defined between the backward process whose support is the entire space and the target distribution with compact support. (2) In high (towards infinite) dimension settings, it becomes necessary to compactify the forward process. For instance, by setting C , the covariance matrix (operator) in the forward process [\(1\)](#page-2-0), with finite trace, the invariant distribution of forward process thus has finite second order moment. The Wasserstein distance scales with the trace of C and hence consistent with the compactification towards infinite dimension generative models. While the KL divergence scales with the dimension and hence cannot be simply generalized to a dimensionless result. In Appendix [B.1](#page-23-0) we provide a motivating example to illustrate.

Related work

Complexity bounds [\[9\]](#page-10-6) obtains the first convergence guarantees in 1-Wasserstein distance assuming that the data distribution satisfies the so-called manifold hypothesis. Some recent works [\[24,](#page-11-3) [14,](#page-11-4) [13\]](#page-11-1) established convergence guarantees under Gaussian and log-concave distributions.

We are also aware several complexity bounds under KL divergence, for instance [\[6,](#page-10-3) [3,](#page-10-2) [7\]](#page-10-4). A common point of these approaches is the utilizing the chain rule of KL divergence, which is a consequence of Girsanov theorem, to separate the global error into local truncation errors. To bound local ones, the probabilistic view point, which estimates the score function under the distribution of forward process, kicks in. In this work, we instead adopt a PDE approach, which provides point-wise estimate of the score. Such estimate also facilitates analysis of score under the approximated backward process and hence provides an Wasserstein bound. In Table[.1,](#page-0-0) we present the comparison of the complexity bounds. Diffusion model in infinite dimension

Typically, diffusion models operates on finite-dimensional spaces. However, in many domains, the underlying signal is infinite-dimensional, and the observed data is a collection of discrete observations of some underlying function, for instance, Bayesian inverse problems [\[26\]](#page-11-5). Up to now, there are many studies applying diffusion models to functional spaces [\[18,](#page-11-6) [16,](#page-11-7) [22,](#page-11-8) [11\]](#page-11-9).

While many theoretical studies $[10, 6, 3]$ $[10, 6, 3]$ $[10, 6, 3]$ suggest that performance guarantees deteriorate with increasing dimension, [\[22\]](#page-11-8) established a dimension-independent convergence rate of the distance from the samples to the target measure in the Wasserstein-2-distance. However, the distance is unideal since it grows exponentially over the running time T . We follow the main steps of the proof, which is a standard

Lyapunov type error estimate [\[17\]](#page-11-10). While with the Gaussian tail assumption, we get a better bound for the non-linear perturbation part (see Corollary [3.2\)](#page-5-2) which decrease exponentially over the running time T in finite dimensional cases. This and together with scheme (6) ensure the accumulation error of discretization does not grow exponential in time, compared with standard estimates. Our approach can also be extended to the infinite-dimensional framework proposed in [\[22\]](#page-11-8).

Lipschitz bounds of the score The score functions in the SGMs are related to the gradient of logdensity ($\log p$) of the forward process. It is well known that the function $\log p$ itself follows a viscous Hamilton Jacobi (vHJ) equation, see [\(15\)](#page-4-0) in later discussion. Then the Lipchitz bounds of the scores are equivalent to Hessian bounds for a vHJ equation. There are various regularity results in the literature for the original Fokker Planck equation or the transformed vHJ, see [\[5,](#page-10-7) [12\]](#page-11-11) and most recent results in [\[20,](#page-11-12) [4,](#page-10-8) [21\]](#page-11-13). We would point out that except [\[21\]](#page-11-13), most results are seeking for a spatially global Hessian bound which only last finite time without the Gaussian tail assumption (Assumption [2\)](#page-3-0) in this work. [\[21\]](#page-11-13) also provides a local-in-space and global-in-time bound while only polynomial in dimension $(d³)$ complexity can be shown from it due to the unknown tail behavior.

2 Preliminaries

2.1 Background and Setting the Stage

A large class of generative diffusion models can be analyzed under the SDE framework, it contains two processes: forward and backward. The forward process, which gradually transforms the data distribution into white noise is an OU process as follows:

$$
d\vec{X}_t = -\frac{1}{2}\vec{X}_t dt + \sqrt{C} dB_t, \quad 0 \le t \le T. \tag{1}
$$

where B_t is a standard Brownian motion, C is a symmetric, positive-definite covariance matrix and T is the final time such that the distribution of X_T is close to the Gaussian distribution $\mathcal{N}(0, C)$. The initial \vec{X}_0 follows the target(data) distribution, denoted as p_0 . We use $p_t(x)$ to denote the density of \vec{X}_t . Then with time reversal $t \to T - t$, the backward process is defined as:

$$
d\overleftarrow{X}_t = (\frac{1}{2}\overleftarrow{X}_t + C\nabla \log p_{T-t}(\overleftarrow{X}_t))dt + \sqrt{C}d\tilde{B}_t, \tag{2}
$$

where \tilde{B}_t is also a standard Brownian motion (may not being the same as B_t . The term,

$$
s(t,x) = C \nabla \log p_t(x),\tag{3}
$$

is generally referred as the score function. The process $(X_t)_{0 \leq t \leq T}$ transforms noise into samples follows p_0 .

To be specific, p_t will solve the Fokker-Planck equation with Cauchy data p_0 :

$$
\begin{cases} \partial_t p = \frac{1}{2} (\nabla \cdot (xp) + \nabla \cdot C \nabla p), \\ p(0, x) = p_0(x). \end{cases} \tag{4}
$$

We also denote \overrightarrow{P}_t (correspondingly \overleftarrow{P}_t) as the marginal distribution of \overrightarrow{X}_t in $(1)(\overleftarrow{X}_t$ in $(2))$. Given initial distribution for [\(2\)](#page-2-1) $\overleftrightarrow{P}_0 = \overrightarrow{P}_T$, then $\forall 0 \le t \le T$, $\overleftrightarrow{P}_t = \overrightarrow{P}_{T-t}$, especially $\overleftrightarrow{P}_T = \overrightarrow{P}_0 = p_0$.

However, in practice the score function $s(t, x) = C \nabla \log p_t(x)$ is not available since no closed form expression of p_0 is known. Thus we learn the score function by a neural network $s_{\theta}(t, x)$, where θ denotes latent variables of neural network. We training the network by optimizing an L_2 estimation loss,

$$
\mathbb{E}_{p_t} || s_{\theta}(t, x) - C \nabla \log p_t(x) ||^2.
$$

Given the estimated score s_{θ} (assumed to be ϵ accurate specified in Assumption [3,](#page-5-3) we can generate samples by an approximation of the backward process starting from the Gaussian distribution $\mathcal{N}(0, C)$:

$$
d\dot{\overline{Y}}_t = \left(\frac{1}{2}\dot{\overline{Y}}_t + s_\theta(T - t, \dot{\overline{Y}}_t)\right)dt + \sqrt{C}d\tilde{B}_t,\tag{5}
$$

Time discretization We employ an Euler-type discretization of the continuous-time stochastic process which will be helpful to the convergence theorems. Let $\delta = t_0 \leq t_1 \leq \cdots \leq t_N = T$ be the discretization points, $\delta = 0$ for the normal setting and $\delta > 0$ for the early-stopping setting. The process with initial distribution $\mathcal{N}(0, C)$ in the discrete scheme is as follows:

$$
d\overleftarrow{Y}_t = \left(-\frac{1}{2}\overleftarrow{Y}_t + \overleftarrow{Y}_{t_k} + s_\theta(T - t_k, \overleftarrow{Y}_{t_k})\right)dt + \sqrt{C}d\hat{B}_t
$$

\n
$$
t \in [t_k, t_{k+1}], \quad k = 0, 1, \cdots, N - 1.
$$
\n(6)

We denote \overleftarrow{Q}_t as the marginal distribution of \overleftrightarrow{Y}_t . This process admits an explicit solution, with $z_k \sim$ $\mathcal{N}(0, C),$

$$
\tilde{Y}_{t_{k+1}} = (2 - e^{\frac{t_k - t_{k+1}}{2}}) \tilde{Y}_{t_k} \n+ 2(1 - e^{\frac{t_k - t_{k+1}}{2}}) s_{\theta} (T - t_k, \tilde{Y}_{t_k}) \n+ \sqrt{1 - e^{t_k - t_{k+1}}} z_k.
$$
\n(7)

Remark 2.1. The discretization [\(7\)](#page-3-2) approximately corresponds to the discrete-time scheme introduced in [\[15\]](#page-11-14), it can also be found in [\[10\]](#page-10-1).

General Notations Let γ_C be the density of Gaussian measure $\mathcal{N}(0, C)$. For an $n \times n$ matrix A, we use the operator norm $\|\cdot\|$:

$$
||A|| = \sup_{v \neq 0} \frac{|Av|}{|v|} :=
$$
 the largest eigenvalue of $\sqrt{A^T A}$.

For a symmetric, positive-definite $n \times n$ matrix A, we use $|\cdot|_A$ to denote weighted l_2 norm in \mathbb{R}^n such that

$$
|x|_A^2 := \langle x, A^{-1}x \rangle.
$$

When A is identity matrix, we neglect the letter for simplicity and $|\cdot|$ is the standard l_2 norm in \mathbb{R}^n .

For vector (matrix, correspondingly) value function f with x as variable, $|f|_{\infty} := \sup_x |f(x)|$ (correspondingly, $||f||_{\infty} := \sup_x ||f(x)||$.

For compactness of the process, we made the following assumption on finite second order moment of the forward process.

Assumption 1. The data distribution has a bounded second moment,

$$
M_2 := \mathbb{E}_{p_0}|x|^2 < \infty. \tag{8}
$$

We further denote,

$$
M_0 = \max\{\text{Tr}(C), M_2\},\tag{9}
$$

relates to the maximum second order moment during the forward process.

2.2 Viscous Hamilton Jacobi Equation approaches

The foundation of our analysis is investigating the behaviour of $\log p$ as the solution of a viscous Hamilton Jacobi Equation(vHJ). Before that, we need the following assumption to derive reasonable point-wise estimates of gradients of the score function,

Assumption 2. The density of target distribution $p_0 \in C^2(\mathbb{R}^d)$ and has the following tail decomposition,

$$
p_0(x) = \exp(-\frac{|x|_A^2}{2}) \exp(h(x)),
$$
\n(10)

where there are independent of dimension constants such that,

 (i) A is a symmetric, positive-definite matrix which can be simultaneously diagonalized with C and

$$
||AC^{-1}|| < \infty, \quad ||CA^{-1}|| < \infty.
$$
 (11)

(ii) the remainder term h follows

$$
|\sqrt{C}\nabla h|_{\infty} < \infty, \quad ||C\nabla^2 h||_{\infty} < \infty.
$$
 (12)

[\(10\)](#page-3-3) is equivalent to the following in log density form,

$$
\log p_0(x) = h(x) - \frac{|x|_A^2}{2}.\tag{13}
$$

Remark 2.2. The assumption assumes the tail distribution of target is similar to a Gaussian distribution with covariance matrix A. This is a stronger assumption (less general) than L-Lipchitz of $\nabla \log p_0$. As suggested by Example 3.4 of [\[21\]](#page-11-13) L-Lipchitz of $\nabla \log p_0$ does not guarantee a Lipchitz bound of $\nabla \log p_t$. While the Gaussian tail is more general than the log-concaveness assumption, as one in [\[13\]](#page-11-1) and [\[14\]](#page-11-4).

In later Theorem [3.8,](#page-6-0) we show that density of forward process at the early stopping time δ follows the assumption with $A = (1 - \exp(-\delta))I_d$.

Our PDE approach considers the score potential function after the following transform,

$$
q(t,x) = -\log p_t(x) - \frac{x^T \bar{A}_t^{-1} x}{2},\tag{14}
$$

where $\bar{A}_t = Ae^{-t} + C(1 - e^{-t})$. Then q satisfies the PDE,

$$
\begin{cases} \n\frac{\partial_t q - \frac{1}{2} \nabla \cdot C \nabla q + \frac{1}{2} |\nabla q|_C^2}{\n+ (C \bar{A}_t^{-1} - \frac{1}{2} I) x \cdot \nabla q = \frac{1}{2} \text{Tr}(C \bar{A}_t^{-1} - I), \\
q(0, x) = h(x). \n\end{cases} \tag{15}
$$

To simplify [\(15\)](#page-4-0), we let $f(t) = \frac{1}{2} \int_0^t \text{Tr}(C \overline{A}_s^{-1} - I) ds$ and make a two step change of variables: let $K(t) = (A\bar{A}_t^{-1})e^{-\frac{t}{2}}, \text{ then}$

$$
\bar{q}(t,x) = q(t, K(t)^{-1}\sqrt{C}x) + f(t),
$$
\n(16)

satisfies the PDE:

$$
\begin{cases} \partial_t \bar{q} - \frac{1}{2} \nabla \cdot K(t)^2 \nabla \bar{q} + \frac{1}{2} |K(t) \nabla \bar{q}|^2 = 0, \\ \bar{q}(0, x) = h(\sqrt{C}x) := \bar{h}(x). \end{cases} \tag{17}
$$

Lastly, we define

$$
\tilde{q}(t,x) = e^{-\bar{q}(t,x)},\tag{18}
$$

which satisfies

$$
\begin{cases} \partial_t \tilde{q} - \frac{1}{2} \nabla \cdot K(t)^2 \nabla \tilde{q} = 0, \quad on \ (0, \infty) \times \mathbb{R}^n, \\ \tilde{q}(0, x) = e^{-\bar{h}(x)}. \end{cases} \tag{19}
$$

The solution is given by

$$
\frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \frac{1}{\sqrt{\det B(t)}} \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy,
$$
 (20)

where $B(t) = \int_0^t K(s)^2 ds = (e^{\frac{t}{2}} - e^{-\frac{t}{2}})K(t)$.

3 Results

In this section, we list the theoretical results and their detailed proofs are provided in Appendix [A.](#page-12-0) Section [3.1](#page-4-1) devotes to the PDE approach to bound of score with Gaussian tail assumption. Section [3.2](#page-5-4) listed the fundamental convergence result in Wasserstein metric. Section [3.3](#page-6-1) and Section [3.4](#page-8-0) are two applications of the convergence result under manifold assumption and Bayesian inverse problem assumption correspondingly. Additionally in Section [3.5,](#page-8-1) we discussed migration of our results to probability flow ODE as the sampling process.

3.1 Lipschitz Bound of Score Function

 \overline{a}

First, we give a Lipschitz bound of the score function under the tail Assumption [2,](#page-3-0) which plays a key role in our convergence theorem.

Theorem 3.1. Suppose that Assumption [2](#page-3-0) holds, then $\bar{q}(t, x)$ in [\(17\)](#page-4-2) satisfies, for $\forall t \in [0, \infty)$,

$$
|\nabla \bar{q}(t,\cdot)|_{\infty} \le |\nabla \bar{q}(0,\cdot)|_{\infty},\tag{21}
$$

$$
|\nabla^2 \bar{q}(t, \cdot)||_{\infty} \le ||\nabla^2 \bar{q}(0, \cdot)||_{\infty} + ||\nabla \bar{q}(0, \cdot)||_{\infty}^2.
$$
\n(22)

Proof see Appendix [A.1.](#page-12-1)

Corollary 3.2. Suppose that Assumption [2](#page-3-0) holds, consider the modified score function,

$$
\tilde{s}(t,x) := C \nabla \log p_t(x) + C \bar{A}_t^{-1} x,\tag{23}
$$

we have the following estimates,

$$
\|\nabla \tilde{s}\|_{\infty} \le e^{-t} L_0, \quad |s|_{\infty} \le e^{-\frac{t}{2}} L_1,\tag{24}
$$

where

$$
K = \sup_{t \ge 0} \|A\bar{A}_t^{-1}\| \le \max\{1, \|AC^{-1}\|\},\tag{25}
$$

$$
L_0 = K^2(||C\nabla^2 h||_{\infty} + |\sqrt{C}\nabla h|_{\infty}^2),\tag{26}
$$

$$
L_1 = K \|C\|^{\frac{1}{2}} |\sqrt{C} \nabla h|_{\infty},\tag{27}
$$

are bounded constants independent of dimension as per Assumption [2.](#page-3-0)

Proof see Appendix [A.2.](#page-12-2)

3.2 Main Convergence theories

We now study how far the samples generated by the scheme [\(6\)](#page-3-1) lie from the true target distribution p_0 . We do this in the Wasserstein-2-distance,

$$
\mathcal{W}_2(\mu,\nu) = \left(\inf_{\kappa} \int |x-y|^2 d\kappa(x,y)\right)^{1/2},
$$

where κ runs over all measures which have marginals μ and ν . To obtain the convergence results, we assume the following bound of score approximation at the discretization points,

Assumption 3. For each time discretization point t_k ,

$$
E_{\overrightarrow{P}_{T-t_k}} |s(T-t_k, x) - s_\theta(T-t_k, x)|^2 \le \epsilon^2.
$$

We would point out due to the limited access of the score approximation error, when deriving the complexity bounds, we always assume for sufficiently accurate score approximation. In Remark [3.6](#page-6-2) , we provide a slightly more general assumption.

In addition to score approximation error, we made the following assumption on gradient of the learned score to regulate the approximated backward process.

Assumption 4. Denote $\tilde{s}_{\theta}(t, x) = s_{\theta}(t, x) + C\overline{A}(t)x$, we assume that $\tilde{s}_{\theta}(t, x)$ shares the same properties as $\tilde{s}(t, x)$ in Corollary [3.2,](#page-5-2) i.e.

$$
\|\nabla \tilde{s}_{\theta}(t,\cdot)\|_{\infty} \le L_0 e^{-t}.\tag{28}
$$

The assumption [4](#page-5-5) can be relaxed to bounded gradient of score and yield a similar complexity bound with a different logarithmic constant, see Remark [3.7](#page-6-3)

Now we present the main theorem in the convergence analysis.

Theorem 3.3. Suppose Assumption [1,](#page-3-4) [2,](#page-3-0) [3,](#page-5-3) [4](#page-5-5) hold, using uniform discretization points in (6) with step size $h = t_{k+1} - t_k \leq 1$ yields,

$$
\mathcal{W}_2(\vec{P}_\delta, \overleftarrow{Q}_{T-\delta}) \le \exp(\tilde{K}_2 e^{-\delta}) \Big(e^{-T + \frac{\delta}{2}} \sqrt{M_2 + \text{Tr}(C)} + \tilde{K}_1 \sqrt{M_0} h + 2\epsilon \Big). \tag{29}
$$

where M_0 is defined in [\(9\)](#page-3-5), \tilde{K}_1, \tilde{K}_2 are some constants which will be specified in the proof.

Proof see Appendix [A.3.](#page-13-0)

With the result in Theorem [3.3,](#page-5-0) we can get the complexity bound by directly computation:

Corollary 3.4. Suppose Assumption [1,](#page-3-4) [2,](#page-3-0) [3,](#page-5-3) [4](#page-5-5) hold, using uniform discretization points in [\(6\)](#page-3-1) with step size $t_{k+1} - t_k \leq 1$ to reach a distribution \overleftarrow{Q}_T such that $\mathcal{W}_2(\overrightarrow{P}_0, \overleftarrow{Q}_T) = O(\epsilon_0)$ requires:

$$
T = \mathcal{O}(\log \frac{M_2 + Tr(C)}{\epsilon_0^2}),\tag{30}
$$

$$
N = \frac{T}{h} = \mathcal{O}(\frac{T}{\epsilon_0}\sqrt{M_0}),\tag{31}
$$

and Assumption [3](#page-5-3) to hold with

$$
\epsilon = \mathcal{O}(\epsilon_0). \tag{32}
$$

The complexity bound in Corollary [3.4](#page-5-1) does not depends on the dimension and hence naturally extends to infinite dimensional generative models. See discussion on definition in Appendix [B.2](#page-24-0) and an example application as Bayesian inverse problems in function space in Section [3.4.](#page-8-0)

Now we provide three remarks, first on optimality of the result and two on the relaxation of assumptions on score approximations.

Remark 3.5. The complexity bound in Corollary [3.4](#page-5-1) is $\mathcal{O}(\frac{\sqrt{d}}{\epsilon})$ with an logarithmic constant for Gaussian tail target. In [\[13\]](#page-11-1) Proposition 8 shows under standard Gaussian distribution, such complexity bound is optimal.

Remark 3.6. In our main Theorem [3.3,](#page-5-0) the Assumption [3](#page-5-3) can be relaxed to the following,

$$
\sum_{k=0}^{N-1} \frac{t_{k+1} - t_k}{T} \sqrt{\mathbb{E}_{\overrightarrow{P}_{T-t_k}} |s(T - t_k, x) - s_\theta(T - t_k, x)|^2} \le \epsilon.
$$
 (33)

We will list the resulting main Theorem [3.3](#page-5-0) in Appendix [A.4](#page-18-0) where complexity bound with respect to dimension (or variance) does not change by switching the Assumption [3](#page-5-3) with [\(33\)](#page-6-4). The complexity bounds for the subsequent applications are direct consequences.

Remark 3.7. We can replace the Assumption [4](#page-5-5) by the following,

$$
\forall t > 0, \quad \|\nabla \tilde{s}_{\theta}(t, \cdot)\|_{\infty} \le \frac{1}{2}.
$$
\n(34)

The complexity bound turns to,

$$
N = \mathcal{O}\left(\frac{\sqrt{M_0}}{\epsilon_0} (\log \frac{M_2 + Tr(C)}{\epsilon_0^2})^2\right),\tag{35}
$$

Detail discussion and proofs, see Appendix [A.5.](#page-19-0)

3.3 Convergence under manifold assumption

To analyze the Lipschitz bound of the score under manifold assumption, we first need the following theorem:

Theorem 3.8. Let $q_{\sigma} = \exp(-\frac{|x|^2}{2\sigma^2}) * q_0$, where q_0 is a distribution with compact support, Diam(Supp q_0) \leq R. Consider $g(x) = \log p_{\sigma}(x) + \frac{|x|^2}{2\sigma^2}$, then

$$
|\nabla g|_{\infty} \le \frac{R}{\sigma^2}, \quad \|\nabla^2 g\|_{\infty} \le \frac{2R^2}{\sigma^4}.
$$
\n(36)

Proof see Appendix [A.6.](#page-20-0)

Remark 3.9. Similar estimates in Theorem [3.8](#page-6-0) can be found in [\[9,](#page-10-6) [21\]](#page-11-13). In the current form [\(36\)](#page-6-5), we extract the spatial growing part to ensure uniform boundedness in space to be consistent with Assumption [2](#page-3-0) .

Applying Theorem [3.8](#page-6-0) to the forward OU-process [\(1\)](#page-2-0) at stopping time δ , then σ^2 corresponds to $1-e^{-\delta}$, $q_0(x)$ corresponds to $e^{\frac{\delta}{2}}p_0(e^{\frac{\delta}{2}}x)$. Then by Corollary [3.2,](#page-5-2) we get the following Lipschitz bound of the score under manifold assumption.

Corollary 3.10. Suppose p_0 is a distribution with compact support, $Diam(Supp_0) \leq R$, take $C = I_d$, $A = (1 - e^{-\delta})I_d$, then the corresponding $\overline{A}_t^{-1} = (1 - e^{-t})I_d$ and for $\forall t \ge \delta$, we have

$$
\|\nabla^2 \log p_t(x) + \bar{A}_t^{-1}\|_{\infty} \le 3(\frac{R}{1 - e^{-\delta}})^2 e^{-t}
$$
\n(37)

$$
|\nabla \log p_t(x) + \bar{A}_t^{-1} x|_{\infty} \le \frac{R}{1 - e^{-\delta}} e^{-\frac{t}{2}}.
$$
\n(38)

Proof see Appendix [A.2.](#page-12-2) With the Lipschitz bound in Corollary [3.10,](#page-6-6) we can get a Wasserstein-2distance bound as in Theorem [3.3:](#page-5-0)

Theorem 3.11. Take $C = I_d$. Suppose p_0 is a distribution with compact support, $Diam(Supp p_0) \leq R$, $\delta \leq 1$ and that Assumption [3](#page-5-3) holds, then we have

$$
\mathcal{W}_2(\vec{P}_\delta, \vec{Q}_{T-\delta})
$$

\n
$$
\leq \exp\left(3\frac{R^2}{\delta^2} + \frac{1}{\delta}\right) \left(e^{-T+\frac{\delta}{2}}\sqrt{M_2 + d}\right)
$$

\n
$$
+ 2(\tilde{K}_3\sqrt{M_0}h + \epsilon).
$$
\n(39)

where $\tilde{K}_3 = \mathcal{O}(\frac{R^2}{\delta^2})$ $\frac{R^2}{\delta^2}$), M_2 is the finite second moment of \overrightarrow{X}_0 , notice that $M_2 \leq M^2$.

Proof see Appendix [A.7.](#page-21-0)

With the result in Theorem [3.11,](#page-7-2) we can get the complexity bound with early stopping setting under manifold assumption by directly computation:

Corollary 3.12. To reach a distribution $\overleftarrow{Q}_{T-\delta}$ such that $\mathcal{W}_2(\overrightarrow{P}_\delta, \overleftarrow{Q}_{T-\delta}) = O(\epsilon_0)$ requires:

$$
T = \mathcal{O}\left(\frac{R^2}{\delta^2} + \log \frac{M_2 + d}{\epsilon_0^2}\right),\tag{40}
$$

$$
N = \frac{T - \delta}{h} = \mathcal{O}\left(\frac{R^2 \sqrt{M_2 \vee d}T}{\delta^2 \epsilon_0} \exp\left(\frac{1}{\delta} + 3\frac{R^2}{\delta^2}\right)\right),\tag{41}
$$

and Assumption [3](#page-5-3) to hold with

$$
\epsilon = \mathcal{O}(\epsilon_0 \exp(-\frac{1}{\delta} - 3\frac{R^2}{\delta^2})).
$$
\n(42)

With noticing that,

$$
\mathcal{W}_2(\vec{P}_{\delta}, \vec{P}_0) \le \sqrt{\mathbb{E}|\vec{X}_{\delta} - \vec{X}_0|^2} \le 2\sqrt{M_0\delta},
$$

we have the following complexity bound with respect to \overrightarrow{P}_0 .

Corollary 3.13. Some assumption as Corollary [3.12,](#page-7-1) to reach a distribution $\overleftarrow{Q}_{T-\delta}$ such that $\mathcal{W}_2(\overrightarrow{P}_0, \overrightarrow{Q}_{T-\delta}) =$ $\mathcal{O}(\epsilon_0)$ requires:

$$
\delta = \mathcal{O}\left(\frac{\epsilon_0^2}{M_0}\right), \quad \log N = \mathcal{O}\left(\frac{R^2 M_0^2}{\epsilon_0^4}\right). \tag{43}
$$

Furthermore, if one consider the dimension dependence for the radius of support under l_2 metric, R , and the second order moment, M_2 , we have the following calculations.

Remark 3.14. In practical application such as image processing, the images are encoded on a finite range. Then the support of p_0 is an hypercube, e.g., Supp $p_0 \n\t\subseteq [0, 255]^d$ which implies a natural assumption,

$$
R^2 = \mathcal{O}(d).
$$

Since $M_2 \leq R^2$, we also have $M_2 = \mathcal{O}(d)$. Then the complexity bound to reach $\mathcal{W}_2(\vec{P}_0, \vec{Q}_{T-\delta}) = \mathcal{O}(\epsilon_0)$ turns to,

$$
T = \mathcal{O}\left(\frac{d^3}{\epsilon_0^4}\right), N = \mathcal{O}\left(\frac{d^{\frac{13}{2}}}{\epsilon_0^9} \exp\left(\frac{1}{\delta} + 3\frac{d}{\delta^2}\right)\right),\tag{44}
$$

then,
$$
\log N = \mathcal{O}(\frac{d}{\delta^2}) = \mathcal{O}(\frac{d^3}{\epsilon_0^4}).
$$
 (45)

3.4 Convergence in the Bayesian Inverse problems

Another important application of generative models is to generative the posterior distribution in the Bayesian inverse problems. See [\[26\]](#page-11-5) for a detailed review. Here we restrict our theories to the following type of applicative scenario where the target distribution p_0 is the posterior distribution. More precisely,

$$
p_0(x) = D_0 \exp\left(-\frac{|x|_C^2}{2}\right) \exp\left(-\frac{|G(x) - y|_{\Sigma}^2}{2}\right),\tag{46}
$$

where D_0 is some constant, C and Σ denotes the covariance matrices of the Gaussian type prior distribution and observation noise distribution. G is a non-linear operator in $C_b^2(R^d, R^m)$ and $y \in R^m$ is an observation. During training, we adopt Gaussian type prior distribution with covariance C in (46) as the invariant measure of forward process [\(1\)](#page-2-0). A conditioned score is trained with the following loss,

$$
\mathbb{E}_{(x,y)\sim p_t} |s_\theta(t,x,y) - C\nabla_x \log p_t(x,y)|^2.
$$
 (47)

where p_t is then the joint distribution of (X_t, Y) in which Y follows $G(X_0) + \mathcal{N}(I, \Sigma)$. For the generation process of posterior distribution with observation y, we assume the following score approximation error.

Assumption 5. Fixing observation y, For each time discretization point t_k ,

$$
E_{\vec{P}_{T-t_k,y}} |s(T-t_k,x,y) - s_{\theta}(T-t_k,x,y)|^2 \le \epsilon^2,
$$

We then have the following theorem.

Theorem 3.15. Suppose X_0 has second moment $M_2 < \infty$, then using uniform discretization points in [\(6\)](#page-3-1) yields

$$
\mathcal{W}_2(\vec{P}_0(\cdot, y), \overleftarrow{Q}_T(\cdot, y))
$$

\n
$$
\leq \exp(\tilde{K}_4) \Big(e^{-T} \sqrt{M_2 + \text{Tr}(C)} \tag{48}
$$

$$
+2(\tilde{K}_5\sqrt{M_0}\frac{T}{N}+\epsilon)\Big). \t\t(49)
$$

where \tilde{K}_4 and \tilde{K}_5 are some dimensionless constant depending on $(||C||, ||\Sigma||, G, y)$, specified in [\(144\)](#page-22-0).

Proof see Appendix [A.8.](#page-21-1) Remark 3.16. With fixed \tilde{K}_4 and \tilde{K}_5 , for ϵ_0 accuracy in Wasserstein distance for [\(49\)](#page-8-3), one requires, $\epsilon = \mathcal{O}(\epsilon_0), T = \mathcal{O}(\log \frac{1}{\epsilon_0})$ $\sqrt{M_2 + \text{Tr}(C)}$ $\frac{\overline{P_{\epsilon_0}(\Gamma)}(C)}{\epsilon_0}$ and $N = \mathcal{O}(\frac{T\sqrt{M_0}}{\epsilon_0^2})$ $\frac{(\overline{M_0})}{\epsilon_0^2}).$

3.5 Migration to probabilistic flow ODE

The probabilistic flow ODE refers to Equation (13) in [\[25\]](#page-11-0). With C as the covariance matrix, the backward process is then,

$$
d\overleftarrow{X}_t = \frac{1}{2}(\overleftarrow{X}_t + C\nabla \log p_{T-t}(\overleftarrow{X}_t))dt, \quad \overleftarrow{X}_0 \sim \mathcal{N}(0, C). \tag{50}
$$

The Lipschitz bound (Corollary [3.2\)](#page-5-2) still holds since the forward process is OU. For the generation process, we revert to the usual Euler-Maruyama scheme:

$$
\overline{dY}_t = \frac{1}{2} (\overleftarrow{Y}_{t_k} + s_\theta (T - t_k, \overleftarrow{Y}_{t_k})) dt, \quad t \in [t_k, t_{k+1}]. \tag{51}
$$

which starts from Gaussian distribution $\overleftarrow{Y}_0 \sim \mathcal{N}(0, C)$. With the Lipschitz bound, our method still work with the ODE flow since we don't rely on Girsanov type theorem in the proof. And we can get the following Wasserstein-2-distance bound with linear complexity.

Theorem 3.17. Suppose Assumption [1,](#page-3-4) [2,](#page-3-0) [3,](#page-5-3) [4,](#page-5-5) hold, using the Euler-Maruyama to the ode flow with uniform step size $h = t_{k+1} - t_k \leq 1$ yields

$$
\mathcal{W}_2(\vec{P}_0, \overleftarrow{Q}_T) \le \exp(\frac{\tilde{K}_2}{2})
$$

$$
\cdot \left(\exp(-\frac{T}{2})(M_2 + \text{Tr}(C)) + \tilde{K}_6 \sqrt{\frac{T^3 M_0}{N}} + \frac{1}{2} \epsilon T\right).
$$
\n(52)

where \tilde{K}_1 and M_0 are the same constants as in Theorem [3.3,](#page-5-0) \tilde{K}_6 will be specified in the proof.

Proof see Appendix [A.9.](#page-22-1) Complexity bounds are accessed with similar procedure and listed below.

• In $C²$ distribution with Gaussian tail, same assumption as Corollary [3.4,](#page-5-1)

$$
N = \mathcal{O}\Big(\frac{M_0}{\epsilon_0^2} (\log \frac{M_2 + Tr(C)}{\epsilon_0^2})^3\Big).
$$

• Under manifold hypothesis with early stopping regime, same assumption as Corollary [3.13,](#page-7-0)

$$
\log N = \mathcal{O}(\frac{R^2M_0^2}{\epsilon_0^4}).
$$

4 Proof Sketch of Theorem [3.3](#page-5-0)

Basic idea We define two strong SDE solutions: \overleftarrow{X}_t which solves [\(2\)](#page-2-1) starting from \overrightarrow{P}_T and and \overleftarrow{Y}_t which solves [\(6\)](#page-3-1) starting from $\mathcal{N}(0, C)$. We couple them by starting from Wasserstein-2 optimally coupled initial conditions and using the same Brownian motion. Running them to time $T - \delta$, we get $\overleftarrow{X}_{T-\delta} \sim \overrightarrow{P}_{\delta}$ and $\overleftarrow{V}_{T-\delta} \sim \overrightarrow{Q}_{T-\delta}$. This is a coupling between $\overrightarrow{P}_{\delta}$ and $\overleftarrow{Q}_{T-\delta}$, by definition, $(\mathbb{E}|X_{$ bounds the Wasserstein-2 distance $\mathcal{W}_2(\overrightarrow{P}_\delta, \overleftarrow{Q}_{T-\delta})$.

Detailed computation Our computation consists of four steps: In step 1, we use Itô's formula and Cauchy's inequality to get

$$
\frac{d\sqrt{e^t \mathbb{E}|\tilde{X}_t - \tilde{Y}_t|^2}}{dt}
$$
\n
$$
\leq e^{\frac{t}{2}} (\mathbb{E}|\tilde{s}(T - t, \tilde{X}_t))
$$
\n
$$
+ (I - C\bar{A}_{T-t}^{-1})\tilde{\tilde{X}}_t - \tilde{s}_{\theta}(T - t_k, \tilde{Y}_{t_k})
$$
\n
$$
- (I - C\bar{A}_{T-t_k}^{-1})\tilde{Y}_{t_k}|^2)^{\frac{1}{2}}, \quad t \in [t_k, t_{k+1}].
$$
\n(53)

In step 2, by significant technical work we get an approximation for the RHS of [\(53\)](#page-9-0):

$$
RHS \ of (53) \le K_1 \sqrt{M_0 (1 - e^{-(t - t_k)})} + \epsilon + K_2 e^{-T + t_k} \sqrt{\mathbb{E} |\overleftarrow{X}_{t_k} - \overleftarrow{Y}_{t_k}|^2}.
$$
 (54)

where K_1, K_2 are some constants.

In step 3, by performing integration over time on [\(54\)](#page-9-1), we get an Iterative formula:

$$
\sqrt{e^{t_{k+1}}\mathbb{E}|\overleftarrow{X}_{t_{k+1}} - \overleftarrow{Y}_{t_{k+1}}|^2} \exp(-K_2 e^{-T+t_{k+1}})
$$
\n
$$
\leq \sqrt{e^{t_k}\mathbb{E}|\overleftarrow{X}_{t_k} - \overleftarrow{Y}_{t_k}|^2} \exp(-K_2 e^{-T+t_k})
$$
\n
$$
+ 2(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon)(e^{\frac{t_{k+1}}{2}} - e^{\frac{t_k}{2}}),
$$
\n(55)

which leads to

$$
\sqrt{\mathbb{E}|X_{T-\delta} - Y_{T-\delta}|^2} \leq e^{K_2} (e^{-\frac{T-\delta}{2}} \sqrt{\mathbb{E}|X_0 - Y_0|^2} + 2(\frac{8}{3}K_1 \sqrt{M_0} h + \epsilon)).
$$
\n(56)

In step 4, by choosing Wasserstein-2 optimally coupled initial conditions and using the fact (see Lemma [A.3\)](#page-23-1):

$$
\mathcal{W}_2^2(\overrightarrow{P}_{T-\delta}, \gamma_C) \leq e^{-(T-\delta)} \mathcal{W}_2^2(\overrightarrow{P}_0, \gamma_C)
$$

$$
\leq e^{-(T-\delta)} (M_2 + \text{Tr}(C)),
$$

we get the bound [\(29\)](#page-5-6) in Theorem [3.3.](#page-5-0)

Limitation

We sketched the limitations of our paper below and for detail discussion please refer to the linked context. We will leave these problems as future directions.

- The paper only analyzed the case when forward process is OU. For general cases when non-linear drifts are involved in the forward process, our analysis framework may not be easily generalized due to lack of explicit expression for solution of forward process.
- An important application of the infinite dimension generative modeling is the Bayesian approach to inverse problems in the functional space. At the moment, our assumption of trace class base distribution with a bounded perturbation can only be derived from the C_b^2 bounded observation operator. See discussion in Section [3.4.](#page-8-0)
- During derivation we have made the standard assumption on the score approximation Assumption [3.](#page-5-3) While this assumptions are not easy to be derived from the operator approximation theories or to be validated through the numerical experiment. Comparing with KL bounds, we also introduce Assumption [4](#page-5-5) to regulate the backward process. The assumption can released as discussed in Remark [3.7](#page-6-3) and logarithmic constant of complexity bound changes.
- Due to the limitation of unknown coefficients in the derivation as well as it is only an upper bound result, our analysis in Corollary [3.12](#page-7-1) cannot give the optimal stopping time for the bounded support case in practical application. Also we restrict ourselves to uniform time discretization. The complexity bound may be improved by adapting the discretization with a better estimate of Lipschitz of score.

Broader Impact

The paper provides insights over the regularity and convergence of one of the state-of-the-art generative models, which the GenAI community may benefit from. There are many potential societal consequences of our work, none which we feel must be specifically highlighted here.

References

- [1] B. Anderson. Reverse-time diffusion equation models. Stoch. Process. Appl., 12(3):313–326, 1982.
- [2] Yoshua Bengio, Ian Goodfellow, and Aaron Courville. Deep learning, volume 1. MIT press Cambridge, MA, USA, 2017.
- [3] Joe Benton, Valentin De Bortoli, Arnaud Doucet, and George Deligiannidis. Linear convergence bounds for diffusion models via stochastic localization. arXiv preprint arXiv:2308.03686, 2023.
- [4] Jonas Blessing and Michael Kupper. Viscous hamilton–jacobi equations in exponential orlicz hearts. Journal de Mathématiques Pures et Appliquées, 163:654–672, 2022.
- [5] L. Caffarelli. Monotonicity properties of optimal transportation¶ and the fkg and related inequalities. Communications in Mathematical Physics, 214:547–563, 2000.
- [6] Hongrui Chen, Holden Lee, and Jianfeng Lu. Improved analysis of score-based generative modeling: User-friendly bounds under minimal smoothness assumptions, 2023.
- [7] Giovanni Conforti, Alain Durmus, and Marta Gentiloni Silveri. Kl convergence guarantees for score diffusion models under minimal data assumptions, 2024.
- [8] Giuseppe Da Prato and Jerzy Zabczyk. Stochastic equations in infinite dimensions, volume 152. Cambridge university press, 2014.
- [9] Valentin De Bortoli. Convergence of denoising diffusion models under the manifold hypothesis. Transactions on Machine Learning Research, 2022.
- [10] Valentin De Bortoli. Convergence of denoising diffusion models under the manifold hypothesis, May 2023. arXiv:2208.05314 [cs, math, stat].
- [11] Giulio Franzese, Giulio Corallo, Simone Rossi, Markus Heinonen, Maurizio Filippone, and Pietro Michiardi. Continuous-time functional diffusion processes. Advances in Neural Information Processing Systems, 36, 2024.
- [12] Yasuhiro Fujita, Hitoshi Ishii, and Paola Loreti. Asymptotic solutions of viscous hamilton–jacobi equations with ornstein–uhlenbeck operator. Communications in Partial Differential Equations, 31(6):827–848, 2006.
- [13] Xuefeng Gao, Hoang M. Nguyen, and Lingjiong Zhu. Wasserstein convergence guarantees for a general class of score-based generative models, 2023.
- [14] Xuefeng Gao and Lingjiong Zhu. Convergence analysis for general probability flow odes of diffusion models in wasserstein distances, 2024.
- [15] Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. Advances in neural information processing systems, 33:6840–6851, 2020.
- [16] Gavin Kerrigan, Justin Ley, and Padhraic Smyth. Diffusion Generative Models in Infinite Dimensions, February 2023. arXiv:2212.00886 [cs, stat].
- [17] Peter E Kloeden and Eckhard Platen. Numerical Solution of Stochastic Differential Equations. Springer, 1992.
- [18] Jae Hyun Lim, Nikola B. Kovachki, Ricardo Baptista, Christopher Beckham, Kamyar Azizzadenesheli, Jean Kossaifi, Vikram Voleti, Jiaming Song, Karsten Kreis, Jan Kautz, Christopher Pal, Arash Vahdat, and Anima Anandkumar. Score-based Diffusion Models in Function Space, November 2023. arXiv:2302.07400 [cs, math, stat].
- [19] Yubin Lu, Zhongjian Wang, and Guillaume Bal. Mathematical analysis of singularities in the diffusion model under the submanifold assumption, 2023.
- [20] Dan Mikulincer and Yair Shenfeld. The brownian transport map. Probability Theory and Related Fields, 2024.
- [21] Connor Mooney, Zhongjian Wang, Jack Xin, and Yifeng Yu. Global well-posedness and convergence analysis of score-based generative models via sharp lipschitz estimates. $arXiv$ preprint arXiv:2405.16104, 2024.
- [22] Jakiw Pidstrigach, Youssef Marzouk, Sebastian Reich, and Sven Wang. Infinite-Dimensional Diffusion Models, October 2023. arXiv:2302.10130 [cs, math, stat].
- [23] Jakiw Pidstrigach, Youssef Marzouk, Sebastian Reich, and Sven Wang. Infinite-dimensional diffusion models. arXiv preprint arXiv:2302.10130, 2023.
- [24] Emile Pierret and Bruno Galerne. Diffusion models for gaussian distributions: Exact solutions and wasserstein errors, 2024.
- [25] Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. arXiv preprint arXiv:2011.13456, 2020.
- [26] Andrew M Stuart. Inverse problems: a bayesian perspective. Acta numerica, 19:451–559, 2010.
- [27] Joshua B Tenenbaum, Vin de Silva, and John C Langford. A global geometric framework for nonlinear dimensionality reduction. science, 290(5500):2319–2323, 2000.
- [28] Cédric Villani. *Topics in optimal transportation*, volume 58. American Mathematical Soc., 2021.
- [29] Cédric Villani et al. *Optimal transport: old and new*, volume 338. Springer, 2009.

The appendix consists of two parts. In Section [A,](#page-12-0) we present all the detailed proofs. In Section [B,](#page-23-2) we discuss generalizing our theories to infinite dimensions.

A Proofs of Theorems

Here we present detail proofs.

A.1 Proof of Theorem [3.1](#page-4-3)

Proof. Consider the solution of [\(19\)](#page-4-4) given by [\(20\)](#page-4-5):

$$
\tilde{q}(t,x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \frac{1}{\sqrt{\det B(t)}} \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy, \quad (t,x) \in (0,\infty) \times \mathbb{R}^n.
$$

So $\bar{q}(t, x) = -\log \tilde{q}(t, x)$ satisfies

$$
\nabla_x \bar{q}(t,x) = -\frac{\nabla_x \tilde{q}(t,x)}{\tilde{q}(t,x)} = -\frac{\int_{\mathbb{R}^n} (\nabla_x \exp(\frac{-|x-y|_{B(t)}^2}{2})) \exp(-\bar{h}(y)) dy}{\int_{\mathbb{R}^n} \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy}
$$

$$
= \frac{\int_{\mathbb{R}^n} (\nabla_y \exp(\frac{-|x-y|_{B(t)}^2}{2})) \exp(-\bar{h}(y)) dy}{\int_{\mathbb{R}^n} \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy}
$$

$$
= \frac{\int_{\mathbb{R}^n} (\nabla_y \bar{h}(y)) \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy}{\int_{\mathbb{R}^n} \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy}.
$$

Here the third equal sign is derived from integration by part.

Since $\exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) \ge 0$, taking absolute value we get $|\nabla_x \bar{q}(t,x)| \le |\nabla h|_{\infty}$, thus

$$
|\nabla \bar{q}(t,\cdot)|_{\infty} \leq |\nabla \bar{h}|_{\infty}.
$$

For any unit direction z, taking derivative twice along that direction and using the same method as above we get :

$$
\nabla_z \cdot \nabla_z \bar{q}(t,x) = \frac{\int_{\mathbb{R}^n} (\nabla_z^2 \bar{h} - (\nabla_z \bar{h}^2))(y) \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy}{\int_{\mathbb{R}^n} \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy} + (\frac{\int_{\mathbb{R}^n} (\nabla_y \bar{h}(y)) \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy}{\int_{\mathbb{R}^n} \exp(\frac{-|x-y|_{B(t)}^2}{2}) \exp(-\bar{h}(y)) dy})^2.
$$

Taking absolute value again we get

$$
|\nabla_z \cdot \nabla_z \bar{q}(t,x)| \leq |\nabla_z^2 \bar{h}|_{\infty} + |\nabla_z \bar{h}|_{\infty}^2,
$$
\n(57)

thus

$$
|\nabla^2 \bar{q}(t,\cdot)\|_{\infty} \leq \|\nabla^2 \bar{h}\|_{\infty} + |\nabla \bar{h}|_{\infty}^2.
$$

 \Box

A.2 Proof of Corollary [3.2](#page-5-2) and Corollary [3.10](#page-6-6)

 $\overline{}$

Proof. By the construction of \bar{q} [\(16\)](#page-4-6), we have

$$
K(t)\sqrt{C^{-1}}\nabla\bar{q}(t,x) = \nabla q(t, K(t)^{-1}\sqrt{C}x),
$$

\n
$$
K(t)^{2}C^{-1}\nabla^{2}\bar{q}(t,x) = \nabla^{2}q(t, K(t)^{-1}\sqrt{C}x).
$$
\n(58)

Notice that

$$
\tilde{s}(t,x) = C\nabla q(t,x),
$$

\n
$$
\nabla \tilde{s}(t,x) = C\nabla^2 q(t,x).
$$
\n(59)

We have

$$
|\tilde{s}(t,\cdot)|_{\infty} = |C\nabla q(t,\cdot)|_{\infty} = |K(t)\sqrt{C}\nabla \bar{q}(t,\cdot)|_{\infty} \le e^{-\frac{t}{2}} \|e^{\frac{t}{2}}K(t)\| \|\sqrt{C}\| |\nabla \bar{q}(t,\cdot)|_{\infty},
$$

$$
\|\nabla \tilde{s}(t,\cdot)\|_{\infty} = \|C\nabla^2 q(t,\cdot)\|_{\infty} = \|K(t)^2 \nabla^2 \bar{q}(t,\cdot)\|_{\infty} \le e^{-t} \|e^{\frac{t}{2}}K(t)\|^2 \|\nabla^2 \bar{q}(t,\cdot)\|_{\infty}.
$$
 (60)

Let $K = \sup_{t\geq 0} ||e^{\frac{t}{2}}K(t)|| = \sup_{t\geq 0} ||A\overline{A}_t^{-1}||$, then with Theorem [3.1](#page-4-3) and the initial value $\overline{q}(0, x) =$ $h(\sqrt{C}x)$, we get

$$
|\tilde{s}(t,\cdot)| \leq e^{-\frac{t}{2}} K ||C||^{\frac{1}{2}} |\sqrt{C} \nabla h|_{\infty},
$$

$$
||\nabla \tilde{s}(t,\cdot)||_{\infty} \leq e^{-t} K^2 (||C \nabla h||_{\infty} + |\sqrt{C} \nabla h|_{\infty}^2).
$$
 (61)

Let

$$
L_0 = K^2(||C\nabla h||_{\infty} + |\sqrt{C}\nabla h|_{\infty}^2),
$$

\n
$$
L_1 = K||C||^{\frac{1}{2}}|\sqrt{C}\nabla h|_{\infty},
$$
\n(62)

then we get the result in Corollary [3.2.](#page-5-2)

In Corollary [3.10,](#page-6-6) we take

$$
C = I_d,
$$

\n
$$
A = (1 - e^{-\delta})I_d,
$$

\n
$$
\bar{A}_t = (1 - e^{-t})^{-1}I_d,
$$
\n(63)

and

$$
\vec{P}_{\delta} = \exp\left(-\frac{|x|^2}{2(1 - e^{-\delta})}\right) * p_{\delta},\tag{64}
$$

where $p_{\delta}(x) := e^{\frac{\delta}{2}} p_0(e^{\frac{\delta}{2}}x)$ satisfies $\text{Diam}(\text{Supp }p_{\delta}) \leq e^{-\frac{\delta}{2}}R$. Then by Theorem [3.8,](#page-6-0) the corresponding $h(x) := \log P_{\delta}(x) + \frac{|x|^2}{2(1-e^{\delta})}$ $rac{|x|}{2(1-e^{\delta})}$ has

$$
|\nabla h|_{\infty} \le \frac{Re^{-\frac{\delta}{2}}}{1 - e^{-\delta}}, \quad \|\nabla^2 h\| \le 2 \frac{R^2 e^{-\delta}}{(1 - e^{-\delta})^2}.
$$
\n(65)

The constants then can be computed directly:

$$
K = \sup_{t \ge \delta} \|A\bar{A}_t^{-1}\| = 1,
$$

\n
$$
L_0 = K^2 (\|\nabla h\|_{\infty} + |\nabla h|_{\infty}^2) \le 3 \frac{R^2 e^{-\delta}}{(1 - e^{-\delta})^2},
$$

\n
$$
L_1 = K |\nabla h|_{\infty} \le \frac{Re^{-\frac{\delta}{2}}}{1 - e^{-\delta}}.
$$
\n(66)

Then by Corollary [3.2,](#page-5-2) for $\forall t \geq \delta$, we have

$$
\|\nabla^2 \log \vec{P}_t(x) + \vec{A}_t^{-1}\|_{\infty} \le L_0 e^{-(t-\delta)} = 3 \frac{R^2}{(1 - e^{-\delta})^2} e^{-t}
$$

$$
|\nabla^2 \log \vec{P}_t(x) + \vec{A}_t^{-1}x|_{\infty} \le L_1 e^{-\frac{t-\delta}{2}} = \frac{R}{1 - e^{-\delta}} e^{-\frac{t}{2}}
$$

A.3 Proof of Theorem [3.3](#page-5-0)

Proof. By Corollary [3.2,](#page-5-2) we have $\forall t \geq 0$,

$$
\|\nabla \tilde{s}(t, \cdot)\|_{\infty} \le L_0 e^{-t},
$$

$$
|\tilde{s}(t, \cdot)|_{\infty} \le L_1 e^{-\frac{t}{2}},
$$
\n(68)

where we recall,

$$
\tilde{s}(t,x) = s(t,x) + C\bar{A}_t^{-1}x,\tag{69}
$$

$$
\tilde{s}_{\theta}(t,x) = s_{\theta}(t,x) + C\bar{A}_t^{-1}x,\tag{70}
$$

$$
K = \sup_{t \ge 0} \|A\bar{A}_t^{-1}\|,\tag{71}
$$

$$
L_0 = K^2(||C\nabla^2 h||_{\infty} + |\sqrt{C}\nabla h|_{\infty}^2),\tag{72}
$$

$$
L_1 = K \|C\|^{\frac{1}{2}} |\sqrt{C} \nabla h|_{\infty}.
$$
\n
$$
(73)
$$

Let
\n
$$
L_2 = \sup_{t \ge 0} ||e^t (I - C\bar{A}_t^{-1})|| = \sup_{t \ge 0} ||(A - C)(Ae^{-t} + C(1 - e^{-t}))^{-1}||,
$$
\n
$$
L_3 = \sup_{t \ge s \ge 0} ||(e^{-t} - e^{-s})^{-1}C(\bar{A}_s^{-1} - \bar{A}_t^{-1})|| = \sup_{t \ge s \ge 0} ||(A - C)C(Ae^{-t} + C(1 - e^{-t}))^{-1}(Ae^{-s} + C(1 - e^{-s}))^{-1}||.
$$
\n(74)

More precisely,

$$
L_2 \le \max\{\|I - CA^{-1}\|, \|AC^{-1} - I\|\},
$$

\n
$$
L_3 \le L_2 \max\{1, \|CA^{-1}\|\}.
$$
\n(75)

Then by [\(74\)](#page-14-0), we have

$$
||I - C\bar{A}_t^{-1}|| \le L_2 e^{-t},
$$

\n
$$
||C\bar{A}_t^{-1}|| \le ||I|| + ||I - C\bar{A}_t^{-1}|| \le 1 + L_2, \quad \forall t \ge 0,
$$

\n
$$
||C\bar{A}_s^{-1} - C\bar{A}_t^{-1}|| \le L_3(e^{-t} - e^{-s}), \quad \forall 0 \le t \le s.
$$
\n(76)

Let $M_0 = \max\{\text{Tr}(C), M_2\}$, then for $\forall t \geq 0$

$$
\mathbb{E}|\vec{X}_t|^2 = \mathbb{E}|\vec{X}_t - e^{-\frac{t}{2}}\vec{X}_0|^2 + \mathbb{E}|e^{-\frac{t}{2}}\vec{X}_0|^2
$$

= $(1 - e^{-t})\text{Tr}(C) + e^{-t}M_2$
 $\leq M_0.$ (77)

For any $0 \le t \le s \le T$,

$$
\mathbb{E}|\overrightarrow{X}_t - \overrightarrow{X}_s|^2 = \mathbb{E}|\overrightarrow{X}_s - e^{-\frac{s-t}{2}}\overrightarrow{X}_t|^2 + (1 - e^{-\frac{s-t}{2}})^2 \mathbb{E}|\overrightarrow{X}_t|^2
$$

\n
$$
\leq (1 - e^{-s+t})\text{Tr}(C) + (1 - e^{-\frac{s-t}{2}})^2 M_0
$$

\n
$$
\leq 2M_0(s-t).
$$
\n(78)

For simplicity of notation we take $\delta = 0$ (no early stopping), while the case $\delta > 0$ can be handle with the same method and we will present the result at the end of the proof.

Step 1. Coupling two solutions of \overleftarrow{X}_t and \overleftarrow{Y}_t for the same Brownian motion. By Itô's formula, we have $for t \in [t_k, t_{k+1}]$

$$
\frac{d|\overleftarrow{X}_t - \overleftarrow{Y}_t|^2}{dt} = -|\overleftarrow{X}_t - \overleftarrow{Y}_t|^2
$$
\n
$$
+ 2\langle \overleftarrow{X}_t - \overleftarrow{Y}_t, \tilde{s}(T - t, \overleftarrow{X}_t) + (I - C\overleftarrow{A}_{T-t}^{-1})\overleftarrow{X}_t - \tilde{s}_\theta(T - t_k, \overleftarrow{Y}_{t_k}) - (I - C\overleftarrow{A}_{T-t_k})\overleftarrow{Y}_{t_k}\rangle.
$$

Take expectation and use Cauchy's inequality we get:

$$
\frac{d\mathbb{E}|\overleftarrow{X}_t - \overleftarrow{Y}_t|^2}{dt} \le -\mathbb{E}|\overleftarrow{X}_t - \overleftarrow{Y}_t|^2
$$
\n
$$
+ 2\mathbb{E}\langle \overleftarrow{X}_t - \overleftarrow{Y}_t, \tilde{s}(T - t, \overleftarrow{X}_t) + (I - C\overleftarrow{A}_{T-t}^{-1})\overleftarrow{X}_t - \tilde{s}_\theta(T - t_k, \overleftarrow{Y}_{t_k}) - (I - C\overleftarrow{A}_{T-t_k})\overleftarrow{Y}_{t_k}\rangle
$$
\n
$$
\le -\mathbb{E}|\overleftarrow{X}_t - \overleftarrow{Y}_t|^2
$$
\n
$$
+ 2\sqrt{(\mathbb{E}|\overleftarrow{X}_t - \overleftarrow{Y}_t|^2)(\mathbb{E}|\tilde{s}(T - t, \overleftarrow{X}_t) + (I - C\overleftarrow{A}_{T-t}^{-1})\overleftarrow{X}_t - \tilde{s}_\theta(T - t_k, \overleftarrow{Y}_{t_k}) - (I - C\overleftarrow{A}_{T-t_k})\overleftarrow{Y}_{t_k}|^2).
$$

Denote $E(t) = \sqrt{e^t \mathbb{E} |\dot{X}_t - \dot{Y}_t|^2}$, then we have

$$
\frac{dE(t)}{dt} \le \sqrt{e^t(\mathbb{E}|\tilde{s}(T-t,\tilde{X}_t) + (I - C\bar{A}_{T-t}^{-1})\tilde{X}_t - \tilde{s}_\theta(T-t_k,\tilde{Y}_{t_k}) - (I - C\bar{A}_{T-t_k}^{-1})\tilde{Y}_{t_k}|^2)}.
$$
\n(79)

Step 2. By mean inequality, RHS of (79) can be bounded by

$$
\sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_t) + (I - C\bar{A}_{T-t}^{-1})\tilde{X}_t - \tilde{s}_{\theta}(T-t_k,\tilde{Y}_{t_k}) - (I - C\bar{A}_{T-t_k}^{-1})\tilde{Y}_{t_k}|^2}
$$
\n
$$
\leq \sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_t) - \tilde{s}_{\theta}(T-t_k,\tilde{Y}_{t_k})|^2} + \sqrt{\mathbb{E}[(I - C\bar{A}_{T-t_k}^{-1}\tilde{X}_t - (I - C\bar{A}_{T-t_k})\tilde{Y}_{t_k}]^2}.
$$
\n(80)

The first term of [\(80\)](#page-14-2):

$$
\sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_t) - \tilde{s}_{\theta}(T-t,\tilde{Y}_{t_k})|^2}
$$
\n
$$
\leq \sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_t) - \tilde{s}(T-t_k,\tilde{X}_{t_k})|^2} + \sqrt{\mathbb{E}|\tilde{s}(T-t_k,\tilde{X}_{t_k}) - \tilde{s}_{\theta}(T-t_k,\tilde{X}_{t_k})|^2}
$$
\n
$$
+ \sqrt{\mathbb{E}|\tilde{s}_{\theta}(T-t_k,\tilde{X}_{t_k}) - \tilde{s}_{\theta}(T-t_k,\tilde{Y}_{t_k})|^2}
$$
\n
$$
=:I_1 + I_2 + I_3.
$$
\n(81)

To bound $I_1 = \sqrt{\mathbb{E}|\tilde{s}(T-t, \overleftarrow{X}_t) - \tilde{s}(T-t_k, \overleftarrow{X}_{t_k})|^2}$, we use the following lemma. **Lemma A.1** (Rmk 3.1 in [\[19\]](#page-11-15)). For $0 \le t < s \le T$, the forward process satisfies:

$$
\nabla \log \vec{P}_s(x_s) = e^{\frac{s-t}{2}} \mathbb{E}_{\vec{P}_{t|s}} [\nabla_y \log \vec{P}_t(y)|x_s]. \tag{82}
$$

Thus,

$$
\tilde{s}(s,\overrightarrow{X}_{s}) = e^{\frac{s-t}{2}} \mathbb{E}_{\overrightarrow{P}_{t|s}}[\tilde{s}(t,\overrightarrow{X}_{t}) - C\overrightarrow{A}(t)\overrightarrow{X}_{t}|\overrightarrow{X}_{s}] + C\overrightarrow{A}(s)\overrightarrow{X}_{s}.
$$
\n(83)

By lemma [A.1,](#page-15-0) we have

$$
I_{1} \leq \sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_{t}) - \tilde{s}(T-t,e^{\frac{t-t_{k}}{2}}\tilde{X}_{t_{k}})|^{2}} + \sqrt{\mathbb{E}|\tilde{s}(T-t,e^{\frac{t-t_{k}}{2}}\tilde{X}_{t_{k}}) - \tilde{s}(T-t_{k},\tilde{X}_{t_{k}})|^{2}}
$$

\n
$$
\leq \sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_{t}) - \tilde{s}(T-t,e^{\frac{t-t_{k}}{2}}\tilde{X}_{t_{k}})|^{2}}
$$

\n
$$
+ \sqrt{\mathbb{E}|\tilde{s}(T-t,e^{\frac{t-t_{k}}{2}}\tilde{X}_{t_{k}}) - e^{\frac{t-t_{k}}{2}}\tilde{s}(T-t,\tilde{X}_{t}) + e^{\frac{t-t_{k}}{2}}C\bar{A}(T-t)\tilde{X}_{t} - C\bar{A}(T-t_{k})\tilde{X}_{t_{k}}|^{2}}
$$

\n
$$
\leq 2\sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_{t}) - \tilde{s}(T-t,e^{\frac{t-t_{k}}{2}}\tilde{X}_{t_{k}})|^{2}}
$$

\n
$$
+ \sqrt{\mathbb{E}|(1-e^{\frac{t-t_{k}}{2}})\tilde{s}(T-t,\tilde{X}_{t}) + e^{\frac{t-t_{k}}{2}}C\bar{A}(T-t)\tilde{X}_{t} - C\bar{A}(T-t_{k})\tilde{X}_{t_{k}}|^{2}}
$$

\n
$$
\leq 2L_{0}e^{-(T-t)}\sqrt{\mathbb{E}|\tilde{X}_{t} - e^{\frac{t-t_{k}}{2}}\tilde{X}_{t_{k}}|^{2}} + \sqrt{\mathbb{E}|(1-e^{\frac{t-t_{k}}{2}})\tilde{s}(T-t,\tilde{X}_{t})|^{2}}
$$

\n
$$
+ \sqrt{\mathbb{E}|e^{\frac{t-t_{k}}{2}}C\bar{A}(T-t)\tilde{X}_{t} - C\bar{A}(T-t_{k})\tilde{X}_{t_{k}}|^{2}}.
$$

\n(84)

By the transition property of the forward OU-process and the fact $t - t_k \leq h \leq 1$, for the first term in [\(84\)](#page-15-1) we have,

$$
\mathbb{E}|\overleftarrow{X}_t - e^{\frac{t - t_k}{2}} \overleftarrow{X}_{t_k}|^2 = e^{t - t_k} \mathbb{E}|\overleftarrow{X}_{t_k} - e^{-\frac{(T - t_k) - (T - t)}{2}} \overleftarrow{X}_t|^2
$$
\n
$$
= e^{t - t_k} \mathbb{E}|\sqrt{1 - e^{-((T - t_k) - (T - t))}} z|^2 \quad z \sim \mathcal{N}(0, C)
$$
\n
$$
= e^{t - t_k} (1 - e^{t_k - t}) \text{Tr}(C)
$$
\n
$$
\leq 2(t - t_k) \text{Tr}(C).
$$
\n(85)

By [\(68\)](#page-13-1), the second term in [\(84\)](#page-15-1) is bounded by,

$$
\mathbb{E}|(1 - e^{\frac{t - t_k}{2}})\tilde{s}(T - t, \overleftarrow{X}_t)|^2 \le (1 - e^{\frac{t - t_k}{2}})^2 |\tilde{s}(T - t, \cdot)|_{\infty}^2
$$

\n
$$
\le (1 - e^{\frac{t - t_k}{2}})^2 L_1^2 e^{-(T - t)}
$$

\n
$$
\le e^{-(T - t)} L_1^2 (t - t_k)^2.
$$
\n(86)

By [\(68\)](#page-13-1), [\(76\)](#page-14-3),[\(77\)](#page-14-4),[\(78\)](#page-14-5), the third term in [\(84\)](#page-15-1) follows,

$$
\sqrt{\mathbb{E}|e^{\frac{t-t_{k}}{2}}C\bar{A}_{T-t}^{-1}\tilde{X}_{t}-C\bar{A}_{T-t_{k}}^{-1}\tilde{X}_{t_{k}}|^{2}}\n\leq\sqrt{\mathbb{E}|(1-e^{\frac{t-t_{k}}{2}})C\bar{A}_{T-t}^{-1}\tilde{X}_{t}|^{2}}+\sqrt{\mathbb{E}|C\bar{A}_{T-t}^{-1}\tilde{X}_{t}-C\bar{A}_{T-t_{k}}^{-1}\tilde{X}_{t_{k}}|^{2}}\n\leq\sqrt{\mathbb{E}|(1-e^{\frac{t-t_{k}}{2}})C\bar{A}_{T-t}^{-1}\tilde{X}_{t}|^{2}}+\sqrt{\mathbb{E}|C(\bar{A}_{T-t}^{-1}-\bar{A}_{T-t_{k}}^{-1})\tilde{X}_{t}|^{2}}\n+\sqrt{\mathbb{E}|C\bar{A}_{T-t_{k}}^{-1}(\tilde{X}_{t}-\tilde{X}_{t_{k}})|^{2}}\n\leq((e^{\frac{t-t_{k}}{2}}-1)\|C\bar{A}_{T-t}^{-1}\|+\|C(\bar{A}_{T-t}^{-1}-\bar{A}_{T-t_{k}}^{-1})\|)\sqrt{\mathbb{E}|\tilde{X}_{t}|^{2}}\n+\|C\bar{A}_{T-t_{k}}^{-1}\|\sqrt{\mathbb{E}|\tilde{X}_{t}-\tilde{X}_{t_{k}}|^{2}}\n\leq((e^{\frac{t-t_{k}}{2}}-1)(1+L_{2}))+L_{3}(e^{-T+t}-e^{-T+t_{k}}))\sqrt{M_{0}}+(1+L_{2})\sqrt{2M_{0}(t-t_{k})}\n\leq((1+L_{2})+2L_{3}e^{-T+t_{k}})(t-t_{k})\sqrt{M_{0}}+(1+L_{2})\sqrt{2M_{0}(t-t_{k})}.
$$
\n(1+L_{3})+2L_{4}e^{-T+t_{k}}+L_{5}(\sqrt{M_{0}}+L_{4}^{-1}+L_{5})\sqrt{2M_{0}(t-t_{k})}.

So combining these three parts of I_1 , we get

$$
I_1 \le 2\sqrt{2}L_0 e^{-(T-t)} \sqrt{Tr(C)(t-t_k)} + L_1 e^{-\frac{(T-t)}{2}} (t-t_k)
$$

+ $(1+\sqrt{2})(1+L_2)\sqrt{(t-t_k)M_0} + 2L_3 e^{-(T-t_k)} \sqrt{M_0(t-t_k)}.$ (88)

By Assumption [3](#page-5-3) with $\overleftrightarrow{X}_{t_k} \sim \overrightarrow{P}_{T-t_k}$,

$$
I_2 = \sqrt{\mathbb{E}|\tilde{s}(T - t_k, \overleftarrow{X}_{t_k}) - \tilde{s}_{\theta}(T - t_k, \overleftarrow{X}_{t_k})|^2} \le \epsilon.
$$
\n(89)

By Assumption [4,](#page-5-5)

$$
I_3 = \sqrt{\mathbb{E}|\tilde{s}_{\theta}(T - t_k, \overleftarrow{X}_{t_k}) - \tilde{s}_{\theta}(T - t_k, \overleftarrow{Y}_{t_k})|^2} \le L_0 e^{-(T - t_k)} \sqrt{\mathbb{E}|\overleftarrow{X}_{t_k} - \overleftarrow{Y}_{t_k}|^2}.
$$
(90)

By (68) , (76) , (77) , (78) , we have that the second term of (80) satisfies:

$$
\sqrt{\mathbb{E}|(I - C\bar{A}_{T-t}^{-1})\overleftarrow{X}_{t} - (I - C\bar{A}_{T-t_{k}}^{-1})\overleftarrow{Y}_{t_{k}}|^{2}}\n\n
$$
\leq \sqrt{\mathbb{E}|(I - C\bar{A}_{T-t}^{-1})\overleftarrow{X}_{t} - (I - C\bar{A}_{T-t_{k}}^{-1})\overleftarrow{X}_{t_{k}}|^{2}} + \sqrt{\mathbb{E}|(I - C\bar{A}_{T-t_{k}}^{-1})\overleftarrow{X}_{t_{k}} - (I - C\bar{A}_{T-t_{k}}^{-1})\overleftarrow{Y}_{t_{k}}|^{2}}\n\n
$$
\leq \sqrt{\mathbb{E}|(I - C\bar{A}_{T-t_{k}}^{-1})\overleftarrow{X}_{t} - (I - C\bar{A}_{T-t_{k}}^{-1})\overleftarrow{X}_{t}|^{2} + \sqrt{\mathbb{E}|(I - C\bar{A}_{T-t_{k}}^{-1})(\overleftarrow{X}_{t} - \overleftarrow{X}_{t_{k}})|^{2}} + \sqrt{\mathbb{E}|(I - C\bar{A}_{T-t_{k}}^{-1})\overleftarrow{X}_{t_{k}} - (I - C\bar{A}_{T-t_{k}}^{-1})\overleftarrow{Y}_{t_{k}}|^{2}}\n\n
$$
\leq ||C\bar{A}_{T-t}^{-1} - C\bar{A}_{T-t_{k}}^{-1}||\sqrt{\mathbb{E}|\overleftarrow{X}_{t}|^{2} + ||I - C\bar{A}_{T-t_{k}}^{-1}||\sqrt{\mathbb{E}|\overleftarrow{X}_{t} - \overleftarrow{X}_{t_{k}}|^{2}} + ||I - C\bar{A}_{T-t_{k}}^{-1}||\sqrt{\mathbb{E}|\overleftarrow{X}_{t} - \overleftarrow{X}_{t_{k}}|^{2}} + ||I - C\bar{A}_{T-t_{k}}^{-1}||\sqrt{\mathbb{E}|\overleftarrow{X}_{t} - \overleftarrow{X}_{t_{k}}|^{2}}\n\n
$$
\leq L_{3}(e^{-T+t} - e^{-T+t_{k}})\sqrt{M_{0}} + L_{2}e^{-T+t_{k}}\sqrt{2M_{0}(t - t_{k})} + L_{2}e^{-T+t_{k}}\sqrt{\mathbb{E}|\overleftarrow{X}_{t_{k}} - \overleftarrow{Y}_{t_{k}}|^{2}}\n\n
$$
\leq 2L_{3}e
$$
$$
$$
$$
$$
$$

Combine I_1 by [\(88\)](#page-16-0), I_2 by [\(89\)](#page-16-1), I_3 by [\(90\)](#page-16-2) and the second term [\(91\)](#page-16-3) of [\(80\)](#page-14-2) we get

$$
\sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_t) + (I - C\bar{A}_{T-t}^{-1})\tilde{X}_t - \tilde{s}_{\theta}(T-t_k,\tilde{Y}_{t_k}) - (I - C\bar{A}_{T-t_k}^{-1})\tilde{Y}_{t_k}|^2}
$$
\n
$$
\leq (2\sqrt{2}L_0e^{-T+t} + (1+\sqrt{2})(1+L_2) + 4L_3e^{-T+t_k} + \sqrt{2}L_2e^{-T+t_k} + L_1e^{-\frac{T-t}{2}})\sqrt{M_0(t-t_k)}
$$
\n
$$
+ \epsilon + (L_0 + L_2)e^{-T+t_k}\sqrt{\mathbb{E}|\tilde{X}_{t_k} - \tilde{Y}_{t_k}|^2}.
$$
\n
$$
(92)
$$

For $0 \le t - t_k \le 1$, we have by direct calculation

$$
t - t_k \le e(1 - e^{-(t - t_k)}),\tag{93}
$$

Thus by [\(92\)](#page-16-4) and [\(93\)](#page-17-0), we have

$$
\sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_t) + (I - C\bar{A}_{T-t}^{-1})\tilde{X}_t - \tilde{s}_{\theta}(T-t_k,\tilde{Y}_{t_k}) - (I - C\bar{A}_{T-t_k}^{-1})\tilde{Y}_{t_k}|^2}
$$
\n
$$
\leq K_1 \sqrt{M_0(1 - e^{-(t-t_k)})} + \epsilon + K_2 e^{-T+t_k} \sqrt{\mathbb{E}|\tilde{X}_{t_k} - \tilde{Y}_{t_k}|^2}.
$$
\n(94)

where

$$
K_1 = \sqrt{e(2\sqrt{2}L_0 + (1+\sqrt{2})(1+L_2) + 4L_3 + \sqrt{2}L_2 + L_1)},
$$

\n
$$
K_2 = L_0 + L_2.
$$
\n(95)

Step 3. Going back to [\(79\)](#page-14-1) we get: $\forall t \in [t_k, t_{k+1}],$

$$
\frac{dE(t)}{dt} \le K_1 \sqrt{M_0(e^t - e^{t_k})} + e^{\frac{t}{2}} (\epsilon + K_2 e^{-T + t_k} \sqrt{\mathbb{E} \|\tilde{X}_{t_k} - \tilde{Y}_{t_k}\|^2}).
$$
\n(96)

Since $t_{k+1} - t_k = h \leq 1$, we have

$$
\int_{t_k}^{t_{k+1}} \sqrt{e^t - e^{t_k}} dt = e^{\frac{t_k}{2}} \int_0^{t_{k+1}-t_k} \sqrt{e^s - 1} ds
$$

\n
$$
= e^{\frac{t_k}{2}} 2(\sqrt{e^{t_{k+1}-t_k} - 1} - \arctan \sqrt{e^{t_{k+1}-t_k} - 1})
$$

\n
$$
\leq \frac{2}{3} e^{\frac{t_k}{2}} (e^{t_{k+1}-t_k} - 1)^{\frac{3}{2}}
$$

\n
$$
= \frac{2}{3} (e^h - 1) \sqrt{e^{t_{k+1}} - e^{t_k}}
$$

\n
$$
\leq \frac{4}{3} h \sqrt{e^{t_{k+1}} - e^{t_k}}
$$

\n
$$
\leq \frac{16}{3} h (e^{\frac{t_{k+1}}{2}} - e^{\frac{t_k}{2}}).
$$
\n(97)

Thus, integration [\(96\)](#page-17-1) over time yields

$$
E(t_{k+1}) \leq E(t_k) + 2(e^{\frac{t_{k+1}}{2}} - e^{\frac{t_k}{2}})(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon + K_2e^{-T + \frac{t_k}{2}}E(t_k))
$$

$$
\leq E(t_k) \exp(K_2e^{-T}(e^{t_{k+1}} - e^{t_k})) + 2(e^{\frac{t_{k+1}}{2}} - e^{\frac{t_k}{2}})(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon) \exp(K_2e^{-T + t_{k+1}}).
$$
\n(98)

This gives us an Iterative formula:

$$
E(t_{k+1}) \exp(-K_2 e^{-T+t_{k+1}}) - 2e^{\frac{t_{k+1}}{2}} \left(\frac{8}{3}K_1 \sqrt{M_0}h + \epsilon\right) \le E(t_k) \exp(-K_2 e^{-T+t_k}) - 2e^{\frac{t_k}{2}} \left(\frac{8}{3}K_1 \sqrt{M_0}h + \epsilon\right). \tag{99}
$$

which leads to

$$
E(T) \exp(-K_2) - 2e^{\frac{T}{2}}(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon) \le E(0) \exp(-K_2e^{-T}) - 2(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon).
$$

Therefore we get

$$
\sqrt{E|X_T - Y_T|^2} \le e^{K_2} \left(e^{-\frac{T}{2}} \sqrt{\mathbb{E}|X_0 - Y_0|^2} + 2\left(\frac{8}{3}K_1 \sqrt{M_0}h + \epsilon\right)\right). \tag{100}
$$

Step 4. Finally we pick a ξ -optimal coupling of $\overleftrightarrow{X}_0 \sim \overrightarrow{P}_T$ and $\overleftrightarrow{Y}_0 \sim \overleftrightarrow{Q}_0 = \gamma_C$ in the Wasserstein distance, i.e

$$
\sqrt{\mathbb{E}|X_0 - Y_0|^2} \le \mathcal{W}_2(\vec{P}_T, \overleftarrow{Q}_0) + \xi,\tag{101}
$$

and obtain

$$
\mathcal{W}_2(\vec{P}_0, \overleftrightarrow{Q}_T) \le \sqrt{\mathbb{E}|\overleftarrow{X}_T - \overleftarrow{Y}_T|^2} \le e^{K_2} \left(e^{-\frac{T}{2}} (\mathcal{W}_2(\vec{P}_T, \overleftrightarrow{Q}_0) + \xi) + 2(\frac{8}{3}K_1 \sqrt{M_0}h + \epsilon)\right).
$$

Since ξ is arbitrary, the bound will be

$$
\mathcal{W}_2(\overset{\leftrightarrow}{P}_0,\overset{\leftarrow}{Q}_T)\leq \sqrt{\mathbb{E}|X_T-\overset{\leftarrow}{Y}_T|^2}\leq e^{K_2}(e^{-\frac{T}{2}}\mathcal{W}_2(\vec{P}_T,\overset{\leftarrow}{Q}_0)+2(\frac{8}{3}K_1\sqrt{M_0}h+\epsilon)).
$$

Noticing that

$$
\mathcal{W}_2^2(\vec{P}_0, \gamma_C) \leq \mathbb{E}_{\vec{P}_0 \otimes \gamma_C} |x - y|^2 = \mathbb{E}_{\vec{P}_0} |x|^2 + \mathbb{E}_{\gamma_C} |y|^2 = M_2 + \text{Tr}(C),
$$

and by Lemma [A.3,](#page-23-1)

$$
\mathcal{W}_2^2(\overrightarrow{P}_t, \gamma_C) \le e^{-t} \mathcal{W}_2^2(\overrightarrow{P}_0, \gamma_C).
$$

We get

$$
\mathcal{W}_2(\vec{P}_0, \overleftarrow{Q}_T) \le e^{K_2} (e^{-T} \sqrt{M_2 + \text{Tr}(C)} + 2(\frac{8}{3} K_1 \sqrt{M_0} h + \epsilon)).
$$
\n(102)

Remark A.2. For the early stop setting, use the same method we can get

$$
\mathcal{W}_2(\vec{P}_\delta, \overleftarrow{Q}_{T-\delta}) \le \exp(K_2 e^{-\delta}) \left(e^{-T+\frac{\delta}{2}} \sqrt{M_2 + \text{Tr}(C)} + 2\left(\frac{8}{3}K_1 \sqrt{M_0}h + \epsilon\right)\right). \tag{103}
$$

Let $\tilde{K}_1 = \frac{16}{3} K_1$ and $\tilde{K}_2 = K_2$ in Theorem [3.3](#page-5-0) then we complete our proof.

A.4 Result under the assumption in Remark [3.6](#page-6-2)

We only need to replace the term I_2 in the analysis of [A.3](#page-13-0) by

$$
\epsilon_k := \sqrt{\mathbb{E}|s(T - t_k, \overleftarrow{X}_{t_k}) - s_\theta(T - t_k, \overleftarrow{X}_{t_k})|^2}.
$$
\n(104)

 \Box

Then in step 3 the Iterative formula [\(99\)](#page-17-2) turns to

$$
E(t_{k+1}) \exp(-K_2 e^{-T+t_{k+1}}) - 2e^{\frac{t_{k+1}}{2}} \left(\frac{8}{3} K_1 \sqrt{M_0} h + \epsilon_k\right) \le E(t_k) \exp(-K_2 e^{-T+t_k}) - 2e^{\frac{t_k}{2}} \left(\frac{8}{3} K_1 \sqrt{M_0} h + \epsilon_k\right),\tag{105}
$$

or equivalently:

$$
E(t_{k+1}) \exp(-K_2 e^{-T+t_{k+1}}) \le E(t_k) \exp(-K_2 e^{-T+t_k}) + 2(e^{\frac{t_{k+1}}{2}} - e^{\frac{t_k}{2}}) (\frac{8}{3} K_1 \sqrt{M_0} h + \epsilon_k), \tag{106}
$$

The relaxed assumption [\(33\)](#page-6-4) gives us:

$$
\sum_{k=0}^{N-1} \frac{e^{\frac{t_{k+1}}{2}} - e^{\frac{t_k}{2}}}{e^{\frac{T}{2}}} \epsilon_k = \sum_{k=0}^{N-1} \frac{e^{\frac{t_k}{2}} (e^{\frac{t_{k+1} - t_k}{2}} - 1)}{e^{\frac{T}{2}}} \epsilon_k
$$
\n
$$
\leq \sum_{k=0}^{N-1} (e^{\frac{t_{k+1} - t_k}{2}} - 1) \epsilon_k
$$
\n
$$
\leq \sum_{k=0}^{N-1} (t_{k+1} - t_k) \epsilon_k
$$
\n
$$
\leq \epsilon T,
$$
\n(107)

so we get

$$
E(T) \exp(-K_2) \le E(0) \exp(-K_2 e^{-T}) + 2(e^{\frac{T}{2}} - 1)\frac{8}{3}K_1 \sqrt{M_0}h + 2e^{\frac{T}{2}}\epsilon T
$$

$$
\le E(0) + 2e^{\frac{T}{2}}(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon T).
$$
 (108)

or equivalently:

$$
\sqrt{\mathbb{E}|X_T - Y_T|^2} \le e^{K_2} \left(e^{-\frac{T}{2}} \sqrt{\mathbb{E}|X_0 - Y_0|^2} + 2\left(\frac{8}{3}K_1 \sqrt{M_0}h + \epsilon T\right)\right). \tag{109}
$$

Then apply the same argument as in [A.3](#page-13-0) step 4, we will get a similar result as (102) :

$$
\mathcal{W}_2(\vec{P}_0, \overleftarrow{Q}_T) \le e^{K_2} (e^{-T} \sqrt{M_2 + \text{Tr}(C)} + 2(\frac{8}{3}K_1 \sqrt{M_0}h + \epsilon T)).
$$
\n(110)

The only difference is the term ϵ here is replaced by ϵT .

By direct computation, the running time T and number of steps N to reach a distribution \overleftarrow{Q}_T such that $W_2(\vec{P}_0, \vec{Q}_T) = O(\epsilon_0)$ are still:

$$
T = \mathcal{O}(\log \frac{M_2 + Tr(C)}{\epsilon_0^2}),\tag{111}
$$

$$
N = \mathcal{O}\left(\frac{T}{\epsilon_0}\sqrt{M_0}\right),\tag{112}
$$

while the assumption in Remark [3.6](#page-6-2) changes to

$$
\epsilon = \mathcal{O}\left(\frac{\epsilon_0}{T}\right) = \mathcal{O}\left(\epsilon_0 \left(\log \frac{M_2 + Tr(C)}{\epsilon_0^2}\right)^{-1}\right). \tag{113}
$$

A.5 Result under the assumption in Remark [3.7](#page-6-3)

Here We need to replace the term I_3 in the analysis of [A.3](#page-13-0) step 2 by

$$
I_3 = \sqrt{\mathbb{E}|\tilde{s}_{\theta}(T - t_k, \overleftarrow{X}_{t_k}) - \tilde{s}_{\theta}(T - t_k, \overleftarrow{Y}_{t_k})|^2} \le \frac{1}{2} \sqrt{\mathbb{E}|\overleftarrow{X}_{t_k} - \overleftarrow{Y}_{t_k}|^2}.
$$
\n(114)

Then the last equation [\(94\)](#page-17-3) becomes

$$
\sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_t) + (I - C\bar{A}_{T-t}^{-1})\tilde{X}_t - \tilde{s}_{\theta}(T-t_k,\tilde{Y}_{t_k}) - (I - C\bar{A}_{T-t_k}^{-1})\tilde{Y}_{t_k}|^2}
$$
\n
$$
\leq K_1 \sqrt{M_0(1 - e^{-(t-t_k)})} + \epsilon + (L_2 e^{-T+t_k} + \frac{1}{2})\sqrt{\mathbb{E}|\tilde{X}_{t_k} - \tilde{Y}_{t_k}|^2}.
$$
\n(115)

in which the term $K_2e^{-T+t_k}$ is replaced by $L_2e^{-T+t_k} + \frac{1}{2}$. Then in step 3, the equation [\(96\)](#page-17-1) becomes

$$
\frac{dE(t)}{dt} \le K_1 \sqrt{M_0(e^t - e^{t_k})} + e^{\frac{t}{2}} (\epsilon + (L_2 e^{-T + t_k} + \frac{1}{2}) \sqrt{\mathbb{E} \|\tilde{X}_{t_k} - \tilde{Y}_{t_k}\|^2}).
$$
\n(116)

Again, using [\(97\)](#page-17-4) integration over time yields

$$
E(t_{k+1}) \leq E(t_k) + 2(e^{\frac{t_{k+1}}{2}} - e^{\frac{t_k}{2}})(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon + (L_2e^{-T + \frac{t_k}{2}} + \frac{1}{2}e^{-\frac{t_k}{2}})E(t_k))
$$

\n
$$
\leq E(t_k) \exp(L_2e^{-T}(e^{t_{k+1}} - e^{t_k}) + \frac{t_{k+1} - t_k}{2})
$$

\n
$$
+ 2(1 - e^{\frac{t_k - t_{k+1}}{2}})(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon) \exp(L_2e^{-T + t_{k+1}} + \frac{t_{k+1}}{2})
$$

\n
$$
\leq E(t_k) \exp(L_2e^{-T}(e^{t_{k+1}} - e^{t_k}) + \frac{t_{k+1} - t_k}{2})
$$

\n
$$
+ (t_{k+1} - t_k)(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon) \exp(L_2e^{-T + t_{k+1}} + \frac{t_{k+1}}{2}).
$$
\n(117)

Then in step 3 the Iterative formula [\(99\)](#page-17-2) turns to

$$
E(t_{k+1}) \exp(-L_2 e^{-T+t_{k+1}} - \frac{t_{k+1}}{2}) \le E(t_k) \exp(-K_2 e^{-T+t_k} - \frac{t_k}{2}) + (t_{k+1} - t_k) (\frac{8}{3} K_1 \sqrt{M_0} h + \epsilon),
$$
\n(118)

which lead to

$$
E(T) \exp(-L_2 - \frac{T}{2}) \le E(0) \exp(-L_2 e^{-T}) + T(\frac{8}{3}K_1 \sqrt{M_0} h + \epsilon).
$$
 (119)

or equivalently:

$$
\sqrt{\mathbb{E}|X_T - Y_T|^2} \le e^{L_2} \left(\sqrt{\mathbb{E}|X_0 - Y_0|^2} + T(\frac{8}{3}K_1\sqrt{M_0}h + \epsilon)\right).
$$
\n(120)

Then apply the same argument as in [A.3](#page-13-0) step 4, we will still a bound

$$
\mathcal{W}_2(\vec{P}_0, \vec{Q}_T) \le e^{L_2} (e^{-\frac{T}{2}} \sqrt{M_2 + \text{Tr}(C)} + T(\frac{8}{3}K_1 \sqrt{M_0}h + \epsilon)).
$$
\n(121)

By direct computation, to reach a distribution \overleftarrow{Q}_T such that $\mathcal{W}_2(\overrightarrow{P}_0, \overleftarrow{Q}_T) = O(\epsilon_0)$ requires:

$$
T = \mathcal{O}(\log \frac{M_2 + Tr(C)}{\epsilon_0^2}),\tag{122}
$$

$$
N = \frac{T}{h} = \mathcal{O}\left(\frac{T^2}{\epsilon_0}\sqrt{M_0}\right) = \mathcal{O}\left(\frac{\sqrt{M_0}}{\epsilon_0}\left(\log\frac{M_2 + Tr(C)}{\epsilon_0^2}\right)^2\right),\tag{123}
$$

and Assumption [3](#page-5-3) to hold with

$$
\epsilon = \mathcal{O}\left(\frac{\epsilon_0}{T}\right) = \mathcal{O}\left(\epsilon_0 \left(\log \frac{M_2 + Tr(C)}{\epsilon_0^2}\right)^{-1}\right). \tag{124}
$$

A.6 Proof of Theorem [3.8](#page-6-0)

Proof. Recall that

$$
q_{\sigma}(x) = \int_{\mathbb{R}^d} \exp(-\frac{|x - y|^2}{2\sigma^2}) q_0(y) dy, := \int_{\mathbb{R}^d} V(x, y) dy.
$$
 (125)

We observe that,

$$
\forall x \in \mathbb{R}^d, \ \forall |y| > R, \quad V(x, y) = 0. \tag{126}
$$

Fixing x , direct computation shows

$$
\nabla \log q_{\sigma}(x) + \frac{x}{\sigma^2} = \frac{\int_{\mathbb{R}^d} \left(-\frac{x-y}{\sigma^2}\right) V(x, y) dy}{\int_{\mathbb{R}^d} V(x, y) dy} + \frac{x}{\sigma^2}
$$
\n(127)

$$
=\frac{\int_{\mathbb{R}^d} \frac{y}{\sigma^2} V(x, y) dy}{\int_{\mathbb{R}^d} V(x, y) dy}.
$$
\n(128)

Taking absolute value and utilizing [\(126\)](#page-20-1), we got

$$
|\nabla g(x)| = |\nabla \log q_{\sigma}(x) + \frac{x}{\sigma^2}| \le \frac{R}{\sigma^2}.
$$

For any unit direction z, taking derivative twice

$$
\nabla_{z} \cdot \nabla_{z} (\log p_{\sigma} + \frac{|x|^{2}}{2\sigma^{2}}) = \frac{1}{\sigma^{2}} \nabla_{z} (\frac{\int_{\mathbb{R}^{d}} (y \cdot z) V(x, y) dy}{\int_{\mathbb{R}^{d}} V(x, y) dy})
$$
\n
$$
= \frac{1}{\sigma^{2}} \frac{\int_{\mathbb{R}^{d}} (y \cdot z) (-\frac{x - y}{\sigma^{2}} \cdot z) V(x, y) dy \int_{\mathbb{R}^{d}} V(x, y) dy - \int_{\mathbb{R}^{d}} (y \cdot z) V(x, y) dy \int_{\mathbb{R}^{d}} (-\frac{x - y}{\sigma^{2}} \cdot z) V(x, y) dy}{(\int_{\mathbb{R}^{d}} V(x, y) dy)^{2}}
$$
\n
$$
= \frac{1}{\sigma^{4}} \frac{\int_{\mathbb{R}^{d}} (y \cdot z)^{2} V(x, y) dy \int_{\mathbb{R}^{d}} V(x, y) dy - (\int_{\mathbb{R}^{d}} (y \cdot z) V(x, y) dy)^{2}}{(\int_{\mathbb{R}^{d}} V(x, y) dy)^{2}}.
$$
\n(129)

Taking absolute value and utilizing [\(126\)](#page-20-1) again, we got,

$$
|\nabla_z \cdot \nabla_z g(x)| = |\nabla_z \cdot \nabla_z (\log q_\sigma + \frac{|x|^2}{2\sigma^2})| \le \frac{2R^2}{\sigma^4}.
$$
 (130)

Thus

$$
\|\nabla^2 g\|_{\infty} \le \frac{2R^2}{\sigma^4}.\tag{131}
$$

 \Box

A.7 proof of Theorem [3.11](#page-7-2)

Proof. Use the same way as in proof [A.3.](#page-13-0) The corresponding constants (see Corollary [3.10\)](#page-6-6)should be replaced by

$$
K = \sup_{t \ge \delta} \|A\overline{A}^{-1}_t\| = 1,
$$

\n
$$
L_0 = 3 \frac{R^2}{(1 - e^{-\delta})^2},
$$

\n
$$
L_1 = \frac{R}{1 - e^{-\delta}},
$$

\n
$$
L_2 = \sup_{t \ge \delta} \|e^t (I - C\overline{A}^{-1}_t)\| = \frac{1}{1 - e^{-\delta}},
$$

\n
$$
L_3 = \sup_{t \ge s \ge \delta} \| (e^{-t} - e^{-s})^{-1} C(\overline{A}^{-1}_s - \overline{A}^{-1}_t) \| = \frac{1}{(1 - e^{-\delta})^2},
$$

\n
$$
M_0 = \max \{ M_2, \text{Tr}(C) \},
$$

\n
$$
K_1 = \sqrt{e} (2\sqrt{2}L_0 + (1 + \sqrt{2})(1 + L_2) + 4L_3 + \sqrt{2}L_2 + L_1),
$$

\n
$$
K_2 = L_0 + L_2.
$$

\n(132)

By Remark [A.2,](#page-18-2) we get

$$
\mathcal{W}_2(\vec{P}_{\delta}, \vec{Q}_{T-\delta}) \le \exp\left((3\frac{R^2}{(1 - e^{-\delta})^2} + \frac{1}{1 - e^{-\delta}})e^{-\delta}\right)\left(e^{-T + \frac{\delta}{2}}\sqrt{R_2 + \text{Tr}(C)} + 2(\frac{8}{3}K_1\sqrt{M_0h} + \epsilon)\right) \le \exp\left((3\frac{M^2}{\delta^2} + \frac{1}{\delta})\left(e^{-T + \frac{\delta}{2}}\sqrt{M_2 + \text{Tr}(C)} + 2(\frac{8}{3}K_1\sqrt{M_0h} + \epsilon)\right)\right).
$$
\n(133)

Let $\tilde{K}_3 = \frac{16}{3} K_1 = \frac{16}{3} \sqrt{e} \left(6 \sqrt{2} \frac{R^2}{(1 - e^{-1})^2}\right)$ $\frac{R^2}{(1-e^{-\delta})^2} + \frac{R}{1-e^{-\delta}} + 4\frac{1}{(1-e^{-\delta})^2} + (1+2\sqrt{2})\frac{1}{1-e^{-\delta}} + 1+\sqrt{2}$ here, then we complete the proof.

A.8 Proof of Theorem [3.15](#page-8-4)

Proof. Use the same way as in proof [A.3.](#page-13-0) Take $A = C$, and $h(x) = -\frac{|G(x) - y|_2^2}{2}$, then $\bar{A}(t) \equiv C^{-1}$ and $h(x)$ satisfies

$$
|\sqrt{C}\nabla h(x)| = |\sqrt{C}\nabla G(x)\Sigma^{-1}(G(x) - y)| \le ||C||^{\frac{1}{2}}(|G|_{\infty} + y)||\Sigma||^{-1}||\nabla G||_{\infty},
$$

\n
$$
||C\nabla^2 h(x)|| = ||C\nabla^2 G(x)\Sigma^{-1}(G(x) - y) + C\nabla G(x)\Sigma^{-1}\nabla G(x)^T|| \le ||C|| ||\Sigma||^{-1}(||\nabla^2 G||_{\infty}(|G|_{\infty} + |y||\nabla G||_{\infty}^2).
$$
\n(134)

The corresponding constants (see Corollary [3.2\)](#page-5-2)should be replaced by

$$
K = \sup_{t \ge 0} \|\bar{A}(t)A\| = 1,
$$
\n(135)

$$
L_0 = K^2(||C\nabla^2 h||_{\infty} + |\sqrt{C}\nabla h|_{\infty}^2) \le ||C||(||\Sigma||^{-1}(||\nabla G||_{\infty}^2 + (|G|_{\infty} + |y|) ||\nabla^2 G||_{\infty}) + ||\Sigma||^{-2} ||\nabla G||_{\infty}^2 (|G|_{\infty} + |y|)^2),
$$
\n(136)

$$
L_1 = K||C||^{\frac{1}{2}}|\sqrt{C}\nabla h| + \infty \le ||C|| ||\Sigma||^{-1}||\nabla G||_{\infty}(|G|_{\infty} + |y|),
$$
\n(137)

$$
L_2 = \sup_{t \ge \delta} \|e^t (I - C\bar{A}(t))\| = 0,\tag{138}
$$

$$
L_3 = \sup_{t \ge s \ge \delta} \|(e^{-t} - e^{-s})^{-1} C(\bar{A}(s) - \bar{A}(t))\| = 0,
$$
\n(139)

$$
M_0 = \max\{M_2, \text{Tr}(C)\},\tag{140}
$$

$$
K_1 = 2\sqrt{2}L_0 + (1+\sqrt{2})(1+L_2) + 4L_3 + \sqrt{2}L_2 + L_1,
$$
\n(141)

$$
K_2 = L_0 + L_2. \tag{142}
$$

By [\(102\)](#page-18-1) in [A.3](#page-13-0) and the constants listed above we get the bound

$$
\mathcal{W}_2(\vec{P}_0, \overleftarrow{Q}_T) \le e^{K_2} (e^{-T} \sqrt{M_2 + \text{Tr}(C)} + 2(\frac{8}{3}K_1 \sqrt{M_0}h + \epsilon)).\tag{143}
$$

Let

$$
\tilde{K}_4 = ||C|| ||\Sigma||^{-1} (||\nabla G||_{\infty}^2 + (|G|_{\infty} + |y|) ||\nabla^2 G||_{\infty})
$$

+
$$
||C|| ||\Sigma||^{-2} ||\nabla G||_{\infty}^2 (|G|_{\infty} + |y|)^2,
$$

$$
\tilde{K}_5 = \frac{8}{3} \sqrt{e} (2\sqrt{2}\tilde{K}_4 + ||C|| ||\Sigma||^{-1} ||\nabla G||_{\infty} (|G|_{\infty} + |y|)
$$

+
$$
1 + \sqrt{2}),
$$

then

$$
K_2 = L_2 \le \tilde{K}_4,
$$

\n
$$
\frac{8}{3} K_1 \le \tilde{K}_5,
$$
\n(144)

$$
\Box
$$

and we get the result

$$
\mathcal{W}_2(\vec{P}_0, \vec{Q}_T) \le e^{\tilde{K}_4} (e^{-T} \sqrt{M_2 + \text{Tr}(C)} + 2(\tilde{K}_5 \sqrt{M_0} h + \epsilon)).
$$
\n(145)

A.9 proof of Theorem [3.17](#page-8-5)

Proof. We follow the steps in [A.3.](#page-13-0) Step 1.For $t \in [t_k, t_{k+1}]$

$$
\frac{d|\overleftarrow{X}_t - \overleftarrow{Y}_t|^2}{dt} = \langle \overleftarrow{X}_t - \overleftarrow{Y}_t, \overleftarrow{X}_t + s(T - t, \overleftarrow{X}_t) - \overleftarrow{Y}_{t_k} - s_\theta(T - t_k, \overleftarrow{Y}_{t_k}) \rangle
$$

\n
$$
= \langle \overleftarrow{X}_t - \overleftarrow{Y}_t, \tilde{s}(T - t, \overleftarrow{X}_t) + (I - C\overleftarrow{A}(T - t))\overleftarrow{X}_t - \tilde{s}_\theta(T - t_k, \overleftarrow{Y}_{t_k}) - (I - C\overleftarrow{A}(T - t_k))\overleftarrow{Y}_{t_k} \rangle,
$$

Take expectation and use Cauchy's inequality we get

$$
\frac{d\mathbb{E}|\overleftarrow{X}_t-\overleftarrow{Y}_t|^2}{dt}\leq \sqrt{\mathbb{E}|\overleftarrow{X}_t-\overleftarrow{Y}_t|^2}\sqrt{\mathbb{E}|\tilde{s}(T-t,\overleftarrow{X}_t)+(I-C\bar{A}_{T-t}^{-1})\overleftarrow{X}_t-\tilde{s}_\theta(T-t_k,\overleftarrow{Y}_{t_k})-(I-C\bar{A}_{T-t_k}^{-1})\overleftarrow{Y}_{t_k}|^2}.
$$
 Therefore

Therefore

$$
\frac{d\sqrt{\mathbb{E}|X_t-\overset{\leftarrow}{Y}_t|^2}}{dt}\leq \frac{1}{2}\sqrt{\mathbb{E}|\tilde{s}(T-t,\overset{\leftarrow}{X}_t)+(I-C\bar{A}_{T-t}^{-1})\overset{\leftarrow}{X}_t-\tilde{s}_{\theta}(T-t_k,\overset{\leftarrow}{Y}_{t_k})-(I-C\bar{A}_{T-t_k}^{-1})\overset{\leftarrow}{Y}_{t_k}|^2}.
$$

Step 2. Recall [\(92\)](#page-16-4) :

$$
\sqrt{\mathbb{E}|\tilde{s}(T-t,\tilde{X}_t) + (I - C\bar{A}_{T-t}^{-1})\tilde{X}_t - \tilde{s}_{\theta}(T-t_k,\tilde{Y}_{t_k}) - (I - C\bar{A}_{T-t_k}^{-1})\tilde{Y}_{t_k}|^2}
$$

\n
$$
\leq (2\sqrt{2}L_0e^{-T+t} + (1+\sqrt{2})(1+L_2) + 4L_3e^{-T+t_k} + \sqrt{2}L_2e^{-T+t_k} + L_1e^{-\frac{T-t}{2}})\sqrt{M_0(t-t_k)}
$$

\n
$$
+ \epsilon + (L_0 + L_2)e^{-T+t_k}\sqrt{\mathbb{E}|\tilde{X}_{t_k} - \tilde{Y}_{t_k}|^2}
$$

\n
$$
= \frac{K_1}{\sqrt{e}}\sqrt{M_0(t-t_k)} + \epsilon + K_2e^{-T+t_k}\sqrt{\mathbb{E}|\tilde{X}_{t_k} - \tilde{Y}_{t_k}|^2}.
$$
\n(146)

Thus we have:

$$
\frac{d\sqrt{\mathbb{E}|X_t - Y_t|^2}}{dt} \le \frac{1}{2} \left(\frac{K_1}{\sqrt{e}} \sqrt{M_0(t - t_k)} + \epsilon + K_2 e^{-T + t_k} \sqrt{\mathbb{E}|X_t - Y_t|}\right)
$$

$$
\le \frac{1}{2} \left(\frac{K_1}{\sqrt{e}} \sqrt{M_0 h} + \epsilon + K_2 e^{-T + t_k} \sqrt{\mathbb{E}|X_t - Y_t|}\right).
$$

 $\mathcal{M}_0,\,\mathcal{K}_1$ and \mathcal{K}_2 are the same constants as in [A.3.](#page-13-0) Step 3.Integration over time yields

$$
\sqrt{\mathbb{E}|X_{t_{k+1}} - Y_{t_{k+1}}|^2} \le \sqrt{\mathbb{E}|X_{t_k} - Y_{t_k}|^2} + \frac{t_{k+1} - t_k}{2} (\frac{K_1}{\sqrt{e}} \sqrt{M_0 h} + \epsilon + K_2 e^{-T + t_k} \sqrt{\mathbb{E}|X_{t_k} - Y_{t_k}|^2}).
$$

$$
\le (\sqrt{\mathbb{E}|X_{t_k} - Y_{t_k}|^2} + \frac{t_{k+1} - t_k}{2} (\frac{K_1}{\sqrt{e}} \sqrt{M_0 h} + \epsilon)) \exp(\frac{K_2}{2} e^{-T} (e^{t_{k+1}} - e^{t_k})).
$$

This gives us an Iterative formula:

$$
\sqrt{\mathbb{E}|\overleftarrow{X}_{t_{k+1}} - \overleftarrow{Y}_{t_{k+1}}|^2} \exp\left(-\frac{K_2}{2}e^{-T+t_{k+1}}\right) - \frac{t_{k+1}}{2} \left(\frac{K_1}{\sqrt{e}}\sqrt{M_0h} + \epsilon\right) \le \sqrt{\mathbb{E}|\overleftarrow{X}_{t_k} - \overleftarrow{Y}_{t_k}|^2} \exp\left(-\frac{K_2}{2}e^{-T+t_k}\right) - \frac{t_k}{2} \left(\frac{K_1}{\sqrt{e}}\sqrt{M_0h} + \epsilon\right).
$$

which leads to

$$
\sqrt{\mathbb{E}|X_T - Y_T|^2} \exp\left(-\frac{K_2}{2}\right) - \frac{T}{2} \left(\frac{K_1}{\sqrt{e}} \sqrt{M_0 h} + \epsilon\right) \le \sqrt{\mathbb{E}|X_0 - Y_0|^2} \exp\left(-\frac{K_2}{2}e^{-T}\right).
$$

Therefore

$$
\sqrt{\mathbb{E}|X_T - Y_T|^2} \le e^{\frac{K_2}{2}} (\sqrt{\mathbb{E}|X_0 - Y_0|^2} + \frac{T}{2} (\frac{K_1}{\sqrt{e}} \sqrt{M_0 h} + \epsilon)).
$$
\n(147)

Step 4. Using the same argument in [A.3](#page-13-0) Step 4, we get

$$
\mathcal{W}_2(\vec{P}_0, \vec{Q}_T) \le e^{\frac{K_2}{2}} (e^{-\frac{T}{2}} \sqrt{M_2 + \text{Tr}(C)} + \frac{T}{2} (\frac{K_1}{\sqrt{e}} \sqrt{M_0 h} + \epsilon)). \tag{148}
$$

Let $\tilde{K}_6 = \frac{K_1}{2\sqrt{e}}$ then we complete the proof.

Here we present a lemma relates to convergence to equilibrium for the OU process, which is known by PDE experts.

Lemma A.3 (Theorem 23.26 [\[29\]](#page-11-16) and Exercise 9.10 [\[28\]](#page-11-17)). Let V be λ -uniformly convex C^2 potential. Consider Langevin process,

$$
dX_t = -\frac{1}{2}\nabla V(X_t)dt + dW_t,
$$
\n(149)

 \Box

with two initial measure μ_0 and ν_0 .

$$
W_2(\mu_t, \nu_t) \le W_2(\mu_0, \nu_0) e^{-\frac{\lambda t}{2}}.
$$
\n(150)

The convergence of OU is direct consequence, with $\lambda = 1$.

B Theories towards the generative diffusion model in infinite dimension

B.1 Motivating example towards the infinite dimensional result

We consider the following target distribution,

$$
p_0(x) = \frac{1}{Z} \Pi_{i=1}^d \left(\frac{1}{2} \exp\left(-\frac{|x - \sqrt{C_i}|^2}{2C_i}\right) + \frac{1}{2} \exp\left(-\frac{|x + \sqrt{C_i}|^2}{2C_i}\right)\right),\tag{151}
$$

where C is an $d \times d$ matrix diagonal matrix with $\{C_i\}_i$ as diagonal entries. Denote the case $d = 1$ and $C_1 = 1$ as

$$
p_0^1(x) = \frac{1}{\sqrt{2\pi}} \frac{1}{2} \left(\exp\left(-\frac{(x-1)^2}{2}\right) + \exp\left(-\frac{(x+1)^2}{2}\right) \right),\,
$$

and if one considers to apply the forward process (1) , one can get the distribution at time t

$$
p_t^1(x) = \frac{1}{\sqrt{2\pi}} \frac{1}{2} \left(\exp\left(-\frac{(x - e^{-\frac{t}{2}})^2}{2}\right) + \exp\left(-\frac{(x + e^{-\frac{t}{2}})^2}{2}\right) \right).
$$

Simple calculation shows,

$$
W_2(p_0, \mathcal{N}(0, C)) \le \sqrt{2\text{Tr}(C)}, \quad KL(p_0||\mathcal{N}(0, C)) = d \cdot KL(p_0^1||\mathcal{N}(0, 1)).\tag{152}
$$

And if one considers to apply the forward process [\(1\)](#page-2-0), one can also show that,

$$
W_2(p_t, N(0, C)) \le e^{-\frac{t}{2}} \sqrt{2\text{Tr}(C)}, \quad KL(p_t||\mathcal{N}(0, C)) = d \cdot KL(p_t^1||\mathcal{N}(0, 1)).
$$
\n(153)

From [\(152\)](#page-23-3) and [\(153\)](#page-23-4) we observe that when increasing dimension of C while keeping $\text{Tr}(C)$ fixed, the Wasserstein-2 bounds only scales with trace of C while the KL bounds scale with d .

B.2 Defining diffusion model in infinite dimension

Nowwe consider the forward and backward process in a separable Hilbert space H . [[\[23\]](#page-11-18). Section 3] The forward process has the same form as [\(1\)](#page-2-0)

$$
d\vec{X}_t = -\frac{1}{2}\vec{X}_t dt + \sqrt{C} dB_t, \quad 0 \le t \le T,
$$
\n(154)

where C becomes a trace-class. We still denote the marginal distributions of \vec{X}_t by \vec{P}_t , then \vec{P}_t will still converge to the stationary distribution $\mathcal{N}(0, C)$ as $t \to \infty$. ([\[8\]](#page-10-9) Theorem 11.11). The corresponding backward process is defined as:

$$
d\overleftarrow{X}_t = \left(\frac{1}{2}\overleftarrow{X}_t + s(T - t, \overleftarrow{X}_t)\right)dt + \sqrt{C}d\tilde{B}_t. \tag{155}
$$

The score function $s(t, x)$ is defined as:

$$
s(t,x) = -\frac{1}{1 - e^{-t}} \mathbb{E}[\overrightarrow{X}_t - e^{-\frac{t}{2}} \overrightarrow{X}_0 | \overrightarrow{X}_t = x],
$$
\n(156)

which will be same as $C\nabla \log \vec{P}_t(x)$ when $H = \mathbb{R}^d$. We still use a neural network $s_{\theta}(t, x)$ to approximate $s(t, x)$ and use our scheme [\(6\)](#page-3-1) for the sampling process. We assume that our Assumption [3](#page-5-3) for the training loss still holds. We will still focus on the Gaussian tail case(Assumtion [2\)](#page-3-0) with a initial Gaussian $\mathcal{N}(0, A)$:

$$
d\vec{P}_0 \propto \exp(h(x))d\mathcal{N}(0, A),\tag{157}
$$

where A shares the same eigen space with C and that both AC^{-1} and $A^{-1}C$ are bounded linear maps and h relates to some log-likelihood and detailed assumptions for it will be specified later in Assumption [6.](#page-24-1) To extend our Theorem [3.3](#page-5-0) to the infinite-dimensional case, we first project H onto a finite-dimensional subspace H^D , and then approximate the infinite-dimensional case using our results on H^D .

Spectral approximation of C [[\[23\]](#page-11-18) Appendix E] Suppose $C(A)$ has an orthonormal basis e_i of eigenvectors and corresponding non-negative eigenvalues $c_i \geq 0$ ($a_i \geq 0$), we define the linear span of the first D eigenvectors as

$$
H^D = span\{e_1, e_2, \cdots, e_D\}.
$$
\n
$$
(158)
$$

Let $P^D: H \to H^D$ be the orthogonal projection onto H^D . We define the finite-dimensional approximations of μ_{data} by $\mu_{data}^D = P^D_{\#}(\mu_{data})$ and discretize the forward process by $\overrightarrow{X_t^D} = P^D \overrightarrow{X}_t$, then $(\overrightarrow{X_t^D})_{t\geq 0}$ will satisfy

$$
d\vec{X}_t^D = -\frac{1}{2}\vec{X}_t^D dt + \sqrt{P^D C P^D} dB_t.
$$
\n(159)

We denote the marginal distributions of $\overrightarrow{X_t^D}$ by $\overrightarrow{P_t^D}$. The corresponding backward process will be

$$
dX_t^D = \left(\frac{1}{2}\overline{X}_t^D + s^D(T - t, \overline{X}_t^D)\right)dt + \sqrt{P^D CP^D}d\tilde{B}_t.
$$
\n(160)

where $s^D(t, x) = P^D \mathbb{E}[s(t, \vec{X}_t) | P^D \vec{X}_t = x] = C^D \nabla \log \vec{P}_t^D(x)$ ([\[23\]](#page-11-18) Appendix E). To apply our PDE method to get the Hessian bound, we need the following assumption:

Assumption 6. $\forall D$, the orthogonal projection P^D satisfies,

1. \overrightarrow{P}_0^D $\frac{D}{0}$ is still a Gaussian tail: $\overrightarrow{P}_{0}^{D}$ $\sum_{0}^{D} (x) = \exp(-\frac{|x|_{AD}^{2}}{2}) \exp(h^{D}(x)),$ 2. $h^D(x)$ satisfies $|\nabla h^D|_{H^D,\infty} \leq |\nabla ||_{H,\infty}$ and $||\nabla^2 h^D||_{H^D,\infty} \leq ||\nabla^2 h||_{H,\infty}$.

Define $C^D \bar{A}^D(t)$ like before in the main context and $\tilde{s}^D(t,x) = s^D(t,x) + C^D \bar{A}^D(t)x$, with Assumption [6](#page-24-1) we have that $\tilde{s}^{D}(t, x)$ still satisfies Corollary [3.2:](#page-5-2)there exist constants $L_0, L_1 \geq 0$ such that

$$
\begin{aligned} \|\nabla \tilde{s}^D(t, \cdot)\|_{\infty} &\le L_0 e^{-t}, \\ \|\tilde{s}^D(t, \cdot)\|_{\infty} &\le L_1 e^{-\frac{t}{2}}, \quad \forall t \ge 0. \end{aligned}
$$

and L_0, L_1 can be independent of the dimension because of the second part of Assumption [6.](#page-24-1) By [\[23\]](#page-11-18) lemma 5 (listed as Proposition [B.1\)](#page-25-0), we have $\tilde{s}^D(t, \vec{X}^D_t) \to \tilde{s}(t, \vec{X}_t) = s(t, \vec{X}_t) + C \vec{A}(t) \vec{X}_t$ both a.s. and L_2 , as $D \to \infty$, thus we may also assume that

Assumption 7. Both $\tilde{s}(t,x)$ and $\tilde{s}_{\theta}(t,x)$ are L_0e^{-t} -Lipschitz.

With Assumption [6](#page-24-1) and [7](#page-24-2) and using the same technical skills in [A.3,](#page-13-0) we can get the same result as Theorem [3.3](#page-5-0) in infinite space:

$$
\mathcal{W}_2(\vec{P}_0, \vec{Q}_T) \le \tilde{K}_2(\exp(-T)(M_2 + \text{Tr}(C)) + \tilde{K}_1 \sqrt{\frac{T}{N} M_0} + 2\epsilon),\tag{161}
$$

where $M_0 = \max\{\text{Tr}(C), M_2\}, \tilde{K}_1, \tilde{K}_2$ are some constants independent of dimension.

Here we listed the lemma that is used to extend our analysis to infinite dimension.

Proposition B.1. ([\[23\]](#page-11-18) lemma 5) Let H be a separable Hilbert space and Z, \tilde{Z} be two random variables taking values in H Let e_i be an orthonormal basis of H. Denote by $H^D = span\{e_1, \dots, e_D\}$ and by P^D the projection onto H^D . Furthermore, let $Z^D = P^D \mathbb{E}[Z|P^D\tilde{Z}]$ Then, if $\mathbb{E}[\|\mathbb{E}[Z|\tilde{Z}]\|_H^2] < \infty$, then $Z^D \to \mathbb{E}[Z|\tilde{Z}]$ in L^2 and almost surely.