Normalized solutions to a quasilinear equation involving critical Sobolev exponent

Nidhi* and K. Sreenadh[†]

Abstract

In this paper we study the existence and regularity results of normalized solutions to the following quasilinear elliptic Choquard equation with critical Sobolev exponent and mixed diffusion type operators:

$$-\Delta_p u + (-\Delta_p)^s u = \lambda |u|^{p-2} u + |u|^{p^*-2} u + \mu (I_\alpha * |u|^q) |u|^{q-2} u \text{ in } \mathbb{R}^N,$$
$$\int_{\mathbb{D}^N} |u|^p dx = \tau,$$

where $N \geq 3$, $\tau > 0$, $\frac{p}{2}(\frac{N+\alpha}{N}) < q < \frac{p}{2}(\frac{N+\alpha}{N-p})$, I_{α} is the Riesz potential of order $\alpha \in (0, N)$, $\mu > 0$ is a parameter, $(-\Delta_p)^s$ is the fractional p-laplacian operator, $p^* = \frac{Np}{N-p}$ is the critical Sobolev exponent and λ appears as a Lagrange multiplier.

Keywords: Normalized solutions, Choquard equation, critical growth, local and nonlocal operator, existence results, Hölder regularity.

1 Introduction

This work is mainly focused on the existence and regularity results of the following quasilinear equation with critical Sobolev exponent, Choquard nonlinearity and a mixed diffusion type operator:

$$-\Delta_p u + (-\Delta_p)^s u = \lambda |u|^{p-2} u + |u|^{p^*-2} u + \mu (I_\alpha * |u|^q) |u|^{q-2} u \text{ in } \mathbb{R}^N,$$
 (1.1)

with a fixed L^p norm

$$\int_{\mathbb{R}^N} |u|^p dx = \tau,\tag{1.2}$$

where $N \ge 3$, 0 < s < 1 < p < N, $\tau > 0$, $p^* = \frac{Np}{N-p}$ is the critical Sobolev exponent. The operators p-laplacian (Δ_p) and fractional p-laplacian $(-\Delta_p)^s$ are defined as:

$$\Delta_p u = div(|\nabla u|^{p-2} \nabla u),$$

and

$$(-\Delta_p)^s u(x) = \lim_{\epsilon \to 0} \int_{\mathbb{R}^N \setminus B_{\epsilon}(0)} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} dy \text{ for } s \in (0, 1).$$

^{*}Department of Mathematics, Indian Institute of Technology, Delhi, Hauz Khas, New Delhi-110016, India. e-mail: nidhi.nidhi@maths.iitd.ac.in

 $^{^\}dagger$ Department of Mathematics, Indian Institute of Technology, Delhi, Hauz Khas, New Delhi-110016, India. e-mail: sreenadh@maths.iitd.ac.in

 I_{α} is the Riesz potential of order $\alpha \in (0, N)$ given by

$$I_{\alpha}(x) = \frac{A_{N,\alpha}}{|x|^{N-\alpha}} \text{ with } A_{N,\alpha} = \frac{\Gamma(\frac{N-2}{2})}{\pi^{\frac{N}{2}} 2^{\alpha} \Gamma(\frac{\alpha}{2})} \text{ for every } x \in \mathbb{R}^{N} \setminus \{0\}.$$
 (1.3)

The problem is well defined for all $\frac{p}{2}(\frac{N+\alpha}{N}) \leq q \leq \frac{p}{2}(\frac{N+\alpha}{N-p})$ due to the following well known Hardy-Littlewood-Sobolev inequality:

Proposition 1.1. Let t, r > 1 and $0 < \alpha < N$ with $1/t + 1/r = 1 + \alpha/N$, $f \in L^t(\mathbb{R}^N)$ and $h \in L^r(\mathbb{R}^N)$. There exists a sharp constant $C(t, r, \alpha, N)$ independent of f, h, such that

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{f(x)h(y)}{|x-y|^{N-\alpha}} \, dxdy \le C(t,r,\alpha,N) \|f\|_{L^t} \|h\|_{L^r}. \tag{1.4}$$

If $t = r = 2N/(N + \alpha)$, then

$$C(t, r, \alpha, N) = C(N, \alpha) = \pi^{\frac{N-\alpha}{2}} \frac{\Gamma(\frac{\alpha}{2})}{\Gamma(\frac{N+\alpha}{2})} \left\{ \frac{\Gamma(\frac{N}{2})}{\Gamma(N)} \right\}^{-\frac{\alpha}{N}}.$$
 (1.5)

Equality holds in (1.4) if and only if $\frac{f}{h} \equiv constant$ and $h(x) = A(\gamma^2 + |x - a|^2)^{(N+\alpha)/2}$ for some $A \in \mathbb{C}, 0 \neq \gamma \in \mathbb{R}$ and $a \in \mathbb{R}^N$.

It follows from Proposition 1.1 that

$$\mathcal{A}_q(u) := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x)|^q |u(y)|^q}{|x - y|^{N - \alpha}} dx dy$$

is well defined if $\frac{p}{2}\left(\frac{N+\alpha}{N}\right) \leq q \leq \frac{p}{2}\left(\frac{N+\alpha}{N-p}\right)$. The expression $(I_{\alpha}*|u|^q)|u|^{q-2}u$ is referred to as Choquard-type nonlinearity, as it was employed by Choquard in the examination of the Hartree-Fock theory of a one-component plasma. At the Symposium on Coulomb Systems, Choquard explained how the energy functional associated with the problem:

$$\begin{cases}
-\Delta u + u = (I_2 * |u|^2)u \text{ in } \mathbb{R}^3, \\
u \in H^1(\mathbb{R}^3).
\end{cases}$$
(1.6)

can be used in this regard, see [17]. Also, (1.6) was used by Pekar in [28] to study the quantum theory model of a stationary polaron. The equations of the type

$$-\Delta u + \lambda u = \mu(I_{\alpha} * |u|^p)|u|^{p-2}u \text{ in } \mathbb{R}^N,$$
(1.7)

are called the Choquard equation. Such equations mostly occur in the field of quantum physics, see [30], and have been extensively studied. We refer the readers to go through [8], [19], and [21] to see how the existence, multiplicity, and qualitative characteristics of the solution to (1.7) are deduced. This study is devoted to the normalized solution for a category of quasilinear elliptic Choquard equation associated with the operator $\mathcal{L} = -\Delta_p + (-\Delta_p)^s$. To the best of our knowledge, there hasn't been much research on such problems. Formally speaking, the solution to the class of problems of the form:

$$\begin{cases}
-\Delta u = \lambda u + g(u) \text{ in } \mathbb{R}^N, \\
\int_{\mathbb{R}^N} |u|^2 dx = c.
\end{cases}$$
(1.8)

is called a normalized solution. The solution of (1.8) gives us the stationary state of a nonlinear Schrödinger equation with a predetermined fixed L^2 -norm. Jeanjean studied (1.8) in [14] and deduced the existence of a radially symmetric solution under some assumptions on g. Further, Bartsch and De Valeriola proved the existence of infinitely many solutions (see [2]). Moreover, taking $g(t) = |t|^{p-1}t$, (1.8) has been studied for bounded domains along with some boundary conditions; interested readers can go through [25] and [31]. The existence of normalized solutions for nonlinear Schrödinger systems has been thoroughly investigated. Readers seeking further information may consult the references [3–5,13,24,26]. The study of quadratic ergodic mean field games systems also investigates normalized solution types, as discussed in [29].

Recently, the study of normalized solutions to the Choquard equation has attracted the researchers. The linear case, that is p=2, has been dealt with by a lot of authors; for instance, see [16,33,35]. The authors in [32] and [20] have studied the existence of normalized solutions to the critical Choquard equation, where the critical exponent is due to the above-mentioned Hardy-Littlewood-Sobolev inequality, involving classical and fractional Laplacian operators, respectively and in [12], such critical growth Choquard equation has been seen studied with the presence of mixed diffusion type operator.

Motivated by the previously mentioned literature, we aim to investigate the existence and regularity results for normalized solutions to a broader category of quasilinear elliptic equations featuring critical Sobolev exponents and mixed diffusion-type operators. In addition to being a critical case, the presence of classical p-laplacian and fractional p-laplacian operators in the equation intensifies the complexity and interest of our study.

We will initiate our study by discussing the regularity properties of a radially symmetric positive weak solution of (1.1)-(1.2). Here, the space framework is the Banach space $W^{1,p}(\mathbb{R}^N)$ equipped with the following equivalent norm:

$$||u|| = (||\nabla u||_p^p + ||u||_p^p + [u]_{s,p}^p)^{\frac{1}{p}},$$

where

$$[u]_{s,p}^p := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy,$$

and $W_r^{1,p}(\mathbb{R}^N) = \{u \in W^{1,p}(\mathbb{R}^N) : u \text{ is radial}\}$. Morever, the space $\mathbb{W} = W^{1,p}(\mathbb{R}^N) \times \mathbb{R}$, with $\|(u,t)\|_{\mathbb{W}}^p := \|u\|^p + |t|^p$, will also be used. The notion of weak solution for (1.1)-(1.2) is as follows:

Definition 1.1. A function $u \in W^{1,p}(\mathbb{R}^N)$ is said to be the weak solution of (1.1)-(1.2) if $||u||_p^p = \tau$ and

$$\int_{\mathbb{R}^{N}} |\nabla u|^{p-2} \nabla u \nabla v + \ll u, v \gg_{s,p} = \lambda \int_{\mathbb{R}^{N}} |u|^{p-2} uv + \int_{\mathbb{R}^{N}} |u|^{p^{*}-2} uv + \mu \int_{\mathbb{R}^{N}} (I_{\alpha} * |u|^{q}) |u|^{q-2} uv,$$

for every $v \in W^{1,p}(\mathbb{R}^N)$, where $\ll u, v \gg_{s,p} := \frac{1}{2} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y)) (v(x) - v(y))}{|x - y|^{N+2s}} dx dy$.

Defining $A := \mu \mathcal{A}_q$, the energy functional corresponding to the problem (1.1) is given by

$$F(u) = \frac{1}{p} \|\nabla u\|_p^p + \frac{\lambda}{p} \|u\|_p^p + \frac{1}{p} [u]_{s,p}^p - \frac{1}{p^*} \|u\|_{p^*}^{p^*} - \frac{1}{2q} A(u),$$

that is, a critical point of F which satisfies the constraint turns out to be the weak solution of (1.1)-(1.2). Precisely, we can say the following about a solution of (1.1)-(1.2):

Theorem 1.1. Let $p \geq 2$, $N - \alpha \leq 2p$ and $p \leq q \leq \frac{p}{2}(\frac{N+\alpha}{N-p})$. If $0 < u \in W^{1,p}(\mathbb{R}^n)$ is a solution of (1.1)-(1.2), corresponding to some $\lambda < 0$, then $u \in L^{\infty}_{loc}(\mathbb{R}^N)$. Moreover, if u is radially symmetric, then $u \in L^{\infty}(\mathbb{R}^N) \cap C^{\delta}_{loc}(\mathbb{R}^N)$ for every $0 < \delta < \Theta = \min\{\frac{sp}{n-1}, 1\}$.

Following the work of R. Biswas and S. Tiwari in [6], an iterative scheme was constructed to show that a positive solution must lie in $L^{\infty}_{loc}(\mathbb{R}^N)$. Moreover, using the radial lemma [34], it is evident that a radially symmetric solution would lie in $L^{\infty}(\mathbb{R}^N)$, and hence the results of [10] can be used to figure out the Hölder regularity.

After obtaining the regularity results, we investigated the existence of the normalized solution. Inspired by the work of X. Shang and P. Ma in [32], existence results for the cases: $\frac{p}{2} \left(\frac{N+\alpha}{N} \right) < q < \frac{p}{2} \left(\frac{N+\alpha+sp}{N} \right)$ and $\frac{p}{2} \left(\frac{N+\alpha+sp}{N} \right) \le q < \frac{p}{2} \left(\frac{N+\alpha}{N-p} \right)$ is discussed separately. Settling the set of functions satisfying the constraint by:

$$S(\tau) := \left\{ u \in W^{1,p}(\mathbb{R}^N) : ||u||_p^p = \tau \right\}.$$

and using the fiber map defined as

$$t \star u(x) := e^{\frac{Nt}{p}} u(e^t x),$$

the idea is to work on the natural Pohozaev manifold, given by

$$P(\tau) := S(\tau) \cap \bar{M} = S(\tau) \cap \left\{ u \in W^{1,p}(\mathbb{R}^N) : M(u) = 0 \right\},$$

where

$$M(u) := \|\nabla u\|_p^p + s[u]_{s,p}^p - \|u\|_{p^*}^{p^*} - \gamma_q A(u),$$

with $\gamma_q = \frac{N}{p} - \frac{N+\alpha}{2q}$. Using the concentration compactness principles by Lions in [18] we deduce the following for $\frac{p}{2} \left(\frac{N+\alpha}{N} \right) < q < \bar{q}_s := \frac{p}{2} \left(\frac{N+\alpha+sp}{N} \right)$:

Theorem 1.2. For 0 < s < 1 < p < N, and $\frac{p}{2} \left(\frac{N+\alpha}{N} \right) < q < \bar{q}_s$, there exists $\tau_0 > 0$ such that, for all $\tau < \tau_0$, the problem (1.1)-(1.2) has a radially symmetric ground state solution $u_{\tau}^* \in W^{1,p}(\mathbb{R}^N)$ corresponding to some $\lambda < 0$ with $E(u_{\tau}^*) < 0$.

Further, for the case of $\bar{q}_s \leq q < \frac{p}{2} \left(\frac{N+\alpha}{N-p} \right)$, ϵ - blow up analysis was used to conclude the following:

Theorem 1.3. Let 0 < s < 1 < p < N and $\mu > 0$ is sufficiently large, then we have the following:

- 1. if $q = \bar{q}_s$, then there exists, $\bar{\tau}_s > 0$ such that for all $\tau \in (0, \bar{\tau}_s)$, (1.1)-(1.2) has a solution $(u_\tau, \lambda_\tau) \in W^{1,p}_r(\mathbb{R}^N) \times \mathbb{R}$ for some $\lambda < 0$,
- 2. if $\bar{q}_s < q \leq \bar{q} := \frac{p}{2} \left(\frac{N + \alpha + p}{N} \right)$, then there exists, $\bar{\tau}_q > 0$ such that for all $\tau \in (0, \bar{\tau}_q)$, (1.1)-(1.2) has a solution $(u_\tau, \lambda_\tau) \in W^{1,p}_r(\mathbb{R}^N) \times \mathbb{R}$ for some $\lambda < 0$,
- 3. for all $\bar{q} < q < \frac{p}{2} \left(\frac{N+\alpha}{N-p} \right)$, (1.1)-(1.2) has a solution $(u_{\tau}, \lambda_{\tau}) \in W_r^{1,p}(\mathbb{R}^N) \times \mathbb{R}$ for some $\lambda < 0$.

Notations

We have used the following notations throughout the paper:

- $T(u) = \|\nabla u\|_p^p + [u]_{s,p}^p$
- S is the best Sobolev constant, see [36], given by:

$$S = \inf_{u \in W^{1,p}(\mathbb{R}^N) \setminus \{0\}} \frac{\|\nabla u\|_p^p}{\|u\|_{p^*}^p}$$
 (1.9)

and is achieved by the family of functions of the form:

$$U_{\epsilon,x_0}(x) = \frac{K_{N,p} \epsilon^{\frac{N-p}{p(p-1)}}}{(\epsilon^{\frac{p}{p-1}} + |x - x_0|^{\frac{p}{p-1}})^{\frac{N-p}{p}}}, \text{ for } x_0 \in \mathbb{R}^N \text{ and } \epsilon > 0.$$
 (1.10)

• $p_{\alpha}^* := \frac{p}{2} \left(\frac{N+\alpha}{N-p} \right)$.

2 Regularity Result

Let us start with discussing the regularity properties of a positive radial solution to (1.1)-(1.2).

Proof of Theorem 1.1 : For $0 < \epsilon < 1$, define $h_{\epsilon}(t) := \sqrt{\epsilon^2 + t^2}$, clearly, $g_{\epsilon} := h'_{\epsilon}$ is a continuously differentiable convex function, with $g_{\epsilon}(0) = 0$ and $|g'_{\epsilon}(t)| \leq M_{\epsilon}$ for some $M_{\epsilon} > 0$, thus by Theorem 2.2.3 of [15], $g_{\epsilon}(u) \in W^{1,p}(\mathbb{R}^N)$. Let $0 < \phi \in C_c^{\infty}(\mathbb{R}^N)$ be arbitrary, now define $\zeta := \phi |g_{\epsilon}(u)|^{p-2} g_{\epsilon}(u) \in W^{1,p}(\mathbb{R}^N)$, then we have:

$$\int_{\mathbb{R}^N} |\nabla u|^{p-2} \nabla u \nabla \zeta = (p-1) \int_{\mathbb{R}^N} |\nabla u|^p \phi |g_{\epsilon}(u)|^{p-2} g_{\epsilon}' + \int_{\mathbb{R}^N} |\nabla u|^{p-2} |g_{\epsilon}(u)|^{p-2} g_{\epsilon}(u) \nabla u \nabla \phi := I_1^{\epsilon},$$
(2.1)

$$\ll u, \zeta \gg_{s,p} = \int_{\mathbb{R}^{N}} \int_{\mathbb{R}^{N}} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y)) (\zeta(x) - \zeta(y))}{|x - y|^{N+sp}} dx dy
\geq \int_{\mathbb{R}^{N}} \int_{\mathbb{R}^{N}} \frac{|h_{\epsilon}(u(x)) - h_{\epsilon}(u(y))|^{p-2} (h_{\epsilon}(u(x)) - h_{\epsilon}(u(y))) (\phi(x) - \phi(y))}{|x - y|^{N+sp}}
:= I_{2}^{\epsilon}$$
(2.2)

by Lemma 3.2 of [6]. Taking ζ as test function, by (2.1) and (2.2) we get:

$$I_{1}^{\epsilon} + I_{2}^{\epsilon} + (-\lambda) \int_{\mathbb{R}^{N}} |u|^{p-2} u \zeta \leq \int_{\mathbb{R}^{N}} |\nabla u|^{p-2} \nabla u \nabla \zeta + \ll u, \zeta \gg_{s,p} -\lambda \int_{\mathbb{R}^{N}} |u|^{p-2} u \zeta$$

$$= \int_{\mathbb{R}^{N}} |u|^{p^{*}-2} u \zeta + \mu \int_{\mathbb{R}^{N}} (I_{\alpha} * |u|^{q}) |u|^{q-2} u \zeta$$

$$\leq \int_{\mathbb{R}^{N}} |u|^{p^{*}-2} u \phi + \mu \int_{\mathbb{R}^{N}} (I_{\alpha} * |u|^{q}) |u|^{q-2} u \phi, \qquad (2.3)$$

since $|g_{\epsilon}(t)| < 1$. Now, taking $\epsilon \to 0$ in (2.3), by dominated convergence theorem, we get:

$$\int_{\mathbb{R}^{N}} |\nabla u|^{p-2} \nabla u \nabla \phi + \ll |u|, \phi \gg_{s,p} \leq \lambda \int_{\mathbb{R}^{N}} |u|^{p-2} u \phi + \int_{\mathbb{R}^{N}} |u|^{p^{*}-2} u \phi + \mu \int_{\mathbb{R}^{N}} (I_{\alpha} * |u|^{q}) |u|^{q-2} u \phi$$
(2.4)

for all $0 < \phi \in C_c^{\infty}(\mathbb{R}^N)$ and hence by density, for all $0 < \phi \in W^{1,p}(\mathbb{R}^N)$. Now, for a fixed $\gamma > 0$, define $u_{\gamma} := \min\{\gamma, |u|\}$ and for some k > 1, set $\beta = kp - p + 1$. Replacing ϕ by u_{γ}^{β} in (2.4), then by Lemma 3.1 of [6] we get:

$$\begin{split} \frac{\beta p^p}{(\beta+p-1)^p} \left(\left\| \nabla (u_\gamma^k) \right\|_p^p + \left[u_\gamma^k \right]_{s,p}^p \right) & \leq & \beta \int_{\{x:|u(x)|<\gamma\}} u_\gamma^{\beta-1} |\nabla u_\gamma|^p + \ll |u|, u_\gamma^\beta \gg_{s,p} \\ & = & \int_{\mathbb{R}^N} |\nabla u|^{p-2} \nabla u \nabla u_\gamma^\beta + \ll |u|, u_\gamma^\beta \gg_{s,p} \\ & \leq & \lambda \int_{\mathbb{R}^N} |u|^{p-2} u u_\gamma^\beta + \int_{\mathbb{R}^N} |u|^{p^*-2} u u_\gamma^\beta \\ & + \mu \int_{\mathbb{R}^N} (I_\alpha * |u|^q) |u|^{q-2} u u_\gamma^\beta. \end{split}$$

Now, since u > 0 and $\lambda < 0$ we get

$$\frac{\beta}{k^p} \left(\left\| \nabla u_\gamma^k \right\|_p^p + [u_\gamma^k]_{s,p}^p \right) + (-\lambda) \left\| u_\gamma^k \right\|_p^p \leq \int_{\mathbb{R}^N} |u|^{p^*-2} u u_\gamma^\beta + \mu \int_{\mathbb{R}^N} (I_\alpha * |u|^q) |u|^{q-2} u u_\gamma^\beta,$$

and hence

$$\frac{C\beta}{k^p} \left\| u_\gamma^k \right\|^p \le \int_{\mathbb{R}^N} |u|^{p^*-2} u u_\gamma^\beta + \mu \int_{\mathbb{R}^N} (I_\alpha * |u|^q) |u|^{q-2} u u_\gamma^\beta,$$

here $C = \min\{1, \frac{(-\lambda)k^p}{\beta}\}$. Thus, by the imbedding $W^{1,p}(\mathbb{R}^N) \hookrightarrow L^{p^*}(\mathbb{R}^N)$, there exists a constant C_1 such that

$$\left\| u_{\gamma}^{k} \right\|_{p^{*}}^{p} \leq C_{1} k^{p-1} \left(\int_{\mathbb{R}^{N}} |u|^{p^{*}-2} u u_{\gamma}^{\beta} + \mu \int_{\mathbb{R}^{N}} (I_{\alpha} * |u|^{q}) |u|^{q-2} u u_{\gamma}^{\beta} \right), \tag{2.5}$$

because, $\beta > k$ for k > 1. Now, for some $\delta > 1$, by Proposition 1.1, Hölder's inequality, and the fact that $(a + b)^c \le a^c + b^c$ whenever a, b > 0 and c < 1, we have:

$$\mu \int_{\mathbb{R}^{N}} (I_{\alpha} * |u|^{q}) |u|^{q-2} u u_{\gamma}^{\beta} \\
\leq \mu C_{N} \left(\int_{\mathbb{R}^{N}} |u|^{\frac{2qN}{N+\alpha}} \right)^{\frac{N+\alpha}{2N}} \left(\int_{\mathbb{R}^{N}} (|u|^{q-2} |u u_{\gamma}^{\beta}|)^{\frac{2N}{N+\alpha}} \right)^{\frac{N+\alpha}{2N}} \\
\leq \bar{C}_{N} \left(\int_{\{x:|u(x)|<\delta\}} (|u(x)|^{q-2} |u u_{\gamma}^{\beta}|)^{\frac{2N}{N+\alpha}} + \int_{\{x|u(x)|\geq\delta\}} (|u|^{q-2} |u u_{\gamma}^{\beta}|)^{\frac{2N}{N+\alpha}} \right)^{\frac{N+\alpha}{2N}} \\
\leq \bar{C}_{N} \left(\int_{\{x:|u(x)|<\delta\}} (|u|^{q-2} |u u_{\gamma}^{\beta}|)^{\frac{2N}{N+\alpha}} \right)^{\frac{N+\alpha}{2N}} + \bar{C}_{N} \left(\int_{\{x:|u(x)|\geq\delta\}} (|u|^{q-2} |u u_{\gamma}^{\beta}|)^{\frac{2N}{N+\alpha}} \right)^{\frac{N+\alpha}{2N}} \\
\leq \bar{C}_{N} \left(\int_{\{x:|u(x)|<\delta\}} (|u|^{q+kp-p})^{\frac{2N}{N+\alpha}} \right)^{\frac{N+\alpha}{2N}} + \bar{C}_{N} \left(\int_{\{x:|u(x)|\geq\delta\}} (|u|^{p^{*}+kp-p}|)^{\frac{2N}{N+\alpha}} \right)^{\frac{N+\alpha}{2N}} \\
\leq \bar{C}_{N} \delta^{q-p} \left(\int_{\{x:|u(x)|<\delta\}} |u|^{\frac{kpp^{*}}{p^{*}_{\alpha}}} \right)^{\frac{p^{*}_{\alpha}}{p^{*}_{\alpha}}} + \bar{C}_{N} \left(\int_{\{x|u(x)|\geq\delta\}} |u|^{kp^{*}} \right)^{\frac{p^{*}_{\alpha}-p}{p^{*}_{\alpha}}} \left(\int_{\{x|u(x)|\geq\delta\}} |u|^{p^{*}} \right)^{\frac{p^{*}_{\alpha}-p}{p^{*}_{\alpha}}} \\
\leq \bar{C}_{N} \delta^{q-p} \|u\|_{\frac{kpp^{*}}{p^{*}_{\alpha}}}^{kp} + \bar{C}_{N} \left(\int_{\{x|u(x)|\geq\delta\}} |u|^{kp^{*}} \right)^{\frac{p^{*}_{\alpha}-p}{p^{*}_{\alpha}}} \left(\int_{\{x|u(x)|\geq\delta\}} |u|^{p^{*}} \right)^{\frac{p^{*}_{\alpha}-p}{p^{*}_{\alpha}}} \\
\leq \bar{C}_{N} \delta^{q-p} \|u\|_{\frac{kpp^{*}}{p^{*}_{\alpha}}}^{kp} + \bar{C}_{N} C(\delta)^{\frac{p^{*}_{\alpha}-p}{p^{*}_{\alpha}}} \|u\|_{kp^{*}}^{kp^{*}}, \tag{2.6}$$

and

$$\int_{\mathbb{R}^{N}} |u|^{p^{*}-2} u u_{\gamma}^{\beta} \leq \int_{\mathbb{R}^{N}} |u|^{p^{*}+\beta-1} = \int_{\mathbb{R}^{N}} |u|^{p^{*}+kp-p}
= \int_{\{x:|u(x)|<\delta\}} |u|^{p^{*}+kp-p} + \int_{\{x:|u(x)|\geq\delta\}} |u|^{p^{*}+kp-p}
\leq \left(\int_{\{x:|u(x)|<\delta\}} |u|^{\frac{kpp^{*}}{p^{*}_{\alpha}}}\right)^{\frac{p^{*}_{\alpha}}{p^{*}}} \left(\int_{\{x:|u(x)|<\delta\}} |u|^{\frac{(p^{*}-p)p^{*}}{p^{*}-p^{*}_{\alpha}}}\right)^{\frac{p^{*}-p^{*}_{\alpha}}{p^{*}}}
+ \left(\int_{\{x|u(x)|\geq\delta\}} |u|^{kp^{*}}\right)^{\frac{p}{p^{*}}} \left(\int_{\{x|u(x)|\geq\delta\}} |u|^{p^{*}}\right)^{\frac{p^{*}-p}{p^{*}-p^{*}_{\alpha}}}
\leq ||u||^{\frac{kp}{kpp^{*}}} \left(\int_{\{x:|u(x)|<\delta\}} |u|^{\frac{(p^{*}-p^{*}_{\alpha}+p^{*}_{\alpha}-p)p^{*}}{p^{*}-p^{*}_{\alpha}}}\right)^{\frac{p^{*}-p^{*}_{\alpha}}{p^{*}}} + C(\delta)^{\frac{p^{*}-p}{p^{*}}} ||u||^{kp}_{kp^{*}}.$$

$$\leq C'\delta^{p^{*}_{\alpha}-p} ||u||^{\frac{kp}{kpp^{*}}} + C(\delta)^{\frac{p^{*}-p}{p^{*}}} ||u||^{kp}_{kp^{*}}.$$

$$(2.7)$$

Using (2.6) and (2.7) in (2.5) we get :

$$\begin{aligned} \left\| u_{\gamma}^{k} \right\|_{p^{*}}^{p} & \leq C_{1} k^{p-1} \left(\bar{C}_{N} \delta^{q-p} + C' \delta^{p_{\alpha}^{*}-p} \right) \left\| u \right\|_{\frac{kpp^{*}}{p_{\alpha}^{*}}}^{kp} \\ & + C_{1} k^{p-1} \left(\bar{C}_{N} C(\delta)^{\frac{p_{\alpha}^{*}-p}{p^{*}}} + C(\delta)^{\frac{p^{*}-p}{p^{*}}} \right) \left\| u \right\|_{kp^{*}}^{kp}, \end{aligned}$$

taking $\delta > 1$ large enough so that $C'' := C_1 k^{p-1} \left(\bar{C}_N C(\delta)^{\frac{p_\alpha^* - p}{p^*}} + C(\delta)^{\frac{p^* - p}{p^*}} \right) < 1$. Thus, Fatou's lemma gives us the following :

$$||u||_{kp^*}^{kp} = ||u^k||_{p^*}^p \le \liminf_{\gamma \to \infty} ||u_{\gamma}^k||_{p^*}^p \le C_1 k^{p-1} \left(\bar{C}_N \delta^{q-p} + C' \delta^{p_{\alpha}^*-p} \right) ||u||_{\frac{kpp^*}{p_{\alpha}^*}}^{kp} + C'' ||u||_{kp^*}^{kp},$$

therefore,

$$||u||_{kp^*} \le \hat{C}^{\frac{1}{kp}} k^{\frac{p-1}{kp}} ||u||_{\frac{kpp^*}{p_{\alpha}^*}} \le k^{\frac{p-1}{kp}} \hat{C}^{\frac{1}{kp}} \left(1 + ||u||_{\frac{kpp^*}{p_{\alpha}^*}}^{kp}\right)^{\frac{1}{kp}}, \tag{2.8}$$

where $\hat{C} = \frac{C_1 k^{p-1} \left(\bar{C}_N \delta^{q-p} + C' \delta^{p_{\alpha}^* - p}\right)}{1 - C''} > 1$. If there exists a sequence $\{k_n\} \to \infty$ as $n \to \infty$ such that $\|u\|_{\frac{k_n p p^*}{p_{\alpha}^*}} \le 1$ for all $n \in \mathbb{N}$. Then, for any bounded domain $\Omega \subset \mathbb{R}^N$, we have

$$||u||_{L^{\frac{k_npp^*}{p_{\alpha}^*}}(\Omega)} \le ||u||_{\frac{k_npp^*}{p_{\alpha}^*}} \le 1 \text{ for all } n \in \mathbb{N},$$

taking $n \to \infty$, we get $u \in L^{\infty}(\Omega)$ and hence $u \in L^{\infty}_{loc}(\mathbb{R}^N)$. Now, suppose there does not exists any such sequence, then there exists $k_0 > 0$ such that

$$||u||_{\frac{kpp^*}{p_{\alpha}^*}}^{\frac{kpp^*}{p_{\alpha}^*}} = \int_{\mathbb{R}^N} |u|^{\frac{kpp^*}{p_{\alpha}^*}} > 1 \text{ for all } k \ge k_0,$$

and hence by (2.8) we get

$$||u||_{kp^*} \le (2\hat{C})^{\frac{1}{kp}} (k^{\frac{1}{k}})^{\frac{p-1}{p}} ||u||_{\frac{kpp^*}{p_*^*}} = (C_*^{\frac{1}{k}})^{\frac{1}{p}} (k^{\frac{1}{k}})^{\frac{p-1}{p}} ||u||_{\frac{kpp^*}{p_*^*}} \text{ for all } k \ge k_0,$$
 (2.9)

where $C_* = 2\hat{C} > 1$. For $k_1 = \frac{k_0 p_{\alpha}^*}{p} > k_0$, (2.9) gives us

$$||u||_{k_1p^*} \le (C_*^{\frac{1}{k_1}})^{\frac{1}{p}} (k_1^{\frac{1}{k_1}})^{\frac{p-1}{p}} ||u||_{k_0p^*},$$

similarly for $k_2 = \frac{k_1 p_{\alpha}^*}{p}$ we have:

$$||u||_{k_2p^*} \le (c_*^{\frac{1}{k_1} + \frac{1}{k_2}})^{\frac{1}{p}} (k_1^{\frac{1}{k_1}} k_2^{\frac{1}{k_2}})^{\frac{p-1}{p}} ||u||_{k_0p^*},$$

proceeding in this manner, for $k_n = \frac{k_{n-1}p_{\alpha}^*}{p} = (\frac{p_{\alpha}^*}{p})^n k_0$ we have:

$$||u||_{k_n p^*} \le \left(C_*^{\sum_{j=1}^n \frac{1}{k_j}}\right)^{\frac{1}{p}} \left(\prod_{j=1}^n k_j^{\frac{1}{k_j}}\right)^{\frac{p-1}{p}} ||u||_{k_0 p^*}. \tag{2.10}$$

Now, since $k_j > 1$ for all $j \in \mathbb{N}$, and $\lim_{j \to \infty} k_j^{\sqrt{\frac{1}{k_j}}} = 1$, there must exist some $C^* > 1$ such that $k_j^{\sqrt{\frac{1}{k_j}}} < c^*$ for all $j \in \mathbb{N}$, thus (2.10), we get:

$$\begin{aligned} \|u\|_{k_n p^*} & \leq \left(C_*^{\sum_{j=1}^n \frac{1}{k_j}} \right)^{\frac{1}{p}} \left(\left(C^* \right)^{\sum_{j=1}^n \frac{1}{\sqrt{k_j}}} \right)^{\frac{p-1}{p}} \|u\|_{k_0 p^*} \\ & \leq \left(C_*^{\frac{k_0 p}{p_\alpha^* - p}} \right)^{\frac{1}{p}} \left(\left(C^* \right)^{\frac{\sqrt{k_0 p}}{\sqrt{p_\alpha^*} - \sqrt{p}}} \right)^{\frac{p-1}{p}} \|u\|_{k_0 p^*} = \tilde{C} \|u\|_{k_0 p^*}, \end{aligned}$$

thus, for any bounded domain Ω in \mathbb{R}^N , we have:

$$||u||_{L^{k_n p^*}(\Omega)} \le ||u||_{k_n p^*} \le \tilde{C} ||u||_{k_0 p^*} \text{ for all } n \in \mathbb{N}.$$
 (2.11)

Now, since $k_n \to \infty$ as $n \to \infty$, then we must have $u \in L^{\infty}(\Omega)$, because, if not so, then there must exists a set $S \subset \Omega$ with positive measure and $\omega > 0$, such that

$$u(x) > \tilde{C} \|u\|_{k_0 p^*} + \omega$$
 for all $x \in S$,

and hence,

$$\lim_{n \to \infty} \|u\|_{L^{k_n p^*}(\Omega)} = \lim_{n \to \infty} \left(\int_{\Omega} |u|^{k_n p^*} \right)^{\frac{1}{k_n p^*}} \ge \lim_{n \to \infty} \left(\int_{S} |u|^{k_n p^*} \right)^{\frac{1}{k_n p^*}} \\
> \lim_{n \to \infty} \left(\int_{S} |\tilde{C}| \|u\|_{k_0 p} + \omega|^{k_n p^*} \right)^{\frac{1}{k_n p^*}} = (\tilde{C}| \|u\|_{k_0 p^*} + \omega) |S|^{\frac{1}{k_n p^*}} \\
= \tilde{C}| \|u\|_{k_0 p^*} + \omega,$$

but this is a contradiction to (2.11). Thus, $u \in L^{\infty}_{loc}(\mathbb{R}^N)$. Now, if u is radial, then by Strauss lemma [34] we have

$$|u(x)| \le \frac{K \|u\|_{W^{1,p}(\mathbb{R}^N)}}{|x|^{\frac{N-1}{p}}} \text{ almost everywhere in } \mathbb{R}^N.$$
 (2.12)

The estimate in (2.12) tells us that $|u(x)| \to 0$ as $|x| \to \infty$ and hence $u \in L^{\infty}(\mathbb{R}^N)$. Moreover, using (2.12) we can see that, $u \in L^t_{loc}(\mathbb{R}^N)$ for all $t \ge 1$ and $u \in L^t(\mathbb{R}^N)$ for all $t \ge p$. Now, as done in [11], one can see that $(I_{\alpha} * |u|^q) \in L^{\infty}(\mathbb{R}^N)$ and hence

$$f := \lambda |u|^{p-2}u + |u|^{p^*-2}u + (I_\alpha * |u|^q)|u|^{q-2}u \in L^t(\mathbb{R}^N) \text{ for all } t \ge \max\{\frac{p}{p-1}, \frac{p}{q-1}\}.$$

Thus, by theorem 1.4 of [10] $u \in C^{\delta}_{loc}(\mathbb{R}^N)$ for every $0 < \delta < \Theta = \min\{\frac{sp}{p-1}, 1\}$. \square Following the establishment of the regularity of a radial solution, we shall examine its existence. To proceed, we require the following preliminary results:

3 Preliminaries for existence results

Proposition 3.1. For all $u \in W^{1,p}(\mathbb{R}^N)$, we have:

$$A(u) \le C_N \|\nabla u\|_p^{2q\gamma_q} \|u\|_p^{2q(1-\gamma_q)}, \tag{3.1}$$

and

$$A(u) \le C_N[u]_{s,p}^{\frac{2q\gamma_q}{s}} \|u\|_p^{\frac{2q(s-\gamma_q)}{s}}.$$
 (3.2)

Proof. For r, t > 1 such that $\frac{1}{r} + \frac{1}{t} = 1 + \frac{\alpha}{N}$, by Proposition 1.1 and the interpolation inequality [9], we get:

$$\begin{split} A(u) & \leq \mu C(N,r,t) \|u^q\|_r \|u^q\|_t \\ & \leq \mu \bar{C}(N,r,t) \left(\|\nabla u\|_p^{\frac{N(qr-p)}{p}} \|u\|_p^{qr-\frac{N(qr-p)}{p}} \right)^{\frac{1}{r}} \left(\|\nabla u\|_p^{\frac{N(qt-p)}{p}} \|u\|_p^{qt-\frac{N(qt-p)}{p}} \right)^{\frac{1}{t}} \\ & = C_N \|\nabla u\|_p^{2q\gamma_q} \|u\|_p^{2q(1-\gamma_q)}, \end{split}$$

where $C_N := \mu \bar{C}(N, r, t)$. Similarly, by the fractional Gagliardo-Nirenberg inequality [27],

$$\begin{split} A(u) & \leq & \mu C(N,r,t) \, \|u^q\|_r \, \|u^q\|_t \\ & \leq & \mu \bar{C}(N,r,t) \left([u]_{s,p}^{\frac{N(qr-p)}{ps}} \, \|u\|_p^{qr-\frac{N(qr-p)}{ps}} \right)^{\frac{1}{r}} \left([u]_{s,p}^{\frac{N(qt-p)}{ps}} \, \|u\|_p^{qr-\frac{N(qt-p)}{ps}} \right)^{\frac{1}{t}} \\ & = & C_N[u]_{s,p}^{\frac{2q\gamma_q}{s}} \, \|u\|_p^{\frac{2q(s-\gamma_q)}{s}} \, , \end{split}$$

where $C_N := \mu \bar{C}(N, r, t)$.

Lemma 3.1. Let $N \geq 3$, $(u, \lambda) \in S(\tau) \times \mathbb{R}$ be a weak solution of (1.1)-(1.2), then $u \in P(\tau)$.

Proof. By, [1] and [22] we can see that u satisfies the Pohozaev identity:

$$\left(\frac{N-p}{p}\right) \|\nabla u\|_{p}^{p} + \left(\frac{N-sp}{p}\right) [u]_{s,p}^{p} = \frac{N\lambda}{p} \|u\|_{p}^{p} + \frac{N}{p^{*}} \|u\|_{p^{*}}^{p^{*}} + \left(\frac{N+\alpha}{2q}\right) A(u), \tag{3.3}$$

and since u is a solution of (1.1), we have:

$$||u||_{p}^{p} + [u]_{s,p}^{p} - \lambda ||u||_{p}^{p} - ||u||_{p^{*}}^{p^{*}} - A(u) = 0.$$
(3.4)

By (3.3) and (3.4) we get

$$M(u) = \|\nabla u\|_p^p + s[u]_{s,p}^p - \|u\|_{p^*}^{p^*} - \gamma_q A(u) = 0.$$

For $u \in W^{1,p}(\mathbb{R}^N)$ define

$$I_u(t) := E(t \star u) = \frac{e^{pt}}{p} \|\nabla u\|_p^p + \frac{e^{spt}}{p} [u]_{s,p}^p - \frac{e^{p^*t}}{p^*} \|u\|_{p^*}^{p^*} - \frac{e^{2q\gamma_q}}{2q} A(u).$$

Let us examine the critical points of I_u .

Lemma 3.2. For $u \in S(\tau)$, $t \in \mathbb{R}$ is a critical point of I_u if and only if $t \star u \in P(\tau)$.

Proof. For any $u \in S(\tau)$ we have:

$$I'_{u}(t) = e^{pt} \|\nabla u\|_{p}^{p} + se^{spt}[u]_{s,p}^{p} - e^{p^{*}t} \|u\|_{p^{*}}^{p^{*}} - \gamma_{q}e^{2q\gamma_{q}t}A(u) = M(t \star u),$$

and $||t \star u||_p^p = ||u||_p^p = \tau$. Therefore, $t \in \mathbb{R}$ is a critical point of I_u if and only if $t \star u \in P(\tau)$. \square

4 Case 1:
$$\frac{p}{2} \left(\frac{N+\alpha}{N} \right) < q < \bar{q}_s := \frac{p}{2} \left(\frac{N+\alpha+sp}{N} \right)$$

By (1.9) and Proposition 3.1, for $u \in S(\tau)$, we have:

$$E(u) \geq \frac{1}{p}T(u) - \frac{1}{p^*S^{\frac{p^*}{p}}}T(u)^{\frac{p^*}{p}} - \frac{C_N}{2q}T(u)^{\frac{2q\gamma_q}{p}}\tau^{\frac{2q(1-\gamma_q)}{p}}$$

$$= T(u)\left(\frac{1}{p} - \frac{1}{p^*S^{\frac{p^*}{p}}}T(u)^{\frac{p^*}{N}} - \frac{C_N}{2q}T(u)^{\frac{2q\gamma_q-p}{p}}\tau^{\frac{2q(1-\gamma_q)}{p}}\right). \tag{4.1}$$

Let us define

$$h_{\tau}(r) := \frac{1}{p} - \frac{1}{n^* S^{\frac{p^*}{p}}} r^{\frac{p^*}{N}} - \frac{C_N}{2q} r^{\frac{2q\gamma_q - p}{p}} \tau^{\frac{2q(1-\gamma_q)}{p}} \text{ for all } r > 0.$$
 (4.2)

and look for a subset $\Lambda(\tau)$ of $S(\tau)$, such that $\min_{u \in P(\tau)} E(u) = \min_{u \in \Lambda(\tau)} E(u)$, using the following property of h_{τ} .

Lemma 4.1. There exists $\tau_0 > 0$ such that, for every $\tau \in (0, \tau_0)$, $h_{\tau}(r)$ has unique global maximum at some $r_{\tau} > 0$, and there exist $r_0 > 0$ such that $h_{\tau}(r) \geq 0$ for all $r \in [\frac{\tau}{\tau_0} r_0, r_0]$.

Proof. Since, $2q\gamma_q - p < 0$, we can see that $h_\tau(r) \to -\infty$ as $r \to +\infty$, $h_\tau(r) \to -\infty$ as $r \to 0$, and

$$h'_{\tau}(r) = -\frac{1}{NS^{\frac{p^*}{p}}} r^{\frac{p^*}{N}-1} - \frac{C_N(2q\gamma_q - p)\tau^{\frac{2q(1-\gamma_q)}{p}}}{2qp} r^{\frac{2q\gamma_q}{p}-2},$$

then clearly,

$$r_{\tau} = \left(\frac{NC_N(p - 2q\gamma_q)S^{\frac{p^*}{p}}}{2pq}\right)^{\frac{p}{p^* - 2q\gamma_q}} \tau^{\frac{2q(1-\gamma_q)}{p^* - 2q\gamma_q}} > 0$$
(4.3)

is the global maxima for h_{τ} with

$$\max_{r \in \mathbb{R}} h_{\tau}(r) = h_{\tau}(r_{\tau}) = \frac{1}{p} - K\tau^{\frac{2q(1-\gamma_q)(p^*-p)}{p(p^*-2q\gamma_q)}}$$

where $K = \frac{C_N}{2q} \left(\frac{NC_N(p-2q\gamma_q)S^{\frac{p^*}{p}}}{2pq} \right) \left(1 + \frac{N(p-2q\gamma_q)S^{\frac{p^*}{p}}}{pp^*} \right) > 0$. Let $\tau_0 = (pK)^{\frac{N(2q\gamma_q-p^*)}{2q(1-\gamma_q)p^*}}$ and $r_0 = r_{\tau_0}$, then

$$\max_{r \in \mathbb{R}} h_{\tau_0}(r) = h_{\tau_0}(r_0) = \frac{1}{p} - K \tau_0^{\frac{2q(1 - \gamma_q)(p^* - p)}{p(p^* - 2q\gamma_q)}} = 0,$$

and hence, for every $\tau < \tau_0$

$$\max_{r \in \mathbb{R}} h_{\tau}(r) = h_{\tau}(r_{\tau}) = \frac{1}{p} - K\tau^{\frac{2q(1-\gamma_q)(p^*-p)}{p(p^*-2q\gamma_q)}} > \frac{1}{p} - K\tau^{\frac{2q(1-\gamma_q)(p^*-p)}{p(p^*-2q\gamma_q)}} = h_{\tau_0}(r_0) = 0.$$
 (4.4)

Now, for every r, define $\phi_r:(0,\tau_0)\to\mathbb{R}$ as $\phi_r(\tau):=h_\tau(r)$. Clearly, ϕ_r is non-increasing for every r, and hence $h_\tau(r_0)>h_{\tau_0}(r_0)=0$ for each $\tau\in(0,\tau_0)$. Also, since 2q-p>0, then for every $\tau<\tau_0$ we have:

$$h_{\tau}(\frac{\tau}{\tau_{0}}r_{0}) = \frac{1}{p} - \frac{(\tau r_{0})^{\frac{p^{*}}{N}}}{p^{*}\tau_{0}^{\frac{p^{*}}{N}}S^{\frac{p^{*}}{p}}} - \frac{C_{N}\tau^{\frac{2q-p}{p}}r_{0}^{\frac{2q\gamma_{q}-p}{p}}}{2q\tau_{0}^{\frac{2q\gamma_{q}-p}{p}}}$$

$$\geq \frac{1}{p} - \frac{r_{0}^{\frac{p^{*}}{N}}}{p^{*}S^{\frac{p^{*}}{p}}} - \frac{C_{N}\tau_{0}^{\frac{2q(1-\gamma_{q})}{p}}r_{0}^{\frac{2q\gamma_{q}-p}{p}}}{2q} = h_{\tau_{0}}(r_{0}) = 0.$$

Therefore, $h_{\tau}(r) \geq 0$ for all $r \in \left[\frac{\tau}{\tau_0} r_0, r_0\right]$.

The above lemma motivates us to define:

$$\Lambda(\tau) := \{ u \in S(\tau) : T(u) < r_0 \}.$$

$$\partial \Lambda(\tau) := \{ u \in S(\tau) : T(u) = r_0 \},$$

so, that we can study $m(\tau) := \min_{u \in \Lambda(\tau)} E(u)$. Considering the notations:

$$P(\tau)^+ := \{ u \in P(\tau) : E(u) < 0 \},$$

$$P(\tau)^- := \{ u \in P(\tau) : E(u) > 0 \},$$

we move forward to discuss some properties of $m(\tau)$ that will help us deduce the existence result.

Lemma 4.2. For every $\tau \in (0, \tau_0)$,

$$m(\tau) < 0 < \min_{u \in \partial \Lambda(\tau)} E(u). \tag{4.5}$$

Proof. For $u \in \partial \Lambda(\tau)$, by (4.1) and (4.4), we have:

$$E(u) \ge r_0 h_\tau(r_0) > r_0 h_{\tau_0}(r_0) = 0.$$

Now, since $2q\gamma_q < sp < p < p^*$, we get

$$E(t \star u) = \frac{e^{pt}}{p} \|\nabla u\|_p^p + \frac{e^{spt}}{p} [u]_{s,p}^p - \frac{e^{p^*t}}{p^*} \|u\|_{p^*}^{p^*} - \frac{e^{2q\gamma_q t}}{2q} A(u) \to 0^- \text{ as } t \to -\infty,$$

and hence, there exists $t_0 < -1$ such that $E(t_0 \star u) < 0$ and $T(t_0 \star u) < r_0$. Therefore, we get (4.5).

Lemma 4.3. $m(\tau) = \min_{u \in P(\tau)} E(u)$.

Proof. For $u \in S(\tau)$, as done in previous lemma we can see that, $I_u(t) \to 0^-$, $T(t \star u) \to 0$ as $t \to -\infty$ and $I_u(t) \to -\infty$ as $t \to +\infty$. Also, $I_u(t) > 0$ for all t such that $t \star u \in \partial \Lambda(\tau)$. Thus, I_u must have at least two crtical points, $\underline{t}_u \leq 0 \leq \overline{t}_u$ where \underline{t}_u corresponds to a local minima with $I_u(\underline{t}_u) < 0$ and \overline{t}_u to a local maxima with $I_u(\overline{t}_u) > 0$. Thus by Lemma 3.2, $\underline{t}_u \star u \in P(\tau)^-$ and $\overline{t}_u \star u \in P(\tau)^+$.

Claim 1: I_u has exactly two roots.

Let if possible, I_u has more than two roots, then g must attain C_q at more than two points where

$$g(t) := e^{(p-2q\gamma_q)t} \|\nabla u\|_p^p + se^{(sp-2q\gamma_q)t} [u]_{s,p}^p - e^{(p^*-2q\gamma_q)t} \|u\|_{p^*}^{p^*},$$

and $C_q = \gamma_q A(u)$. That is, g must have at least two critical points. Moreover, since every critical point of g is a root of:

$$f(t) := (p^* - 2q\gamma_q)e^{(p^* - sp)t} \|u\|_{p^*}^{p^*} - (p - 2q\gamma_q)e^{(p - sp)t} \|\nabla u\|_p^p - s(sp - 2q\gamma_q)[u]_{s,p}^p,$$

thus, f must have at least two roots. Now, let us examine the function f. We have:

$$f'(t) = (p^* - sp)(p^* - 2q\gamma_q)e^{(p^* - sp)t} \|u\|_{p^*}^{p^*} - (p - sp)(p - 2q\gamma_q)e^{(p - sp)t} \|\nabla u\|_{p}^{p},$$

and $t_0 = \frac{1}{p^* - p} \ln \left| \frac{(p - sp)(p - 2q\gamma_q) \|\nabla u\|_p^p}{(p^* - sp)(p^* - 2q\gamma_q) \|u\|_{p^*}^{p^*}} \right|$ is the unique critical point and global minima of f. Since $p^* > p$, we can see that $f(t) \to 0^-$ as $t \to -\infty$, thus $f(t_0) < 0$ and f can have at most one root at some $t > t_0$. This gives us a contradiction, hence I_u has exactly two roots \underline{t}_u and \overline{t}_u .

Claim 2 : $\underline{t}_u \star u \in \Lambda(\tau)$.

Let if possible $T(\underline{t}_u \star u) \geq r_0$. Now, since $T(t \star u) \to 0$ as $t \to -\infty$ and $t \mapsto T(t \star u)$ is non-decreasing, therefore, there exists some $t_0 < \underline{t}_u$ such that $T(t_0 \star u) = r_0$, that is $t_0 \star u \in \partial \Lambda(\tau)$. Thus, we get $I_u(t_0) = E(t_0 \star u) > 0$. But, since $t_0 < \underline{t}_u$ and I_u is non-increasing on the interval $(-\infty, \underline{t}_u]$ with $\sup_{t \in (-\infty, \underline{t}_u]} I_u(t) = \lim_{t \to -\infty} I_u(t) = 0^-$, thus $I_u(t_0) < 0$. Hence, by contradiction, $\underline{t}_u \star u \in \Lambda(\tau)$.

Now, for $u \in P(\tau)^-$ and t < 0, define $v(x) := e^{-\frac{Nt}{p}} u(e^{-t}x)$. Then $t \star v = u \in P(\tau)^- \subset P(\tau)$, hence $I'_v(t) = 0$. Thus, proceeding as in claim 2, we can see that $u = t \star v \in \Lambda(\tau)$. Therefore, $P(\tau)^- \subset \Lambda(\tau)$, which gives us:

$$m(\tau) = \min_{u \in \Lambda(\tau)} E(u) \le \min_{u \in P(\tau)^{-}} E(u) = \min_{u \in P(\tau)} E(u).$$

Also, if u is a minimizer of E on $\Lambda(\tau)$, then $I_u(t) \geq I_u(0)$ for all $t \leq \underline{t}_u$, since $t \star u \in \Lambda(\tau)$ for all $t \leq \underline{t}_u$. Moreover, since I_u is increasing in $[\underline{t}_u, \overline{t}_u]$, we get $I_u(\underline{t}_u) = I_u(0)$ and hence $\underline{t}_u = 0$. Therefore, $u = 0 \star u = \underline{t}_u \star u \in P(\tau)^-$. Thus,

$$m(\tau) = E(u) \ge \min_{u \in P(\tau)^{-}} E(u) = \min_{u \in P(\tau)} E(u).$$

Lemma 4.4. For every, τ_1 , $\tau_2 > 0$ such that $\tau_1 + \tau_2 = \tau \in (0, \tau_0)$, we have:

$$m(\tau) \le m(\tau_1) + m(\tau_2).$$

Further, if either $m(\tau_1)$ or $m(\tau_2)$ is achieved, then $m(\tau) < m(\tau_1) + m(\tau_2)$.

Proof. Without loss of generality, let $\tau_1 \leq \tau_2$. By definition of $m(\tau)$ and Lemma 4.2, we can find $u \in \Lambda(\tau_1)$ such that

$$E(u) < m(\tau_1) + \epsilon \text{ and } E(u) < 0. \tag{4.6}$$

Then, by (4.1), we get $0 > E(u) > T(u)h_{\tau_1}(T(u))$, hence by Lemma 4.1 and the fact that $u \in \Lambda(\tau_1)$, we get

$$T(u) < \frac{\tau_1}{\tau} r_0.$$

For $\delta = \frac{\tau_2}{\tau_1} \ge 1$, let us define $\bar{u}(x) := u(\delta^{-\frac{1}{N}}x)$, then

$$T(\bar{u}) = \delta(\delta^{-\frac{p}{N}} \|\nabla\|_p^p + \delta^{-\frac{sp}{N}} [u]_{s,p}^p) < \delta T(u) < \frac{\tau_2}{\tau} r_0 < r_0.$$

Thus, by (4.6): $m(\tau_2) \leq E(\bar{u}) \leq \delta E(u) < \delta(m(\tau_1) + \epsilon)$, since $\epsilon > 0$ is arbitrary, we get,

$$m(\tau_2) \le \frac{\tau_2}{\tau_1} m(\tau_1). \tag{4.7}$$

Also, since $\tau_2 \leq \tau_1 + \tau_2$, we can follow the same procedure to deduce

$$m(\tau) = m(\tau_1 + \tau_2) \le \frac{(\tau_1 + \tau_2)}{\tau_2} m(\tau_2)$$
 (4.8)

and hence by (4.7), $m(\tau) \leq m(\tau_1) + m(\tau_2)$. Further, if $m(\tau_1)$ is achieved, then we can take $\epsilon = 0$ in (4.6) will give us the required result. Similarly if $m(\tau_2)$ is achieved then we will get a strict inequality while proving (4.8) and hence the required result.

Lemma 4.5. The function $\phi:(0,\tau_0)\to\mathbb{R}$, defined as:

$$\phi(\tau) = m(\tau)$$

is continuous.

Proof. Let $\{\tau_n\} \to \tau$ in $(0, \tau_0)$. For every $n \in \mathbb{N}$, let $u_n \in \Lambda(\tau_n)$ be such that:

$$E(u_n) < m(\tau_n) + \frac{1}{n} \text{ and } E(u_n) < 0.$$
 (4.9)

Define $v_n := \left(\frac{\tau}{\tau_n}\right)^{\frac{1}{p}} u_n$, clearly, $\{v_n\}$ is a sequence in $S(\tau)$, and $T(v_n) = \frac{\tau}{\tau_n} T(u_n)$ for every $n \in \mathbb{N}$. Now, since $E(u_n) < 0$, then by (4.1) and Lemma 4.1, we get $T(u_n) < \frac{\tau_n}{\tau_0} r_0$, and hence

$$T(v_n) = \frac{\tau}{\tau_n} T(u_n) < \frac{\tau}{\tau_0} < r.$$

Therefore, $\{v_n\}$ is a sequence in $\Lambda(\tau)$. Then, by (4.9) we get

$$m(\tau) \leq E(v_n) = \frac{\tau}{p\tau_n} \|\nabla u_n\|_p^p + \frac{\tau}{p\tau_n} [u_n]_{s,p}^p - \left(\frac{\tau}{\tau_n}\right)^{\frac{p^*}{p}} \frac{\|u_n\|_{p^*}^p}{p^*} - \left(\frac{\tau}{\tau_n}\right)^{\frac{2q}{p}} \frac{A(u_n)}{2q}$$

$$= E(u_n) + \left(\frac{\tau}{\tau_n} - 1\right) \frac{\|\nabla u_n\|}{p} + \left(\frac{\tau}{\tau_n} - 1\right) \frac{[u_n]_{s,p}^p}{p} - \left(\left(\frac{\tau}{\tau_n}\right)^{\frac{p^*}{p}} - 1\right) \frac{\|u_n\|_{p^*}^p}{p^*}$$

$$- \left(\left(\frac{\tau}{\tau_n}\right)^{\frac{2q}{p}} - 1\right) \frac{A(u_n)}{2q} = E(u_n) + o_n(1)$$

$$< m(\tau_n) + \frac{1}{n} + o_n(1), \tag{4.10}$$

hence $m(\tau) \leq \liminf m(\tau_n)$. Now, let $\{w_n\} \subset \Lambda(\tau)$ be a minimizing sequence for $m(\tau)$ with $E(w_n) < 0$. Setting $\bar{w}_n = \left(\frac{\tau_n}{\tau}\right)^{\frac{1}{p}} w_n$, by (4.1) and Lemma 4.1 we get $\bar{w}_n \in \Lambda(\tau_n)$. Thus, as done in (4.10) we get:

$$m(\tau_n) \le E(\bar{w}_n) = E(w_n) + o_n(1) = m(\tau) + o_n(1),$$

hence $m(\tau) \ge \limsup m(\tau_n)$. Therefore, $\phi(\tau) = m(\tau) = \lim_{n \to \infty} m(\tau_n) = \lim_{n \to \infty} \phi(\tau_n)$ and since, $\tau \in (0, \tau_0)$ is arbitrary, ϕ is continuous.

Proposition 4.1. If $\{u_n\} \subset \Lambda(\tau)$ is a minimizing sequence for E on $\Lambda(\tau)$, then either of the following holds:

(1)
$$\limsup_{n \to \infty} \left(\sup_{z \in \mathbb{R}^N} \int_{B_1(z)} |u_n|^2 dx \right) = 0.$$

(2) There exists $u \in \Lambda(\tau)$ and a sequence $\{y_n\} \subset \mathbb{R}^N$ such that $u_n(.-y_n) \to u$ in $W^{1,p}(\mathbb{R}^N)$ as $n \to \infty$ and $E(u) = m(\tau)$.

Proof. Let $\{u_n\} \subset \Lambda(\tau)$ be such that $E(u_n) \to m(\tau)$ as $n \to \infty$. Suppose that (1) does not hold, then proving (2) will suffice. Now since $||u_n||_p^p = \tau > 0$ for all $n \in \mathbb{N}$, then we can find a sequence $\{y_n\}$ in \mathbb{R}^N such that

$$0 < \lim_{n \to \infty} \int_{B_1(0)} |u_n(x - y_n)|^p dx \le \tau. \tag{4.11}$$

Claim 1 : $\{u_n\}$ is bounded in $W^{1,p}(\mathbb{R}^N)$.

Since $\{u_n\} \subset \Lambda(\tau)$, we have $||u_n||^p = T(u_n) + ||u_n||_p^p < r_0 + \tau$ for all $n \in \mathbb{N}$. Hence, $\{u_n\}$ is bounded in $W^{1,p}(\mathbb{R}^N)$.

Now, let $u \in W^{1,p}(\mathbb{R}^N)$ be such that $\{u_n(.-y_n)\} \rightharpoonup u$, weakly in $W^{1,p}(\mathbb{R}^N)$, upto a subsequence. Then, we have the following:

$$\begin{cases}
\{u_n(.-y_n)\} \to u & \text{in } L^r_{loc}(\mathbb{R}^N) \text{ for all } r \in [p, p^*) \\
\{u_n(.-y_n)\} \to u & \text{almost everywhere in } \mathbb{R}^N.
\end{cases}$$
(4.12)

By (4.11) and (4.12), it is clear that $u \neq 0$ and hence $||u||_p^p > 0$. Setting $v_n := u_n(.-y_n) - u$, by Brezis Lieb lemma, we get

$$\begin{cases}
 \|v_n + u\|_p^p &= \|v_n\|_p^p + \|u\|_p^p + o_n(1) \\
 \|v_n + u\|_{p^*}^{p^*} &= \|v_n\|_{p^*}^{p^*} + \|u\|_{p^*}^{p^*} + o_n(1).
\end{cases}$$
(4.13)

Definig $B: W^{1,p}(\mathbb{R}^N) \to \mathbb{R}$ such that $B(u) := \frac{\|u\|^p}{p}$, then by (4.13), we get:

$$T(v_n + u) = T(v_n) + T(u) + o_n(1), (4.14)$$

also, by Brezis-lieb lemma for Reisz potential [23], we have:

$$A(v_n + u) = A(v_n) + A(u) + o_n(1). (4.15)$$

Hence, by (4.13)-(4.15) we get

$$E(u_n) = E(u_n(.-y_n)) = \frac{T(v_n + u)}{p} - \frac{\|v_n + u\|_{p^*}^{p^*}}{p^*} - \frac{A(v_n + u)}{2q}$$

$$= E(v_n) + E(u) + o_n(1). \tag{4.16}$$

Claim 2: $\limsup_{n\to\infty} \sup_{z\in\mathbb{R}^N} \int_{B_1(z)} |v_n|^p dx = 0.$

Let if possible

$$\limsup_{n \to \infty} \sup_{z \in \mathbb{R}^N} \int_{B_1(z)} |v_n|^p dx \neq 0,$$

then by boundedness of $\{v_n\}$, as done above, we can find a sequence $\{z_n\} \subset \mathbb{R}^N$ and $v \in W^{1,p}(\mathbb{R}^N)$ such that upto a subsequence

$$\{v_n(.-z_n)\} \to v$$
 weakly in $W^{1,p}(\mathbb{R}^N)$,
 $\{v_n(.-z_n)\} \to v$ in $L^r_{loc}(\mathbb{R}^N)$ for all $r \in [p, p^*)$,
 $\{v_n(.-z_n)\} \to v$ almost everywhere in \mathbb{R}^N .

Moreover, since $v \neq 0$, defining $w_n(x) := v_n(x - z_n) - v(x)$ and following the same procedure as done in (4.13)-(4.16):

$$\begin{cases}
 \|v_n\|_p^p &= \|v\|_p^p + \|w_n\|_p^p + o_n(1) \\
 T(v_n) &= T(v) + T(w_n) + o_n(1) \\
 E(v_n) &= E(v) + E(w_n) + o_n(1).
\end{cases}$$
(4.17)

Let $d := ||u||_p^p$, $b := ||v||_p^p$ and $b := \tau - d - b$. Since $\{u_n\} \subset \Lambda(\tau)$, then by (4.14), we have $T(u) \leq T(v_n + u) = T(u_n) < r_0$, thus $u \in \Lambda(d)$, similarly, $v \in \Lambda(b)$. Now, using (4.17) in (4.13) and (4.16) we get:

$$\begin{cases}
 \|u_n\|_p^p = \|v\|_p^p + \|w_n\|_p^p + \|u\|_p^p + o_n(1) \\
 E(u_n) = E(v) + E(w_n) + E(u) + o_n(1).
\end{cases}$$
(4.18)

Thus $0 \le \lim_{n \to \infty} ||w_n||_p^p = \tau - b - d = l$.

Case 1: l > 0. Clearly, by (4.17) and (4.14), one can see that $w_n \in \Lambda(\|w_n\|_p^p)$ for all $n \in \mathbb{N}$ and hence

$$E(w_n) \ge m(\|w_n\|_p^p).$$

Thus by (4.18), Lemma 4.4 and Lemma 4.5

$$m(\tau) = E(u_n) + o_n(1) = E(v) + E(w_n) + E(u) + o_n(1)$$

$$\geq E(v) + E(u) + m(\|w_n\|_p^p) + o_n(1) \geq m(b) + m(d) + m(\|w_n\|_p^p) + o_n(1)$$

$$= m(b) + m(d) + m(l) \geq m(\tau),$$

hence m(b) and m(d) are achieved by v and u respectively. Therefore, by Lemma 4.4 we get:

$$m(\tau) \ge m(b) + m(d) + m(l) > m(b+d) + m(l) \ge m(b+d+l) = m(\tau),$$

now this is a contradiction.

Case 2: l = 0.

Now, by (3.1) we have:

$$|A(w_n)| \le C_N \|\nabla w_n\|_p^{2q\gamma_q} \|w_n\|_p^{2q(1-\gamma_q)} \le C' \|w_n\|_p^{2q(1-\gamma_q)} \to 0 \text{ as } n \to \infty,$$

since $\{w_n\}$ is bounded in $W^{1,p}(\mathbb{R}^N)$ and $\lim_{n\to\infty}\|w_n\|_p^p=l=0$. Thus by (1.9) we get

$$E(w_n) = \frac{T(w_n)}{p} - \frac{\|w_n\|_{p^*}^{p^*}}{p^*} - \frac{A(w_n)}{2q} \ge T(w_n) \left(\frac{1}{p} - \frac{T(u)^{\frac{p^*-p}{p}}}{p^*S^{\frac{p^*}{p}}}\right) + o_n(1)$$

$$> T(w_n) \left(\frac{1}{p} - \frac{\frac{p^*-p}{p}}{p^*S^{\frac{p^*}{p}}}\right) + o_n(1) > T(w_n)h_{\tau_0}(r_0) + o_n(1) \text{ for every } n \in \mathbb{N},$$

and hence

$$\lim_{n \to \infty} E(w_n) \ge 0. \tag{4.19}$$

Proceeding as in case 1, we have:

$$m(\tau) = E(u_n) + o_n(1) \ge E(v) + E(u) + E(w_n) + o_n(1) \ge m(b) + m(d) \ge m(\tau), \quad (4.20)$$

since l = 0, (4.20) tells us that m(b) and m(d) are achieved and hence by Lemma 4.4 we have:

$$m(\tau) > m(b) + m(d) > m(b+d) = m(\tau).$$

This leads us to a contradiction. Thus, we have

$$\limsup_{n \to \infty} \sup_{z \in \mathbb{R}^N} \int_{B_1(z)} |v_n|^p dx = 0.$$

Therefore, by Lemma 1.1 of [18] we get $\{v_n\} \to 0$ in $L^r(\mathbb{R}^N)$ for all $r \in (p, p^*)$ and hence by Proposition 1.1 we have

$$A(v_n) \le \mu A_{\alpha} C_N \left(\int_{\mathbb{R}^N} |v_n|^{\frac{2Nq}{N+\alpha}} \right)^{\frac{N+\alpha}{N}} \to 0 \text{ as } n \to \infty, \text{ for all } q \in \left(\frac{p}{2} \left(\frac{N+\alpha}{N} \right), \bar{q_s} \right). \tag{4.21}$$

Claim 3: $\lim_{n\to\infty} ||v_n||_p^p = 0.$

Clearly, if $d = \tau$, then by (4.13), $||v_n||_p^p = ||u_n||_p^p - ||u||_p^p + o_n(1) = \tau - d + o_n(1) = o_n(1)$ and

we are done. Now, let if possible, $d \neq \tau$, without loss of generality, we can assume $d < \tau$. By, (4.14), one can see that $v_n \in \Lambda(\|v_n\|_p^p)$ and hence $E(v_n) \geq m(\|v_n\|_p^p)$ for all $n \in \mathbb{N}$. Thus, by (4.16), Lemma 4.5 and Lemma 4.4 we get:

$$m(\tau) = E(u_n) + o_n(1) = E(v_n) + E(u) + o_n(1) \ge m(\|v_n\|_p^p) + E(u) + o_n(1)$$

= $m(\tau - d) + E(u) + o_n(1)$ for all $n \in \mathbb{N}$,

and hence

$$m(\tau) \ge m(\tau - d) + E(u). \tag{4.22}$$

Since, $u \in \Lambda(d)$, we have $E(u) \ge m(d)$. Now, if E(u) > m(d), then by (4.22) and Lemma 4.4 we get

$$m(\tau) > m(\tau - d) + m(d) \ge m(\tau).$$

Therefore, by contradiction, we must have E(u) = m(d). Now, since m(d) is achieved, then again by (4.22) and Lemma 4.4 we get:

$$m(\tau) \ge m(\tau - d) + E(u) = m(\tau - d) + m(d) > m(\tau).$$

Thus, our assumption that $d < \tau$ was wrong, similarly we can see that $d > \tau$ cannot hold and hence $\lim_{n \to \infty} ||v_n||_p^p = 0$.

Claim 4: $\lim_{n\to\infty} T(v_n) = 0$.

Clearly, if claim 4 holds, then by claim 3 and 4, we can see that $\{v_n\} \to 0$ in $W^{1,p}(\mathbb{R}^N)$ and hence $u_n(.-y_n) \to u$. Thus by continuity of E, we will get $E(u) = m(\tau)$. Therefore, proving claim 4 will suffice.

Now, since $d = \tau$, we have $E(u) \ge m(\tau)$, also,

$$m(\tau) = E(u_n) + o_n(1) = E(u) + E(v_n) + o_n(1),$$

thus,

$$E(v_n) = m(\tau) - E(u) + o_n(1) \le o_n(1),$$

hence

$$\limsup_{n \to \infty} E(v_n) \le 0.$$
(4.23)

By Lemma 4.1 we know that $h_{\tau_0}(r_0) = 0$, also, $T(v_n) < r_0$, this implies

$$\frac{T(v_n)}{p} - \frac{\|v_n\|_{p^*}^{p^*}}{p^*} \ge \frac{T(v_n)}{p} - \frac{\|\nabla v_n\|_p^p}{p^* S^{\frac{p^*}{p}}} \ge T(v_n) \left(\frac{1}{p} - \frac{T(v_n)^{\frac{p^* - p}{p}}}{p^* S^{\frac{p^*}{p}}}\right) > \beta_0 T(v_n),$$

where

$$\beta_0 = \left(\frac{1}{p} - \frac{r_0^{\frac{p^* - p}{p}}}{p^* S^{\frac{p^*}{p}}}\right) = h_{\tau_0(r_0)} + \frac{C_N \tau_0^{\frac{2q(1 - \gamma_q)}{p}} r_0^{\frac{2q\gamma_q - p}{p}}}{2q} > 0.$$

Thus,

$$\limsup_{n \to \infty} \beta_0 T(v_n) \le \limsup_{n \to \infty} \left(\frac{T(v_n)}{p} - \frac{\|v_n\|_{p^*}^{p^*}}{p^*} \right) = \limsup_{n \to \infty} \left(E(v_n) + \frac{A(v_n)}{2q} \right) \le 0,$$

by (4.21) and (4.23). Therefore, $\lim_{n\to\infty}T(v_n)=0$ and we are done.

Using the above proposition and properties of $m(\tau)$, let us prove the main Existence result for $\frac{p}{2}\left(\frac{N+\alpha}{N}\right) < q < \bar{q}_s$.

Proof of Theorem 1.2: Let $\{u_n\} \subset \Lambda(\tau)$ be a minimizing sequence for $m(\tau)$. By Lemma 4.2 we have $m(\tau) < 0$, now if $\limsup_{n \to \infty} \left(\sup_{z \in \mathbb{R}^N} \int_{B_1(z)} |u_n|^2 dx \right) = 0$, then as done in Proposition 4.1, we will get $\{u_n\} \to 0$ in $L^r(\mathbb{R}^N)$ for all $r \in (p, p^*)$ and hence $\{A(u_n)\} \to 0$ as $n \to \infty$. Further, proceeding as in (4.19), we will end up getting $m(\tau) \geq 0$, which is a contradiction. Thus

$$\limsup_{n \to \infty} \left(\sup_{z \in \mathbb{R}^N} \int_{B_1(z)} |u_n|^2 dx \right) \neq 0.$$

Now, Proposition 4.1 gives us the existence of a sequence $\{y_n\}$ in \mathbb{R}^N and $u_{\tau} \in \Lambda(\tau)$ such that

$$\{u_n(.-y_n)\} \to u_\tau \text{ in } W^{1,p}(\mathbb{R}^N) \text{ as } n \to \infty,$$

and $E(u) = m(\tau)$. Now, let u_{τ}^* be the symmetric decreasing rearrangement of u_{τ} . Using classical rearrangement inequalities [12], we have :

$$[u_{\tau}^{*}]_{s,p}^{p} = \int_{\mathbb{R}^{N}} \int_{\mathbb{R}^{N}} \frac{|u_{\tau}^{*}(x) - u_{\tau}^{*}(y)|^{p}}{|x - y|^{N + sp}} dx dy = \int_{\mathbb{R}^{N}} \frac{1}{|z|^{N + sp}} \left(\int_{\mathbb{R}^{N}} |u_{\tau}^{*}(x + z) - u_{\tau}^{*}(x)|^{p} dx \right) dz$$

$$\leq \int_{\mathbb{R}^{N}} \frac{1}{|z|^{N + sp}} \left(\int_{\mathbb{R}^{N}} \frac{|u_{\tau}(x + z) - u_{\tau}(x)|^{p}}{|x - y|^{N + sp}} dx \right) dz = [u_{\tau}]_{s,p}^{p},$$

 $\|u_{\tau}^*\|_{p^*}^{p^*} = \|u_{\tau}\|_{p^*}^{p^*}$; $\|\nabla u_{\tau}^*\|_{p}^{p} \le \|\nabla u_{\tau}\|_{p}^{p}$ and $A(u_{\tau}^*) \ge A(u_{\tau})$. Therefore, $T(u_{\tau}^*) \le T(u_{\tau}) < r_0$, hence $E(u_{\tau}^*) \ge m(\tau)$, since $u_{\tau}^* \in \Lambda(\tau)$. Thus,

$$m(\tau) \le E(u_{\tau}^*) = \frac{T(u_{\tau}^*)}{p} - \frac{\|u_{\tau}^*\|_{p^*}^{p^*}}{p^*} - \frac{A(u_{\tau}^*)}{2a} \le \frac{T(u_{\tau})}{p} - \frac{\|u_{\tau}\|_{p^*}^{p^*}}{p^*} - \frac{A(u_{\tau})}{2a} = m(\tau).$$

Above calculations tell us that $E(u_{\tau}^*) = m(\tau)$, and hence $m(\tau)$ is achieved by a radially symmetric function in $W^{1,p}(\mathbb{R}^N)$. Moreover, since $E(u_{\tau}^*) = m(\tau) < 0$ and u_{τ}^* solves (1.1) for $\lambda = \lambda_{\tau}$, we get

$$\lambda_{\tau} \|u_{\tau}^{*}\|_{p}^{p} = T(u_{\tau}^{*}) - A(u_{\tau}^{*}) - \|u_{\tau}^{*}\|_{p^{*}}^{p^{*}} = pm(\tau) + \frac{pA(u_{\tau}^{*})}{2q} + \frac{p\|u_{\tau}^{*}\|_{p^{*}}^{p^{*}}}{p^{*}} - A(u_{\tau}^{*}) - \|u_{\tau}^{*}\|_{p^{*}}^{p^{*}}$$

$$= pm(\tau) + \left(\frac{p}{2q} - 1\right)A(u_{\tau}^{*}) + \left(\frac{p}{p^{*}} - 1\right)\|u_{\tau}^{*}\|_{p^{*}}^{p^{*}} < 0,$$

therefore, $\lambda_{\tau} < 0$.

5 Case 2: $\bar{q}_s \leq q < \frac{p}{2} \left(\frac{N+\alpha}{N-p} \right)$

Let $S_r(\tau) = S(\tau) \cap W_r^{1,p}(\mathbb{R}^N)$, $P_r(\tau) = P(\tau) \cap W_r^{1,p}(\mathbb{R}^N)$ and $m_r(\tau) = \inf_{u \in P_r(\tau)} E(u)$. Define $\Phi: W^{1,p}(\mathbb{R}^N) \times \mathbb{R} \to \mathbb{R}$ as

$$\Phi(u,t) := E(t \star u) = \frac{e^{pt}}{n} \|\nabla u\|_p^p + \frac{e^{spt}}{n} [u]_{s,p}^p - \frac{e^{p^*t}}{n^*} \|u\|_{p^*}^{p^*} - \frac{e^{2q\gamma_q t}}{2a} A(u),$$

clearly, $\Phi \in C^1$.

Lemma 5.1. Let $\tau > 0$, then for any fixed $u \in S_r(\tau)$, we have:

1.
$$T(t \star u) \to 0$$
 and $\Phi(u, t) \to 0^+$ as $t \to -\infty$,

2.
$$T(t \star u) \to +\infty$$
 and $\Phi(u,t) \to -\infty$ as $t \to +\infty$.

In particular, for $q = \bar{q}_s$, the above result holds for all $\tau \in (0, \bar{\tau}_s)$ where $\bar{\tau}_s = (\frac{2\bar{q}_s}{pC_N})^{\frac{N}{\alpha+sp}}$.

Proof. Since,

$$T(t \star u) = e^{pt} \|\nabla u\|_p^p + e^{spt} [u]_{s,p}^p,$$

clearly, $T(t \star u) \to 0$ as $t \to -\infty$ and $T(t \star u) \to +\infty$ as $t \to +\infty$. For $q = \bar{q}_s$, by (3.2) we have

$$\begin{split} \Phi(u,t) &= \frac{e^{pt}}{p} \|\nabla u\|_p^p + \frac{e^{spt}}{p} [u]_{s,p}^p - \frac{e^{p^*t}}{p^*} \|u\|_{p^*}^{p^*} - \frac{e^{spt}}{2\bar{q}} A(u) \\ &\geq \frac{e^{spt}}{p} [u]_{s,p}^p - \frac{e^{spt}C_N}{2\bar{q}} [u]_{s,p}^p \tau^{\frac{\alpha+sp}{N}} - \frac{e^{p^*t}}{p^*} \|u\|_{p^*}^{p^*} \\ &= e^{spt} [u]_{s,p} \left(\frac{1}{p} - \frac{C_N \tau^{\frac{\alpha+sp}{N}}}{2\bar{q}} \right) - \frac{e^{p^*t}}{p^*} \|u\|_{p^*}^{p^*}. \end{split}$$

Therefore,

$$\Phi(u,t) \to \begin{cases}
0^+ & \text{as } t \to -\infty, \\
-\infty & \text{as } t \to +\infty,
\end{cases}$$

for all $\tau < \bar{\tau}_s$. Also, since for $\bar{q}_s < q < \frac{p}{2}(\frac{N+\alpha}{N-p})$, we have $sp < 2q\gamma_q$ and $p < p^*$ we get the required result.

Define

$$A_k := \{ u \in S_r(\tau) : T(u) < k \}$$

and

$$\partial A_k := \{ u \in S_r(\tau) : T(u) = k \}.$$

Lemma 5.2. There exists $k_2 > k_1 > 0$ such that

$$0 < \sup_{u \in A_{k_1}} E(u) < \inf_{u \in \partial A_{k_2}} E(u), \tag{5.1}$$

and E(u), M(u) > 0 for all $u \in A_{k_2}$, whenever $\bar{q}_s < q < \frac{p}{2}(\frac{N+\alpha}{N-p})$ with $\tau > 0$ and $q = \bar{q}_s$ with $\tau \in (0, \bar{\tau}_s)$.

Proof. For $q = \bar{q}_s$, by (3.2) and (1.9) we have:

$$E(u) \ge \frac{T(u)}{p} - \frac{T(u)^{\frac{p^*}{p}}}{p^* S^{\frac{p^*}{p}}} - \frac{C_N T(u) \tau^{\frac{\alpha + sp}{N}}}{2\bar{q}} = \frac{T(u)}{p} \left(1 - \frac{pC_N \tau^{\frac{\alpha + sp}{N}}}{2\bar{q}} \right) - \frac{T(u)^{\frac{p^*}{p}}}{p^* S^{\frac{p^*}{p}}},$$

and

$$M(u) \ge sT(u) - \frac{T(u)^{\frac{p^*}{p}}}{S^{\frac{p^*}{p}}} - \gamma_{\bar{q}} C_N T(u) \tau^{\frac{\alpha + sp}{N}} = s \left(1 - \frac{pC_N \tau^{\frac{\alpha + sp}{N}}}{2\bar{q}}\right) T(u) - \frac{T(u)^{\frac{p^*}{p}}}{S^{\frac{p^*}{p}}}.$$

Now, since $p^* > p$, then for $\tau < \bar{\tau}_s$, we can find $k_2 > 0$ small enough so that $E(u) \ge \rho > 0$ for all $u \in \partial A_{k_2}$, M(u), E(u) > 0 for all $u \in A_{k_2}$ and

$$\inf_{u \in \partial A_{k_2}} E(u) \ge \rho > 0.$$

Also, since

$$E(u) \le \frac{T(u)}{p},$$

we can find $0 < k_1 < k_2$ such that

$$0 < \sup_{u \in A_{k_1}} E(u) < \rho \le \inf_{u \in \partial A_{k_2}} E(u).$$

Now, for $\bar{q}_s < q < \frac{p}{2} \left(\frac{N+\alpha}{N-p} \right)$, we have:

$$E(u) \ge \frac{T(u)}{p} - \frac{T(u)^{\frac{p^*}{p}}}{p^* S^{\frac{p^*}{p}}} - \frac{C_N}{2q} T(u)^{\frac{2q\gamma_q}{ps}} \tau^{\frac{2q(1-\gamma_q)}{sp}},$$

$$M(u) \ge sT(u) - \frac{T(u)^{\frac{p^*}{p}}}{S^{\frac{p^*}{p}}} - C_N \gamma_q T(u)^{\frac{2q\gamma_q}{sp}} \tau^{\frac{2q(1-\gamma_q)}{sp}},$$

and

$$E(u) \le \frac{T(u)}{p}.$$

Since $p^* > p$ and $2q\gamma_q > sp$, following the same procedure as done for the case $q = \bar{q}_s$, we can find $0 < k_1 < k_2$ satisfying (5.1) and E(u), M(u) > 0 for all $u \in A_{k_2}$.

By Lemma 5.1 and Lemma 5.2, we can find $u_1, u_2 \in S_r(\tau)$ such that

$$T(u_1) \le k_1 < k_2 < T(u_2); \quad E(u_1) > 0 > E(u_2) \text{ and } M(u_2) < 0.$$
 (5.2)

Define

$$r(\tau) := \inf_{\eta \in \Gamma(\tau)} \max_{z \in [0,1]} E(\eta(z)),$$

where

$$\Gamma(\tau) = \{ \eta \in C([0,1], S_r(\tau)) : \eta(0) = u_1 \text{ and } \eta(1) = u_2 \},$$

and

$$\tilde{r}(\tau) := \inf_{\tilde{\eta} \in \tilde{\Gamma}(\tau)} \max_{z \in [0,1]} \Phi(\tilde{\eta}(z)),$$

where

$$\tilde{\Gamma}(\tau) = \{ \tilde{\eta} \in C([0,1], S_r(\tau) \times \mathbb{R}) : \tilde{\eta}(0) = (u_1, 0) \text{ and } \tilde{\eta}(1) = (u_2, 0) \}.$$

Clearly, $r(\tau) \geq \max\{E(u_1), E(u_2)\} := \sigma_{\tau}$.

Lemma 5.3. For any fixed $u \in S_r(\tau)$, the function $I_u(t)$ has a unique critical point $t_u \in \mathbb{R}$ such that $t_u \star u \in P(\tau)$ corresponding to the maxima of I_u . In particular, for $q = \bar{q}_s$, the above statement is true for all $\tau \in (0, \bar{\tau}_s)$.

Proof. Let $u \in S_r(\tau)$ and $q = \bar{q}_s$. By Lemma 3.2, we know that $t_u \in \mathbb{R}$ is a critical point of I_u if and only if $t_u \star u \in P_r(\tau)$. Now, since

$$I'_{u}(t) = e^{pt} \|\nabla u\|_{p}^{p} + se^{spt} [u]_{s,p}^{p} - e^{p^{*}t} \|u\|_{p^{*}}^{p^{*}} - \gamma_{\bar{q}_{s}} e^{spt} A(u),$$

if t_u is a critical point of I_u , then $g(t_u) = \gamma_{\bar{q}_s} A(u) = C_u$ where

$$g(t) = e^{(p-sp)t_u} \|\nabla u\|_p^p + s[u]_{s,p}^p - e^{(p^*-sp)t_u} \|u\|_{p^*}^{p^*}.$$

Claim: I_u has unique critical point.

Let if possible, I_u has more than one critical points. Now, by Lemma 5.1, we know that $I_u(t) \to 0^+$ as $t \to -\infty$ and $I_u(t) \to -\infty$ as $t \to +\infty$, one can easily see that in this case, I_u must have at least three critical. Then g must attain C_u at more than two points, hence g must have at least two critical points. But

$$t_0 = \frac{1}{p^* - p} \ln \left(\frac{p(1-s) \|\nabla u\|_p^p}{(p^* - sp) \|u\|_{p^*}^{p^*}} \right),$$

is the only critical point of g. Therefore, by contradiction, I_u has exactly one critical point t_u corresponding to its global maxima.

Similarly, for $\bar{q}_s < q < \frac{p}{2}(\frac{N+\alpha}{N-n})$, if I_u has more than one critical points, then \tilde{g} defined as:

$$\tilde{g}(t) = e^{(p-2q\gamma_q)} \|\nabla u\|_p^p + se^{(sp-2q\gamma_q)t} [u]_{s,p}^p - e^{(p^*-2q\gamma_q)t} \|u\|_{p^*}^{p^*}$$

must attain $\gamma_q A(u) = \tilde{C}_u$ at more than two points and hence have at least two critical points. That is,

$$\tilde{g}'(t) = e^{(p-2q\gamma_q)t} \left((p-2q\gamma_q) \|\nabla u\|_p^p + s(sp-2q\gamma_q)e^{(sp-p)t} [u]_{s,p}^p - (p^*-2q\gamma_q)e^{(p^*-p)t} \|u\|_{p^*}^p \right),$$

has atleast two roots, hence

$$f(t) = (p - 2q\gamma_q) \|\nabla u\|_p^p + s(sp - 2q\gamma_q)e^{(sp - p)t}[u]_{s,p}^p - (p^* - 2q\gamma_q)e^{(p^* - p)t} \|u\|_{p^*}^{p^*}$$

has at least one critical point. Now, if t_0 is a critical point of f, then we must have

$$(p^* - 2q\gamma_q) = \frac{s(sp - 2q\gamma_q)(sp - p)e^{(sp - p)t}[u]_{s,p}^p}{(p^* - p)e^{(p^* - p)t}\|u\|_{p^*}^{p^*}} > 0 \text{ for all } \bar{q} < q < \frac{p}{2} \left(\frac{N + \alpha}{N - p}\right),$$

that is,

$$2q\gamma_q < p^* \text{ for all } \bar{q}_s < q < \frac{p}{2} \left(\frac{N+\alpha}{N-p} \right),$$

hence we must have $\frac{N+\alpha}{N-p} < \frac{N+\alpha+p^*}{N}$, but since $p^* = \frac{N}{p}(p^*-p)$, we have:

$$(N-p)(p^* + N + \alpha) - N(N + \alpha) = N(p^* - p) - \alpha p - p^*p = -\alpha p < 0.$$

Therefore, f does not have any critical point and hence I_u has unique critical point, which corresponds to its maxima.

Lemma 5.4.
$$r(\tau) = \tilde{r}(\tau) = m_r(\tau)$$
.

Proof. Claim $1: r(\tau) = \tilde{r}(\tau)$.

For any $(\tilde{\eta}_1, \tilde{\eta}_2) = \tilde{\eta} \in \tilde{\Gamma}(\tau)$, define $g(z) := \tilde{\eta}_2(z) \star \tilde{\eta}_1(z)$ for every $z \in [0, 1]$. Clearly, $g \in \Gamma(\tau)$ and hence

$$r(\tau) \le \max_{z \in [0,1]} E(g(z)) = \max_{z \in [0,1]} \Phi(\tilde{\eta}(z)),$$

since, $\tilde{\eta} \in \Gamma(\tau)$ is arbitrary, we get $r(\tau) \leq \tilde{r}(\tau)$. On the other hand, for any $\eta \in \Gamma(\tau)$, we can take $\tilde{\eta} = (\tilde{\eta}_1, 0) \in \tilde{\Gamma}(\tau)$ and deduce that:

$$\tilde{r}(\tau) \leq \max_{z \in [0,1]} \Phi(\tilde{\eta}(z)) = \max_{z \in [0,1]} E(\eta(z)),$$

hence $\tilde{r}(\tau) \leq r(\tau)$. Thus we are done with claim 1.

Claim 2 : $r(\tau) = m_r(\tau)$.

For any $u \in P_r(\tau)$, by Lemma 5.1 we can find $t_1 < 0 < t_2$ such that $T(t_1 \star u) < k_1 < k_2 < T(t_2 \star u)$, $\Phi(u, t_2) < 0 < \Phi(u, t_1)$ and $M(t_2 \star u) = I'_u(t_2) < 0$. By the definition of u_1 and u_2 given in (5.2), we can take $u_1 = t_1 \star u$ and $u_2 = t_2 \star u$. Setting $\eta(z) := ((1 - z)t_1 + zt_2) \star u$ for all $z \in [0, 1]$, clearly $\eta \in \Gamma(\tau)$ and since $I'_u(0) = 0$ by Lemma 5.3 we get

$$r(\tau) \le \max_{z \in [0,1]} E(\eta(z)) = \max_{t \in [t_1, t_2]} E(t \star u) = \max_{t \in [t_1, t_2]} I_u(t) = I_u(0) = E(u). \tag{5.3}$$

as (5.3) is true for every $u \in P_r(\tau)$, we can deduce that $r(\tau) \leq m_r(\tau)$.

Now, if we define $\tilde{h}(z) := M(\tilde{\eta}_2(z) \star \tilde{\eta}_1(z))$ on [0,1], for some $(\tilde{\eta}_1, \tilde{\eta}_2) = \tilde{\eta} \in \tilde{\Gamma}(\tau)$, we get

$$\tilde{h}(0) = M(0 \star u_1) = M(u_1) > 0,$$

by Lemma 5.2, and

$$\tilde{h}(1) = M(0 \star u_2) = M(u_2) < 0.$$

Thus, there exists $z_0 \in (0,1)$ such that $\tilde{h}(z_0) = 0$ and hence $\tilde{\eta}_2(z_0) \star \tilde{\eta}_1(z_0) \in P_r(\tau)$. Therefore,

$$m_r(\tau) \le E(\tilde{\eta}_2(z_0) \star \tilde{\eta}_1(z_0)) = \Phi(\tilde{\eta}(z_0)) \le \max_{z \in [0,1]} \Phi(\tilde{\eta}(z_0)).$$

Since $\tilde{\eta} \in \tilde{\Gamma}(\tau)$ is arbitrary, we are done.

Lemma 5.5. For $u \in S_r(\tau)$, defining

$$T_u = \{ w \in W^{1,p}(\mathbb{R}^N) : \int_{\mathbb{R}^N} |u|^{p-2} uw = 0 \},$$

then we can find a sequence $\{u_n\} \subset S_r(\tau)$ such that

- $1 \lim_{n \to \infty} E(u_n) = m_r(\tau),$
- $2 \lim_{n \to \infty} M(u_n) = 0,$
- $3\lim_{n\to\infty} E'_{S_r(\tau)}=0$, that is $E'(u_n)(w)\to 0$ uniformly for all $w\in T_{u_n}$ with $||w||\leq 1$.

Proof. Following the proof of Lemma 2.3 of [14], we can find the sequence $\{(v_n, t_n)\}\subset S_r(\tau)\times\mathbb{R}$ such that

$$\{\Phi(v_n, t_n)\} \to \tilde{r}(\tau),\tag{5.4}$$

and

$$\left\|\Phi'_{S_r(\tau)\times\mathbb{R}}(v_n,t_n)\right\| \le \frac{2}{\sqrt{n}} \to 0 \text{ as } n \to \infty,$$

that is,

$$|\Phi'(v_n, t_n)(w_n)| \le \frac{2}{\sqrt{n}} \|w\|_{\mathbb{W}} \text{ for all } w_n \in \tilde{T}_{(v_n, t_n)}$$
 (5.5)

where $\tilde{T}_{(v_n,t_n)} = \{(w_1, w_2) \in S_r(\tau) \times \mathbb{R} : \int_{\mathbb{R}^N} |v_n|^{p-2} v_n w_1 = 0\}$. Set $u_n = t_n \star v_n \in S_r(\tau)$. Thus we get 1 by (5.4) and Lemma 5.4. Now, for any $w = (w_1, w_2) \in \mathbb{W}$, we have:

$$\Phi'(v_n, t_n)(w_1, w_2) = e^{pt_n} \int_{\mathbb{R}^N} |\nabla v_n|^{p-2} \nabla v \nabla w_1 + e^{pt_n} w_2 \|\nabla v_n\|_p^p + e^{spt_n} \ll v_n, w_1 \gg
+ sw_2 e^{spt_n} [v_n]_{s,p}^p - e^{p^*t_n} \int_{\mathbb{R}^N} |v_n|^{p^*-2} v_n w_1 - w_2 e^{p^*t_n} \|v_n\|_{p^*}^{p^*}
- e^{2q\gamma_q t_n} \mu \int_{\mathbb{R}^N} (I_\alpha * |v_n|^q) |v_n|^{q-2} v_n w_1 - \gamma_q w_2 e^{2q\gamma_q t_n} A(v_n),$$

taking w = (0,1) we get $w \in \tilde{T}_{(v_n,t_n)}$ and hence by (5.5)

$$|M(u_n)| = |M(t_n \star v_n)| = |e^{pt_n} \|\nabla v_n\|_p^p + se^{spt_n} [v_n]_{s,p}^p - e^{p^*t_n} \|v_n\|_{p^*}^{p^*} - \gamma_q e^{2q\gamma_q t_n} A(v_n)|$$

$$= |\Phi'(v_n, t_n)(w)| \le \frac{2}{\sqrt{n}} \to 0 \text{ as } n \to \infty.$$

Thus we are done with 2. Further, taking $w \in W^{1,p}(\mathbb{R}^N)$ such that $\int_{\mathbb{R}^N} |u_n|^{p-2}u_nw = 0$, we get:

$$E'(u_{n})(w) = \int_{\mathbb{R}^{N}} |\nabla u_{n}|^{p-2} \nabla u_{n} \nabla w + \ll u_{n}, w \gg -\int_{\mathbb{R}^{N}} |u_{n}|^{p^{*}-2} u_{n} w$$

$$-\mu \int_{\mathbb{R}^{N}} (I_{\alpha} * |u_{n}|^{q}) |u_{n}|^{q-2} u_{n} w$$

$$= e^{pt_{n}} \int_{\mathbb{R}^{N}} |\nabla v_{n}|^{p-2} \nabla v_{n} \nabla \tilde{w} + e^{spt_{n}} \ll v_{n}, \tilde{w} \gg -e^{p^{*}t_{n}} \int_{\mathbb{R}^{N}} |v_{n}|^{p^{*}-2} v_{n} \tilde{w}$$

$$-e^{2q\gamma_{q}t_{n}} \mu \int_{\mathbb{R}^{N}} (I_{\alpha} * |v_{n}|^{q}) |v_{n}|^{q-2} v_{n} \tilde{w} = \Phi'(v_{n}, t_{n})(\tilde{w}, 0),$$

where $\tilde{w}(x) = e^{-\frac{Nt_n}{p}} w(e^{-t_n}x)$. Clearly, $(\tilde{w}, 0) \in \tilde{T}_{(v_n, t_n)}$, therefore, by (5.5) we get:

$$|E'(u_n)(w)| = |\Phi'(v_n, t_n)(\tilde{w}, 0)| \le \frac{2}{\sqrt{n}} \|(\tilde{w}, 0)\|_{\mathbb{W}} \le \frac{C}{\sqrt{n}} \|w\| \text{ for large } n.$$

Since, $w \in T_{u_n}$ is arbitrary,

$$\sup\{E'(u_n)(w): w \in T_{u_n} \text{ with } ||w|| \le 1\} \le \frac{C}{\sqrt{n}} \to 0 \text{ as } n \to \infty.$$

Lemma 5.6. If $0 \neq m_r(\tau) < \frac{S^{\frac{N}{p}}}{N}$ and $\{u_n\}$ is the sequence deduced in Lemma 5.5, then for large enough $\mu > 0$, either of the following holds true:

a) Upto a subsequence, $\{u_n\} \rightharpoonup u$ weakly in $W^{1,p}(\mathbb{R}^N)$ but not strongly and u is a solution of (1.1) for some parameter $\lambda \in \mathbb{R}$ with

$$E(u) \le m_r(\tau) - \frac{S^{\frac{N}{p}}}{N}.$$

b) Upto a subsequence, $\{u_n\} \to u$ strongly in $W^{1,p}(\mathbb{R}^N)$ and u solves (1.1)-(1.2) for some $\lambda < 0$, with $E(u) = m_r(\tau)$.

In particular, for $q = \bar{q}_s$, we will take $\tau \in (0, \bar{\tau}_s)$ and for $\bar{q}_s < q \leq \bar{q} := \frac{p}{2}(\frac{N + \alpha + p}{N})$, we take $\tau < \bar{\tau}_q = (\frac{2qp^*}{NC_N(p^* - 2q\gamma_q)})^{\frac{p}{2q(1 - \gamma_q)}}$.

Proof. Let $\{u_n\}$ be the sequence deduced in Lemma 5.5.

Claim 1: $\{u_n\}$ is bounded.

For $q = \bar{q}_s$, since $E(u_n) \to m_r(\tau)$ and $M(u_n) \to 0$ then for large n, using (3.2) we have

$$m_{r}(\tau) + 1 \geq E(u_{n}) = E(u_{n}) - \frac{1}{p^{*}}M(u_{n}) + o_{n}(1)$$

$$= \left(\frac{1}{p} - \frac{1}{p^{*}}\right) \|\nabla u_{n}\|_{p}^{p} + \left(\frac{1}{p} - \frac{s}{p^{*}}\right) [u_{n}]_{s,p}^{p} - \left(\frac{1}{2\bar{q}_{s}} - \frac{\gamma_{\bar{q}_{s}}}{p^{*}}\right) A(u_{n}) + o_{n}(1)$$

$$= \frac{\|\nabla u_{n}\|_{p}^{p}}{N} + \left(\frac{1}{p} - \frac{s}{p^{*}}\right) [u_{n}]_{s,p}^{p} - \frac{(p^{*} - sp)}{2p^{*}\bar{q}_{s}} A(u_{n}) + o_{n}(1)$$

$$\geq \frac{\|\nabla u_{n}\|_{p}^{p}}{N} + \left(\frac{1}{p} - \frac{s}{p^{*}} - \left(\frac{(p^{*} - sp)}{2p^{*}\bar{q}_{s}}\right) C_{N} \tau^{\frac{\alpha + sp}{N}}\right) [u_{n}]_{s,p}^{p} \geq KT(u_{n}),$$

where $K = \min\{\frac{1}{N}, \frac{1}{p} - \frac{s}{p^*} - \left(\frac{(p^* - sp)}{2p^* \bar{q}_s}\right) C_N \tau^{\frac{\alpha + sp}{N}}\} > 0$ for all $\tau < \bar{\tau}_s$. Similarly, for $\bar{q}_s < q \leq \bar{q}$, using (3.1), for large n we get:

$$m_r(\tau) + 1 \ge E(u_n) = E(u_n) - \frac{1}{p^*} M(u_n) + o_n(1)$$

 $\ge \frac{T(u_n)}{N} + \frac{(2q\gamma_q - p^*)}{2qp^*} C_N \tau^{\frac{2q(1-\gamma_q)}{p}} T(u_n),$

since, $2q\gamma_q and <math>T(u_n) \ge 1$ for large n (if not, then directly $\{u_n\}$ becomes bounded). Hence $\{u_n\}$ is bounded, for $\tau < \bar{\tau}_q$, and for $\bar{q} < q < \frac{p}{2}(\frac{N+\alpha}{N-p})$, we have:

$$m_r(\tau) + 1 \ge E(u_n) = E(u_n) + \frac{M(u_n)}{p} + o_n(1)$$

 $\ge \left(\frac{1}{p} - \frac{1}{p^*}\right) \|u\|_{p^*}^{p^*} + \left(\frac{\gamma_q}{p} - \frac{1}{2q}\right) A(u_n) + o_n(1).$

This tells us that $\{\|u_n\|_{p^*}^{p^*}\}$ and $\{A(u_n)\}$ are bounded, thus

$$sT(u_n) \le \|\nabla u_n\|_p^p + s[u_n]_{s,p}^p = \|u_n\|_{p^*}^{p^*} + \gamma_q A(u_n) + o_n(1) < M,$$

for some M > 0. Therefore, $\{u_n\}$ is bounded in $S_r(\tau) \subset W_r^{1,p}(\mathbb{R}^N)$. Since $W_r^{1,p}(\mathbb{R}^N)$ is compactly imbedded in $L^t(\mathbb{R}^N)$ for all $t \in (p, p^*)$, we can find $u \in W_r^{1,p}(\mathbb{R}^N)$ such that upto a subsequence

$$\{u_n\} \longrightarrow u$$
, weakly in $W_r^{1,p}(\mathbb{R}^N)$,

$$\{u_n\} \to u$$
, strongly in $L^t(\mathbb{R}^N)$ for all $t \in (p, p^*)$, $\{u_n\} \to u$ almost everywhere in \mathbb{R}^N .

Following the proof of Lemma 2.5 of [14], one can easily see that, for every $w \in W^{1,p}(\mathbb{R}^N)$

$$E'(u_n)(w) - \lambda_n \int_{\mathbb{R}^N} |u_n|^{p-2} u_n w = o_n(1)$$
 (5.6)

where

$$\lambda_n = \frac{E'(u_n)(u_n)}{\|u_n\|_p^p} = \frac{1}{\tau} \left(\|\nabla u_n\|_p^p + [u_n]_{s,p}^p - \|u_n\|_{p^*}^{p^*} - A(u_n) \right).$$

Since $\{u_n\}$ is bounded in $W^{1,p}(\mathbb{R}^N)$, clearly, $\{\lambda_n\}$ must be bounded and convergent upto a subsequence to some $\lambda_{\tau} \in \mathbb{R}$.

Claim 2: $u \neq 0$.

Let if possible, u(x) = 0 for all $x \in \mathbb{R}^N$. Then by (1.4), we have

$$A(u_n) \le \mu C_N \left(\int_{\mathbb{R}^N} |u_n|^{\frac{2Nq}{N+\alpha}} \right)^{\frac{N+\alpha}{N}} \to \mu C_N \left(\int_{\mathbb{R}^N} |u|^{\frac{2Nq}{N+\alpha}} \right)^{\frac{N+\alpha}{N}} = 0, \tag{5.7}$$

and since, $M(u_n) = o_n(1)$, we get $\lim_{n \to \infty} (\|\nabla u_n\|_p^p + s[u_n]_{s,p}^p) = \lim_{n \to \infty} \|u_n\|_{p^*}^p$. Let $l \ge 0$ be such that

$$l = \lim_{n \to \infty} (\|\nabla u_n\|_p^p + s[u_n]_{s,p}^p) = \lim_{n \to \infty} \|u_n\|_{p^*}^{p^*}.$$

Then by (1.9), we get:

$$S \le \frac{l}{l^{\frac{p}{p^*}}} \Rightarrow l(Sl^{\frac{p}{p^*}-1} - 1) \le 0,$$

Therefore, either l = 0 or $l \ge S^{\frac{N}{p}}$. Now, if $l \ge S^{\frac{N}{p}}$, then by (5.7)

$$m_r(\tau) = E(u_n) + o_n(1) = E(u_n) - \frac{1}{p^*} M(u_n) + o_n(1)$$

$$\geq \frac{1}{N} \|\nabla u_n\|_p^p + \frac{s}{N} [u_n]_{s,p}^p + o_n(1) = \frac{l}{N} + o_n(1) \geq \frac{S^{\frac{N}{p}}}{N},$$

but we are given that $m_r(\tau) < \frac{S^{\frac{N}{p}}}{N}$, and if l = 0, then we will get $m_r(\tau) = \lim_{n \to \infty} E(u_n) = 0$. Thus, by contradiction, we must have $u \neq 0$. Also, since $\{u_n\} \to u$, we can see that,

$$\int_{\mathbb{R}^N} |\nabla u|^{p-2} \nabla u \nabla \zeta + \ll u, \zeta \gg = \lambda \int_{\mathbb{R}^N} |u|^{p-2} u \zeta + \int_{\mathbb{R}^N} |u|^{p^*-2} u \zeta + \mu \int_{\mathbb{R}^N} (I_\alpha * |u|^q) |u|^{q-2} u \zeta,$$

for all $\zeta \in C_c^{\infty}(\mathbb{R}^N)$. Thus, u is a weak solution of

$$-\Delta_p u + (-\Delta_p)^s u = \lambda |u|^{p-2} u + \mu (I_\alpha * |u|^q) |u|^{q-2} u + |u|^{p^*-2} u \text{ in } \mathbb{R}^N,$$
 (5.8)

and hence by Lemma 3.1 M(u) = 0. Now, let $v_n = (u_n - u) \rightharpoonup 0$, weakly in $W^{1,p}(\mathbb{R}^N)$, then as done in Proposition 4.1, we have:

$$\begin{cases}
 \|u_n\|_p^p &= \|v_n\|_p^p + \|u\|_p^p + o_n(1) \\
 \|u_n\|_{p^*}^{p^*} &= \|v_n\|_{p^*}^{p^*} + \|u\|_{p^*}^{p^*} + o_n(1) \\
 T_s(u_n) &= T_s(v_n) + T_s(u) + o_n(1) \\
 A(u_n) &= A(v_n) + A(u) + o_n(1)
\end{cases}$$
(5.9)

where $T_s(u) = \|\nabla u\|_p^p + s[u]_{s,p}^p$. Since, $\{v_n\} \to 0$ in $L^t(\mathbb{R}^N)$ for all $t \in (p, p^*)$, by (1.4) we get $A(v_n) = o_n(1)$, thus by (5.9) we have

$$o_n(1) = M(u_n) = T_s(u_n) - \|u_n\|_{p^*}^{p^*} - \gamma_q A(u_n)$$

= $T_s(v_n) - \|v_n\|_{p^*}^{p^*} - A(v_n) + M(u) = T_s(v_n) - \|v_n\|_{p^*}^{p^*} + o_n(1),$

that is, $T_s(v_n) = \|v_n\|_{p^*}^{p^*} + o_n(1)$. Let $l = \lim_{n \to \infty} T_s(v_n) = \lim_{n \to \infty} \|v_n\|_{p^*}^{p^*} \ge 0$, then again by (1.9) we have, either l = 0 or $l \ge S^{\frac{N}{p}}$. Now, if $l \ge S^{\frac{N}{p}}$, then by (5.9) and the fact that $A(v_n) = o_n(1)$, we will get

$$m_{r}(\tau) = \lim_{n \to \infty} E(u_{n}) = \lim_{n \to \infty} \left(\frac{T(u_{n})}{p} - \frac{\|u_{n}\|_{p^{*}}^{p^{*}}}{p^{*}} - \frac{A(u_{n})}{2q} \right)$$

$$= \lim_{n \to \infty} \left(\frac{T(v_{n})}{p} - \frac{\|v_{n}\|_{p^{*}}^{p^{*}}}{p^{*}} \right) + E(u) \ge \lim_{n \to \infty} \left(\frac{T_{s}(v_{n})}{p} - \frac{\|v_{n}\|_{p^{*}}^{p^{*}}}{p^{*}} \right) + E(u)$$

$$= \frac{l}{N} + E(u) \ge \frac{S^{\frac{N}{p}}}{N} + E(u).$$

Hence a) holds true, and with l=0 we get strong convergence of $\{\|\nabla v_n\|_p^p\}$, $\{[v_n]_{s,p}^p\}$ and $\{\|u_n\|_{p^*}^{p^*}\}$. Then $\lim_{n\to\infty} E'(u_n)(u_n) = E(u)(u)$, and hence

$$\lambda = \lim_{n \to \infty} \lambda_n = \lim_{n \to \infty} \frac{E'(u_n)(u_n)}{\tau} = \frac{E'(u)(u)}{\tau} = \frac{1}{\tau} \left(E'(u)(u) - M(u) \right) = \frac{(1-s)}{\tau} [u]_{s,p}^p + \frac{(\gamma_q - 1)}{\tau} \mu \mathcal{A}_q(u) < 0,$$

for sufficiently large $\mu > 0$. Moreover, by (5.9), (5.6) and (5.8) we get

$$\lambda \|v_n\|_p^p = \lambda_n \|u_n\|_p^p - \lambda \|u\|_p^p + o_n(1) = E'(u_n)(u_n) - E'(u)(u) + o_n(1)$$

$$= -A(u_n) + A(u) + o_n(1) = -A(v_n) + o_n(1) = o_n(1).$$

Therefore $\{u_n\} \to u$ strongly in $W^{1,p}(\mathbb{R}^N)$ and hence b) holds.

Lemma 5.7. For all $\bar{q_s} \leq q < \frac{p}{2} \left(\frac{N+\alpha}{N-p} \right)$,

$$m_r(\tau) < \frac{S^{\frac{N}{p}}}{N}.\tag{5.10}$$

Proof. Let $\psi \in C_c^{\infty}(\mathbb{R}^N, [0, 1])$ be a radial cut-off function such that $\psi = 1$ in $B_1(0)$ and $\psi = 0$ in $\mathbb{R}^N \setminus B_2(0)$. For $\epsilon > 0$, define $u_{\epsilon}(x) := \psi(x)U_{\epsilon}(x)$ where U_{ϵ} is as defined in (1.10) with $x_0 = 0$, then by [1] we have:

$$\|\nabla u_{\epsilon}\|_{p}^{p} = K_{1} + O(\epsilon^{\frac{N-p}{p-1}}),$$
 (5.11)

$$[u_{\epsilon}]_{s,p}^{p} = O(\epsilon^{m_{N,p,s}}) \text{ where } m_{N,p,s} = \min\left\{\frac{N-p}{p-1}, p(1-s)\right\},$$
 (5.12)

$$\|u_{\epsilon}\|_{p^*}^{p^*} = K_2 + O(\epsilon^{-\frac{N}{p-1}}),$$
 (5.13)

with $S = K_1/K_2^{\frac{p}{p^*}}$. Since

$$||u_{\epsilon}||_{p}^{p} = \int_{B_{1}(0)} |U_{\epsilon}(x)|^{p} dx + \int_{\mathbb{R}^{N} \setminus B_{1}(0)} |\psi(x)|^{p} |U_{\epsilon}(x)|^{p} dx = \int_{B_{1}(0)} |U_{\epsilon}(x)| dx + O(\epsilon^{\frac{N-p}{p-1}}),$$

following the work of Brezis and Nirenberg in [7], we have:

$$||u_{\epsilon}||_{p}^{p} = \begin{cases} K_{3}\epsilon^{p} + O(\epsilon^{\frac{N-p}{p-1}}), & \text{for } N > p^{2}, \\ K_{4}\epsilon^{p}|\ln(\epsilon)| + O(\epsilon^{\frac{N-p}{p-1}}) & \text{for } N = p^{2}, \\ K_{5}\epsilon^{p} + O(\epsilon^{\frac{N-p}{p-1}}) & \text{for } p < N < p^{2}. \end{cases}$$
(5.14)

Also, a direct computation give us:

$$A(u_{\epsilon}) \ge K_6 \epsilon^{2\bar{q}_s(1-\gamma_{\bar{q}_s})}. \tag{5.15}$$

Set $v_{\epsilon}(x) := \frac{\tau^{\frac{1}{p}} u_{\epsilon}(x)}{\|u_{\epsilon}\|_{p}}$, clearly $v_{\epsilon} \in S_{r}(\tau)$ and hence by Lemma 5.3, there exists $t_{\epsilon} \in \mathbb{R}^{N}$ such that $t_{\epsilon} \star v_{\epsilon} \in P_{r}(\tau)$ and $I_{v_{\epsilon}}(t_{\epsilon}) = \max_{t \in \mathbb{R}} I_{v_{\epsilon}}(t)$. Considering the sequence $\{t_{\epsilon}\}$, one can observe that, if $\{t_{\epsilon}\} \to -\infty$ as $\epsilon \to 0$, then by Lemma 5.1, we must have

$$m_r(\tau) \le E(t_{\epsilon} \star v_{\epsilon}) \to 0^+ \text{ as } \epsilon \to 0,$$

thus, $m_r(\tau) \leq 0 < \frac{S^{N/p}}{N}$. Similarly, if $\{t_{\epsilon}\} \to +\infty$ as $\epsilon \to 0$, using Lemma 5.1, we get, $m_r(\tau) < 0 < \frac{S^{N/p}}{N}$. In fact, by Lemma 5.4, we have $m_r(\tau) = r(\tau) > 0$, thus the above cases are not possible, that is, $\{t_{\epsilon}\}$ must be bounded.

Now, let a < b, be such that such that $a < t_{\epsilon} < b$ for all $\epsilon > 0$. Since $M(t_{\epsilon} \star v_{\epsilon}) = 0$, we have

$$e^{(p^*-p)t_{\epsilon}} \|v_{\epsilon}\|_{p^*}^{p^*} = B_{\epsilon}(v_{\epsilon}) - \gamma_q e^{(2q\gamma_q - p)t_{\epsilon}} A(v_{\epsilon}),$$

where $B_{\epsilon}(u) = \|\nabla u\|_p^p + se^{(sp-p)t_{\epsilon}}[u]_{s,p}^p$. Thus, we have:

$$e^{pt_{\epsilon}} \le \left(\frac{B_{\epsilon}(v_{\epsilon})}{\|v_{\epsilon}\|_{p^{*}}^{p^{*}}}\right)^{\frac{p}{p^{*}-p}} = \frac{\|u_{\epsilon}\|_{p}^{p}}{\tau} \left(\frac{B_{\epsilon}(u_{\epsilon})}{\|u_{\epsilon}\|_{p^{*}}^{p^{*}}}\right)^{\frac{p}{p^{*}-p}}.$$

$$(5.16)$$

Now, define

$$g(t) := \frac{e^{pt} \|\nabla v_{\epsilon}\|_{p}^{p}}{p} - \frac{e^{p^{*}t} \|v_{\epsilon}\|_{p^{*}}^{p^{*}}}{p^{*}}$$

clearly g has global maxima at $t = t_0$ such that

$$e^{t_0} = \left(\frac{\|\nabla v_{\epsilon}\|_p^p}{\|v_{\epsilon}\|_{p^*}^{p^*}}\right)^{\frac{1}{p^*-p}},$$

with

$$\max_{t \in \mathbb{R}} g(t) = g(t_0) = \frac{1}{N} \left(\frac{\|\nabla v_{\epsilon}\|_{p}^{p}}{\|v_{\epsilon}\|_{p}^{p}} \right)^{\frac{p^*}{p^* - p}} = \frac{1}{N} \left(\frac{\|\nabla u_{\epsilon}\|_{p}^{p}}{\|u_{\epsilon}\|_{p}^{p}} \right)^{\frac{p^*}{p^* - p}} \\
= \frac{1}{N} S^{\frac{N}{p}} + O(\epsilon^{\frac{N - p}{p - 1}}), \tag{5.17}$$

by (5.11) and (5.13). This gives us:

$$E(t_{\epsilon} \star v_{\epsilon}) = g(t_{\epsilon}) + \frac{e^{spt_{\epsilon}}[v_{\epsilon}]_{s,p}^{p}}{p} - \frac{e^{2q\gamma_{q}t_{\epsilon}}A(v_{\epsilon})}{2q}$$

$$\leq \frac{1}{N}S^{\frac{N}{p}} + O(\epsilon^{\frac{N-p}{p-1}}) + \frac{e^{spt_{\epsilon}}[v_{\epsilon}]_{s,p}^{p}}{p} - \frac{e^{2q\gamma_{q}t_{\epsilon}}A(v_{\epsilon})}{2q}.$$
(5.18)

Now, for $N > p^2$, by (5.11)-(5.15) and (5.16) we have:

$$\frac{e^{pt_{\epsilon}}[v_{\epsilon}]_{s,p}^{p}}{A(v_{\epsilon})} = \frac{e^{pt_{\epsilon}} \|u_{\epsilon}\|_{p}^{2q-p} [u_{\epsilon}]_{s,p}^{p}}{\tau^{\frac{2q-p}{p}} A(u_{\epsilon})} \leq \left(\frac{e^{pt_{\epsilon}} \|u_{\epsilon}\|_{p}^{2q-p} [u_{\epsilon}]_{s,p}^{p}}{\tau^{\frac{2q-p}{p}} A(u_{\epsilon})}\right) \left(\frac{B_{\epsilon}(u_{\epsilon})}{\|u_{\epsilon}\|_{p^{*}}^{p^{*}}}\right)^{\frac{p}{p^{*}-p}} \frac{\|u_{\epsilon}\|_{p}^{p}}{\tau}$$

$$= \frac{\|u_{\epsilon}\|_{p}^{2q} [u_{\epsilon}]_{s,p}^{p} B_{\epsilon}(u_{\epsilon})^{\frac{p}{p^{*}-p}}}{\tau^{2q} A(u_{\epsilon})(\|u_{\epsilon}\|_{p^{*}}^{p^{*}})^{\frac{p}{p^{*}-p}}}$$

$$\leq \frac{K' \epsilon^{(m_{N,p,s}+2q\gamma_{q})} (K_{3} + O(\epsilon^{\frac{N-p}{p-1}}))^{2q} (K_{1} + O(\epsilon^{\frac{N-p}{p-1}} + O(\epsilon^{m_{N,ps}}))^{\frac{p}{p^{*}-p}}}{\tau^{2q} (K_{2} + O(\epsilon^{-\frac{N}{p-1}}))^{\frac{p}{p^{*}-p}}}$$

$$\to 0 \text{ as } \epsilon \to 0.$$

Therefore, $\frac{[v_{\epsilon}]_{s,p}^p}{A(v_{\epsilon})} \to 0$ as $\epsilon \to 0$, since, $0 < e^{ap} < e^{pt_{\epsilon}} < e^{bp}$. Thus, we can find $\epsilon_0 > 0$ small enough, such that

$$\frac{e^{spt_{\epsilon}}[v_{\epsilon}]_{s,p}^{p}}{p} - \frac{e^{2q\gamma_{q}t_{\epsilon}}A(v_{\epsilon})}{2q} < 0 \text{ for all } \epsilon < \epsilon_{0}.$$

Hence, by (5.18), we can conclude (5.10). Following the same procedure, one can easily deduce (5.10) for $p < N \le p^2$.

Proof of Theorem 1.3: Let $\{u_n\}$ be the sequence deduced in Lemma 5.5. Suppose a) of Lemma 5.6 holds, then there exists $u \in W^{1,p}(\mathbb{R}^N)$ such that $\{u_n\} \rightharpoonup u$ in $W^{1,p}(\mathbb{R}^N)$, and u solves (1.1) with

$$E(u) \le m_r(\tau) - \frac{S^{N/p}}{N} < 0,$$

by Lemma 5.7. Thus, by Lemma 5.1 and Lemma 5.3 we have, $I'_u(0) < 0$ and hence $M(u) = M(0 \star u) = I'_u(0) < 0$. But since u solve (1.1), we must have M(u) = 0. Therefore, b) of Lemma 5.6 holds and hence the required result.

References

- [1] P. d'Avenia, G Siciliano and M. Squassina, On fractional Choquard equations, Mathematical Models and Methods in Applied Sciences 25(2015), no. 08, 1447-76.
- [2] T. Bartsch and S. de Valeriola, Normalized solutions of nonlinear Schrödinger equations, Arch. Math. 100 (2013), 75-83.
- [3] T. Bartsch and L. Jeanjean, *Normalized solutions for nonlinear Schrödinger systems*, Proceedings of the Royal Society of Edinburgh Section A: Mathematics 148 (2018), no. 2, 225-242.

- [4] T. Bartsch, L. Jeanjean and N. Soave, Normalized solutions for a system of coupled cubic Schrödinger equations on \mathbb{R}^3 , Journal de Mathématiques Pures et Appliquées 106 (2016), no. 4, 583-614.
- [5] T. Bartsch and N. Soave, Multiple normalized solutions for a competing system of Schrödinger equations, Calculus of Variations and Partial Differential Equations 58 (2019), 1-24.
- [6] R. Biswas and S. Tiwari, Regularity results for Choquard equations involving fractional p-Laplacian, Mathematische Nachrichten 296(2023), no. 09, 4060-85.
- [7] H. Brézis and L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Communications on pure and applied mathematics 36(1983), no. 4, 437-77.
- [8] R. Filippucci and M. Ghergu, Singular solutions for coercive quasilinear elliptic inequalities with nonlocal terms, Nonlinear Analysis 197 (2020), 111857.
- [9] A. Fiorenza, M.R. Formica, T.G. Roskovec and F. Soudský, *Detailed proof of classi-cal Gagliardo-Nirenberg interpolation inequality with historical remarks*, Zeitschrift für Analysis und ihre Anwendungen 40(2021), no. 02, 217-36.
- [10] P. Garain and E. Lindgren Higher Hölder regularity for mixed local and nonlocal degenerate elliptic equations, Calculus of Variations and Partial Differential Equations 62(2023), no. 2, 67.
- [11] J. Giacomoni, N. Nidhi and K. Sreenadh, Normalized solution to a Choquard equation involving mixed local and non local operators, preprint.
- [12] J. Giacomoni, N. Nidhi and K. Sreenadh, Normalized solutions to a critical growth Choquard equation involving mixed operators, Asymptotic Analysis 2024(Preprint), 1-34.
- [13] T. Gou and L. Jeanjean Multiple positive normalized solutions for nonlinear Schrödinger systems, Nonlinearity 31 (2018), no. 5, 2319.
- [14] L. Jeanjean, Existence of solutions with prescribed norm for semilinear elliptic equations, Nonlinear Analysis: Theory, Methods and Applications 28(1997), no. 10, 1633-59.
- [15] S. Kesavan, *Topics in Functional Analysis and Applications*, New Age International Publishers, New Delhi, vol. 23 (2019).
- [16] C. Lei, M. Yang and B. Zhang, Sufficient and Necessary Conditions for Normalized Solutions to a Choquard Equation, The Journal of Geometric Analysis 33(2023), no. 4, 109.
- [17] E.H. Lieb, Existence and uniqueness of the minimizing solution of Choquard's nonlinear equation, Studies in Applied Mathematics 57 (1977), no. 2, 93-105.

- [18] P.L. Lions, The concentration-compactness principle in the calculus of variations. The locally compact case, part 2, InAnnales de l'Institut Henri Poincaré C, Analyse non linéaire 1(1984), No. 4, 223-283.
- [19] Z. Liu, V.D. Rădulescu, C. Tang and J. Zhang, Another look at planar Schrödinger-Newton systems, Journal of Differential Equations 328 (2022), 65-104.
- [20] Y. Meng and X. He, Normalized Solutions for the Fractional Choquard Equations with Hardy-Littlewood-Sobolev Upper Critical Exponent, Qualitative Theory of Dynamical Systems 23(2024), no. 1, 19.
- [21] V. Moroz and J. Van Schaftingen, Groundstates of nonlinear Choquard equations: existence, qualitative properties and decay asymptotics, Journal of Functional Analysis 265 (2013), no. 2, 153-184.
- [22] V Moroz and J. Van Schaftingen, Existence of groundstates for a class of nonlinear Choquard equations, Transactions of the American Mathematical Society 367(2015), no. 9, 6557-79.
- [23] V. Moroz and J. Van Schaftingen, A guide to the Choquard equation, Journal of Fixed Point Theory and Applications 19(2017), 773-813.
- [24] B. Noris, H. Tavares and G. Verzini, Stable solitary waves with prescribed L²-mass for the cubic Schrödinger system with trapping potentials, Discrete and Continuous Dynamical Systems 35 (2015), no. 12, 6085-6112.
- [25] B. Noris, H. Tavares and G. Verzini, Existence and orbital stability of the ground states with prescribed mass for the L^2 -critical and supercritical NLS on bounded domains, Analysis and PDE 7 (2015), no. 8, 1807-1838.
- [26] B. Noris, H. Tavares and G. Verzini, Normalized solutions for nonlinear Schrödinger systems on bounded domains, Nonlinearity 32 (2019), no. 3, 1044.
- [27] Y.J. Park, Fractional Gagliardo-Nirenberg inequality, Journal of the Chungcheong Mathematical Society 24(2011), no.3, 583-586.
- [28] S.I.Pekar, Untersuchungen über die Elektronentheorie der Kristalle, De Gruyter (1954).
- [29] B. Pellacci, A. Pistoia, G. Vaira and G. Verzini, Normalized concentrating solutions to nonlinear elliptic problems, Journal of Differential Equations 275 (2021), 882-919.
- [30] R. Penrose, On gravity's role in quantum state reduction, General relativity and gravitation 28 (1996), 581-600.
- [31] D. Pierotti and G. Verzini, Normalized bound states for the nonlinear Schrödinger equation in bounded domains, Calculus of Variations and Partial Differential Equations 56 (2017), 1-27.

- [32] X. Shang and P. Ma, Normalized solutions to the nonlinear Choquard equations with Hardy-Littlewood-Sobolev upper critical exponent, Journal of Mathematical Analysis and Applications 521(2023), no. 2, 126916.
- [33] X. Shen, Y. Lv and Z. Ou, Normalized Solutions to the Fractional Schrödinger Equation with Critical Growth, Qualitative Theory of Dynamical Systems 23 (2024), no. 3, 145.
- [34] W. Sickel, D. Yang and W. Yuan, The radial lemma of Strauss in the context of Morrey spaces. Annales Fennici Mathematici 39 (2014), 417–442.
- [35] N. Soave, Normalized ground states for the NLS equation with combined nonlinearities, Journal of Differential Equations 269 (2020), no. 9, 6941-6987.
- [36] G. Talenti, Best constant in Sobolev inequality, Annali di Matematica pura ed Applicata. 110(1976), 353-372.