CONVEX WAVES GRAZING CONVEX OBSTACLES TO HIGH ORDER

JIAN WANG AND MARK WILLIAMS

ABSTRACT. For large classes of convex incoming waves and convex obstacles, we verify the grazing sets are C^1 co-dimensional two submanifolds, and the reflected flow maps are diffeomorphisms in the illuminable region and homeomorphisms up to the shadow boundaries. We also construct explicit examples of convex obstacles with smooth boundaries, for which the grazing set for planar or spherical incident waves are not C^1 . As an application, we state a theorem on the propagation of semilinear wave oscillations grazing a convex obstacle to arbitrary high finite or infinite order in low regularity spaces.

Contents

1. Introduction	
2. Definitions, results, and applications	ć.
2.1. Incoming and reflected phases	4
2.2. Orders of grazing	7
2.3. Main results	(
2.4. Applications	10
3. Reflected flow maps for spherical incoming waves	11
4. The grazing set for spherical waves hitting a convex obsta	cle 17
4.1. Branching does not occur.	17
4.2. Examples of good grazing sets	21
4.3. Examples of bad grazing sets	25
5. The reflected flow map for general convex incoming waves	28
References	34

1. Introduction

In the paper [WW23] we provided a geometric optics description in spaces of low regularity of the transport of oscillations in solutions to linear and some semilinear second-order hyperbolic boundary problems along rays that graze the boundary of a convex obstacle to arbitrarily high finite or infinite order. The fundamental motivating example is the case where the spacetime manifold is $M = (\mathbb{R}^n \setminus \mathcal{O}) \times \mathbb{R}_t$, where $\mathcal{O} \subset \mathbb{R}^n$ is an open convex obstacle with C^{∞} boundary, and the governing hyperbolic operator is the wave operator $\square := \Delta - \partial_t^2$.

The main theorem of [WW23], which applies to quite general second-order hyperbolic operators, obstacles, and incoming waves, showed that high frequency exact solutions with high order grazing rays are well approximated in spaces of low regularity by approximate solutions constructed from fairly explicit solutions to simple profile equations. The theorem has two main assumptions, both of a geometric nature, which need to be verified for each choice of hyperbolic operator, obstacle, and incoming wave. The first is that the grazing set, that is, the set of points on the spacetime boundary at which incoming characteristics meet the boundary tangentially, is a codimension two, C^1 submanifold of spacetime. The second is that the reflected flow map, which sends points on the spacetime boundary forward in time to points on reflected and grazing rays, is injective and has appropriate regularity properties near the grazing set. These two assumptions are in general hard to verify.

In §2 we explain how to distinguish different orders of grazing. The orders may be $2,4,6,\ldots$ or ∞ . We showed in [WW23] that when the order is 2, the reflected flow map (RFM) and grazing set (GS) assumptions always hold for very general second order hyperbolic operators, obstacles, and incoming waves. When the order of grazing is higher than two, the GS assumption can fail even for \square , strictly convex obstacles, and incoming plane waves, and the RFM assumption becomes much harder to verify even in such cases. In [WW23] we identified large classes of strictly convex obstacles in all dimensions for which both assumptions are satisfied when incoming plane waves for \square graze the obstacle to arbitrarily high finite or infinite order.

Plane waves may be thought of as being produced by a point source at infinity. Much of this paper is concerned with verifying the RFM and GS assumptions for incoming spherical waves for \square , which may be viewed as arising from a point source at a finite distance from the obstacle. This is done in §§3–4. The incoming waves we consider in this paper are associated to phase functions of the form

$$\phi_i(t,x) = -t + \psi_i(x),$$

where $\psi_i: \Omega \to \mathbb{R}$, $\Omega \subset \mathbb{R}^n$, is a smooth convex function that satisfies the eikonal equation

$$(\partial_{x_1}\psi_i)^2 + |\nabla\psi_i|^2 = 1, \ \nabla := (\partial_{x_2}, \cdots, \partial_{x_n}).$$

These are the *convex waves* referred to in the title. When $\psi_i(x) = \theta \cdot x$ for $\theta \in S^{n-1}$ we obtain plane waves; when $\psi_i(x) = |x - b|$ we obtain spherical waves associated to a source at $b \in \mathbb{R}^n$. §5 verifies the RFM assumption for general convex incoming waves. Many examples of such waves are provided in Remark after Assumption 5.1.

While the RFM assumption turns out to hold for general convex waves and convex obstacles, this is not the case for the GS assumption. In §4 we verify the GS assumption for incoming spherical waves for a large class of convex obstacles, but we also present examples for which the assumption fails for incoming plane and spherical waves; in these examples the grazing set is cusped; roughly, it is the graph of a function that is continuous, but not C^1 . We also present examples for which the grazing set is C^1 but not twice differentiable, so the GS assumption holds, but just barely. It is perhaps surprising that these examples of nonsmooth grazing sets occur even in problems where the convex obstacle has C^{∞} boundary and the incoming waves are plane waves or spherical waves. Moreover, for a given obstacle the GS assumption can fail or not depending on the location of the source b of spherical waves with phase -t + |x - b|.

We think that the study of the reflected flow map and the grazing set are of independent interest, apart from the connection to the main theorem of [WW23]. So we have tried to write this paper so that it can be read independently of [WW23].

Remark on notations.

- 1. For the dot product of vectors $u, v \in \mathbb{R}^k$ we use both $u \cdot v$ and $\langle u, v \rangle$.
- 2. We denote $x = (x_1, \overline{x}) \in \mathbb{R}^{1+(n-1)}$. Except for Definition 2.1, we denote $\nabla = \nabla_{\overline{x}} = (\partial_{x_2}, \dots, \partial_{x_n})$.

2. Definitions, results, and applications

In this section we explain the RFM and GS assumptions. We begin by clarifying the definition of C^1 convex (concave) functions and strictly convex (concave) functions.

Definition 2.1. Let $U \subset \mathbb{R}^n$ be an open convex set.¹

(1) A C^1 function $f: U \to \mathbb{R}$ is said to be convex (or convex on U) if for any $z_1, z_2 \in U$,

$$f(z_2) - f(z_1) \ge \langle \nabla f(z_1), z_2 - z_1 \rangle; \tag{2.1}$$

here $\nabla f = (\partial_{x_1} f, \dots, \partial_{x_n} f)$. A C^1 function is said to be concave if \geq in (2.1) is replaced by \leq .

(2) A C^1 function $f: U \to \mathbb{R}$ is said to be strictly convex/convave, if (2.1) (resp. (2.1) with \leq) holds for all $z_1, z_2 \in U$ with equality holding if and only if $z_1 = z_2$.

¹This means that if $x, y \in U$, then the segment [x, y] is contained in U.

Remarks. 1. This definition appears to be standard. For example, it is the one given in [RV73]. For $f:U\to\mathbb{R}$ as above and any subset $V\subset U$, we say that f is convex/concave on V if it is convex/concave on U.

- 2. A C^2 function on U is convex/concave if and only if its Hessian matrix is positive/negative semi-definite.
- 3. If a C^2 function has a positive/negative definite Hessian matrix, then it is strictly convex/concave. The converse is not true.
- 4. The function, $f(z) = e^{-\frac{1}{|z|}}, z \neq 0$; f(0) = 0, is strictly convex but has vanishing Hessian at z=0. The function, $g(z)=|z|^a$, a>1, $a\in\mathbb{R}$, is C^1 and strictly convex.

Denote by $\mathcal{O} \subset \mathbb{R}^n$ a convex obstacle given by

$$\mathcal{O} := \{ (x_1, \overline{x}) | \ x_1 < F(\overline{x}), \ \overline{x} \in \mathbb{R}^{n-1} \}$$
 (2.2)

where $F: \mathbb{R}^{n-1} \to \mathbb{R}$ is a C^{∞} strictly concave function, and set $M:=(\mathbb{R}^n \setminus \mathcal{O}) \times \mathbb{R}_t$. We can arrange by translation and rotation of \mathcal{O} so that F has a max at $\overline{x} = 0$ with F(0) = 1. The tangent plane to $\partial \mathcal{O}$ at $x = (x_1, \overline{x}) = (1, 0)$ is then $x_1 = 1$.

2.1. Incoming and reflected phases. The "waves" that we consider are solutions to a continuation/reflection problem on $M_T := (\mathbb{R}^n \setminus \mathcal{O}) \times [-T, T]$ of the form²

$$\bigcap \Box u^{\epsilon} = f(x, t, u^{\epsilon}, \nabla_{x,t} u^{\epsilon}) \qquad \text{in } M_T, \tag{2.3a}$$

$$\begin{cases}
\Box u^{\epsilon} = f(x, t, u^{\epsilon}, \nabla_{x, t} u^{\epsilon}) & \text{in } M_{T}, \\
u^{\epsilon} = 0 & \text{on } \partial \mathcal{O} \times [-T, T], \\
u^{\epsilon} \sim u^{1}(x, t) + \epsilon U_{1}(x, t, \phi_{i}/\epsilon) & \text{in } M_{T} \cap \{t < -T + \delta\}.
\end{cases} (2.3a)$$
(2.3b)

$$\left(u^{\epsilon} \sim u^{1}(x,t) + \epsilon U_{1}(x,t,\phi_{i}/\epsilon) \quad \text{in } M_{T} \cap \{t < -T + \delta\} \right). \tag{2.3c}$$

Here the functions u^1 and $U_1(x,t,\theta_i)$ have (x,t)-support strictly away from ∂M , the function $U_1(x,t,\theta_i)$, called the incoming profile, is 2π -periodic in θ_i , and $\phi_i(x,t) =$ $-t + \psi_i(x)$, where $\psi_i : U \to \mathbb{R}$ is a smooth convex function that satisfies the eikonal equation

$$(\partial_{x_1}\psi_i)^2 + |\nabla\psi_i|^2 = 1$$

where $\nabla := (\partial_{x_2}, \cdots, \partial_{x_n})$, on some convex open set $U \subset \mathbb{R}^n$, $U \ni x = (1,0)$. We express condition (2.3c) by saying that the wave u^{ϵ} oscillates with the incoming phase ϕ_i in $t < -T + \delta$ (the past). The function f(x, t, p, q) is Lipschitz in the argument

The reflected phase ϕ_r needed to describe u^{ϵ} in the future is constructed using null bicharacteristics of \square . To describe these we let $\xi = (\xi_1, \overline{\xi}) \in \mathbb{R}^n$, $\tau \in \mathbb{R}$ and write the symbol of \square as $p(x,t,\xi,\tau)=\xi_1^2+|\overline{\xi}|^2-\tau^2$. Null bicharacteristics of \square are integral curves $\gamma: \mathbb{R} \to T^*M$ of the Hamiltonian vector field $H_p = 2\xi_1 \partial_{x_1} + 2\overline{\xi} \partial_{\overline{x}} - 2\tau \partial_t$ such

²The exact meaning of "~" in (2.3c) is not important for this paper and can be found in [WW23, Definition 1.3].

that $\gamma(s) \in p^{-1}(0)$ for all s. Thus, the components of $\gamma(s) = (x(s), t(s), \xi(s), \tau(s))$ satisfy

$$\dot{x}(s) = 2\xi, \ \dot{t}(s) = -2\tau, \ \dot{\xi}(s) = 0, \ \dot{\tau}(s) = 0; \quad |\xi(s)| = |\tau(s)|.$$

Note that null bicharacteristics are straight lines, and that the direction of (x(s), t(s)) is given by $(2\xi(0), -2\tau(0))$ for all s. Let us set

$$\xi_1^i(\overline{x}) := \partial_{x_1} \psi_i(F(\overline{x}), \overline{x}), \ \overline{\xi}^i(\overline{x}) := \partial_{\overline{x}} \psi_i(F(\overline{x}), \overline{x}).$$

For points \overline{x} such that

$$\xi_1^i(\overline{x}) - \nabla F(\overline{x}) \cdot \overline{\xi}^i(\overline{x}) \le 0 \tag{2.4}$$

we define the forward incoming null bicharacteristic associated to ϕ_i , which hits ∂M at $(F(\overline{x}), \overline{x}, t)$ when s = 0, to be³

$$\gamma_i(s; \overline{x}, t) = (F(\overline{x}) + 2s\xi_i^1(\overline{x}), \overline{x} + 2s\overline{\xi}^i(\overline{x}), t + 2s, \xi_i^1(\overline{x}), \overline{\xi}^i(\overline{x}), -1),$$

and the associated forward incoming characteristic to be $\pi_{(x,t)}\gamma_i$, the projection onto the (x,t) coordinates. In terms of ϕ_i the direction of $\pi_{(x,t)}\gamma_i$ is thus

$$2(\partial_x \phi_i(F(\overline{x}), \overline{x}, t), -\partial_t \phi_i(F(\overline{x}), \overline{x}).$$

In the case < the condition (2.4) means that $\pi_{(x,t)}\gamma_i$ enters $M^c := \mathbb{R}^{n+1} \setminus M$ for small s > 0, while in the case = the condition means that $\pi_{(x,t)}\gamma_i$ is tangent to ∂M at s = 0.

To identify reflected null bicharacteristics we introduce the inclusion map $i: \partial M \to M$, which induces the pullback map $i^*: \partial T^*M \to T^*\partial M$. If we use the map $(\overline{x}, t) \to (F(\overline{x}), \overline{x}, t)$ to parametrize ∂M , then i^* is given in coordinates by

$$i^*(F(\overline{x}), \overline{x}, t, \xi_1, \overline{\xi}, \tau) = (\overline{x}, t, \xi_1 \nabla F(\overline{x}) + \overline{\xi}, \tau).$$

Thus, the kernel of i^* is $N^*\partial M$, that is:

$$(\xi_1 \nabla F(\overline{x}) + \overline{\xi}, \tau) = 0 \iff (\xi_1, \overline{\xi}, \tau) \parallel (1, -\nabla F(\overline{x}), 0)$$

$$\Leftrightarrow (F(\overline{x}), \overline{x}, t, \xi_1, \overline{\xi}, \tau) \in N^* \partial M.$$
 (2.5)

Now set $\phi_0(\overline{x},t) := \phi_i(F(\overline{x}),\overline{x},t)$ and observe that γ_i "passes over" the point

$$(\overline{x}, t, \partial_{\overline{x}, t} \phi_0(\overline{x}, t)) \in T^* \partial M$$

in the sense that

$$i^*\gamma_i(0;\overline{x},t) = (\overline{x},t,\xi_1^i(\overline{x})\nabla F(\overline{x}) + \overline{\xi}^i(\overline{x}), -1) = (\overline{x},t,\partial_{\overline{x},t}\phi_0(\overline{x},t)) := \sigma(\overline{x},t).$$

From (2.5) we see that for \overline{x} such that (2.4) holds with <, there is another forward null bicharacteristic that passes over $\sigma(\overline{x}, t)$ at s = 0, namely

$$\gamma_r(s; \overline{x}, t) := (F(\overline{x}) + 2s\xi_1^r(\overline{x}), \overline{x} + 2s\overline{\xi}^r(\overline{x}), t + 2s, \xi_1^r(\overline{x}), \overline{\xi}^r(\overline{x}), -1),$$

³Here "forward" means that t increases as s increases. We sometimes suppress the (\overline{x}, t) dependence and write just $\gamma_i(s)$.

where $\xi^r(\overline{x})$ satisfies for some $c(\overline{x}) \neq 0$:

$$\xi^r(\overline{x}) - \xi^i(\overline{x}) = c(\overline{x})(1, -\nabla F(\overline{x})) \text{ and } |\xi^r(\overline{x})| = 1.$$
 (2.6)

The equations (2.6) are readily solved to yield

$$\xi_1^r = \xi_1^i - \frac{2}{1 + |\nabla F|^2} (\xi_1^i - \langle \overline{\xi}^i, \nabla F \rangle),$$

$$\overline{\xi}^r = \overline{\xi}^i + \frac{2\nabla F}{1 + |\nabla F|^2} (\xi_1^i - \langle \overline{\xi}^i, \nabla F \rangle).$$

Observe that equality holds in $(2.4) \Leftrightarrow \overline{\xi}^r(\overline{x}) = \overline{\xi}^i(\overline{x})$, the case where both $\pi_{(x,t)}\gamma_i$ and $\pi_{(x,t)}\gamma_r$ are tangent to ∂M at $(F(\overline{x}), \overline{x}, t)$. We call γ_r the reflected null bicharacteristic corresponding to γ_i . This discussion motivates the following definition.

Definition 2.2. (a) The grazing set and the illuminable region are respectively

$$G_{\phi_i} := \{ (F(\overline{x}), \overline{x}, t) | \langle \overline{\xi}^i, \nabla F \rangle - \xi_1^i = 0 \},$$

$$I_- := \{ (F(\overline{x}), \overline{x}, t) | \langle \overline{\xi}^i, \nabla F \rangle - \xi_1^i > 0 \}.$$

$$(2.7)$$

In (2.7) we require that $\overline{x} \in \overline{B(0,R)}$ and $t \in [-T,T]$ for some positive R,T that may vary from one example (that is, choice of \mathcal{O} and ϕ_i) to another.

(b) For $s_0 > 0$ the reflected flow map is the map⁴

$$Z_r: [0, s_0] \times (G_{\phi_i} \cup I_-) \to M \text{ where}$$

$$Z_r(s, \overline{x}, t) := \pi_{(x,t)} \gamma_r(s; \overline{x}, t) = (F(\overline{x}) + 2s\xi_1^r(\overline{x}), \overline{x} + 2s\overline{\xi}^r(\overline{x}), t + 2s).$$
(2.8)

With our normalization that F has a max at $\overline{x} = 0$ with F(0) = 1, we usually choose the incoming phase ϕ_i so that $(1,0,t) \in G_{\phi_i}$ for all t. The incoming characteristic $\pi_{(x,t)}\gamma_i(s;0,t)$ is then tangent to ∂M at (1,0,t).

Assumption 2.3 (Reflected flow map assumption (RFM)). There exist $s_0 > 0$ and positive constants R, T as in Definition 2.2 such that the map Z_r is a C^{∞} diffeomorphism from $[0, s_0] \times I_-$ onto its range, which extends to a homeomorphism from $[0, s_0] \times (G_{\phi_i} \cup I_-)$ onto its range.

Let us denote the ranges of Z_r on $[0, s_0] \times (G_{\phi_i} \cup I_-)$ and $[0, s_0] \times I_-$ by \mathcal{J}_r and \mathcal{J}'_r respectively. Write the map Z_r^{-1} defined on \mathcal{J}_r as

$$Z_r^{-1}(y_1, \overline{y}, t') = (s, \overline{x}, t) \in [0, s_0] \times (G_{\phi_i} \cup I_-).$$
 (2.9)

The method of characteristics (see [Wil22], for example) constructs the reflected phase ϕ_r on \mathcal{J}_r so that the following properties hold when (2.9) holds:

$$\phi_r(y_1, \overline{y}, t') = \phi_i(F(\overline{x}), \overline{x}, t), \ d\phi_r(y_1, \overline{y}, t') = (\xi^r(\overline{x}), -1). \tag{2.10}$$

⁴Here we use the map $(\overline{x},t) \to (F(\overline{x}),\overline{x},t)$ to identify $G_{\phi_i} \cup I_-$ with a subset of \overline{x},t space.

So ϕ_r is a characteristic phase (that is, $p(y, t', d\phi_r(y, t')) = 0$) such that both ϕ_r and $d\phi_r$ are constant along any particular reflected characteristic $\pi_{(x,t)}\gamma_r(s; \overline{x}, t)$.

Remark. When Assumption 2.3 holds, the formulas (2.10) show that

$$\phi_r \in C^{\infty}(\mathcal{J}'_r)$$
 but we just have $\phi_r \in C^1(\mathcal{J}_r)$.

In fact, ϕ_r generally fails to be in $C^2(\mathcal{J}_r)$; the singularity in ϕ_r is due to the singularity of Z_r^{-1} at $[0, s_0] \times G_{\phi_i}$.

2.2. Orders of grazing. To formulate the grazing set assumption GS, we first recall the classical decomposition

$$T^*\partial M \setminus 0 = E \cup H \cup G$$

into elliptic, hyperbolic, and glancing sets. If $\sigma \in T^*\partial M \setminus 0$, we say that σ belongs to E, H, or G if the number of elements in $(i^*)^{-1}(\sigma) \cap p^{-1}(0)$ is zero, two, or one respectively. The sets E and H are conic open subsets of $T^*\partial M \setminus 0$, and G is a closed conic hypersurface in $T^*\partial M \setminus 0$.

Below we let $\beta(x,t) := x_1 - F(\overline{x})$ denote the defining function of ∂M .

Definition 2.4 ([MS78]). Let $\sigma = (m, \nu) \in G$ and suppose $(i^*)^{-1}(\sigma) \cap p^{-1}(0) = \{\rho\}$, where $\rho \in T_m^*M$. We say $\sigma \in G^l$, the glancing set of order at least $l \geq 2$, if

$$p(\rho) = 0$$
 and $H_p^j \beta(\rho) = 0$ for $0 \le j < l$.

Thus, $G = G^2 \supset G^3 \supset \cdots \supset G^{\infty}$.

We say $\sigma \in G^l \setminus G^{l+1}$, the set of glancing points of exact order l, if $\sigma \in G^l$ and $H_p^l\beta(\rho) \neq 0$. Consider points $\sigma \in G^{2k} \setminus G^{2k+1}$, $k \geq 1$, such that $H_p^{2k}\beta(\rho) > 0$. When k = 1, such a point σ is a classical diffractive point as studied in [Mel75] or [Che96]. When $k \geq 1$ we refer to σ as a diffractive point of order 2k, and we write

$$\sigma \in G_d^{2k} \setminus G^{2k+1} \iff p(\rho) = 0, \ H_p^j \beta(\rho) = 0 \ \text{for} \ 0 \le j < 2k, \ \text{and} \ H_p^{2k} \beta(\rho) > 0.$$

Remarks. 1. If $\sigma \in G_d^{2k} \setminus G^{2k+1}$, let $\gamma(s)$ denote the bicharacteristic of p such that $\gamma(0) = \rho$. Then γ is tangent to ∂T^*M at ρ and lies $T^*\mathring{M}$ for small $s \neq 0$.

2. Gliding points of order 2k, $\sigma \in G_g^{2k} \setminus G^{2k+1}$, are defined as in (5.15) with the single change $H_p^{2k}\beta(\rho) < 0$. If $\sigma \in G^l \setminus G^{l+1}$ for some odd l, we call σ an inflection point of order l.

Definition 2.5 (Diffractive points of order ∞). Let $\sigma \in G^{\infty}$ and suppose $(i^*)^{-1}(\sigma) \cap p^{-1}(0) = \{\rho\}$. We say that σ is a diffractive point of order ∞ and write $\sigma \in G_d^{\infty}$ if the bicharacteristic $\gamma(s)$ of p such that $\gamma(0) = \rho$ lies T^*M for small $s \neq 0$.

Definition 2.6 (Glancing points of diffractive type). We denote by

$$\mathcal{G}_d := \cup_{k=1}^{\infty} \left(G_d^{2k} \setminus G^{2k+1} \right) \cup G_d^{\infty}$$

the set of glancing points of diffractive type.

For a fixed $t \in \mathbb{R}$ let $\rho = (1, 0, t, \xi_1^i(0), \overline{\xi}_i(0), -1) \in \partial T^*M \cap p^{-1}(0)$ and observe that

$$(1,0,t) \in G_{\phi_{i}} \iff \overline{\xi}_{i}(0) \cdot \nabla F(0) - \xi_{1}^{i}(0) = 0$$

$$\Leftrightarrow 0 = H_{p}\beta(\underline{\rho}) = 2\xi_{1}^{i}(0) - \nabla F(0) \cdot 2\overline{\xi}^{i}(0)$$

$$\Leftrightarrow i^{*}\underline{\rho} = (0,t,\xi_{1}^{i}(0)\nabla F(0) + \overline{\xi}^{i}(0),-1)$$

$$= (0,t,d\phi_{0}(0,t)) := \underline{\sigma} \in G.$$

$$(2.11)$$

We now state the grazing set assumption, letting β , (1,0,t), $\underline{\sigma}$ and $\underline{\rho}$ be as in (2.11) except that now we require $\underline{\sigma} \in \mathcal{G}_d$ (Definition 2.6); recall $\phi_0 = \phi_i \circ i : \partial M \to \mathbb{R}$.

Assumption 2.7 (Grazing set assumption (GS)). Suppose $\underline{\sigma} \in \mathcal{G}_d$. There exists an open set $U \ni (1,0,t)$ in $\mathbb{R}^n \times \mathbb{R}_t$ and a C^1 function $\zeta : U \to \mathbb{R}$ with $d\zeta \neq 0$ on U such that:

- (a) $G_{\phi_i} \cap U = \{(x,t) \in U : \zeta(x,t) = 0 \text{ and } \beta(x,t) = 0\};$
- (b) $H_p\zeta(\rho) \neq 0$;
- (c) There exists an open set $V \ni \underline{\sigma}$ in $T^*\partial M$ such that $G \cap Graph \ d\phi_0 \cap V \subset \mathcal{G}_d$.

Thus, the assumption requires that G_{ϕ_i} be a C^1 codimension two submanifold of $\mathbb{R}^n \cap \mathbb{R}_t$ near (1,0,t). In various examples of §4 where the assumption holds, condition (c) follows from convexity of \mathcal{O} , and the transversality condition (b) needs to be checked in each case.

For $\underline{\sigma} \in G$ as in (2.11) it will be convenient to have a simple reformulation of the condition that $\underline{\sigma} \in G_d^{2k} \setminus G^{2k+1}$ in terms of F. Let

$$\gamma_i(s;0,t) = (1 + 2s\xi_1^i(0), 2s\overline{\xi}^i(0), t + 2s, \xi_1^i(0), \overline{\xi}^i(0), -1)$$

be the incoming null bicharacteristic such that $\gamma_i(0;0,t)=\underline{\rho}$. From (2.11) we see that

$$\sigma \in G \Leftrightarrow 0 = H_p\beta(\underline{\rho}) = \frac{d}{ds}\beta(\gamma_i(s))|_{s=0}.$$

A direct computation using $H_p^j \beta(\gamma_i(s)) = \left(\frac{d}{ds}\right)^j \left[(1 + 2s\xi_1^i(0)) - F(2s\overline{\xi}^i(0)) \right]$ now shows that⁵

(a)
$$H_n^j \beta(\rho) = 0$$
 for $1 \le j < 2k$ and $H_n^{2k} \beta(\rho) > 0$ if and only if

(b)
$$\xi_1^i(0) - \nabla F(0) \cdot \overline{\xi}^i(0) = 0$$
, $\nabla^j F(0) (\overline{\xi}^i(0), \dots, \overline{\xi}^i(0))_j = 0$ for $2 \le j < 2k$ (2.12) and $\nabla^{2k} F(0) (\overline{\xi}^i(0), \dots, \overline{\xi}^i(0))_{2k} < 0$.

⁵Here the subscript j in (2.12)(b) indicates that $\overline{\xi}^i(0)$ occurs j times in the argument of the j-multilinear form $\nabla^j F(0)$.

2.3. Main results. §3 proves a strengthened, global version of the reflected flow map assumption (RFM) for spherical incoming waves.

Theorem 1. Let the convex obstacle $\mathcal{O} \subset \mathbb{R}^n$ be as (2.2), and consider incoming waves associated to the spherical phase function $\phi_i(x,t) = -t + |x-b|$, where $b \in \mathbb{R}^n \setminus \overline{\mathcal{O}}$. In the definition of the grazing set G_{ϕ_i} and the illuminable region I_- , Definition 3.2, allow R and T to be any positive numbers. Then the RFM assumption, Assumption 2.3, holds with s_0 any positive number.

The main result of §5 is a verification of the RFM assumption for general convex incoming waves.

Theorem 2. Let the convex obstacle $\mathcal{O} \subset \mathbb{R}^n$ be as (2.2), and consider incoming waves associated to any incoming phase function of the form $\phi_i(x,t) = -t + \psi_i$, where $\psi_i : \Omega \to \mathbb{R}$ is C^{∞} , convex, and satisfies

$$(\partial_{x_1}\psi_i)^2 + |\nabla\psi_i|^2 = 1$$

on some \mathbb{R}^n -open set $\Omega \ni (1,0)$. Then the RFM assumption, Assumption 2.3, holds with s_0 and T any positive numbers provided R is taken sufficiently small.⁶

Unlike the RFM assumption the GS assumption becomes more difficult to study and is more likely to fail as the dimension n of the obstacle increases.

In §4 we study properties of the grazing set when n=3 for both spherical and plane incoming waves, mostly spherical. One might hope that the GS assumption is always satisfied; after all, the obstacles are convex with C^{∞} boundaries and the incoming waves are spherical. But we provide examples to show this is not the case. We are mainly concerned in §4 to provide large classes of examples where the GS assumption holds.

Proposition 4.2 rules out branching for a class of convex obstacles and spherical incoming waves for which grazing can occur to any finite or infinite order at x = (1,0). Branching occurs when $\pi_x G_{\phi_i}$ consists of more than two smooth curves which meet and terminate at x = (1,0). In any such case the GS assumption fails of course. Even when branching does not occur, the GS assumption may fail when the set $\pi_{\overline{x}} G_{\phi_i}$ has a cusp; such examples are given in §4.3 for both spherical and planar incoming waves.

In Proposition 4.3 we write $\overline{x} = (x_2, x_3) = (u, v)$ and suppose that the obstacle \mathcal{O} as in (2.2) is defined by a function $F : \mathbb{R}^2 \to \mathbb{R}$ of the form

$$F(u,v) = 1 - (c_0 u^{2k} + c_1 u^{2k-2} v^2 + c_2 u^{2k-4} v^4 + \dots + c_{k-1} u^2 v^{2k-2} + c_k v^{2k})$$

$$:= 1 - G(u,v), \quad k \ge 1,$$
(2.13)

⁶Here again R and T are the parameters appearing in Definition 3.2.

where $\nabla^2 G(u,v) > 0$ for $(u,v) \neq 0$ and all $c_j > 0$. The proposition shows that when the incoming phase is taken to be $\phi_i(x,t) = -t + |x-b|$ with $b = (1,\overline{b}) = (1,-1,0)$, the GS assumption is satisfied with $\underline{\sigma} = (0,t,-\overline{b},-1) \in \mathcal{G}_d$. Using the formula for Fand (2.12) we see that $\underline{\sigma} \in G_d^{2k} \setminus G^{2k+1}$. Proposition 4.4 shows that the GS assumption continues to hold for $\phi_i(x,t) = -t + |x-b|$ when F as in (2.13) is modified by adding any finite number of polynomial terms of order > 2k + 1.

2.4. **Applications.** The results of this paper extend the range of application of the main result of [WW23]. [WW23, Theorem 2] gives an explicit geometric optics description of solutions to the initial boundary value problem (2.3a)–(2.3c) whenever the RFM and GS assumptions are satisfied by the obstacle \mathcal{O} and incoming phase ϕ_i . In [WW23] we verified these assumptions for some families of convex obstacles \mathcal{O} , but only for planar incoming waves associated to linear phases $\phi_i(x,t) = -t + \theta \cdot x$, $|\theta| = 1$. The results of §§3–4 of this paper show that both assumptions hold for certain families of convex obstacles with grazing points of arbitrarily high order, when the incoming phases are the nonlinear phases that describe spherical waves, $\phi_i(x,t) = -t + |x - b|$.

We now restate, somewhat informally, the main result of [WW23] for the problems considered here, referring the reader to that paper for a more precise and general formulation.

Theorem 3 ([WW23, Theorem 2]). Consider the initial boundary value problem (2.3a)–(2.3c) under the RFM assumption, Assumptions 2.3. Suppose the initial oscillatory profile $U_1(x,t,\theta_i)$ is supported near a characteristic of ϕ_i that grazes ∂M (possibly to high order) at the point $(x_1, \overline{x}, t) = (1,0,0) \in G_{\phi_i}$. Assume that the GS assumption, Assumption 2.7, is satisfied near $\underline{\sigma} = (0,0,d\phi_i(0,0)) \in \mathcal{G}_d$. Then if T > 0 is small enough, the solution $u^{\epsilon} \in H^1(M_T)$ satisfies

$$u^{\epsilon}(x,t)|_{M_T} \sim_{H^1} u(x,t) + \epsilon U_r(x,t,\phi_r/\epsilon) + \epsilon U_i(x,t,\phi_i/\epsilon). \tag{2.14}$$

Here $U_k(x,t,\theta_k)$ for k=r,i is periodic in θ_k with mean zero, and the functions

$$u \in H^1(M_T), \ \partial_{\theta_r} U_r \in L^2(M_T \times \mathbb{T}), \ \partial_{\theta_i} U_i \in L^2(M_T \times \mathbb{T})$$
 (2.15)

are constructed to satisfy the profile equations [WW23, (4.4)–(4.6)].

Remarks. 1. The meaning of " \sim_{H^1} " in (2.14) is explained in [WW23, Definition 1.3]. Roughly, it means that the leading part of u^{ϵ} has the given form up to an error that is small in H^1 when the wavelength ϵ is small.

⁷The condition $\nabla^2 G(u,v) > 0$ for $(u,v) \neq 0$ implies G(u,v) > 0 for $(u,v) \neq 0$. The property G(u,v) > 0 implies that c_0 and c_k are positive (so take $c_0 = 1$).

⁸When $\underline{\sigma} \in G_d^2 \setminus G^3$, the results of [Che96] and [WW23] show that the RFM and GS assumptions are satisfied for any incoming characteristic phase; moreover, the grazing set is C^{∞} , not just C^1 .

2. The profile equations referred to in the theorem are a coupled system for the functions u, U_r , U_i in (2.15) in which the equations for U_r , U_i describe transport along the characteristic vector fields associated to ϕ_r and ϕ_i :

$$T_{\phi_k} = 2\nabla_x \phi_k \partial_x - 2\partial_t \phi_k \partial_t, \quad k = r, i.$$

3. We have seen that the RFM assumption allows us to construct the reflected phase ϕ_r as a C^1 function on \mathcal{J}_r , the range of Z_r as in (2.8). The GS assumption implies that the union of the forward and backward flowouts of the grazing set G_{ϕ_i} under T_{ϕ_i} is a C^1 hypersurface in $\mathbb{R}^{n+1}_{x,t}$. We write this union as $SB = SB_+ \cup SB_-$ and refer to SB_+ , the forward flowout, as the "shadow boundary". From the profile equations [WW23, (4.4)–(4.6)] we can read off the property that $U_r(x,t,\theta_r)$ and $U_i(x,t,\theta_i)$ have their supports inside the set of points of M_T that can be reached by starting from points (x,t) in the support of the initial profile $U_1(x,t,\theta_i)$ as in (2.3c), then flowing forward along T_{ϕ_i} , and continuing along T_{ϕ_r} in any case where a characteristic of T_{ϕ_i} hits the boundary. With (2.14) this shows that the exact solution u^{ϵ} has no high frequency oscillations in the shadow region adjacent to SB_+ that are detectable in the H^1 norm.

3. Reflected flow maps for spherical incoming waves

As in $\S 2$, we assume \mathcal{O} is a convex obstacle given by

$$\mathcal{O} := \{ (x_1, \overline{x}) \mid x_1 < F(\overline{x}), \ \overline{x} \in \mathbb{R}^{n-1} \}$$

where $F: \mathbb{R}^{n-1} \to \mathbb{R}$ is a C^{∞} strictly concave function.⁹ We consider the following phase function for spherical incoming waves

$$\phi_i(x,t) = -t + |x-b|, \ b \in \mathbb{R}^n \setminus \overline{\mathcal{O}}.$$

We define

$$\rho(\overline{x}) := |(F(\overline{x}), \overline{x}) - b|$$

and

$$\alpha(\overline{x}) := (\alpha_1(\overline{x}), \overline{\alpha}(\overline{x})) \in \mathbb{S}^{n-1}, \ \alpha_1(\overline{x}) := \frac{F(\overline{x}) - b_1}{\rho(\overline{x})}, \ \overline{\alpha}(\overline{x}) := \frac{\overline{x} - \overline{b}}{\rho(\overline{x})}. \tag{3.1}$$

According to Definition 2.2, the grazing set and the illuminable regions are given by

$$G_{\phi_i} = \{ (F(\overline{x}), \overline{x}, t) \mid \alpha_1(\overline{x}) = \langle \nabla F(\overline{x}), \overline{\alpha}(\overline{x}) \rangle \},$$

$$I_- = \{ (F(\overline{x}), \overline{x}, t) \mid \alpha_1(\overline{x}) < \langle \nabla F(\overline{x}), \overline{\alpha}(\overline{x}) \rangle \}.$$

$$(3.2)$$

Fix any R > 0 and any T > 0. In (3.2) and throughout §3 we allow $\overline{x} \in \overline{B(0,R)} \subset \mathbb{R}^{n-1}$ and $t \in [-T,T]$.

⁹The normalization that F has a max at $\overline{x} = 0$ with F(0) = 1 is actually not used in §3, but the reader is free to assume that.

For $t^0 \in \mathbb{R}$, $x^0 \in \mathbb{R}^n \setminus (\mathcal{O} \cup \{b\})$, we have

$$\nabla_{x,t}\phi_i(x^0,t^0) = \left(\frac{x^0 - b}{|x^0 - b|}, -1\right).$$

The incoming bicharacteristic passing $(x^0, t^0, \frac{x^0-b}{|x^0-b|}, -1)$ is

$$\gamma_i(s) = \left(x_1^0 + 2s \frac{x_1^0 - b_1}{|x^0 - b|}, \overline{x}^0 + 2s \frac{\overline{x}^0 - \overline{b}}{|x^0 - b|}, t^0 - 2s\tau, \frac{x_1^0 - b_1}{|x^0 - b|}, \frac{\overline{x}^0 - \overline{b}}{|x^0 - b|}, -1\right).$$

Suppose $\gamma_i(s)$ intersects $\partial T^*((\mathbb{R}^n \setminus \mathcal{O}) \times \mathbb{R})$ at $(F(\overline{x}), \overline{x}, t', \xi_1^i, \overline{\xi}^i, -1)$, then $\xi^i(\overline{x}) = \frac{x^0 - b}{|x^0 - b|} = \alpha(\overline{x})$. Let $\xi^r(\overline{x})$ be the reflected direction, then

$$(\xi^i, -1) - (\xi^r, \tau^r) = c(1, -\nabla F, 0), |\xi^r| = |\tau^r|.$$

From here we solve

$$\xi_{1}^{r}(\overline{x}) = \alpha_{1}(\overline{x}) + \frac{2}{1 + |\nabla F(\overline{x})|^{2}} \left(\langle \nabla F(\overline{x}), \overline{\alpha}(\overline{x}) \rangle - \alpha_{1}(\overline{x}) \right),$$

$$\overline{\xi}^{r}(\overline{x}) = \overline{\alpha}(\overline{x}) - \frac{2\nabla F(\overline{x})}{1 + |\nabla F(\overline{x})|^{2}} \left(\langle \nabla F(\overline{x}), \overline{\alpha}(\overline{x}) \rangle - \alpha_{1}(\overline{x}) \right).$$
(3.3)

The reflected flow map is then given by

$$Z_r(s, \overline{x}, t) = (F(\overline{x}) + 2s\xi_1^r(\overline{x}), \overline{x} + 2s\overline{\xi}^r(\overline{x}), t + 2s).$$

Proposition 3.1. Fix any $s_0 > 0$ and take $\overline{x} \in \overline{B(0,R)}$, $t \in [-T,T]$ for any positive R and T. The reflected flow map Z_r is a diffeomorphism from $[0,s_0] \times I_-$ onto its range and extends to a homeomorphism from $[0,s_0] \times (G_{\phi_i} \cup I_-)$ onto its range.

Proof. It suffices to show the following two claims

- 1. Z_r is injective on $[0, s_0] \times (G_{\phi_i} \cup I_-)$;
- 2. The Jacobian j of Z_r is non-zero on $[0, s_0] \times I_-$.
- 1. Injectivity. We claim that the following holds when $\overline{x}^* \neq \overline{x}$

$$\langle \xi^r(\overline{x}^*) - \xi^r(\overline{x}), (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}) \rangle > 0.$$

Indeed, using (3.1) we have

$$\overline{\xi}^r(\overline{x}) = \overline{\alpha}(\overline{x}) - \nabla F(\overline{x})(\xi_1^r(\overline{x}) - \alpha_1(\overline{x})) \tag{3.4}$$

and the same equality holds when \overline{x} is replaced by \overline{x}^* . Equation (3.4) implies that

$$\begin{split} &\langle \overline{\xi}^r(\overline{x}^*) - \overline{\xi}^r(\overline{x}), \overline{x}^* - \overline{x} \rangle \\ = &\langle \overline{\alpha}(\overline{x}^*) - \overline{\alpha}(\overline{x}), \overline{x}^* - \overline{x} \rangle \\ &+ \langle \nabla F(\overline{x}^*), \overline{x} - \overline{x}^* \rangle (\xi_1^r(\overline{x}^*) - \alpha_1(\overline{x}^*)) + \langle \nabla F(\overline{x}), \overline{x}^* - \overline{x} \rangle (\xi_1^r(\overline{x}) - \alpha_1(\overline{x})) \\ \geq &\langle \overline{\alpha}(\overline{x}^*) - \overline{\alpha}(\overline{x}), \overline{x}^* - \overline{x} \rangle \\ &+ (F(\overline{x}) - F(\overline{x}^*)) (\xi_1^r(\overline{x}^*) - \alpha_1(\overline{x}^*)) + (F(\overline{x}^*) - F(\overline{x})) (\xi_1^r(\overline{x}) - \alpha_1(\overline{x})) \\ = &- (F(\overline{x}^*) - F(\overline{x})) (\xi_1^r(\overline{x}^*) - \xi_1^r(\overline{x})) \\ &+ (F(\overline{x}^*) - F(\overline{x})) (\alpha_1(\overline{x}^*) - \alpha_1(\overline{x})) + \langle \overline{\alpha}(\overline{x}^*) - \overline{\alpha}(\overline{x}), \overline{x}^* - \overline{x} \rangle. \end{split}$$

Since F is strictly concave, we know the equality holds if and only if

$$\xi_1^r(\overline{x}^*) = \alpha_1(\overline{x}^*) \text{ and } \xi_1^r(\overline{x}) = \alpha_1(\overline{x}).$$
 (3.5)

Next we show that

$$(F(\overline{x}^*) - F(\overline{x}))(\alpha_1(\overline{x}^*) - \alpha_1(\overline{x})) + \langle \overline{\alpha}(\overline{x}^*) - \overline{\alpha}(\overline{x}), \overline{x}^* - \overline{x} \rangle \ge 0,$$

that is,

$$\langle \alpha(\overline{x}^*) - \alpha(\overline{x}), (F(\overline{x}^*), \overline{x}^*) - (F(\overline{x}), \overline{x}) \rangle \ge 0.$$

Using definitions of α and ρ , we need to show

$$\langle \alpha(\overline{x}^*) - \alpha(\overline{x}), \rho(\overline{x}^*)\alpha(\overline{x}^*) - \rho(\overline{x})\alpha(\overline{x}) \rangle \ge 0.$$

A direct computation shows that the left-hand side is

$$(\rho(\overline{x}^*) + \rho(\overline{x}))(1 - \langle \alpha(\overline{x}^*), \alpha(\overline{x}) \rangle)$$

which is non-negative and equals zero if and only if

$$\alpha(\overline{x}^*) = \alpha(\overline{x}). \tag{3.6}$$

Thus we have shown that

$$\langle \xi^r(\overline{x}^*) - \xi^r(\overline{x}), (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}) \rangle \ge 0$$

and equality holds if and only if both (3.5) and (3.6) hold. It remains to show that the equality can not hold. Otherwise, combining (3.5) and (3.6), we know

$$\alpha(\overline{x}^*) = \alpha(\overline{x}) \text{ and } \xi^r(\overline{x}^*) = \xi^r(\overline{x}).$$

But this implies that

$$\nabla F(\overline{x}^*) = \nabla F(\overline{x})$$

which is impossible when $\overline{x}^* \neq \overline{x}$.

The map Z_r is continuous and injective on the compact domain $[0, s_0] \times (G_{\phi_i} \cup I_-)$, and hence is a homeomorphism on this domain.

2. Diffeomorphism. As in the plane wave case [WW23, Proposition 8.10], we have

$$j(s, \overline{x}, t) = 2\xi_1^r \det(A), \ A := I - \frac{\overline{\xi}^r \otimes \nabla F}{\xi_1^r} + 2s \left(\frac{\partial \overline{\xi}^r}{\partial \overline{x}} - \frac{\overline{\xi}^r \otimes \nabla \xi_1^r}{\xi_1^r} \right).$$

We first differentiate the identity $(\xi_1^r)^2 + |\overline{\xi}^r|^2 = 1$ with respect to \overline{x} to obtain

$$\xi_1^r \nabla \xi_1^r + \overline{\xi}^r \frac{\partial \overline{\xi}^r}{\partial \overline{x}} = 0 \implies \nabla \xi_1^r = -\frac{\overline{\xi}^r}{\xi_1^r} \frac{\partial \overline{\xi}^r}{\partial \overline{x}}.$$

Thus

$$\frac{\partial \overline{\xi}^r}{\partial \overline{x}} - \frac{\overline{\xi}^r \otimes \nabla \xi_1^r}{\xi_1^r} = \left(I + \frac{\overline{\xi}^r \otimes \overline{\xi}^r}{(\xi_1^r)^2}\right) \frac{\partial \overline{\xi}^r}{\partial \overline{x}}.$$

In the following we denote

$$B := I - \frac{\overline{\xi}^r \otimes \nabla F}{\xi_1^r}, \ C := I + \frac{\overline{\xi}^r \otimes \overline{\xi}^r}{(\xi_1^r)^2}.$$

Then

$$A = B + 2sC \frac{\partial \overline{\xi}^r}{\partial \overline{x}}.$$

2.1 Computation of $\frac{\partial \overline{\xi}^r}{\partial \overline{x}}$. Differentiate (3.4) with respect to \overline{x} and we find

$$\frac{\partial \overline{\xi}^r}{\partial \overline{x}} = \frac{\partial \overline{\alpha}}{\partial \overline{x}} + (\alpha_1 - \xi_1^r) \nabla^2 F + \nabla F \otimes \nabla (\alpha_1 - \xi_1^r). \tag{3.7}$$

Differentiating the first equation in (3.3) with respect to \overline{x} gives

$$\nabla(\alpha_1 - \xi_1^r) = -\frac{2\overline{\xi}^r}{1 + |\nabla F|^2} \nabla^2 F - \frac{2}{1 + |\nabla F|^2} \left(\nabla F + \frac{\overline{\alpha}}{\alpha_1}\right) \frac{\partial \overline{\alpha}}{\partial \overline{x}}.$$
 (3.8)

In the computation we used $|\alpha| = 1$, which implies that

$$\nabla \alpha_1 = -\frac{\overline{\alpha}}{\alpha_1} \frac{\partial \overline{\alpha}}{\partial \overline{x}}.$$

Putting (3.7) and (3.8) together gives

$$\frac{\partial \overline{\xi}^r}{\partial \overline{x}} = \left(I - \frac{2\nabla F \otimes (\alpha_1 \nabla F + \overline{\alpha})}{\alpha_1 (1 + |\nabla F|^2)}\right) \frac{\partial \overline{\alpha}}{\partial \overline{x}} + \left((\alpha_1 - \xi_1^r)I - \frac{2\nabla F \otimes \overline{\xi}^r}{1 + |\nabla F|^2}\right) \nabla^2 F.$$

In the following we denote

$$K := \left(I - \frac{2\nabla F \otimes (\alpha_1 \nabla F + \overline{\alpha})}{\alpha_1 (1 + |\nabla F|^2)}\right) \frac{\partial \overline{\alpha}}{\partial \overline{x}}, \ L := \left((\alpha_1 - \xi_1^r)I - \frac{2\nabla F \otimes \overline{\xi}^r}{1 + |\nabla F|^2}\right) \nabla^2 F.$$

A direct computation using the definition (3.1) of $\overline{\alpha}$ gives

$$\frac{\partial \overline{\alpha}}{\partial \overline{x}} = \frac{I - \overline{\alpha} \otimes (\alpha_1 \nabla F + \overline{\alpha})}{\rho}.$$

Inserting this in the expression of K gives

$$\begin{split} K = & \frac{1}{\rho} \left(I - \frac{2\nabla F \otimes (\alpha_1 \nabla F + \overline{\alpha})}{\alpha_1 (1 + |\nabla F|^2)} \right) \left(I - \overline{\alpha} \otimes (\alpha_1 \nabla F + \overline{\alpha}) \right) \\ = & \frac{1}{\rho} \left(I - \frac{2\nabla F \otimes (\alpha_1 \nabla F + \overline{\alpha})}{\alpha_1 (1 + |\nabla F|^2)} - \overline{\alpha} \otimes (\alpha_1 \nabla F + \overline{\alpha}) + \frac{2\langle \alpha_1 \nabla F + \overline{\alpha}, \overline{\alpha} \rangle \nabla F \otimes (\alpha_1 \nabla F + \overline{\alpha})}{\alpha_1 (1 + |\nabla F|^2)} \right) \\ = & \frac{1}{\rho} \left(I - \frac{2\nabla F \otimes (\alpha_1 \nabla F + \overline{\alpha})}{\alpha_1 (1 + |\nabla F|^2)} - \overline{\alpha} \otimes (\alpha_1 \nabla F + \overline{\alpha}) + \frac{2(1 + \alpha_1 (\langle \nabla F, \overline{\alpha} \rangle - \alpha_1)) \nabla F \otimes (\alpha_1 \nabla F + \overline{\alpha})}{\alpha_1 (1 + |\nabla F|^2)} \right) \\ = & \frac{1}{\rho} \left(I - \overline{\alpha} \otimes (\alpha_1 \nabla F + \overline{\alpha}) + \frac{2(\langle \nabla F, \overline{\alpha} \rangle - \alpha_1) \nabla F \otimes (\alpha_1 \nabla F + \overline{\alpha})}{(1 + |\nabla F|^2)} \right) \\ = & \frac{I - \overline{\xi}^r \otimes (\alpha_1 \nabla F + \overline{\alpha})}{\rho}. \end{split}$$

Now for L, we write

$$L = (\alpha_1 - \xi_1^r) \left(I - \frac{2\nabla F \otimes \overline{\xi}^r}{(\alpha_1 - \xi_1^r)(1 + |\nabla F|^2)} \right) \nabla^2 F.$$

Use the first equation in (3.3) and we have

$$L = \left(I + \frac{\nabla F \otimes \overline{\xi}^r}{\langle \nabla F, \overline{\alpha} \rangle - \alpha_1}\right) \frac{2(\langle \nabla F, \overline{\alpha} \rangle - \alpha_1)}{1 + |\nabla F|^2} \times (-\nabla^2 F).$$

We need the following lemma to proceed.

Lemma 3.2. Let $a, b \in \mathbb{R}^{n-1}$ be row vectors, then

$$\det(I + a \otimes b) = 1 + \langle a, b \rangle.$$

Moreover,

$$(I + a \otimes b)^{-1} = I - \frac{a \otimes b}{1 + \langle a, b \rangle}$$

as long as $\langle a, b \rangle \neq -1$.

Proof. For the first part, see [WW23, (8.50)]. For the second identity, we compute directly

$$(I+a\otimes b)\left(I-\frac{a\otimes b}{1+\langle a,b\rangle}\right)=I+\left(1-\frac{1}{1+\langle a,b\rangle}\right)a\otimes b-\frac{\langle a,b\rangle a\otimes b}{1+\langle a,b\rangle}=I.$$

This completes the proof.

Apply Lemma 3.2 to B and we find that

$$\det(B) = \frac{\xi_1^r - \langle \overline{\xi}^r, \nabla F \rangle}{\xi_1^r} = \frac{\langle \nabla F, \overline{\alpha} \rangle - \alpha_1}{\xi_1^r}.$$

Moreover,

$$B^{T} = I - \frac{\nabla F \otimes \overline{\xi}^{r}}{\xi_{1}^{r}}, \ (B^{T})^{-1} = I + \frac{\nabla F \otimes \overline{\xi}^{r}}{\xi_{1} - \langle \nabla F, \overline{\xi}^{r} \rangle} = I + \frac{\nabla F \otimes \overline{\xi}^{r}}{\langle \nabla F, \overline{\alpha} \rangle - \alpha_{1}}.$$

As a result, we have 10

$$B^{T}K = \frac{1}{\rho} \left(I - \frac{\nabla F \otimes \overline{\xi}^{T}}{\xi_{1}^{T}} \right) \left(I - \overline{\xi}^{r} \otimes (\alpha_{1}\nabla F + \overline{\alpha}) \right)$$

$$= \frac{1}{\rho} \left(I - \frac{\nabla F \otimes \overline{\xi}^{T}}{\xi_{1}^{T}} - \overline{\xi}^{r} \otimes (\alpha_{1}\nabla F + \overline{\alpha}) + \frac{|\overline{\xi}^{r}|^{2}\nabla F \otimes (\alpha_{1}\nabla F + \overline{\alpha})}{\xi_{1}^{T}} \right)$$

$$= \frac{1}{\rho} \left(I - \frac{\nabla F \otimes \overline{\xi}^{T}}{\xi_{1}^{T}} - \overline{\xi}^{r} \otimes (\alpha_{1}\nabla F + \overline{\alpha}) + \frac{(1 - (\xi_{1}^{r})^{2})\nabla F \otimes (\alpha_{1}\nabla F + \overline{\alpha})}{\xi_{1}^{T}} \right)$$

$$= \frac{1}{\rho} \left(I + \frac{\nabla F \otimes (\alpha_{1}\nabla F + \overline{\alpha} - \overline{\xi}^{r})}{\xi_{1}^{T}} - (\xi_{1}\nabla F + \overline{\xi}^{r}) \otimes (\alpha_{1}\nabla F + \overline{\alpha}) \right)$$

$$= \frac{I + \nabla F \otimes \nabla F - (\alpha_{1}\nabla F + \overline{\alpha}) \otimes (\alpha_{1}\nabla F + \overline{\alpha})}{\rho}.$$
(3.9)

From here we can see that B^TK is symmetric. We claim that B^TK is also positive definite. Indeed, for any $v \in \mathbb{R}^{n-1}$, there holds

$$\rho \langle B^T K v, v \rangle = |v|^2 + |\langle \nabla F, v \rangle|^2 - |\langle \alpha_1 \nabla F + \overline{\alpha}, v \rangle|^2$$
$$= |v|^2 + |\langle \nabla F, v \rangle|^2 - |\langle (\alpha_1, \overline{\alpha}), (\langle \nabla F, v \rangle, v) \rangle|^2 \ge 0.$$

Equality holds if and only if for some $c \neq 0$

$$\alpha_1 = c\langle \nabla F, v \rangle$$
 and $\overline{\alpha} = cv$,

which implies

$$\alpha_1 = \langle \nabla F, \overline{\alpha} \rangle.$$

But the latter can not happen in I_{-} according to (3.2).

The formulæ for $(B^{-1})^T$ and L tell us

$$B^{T}L = -\frac{2(\langle \nabla F, \overline{\alpha} \rangle - \alpha_{1})}{1 + |\nabla F|^{2}} \nabla^{2} F.$$

Notice that $\langle \nabla F, \overline{\alpha} \rangle - \alpha_1 > 0$ on I_- and $-\nabla^2 F$ is positive semi-definite, thus $B^T L$ is symmetric and positive semi-definite.

Finally, we conclude that

$$j(s, \overline{x}, t) = 2\xi_1^r \det(A) = 2\xi_1^r \det(B + 2sC(K + L))$$

= $2\xi_1^r \det(B) \det(I + 2sB^{-1}C(B^{-1})^T(B^TK + B^TL))$
= $2(\langle \nabla F, \overline{\alpha} \rangle - \alpha_1) \det(I + 2sB^{-1}C(B^{-1})^T(B^TK + B^TL)).$

Notice now that both $B^{-1}C(B^{-1})^T$ and $B^TK + B^TL$ are symmetric and positive definite. A similar argument as in the plane wave case [WW23, Proposition 8.10] shows that the product of these two matrices has positive eigenvalues. Thus we conclude that

$$j(s, \overline{x}, t) \ge 2(\langle \nabla F, \overline{\alpha} \rangle - \alpha_1) > 0.$$

This completes the proof.

The last equality in (3.9) we used $\xi_1 \nabla F + \overline{\xi}^r = \alpha_1 \nabla F + \overline{\alpha}^r$.

4. The grazing set for spherical waves hitting a convex obstacle

In this section we present a number of examples of the diffraction of spherical waves by strictly convex obstacles in 3D where we can verify that the grazing set satisfies [WW23, Assumption 2.10], including examples where the grazing set is C^1 but not twice differentiable. The operator in question is the operator on \mathbb{R}^{n+1} given by

$$\Box = \partial_{x_1}^2 + \dots + \partial_{x_n}^2 - \partial_t^2.$$

As before we allow arbitrarily high order grazing. We also present examples where the grazing set is continuous but not differentiable, so [WW23, Assumption 2.10] fails to hold. In general it is challenging to understand the nature (regularity, branching,...) of the grazing set near a point of high order grazing because the implicit function theorem cannot be used.

4.1. Branching does not occur. The first proposition rules out *branching* of the grazing set for a large class of strictly convex obstacles $\mathcal{O} \subset \mathbb{R}^n$ defined by $x_1 = F(\overline{x})$ as in [WW23, Proposition 8.1]. Let us explain what this means.

We write $x=(x_1,\overline{x})$, where $\overline{x}=(x_2,x_3)\in B(0,r)$ for some small enough ball $B(0,r)=\{\overline{x}:|\overline{x}|< r\}$. We take a "spherical" incoming phase $\phi_i=-t+|x-b|$, where $b=(1,\overline{b})=(1,-1,0)$. The obstacle corresponding to F has $\underline{\sigma}=i^*\underline{\rho}\in G_d^{2k}\setminus G^{2k+1}$, where $\underline{\rho}=(1,0,0,t^0,0,-\overline{b},-1)$. To see this note that with $x^0=(1,0,0)$, the incoming bicharacteristic passing through $(x^0,t^0,\frac{x^0-b}{|x^0-b|},-1)$ at s=0 is

$$\gamma_i(s) = (1, 2s(-\overline{b}), t^0 + 2s, 0, -\overline{b}, -1),$$

and repeat the computation [WW23, (8.5)] with $-\overline{b}$ in place of $\overline{\theta}$.

Assumption 4.1. We take F as in [WW23, Definition 8.1], but require also that for \overline{x} near 0,

(a)
$$\nabla^2 F(\overline{x}) < 0 \text{ for } \overline{x} \neq 0,$$

(b) $\partial_{x_3} F(\overline{x}) = 0 \iff x_3 = 0.$ (4.1)

Later in this section we provide many examples of functions F satisfying Assumption 4.1.

The grazing set near (x^0, t^0) is

$$G_{\phi_i} = \{ (F(\overline{x}), \overline{x}, t) \mid F(\overline{x}) - 1 - \nabla F(\overline{x}) \cdot (\overline{x} - \overline{b}) = 0, \ \overline{x} \ \text{near} \ 0, \ t \in \mathbb{R} \}.$$

We claim that for F as above, the set

$$\pi_{\overline{x}}G_{\phi_i} := \{ \overline{x} \text{ near } 0 \mid \mathcal{H}(\overline{x}) := F(\overline{x}) - 1 - \nabla F(\overline{x}) \cdot (\overline{x} - \overline{b}) = 0 \}$$

consists of a single continuous curve through $\overline{x} = 0$.

Proposition 4.2. Let F be as in [WW23, Definition 8.1] and also satisfy Assumption 4.1. Then the set $\pi_{\overline{x}}G_{\phi_i} \setminus \{0\}$ consists of two C^{∞} curves, one in $x_3 > 0$ and one in $x_3 < 0$. The set $\pi_{\overline{x}}G_{\phi_i}$ is a single continuous curve through $\overline{x} = 0$.

Each curve in $\pi_{\overline{x}}G_{\phi_i}\setminus\{0\}$ lifts under the map $\overline{x}\mapsto(F(\overline{x}),\overline{x})$ to a curve in $\pi_xG_{\phi_i}$. The proposition rules out the possibility that the set $\pi_{\overline{x}}G_{\phi_i}\setminus\{0\}$ consists of more than two smooth curves that meet and terminate at $\overline{x}=0$. That possibility is what we refer to as *branching*.

Proposition 4.2 leads us to expect, but does not claim, that near $\overline{x} = 0$ the set $\pi_{\overline{x}}G_{\phi_i}$ is either the graph of a continuous function $x_2 = x_2(x_3)$ such that $x_2(0) = 0$, or the graph of a continuous function $x_3 = x_3(x_2)$ such that $x_3(0) = 0$.

Proof of Proposition 4.2. 1. Smoothness of $\pi_{\overline{x}}G_{\phi_i}$ away from $\overline{x} = 0$. For \overline{x} near 0, $\overline{x} \neq 0$ we compute

$$\nabla \mathcal{H}(\overline{x}) = \nabla F(\overline{x}) - \nabla^2 F(\overline{x})(\overline{x} - \overline{b}) - \nabla F(\overline{x}) = -\nabla^2 F(\overline{x})(\overline{x} - \overline{b}) \neq 0, \tag{4.2}$$

since $\nabla^2 F < 0$ for $\overline{x} \neq 0$.

2. The curve $C(A^*)$. In the x_1, x_2 plane fix a point $A^* = (F(x_2^*, 0), x_2^*)$ near $(x_1, x_2) = (1, 0)$ with $x_2^* < 0$ on the strictly concave curve $x_1 = F(x_2, 0)$, and draw the infinite ray from (1, -1) toward A^* . This ray intersects $x_1 = F(x_2, 0)$ at A^* and at a second point $B^* = (F(x_2^{**}, 0), x_2^{**})$ in $x_2 > 0$; see Figure Let $P(A^*)$ denote the half-plane in x-space consisting of all points that project to this ray under $(x_1, x_2, x_3) \to (x_1, x_2)$. We have

$$P(A^*) = \{(x_1, x_2, x_3) : (x_1 - 1)(x_2^* + 1) + (x_2 + 1)(1 - F(x_2^*, 0)) = 0.\}$$
(4.3)

Recall from Definition 8.1 of [WW23] that the obstacle \mathcal{O} has $\partial \mathcal{O} = \{(F(\overline{x}), \overline{x}) : \overline{x} \in B(0,r)\}$ for some r > 0. Set

$$C(A^*) = P(A^*) \cap \partial \mathcal{O} = \{ (F(\overline{x}), \overline{x}) : K(\overline{x}) := (F(\overline{x}) - 1)(x_2^* + 1) + (x_2 + 1)(1 - F(x_2^*, 0)) = 0 \}.$$
(4.4)

We will show that $C(A^*)$ is a C^{∞} closed strictly convex curve that contains exactly two points in $\pi_x G_{\phi_i}$.

3. Smoothness of $C(A^*)$ **.** We show that

$$\nabla K(\overline{x}) = (F_{x_2}(\overline{x})(x_2^* + 1) + (1 - F(x_2^*, 0)), F_{x_3}(\overline{x})(x_2^* + 1)$$
(4.5)

is nonzero at any \overline{x} where $K(\overline{x}) = 0$. If $F_{x_3}(\overline{x}) \neq 0$, then $\nabla K(\overline{x}) \neq 0$, so suppose $F_{x_3}(\overline{x}) = 0$. By (4.1)(b) this implies $x_3 = 0$. We show now that the conditions

$$(a)x_3 = 0, (b)K(\overline{x}) = 0, (c)K_{x_2}(\overline{x}) = 0$$
 (4.6)

 $[\]overline{{}^{11}\text{Recall }b} = (1, -1, 0).$

are incompatible. Conditions 4.6(b), (c) imply

$$F_{x_2}(\overline{x}) = \frac{F(x_2^*, 0) - 1}{x_2^* + 1} = \frac{F(\overline{x}) - 1}{x_2 + 1} < 0. \tag{4.7}$$

A Taylor expansion gives

$$1 - F(\overline{x}) = F(0) - F(\overline{x}) = -\nabla F(\overline{x})\overline{x} + \int_0^1 (1 - s)\langle \nabla^2 F(\overline{x}(1 - s))ds \ \overline{x}, \overline{x} \rangle =$$

$$- F_{x_2}(\overline{x})x_2 + \int_0^1 (1 - s)F_{x_2x_2}(\overline{x}(1 - s))ds \ x_2^2,$$

$$(4.8)$$

where we used (4.6)(a) to get the third equality. From (4.7) and (4.8) we obtain

$$F_{x_2}(\overline{x})(x_2+1) = F_{x_2}(\overline{x})x_2 - \int_0^1 (1-s)F_{x_2x_2}(\overline{x}(1-s))ds \ x_2^2 \Rightarrow$$

$$F_{x_2}(\overline{x}) = -\int_0^1 (1-s)F_{x_2x_2}(\overline{x}(1-s))ds \ x_2^2 > 0,$$
(4.9)

which contradicts (4.7).

4. Nonvanishing curvature of $C(A^*)$. Let $Q = (F(\overline{x}), \overline{x}) \in C(A^*)$ be a point where $K_{x_2}(\overline{x}) \neq 0$. Near \overline{x} the curve $\pi_{\overline{x}}C(A^*)$ is the graph of a C^{∞} function $x_2 = x_2(x_3)$.. Differentiating $K(x_2(x_3), x_3) = 0$ twice with respect to x_3 we obtain

$$K_{x_2}x_2' + K_{x_3} = 0$$

$$\langle \nabla^2 K(x_2(x_3), x_3)(x_2', 1), (x_2', 1) \rangle + K_{x_2}x_2'' = 0 \Rightarrow x_2'' \neq 0.$$
(4.10)

Here we used $\nabla^2 K(\overline{x}) = \nabla^2 F(\overline{x})(x_2^* + 1) < 0$. Near Q we can parametrize $C(A^*)$ by

$$r(x_3) = (F(x_2(x_3), x_3), x_2(x_3), x_3), \tag{4.11}$$

with which we compute

$$r'(x_3) = (F_{x_2}x_2' + F_{x_3}, x_2', 1) := (a, x_2', 1)$$

$$r''(x_3) := (b, x_2'', 0),$$
(4.12)

so $r' \times r'' = (-x_2'', b, *) \neq 0.$ Now apply the curvature formula for a parametrized curve

$$\kappa(x_3) = \frac{|r'(x_3) \times r''(x_3)|}{|r'(x_3)|^3} \neq 0.$$
(4.13)

5. Grazing points on $C(A^*)$. We now show that $C(A^*)$ contains one grazing point in $x_3 > 0$ and one in $x_3 < 0$. The infinite ray from (1, -1) toward A^* described in step 1, ray (A^*) , makes an angle of $\theta = 0$ with the x_1x_2 plane. Keeping b = (1, -1, 0) fixed, we now increase the angle θ that ray (A^*) makes with the x_1x_2 plane, always keeping

¹²Points where $K_{x_3}(\overline{x}) \neq 0$ are treated similarly.

¹³We have $F_{x_2}(x_2^*,0) = \int_0^1 F_{x_2x_2}(s(x_2^*,0))ds \cdot x_2 > 0$, so $K_{x_2}(x_2^*,0) > 0$ by (4.5). This implies $x_2'' > 0$ at $(x_2^*,0)$ by (4.10).

the rotated ray, $\operatorname{ray}(\theta)$, in the plane $P(A^*)$. As θ increases let $A(\theta)$ denote the point closer to b at which $\operatorname{ray}(\theta)$ intersects $C(A^*)$. Denote by $m(\theta)$ the slope of the tangent vector to $C(A^*)$ at $A(\theta)$.¹⁴ So m(0) > 0 and by step 4 $m(\theta)$ strictly decreases as θ increases.¹⁵ The strict monotonicity of $m(\theta)$ implies that if there is a positive angle θ , call it θ_g , where $\tan \theta_g = m(\theta_g)$, then θ_g is unique. The point $A(\theta_g)$ is then the unique grazing point on $C(A^*)$ in $x_3 > 0$.

To see that θ_q exists, we use Assumption 4.1 and (4.4) to write

$$C(A^*) \cap \{x_3 > 0\} = \{(F(x_2, x_3(x_2)), x_2, x_3(x_2)) : x_2^* < x_2 < x_2^{**}\},$$
 (4.14)

where $x_3(x_2)$ is a C^{∞} function constructed by concatenation using the implicit function theorem. Setting $r(x_2) = (F(x_2, x_3(x_2)), x_2)$ we obtain from (4.4) that

$$r'(x_2) = \left(\frac{F(x_2^*, 0) - 1}{x_2^* + 1}, 1\right) := (C^*, 1). \tag{4.15}$$

Let $s(x_2) = \int_{x_2^*}^{x_2} |r'(x_2|dx_2 = |(C^*, 1)| (x_2 - x_2^*)$ and define the open s-interval

$$I = (0, |(C^*, 1)| (x_2^{**} - x_2^*)).$$

We can view $C(A^*) \cap \{x_3 > 0\}$ as a graph above the interval I of the function

$$x_3 = f(s)$$
, where $f(s) := x_3(x_2(s))$. (4.16)

If ray(θ) intersects $C(A^*)$ at a point corresponding to (s, f(s)), then $\tan(\theta) = \frac{f(s)}{s+a} := g(s)$, where a is the length of the segment from (1, -1) to A^* . The existence of θ_g is equivalent to the existence of s_g such that

$$\frac{f(s_g)}{s_g + a} = f'(s_g). {(4.17)}$$

Now $g'(s) = 0 \Leftrightarrow f'(s) = \frac{f(s)}{s+a}$. Since g vanishes at the endpoints of I, Rolle's theorem implies the existence of $s_g \in I$.

A similar argument yields a unique grazing point on $C(A^*)$ in $x_3 < 0$.

6. No branching. With (4.4) we see that $P(A^*)$ contains exactly two points of $\pi_x G_{\phi_i}$, one in $x_3 > 0$ and one in $x_3 < 0$. Moreover,

$$\cup_{A^* \text{ near } (1,0), \ x_2^* \le 0} \ (P(A^*) \cap \pi_x G_{\phi_i})$$

is equal to the set of points in $\pi_x G_{\phi_i}$ near (1,0,0). Step **1** implies that as $x_2^* \nearrow 0$, the points of $P(A^*) \cap \pi_x G_{\phi_i}$ trace out two curves that are C^{∞} away from (1,0,0). The curves approach (1,0,0) continuously as $x_2^* \nearrow 0$, since $P(A^*)$ approaches the plane $x_1 = 0$.

^{14 &}quot;Slope" here means the change in x_3 divided by the magnitude of the change in (x_1, x_2) along the tangent vector.

 $^{^{15}}$ To see this use (4.12).

¹⁶The map $s \to (F(x_2(s), x_3(x_2(s))), x_2(s), x_3(x_2(s)))$ is bijective from I onto $C(A^*) \cap \{x_3 > 0\}$.

4.2. Examples of good grazing sets. The next Proposition gives a large class of examples that satisfy the conditions of Proposition 4.2 as well as [WW23, Assumption 2.10]. We then extend this class much further in Proposition 4.4. It will be convenient below sometimes to write $\overline{x} = (x_2, x_3) = (u, v)$. In these examples the set $\pi_{\overline{x}}G_{\phi_i}$ turns out to be given near $\overline{x} = 0$ by the graph of a C^1 function u = u(v) with u = u(0).

Proposition 4.3. Suppose

$$F(u,v) = 1 - (c_0 u^{2k} + c_1 u^{2k-2} v^2 + c_2 u^{2k-4} v^4 + \dots + c_{k-1} u^2 v^{2k-2} + c_k v^{2k})$$

$$:= 1 - G(u,v), \quad k > 1,$$
(4.18)

where $\nabla^2 G(u,v) > 0$ for $(u,v) \neq 0$.¹⁷ Suppose all $c_j > 0$. Define the obstacle \mathcal{O} as in [WW23, Definition 8.1] by $x_1 = F(\overline{x})$, and take the incoming phase to be $\phi_i = -t + |x-b|$, b = (1,-1,0) as before. This class of problems satisfies [WW23, Assumption 2.10].

- *Proof.* 1. We know that branching does not occur, but we do not know yet that the continuous curve described in Proposition 4.2 can be realized as the graph of a function u = u(v) or v = v(u). For the moment we regard $\pi_{\overline{x}}G_{\phi_i}$ as defining a relation.
- **2. Estimate of** |u| **for points in** $\pi_{\overline{x}}G_{\phi_i}$. The next step is to estimate |u| in terms of |v| for points $(u,v) \in \pi_{\overline{x}}G_{\phi_i}$. I'll write the details for the case k=3; but the same argument clearly works for any k. So we take

$$G(u,v) = u^6 + c_1 u^4 v^2 + c_2 u^2 v^4 + c_3 v^6, c_i > 0$$

The grazing set is defined by the equation

$$\mathcal{H}(u,v) := (F(u,v) - 1) - \nabla F(u,v) \cdot ((u,v) - (-1,0)) = 0 \tag{4.19}$$

or equivalently

$$(2k-1)G(u,v) + G_u(u,v)$$

= $5(u^6 + c_1u^4v^2 + c_2u^2v^4 + c_3v^6) + (6u^5 + 4c_1u^3v^2 + 2c_2uv^4) = 0.$

Rewrite this as

$$u\left[u^{4}(6+5u)+v^{4}(2c_{2}+5c_{2}u)+u^{2}v^{2}(4c_{1}+5c_{1}u)\right]=-5c_{3}v^{6},$$
(4.20)

where $c_2, c_3 > 0$. The quantity in brackets is $\gtrsim u^4 + v^4$ for |(u, v)| small, so for |(u, v)| small, $v \neq 0$ we have¹⁸

$$|u| \lesssim \frac{|v|^6}{u^4 + v^4} \lesssim v^2.$$
 (4.21)

¹⁷The condition $\nabla^2 G(u,v) > 0$ for $(u,v) \neq 0$ implies G(u,v) > 0 for $(u,v) \neq 0$. The property G(u,v) > 0 implies that c_0 and c_k are positive (so take $c_0 = 1$).

¹⁸Note that for solutions (u, v), $u = 0 \Leftrightarrow v = 0$.

3. A relational formula. Dividing (4.20) by v^4 and rearranging slightly we obtain

$$u(2c_2 + 5c_2u) + 5c_3v^2 + r_4(u, v) = 0, \text{ where}$$

$$r_4(u, v) = \frac{u^5}{v^4}(6 + 5u) + \frac{u^3}{v^2}(4c_1 + 5c_1u) = O(v^6) + O(v^4) = O(v^4)$$
(4.22)

by step 2. From (4.22) and (4.21) we immediately obtain

$$u = \frac{-5c_3v^2}{2c_2} + O(v^4). (4.23)$$

At this point we know that the $O(v^4)$ term is a function of (u, v), not necessarily just of v.

4. The relation is a function. Now we use step **3** to show that $\pi_{\overline{x}}G_{\phi_i}$ is given by a function u = u(v) near $\overline{x} = 0$. We show that there exists r > 0 and a function u = u(v) such that

$$\pi_{\overline{x}}G_{\phi_i} \cap B(0,r) = \text{graph of } u.$$

Observe first that for $(u, v) \in \pi_{\overline{x}}G_{\phi_i}$, $u = 0 \Leftrightarrow v = 0$. For a given small r > 0 suppose there exist points (u_1, v) , (u_2, v) in $\pi_{\overline{x}}G_{\phi_i}$ with $u_1 > u_2$ and (say) v > 0. Then

$$0 = \mathcal{H}(u_{1}, v) - \mathcal{H}(u_{2}, v)$$

$$= \int_{0}^{1} \nabla \mathcal{H}(u_{2} + t(u_{1} - u_{2}), v) dt \cdot (u_{1} - u_{2}, 0)$$

$$= (u_{1} - u_{2}) \int_{0}^{1} \langle -\nabla^{2} F(u_{2} + t(u_{1} - u_{2}), v) (u_{2} + t(u_{1} - u_{2}) + 1, v), (1, 0) \rangle dt$$

$$= (u_{1} - u_{2}) \int_{0}^{1} \langle -\nabla^{2} F(u_{2} + t(u_{1} - u_{2}), v) (1, 0), (1, 0) \rangle dt$$

$$+ (u_{1} - u_{2}) \int_{0}^{1} \langle -\nabla^{2} F(u_{2} + t(u_{1} - u_{2}), v) (u_{2} + t(u_{1} - u_{2}), v), (1, 0) \rangle dt$$

$$= : A_{1} + A_{2},$$

$$(4.24)$$

where we used (4.2) for the third equality. We have

$$A_1 = -(u_1 - u_2) \int_0^1 F_{uu}(u_2 + t(u_1 - u_2), v) dt > 0.$$
 (4.25)

By step **3** we can write $(u_2 + t(u_1 - u_2), v) = v(O(v), 1)$, so

$$A_{2} = -(u_{1} - u_{2})v \int_{0}^{1} [F_{uu}(u_{2} + t(u_{1} - u_{2}), v)O(v) + F_{uv}(u_{2} + t(u_{1} - u_{2}), v)]dt.$$
(4.26)

For functions F(u, v) satisfying the assumptions of Proposition 4.3, it is easily checked that there exists C > 0 such that

$$|F_{uu}(u,v)| \ge C|v||F_{uv}(u,v)|$$
 (4.27)

for |(u,v)| small. Thus, if r>0 is small enough, the formulas (4.25), (4.26) imply

$$|A_2| \le \frac{1}{2}|A_1|.$$

Hence $A_1 + A_2 > 0$; so (4.24) implies 0 > 0, a contradiction.

5. Check u(v) is C^1 . With slight abuse we now write $r_4(u(v), v) = r_4(v)$. Using (4.2), (4.27), and the implicit function theorem we see that u(v) is C^{∞} in $v \neq 0$. Computing $u'(0) = r'_4(0) = 0$ using difference quotients, (4.22), and (4.23), we obtain that u and r_4 are differentiable on an open v interval containing 0 and C^{∞} away from v = 0. To show u(v) is C^1 near zero we now estimate u'(v) by differentiating (4.20) with respect to v, using the positivity of the bracketed term on the left, and using (4.21). After a straightforward computation this gives

$$|u'(v)| = O(|v|),$$

so u(v) (and hence $r_4(v)$) are C^1 near v=0.

5. Check $H_p(\zeta)(\rho) \neq 0$. We have

$$H_p = 2\xi_1 \partial_{x_1} + 2\overline{\xi} \partial_{\overline{x}} - 2\tau \partial_t \text{ and } \underline{\rho} = (1, 0, 0, t^0, 0, -\overline{b}, -1),$$

where $\overline{b}=(-1,0)$. Let $\zeta(x,t):x_2-u(x_3)$, where $u(\cdot)$ is the C^1 function just constructed. Then

$$H_p\zeta(\rho) = (2\partial_{x_2} + 2\partial_t)(x_2 - u(x_3))|_{\rho} = 2.$$

6. Points near $\underline{\sigma}$ in $G \cap \text{Graph } d\phi_0$ lie in \mathcal{G}_d . We have

$$\phi_i(x,t) = -t + |x-b|$$
, so $\phi_0(\overline{x},t) = -t + |(F(\overline{x}) - b_1, \overline{x} - \overline{b})| := -t + D(\overline{x})$.

Points $\sigma \in \text{Graph } d\phi_0 \text{ near } \underline{\sigma} = (0, t^0, -\overline{b}, -1) \text{ have the form } \sigma = (\overline{x}, t, \nabla D(\overline{x}), -1),$ where \overline{x} , t lie near $0, t^0$ respectively. Such a point lies in G if and only if

$$(i^*)^{-1}(\sigma) \cap p^{-1}(0)$$

consists of a single point. The possible elements of this set were computed just before Proposition 3.1 to be

$$(F(\overline{x}), \overline{x}, t, \xi^i(\overline{x}), -1)$$
 and $(F(\overline{x}), \overline{x}, t, \xi^r(\overline{x}), -1),$

¹⁹These properties and the estimate (4.22) do not imply that r_4 is C^1 near v = 0. Consider the example $p(v) = \begin{cases} v^4 \sin(1/v^4), v \neq 0 \\ 0, v = 0 \end{cases}$.

where $\xi^i(\overline{x}) = \alpha(\overline{x})$ and $\xi^r(\overline{x})$ are as in (3.3).²⁰ Clearly, these points coincide if and only if $\overline{x} = (u, v)$ satisfies the equation (4.19). In that case we set $\rho := (F(\overline{x}), \overline{x}, t, \alpha(\overline{x}), -1)$.

If $\overline{x} = 0$ then obviously $\sigma \in G_d^{2k} \setminus G^{2k+1}$, just like $\underline{\sigma}$. If $\overline{x} \neq 0$, setting $\beta := x_1 - F(\overline{x})$ we compute

$$H_p\beta(\rho) = \left[2\frac{F-1}{D}\partial_{x_1} + 2\frac{\overline{x}-\overline{b}}{D} \cdot \nabla_{\overline{x}} + 2\partial_t\right](x_1 - F)(\rho)$$
$$= \frac{2}{D}[F(\overline{x}) - 1 - (\overline{x} - \overline{b}) \cdot \nabla F(\overline{x})] = \frac{2\mathcal{H}}{D}(\overline{x}) = 0$$

and

$$H_p^2 \beta(\rho) = 2 \frac{\overline{x} - \overline{b}}{D} \cdot \nabla_{\overline{x}} \left(\frac{2\mathcal{H}}{D} \right) (\overline{x}) = 4 \frac{\overline{x} - \overline{b}}{D} \cdot \frac{\nabla_{\overline{x}} \mathcal{H}}{D} (\overline{x})$$

$$= -\frac{4}{D^2} \nabla^2 F(\overline{x}) (\overline{x} - \overline{b}) \cdot (\overline{x} - \overline{b}) > 0.$$
(4.28)

Here we used $\mathcal{H}(\overline{x}) = 0$ to get the second equality in (4.28). This shows $\sigma \in G_d^2 \setminus G^3$ when $\overline{x} \neq 0$.

Remark. The condition that $c_j > 0$ for all the coefficients c_j that appear in G can clearly be relaxed, but it must be relaxed carefully. For example, if $c_1 < 0$ with $|c_1|$ too large, the bracketed factor on the left in (4.20) will no longer be $\geq u^4 + v^4$ for |(u,v)| small. But such properties of c_1 might be incompatible with the requirement that $\nabla^2 G > 0$ for $(u,v) \neq 0$. We shall not pursue this direction any further at the moment. Recall that the coefficients of u^{2k} and v^{2k} must be positive, since G(u,v) > 0 for $(u,v) \neq 0$.

The next proposition shows that adding higher order terms to a function F as in Proposition 4.3 does not affect the conclusion of that proposition.

Proposition 4.4. Suppose F = 1 - G is exactly as in Proposition 4.3. For some $N \in \mathbb{N}$ let

$$G_{mod} = G + G_{2k+1} + \dots + G_{2k+N},$$

where for $j \geq 2k + 1$ the function $G_j(u, v)$ is a homogeneous polynomial of degree j. Define the obstacle \mathcal{O} as in Definition 8.1 of [WW23] by $x_1 = F_{mod}(\overline{x}) = 1 - G_{mod}(\overline{x})$, and take the incoming phase to be $\phi_i = -t + |x - b|$, b = (1, -1, 0) as before.

This class of problems satisfies [WW23, Assumption 2.10].

Sketch of proof. The proof is a straightforward repetition of the proof of Proposition 4.3, where one checks at each step that the terms coming from G in G_{mod} dominate the terms coming from $G_{2k+1} + \cdots + G_{2k+N}$ when \overline{x} is small. One point to note is that the

²⁰Recall $\alpha(\overline{x}) = \frac{1}{D(\overline{x})} (F(\overline{x}) - b_1, \overline{x} - \overline{b}).$

condition (4.1)(b) in Assumption 4.1 may fail now. For example, if $F_{mod} = 1 - G_{mod}$, where

$$G_{mod}(u, v) = (u^6 + c_1 u^4 v^2 + c_2 u^2 v^4 + c_3 v^6) - v u^6, c_i > 0,$$

then F_{mod} does not satisfy (4.1)(b). This causes no difficulty, since Proposition 4.2 was not used in the proof of Proposition 4.3. Step **4** of that proof proves not only that branching does not occur, but that $\pi_{\overline{x}}G_{\phi_i}$ is the graph of a function u = u(v) near $\overline{x} = 0$.

Remark. The argument of step 4 of the proof of Proposition 4.3, or a slight modification thereof, can be used to show that $\pi_{\overline{x}}G_{\phi_i}$ is the graph of a function u=u(v) near $\overline{x}=0$ for spherical waves diffracted by convex obstacles that are defined by some functions F(u,v) as in [WW23, Definition 8.1], which are not necessarily of the form (4.18), but which satisfy $|F_{uu}| \geq C|v||F_{uv}|$ near $\overline{x}=0$. We omit such verifications in the examples of the next section.

4.3. Examples of bad grazing sets. We now present examples showing that there exist C^{∞} strictly convex obstacles for which the grazing set for incoming spherical waves is continuous but not differentiable at a high-order grazing point. We also give such examples for incoming planar waves.




FIGURE 1. Cusped grazing set for the obstacle given by $x_1 = 1 - (x_2^4 + x_3^2)$ with light source b = (1, -1, 0).




FIGURE 2. Cusped grazing set for the obstacle given by $x_1 = 1 - (x_2^4 + x_2^2 x_3^2 + x_3^2)$ with light source b = (1, -1, 0).

4.3.1. Cusped grazing sets: examples where the grazing curve is continuous but not differentiable at v=0: spherical waves. The simplest example is probably F(u,v)=1-G(u,v), where $G(u,v)=u^4+v^2$. In this case the obstacle is strictly convex near x=(1,0,0), so F satisfies [WW23, Definition 8.1], but we don't have $\nabla^2 G(u,v)>0$ for $(u,v)\neq (0,0)$. Again we take the incoming phase to be $\phi_i=-t+|x-b|$, where $b=(1,\overline{b}), \overline{b}=(-1,0)$.

We have

$$\rho = (1, 0, 0, t^0, 0, -\overline{b}, -1) \in G_d^4 \setminus G^5.$$

The equation (4.19) is

$$3u^4 + 4u^3 + v^2 = 0.$$

This gives $u = -v^{2/3}(4+3u)^{-1/3} \Rightarrow |u| \sim v^{2/3}$ for (u,v) small. Expanding we obtain

$$u = -4^{-1/3}v^{2/3} + O(v^{4/3}),$$

which implies u'(0) does not exist.

Here is another example of a cusped grazing set, but where $\nabla^2 G(u,v) > 0$ for $(u,v) \neq (0,0)$, Take F(u,v) = 1 - G(u,v), where $G(u,v) = u^4 + u^2v^2 + v^2$. The function F satisfies Assumption 4.1. Here again $\underline{\rho} \in G_d^4 \setminus G^5$. Now the equation (4.19) is

$$3(u^4 + u^2v^2) + 4u^3 + 2uv^2 + v^2 = 0, (4.29)$$

SO

$$u^{3}(4+3u) = -v^{2}(1+2u+3u^{2}). (4.30)$$


Arguing as above, we find


$$u = -4^{-1/3}v^{2/3} + O(v^{4/3}). (4.31)$$

From (4.31) it is clear that the limit defining u'(0) does not exist. We can get more information if we rewrite (4.29) as

$$u[(4u^{2} + 2v^{2}) + 3(u^{3} + uv^{2}] := uB(u, v) = -v^{2},$$
(4.32)

and note that $B(u,v) \gtrsim u^2 + v^2 \sim v^{4/3}$ for (u,v) small. Now differentiate (4.32) with respect to v, divide by B, and use $|v/u^2| \sim v^{-1/3}$ to obtain $|u'(v)| \sim |v|^{-1/3}$ for $v \neq 0$. One can analyze u''(v) in a similar way to see that $0 < u'' \sim v^{-4/3}$ for $v \neq 0$, so the graph of u = u(v) is concave up viewed from the positive u-axis. Thus, u(v) "behaves" near v = 0 just like the function $w(v) = v^{2/3}$.








FIGURE 3. Cusped grazing set for the obstacle given by $x_1 = 1 - (x_2^4 + 2x_2^2x_3^2 + x_2x_3^2 + x_3^2)$ with light source at infinity.

4.3.2. Effect of changing source location on regularity of the grazing set. It is clear that the grazing set depends on source location. Here we show that the regularity of the grazing set can also depend on source location.

Again take F = 1 - G(u, v) where $G(u, v) = u^4 + v^2$, but now put the source at $c = (1, 0, 1) := (1, \overline{c})$, so the incoming phase is $\phi_i = -t + |x - c|$ and

$$\rho^* = (1, 0, 0, t^0, 0, -\overline{c}, -1) \in G_d^4 \setminus G^5.$$

The equation (4.19) is now

$$3u^4 + v^2 - 2v = 0,$$

which implies the grazing set is given near (u, v) = (0, 0) by $v = 2 - \sqrt{4 - 12u^4}$, a C^{∞} function. Recall that when b = (1, -1, 0) we get a cusped grazing set for this obstacle.²¹

4.3.3. Grazing curves that are continuous, but not differentiable at v=0: source at infinity. We return for the moment to the case of a wave source at infinity. Take F(u,v)=1-G(u,v), where $G(u,v)=u^4+2u^2v^2+uv^2+v^2$. The incoming phase is $\phi_i=-t+(u,v)\cdot\overline{\theta}$, where $\overline{\theta}=(1,0)$. Here the point


$$\rho = (1, 0, 0, t^0, 0, -\overline{b}, -1) \in G_d^4 \setminus G^5$$


and $\nabla^2 G(u,v) > 0$ for $(u,v) \neq (0,0)$. When the source is at infinity the grazing curve is defined by $\nabla G(u,v) \cdot \overline{\theta} = G_u = 0$ instead of (4.19). In this case we obtain

$$v^{2} + 4uv^{2} + 4u^{3} = 0$$
; that is $4u^{3} = -v^{2}(1 + 4u)$.

This is similar to (4.30). The same analysis shows we have a cusped grazing curve.

The grazing set for b = (1, 1, 0) has a cusp similar to that for b = (1, -1, 0).








FIGURE 4. C^1 grazing set for the obstacle given by $x_1 = 1 - (x_2^4 + x_2 x_3^4 + x_2^2 x_3^4 + x_3^2)$ with light source at infinity.

4.3.4. Grazing sets that are C^1 but not twice differentiable at v=0: source at infinity. Take F(u,v)=1-G(u,v), where $G(u,v)=u^4+uv^4+u^2v^4+v^2$. The incoming phase is $\phi_i=-t+(u,v)\cdot \overline{\theta}$, where $\overline{\theta}=(1,0)$. Here again $\underline{\rho}\in G_d^4\setminus G^5$ and $\nabla^2 G(u,v)>0$ for $(u,v)\neq (0,0)$. In this case the grazing curve is

$$G_u = 4u^3 + v^4 + 2uv^4 = 0 \Rightarrow 4u^3 = -v^4(1+2u) \Rightarrow u = 4^{-1/3}v^{4/3} + O(v^{8/3}),$$
 (4.33) so $|u| \sim |v|^{4/3}$.

Since u(0) = 0, this implies u'(0) = 0. We know u is C^{∞} for $v \neq 0$ (by the implicit function theorem), so now we have u is differentiable near v = 0.

To see that u is C^1 rewrite $G_u = 0$ as $u(4u^2 + 2v^4) = -v^4$, and differentiate with respect to v obtaining

$$u'(4u^{2} + 2v^{4}) + u(8uu' + 8v^{3}) = -4v^{3}.$$
(4.34)

Divide by $4u^2 + 2v^4$ and use (4.33) to see that

$$|u'(v)| \sim |v^3/u^2| \sim v^{1/3}.$$
 (4.35)

Thus u is C^1 .

From (4.35) it is clear that the limit defining u''(0) does not exist. We can get more information by differentiating (4.34) with respect to v to obtain

$$|u''(v)| \sim v^{-2/3}$$
 for $v \neq 0$.

5. The reflected flow map for general convex incoming waves

We show in this section that the computations in the previous section can be generalized to a larger class of incoming waves. Suppose \mathcal{O} is a convex obstacle given by

$$\mathcal{O} := \{ (x_1, \overline{x}) | x_1 < F(\overline{x}), \ \overline{x} \in \mathbb{R}^{n-1} \}$$

where $F: \mathbb{R}^{n-1} \to \mathbb{R}$ is a C^{∞} strictly concave function. We can arrange by translation and rotation of \mathcal{O} so that F has a max at $\overline{x} = 0$ with F(0) = 1. The tangent plane to $\partial \mathcal{O}$ at $(x_1, \overline{x}) = (1, 0)$ is then $x_1 = 1$.

We consider incoming phases of the form

$$\phi_i(x) = -t + \psi_i(x), \ (\partial_{x_1} \psi_i)^2 + |\nabla \psi_i|^2 = 1, \tag{5.1}$$

where ψ_i is a convex function. In what follows the notation ∇ always means $\nabla_{\overline{x}}$.

Assumption 5.1. Let F and \mathcal{O} be as just described. Assume there exists r > 0, an \mathbb{R}^n -open set $\Omega \ni (1,0)$, and a C^{∞} function $\psi : \Omega \to \mathbb{R}$ such that

(a)
$$\Omega \cap \partial \mathcal{O} = \{ (F(\overline{x}), \overline{x}) : \overline{x} \in B(0, r) \}$$

(b) $|\nabla \psi_i| = 1$ and ψ_i is convex on Ω .

Remark. We can obtain a large supply of functions ψ_i satisfying the conditions of Assumption 5.1 as follows. Let $\mathcal{C} \subset \mathbb{R}^n$ be any open convex set with C^{∞} boundary $\partial \mathcal{C}$. It is well-known that the distance-to-the-boundary function $\psi_i(x) = -\text{dist}(x, \partial C)$ is smooth, convex, and satisfies $|\nabla \psi_i| = 1$ for points of \mathcal{C} sufficiently near $\partial \mathcal{C}$. When $\partial \mathcal{C}$ is a plane, the incoming waves determined by (5.1) are plane waves; when $\partial \mathcal{C}$ is a sphere, the incoming waves are spherical waves, and so on.

If the incoming bicharacteristics γ_i hit the cotangent bundle on the space-time boundary at $(F(\overline{x}), \overline{x}, t, \xi_1^i(\overline{x}), \overline{\xi}^i(\overline{x}), -1)$, then there must hold

$$\xi_1^i(\overline{x}) = \partial_{x_1} \psi_i(F(\overline{x}), \overline{x}), \ \overline{\xi}^i(\overline{x}) = \nabla \psi_i(F(\overline{x}), \overline{x}).$$

The grazing set and the illuminable set are

$$G_{\phi_i} = \{ (F(\overline{x}), \overline{x}, t) | \langle \overline{\xi}^i, \nabla F \rangle - \xi_1^i = 0 \}, \ I_- = \{ (F(\overline{x}), \overline{x}, t) | \langle \overline{\xi}^i, \nabla F \rangle - \xi_1^i > 0 \}.$$
 (5.3)

Fix any T > 0 and any $r_1 < r$ for r as in Assumption 5.1. In (5.3) and the rest of this section we take $\overline{x} \in \overline{B(0,r_1)}$ and $t \in [-T,T]$.

Lemma 5.2. Let $\Psi(\overline{x}) := \psi_i(F(\overline{x}), \overline{x})$. If the incoming phase satisfies Assumption 5.1, then:

1. The matrix

$$\nabla^2 \Psi - \xi_1^i \nabla^2 F = \frac{\partial \overline{\xi}^i}{\partial \overline{x}} + \nabla F \otimes \nabla \xi_1^i$$

is symmetric and positive semi-definite.

2. The incoming directions ξ^i are non-focusing: for \overline{x}^* , \overline{x} in the \overline{x} -projection of $G_{\phi_i} \cup I$, the following holds

$$\langle (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}), \xi^i(\overline{x}^*) - \xi^i(\overline{x}) \rangle \ge 0.$$

Proof. 1. We have $\nabla \Psi = \xi_1^i \nabla F + \overline{\xi}^i$. Differentiate this to obtain the given identity. On the other hand a direct computation shows

$$\nabla^{2}\Psi = (\partial_{x_{1}}^{2}\psi_{i})\nabla F \otimes \nabla F + \nabla F \otimes \nabla \partial_{x_{1}}\psi_{i} + \nabla \partial_{x_{1}}\psi_{i} \otimes \nabla F + \nabla^{2}\psi_{i} + (\partial_{x_{1}}\psi_{i})\nabla^{2}F$$

$$= ((\nabla F)^{T} \quad I)\begin{pmatrix} \partial_{x_{1}}^{2}\psi_{i} & \nabla \partial_{x_{1}}\psi_{i} \\ (\nabla \partial_{x_{1}}\psi_{i})^{T} & \nabla^{2}\psi_{i} \end{pmatrix}\begin{pmatrix} \nabla F \\ I \end{pmatrix} + (\partial_{x_{1}}\psi_{i})\nabla^{2}F.$$

The result follows since the first matrix on the right is symmetric and positive semidefinite.

2. Using the formula $\xi^i(\overline{x}) = (\partial_{x_1}\psi_i, \nabla\psi_i)(F(\overline{x}), \overline{x})$ and the convexity of ψ_i , we find

$$\begin{split} &\langle (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}), \xi^i(\overline{x}^*) - \xi^i(\overline{x}) \rangle \\ &= -\langle (\partial_{x_1} \psi_i, \nabla \psi_i)(F(\overline{x}^*), \overline{x}^*), (F(\overline{x}), \overline{x}) - (F(\overline{x}^*), \overline{x}^*) \rangle \\ &- \langle (\partial_{x_1} \psi_i, \nabla \psi_i)(F(\overline{x}), \overline{x}), (F(\overline{x}^*), \overline{x}^*) - (F(\overline{x}), \overline{x}) \rangle \\ &\geq -(\psi_i(F(\overline{x}), \overline{x}) - \psi_i(F(\overline{x}^*), \overline{x}^*)) - (\psi_i(F(\overline{x}^*), \overline{x}^*) - \psi_i(F(\overline{x}), \overline{x})) = 0. \end{split}$$

This completes the proof.

Let $\xi^r(\overline{x})$ be the reflected covector field on the boundary, then

$$\xi^r - \xi^i = c(1, -\nabla F), \ |\xi^r| = 1.$$

From here we solve

$$\xi_1^r = \xi_1^i - \frac{2}{1 + |\nabla F|^2} (\xi_1^i - \langle \overline{\xi}^i, \nabla F \rangle),$$

$$\overline{\xi}^r = \overline{\xi}^i + \frac{2\nabla F}{1 + |\nabla F|^2} (\xi_1^i - \langle \overline{\xi}^i, \nabla F \rangle).$$
(5.4)

We obtain the following identities from (5.4):

$$\xi_1^i \nabla F + \overline{\xi}^i = \xi_1^r \nabla F + \overline{\xi}^r, \ \xi_1^r - \langle \nabla F, \overline{\xi}^r \rangle = -(\xi_1^i - \langle \overline{\xi}^i, \nabla F \rangle). \tag{5.5}$$

The reflected flow map is then given by

$$Z_r(s,\overline{x},t) = (F(\overline{x}) + 2s\xi_1^r(\overline{x}), \overline{x} + 2s\overline{\xi}^r(\overline{x}), t + 2s) \text{ for } s \in [0,\infty), \ \overline{x} \in B(0,r), \ t \in \mathbb{R}.$$

Proposition 5.3. Fix any $s_0 > 0$. The reflected flow map Z_r is a diffeomorphism from $[0, s_0] \times I_-$ onto its range and extends to a homeomorphism from $[0, s_0] \times (G_{\phi_i} \cup I_-)$ onto its range.

Proof. Injectivity. It suffices to show that

$$z(s, \overline{x}) := (F(\overline{x}), \overline{x}) + 2s\xi^r(\overline{x})$$

is injective from the (s, \overline{x}) -projection of $G_{\phi_i} \cup I_-$ to its range. Suppose the contrary, then there exists $(s, \overline{x}) \neq (s^*, \overline{x}^*)$ in the (s, \overline{x}) -projection of $G_{\phi_i} \cup I_-$ such that²²

$$z(s, \overline{x}) = z(s^*, \overline{x}^*), \text{ i.e., } (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}) = 2s\xi^r(\overline{x}) - 2s^*\xi^r(\overline{x}^*).$$
 (5.6)

Using the second equation in (5.6) we find

$$\langle (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}), \xi^r(\overline{x}^*) - \xi^r(\overline{x}) \rangle = -2(s + s^*)(1 - \langle \xi^r(\overline{x}^*), \xi^r(\overline{x}) \rangle) \le 0 \quad (5.7)$$

This assumption implies $\overline{x} \neq \overline{x}^*$.

with equality holding if and only if

$$\xi^r(\overline{x}^*) = \xi^r(\overline{x}). \tag{5.8}$$

On the other hand, we show that for all $\overline{x}, \overline{x}^*$, there holds

$$\langle (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}), \xi^r(\overline{x}^*) - \xi^r(\overline{x}) \rangle \ge 0, \tag{5.9}$$

with equality holding if and only if

$$\xi_1^i(\overline{x}) - \langle \overline{\xi}^i(\overline{x}), \nabla F(\overline{x}) \rangle = \xi_1^i(\overline{x}^*) - \langle \overline{\xi}^i(\overline{x}^*), \nabla F(\overline{x}^*) \rangle = 0, \text{ and }$$

$$\langle (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}), \xi^i(\overline{x}^*) - \xi^i(\overline{x}) \rangle = 0.$$
(5.10)

Indeed, we have

$$\langle (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}), \xi^r(\overline{x}^*) - \xi^r(\overline{x}) \rangle$$

$$\stackrel{(5.5)}{=} (F(\overline{x}^*) - F(\overline{x}))(\xi_1^r(\overline{x}^*) - \xi_1^r(\overline{x})) + \langle \overline{x}^* - \overline{x}, \overline{\xi}^i(\overline{x}^*) - \overline{\xi}^i(\overline{x}) \rangle + (\xi_1^r(\overline{x}^*) - \xi_1^i(\overline{x}^*))\langle \nabla F(\overline{x}^*), \overline{x} - \overline{x}^* \rangle + (\xi_1^r(\overline{x}) - \xi_1^i(\overline{x}))\langle \nabla F(\overline{x}), \overline{x}^* - \overline{x} \rangle \geq (F(\overline{x}^*) - F(\overline{x}))(\xi_1^r(\overline{x}^*) - \xi_1^r(\overline{x})) + \langle \overline{x}^* - \overline{x}, \overline{\xi}^i(\overline{x}^*) - \overline{\xi}^i(\overline{x}) \rangle + (\xi_1^r(\overline{x}^*) - \xi_1^i(\overline{x}^*))(F(\overline{x}) - F(\overline{x}^*)) + (\xi_1^r(\overline{x}) - \xi_1^i(\overline{x}))(F(\overline{x}^*) - F(\overline{x})) = \langle (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}), \xi^i(\overline{x}^*) - \xi^i(\overline{x}) \rangle \geq 0.$$
 (5.11)

Here we used (5.4) and the definition of $G_{\phi_i} \cup I_-$, which imply that

$$\xi_{1}^{r}(\overline{x}^{*}) - \xi_{1}^{i}(\overline{x}^{*}) = \frac{2}{1 + |\nabla F(\overline{x}^{*})|^{2}} \left(\langle \overline{\xi}^{i}(\overline{x}^{*}), \nabla F(\overline{x}^{*}) \rangle - \xi_{1}^{i}(\overline{x}^{*}) \right) \ge 0,$$

$$\xi_{1}^{r}(\overline{x}) - \xi_{1}^{i}(\overline{x}) = \frac{2}{1 + |\nabla F(\overline{x})|^{2}} \left(\langle \overline{\xi}^{i}(\overline{x}), \nabla F(\overline{x}) \rangle - \xi_{1}^{i}(\overline{x}) \right) \ge 0.$$
(5.12)

Since F is strictly concave, with (5.12) we conclude that overall equality holds in (5.11) if and only if

$$\xi_1^i(\overline{x}) - \langle \overline{\xi}^i(\overline{x}), \nabla F(\overline{x}) \rangle = \xi_1^i(\overline{x}^*) - \langle \overline{\xi}^i(\overline{x}^*), \nabla F(\overline{x}^*) \rangle = 0, \text{ and } \\ \langle (F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}), \xi^i(\overline{x}^*) - \xi^i(\overline{x}) \rangle = 0.$$

Combining (5.7) and (5.9), we conclude that (5.8) and (5.10) must be true. The first condition in (5.10), (5.4), and (5.8) together imply that

$$\xi^{i}(\overline{x}^{*}) = \xi^{r}(\overline{x}^{*}) = \xi^{r}(\overline{x}) = \xi^{i}(\overline{x}). \tag{5.13}$$

This and (5.6) give

$$(F(\overline{x}^*) - F(\overline{x}), \overline{x}^* - \overline{x}) = 2(s - s^*)\xi^i(\overline{x}). \tag{5.14}$$

From (5.10) we have $\xi_1^i(\overline{x}) = \langle \overline{\xi}^i(\overline{x}), \nabla F(\overline{x}) \rangle$. Multiplying this equation by $2(s - s^*)$ and using (5.14), we obtain

$$F(\overline{x}^*) - F(\overline{x}) = \langle \nabla F(\overline{x}), \overline{x}^* - \overline{x} \rangle.$$

Since F is strictly concave, this can be true only if $\overline{x} = \overline{x}^*$, which implies that $(s, \overline{x}) = (s^*, \overline{x}^*)$ by (5.6). This contradicts the assumption.

The map Z_r is continuous and injective on the compact domain $[0, s_0] \times (G_{\phi_i} \cup I_-)$, and hence is a homeomorphism on this domain.

Diffeomorphism. Recall that

$$A = I - \frac{\overline{\xi}^r \otimes \nabla F}{\xi_1^r} + 2s \left(\frac{\partial \overline{\xi}^r}{\partial \overline{x}} - \frac{\overline{\xi}^r \otimes \nabla \xi_1^r}{\xi_1^r} \right).$$

Differentiating $|\xi^r|^2 = 1$ we find

$$A = I - \frac{\overline{\xi}^r \otimes \nabla F}{\xi_1^r} + 2s \left(I + \frac{\overline{\xi}^r \otimes \overline{\xi}^r}{(\xi_1^r)^2} \right) \frac{\partial \overline{\xi}^r}{\partial \overline{x}}.$$
 (5.15)

Recalling (5.5), we differentiate

$$\overline{\xi}^r = \overline{\xi}^i + (\xi_1^i - \xi_1^r) \nabla F$$

to obtain

$$\frac{\partial \overline{\xi}^r}{\partial \overline{x}} = \frac{\partial \overline{\xi}^i}{\partial \overline{x}} + \nabla F \otimes \nabla (\xi_1^i - \xi_1^r) + (\xi_1^i - \xi_1^r) \nabla^2 F.$$

Differentiate the defining formula (5.4) of ξ_1^r to find

$$\begin{split} \nabla(\xi_1^i - \xi_1^r) = & 2\frac{\nabla \xi_1^i - \nabla F \frac{\partial \overline{\xi}^i}{\partial \overline{x}} - \overline{\xi}^i \nabla^2 F}{1 + |\nabla F|^2} - 2\frac{\overline{\xi}^r - \overline{\xi}^i}{1 + |\nabla F|^2} \nabla^2 F \\ = & - 2\frac{\overline{\xi}^i + \xi_1^i \nabla F}{\xi_1^i (1 + |\nabla F|^2)} \frac{\partial \overline{\xi}^i}{\partial \overline{x}} - \frac{2\overline{\xi}^r \nabla^2 F}{1 + |\nabla F|^2}. \end{split}$$

Therefore we find

$$\begin{split} \frac{\partial \overline{\xi}^r}{\partial \overline{x}} = & \frac{\partial \overline{\xi}^i}{\partial \overline{x}} - \frac{2\nabla F \otimes (\overline{\xi}^i + \xi_1^i \nabla F)}{\xi_1^i (1 + |\nabla F|^2)} \frac{\partial \overline{\xi}^i}{\partial \overline{x}} - \frac{2}{1 + |\nabla F|^2} (\nabla F \otimes \overline{\xi}^r) \nabla^2 F + (\xi_1^i - \xi_1^r) \nabla^2 F \\ = & \left(I - \frac{2\nabla F \otimes (\overline{\xi}^i + \xi_1^i \nabla F)}{\xi_1^i (1 + |\nabla F|^2)} \right) \frac{\partial \overline{\xi}^i}{\partial \overline{x}} + (\xi_1^i - \xi_1^r) \left(I - \frac{2\nabla F \otimes \overline{\xi}^r}{(1 + |\nabla F|^2)(\xi_1^i - \xi_1^r)} \right) \nabla^2 F. \end{split}$$

In the following we denote

$$B := I - \frac{\overline{\xi}^r \otimes \nabla F}{\xi_1^r}, \ C := I + \frac{\overline{\xi}^r \otimes \overline{\xi}^r}{(\xi_1^r)^2},$$

$$K := \left(I - \frac{2\nabla F \otimes (\overline{\xi}^i + \xi_1^i \nabla F)}{\xi_1^i (1 + |\nabla F|^2)}\right) \frac{\partial \overline{\xi}^i}{\partial \overline{x}},$$

$$L := (\xi_1^i - \xi_1^r) \left(I - \frac{2\nabla F \otimes \overline{\xi}^r}{(1 + |\nabla F|^2)(\xi_1^i - \xi_1^r)}\right) \nabla^2 F.$$

This allows us to rewrite A as in (5.15) as

$$A = B + 2sC\frac{\partial \overline{\xi}^{r}}{\partial \overline{x}} = B + 2sC(K + L) = B[I + 2sB^{-1}C(B^{-1})^{T}(B^{T}K + B^{T}L)].$$

We compute

$$B^{T}K = \left(I - \frac{\nabla F \otimes \overline{\xi}^{r}}{\xi_{1}^{r}} - \frac{2\nabla F \otimes (\overline{\xi}^{i} + \xi_{1}^{i}\nabla F)}{\xi_{1}^{i}(1 + |\nabla F|^{2})} + \frac{2\langle \overline{\xi}^{r}, \nabla F \rangle \nabla F \otimes (\overline{\xi}^{i} + \xi_{1}^{i}\nabla F)}{\xi_{1}^{i}\xi_{1}^{r}(1 + |\nabla F|^{2})}\right) \frac{\partial \overline{\xi}^{i}}{\partial \overline{x}}.$$

By the definition of $\overline{\xi}^r$ (5.4) we have

$$\begin{split} \langle \overline{\xi}^r, \nabla F \rangle = & \langle \overline{\xi}^i, \nabla F \rangle + |\nabla F|^2 (\xi_1^i - \xi_1^r) \\ = & \frac{1 + |\nabla F|^2}{2} (\xi_1^r - \xi_1^i) + |\nabla F|^2 (\xi_1^i - \xi_1^r) + \xi_1^i \\ = & \frac{1 + |\nabla F|^2}{2} \xi_1^i + \frac{1 - |\nabla F|^2}{2} \xi_1^r. \end{split}$$

This implies²³

$$B^{T}K = \left(I - \frac{\nabla F \otimes \overline{\xi}^{r}}{\xi_{1}^{r}} + \frac{\xi_{1}^{i} - \xi_{1}^{r}}{\xi_{1}^{i}\xi_{1}^{r}} \nabla F \otimes (\overline{\xi}^{i} + \xi_{1}^{i}\nabla F)\right) \frac{\partial \overline{\xi}^{i}}{\partial \overline{x}}$$

$$= \left(I + \frac{\nabla F \otimes ((\xi_{1}^{i} - \xi_{1}^{r})\nabla F - \overline{\xi}^{r})}{\xi_{1}^{i}}\right) \frac{\partial \overline{\xi}^{i}}{\partial \overline{x}}$$

$$= \left(I - \frac{\nabla F \otimes \overline{\xi}^{i}}{\xi_{1}^{i}}\right) \frac{\partial \overline{\xi}^{i}}{\partial \overline{x}} = \frac{\partial \overline{\xi}^{i}}{\partial \overline{x}} + \nabla F \otimes \nabla \xi_{1}^{i} = \nabla^{2}\Psi - \xi_{1}^{i}\nabla^{2}F.$$

$$(5.16)$$

Now apply Lemma 5.2 to see that B^TK is symmetric and positive semi-definite.

We now simplify B^TL . Notice that

$$(B^{-1})^T = I + \frac{\nabla F \otimes \overline{\xi}^r}{\xi_1^r - \langle \nabla F, \overline{\xi}^r \rangle} = I - \frac{\nabla F \otimes \overline{\xi}^r}{\xi_1^i - \langle \nabla F, \overline{\xi}^i \rangle}.$$

On the other hand, using the definition (5.4) of ξ_1^r , we have

$$L = (\xi_1^i - \xi_1^r) \left(I - \frac{\nabla F \otimes \overline{\xi}^r}{\xi_1^i - \langle \overline{\xi}^i, \nabla F \rangle} \right) \nabla^2 F = (\xi_1^i - \xi_1^r) B^{-T} \nabla^2 F.$$

Thus we have

$$B^{T}L = (\xi_{1}^{i} - \xi_{1}^{r})\nabla^{2}F \ge 0.$$
 (5.17)

²³To obtain the second equality in (5.16), use (5.5) to replace $\overline{\xi}^i + \xi_1^i \nabla F$ by $\overline{\xi}^r + \xi_1^r \nabla F$.

Arguing as in the case of spherical waves, we obtain

$$\begin{split} j(s,\overline{x},t) = & 2\xi_1^r \det A = 2\xi_1^r \det(B) \det(I + 2sB^{-1}C(K+L)) \\ = & 2(\langle \overline{\xi}^i, \nabla F \rangle - \xi_1^i) \det(I + 2sB^{-1}C(B^{-1})^T(B^TK + B^TL)) \\ \geq & 2(\langle \overline{\xi}^i, \nabla F \rangle - \xi_1^i) > 0. \end{split}$$

This completes the proof.

References

- [Che96] Christophe Cheverry. Propagation d'oscillations près d'un point diffractif. J. Math. Pures Appl., 75:419–467, 1996.
- [Mel75] Richard B. Melrose. Microlocal parametrices for diffractive boundary value problems. Duke Math. J., 42:605–635, 1975.
- [MS78] Richard B. Melrose and Johannes Sjöstrand. Singularities of boundary value problems. I. Communications on Pure and Applied Mathematics, 31:593–617, 1978.
- [RV73] A. Wayne Roberts and Dale E. Varberg. Convex Functions. Academic Press, 1973.
- [Wil22] Mark Williams. Solving eikonal equations by the method of characteristics. https://markwilliams.web.unc.edu/wp-content/uploads/sites/19674/2022/01/eikonal.pdf, 2022.
- [WW23] Jian Wang and Mark Williams. Transport of nonlinear oscillations along rays that graze a convex obstacle to any order. *Preprint*, 2023.

Email address: wangjian@ihes.fr

Institut des Hautes Études Scientifiques, Bures-sur-Yvette, France 91893

 $Email\ address: {\tt williams@math.unc.edu}$

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA, CHAPEL HILL, NC 27514