ON THE NIELSEN-THOMSEN SEQUENCE

LAURENT CANTIER

ABSTRACT. The Nielsen-Thomsen sequence plays a pivotal role in refining invariants for C*-algebras beyond the Elliott classification framework. This paper revisits the sequence, introducing the concepts of Nielsen-Thomsen bases, rotation maps and diagonalisable morphisms, to better understand its unnatural splitting. These insights enable novel comparison methods for *-homomorphisms at the level of the Hausdorffized algebraic K_1 -groups, and subsequently the Hausdorffized unitary Cuntz group.

We apply our methods to classification via the Hausdorffized unitary Cuntz semigroup. In particular, we present a new proof of the non-isomorphism between the two AT-algebras of [19]. We also exhibit several pairs of (non-unitarily equivalent) *-homomorphisms with domain $C(\mathbb{T})$.

1. Introduction

The classification program for nuclear C*-algebras has seen remarkable progress in the past decade, building on Elliott's groundbreaking conjecture that simple, separable, unital, nuclear C*algebras could be classified by K-theoretic and tracial invariants. This vision was fully realized for the class of Z-stable algebras satisfying the Universal Coefficient Theorem (UCT) through the combined efforts of many researchers, including the breakthrough works of [14], or [20, 21]. These developments culminated in the classification of simple, separable, unital, nuclear, Z-stable C*-algebras satisfying the UCT. See [27] for a general overview and [9] for a remarkable detailed and innovative exposition on the matter. Nevertheless, classification outside the simple and \mathcal{Z} -stable setting presents additional challenges that have motivated the development of refined invariants. For instance, all the way back to the classification of certain *-homomorphisms between circle algebras done in [22], Nielsen and Thomsen present a split-exact sequence referred now to as the Nielsen-Thomsen sequence. This sequence, introduced through the study of the Hausdorffized algebraic K₁-group and its relationship with the de la Harpe-Skandalis determinant, emerged as a particularly intriguing tool in this context. While their seminal work revealed its potential for distinguishing *-homomorphisms that share identical Elliott invariants, several fundamental questions and applications, e.g. on the unnatural splitting of the sequence, have remained unexplored.

The past ten years have seen significant advances in the classification of non-simple C^* -algebras. We recall Robert's classification of inductive limit of one-dimensional NCCW-complexes with trivial K_1 -groups by means of the Cuntz semigroup. See [23]. Another approach began with classification of AT algebras of real rank zero by means of the traditional Elliott invariant. See [15, 16]. This approach and its techniques have been expanded over the years and the work of many, resulting in Gong, Jiang, and Li's classification of AH algebras with the ideal property by means of a refined version of the Elliott invariant. See [19]. These works highlighted the crucial role played by the

1

The author was supported by the Spanish Ministry of Universities and the European Union-NextGenerationEU through a Margarita Salas grant. partially supported by MINECO (grant No. PID2020-113047GB-I00), and by the Departament de Recerca i Universitats de la Generalitat de Catalunya (grant No. 2021-SGR-01015).

ideal structure and the information of the Nielsen-Thomsen sequence for classification. As a result, the recent developments of the Cuntz semigroup and its (Hausdorffized) unitary version(s) as classification invariants have provided new perspectives on these challenges. See [5, 6, 7].

In this paper, we explore the structural and internal properties of the Hausdorffized unitary Cuntz semigroup. Our investigation begins with a systematic study of the Nielsen-Thomsen sequence to better understand the information encoded at the morphism level. To achieve this, we introduce new concepts, including Nielsen-Thomsen bases and rotation maps. These tools enable us to reinterpret the 'unnatural' splitting phenomenon through matrix representations of *-homomorphisms at the level of the Hausdorffized K_1 -group and diagonalisable morphisms. This approach quantifies the effect of *-homomorphisms on the Nielsen-Thomsen bases by measuring the disturbance caused by these maps, allowing for construction of a metric \mathfrak{d} to compare *-homomorphisms at the level of the Hausdorffized algebraic K_1 -group. While this framework provides a new perspective on distinguishing *-homomorphisms between C*-algebras that agree on the traditional Elliott invariant, it also establishes in turn, novel comparison methods the Hausdorffized unitary Cuntz group and its morphisms. Finally, we end the manuscript by applying the proposed methods to distinguishing C*-algebras and *-homomorphisms with domain $C(\mathbb{T})$ by means of the Hausdorffized unitary Cuntz group. We gather them all in the following theorem.

Theorem. (i) The (non-simple) A \mathbb{T} -algebras constructed in [19] agree on the Elliott invariant, the Hausdorffized algebraic K_1 -group the Cuntz semigroup and its unitary version.

Yet, they are distinguished by the Hausdorffized unitary Cuntz semigroup.

(ii) There exists a family $\{\varphi_k \colon C(\mathbb{T}) \longrightarrow C([0,1]) \otimes M_{2^{\infty}}\}_{k \in \mathbb{N}}$ of *-homomorphisms agreeing on the Elliott invariant, the Cuntz semigroup and its unitary version.

Yet, we compute that $\mathfrak{d}(\overline{K}_1(\varphi_k), \overline{K}_1(\varphi_l)) = \frac{|k-l|}{2}$ and hence, they are pairwise distinguished by the Hausdorffized algebraic K_1 -group.

(iii) There exist two *-homomorphisms $\varphi_u, \varphi_v \colon C(\mathbb{T}) \longrightarrow A$, where A is an AI-algebra, agreeing on the Elliott invariant, the Cuntz semigroup and its unitary version and the \overline{K}_1 -group.

Yet, they are distinguished by the Hausdorffized unitary Cuntz semigroup.

Organization of the paper. The paper is organized as follows. In Section 2, we introduce the necessary background on Nielsen-Thomsen bases and develop the theory of rotation maps for *-homomorphisms. This yields to the construction of a relevant metric to compare *-homomorphisms at the level of their Hausdorffized algebraic unitary group. Section 3 is devoted to refined Cuntz semigroups and the comparison theory of morphisms, building on recent developments in the theory of Cu-semigroups. Finally, in Section 4, we demonstrate the effectiveness of our methods by providing alternative proofs of the aforementioned classification results, illustrating how our framework naturally captures and explains these phenomena through the lens of the Nielsen-Thomsen sequence and its associated invariants.

Acknowledgments. The author would like to thank B. Jacelon for inspiring discussions around unitary elements of C^* -algebras and the de la Harpe-Skandalis determinant.

2. The Rotation map of *-homomorphisms

In [22], Nielsen and Thomsen pointed out that the Hausdorffized algebraic K_1 -group contains information unseen by the Elliott invariant, and hence, is essential for classifying *-homomorphisms. They observed that this group splits between the K_1 -group and traces together with projections, giving rise to the *Nielsen-Thomsen sequence*. Notably, they showed that this sequence splits 'unnaturally' at the morphism level. In this section, we delve deeper into this phenomenon by introducing the notions of Nielsen-Thomsen bases, rotation maps and diagonalisable morphisms. As a consequence, we are then able to quantify how *-homomorphisms affect the splitting behaviour via metrics that we construct. Let us first recall key preliminary facts around this sequence.

• The de la Harpe-Skandalis determinant. For a unital C*-algebra A, we denote the subgroup of A generated by additive commutators [a,b] := ab - ba by [A,A] and we define the *universal trace on* A to be the quotient map $\operatorname{Tr}: A \longrightarrow A/\overline{[A,A]}$. We extend this (continuous) tracial linear map to $M_{\infty}(A)$ in a canonical way that we also denote $\operatorname{Tr}: M_{\infty}(A) \longrightarrow A/\overline{[A,A]}$.

Following the seminal work of [13], for any smooth path $\xi:[0,1]\longrightarrow U_0^\infty(A)$, we define

$$\widehat{\Delta}(\xi) := \frac{1}{2i\pi} \int_0^1 \operatorname{Tr}(\xi'(t)\xi^{-1}(t)) dt$$

Note that the image $\widehat{\Delta}(\xi)$ lies in fact in the quotient $A_{\rm sa}/A_0$ where $A_0 := \{h \in A_{\rm sa} \mid {\rm Tr}(h) = 0\}$ and it is well-known that $A_{\rm sa}/A_0$ can be identified with Aff $T_1(A)$ by sending [h] to ${\rm ev}_h : \tau \in T_1(A) \mapsto \tau(h) \in \mathbb{R}$, where $T_1(A)$ denotes the (bounded) tracial states of A.

Therefore, the application $\widehat{\Delta}$ naturally induces a group morphism $\underline{\operatorname{Tr}}: \mathrm{K}_0(A) \longrightarrow A_{\mathrm{sa}}/A_0 \simeq \mathrm{Aff}\, T_1(A)$ by sending $[p] \mapsto \mathrm{Tr}(p) \mapsto \mathrm{ev}_p$ for any projection $p \in M_\infty(A)$. (This is done via the identification $\mathrm{K}_0(A) \simeq \pi_1(U_\infty^0(A))$.) We now define the de la Harpe-Skandalis determinant as

$$\Delta: U_0^{\infty}(A) \longrightarrow \operatorname{Aff} T_1(A) / \operatorname{im}(\underline{\operatorname{Tr}})$$

$$u \longmapsto [\widehat{\Delta}(\xi_u)]$$

where ξ_u is any smooth path connecting the identity to u. In particular, $\Delta(e^{2i\pi h}) = [\operatorname{ev}_h]$ for any self-adjoint $h \in M_{\infty}(A)$.

Even though it was observed in the original work de la Harpe and Skandalis that $\overline{\ker \Delta} \subseteq \overline{DU_0^\infty(A)}$ (see [13, Proposition 4]), there exist cases where $DU_0^\infty(A) \subsetneq \ker \Delta \subsetneq \overline{DU_0^\infty(A)}$. A fortiori, the group morphism Δ is not continuous in general. However, Thomsen introduced a non-stable version of the determinant to obtain a continuous version. Let $\overline{K_0(A)} := \overline{\operatorname{im}(\underline{\operatorname{Tr}})}$ denotes the closure of the image of $\underline{\operatorname{Tr}}$ in $A_{\operatorname{sa}}/A_0$. We now define the non-stable de la Harpe-Skandalis determinant

$$\overline{\Delta}: U_0^{\infty}(A) \longrightarrow \operatorname{Aff} T_1(A)/\overline{\mathrm{K}_0(A)}$$

$$u \longmapsto [\Delta(u)]$$

to be the composition of Δ with the canonical projection of Aff $T_1(A)/\operatorname{im}(\underline{\operatorname{Tr}})$ \longrightarrow Aff $T_1(A)/\overline{\operatorname{im}(\underline{\operatorname{Tr}})}$. It is shown in [26, Theorem 3.2] that $\overline{\Delta}$ is a surjective continuous group homomorphism satisfying that

$$U_0^{\infty}(A)/\overline{DU_0^{\infty}(A)} \stackrel{\overline{\Delta}}{\simeq} \operatorname{Aff} T_1(A)/\overline{\mathrm{K}_0(A)}.$$

For convenience, we may write $H(A) := Aff T_1(A) / \overline{K_0(A)}$.

• The Nielsen-Thomsen sequence. Let A be a unital C*-algebra. It is known that $DU(A) \subseteq DU_0(M_3(A)) \subseteq U_0^{\infty}(A)$. We deduce that $\overline{DU^{\infty}(A)}$ is a normal subgroup of $U_0^{\infty}(A)$ and we obtain a canonical extension

$$0 \longrightarrow U_0^{\infty}(A)/\overline{DU^{\infty}(A)} \stackrel{i}{\longrightarrow} U^{\infty}(A)/\overline{DU^{\infty}(A)} \stackrel{\pi}{\longrightarrow} U^{\infty}(A)/\overline{U_0^{\infty}(A)} \longrightarrow 0$$

The middle term is often referred to as the *Hausdorffized algebraic* K_1 -group that we will denote by \overline{K}_1 to ease the notations.

It is readily checked that $U_0^{\infty}(A)/\overline{DU^{\infty}(A)}$ is a divisible subgroup of $U^{\infty}(A)/\overline{DU^{\infty}(A)}$. Therefore, the latter extension splits. Finally, by using the non-stable de la Harpe-Skandalis determinant, we obtain the extension referred to as the *Nielsen-Thomsen sequence*

$$0 \longrightarrow \mathrm{H}(A) \stackrel{i}{\longrightarrow} \overline{\mathrm{K}}_1(A) \stackrel{\pi}{\longrightarrow} \mathrm{K}_1(A) \longrightarrow 0$$

where $H(A) := Aff T_1(A) / \overline{K_0(A)}$.

It is worth mentioning that under a K₁-surjectivity assumption (e.g. in the stable rank one case), we may work in the C*-algebra itself instead of matrices since we have that $H(A) \simeq U_0(A)/\overline{DU(A)}$, $\overline{K}_1(A) \simeq U(A)/\overline{DU(A)}$ and $K_1(A) \simeq U(A)/U_0(A)$.

We have seen that the Hausdorffized algebraic K_1 -group of any C^* -algebra A is entirely recovered by the original Elliott invariant, since we have an isomorphism $\overline{K}_1(A) \simeq H(A) \oplus K_1(A)$. Nevertheless, Nielsen and Thomsen showed in [22] that the sequence splits 'unnaturally' at the level of morphisms. In other words, given a *-homomorphism $\phi \colon A \longrightarrow B$, the data of $H(\phi)$ and $K_1(\phi)$, where $H(\phi) \colon \operatorname{Aff} T_1(A)/\overline{K_0(A)} \longrightarrow \operatorname{Aff} T_1(B)/\overline{K_0(B)}$ is the induced morphism by ϕ , is (in general) not enough to recover $\overline{K}_1(\phi)$. We can restate the above as follows.

- (i) For any C*-algebra A, the group $\overline{K}_1(A)$ is split.
- (ii) For any *-homomorphism $\phi \colon A \longrightarrow B$, the group morphism $\overline{K}_1(\phi)$ need not be diagonalisable. As a conclusion, it is necessary to incorporate the functor \overline{K}_1 to classify a larger class of *-homomorphisms, and C*-algebras that cannot be classified via the original Elliott invariant.
- Nielsen-Thomsen bases and the rotation map. We shed new light on the unnaturalness of the Nielsen-Thomsen sequence through *Nielsen-Thomsen bases*. These bases provide a systematic way to analyse the splitting behavior of the sequence at the morphism level. We are able to explicitly compute a matrix representation of *-homomorphisms at the level of the \overline{K}_1 -group in terms of their value on traces, K_1 -group and what we refer to as rotation maps with respect to the bases. This approach not only clarifies why the sequence splits unnaturally, but also allows us to quantify and interpret the deviations caused by *-homomorphisms.

Notation 2.1. For any unitary element u of a C*-algebra A, we denote its respective unitary equivalence classes by $\overline{u} := [u]_{\overline{K}_1(A)}$ and $[u] := [u]_{K_1(A)}$.

Definition 2.2. Let A be a unital C*-algebra. A Nielsen-Thomsen basis of A is a set $\{c_k \in U^{\infty}(A)\}_{k \in K_1(A)}$ of elements in $U^{\infty}(A)$ indexed by $K_1(A)$ such that the mapping

$$s_A \colon \mathrm{K}_1(A) \longrightarrow \overline{\mathrm{K}}_1(A)$$

$$k \longmapsto \overline{c_k}$$

is a group morphism satisfying $\pi_A \circ s_A = \mathrm{id}_{\mathrm{K}_1(A)}$. By convention, we always fix $c_0 := 1_A$.

Proposition 2.3. Let A be a unital C^* -algebra. Then

- (i) A admits Nielsen-Thomsen bases.
- (ii) Any Nielsen-Thomsen basis $\mathcal{C} := \{c_k\}_{k \in \mathrm{K}_1(A)}$ of A induces a unique group isomorphism $\nu_{\mathcal{C}} \colon \overline{\mathrm{K}}_1(A) \simeq \mathrm{H}(A) \oplus \mathrm{K}_1(A)$ sending $\overline{u} \mapsto (\overline{\Delta} \left(\begin{smallmatrix} u & 0 \\ 0 & c_{[u]}^* \end{smallmatrix}\right), [u])$.
 - (ii') Any Nielsen-Thomsen basis of A induces a unique retract $r \colon \overline{K}_1(A) \longrightarrow H(A)$.

Proof. These follow from standard results about split-extensions of abelian groups and are left for the reader to check. \Box

Conversely, it is readily checked that any group isomorphism $\nu \colon \overline{\mathrm{K}}_1(A) \simeq \mathrm{H}(A) \oplus \mathrm{K}_1(A)$ or equivalently, any retract $r \colon \overline{\mathrm{K}}_1(A) \longrightarrow \mathrm{H}(A)$ of $i \colon \mathrm{H}(A) \longrightarrow \overline{\mathrm{K}}_1(A)$ induces a unique Nielsen-Thomsen basis.

Definition 2.4. Let A, B be C*-algebras and $C = \{c_k\}_{k \in K_1(A)}, \mathcal{D} = \{d_k\}_{k \in K_1(B)}$ be Nielsen-Thomsen bases of A, B.

For any group morphism $\beta \colon \overline{\mathrm{K}}_1(A) \longrightarrow \overline{\mathrm{K}}_1(B)$, we define the matrix of β in bases \mathcal{CD} to be the unique 2-by-2 matrix (with group morphisms entries) such that the following diagram commutes

$$\begin{array}{c|c} \overline{\mathrm{K}}_{1}(A) & \xrightarrow{\nu_{\mathcal{C}}} & \mathrm{H}(A) \oplus \mathrm{K}_{1}(A) \\ \beta & & & & & \mathrm{Mat}_{\mathcal{CD}}(\beta) \\ \hline \overline{\mathrm{K}}_{1}(B) & \xrightarrow{\nu_{\mathcal{D}}} & \mathrm{H}(B) \oplus \mathrm{K}_{1}(B) \end{array}$$

where $\nu_{\mathcal{C}}, \nu_{\mathcal{D}}$ are the unique isomorphisms induced by the respective Nielsen-Thomsen bases.

We say that β is Nielsen-Thomsen diagonalisable if, there exist Nielsen-Thomsen bases such that the matrix of β is diagonal.

Theorem 2.5. Let $\phi: A \longrightarrow B$ be a *-homomorphism between unital C*-algebras. Let $\mathcal{C} = \{c_k\}_{k \in \mathrm{K}_1(A)}, \mathcal{D} = \{d_k\}_{k \in \mathrm{K}_1(B)}$ be Nielsen-Thomsen bases of A, B.

Let s_A, r_B be the respective section and retract induced by the bases. We compute that

$$r_B \circ \overline{\mathrm{K}}_1(\phi) \circ s_A \colon \mathrm{K}_1(A) \longrightarrow \mathrm{H}(B)$$

$$k \longmapsto \overline{\Delta} \begin{pmatrix} \phi(c_k) & 0 \\ 0 & d^*_{[\mathrm{K}_1(\phi)(k)]} \end{pmatrix}$$

which we write $R_{\mathcal{CD}}(\phi)$ and refer to as the rotation map of ϕ with respect to the bases \mathcal{CD} .

Furthermore, we have that

$$\operatorname{Mat}_{\mathcal{CD}}(\overline{K}_1(\phi)) = \begin{pmatrix} H(\phi) & R_{\mathcal{CD}}(\phi) \\ 0 & K_1(\phi) \end{pmatrix}$$

Proof. We know that ϕ induces a morphism $(H(\phi), \overline{K}_1(\phi), K_1(\phi))$ between the Nielsen-Thomsen sequences of A and B. (Observe that this triple is a priori not compatible with the chosen splits.) This is equivalent to have that $\overline{K}_1(\phi) \circ i_A = i_B \circ H(\phi)$ and $K_1(\phi) \circ \pi_A = \pi_B \circ \overline{K}_1(\phi)$.

As a result, we obtain that

$$\begin{cases} H(\phi) = r_B \circ i_B \circ H(\phi) = r_B \circ \overline{K}_1(\phi) \circ i_A \\ K_1(\phi) = K_1(\phi) \circ \pi_A \circ s_A = \pi_B \circ \overline{K}_1(\phi) \circ s_A \end{cases}$$

from which the computations of the entries of $\operatorname{Mat}_{\mathcal{CD}}(\overline{K}_1(\phi))$ follow.

Remark 2.6. (i) The rotation map quantifies how ϕ 'affects' the Nielsen-Thomsen bases chosen.

(ii) It can be shown that rotation maps of ϕ with respect to any bases all agree on $\text{Tor}(K_1(A))$. However, we will be interested in AT-algebras which have torsion-free K_1 -groups, and hence, we do not pursue these ideas here.

Corollary 2.7. Let A, B be C^* -algebras. Let $\beta \colon \overline{K}_1(A) \longrightarrow \overline{K}_1(B)$ be a morphism. The following are equivalent

- (i) β is Nielsen-Thomsen diagonalisable.
- (ii) There exist a section $s_A \colon K_1(A) \to \overline{K}_1(A)$ and a retract $r_B \colon \overline{K}_1(B) \to H(B)$ such that $r_B \circ \beta \circ s_A = 0$ and $\beta \circ i_A \subseteq \operatorname{im}(i_B)$.

If moreover $\beta = \overline{K}_1(\phi)$ for some *-homomorphism $\phi: A \longrightarrow B$, the above are also equivalent to

- (iii) The triple $(H(\phi), \overline{K}_1(\phi), K_1(\phi))$ between the Nielsen-Thomsen sequences of A and B is split.
- (iv) There exist sections s_A, s_B of π_A, π_B respectively such that the following square commutes

$$\overline{K}_{1}(A) \xrightarrow{\pi_{A}} K_{1}(A)$$

$$\overline{K}_{1}(\phi) \downarrow \qquad \qquad \downarrow K_{1}(\phi)$$

$$\overline{K}_{1}(B) \xrightarrow{\pi_{b}} K_{1}(B)$$

Proof. These follow from standard results about split-extensions of abelian groups and are left for the reader to check. \Box

• Extended de la Harpe-Skandalis determinant. Let A be a unital C*-algebra. It is well-known that its unitary elements are in bijective correspondence with $\operatorname{Hom}_{\mathbb{C}^*,1}(C(\mathbb{T}),A)$ by sending $u \in U(A)$ to $\varphi_u \colon C(\mathbb{T}) \longrightarrow A$, where $\varphi_u(\operatorname{id}_{\mathbb{T}}) = u$. We also recall that $\operatorname{K}_1(C(\mathbb{T})) \simeq \mathbb{Z}$ is generated by $[\operatorname{id}_{\mathbb{T}}] = 1_{\mathbb{Z}}$.

We consider the canonical Nielsen-Thomsen basis of $C(\mathbb{T})$ to be the set

$$\mathcal{C}_0 := \bigg\{ 1_A, \binom{\mathrm{id}_{\mathbb{T}}}{\ddots}, \binom{\mathrm{id}_{\mathbb{T}}^*}{\ddots}, \binom{\mathrm{id}_{\mathbb{T}}^*}{\ddots} \mid k \geq 1 \bigg\}.$$

Definition 2.8. Let A be a unital C*-algebra. Fix a Nielsen-Thomsen basis $\mathcal{C} := \{c_k\}_{k \in \mathrm{K}_1(A)}$ of A. We define the extended de la Harpe-Skandalis determinant with respect to \mathcal{C} as

$$\overline{\Delta}_{\mathcal{C}} \colon U^{\infty}(A) \longrightarrow \mathrm{H}(A)$$

$$u \longmapsto \overline{\Delta} \left(\begin{smallmatrix} u & 0 \\ 0 & c^*_{[u]} \end{smallmatrix}\right)$$

Remark 2.9. (i) Observe that $R_{\mathcal{C}_0\mathcal{C}}(\varphi_u) = \overline{\Delta}_{\mathcal{C}}(u)$. In particular, we compute that

$$\operatorname{Mat}_{\mathcal{C}_0\mathcal{C}}(\overline{\mathrm{K}}_1(\varphi_u)) = \begin{pmatrix} \mathrm{H}(\varphi_u) & 1_{\mathbb{Z}} \mapsto \overline{\Delta}_{\mathcal{C}}(u) \\ 0 & [u] \end{pmatrix}.$$

(ii) Whenever $u \in U_0^{\infty}(A)$, we recover the original determinant. That is, $\overline{\Delta}_{\mathcal{C}}(u) = \overline{\Delta}(u)$.

The above allows us to exhibit an example of a *-homomorphism that is not Nielsen-Thomsen diagonalisable

Example 2.10. (Non-diagonalisable morphism.) Let $u \in C([0,1])$ be the unitary element defined by $t \mapsto \exp(2i\pi t)$ and let $\varphi_u \colon C(\mathbb{T}) \to C([0,1])$ be its associated *-homomorphism sending $\mathrm{id}_{\mathbb{T}} \mapsto u$. We know that $\mathrm{K}_1(C([0,1]))$ is trivial. Therefore, there exists a unique Nielsen-Thomsen basis \mathcal{D}_0 for C([0,1]), which is the trivial one. Moreover C([0,1]) is commutative and has stable rank one, from which we deduce that any unitary $v \in C([0,1])$ satisfies $\overline{\Delta}(v) = 0$ if and only if $v = \mathbb{1}_{[0,1]}$.

On the other hand, we know that $K_1(C(\mathbb{T})) \simeq \mathbb{Z}$. Let $\mathcal{C} := \{c_k\}_{k \in \mathbb{Z}}$ be a Nielsen-Thomsen basis for $C(\mathbb{T})$. We know that $\varphi_u(c_1) = c_1 \circ u$ and that $R_{\mathcal{CD}_0}(\varphi_u) : \mathbb{Z} \to C_{\mathbb{T}}([0,1])$ sends $1_{\mathbb{Z}} \mapsto \overline{\Delta}(\varphi_u(c_1))$. Now, it is readily checked that $c_1 \circ u = \mathbb{1}_{[0,1]}$ if and only if $c_1 = \mathbb{1}_{[0,1]}$. However, $[c_1]_{K_1} = 1$ and hence, we conclude that $K_1(\varphi_u)$ is not Nielsen-Thomsen diagonalisable.

• Comparison of *-homomorphisms at the level of the \overline{K}_1 -group. The category of abelian groups can be equipped with the discrete topology, which induces the trivial metric d_{triv} on the set of morphisms. While this metric is the finest possible (as it arises from the discrete topology), it is also somewhat 'pathological', in the sense that it only distinguishes whether two group morphisms are the same or different. In what follows, we exploit the matrix representations in Nielsen-Thomsen bases to define a more informative, and thus more useful, metric \mathfrak{d} that compares *-homomorphisms at the level of their \overline{K}_1 -groups.

Let A, B be C*-algebras. Let \mathcal{C}, \mathcal{D} be Nielsen-Thomsen bases of A, B respectively. Let $\phi, \psi \colon A \longrightarrow B$ be any *-homomorphisms. Recall that we have

$$\operatorname{Mat}_{\mathcal{CD}}(\overline{K}_{1}(\phi)) = \begin{pmatrix} H(\phi) & R_{\mathcal{CD}}(\phi) \\ 0 & K_{1}(\phi) \end{pmatrix} \qquad \operatorname{Mat}_{\mathcal{CD}}(\overline{K}_{1}(\psi)) = \begin{pmatrix} H(\psi) & R_{\mathcal{CD}}(\psi) \\ 0 & K_{1}(\psi) \end{pmatrix}$$

We aim to construct a metric \mathfrak{d} that compares $\overline{K}_1(\phi)$ and $\overline{K}_1(\psi)$ based on on entry-wise comparison. A priori, such a comparison depends on the Nielsen-Thomsen bases involved. However, we will see that any Nielsen-Thomsen bases chosen yield equivalent metrics.

• Comparison of $H(\phi)$ and $H(\psi)$. Recall that $Aff T_1(A) \simeq A_{sa}/A_0$ and we denote its elements by $\hat{h} := ev_h$, where $h \in A_{sa}$. Also, $Aff T_1(A)$ is a complete order unit space equipped with metric as follows. For $\hat{h}, \hat{l} \in Aff T_1(A)$, we have $\|\hat{h} - \hat{l}\| = \sup_{\tau \in T_1(A)} \{|\tau(h) - \tau(l)|\}$. Therefore,

$$d(\operatorname{Aff} T_1(\phi), \operatorname{Aff} T_1(\psi)) = \sup_{h \in A_{\operatorname{sa}}} \{ \|\widehat{\phi(h)} - \widehat{\psi(h)}\| \}$$

Further, we equip $\mathrm{H}(A)$ with the quotient metric as follows. For $[\widehat{h}], [\widehat{l}] \in \mathrm{H}(A)$, we have $d([\widehat{h}], [\widehat{l}]) = \inf_{\widehat{p} \in \overline{\mathrm{K}_0(A)}} \{ \|\widehat{h} - \widehat{l} - \widehat{p}\| \}$. Therefore, we have $d(\mathrm{H}(\phi), \mathrm{H}(\psi)) := \sup_{h \in A_{\mathrm{sa}}} \{ d([\widehat{\phi(h)}], [\widehat{\psi(l)}]) \}$ and immediately deduce that

$$d(H(\phi), H(\psi)) \leq d(Aff T_1(\phi), Aff T_1(\psi)).$$

• Comparison of $K_1(\phi)$ and $K_1(\psi)$. We use the trivial metric on this entry. That is,

$$d_{\mathrm{triv}}(\mathrm{K}_1(\phi),\mathrm{K}_1(\psi)) = \left\{ \begin{array}{l} 0 \text{ whenever } \mathrm{K}_1(\phi) = \mathrm{K}_1(\psi) \\ \infty \text{ else} \end{array} \right.$$

• Comparison of the rotation maps of ϕ and ψ . Recall that the rotation maps of ϕ and ψ are group morphisms from $K_1(A) \to H(B)$ depending on the Nielsen-Thomsen bases chosen.

More particularly, we have

$$R_{\mathcal{CD}}(\phi) \colon k \longmapsto \overline{\Delta} \begin{pmatrix} \phi(c_k) & 0 \\ 0 & d^*_{[\mathrm{K}_1(\phi)(k)]} \end{pmatrix} \quad \text{and} \quad R_{\mathcal{CD}}(\psi) \colon k \longmapsto \overline{\Delta} \begin{pmatrix} \psi(c_k) & 0 \\ 0 & d^*_{[\mathrm{K}_1(\phi)(k)]} \end{pmatrix}$$

where $\mathcal{C} = \{c_k\}_{k \in \mathrm{K}_1(A)}, \mathcal{D} = \{d_k\}_{k \in \mathrm{K}_1(B)}$ are fixed Nielsen-Thomsen bases.

We know that the codomain of these maps is equipped with a metric. As a result, we are able to compare them in the following way. To ease notations, let us first introduce the *relative rotation* map of (ϕ, ψ) in the bases \mathcal{CD} that we denote

$$R_{\mathcal{CD}}(\phi, \psi) \colon \mathrm{K}_1(A) \longrightarrow \mathrm{H}(B)$$

 $k \longmapsto R_{\mathcal{CD}}(\phi)(k) - R_{\mathcal{CD}}(\psi)(k)$

and define the following associated metric

$$d_{R,\mathcal{CD}}(\phi,\psi) := \sup_{k \in K_1(A)} \{ \| R_{\mathcal{CD}}(\phi,\psi)(k) \| \}$$

It need not be true that the metrics $d_{R,\mathcal{C}\mathcal{D}}$ and $d_{R,\mathcal{C}'\mathcal{D}'}$ are equivalent for different pairs of bases. Nevertheless, the following lemma shows how they interplay with one another and will be crucial to prove that the comparisons obtained at the level of the \overline{K}_1 -group are indeed equivalent.

Lemma 2.11. Let $\phi, \psi: A \longrightarrow B$ be *-homomorphisms. Let $\mathcal{C} = \{c_k\}_{k \in \mathrm{K}_1(A)}, \mathcal{D}$ be Nielsen-Thomsen bases of A, B.

(i) Assume that $K_1(\phi) = K_1(\psi)$. For any $k \in K_1(A)$, we compute

$$R_{\mathcal{CD}}(\phi, \psi)(k) = \overline{\Delta} \begin{pmatrix} \phi(c_k) & 0 \\ 0 & \psi^*(c_k) \end{pmatrix}.$$

In particular, the relative rotation map of (ϕ, ψ) only depends on the basis chosen for A.

(ii) Let $C' = \{c'_k\}_{k \in K_1(A)}$ and D' be other Nielsen-Thomsen bases of A and B. For any $k \in K_1(A)$, we compute

$$R_{\mathcal{CD}}(\phi, \psi)(k) - R_{\mathcal{C}'\mathcal{D}'}(\phi, \psi)(k) = \widehat{\phi(h_k)} - \widehat{\psi(h_k)}$$

where $\widehat{h_k} := \overline{\Delta} \begin{pmatrix} c_k & 0 \\ 0 & (c'_k)^* \end{pmatrix}$. In particular, the map $R_{\mathcal{CD}}(\phi, \psi)(k) - R_{\mathcal{C'D'}}(\phi, \psi)$ only depends on the bases chosen for A.

Proof. (i) Follows from the fact that $d_{[K_1(\phi)(k)]} = d_{[K_1(\psi)(k)]}$ and that $\overline{\Delta}(u) + \overline{\Delta}(u^*) = 0$.

(ii) Using that $\overline{\Delta}(u) + \overline{\Delta}(v) = \overline{\Delta}(\begin{smallmatrix} u & 0 \\ 0 & v \end{smallmatrix})$, and again that that $\overline{\Delta}(u) + \overline{\Delta}(u^*) = 0$, we obtain that for any $k \in \mathrm{K}_1(A)$

$$R_{\mathcal{CD}}(\phi,\psi)(k) - R_{\mathcal{C'D'}}(\phi,\psi)(k) = \overline{\Delta} \begin{pmatrix} \phi(c_k) & 0 \\ 0 & \phi(c_k')^* \end{pmatrix} - \overline{\Delta} \begin{pmatrix} \psi(c_k) & 0 \\ 0 & \psi(c_k')^* \end{pmatrix}.$$

and the result follows.

Gathering all the above, we are now ready to construct a (non-trivial) metric to compare the *-homomorphisms at the level of the Hausdorffized algebraic unitary group. Most importantly, we will see that the topology induced by this metric does not depend on the Nielsen-Thomsen bases.

Definition 2.12. Let $\phi, \psi \colon A \longrightarrow B$ be *-homomorphisms between C*-algebras A, B. Let \mathcal{C}, \mathcal{D} be Nielsen-Thomsen bases of A, B. We define the following metric

$$\mathfrak{d}_{\mathcal{CD}}(\overline{\mathrm{K}}_{1}(\phi),\overline{\mathrm{K}}_{1}(\psi)):=d(\mathrm{H}(\phi),\mathrm{H}(\psi))+d_{R,\mathcal{CD}}(\phi,\psi)+d_{\mathrm{triv}}(\mathrm{K}_{1}(\phi),\mathrm{K}_{1}(\psi)).$$

Theorem 2.13. Let C, C' be Nielsen-Thomsen bases of A and D, D' be Nielsen-Thomsen bases of B. Then, for any *-homomorphisms $\phi, \psi \colon A \longrightarrow B$, we have

$$\mathfrak{d}_{\mathcal{CD}}(\overline{K}_1(\phi),\overline{K}_1(\psi)) \leq 2\mathfrak{d}_{\mathcal{C}'\mathcal{D}'}(\overline{K}_1(\phi),\overline{K}_1(\psi))$$

In particular, $\mathfrak{d}_{\mathcal{CD}}$ and $\mathfrak{d}_{\mathcal{C}'\mathcal{D}'}$ are equivalent metrics.

Proof. First, we observe that $d(\mathbf{H}(\phi), \mathbf{H}(\psi)) \leq d_{R,\mathcal{C}'\mathcal{D}'}(\phi, \psi)$. Also, we know that $\|R_{\mathcal{C}\mathcal{D}}(\phi, \psi)(k)\| \leq \|R_{\mathcal{C}'\mathcal{D}'}(\phi, \psi)(k)\| + \|\widehat{\phi(h_k)} - \widehat{\psi(h_k)}\|$ for any $k \in K_1(A)$. (This is essentially (ii) of Lemma 2.11.) This implies that $d_{R,\mathcal{C}\mathcal{D}}(\phi, \psi) \leq d_{R,\mathcal{C}'\mathcal{D}'}(\phi, \psi) + d(\mathbf{H}(\phi), \mathbf{H}(\psi))$. As a result, we compute that

$$\begin{split} \mathfrak{d}_{\mathcal{CD}}(\overline{K}_1(\phi),\overline{K}_1(\psi)) &\leq \mathfrak{d}_{\mathcal{C}'\mathcal{D}'}(\overline{K}_1(\phi),\overline{K}_1(\psi)) + d(H(\phi),H(\psi)) \\ &\leq 2\mathfrak{d}_{\mathcal{C}'\mathcal{D}'}(\overline{K}_1(\phi),\overline{K}_1(\psi)). \end{split}$$

Even though the Nielsen-Thomsen bases involved will often be unambiguous, we may abusively denote any such (equivalent) metrics by \mathfrak{d} . Finally, it is worth mentioning that $\mathfrak{d}(\overline{K}_1(\phi), \overline{K}_1(\psi)) \leq d_{\mathrm{triv}}((\overline{K}_1(\phi), \overline{K}_1(\psi)))$. In the last part of this manuscript, we expose cases where $0 < \mathfrak{d}(\overline{K}_1(\phi), \overline{K}_1(\psi)) < \infty$ showcasing the usefulness of \mathfrak{d} .

3. Refined Cuntz semigroups and comparison of morphisms

A common strategy for classifying large classes of C*-algebras involves first classifying adequate classes of *-homomorphisms via uniqueness and existence results. In practice, this requires methods to approximately distinguish concrete morphisms and approximately lift abstract morphisms within the target category. To this end, the choice of invariants with suitable codomains equipped with suitable metrics to compare morphisms play a pivotal role in facilitating classification. This section exhibits such tools for refined Cuntz semigroups, such as the unitary Cuntz semigroup and its Hausdorffized version, and explores how these refinements enable more informative comparisons of morphisms.

A. The Cuntz semigroup. The Cuntz semigroup was introduced in [12] and has been a powerful tool for classification since the milestone paper [11]. We refer the reader [18] for a detailed survey around the Cuntz semigroup, where they shall find all the basics they might need, and to [1] for state of the art results.

Let (S, \leq) be an ordered monoid and let x, y in S. We say that x is way-below y and we write $x \ll y$ if, for all increasing sequences $(z_n)_{n \in \mathbb{N}}$ in S that have a supremum, if $\sup_{n \in \mathbb{N}} z_n \geq y$, then there exists k such that $z_k \geq x$.

We say that S is an abstract Cuntz semigroup, or a Cu-semigroup, if S satisfies the above axioms.

- (O1) Every increasing sequence of elements in S has a supremum.
- (O2) For any $x \in S$, there exists a \ll -increasing sequence $(x_n)_{n \in \mathbb{N}}$ in S such that $\sup_n x_n = x$.
- (O3) Addition and the compact containment relation are compatible.
- (O4) Addition and suprema of increasing sequences are compatible.

We say that a map $\alpha: S \longrightarrow T$ is a Cu-morphism if α is an ordered monoid morphism preserving the compact-containment relation and suprema of increasing sequences.

Let A be a C*-algebra and let $a, b \in A_+$. We say that a Cuntz subequivalent to b in A, denoted $a \leq_{\text{Cu}} b$, if for every $\epsilon > 0$ there exists $r \in A$ such that $||rbr^* - a|| < \epsilon$. We antisymmetrize this subequivalence to obtain a an equivalence relation \sim_{Cu} called the Cuntz equivalence relation. The Cuntz semigroup of A is

$$Cu(A) = (A \otimes \mathcal{K})_+ / \sim_{Cu}$$

We equip Cu(A) with the addition induced by diagonal addition and the order induced by \leq , which makes it a Cu-semigroup.

It has been shown in [11] and [2] that the functor $Cu: C^* \longrightarrow Cu$ is well-defined and continuous.

• Comparison of Cu-morphisms. We recall that several metrics on the set $\operatorname{Hom}_{\operatorname{Cu}}(S,T)$ of morphisms between abstract Cu-semigroups S,T have been studied. Also, that a 'comparison on finite sets' has played a significant role in classification of *-homomorphisms by means of the Cuntz semigroup. (We refer the reader to [8, Section 5] for a general overview.)

The first meaningful metric has appeared in the classification of *-homomorphisms with domain $C_0((0,1])$. (See [10] and [24].) Recall that $\operatorname{Cu}(C(]0,1])) \simeq \operatorname{Lsc}((0,1],\overline{\mathbb{N}})$. More generally, it is known that $\operatorname{Lsc}(X,\overline{\mathbb{N}})$ is a Cu-semigroup for any compact Hausdorff space X that may differ from $\operatorname{Cu}(C(X))$. Let X be a (locally) compact Hausdorff space and let $T \in \operatorname{Cu}$. For any Cu-morphisms $\alpha,\beta \colon \operatorname{Lsc}(X,\overline{\mathbb{N}}) \longrightarrow T$, we define

$$d_{\mathrm{Cu}}(\alpha,\beta) = \inf \Big\{ r > 0 \mid \alpha(\mathbb{1}_U) \le \beta(\mathbb{1}_{U_r}) \text{ and } \beta(\mathbb{1}_U) \le \alpha(\mathbb{1}_{U_r}) \text{ for any } U \in \mathcal{O}(X) \Big\}$$

where $\mathcal{O}(X)$ denotes the open sets of X and $U_r := \bigcup_{x \in U} B_r(x)$. This is a metric on the set $\mathrm{Hom}_{\mathrm{Cu}}(\mathrm{Lsc}(X,\overline{\mathbb{N}}),T)$ referred to as the Cu-metric.

Later on, other types of comparison have been studied in [8]. We present a couple of abstract approaches that have been developed. In what follows, $\alpha, \beta \colon S \longrightarrow T$ are Cu-morphisms.

[Finite-set comparison] Let $F \subseteq S$ be a (finite) set. We say that α and β compare on F and we write $\alpha \simeq_F \beta$ if, for any $g, h \in F$ such that $g \ll h$ (in S), we have that $\alpha(g) \leq \beta(h)$ and $\beta(g) \leq \alpha(h)$.

Observe that $\alpha = \beta$ if and only if $\alpha \simeq_F \beta$ for any finite set $F \subseteq S$ if and only if $\alpha \simeq_{\{g,h|g \ll h\}} \beta$.

[Thomsen comparison] We define Thomsen semigroup of a Cu-semigroup S defined by Th(S) := $\text{Hom}_{\text{Cu}}(\mathbb{G}, S)$, where $\mathbb{G} := \{ f \in \text{Lsc}([0, 1], \overline{\mathbb{N}}) \mid f(0) = 0 \text{ and } f \text{ is increasing} \}$ is the generator of the category Cu.

Let S, T be Cu-semigroups and let $N \subseteq Th(S)$. For any Cu-morphisms $\alpha, \beta \colon S \longrightarrow T$, we define

$$d_{\mathcal{N}}(\alpha,\beta) := \sup_{\nu \in \mathcal{N}} \{ d_{\mathcal{C}u}(\alpha \circ \nu, \beta \circ \nu) \}.$$

This is a metric whenever N has a generating image, that is, whenever the monoid generated by $\{\nu(\mathbb{1}_{(t,1]}) \mid \nu \in \mathbb{N}, t \in (0,1]\}$ is sup-dense in S. Roughly speaking, we require N to be large enough. In such a case, we refer to the above as the Cu-metric induced by N

Remark 3.1. (i) The finite-set comparison and the Thomsen comparison are intrinsically related to one another. See [8, Lemma 5.15].

- (ii) The Cu-metric on $\operatorname{Hom}_{\operatorname{Cu}}(\operatorname{Lsc}(X,\overline{\mathbb{N}}))$ is in particular a Cu-metric induced by an adequate N_X . See [8, Examples 5.13 5.14]. (In fact, any Cu-metric considered in the past can be realized from an adequate Thomsen comparison.)
- (iii) If the set N is too large, e.g. N is Th(S), then the induced Cu-metric may be the trivial one. See [8, Example 5.10].
- B. Refined Cuntz semigroups. In [7], the author proposed a systematic construction for incorporating additional invariants into the Cuntz semigroup, resulting in refined versions that capture more structural information about C*-algebras and their ideals. By leveraging categorical properties of the Cuntz semigroup and its target category Cu, these refined invariants provide a robust context for classification. Here, we present several methods to compare abstract and concrete morphisms, crucial for distinguishing and lifting morphisms. Let us first recall the process and we refer the reader to [7, Section 4.A] for more details.
- The Cu_K -construction. Let A be C^* -algebra and let $K: C^* \to AbGp$ be a continuous functor. We view Cu(A) as a partially ordered set which induces a small category. (The objects are its elements and morphisms are induced by the order.) Therefore, we can consider the functor

$$K_A : Cu(A) \longrightarrow AbGp$$

 $x \longmapsto K(I_x)$
 $x \le y \longmapsto K(I_x \xrightarrow{\subseteq} I_y)$

where $I_x \in \operatorname{Lat}_f(A)$ is the corresponding closed two-sided ideal of A through the isomorphism $\operatorname{Lat}_f(\operatorname{Cu}(A)) \simeq \operatorname{Lat}_f(A)$.

(i) The Cu_K -construction of a C^* -algebra A is an ordered monoid $Cu_K(A)$ consisting of pairs (x, g), where $x \in Cu(A)$ and $g \in K(I_x)$.

The addition is given by

$$(x,g) + (y,h) := (x+y, K_A(x \le x+y)(g) + K_A(y \le x+y)(h)).$$

The order is given by

$$(x,g) \leq (y,h)$$
 whenever $x \leq y$ in $Cu(A)$ and $K_A(x \leq y)(g) = h$ in $K_A(I_y)$.

(ii) The Cu_K -construction of a *-homomorphism $\phi: A \to B$ is an ordered monoid morphism $Cu_K(\phi)$ sending

$$\operatorname{Cu}_{\mathrm{K}}(\phi) \colon (x,g) \mapsto (\operatorname{Cu}(\phi)(x), \operatorname{K}(I_x \xrightarrow{\phi} (I_x)_{\phi})(g))$$

where $I_{\phi} \in \text{Lat}(B)$ denotes the smallest ideal of B containing $\phi(I)$. Therefore, $\text{Cu}_{\mathbf{K}}(\phi)$ is entirely determined by the data $(\text{Cu}(\phi), \{\overline{\mathbf{K}}_1(I \xrightarrow{\phi} I_{\phi})\}_{I \in \text{Lat}_f(A)})$.

We briefly recall some definitions and properties that will be of use later on about Cu_K -constructions. We refer the reader to [7] for more details and results on the matter.

Theorem 3.2. (i) The assignment $Cu_K: C^* \longrightarrow Cu^*$ is a well-defined continuous functor.

(The category Cu* resembles the category Cu, but the ordered monoids involved need not be positively ordered.)

(ii) Let $\phi: A \to B$ be a *-homomorphism between C^* -algebras and let $I \in \operatorname{Lat}_f(A)$. Then for any $J \in \operatorname{Lat}(B)$ such that $J \supseteq I_{\phi}$, the following diagram is commutative with exact rows in Cu^*

$$D(I, \phi, J) \colon 0 \longrightarrow \operatorname{Cu}(I) \xrightarrow{i_1} \operatorname{Cu}_{\mathsf{K}}(I) \xrightarrow{j_1} \mathsf{K}(I) \longrightarrow 0$$

$$\operatorname{Cu}(\phi) \downarrow \qquad \operatorname{Cu}_{\mathsf{K}}(\phi) \downarrow \qquad q_1 \qquad \bigvee_{\mathsf{K}(I \xrightarrow{\phi} I_{\phi} \subseteq J)} \mathsf{K}(J) \xrightarrow{i_2} \operatorname{Cu}_{\mathsf{K}}(J) \xrightarrow{j_2} \mathsf{K}(J) \longrightarrow 0$$

where I_{ϕ} is the smallest element of $\operatorname{Lat}(B)$ obtained by I through ϕ , i.e. $I_{\phi} := \overline{B\phi(I)B}$.

- (ii') The canonical diagram $D(I, \phi, J) \stackrel{\iota}{\longrightarrow} D(A, \phi, B)$ is also commutative with exact rows.
- **3.3.** (The category Cu_w .) The general procedure of merging a Cu-semigroup S with a functor $G: S \to AbGp$ has been introduced in [7] and referred to as a 'webbing transformation'. Roughly speaking, given S, T and $\alpha_0: S \to T$ in the category Cu, together with functors G, H from S, T respectively into the category AbGp and a natural transformation from G to H, we obtain objects and morphisms in the category Cu^* in the following form

$$S_G, T_H$$
 and $\alpha = (\alpha_0, \{\alpha_I\}_{I \in \operatorname{Lat}_f(S)}) \colon S_G \to T_H.$

The webbing transformations form a subcategory $Cu_w \subseteq Cu^*$ closed under inductive limits, which need not be full. Nevertheless, having classification in mind, it may be more adequate to consider the restriction $Cu_K \colon C^* \longrightarrow Cu_w$ which is again a continuous functor, since any Cu_K -construction (of both C^* -algebras and *-homomorphisms) belongs to Cu_w .

• Comparison of Cu_K -morphisms. We introduce several ways to compare $Cu_{\mathfrak{w}}$ -morphisms between Cu_K -constructions, that we naturally call Cu_K -morphisms. These comparisons are based on a Cu-comparison in the first component and a group comparison in the second component.

We need not look to more general webbing transformations for our work. Therefore, we will only look at Cu_K -morphisms for the clarity of the exposition, where K stands for either K_1 or \overline{K}_1 . However, we mention that the following remains true for the entire category $Cu_{\mathfrak{w}}$.

In what follows and up until the next section, we let A, B be C*-algebras, K: C* \to AbGp be a continuous functor and $\alpha := (\alpha_0, \{\alpha_I\}), \beta := (\beta_0, \{\beta_I\})$: Cu_K $(A) \longrightarrow$ Cu_K(B) be Cu_K-morphisms.

[Finite-set comparison] Let $F \subseteq \text{Cu}_K(A)$ be a (finite) set. We say that α and β compare on F and we write $\alpha \simeq_F \beta$ if, for any $(x,g),(y,h) \in F$ such that $(x,g) \ll (y,h)$ we have that $\alpha(x,g) \leq \beta(y,h)$ and $\beta(x,g) \leq \alpha(y,h)$.

Proposition 3.4. Retain the above setting and let $F_0 := \{x \in S \mid (x,g) \in F \text{ for some } g \in K(x)\}$. Then $\alpha \simeq_F \beta$ if and only if the following conditions are satisfied

- (i) $\alpha_0 \simeq_{F_0} \beta_0$
- (ii) For any $x, y \in F_0$ such that $x \ll y$, and for any $g \in K(x)$ such that $(x, g) \in F$, we have that

$$\left\{ \begin{array}{l} \mathrm{K}(\alpha_0(x) \leq \alpha_0(y)) \circ \alpha_{I_x}(g) = \mathrm{K}(\beta_0(x) \leq \alpha_0(y)) \circ \beta_{I_x}(g) \ \ in \ \mathrm{K}(\alpha_0(y)) \\ \mathrm{K}(\beta_0(x) \leq \beta_0(y)) \circ \beta_{I_x}(g) = \mathrm{K}(\alpha_0(x) \leq \beta_0(y)) \circ \alpha_{I_x}(g) \ \ in \ \mathrm{K}(\beta_0(y)) \end{array} \right.$$

Proof. This follows from the characterization obtained [7, Proposition 2.12], stating that $(x, g) \ll (y, h)$ if and only if $x \ll y \in S$ and $K(x \leq y)(g) = h$.

[Metric comparison] Let us now use Thomsen comparison (and a fortiori, the Cu-metric in specific cases) available on the first components α_0, β_0 of α, β to get a 'metric-type' comparison. The main obstacle is that for a given ideal $I \in \text{Lat}_f(S)$, the morphisms α_I, β_I need not have the same codomain. Nevertheless, we can always compare them in a larger group. How larger the group need be will depend on the Cu-part of the morphisms.

Definition 3.5. (The fiber diagrams.) Let $x, y \in S$ be such that $x \ll y$. We define the fiber diagram of α and β at coordinates (x, y) as follows

$$\mathrm{K}(\beta_0(y)) \underbrace{\overset{\mathrm{K}(\leq)}{\underset{\mathrm{K}_{\delta}(\leq)}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}_{\delta}(\leq)}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y))}{\underset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y)})}{\overset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y)})}{\overset{\mathrm{K}(\beta_0(y))}{\overset{\mathrm{K}(\beta_0(y)})}{\overset{\mathrm{K}(\beta_0(y)})}{\overset{\mathrm{K}(\beta_0(y)})}$$

where the maps $K_{\delta}(t_1 \leq t_2)$ are defined to be $K(t_1 \leq t_2)$ whenever $t_1 \leq t_2$ and the empty morphism (i.e., no arrows) otherwise.

We denote the above diagram by $\mathcal{F}_{\{\alpha,\beta\}}(x,y)$, or simply $\mathcal{F}(x,y)$ when the context is clear.

Remark 3.6. The compositions $K(\leq) \circ \alpha_{I_x} \colon K(x) \to K(\alpha_0(y))$ and $K(\leq) \circ \beta_{I_x} \colon K(x) \to K(\beta_0(y))$ factorize through $K(I_y)$. Hence the fiber diagram $\mathcal{F}(x,y)$ is equivalent to

$$\begin{split} & \mathrm{K}(\beta_0(x)) \xleftarrow{\beta_{I_x}} & \mathrm{K}(x) \xrightarrow{\alpha_{I_x}} & \mathrm{K}(\alpha_0(x)) \\ & \mathrm{K}_{\delta}(\leq) \bigvee & \mathrm{K}(\leq) \bigvee & \bigvee & \mathrm{K}_{\delta}(\leq) \\ & \mathrm{K}(\alpha_0(y)) \xleftarrow{\alpha_{I_y}} & \mathrm{K}(y) \xrightarrow{\beta_{I_x}} & \mathrm{K}(\beta_0(y)) \end{split}$$

We highlight that all the group morphisms involved in the fiber diagrams $\mathcal{F}(x,y)$ of α and β (that is, group morphisms which belong to either $\{\alpha_I\}, \{\beta_I\}$ or $\{K(s' \leq s)\}, \{K(t' \leq t)\}$) can always be equipped with a metric d. (E.g., take d to be the trivial metric.) Naturally, we define the norm of a diagram $\mathcal{F}(x,y)$ associated to d by

$$\|\mathcal{F}(x,y)\|_d := \max_{(c,c')\in\Gamma} \{d(c,c')\}$$

where Γ denotes all the pairs (c, c') of paths in $\mathcal{F}(x, y)$ with the same endpoints.

We are now ready to define several metrics to compare α and β as follows.

Definition 3.7. Let d be a metric on the set of fiber diagrams of α and β and let $N \subseteq Th(Cu(A))$ be a set with a generating image. We define

$$d_{\mathrm{N},d}^*(\alpha,\beta) := \sup_{\nu \in \mathrm{N}} \inf \left\{ r > \epsilon_{\nu} \mid \| \mathcal{F}(\nu(\mathbb{1}_{(t+r,1]}),\nu(\mathbb{1}_{(t,1]}) \|_d \leq 4r \text{ for any } t \in [0,1] \right\}$$

where $\epsilon_{\nu} := d_{\text{Cu}}(\alpha_0 \circ \nu, \beta_0 \circ \nu).$

In the particular case where $Cu(A) \simeq Lsc(X, \overline{\mathbb{N}})$, we define

$$d_{\mathrm{Cu},d}^*(\alpha,\beta) := \inf \Big\{ r > \epsilon_0 \mid \|\mathcal{F}(\mathbb{1}_U,\mathbb{1}_{U_r})\|_d \le 4r \text{ for any } U \in \mathcal{O}(X) \Big\}$$

where $\epsilon_0 := d_{Cu}(\alpha_0, \beta_0)$.

These are well-defined metrics on the set $\mathrm{Hom}_{\mathrm{Cu}_{\mathtt{w}}}(\mathrm{Cu}_{\mathrm{K}}(A),\mathrm{Cu}_{\mathrm{K}}(B)).$

It is immediate that these metrics are finer than their counter-part in the category Cu. Concretely, $d_{\text{Cu}}(\alpha_0, \beta_0) \leq d_{\text{Cu},d}^*(\alpha, \beta)$. (Similarly for metrics induced by any $N \subseteq \text{Th}(\text{Cu}(A))$.) Also, the 'factor 4' involved on the fiber diagrams comparison is mostly for practical reasons whenever using the Hausdorffized unitary Cuntz semigroup with the metric \mathfrak{d} on the fiber diagrams. See Question 3.20 and explicit computations in Section 4.

Finally, we mention that one could imagine a metric where the measures of the 'Cu-components' and 'the fiber diagrams components' are uncorrelated but we do not pursue this idea.

Proposition 3.8. Assume that $Cu(A) \simeq Lsc(X, \overline{\mathbb{N}})$ for a compact Hausdorff space X.

(i) Let d be a metric on the set of fiber diagram. For any $x \in Cu(A)$ and any $z \in Cu(B)$ with $\alpha_0(x), \beta_0(x) \leq z$, we have

$$d(K(\alpha_0(x) \leq z) \circ \alpha_I, K(\beta_0(x) \leq z) \circ \beta_I) \leq 4d^*_{Cu,d}(\alpha,\beta).$$

(ii) Let $d_{\rm triv}$ be the trivial metric on the set of fiber diagram. We compute

$$d_{\mathrm{Cu}}^*(\alpha,\beta) = \inf \Big\{ r > \epsilon_0 \mid \mathcal{F}(\mathbb{1}_U,\mathbb{1}_{U_r}) \text{ commutes for any } U \in \mathcal{O}(X) \Big\}$$

where $\epsilon_0 := d_{Cu}(\alpha_0, \beta_0)$. We may write d_{Cu}^* instead of $d_{Cu, d_{triv}}^*$.

Analogous statements hold for any C^* -algebra A and the metrics induced by some $N \subseteq Cu(A)$.

Proof. Both computations follow from construction and are left for the reader to check.

3.9. (The unitary Cuntz semigroup.) Let us introduce a first refined version of the Cuntz semigroup obtained via the Cu_K -construction applied to the unitary group $K_1 \colon C^* \to AbGp$.

Definition 3.10. [3] - [7, 4.B]. Let A be a C*-algebra. We define The unitary Cuntz semigroup of A, denoted by $Cu_{K_1}(A)$, to be the merging of the Cuntz semigroup with the K_1 -group.

We remark that the unitary Cuntz semigroup had been studied intensively by the author, and therebefore termed Cu₁ instead of Cu_{K1}. Further, we now have a metric d_N^* (or d_{Cu}^* in particular cases) on the set $\operatorname{Hom}_{Cu_m}(\operatorname{Cu}_{K_1}(A), \operatorname{Cu}_{K_1}(B))$.

C. The Hausdorffized unitary Cuntz semigroup and its associated metric(s). In what follows, we introduce and study a second refined version of the Cuntz semigroup obtained via the Cu_K-construction applied to the Hausdorffized algebraic unitary group $\overline{K}_1 \colon C^* \to AbGp$

Definition 3.11. [7, 4.11] Let A be a C*-algebra. We define the *Hausdorffized unitary Cuntz semi-group* of A, denoted by $Cu_{\overline{K}_1}(A)$, to be the merging of the Cuntz semigroup with the Hausdorffized algebraic K_1 -group.

Let A, B be C*-algebras. Consider a set $N \subseteq Th(Cu(A))$ with a generating image. Similarly as for the unitary Cuntz semigroup, we now have a metric d_N^* (or d_{Cu}^* in particular cases) on the set $Hom_{Cu_{\overline{W}}}(Cu_{\overline{K}_1}(A), Cu_{\overline{K}_1}(B))$.

Nevertheless, we have built a more informative metric \mathfrak{d} to compare *-homomorphisms $\phi, \psi \colon A \longrightarrow B$ on their \overline{K}_1 -group. As a consequence, whenever we wish to compare concrete morphisms $\operatorname{Cu}_{\overline{K}_1}(\psi)$ and $\operatorname{Cu}_{\overline{K}_1}(\phi)$, we can equip the set of fiber diagrams of \overline{K}_1 -groups with the metric \mathfrak{d} instead to obtain a more informative comparison that we may abusively write \mathfrak{d}_N^* . More concretely, we have

Proposition 3.12. Let X be a compact Hausdorff space and let $\phi, \psi \colon C(X) \longrightarrow B$ be *-homomorphisms between C*-algebras. We compute

$$\mathfrak{d}_{\mathrm{Cu}}^*(\alpha,\beta) = \inf \Big\{ r > \epsilon_0 \mid \|\mathcal{F}(\mathbb{1}_U,\mathbb{1}_{U_r})\|_{\mathfrak{d}} \le 4r \text{ for any } U \in \mathcal{O}(X) \Big\}$$

where $\epsilon_0 := d_{Cu}(\alpha_0, \beta_0)$ and the fiber diagrams involve \overline{K}_1 -groups.

Analogous result holds for any C^* -algebra A and any metric \mathfrak{d}_N^* induced by some $N \subseteq Cu(A)$.

Proof. This is immediate from construction.

We end this part by summarizing and relating all the comparisons of *-homomorphisms at the level of the Cuntz semigroup and its refined versions that we have considered so far.

Proposition 3.13. Let X be a compact Hausdorff space and let $\phi, \psi \colon C(X) \longrightarrow B$ be *-homomorphisms between C*-algebras. We have

$$\begin{split} d_{\mathrm{Cu}}(\mathrm{Cu}(\psi),\mathrm{Cu}(\phi)) &\leq d_{\mathrm{Cu}}^*(\mathrm{Cu}_{\mathrm{K}_1}(\psi),\mathrm{Cu}_{\mathrm{K}_1}(\phi)) \\ &\leq \mathfrak{d}_{\mathrm{Cu}}^*(\mathrm{Cu}_{\overline{\mathrm{K}}_1}(\psi),\mathrm{Cu}_{\overline{\mathrm{K}}_1}(\phi)) \leq d_{\mathrm{Cu}}^*(\mathrm{Cu}_{\overline{\mathrm{K}}_1}(\psi),\mathrm{Cu}_{\overline{\mathrm{K}}_1}(\phi)) \leq d_U(\psi,\phi) \end{split}$$

where d_U denotes the unitary distance between ϕ and ψ .

Analogous inequalities hold for any C*-algebra A and the metrics induced by some $N \subseteq Cu(A)$.

Proof. The result follows from the observations that $d_{\text{triv}}(K_1(\phi), K_1(\psi)) \leq \mathfrak{d}(\overline{K}_1(\phi), \overline{K}_1(\psi)) \leq d_{\text{triv}}(\overline{K}_1(\phi), \overline{K}_1(\psi))$ and that $d_{\text{triv}}(K_1(\phi), K_1(\psi)) = 0$ if and only if $\mathfrak{d}(\overline{K}_1(\phi), \overline{K}_1(\psi)) < \infty$.

D. Connections between Cu-metrics and comparison on Traces. We have introduced several ways to compare *-homomorphisms $\phi, \psi \colon A \longrightarrow B$ on distinct invariants. The comparison \mathfrak{d}_{N}^{*} (or its analogue \mathfrak{d}_{Cu}^{*}) at the level of the Hausdorffized unitary Cuntz semigroup is, in particular, comparing ϕ and ψ at the level of their Cuntz semigroup and their traces. (Or more particularly, on the functor H.)

Now, it is shown in [17] that there is a functorial equivalence $F(Cu(.)) \simeq T(.)$ between the cone of (not necessarily bounded) traces and the cone of functionals. (Recall that a functional of a Cusemigroup S is monoid morphism $\lambda \colon S \to \overline{\mathbb{R}}_+$ preserving suprema of increasing sequences.) Roughly speaking, this amounts to the fact that the cone of functional of the Cuntz semigroup of a C*-algebra recovers the cone of its tracial states. It would then seem enough to compare the morphisms at the level of the Cuntz semigroup to get a comparison at the level of tracial states. As a first step towards this conjecture, let us prove the following.

Theorem 3.14. Let $\phi, \psi \colon A \longrightarrow B$ be *-homomorphisms between C*-algebras. Assume that ϕ and ψ agree at the level of the Cuntz semigroup.

Then Aff $T_1(\phi) = \text{Aff } T_1(\psi)$. A fortiori, we have $H(\phi) = H(\psi)$.

Proof. Let $\tau \in T_1(A)$ and let $\lambda_{\tau} \in F(Cu(A))$ be its corresponding functional. Let $a \in A_+$. Using [17] and the fact that *-homomorphisms preserve continuous functional calculus, we obtain that

$$\tau(\phi(a)) = \int_0^\infty \lambda_\tau([(\phi(a) - t)_+])dt = \int_0^\infty \lambda_\tau([\phi((a - t)_+)])dt$$
$$= \int_0^\infty \lambda_\tau \circ \operatorname{Cu}(\phi)([(a - t)_+])dt = \int_0^\infty \lambda_\tau \circ \operatorname{Cu}(\psi)([(a - t)_+])dt$$
$$= \tau(\psi(a)).$$

П

In view of classifying *-homomorphisms, we wonder whether we can obtain an approximate version of the latter result. For instance, in the particular case where A = C([0,1]) and B is a C*-algebra of stable rank one, the classification theorem obtained in [23] immediately tells us that $d(\operatorname{Aff} T_1(\phi), \operatorname{Aff} T_1(\psi)) \leq d_{\operatorname{Cu}}(\operatorname{Cu}(\phi), \operatorname{Cu}(\psi))$. More generally, it is natural to ask the following.

Question 3.15. (i) Does there always exist an adequate metric d_N on the set of Cu-morphims? In such case, do we always have $d(\text{Aff } T_1(\phi), \text{Aff } T_1(\psi)) \leq d_N(\text{Cu}(\phi), \text{Cu}(\psi))$?

- (ii) Is the Cu-comparison on finite sets finer than the Aff T_1 -comparison on finite set?
- **3.16.** Let us investigate more closely and partially solve the above questions. We first settle some preliminary observations and notations that will be needed for this task.
- (i) It is well-known that the positive elements of a C*-algebra A are in bijective correspondence with $\operatorname{Hom}_{C^*}(C((0,1]),A)$ by sending $a \in A_+$ to $\varphi_a \colon C_0((0,1]) \longrightarrow A$, where $\varphi_a(\operatorname{id}_{(0,1]}) = a$.

Therefore, we have the following mappings

$$A_+ \longrightarrow \operatorname{Hom}_{\mathrm{C}^*}(C((0,1]), A) / \sim_{\operatorname{aue}} \longrightarrow \operatorname{Th}(\operatorname{Cu}(A))$$

 $a \longmapsto [\varphi_a] \longmapsto \nu_a := \operatorname{Cu}(\varphi_a)_{|\mathbb{G}}$

(ii) Under the stable rank one hypothesis, we know that $\operatorname{Th}(\operatorname{Cu}(A))$ is isomorphic (as a monoid) to $\operatorname{Hom}_{\mathrm{C}^*}(C_0((0,1]),A)/\sim_{\operatorname{aue}}$. Moreover, for any $\varphi_1,\varphi_2\in\operatorname{Hom}_{\mathrm{C}^*}(C_0((0,1]),A)/\sim_{\operatorname{aue}}$ satisfying $\varphi_1\sim_{\operatorname{aue}}\varphi_2$, it is readily checked that $\widehat{a_1}=\widehat{a_2}$ in $\operatorname{Aff} T_1(A)$, where $a_1:=\varphi_1(\operatorname{id}_{(0,1]}), a_2:=\varphi_1(\operatorname{id}_{(0,1]}).$ Therefore, we have the following mappings

$$\begin{array}{ccc} \operatorname{Th}(\operatorname{Cu}(A)) & \xrightarrow{\simeq} \operatorname{Hom}_{\operatorname{C}^*}(C((0,1]),A)/\sim_{\operatorname{aue}} & \operatorname{Aff} T_1(A) \\ \nu & \longmapsto & [\varphi_{\nu}] & \longmapsto \widehat{a_{\nu}} := \widehat{\varphi_{\nu}(\operatorname{id})} \end{array}$$

(iii) For any $\nu \in \operatorname{Th}(S)$ and any $\delta > 0$, we construct $\nu_{\delta} \colon \mathbb{G} \longrightarrow S$ which sends $\mathbb{1}_{(t,1]} \mapsto \nu(\mathbb{1}_{(t+\delta,1]})$. It is readily checked that $\nu_{\delta} \in \operatorname{Th}(S)$. Also, whenever $S = \operatorname{Cu}(A)$ is the Cuntz semigroup of a stable rank one C*-algebra, we have that $\widehat{a_{\nu_{\delta}}} = \widehat{(a-\delta)}_+$.

Lemma 3.17. Let A be a C*-algebra of stable rank one. Let $\epsilon > 0$.

- (i) For any $a, b \in A_+$, we have that $\|\widehat{a} \widehat{b}\| \le \epsilon$ whenever $d_{Cu}(\nu_a, \nu_b) \le \epsilon$.
- (ii) For any $\nu, \eta \in Th(Cu(A))$ and any $\delta > 0$, we have that $d_{Cu}(\nu_{\delta}, \eta_{\delta}) \leq \epsilon$ whenever $d_{Cu}(\nu, \eta) \leq \epsilon$.

Proof. (i) Let $\tau \in T_1(A)$ and let $\lambda_{\tau} \in F(Cu(A))$ be its corresponding functional. Again, by [17], we have that

$$\tau(a) = \int_0^{\|a\|} \lambda_{\tau}([(a-t)_+])dt$$

$$= \int_0^{\|a\|} \lambda_{\tau}([(a-t+\epsilon)_+])dt + \int_0^{\epsilon} \lambda_{\tau}([(a-t)_+])dt$$

$$\leq \int_0^{\|a\|} \lambda_{\tau}([(b-t)_+])dt + \int_0^{\epsilon} \lambda_{\tau}([(1_A])dt$$

$$\leq \tau(b) + \epsilon$$

which proves the first statement.

(ii) We know that $\nu(\mathbb{1}_{(t,1]}), \eta(\mathbb{1}_{(t,1]}) \leq \eta(\mathbb{1}_{(t-\epsilon,1]}), \nu(\mathbb{1}_{(t-\epsilon,1]})$ for any $t \in [0,1]$. In particular, $\nu(\mathbb{1}_{(t-\delta,1]}), \eta(\mathbb{1}_{(t-\delta,1]}) \leq \eta(\mathbb{1}_{(t-\delta-\epsilon,1]}), \nu(\mathbb{1}_{(t-\delta-\epsilon,1]})$.

Theorem 3.18. Let A be a C^* -algebra of stable rank one. For any finite set $H \subseteq Aff T_1(A)$, there exists a finite set $N \subseteq Th(Cu(A))$ satisfying the following.

For any C*-algebra B of stable rank one, any *-homomorphisms $\phi, \psi \colon A \longrightarrow B$ and any $\epsilon > 0$, we have that $\max_{\hat{h} \in H} \{ \| \operatorname{Aff} T_1(\phi)(\hat{h}) - \operatorname{Aff} T_1(\psi)(\hat{h}) \| \} < \epsilon \text{ whenever } d_N(\operatorname{Cu}(\phi), \operatorname{Cu}(\psi)) < \epsilon/2$

Proof. Since Aff $T_1(A) \simeq A_{\rm sa}/A_0$, we can form a finite subset $H' \subseteq A_{\rm sa}$ of representatives of H. (That is, the assignment $H' \to H$ sending $h \mapsto \hat{h}$ is well-defined and surjective.) Furthermore, we know that there exist unique $h_+, h_- \in A_+$ such that $h = h_+ - h_-$.

Now consider the following finite subset of Th(Cu(A))

$$N := \{ \nu_{(h_+)}, \nu_{(h_-)} \mid h_+ - h_- \in H' \}.$$

Then for $\hat{h} \in H$ we compute that

$$\|\operatorname{Aff} T_{1}(\phi)(\widehat{h}) - \operatorname{Aff} T_{1}(\psi)(\widehat{h})\| = \|\widehat{\phi(h)} - \widehat{\psi(h)}\|$$

$$\leq \|(\widehat{\phi(h_{+})} - \widehat{\psi(h_{+})}) - (\widehat{\phi(h_{-})} - \widehat{\psi(h_{-})})\|$$

$$\leq \|\widehat{\phi(h_{+})} - \widehat{\psi(h_{+})}\| + \|\widehat{\phi(h_{-})} - \widehat{\psi(h_{-})}\|$$

$$\leq 2\epsilon/2 = \epsilon$$

which ends the proof.

Theorem 3.19. Let A be a C^* -algebra of stable rank one. Let $N \subseteq Th(Cu(A))$ be a set with a generating image. For any C^* -algebra B of stable rank one, any *-homomorphisms $\phi, \psi \colon A \longrightarrow B$, we have that

$$d(H(\phi), H(\psi)) \le d(Aff T_1(\phi), Aff T_1(\psi)) \le 2d_N(Cu(\phi), Cu(\psi))$$

whenever the Grothendieck group of the monoid generated by $\{\widehat{a_{(\nu_{\delta})}} \mid \nu \in \mathbb{N}, \delta \geq 0\}$ is dense in Aff $T_1(A)$.

Proof. Let us start by proving the following result.

Claim: For any $\nu \in \text{Th}(\text{Cu}(A))$ and any $\delta \geq 0$, we have $\|\operatorname{Aff} T_1(\phi)(\widehat{a_{(\nu_{\delta})}}) - \operatorname{Aff} T_1(\psi)(\widehat{a_{(\nu_{\delta})}})\| < \epsilon$ whenever $d_{\text{Cu}}(\operatorname{Cu}(\phi) \circ \nu, \operatorname{Cu}(\psi) \circ \nu) < \epsilon$.

Proof of the Claim: Let a_{ν} , a'_{ν} be a representatives of $\widehat{a_{\nu}}$. Define $b := \phi(a_{\nu})$ and $b' := \psi(a'_{\nu})$. Observe that $\nu_b = \operatorname{Cu}(\phi) \circ \nu$ and $\nu_{b'} = \operatorname{Cu}(\psi) \circ \nu$. Also that $\widehat{b} = \operatorname{Aff} T_1(\phi)(\widehat{a_{\nu}})$ and $\widehat{b'} = \operatorname{Aff} T_1(\psi)(\widehat{a_{\nu}})$. We get the result for the particular case $\delta = 0$ after applying (i) of Lemma 3.17 to b and b'. The general case follows by (ii) of Lemma 3.17.

Now, write \tilde{N} to be the monoid generated by N. It is readily checked that $d_{\tilde{N}}(Cu(\phi), Cu(\psi)) \leq d_{N}(Cu(\phi), Cu(\psi))$. Let $\hat{h} \in Aff T_{1}(A)$ and let $\epsilon > 0$. We know that there exists $\delta_{+}, \delta_{-} \geq 0$ and $\nu, \eta \in \tilde{N}$ such that $\|\hat{h} - (a_{(\nu\delta_{+})} - a_{(\eta\delta_{-})})\| < \epsilon$. Denote $h' := (a_{(\nu\delta_{+})} - a_{(\eta\delta_{-})})$. By the previous theorem (or rather, its proof), we know that for any $\hat{h} \in Aff T_{1}(A)$ and any $\epsilon > 0$, we have that

$$\|\operatorname{Aff} T_1(\phi)(\widehat{h}) - \operatorname{Aff} T_1(\psi)(\widehat{h})\| = 2\epsilon + \|\widehat{\phi(h')} - \widehat{\psi(h')}\|$$

$$\leq 2\epsilon + 2d_{\tilde{N}}(\operatorname{Cu}(\phi), \operatorname{Cu}(\psi))$$

$$\leq 2\epsilon + 2d_{\tilde{N}}(\operatorname{Cu}(\phi), \operatorname{Cu}(\psi)).$$

from which the result follows.

Question 3.20. Let X be a one-dimensional compact metric space and let B be a C*-algebra of a stable rank one. Let $\phi, \psi \colon C(X) \longrightarrow B$ be *-homomorphisms.

Is the monoid generated by $\{\widehat{a_{(\nu_{\delta})}} \mid \nu \in \mathcal{N}_X, \delta \geq 0\}$ dense in $C_{\mathbb{R}}(X)$ where \mathcal{N}_X induces the Cu-metric? Does there exist a weaker assumption providing the following inequality

$$d(H(\phi), H(\psi)) \le 2d_{Cu}(Cu(\phi), Cu(\psi))$$
?

4. Distinguishing C*-algebras and *-homomorphisms

We apply the above to distinguish concrete examples of C*-algebras and *-homomorphisms that neither the original Elliott invariant nor the (unitary) Cuntz semigroup are able to distinguish. These examples highlight the neatness and applicability of our methods to classification problems.

A. The Gong-Jiang-Li example revisited. In [19], the authors have built (non-simple) AT-algebras A and B that are distinguished by a refined version of the Elliott invariant. More particularly, these C*-algebras differ from an internal property, called uniformly varied determinants of connecting maps, which in turn implies a system of splitting between specific corner algebras. In the original manuscript, both the invariant and the distinguishing arguments are quite long to unravel. Also, it is stated without any proof that these C*-algebras cannot be distinguished by their Cuntz semigroup.

We aim to distinguish these C^* -algebras via the Hausdorffized unitary Cuntz semigroup and our methods, providing a more conceptual and streamlined proof of the non-isomorphism between A and B. Let us start by recalling their constructions.

• Construction of the blocks.

Let $(p_n)_{n\geq 1}$ denotes the sequence of all the prime numbers in increasing order.

Let $(k_n)_{n\geq 1}$ be a strictly increasing sequence of natural numbers such that $k_1\geq 2$.

The building blocks of the inductive system for A and B are defined by

$$\begin{split} A_1 &= B_1 = C(\mathbb{T}) \\ A_2 &= B_2 = M_{p_1^{k_1}}(C([0,1]]) \oplus M_{p_1^{k_1}}(C(\mathbb{T})) \\ A_3 &= B_3 = M_{p_1^{k_1}p_1^{k_2}}(C([0,1]]) \oplus M_{p_1^{k_1}p_2^{k_2}}(C([0,1]]) \oplus M_{p_1^{k_1}p_2^{k_2}}(C(\mathbb{T})) \\ A_4 &= B_4 = M_{p_1^{k_1}p_1^{k_2}p_1^{k_3}}(C([0,1]]) \oplus M_{p_1^{k_1}p_2^{k_2}p_2^{k_3}}(C([0,1]]) \oplus M_{p_1^{k_1}p_2^{k_2}p_3^{k_3}}(C([0,1])) \oplus M_{p_1^{k_1}p_2^{k_2}p_3^{k_3}}(C([0,1])) \\ &\vdots \end{split}$$

Let $n \in \mathbb{N}$. Write $[n, i] := \prod_{j=1}^{i} p_j^{k_j} \prod_{j=i+1}^{n-1} p_i^{k_j}$, for any $1 \le i \le n-1$ and [n, n] := [n, n-1]. (Notice that $[n+1, i] = p_i^{k_n} [n, i]$ for any $1 \le i \le n-1$.) We construct inductively

$$A_n = B_n = \bigoplus_{i=1}^{n-1} M_{[n,i]}(C([0,1])) \oplus M_{[n,n]}(C(\mathbb{T})).$$

• Construction of the connecting maps.

Let $(t_n)_{n\geq 1}$ be a countable dense subset of [0,1].

Let $(z_n)_{n>1}$ be a countable dense subset of \mathbb{T} .

Let $n \in \mathbb{N}$. The connecting maps $\phi_{nn+1}: A_n \longrightarrow A_{n+1}$ and $\psi_{nn+1}: B_n \longrightarrow B_{n+1}$ of the inductive systems are defined also inductively, via partial *-homomorphisms as follows.

• The n-1 first partial *-homomorphisms for ϕ_{nn+1} and ψ_{nn+1} . Let $1 \le i \le n-1$.

$$\phi_{nn+1}^i = \psi_{nn+1}^i \colon M_{[n,i]}(C([0,1])) \longrightarrow M_{[n+1,i]}(C([0,1]))$$

$$f \longmapsto \begin{pmatrix} f & & \\ & \ddots & \\ & & f_{(t_n)} \end{pmatrix}$$

• The n-th partial *-homomorphism for ϕ_{nn+1} and ψ_{nn+1} . Let exp: $[0,1] \to \mathbb{T}$ be the exponential map sending $t \mapsto e^{2i\pi t}$.

$$\phi_{nn+1}^n \colon M_{[n,n]}(C(\mathbb{T})) \longrightarrow M_{[n+1,n]}(C([0,1]))$$

$$f \longmapsto \begin{pmatrix} f \circ \exp^{-1} & & & \\ & f \circ \exp(1/r_n) & & \\ & & \ddots & \\ & & & f \circ \exp(r_n - 1/r_n) \end{pmatrix}$$

where $r_n := p_n^{k_n} - 1$. Now let $l_n := 4^n [n, n+1]$.

$$\psi_{nn+1}^{n} \colon M_{[n,n]}(C(\mathbb{T})) \longrightarrow M_{[n+1,n]}(C([0,1]))$$

$$f \longmapsto \begin{pmatrix} f \circ \exp^{t_{n}} & & & \\ & f \circ \exp(t) & & & \\ & & f \circ \exp(t/r_{n}) & & \\ & & \ddots & & \\ & & & f \circ \exp(r_{n}-1/r_{n}) \end{pmatrix}$$

• The (n+1)-th partial *-homomorphism for ϕ_{nn+1} and ψ_{nn+1} .

$$\phi_{nn+1}^{n+1} = \psi_{nn+1}^{n+1} \colon M_{[n,n]}(C(\mathbb{T})) \longrightarrow M_{[n+1,n+1]}(C(\mathbb{T}))$$

$$f \longmapsto \begin{pmatrix} f \\ f(z_n) \\ & \ddots \\ & f(z_n) \end{pmatrix}$$

We finally define

$$A := \underset{\stackrel{\longrightarrow}{\lim}}{\underset{n}{\lim}} (A_n, \phi_{nn+1})$$
 and $B := \underset{\stackrel{\longrightarrow}{\lim}}{\underset{n}{\lim}} (B_n, \psi_{nn+1})$

where $\phi_{nn+1} = (\phi_{nn+1}^1, ..., \phi_{nn+1}^{n-1}, (\phi_{nn+1}^n, \phi_{nn+1}^{n+1}))$ and $\psi_{nn+1} = (\psi_{nn+1}^1, ..., \psi_{nn+1}^{n-1}, (\psi_{nn+1}^n, \psi_{nn+1}^{n+1}))$

Proposition 4.1. Both A and B are separable unital stable rank one C^* -algebras.

Proof. Since all C*-algebras of the inductive systems are separable and unital, together with the fact that all morphisms are also unital, we easily obtain that A and B are unital separable C*-algebras. In addition, the stable rank one property is preserved by inductive limits and any interval or circle algebra has stable rank one.

Next, we dive into the lattice of ideals of both A and B. More particularly, we exhibit the set of simple ideals of these C^* -algebras together with another set of peculiar ideals that will be of use to distinguish A and B by means of the Hausdorffized unitary Cuntz semigroup.

Lemma 4.2. Let $j \geq 1$. We consider the following inductive systems

$$\left\{ \begin{array}{ll} \mathfrak{s}_j := \varinjlim_{n \geq j} (A_n^j), \phi_{nm_{\mid A_n^j}}), & \quad \mathfrak{p}_j := \varinjlim_{n \geq j} (I_{n,j}, \phi_{nm_{\mid I_{n,j}}}) & \text{ and } \quad \mathfrak{a}_j := \varinjlim_{n \geq j} (I_{n,j}^c, \phi_{nm_{\mid I_{n,j}^c}}). \\ \mathfrak{t}_j := \varinjlim_{n \geq j} (B_n^j), \phi_{nm_{\mid A_n^j}}), & \quad \mathfrak{q}_j := \varinjlim_{n \geq j} (I_{n,j}, \psi_{nm_{\mid I_{n,j}}}) & \text{ and } \quad \mathfrak{b}_j := \varinjlim_{n \geq j} (I_{n,j}^c, \psi_{nm_{\mid I_{n,j}^c}}). \end{array} \right.$$

where $I_{n,j} = \bigoplus_{i \geq j}^{n-1} M_{[n,i]}(C([0,1])) \oplus M_{[n,n]}(C(\mathbb{T}))$ and $I_{n,j}^c = \bigoplus_{i=1}^{j-1} M_{[n,i]}(C([0,1]))$ are complementary of one another in A_n and B_n .

- (i) The sets of simple ideals of A and B are respectively $\{\mathfrak{s}_j\}_{j\geq 1}$ and $\{\mathfrak{t}_j\}_{j\geq 1}$.
- (ii) We have $\mathfrak{p}_j = \bigoplus_{i=1}^{j-1} \mathfrak{s}_i$ and $\mathfrak{q}_j = \bigoplus_{i=1}^{j-1} \mathfrak{t}_i$. (iii) Both \mathfrak{q}_j and \mathfrak{b}_j are ideals of A and B respectively and

$$\begin{cases} A = \mathfrak{a}_j \oplus \mathfrak{p}_j. \\ B = \mathfrak{b}_j \oplus \mathfrak{q}_j. \end{cases}$$

Proof. This is done similarly as in the proof of [5, Theorem 3.2].

Not only has it been proved in [19] that A and B agree at the level of the (original) Elliott invariant, but also it has been stated that they agree at the level of the Cuntz semigroup. Even though this statement is true, we would like to offer a proof of this, via an approximate intertwining argument.

П

In [4], an approximate intertwining theorem has been developed only for specific Cu-semigroups termed uniformly-based. Later on in [8], this technique has been generalized to the entire category Cu and also to the category Cu*. Even though the theorems have been stated in great generality using the finite-set comparison, we can restate them in our specific case as follows.

Theorem 4.3. [4, Theorem 3.16] - [8, Theorem 3.17]. Let $(S_i, \sigma_{ij}), (T_i, \tau_{ij})$ be inductive sequences in the category Cu*.

Assume that there are Cu^* -morphisms $\alpha_i : S_i \to T_i$ and $\beta_i : T_i \to S_{i+1}$ together with a metric d^* such that

$$d^*(\beta_i \circ \alpha_i, \sigma_{ii+1}) < 1/2^i$$
 and $d^*(\alpha_{i+1} \circ \beta_i, \tau_{ii+1}) < 1/2^i$

Then there exists a Cu^* -isomorphism between $\lim_{i \to \infty} (S_i, \sigma_{ij}) \simeq \lim_{i \to \infty} (T_i, \tau_{ij})$.

Lemma 4.4. Let n > 1. We compute that

$$d_{\text{Cu}}(\text{Cu}(\phi_{nn+1}^n), \text{Cu}(\psi_{nn+1}^n)) = d_{\text{Cu}}^*(\text{Cu}_{\text{K}_1}(\phi_{nn+1}^n), \text{Cu}_{\text{K}_1}(\psi_{nn+1}^n)) \le 1/r_n.$$

Proof. First, let us compute the Cu-distance. Let $U \subseteq \mathbb{T}$ be an open set. Let $k_U \in \mathbb{N}$ be the number of elements of the set $\{\exp(i/r_n) \mid 1 \le i \le r_n - 1\}$ belonging to U. We compute that

$$k_U \le \mathrm{Cu}(\phi_{nn+1}^n)(1_U)(t), \mathrm{Cu}(\psi_{nn+1}^n)(1_U)(t) \le k_U + 2.$$

Further, we either have that $U_{1/r_n} = \mathbb{T}$ or else, $k_{U_{1/r_n}} = k_U + 2$. In both cases, we obtain the following inequalities for any open set $U \subseteq \mathbb{T}$

$$Cu(\phi_{nn+1}^n)(1_U) \le Cu(\psi_{nn+1}^n)(1_{U_{r_n}})$$
 and $Cu(\psi_{nn+1}^n)(1_U) \le Cu(\phi_{nn+1}^n)(1_{U_{r_n}})$.

Second, let us compute the Cu*-distance. Let $U \subseteq \mathbb{T}$ be an open set. We easily see that $k_{U_1/r_n} > 0$. Therefore, we compute that for any $U \subseteq \mathbb{T}$

$$Cu(\phi)(1_{U_{1/r}}), Cu(\psi)(1_{U_{1/r}}) \geq \mathbb{1}_{[0,1]}.$$

Let us write $x_A = \mathrm{Cu}(\phi)(1_U), y_A = \mathrm{Cu}(\phi)(1_{U_{1/r_n}})$ and similarly we write $x_B = \mathrm{Cu}(\psi)(1_U), y_B = \mathrm{Cu}(\psi)(1_U)$ $\mathrm{Cu}(\phi)(1_{U_{1/r_n}})$. We know that $\mathrm{K}_1(I_{y_A}) \simeq \mathrm{K}_1(I_{y_B}) \simeq \{0\}$. Therefore, the fiber diagram of $\mathrm{Cu}(\phi)$ and $Cu(\psi)$ at coordinates $(1_U, 1_{U_{r_n}})$ trivially commutes, which ends the proof. **Theorem 4.5.** We have the following isomorphisms.

- (i) $Cu(A) \simeq Cu(B)$.
- (ii) $Cu_{K_1}(A) \simeq Cu_{K_1}(B)$. In particular, A and B agree at the level of K_0 and K_1 . Furthermore, any (scaled) Cu-isomorphism $\alpha_0 \colon Cu(A) \simeq Cu(B)$ maps $\mathfrak{p}_j \mapsto \mathfrak{q}_j$, for all $j \geq 1$.

Proof. To prove (i) and (ii), we note that $A_n = B_n$ and that $\phi_{nn+1}^i = \psi_{nn+1}^i$ for all $n \in \mathbb{N}$ and all $i \leq n+1$ such that $i \neq n$. Therefore we can apply the approximate intertwining theorem with identity maps between the sequences involved and the computation obtained in the previous lemma.

Now, consider any scaled Cu-isomorphism α_0 : $\operatorname{Cu}(A) \simeq \operatorname{Cu}(B)$. Observe that $\operatorname{Cu}(I) \simeq \operatorname{Cu}(I_{\alpha_0})$. In particular, we have an isomorphism $\operatorname{Cu}(I)_c \simeq \operatorname{Cu}(I_{\alpha_0})_c$ between their monoids of compact elements. In the stable rank one context, this implies that $\operatorname{K}_0(I) \simeq \operatorname{K}_1(I_{\alpha_0})$. Furthermore, simple ideals of A are mapped to simple ideals of B and it is readily computed that for all $j \geq 1$

$$\mathrm{K}_0(\mathfrak{s}_j) \simeq \mathrm{K}_0(\mathfrak{t}_j) \simeq \mathbb{Z}[\frac{1}{p_j}].$$

Therefore, θ_{α_0} maps $\mathfrak{s}_j \mapsto \mathfrak{t}_j$ and α_0 maps $[1_{\mathfrak{s}_i}] \mapsto [1_{\mathfrak{t}_i}]$.

Finally, we observe that $[1_A] = \sum_{k=1}^{j-1} [1_{\mathfrak{s}_k}] + [1_{\mathfrak{p}_j}]$ for any j. Similarly for B. Again, in the stable rank one context, it is well-known that $\mathrm{Cu}(A)$ and $\mathrm{Cu}(B)$ have cancellation of compact elements. (See e.g. [25, Theorem 4.3].) As a result, we deduce that $\alpha_0([1_{\mathfrak{p}_j}]) = [1_{\mathfrak{q}_j}]$ for any $j \geq 1$.

Theorem 4.6. Let $j \geq 1$. Consider the canonical embeddings $\iota_j^A : \mathfrak{p}_j \hookrightarrow A$ and $\iota_j^B : \mathfrak{q}_j \hookrightarrow B$.

- (i) $\{\overline{K}_1(\iota_i^A)\}_{j\geq 1}$ are simultaneously Nielsen-Thomsen diagonalisable.
- (ii) $\{\overline{K}_1(\iota_i^B)\}_{j\geq 1}$ are not simultaneously Nielsen-Thomsen diagonalisable.

$$[j,j]-1$$
-times

Proof. (i) Let $j \geq 1$. We define $u_j := \operatorname{diag}(\operatorname{id}_{\mathbb{T}}, 1, \ldots, 1) \in M_{[j,j]}(C(\mathbb{T}))$ and we denote by C_j the Nielsen-Thomsen basis of \mathfrak{p}_j induced by the section $s_j : 1_{\mathbb{Z}} \mapsto \phi_{j\infty}(u_j)$. Let us check that $\overline{K}_1(\iota_j^A)$ is diagonalisable in the bases C_j , C_1 . By Corollary 2.7, it is enough to have that


$$\overline{\Delta}\left(\begin{smallmatrix}\phi_{j\infty}(u_j) & 0\\ 0 & \phi_{1\infty}^*(u_1)\end{smallmatrix}\right) = 0 \text{ in } H(B).$$

This is readily deduced from the partial *-homomorphisms defining the connecting maps of A.

(ii) Let $j \geq 1$. Observe that $B = \mathfrak{q}_1 \supseteq \mathfrak{q}_j \supseteq \mathfrak{q}_{j+1}$ and that $0 = \mathfrak{b}_1 \subseteq \mathfrak{b}_j \subseteq \mathfrak{b}_{j+1}$. Combined with Lemma 4.2 (ii), we deduce that

$$\theta_j^i \colon \mathfrak{q}_j \xrightarrow{\iota_j^B} B \xrightarrow{\pi_i} \mathfrak{b}_i$$
 is the zero morphism for any $1 \le i \le j$.

Now, assume that $\overline{K}_1(\iota_j^B)$ is diagonalisable. By Corollary 2.7, there exist sections s_j, s_1 of $\overline{K}_1(\mathfrak{q}_j) \twoheadrightarrow K_1(\mathfrak{q}_j), \overline{K}_1(B) \twoheadrightarrow K_1(B)$ respectively, such that the following diagram commutes for any $1 \le i \le j$



Therefore, if we assume that $\{\overline{K}_1(\iota_j^B)\}_{j\geq 1}$ are simultaneously diagonalisable in Nielsen-Thomsen bases $\{\mathcal{C}_j\}_j$, this would imply that $\overline{K}_1(\pi_i)\circ s_1$ is trivial for all $i\geq 1$, where s_1 is the section induced by \mathcal{C}_1 . However, it has been computed in [19, pp. 73-74], that for any section s_1 and any representative $u\in B$ of $s_1(1_{\mathbb{Z}})$, we can find $m\in\mathbb{N}$ big enough such that $\|\overline{K}_1(\pi_m)\circ s_1(1_{\mathbb{Z}})\|\geq 3$ which leads to a contradiction.

Corollary 4.7. There is no Cu^* -isomorphism between $Cu_{\overline{K}_1}(A)$ and $Cu_{\overline{K}_1}(B)$.

Proof. Assume that there exists a Cu^{\sim} -isomorphism $\alpha \colon \operatorname{Cu}_{\overline{K}_1}(A) \longrightarrow \operatorname{Cu}_{\overline{K}_1}(A)$. We know that α induces a Cu-isomorphism $\alpha_0 \colon \operatorname{Cu}(A) \longrightarrow \operatorname{Cu}(B)$, which in turn induces a lattice isomorphism $\theta_{\alpha_0} \colon \operatorname{Lat}(A) \simeq \operatorname{Lat}(B)$. Furthermore, by Theorem 4.5, we know that $\theta_{\alpha_0}(\mathfrak{p}_j) = \mathfrak{q}_j$ for any $j \geq 1$.

Now, by the properties of the Cu_K -constructions recalled in Theorem 3.2 (ii)/(ii') -see also [7, Theorem 4.5]-, we know that the following diagram is commutative with exact rows

$$0 \longrightarrow \operatorname{Cu}(\mathfrak{p}_{j}) \longrightarrow \operatorname{Cu}_{\overline{K}_{1}}(\mathfrak{p}_{j}) \longrightarrow \overline{K}_{1}(\mathfrak{p}_{j}) \longrightarrow 0$$

$$\downarrow^{\alpha_{\max}} \qquad \qquad \downarrow^{\alpha_{\max}} \qquad \downarrow^{\alpha_{\max}}$$

where the vertical arrows are isomorphisms in their respective categories obtained via restriction. In particular, we obtain that

$$\overline{\mathrm{K}}_1(\mathfrak{p}_i) \simeq \overline{\mathrm{K}}_1(\mathfrak{q}_i)$$
 for any $j \geq 1$

from which we deduce that

$$\overline{K}_{1}(\mathfrak{p}_{j}) \xrightarrow{\simeq} \overline{K}_{1}(\mathfrak{q}_{j})$$

$$\overline{K}_{1}(\iota_{j}^{A}) \downarrow \qquad \qquad \downarrow \overline{K}_{1}(\iota_{j}^{B})$$

$$\overline{K}_{1}(A) \xrightarrow{\simeq} \overline{K}_{1}(B)$$

commutes. Nevertheless, by Theorem 4.6 we know that $\{\overline{K}_1(\iota_j^A)\}_j$ are simultaneously Nielsen-Thomsen diagonalisable while $\{\overline{K}_1(\iota_j^B)\}_j$ are not, which leads to a contradiction.

B. The Robert example(s) revisited. In [6] the author has exposed a pair of non-unitarily equivalent *-homomorphisms from $C(\mathbb{T})$. (This was based on private communication with L. Robert.) The original distinction relied on the information given by de la Harpe-Skandalis determinant of the identity map, we intend to generalize these constructions and (re)state similar results via our methods. We also obtain additional information on this family by measuring how far these *-homomorphisms are from one another with respect to the metric \mathfrak{d} .

We start by recalling the construction of the first pair *-homomorphisms exposed in [6, Section 4.A].

• Construction of the unitary elements of $C([0,1]) \otimes M_{2^{\infty}}$. Recall that $M_{2^{\infty}}$ can be written as the inductive limit of $(M_{2^n}, \phi_{nm})_n$ where $\phi_{nn+1} : M_{2^n} \longrightarrow M_{2^{n+1}}$ sends $a \longmapsto \binom{a}{a}$.

For any $n \in \mathbb{N}$ we consider the following unitary element of M_{2^n}

$$w_n := \begin{pmatrix} 1 & e^{2i\pi/2^n} & & & \\ & \ddots & & & \\ & & \ddots & & \\ & & & e^{2i\pi(2^n - 1)/2^n} \end{pmatrix}.$$

It can be argued that the sequence $(\phi_{n\infty}(w_n))_n$ converges towards a unitary $w \in M_{2^{\infty}}$ with full spectrum. (See [6, 4.A].) Finally, we define unitary elements of $C([0,1]) \otimes M_{2^{\infty}}$ for any $k \in \mathbb{N}$ as follows

$$u_0 := 1_{[0,1]} \otimes w$$
 and $u_k := e^{2i\pi k \operatorname{id}_{[0,1]}} \otimes w$

inducing *-homomorphisms $\varphi_{u_k} : C(\mathbb{T}) \longrightarrow C([0,1]) \otimes M_{2^{\infty}}$ mapping $\mathrm{id}_{\mathbb{T}}$ to u_k respectively.

Theorem 4.8. We compute that

$$d_{\operatorname{Cu}}^*(\operatorname{Cu}_{\operatorname{K}_1}(\varphi_{u_k}),\operatorname{Cu}_{\operatorname{K}_1}(\varphi_{u_l})) = 0 \quad and \quad \mathfrak{d}(\overline{\operatorname{K}}_1(\varphi_{u_k}),\overline{\operatorname{K}}_1(\varphi_{u_l})) = \frac{|k-l|}{2}.$$

Proof. Let $k, l \in \mathbb{N}$ and $n \in \mathbb{N}$. We consider $u_{k,n} := e^{2i\pi k \operatorname{id}_{[0,1]}} \otimes w_n$ and $u_{l,n} := e^{2i\pi l \operatorname{id}_{[0,1]}} \otimes w_n$. We proceed similarly as in the proof of [6, 4.A] to compute that

$$d_{\mathrm{Cu}}(\mathrm{Cu}(\varphi_{u_{k,n}}),\mathrm{Cu}(\varphi_{u_{l,n}})) \le 1/2^n.$$

Now, let $U \subseteq \mathbb{T}$ be an open set. It is readily observed that the ideals generated by $\operatorname{Cu}(\varphi_{u_{k,n}})(\mathbb{1}_{U_{\frac{1}{2^n}}})$ and $\operatorname{Cu}(\varphi_{u_{l,n}})(\mathbb{1}_{U_{\frac{1}{2^n}}})$ are equal to A. Combined with the fact that A has trivial K_1 -group, we deduce that the fiber diagram of $\operatorname{Cu}(\varphi_{u_{k,n}})$ and $\operatorname{Cu}(\varphi_{u_{l,n}})$ at coordinates $(1_U,1_{U_{\frac{1}{2^n}}})$ trivially commutes. This yields

$$d_{\operatorname{Cu}}^*(\operatorname{Cu}_{\operatorname{K}_1}(\varphi_{u_{k,n}}), \operatorname{Cu}_{\operatorname{K}_1}(\varphi_{u_{l,n}})) \le 1/2^n$$

We remark that for any $k \in \mathbb{N}$, $d_U((\mathrm{id} \otimes \phi_{n\infty})(u_{k,n}), u_k) \xrightarrow[n \to \infty]{} 0$. We now deduce the first computation by a standard argument.

Next, let us compute the distance \mathfrak{d} between $\overline{\mathrm{K}}_1(\varphi_{u_k})$ and $\overline{\mathrm{K}}_1(\varphi_{u_l})$. First, it is immediate that $\mathrm{K}_1(\varphi_{u_k}) = \mathrm{K}_1(\varphi_{u_l})$ is the zero-morphism. Also, we know that $d_{\mathrm{Cu}}(\mathrm{Cu}(\varphi_{u_k}), \mathrm{Cu}(\varphi_{u_l})) = 0$, which implies that $d(\mathrm{H}(\varphi_{u_k}), \mathrm{H}(\varphi_{u_l})) = 0$ by Theorem 3.14. Now, let us fix \mathcal{C}_0 to be the canonical Nielsen-Thomsen basis of $C(\mathbb{T})$ and let \mathcal{D} be any Nielsen-Thomsen basis of A.

Observe that $R_{\mathcal{C}_0\mathcal{D}}(\varphi_{u_k}, \varphi_{u_l}) \colon \mathbb{Z} \longrightarrow \mathrm{H}(A)$ sends $1_{\mathbb{Z}} \longmapsto \overline{\Delta} \begin{pmatrix} u_k & 0 \\ 0 & u_l \end{pmatrix} = \overline{\Delta}(u_k) - \overline{\Delta}(u_l)$. Furthermore, it is well-known that Aff $T_1(A) \simeq C([0,1],\mathbb{R})$ and that $\mathrm{K}_0(A) \simeq \mathbb{N}[\frac{1}{2}]$. We deduce that

$$H(A) \simeq C([0,1],\mathbb{R})/\{\text{constant functions}\}.$$

Finally, standard arguments show that the quotient-norm in $C([0,1],\mathbb{R})/\{\text{constant functions}\}$ is given by $||[f]|| = \frac{1}{2}(\max f(t) - \min f(t))$ for any $f \in C([0,1],\mathbb{R})$.

Now, let $C := \tau_M(h_w) \in \mathbb{R}$, where τ_M is the unique trace on $M_{2^{\infty}}$ and $h_w \in M_{2^{\infty}}$ is any self-adjoint element such that $e^{2i\pi h_w} = w$. The computations done in [6, 4.A] give us that

$$\overline{\Delta}(u_k) = [t \longmapsto C + kt]_{H(A)}$$

for any $k \in \mathbb{N}$ from which we deduce the second computation.

As an immediate consequence, we conclude the following behaviour for the family $\{\varphi_{u_k}\}_k$ of *-homomorphisms from $\mathbb{C}(\mathbb{T})$ to A.

Corollary 4.9. The *-homomorphisms $\{\varphi_{u_k}\}_k$ all agree on the Cuntz semigroup and the unitary Cuntz semigroup. Yet, they are pairwise not approximately unitarily equivalent.

More particularly, for any distinct $k, l \in \mathbb{N}$, the *-homomorphisms φ_{u_k} and φ_{u_l} are distinguished by the Hausdorffized algebraic K_1 -group.

(And a fortiori, by the Hausdorffized unitary Cuntz semigroup.)

- C. A novel example. We end the manuscript with a novel example that illustrates the necessity of the Hausdorffized unitary Cuntz semigroup in order to classify *-homomorphisms from $C(\mathbb{T})$. We exhibit a pair of unitary elements u, v of an AI-algebra A whose induced *-homomorphisms φ_u, φ_v are shown to agree on both the unitary Cuntz semigroup and the Hausdorffized algebraic K_1 -group. (This, in turn, implies that they coincide on the Cuntz semigroup and the K_1 -group.) Nevertheless, φ_u and φ_v are not approximately unitarily equivalent, as they are distinguished by the Hausdorffized unitary Cuntz semigroup and thereby, showcases its ability to distinguish *-homomorphisms that cannot be separated by any of the other invariants considered in this study.
- Construction of A and its unitary elements. Let us consider A to be the unital AI-algebra obtained as the inductive limit of $(C([0,1]) \otimes M_{2^n}, \phi_{nn+1})_{n \in \mathbb{N}}$ where $\phi_{nn+1} \colon C([0,1]) \otimes M_{2^n} \longrightarrow C([0,1]) \otimes M_{2^{n+1}}$ sends $f \longrightarrow \begin{pmatrix} f \\ f(0) \end{pmatrix}$.

Let us consider two piecewise-linear functions $f,g\colon [0,1] \longrightarrow \mathbb{R}$ as follows.

- (i) Set f(0) = 0, $f(\frac{1}{2}) = \frac{1}{4}$, f(1) = 0 and define f to be linear between these points.
- (ii) Set g(0) = 0, $g(\frac{1}{2}) = 0$, g(1) = 0, $g(\frac{1}{4}) = \frac{1}{4}$, $g(\frac{3}{4}) = \frac{1}{4}$ and define g to be linear between these points.

For any $n \in \mathbb{N}$ we consider the following diagonal unitary elements of $C([0,1]) \otimes M_{2^n}$.

$$u_n := \operatorname{diag}(e^{2i\pi f}, \lambda_{1,n}, \dots, \lambda_{2^{n-1}-1,n}, e^{i\pi}e^{2i\pi g}, e^{i\pi}\lambda_1, \dots, e^{i\pi}\lambda_{2^{n-1}-1,n})$$
 and $v_n := e^{i\pi}u_n$
where $\lambda_{k,n} := (e^{2i\pi k/2^{n-1}})^{\frac{1}{4}}$ for any $1 \le k \le 2^{n-1} - 1$.

As in the previous example, it can be argued that $d_U(\phi_{nm}(u_n), u_m) \leq 1/2^{n-1} - 1/2^{m-1}$. (See also the original argument in [6, 4.A]. Roughly speaking, both $\phi_{nm}(u_n)$ and u_m have the same two 'moving' eigenvalues and their 'fixed' eigenvalues can be paired to be at distance at most $1/2^{n-1} - 1/2^{m-1}$.)

Consequently, the sequence $(\phi_{n\infty}(u_n))_n$ is Cauchy for d_U and hence, converges towards a unitary element u in A. Similarly the sequence $(\phi_{n\infty}(v_n))_n$ converges towards a unitary element v in A and we observe that $\operatorname{sp}(u) = \operatorname{sp}(v) = [1, e^{2i\pi/4}] \cup [-1, e^{-2i\pi/4}] \subseteq \mathbb{T}$.

Theorem 4.10. We compute that

$$\begin{cases} d_{\operatorname{Cu}}^*(\operatorname{Cu}_{K_1}(\varphi_u), \operatorname{Cu}_{K_1}(\varphi_v)) = 0 \\ \mathfrak{d}(\overline{K}_1(\varphi_u), \overline{K}_1(\varphi_v)) = 0 \\ \mathfrak{d}_{\operatorname{Cu}}^*(\operatorname{Cu}_{\overline{K}_1}(\varphi_u), \operatorname{Cu}_{\overline{K}_1}(\varphi_v)) \ge 1/8 \end{cases}$$

As a consequence, φ_u and φ_v are not approximately unitarily equivalent even though they agree on the unitary Cuntz semigroup and the Hausdorffized algebraic K_1 -group.

Proof. We proceed similarly as in the previous example and the proof of [6, 4.A] to compute that

$$d_{\mathrm{Cu}}(\mathrm{Cu}(\varphi_{u_n}),\mathrm{Cu}(\varphi_{v_n})) \le 1/2^{n-1}.$$

Let $U \subseteq \mathbb{T}$ be an open set. We write $V := U_{\frac{1}{2^{n-1}}}$. It is readily observed that the ideals generated by $Cu(\varphi_{u_{k,n}})(\mathbb{1}_V)$ and $Cu(\varphi_{u_{l,n}})(\mathbb{1}_V)$ are either trivial or equal to A. Combined with the fact both the trivial algebra and A have trivial K_1 -groups, we obtain that

$$d_{\operatorname{Cu}}^*(\operatorname{Cu}_{\operatorname{K}_1}(\varphi_{u_n}), \operatorname{Cu}_{\operatorname{K}_1}(\varphi_{v_n})) \le 1/2^{n-1}$$

from which the first computation follows.

For the second computation, observe that $u = e^{i\pi}v$ and therefore we immediately obtain that $\overline{\Delta}(u) = \overline{\Delta}(v)$. Combined with the fact that $K_1(\varphi_u)$ and $K_1(\varphi_v)$ are trivial and that $d(H(\varphi_u), H(\varphi_v)) = 0$ since $d_{\text{Cu}}(\text{Cu}(\varphi_u), \text{Cu}(\varphi_v)) = 0$, then we conclude that

$$\mathfrak{d}(\overline{\mathrm{K}}_{1}(\varphi_{u}),\overline{\mathrm{K}}_{1}(\varphi_{v}))=0.$$

For the third computation, let us write $\alpha := \operatorname{Cu}_{\overline{K}_1}(\varphi_u)$ and $\beta := \operatorname{Cu}_{\overline{K}_1}(\varphi_v)$. Similarly, we write $\alpha_n := \operatorname{Cu}_{\overline{K}_1}(\varphi_{u_n})$ and $\beta_n := \operatorname{Cu}_{\overline{K}_1}(\varphi_{v_n})$. Recall that $\alpha = (\alpha_0, \{\alpha_I\}_{I \in \operatorname{Lat}_f(A)})$ where $\alpha_0 := \operatorname{Cu}(\varphi_u)$ and $\alpha_I := \overline{K}_1(I \xrightarrow{\varphi_u} I_{\varphi_u})$. Similarly for β , α_n and β_n .

We consider the ideal of $C(\mathbb{T})$ generated by the element $x:=\mathbbm{1}_{(1,e^{2i\pi/4})}\in \mathrm{Lsc}(\mathbb{T},\overline{\mathbb{N}})$ that we write I. Observe that $I_{\varphi_u}=I_{\varphi_v}=I_{\varphi_{u_n}}=I_{\varphi_{v_n}}=A$. Therefore, for any $y\in \mathrm{Lsc}(\mathbb{T},\overline{\mathbb{N}})$ such that $x\ll y$, the fiber diagram of α and β at coordinates (x,y) falls down to

$$\overline{\mathrm{K}}_{1}(I) \xrightarrow{\alpha_{I}} \overline{\mathrm{K}}_{1}(A)$$

Similarly for the fiber diagram of α_n and β_n at coordinates (x, y).

We aim to compute a (non-zero) lower-bound m > 0 for $\mathfrak{d}(\alpha_I, \beta_I)$. By Proposition 3.8, this will in turn imply that $d^*_{\mathrm{Cu},\mathfrak{d}}(\mathrm{Cu}_{\overline{\mathrm{K}}_1}(\varphi_u), \mathrm{Cu}_{\overline{\mathrm{K}}_1}(\varphi_v)) \geq m/4$. First, observe that it is enough to find a lower-bound for $\mathfrak{d}((\alpha_n)_I, (\beta_n)_I)$ for some $n \in \mathbb{N}$ big enough, since we have that

$$\begin{split} \mathfrak{d}((\alpha_n)_I,(\beta_n)_I) &\leq \mathfrak{d}(\overline{\mathrm{K}}_1(\phi_{n\infty}) \circ (\alpha_n)_I, \overline{\mathrm{K}}_1(\phi_{n\infty}) \circ (\beta_n)_I) \\ &\leq \mathfrak{d}(\alpha_I, \overline{\mathrm{K}}_1(\phi_{n\infty}) \circ (\alpha_n)_I) + \mathfrak{d}(\alpha_I,\beta_I) + \mathfrak{d}(\beta_I, \overline{\mathrm{K}}_1(\phi_{n\infty}) \circ (\beta_n)_I) \\ &\leq 2\epsilon + \mathfrak{d}(\alpha_I,\beta_I). \end{split}$$

Then, to compute $\mathfrak{d}((\alpha_n)_I, (\beta_n)_I)$, we proceed similarly as in the previous example to compute that $\overline{\Delta}((u_n)_I) = [t \mapsto C + 4f]_{\mathrm{H}(A)}$ and $\overline{\Delta}((v_n)_I) = [t \mapsto C' + 4g]_{\mathrm{H}(A)}$, for some constants $C, C' \in \mathbb{R}$. We finally deduce that $\mathfrak{d}((\alpha_n)_I, (\beta_n)_I) \geq \frac{1}{2}(\max 4(f-g)(t) - \min 4(f-g)(t)) = 1/2$ and hence, the last computation.

References

- [1] R. Antoine, F. Perera, L. Robert, and H. Thiel. C*-algebras of stable rank one and their Cuntz semigroups. *Duke Math. J.*, 171(1):33–99, 2022.
- [2] R. Antoine, F. Perera, and H. Thiel. Tensor products and regularity properties of Cuntz semigroups. *Mem. Amer. Math. Soc.*, 251(1199):viii+191, 2018.
- [3] L. Cantier. A unitary Cuntz semigroup for C*-algebras of stable rank one. J. Funct. Anal., 281(9):109175, 2021.
- [4] L. Cantier. Uniformly based Cuntz semigroups and approximate intertwinings. *Int. J. Math.*, 33(9):Paper No. 2250062, 36, 2022.
- [5] L. Cantier. The unitary Cuntz semigroup on the classification of non-simple C*-algebras. J. Math. Anal. Appl., 522(2):127003, 2023.
- [6] L. Cantier. Towards a classification of unitary elements of C*-algebras. Preprint. arXiv:2304.01761, 2023.
- [7] L. Cantier. A systematic approach for invariants of C*-algebras. Studia Math., 273, pp. 63-99. 2023.
- [8] L. Cantier and E.Vilalta. Fraissé theory for Cuntz semigroups. J. Algebra 658, pp. 319-364, 2024.
- [9] J. R. Carrión, J. Gabe, C. Schafhauser, A. Tikuisis and S. White Classifying *-homomorphisms I: Unital simple nuclear C*-algebras Preprint. arXiv:2307.06480, 2023.
- [10] A. Ciuperca and G. A. Elliott. A remark on invariants for C*-algebras of stable rank one. *Int. Math. Res. Not. IMRN*, (5):Art. ID rnm 158, 33, 2008.

- [11] K. T. Coward, G. A. Elliott, and C. Ivanescu. The Cuntz semigroup as an invariant for C*-algebras. J. Reine Angew. Math., 623:161–193, 2008.
- [12] J. Cuntz. Dimension functions on simple C*-algebras. Math. Ann., 233(2):145–153, 1978.
- [13] P. de la Harpe and G. Skandalis. Determinant associe a une trace sur une algèbre de Banach. Ann. Inst. Fourier (Grenoble) 34, no. 1, pp. 241-260 (French, with English summary), 1984.
- [14] G. A. Elliott, G. Gong, H. Lin and Z. Niu. On the classification of simple amenable C*-algebras with finite decomposition rank, II. J. Noncommut. Geom.. In press. 10.4171/JNCG/560, 2024.
- [15] G. A. Elliott. On the classification of C*-algebras of real rank zero. J. Reine Angew. Math., 443, pp. 179-219, 1993.
- [16] G. A. Elliott and G. Gong. On the classification of C*-algebras of real rank zero, II. Ann. of Math., 144, no. 3., pp. 497-610, 1996.
- [17] G. A. Elliott, L. Robert, and L. Santiago. The cone of lower semicontinuous traces on a C*-algebra. Amer. J. Math., 133(4):969–1005, 2011.
- [18] E. Gardella and F. Perera. The modern theory of Cuntz semigroups of C*-algebras. Preprint. arXiv:2212.02290, 2022.
- [19] G. Gong, C. Jiang and L. Li. Hausdorffified algebraic K_1 -group and invariants for C*- algebras with the ideal property. *Ann. K-Theory*, vol. 5 no. 1, pp. 43-78, 2020.
- [20] G. Gong, H. Lin, and Z. Niu. A classification of finite simple amenable Z-stable C*-algebras. I: C*-algebras with generalized tracial rank one. C. R. Math. Acad. Sci. Soc. R. Canada, 42, pp. 63-450, 2020.
- [21] G. Gong, H. Lin and Z. Niu. A classification of finite simple amenable Z-stable C*-algebras, II: C*-algebras with rational generalized tracial rank one. C. R. Math. Acad. Sci. Soc. R. Canada, 42, pp. 451-539, 2020.
- [22] K. E. Nielsen and K. Thomsen. Limits of circle algebras. Expo. Math., 14, pp. 17-56, 1996.
- [23] L. Robert. Classification of inductive limits of 1-dimensional NCCW complexes. Adv. Math., 231(5):2802–2836, 2012.
- [24] L. Robert and L. Santiago. Classification of C^* -homomorphisms from $C_0(0,1]$ to a C^* -algebra. J. Funct. Anal., 258(3):869–892, 2010.
- [25] M. Rordam and W. Winter. The Jiang-Su algebra revisited. J. Reine Angew. Math. 642, 129-155, 2010.
- [26] K. Thomsen Traces, unitary characters and crossed products by Z. Publ. Res. Inst. Math. Sci., 31(6), pp. 1011-1029, 1995.
- [27] W. Winter. Structure of nuclear C*-algebras: from quasidiagonality to classification and back again. *Proceedings of the International Congress of Mathematicians-Rio de Janeiro*, Vol. III. Invited lectures, pp 1801-1823, 2018.

LAURENT CANTIER,

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BELLATERRA, SPAIN Email address: laurent.cantier@uab.cat

 URL : www.laurentcantier.fr