

The entropic optimal (self-)transport problem:

Limit distributions for decreasing regularization with application to score function estimation

Gilles Mordant*

IMS, Universität Göttingen

Abstract: We study the statistical properties of the entropic optimal (self) transport problem for smooth probability measures. We provide an accurate description of the limit distribution for entropic (self-)potentials and plans for *shrinking regularization parameter*, which strongly contrasts prior work where the regularization parameter is held fix. Additionally, we show that a rescaling of the barycentric projection of the empirical entropic optimal self-transport plans converges to the score function, a central object for diffusion models, and characterize the asymptotic fluctuations both pointwise and in L^2 .

Finally, we describe under what conditions the methods used enable to derive (pointwise) limiting distribution results for the empirical entropic optimal transport potentials in the case of two different measures and appropriately chosen shrinking regularization parameter. This endeavour requires better understanding the composition of Sinkhorn operators, a result of independent interest.

1. Introduction

The main purpose of this work is to understand statistical properties of the maximizer of the *entropic optimal self-transport problem*

$$\text{OT}_\varepsilon(\rho, \rho) := \sup_{f \in L^1(\rho)} 2 \int f \, d\rho - \varepsilon \iint e^{(f(x)+f(y)-\frac{1}{2}\|x-y\|^2)/\varepsilon} \, d\rho(x) \, d\rho(y) + \varepsilon, \quad (1)$$

where $\varepsilon > 0$ is a regularization parameter. We call the maximizer the *entropic self-potential*, which we denote by f_ε . In practice, Eq. (1) can be efficiently computed on the basis of i.i.d. samples $X_1, \dots, X_n \sim \rho$ by iterating a fixed-point equation; we denote the empirical maximizer by \hat{f}_ε .

Equation (1) is a variant of the entropic optimal transport problem for which both initial and terminal measures are the same (see Section 2). Unlike the optimal transport problem (see Villani (2009); Santambrogio (2015)), it is well-known that $\text{OT}_\varepsilon(\rho, \rho) \neq 0$ for positive regularization. This discrepancy leads to further non-trivial differences at both computational and statistical fronts;

*gilles.mordant@uni-goettingen.de

see for example (Feydy et al., 2019; Genevay et al., 2019; Chizat et al., 2020; González-Sanz et al., 2022; Goldfeld et al., 2022). In these works, the following fluctuations appear (either implicitly or explicitly):

$$D(f_\varepsilon) := \hat{f}_\varepsilon(x) - f_\varepsilon(x), \quad \text{and} \quad D(\nabla f_\varepsilon) := \nabla \hat{f}_\varepsilon(x) - \nabla f_\varepsilon(x).$$

For example, $D(f_\varepsilon)$ arises in estimating the *Sinkhorn divergence*, a debiased version of the entropic optimal transport cost Feydy et al. (2019); Lavenant et al. (2024). Meanwhile, $D(\nabla f_\varepsilon)$ arises when debiasing entropic transport maps Pooladian et al. (2022).

In these aforementioned works, the regularization parameter for entropic optimal transport is kept fixed. This yields statistical results of a parametric flavor, as the potentials are known to lie in a reproducing kernel Hilbert space (Genevay et al., 2019). Though, Pooladian and Niles-Weed (2021) demonstrated that entropic optimal transport can be used to estimate optimal transport maps when the regularization parameter shrinks with the number of samples growing, i.e., $\varepsilon = \varepsilon(n) \rightarrow 0$ as $n \rightarrow \infty$. Still, a concrete understanding of the statistical impact of sample size-dependent regularization has not been carried out before in the continuous setting and the optimal rates of convergence remain elusive.

The self-transport problem has another remarkable feature when taking the regularization parameter to zero: *the gradient of the potential converges to the score function*. Given the importance of the latter in diffusion models, it is interesting per se to understand the properties of a score function estimator arising from the optimal transport theory.

To see why entropic optimal self-transport might give rise to a reasonable estimator of the score function, we turn to the Gaussian case, where exact and explicit calculations can be performed. Let $\rho_A := \mathcal{N}(0, A)$, a centered Gaussian distribution with covariance matrix $A \succ 0$. The closed-form expression for the self-potential $f_{\varepsilon, A}$, i.e., the optimizer of $\text{OT}_\varepsilon(\rho_A, \rho_A)$, is known. See, e.g., Pooladian et al. (2022, Proposition 4). A series expansion in ε precisely yields

$$f_{\varepsilon, A}(x) = -\frac{\varepsilon}{2} \log \rho_A(x) - \frac{\varepsilon^2}{16} x^\top A^{-2} x + x^\top (\text{O}(\varepsilon^4 I)) x, \quad \text{as } \varepsilon \rightarrow 0. \quad (2)$$

Taking gradients with respect to x and re-arranging, we see that

$$-\frac{2}{\varepsilon} \nabla f_{\varepsilon, A}(x) = \nabla \log \rho_A(x) + \frac{\varepsilon}{8} A^{-2} x + \text{O}(\varepsilon^3 I) x, \quad \text{as } \varepsilon \rightarrow 0.$$

The first term in the right-hand side is the announced score function.

In light of this calculation, slightly anticipating the subsequent developments of the paper, we can define an estimator of the score function $\nabla \log \rho$ as

$$\hat{s}_\varepsilon(x) := -\frac{2}{\varepsilon} \nabla \hat{f}_\varepsilon(x) = \frac{2}{\varepsilon} \left(x - \frac{\sum_{i=1}^n X_i e^{\hat{f}_\varepsilon(X_i)/\varepsilon} e^{-\frac{1}{2\varepsilon} \|x - X_i\|^2}}{\sum_{j=1}^n e^{\hat{f}_\varepsilon(X_j)/\varepsilon} e^{-\frac{1}{2\varepsilon} \|x - X_j\|^2}} \right), \quad (3)$$

where X_1, \dots, X_n is an i.i.d. sample as above. The second equality is the discrete counterpart to Equation (9) below.

Our estimator can thus be viewed as a *weighted kernel density estimator*, where the regularization parameter ε takes the role of the bandwidth parameter. Indeed, one can check that if the estimated density has the form $\hat{\rho}(x) \propto \sum_{i=1}^n K((x - X_i)/\sqrt{\varepsilon})$, with $K(\cdot)$ being the Gaussian kernel, then

$$\nabla \log \hat{\rho}(x) = \frac{1}{\varepsilon} \left(x - \frac{\sum_{i=1}^n X_i e^{-\frac{1}{2\varepsilon} \|x - X_i\|^2}}{\sum_{j=1}^n e^{-\frac{1}{2\varepsilon} \|x - X_j\|^2}} \right). \quad (4)$$

Up to the factors $e^{\hat{f}_\varepsilon(X_i)/\varepsilon}$ (as well as the multiplicative factor of 2), $\nabla \log \hat{\rho}$ coincides with \hat{s}_ε .

We argue that this estimator of the score function based on optimal transport might be better than kernel density estimation in a variety of settings, in particular under the hypothesis that the distribution concentrates on a manifold of low intrinsic dimension. Simulation results back this view, as Figure 1 shows. This is further in line with the low complexity adaptation principle recently uncovered by [Hundrieser et al. \(2024\)](#). It states that optimal transport adapts to smallest complexity of the class of dual potentials. Roughly sketched, this implies that if *one* measure is supported on a low dimensional manifold, the convergence rate will depend on the intrinsic and not the ambient dimension. Since then, extensions have been developed in [Groppe and Hundrieser \(2023\)](#) or [Stromme \(2023\)](#). As this paper is the first of this kind, we restrict ourselves to the case where the data is fully supported on Euclidean space, further work in the case of the manifold hypothesis will follow.

Finally, the understanding the statistical fluctuations of the self-transport problem we develop provides a framework to understand the fluctuations of general entropy regularised optimal transport problems, which is a major contribution to this very active field.

Contributions

With this background in mind, our three primary contributions are summarized as follows. First, we provide a second-order expansion of the entropic self-potentials. This allows us to achieve tight asymptotic control on $\sqrt{n}D(f_{\varepsilon_n})$ and $\sqrt{n}D(\nabla f_{\varepsilon_n})$ as $n \rightarrow \infty$ with $\varepsilon_n \rightarrow 0$. With these asymptotic results, we are able to derive (pointwise) limiting distribution results for the potentials *for shrinking regularization parameter*, which is new. As a further application, we show that a rescaling of ∇f_ε is a reasonable estimator of the score function of ρ , written $\nabla \log \rho$.

In the second part of the paper, we extend the scope to the statistical aspects of regularized optimal transport between two empirical measures when the regularization parameter goes to zero at a properly chosen rate. We exhibit the conditions under which limiting distribution results can be obtained and show the link with the linearization of Monge-Ampère approach in [Manole et al. \(2023\)](#).

In short, in this paper, we

- show a linearization result for the fluctuations of the potentials and maps of self EOT even for decreasing epsilon;
- derive limiting distribution results and asymptotic rates;
- pave the way towards limiting distribution results for the two measure case, finely understanding the Sinkhorn operators when the regularization parameter goes to zero and linking such operator to elliptic differential operators.

Related work

The entropic optimal (self-)transport problem. Entropic optimal transport has received much attention in recent years due to its computational tractability in large-scale tasks. Statistical and theoretical analyses have been performed in tandem. See [Chewi et al. \(2024\)](#) for a thorough overview of recent works. The entropic map was statistically analysed by [Pooladian and Niles-Weed \(2021\)](#); [Pooladian et al. \(2023\)](#) as a computationally tractable estimator of the optimal transport map for the quadratic cost ([Brenier, 1991](#)), following the work of [Hütter and Rigollet \(2021\)](#). This map was subsequently analysed in [Rigollet and Stromme \(2022\)](#) using different techniques.

The entropic optimal self-transport problem has been studied independently by, e.g., [Genevay et al. \(2019\)](#); [Feydy et al. \(2019\)](#) in general-purpose machine learning, and by [Marshall and Coifman \(2019\)](#) in the context of manifold learning. For example, in machine learning, the entropic self-OT problem was used to define the *Sinkhorn divergence*, written

$$S_\varepsilon(\rho, \mu) := \text{OT}_\varepsilon(\rho, \mu) - \frac{1}{2}(\text{OT}_\varepsilon(\rho, \rho) + \text{OT}_\varepsilon(\mu, \mu)),$$

which leads to $S_\varepsilon(\rho, \mu) \geq 0$ with equality if and only if $\rho = \mu$ ([Feydy et al., 2019](#), Theorem 1). In a sense, $S_\varepsilon(\rho, \mu)$ is a *debiased* version of $\text{OT}_\varepsilon(\rho, \mu)$. Statistical and algorithmic implications of the Sinkhorn divergence were considered in [Feydy et al. \(2019\)](#); [Pooladian et al. \(2022\)](#), among others. [Marshall and Coifman \(2019\)](#) proposed to use self-transport to perform manifold learning, which was further extended to the empirical case by [Landa et al. \(2021\)](#). In a similar vein, the work of [Landa and Cheng \(2023\)](#) used self-transport as a robust tool for density estimation.

Finally, we note that the Kullback–Leibler regularization is not the only interesting one. Optimal transport with L^2 regularization has been considered by ([Blondel et al., 2018](#)) where they noticed that it gives sparse transport plans. Building up on that fact, [Zhang et al. \(2023\)](#) proposed to use to carry out manifold learning relying on L^2 regularized self-transport; the approach is robust to heteroskedastic noise in high-dimensional spaces.

The estimation of the score functions. For a probability measure $\rho \propto e^{-V}$, the estimation of score functions has attracted much less attention than

the estimation of densities or gradient of densities, which is a standard statistical topic (Silverman, 2018; Tsybakov, 2009). A procedure to estimate of log-concave densities is explained in the recent review paper by Samworth (2018). The methodology therein relies on the so-called log-concave projection; an optimisation problem that is quite intricate to solve in higher dimensions. Even though minimax rates are known in that case, they only apply to the estimation of the density and not its gradient. Seregin and Wellner (2010) obtain $n^{-2/(d+4)}$ as pointwise minimax rate with respect to the absolute loss function at a point of sufficient regularity. Still, the global picture is far from complete. Another view on the problem could be that of estimation of the gradient of a shape constrained function. To the best of our knowledge, that approach hasn't been pursued in general dimensions.

An important line of research relies on the idea of score matching put forth by Hyvärinen and Dayan (2005) in which the estimated score function is taken to be an element of a parametric class. Sriperumbudur et al. (2017) then extended it to a reproducing kernel Hilbert space (RKHS) framework. An extension was proposed in Zhou et al. (2020). More recently, Song et al. (2020) proposed a sliced version of score matching.

While completing this work, we became aware of the paper by Wibisono et al. (2024), where they propose an estimator of the score based on (truncated) kernel density estimation (KDE). By deriving suitable lower bounds, they show that their estimator is minimax optimal up to logarithmic factors, though our analyses and estimator are substantially different.

Limiting distribution results in optimal transport and variants. A plethora of work has studied distribution limits and the convergence rates in variations of the optimal transport problem. Listing in detail the substance of these contributions does not align with the goal of keeping this paper concise; we warmly invite the reader to read the following references and the work they cite. For the cost, see Sommerfeld and Munk (2018); del Barrio and Loubes (2019); Hundrieser et al. (2022); del Barrio et al. (2023); Hundrieser et al. (2023); Goldfeld et al. (2024a,b). For the plan, Klatt et al. (2022); Manole et al. (2023); González-Sanz et al. (2022); Goldfeld et al. (2024a) are relevant references. Note that the understanding of the limit laws provided by González-Sanz et al. (2022) is fundamental and is the sole reference providing a clear, detailed description of the asymptotic fluctuations in terms of Sinkhorn operators. Finally, a completely different approach to estimate the transport map, based on a sum-of-squares reformulation of the semi-dual formulation, is provided in Vacher et al. (2024). Their method is both computationally attractive and nearly achieves the minimax rate.

In all that we do, the only cost of interest is the square quadratic cost, as the theory is particularly neat in this case. Just like in Pal (2024) the approach might work for more general costs provided that the divergence based on the optimal transport potentials is sufficiently well behaved. The interest for more general costs and even estimated ones is clearly important for recent developments in OT, see Grave et al. (2019); Pooladian et al. (2024); Hundrieser et al. (2024).

Notation

In the sequel, all random variables will be defined on a common, sufficiently rich probability space $(\Omega, \mathcal{A}, \mathbb{P})$. We will slightly abuse notation and use both ρ as a probability measure and to mean its density $d\rho/d\lambda$, where λ is the Lebesgue measure. For a probability measure ρ , we write ρ_n as the empirical measure of ρ based on n i.i.d. samples. Convergence in distribution is denoted by \rightsquigarrow . As usual, $a \lesssim b$, means that there exists a constant $C > 0$ such that $a \leq Cb$. We sometimes use $a_n \asymp b_n$ when $a_n = b_n + o(1)$ as $n \rightarrow \infty$. We set $\mathbb{G}_n = \sqrt{n}(\rho_n - \rho)$. For a function $f \in L^1(\rho)$, we use $\rho f := \int f d\rho$. The Fenchel–Legendre of a function $\varphi(y)$ defined on a domain \mathcal{Y} is $\varphi^*(x) := \sup_{y \in \mathcal{Y}} (\langle x, y \rangle - \varphi(y))$. The set of k times continuously differentiable functions is denoted \mathcal{C}^k . The Hessian of a function f at a point x will be denoted $\nabla^2 f(x)$. Finally, we set

$$k_\varepsilon(x, y) := \exp\left(-\frac{1}{2\varepsilon}\|x - y\|^2\right).$$

2. Background on entropic optimal transport

For two Borel probability measures with finite second moment μ, ν on \mathbb{R}^d , and regularization parameter $\varepsilon > 0$, the *entropic optimal transport* problem is given by

$$\text{OT}_\varepsilon(\rho, \mu) := \min_{\pi \in \Gamma(\mu, \nu)} \iint \frac{1}{2}\|x - y\|^2 d\pi(x, y) + \varepsilon \text{KL}(\pi \| \mu \otimes \nu), \quad (5)$$

where $\Gamma(\mu, \nu)$ is the set of joint measures with marginals ρ and μ , and

$$\text{KL}(\pi \| \mu \otimes \nu) = \iint \log\left(\frac{d\pi}{d(\mu \otimes \nu)}\right) d\pi,$$

whenever π has a density with respect to the product measure $\mu \otimes \nu$, and is $+\infty$ otherwise. Due to the strict convexity of the objective, a unique minimizer, the *entropic optimal (transport) plan*, written π_ε , always exists and is unique.

Entropic regularization was first introduced as an algorithmic tool to approximate the *unregularized* optimal transport cost, i.e., when $\varepsilon = 0$ (Cuturi, 2013; Peyré and Cuturi, 2019). We refer the reader to Villani (2009); Santambrogio (2015); Villani (2021) for in-depth introduction to the unregularized case.

2.1. Dual formulation and barycentric projections

The primal problem (5) admits a dual formulation which is extremely useful, both from an analytical and computational perspective. The dual formulation reads (see e.g., Genevay (2019))

$$\begin{aligned} \text{OT}_\varepsilon(\rho, \mu) = \\ \sup_{(f, g) \in L^1(\mu \otimes \nu)} \int f d\mu + \int g d\nu - \varepsilon \iint e^{(f(x) + g(y) - \frac{1}{2}\|x - y\|^2)/\varepsilon} d\mu(x) d\nu(y) + \varepsilon. \end{aligned} \quad (6)$$

We call the maximizers to Eq. (6) the *entropic optimal dual potentials*, written $(f_\varepsilon, g_\varepsilon)$. Moreover, the entropic optimal plan can be readily expressed as a function of the entropic optimal potentials through a primal-dual recovery relationship (Csiszár, 1975), i.e.,

$$d\pi_\varepsilon(x, y) = \exp\left(\frac{f_\varepsilon(x) + g_\varepsilon(y) - \frac{1}{2}\|x - y\|^2}{\varepsilon}\right) d\mu(x) d\nu(y),$$

which implies that, at optimality,

$$\text{OT}_\varepsilon(\mu, \nu) = \int f_\varepsilon d\mu + \int g_\varepsilon d\nu.$$

Moreover, the entropic optimal dual potentials are known to satisfy the following relationships, see Mena and Niles-Weed (2019); Nutz and Wiesel (2021).

$$\begin{aligned} \int e^{(f_\varepsilon(x) + g_\varepsilon(y) - \frac{1}{2}\|x - y\|^2)/\varepsilon} d\mu(x) &= 1, \quad \forall y \in \mathbb{R}^d, \\ \int e^{(f_\varepsilon(x) + g_\varepsilon(y) - \frac{1}{2}\|x - y\|^2)/\varepsilon} d\nu(y) &= 1, \quad \forall x \in \mathbb{R}^d. \end{aligned} \quad (7)$$

These relations are crucial for the rest of our work.

The *entropic map* between μ and ν is simply the barycentric projection of π_ε , which, for $x \in \mathbb{R}^d$, is given by

$$T_\varepsilon(x) := \int y d\pi_\varepsilon^x(y) = \mathbb{E}_{\pi_\varepsilon}[Y \mid X = x]. \quad (8)$$

Alternatively, by Eq. (7),

$$T_\varepsilon(x) := \frac{\int y e^{\frac{1}{\varepsilon}(g_\varepsilon(y) - \frac{1}{2}\|x - y\|^2)} d\nu(y)}{\int e^{\frac{1}{\varepsilon}(g_\varepsilon(y) - \frac{1}{2}\|x - y\|^2)} d\nu(y)}. \quad (9)$$

It can also be verified from these optimality conditions that $T_\varepsilon = \text{id} - \nabla f_\varepsilon$ as in Pooladian and Niles-Weed (2021, Proposition 2).

2.2. The entropic optimal self-transport problem

An important fact about entropic optimal transport is that it is inherently biased. That is to say, for a fixed probability measure ρ , the entropic optimal self-transport problem yields $\text{OT}_\varepsilon(\rho, \rho) \neq 0$; see Feydy et al. (2019); Genevay (2019); Marshall and Coifman (2019).

Viewed as a special case of both Eq. (5) and Eq. (6), we see that the following change of variables

$$u_\varepsilon(x) := \exp(-f_\varepsilon(x)/\varepsilon), \quad (10)$$

allows the the system (7) to become a single fixed point equation (with $g_\varepsilon \equiv f_\varepsilon$)

$$u_\varepsilon(x) = \int \frac{1}{u_\varepsilon(y)} \exp\left(-\frac{1}{2\varepsilon}\|x - y\|^2\right) d\rho(y). \quad (11)$$

3. Main results for self-transport

We will show that the entropic optimal self-transport problem can be used to estimate *score functions* of probability measure i.e., the gradient of the logarithm of the density. Score functions arise in the Langevin diffusion, and the widely used Langevin Monte Carlo algorithm for sampling from a distribution ρ ; see [Chewi \(2023\)](#) for a recent survey. Langevin Diffusion is also at the core of denoising diffusion models, see [Peyré \(2023\)](#) and the reference therein.

Throughout this work, we will require the following assumption on ρ .

(Supp) The density ρ is supported on a compact set K and C^2 on its domain. Furthermore, $\forall x \in K$, we have that $\ell \leq \rho(x) \leq L$, for $0 < \ell \leq L$.

Assumption **(Supp)** is a standard, it is made in a large variety of reference works such as in [Hütter and Rigollet \(2021\)](#); [Pooladian and Niles-Weed \(2021\)](#); [Manole et al. \(2024\)](#), to cite but a few.

3.1. Second-order expansion of the potentials and score functions

The expansion in the Gaussian case mentioned in the introduction naturally asks the question whether such a development holds for other cases. In that vein, we prove the following proposition, which is inspired by the work of [Marshall and Coifman \(2019\)](#). To do so, we need the following regularity assumption.

(Tay) The density ρ is such that there exists $\varepsilon_0 > 0$ and a constant C such that for all $\varepsilon \leq \varepsilon_0$ and all $x, z \in \text{int}(\text{supp}(\rho))$,

$$\left| \frac{u_\varepsilon(x + \sqrt{\varepsilon}z)}{u_\varepsilon(x)} - 1 - \sqrt{\varepsilon} \nabla \log(u_\varepsilon(x))^\top z - \frac{\varepsilon z^\top \nabla^2 u_\varepsilon(x) z}{2u_\varepsilon(x)} \right| \leq \varepsilon^{3/2} C \|z\|^2.$$

Assumption **(Tay)** is requiring a second order Taylor expansion of the potentials with a controlled remainder that is uniform in ε . It is a technical assumption that only serves in the proof of [Proposition 1](#). Furthermore, the validity of the expansion in [Proposition 1](#) can also be deduced by other means. In [Remark 4](#), we provide a complimentary view based on Fourier analysis. In both cases, the unknown regularity of the potentials intervene, yet in slightly different ways. Both view complement each other.

Proposition 1 (Second-order expansion). Let f_ε be the optimal entropic self-potential for ρ satisfying **(Supp)**-**(Tay)**. For any fixed point $x \in \text{int}(\text{supp}(\rho))$, as $\varepsilon \rightarrow 0$, the following holds:

$$\begin{aligned} & \exp(-2f_\varepsilon(x)/\varepsilon) \\ &= \rho(x)(2\pi\varepsilon)^{d/2} \left(1 + \varepsilon \text{tr} \left(\frac{\nabla^2 \rho(x)}{4\rho(x)} - \frac{1}{8} (\nabla \log \rho(x)) (\nabla \log \rho(x))^\top \right) + o(\varepsilon) \right). \end{aligned} \tag{12}$$

The proof is provided in the [Appendix B.1](#).

Remark 2 (Leading bias and harmonic characteristic). As noted in [Agarwal et al. \(2024\)](#), the term of order ε in the expansion above has an interpretation. It is called the harmonic characteristic and can be interpreted as a “mean acceleration” of a Langevin bridge. We refer to the aforementioned reference for more background.

(Reg) For the rest of the paper, we will make the slightly stronger assumption that there exists constant C_{f, ε_0} such that

$$\left| f_\varepsilon(x) + \frac{\varepsilon}{2} \log \rho(x) + \frac{\varepsilon d}{4} \log(2\pi\varepsilon) \right| \leq \varepsilon^2 C_f \|x\|^2,$$

for every $\varepsilon \leq \varepsilon_0$ uniformly in $x \in \text{supp}(\rho)$.

Remark 3 (On assumption **(Reg)**). Note that this assumption holds when ρ is a Gaussian density. Further, for C^6 distributions supported on compact manifold of dimension smaller than 5, a similar expansion holds, see [Landa and Cheng \(2023, Theorem 2.2, Eq. \(11\)\)](#). In a slightly different setting, the same type of expansion has also been obtained in [Deb et al. \(2023\)](#) also relying on a Laplace Method. This is thus a mild assumption.

Remark 4 (Another view on the fix point equation). Up to shifting the cost function by $\varepsilon d \log(2\pi\varepsilon)$, we can rewrite (11) as

$$u_\varepsilon = \gamma_\varepsilon * \left(\frac{\rho}{u} \right),$$

with γ_ε the isotropic centred Gaussian density with variance ε . Assuming that the Fourier transform of both sides is well defined, expanding in ε the Fourier transform of γ_ε , we get

$$\mathcal{F}(u_\varepsilon)[\xi] = \mathcal{F}\left(\frac{\rho}{u_\varepsilon}\right)[\xi] \left(1 - \frac{\varepsilon}{2} \|\xi\|^2 + o(\varepsilon)\right).$$

Setting $u_\varepsilon = u_0 + \varepsilon u_1 + o(\varepsilon)$, a further expansion gives

$$\mathcal{F}(u_0)[\xi] + \varepsilon \mathcal{F}(u_1)[\xi] = \mathcal{F}\left(\frac{\rho}{u_0}\right)[\xi] - \varepsilon \mathcal{F}\left(\frac{\rho u_1}{u_0^2}\right)[\xi] - \frac{\varepsilon}{2} \|\xi\|^2 \mathcal{F}\left(\frac{\rho}{u_0}\right)[\xi] + o(\varepsilon).$$

At the constant scale, $\mathcal{F}(u_0)[\xi] = \mathcal{F}(\rho/u_0)[\xi]$ enforces $u_0 = \sqrt{\rho}$. At the ε scale, one thus gets the equation

$$\mathcal{F}(u_1)[\xi] = -\frac{1}{4} \|\xi\|^2 \mathcal{F}(\sqrt{\rho})[\xi],$$

which is characteristic for elliptic differential equations, as expected from the trace term in (12). Note that this view on the problem only requires the Fourier transform of the various quantities of interest to be defined, which is quite mild.

3.2. Statistical fluctuations via linearization

The main result relies on Assumption **(Conv)**. Even though this assumption might appear strong at first reading, it is not, as we discuss it in detail in Section 3.4. Remark also that this assumption is only sufficient, it is however much more transparent than any other weaker assumption the proof suggests.

(Conv) Assume that, taking $n \rightarrow \infty$, $\varepsilon_n \rightarrow 0$ and $\sqrt{n}\varepsilon_n^{d/4}/\sqrt{\log n} \rightarrow \infty$,

$$\frac{1}{\varepsilon_n} \|\hat{f}_{\varepsilon_n} - f_{\varepsilon_n}\|_{\infty} \rightarrow 0,$$

in probability.

To better understand this assumption, by the discussion of the previous section, $f_{\varepsilon_n}/\varepsilon_n$ is converging. Assumption **(Conv)** thus only requires convergence of an empirical quantity to a population quantity for “bandwidth size” decreasing slowly, no rate of convergence is assume here. Section 3.4 provides a much longer discussion on why such an assumption is reasonable.

For the statistical fluctuations, we need to introduce the following operator

$$h \mapsto K_{\varepsilon}[h](x) := \int h(y)\pi_{\varepsilon}(x, y)d\rho(y),$$

which is compact from $L^2(\rho) \rightarrow L^2(\rho)$ for any $\varepsilon > 0$. Indeed, this is a Markov operator with invariant distribution ρ , by construction of the coupling. With these elements at hand, we can state the two main linearization results.

Proposition 5 (Fluctuations of entropic self-potentials). Let f_{ε_n} be the entropic self-potential for ρ satisfying **(Supp)** and **(Reg)**, and let \hat{f}_{ε_n} be its empirical counterpart based on n i.i.d. samples from the distribution ρ . Assume **(Conv)** and fix $x \in \text{int}(\text{supp}(\rho))$. Choosing $\sqrt{n}\varepsilon_n^{d/4}/\sqrt{\log n} \rightarrow \infty$, as $n \rightarrow \infty$, there exists a sequence a_n with $a_n\varepsilon_n \rightarrow \infty$, as $n \rightarrow \infty$, such that

$$\begin{aligned} & a_n(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x)) + a_n K_{\varepsilon_n}[(\hat{f}_{\varepsilon_n} - f_{\varepsilon_n})(1 + o(1))](x) \\ &= -\frac{a_n\varepsilon_n}{\sqrt{n}} \int \frac{\exp\left(-\frac{1}{2\varepsilon_n}\|y-x\|^2\right)}{(2\pi\varepsilon_n)^{d/2}\rho^{1/2}(y)\rho^{1/2}(x)(1+o(1))} d\mathbb{G}_n(y) + o_p(1). \end{aligned}$$

Proposition 6 (Fluctuations of gradients of entropic self-potentials). Let us consider the exact same setting as in Proposition 5. Then, for $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$ but decaying slower than $n^{-\frac{1}{d-3}}$, it holds that

$$\begin{aligned} & a_n\varepsilon_n \left(\nabla f_{\varepsilon_n}(x) - \nabla \hat{f}_{\varepsilon_n}(x) \right) \\ &= -\frac{a_n\varepsilon_n}{\sqrt{n}} \int y \frac{\exp\left(-\frac{1}{2\varepsilon_n}\|y-x\|^2\right)}{(2\pi\varepsilon_n)^{d/2}\rho^{1/2}(y)\rho^{1/2}(x)(1+o(1))} d\mathbb{G}_n(y) \\ & - a_n \left(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x) \right) K_{\varepsilon_n}[\text{id}](x) - K_{\varepsilon_n} \left[\text{id} \times a_n \left(\hat{f}_{\varepsilon_n}(\cdot) - f_{\varepsilon_n}(\cdot) \right) \right] (x) \\ & + o_p(1). \end{aligned}$$

The proofs of both of these results are deferred to Appendix B. The two propositions above are fundamental to the understanding of the fluctuations of the potentials and plans.

3.3. Application 1: Limit distributions

As a first application, we prove the following (pointwise) limit distribution for the entropic self-potentials. This result is similar in flavor to those present in e.g., González-Sanz et al. (2022) and Goldfeld et al. (2022), though with a key differences: their results are restricted to the case where $\varepsilon > 0$ is fixed, whereas we permit $\varepsilon_n \rightarrow 0$. The proofs are deferred to Appendix B.4.

Theorem 7 (Limit distribution for the empirical potentials). *Consider an i.i.d. sample $X_1, \dots, X_n \sim \rho$ as above and suppose (Supp) and (Reg) and (Conv) hold. Then, for any $x_1, \dots, x_m \in \text{int}(\text{supp}(\rho))$ with $m \in \mathbb{N}$ fixed,*

$$\sqrt{n} \varepsilon_n^{-1+d/4} \begin{pmatrix} (\hat{f}_{\varepsilon_n}(x_1) - f_{\varepsilon_n}(x_1)) \\ \vdots \\ (\hat{f}_{\varepsilon_n}(x_m) - f_{\varepsilon_n}(x_m)) \end{pmatrix} \rightsquigarrow \mathcal{N} \left(0_m, C_3 \text{diag} \left(\rho(x_1)^{-1}, \dots, \rho(x_m)^{-1} \right) \right),$$

as $n \rightarrow \infty$ provided that $\sqrt{n} \varepsilon_n^{d/4} / \sqrt{\log n} \rightarrow \infty$, where

$$C_3 := \sum_{0 \leq \kappa, \kappa' \leq \infty} 2^{-\kappa - \kappa'} \sum_{\eta=0}^{\kappa} \sum_{\eta'=0}^{\kappa'} \binom{\kappa}{\eta} (-1)^\eta (-1)^{\eta'} \binom{\kappa'}{\eta'} (\eta + \eta' + 2)^{-d/2}.$$

Note that C_3 is related to binomial transform and Roman harmonic numbers Sesma (2017). We deem the full simplification of this quantity beyond the scope of the current paper. However, a first simplification for d even is the content of the following remark.

Remark 8. For $d/2 \in \mathbb{N}$, using the equality (Kim et al., 2023, Lemma 2.1)

$$\int_0^1 (1-t)^{\kappa'} (\log t)^{d/2} t^{\eta+1} dt = (d/2 - 1)! \sum_{\eta'=0}^{\kappa'} \binom{\kappa'}{\eta'} (-1)^{\eta' - (d/2 - 1)},$$

we can simplify

$$\begin{aligned} \sum_{\eta=0}^{\kappa} \sum_{\eta'=0}^{\kappa'} \binom{\kappa}{\eta} (-1)^\eta (-1)^{\eta'} \binom{\kappa'}{\eta'} (\eta + \eta' + 2)^{-d/2} \\ = \frac{(-1)^{1-d/2}}{(d/2 - 1)!} \int_0^1 (1-t)^{\kappa + \kappa'} t (\log t)^{d/2 - 1} dt, \end{aligned}$$

for $\kappa, \kappa' > 0$.

Theorem 9. Consider the setting of Theorem 7 again. Then, for any fixed $x \in \text{int}(\text{supp}(\rho))$ provided that $\sqrt{n} \varepsilon_n^{d/4} / \sqrt{\log n} \rightarrow \infty$, as $n \rightarrow \infty$,

$$\sqrt{n} \varepsilon_n^{d/4} \left(\nabla \hat{f}_{\varepsilon_n}(x) - \nabla f_{\varepsilon_n}(x) \right) \rightsquigarrow \mathcal{N}(0_d, \Sigma_d(x)),$$

for some covariance matrix $\Sigma_d(x)$.

Corollary 10. Consider the setting of Theorem 9 and fix $x \in \text{int}(\text{supp}(\rho))$. Then,

$$\sqrt{n} \varepsilon_n^{d/4+1} \left(\frac{2\nabla \hat{f}_{\varepsilon_n}(x)}{\varepsilon_n} - \nabla \log \rho(x) \right) \rightsquigarrow \mathcal{N}(0_d, \Sigma_d(x)),$$

for $\varepsilon_n = o\left(n^{-\frac{2}{d+8}}\right)$.

In this subsection, we exhibit rates of convergence in the $L^2(\rho)$ metric, which is the one of interest when it comes to studying the convergence diffusion schemes. To continue, we require the following additional assumptions. By analogy with the classical notation used for log-concave measures, let us introduce the notation $V(x) := -\log \rho(x)$, $x \in \text{supp}(\rho)$.

(Reg') There exists a function $R(x) \in \mathcal{C}^1(\text{supp}(\rho))$ such that, for $x \in \text{int}(\text{supp}(\rho))$

$$\left| -2f_\varepsilon(x)/\varepsilon - \log \rho(x) - \frac{d}{2} \log(2\pi\varepsilon) - \varepsilon \text{tr} \left(\frac{\nabla^2 \rho(x)}{4\rho(x)} - \frac{1}{8} (\nabla \log \rho(x)) (\nabla \log \rho(x))^\top \right) \right| = \varepsilon^2 R(x).$$

(C) The Hessian of V is Lipschitz, i.e., there exists $M > 0$ such that

$$\|\nabla^2 V(x) - \nabla^2 V(y)\| \leq M \|x - y\|.$$

Furthermore, $\nabla^2 V \preceq \beta I_d$.

Theorem 11 ($L^2(\rho)$ convergence rates). Suppose **(Supp)**, **(Reg')**, **(Conv)** and **(C)** hold. Further assume that the support is bounded. Setting $\varepsilon_n \asymp n^{-1/(d+4)}$, the estimator Eq. (3) achieves the following rate of estimation

$$\mathbb{E} \|\hat{s}_\varepsilon - \nabla \log \rho\|_{L^2(\rho)}^2 \lesssim n^{-2/(d+4)}. \quad (13)$$

Remark 12. Our proofs, and thus our results, are asymptotic. Yet, they can be converted into finite sample bounds at the price of multiplying by sufficiently large (but universal) constants and taking n larger than some $n_0 \in \mathbb{N}$.

The proof of Theorem 11 follows a traditional bias-variance trade-off arising from the inequality

$$\mathbb{E} \|\hat{s}_\varepsilon - \nabla \log \rho\|_{L^2(\rho)}^2 \leq 2\mathbb{E} \|\hat{s}_\varepsilon - s_\varepsilon\|_{L^2(\rho)}^2 + 2\|s_\varepsilon - \nabla \log \rho\|_{L^2(\rho)}^2, \quad (14)$$

where s_ε is the population counterpart to our estimator. The proof follows directly from the following two propositions. Proposition 13 makes Eq. (2) rigorous

in the non-Gaussian case, which follows from the precise asymptotic expansion given in Proposition 1. We then control the stochastic fluctuations in Proposition 6, from which a trade-off in ε_n yields the final result; we omit this final step.

Proposition 13 (Population bias). Under **(Reg')** and **(C)** it holds that,

$$\|s_\varepsilon - \nabla \log \rho\|_{L^2(\rho)}^2 \lesssim \varepsilon^2 (M^2 d + \beta^2 I_0(\rho)),$$

where $I_0(\rho)$ is the Fisher information of ρ , and we otherwise omit universal constants.

Proposition 14 (Statistical fluctuations). Consider a sequence $(\varepsilon_n)_{n \geq 1}$ with $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$. Under **(Supp)** and a compact support assumption, it holds that

$$\mathbb{E} \|\hat{s}_{\varepsilon_n} - s_{\varepsilon_n}\|_{L^2(\rho)}^2 \lesssim n^{-1} \varepsilon_n^{-d-2}.$$

3.4. About the uniform convergence hypothesis

In **(Conv)**, we assume that $\varepsilon_n^{-1} \|\hat{f}_{\varepsilon_n} - f_{\varepsilon_n}\|_\infty$ converges to zero in probability. We will now explain why this assumption is quite mild. The difference between the actual potential and the first term of the expansion, $f(x)/\varepsilon_n + \log(\rho(x))/2$, is well understood thanks to Proposition 1 or Remark 4. Therefore, we will focus on the empirical part. Recall the fact that the empirical potential satisfies the fixed point equation

$$\frac{\hat{f}(x)}{\varepsilon_n} = -\log \frac{1}{n} \sum_{i=1}^n \frac{\exp(-\frac{\|x-X_i\|^2}{2\varepsilon_n})}{(2\pi\varepsilon_n)^{d/2}} \exp\left(\frac{\hat{f}(X_i)}{\varepsilon_n}\right).$$

Now, let us replace \hat{f}/ε_n by $-\log(\rho)/2$ in the identity above. This raises the question whether

$$\rho^{1/2}(x) \text{ is approximately } \frac{1}{n} \sum_{i=1}^n \frac{\exp(-\frac{\|x-X_i\|^2}{2\varepsilon_n})}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(X_i)},$$

which we now establish. First, observe that

$$\mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n \frac{\exp(-\frac{\|x-X_i\|^2}{\varepsilon_n})}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(X_i)} \right] = \rho^{1/2}(x)(1 + o(1)), \text{ as } \varepsilon_n \rightarrow 0.$$

The above result holds true with a uniform control on the remainder for compactly supported distributions. The form of the remainder in the Gaussian case can even be made explicit, as it requires computing a Gaussian convolution. Now, to claim that the *ansatz* we make is reasonable, we have to show that

$$x \mapsto \frac{1}{n} \sum_{i=1}^n \frac{\exp(-\frac{\|x-X_i\|^2}{\varepsilon_n})}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(X_i)}$$

concentrates around its expectation in the L^∞ norm. The classical proof of convergence of kernel density estimators in the uniform norm relies on the fact that the class of functions

$$\mathcal{F}_0 := \left\{ x \mapsto \exp\left(-\frac{\|x-y\|^2}{h}\right); h > 0, y \in \mathbb{R}^d \right\}$$

is a VC class. Therefore, it is clear that for any measure Q , and $f, f' \in \mathcal{F}_0$,

$$\mathbb{E}_{Y \sim Q} \left[\left(\frac{f(Y)}{\rho^{1/2}(Y)} - \frac{f'(Y)}{\rho^{1/2}(Y)} \right)^2 \right] \leq \ell^{-1} \mathbb{E}_{Y \sim Q} \left[(f(Y) - f'(Y))^2 \right],$$

that is, dividing by the square root of the density does not change the complexity of the function class. Because of this, we can rely on the same arguments as in [Einmahl and Mason \(2005\)](#) to conclude that

$$\left\| \frac{1}{n} \sum_{i=1}^n \frac{\exp(-\frac{\|x-X_i\|^2}{\varepsilon_n})}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(X_i)} - \mathbb{E} \left[\frac{\exp(-\frac{\|x-X\|^2}{\varepsilon_n})}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(X)} \right] \right\|_\infty \rightarrow 0 \quad a.s.,$$

as $n \rightarrow \infty, \varepsilon_n \rightarrow 0$ and $\sqrt{n\varepsilon_n^{d/4}}/\sqrt{\log n} \rightarrow \infty$. Thus, we show that setting $\hat{f}/\varepsilon_n = -\log(\rho)/2$ asymptotically fulfils the fixed-point equation in a suitably strong sense. This argument further shows why we require a lower bounded density on a compact set: Assumption [\(Supp\)](#). We believe this requirement to be an artifact of the proof.

4. About extending the results to two different measures

A natural question is whether our results can be extended to two different measures and whether the rates of convergence will be the same in that case. The situation in the two-measure case is more involved than in the one measure case. The objective of this section is thus to identify the principles at play while remaining not too technical. *We only brush the big picture and hope to generalize the results in a near-future. Our main contribution here is thus a qualitative view.* We shall show that the structure of the problem is different in the one and in the two-measure case. This difference can be roughly explained as follows.

In the self-transport case, as no transport happens for the unregularized problem, the leading behavior of the entropy-regularized counterpart is that of the blurring. In the two measure case however, the transport is the leading behavior for small regularization parameter, which explains why the operators appearing will be reminiscent of the linearisation of the Monge–Ampère equation.

Making sense of this claim is the purpose of the following subsections. Before developing further, let us introduce $K_{\varepsilon_n}^\mu[h]$, the operator defined as

$$h \mapsto \int \pi_{\varepsilon_n}(\cdot, y)h(y)d\mu(y).$$

Note that the notation has been adapted to reflect the measure against which one integrates, which is necessary as the situation is not symmetric anymore. Such operators (for μ and ν) are called the Sinkhorn operators. Also, let us recall a few facts that will turn useful. First, recall that the optimization problem 6 is the dual of the classical optimal transport problem. Any pair of optimizers (f_0, g_0) admits a particular form, arising from the c -transformation. More explicitly, one first can write

$$(f_0(x), g_0(y)) = \left(\frac{\|x\|^2}{2} - \psi_0(x), \frac{\|y\|^2}{2} - \varphi_0(y) \right),$$

where ψ_0 and φ_0 are Fenchel–Legendre conjugates of one another. An important element, is that the support of the optimal (unregularized) transport plan is the set

$$\{(x, y) : \psi_0(x) + \varphi_0(y) - \langle x, y \rangle = 0\}.$$

Finally, note that $\psi_0(x) + \varphi_0(y) - \langle x, y \rangle$ is locally quadratic in each argument, which is a central element and appears, for instance, in Assumption (Reg2). Let us set

$$D(x, y) := \psi_0(x) + \varphi_0(y) - \langle x, y \rangle.$$

In the sequel, we will denote by x^* the image of the point x by the optimal transport map¹, i.e., $x^* := T_0(x)$.

To simplify greatly the exposition and as the objective of this section is mostly to understand what happens we make the following assumption on the fluctuations of entropic potentials in the two-measure case.

- (Lin)** Consider a_n is a diverging sequence such that $a_n\varepsilon_n \rightarrow \infty$, as $n \rightarrow \infty$. Let $(f_{\varepsilon_n}, g_{\varepsilon_n})$ be a pair of entropic potentials and let $(\hat{f}_{\varepsilon_n}, \hat{g}_{\varepsilon_n})$ be their empirical counterpart based on n i.i.d. samples from μ, ν . We assume that the asymptotic expansion

$$\begin{aligned} a_n(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x)) + K_{\varepsilon_n}^\nu [a_n(\hat{g}_{\varepsilon_n}(\cdot) - g_{\varepsilon_n}(\cdot))] (x) \\ = -\frac{a_n\varepsilon_n}{\sqrt{n}} \int \pi_{\varepsilon_n}(x, y)d\mathbb{G}_n^\nu(y) + o_p(1) \end{aligned}$$

holds as well as its symmetric version.

Remark 15 (Fluctuations of entropic potentials in the two-measure case). The proof of this result would follow the same lines as Proposition 5 replacing the self-transport dual condition by the general dual fixed point system. There however, we would require the assumption

$$\frac{1}{\varepsilon_n} \|\hat{g}_{\varepsilon_n} - g_{\varepsilon_n}\|_\infty \rightarrow 0,$$

¹The latter exists owing to our assumptions on the densities.

as $n \rightarrow \infty, \varepsilon_n \rightarrow 0$ to conclude. There is currently no handle on the convergence above and it seems beyond the horizon to even argue along the lines of Section 3.4.

Proposition 16. Under the same setting as Assumption **(Lin)**, it holds

$$\begin{aligned} & \begin{pmatrix} \hat{f}_{\varepsilon_n} - f_{\varepsilon_n} \\ \hat{g}_{\varepsilon_n} - g_{\varepsilon_n} \end{pmatrix} \\ &= -\frac{\varepsilon_n}{\sqrt{n}} \begin{pmatrix} (\text{id} - K_{\varepsilon_n}^{\nu} K_{\varepsilon_n}^{\mu})^{-1} & -K_{\varepsilon_n}^{\nu} (\text{id} - K_{\varepsilon_n}^{\mu} K_{\varepsilon_n}^{\nu})^{-1} \\ -(\text{id} - K_{\varepsilon_n}^{\mu} K_{\varepsilon_n}^{\nu})^{-1} K_{\varepsilon_n}^{\mu} & (\text{id} - K_{\varepsilon_n}^{\mu} K_{\varepsilon_n}^{\nu})^{-1} \end{pmatrix} \begin{bmatrix} \int \pi_{\varepsilon_n}(\cdot, y) dG_n^{\nu}(y) \\ \int \pi_{\varepsilon_n}(x, \cdot) dG_n^{\mu}(x) \end{bmatrix} \\ &+ o_p(a_n^{-1}). \end{aligned}$$

Note that this result is reminiscent of the fundamental work of [González-Sanz et al. \(2022\)](#) with the difference that the regularization parameter is fixed therein.

4.1. On the limit of the Sinkhorn operators

As appears in developments above, a key object to understand the limiting distribution is the inverse of the identity minus the successive application of the two Sinkhorn operators, i.e., $(\text{id} - K_{\varepsilon_n}^{\nu} K_{\varepsilon_n}^{\mu})^{-1}$. The composition of the Sinkhorn operators is the object of Theorem 19, which requires a few prerequisites.

We start by requiring regularity of the unregularized optimal transport potentials.

(Reg2) For fixed $x \in \mathbb{R}^d$, there exists $C \geq 0$ such that

$$\left\| \varphi_0(y) + \psi_0(x) - \langle x, y \rangle - \frac{1}{2}(y - x^*)^{\top} \nabla^2 \varphi_0(x)(y - x^*) \right\| \leq C \|y - x^*\|^3,$$

and similarly in the other variable.

(Reg2') The convex potential φ_0^* is at least C^4 with the following bounds on the eigenvalues $0 < \lambda_{\min}(\nabla^2 \varphi_0^*) \leq \lambda_{\max}(\nabla^2 \varphi_0^*) < +\infty$. Furthermore, the Hessian $\nabla^2 \varphi_0^*$ is itself L -Lipschitz.

Remark 17. Note that both these assumptions hold for Gaussian measures. Indeed, $C = 0$ in **(Reg2)** and $L = 0$ in **(Reg2')**. Furthermore, the regularity theory for the Monge–Ampère equation ([Caffarelli, 1996](#); [Urbas, 1997](#)) provides conditions under which **(Reg2')**, **(Reg2)** hold. Yet, such conditions appear very restrictive, which is why we prefer stating a more general condition.

We additionally require the following assumption, which is an expansion of the potentials in the two measure case.

(Exp) Assume that there exist constants $C_f, C_g, \varepsilon_0 > 0$, such that

$$\begin{aligned} \left| f_{\varepsilon}(x) - f_0(x) + \frac{\varepsilon}{2} \log \mu(x) + \frac{\varepsilon d}{4} \log(2\pi\varepsilon) \right| &\leq \varepsilon^2 C_f \|x\|^2 \\ \left| g_{\varepsilon}(y) - g_0(y) + \frac{\varepsilon}{2} \log \nu(y) + \frac{\varepsilon d}{4} \log(2\pi\varepsilon) \right| &\leq \varepsilon^2 C_g \|y\|^2, \end{aligned}$$

for all $\varepsilon \leq \varepsilon_0$.

This type of expansions has been presented at a BIRS Workshop by Léger (2021) and are since well-accepted. Unfortunately, these results have never been written down, to the best of our knowledge. Remark also that this type of developments further hold true in the Gaussian case, see Mallasto et al. (2022), which is why, we deem this assumption to be reasonable.

Additionally, we exhibit in Remark 18 the fact that such expansions fulfil the duality constraints in the limit $\varepsilon \rightarrow 0$, showing why such development make sense.

Remark 18 (On the expansion of the potentials in the two-measure case). We show that the expansion of the potentials is indeed legitimate. Start from the dual condition

$$1 = \int \pi_\varepsilon(x, y) \nu(y) dy.$$

Then plugging-in the assumed expansion, we get

$$\begin{aligned} 1 &= \int \exp\left(\frac{1}{\varepsilon_n} [f_\varepsilon(x) + g_\varepsilon(y) - \frac{1}{2}\|x - y\|^2]\right) \nu(y) dy \\ &= \int \frac{1}{(2\pi\varepsilon)^{d/2} \sqrt{\mu(x)\nu(y)}} \exp\left(\frac{1}{\varepsilon} [f_0(x) + g_0(y) - \frac{1}{2}\|x - y\|^2] + o(1)\right) \nu(y) dy \\ &= \int \frac{1}{(2\pi\varepsilon)^{d/2} \sqrt{\mu(x)\nu(y)}} \exp\left(-\frac{1}{2\varepsilon} [(y - x^*)^\top \nabla^2 \varphi_0(x)(y - x^*)] + o(1)\right) \nu(y) dy \\ &= \int \frac{\sqrt{\det[\nabla^2 \varphi_0(x)]}}{(2\pi\varepsilon)^{d/2}} \exp\left(-\frac{1}{2\varepsilon} [(y - x^*)^\top \nabla^2 \varphi_0(x)(y - x^*)] + o(1)\right) \frac{\nu^{1/2}(y)}{\sqrt{\nu(x^*)}} dy, \end{aligned}$$

where we used the regularity of the optimal transport and the Monge–Ampère equation. The little o in the above is thus expressing the departure from quadraticity of the metric tensor as well as the remainder in the expansion of the potentials. All in all, we see that the expansion satisfies the dual condition as $\varepsilon \rightarrow 0$. The same holds for the other dual condition using a symmetric reasoning.

Theorem 19. *Under Assumptions (Exp), (Reg2) and (Reg2'), it holds that*

$$\begin{aligned} h &\mapsto K_{\varepsilon_n}^\mu \left[K_{\varepsilon_n}^\nu [h] \right] (y_2) \\ &= \frac{1 + o(1)}{\nu^{1/2}(y_2)} \int \frac{1}{(2\pi\varepsilon_n)^{d/2}} h(y) \sqrt{\frac{\det[\nabla^2 \varphi_0^*(y_2)] \det[\nabla^2 \varphi_0^*(y)]}{\det[\nabla^2 \varphi_0^*(y_2) + \nabla^2 \varphi_0^*(y)]}} \\ &\quad \times \exp\left(-\frac{1}{4\varepsilon_n} (y - y_2)^\top [\nabla^2 \varphi_0^*(y_2)] (y - y_2) + o\left(\frac{\|y - y_2\|^2}{\varepsilon_n}\right)\right) \nu^{1/2}(y) dy, \end{aligned} \tag{15}$$

as $\varepsilon_n \rightarrow 0$.

The meaning of Theorem 19, is the composition of the two Sinkhorn operators is again an integral operator which behaves like a convolution with a Gaussian-type density whose quadratic form matrix depends on the Kim–McCann geometry. Note also the presence of the square root of the determinant inside the integral. By analogy, recall that for a Riemannian manifold (\mathcal{M}, g) , $\sqrt{\det g}$ is the volume element. A Taylor expansion in y around y_2 of this square root will thus provide the local fluctuations of the volume form of the metric tensor.

4.2. Towards limiting distribution results

To simplify the exposition, we will place ourselves in the one-sample case. In that setting, we get that

$$a_n(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x)) = -\frac{a_n \varepsilon_n}{\sqrt{n}} \int (\text{id} - K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu)^{-1} [\pi_{\varepsilon_n}(x, y)] d\mathbb{G}_n^\mu(y) + o_p(1), \tag{16}$$

using the Neumann series expansion of the inverse and noticing that each term is an integral operator.

The main claim of this section is the following.

Theorem 20. *Under Assumptions (Lin), (Exp), (Reg2) and (Reg2'), for fixed $x \in \text{supp}(\mu)$, there exist some matrices $\Sigma(x)$ and $\Sigma'(x)$ such that*

$$\sqrt{n} \varepsilon_n^{d/4} \left(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x) \right) \rightsquigarrow \mathcal{N}(0, \Sigma(x)), \quad \text{as } \varepsilon_n \rightarrow 0, n \rightarrow \infty.$$

Further,

$$\sqrt{n} \varepsilon_n^{d/4-1/2} \left(\nabla \hat{f}_{\varepsilon_n}(x) - \nabla f_{\varepsilon_n}(x) \right) \rightsquigarrow \mathcal{N}(0, \Sigma'(x)), \quad \text{as } \varepsilon_n \rightarrow 0, n \rightarrow \infty.$$

In the case where the Kim–McCann tensor is constant and equal to A , it further holds that

$$\Sigma(x) = \lim_{n \rightarrow \infty} \frac{1}{\mu(x)} \int_{\mathbb{R}^d} \left(\frac{(2\pi\varepsilon_n)^{-d/4} \mathcal{F} \left[e^{-\frac{1}{\varepsilon_n} D(x, \cdot)} \right] (\xi)}{\xi^\top A^{-1} \xi} \right)^2 d\xi,$$

where the last limit exists as $D(x, \cdot)$ has a quadratic behavior.

Remark 21. In the recent work of Manole et al. (2023), they study the statistical impacts of estimating the optimal transport map between μ and ν by the optimal transport between kernel density estimators of empirical measures. They rely on a linearisation of the Monge–Ampère equation which gives rise to studying the solutions of an elliptic with random functions. Interestingly, the structure of the asymptotic variance is very close to that we obtain above; see Theorem 5 in that reference for instance. To help the reader, we provide the gist of their approach and their main result in Section A.2.

Proof of Theorem 20. The proof will follow from the Lindeberg–Feller central limit theorem, using the representation (16). To this end, we first aim at understanding the behavior of the “Sinkhorn fluctuation kernel”

$$(\text{id} - K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu)^{-1}[\pi_{\varepsilon_n}(x, y)],$$

which we carry out in detail below. The latter kernel is further regular under the set of assumptions considered. For the statement on the transport plans, one can take the gradient under the integral sign and one again invokes the Lindeberg–Feller central limit theorem. Explicit forms of the asymptotic variance can be obtained in certain cases and rely on the Plancherel theorem.

We are thus left with the problem of understanding

$$\begin{aligned} y &\mapsto (\text{id} - K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu)^{-1}[\pi_{\varepsilon_n}(x, y)] \\ &= y \mapsto (\text{id} - K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu)^{-1} \left[\frac{\exp\left(-\frac{1}{\varepsilon_n}D(x, y) + o(1)\right)}{(2\pi\varepsilon_n)^{d/2}\nu^{1/2}(y)\mu^{1/2}(x)} \right]. \end{aligned}$$

Recall that $K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu$. The correct order of the spectral gap is $1 - O(\varepsilon_n)$ so that $\varepsilon_n(\text{id} - K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu)^{-1}$ is a bounded operator for $\varepsilon_n < \varepsilon_0$, for some $\varepsilon_0 > 0$. The quantity

$$(\text{id} - K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu)^{-1}[\pi_{\varepsilon_n}(x, y)]$$

is actually equivalent to finding $y \mapsto h_x(y)$ such that

$$h_x(y) - K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu [h_x(\cdot)](y) = \left[\frac{\exp\left(-\frac{1}{\varepsilon_n}D(x, y) + o(1)\right)}{(2\pi\varepsilon_n)^{d/2}\nu^{1/2}(y)\mu^{1/2}(x)} \right].$$

Finally, set

$$h_x(y) = \frac{\tilde{h}_x(y)}{\nu^{1/2}(y)\mu^{1/2}(x)}.$$

We first consider the case of a constant Kim–McCann curvature before turning to the more general case, as it provides a particularly neat picture.

Constant Kim–McCann curvature. Owing to the explicit characterisation of $K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu$, assuming for the rest of this section that the Kim–McCann metric tensor is constant, i.e., $\nabla^2\varphi_0^*(y_2) = A$, $\forall y_2$, we can write

$$\tilde{h}_x(y) - (\gamma_2(y) * \tilde{h}_x)(y) \asymp \left[\frac{1}{(2\pi\varepsilon_n)^{d/2}} \exp\left(-\frac{1}{\varepsilon_n}D(x, y) + o(1)\right) \right],$$

where

$$\gamma_2(y) := \frac{\sqrt{\det A}}{(4\pi\varepsilon_n)^{d/2}} \exp\left(-\frac{1}{4\varepsilon_n}y^\top Ay\right).$$

Denoting by \mathcal{F} the Fourier transform, we thus get that

$$h_x(y) \asymp \frac{1}{\nu^{1/2}(y)\mu^{1/2}(x)} \mathcal{F}^{-1} \left[\frac{(2\pi\varepsilon_n)^{-d/2} \mathcal{F} \left[e^{-\frac{1}{\varepsilon_n} D(x, \cdot) + o(1)} \right]}{1 - \mathcal{F}[\gamma_2(\cdot)]} \right] (y). \quad (17)$$

Noticing that

$$\mathcal{F}[\gamma_2](\xi) = \exp(-\varepsilon_n \xi^\top A^{-1} \xi),$$

we can even rewrite

$$h_x(y) \asymp \frac{1}{\nu^{1/2}(y)\mu^{1/2}(x)} \mathcal{F}^{-1} \left[\frac{(2\pi\varepsilon_n)^{-d/2} \mathcal{F} \left[e^{-\frac{1}{\varepsilon_n} D(x, \cdot) + o(1)} \right]}{\varepsilon_n \xi^\top A^{-1} \xi + o(\varepsilon_n)} \right] (y)$$

The fluctuations of the Sinkhorn divergence, for $\varepsilon_n \rightarrow 0$, behaves like a kernel density estimator, where the kernel is the canonical solution of an elliptic equation with Gaussian right hand side in the case where the Kim-McCann metric tensor has a constant curvature, i.e., when γ_2 is Gaussian. Remark that the spectral gap appears at the denominator.

Nonconstant Kim–McCann curvature In the previous section, we assumed that the Kim–McCann curvature was constant. We thus now describe the more general case. Then,

$$\begin{aligned} & \mathcal{F} \left[K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu [\tilde{h}_x] \right] (\xi) \\ & \asymp \int \tilde{h}_x(y) e^{-i2\pi \langle \xi, y \rangle} \left[\int \frac{1}{(4\pi)^{d/2}} \sqrt{\frac{\det[\nabla^2 \varphi_0^*(y + \sqrt{\varepsilon_n} u)] \det[\nabla^2 \varphi_0^*(y)]}{\det[\nabla^2 \varphi_0^*(y + \sqrt{\varepsilon_n} u) + \nabla^2 \varphi_0^*(y)]}} \right. \\ & \quad \left. \times \exp \left(-\frac{1}{4} u^\top \nabla^2 \varphi_0^*(y + \sqrt{\varepsilon_n} u) u \right) e^{-i2\pi \langle \sqrt{\varepsilon_n} \xi, u \rangle} du \right] dy. \end{aligned}$$

All in all,

$$\begin{aligned}
 & \mathcal{F}[\tilde{h}_x](\xi) - \mathcal{F}\left[K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu[\tilde{h}_x]\right](\xi) \\
 & \asymp \int \tilde{h}_x(y) e^{-i2\pi\langle \xi, y \rangle} \left[1 - \int \frac{1}{(2\pi)^{d/2}} \sqrt{\frac{\det[\nabla^2 \varphi_0^*(y + \sqrt{\varepsilon_n}u)] \det[\nabla^2 \varphi_0^*(y)]}{\det[\nabla^2 \varphi_0^*(y + \sqrt{\varepsilon_n}u) + \nabla^2 \varphi_0^*(y)]}} \right. \\
 & \quad \left. \times \exp\left(-\frac{1}{4}u^\top \nabla^2 \varphi_0^*(y + \sqrt{\varepsilon_n}u)u\right) e^{-i2\pi\langle \sqrt{\varepsilon_n}\xi, u \rangle} du \right] dy \\
 & \asymp \int \tilde{h}_x(y) e^{-i2\pi\langle \xi, y \rangle} \left[\int \frac{e^{-\frac{u^\top \nabla^2 \varphi_0^*(y)u}{4}}}{(4\pi)^{d/2} \det^{-1/2}[\nabla^2 \varphi_0^*(y)]} \right. \\
 & \quad \left. \times \left(\sqrt{\varepsilon_n}L(y, \xi)^\top u + \varepsilon_n u^\top M(y, \xi)u + o(\varepsilon_n)\right) du \right] dy \\
 & \asymp \varepsilon_n \int \tilde{h}_x(y) e^{-i2\pi\langle \xi, y \rangle} \operatorname{tr}\left[M(y, \xi)(\nabla^2 \varphi_0^*(y))^{-1}(1 + o(1))\right] dy,
 \end{aligned}$$

where $L(y, \xi), M(y, \xi)$ are given by a second order Taylor development. This ultimately leads to the relationship

$$\begin{aligned}
 & \varepsilon_n \int \tilde{h}_x(y) e^{-i2\pi\langle \xi, y \rangle} \operatorname{tr}\left[M(y, \xi)(\nabla^2 \varphi_0^*(y))^{-1}(1 + o(1))\right] dy \\
 & = (2\pi\varepsilon_n)^{-d/2} \mathcal{F}\left[e^{-\frac{1}{\varepsilon_n}D(x, \cdot) + o(1)}\right].
 \end{aligned}$$

This implies that the solution \tilde{h}_x will converge to the canonical solution of some elliptic partial differential equation. \square

4.3. On estimating the true transport map

Let us denote by $\hat{T}_{\varepsilon_n}(x)$ the empirical barycentric projection at x ; recall Equation (9). In the setting of Theorem 20, one would like to get for fixed x

$$\sqrt{n}\varepsilon_n^{d/4-1/2} \left(\hat{T}_{\varepsilon_n}(x) - T(x)\right) \rightsquigarrow \mathcal{N}(0, \Sigma'(x)), \quad \text{as } \varepsilon_n \rightarrow 0, n \rightarrow \infty,$$

provided that one chooses ε_n in a suitable way. To do so, in view of Theorem 20, it would remain to control the bias. The barycentric projection can conveniently be written as $K_{\varepsilon_n}[\text{id}](x)$. Recall the proof of Theorem 19, where we now are setting $h(y) = y$. Carrying out the same analysis to the second order, we get under our regularity assumptions that $T_{\varepsilon_n}(x) = T_0(x) + O(\varepsilon_n^\delta)$, where delta will depend on the rate of convergence of the potential expansions, which is the content of Assumption (Exp), and the regularity of the Kim–McCann metric tensor to the fourth order, which is covered by Assumption (Reg2). Note that in the Gaussian case explicit formulas exist (Pooladian et al., 2022, Proposition 3) and give

$$T_\varepsilon(x) = T_0(x) - \frac{\varepsilon}{2} \Sigma_0^{-1} x + o(\varepsilon)x,$$

where Σ_0 is the covariance matrix of the source distribution. It would remain to choose the relationship between n and ε_n such that the bias vanishes. Getting rigorous higher order expansion like those of Assumption **(Exp)** is still an open question for the analysis community.

5. Empirical estimation of score functions

5.1. On the computation of our estimator

We now show that the computation of the self-transport potentials is easy as it is based on the fixed point relationship 11. Iterating this process on the basis of samples usually converges in a handful of iterations (typically in two to three iterations, as also observed by Feydy et al. (2019)). We provide pseudo-code in Algorithm 1 (“LSE” refers to the ubiquitous log-sum-exp operator, which leads to well-conditioned updates).

Algorithm 1 Computation of the self-OT potential

Input: $K_{\max} \geq 0$, data $(x_i)_{i=1}^n \sim \rho$, regularization parameter ε
Initialize: $f^{(0)} \leftarrow 0_n \in \mathbb{R}^n$ and $k \leftarrow 1$ $\triangleright f_i^{(k)} := f^{(k)}(x_i)$
while $k \leq K_{\max}$ **do**
 $f_i^{(k)} \leftarrow \frac{1}{2} f_i^{(k-1)} - \frac{\varepsilon}{2} \text{LSE}_{j=1}^n \left[\log\left(\frac{1}{n}\right) + \frac{1}{\varepsilon} f_j^{(k-1)} - \frac{1}{2\varepsilon} \|x_i - x_j\|^2 \right]$
 $k \leftarrow k + 1$
end while
Return: $f^{(K_{\max})}$

5.2. Adaptation to intrinsic dimension

We have showed that the gradient of the self-transport potentials can serve as estimators of the score function, a central object for diffusion models. In this experiment, we consider a Gaussian random variable that is supported on a 3-dimensional subspace in the case where the observed data is of dimension d , for $d \in \{4, 6, 8, 10\}$. The L^2 error is estimated based on Monte-Carlo integration with 50000 points and the results are averaged over 5 replicates for more robustness of the results. The results are presented in Figure 1. From that graph, it is clear that estimation via self-transport, using the bandwidth size that is optimal for the ambient dimension is performing much better than the kernel density estimator with the same bandwidth choice. This is an instance of the adaptivity property hinted at in the introduction. In practice, for machine learning application where the intrinsic dimension is unknown, this feature of self-transport is clear advantage of kernel density estimators.

6. Conclusion

In this paper, we developed a statistical theory for the maximizer f_ε to the entropic optimal self-transport problem. We established (pointwise) limit dis-




Fig 1: Estimation error (MC-approximated) of the self-ot scores and KDE in a Gaussian model for various dimensions $d \in \{4, 6, 8, 10\}$ with the naïve equivalent bandwidth choice $n^{-\frac{1}{d+4}}$. The covariance matrix is diagonal on the subspace on which it concentrates with diagonal $(0.5, 1, 1.5)$.

tribution results for f_ε for shrinking regularization parameter. Additionally, we showed how the maximizer can be used to easily estimate the score function of a smooth distribution ρ on the basis of i.i.d. samples.

We extended the techniques developed in the first part of this paper to study the (pointwise) fluctuations of the entropic potentials and plans in the non-self transport case for vanishing ε , which opens the door to a complete understanding of the regularized optimal transport.

Acknowledgements

I am thankful to Alberto González-Sanz, Shayan Hundrieser, Jonathan Niles-Weed, Aram-Alexandre Pooladian, Tudor Manole and Thomas Staudt for helpful discussions. The support of the Deutsche Forschungsgemeinschaft through SFB1456 is further acknowledged.

References

- M. Agarwal, Z. Harchaoui, G. Mulcahy, and S. Pal. Iterated Schrödinger bridge approximation to Wasserstein gradient flows. *arXiv:2406.10823*, 2024.
- M. Blondel, V. Seguy, and A. Rolet. Smooth and sparse optimal transport. In *International conference on artificial intelligence and statistics*, pages 880–889. PMLR, 2018.
- Y. Brenier. Polar factorization and monotone rearrangement of vector-valued functions. *Comm. Pure Appl. Math.*, 44(4):375–417, 1991. ISSN 0010-3640. . URL <https://doi.org/10.1002/cpa.3160440402>.

- L. A. Caffarelli. Boundary regularity of maps with convex potentials–ii. *The Annals of Mathematics*, 144(3):453, 1996. .
- G. Carlier and M. Laborde. A differential approach to the multi-marginal schrödinger system. *SIAM Journal on Mathematical Analysis*, 52(1):709–717, 2020.
- S. Chewi. Log-concave sampling. Book draft available at <https://chewisinho.github.io>, 2023.
- S. Chewi, J. Niles-Weed, and P. Rigollet. Statistical optimal transport. *arXiv preprint arXiv:2407.18163*, 2024.
- L. Chizat, P. Roussillon, F. Léger, F.-X. Vialard, and G. Peyré. Faster wasserstein distance estimation with the sinkhorn divergence. *Advances in Neural Information Processing Systems*, 33:2257–2269, 2020.
- I. Csiszár. I -divergence geometry of probability distributions and minimization problems. *Ann. Probability*, 3:146–158, 1975.
- M. Cuturi. Sinkhorn distances: lightspeed computation of optimal transport. *Advances in Neural Information Processing Systems*, 26, 2013.
- N. Deb, Y.-H. Kim, S. Pal, and G. Schiebinger. Wasserstein mirror gradient flow as the limit of the sinkhorn algorithm. *arXiv preprint arXiv:2307.16421*, 2023.
- E. del Barrio and J.-M. Loubes. Central limit theorems for empirical transportation cost in general dimension. *The Annals of Probability*, 47(2):926 – 951, 2019. . URL <https://doi.org/10.1214/18-AOP1275>.
- E. del Barrio, A. G.-S. Sanz, J.-M. Loubes, and J. Niles-Weed. An improved central limit theorem and fast convergence rates for entropic transportation costs. *SIAM Journal on Mathematics of Data Science*, 5(3):639–669, 2023.
- U. Einmahl and D. M. Mason. Uniform in bandwidth consistency of kernel-type function estimators. *The Annals of Statistics*, 33(3):1380 – 1403, 2005.
- J. Feydy, T. Séjourné, F.-X. Vialard, S.-i. Amari, A. Trounev, and G. Peyré. Interpolating between optimal transport and mmd using sinkhorn divergences. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 2681–2690. PMLR, 2019.
- A. Genevay. *Entropy-regularized optimal transport for machine learning*. PhD thesis, Paris Sciences et Lettres (ComUE), 2019.
- A. Genevay, L. Chizat, F. Bach, M. Cuturi, and G. Peyré. Sample complexity of Sinkhorn divergences. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1574–1583. PMLR, 2019.
- Z. Goldfeld, K. Kato, G. Rioux, and R. Sadhu. Limit theorems for entropic optimal transport maps and the Sinkhorn divergence. *arXiv preprint arXiv:2207.08683*, 2022.
- Z. Goldfeld, K. Kato, G. Rioux, and R. Sadhu. Limit theorems for entropic optimal transport maps and sinkhorn divergence. *Electronic Journal of Statistics*, 18(1):980–1041, 2024a.
- Z. Goldfeld, K. Kato, G. Rioux, and R. Sadhu. Statistical inference with regularized optimal transport. *Information and Inference: A Journal of the IMA*, 13(1):iaad056, 2024b.
- A. González-Sanz and S. Hundrieser. Weak limits for empirical entropic optimal

- transport: Beyond smooth costs. *arXiv preprint arXiv:2305.09745*, 2023.
- A. González-Sanz, J.-M. Loubes, and J. Niles-Weed. Weak limits of entropy regularized optimal transport; potentials, plans and divergences. *arXiv preprint arXiv:2207.07427*, 2022.
- E. Grave, A. Joulin, and Q. Berthet. Unsupervised alignment of embeddings with wasserstein procrustes. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1880–1890. PMLR, 2019.
- M. Groppe and S. Hundrieser. Lower complexity adaptation for empirical entropic optimal transport. *arXiv preprint arXiv:2306.13580*, 2023.
- S. Hundrieser, M. Klatt, T. Staudt, and A. Munk. A unifying approach to distributional limits for empirical optimal transport. *arXiv preprint arXiv:2202.12790*, 2022.
- S. Hundrieser, G. Mordant, C. A. Weitkamp, and A. Munk. Empirical optimal transport under estimated costs: Distributional limits and statistical applications, 2023.
- S. Hundrieser, G. Mordant, C. A. Weitkamp, and A. Munk. Empirical optimal transport under estimated costs: Distributional limits and statistical applications. *Stochastic Processes and their Applications*, 178:104462, 2024.
- J.-C. Hütter and P. Rigollet. Minimax estimation of smooth optimal transport maps. *The Annals of Statistics*, 49(2):1166 – 1194, 2021. . URL <https://doi.org/10.1214/20-AOS1997>.
- J.-C. Hütter and P. Rigollet. Minimax estimation of smooth optimal transport maps. *Ann. Statist.*, 49(2):1166–1194, 2021.
- A. Hyvärinen and P. Dayan. Estimation of non-normalized statistical models by score matching. *Journal of Machine Learning Research*, 6(4), 2005.
- D. S. Kim, H. Kim, and T. Kim. Some identities on generalized harmonic numbers and generalized harmonic functions. *Demonstratio Mathematica*, 56(1):20220229, 2023.
- M. Klatt, A. Munk, and Y. Zemel. Limit laws for empirical optimal solutions in random linear programs. *Annals of Operations Research*, 315(1):251–278, 2022.
- B. Landa and X. Cheng. Robust inference of manifold density and geometry by doubly stochastic scaling. *SIAM Journal on Mathematics of Data Science*, 5(3):589–614, 2023. .
- B. Landa, R. R. Coifman, and Y. Kluger. Doubly stochastic normalization of the Gaussian kernel is robust to heteroskedastic noise. *SIAM journal on mathematics of data science*, 3(1):388–413, 2021.
- H. Lavenant, J. Luckhardt, G. Mordant, B. Schmitzer, and L. Tamanini. The Riemannian geometry of Sinkhorn divergences. *arXiv preprint arXiv:2405.04987*, 2024.
- F. Léger. BIRS workshop: Entropic regularization of optimal transport and applications (21w5120), 2021. URL <https://www.birs.ca/events/2021/5-day-workshops/21w5120>.
- A. Mallasto, A. Gerolin, and H. Q. Minh. Entropy-regularized 2-Wasserstein distance between Gaussian measures. *Information Geometry*, pages 1–35, 2021.

- A. Mallasto, A. Gerolin, and H. Q. Minh. Entropy-regularized 2-Wasserstein distance between Gaussian measures. *Information Geometry*, 5(1):289–323, 2022.
- T. Manole, S. Balakrishnan, J. Niles-Weed, and L. Wasserman. Central limit theorems for smooth optimal transport maps. *arXiv:2312.12407*, 2023.
- T. Manole, S. Balakrishnan, J. Niles-Weed, and L. Wasserman. Plugin estimation of smooth optimal transport maps. *The Annals of Statistics*, 52(3):966–998, 2024.
- N. F. Marshall and R. R. Coifman. Manifold learning with bi-stochastic kernels. *IMA Journal of Applied Mathematics*, 84(3):455–482, 2019.
- G. Mena and J. Niles-Weed. Statistical bounds for entropic optimal transport: sample complexity and the central limit theorem. *Advances in Neural Information Processing Systems*, 32, 2019.
- M. Nutz and J. Wiesel. Entropic optimal transport: convergence of potentials. *Probability Theory and Related Fields*, pages 1–24, 2021.
- S. Pal. On the difference between entropic cost and the optimal transport cost. *The Annals of Applied Probability*, 34(1B):1003–1028, 2024.
- G. Peyré. Denoising diffusion models, 2023.
- G. Peyré and M. Cuturi. Computational optimal transport. *Foundations and Trends® in Machine Learning*, 11(5-6):355–607, 2019.
- A.-A. Pooladian and J. Niles-Weed. Entropic estimation of optimal transport maps. *arXiv preprint arXiv:2109.12004*, 2021.
- A.-A. Pooladian, M. Cuturi, and J. Niles-Weed. Debiasser beware: Pitfalls of centering regularized transport maps. In *International Conference on Machine Learning*, pages 17830–17847. PMLR, 2022.
- A.-A. Pooladian, V. Divol, and J. Niles-Weed. Minimax estimation of discontinuous optimal transport maps: The semi-discrete case. *arXiv preprint arXiv:2301.11302*, 2023.
- A.-A. Pooladian, C. Domingo-Enrich, R. T. Chen, and B. Amos. Neural optimal transport with Lagrangian costs. *arXiv preprint arXiv:2406.00288*, 2024.
- P. Rigollet and A. J. Stromme. On the sample complexity of entropic optimal transport. *arXiv preprint arXiv:2206.13472*, 2022.
- R. J. Samworth. Recent Progress in Log-Concave Density Estimation. *Statistical Science*, 33(4):493 – 509, 2018. . URL <https://doi.org/10.1214/18-STS666>.
- F. Santambrogio. Optimal transport for applied mathematicians. *Birkhäuser, NY*, 55(58-63):94, 2015.
- A. SerEGIN and J. A. Wellner. Nonparametric estimation of multivariate convex-transformed densities. *Annals of statistics*, 38(6):3751, 2010.
- J. Sesma. The Roman harmonic numbers revisited. *Journal of Number Theory*, 180:544–565, 2017.
- B. W. Silverman. *Density estimation for statistics and data analysis*. Routledge, 2018.
- M. Sommerfeld and A. Munk. Inference for empirical Wasserstein distances on finite spaces. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 80(1):219–238, 2018.
- Y. Song, S. Garg, J. Shi, and S. Ermon. Sliced score matching: A scalable

- approach to density and score estimation. In *Uncertainty in Artificial Intelligence*, pages 574–584. PMLR, 2020.
- B. Sriperumbudur, K. Fukumizu, A. Gretton, A. Hyv, R. Kumar, et al. Density estimation in infinite dimensional exponential families. *Journal of Machine Learning Research*, 18(57):1–59, 2017.
- A. J. Stromme. Minimum intrinsic dimension scaling for entropic optimal transport, 2023.
- A. B. Tsybakov. *Introduction to nonparametric estimation*. Springer Series in Statistics. Springer, New York, 2009.
- J. Urbas. On the second boundary value problem for equations of Monge-Ampère type. *Journal für die reine und angewandte Mathematik*, 487:115–124, 1997. URL <http://eudml.org/doc/153916>.
- A. Vacher, B. Muzellec, F. Bach, F.-X. Vialard, and A. Rudi. Optimal estimation of smooth transport maps with kernel SoS. *SIAM Journal on Mathematics of Data Science*, 6(2):311–342, 2024.
- A. W. Van der Vaart. *Asymptotic statistics*, volume 3. Cambridge university press, 2000.
- C. Villani. *Optimal transport: old and new*, volume 338. Springer, 2009.
- C. Villani. *Topics in optimal transportation*, volume 58. American Mathematical Soc., 2021.
- A. Wibisono, Y. Wu, and K. Y. Yang. Optimal score estimation via empirical Bayes smoothing, 2024.
- S. Zhang, G. Mordant, T. Matsumoto, and G. Schiebinger. Manifold learning with sparse regularised optimal transport. *arXiv preprint arXiv:2307.09816*, 2023.
- Y. Zhou, J. Shi, and J. Zhu. Nonparametric score estimators. In *International Conference on Machine Learning*, pages 11513–11522. PMLR, 2020.

Appendix A: Additional background

A.1. EOT in the Gaussian setting

In this section we first state the results about the potentials in the Gaussian case.

This is the content of Theorem 2 in [Mallasto et al. \(2021\)](#), which we slightly rephrased and took out the part of interest to us.

Theorem 22. *Let $\mu_i = \mathcal{N}(0, K_i)$ for $i \in 0, 1$ be two centred Gaussian distributions in \mathbb{R}^d , write $N_{i,j}^\varepsilon = \left(I + \frac{16}{\varepsilon^2} K_i^{1/2} K_j K_i^{1/2}\right)^{1/2}$ and $M^\varepsilon = I + \left(I + \frac{16}{\varepsilon^2} K_0 K_1\right)^{1/2}$. Then, the density of the optimal transport plan is given by*

$$\pi^\varepsilon(x, y) = \alpha^\varepsilon(x) \alpha^\varepsilon(y) \exp\left(-\frac{\|x - y\|^2}{\varepsilon}\right) \mu_0(x) \mu_1(y),$$

where $\alpha^\varepsilon(x) = \exp(x^\top A x + a)$, $\beta^\varepsilon(y) = \exp(y^\top B y + b)$ and

$$\begin{aligned} A &= \frac{1}{4} K_0^{-1/2} \left(I + \frac{4}{\varepsilon} K_0 - N_{01}^\varepsilon \right) K_0^{-1/2} \\ B &= \frac{1}{4} K_1^{-1/2} \left(I + \frac{4}{\varepsilon} K_1 - N_{10}^\varepsilon \right) K_1^{-1/2} \\ \exp(a + b) &= \sqrt{2^{-n} \det M^\varepsilon}. \end{aligned}$$

A.2. The approach of Manole et al. to estimate the transport map.

In the recent work by [Manole et al. \(2024\)](#), the authors propose to rely on kernel density estimators the Monge–Ampère equation to estimate transport plans given, as above, an i.i.d. sample. In this appendix we will just explain the ideas so that the reader can compare their results to the picture suggested by Section 4.

Start from the Monge–Ampère equation, assuming two probability densities p, q nonvanishing on the torus,

$$\det \nabla^2 \varphi_0 = \frac{p}{g \circ \nabla \varphi_0}.$$

Then, if one replaces q by a kernel density estimator \hat{q}_h with bandwidth h in the equation above, one has

$$\det \nabla^2 \hat{\varphi} = \frac{p}{\hat{g}_h \circ \nabla \hat{\varphi}}.$$

As \hat{p}_h converges to p as the sample size increases and the bandwidth decreases, [Manole et al. \(2024\)](#) prove that

$$\hat{\varphi} - \varphi_0 \asymp L^{-1}[\hat{q}_h - a],$$

where

$$L[u] = -\operatorname{div} (q \nabla u (\nabla \varphi_0^*)),$$

where div is the divergence operator and the asymptotic equivalence is in a suitable topology. Their theorem 5 then shows that, upon choosing properly the bandwidth h_n as a function of the sample size and for sufficiently smooth densities on the torus,

$$\sqrt{nh_n^{d-2}} \left(\hat{T}_n(x) - T_0(x) \right) \rightsquigarrow \mathcal{N}(0, \Sigma(x)), \quad \text{as } n \rightarrow \infty,$$

where

$$\Sigma(x) = \frac{1}{p(x)} \int_{\mathbb{R}^d} \xi \xi^\top \left(\frac{\mathcal{F}[K](\mathcal{M}(x)\xi)}{2\pi \langle \mathcal{M}(x)\xi, \xi \rangle} \right) d\xi, \quad \text{with } \mathcal{M}(x) := \nabla^2 \varphi_0^* (\nabla^2 \varphi_0(x)),$$

where K is the kernel used.

Appendix B: Proofs

B.1. Proof of Proposition 1

Proof of Proposition 1. It holds that

$$\begin{aligned} u(x) &= \int \frac{k_\varepsilon(x, y)}{u(y)} d\rho(y) \\ &= \varepsilon^{d/2} \int \frac{\rho(x + \sqrt{\varepsilon}z)}{u(x + \sqrt{\varepsilon}z)} \varphi(z) dz. \end{aligned}$$

where $\varphi(z) = \exp(-\|z\|^2/2)$. Then, using **(Tay)**,

$$u(x) = \frac{\rho(x)}{u(x)} \varepsilon^{d/2} \int \frac{1 + \sqrt{\varepsilon} \nabla \log \rho(x)^\top z + \varepsilon z^\top \nabla^2 \rho(x) z / (2\rho(x)) + o(\varepsilon)}{1 + \sqrt{\varepsilon} \nabla \log u(x)^\top z + \varepsilon z^\top \nabla^2 u(x) z / (2u(x)) + o(\varepsilon)} \varphi(z) dz.$$

Let us focus on the first summand in the display above and develop the inner fraction to get

$$\begin{aligned} & \frac{1 + \sqrt{\varepsilon} \nabla \log \rho(x)^\top z + \varepsilon z^\top \nabla^2 \rho(x) z / (2\rho(x)) + o(\varepsilon)}{1 + \sqrt{\varepsilon} \nabla \log u(x)^\top z + \varepsilon z^\top \nabla^2 u(x) z / (2u(x)) + o(\varepsilon)} \\ &= (1 + \sqrt{\varepsilon} \nabla \log \rho(x)^\top z + \varepsilon z^\top \nabla^2 \rho(x) z / (2\rho(x)) + o(\varepsilon)) \\ & \times \left(1 - \sqrt{\varepsilon} \nabla \log u(x)^\top z + \varepsilon z^\top \nabla^2 u(x) z / (2u(x)) \right. \\ & \quad \left. + \left(\sqrt{\varepsilon} \nabla \log u(x)^\top z \right)^2 + O(\varepsilon^{3/2}) + o(\varepsilon) \right). \end{aligned}$$

Above, the term of order $O(\varepsilon^{3/2})$ behaves like

$$\varepsilon^{3/2} \nabla \log \rho(x)^\top z \frac{z^\top \nabla^2 \rho(x) z}{2\rho(x)} + C \varepsilon^{3/2} \|z\|^2,$$

owing to the relationship between u and ρ , which we show below. It is thus negligible once integrated against a standard Gaussian. Also, the integral of the second-order Taylor residual of ρ with respect to the Gaussian measure is well-controlled; it has the form

$$\sum_{|\alpha|=2} \int h_\alpha(x + \sqrt{\varepsilon}z)(\sqrt{\varepsilon}z)^\alpha e^{-\frac{1}{2}\|z\|^2} dz,$$

for functions h_α such that $\lim_{z \rightarrow x} h_\alpha(z) = 0$. Clearly,

$$\varepsilon \int h_\alpha(x + \sqrt{\varepsilon}z) z^\alpha e^{-\frac{1}{2}\|z\|^2} dz = o(\varepsilon).$$

Altogether,

$$\begin{aligned} u(x)^2 &= \rho(x)\varepsilon^{d/2} \\ &\times \int \left(1 + \varepsilon z^\top \left(\frac{\nabla^2 \rho(x)}{2\rho(x)} - \frac{\nabla^2 u(x)}{2u(x)} - \nabla \log \rho(x) \nabla \log u(x)^\top \right. \right. \\ &\quad \left. \left. + \nabla \log u(x) \nabla \log u(x)^\top \right) z + o(\varepsilon) \right) \varphi(z) dz \end{aligned}$$

Finally,

$$u(x)^2 = \rho(x)(2\pi\varepsilon)^{d/2} \left(1 + \varepsilon \operatorname{tr} \left(\frac{\nabla^2 \rho(x)}{2\rho(x)} - \frac{\nabla^2 u(x)}{2u(x)} \right) + o(\varepsilon) \right).$$

We can further refine this result by “bootstrapping”, i.e., observing that the asymptotics expansion $\nabla \log u(x) = \nabla \log \rho^{1/2} + o(1)$ can be plugged-in. Therefore,

$$\nabla \log \rho(x) \nabla \log u(x)^\top = \frac{1}{2} \nabla \log \rho(x) (\nabla \log \rho(x))^\top + o(1).$$

Also,

$$\begin{aligned} \nabla^2 u(x) &= (2\pi\varepsilon)^{d/4} \rho^{1/2}(x) (1 + o(1)) \\ &= (2\pi\varepsilon)^{d/4} \nabla \left(\frac{1}{2} \rho^{-1/2}(x) \nabla \rho(x) \right) (1 + o(1)) \\ &= (2\pi\varepsilon)^{d/4} \frac{1}{2} \left(-\frac{1}{2} \rho^{-3/2}(x) \nabla \rho(x) (\nabla \rho(x))^\top + \rho^{-1/2}(x) \nabla^2 \rho(x) \right) (1 + o(1)) \end{aligned}$$

Thus,

$$\begin{aligned} &\operatorname{tr} \left(\frac{\nabla^2 \rho(x)}{2\rho(x)} - \frac{\nabla^2 u(x)}{2u(x)} - \nabla \log \rho(x) \nabla \log u(x)^\top \right) \\ &= \operatorname{tr} \left(\frac{\nabla^2 \rho(x)}{2\rho(x)} + \frac{1}{8} \nabla \log \rho(x) (\nabla \log \rho(x))^\top - \frac{\nabla^2 \rho(x)}{4\rho(x)} \right. \\ &\quad \left. - \frac{1}{4} \nabla \log \rho(x) (\nabla \log \rho(x))^\top \right) + o(1) \end{aligned}$$

and the claim follows. \square

B.2. Proof of Proposition 5

Before turning to the proof, we remark that the Laplace method combined with the expansions on the potentials give

$$\begin{aligned} K_{\varepsilon_n}[h] &= \int \frac{\exp\left(-\frac{1}{2\varepsilon_n}\|y-x\|^2\right)}{(2\pi\varepsilon_n)^{d/2}\rho^{1/2}(y)\rho^{1/2}(x)(1+o(1))} h(y) d\rho(y) \\ &= \frac{\rho^{1/2}(x)}{\rho^{1/2}(x)} h(x)(1+o(1)) = h(x)(1+o(1)). \end{aligned}$$

Proof of Proposition 5. To simplify the notation, we replace f_{ε_n} by f below. We start by re-writing the fluctuations using the fixed point equation characterization as

$$\begin{aligned} &a_n(\hat{f}(x) - f(x)) \\ &= a_n\varepsilon_n \left(\log \int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho - \log \int \exp\left(\frac{1}{\varepsilon_n}\hat{f}\right) k_{\varepsilon_n}(x, \cdot) d\rho_n \right) \\ &= a_n\varepsilon_n \log \left(\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho / \int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho_n \right) \\ &+ a_n\varepsilon_n \log \left(\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho_n / \int \exp\left(\frac{1}{\varepsilon_n}\hat{f}\right) k_{\varepsilon_n}(x, \cdot) d\rho_n \right) \\ &=: C(x) + D(x). \end{aligned}$$

Then,

$$\begin{aligned} C(x) &= -a_n\varepsilon_n \log \left(\frac{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho_n}{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho} \right) \\ &= -a_n\varepsilon_n \log \left(\frac{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho}{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho} \right. \\ &\quad \left. + \frac{1}{\sqrt{n}} \frac{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d(\sqrt{n}(\rho_n - \rho))}{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho} \right) \\ &= -\frac{a_n\varepsilon_n}{\sqrt{n}} \frac{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\mathbb{G}_n}{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho} \\ &\quad + o \left(-\frac{a_n\varepsilon_n}{\sqrt{n}} \frac{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\mathbb{G}_n}{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho} \right) \\ &= -\frac{a_n\varepsilon_n}{\sqrt{n}} \int \pi_{\varepsilon_n} d\mathbb{G}_n + o \left(-\frac{a_n\varepsilon_n}{\sqrt{n}} \frac{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\mathbb{G}_n}{\int \exp\left(\frac{1}{\varepsilon_n}f\right) k_{\varepsilon_n}(x, \cdot) d\rho} \right), \end{aligned}$$

where $\mathbb{G}_n := \sqrt{n}(\rho_n - \rho)$. Because of Assumption **(Reg)**, we get

$$\frac{a_n \varepsilon_n}{\sqrt{n}} \int \pi_{\varepsilon_n}(x, \cdot) d\mathbb{G}_n = \frac{a_n \varepsilon_n}{\sqrt{n}} \int \frac{\exp\left(\frac{-1}{2\varepsilon_n} \|y - x\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(y) \rho^{1/2}(x) (1 + o(1))} d\mathbb{G}_n(y).$$

On the other hand, as $a_n \varepsilon_n \rightarrow \infty$ as $n \rightarrow \infty$,

$$\begin{aligned} D(x) &= -a_n \varepsilon_n \log \left(\frac{\int \exp\left(\frac{1}{\varepsilon_n} \hat{f}\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n}{\int \exp\left(\frac{1}{\varepsilon_n} f\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n} \right) \\ &= -a_n \varepsilon_n \log \left(\frac{\int \exp\left(\frac{1}{\varepsilon_n} f + \frac{a_n}{\varepsilon_n a_n} (\hat{f} - f)\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n}{\int \exp\left(\frac{1}{\varepsilon_n} f\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n} \right) \\ &= -a_n \varepsilon_n \log \left(\frac{\int \exp\left(\frac{1}{\varepsilon_n} f\right) \left(1 + \frac{1}{\varepsilon_n} (\hat{f} - f) + O(\varepsilon_n^{-2} (\hat{f} - f)^2)\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n}{\int \exp\left(\frac{1}{\varepsilon_n} f\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n} \right) \\ &= -a_n \varepsilon_n \log \left(1 + \frac{1}{\varepsilon_n a_n} \frac{\int \exp\left(\frac{1}{\varepsilon_n} f\right) \left(a_n (\hat{f} - f) + O(a_n \varepsilon_n^{-1} (\hat{f} - f)^2)\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n}{\int \exp\left(\frac{1}{\varepsilon_n} f\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n} \right) \\ &=: -a_n \varepsilon_n \log \left(1 + \frac{1}{\varepsilon_n a_n} \Xi_n(x) \right), \end{aligned}$$

say. Now, let us use the fact that $\log(1 + x) = x + O(x^2)$ as $x \rightarrow 0$. Further, we aim at making ρ appear instead of $\hat{\rho}_n$ with a “plus-minus” trick. Also, we recall that

$$\hat{\rho}_n = \rho + \frac{1}{\sqrt{n}} \sqrt{n}(\hat{\rho}_n - \rho) = \rho + \frac{1}{\sqrt{n}} \mathbb{G}_n.$$

This will be used at the denominator as well so that we can make the optimal

coupling appear. All in all,

$$\begin{aligned}
& D(x) \\
&= - \frac{\int \exp\left(\frac{1}{\varepsilon_n} f\right) \left(a_n(\hat{f} - f) + O(a_n \varepsilon_n^{-1}(\hat{f} - f)^2)\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n}{\int \exp\left(\frac{1}{\varepsilon_n} f\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n} \\
&+ O(\varepsilon_n^{-1} a_n^{-1} \Xi_n(x)^2) \\
&= - \frac{\int \exp\left(\frac{1}{\varepsilon_n} f\right) \left(a_n(\hat{f} - f) + O(a_n \varepsilon_n^{-1}(\hat{f} - f)^2)\right) k_{\varepsilon_n}(x, \cdot) d\hat{\rho}_n}{\int \exp\left(\frac{1}{\varepsilon_n} f\right) k_{\varepsilon_n}(x, \cdot) d\rho + \frac{1}{\sqrt{n}} \int \exp\left(\frac{1}{\varepsilon_n} f\right) k_{\varepsilon_n}(x, \cdot) d\mathbb{G}_n} \\
&+ O(\varepsilon_n^{-1} a_n^{-1} \Xi_n(x)^2) \\
&= -(1 + O(\zeta_n)) \int \exp\left(\frac{1}{\varepsilon_n} (f(x) + f(y))\right) \\
&\quad \times \left(a_n(\hat{f}(y) - f(y)) + O(a_n \varepsilon_n^{-1}(\hat{f} - f)^2)\right) k_{\varepsilon_n}(x, y) d\rho(y) \\
&- (1 + O(\zeta_n)) \frac{1}{\sqrt{n}} \int \exp\left(\frac{1}{\varepsilon_n} f(x) + f(y)\right) \\
&\quad \times \left(a_n(\hat{f}(y) - f(y)) + O(a_n \varepsilon_n^{-1}(\hat{f} - f)^2)\right) k_{\varepsilon_n}(x, y) d\mathbb{G}_n(y) \\
&+ O(\varepsilon_n^{-1} a_n^{-1} \Xi_n(x)^2) \\
&= -(1 + O(\zeta_n)) K_{\varepsilon_n} [a_n(\hat{f} - f)] - (1 + O(\zeta_n)) O(K_{\varepsilon_n} [a_n \varepsilon_n^{-1} (\hat{f} - f)^2]) \\
&+ O(\varepsilon_n^{-1} a_n^{-1} \Xi_n(x)^2) \\
&- \frac{(1 + O(\zeta_n))}{\sqrt{n}} \int \exp\left(\frac{1}{\varepsilon_n} f(x) + f(y)\right) \\
&\quad \times \left(a_n(\hat{f}(y) - f(y)) + O(a_n \varepsilon_n^{-1}(\hat{f} - f)^2)\right) k_{\varepsilon_n}(x, y) d\mathbb{G}_n(y),
\end{aligned}$$

where we set

$$\zeta_n := \frac{1}{\sqrt{n}} \int \pi_{\varepsilon_n}(x, y) d\mathbb{G}_n(y).$$

We will now show that $\zeta_n = o(1)$. Using Proposition 1, we get

$$\zeta_n = \frac{1}{\sqrt{n\rho(x)}} \int \frac{k_{\varepsilon_n}(x, y)}{(2\pi\varepsilon_n)^{d/2} \sqrt{\rho(y)}} d\mathbb{G}_n(y) + o(1),$$

which converges to zero if $\sqrt{n}\varepsilon_n^{d/4} \rightarrow \infty$, as

$$\int \left(\frac{k_{\varepsilon_n}(x, y)}{(2\pi\varepsilon_n)^{d/4} \sqrt{\rho(y)}(1 + o(1))} \right)^2 d\rho(y) \rightarrow c,$$

for some constant c . The term

$$\frac{1}{\sqrt{n}} \int \exp\left(\frac{1}{\varepsilon_n} f(x) + f(y)\right) \left(a_n(\hat{f}(y) - f(y)) + o_p(1)\right) k_{\varepsilon_n}(x, y) d\mathbb{G}_n(y)$$

is also negligible. It can be rewritten

$$R(x) := \frac{a_n \varepsilon_n}{\sqrt{n} \varepsilon_n^{d/4}} \int \frac{k_{\varepsilon_n}(x, y)}{(2\pi)^{d/2} \varepsilon_n^{d/4} \sqrt{\rho(x)\rho(y)}(1 + o(1))} \frac{(\hat{f}(y) - f(y)) + o_p(1)}{\varepsilon_n} d\mathbb{G}_n(y).$$

Now, remark that one will choose $a_n \varepsilon_n / (\sqrt{n} \varepsilon_n^{d/4}) = 1$ for $C(x)$ to admit a limit. Thus, owing to Assumption **(Conv)**, as

$$\int \frac{k_{\varepsilon_n}(x, y)}{(2\pi)^{d/2} \varepsilon_n^{d/4} \sqrt{\rho(x)\rho(y)}(1 + o(1))} d\mathbb{G}_n(y)$$

converges to a Gaussian random variable, $R(x)$ converges to 0 in probability. \square

B.3. Proof of Proposition 6

Proof of Proposition 6. By Pooladian and Niles-Weed (2021, Proposition 2), we can express the gradient of the potentials as barycentric projections, so that

$$\begin{aligned} & a_n \varepsilon_n \left(\nabla f_{\varepsilon_n}(x) - \nabla \hat{f}_{\varepsilon_n}(x) \right) \\ &= a_n \varepsilon_n \int y d\pi_{\varepsilon_n}^x(y) - \int y d\hat{\pi}_{\varepsilon_n}^x(y) \\ &= -a_n \varepsilon_n \int y \exp\left(\frac{1}{\varepsilon_n} \left(f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y) - \frac{1}{2} \|x - y\|^2 \right)\right) d(\hat{\rho}_n - \rho)(y) \\ &+ a_n \varepsilon_n \int y k_{\varepsilon_n}(x, y) \\ &\quad \times \left(\exp\left(\frac{1}{\varepsilon_n} (f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y))\right) - \exp\left(\frac{1}{\varepsilon_n} (\hat{f}_{\varepsilon_n}(x) + \hat{f}_{\varepsilon_n}(y))\right) \right) d\hat{\rho}(y) \\ &=: A + B. \end{aligned}$$

A is dealt with using the expansion for the potentials and we get,

$$A = -\frac{a_n \varepsilon_n}{\sqrt{n}} \int y \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y - x\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(y) \rho^{1/2}(x) (1 + o(1))} d\mathbb{G}_n(y).$$

Let us now turn to B . It holds that

$$\begin{aligned}
 B(x) &= \\
 & a_n \varepsilon_n \int y k_{\varepsilon_n}(x, y) \left(\exp\left(\frac{1}{\varepsilon_n}(f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y))\right) - \exp\left(\frac{1}{\varepsilon_n}(\hat{f}_{\varepsilon_n}(x) + \hat{f}_{\varepsilon_n}(y))\right) \right) d\hat{\rho}(y) \\
 &= a_n \varepsilon_n \int y k_{\varepsilon_n}(x, y) \exp\left(\frac{f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y)}{\varepsilon_n}\right) \\
 &\quad \times \left(1 - \exp\left(\frac{a_n}{\varepsilon_n a_n}(\hat{f}_{\varepsilon_n}(x) + \hat{f}_{\varepsilon_n}(y) - f_{\varepsilon_n}(x) - f_{\varepsilon_n}(y))\right)\right) d\hat{\rho}(y) \\
 &= a_n \varepsilon_n \int y k_{\varepsilon_n}(x, y) \exp\left(\frac{1}{\varepsilon_n}(f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y))\right) \\
 &\quad \times \left(\frac{-a_n}{\varepsilon_n a_n}(\hat{f}_{\varepsilon_n}(x) + \hat{f}_{\varepsilon_n}(y) - f_{\varepsilon_n}(x) - f_{\varepsilon_n}(y))\right) d\hat{\rho}(y) \\
 &+ o_p(1) \quad \text{(owing to **(Conv)**)} \\
 &= -a_n (\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x)) \int y k_{\varepsilon_n}(x, y) \exp\left(\frac{1}{\varepsilon_n}(f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y))\right) d\hat{\rho}(y) + o_p(1) \\
 &+ \int y k_{\varepsilon_n}(x, y) \exp\left(\frac{1}{\varepsilon_n}(f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y))\right) \left(-a_n(\hat{f}_{\varepsilon_n}(y) - f_{\varepsilon_n}(y))\right) d\hat{\rho}(y) \\
 &= -a_n (\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x)) K_{\varepsilon_n}[x] - K_{\varepsilon_n}\left[x a_n (\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x))\right] + o_p(1) \\
 &- \frac{a_n}{\sqrt{n}} (\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x)) \int y k_{\varepsilon_n}(x, y) \exp\left(\frac{1}{\varepsilon_n}(f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y))\right) d\mathbb{G}_n(y) \\
 &+ \int y k_{\varepsilon_n}(x, y) \exp\left(\frac{1}{\varepsilon_n}(f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y))\right) \left(-\frac{a_n}{\sqrt{n}}(\hat{f}_{\varepsilon_n}(y) - f_{\varepsilon_n}(y))\right) d\mathbb{G}_n(y).
 \end{aligned}$$

For the two last summands, notice first that while a_n is chosen such that

$$-\frac{a_n \varepsilon_n}{\sqrt{n}} \int y k_{\varepsilon_n}(x, y) \exp\left(\frac{1}{\varepsilon_n}(f_{\varepsilon_n}(x) + f_{\varepsilon_n}(y))\right) d\mathbb{G}_n(y)$$

converges to a random variable, while $\varepsilon_n^{-1}(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x)) \rightarrow 0$ as in the proof of Proposition 5. \square

B.4. Proof of Theorem 7

Proof of Theorem 7. As announced, the proof builds on the linearization in Proposition 5.

Step 1. As it shall become important in the sequel, we first show limiting distribution results for quantities of the type

$$\left(\int_{\mathbb{R}^d} \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y - x_j\|^2\right)}{\rho^{1/2}(y)} d\mathbb{G}_n(y) \right)_{j=1, \dots, m} \quad (18)$$

up to choosing an appropriate scaling. Then, we need to control the limiting behavior of $(\text{id} + K_{\varepsilon_n})^{-1}$ when the latter acts on a particular sequence of (ran-

dom) functions.

To obtain the convergence in distribution of (18) after proper rescaling, we use the Lindeberg–Feller theorem (Van der Vaart, 2000, Proposition 2.27). One remarks that

$$\rho \left[\varepsilon_n^{-d/4} \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y-x\|^2\right)}{\rho^{1/2}(y)} \right] < \infty$$

as well as

$$\begin{aligned} \rho \left[\left(\varepsilon_n^{-d/4} \exp\left(-\frac{1}{2\varepsilon_n} \|y-x\|^2\right) \rho^{-1/2}(y) \right)^2 \right] &= \int_{\mathbb{R}^d} \varepsilon_n^{-d/2} \exp\left(-\frac{1}{\varepsilon_n} \|y-x\|^2\right) dy \\ &= \pi^{d/2}. \end{aligned}$$

Further,

$$\begin{aligned} &\rho \left[\varepsilon_n^{-d/2} \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y-x_1\|^2\right) \exp\left(-\frac{1}{2\varepsilon_n} \|y-x_2\|^2\right)}{\rho(y)} \right] \\ &= \exp\left(-\frac{1}{4\varepsilon_n} \|x_1-x_2\|^2\right) \int_{\mathbb{R}^d} \varepsilon_n^{-d/2} \exp\left(-\frac{1}{\varepsilon_n} \|y-\frac{x_1+x_2}{2}\|^2\right) dy \\ &= \exp\left(-\frac{1}{4\varepsilon_n} \|x_1-x_2\|^2\right) \pi^{d/2}. \end{aligned}$$

It is then clear that the limiting covariance matrix (as $\varepsilon_n \rightarrow 0$) is well behaved. Setting

$$Y_{n,i} := \left(\varepsilon_n^{-d/4} \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|X_i-x_1\|^2\right)}{\sqrt{n}\rho^{1/2}(X_i)}, \dots, \varepsilon_n^{-d/4} \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|X_i-x_m\|^2\right)}{\sqrt{n}\rho^{1/2}(X_i)} \right)^\top,$$

it is then a routine task to check that

$$\sum_{i=1}^n \mathbb{E} \|Y_{n,i}\|^2 \mathbf{1}\{\|Y_{n,i}\| > \delta\} \rightarrow 0, \text{ for every } \delta > 0.$$

because of the assumption $\sqrt{n}\varepsilon_n^{d/4} \sqrt{\log n} \rightarrow \infty$ as $n \rightarrow \infty$. Indeed,

$$\begin{aligned} &\sum_{i=1}^n \mathbb{E} \|Y_{n,i}\|^2 \mathbf{1}\{\|Y_{n,i}\| > \delta\} \\ &= n \int \sum_{j=1}^m \varepsilon_n^{-d/2} \frac{\exp\left(-\frac{1}{\varepsilon_n} \|y-x_1\|^2\right)}{n} \\ &\quad \times \mathbf{1} \left\{ \sum_{j=1}^m \exp\left(-\frac{1}{\varepsilon_n} \|y-x_1\|^2 + V(y)\right) > \delta^2 \varepsilon_n^{d/2} n \right\} dy. \end{aligned}$$

From this, we conclude that

$$\sum_{i=1}^n (Y_{n,i} - \mathbb{E}Y_{n,i}, \dots, Y_{n,m} - \mathbb{E}Y_{n,m})^\top \rightsquigarrow \mathcal{N}(0_m, \pi^{d/2} I_m),$$

which is the sought limit for the quantity in Equation (18) after rescaling by $\varepsilon_n^{-d/4}$.

Step 2. Even though the operator K_{ε_n} converges to the identity, one cannot simply replace it by its limit. We thus now consider the questions arising from the presence of this (sequence of) inverse operator(s). First remark that $(2+x)^{-1}$ is an analytic function of x in a neighbourhood of zero. Thus, one can use a series development for operators and get

$$(2 \text{id} + (K_{\varepsilon_n} - \text{id}))^{-1} = \frac{1}{2} \sum_{\kappa=0}^{\infty} (\text{id} - K_{\varepsilon_n})^\kappa 2^{-\kappa}$$

Looking at the above, we see that terms of the form

$$(\text{id} - K_{\varepsilon_n}) \left[\int \pi_{\varepsilon_n}(\cdot, y) d\mathbb{G}_n(y) \right] (x) =: A + B,$$

will play a crucial role. As such computations will occur over and over again, observe that

$$\begin{aligned} B &= K_{\varepsilon_n} \left[\int \pi_{\varepsilon_n}(\cdot, y) d\mathbb{G}_n(y) \right] (x) \\ &= \frac{1}{\rho^{1/2}(x)(1 + o_x(1))} \int \frac{e^{-\frac{1}{4\varepsilon_n} \|x-y\|^2}}{(4\pi\varepsilon_n)^{d/2} \rho^{1/2}(y)(1 + o_{y,z}(1))} \int \frac{e^{-\frac{1}{\varepsilon_n} \|x-z\|^2}}{(\pi\varepsilon_n)^{d/2}} dz d\mathbb{G}_n(y) \\ &= \frac{1}{\rho^{1/2}(x)(1 + o_x(1))} \int \frac{e^{-\frac{1}{4\varepsilon_n} \|x-y\|^2}}{(4\pi\varepsilon_n)^{d/2} \rho^{1/2}(y)(1 + o_y(1))} d\mathbb{G}_n(y) + o_p(1). \end{aligned}$$

Its limiting distribution for fixed x , after multiplication by $\varepsilon_n^{d/4}$, using the same argument as above is a Gaussian random variable with variance $2^{-3d/2} \pi^{-d/2} \rho^{-1}(x)$. We further recall that the integrals of the $o(1)$ terms are actually negligible owing to Assumption **(Reg)**.

In general,

$$\begin{aligned} &K_{\varepsilon_n}^\kappa [h](x_{\kappa+1}) \\ &= (2\pi\varepsilon_n)^{d\kappa/2} \rho^{-1/2}(x_{\kappa+1}) \\ &\times \int \cdots \int \exp \left(-\frac{1}{2\varepsilon_n} \sum_{j=2}^{\kappa+1} \|x_j - x_{j-1}\|^2 \right) \left(1 + \sum_{j=1}^{\kappa+1} o_{x_j}(1) \right) \rho^{1/2}(x_1) h(x_1) dx_1 \cdots dx_\kappa. \end{aligned}$$

This multiple integral is a series of Gaussian convolution except for the integral

with respect to x_1 . We thus have, up to a negligible term,

$$\begin{aligned} & K_{\varepsilon_n}^\kappa[h](x_{\kappa+1}) \\ &= (2\kappa\pi\varepsilon_n)^{-d/2}\rho^{-1/2}(x_{\kappa+1}) \int \exp\left(-\frac{1}{2\kappa\varepsilon_n}\|x_{\kappa+1}-x_1\|^2\right)\rho^{1/2}(x_1)h(x_1)dx_1. \end{aligned} \tag{19}$$

Now, for any fixed J ,

$$\begin{aligned} & \mathbb{E}\left[\left(\sum_{\kappa=0}^J 2^{-\kappa}(\text{id}-K_{\varepsilon_n})^\kappa[h_n]\right)^2\right] \\ &= \sum_{0\leq\kappa,\kappa'\leq J} 2^{-\kappa-\kappa'}\mathbb{E}\left[(\text{id}-K_{\varepsilon_n})^\kappa[h_n](\text{id}-K_{\varepsilon_n})^{\kappa'}[h_n]\right] \\ &= \sum_{0\leq\kappa,\kappa'\leq J} 2^{-\kappa-\kappa'}\mathbb{E}\left[\sum_{\eta=0}^{\kappa}\binom{\kappa}{\eta}(-K_{\varepsilon_n})^\eta[h_n]\sum_{\eta'=0}^{\kappa'}\binom{\kappa'}{\eta'}(-K_{\varepsilon_n})^{\eta'}[h_n]\right] \\ &= \sum_{0\leq\kappa,\kappa'\leq J} 2^{-\kappa-\kappa'}\sum_{\eta=0}^{\kappa}\sum_{\eta'=0}^{\kappa'}\binom{\kappa}{\eta}(-1)^\eta(-1)^{\eta'}\binom{\kappa'}{\eta'}\mathbb{E}\left[K_{\varepsilon_n}^\eta[h_n]K_{\varepsilon_n}^{\eta'}[h_n]\right] \end{aligned}$$

By a direct computation similar to those above, we get

$$\mathbb{E}\left[K_{\varepsilon_n}^\eta[h_n]K_{\varepsilon_n}^{\eta'}[h_n]\right]=\rho^{-1}(\cdot)(2\pi\varepsilon_n)^{-d/2}(\eta+\eta'+2)^{-d/2}(1+o(1)).$$

By the development above, setting

$$S_J(n):=\varepsilon_n^{d/4}\sum_{\kappa=0}^J 2^{-\kappa}(\text{id}-K_{\varepsilon_n})^\kappa[h_n](x),$$

one notices readily that $(S_J(n))_{J\in\mathbb{N}}$ is Cauchy in L^2 with constants that can be made uniform in n, ε_n — recall that x is fixed. For each fixed J , the distributional limit of the truncated series exists as a sum of finitely many random functions. Denote by \mathfrak{d} any distance metrizing convergence in distribution and by X^J the limit as $n\rightarrow\infty$ of the truncated series $S_J(n)$. X_J has a Gaussian distribution, one can thus add to it an independent Gaussian random variable Y such that X^J+Y has the Gaussian distribution which is the limiting distribution in Theorem 4; call a random variable with that distribution X^∞ . Therefore, by the triangle inequality, it holds

$$\mathfrak{d}(S_\infty(n), X^\infty)\leq\mathfrak{d}(X^\infty, X^J)+\mathfrak{d}(S_J(n), X^J)+\mathfrak{d}(S_J(n), S_\infty(n)).$$

As convergence in L^2 implies convergence in distribution, for all $\varepsilon>0$, there exists J such that

$$\mathfrak{d}(S_\infty(n), X^\infty)\leq\varepsilon/3+\mathfrak{d}(S_J(n), X^J)+\varepsilon/3.$$

Now, for n sufficiently large, the middle term in the above display can be made smaller than $\varepsilon/3$. Finally, recall the exact form of the asymptotically negligible terms in the statement of Proposition 5 obtained in the proof of that Proposition. As these terms also involve

$$\frac{k_{\varepsilon_n}(x, y)}{(2\pi\varepsilon_n)^{d/4}\sqrt{\rho(x)(1 + o(1))}},$$

applying $(\text{id} + K_{\varepsilon_n})^{-1}$ to these terms is valid and they remain asymptotically negligible after applying that operator. \square

Lemma 23. Consider class of function

$$\mathcal{F} := \left\{ \varepsilon^{d/4} \frac{\exp\left(-\frac{1}{2\varepsilon k} \|x - \cdot\|^2\right)}{(2\pi\varepsilon k)^{d/2} \rho^{1/2}(\cdot)}; \varepsilon > 0, k \in \mathbb{N} \right\}.$$

It is such that for $f_{\varepsilon, k} \in \mathcal{F}$, it holds

$$\mathbb{P} f_{\varepsilon, k}^2 = \int \frac{\exp\left(-\frac{1}{k\varepsilon} \|x - y\|^2\right)}{(2\pi k)^d \varepsilon^{d/2}} dy = 2^{-d} (\pi k)^{-d/2},$$

irrespective of ε .

The proof of the above lemma is a mere computation.

B.5. Proof of Theorem 9 and Corollary 10

Proof of Corollary 10. The proof directly follows from Theorem 9, once we establish that Proposition 1 still holds after taking the logarithm and the gradient. The proof follows the same spirit as that of Proposition 1 and we thus leave it to the reader. \square

Proof of Theorem 9. The proof relies on the representation derived from Proposition 6. The proof is split into three parts.

Step 1. We start by stating the limit of

$$A_1(x) := \frac{a_n \varepsilon_n}{\sqrt{n}} \int y \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y - x\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(y) \rho^{1/2}(x) (1 + o(1))} d\mathbb{G}_n(y).$$

We will invoke the Lindeberg–Feller central limit theorem. Upon choosing the sequence $a_n \varepsilon_n^{1+d/4} / \sqrt{n}$ converging to a constant, we remark that

$$\int y y^\top \frac{\exp\left(-\frac{1}{\varepsilon_n} \|y - x\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho(y) \rho(x) (1 + o(1))} d\rho(y) = \rho^{-1}(x) 2^{-d} \pi^{-d/2} x x^\top + o(1),$$

which is the first condition pertaining to the variance. The tail condition also holds along the lines of the argument in the Proof of Theorem 7. This yields, choosing a_n as above,

$$\int y \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y - x\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(y) \rho^{1/2}(x) (1 + o(1))} d\mathbb{G}_n(y) \rightsquigarrow \mathcal{N}(0_d, \rho^{-1}(x) 2^{-d} \pi^{-d/2} x x^\top)$$

Step 2. We now study the term

$$A_2(x) := K_{\varepsilon_n} \left[\text{id } a_n \left(\hat{f}_{\varepsilon_n}(\cdot) - f_{\varepsilon_n}(\cdot) \right) \right] (x)$$

Observe that

$$\begin{aligned} A_2(x) &= -\frac{a_n \varepsilon_n}{\sqrt{n}} \int y \pi_{\varepsilon_n}(x, y) \\ &\quad (\text{id} + K_{\varepsilon_n})^{-1} \left[\int \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|\cdot - z\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(z) \rho^{1/2}(\cdot) (1 + o(1))} d\mathbb{G}(z) \right] (y) d\rho(y) \\ &= -\frac{a_n \varepsilon_n}{\sqrt{n}} \int (\text{id} + K_{\varepsilon_n})^{-1} \left[\text{id} \times \pi_{\varepsilon_n}(x, \cdot) \right] (y) \\ &\quad \int \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y - z\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(z) \rho^{1/2}(y) (1 + o(1))} d\mathbb{G}(z) d\rho(y), \end{aligned}$$

as for each ε_n , $(\text{id} + K_{\varepsilon_n})^{-1}$ is a self-adjoint operator from $L^2(\rho)$ to $L^2(\rho)$. Also, let us observe that

$$\begin{aligned} K_{\varepsilon_n} \left[\text{id} \times \pi_{\varepsilon_n}(x, \cdot) \right] (y) &= \int z \pi_{\varepsilon_n}(y, z) \pi_{\varepsilon_n}(x, z) \rho(z) dz \\ &= \frac{1}{(4\pi\varepsilon_n)^{d/2} \rho^{1/2}(x) \rho^{1/2}(y)} \int z \frac{1}{(\pi\varepsilon_n)^{d/2} (1 + o(1))} \exp\left(-\frac{\|x - z\|^2 + \|y - z\|^2}{2\varepsilon_n}\right) dz \\ &= \frac{\exp\left(-\frac{1}{4\varepsilon_n} \|x - y\|^2\right)}{(4\pi\varepsilon_n)^{d/2} \rho^{1/2}(x) \rho^{1/2}(y)} \int z \frac{1}{(\pi\varepsilon_n)^{d/2} (1 + o(1))} \exp\left(-\frac{1}{\varepsilon_n} \left\|z - \frac{x + y}{2}\right\|^2\right) dz \\ &= \frac{\exp\left(-\frac{1}{4\varepsilon_n} \|x - y\|^2\right)}{(4\pi\varepsilon_n)^{d/2} \rho^{1/2}(x) \rho^{1/2}(y)} \frac{x + y}{2} (1 + o(1)). \end{aligned}$$

In full generality, for $\kappa > 1$,

$$\begin{aligned}
& K_{\varepsilon_n}^\kappa \left[\text{id} \times \pi_{\varepsilon_n}(x, \cdot) \right] (y) \\
&= K_{\varepsilon_n}^{\kappa-1} \left[\frac{\exp\left(-\frac{1}{4\varepsilon_n} \|\cdot - x\|^2\right)}{(4\pi\varepsilon_n)^{d/2} \rho^{1/2}(\cdot) \rho^{1/2}(x)} \cdot + x (1 + o(1)) \right] (y) \\
&= \frac{\rho^{-1/2}(x) \rho^{-1/2}(y)}{(2\pi(\kappa-1)\varepsilon_n)^{d/2} (4\pi\varepsilon_n)^{d/2}} \int \frac{z+x}{2(1+o(1))} \exp\left(-\frac{\|x-z\|^2}{4\varepsilon_n} - \frac{\|y-z\|^2}{2(\kappa-1)\varepsilon_n}\right) dz \\
&= \frac{\exp\left(-\frac{1}{2\varepsilon_n(\kappa-1)} \|x-y\|^2\right)}{(2\pi(\kappa-1)\varepsilon_n)^{d/2} (4\pi\varepsilon_n)^{d/2} \rho^{1/2}(x) \rho^{1/2}(y)} \\
&\quad \times \int \frac{z+x}{2(1+o(1))} \exp\left(-\frac{\kappa+1}{4\varepsilon_n(\kappa-1)} \left\| z - \frac{(\kappa-1)x+2y}{\kappa+1} \right\|^2\right) dz \\
&= \frac{\exp\left(-\frac{1}{2\varepsilon_n(\kappa+1)} \|x-y\|^2\right)}{(4\pi\varepsilon_n)^{d/2} \rho^{1/2}(x) \rho^{1/2}(y)} \left(\frac{2}{\kappa+1}\right)^{d/2} \left(\frac{x}{2} + \frac{(\kappa-1)x+2y}{2(\kappa+1)}\right) (1+o(1)) \\
&= \frac{\exp\left(-\frac{1}{2\varepsilon_n(\kappa+1)} \|x-y\|^2\right)}{(4\pi\varepsilon_n)^{d/2} \rho^{1/2}(x) \rho^{1/2}(y)} \left(\frac{2}{\kappa+1}\right)^{d/2} \left(\frac{x}{2} + \frac{(\kappa-1)x+2y}{2(\kappa+1)}\right) (1+o(1)) \\
&= \frac{\exp\left(-\frac{1}{2\varepsilon_n(\kappa+1)} \|x-y\|^2\right)}{(2\pi(\kappa+1)\varepsilon_n)^{d/2} \rho^{1/2}(x) \rho^{1/2}(y)} \left(x + \frac{y-x}{\kappa+1}\right) (1+o(1))
\end{aligned}$$

where the second equality follows from (19).

As above,

$$(\text{id} + K_{\varepsilon_n})^{-1} \left[\text{id} \times \pi_{\varepsilon_n}(x, \cdot) \right] (y) = \frac{1}{2} \sum_{\kappa=0}^{\infty} 2^{-\kappa} \sum_{\eta=0}^{\kappa} \binom{\kappa}{\eta} (-K_{\varepsilon_n})^\eta \left[\text{id} \times \pi_{\varepsilon_n}(x, \cdot) \right] (y),$$

so that

$$\begin{aligned}
& A_2(x) \\
&= \frac{1}{2} \sum_{\kappa=0}^{\infty} 2^{-\kappa} \sum_{\eta=0}^{\kappa} \binom{\kappa}{\eta} (-1)^\eta \iint K_{\varepsilon_n}^\eta [\text{id} \times \pi_{\varepsilon_n}(x, \cdot)](y) \\
&\quad \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y - z\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(z) \rho^{1/2}(y)} d\mathbb{G}(z) d\rho(y) \\
&= \frac{1}{2} \sum_{\kappa=0}^{\infty} 2^{-\kappa} \sum_{\eta=0}^{\kappa} \binom{\kappa}{\eta} (-1)^\eta \\
&\quad \times \iint \frac{\exp\left(-\frac{1}{2\varepsilon_n(\eta+1)} \|x - y\|^2\right)}{(2\pi(\eta+1)\varepsilon_n)^{d/2} \rho^{1/2}(x) \rho^{1/2}(y)} \left(x + \frac{y-x}{\eta+1}\right) \\
&\quad \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y - z\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(z) \rho^{1/2}(y)} d\mathbb{G}(z) d\rho(y) \\
&= \frac{1}{2} \sum_{\kappa=0}^{\infty} 2^{-\kappa} \sum_{\eta=0}^{\kappa} \binom{\kappa}{\eta} (-1)^\eta \\
&\quad \times \iint \frac{\exp\left(-\frac{1}{2\varepsilon_n(\eta+1)} \|x - y\|^2\right)}{(2\pi(\eta+1)\varepsilon_n)^{d/2} \rho^{1/2}(x)} \left(x + \frac{y-x}{\eta+1}\right) \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y - z\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(z)} d\mathbb{G}(z) dy.
\end{aligned}$$

Finally, using the same type of arguments as before, we get

$$\begin{aligned}
& A_2(x) \\
&= \frac{1}{2} \sum_{\kappa=0}^{\infty} 2^{-\kappa} \sum_{\eta=0}^{\kappa} \binom{\kappa}{\eta} (-1)^\eta \\
&\quad \times \int \frac{\exp\left(-\frac{1}{2\varepsilon_n(\eta+z)} \|x - z\|^2\right)}{(2\pi(\eta+1)\varepsilon_n)^{d/2} \rho^{1/2}(x) \rho^{1/2}(z)} \left(x + \frac{x}{(\eta+1)(\eta+2)} - \frac{z(\eta+1)}{\eta+2}\right) d\mathbb{G}(z) dy.
\end{aligned}$$

Step 3. It remains to combine the elements. Setting

$$A_3(x) = a_n \left(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x) \right) K_{\varepsilon_n}[\text{id}](x),$$

both steps above along with Theorem 7, guarantee the joint convergence, for fixed $x \in \mathbb{R}^d$, of the random vector $(A_1(x), A_2(x), A_3(x))^\top$ to a multivariate, centered Gaussian distribution. As

$$\sqrt{n} \varepsilon_n^{d/4} \left(\nabla \hat{f}_{\varepsilon_n} - \nabla f_{\varepsilon_n} \right)$$

is asymptotically equivalent to a linear combination of $(A_1(x), A_2(x), A_3(x))$, the claim follows. \square

B.6. Proof of Propositions 13 and 14

Proof of Proposition 13. From Proposition 1, the following holds after a simple rearrangement

$$\begin{aligned} & -2f_\varepsilon(x)/\varepsilon - \log \rho(x) \\ &= \frac{d}{2} \log(2\pi\varepsilon) + \varepsilon \operatorname{tr} \left(\frac{\nabla^2 \rho(x)}{4\rho(x)} - \frac{1}{8} (\nabla \log \rho(x)) (\nabla \log \rho(x))^\top \right) + o(\varepsilon), \end{aligned}$$

Then, remark that

$$\frac{\nabla^2 \rho(x)}{\rho(x)} = \exp(V) \nabla^2 (\exp(-V)) = -\nabla^2 V + (\nabla V)(\nabla V)^\top.$$

Putting everything together, we have

$$-2f_\varepsilon(x)/\varepsilon - \log \rho(x) = \frac{d}{2} \log(2\pi\varepsilon) + \frac{\varepsilon}{4} (-\Delta V(x) + \frac{1}{2} \|\nabla V(x)\|^2) + o(\varepsilon).$$

Taking the gradient on both sides in the display above yields

$$s_\varepsilon(x) - \nabla \log \rho(x) = \frac{\varepsilon}{4} (-\nabla \Delta V(x) + \nabla^2 V(x) \nabla V(x)) + o(\varepsilon),$$

which follows from the chain rule. Thus, for ε small enough, we arrive at

$$\begin{aligned} & \|s_\varepsilon - \nabla \log \rho\|_{L^2(\rho)}^2 \\ & \lesssim \varepsilon^2 \left(\int \|\nabla \Delta V(x)\|^2 d\rho(x) + \int \|\nabla^2 V(x) \nabla \log \rho(x)\|^2 d\rho(x) \right). \end{aligned}$$

The claim then follows from Assumption (C). □

Proof of Proposition 14. By Proposition 6, we have that

$$\begin{aligned} \mathbb{E} \|\hat{s}_{\varepsilon_n} - s_{\varepsilon_n}\|_{L^2(\rho)}^2 & \lesssim \mathbb{E} \int \left\| \frac{1}{\varepsilon_n \sqrt{n}} \int y \frac{\exp\left(-\frac{1}{2\varepsilon_n} \|y-x\|^2\right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(y) (1+o(1))} d\mathbb{G}_n(y) \right\|^2 dx \\ & \quad + \mathbb{E} \left\| K_{\varepsilon_n}[x] \varepsilon_n^{-2} \left(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x) \right) \right\|_{L^2(\rho)}^2 \\ & \quad + \mathbb{E} \left\| K_{\varepsilon_n} \left[x \varepsilon_n^{-2} \left(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x) \right) \right] \right\|_{L^2(\rho)}^2, \end{aligned}$$

using the basic inequality $(a+b+c)^2 \leq 3(a^2+b^2+c^2)$. Let us focus the first term. Rewriting, we have

$$\begin{aligned} I_1 & := n^{-1} \varepsilon_n^{-d-2} \\ & \times \int \left\| \int y \exp\left(-\frac{1}{2\varepsilon_n} \|x-y\|^2 + \frac{1}{2}V(x) + \frac{1}{2}V(y)\right) (1+o(1)) d\mathbb{G}_n(y) \right\|^2 d\rho(x). \end{aligned}$$

One can thus fix an arbitrary compact $K \ni 0$ containing the product of the supports and get that

$$I_1 \asymp n^{-1} \varepsilon_n^{-d-2} \int \left\| \int \mathbb{1}_{(x,y) \in Ky} \exp \left(-\frac{1}{2\varepsilon_n} \|x-y\|^2 + \frac{1}{2}V(x) + \frac{1}{2}V(y) \right) (1 + o(1)) d\mathbb{G}_n(y) \right\|^2 \rho(x) dx.$$

On that compact K ,

$$y \exp \left(-\frac{1}{2\varepsilon_n} \|x-y\|^2 + \frac{1}{2}V(x) + \frac{1}{2}V(y) \right) \rightarrow \frac{x}{\rho(x)},$$

uniformly in x, y as $\varepsilon_n \rightarrow 0$. Thus,

$$\int \left\| \int y \exp \left(-\frac{1}{2\varepsilon_n} \|x-y\|^2 + \frac{1}{2}V(x) + \frac{1}{2}V(y) \right) d\mathbb{G}_n(y) \right\|^2 \rho(x) dx = O_p(1).$$

Recalling that, by Proposition 5,

$$\varepsilon_n^{-1} (\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x)) \asymp \frac{1}{2\sqrt{n}} \left[\int \frac{\exp \left(-\frac{1}{2\varepsilon_n} \|y-x\|^2 \right)}{(2\pi\varepsilon_n)^{d/2} \rho^{1/2}(y) \rho^{1/2}(x) (1 + o(1))} d\mathbb{G}_n(y) \right],$$

we have

$$\begin{aligned} & \left\| K_{\varepsilon_n}[x] \varepsilon_n^{-2} \left(\hat{f}_{\varepsilon_n}(x) - f_{\varepsilon_n}(x) \right) \right\|_{L^2(\rho)}^2 \\ & \asymp n^{-1} \varepsilon_n^{-d-2} \int \|x\|^2 \left(\int \frac{\exp \left(-\frac{1}{2\varepsilon_n} \|y-x\|^2 \right)}{\rho^{1/2}(y) \rho^{1/2}(x)} d\mathbb{G}_n(y) \right)^2 \rho(x) dx, \end{aligned}$$

and similarly for the third term. Both terms can be treated in the exact same way as above. \square

B.7. Proof of Proposition 16

Proof of Proposition 16. By symmetry, Assumption **(Lin)** yields

$$\begin{pmatrix} \text{id} & K_{\varepsilon_n}^\nu \\ K_{\varepsilon_n}^\mu & \text{id} \end{pmatrix} \begin{pmatrix} \hat{f}_{\varepsilon_n} - f_{\varepsilon_n} \\ \hat{g}_{\varepsilon_n} - g_{\varepsilon_n} \end{pmatrix} = -\frac{\varepsilon_n}{\sqrt{n}} \begin{pmatrix} \int \pi_{\varepsilon_n}(\cdot, y) d\mathbb{G}_n^\nu(y) \\ \int \pi_{\varepsilon_n}(x, \cdot) d\mathbb{G}_n^\mu(x) \end{pmatrix} + o_p(a_n^{-1}). \quad (20)$$

Using the formula for inversion of block operators

$$\begin{pmatrix} \text{id} & K_{\varepsilon_n}^\nu \\ K_{\varepsilon_n}^\mu & \text{id} \end{pmatrix}^{-1} = \begin{pmatrix} \text{id} + K_{\varepsilon_n}^\nu (\text{id} - K_{\varepsilon_n}^\mu K_{\varepsilon_n}^\nu)^{-1} K_{\varepsilon_n}^\mu & -K_{\varepsilon_n}^\nu (\text{id} - K_{\varepsilon_n}^\mu K_{\varepsilon_n}^\nu)^{-1} \\ -(\text{id} - K_{\varepsilon_n}^\mu K_{\varepsilon_n}^\nu)^{-1} K_{\varepsilon_n}^\mu & (\text{id} - K_{\varepsilon_n}^\mu K_{\varepsilon_n}^\nu)^{-1} \end{pmatrix}.$$

We recall that the inverse operators appearing on the right hand side above exist for each $\varepsilon > 0$ as stated in [Carlier and Laborde \(2020\)](#); [González-Sanz et al.](#)

(2022); [González-Sanz and Hundrieser \(2023\)](#). However, note that the constant function 1 is an eigenfunction with eigenvalue 1 of both $K_{\varepsilon_n}^\nu$ and $K_{\varepsilon_n}^\mu$, as

$$\int \pi_{\varepsilon_n}(\cdot, y) d\nu(y) = 1 = \int \pi_{\varepsilon_n}(x, \cdot) d\mu(x).$$

Thus, invertibility only holds for appropriate, but nonconstant functions, which is the case for the quantities on the right hand side of Equation (20).

For each $\varepsilon_n > 0$, we have a Neumann series expansion of the inverse on the set of non-constant functions. It indeed holds

$$(\text{id} - K_{\varepsilon_n}^\mu K_{\varepsilon_n}^\nu)^{-1} = \sum_{\kappa=0}^{\infty} (K_{\varepsilon_n}^\mu K_{\varepsilon_n}^\nu)^\kappa.$$

In the view of this, we get the simplification

$$\begin{pmatrix} \text{id} & K_{\varepsilon_n}^\nu \\ K_{\varepsilon_n}^\mu & \text{id} \end{pmatrix}^{-1} = \begin{pmatrix} (\text{id} - K_{\varepsilon_n}^\nu K_{\varepsilon_n}^\mu)^{-1} & -K_{\varepsilon_n}^\nu (\text{id} - K_{\varepsilon_n}^\mu K_{\varepsilon_n}^\nu)^{-1} \\ -(\text{id} - K_{\varepsilon_n}^\mu K_{\varepsilon_n}^\nu)^{-1} K_{\varepsilon_n}^\mu & (\text{id} - K_{\varepsilon_n}^\mu K_{\varepsilon_n}^\nu)^{-1} \end{pmatrix}. \quad \square$$

B.8. Proof of Theorem 19

Proof of Theorem 19. We start with preliminary developments helping towards proving the claim. We omit the negligible terms to simplify the exposition; it is however licit because of the assumptions made. Let us use the previous expansions and the Laplace method to get

$$\begin{aligned} & K_{\varepsilon_n}^\nu[h](x) \\ &= \int \exp\left(\frac{1}{\varepsilon_n} [f_{\varepsilon_n}(x) + g_{\varepsilon_n}(y) - \frac{1}{2}\|x - y\|^2]\right) h(y) \nu(y) dy \\ &= \int \frac{1}{(2\pi\varepsilon_n)^{d/2}} \exp\left(\frac{1}{\varepsilon_n} [f_0(x) + g_0(y) - \frac{1}{2}\|x - y\|^2] + o(1)\right) \\ &\quad \times h(y) \nu(y)^{-1/2} \mu^{-1/2}(x) \nu(y) dy \\ &= \int \frac{1}{(2\pi\varepsilon_n)^{d/2}} \exp\left(-\frac{1}{2\varepsilon_n} [(y - x^*)^\top [\nabla^2 \varphi_0(x)]^{-1} (y - x^*) + o(1)] + o(1)\right) \\ &\quad \times h(y) \nu(y)^{-1/2} \mu^{-1/2}(x) \nu(y) dy \\ &= (\det[\nabla^2 \varphi_0(x)])^{1/2} h(x^*) \nu(x^*)^{-1/2} \mu^{-1/2}(x) \nu(x^*) (1 + o(1)) \\ &= \frac{h(x^*) \nu(x^*)}{\nu(x^*)} (1 + o(1)) \\ &= h(x^*) (1 + o(1)), \end{aligned}$$

where we recall the Monge–Ampère equation $\mu(x) = \nu(x^*) \det[\nabla^2 \varphi_0(x)]$.

Also, by symmetry,

$$\begin{aligned}
& K_{\varepsilon_n}^\mu [h](y) \\
&= \int \exp\left(\frac{1}{\varepsilon_n}[f_{\varepsilon_n}(x) + g_{\varepsilon_n}(y) - \frac{1}{2}\|x - y\|^2]\right) h(x)\mu(x)dx \\
&\asymp \int \frac{1}{(2\pi\varepsilon_n)^{d/2}} \exp\left(\frac{1}{\varepsilon_n}[f_0(x) + g_0(y) - \frac{1}{2}\|x - y\|^2]\right) \nu(y)^{-1/2}\mu^{-1/2}(x)h(x)\mu(x)dx \\
&\asymp \int \frac{1}{(2\pi\varepsilon_n)^{d/2}} \exp\left(-\frac{1}{2\varepsilon_n}(x - \nabla\varphi_0^*(y))^\top [\nabla^2\varphi_0^*(y)]^{-1}(x - \nabla\varphi_0^*(y))\right) \\
&\quad \times \nu(y)^{-1/2}\mu^{1/2}(x)h(x)dx.
\end{aligned}$$

We now can turn to the main claim.

$$\begin{aligned}
& K_{\varepsilon_n}^\mu \left[K_{\varepsilon_n}^\nu [h] \right] (y_2) \\
&\asymp \frac{1}{\nu^{1/2}(y_2)} \iint \frac{1}{(2\pi\varepsilon_n)^d} \\
&\quad \times \exp\left(-\frac{1}{2\varepsilon_n}(x - \nabla\varphi_0^*(y_2))^\top [\nabla^2\varphi_0^*(y_2)]^{-1}(x - \nabla\varphi_0^*(y_2))\right) \\
&\quad \times \exp\left(-\frac{1}{2\varepsilon_n}(x - \nabla\varphi_0^*(y))^\top [\nabla^2\varphi_0^*(y)]^{-1}(x - \nabla\varphi_0^*(y))\right) dx h(y)\nu^{1/2}(y)dy \\
&\asymp \frac{1}{\nu^{1/2}(y_2)} \int \frac{1}{(2\pi\varepsilon_n)^d} \sqrt{\frac{\det[2\pi\varepsilon_n\nabla^2\varphi_0^*(y_2)] \det[2\pi\varepsilon_n\nabla^2\varphi_0^*(y)]}{\det[2\pi\varepsilon_n\nabla^2\varphi_0^*(y_2) + \nabla^2\varphi_0^*(y)]}} h(y) \\
&\quad \times \exp\left(-\frac{1}{2\varepsilon_n}(\nabla\varphi_0^*(y) - \nabla\varphi_0^*(y_2))^\top [\nabla^2\varphi_0^*(y_2) + \nabla^2\varphi_0^*(y)]^{-1}(\nabla\varphi_0^*(y) - \nabla\varphi_0^*(y_2))\right) \\
&\quad \nu^{1/2}(y)dy,
\end{aligned}$$

where the last equivalence follows from the fact that the inner integral is a convolution of two Gaussians, one with mean 0 and covariance matrix $\nabla^2\varphi_0^*(y_2)$ and the other one with mean $\nabla\varphi_0^*(y)$ and covariance matrix $\nabla^2\varphi_0^*(y)$. Further,

$$\begin{aligned}
& K_{\varepsilon_n}^\mu \left[K_{\varepsilon_n}^\nu [h] \right] (y_2) \\
&\asymp \frac{1}{\nu^{1/2}(y_2)} \int \frac{1}{(2\pi\varepsilon_n)^{d/2}} \sqrt{\frac{\det[\nabla^2\varphi_0^*(y_2)] \det[\nabla^2\varphi_0^*(y)]}{\det[\nabla^2\varphi_0^*(y_2) + \nabla^2\varphi_0^*(y)]}} h(y) \\
&\quad \times \exp\left(-\frac{1}{2\varepsilon_n}(\nabla\varphi_0^*(y) - \nabla\varphi_0^*(y_2))^\top [\nabla^2\varphi_0^*(y_2) + \nabla^2\varphi_0^*(y)]^{-1}(\nabla\varphi_0^*(y) - \nabla\varphi_0^*(y_2))\right) \\
&\quad \nu^{1/2}(y)dy.
\end{aligned}$$

Thus, owing to the assumed Lipschitzianity of the metric tensor,

$$\begin{aligned}
& K_{\varepsilon_n}^\mu \left[K_{\varepsilon_n}^\nu [h] \right] (y_2) \\
& \asymp \frac{1}{\nu^{1/2}(y_2)} \int \frac{1}{(2\pi\varepsilon_n)^{d/2}} h(y) \sqrt{\frac{\det[\nabla^2\varphi_0^*(y_2)] \det[\nabla^2\varphi_0^*(y)]}{\det[\nabla^2\varphi_0^*(y_2) + \nabla^2\varphi_0^*(y)]}} \\
& \times \exp \left(-\frac{1}{4\varepsilon_n} (\nabla\varphi_0^*(y) - \nabla\varphi_0^*(y_2))^\top [\nabla^2\varphi_0^*(y_2) + o(1)]^{-1} (\nabla\varphi_0^*(y) - \nabla\varphi_0^*(y_2)) \right) \\
& \nu^{1/2}(y) dy,
\end{aligned}$$

where the $o(1)$ in the display above is $O(\|y - y_2\|)$. The claim follows owing to the Bregman duality, i.e.,

$$\begin{aligned}
& (\nabla\varphi_0^*(y) - \nabla\varphi_0^*(y_2))^\top [\nabla^2\varphi_0^*(y_2) + o(1)]^{-1} (\nabla\varphi_0^*(y) - \nabla\varphi_0^*(y_2)) \\
& = (y - y_2)^\top [\nabla^2\varphi_0^*(y_2)] (y - y_2) + o(\|y - y_2\|^2),
\end{aligned}$$

which comes from the fact that both terms of the equality above are quadratic approximations of the Bregman divergence $\varphi_0(y) + \psi_0(x) - \langle x, y \rangle$ in one or the other variable. \square