DIGITAL *n*-MANIFOLDS WITH OR WITHOUT BOUNDARIES

MELİH İS AND İSMET KARACA

ABSTRACT. This work aims to define the concept of manifold, which has a very important place in the topology, on digital images. So, a general perspective is provided for two and three-dimensional imaging studies on digital curves and digital surfaces. Throughout the study, the features present in topological manifolds but that are not satisfied in the discrete version are specifically underlined. In addition, other concepts closely related to manifolds such as submanifold, orientation, and partition of unity are also discussed in digital images.

1. INTRODUCTION

A topological space locally homeomorphic to n-dimensional Euclidean space is called a topological n-manifold. It provides for a large variety of global structures while having a local resemblance to the Euclidean space. As a fundamental idea in modern mathematics and physics, a manifold takes the intuitive idea of curves and surfaces to higher dimensions [21]. The concept of a manifold is first proposed by Bernhard Riemann in the 19th century and differentiable manifolds are his main area of interest, especially in the setting of geometry and analysis [23]. As topology advanced in the early 20th century, the definition of topological manifolds is presented by the work of mathematicians like Henri Poincaré and Felix Hausdorff [14,22].

Topological manifolds have some essential characteristics that make them essential to science and mathematics. Depending on the situation, they can impose differentiable, Riemannian, or symplectic structures, which is one of their most remarkable characteristics. As an example in physics, especially in general relativity, spacetime is represented as a four-dimensional differentiable manifold with a Lorentzian metric. For another, data science and manifold studies are related, as manifold learning techniques seek to reveal low-dimensional manifold structures hidden behind high-dimensional data [28].

Manifolds facilitate the simultaneous examination of both local and global properties. For example, the fact that a manifold is locally flat (differentiable) allows some analysis operations such as differentiation and integration to be performed on it. However, the global topological structure of the manifold dictates the fundamental behavior of these operations. For instance, a 2-dimensional sphere (S^2) shares the local properties of the plane (\mathbb{R}^2) but has a different topological structure globally. In algebraic topology and homotopy theory, invariants such as homology

Date: December 17, 2024.

²⁰¹⁰ Mathematics Subject Classification. 57N65, 58K65, 32Q55, 68U05.

Key words and phrases. digital topology, manifold, manifold with boundary, orientation, partition of unity.

and cohomology groups aid in the classification and structural analysis of manifolds [27]. A manifold's holes, gaps, and other topological characteristics can be expressed numerically using these tools. As an example, the fundamental group of a manifold analyzes its cyclic structures, whereas homology groups offer a broader viewpoint. Such topological and algebraic topological information plays a critical role in the classification of manifolds.

Digital topology investigates the application of traditional topological ideas to digital images and offers numerous significant results on digital curves and surfaces [20]. These structures are so crucial, especially in the fields of digital image processing and computer graphics. Digital surfaces are employed in the structural study of three-dimensional models, and digital curves are frequently utilized to depict boundaries in digital images [16,24,25]. A digital curve is commonly described as a simple closed path [24]. According to the Jordan curve theorem in classical topology, a plane is divided into inner and exterior regions by a simple closed curve. This theorem, when applied to the digital plane in digital topology, demonstrates how a digital curve separates a picture into an interior and an exterior region [20]. A digital curve is said to be simple if it simply has the same beginning and ending points and is not closed on itself. The accuracy of algorithms used in digital image processing depends on this feature [24]. On the other hand, in three-dimensional digital images, digital surfaces are used to simulate an object's interior structure or boundary [16]. The homotopic and homological properties of digital surfaces are similar to those of classical surfaces. Specifically, digital surfaces are analyzed using topological invariants such as the Euler characteristic [10]. Applications like volume calculation, boundary detection, and three-dimensional object reconstruction commonly use digital surfaces. Digital surface analysis is essential, particularly in the fields of industrial modeling and medical imaging [16]. These examinations on digital curves and surfaces demonstrate both the theoretical and practical value of digital topology.

Historically, the first ideas about the concept of digital manifold, and the works of formalizing these ideas in a systematic order are carried out by Rosenfeld [24]. Herman [16], [8], and [9]. However, these studies are generally based on parallel moves or cell structures and do not provide an explicit definition of digital manifold via adjacency relations. Mathematically, in this manuscript, we examine the elements that constitute the topological manifold, namely the properties of being Hausdorff, being second-countable, and being local homeomorphism, on digital images and how the concept of digital manifold should be defined using adjacency relations. In Section 2, we examine how some important definitions and results are used in digital images. After that, in Section 3, we present the notion of 'digital n-manifold' (and also 'digital n-manifold with boundary') by interpreting the discrete version of the local homeomorphism property (note that we shall explain the fact that this is enough to define a topological n-manifold in digital topology in the section). Furthermore, we immediately exemplify this new concept by considering the digital intervals, digital circles, or any subsets of \mathbb{Z}^n . Also, we mention some examples of digital images that do not have a digital manifold structure. In addition, the important point that we frequently refer to in this section is to reveal the final state of the manifold properties on the discrete setting, which exist in topological spaces but change in digital images. We also deal with some manifold-related notions such as submanifolds, orientations, and partitions of unities. In Section 4, we present some open problems for many concepts in the previous section and guidance for future digital manifold works.

2. Preliminaries

In this section, we introduce the discrete version of some topological concepts such as digital image, adjacency relation, digital homotopy, digital isomorphism, etc., which are the basic elements of digital topology.

Assume that M is a subset of \mathbb{Z}^n and κ_l is a given adjacency relation over the points of M. Then a *digital image* is a binary form that works on discrete structures and has an adjacency relation rather than a topology [19]. It is generally denoted by the pair (M, κ_l) and the adjacency relation κ_l is defined as follows:

Definition 2.1. [19] Given any two distinct points $m = (m_1, \dots, m_n)$ and $m' = (m'_1, \dots, m'_n)$ of \mathbb{Z}^n , and an integer l with $1 \leq l \leq n$, we say that m is called κ_l -adjacent to m' if the following satisfy:

- There exist the maximum number of l indices $k \in \{1, \dots, n\}$ with the condition $|m_k m'_k| = 1$.
- The equality m_s = m'_s holds for any indices s ∈ {1, · · · , n} with the condition |m_s m'_s| ≠ 1.

For simple illustrations of adjacency relations, we observe that

- $\kappa_1 = 2$ -adjacency in \mathbb{Z} ,
- $\kappa_1 = 4$ -adjacency and $\kappa_2 = 8$ -adjacency in \mathbb{Z}^2 , and
- $\kappa_1 = 6$ -adjacency, $\kappa_2 = 18$ -adjacency, and $\kappa_3 = 26$ -adjacency in \mathbb{Z}^3 .

The fact that m is κ_l -adjacent to m' is shown by the notation $m \sim_{\kappa_l} m'$.

Assume that (M, κ_l) is a digital image and m is any point of M. Then the *digital* neighborhood [6] of m is

$$N_{\kappa_{l}}(m) = \{ m' \in \mathbb{Z}^{n} : m \sim_{\kappa_{l}} m' \}.$$

In addition, $N^*_{\kappa_l}(m)$ is given by $N_{\kappa_l}(m) \cup \{m\}$.

A digital image (M, κ_l) is called *digitally* κ_l -connected [15] if and only if for any distinct points $m, m' \in M$, there exists a subset $\{m_0, m_1, ..., m_s\} \subseteq M$ with the following conditions:

- The initial and final points are m and m', respectively, i.e., $m = m_0$ and $m' = m_s$.
- One has $m_j \sim_{\kappa_l} m_{j+1}$ for each j = 0, 1, ..., s 1.

By recalling the definition of totally disconnected space, we can give the following in digital images in parallel with topological spaces:

Definition 2.2. A digital image is said to be *totally disconnected* provided that it admits only one-point sets as digitally connected subsets.

Assume that $\alpha : (M, \kappa_l) \to (M', \kappa_s)$ is any map of digital images. Then α is digitally (κ_l, κ_s) -continuous [2, 26] if

 M_1 is κ_l -connected in $M \Rightarrow \alpha(M_1)$ is κ_s -connected in M'

for any subset $M_1 \subseteq M$. Proposition 2.5 in [2] emphasizes that when we have two digitally continuous maps, their composition is also a digitally continuous map. A digital (κ_l, κ_s) -isomorphism is a map $\alpha : (M, \kappa_l) \to (M', \kappa_s)$ of digital images provided that each of the following satisfies [4, 12]:

- α is bijective and (κ_l, κ_s) -continuous.
- The inverse of α is (κ_s, κ_l) -continuous.

The digital n-sphere [3], the discrete version of S^n in topological spaces, is presented by the boundary of I_n . More explicitly, S^n in digital images is $[-1,1]_{\mathbb{Z}}^{n+1} \setminus \{(0,\cdots,0)\} \subset \mathbb{Z}^{n+1}$, where $(0,\cdots,0)$ denotes the origin of \mathbb{Z}^{n+1} and sometimes shortly expressed by 0_{n+1} .

The strong adjacency NP, also known as the normal product adjacency, is defined as follows for the Cartesian product created by two digital images.

Definition 2.3. [1] Assume that (M, κ_l) and (M', κ_s) are any digital images (or graphs). Let $m_1, m_2 \in M$ and $m'_1, m'_2 \in M'$ be any points. Then

$$(m_1, m'_1) \sim_{\mathrm{NP}(\kappa_l, \kappa_s)} (m_2, m'_2)$$
 in $M \times M'_1$

if and only if one of the following satisfies:

a) $m_1 = m_2$ and $m'_1 \sim_{\kappa_s} m'_2$, or b) $m_1 \sim_{\kappa_l} m_2$ and $m'_1 = m'_2$, or c) $m_1 \sim_{\kappa_l} m_2$ and $m'_1 \sim_{\kappa_s} m'_2$.

In [5], an important property tells us that a κ_l -adjacency can be differ from a strong adjacency. The NP-adjacency is entirely determined by the adjacencies of the factors. Moreover, one has the fact that shows when the NP-adjacency is equal to the c_k -adjacency.

Proposition 2.4. [5] Assume that (M, κ_l) and (M', κ_s) are any digital images. Then the NP-adjacency and the κ_{l+s} -adjacency are equal to each other in the Cartesian product $M \times M'$.

A digital interval [3], the discrete version of a closed interval in topological spaces, is given by $[m, m']_{\mathbb{Z}} = \{r \in \mathbb{Z} : m \leq r \leq m'\}$. Assume that $[0, j]_{\mathbb{Z}}$ is any digital interval with the 2-adjacency and $f, g: (M, \kappa_l) \to (M', \kappa_s)$ are two (κ_l, κ_s) -continuous maps. Then f and g are called *digitally* (κ_l, κ_s) -homotopic [2] if there is a digitally continuous map $H : M \times [0, j]_{\mathbb{Z}} \to M'$ satisfying H(m, 0) = f(x) and H(m, j) = g(x) for all $m \in M$ with the properties that, for any fixed $m \in M$, the map

$$H_m: [0,j]_{\mathbb{Z}} \to M$$

4

defined by $H_m(t) = H(m, t)$ is digitally continuous for all t, and for any fixed $t \in [0, j]_{\mathbb{Z}}$, the map

$$H_t: M \to M'$$

defined by $H_t(m) = K(m, t)$ is digitally continuous for all x. Moreover, n is the number of steps of H and H is called a *digital homotopy* between f and g in j steps. $f \simeq_{\kappa_l,\kappa_s} g$ indicates that f and g are digitally homotopic to each other. Furthermore, a digital image (M,κ_l) is *digitally* κ_l -contractible [2] if $1_M \simeq_{\kappa_l,\kappa_l} s$ for some constant map $s: M \to M$.

Assume that M' is a κ_s -connected digital image, $m_0 \in M'$ a basepoint, and $f: (M, \kappa_l) \to (M', \kappa_s)$ a (κ_l, κ_s) -continuous map. Then f is a *digital fiber bundle* with digital fiber (N, κ_v) provided that the following are satisfied [18]:

- $f^{-1}(m_0) = N$ and f is surjective.
- Let $m' \in M'$ any point. Then there is a κ_s -connected subset $W \subseteq \mathbb{Z}$ with $m' \in W$ and an isomorphism $\varphi : f^{-1}(W) \to W \times N$ with $\pi_1 \circ \varphi = f$. Here, $\pi_1 : (W \times N, \kappa_*) \to (W, \kappa_s)$ is the first projection, where κ_* is $NP(\kappa_s, \kappa_1)$.

The digital Euler characteristic, the discrete version of Euler characteristic in topological spaces, is defined in [13] but we use the improved version of this notion [17]: Let (M, κ_l) be a digital image. A set of i + 1 unique elements of a digital image (M, κ_l) , each pair of which is κ_l -adjacent, constitutes a κ_l -simplex in M of dimension i. Assume now that α_r is the number of unique r-dimensional κ_l -simplices in X, and n is the greatest integer i such that M has a κ_l -simplex of dimension i. Then the digital Euler characteristic [17] of a digital image (M, κ_l) ,

denoted by $\chi(M, \kappa_l)$ or shortly $\chi(M)$, is defined by the number $\sum_{r=0}^{n} (-1)^r \alpha_r$.

Proposition 2.5. [13, 17] If (M, κ_l) and (M', κ_s) are digitally isomorphic digital images, then their digital Euler characteristics are the same, *i.e.*,

$$\chi(M,\kappa_l) = \chi(M,\kappa_s).$$

3. DIGITAL MANIFOLDS AND SOME COUNTEREXAMPLES

A topological n-manifold M includes three basic properties as follows.

- M is Hausdorff.
- *M* is second-countable.
- M admits a local homeomorphism, i.e., every point $m \in M$ has a neighborhood W such that W is homeomorphic to an open subset of \mathbb{R}^n .

When defining the concept of a manifold in digital topology is desired, it is necessary to examine these three properties separately on digital images. The properties of being a Hausdorff space and being a second-countable space in digital images do not tell us new things mathematically. We need to focus more on the third property.

The discrete topology is a Hausdorff space since every subset is open. Indeed, any two distinct points can be separated by open sets, each containing only that point. Recall that no topology exists on digital images, that is, every subset can be assumed to be open. Therefore, digital images, the discrete construction of topological spaces over \mathbb{Z}^n , always tend to have the Hausdorff property.

Let (M, κ) be a digital image. Then a collection β of subsets of M is called a digital base for the digital image M if and only if every subset $N \subseteq M$ can be written as the union of elements of β . In addition, a digital image M is called second-countable if M has a countable digital basis. Recall that the discrete topological space (\mathbb{R}, τ_d) is not a second-countable space even if $\beta = \{m : m \in \mathbb{R}\}$ is a basis for \mathbb{R} . Because \mathbb{R} is not countable. On the other hand, the collection $\{m : m \in \mathbb{Z}\}$ forms a countable digital basis for the discrete version \mathbb{Z} of real numbers. Thus, digital images, the discrete construction of topological spaces over \mathbb{Z}^n , always tend to have the second-countable property.

Definition 3.1. A digital image (M, κ) is called a digital *n*-manifold provided that for a minimum non-negative integer *n*, each point $m \in X$ has a digital neighborhood $N_{\kappa}(m)$ that is digitally isomorphic to $N_{\lambda}(s)$ of at least one point *s* in \mathbb{Z}^n .

A digital isomorphism $N_{\kappa}(m) \to N_{\lambda}(s)$ in Definition 3.1 is called a digital chart. The inverse of this isomorphism $N_{\lambda}(m) \to N_{\kappa}(s)$ is called a digital local parametrization of M about m. A set of digital charts is said to be a digital atlas provided that M can be covered by their domains.

Example 3.2. • The digital 0-sphere $S^0 = \{-1, 1\}$ is a digital 0-manifold since $N_{\kappa}(-1) = N_{\kappa}(1) = \emptyset$ for -1 and $1 \in S^0$.

- \mathbb{Z} is a digital 1-manifold: Assume that m is any integer. Then we have $N_2(m) = \{m 1, m + 1\}$. Therefore, $N_2(m)$ is digitally isomorphic to $N_2(s) = \{s 1, s + 1\}$ for any s different from m in \mathbb{Z} by the map $\gamma: N_2(m) \to N_2(s), \gamma(m-1) = s 1$ and $\gamma(m+1) = s + 1$.
- S^1 is a digital 1-manifold: Let S^1 have $\kappa_1 = 4$ or $\kappa_2 = 8$ -adjacency. Each point $m \in S^1$ has two adjacent points, say m_1 and m_2 . Therefore, there is a point s in \mathbb{Z} with the property that $P = N_{\kappa_l}(m) = \{m_1, m_2\}$ is digitally isomorphic to $N_2(s) = \{s 1, s + 1\}$ for l = 1, 2.

Example 3.3. It is not possible to say that any subset of a digital n-manifold is a digital n-manifold. For example \mathbb{Z}^2 is a digital 2-manifold but

$$M = \mathbb{Z} \times \{0\} \cup \{0\} \times \mathbb{Z} \subseteq \mathbb{Z}^2$$

is not a digital manifold of any dimension: Suppose that M is a digital 2-manifold with the 4-adjacency. Then, for any point $m \in M - \{(0,0)\}$, we have a digital isomorphism

$$\gamma: N_4(m) \to N_4(s),$$

where $s \in \mathbb{Z}^2$ is any point. However, this is a contradiction because $N_4(m)$ has exactly two totally disconnected points and $N_4(s)$ has exactly four totally disconnected points, i.e., a bijective map cannot be constructed. Similarly, by assuming that M is a digital 1-manifold, we have a digital isomorphism

$$\gamma': N_4((0,0)) \to N_2(s')$$

for any point $s' \in \mathbb{Z}$. This is again a contradiction because $N_4((0,0))$ has four totally disconnected points (1,0), (0,1), (-1,0), and (0,-1) whereas $N_2(s')$ has two totally disconnected points s' - 1 and s' + 1. Moreover, M cannot be a 1-manifold. A similar way can be followed when M has the 8-adjacency. Indeed, (2,0) has exactly two 8-adjacent points ((1,0) and (3,0)) but (1,0) has exactly four 8-adjacent points ((2,0), (0,0), (-1,0), and (1,0)).

The union of two digital manifolds need not be a digital manifold again. For instance, let M_1 and M_2 be two digital images

$$\{(m_1, 0) : 0 \le m_1 \le 4\} \cup \{(4, m_1) : 1 \le m_1 \le 4\} \\ \cup \{(m_1, 4) : 0 \le m_1 \le 3\} \cup \{(0, m_1) : 1 \le m_1 \le 3\}$$

and

 $\{m_1, m_2 \in \mathbb{Z} : m_1 = m_2 \text{ and } 0 \le m_1, m_2 \le 4\}$

with the 4-adjacency, respectively. Even if M_1 is a digital 1-manifold and M_2 is a digital 0-manifold, $M_1 \cup M_2$ is not a digital manifold. Indeed, whenever the point (0, 4) behaves like an element of a digital 1-manifold $(N_4((0, 4)))$ has two totally disconnected points (0, 3) and (1, 4) but (2, 2) behaves like an element of a digital 0-manifold $(N_4((2, 2)) = \emptyset)$.

Another difference from a topological space is in the classification of some manifolds in digital images. For instance, a connected 1-manifold can be homeomorphic to one of the following manifolds: a circle, a closed interval [0, 1], the real numbers, or the half-line. In the discrete version, one can intuitively think that a digitally connected 1-manifold can be digitally isomorphic to a digital interval or a digital 1-sphere since the discretizations of the closed interval, \mathbb{R} , and the half-line are all a digital interval. However, this is false. Indeed, a digital 1-manifold $M = \{(0,0), (0,1), (1,0), (1,1)\}$ with the 4-adjacency is not digitally isomorphic to these two possible digital images (see Figure 3.1).



FIGURE 3.1. As an example of a digital 1-manifold, the digital image $M = \{(0,0), (0,1), (1,0), (1,1)\}$ with the 4-adjacency.

Definition 3.4. A digital image (M, κ) is called a digital *n*-manifold with boundary provided that for a minimum non-negative integer *n*, each point $m \in M$ has a digital neighborhood $N_{\kappa}(m)$ that is digitally isomorphic to $N_{\lambda}(s)$ of at least one point s in a digital image $D_+^n = \{(m_1, \cdots, m_n) \in \mathbb{Z}^n : m_j \ge 0 \text{ for all } j = 1, \cdots, n\}$ or \mathbb{Z}^n .

The boundary ∂M of a digital *n*-manifold (M, κ) is defined by a digital image having all points mapped to $m_j = 0$ by a digital chart for at least one $j \in \{1, \dots, n\}$ and the interior $\operatorname{int}(M)$ of a digital *n*-manifold (M, κ) is defined by a digital image having all points mapped to $m_j > 0$ by some digital charts for all j, i.e.,

$$M = \operatorname{int}(M) \cup \partial M.$$

If $\partial M = \emptyset$, then (M, κ) is a manifold (without boundary), namely that Definition 3.4 corresponds to Definition 3.1.

Example 3.5. Let $M = [1,3]_{\mathbb{Z}} \times [1,3]_{\mathbb{Z}}$ be a digital image in \mathbb{Z}^2 with the 4-adjacency. The point $(2,2) \in M$ admits a digital isomorphism

$$N_4((2,2)) \to N_4(0,0)$$

for the point $(0,0) \in D^2_+$. Some points such as (2,1), (3,2), (2,3), and (1,2) admit a digital isomorphism

$$N_4(-) \to N_4((1,0))$$

for the point $(1,0) \in D^2_+$. The points (1,1), (3,1), (3,3), and (1,3) admit a digital isomorphism

$$N_4(-) \to N_4(5,5)$$

for the point $(5,5) \in D^2_+$. This shows that (M,4) is a digital 2-manifold with boundary. Moreover, we observe that (2,2) is the only interior point of M and $\partial M = \{m \in M : m \neq (2,2)\}$ is the boundary set of M. It follows that int(m) is a digital 0-manifold and ∂M is a digital 1-manifold.

In topological spaces, if M is an n-manifold with boundary, then int(M) is an n-manifold, too. However, this is not always true for digital images. Example 3.5 presents a counterexample of this in digital images.

Proposition 3.6. A digital interval is a digital 1-manifold with two boundaries.

Proof. Let $[a, b]_{\mathbb{Z}}$ be any digital interval. Then $N_2(a)$ and $N_2(b)$ have exactly one totally disconnected points $\{a+1\}$ and $\{b-1\}$, respectively. For any point $0 \in D^1_+$, we have two digital isomorphisms

$$N_2(a) \to N_2(0)$$

with $a + 1 \mapsto 1$ and

$$N_2(b) \to N_2(0)$$

with $b - 1 \mapsto 1$. It follows that $a, b \in \partial M$. Moreover, if $[a, b]_{\mathbb{Z}}$ has any point c such that $c \neq a$ and $c \neq b$, then we have a digital isomorphism

$$N_2(c) \to N_2(s)$$

with $c - 1 \mapsto s - 1$ and $c + 1 \mapsto s + 1$ for any point $s \in D^1_+ - \{0\}$ (namely that $c \in int([a, b]_{\mathbb{Z}}))$. Therefore, $[a, b]_{\mathbb{Z}}$ is a digital 1-manifold with two boundaries a and b.

Proposition 3.7. The digital 2-sphere S^2 is a digital 2-manifold with boundaries with the 6-adjacency, and a digital 2-manifold with the 18 and 26-adjacencies, where D^2_+ has the 4-adjacency.

Proof. First, consider the 6-adjacency. For any point $m \in S^2$, $N_6(m)$ has three or four totally disconnected points (for example $N_6(1, 1, 1)$ has three totally disconnected points and $N_6(0, 0, 1)$ has four totally disconnected points). Then, for any points (1, 0) and (2, 2) in D^2_+ , there exist two digital isomorphisms

$$N_6(m_1) \to N_4((1,0))$$
 and $N_6(m_2) \to N_4((2,2))$,

where m_1 and m_2 can be considered any points of S^2 with the properties that $N_6(m_1)$ has three totally disconnected points and $N_6(m_2)$ has four totally disconnected points, respectively. Thus, S^2 is a digital 2-manifold with boundaries. Moreover, $\partial S^2 = \{(x, y, z) : x, y, z \in \{-1, 1\}\}$ and $\operatorname{int}(S^2) = S^2 - \partial S^2$. Assume now that S^2 has the 18-adjacency. For each point $m_3 \in S^2$, $N_{18}(m_3)$ has exactly four totally disconnected points. It follows that $N_{18}(m_3) \to N_4(s)$ is a digital isomorphism for any point s in \mathbb{Z}^2 . Furthermore, if we consider the 26-adjacency on S^2 , then each point $m_4 \in S^2$ admits that $N_{26}(m_4)$ has four totally disconnected points, too. Thus, by applying a similar process, it is obtained that S^2 with the 26-adjacency is again a digital 2-manifold.

In topological spaces, if M is an n-manifold with boundary, then ∂M is an (n-1)-manifold. However, this is not always true for digital images. Proposition 3.7 presents a counterexample of this in digital images because $(S^2, 6)$ is a digital 2-manifold with boundary and ∂S^2 is a digital 0-manifold, not a digital 1-manifold, with the 6-adjacency.

Proposition 3.8. The digital 2-sphere S^2 is not a digital 2-manifold, where D^2_+ has the 8-adjacency.

Proof. Let $m_1 \in \mathbb{Z}$ be nonzero. Then any points of the form $(m_1, 0)$ in D^2_+ admits that $N_8((m_1, 0))$ has 5 adjacent points and any points of the form (m_1, m_2) in D^2_+ with $m_1, m_2 > 0$ admits that $N_8((m_1, m_2))$ has 8 adjacent points. In addition, (0, 0)admits that $N_8((0, 0))$ has 3 adjacent points. Since each point $m \in S^2$ admits that $N_6(m)$ has three or four totally disconnected points, there is no digital isomorphism from $N_6(m)$ to $N_8(s)$ for any $s \in D^2_+$ or $s \in \mathbb{Z}^2$. In addition, a similar process can be done for S^2 with the 18 and 26-adjacencies.

The proofs of Proposition 3.7 and Proposition 3.8 can be generalized over the number of elements for corresponding $M_{\kappa_l}(m)$. So, we have the following result:

Theorem 3.9. *i)* The digital *n*-sphere is a digital *n*-manifold with boundaries with the κ_1 -adjacency, and for other adjacencies, it is a digital *n*-manifold, where D^n_+ has the κ_1 -adjacency.

ii) The digital n-sphere is not a digital n-manifold, where D^2_+ has the κ_l -adjacency with l > 1.

Note that $[0,1] \times [0,1]$ is a 2-manifold with boundary in topological spaces but $[0,1]_{\mathbb{Z}} \times [0,1]_{\mathbb{Z}}$ is a digital 1-manifold without boundary with the 4-adjacency.

Any open subset in n-dimensional Euclidean space is a topological n-manifold but this cannot be transferred into digital images.

Proposition 3.10. A digital neighborhood $(N_{\kappa}(m), \kappa_1)$ of any point $m \in \mathbb{Z}^n$ is a digital 0-manifold. In addition, $(N^*_{\kappa}(p), \kappa_l)$ need not be a digital manifold.

Proof. Let m be any point in (\mathbb{Z}^n, κ_1) . Then $N_{\kappa_1}(m)$ has totally disconnected κ_1 points. Choose m' as one of these points in $N_{\kappa_1}(m)$. Since m' has no κ_1 -neighbours, the digital neighborhood $N_{\kappa_1}(m)$ of m is emptyset. This means that it is a digital 0-manifold. On the other hand, when we consider $(0,0) \in \mathbb{Z}^2$, the digital image $N_4^*((0,0)) = \{(0,0), (0,1), (-1,0), (0,-1), (1,0)\}$ is not a digital manifold by using the same method with Example 3.3 (observe that (0,1) has only one adjacent point but (0,0) has four adjacent points.).

For topological manifolds, if it is connected, then its interior is also connected. This is not true for digital images. As for the counterexample, consider the digital image

$$M = ([0,4]_{\mathbb{Z}} \times [0,4]_{\mathbb{Z}}) - \{(2,2)\}$$

with the 4-adjacency (see Figure 3.2). Then $int(M) = \{(1, 1), (3, 1), (3, 3), (1, 3)\}$ because the only elements of M that have four adjacent points are (1, 1), (1, 3), (3, 1), and (3, 3). It follows that int(M) is not digitally connected even if M is digitally connected 2-manifold with boundary.



FIGURE 3.2. As an example of a digital 2-manifold with boundary, the digital image $M = ([0,4]_{\mathbb{Z}} \times [0,4]_{\mathbb{Z}}) - \{(2,2)\}$ by considering the 4-adjacency.

Definition 3.11. Let (M, κ) be a digital *n*-manifold. A subset *S* of *M* with the digital *r*-manifold structure for $r \leq n$ is a digital *r*-dimensional submanifold if *S* is the image of an embedding of digital images, i.e., $\gamma : S \to M$ is a digital isomorphism onto its image.

10

It is not always the case that S and M have the same dimension. For example, when we consider \mathbb{Z}^2 as the digital 2-manifold, \mathbb{Z} is a digital 1-dimensional submanifold of \mathbb{Z}^n by using a digital embedding $\gamma : \mathbb{Z} \to \mathbb{Z}^2$, $\gamma(m) = (m, 0)$.

Example 3.12. A digital (n-1)-sphere is a digital n-dimensional submanifold of \mathbb{Z}^n .

In topological spaces, $\chi(M_1 \times M_2) = \chi(M_1)\chi(M_2)$, where χ denotes the Euler characteristic of a space. Moreover, $\chi(M_1) \sqcup \chi(M_2) = \chi(M_1) + \chi(M_2)$, where \sqcup denotes the disjoint union.

Theorem 3.13. Let (M_1, κ_1) and (M_2, κ_2) be any digital manifolds of any dimensions. Then $\chi(M_1 \times M_2)$ does not always equal to $\chi(M_1)\chi(M_2)$ in digital images. Moreover, $\chi(M_1 \sqcup M_2)$ does not always equal to $\chi(M_1) + \chi(M_2)$ in digital images, too.

Proof. By assuming that (M_1, κ_1) and (M_2, κ_2) are both equal to $([0, 1]_{\mathbb{Z}}, 2)$, we have that $\chi(M_1) = \chi(M_2) = 2 - 1 = 1$. Therefore, $\chi(M_1)\chi(M_2) = 1$. On the other hand, we observe that $\chi(M_1 \times M_2, 4) = 4 - 4 - 4 = -4$ with the 4-adjacency. For the second statement, choose M_1 as $\{(0,0), (1,0)\}$ and M_2 as $\{(0,1), (1,1)\}$ by considering the 4-adjacency. Thus, $\chi(M_1) + \chi(M_2) = 2$ but $\chi(M_1 \sqcup M_2) = -4$. \Box

Contractible spaces have Euler characteristic 1 for topological manifolds. In digital images, digital 2–sphere with the 18–adjacency is digitally 18–contractible but its Euler characteristic is equal to -2 [17].

An orientation of a digital 0-manifold M is a function $\epsilon : (M, \kappa) \to (\{\mp 1\}, 2)$. To compute the number of all orientations in a digital 1-manifold, a linear order can be used. On a set M, the system of rays $\{m \in M \mid r < m\}$ for an element $r \in M$ encodes a linear order <.

Proposition 3.14. A digital interval (a type of digital 1-manifold) has exactly two linear orders.

Proof. Let $[a, b]_{\mathbb{Z}}$ be any digital interval. Then two linear orders < and > (we can think of these orders as from a to b and from b to a, respectively) exist on this interval. It follows that there are two rays $\{p \in M \mid r < p\}$ and $\{p \in M \mid p < r\}$, the digitally connected components of $M - \{r\}$. When we consider any linear order << different from <, the rays $\{p \in M \mid r < cp\}$ and $\{p \in M \mid p < r\}$ intersect U and V in $M - \{r\}$ in disjoint open sets. Since U and V are digitally connected, one of these rays is equal to U or V. This means that << must be equal to < or >.

Thus, we have the following by the previous fact:

Theorem 3.15. A digital interval has exactly two orientations.

Recall that a digital bundle with a digital fiber is given in [18]. A special kind of digital fiber bundle is called a digital vector bundle if the digital fiber is a vector space. The default type of a digital vector bundle is a digital real vector bundle, which has a digital fiber that is a real vector space. Let $M \to N$ be a digital real vector bundle of rank s. Then the j-th digital Stiefel-Whitney class is defined as the digital cohomology class $w_j(M) \in H^{j,q}(N; \mathbb{Z}_2)$ for $0 \leq j \leq s$.

Definition 3.16. A digital manifold M is orientable (otherwise non-orientable) if and only if the first digital Stiefel-Whitney class is zero (otherwise nonzero).

Example 3.17. Consider a digital 2-manifold

 $M_1 = \{(1, 1, 0), (0, 2, 0), (-1, 1, 0), (0, 0, 0), (0, 1, -1), (0, 1, 1)\}$

with the 18-adjacency. By Example 4.2 in [11], we have that $H^{1,18}(M_1;\mathbb{Z}_2) = 0$. This shows that $w_1(M_1) = 0$, which concludes that M_1 is an orientable digital 2-manifold.

Example 3.18. Let $M_2 = [0,1]_{\mathbb{Z}} \times [0,1]_{\mathbb{Z}} \times [0,1]_{\mathbb{Z}}$ be a digital 2-manifold with boundary with respect to the 6-adjacency. In [7], we know that $H^{1,6}(M_2;\mathbb{Z}_2) = \mathbb{Z}_2^5$. It follows that $w_1(M_2)$ need not be zero, which concludes that M_2 is a non-orientable digital 2-manifold with boundary.

Definition 3.19. Let (M, κ) be a digital image and $f : M \to \mathbb{Z}$ a function. Then a digital support of f is defined as $\operatorname{sp}(f) = \{p \in M : f(p) \neq 0\}$.

The definition of digital support directly gives some properties:

- $p \notin \operatorname{sp}(f)$ implies that f(p) must be zero.
- The fact sp(f) is the empty set implies that f vanishes.

Proposition 3.20. Let $f, g: (M, \kappa) \to (\mathbb{Z}, 2)$ be two functions. Then

a) $sp(f.g) = sp(f) \cap sp(g)$.

b) $sp(f+g) = sp(f) \cup sp(g)$.

Proof. Let $p \in \mathbb{Z}$ be any point.

a) Then we have

$$\begin{split} p \in \mathrm{sp}(f.g) & \Leftrightarrow \quad (f.g)(p) \neq 0 \Leftrightarrow f(p).g(p) \neq 0 \Leftrightarrow f(p) \neq 0 \land \ g(p) \neq 0 \\ & \Leftrightarrow \quad x \in \mathrm{sp}(f) \land \ p \in \mathrm{sp}(g) \Leftrightarrow p \in \mathrm{sp}(f) \cap \mathrm{sp}(g). \end{split}$$

b) Then we have

$$\begin{array}{ll} p\in \mathrm{sp}(f+g) & \Leftrightarrow & (f+g)(p) \neq 0 \Leftrightarrow f(p)+g(p) \neq 0 \Leftrightarrow f(p) \neq 0 \ \lor \ g(p) \neq 0 \\ & \Leftrightarrow & p\in \mathrm{sp}(f) \ \lor \ p\in \mathrm{sp}(g) \Leftrightarrow p\in \mathrm{sp}(f) \cup \mathrm{sp}(g). \end{array}$$

Proposition 3.21. Let $f : (M, \kappa) \to (\mathbb{Z}, 2)$ be a function and $\alpha : (M', \kappa') \to (M, \kappa)$ a digital isomorphism. Then $sp(f \circ \alpha) = \alpha^{-1}(sp(f))$. *Proof.* Let $p' \in M'$ be any element. Then

$$\alpha(\operatorname{sp}(f \circ \alpha) = \alpha(\{p' \in M' : f(\alpha(p')) \neq 0\}) = \{p \in M : f(p) \neq 0\} = \operatorname{sp}(f).$$

The shows that $\operatorname{sp}(f \circ \alpha) = \alpha^{-1}(\operatorname{sp}(f)).$

Definition 3.22. A digital partition of unity of a digital manifold (M, κ) is the set

 $\{\alpha_i: M \to \mathbb{Z} \mid \alpha_i \text{ is digitally continuous for each } i \in [0, m]_{\mathbb{Z}}\}$

for which the following properties hold:

- 1. Each α_i is nonnegative.
- 2. Every point $p \in M$ admits a digital neighborhood $N_{\kappa}(p)$ with the property that $N_{\kappa}(p) \cap \operatorname{sp}(\alpha_i) \neq \emptyset$ for all *i*.
- 3. For each $p \in M$, $g_1(p) + \cdots + g_m(p) = m$.

Definition 3.23. A digital partition of unity $\{\alpha_i\}_{i \in [0,m]_{\mathbb{Z}}}$ on a digital manifold (M, κ) is subordinate to a cover $\{M_1, M_2, \dots, M_r\}$ of M provided that each α_i admits a digital image M_i of the cover with the property $\operatorname{sp}(\alpha_i) \subset M_i$.

Example 3.24. Consider the digital 2-manifold \mathbb{Z}^2 . Let m > 1 be a positive integer and define a cover $\{M_1, M_2\}$ of \mathbb{Z}^2 by $M_1 = \{(p_1, p_2) : p_1 < m\}$ and $M_2 = \{(p_1, p_2) : p_1 > 0\}$. The intersection of M_1 and M_2 is $\{(p_1, p_2) : 0 < p_1 < m\}$, which is a digital 2-manifold with boundary. Consider two functions $\alpha_1 : \mathbb{Z}^2 \to \mathbb{Z}$ and $\alpha_2 : \mathbb{Z}^2 \to \mathbb{Z}$ defined by

$$\alpha_1(p_1, p_2) = \begin{cases} m, & \text{if } p_1 \le 0\\ m - p_1, & \text{if } 0 < p_1 < m\\ 0, & \text{if } p_1 \ge m \end{cases}$$

and

$$\alpha_2(p_1, p_2) = \begin{cases} 0, & \text{if } p_1 \le 0\\ p_1, & \text{if } 0 < p_1 < m \\ m, & \text{if } p_1 \ge m \end{cases}$$

respectively. Each $\alpha_i \geq 0$ is digitally continuous for $i \in \{1,2\}$ and supported with the corresponding set M_i . When we consider any point $(p_1, p_2) \in \mathbb{Z}^2$, we observe that $\alpha_1(p_1, p_2) + \alpha_2(p_1, p_2) = m$. If we consider any point $(p_1, p_2) \in M_1 \cap M_2$, then we have that $N_4((p_1, p_2)) \cap \operatorname{sp}(\alpha_1)$ and $N_4((p_1, p_2)) \cap \operatorname{sp}(\alpha_2)$ are nonempty. Moreover, we get $\operatorname{sp}(\alpha_1) \subset M_1$ and $\operatorname{sp}(\alpha_2) \subset M_2$. Finally, we conclude that $\{\alpha_1, \alpha_2\}$ forms a digital partition of unity on $M_1 \cap M_2$ subordinate to a cover $\{M_1, M_2\}$.

4. CONCLUSION AND OPEN PROBLEMS

Many important homotopy, homology, and cohomology calculations in digital topology are performed on digital curves and surfaces, so studying digital manifolds takes these to the next level. It also opens the way for classifications of digital manifolds, a generalization of digital curves and digital surfaces often used in applications such as animated films, video games, and medical imaging. In topological robotics, a manifold of configurations is often used to model the space in which robots move. Therefore, the robot's journey from the beginning point to the destination point in motion planning difficulties turns into a path planning problem on the manifold. As an out-of-topology example, high-dimensional data often reside on low-dimensional manifolds. Thus, by examining these structures, manifold learning approaches offer data visualization and dimensionality reduction. These illustrations demonstrate the usage of manifolds in a variety of domains, including digital topology, ranging from pure mathematics to applied sciences.

The primary goal of this paper is to examine the definition of the term "digital manifold" based on digital images, not cells or simplexes. More precisely, the local homeomorphism property is explicitly investigated on digital images based on adjacency relations. After this, the rest of the paper discusses notions, such as submanifold, orientation, and partition of unity, related to the concept of digital manifold. Moreover, some counterexamples of digital images are presented. Besides, a few open problems are left for the readers in this section.

Open Problem 1: Let (M_1, κ) be a digital n_1 -manifold, (M_2, λ) a digital n_2 -manifold, and $n_3 \leq n_1 + n_2$. Then is it true or not that their Cartesian product $(M_1 \times M_2, \operatorname{NP}(\kappa, \lambda))$ is a digital n_3 -manifold?

Open Problem 2: Are the only digital images to which a digitally connected 1-manifold is digitally isomorphic the digital 1-sphere, the digital interval, and the set $M = \{(0,0), (0,1), (1,0), (1,1)\}$? Or are there any other digitally connected 1-manifolds? Do they always have orientations of 2?

Open Problem 3: Based on the solution of Open Problem 2, can we say that a digitally connected 1-manifold is either digitally contractible or digitally homotopy equivalent to a digital 1-sphere?

Open Problem 4: Is it true that every n-dimensional manifold with boundary lies in an n-dimensional manifold?

Open Problem 5: Can we define a "digital smooth (differentiable) manifold"? To answer this, it is valuable to comment on the derivative definitions in digital images. In addition, how do we define the orientation of a digital smooth manifold for n > 1?

References

- [1] C. Berge, Graphs and Hypergraphs, 2nd ed., North-Holland, Amsterdam (1976).
- [2] L. Boxer, A classical construction for the digital fundamental group, Journal of Mathematical Imaging and Vision, 10 (1999), 51-62.
- [3] L. Boxer, Homotopy properties of sphere-like digital images, Journal of Mathematical Imaging and Vision, 24 (2006), 167-175.
- [4] L. Boxer, Digital products, wedges, and covering spaces, Journal of Mathematical Imaging and Vision, 25 (2006), 169-171.
- [5] L. Boxer and İ. Karaca, Fundamental groups for digital products, Advances and Applications in Mathematical Sciences, 11(4) (2012), 161-180.
- [6] L. Boxer, Generalized normal product adjacency in digital topology, Applied General Topology, 18(2) (2017), 401-427.
- [7] G. Burak, Dijital Kohomoloji Grupları, PhD thesis, Adnan Menderes University, Aydın, Türkiye, 2014.
- [8] L. Chen, Gradually varied surfaces on digital manifold, In: Second International Conference on Industrial and Applied Mathematics, Washington, DC (1991).

- [9] L. Chen and J. Zhang, Digital manifolds: An intuitive definition and some properties, In: Proceedings on the Second ACM Symposium on Solid Modeling and Applications, pp. 459-460 (1993).
- [10] L. Chen, Discrete Surfaces and Manifolds, Scientific Practical Computing, Rockville, MD (2004).
- [11] Ö. Ege and I. Karaca, Cohomology theory for digital images, Romanian Journal of Information Science and Technology, 16(1) (2013), 10-28.
- [12] S.E. Han, Digital coverings and their applications, Journal of Applied Mathematics and Computing, 18 (2005), 487-495.
- [13] S.E. Han, Digital fundamental group and Euler characteristic of a connected sum of digital closed surfaces, Information Sciences, 177 (16) (2007), 3314-3326.
- [14] F. Hausdorff, Grundzüge der Mengenlehre, Veit & Company, Leipzig (1914).
- [15] G.T. Herman, Oriented surfaces in digital spaces, CVGIP: Graphical Models and Image Processing, 55 (1993), 381-396.
- [16] G.T. Herman, Geometry of digital spaces, In: Vision Geometry VII, Vol. 3454, SPIE, pp. 2-13 (1998).
- [17] İ. Karaca, L. Boxer, and A. Öztel, Topological invariants in digital images, Journal of Mathematical Sciences: Advances and Applications, 2 (2011), 109-140.
- [18] İ. Karaca and T. Vergili, Fiber bundles in digital images, Proceeding of 2nd International Symposium on Computing in Science and Engineering, 700(67) (2011), 1260-1265.
- [19] T.Y. Kong, A digital fundamental group, Computer Graphics, 13 (1989), 159-166.
- [20] T.Y. Kong and A. Rosenfeld, Digital topology: introduction and survey, Computer Vision, Graphics, and Image Processing, 48(3) (1989), 357-393.
- [21] J.M. Lee, Introduction to Topological Manifolds, Vol. 202, Springer Science & Business Media, (2010).
- [22] H. Poincaré, Analysis situs, Journal de l'École Polytechnique, 2(1) (1895), 1-123.
- [23] B. Riemann, Über die Hypothesen, welche der Geometrie zu Grunde liegen, Physikalische Blatter, 10(7) (1854), 296-306.
- [24] A. Rosenfeld, Connectivity in digital pictures, Journal of the ACM, 17(1) (1970), 146-160.
- [25] A. Rosenfeld, Digital topology, The American Mathematical Monthly, 86(8) (1979), 621-630.
- [26] A. Rosenfeld, Continuous functions on digital pictures, Pattern Recognition Letters, 4 (1986), 177-184.
- [27] E.H. Spanier, Algebraic Topology, Springer New York, NY (1966).
- [28] J.B. Tenenbaum, V.D. Silva, and J.C. Langford, A global geometric framework for nonlinear dimensionality reduction, Science, 290(5500) (2000), 2319-2323.

Melih İs, Ege University, Faculty of Sciences, Department of Mathematics, İzmir, Türkiye

Email address: melih.is@ege.edu.tr

İsmet Karaca, Ege University, Faculty of Sciences, Department of Mathematics, İzmir, Türkiye

Email address: ismet.karaca@ege.edu.tr