Fermi's golden rule in tunneling models with quantum waveguides perturbed by Kato class measures

Sylwia Kondej^{*} and Kacper Ślipko

Institute of Physics, University of Zielona Góra, ul. Szafrana 4a, 65246 Zielona Góra, Poland *e-mail: s.kondej@if.uz.zgora.pl, slipkokacper@gmail.com*

Abstract

In this paper we consider two dimensional quantum system with an infinite waveguide of the width d and a transversally invariant profile. Furthermore, we assume that at a distant ρ there is a perturbation defined by the Kato measure. We show that, under certain conditions, the resolvent of the Hamiltonian has the second sheet pole which reproduces the resonance at $z(\rho)$ with the asymptotics $z(\rho) = \mathcal{E}_{\beta;n} + \mathcal{O}\left(\frac{\exp(-\sqrt{2|\mathcal{E}_{\beta;n}|\rho})}{\rho}\right)$ for ρ large and with the resonant energy $\mathcal{E}_{\beta;n}$. Moreover, we show that the imaginary component of $z(\rho)$ satisfies Fermi's golden rule which we explicitly derive.

Keywords: Resonances in quantum system, tunnelling, resolvent poles, Fermi's golden rule. Mathematics Subject Classification: 47B38, 81Q10, 81Q15, 81Q80

1 Introduction

The problem studied in this paper concerns a two dimensional quantum system with a straight waveguide of width d perturbed by a distant potential, represented by the Kato class measure. The waveguide will be imitated by the potential

$$V_{\alpha} = -\alpha(x_2)\chi_{\Sigma}(x), \qquad (1.1)$$

where

$$\Sigma := \{ x = (x_1, x_2) : x_1 \in \mathbb{R}, x_2 \in [0, d] \},$$
(1.2)

and $\chi_{\mathcal{A}}(\cdot)$ denotes the characteristic function of the set $\mathcal{A} \subset \mathbb{R}^2$, i.e. it is equal 1 on \mathcal{A} and 0 on $\mathcal{A} \setminus \mathbb{R}^2$; moreover, $\alpha : [0,d] \to \mathbb{R}$ stands for a piecewise continuous function. Following the notation introduced in [13], we will call the interaction defined by V_{α} 'soft waveguide' to emphasize the possibility of tunnelling out of Σ . As we have already mentioned, apart from the waveguide Σ , we introduce into the system a trap potential localized on a compact set $\Omega \subset \mathbb{R}^2$ and defined by a Radon measure $\mu_{\Omega} \equiv \mu$ belonging to the Kato class and supported on Ω . We assume that $\Omega \cap \Sigma = \emptyset$ and reserve the special notation ρ for the distance $\rho := \min_{x \in \Sigma, y \in \Omega} |x - y|$. The Hamiltonian of the system can be defined as the operator associated to the sesquilinear form

$$\mathscr{E}_{\alpha,\beta\mu}[f] = \int_{\mathbb{R}^2} |\nabla f(x)|^2 \,\mathrm{d}x - \int_{\Sigma} \alpha(x_2) |f(x)|^2 \,\mathrm{d}x - \beta \int_{\mathbb{R}^2} |I_{\mu}f(x)|^2 \,\mathrm{d}\mu(x) \,, \quad f \in W^{1,2}(\mathbb{R}^2) \,, \tag{1.3}$$

where $W^{1,2}(\mathbb{R}^2)$ stands for the Sobolev space $W^{1,2}(\mathbb{R}^2) = \{f \in L^2(\mathbb{R}^2) : \partial_i f \in L^2(\mathbb{R}^2), i = 1, 2\}^1$ and I_μ denotes the space embedding $I_\mu : W^{1,2}(\mathbb{R}^2) \to L^2_\mu \equiv L^2(\mathbb{R}^2, d\mu)$ and $\beta \in \mathbb{R}$. The role of the parameter β

^{*}Corresponding author.

¹The symbols ∂_i stand for the distributional derivatives labelled by two cartesian variables if \mathbb{R}^2 .




Figure 1: A schematic illustration of the model. Σ represents the structure of the waveguide, and Ω represents the perturbation.

at this general stage is to maintain consistency with further notations. The resulting Hamiltonian will be denoted $H_{\alpha,\beta\mu}$.

The perturbation determined by μ allows various realisations that are interesting from a quantum point of view. For example, we can consider a continuous function $V : \mathbb{R}^2 \to \mathbb{R}$ with the support Ω . Then, for any Borel set $\mathcal{B} \subset \mathbb{R}^2$, we have $\mu(\mathcal{B}) = \int_{\mathcal{B}} V(x) dx$. Note that $\Omega \subset \mathbb{R}^2$ can be defined as a set of a lower dimension. This leads to another important example of the measure μ defined by the delta potential, named also the delta interaction. To explain in more detail, assume that Ω is a finite line (loop or open arc) in \mathbb{R}^2 . Then $\mu(\mathcal{B}) = \lim(\mathcal{B} \cap \Omega)$, where $\lim(\cdot)$ stands for the linear measure. In this situation the perturbation formally corresponds to $-\beta \delta(x - \Omega)$, where $\delta(\cdot - \Omega)$ stands for the Dirac delta supported on Ω . The examples of perturbations belonging to the Radon measure class are shown in Fig. 1. For the clarity of the presentation we will first demonstrate the reasoning for the simplified model, namely $\mu(\mathcal{B}) = \int_{\mathcal{B}\cap\Omega} dx$, where $\Omega \subset \mathbb{R}^2$ is compact and $\alpha(\cdot) = \alpha$, with $\alpha > 0$. For this case we use a special notation for the Hamiltonian:

$$H_{\alpha,\beta} = -\Delta - \alpha \chi_{\Sigma}(x) - \beta \chi_{\Omega}(x) \,. \tag{1.4}$$

Modern nanotechnology allows the fabrication of semiconductors with different shapes and geometries. Leveraging a high mobility material such as $Ga_{1-x}Al_xAs$ we can produce a quasi one or two dimensional electron gas in quantum wires, waveguides or layers. The question of what properties of semiconductor structures affect the quantum transport, is essential from the perspective of the functioning of microelectronic devices. Inspired by the question of the relation between architecture of the semiconductor structure and spectral characteristics, we investigate the resonant quantum model with an open channel and a trap inducing resonances due to the tunneling. Such a system is govened by a certain realization of $H_{\alpha,\beta\mu}$, for example by (1.4) and our main aim is to understand how the distance between Σ and Ω , as a simple parameter of the system architecture, affects the resonances.

To take a closer look at the spectral situation we auxiliary assume that $\beta = 0$ and consider the system governed by $H_{\alpha,0}$, i.e. which is employed only with the straight waveguide. The translational invariance allows to decompose $H_{\alpha,0}$ onto longitudinal and transversal components. The latter is determined by the one dimensional Hamiltionian h_{α} with the potential $-\alpha(\cdot)\chi_{[0,d]}$. Obviously, the essential spectrum of h_{α} ranges from 0 to infinity. In the following we assume that h_{α} admits also discrete spectral point below 0. Let $\{E_{\alpha;1} \leq ... \leq E_{\alpha,N_{\alpha}}\}$ stand for the complete set of the discrete eigenvalues of h_{α} ; we denote also $\mathcal{N}_{\alpha} = \{1, 2, ..., N_{\alpha}\}$. Backing to the two dimensional system with the Hamiltonian $H_{\alpha,0}$ we can state that its spectrum covers

$$\sigma(H_{\alpha,0}) = \sigma_{\rm ess}(H_{\alpha,0}) = [E_{\alpha}, \infty), \qquad (1.5)$$

where $E_{\alpha} = E_{1;\alpha}$ is the lowest eigenvalue of h_{α} . Under certain conditions on μ , we can also show that the essential spectrum remains stable under the perturbation which can be formally written as $-\beta\mu$, i.e. $\sigma_{\rm ess}(H_{\alpha,\beta\mu}) = [E_{\alpha},\infty)$. However, it can produce discrete eigenvalues below the threshold of the essential spectrum.

The main topic of this paper it devoted to the problem of resonances that may emerge if the distance ρ is very large. The resonances and the related metastable states are a kind of spectral memory about the decoupled systems $H_{\alpha,0}$ and $H_{0,\beta\mu}$, which formally corresponds to $\rho = \infty$. To explain in more detail, assume that $\alpha = 0$ and then the corresponding Hamiltonian $H_{0,\beta\mu}$ has the eigenvalues $\{\mathcal{E}_{\beta;1} \leq ... \leq \mathcal{E}_{\beta;N_{\beta}}\}$, $\mathcal{N}_{\beta} := \{1,..,N_{\beta}\}$ with a corresponding eigenfunction $\{\omega_{\beta;n}\}_{n\in\mathcal{N}_{\beta}}$. If for a given *n* there exists $\mathcal{E}_{\beta;n}$ such that $\mathcal{E}_{\beta;n} > E_{\alpha}$ then we may expect that a metastable state with the resonant energy $\mathcal{E}_{\beta;n}$ can appear in the system governed by $H_{\alpha,\beta\mu}$. This will be manifested as a second sheet resolvent pole residing in a certain neighbourhood of $\mathcal{E}_{\beta;n}$.

Main results of the paper. The main result of this paper can be formulated as follows.

• Assume $\mathcal{E}_{\beta;n}$ is a simple eigenvalue of $H_{0,\beta\mu}$ and it belongs to $(E_{\alpha;j}, E_{\alpha;j+1})$ for $j \in \mathcal{N}_{\alpha}$. The resolvent of $H_{\alpha,\beta\mu}$ has a second sheet analytic continuation to the lower half-plane and there exists a unique resolvent pole in a neighbourhood of $\mathcal{E}_{\beta;n}$ that admits the asymptotics

$$z_n(\rho) = \mathcal{E}_{\beta;n} + \mathcal{O}\left(\frac{\mathrm{e}^{-\sqrt{2|\mathcal{E}_{\beta;n}|}\rho}}{\rho}\right).$$
(1.6)

• The dominated term of the imaginary component of pole takes the form

$$\Gamma_{n}(\rho) = -\beta^{2} \sum_{k=1}^{j} \frac{\pi}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \Big[\Big| \int_{\Omega} I_{\mu}(\omega_{\beta;n}(x)^{*}\varphi_{k}(x, p_{1})) d\mu(x) \Big|_{p_{1} = (\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}}^{2} \\ + \Big| \int_{\Omega} I_{\mu}(\omega_{\beta;n}(x)^{*}\varphi_{k}(x, p_{1})) d\mu(x) \Big|_{p_{1} = -(\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}}^{2} \Big],$$
(1.7)

where $\varphi_k(x, p_1)$ are the generalized eigenfunctions of $H_{\alpha,0}$ and let us recall that $\omega_{\beta;n}$ stand for the eigenfunctions of $H_{0,\beta\mu}$; the above integrals depend on the argument p_1 and the indexes mean that they are determined at the specific points $p_1 = \pm (\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}$. For the special case of Hamiltonian defined by (1.4), the dominated term turns to

$$\Gamma_{n}(\rho) = -\beta^{2} \sum_{k=1}^{j} \frac{\pi}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \left[\left| \int_{\Omega} \omega_{\beta;n}(x)^{*} \varphi_{k}(x, p_{1}) \mathrm{d}x \right|_{p_{1} = (\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}}^{2} + \left| \int_{\Omega} \omega_{\beta;n}(x)^{*} \varphi_{k}(x, p_{1}) \mathrm{d}x \right|_{p_{1} = -(\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}}^{2} \right].$$
(1.8)

From the physical point of view, the imaginary component of pole plays the main rule in the time evolution of metastable state. Fermi's golden rule plays an essential role in physics. It describes the rate of transition from an energy eigenstate of a quantum system to an energy continuum as an effect of a weak perturbation. The corresponding formula was, first time, obtained by Dirac [9] and then, after more than twenty years, it was analyzed and called as 'golden rule' by Fermi [20]. From the physical perspective, distantly located trap allows for controlling scattering processes and quantum transport in the waveguide, influencing tunneling and dissipative processes. The distance between the trap and the waveguide can be adjusted to impact the system dynamics. Studying these decay mechanisms, especially through explicit derivations like Fermi's golden rule, is crucial for optimizing such systems. This provides insights into key components of quantum technology such as wires and waveguides, and facilitates the optimization of related processes.

We would like to conclude this introduction by saying a few words about how the results obtained in this paper fit into a wider context. The work belongs to the line of research that is dedicated to the exploration of the spectral properties of quantum systems with soft waveguides. These types of systems have been analyzed mainly from the point of view of the existence of bound states, cf. [13, 14, 15, 18, 19, 22, 30]. The problem of resonances in soft waveguides remains a lot of questions open. It is worth mentioning that the problem of resonances induced by breaking the symmetry in the soft waveguide systems has been analyzed in [26] by the first author of the present paper. In [26] the resonances were memories of the embedded eigenvalues preserved by symmetry of the system. In the current paper, the resonances are induced by the tunneling, which means that the strategy for recovering resonance pole is necessarily not the same, although the main technical concepts are borrowed from [26]. On the other hand, in the present paper we discuss systems with a general class of tunnel traps defined by the Kato class measures. This aspect is interesting from both mathematical and physical perspectives. First, it demonstrates that the resonance techniques (partially developed in [26]) can be extended to a much broader scenario. Moreover, it shows that the trap can be defined not only by a regular potential but also by lower-dimensional structures, such as wires. The latter is interesting, in particular, from the physical point of view. Additionally, the present paper provides an explicit formulation of Fermi's golden rule for this generalized case, which in [26] was presented only implicitly in terms of the spectral resolution, a formulation less familiar to physicists.

Note that the problem of resonances induced by the effect of tunnelling in the model with an infinite straight line and a distant point interaction has been analyzed in [16] and, on the other hand, the resonances in the three dimensional systems with point potentials have been investigated, for example in [8, 31], see also [3]. It is also worth mentioning that problem of resonances in, so called, hard wall (Dirichlet) waveguides has been undertaken in series of papers, cf. [5, 10, 11, 29, 25] see also references in [12]. However, in these systems, contrary to the model discussed in this paper, a particle cannot escape from the waveguide because of the infinite potential barrier at the boundary.

The paper is organized as follows: In Section 2 we provide the description of Hamiltonians $H_{\alpha,\beta}$, $H_{\alpha,\beta}$ and $H_{\alpha,\beta}$ corresponding to (1.4). We state preliminary facts about the spectra and derive the resolvents of the investigated Hamiltonians. In Section 3 we show that, under certain conditions, the resolvent of $H_{\alpha,\beta}$ has a unique second sheet pole and derive its asymptotics. In Section 4 we analyze the imaginary part of the pole and show that Fermi's golden rule is fulfilled in the form of (1.8). Finally, in Section 5 we generalize the result for large class of potential defined by $H_{\alpha,\beta\mu}$ and related to (1.3).

2 Description of simplified model: influence of distant well on a straight waveguide

In this section, we consider a two dimensional quantum mechanical model with the potential supported on a straight, infinite planar waveguide and a distant well potential located on a compact set $\Omega \subset \mathbb{R}^2$ with the smooth boundaries $\partial \Omega$. We assume that the waveguide has a width d, therefore it can be defined as follows

$$\Sigma := \{ x = (x_1, x_2) : x_1 \in \mathbb{R}, x_2 \in [0, d] \}.$$
(2.1)

Furthermore, we assume that $\Omega \cap \Sigma = \emptyset$ and denote the distance

$$\rho := \min_{x \in \Sigma, y \in \Omega} |x - y|.$$
(2.2)

Due to the orientation of Σ we can also write $\rho = \min_{x \in \Sigma, y \in \Omega} |x_2 - y_2|$. The parameter ρ will play the essential role in the further discussion.

Let α and β be positive numbers, and using them, we define the Hamiltonian

$$H_{\alpha,\beta} = -\frac{d^2}{dx_1^2} - \frac{d^2}{dx_2^2} - \alpha \chi_{\Sigma}(x) - \beta \chi_{\Omega}(x) : W^{2,2}(\mathbb{R}^2) \to L^2(\mathbb{R}^2), \qquad (2.3)$$

where the Sobolev space $W^{2,2}(\mathbb{R}^2) = \{f \in L^2(\mathbb{R}^2) : \partial_i \partial_j f \in L^2(\mathbb{R}^2), i, j \in \{1,2\}\}$ determines the maximal domain. In fact, the above Hamiltonian depends on Σ and Ω as well, however, to avoid over-indexing, we do not involve Σ and Ω into the notation for the Hamiltonian. $H_{\alpha,\beta}$ defined by (2.3) determines a particular example of (1.1).

2.1 Insight into a strip model

We auxiliary consider the situation $\beta = 0$, i.e. we remove the well potential supported on Ω , leaving only the strip interaction. The corresponding Hamiltonian $H_{\alpha,0}$ has the translational symmetry, so we can decouple it into transverse and longitudinal components. The latter carries the information only about the kinetic energy. The transverse component of Hamiltonian $H_{\alpha,0}$ is defined by

$$h_{\alpha} = -\frac{d^2}{dx_2^2} - \alpha \chi_{[0,d]}(x_2) \, : \, W^{2,2}(\mathbb{R}) \to L^2(\mathbb{R}) \,, \tag{2.4}$$

which determines the standard one dimensional Hamiltonian with a rectangular well of depth α . The negative eigenvalues and eigenfunctions of h_{α} formed in the ascending order will be denoted

$$\{E_{\alpha;1} \le E_{\alpha;2} \le \dots \le E_{\alpha;N_{\alpha}}\},\tag{2.5}$$

and

$$\{\phi_{\alpha;n}\}_{n\in\mathcal{N}_{\alpha}}, \quad \mathcal{N}_{\alpha}:=\{1,...N_{\alpha}\}$$
(2.6)

stand for the corresponding eigenfunctions. Then $H_{\alpha,0}$ can be represented as

$$H_{\alpha,0} = -\frac{d^2}{dx_1^2} \otimes \mathbf{I} + \mathbf{I} \otimes \mathbf{h}_{\alpha} \,. \tag{2.7}$$

Consequently, the spectrum of $H_{\alpha,0}$ admits only the essential component of the form

$$\sigma_{\rm ess}(H_{\alpha,0}) = [E_{\alpha}, \infty), \quad \text{where} \quad E_{\alpha} = E_{\alpha;1}.$$
(2.8)

Notation. In the following discussion we will use the notation I for the unit operator independently of space where it acts.

2.1.1 Resolvent of $H_{\alpha,0}$

In this section we are going to derive the resolvent of $H_{\alpha,0}$ which will be the basic tool in the analysis of resonances. Let us recall that the 'free' resolvent is given by $R(z) = (-\Delta - z)^{-1}$, and it can be represented as an integral operator with a kernel defined by the Macdonald function $K_0(\cdot)$, also known as the modified Bessel function of the second kind.²

In the present case, when the potential is defined by a negative coupling constant $-\alpha$, the Birman–Schwinger operator is given by $-\alpha\chi_{\Sigma}R(z)\chi_{\Sigma}$: $L^{2}(\Sigma) \rightarrow L^{2}(\Sigma)$. Consequently, for z living in the resolvent set $\varrho(H_{\alpha,0})$ the operator

$$I - \alpha \chi_{\Sigma} R(z) \chi_{\Sigma} : L^{2}(\Sigma) \to L^{2}(\Sigma)$$
(2.10)

is invertible. Using the Birman–Schwinger concept we represent the resolvent $R_{\alpha,0}(z) = (H_{\alpha,0} - z)^{-1}$ as

$$R_{\alpha,0}(z) = R(z) + \alpha R(z)\chi_{\Sigma}[I - \alpha\chi_{\Sigma}R(z)\chi_{\Sigma}]^{-1}\chi_{\Sigma}R(z), \qquad (2.11)$$

cf. [6, 7, 23, 32, 36].

To proceed further we are going to present the resolvent $R_{\alpha,0}(z)$ in an alternative form that reveals transverse symmetry and reflexes the decomposition (2.7); a similar construction has been carried out in [15, 26]. Note that the spectral family (equivalently, the resolution of the identity) associated to the operator h_{α} is determined the eigenprojectors $P_{\alpha;n} = (\phi_{\alpha;n}, \cdot)_{L^2(\mathbb{R})} \phi_{\alpha;n}$ and the component corresponding to the

$$R(z)f = \frac{1}{2\pi} \int_{\mathbb{R}^2} K_0(k_z |\cdot -y|) f(y) \, \mathrm{d}y, \quad f \in L^2(\mathbb{R}^2),$$
(2.9)

where $k_z = -i\sqrt{z}$ with $\Im\sqrt{z} > 0$ and the kernel of R(z) defines the Green's function of the operator $-\Delta - z$.

²Alternatively, the kernel of R(z) can be expressed using the Hankel function as $\frac{i}{4}H_0(\sqrt{z}|\cdot-\cdot|)$. Specifically:

continuum of h_{α} which we denote $E_{\alpha}(\cdot)$. Note that the longitudinal component of $H_{\alpha,0}$, defined by $-\frac{d^2}{dx_1^2} \otimes I$ on $L^2(\mathbb{R}) \otimes L^2(\mathbb{R})$, reproduces the spectral family for the operator $-\frac{d^2}{dx_1^2}$ acting in $L^2(\mathbb{R})$. We denote it as $E_0(\cdot)$, and the explicit form of $E_0(\cdot)$ is given by:

$$E_0(\lambda)f = \frac{1}{(2\pi)} \int_{|\xi|^2 \le \lambda} f(\xi) e^{ix \cdot \xi} \,\mathrm{d}\xi.$$
(2.12)

By means of the introduced notation we can write

$$H_{\alpha,0}\left(\mathbf{I}\otimes P_{\alpha;n}\right) = \int_{[0,\infty)} (\lambda + E_{\alpha;n}) \,\mathrm{d}E_0(\lambda) \otimes P_{\alpha;n} \,. \tag{2.13}$$

This, consequently, allows to define components of $R_{\alpha,0}(z)$ related to the projection onto $L^2(\mathbb{R}) \otimes P_{\alpha}L^2(\mathbb{R})$, where $P_{\alpha} = \sum_{n \in \mathcal{N}_{\alpha}} P_{\alpha,n}$, and $L^2(\mathbb{R}) \otimes P_{\alpha}^{\perp}L^2(\mathbb{R})$, where P_{α}^{\perp} stands for the orthogonal complement of P_{α} . Precisely,

$$R_{\alpha,0}^{d}(z) := R_{\alpha,0}(z) \left(\mathbf{I} \otimes P_{\alpha} \right) = \sum_{n \in \mathcal{N}_{\alpha}} \int_{[0,\infty)} \frac{\mathrm{d}E_{0}(\lambda)}{\lambda + E_{\alpha;n} - z} \otimes P_{\alpha;n}$$
(2.14)

and

$$R_{\alpha,0}^{c}(z) := R_{\alpha,0}(z) \left(\mathbf{I} \otimes P_{\alpha}^{\perp} \right) = \int_{[0,\infty)} \int_{[0,\infty)} \frac{\mathrm{d}E_{0}(\lambda) \otimes \mathrm{d}E_{\alpha}(\lambda')}{\lambda + \lambda' - z} \,.$$
(2.15)

Note that $E_0(\cdot)$ can be also expanded by means of the plane waves $(2\pi)^{-1/2}e^{ix_1p_1}$ which yields

$$\left(-\frac{d^2}{dx_1^2} + E_{\alpha;n} - z\right)^{-1} f(x_1) = \int_{[0,\infty)} \frac{\mathrm{d}E_0(\lambda)}{\lambda + E_{\alpha;n} - z} f(x_1) = \frac{1}{2\pi} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{\mathrm{e}^{ip_1(x_1 - y_1)}}{p_1^2 + E_{\alpha;n} - z} f(y_1) \,\mathrm{d}p_1 \,\mathrm{d}y_1 \,. \tag{2.16}$$

Furthermore, the one dimensional Green's function which appears in the last equivalence admits also representation

$$\frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{ip_1(x_1-y_1)}}{p_1^2 + E_{\alpha;n} - z} \, \mathrm{d}p_1 = \frac{i}{2} \frac{e^{i\tau_{\alpha;n}(z)|x_1-y_1|}}{\tau_{\alpha;n}(z)} \,, \tag{2.17}$$

where $\tau_{\alpha;n}(z) := \sqrt{z - E_{\alpha;n}}$ with the square root determined on the first analytical sheet, i.e. $\Im \tau_{\alpha;n}(z) > 0$. Summarizing, we get the decomposition

$$R_{\alpha,0}(z) = R^d_{\alpha,0}(z) + R^c_{\alpha,0}(z), \qquad (2.18)$$

where

$$R_{\alpha,0}^{d}(z)f(x) = \frac{1}{2\pi} \sum_{n \in \mathcal{N}_{\alpha}} \int_{\mathbb{R}^{2}} \int_{\mathbb{R}} \frac{e^{ip_{1}(x_{1}-y_{1})}}{p_{1}^{2} + E_{\alpha;n} - z} \phi_{\alpha;n}(x_{2})\phi_{\alpha;n}(y_{2})^{*}f(y) \,\mathrm{d}p_{1} \,\mathrm{d}y$$
$$= \frac{i}{2} \sum_{n \in \mathcal{N}_{\alpha}} \int_{\mathbb{R}^{2}} \frac{e^{i\tau_{\alpha;n}(z)|x_{1}-y_{1}|}}{\tau_{\alpha;n}(z)} \phi_{\alpha;n}(x_{2})\phi_{\alpha;n}(y_{2})^{*}f(y) \,\mathrm{d}y \,,$$
(2.19)

where $y = (y_1, y_2)$ and

$$R_{\alpha,0}^{c}(z)f(x) = \frac{i}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{\mathrm{e}^{i\tau_{\alpha}(z,p_{2})|x_{1}-y_{1}|}}{\tau_{\alpha}(z,p_{2})} \psi_{\alpha}(x_{2},p_{2})\psi_{\alpha}(y_{2},p_{2})f(y)\,\mathrm{d}y\,\mathrm{d}p_{2}\,,\tag{2.20}$$

where $\tau_{\alpha}(z, p_2) := \sqrt{z - p_2^2}$ with $\Im\sqrt{z - p_2^2} > 0$, and $\psi_{\alpha}(\cdot, \cdot)$ stand for the generalized eigenfunctions of h_{α} (constructed from plane waves). The similar decomposition were used in [15].

2.2 Quantum well system without straight strip

In order to collect the most important information about the spectral properties of the system governed by $H_{\alpha,\beta}$, in this section we will auxiliary consider $H_{0,\beta}$. Since the potential is negative we can conclude that the negative discrete spectrum is non empty, cf. [33, XIII.11]. The discrete eigenvalues of $H_{0,\beta}$ formed in the ascending order will be denoted as

$$\{\mathcal{E}_{\beta;1} \le \mathcal{E}_{\beta;2} \le \dots \le \mathcal{E}_{\beta;N_{\beta}}\}\tag{2.21}$$

and, let

$$\{\omega_{\beta;n}\}_{n\in\mathcal{N}_{\beta}}, \quad \mathcal{N}_{\beta}:=\{1,\dots N_{\beta}\}, \qquad (2.22)$$

stand for the corresponding eigenfunctions.

We again use the Birman–Schwinger resolvent $R_{0,\beta}(z) = (H_{0,\beta} - z)^{-1}$ which in this case takes the form

$$R_{0,\beta}(z) = R(z) + \beta R(z)\chi_{\Omega}(\mathbf{I} - \beta\chi_{\Omega}R(z)\chi_{\Omega})^{-1}\chi_{\Omega}R(z), \quad z \in \varrho(H_{0,\beta}).$$
(2.23)

The Birman–Schwinger principle allows to associate zeros of $I - \beta \chi_{\Omega} R(\mathcal{E}_{\beta;n}) \chi_{\Omega} : L^2(\Omega) \to L^2(\Omega)$ and eigenfunctions $\omega_{\beta;n}$ of $H_{0,\beta}$ in the following way

$$\omega_{\beta;n} = R(\mathcal{E}_{\beta;n})\chi_{\Omega}w_{\beta;n}, \quad \text{where} \quad w_{\beta;n} \in \ker[\mathbf{I} - \beta\chi_{\Omega}R(\mathcal{E}_{\beta;n})\chi_{\Omega}] \quad \text{and} \quad n \in \mathcal{N}_{\beta}, \tag{2.24}$$

cf. [6, 7, 23, 32]. Let S_n denote a sufficiently small neighbourhood of $\mathcal{E}_{\beta;n}$. With a view to further discussion we prove the following statement.

Lemma 2.1 Assume that $\mathcal{E}_{\beta;n}$ is a simple discrete eigenvalue of $H_{0,\beta}$ and $z \in \mathcal{S}_n \setminus \{\mathcal{E}_{\beta;n}\}$. Then the operator $(I - \beta \chi_\Omega R(z) \chi_\Omega)^{-1} : L^2(\Omega) \to L^2(\Omega)$ admits the representation

$$(\mathbf{I} - \beta \chi_{\Omega} R(z) \chi_{\Omega})^{-1} = \frac{1}{\beta} \frac{1}{\mathcal{E}_{\beta;n} - z} \mathbf{P}_{\beta;n} + A_n(z), \qquad (2.25)$$

where $P_{\beta;n} = (w_{\beta;n}, \cdot)_{L^2(\Omega)} w_{\beta;n}$ and $A_n(z)$ is analytic in S_n .

Proof. Note that $R_{0,\beta}(z)$ is the resolvent of a self adjoint operator with isolated spectral point at $\mathcal{E}_{\beta;n}$ with corresponding eigenvector $\omega_{\beta;n}$ and eigenprojector $P_{\beta;n}$. Thus, for $z \in S_n \setminus \{\mathcal{E}_{\beta;n}\}$ we can write

$$R_{0,\beta}(z) = \frac{1}{\mathcal{E}_{\beta;n} - z} P_{\beta;n} + B_n(z) : L^2(\mathbb{R}^2) \to L^2(\mathbb{R}^2), \qquad (2.26)$$

where $B_n(z)$ acts in $L^2(\mathbb{R}^2)$ and it is analytic in S_n ; the above follows from the spectral resolution of the $R_{0,\beta}(z)$, cf. [24, V-(3.21)]. The singularities coming from the remaining isolated points contribute to $B_n(z)$. On the other hand, since $(I - \beta \chi_\Omega R(z) \chi_\Omega)^{-1}$ has the same pole of the multiplicity one, it can be decomposed as

$$(\mathbf{I} - \beta \chi_{\Omega} R(z) \chi_{\Omega})^{-1} = \frac{1}{\mathcal{E}_{\beta;n} - z} \tilde{P}_n(z) + \tilde{B}_n(z) : L^2(\Omega) \to L^2(\Omega) , \qquad (2.27)$$

where $\tilde{B}_n(z)$ stands for an analytic component and $\tilde{P}_n(z)$ is a projector acting in the space $L^2(\Omega)$, cf. [21]. Comparing the above with (2.26) and taking into account the resolvent formula (2.23), we can write

$$P_{\beta;n} = \beta R(z) \chi_{\Omega} \tilde{P}_n(z) \chi_{\Omega} R(z) .$$
(2.28)

Furthermore, using the analyticity of the resolvent R(z) for $z \in S_n$, i.e.

$$R(z) = R(\mathcal{E}_{\beta;n}) + (\mathcal{E}_{\beta;n} - z)R(z)R(\mathcal{E}_{\beta;n})$$
(2.29)

we obtain from (2.23) that $\beta R(\mathcal{E}_{\beta;n})\chi_{\Omega}\tilde{P}_n(\mathcal{E}_{\beta;n})\chi_{\Omega}R(\mathcal{E}_{\beta;n}) = P_{\beta;n} + \mathcal{O}_{\text{op.u.}}(z - \mathcal{E}_{\beta;n})$, where the error term is understood in the norm operator sense. Implementing $P_{\beta;n} = (w_{\beta;n}, \cdot)_{L^2(\Omega)}w_{\beta;n}$ which, by definition of $\omega_{\beta;n}$ and $w_{\beta,n}$ (see (2.24)) satisfies $P_{\beta;n} = R(\mathcal{E}_{\beta;n})P_{\beta;n}R(\mathcal{E}_{\beta;n})$, we state that $\tilde{P}_n(\mathcal{E}_{\beta;n}) = \beta^{-1}P_{\beta;n}$ and eq. (2.27) turns to

$$(\mathbf{I} - \beta \chi_{\Omega} R(z) \chi_{\Omega})^{-1} = \frac{1}{\beta} \frac{1}{\mathcal{E}_{\beta;n} - z} \mathbf{P}_{\beta;n} + A_n(z), \qquad (2.30)$$

where $A_n(z)$ acting in $L^2(\Omega)$ is analytic in \mathcal{S}_n . This completes the proof.

2.3 Resolvent of $H_{\alpha,\beta}$ and preliminary facts about its spectral properties

Considering as a starting point the system governed by $H_{\alpha,0}$ with corresponding resolvent $R_{\alpha,0}(z)$, given by (2.11), we implement the pertubation determined by $-\beta\chi_{\Omega}$. To apply this strategy we assume that a complex number z is such that the operator $I - \beta\chi_{\Omega}R_{\alpha,0}(z)\chi_{\Omega} : L^2(\Omega) \to L^2(\Omega)$ is invertible. After adding the potential $-\beta\chi_{\Omega}$ to the system governed by $H_{\alpha,0}$ the resolvent takes the form

$$R_{\alpha,\beta}(z) = R_{\alpha,0}(z) + \beta R_{\alpha,0}(z)\chi_{\Omega}(\mathbf{I} - \beta\chi_{\Omega}R_{\alpha,0}(z)\chi_{\Omega})^{-1}\chi_{\Omega}R_{\alpha,0}(z), \qquad (2.31)$$

cf. [6, 7, 23, 32].

Essential spectrum. The stability of the essential spectrum follows from Weyl's Theorem, cf. [33, Thm. XIII.14]. Indeed, assuming the $z \in \rho(H_{\alpha,0})$ and employing compactness of Ω one can show that both $R(z)\chi_{\Omega}$: $L^2(\Omega) \to L^2(\mathbb{R}^2)$ and $\chi_{\Sigma}R(z)\chi_{\Omega} : L^2(\Omega) \to L^2(\Sigma)$ are compact; we postpone a detailed analysis to the next section, see Lemma 3.1. Applying now (2.11) we obtain the compactness of $R_{\alpha,0}(z)\chi_{\Omega}$. This, in view of Weyl's Theorem, yields

$$\sigma_{\rm ess}(H_{\alpha,\beta}) = \sigma_{\rm ess}(H_{\alpha,0}) = [E_{\alpha},\infty).$$
(2.32)

Remark 2.2 Although the discrete spectrum of $H_{\alpha,\beta}$ does not belong to the main line of this paper, it is worth mentioning that the attractive potential $-\beta\chi_{\Omega}$ leads to at least one discrete eigenvalue below E_{α} . One can establish this by using the variational method and constructing a trial function with separate variables, where ϕ_{α} represents the transverse direction and a positive propagating function is defined in the logitudinal direction. The details are presented in [26].

3 Resonances induced by the tunnelling

3.1 Analytic extension of $\chi_{\Omega}R_{\alpha,0}(z)\chi_{\Omega}$

To analyze the resonances we will adopt the strategy of finding the poles of the resolvent (2.31) as zeros of $I - \beta \chi_{\Omega} R_{\alpha,0}(z) \chi_{\Omega}$. Therefore, our first aim is to construct a second sheet continuation of $\chi_{\Omega} R_{\alpha,0}(z) \chi_{\Omega}$. We start our discussion from the analysis of $\tau_{\alpha;n}(z)$ (contributing $\chi_{\Omega} R_{\alpha,0}^d(z) \chi_{\Omega}$) which has been defined so far for the first Riemann sheet, i.e. $\Im \tau_{\alpha;n}(z) > 0$. For each $n \in \mathcal{N}_{\alpha}$, the function $z \mapsto \tau_{\alpha;n}(z)$ admits a second sheet analytic continuation through $[E_{\alpha;n}, \infty)$ and the values of $\Im \tau_{\alpha;n}(z)$ are negative in the lower complex half-plane. Consequently, for each $k \in \mathcal{N}_{\alpha}$, we can construct the lower half-plane analytic continuation of

$$\chi_{\Omega} R^d_{\alpha,0}(z) \chi_{\Omega} = \chi_{\Omega}(x) \Big[\frac{i}{2} \sum_{n \in \mathcal{N}_{\alpha}} \frac{\mathrm{e}^{i\tau_{\alpha;n}(z)|x_1 - y_1|}}{\tau_{\alpha;n}(z)} \phi_{\alpha;n}(x_2) \phi_{\alpha;n}(y_2)^* \Big] \chi_{\Omega}(y) , \qquad (3.1)$$

via the segments $(E_{\alpha;k}, E_{\alpha;k+1})$. More precisely, assuming that $z = \lambda + i\varepsilon$, where $\lambda \in (E_{\alpha;k}, E_{\alpha;k+1})$ we move to the lower half-plane, i.e. varying with ε from positive to negative values and defining $\Im \tau_{\alpha;n}(z) < 0$ if $n \leq k$ and $\Im \tau_{\alpha;n}(z) > 0$ if n > k. We insert it into (3.1) and, in this way, we construct the second sheet component of $\chi_{\Omega} R^d_{\alpha,0}(z)\chi_{\Omega} : L^2(\Omega) \to L^2(\Omega)$. On the other hand, the component $R^c_{\alpha,0}(z)$ does not require any additional analysis, since it is analytic for $z = \lambda + i\varepsilon$, where $\lambda \in (E_{\alpha;k}, E_{\alpha;k+1})$ and ε belongs to a neighbourhood of 0. In view of (2.18), this constitutes, for any $k \in \mathcal{N}_{\alpha}$, the second sheet continuation of $\chi_{\Omega} R_{\alpha,0}(z)\chi_{\Omega}$ via $(E_{\alpha;k}, E_{\alpha;k+1})$ to a certain set $\mathcal{C}_k \subset \mathbb{C}_-$. We remain the same notation for the second sheet operator valued function $\mathcal{C}_k \ni z \mapsto \chi_{\Omega} R_{\alpha,0}(z)\chi_{\Omega}$. The above reasoning has also been adopted in [26].

3.2 Zeros of Birman–Schwinger operator

To find out the poles of resolvent determined by zeros of $I - \beta \chi_{\Omega} R_{\alpha,0}(z) \chi_{\Omega} : L^2(\Omega) \to L^2(\Omega)$ we adopt (2.11) which allows to state that z satisfies ker $[I - \beta \chi_{\Omega} R_{\alpha,0}(z) \chi_{\Omega}] \neq \emptyset$ if and only if

$$\ker[\mathbf{I} - \beta \chi_{\Omega} R(z) \chi_{\Omega} - G_{\alpha,\beta}(z)] \neq \emptyset, \qquad (3.2)$$

where

$$G_{\alpha,\beta}(z) = \beta \, \alpha \, \chi_{\Omega} R(z) \chi_{\Sigma} (\mathbf{I} - \alpha \chi_{\Sigma} R(z) \chi_{\Sigma})^{-1} \chi_{\Sigma} R(z) \chi_{\Omega} \, : \, L^{2}(\Omega) \to L^{2}(\Omega) \, . \tag{3.3}$$

Our aim is to express the condition (3.2) in the terms of a scalar equation. Note that $\beta \chi_{\Omega} R(z) \chi_{\Omega}$ determines the Birman–Schwinger operator for $H_{0,\beta}$. For z living in a certain neighbourhood of the discrete eigenvalue $\mathcal{E}_{\beta;n}$ but $z \neq \mathcal{E}_{\beta;n}$ the operator $I - \beta \chi_{\Omega} R(z) \chi_{\Omega}$ is invertible and its inverse has the representation (2.25). Therefore one gets

$$\mathbf{I} - \beta \chi_{\Omega} R(z) \chi_{\Omega} - G_{\alpha,\beta}(z) = (\mathbf{I} - \beta \chi_{\Omega} R(z) \chi_{\Omega}) \Big[\mathbf{I} - (\mathbf{I} - \beta \chi_{\Omega} R(z) \chi_{\Omega})^{-1} G_{\alpha,\beta}(z) \Big].$$
(3.4)

To proceed further we estimate the norms involved in the above formula.

Lemma 3.1 The operator $\chi_{\Omega}R(z)\chi_{\Sigma}: L^2(\Sigma) \to L^2(\Omega)$ has the Hilbert–Schmidt norm

$$\|\chi_{\Omega} R(z) \chi_{\Sigma}\|_{\mathrm{HS}}^{2} \le C \frac{|\Omega|}{|z|^{1/2} \mathcal{T}_{z}} \frac{\mathrm{e}^{-\sqrt{2}\mathcal{T}_{z}\rho}}{\rho} , \qquad (3.5)$$

where $\mathcal{T}_z := \Im \sqrt{z} > 0$, ρ is defined by (2.2) and C is a positive constant.

Proof. To prove the claim we will employ the representation (2.9) which implies

$$\|\chi_{\Omega}R(z)\chi_{\Sigma}\|_{\mathrm{HS}}^{2} = \frac{1}{(2\pi)^{2}} \int_{\Omega} \int_{\Sigma} \left| K_{0}(k_{z}|x-y|) \right|^{2} \mathrm{d}x \,\mathrm{d}y \,, \tag{3.6}$$

where $k_z = -i\sqrt{z}$; let us remind that $x = (x_1, x_2)$ and the same notation is adopted for y. Developing the above norm, we obtain

$$\begin{aligned} \|\chi_{\Omega}R(z)\chi_{\Sigma}\|_{\mathrm{HS}}^{2} &= \frac{1}{(2\pi)^{2}} \int_{\Omega} \int_{\Sigma} \left| K_{0}(k_{z}((x_{1}-y_{1})^{2}+(x_{2}-y_{2})^{2})^{1/2} \right|^{2} \mathrm{d}x \,\mathrm{d}y \\ &= \frac{d}{(2\pi)^{2}} \int_{\Omega} \int_{\mathbb{R}} \left| K_{0}(k_{z}((x_{1}-y_{1})^{2}+y_{2}^{2})^{1/2} \right|^{2} \mathrm{d}x_{1} \,\mathrm{d}y \,. \end{aligned}$$
(3.7)

Using the asymptotics of the Macdonald function for the large argument, [1, p. 378, 9.7.2], we obtain

$$|K_0(k_z\,\xi)|^2 \le C' \frac{1}{\xi |z|^{1/2}} \mathrm{e}^{-2\mathcal{T}_z\xi}\,,\tag{3.8}$$

for $\xi > \xi_0$, where C' and ξ_0 are positive constants and C' depends on ξ_0 . Combining the above inequality and (3.7) and employing (2.2) together with the monotonicity of the function $\mathbb{R}_+ \ni x \mapsto x^{-1} e^{-cx}$, c > 0 we can continue with estimates for the Hilbert–Schmidt norm:

$$\begin{aligned} \|\chi_{\Omega} R(z)\chi_{\Sigma}\|_{\mathrm{HS}}^{2} &\leq C' \frac{d}{(2\pi)^{2}} \cdot \frac{1}{|z|^{1/2}} \int_{\Omega} \int_{\mathbb{R}} \frac{\mathrm{e}^{-2\mathcal{T}_{z}(x_{1}^{2}+\rho^{2})^{1/2}}}{(x_{1}^{2}+\rho^{2})^{1/2}} \,\mathrm{d}x_{1} \,\mathrm{d}y \\ &= C' \frac{d|\Omega|}{(2\pi)^{2}} \cdot \frac{1}{|z|^{1/2}} \int_{\mathbb{R}} \frac{\mathrm{e}^{-2\mathcal{T}_{z}(x_{1}^{2}+\rho^{2})^{1/2}}}{(x_{1}^{2}+\rho^{2})^{1/2}} \,\mathrm{d}x_{1} \,. \end{aligned}$$
(3.9)

Applying again the monotonicity of the integrand function and the inequality

$$\sqrt{2}(a^2+b^2)^{1/2} \ge |a|+|b|, \qquad a, b \in \mathbb{R},$$

we arrive at

$$\begin{aligned} \|\chi_{\Omega} R(z)\chi_{\Sigma}\|_{\mathrm{HS}}^{2} &\leq C' \frac{\sqrt{2}d \,|\Omega|}{(2\pi)^{2}} \cdot \frac{1}{|z|^{1/2}} \int_{\mathbb{R}} \frac{\mathrm{e}^{-\sqrt{2}\mathcal{T}_{z}(|x_{1}|+\rho)}}{|x_{1}|+\rho} \,\mathrm{d}x_{1} \\ &\leq C \frac{|\Omega|}{|z|^{1/2}\mathcal{T}_{z}} \frac{\mathrm{e}^{-\sqrt{2}\mathcal{T}_{z}\rho}}{\rho} \,, \end{aligned}$$
(3.10)

where $C = C' \frac{d}{2\pi^2}$. This shows (3.5).

In fact the above reasoning shows that the operator $\chi_{\Sigma}R(z)\chi_{\Omega} : L^2(\Omega) \to L^2(\Sigma)$ admits the same Hilbert–Schmidt norm bound given by (3.1). Proceeding in the same way as in (3.9) one can show that $\chi_{\Sigma}R(z)\chi_{\Sigma}$ is Hilbert–Schmidt. Using the Fredholm theorem, cf. [34], we can conclude that $(I - \alpha\chi_{\Sigma}R(z)\chi_{\Sigma})^{-1}$ exists and it is bounded for $z \in [E_{\alpha}, \infty)$ apart from a set of isolated points. Furthermore, applying (3.3) and the boundedness of $(I - \alpha\chi_{\Sigma}R(z)\chi_{\Sigma})^{-1}$ we obtain the following bound for operator $G_{\alpha,\beta}(z) : L^2(\Omega) \to L^2(\Omega)$,

$$\|G_{\alpha,\beta}(z)\|_{\mathrm{HS}} \le C_1 \frac{\mathrm{e}^{-\sqrt{2}\mathcal{T}_z\rho}}{\rho}, \qquad (3.11)$$

where C_1 is a positive constant which depends on z.

Remark 3.2 Analytic continuation of the resolvent $R_{\alpha,\beta}(z)$. In Section 3.1 we constructed analytic continuation of $\chi_{\Omega}R_{\alpha,0}(z)\chi_{\Omega}$ via the segments $(E_{\alpha;k}, E_{\alpha;k+1})$. This means that $I - \beta\chi_{\Omega}R_{\alpha,0}(z)\chi_{\Omega}$ is analytic in C_k . Moreover, applying the estimates (3.5) and (3.11), we conclude that $\chi_{\Omega}R_{\alpha,0}(z)\chi_{\Omega}$ is a Hilbert–Schmidt operator. Using again the Fredholm theorem, cf. [34], we state that $I - \beta\chi_{\Omega}R_{\alpha,0}(z)\chi_{\Omega}$ is invertible in C_k apart from isolated points. For any $g, f \in C_0(\mathbb{R}^2)$ we can also construct the continuation of $gR_{\alpha,0}(z)f$ to C_k applying the reasoning from Section 3.1. This leads to continuation of

$$fR_{\alpha,\beta}(z)g = fR_{\alpha,0}(z)g + \beta fR_{\alpha,0}(z)\chi_{\Omega}(\mathbf{I} - \beta\chi_{\Omega}R_{\alpha,0}(z)\chi_{\Omega})^{-1}\chi_{\Omega}R_{\alpha,0}(z)g, \qquad (3.12)$$

via the segment $(E_{\alpha;k}, E_{\alpha;k+1})$ to \mathcal{C}_k apart from isolated points.

Let us back to the analysis of (3.4). Combining the decomposition (2.25) and the fact that the operator valued function $z \mapsto A_n(z)$ is analytic a certain neighbourhood S_n of $\mathcal{E}_{\beta;n}$ and taking into account the norm estimate (3.11) we state that $I - A_n(z)G_{\alpha,\beta}(z)$ is invertible for ρ large enough. Consequently, the operator contributing to (3.4) can be written

$$\mathbf{I} - (\mathbf{I} - \beta \chi_{\Omega} R(z) \chi_{\Omega})^{-1} G_{\alpha,\beta}(z) = \left[\mathbf{I} - A_n(z) G_{\alpha,\beta}(z) \right] \left[\mathbf{I} + \frac{1}{\beta} \frac{1}{z - \mathcal{E}_{\beta;n}} \left[\mathbf{I} - A_n(z) G_{\alpha,\beta}(z) \right]^{-1} \mathbf{P}_{\beta;n} G_{\alpha,\beta}(z) \right].$$
(3.13)

Applying (3.13) and (3.4) we come to the conclusion that the condition (3.2) is equivalent to

$$\ker\left[z - \mathcal{E}_{\beta;n} + \frac{1}{\beta} \left[\mathbf{I} - A_n(z) G_{\alpha,\beta}(z) \right]^{-1} \mathbf{P}_{\beta;n} G_{\alpha,\beta}(z) \right] \neq \emptyset.$$
(3.14)

Note that the operator $L_n(z) := [I - A_n(z)G_{\alpha,\beta}(z)]^{-1} P_{\beta;n}G_{\alpha,\beta}(z)$ acting in $L^2(\Omega)$ is rank one and it acts as

$$L_{n}(z)f = (w_{\beta;n}, G_{\alpha,\beta}(z)f)_{L^{2}(\Omega)} \left[I - A_{n}(z)G_{\alpha,\beta}(z) \right]^{-1} w_{\beta;n}.$$
(3.15)

Therefore, the condition (3.14) can be transited to the scalar equation

$$z - \mathcal{E}_{\beta;n} + \frac{1}{\beta} (w_{\beta;n}, G_{\alpha,\beta}(z) [I - A_n(z)G_{\alpha,\beta}(z)]^{-1} w_{\beta;n})_{L^2(\Omega)} = 0.$$
(3.16)

With the above results, we are ready to prove the following statement, which establishes the resonances.

Theorem 3.3 Assume that $H_{0,\beta}$ has a simple, discrete eigenvalue $\mathcal{E}_{\beta;n} \in (E_{\alpha;j}, E_{\alpha;j+1})$. Then the second sheet analytic continuation of $R_{\alpha,\beta}(z)$ defined by (3.12) has a unique pole $z_n \in S_n$, i.e. ker[I – $\beta \chi_\Omega R_{\alpha,0}(z_n)\chi_\Omega \neq \emptyset$, with the asymptotics

$$z_n(\rho) = \mathcal{E}_{\beta;n} + \mathcal{O}\left(\frac{\mathrm{e}^{-\sqrt{2|\mathcal{E}_{\beta;n}|}\rho}}{\rho}\right).$$
(3.17)

Proof. Step 1. Auxiliary statement: the Neumann expansion. We adopt the notation $b := \frac{1}{\rho}$, which means that $b \to 0$ for $\rho \to \infty$. Note that the operator $G_{\alpha,\beta}(z)$ depends on b because it is defined by the embeddings acting between spaces $L^2(\Omega)$ and $L^2(\Sigma)$, see (3.3). Since the parameter b will play the essential rule in the further discussion we implement the notation

$$\eta(z,b) := (w_{\beta;n}, G_{\alpha,\beta}(z) [I - A_n(z)G_{\alpha,\beta}(z)]^{-1} w_{\beta;n})_{L^2(\Omega)}.$$
(3.18)

In view of (3.11) we have $\lim_{b\to 0} \eta(z, b) = 0$ for any $z \in S_n$. We continuously extend function η assuming that $\eta(z, 0) = 0$. Note that formula (3.18) can be rephrased in the terms of the Neumann series

$$[I - A_n(z)G_{\alpha,\beta}(z)]^{-1} = I + A_n(z)G_{\alpha,\beta}(z) + [A_n(z)G_{\alpha,\beta}(z)]^2 + \dots$$
(3.19)

which implies

$$\eta(z,b) = \left[(w_{\beta;n}, G_{\alpha,\beta}(z)w_{\beta;n})_{L^{2}(\Omega)} + (w_{\beta;n}, G_{\alpha,\beta}(z)A_{n}(z)G_{\alpha,\beta}(z)w_{\beta;n})_{L^{2}(\Omega)} + (w_{\beta;n}, G_{\alpha,\beta}(z)[A_{n}(z)G_{\alpha,\beta}(z)]^{2}w_{\beta;n})_{L^{2}(\Omega)} + \ldots \right] = 0.$$
(3.20)

Step 2. Unique solution of the spectral equation. To discuss further properties of η we will investigate analycity of $G_{\alpha,\beta}(z)$, which is related to the resolvent $R_{\alpha,0}(z)$ by the equivalence $G_{\alpha,\beta}(z) = \beta [\chi_{\Omega} R_{\alpha,0}(z)\chi_{\Omega} - \chi_{\Omega} R(z)\chi_{\Omega}]$, cf. (3.3) and (2.31). As a consequence of the first resolvent identity

$$\chi_{\Omega}[R(z) - R(z')]\chi_{\Omega} = (z' - z)\chi_{\Omega}R(z)R(z')\chi_{\Omega}, \qquad (3.21)$$

and the analogous identity for $R_{\alpha,0}(z)$, we get

$$G_{\alpha,\beta}(z) - G_{\alpha,\beta}(z') = \beta \chi_{\Omega} \Big[R_{\alpha,0}(z) - R_{\alpha,0}(z') - R(z) + R(z') \Big] \chi_{\Omega}$$

$$= (z'-z)\beta \Big[\chi_{\Omega} R_{\alpha,0}(z) R_{\alpha,0}(z') \chi_{\Omega} - \chi_{\Omega} R(z) R(z') \chi_{\Omega} \Big]$$

$$= (z'-z) \Big[\chi_{\Omega} R(z) \chi_{\Omega} G_{\alpha,\beta}(z') + G_{\alpha,\beta}(z) \chi_{\Omega} R(z') \chi_{\Omega} + \frac{1}{\beta} G_{\alpha,\beta}(z) G_{\alpha,\beta}(z') \Big].$$
(3.22)

Note that the operators in the last line of (3.22) can be bounded by means of (3.11). In view of the analyticity of $A_n(z)$, the same reasoning can be proceeded for operators appearing in further terms of (3.20). Therefore we can state that $\eta(z, b)$ is analytic as the function of $z \in S_n$, and

$$\eta(z,b) - \eta(z',b) = (z - z')f(z,z';b), \qquad (3.23)$$

where $f(\cdot, \cdot; b)$ is analytic and $\lim_{b\to 0} f(z, z'; b) = 0$ uniformly with respect to z and z'. This gives $\partial_z \eta(z; b)|_{b=0} = 0$, where we understand the symbol $|_{b=0}$ as the limiting value for $b \to 0$. The latter follows directly from the estimate (3.11) and eq. (3.22). This means that

$$\frac{\partial \tilde{\eta}(z,b)}{\partial z}\Big|_{b=0} = 1, \quad \text{where} \quad \tilde{\eta}(z,b) := z - \mathcal{E}_{\beta;n} + \beta^{-1} \eta(z,b).$$
(3.24)

Note that the spectral equation (3.16) reads

$$\tilde{\eta}(z,b) = 0. \tag{3.25}$$

Combining (3.24) with the implicit function theorem and the fact that $\tilde{\eta}(\mathcal{E}_{\beta;n}, 0) = 0$, we can conclude that for *b* small enough the equation (3.25) has a unique solution $z_n(b)$ such that $\tilde{\eta}(z_n(b), b) = 0$. The asymptotics of the solution can be recovered from (3.20). Since we are interested in a solutions for $b \neq 0$ we again back to the variable ρ and the asymptotics $\rho \to \infty$. Without a danger of confusion, we keep the same notation for $z_n(\cdot)$ expressed in the terms of ρ . Step 3. Asymptotics of the solution. Combining the spectral equation (3.25) and the expansion (3.20) together with the definition of $\tilde{\eta}(\cdot, \cdot)$ from (3.24), we get

$$z_{n}(\rho) = \mathcal{E}_{\beta;n} - \frac{1}{\beta} \Big[(w_{\beta;n}, G_{\alpha,\beta}(z)w_{\beta;n})_{L^{2}(\Omega)} + (w_{\beta;n}, G_{\alpha,\beta}(z)A_{n}(z)G_{\alpha,\beta}(z)w_{\beta;n})_{L^{2}(\Omega)} + (w_{\beta;n}, G_{\alpha,\beta}(z)[A_{n}(z)G_{\alpha,\beta}(z)]^{2} w_{\beta;n})_{L^{2}(\Omega)} + \dots \Big].$$
(3.26)

As follows from the above discussion, the dominated term of $z_n(\rho)$ comes from the term $-\beta^{-1}(w_{\beta;n}, G_{\alpha,\beta}(z)w_{\beta;n})_{L^2(\Omega)}, z \in S_n$. Therefore we expand

$$(w_{\beta;n}, G_{\alpha,\beta}(z)w_{\beta;n})_{L^{2}(\Omega)} = (w_{\beta;n}, G_{\alpha,\beta}(\mathcal{E}_{\beta;n})w_{\beta;n})_{L^{2}(\Omega)} + (z - \mathcal{E}_{\beta;n})(w_{\beta;n}T_{n}(z)w_{\beta;n})_{L^{2}(\Omega)},$$
(3.27)

where $T_n(z)$ has the norm that can be bounded by the norm of $G_{\alpha,\beta}(z)$, see (3.22). Applying the expansions (3.26) and (3.27) we get

$$z_{n}(\rho) = \mathcal{E}_{\beta;n} - \frac{1}{\beta} \Big[1 + \beta^{-1} (w_{\beta;n}, T_{n}(z)w_{\beta;n})_{L^{2}(\Omega)} \Big]^{-1} \Big[(w_{\beta;n}, G_{\alpha,\beta}(\mathcal{E}_{\beta;n})w_{\beta;n})_{L^{2}(\Omega)} \\ + (w_{\beta;n}, G_{\alpha,\beta}(z)A_{n}(z)G_{\alpha,\beta}(z)w_{\beta;n})_{L^{2}(\Omega)} + (w_{\beta;n}, G_{\alpha,\beta}(z)[A_{n}(z)G_{\alpha,\beta}(z)]^{2} w_{\beta;n})_{L^{2}(\Omega)} + \dots \Big].$$

$$(3.28)$$

The above expansion shows that the dominated term is determined by

$$-\frac{1}{\beta}(w_{\beta;n}, G_{\alpha,\beta}(\mathcal{E}_{\beta;n})w_{\beta;n})_{L^2(\Omega)}, \qquad (3.29)$$

which, in view of (3.11), yields the asymptotics

$$z_n(\rho) = \mathcal{E}_{\beta;n} + \mathcal{O}\left(\frac{\mathrm{e}^{-\sqrt{2|\mathcal{E}_{\beta;n}|}\rho}}{\rho}\right).$$
(3.30)

This completes the proof.

4 Fermi's golden rule

The discussion of this section is devoted to a deeper analysis of the asymptotic of $\Im z_n(\rho)$. First we analyze the component of $\Im z_n(\rho)$ given by

$$\Gamma_n(\rho) := -\frac{1}{\beta} \Im(w_{\beta;n}, G_{\alpha,\beta}(\mathcal{E}_{\beta;n}) w_{\beta;n})_{L^2(\Omega)}, \qquad (4.1)$$

which comes from (3.29); cf. (3.28). Using the Birman-Schwinger principle (2.24) we get

$$\chi_{\Omega} R(\mathcal{E}_{\beta;n}) \chi_{\Omega} w_{\beta;n} = \chi_{\Omega} \omega_{\beta;n} , \quad \chi_{\Omega} R(\mathcal{E}_{\beta;n}) \chi_{\Omega} w_{\beta;n} = \beta^{-1} w_{\beta;n} .$$
(4.2)

This yields

$$(w_{\beta;n}, G_{\alpha,\beta}(\mathcal{E}_{\beta;n})w_{\beta;n})_{L^2(\Omega)} = \beta^2(\chi_\Omega \omega_{\beta;n}, G_{\alpha,\beta}(\mathcal{E}_{\beta;n})\chi_\Omega \omega_{\beta;n})_{L^2(\Omega)}.$$
(4.3)

On the other hand, $\mathcal{E}_{\beta;n} < 0$ belongs to the resolvent sets of $R^c_{\alpha,0}(\cdot)$ and $R(\cdot)$, which are self adjoint operators $(R^c_{\alpha,0}(\cdot)$ is defined by (2.18)). Therefore, we have

$$\Im(\chi_{\Omega}\omega_{\beta;n}, R(\mathcal{E}_{\beta;n})\chi_{\Omega}\omega_{\beta;n})_{L^{2}(\Omega)} = \Im(\chi_{\Omega}\omega_{\beta;n}, R^{c}_{\alpha,0}(\mathcal{E}_{\beta;n})\chi_{\Omega}\omega_{\beta;n})_{L^{2}(\Omega)} = 0.$$

$$(4.4)$$

Using again the equation

$$G_{\alpha,\beta}(z) = \beta [\chi_{\Omega} R_{\alpha,0}(z) \chi_{\Omega} - \chi_{\Omega} R(z) \chi_{\Omega}], \qquad (4.5)$$

together with (4.4), we get

$$\Im(\chi_{\Omega}\omega_{\beta;n}, G_{\alpha,\beta}(\mathcal{E}_{\beta;n})\chi_{\Omega}\omega_{\beta;n})_{L^{2}(\Omega)} = \beta\Im(\chi_{\Omega}\omega_{\beta;n}, R^{d}_{\alpha,0}(\mathcal{E}_{\beta;n})\chi_{\Omega}\omega_{\beta;n})_{L^{2}(\Omega)}.$$
(4.6)

Furthermore, applying (2.19) we have

$$\Im(\chi_{\Omega}\omega_{\beta;n}, R^{d}_{\alpha,0}(\mathcal{E}_{\beta;n})\chi_{\Omega}\omega_{\beta;n})_{L^{2}(\Omega)} = \Im\frac{1}{2\pi} \Big[\lim_{\varepsilon \to 0} \sum_{k \in \mathcal{N}_{\alpha}} \int_{\Omega \times \Omega} \int_{\mathbb{R}} \frac{\mathrm{e}^{ip_{1}(x_{1}-y_{1})}}{p_{1}^{2} + E_{\alpha;k} - (\mathcal{E}_{\beta;n} + i\varepsilon)} \phi_{\alpha;k}(x_{2}) \phi_{\alpha;k}(y_{2})^{*} \omega_{\beta;n}(x)^{*} \omega_{\beta;n}(y) \mathrm{d}p_{1} \,\mathrm{d}x \,\mathrm{d}y\Big].$$

$$(4.7)$$

To deal with the integral

$$\int_{\mathbb{R}} \frac{\mathrm{e}^{ip_1(x_1-y_1)}}{p_1^2 - (b_{k,n} + i\varepsilon)} \mathrm{d}p_1 \,, \quad \text{where} \quad b_{k,n} := -(E_{\alpha;k} - \mathcal{E}_{\beta;n}) \,, \tag{4.8}$$

we distinguish two cases $b_{k,n} > 0$ and $b_{k,n} < 0$. Assume first that $b_{k,n} > 0$; then the above integrand has two poles $\pm(\tilde{b}_{\epsilon} + i\tilde{\varepsilon})$, where $\tilde{b}_{\epsilon} \to \sqrt{b_{k,n}}$ and $\tilde{\varepsilon} \to \text{for } \varepsilon \to 0$. Applying the Sokhotski–Plemelj formula which can be formally written $\lim_{\varepsilon \to 0} \frac{1}{x\pm i\varepsilon} = \mathcal{P}\frac{1}{x} \mp i\pi\delta(x)$, where \mathcal{P} stands for the principle value, we get

$$\lim_{\varepsilon \to 0} \int_{\mathbb{R}} \frac{e^{ip_1(x_1 - y_1)}}{p_1^2 - (b_{k,n} + i\varepsilon)} dp_1 = \lim_{\varepsilon \to 0} \int_{\mathbb{R}} \frac{e^{ip_1(x_1 - y_1)}}{(p_1 - (\tilde{b}_{\varepsilon} + i\tilde{\varepsilon}))(p_1 + (\tilde{b}_{\varepsilon} + i\tilde{\varepsilon}))} dp_1
= \frac{i\pi}{2\sqrt{b_{k,n}}} \Big[e^{i\sqrt{b_{k,n}}(x_1 - y_1)} + e^{-i\sqrt{b_{k,n}}(x_1 - y_1)} \Big] + \mathcal{P} \int_{\mathbb{R}} \frac{e^{ip_1(x_1 - y_1)}}{p_1^2 - b_{k,n}} dp_1 \qquad (4.9)
= \frac{i\pi}{\sqrt{b_{k,n}}} \cos(\sqrt{b_{k,n}}|x_1 - y_1|) + \mathcal{P} \int_{\mathbb{R}} \frac{e^{ip_1(x_1 - y_1)}}{p_1^2 - b_{k,n}} dp_1 .$$

For $b_{k,n} < 0$ the analogous limit reproduces $\pi \frac{\exp(-|b_{k,n}|^{1/2}|x_1-x_2|)}{|b_{k,n}|^{1/2}}$, which contributes to the real component of (4.6). This, in view of (4.7) and (4.9) implies

$$\Im(\chi_{\Omega}\omega_{\beta;n}, R^{d}_{\alpha,0}(\mathcal{E}_{\beta;n})\chi_{\Omega}\omega_{\beta;n})_{L^{2}(\Omega)} = \sum_{k: E_{\alpha;k} < \mathcal{E}_{\beta;n}} \frac{1}{2\sqrt{b_{k,n}}} \int_{\Omega \times \Omega} \cos(\sqrt{b_{k,n}}|x_{1} - y_{1}|)\phi_{\alpha;k}(x_{2})\phi_{\alpha;k}(y_{2})^{*}\omega_{\beta;n}(x)^{*}\omega_{\beta;n}(y) \,\mathrm{d}x \,\mathrm{d}y \\ = \sum_{k=1}^{j} \frac{1}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \int_{\Omega \times \Omega} \cos(\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}|x_{1} - y_{1}|)\phi_{\alpha;k}(x_{2})\phi_{\alpha;k}(y_{2})^{*}\omega_{\beta;n}(x)^{*}\omega_{\beta;n}(y) \,\mathrm{d}x \,\mathrm{d}y \,,$$

$$(4.10)$$

where, in the last step, we have changed the summation from $\sum_{k: E_{\alpha;k} < \mathcal{E}_{\beta;n}}$ to $\sum_{k=1}^{j}$ since $\mathcal{E}_{\beta;n} \in (E_{\alpha;j}, E_{\alpha;j+1})$. Using now (4.1), (4.3), (4.6) and (4.10) we can conclude

$$\Gamma_{n}(\rho) = -\beta^{2} \sum_{k=1}^{\mathcal{I}} \frac{1}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \Big[\int_{\Omega \times \Omega} \cos(\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}} |x_{1} - y_{1}|) \phi_{\alpha;k}(x_{2}) \phi_{\alpha;k}(y_{2})^{*} \\ \times \omega_{\beta;n}(x)^{*} \omega_{\beta;n}(y) \,\mathrm{d}x \,\mathrm{d}y \Big].$$

$$(4.11)$$

In fact, $\Gamma_n(\rho)$ can be expressed by means of the generalized eigenfunctions $\varphi_k(x, p_1) = \frac{1}{\sqrt{2\pi}} e^{ip_1 x_1} \phi_{\alpha;k}(x_2)$ of $H_{\alpha,0}$. Using the first term of the second line of (4.9) and $\varphi_k(x, p_1)$ we can write

$$\Gamma_{n}(\rho) = -\beta^{2} \sum_{k=1}^{j} \frac{\pi}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \left[\left| \int_{\Omega} \omega_{\beta;n}(x)^{*} \varphi_{k}(x, p_{1}) \mathrm{d}x \right|_{p_{1} = (\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}}^{2} + \left| \int_{\Omega} \omega_{\beta;n}(x)^{*} \varphi_{k}(x, p_{1}) \mathrm{d}x \right|_{p_{1} = -(\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}}^{2} \right],$$
(4.12)

where the lower indexes indicate that the values of functions are taken at specified points. The above Formula reveals the well known Fermi's golden rule, cf. [33, XII. 6]. In Appendix 5.1, we show that the further terms contributing to (3.28) possess the stronger asymptotics then (4.12).

Remark 4.1 Note that, $\varphi_k(\cdot, p_1) \notin L^2(\mathbb{R}^2)$, however, the function $\chi_\Omega \varphi_k(\cdot, p_1)$ is well defined as an element of $L^2(\mathbb{R}^2)$. The expressions (4.12) can be reformulated in the form more familiar to physicists

$$\Gamma_{n}(\rho) = -\beta^{2} \sum_{j=1}^{J} \frac{\pi}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \Big[|(\omega_{\beta;n}, \chi_{\Omega}\varphi_{k}(\cdot, p_{1}))|^{2}_{p_{1} = (\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}} + |(\omega_{\beta;n}, \chi_{\Omega}\varphi_{k}(\cdot, p_{1}))|^{2}_{p_{1} = -(\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}} \Big],$$
(4.13)

where the scalar product in the above formulae is defined in the space $L^2(\mathbb{R}^2)$. The resonance width is determined by

$$\gamma_n(\rho) := 2 \left| \Im z_n(\rho) \right|, \tag{4.14}$$

and, therefore the dominated term of $\gamma_n(\cdot)$ takes the form

$$\gamma_{n}(\rho) \approx 2\pi \sum_{k=1}^{j} \frac{1}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \Big[|(\omega_{\beta;n}, V_{\beta}\varphi_{k}(\cdot, p_{1}))|^{2}_{p_{1} = p_{n,k}} + |(\omega_{\beta;n}, V_{\beta}\varphi_{k}(\cdot, p_{1}))|^{2}_{p_{1} = -p_{n,k}} \Big], \qquad (4.15)$$

where $V_{\beta} = -\beta \chi_{\Omega}$ and $p_{n,k} = (\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}$. The above formula represents also the decay rate of the resonant state in the first approximation. It reflexes the characteristic properties, namely the contribution of the transition amplitude $(\omega_{\beta;n}, V_{\beta}\varphi_k(\cdot, p_1))$ and spectral density function $(2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}})^{-1}$. The summation $\sum_{k=1}^{j}$ comes from the fact that we have to take into account all modes which 'catch' the resonant energy $\mathcal{E}_{\beta;n}$.

4.1 Asymptotics of $\Gamma_n(\rho)$

Let us analyze in more detail the asymptotic behaviour of $\Gamma_n(\rho)$ for $\rho \to \infty$. For this aim we use the expression (4.10). Using the fact that for $x_2 \in \mathbb{R} \setminus [0, d]$, we have

$$\phi_{\alpha;k}(x_2) = M_{\alpha,k} e^{-\sqrt{|E_{\alpha;k}|}|x_2|}, \qquad (4.16)$$

where M_{α} is the normalization constant, we can estimate (4.10) by

$$\sum_{k=1}^{j} \frac{|M_{\alpha,k}|^2 \mathrm{e}^{-2\sqrt{|E_{\alpha;k}|}\rho}}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \int_{\Omega \times \Omega} \left| \cos(\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}} |x_1 - y_1|) \,\omega_{\beta;n}(x)^* \omega_{\beta;n}(y) \right| \,\mathrm{d}x \,\mathrm{d}y$$

$$\leq j C_{j,n} \mathrm{e}^{-2\sqrt{|E_{\alpha;j}|}\rho} \,, \tag{4.17}$$

where

$$C_{j,n} := \max_{1 \le k \le j} \left[\frac{|M_{\alpha,k}|^2}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \int_{\Omega \times \Omega} \left| \cos(\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}} |x_1 - y_1|) \,\omega_{\beta;n}(x)^* \omega_{\beta;n}(y) \right| \mathrm{d}x \,\mathrm{d}y \right]; \quad (4.18)$$

in the above reasoning we used the fact that $\sqrt{|\mathcal{E}_{\beta;n}|} < \sqrt{|E_{\alpha;k}|}$ for any k = 1, ..., j. In fact it follows from (4.12) that all the components labeled by 1, ..., j contributing to $\Gamma_n(\rho)$ have the negative sign (they can not vanish because the functions in the scalar products are not orthogonal). Using (4.16) we state

$$\Gamma_n(\rho) \approx e^{-2\sqrt{|E_{\alpha;j}|}\rho}, \quad \text{where} \quad \mathcal{E}_{\beta;n} \in (E_{\alpha;j}, E_{\alpha;j+1}).$$
(4.19)

On the other hand, note that for ρ sufficiently large we have

$$e^{-2\sqrt{|E_{\alpha;j}|}\rho} \le \frac{e^{-\sqrt{2}|\mathcal{E}_{\beta;n}|\rho}}{\rho}; \qquad (4.20)$$

therefore the asymptotics (4.19) enhances (3.17) for the imaginary component of $z_n(\rho)$.

5 Generalization of the system: varying transverse profile and perturbation defined by the Kato class measure

In this section we consider a generalization of the model and assume that the transverse profile of the waveguide is given by a piecewise continuous function $\alpha : [0, d] \to \mathbb{R}$. On the other hand, we generalize the well potential supported on Ω by a Radon measure μ belonging to the Kato class, i.e. satisfying

$$\lim_{\varepsilon \to 0} \sup_{x \in \mathbb{R}^2} \int_{B(x,\varepsilon)} |\log(|x-y|)| \,\mathrm{d}\mu(y) = 0\,, \tag{5.1}$$

where $B(x,\varepsilon) \subset \mathbb{R}^2$ denotes a ball centered at x and with the radius ε . Throughout this section we assume that the support $\Omega \subset \mathbb{R}^2$ of μ is compact; in this part of further discussion where support of μ plays an essential role we will use the notation $\mu_{\Omega} \equiv \mu$. The examples of the Kato measures has been discussed in Introduction.

To maintain the consistency with previous discussion we define the remote perturbation by $-\beta\mu$, where $\beta \in \mathbb{R}$. Then the Hamiltonian is associated to the sesquilinear form

$$\mathscr{E}_{\alpha,\beta\mu}[f] = \int_{\mathbb{R}^2} |\nabla f(x)|^2 \,\mathrm{d}x - \int_{\Sigma} \alpha(x_2) |f(x)|^2 \,\mathrm{d}x - \beta \int_{\mathbb{R}^2} |I_{\mu}f(x)|^2 \,\mathrm{d}\mu(x) \,, \quad f \in W^{1,2}(\mathbb{R}^2) \,, \tag{5.2}$$

where I_{μ} denotes the space embedding $I_{\mu} : W^{1,2}(\mathbb{R}^2) \to L^2_{\mu} \equiv L^2(\mathbb{R}^2, d\mu)$ which is continuous, cf. [2]. For any Kato class measure μ and any small parameter $\varepsilon > 0$ there exists $C(\varepsilon)$ such that

$$\int_{\mathbb{R}^2} |I_{\mu}f(x)|^2 \mathrm{d}\mu(x) \le \varepsilon \int_{\mathbb{R}^2} |\nabla f(x)|^2 \mathrm{d}x + C(\varepsilon) \int_{\mathbb{R}^2} |f(x)|^2 \mathrm{d}x \,, \tag{5.3}$$

cf. [7]. This yields

$$\int_{\mathbb{R}^2} |I_{\mu}f(x)|^2 \mathrm{d}\mu(x) \le \varepsilon \Big[\int_{\mathbb{R}^2} |\nabla f(x)|^2 \mathrm{d}x - \int_{\Sigma} \alpha(x_2) |f(x)|^2 \mathrm{d}x \Big] + C_1(\varepsilon) \int_{\mathbb{R}^2} |f(x)|^2 \mathrm{d}x \,, \tag{5.4}$$

where $C_1(\varepsilon) = C(\varepsilon) + \varepsilon \sup |\alpha(x_2)|$, which implies that the operator associated to $\mathscr{C}_{\alpha,\beta\mu}$ has a unique self adjoint extension in view of the KLMN Theorem, cf. [34]. The resulting operator defines a generalization of $H_{\alpha,\beta}$ given by (2.3).

To reconstruct the resolvent of $H_{0,\beta\mu}$ we define the embeddings of R(z) to L^2_{μ} by

$$R(z)|_{0\mu}f = R(z) * f\mu = \frac{1}{2\pi} \int_{\mathbb{R}^2} K_0(k_z|x-y|)f(y) \,\mathrm{d}\mu(y) \,:\, L^2_\mu \to L^2(\mathbb{R}^2)\,,\tag{5.5}$$

where $k_z = -i\sqrt{z}$. Furthermore, we define $R(z)|_{\mu 0} : L^2(\mathbb{R}^2) \to L^2_{\mu}$ which acts as $R(z)|_{\mu 0} f = I_{\mu}R(z)f$ and, finally, we construct the bilateral embedding by

$$R(z)|_{\mu\mu} := I_{\mu}R(z)|_{0\mu} : L^2_{\mu} \to L^2_{\mu}.$$
(5.6)

We are ready to construct the resolvent of $(H_{0,\beta} - z)^{-1}$ by

$$R_{0,\beta\mu}(z) = R(z) + \beta R(z)|_{0\mu} (\mathbf{I} - \beta R(z)|_{\mu\mu})^{-1} R(z)|_{\mu0}.$$
(5.7)

The similar types of Hamiltonian were discussed, for example in [7, 28].

On the other hand, the resolvent $R_{\alpha,0}(z)$ represented in the Birman–Schwinger form is given by

$$R_{\alpha,0}(z) = R(z) + R(z)\alpha^{1/2}\chi_{\Sigma}(I - |\alpha|^{1/2}\chi_{\Sigma}R(z)\alpha^{1/2}\chi_{\Sigma})^{-1}|\alpha|^{1/2}\chi_{\Sigma}R(z), \qquad (5.8)$$

where $\alpha(\cdot)$ is now a function of x_2 and $\alpha^{1/2} = \operatorname{sgn}(\alpha) |\alpha|^{1/2}$. Using the embeddings of R(z) to L^2_{μ} and the strip resolvent (5.8) we can build the operator

$$\mathbf{I} - \beta R_{\alpha,0}(z)|_{\mu\mu} = \mathbf{I} - \beta \Big[R(z)|_{\mu\mu} + R(z)|_{\mu0} \alpha^{1/2} \chi_{\Sigma} (\mathbf{I} - |\alpha|^{1/2} \chi_{\Sigma} R(z) \alpha^{1/2} \chi_{\Sigma})^{-1} |\alpha|^{1/2} \chi_{\Sigma} R(z)|_{0\mu} \Big] : L^2_{\mu} \to L^2_{\mu},$$

which contributes to the Birman–Schwinger resolvent of $H_{\alpha,\beta}$. Assuming now that z is such that the above operator is invertible, we construct the resolvent of $H_{\alpha,\beta}$ by

$$R_{\alpha,\beta\mu}(z) = R_{\alpha,0}(z) + \beta R_{\alpha,0}(z)|_{0\mu} (\mathbf{I} - \beta R_{\alpha,0}(z)|_{\mu\mu})^{-1} R_{\alpha,0}(z)|_{\mu0} : L^2(\mathbb{R}^2) \to L^2(\mathbb{R}^2).$$
(5.9)

The stability of the essential spectrum

$$\sigma_{\rm ess}(H_{\alpha,\beta\mu}) = \sigma_{\rm ess}(H_{\alpha,0}) = [E_{\alpha},\infty)$$
(5.10)

is a consequence of the following reasoning. Note that the operators $R(z)|_{\mu 0} : L^2(\mathbb{R}^2) \to L^2_{\mu}$ and $R(z)|_{0\mu} : L^2_{\mu} \to L^2(\mathbb{R}^2)$ are compact in view of [7, Le. 3.1]. Therefore, using (5.8) we can state that both

$$R_{\alpha,0}(z)|_{0\mu} : L^2_{\mu} \to L^2(\mathbb{R}^2) \text{ and } R_{\alpha,0}(z)|_{\mu 0} : L^2(\mathbb{R}^2) \to L^2_{\mu}$$
 (5.11)

are also compact since the operator $(I - |\alpha|^{1/2} \chi_{\Sigma} R(z) \alpha^{1/2} \chi_{\Sigma})^{-1}$ is bounded. Consequently, we get the compactness of the second component of (5.9) and, in view of the Weyl's theorem, we come to the conclusion that the essential spectra of $H_{\alpha,0}$ and $H_{\alpha,\beta\mu}$ coincide.

Assumptions on $H_{\alpha,0}$ and $H_{0,\beta\mu}$. As before, let h_{α} stand for the transverse component of the Hamiltonian $H_{\alpha,0}$. In the current framework it takes the form

$$h_{\alpha} = -\frac{d^2}{dx_2^2} - \alpha(x_2)\chi_{[0,d]}(x_2) : W^{2,2}(\mathbb{R}) \to L^2(\mathbb{R}).$$
(5.12)

In the following we assume that h_{α} has at least one negative discrete spectrum point. It happens, for example, if $\int_{[0,d]} \alpha(x_2) dx_2 \geq 0$, cf. [35]. Following the convention of the previous discussion, we will keep the notation $\{E_{\alpha;1} \leq E_{\alpha;2} \leq \ldots \leq E_{\alpha;N_{\alpha}}\}$ and $\{\phi_{\alpha;n}\}_{n \in \mathcal{N}_{\alpha}}$ for the eigenvalues and eigenfunctions of h_{α} , see. (2.5). On the other hand, the eigenvalues and eigenfunctions of $H_{0,\beta\mu}$ will be denoted by $\{\mathcal{E}_{\beta;1} \leq \mathcal{E}_{\beta;2} \leq \ldots \leq \mathcal{E}_{\beta;N_{\beta}}\}$ and $\{\omega_{\beta;n}\}_{n \in \mathcal{N}_{\beta}}$, cf. (2.21). In the following we assume that there exists $n \in \mathcal{N}_{\beta}$ and $j \in \mathcal{N}_{\alpha}$ such that $\mathcal{E}_{\beta;n} \in (E_{\alpha;j}, E_{\alpha;j+1})$.

To investigate the problem of resonances in the generalized potential system we need to estimate

$$G_{\alpha,\beta}(z) = \beta R_{\alpha,0}(z)|_{\mu\mu} (\mathbf{I} - \beta R_{\alpha,0}(z)|_{\mu\mu})^{-1} R_{\alpha,0}(z)|_{\mu\mu} : L^2_{\mu} \to L^2_{\mu},$$
(5.13)

which states an analogue of (3.3). Similarly as before, we start from the estimates of the embeddings of the free resolvent

$$\|R(z)\|_{\mu 0} \chi_{\Sigma}\|_{\mathrm{HS}}^{2} = \frac{1}{2\pi} \int_{\mathbb{R}^{2}} \int_{\Sigma} |I_{\mu} K_{0}(k_{z}|x-y|)|^{2} \mathrm{d}\mu(x) \mathrm{d}y$$

$$\leq \sup_{x \in \Omega} \frac{|\mu|}{2\pi} \int_{\Sigma} |I_{\mu} K_{0}(k_{z}|x-y|)|^{2} \mathrm{d}y \leq C \frac{|\mu|}{|z|^{1/2} \mathcal{T}_{z}} \frac{\mathrm{e}^{-\sqrt{2}\mathcal{T}_{z}\rho}}{\rho},$$
(5.14)

where $|\mu| = \int_{\mathbb{R}^2} d\mu$ and the embedding I_{μ} is applied to the kernel defined by K_0 understood as the function of the variable x. Consequently, taking into account the fact that $(I - \beta R_{\alpha,0}(z)|_{\mu\mu})^{-1}$ is bounded, the operator $G_{\alpha,\beta}(z) : L^2_{\mu} \to L^2_{\mu}$ can be estimated

$$\|G_{\alpha,\beta}(z)\|_{\rm HS} \le C_1 \frac{{\rm e}^{-\sqrt{2}\mathcal{T}_z\rho}}{\rho},$$
(5.15)

where C_1 is a positive constant depending on $|\mu|$ and z. Formula (5.15) is a generalisation of (3.11).

Repeating the reasoning from Section 3.1 we construct the second sheet analytic continuation of $R_{\alpha,0}(z)$ via the segment $(E_{\alpha;j}, E_{\alpha;j+1})$ in the sense of (3.12). Using this result and relying on the same analysis as in the proof of Theorem 3.3 we can prove the that $R_{\alpha,\beta\mu}(z)$ has the second sheet pole located at

$$z_n(\rho) = \mathcal{E}_{\beta,n} + \mathcal{O}\left(\frac{\mathrm{e}^{-\sqrt{2|\mathcal{E}_{\beta,n}|}\rho}}{\rho}\right).$$
(5.16)

Moreover, the dominated term of its imaginary part satisfies

$$\Gamma_{n}(\rho) = -\beta^{2} \sum_{k=1}^{j} \frac{\pi}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \left[\left| \int_{\Omega} I_{\mu}(\omega_{\beta;n}(x)^{*}\varphi_{k}(x, p_{1})) \mathrm{d}\mu(x) \right|^{2}_{p_{1} = (\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}} + \left| \int_{\Omega} I_{\mu}(\omega_{\beta;n}(x)^{*}\varphi_{k}(x, p_{1})) \mathrm{d}\mu(x) \right|^{2}_{p_{1} = -(\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}} \right].$$
(5.17)

The above result states a generalization of Fermi's golden role (4.12) for the Kato class measures.

5.1 A special case: the trap defined by a delta interaction

In this section we consider a special case of the Kato class measure, defined by the Dirac delta potential supported on a curve. Let Ω be a finite C^1 planar curve (open arc or loop) with the length l. Assume that Ω is parameterized by the arc length $s \in I_l := [0, l]$ and is the graph of the function $s \mapsto \varpi(s) \in \mathbb{R}^2$. Our aim is to analyze the resonances problem for the Hamiltonian which can be symbolically written

$$-\Delta - \alpha \chi_{\Sigma}(x) - \beta \delta(x - \Omega), \quad \alpha < 0, \ \beta \in \mathbb{R}.$$
(5.18)

Remark 5.1 Delta interaction: a particular case of Kato class measure. In the analyzed case, the measure μ corresponding to the delta interaction will be denoted as δ . It is defined as follows: for any Borel set $\mathcal{B} \subset \mathbb{R}^2$, we determine

$$\delta(\mathcal{B}) = \ln(\mathcal{B} \cap \Omega),\tag{5.19}$$

where $lin(\cdot)$ denotes the linear measure. Consequently, δ belongs to the Kato class of measures, cf. [7].

To give the expression (5.18) a mathematical meaning, we introduce the corresponding embedding I_{δ} : $W^{1,2}(\mathbb{R}^2) \to L^2(I_l)$ acting as $I_{\delta}f = f(\varpi(\cdot))$. Then, for this specific case, the sesquilinear form (5.2) can be expressed as

$$\mathscr{E}_{\alpha,\beta\delta}[f] = \int_{\mathbb{R}^2} |\nabla f(x)|^2 \,\mathrm{d}x - \alpha \int_{\Sigma} |f(x)|^2 \,\mathrm{d}x - \beta \int_{I_l} |f(\varpi(s))|^2 \,\mathrm{d}s \,, \quad f \in W^{1,2}(\mathbb{R}^2) \,. \tag{5.20}$$

The special case of (5.4) takes the form

$$\int_{I_l} |f(\varpi(s))|^2 \mathrm{d}s \le \varepsilon \Big[\int_{\mathbb{R}^2} |\nabla f(x)|^2 \mathrm{d}x - \alpha \int_{\Sigma} |f(x)|^2 \mathrm{d}x \Big] + C_1(\varepsilon) \int_{\mathbb{R}^2} |f(x)|^2 \mathrm{d}x \,. \tag{5.21}$$

Consequently, the operator $H_{\alpha,\beta\delta}$ associated to $\mathscr{E}_{\alpha,\beta\delta}$ is self adjoint.

Remark 5.2 Negative eigenvalues of $H_{0,\beta\delta}$ for the delta interaction trap. It is worth mentioning that for any negative coupling constant β , the operator $H_{0,\beta\delta}$ always admits at least one negative eigenvalue. However, it should be noted at the same time that in systems of dimensions three or higher, a sufficiently weak coupling constant supported on a finite curve does not generate a discrete spectrum, cf. [7, Thm 4.2(iii)].

The embeddings of R(z) to $L^2(I_l)$ defined by the Dirac delta are given by

$$R(z)|_{0\delta}f = \frac{1}{2\pi} \int_{\mathbb{R}^2} K_0(k_z | x - \varpi(s)|) f(s) \,\mathrm{d}s \, : \, L^2(I_l) \to L^2(\mathbb{R}^2) \,, \tag{5.22}$$

and

$$R(z)|_{\delta\delta} := \frac{1}{2\pi} \int_{I_l} K_0(k_z |\varpi(\cdot) - \varpi(s)|) f(s) \,\mathrm{d}s \, : \, L^2(I_l) \to L^2(I_l) \,, \tag{5.23}$$

moreover, $R(z)|_{\delta 0} = I_{\delta}R(z) : L^2(\mathbb{R}^2) \to L^2(I_l)$. This allows to construct the resolvent $(H_{0,\beta\delta} - z)^{-1}$ by

$$R_{0,\beta\delta}(z) = R(z) + \beta R(z)|_{0\delta} (I - \beta R(z)|_{\delta\delta})^{-1} R(z)|_{\delta0}.$$
(5.24)

Assume that $\beta > 0$. Relying on Remark 5.2 we conclude that the set of negative eigenvalues of $H_{0,\beta\delta}$ is not empty; the same holds for $H_{\alpha,0}$. This means that the assumption formulated for the general measure μ are automatically satisfied. We keep the notation $\mathcal{E}_{\beta;n}$, $n \in \mathcal{N}_{\beta}$ and $\mathcal{E}_{\alpha;k}$, $k \in \mathcal{N}_{\alpha}$. For the delta interaction the inequalities (5.14) take the forms

$$\|R(z)|_{\delta 0}\chi_{\Sigma}\|_{\mathrm{HS}}^{2} \leq \sup_{x \in \Omega} \frac{l}{2\pi} \int_{\Sigma} |K_{0}(k_{z}|x-y|)|^{2} \mathrm{d}y \leq C \frac{l}{|z|^{1/2} \mathcal{T}_{z}} \frac{\mathrm{e}^{-\sqrt{2}\mathcal{T}_{z}\rho}}{\rho} \,.$$
(5.25)

Furthermore, a realization of (5.17) for the discussed special case is given by

$$\Gamma_{n}(\rho) = -\beta^{2} \sum_{k=1}^{j} \frac{\pi}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \left[\left| \int_{I_{l}} (\omega_{\beta;n}(\varpi(s))^{*}\varphi_{k}(\varpi(s)p_{1})) \mathrm{d}s \right|_{p_{1} = (\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}}^{2} + \left| \int_{I_{l}} (\omega_{\beta;n}(\varpi(s))^{*}\varphi_{k}(\varpi(s)), p_{1})) \mathrm{d}s \right|_{p_{1} = -(\mathcal{E}_{\beta;n} - E_{\alpha;k})^{1/2}}^{2} \right].$$
(5.26)

The above formula provides Fermi's golden rule for the resonant pole that appears in a neighborhood of $\mathcal{E}_{\beta;n} \in (E_{\alpha;j}, E_{\alpha;j+1})$, in systems governed by $H_{\alpha,\beta\delta}$.

Remark 5.3 Waveguide with a point interaction. Notably, Fermi's golden rule does not hold in the form analogous to (5.26) for a trap defined by a more singular potential determined by a point interaction. This follows from the fact that a point potential in two dimensional system requires logarithmic regularization, cf. [3], and consequently, the eigenfunctions $\omega_{\beta;n}$ exhibit a corresponding singularity at this point. While we can expect resonances with an exponentially decaying imaginary component, Fermi's golden rule requires modification. We defer the exploration of this topic, along with related concepts, to future research.

Appendix: The asymptotics of further terms contributing to $\Im z_n(\rho)$

First, we discuss the asymptotics of $(w_{\beta;n}, G_{\alpha,\beta}(z)A_n(z)G_{\alpha,\beta}(z)w_{\beta;n})_{L^2(\Omega)}$, which is contributes to (3.28). Repeating the argument from the proof of Theorem 3.3 it suffices to consider this component for $z = \mathcal{E}_{\beta;n}$. Furthermore, since we are interested Fermi's golden rule our aim is to analyze

$$\Im(w_{\beta;n}, G_{\alpha,\beta}(\mathcal{E}_{\beta;n})A_n(\mathcal{E}_{\beta;n})G_{\alpha,\beta}(\mathcal{E}_{\beta;n})w_{\beta;n})_{L^2(\Omega)}.$$
(5.27)

We use the decomposition $G_{\alpha,\beta}(\mathcal{E}_{\beta;n}) = G^R_{\alpha,\beta}(\mathcal{E}_{\beta;n}) + iG^I_{\alpha,\beta}(\mathcal{E}_{\beta;n})$, where $G^R_{\alpha,\beta}(\mathcal{E}_{\beta;n})$ and $G^I_{\alpha,\beta}(\mathcal{E}_{\beta;n})$ are self adjoint. In fact $G^I_{\alpha,\beta}$ is determined by the operator employed in (4.10). Then (5.27) can be written

$$\begin{aligned} \Im(w_{\beta;n}, G_{\alpha,\beta}(\mathcal{E}_{\beta;n})A_n(\mathcal{E}_{\beta;n})G_{\alpha,\beta}(\mathcal{E}_{\beta;n})w_{\beta;n})_{L^2(\Omega)} \\ &= (w_{\beta;n}, G^I_{\alpha,\beta}(\mathcal{E}_{\beta;n})A_n(\mathcal{E}_{\beta;n})G^R_{\alpha,\beta}(\mathcal{E}_{\beta;n})w_{\beta;n})_{L^2(\Omega)} + (w_{\beta;n}, G^R_{\alpha,\beta}(\mathcal{E}_{\beta;n})A_n(\mathcal{E}_{\beta;n})G^I_{\alpha,\beta}(\mathcal{E}_{\beta;n})w_{\beta;n})_{L^2(\Omega)}. \end{aligned}$$
(5.28)

It suffices to estimate one of the above components. Using the expression from (4.10) we get

$$\begin{split} \left|\Im(\chi_{\Omega}\omega_{\beta;n}, G_{\alpha,\beta}^{I}(\mathcal{E}_{\beta;n})A_{n}(\mathcal{E}_{\beta;n})G_{\alpha,\beta}^{R}(\mathcal{E}_{\beta;n})\chi_{\Omega}\omega_{\beta;n})_{L^{2}(\Omega)}\right| \\ &\leq \sum_{k=1}^{j} \frac{\beta}{2\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}} \int_{\Omega \times \Omega} \left|\cos(\sqrt{\mathcal{E}_{\beta;n} - E_{\alpha;k}}|x_{1} - y_{1}|)\phi_{\alpha;k}(x_{2})\phi_{\alpha;k}(y_{2})^{*}\omega_{\beta;n}(x)^{*}\right. \\ &\quad \times (A_{n}(\mathcal{E}_{\beta;n})G_{\alpha,\beta}^{R}(\mathcal{E}_{\beta;n})\omega_{\beta;n})(y)\right| \,\mathrm{d}x \,\mathrm{d}y \\ &\leq \operatorname{const} e^{-2\sqrt{|E_{\alpha,j}|}\rho} |\Omega|^{2} \left\|\omega_{\beta;n}\right\|_{L^{1}(\Omega)} \left\|A_{n}(\mathcal{E}_{\beta;n})G_{\alpha,\beta}^{I}(\mathcal{E}_{\beta;n})\omega_{\beta;n}\right\|_{L^{1}(\Omega)} \\ &\leq \operatorname{const} e^{-2\sqrt{|E_{\alpha,j}|}\rho} |\Omega|^{2} \left\|\omega_{\beta;n}\right\|_{L^{2}(\Omega)} \left\|A_{n}(\mathcal{E}_{\beta;n})G_{\alpha,\beta}^{I}(\mathcal{E}_{\beta;n})\omega_{\beta;n}\right\|_{L^{2}(\Omega)}, \\ &\leq \operatorname{const} \frac{e^{-2\sqrt{|E_{\alpha,j}|}\rho} e^{-\sqrt{2|\mathcal{E}_{\beta;n}|}\rho}}{\rho} |\Omega|^{2} \left\|\omega_{\beta;n}\right\|_{L^{2}(\Omega)}^{2}, \end{split}$$
(5.29)

where we used $||f||_{L^1(\Omega)} \leq |\Omega| ||f||_{L^2(\Omega)}$ for $f \in L^2(\Omega)$, in the last inequality we applied the norm bound for $G_{\alpha,\beta}(\cdot)$, (see (3.11)) and the boundness of $A_n(\mathcal{E}_{\beta;n})$; the constants contained in the symbol 'const' depend on j and coupling constants α and β . The resulting estimate shows that the asymptotics of (5.27) is stronger then (4.19) and, consequently, is does not contribute to the dominated term. The further terms contributing to (3.28) can be estimated analogously.

Acknowledgement

The authors would like to thank the anonymous referees for the careful reading of the paper and for their valuable remarks and comments, which have enhanced its quality. This work was partially supported by a program of the Polish Ministry of Science under the title 'Regional Excellence Initiative', project no. RID/SP/0050/2024/1.

Data Availability Statement

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

Declaration of conflicting interests.

Authors declare that they have no conflicts of interest.

References

- M. Abramowitz and I. Stegun, Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables.
- [2] R. A. Adams, J. J. F. Fournier, Sobolev Spaces, 2nd Edition, Volume 140 June 26, 2003.
- [3] S. Albeverio, F. Gesztesy, R. Hoegh-Krohn and H. Holden, with an appendix by P. Exner, Solvable Models in Quantum Mechanics Second Edition, AMS Chelsea Publishing, Providence, R.I., 2005.
- [4] V. Barrera-Figueroa, V. Rabinovich and S. Loredo-Ramírez, Asymptotic and numerical analysis of slowly varying two-dimensional quantum waveguides, J. Phys. A: Math. Theor. 55 (2022) 095202.
- [5] P. Briet and M. Gharsalli, Stark resonances in a quantum waveguide with analytic curvature, J. Phys. A: Math. Theor. 49 (2016) 495202.
- [6] J. Behrndt, A. F. M. ter Elst and F. Gesztesy, The generalized Birman—Schwinger principle Trans. Amer. Math. Soc. 375 (2022), 799-845.
- [7] J. F. Brasche, P. Exner, Yu. A. Kuperin and P. Šeba, Schrödinger operators with singular interactions, J. Math. Anal. Appl. 184 (1994), 112–139.
- [8] P. L. Christiansen, H. C. Arnbak, A. V. Zolotaryuk, V. N. Ermakov and Y. B. Gaididei, On the existence of resonances in the transmission probability for interactions arising from derivatives of Dirac's delta function, J. Phys. A: Math. Gen. 36 (2003), 7589.
- [9] P. A. M. Dirac, The Quantum Theory of Emission and Absorption of Radiation, Proc. Roy. Soc. A114 243 (1927), 243–265.
- [10] A. Delitsyn and D. S. Grebenkov, Resonance scattering in a waveguide with identical thick perforated barriers, *Applied Mathematics and Computation*, **412** (2022) pp.126592.

- [11] P. Duclos, P. Exner and B. Meller, Exponential bounds on curvature induced resonances in a twodimensional Dirichlet tube, *Helv. Phys. Acta* 71 (1998), 477-492.
- [12] P. Exner and H. Kovařík, Quantum Waveguides, 2015.
- [13] P. Exner, Spectral properties of soft quantum waveguides J. Phys. A: Math. Theor. 53 (2020), No. 355302.
- [14] P. Exner, Soft quantum waveguides in three dimensions, J.Math.Phys. 63(4) (2022), 042103.
- [15] P. Exner, S. Kondej and V. Lotoreichik, Bound states of weakly deformed soft waveguides. Accepted for Asymp. Anal. Preprint on arXiv:2211.01989 [math-ph].
- [16] P. Exner and S. Kondej, Schrödinger operators with singular interactions: a model of tunneling resonances, J. Phys. A: Math. and Gen. 37(34) (2004), 8255.
- [17] P. Exner and V. Lotoreichik, Optimization of the lowest eigenvalue of a soft quantum ring, Lett. Math. Phys. 111(28) (2021).
- [18] P. Exner and D. Spitzkopf, Tunneling in soft waveguides: closing a book, J. Phys. A: Math. Theor. 57 (2024) 125301.
- [19] P. Exner and S. Vugalter, Bound states in bent soft waveguides, accepted for J. Spect. Th., Preprint on arXiv:2304.14776 [math-ph].
- [20] E. Fermi, Nuclear Physics University of Chicago Press 1950.
- [21] F. Gesztesy, H. Holden and R. Nichols On Factorizations of Analytic Operator-Valued Functions and Eigenvalue Multiplicity Questions Integr. Equ. Oper. Theory 8 (2015) 61–94.
- [22] S. Haag, J. Lampart and S. Teufel, Generalised Quantum Waveguides Annales Henri Poincaré 16, (2015) 2535–2568.
- [23] M. Hansmann and D. Krejčiřík, The abstract Birman–Schwinger principle and spectral stability, J. Anal. Math. 148 (2022) 361-398.
- [24] T. Kato, Perturbation Theory for Linear Operators, Springer, 1995.
- [25] C. S. Kim, A. M. Satanin Y. S. Joe and R. M. Cosby, Resonant tunneling in a quantum waveguide: Effect of a finite-size attractive impurity, Phys. Rev. B 60(15) (1999).
- [26] S. Kondej, Quantum soft waveguides with resonances induced by broken symmetry. J. Phys. A: Math. Theor. 57 (2024) 195201.
- [27] S. Kondej, D. Krejčiřík and J. Kříž, Soft quantum waveguides with an explicit cut-locus, J. Phys. A: Math. Theor. 54 (2021).
- [28] S. Kondej, V. Lotoreichik, 2D Schrödinger operators perturbed by finite measures, J. Math. Anal. Appl. 420(2), (2014).
- [29] H. Kovařík and A. Sacchetti, Resonances in twisted quantum waveguides, J. Phys. A: Math. Theor. 40, 29 (2007) 8371.
- [30] D. Krejčiřík, J. Kříž, Bound states in soft quantum layers Publ. RIMS, Kyoto University, 60 (2024) 4.
- [31] J. Lipovský and V. Lotoreichik, Asymptotics of Resonances Induced by Point Interactions Acta. Phys. Pol. A 132 (6) (2017).

- [32] A. Posilicano, A Krein-like Formula for Singular Perturbations of Self-Adjoint Operators and Applications, J. Func. Anal. 183 (2001) 1, 109-147.
- [33] M. Reed, B. Simon. Methods of Modern Mathematical Physics. IV. Analysis of Operators, Academic Press, New York, 1978.
- [34] M. Reed and B. Simon. Methods of Modern Mathematical Physics, vol. I. Functional analysis, Academic Press, New York, 1980.
- [35] B. Simon, The bound state of weakly coupled Schrödinger operators in one and two dimensions, Ann. of Phys. 97, Issue 2, (1976), 279-288.
- [36] B. Simon, Operator Theory. A Comprehensive Course in Analysis, Part 4, American Mathematical Society, Providence, 2015.