LINEARIZATION PROBLEM FOR FINITE SUBGROUPS OF THE PLANE CREMONA GROUP

ANTOINE PINARDIN, ARMAN SARIKYAN, AND EGOR YASINSKY

ABSTRACT. We give a complete solution of the linearization problem in the plane Cremona group over an algebraically closed field of characteristic zero.

Contents

1.	The linearization problem: brief history of previous work		
	1.1.	Finite subgroups of the plane Cremona group	2
	1.2.	The Bogomolov–Prokhorov invariant	3
	1.3.	The Burnside formalism	4
	1.4.	Classification of linearizable actions	5
2.	Rationa	al G -surfaces	7
	2.1.	G-Minimal model program	7
	2.2.	Sarkisov program and birational rigidity	8
	2.3.	G-minimal surfaces	10
3.	Group	theory	11
	3.1.	Klein's classification	11
	3.2.	Blichfeldt's classification	13
	3.3.	Goursat's lemma	14
	3.4.	Actions on sets	15
	3.5.	Generalized dihedral groups	16
4.	G-del F	Pezzo surfaces	17
	4.1.	Degree 4	17
	4.2.	Degree 5	17
	4.3.	Degree 6	18
5.	Finite g	groups acting on smooth quadric surfaces	22
6.	Lineari	zation of finite groups acting on Hirzebruch surfaces	30
	6.1.	G-Hirzebruch surfaces	30
	6.2.	Quadrics with invariant Picard group of rank 1	32
	6.3.	Quadrics with invariant Picard group of rank 2	35
-	-	Magma code	38
Re	ferences		42

²⁰¹⁰ Mathematics Subject Classification. 14E07, 14E05, 14E30, 14J45, 14M22.

1. The linearization problem: Brief History of Previous Work

1.1. Finite subgroups of the plane Cremona group. Given a birational self-map of the projective plane, for example

$$\varphi \colon [x:y:z] \mapsto [x(z-y):z(x-y):xz],$$

it is not at all obvious whether this map is conjugate to a *linear* one within the entire group of birational transformations of the plane, known as the *Cremona group* of rank 2. The map φ is indeed non-regular, as it has the indeterminacy points [1:0:0], [0:1:0], [0:0:1]. As a matter of fact, φ is conjugate over the field $\mathbb{Q}(\sqrt{5})$ to a linear map [BB04]. Letting $\zeta = (1 + \sqrt{5})/2$, the birational involution

$$[x:y:z] \mapsto \left[(x-\zeta y)(y-z)(z-\zeta x): (\zeta^{-1}x-y)(\zeta^{2}y-z)(z-x): (x-y)(\zeta^{2}y-z)(z-\zeta x) \right]$$

conjugates φ to

$$[x:y:z]\mapsto [y-\zeta^{-2}z:y-\zeta^{-1}x:y].$$

Naturally, the problem becomes even more complicated when considering arbitrary finite, not necessarily cyclic, groups and/or specific base fields. Regarding the latter, in this paper we will always work over an algebraically closed field \mathbf{k} of characteristic zero, say $\mathbf{k} = \mathbb{C}$. In this case, the very description of all finite subgroups of the plane Cremona group, denoted $\operatorname{Cr}_2(\mathbf{k})$ hereafter, becomes a challenging task. This problem traces its origins to E. Bertini's work on involutions in $\operatorname{Cr}_2(\mathbb{C})$. Bertini identified three types of conjugacy classes, now referred to as de Jonquières, Geiser, and Bertini involutions. However, his classification was incomplete, and his proofs lacked rigor. Progress continued in 1895 with S. Kantor and A. Wiman, who provided a more detailed description of finite subgroups in $\operatorname{Cr}_2(\mathbb{C})$, though their work was not entirely accurate too.

The modern approach to this problem began with a crucial insight of Yu. Manin and V. Iskovskikh, who discovered a connection between the conjugacy classes of finite subgroups in the Cremona groups and classification of G-varieties up to G-birational equivalence between them (see Section 2). In 2000, L. Bayle and A. Beauville further developed this approach in their study of involutions [BB00]. Later, T. de Fernex extended the classification to subgroups of prime order [dF04], while J. Blanc classified finite abelian subgroups in $\operatorname{Cr}_2(\mathbb{C})$, see [Bla06]. The most comprehensive description of arbitrary finite subgroups in $\operatorname{Cr}_2(\mathbb{C})$ was obtained by I. Dolgachev and V. Iskovskikh in their seminal work [DI09].

What Dolgachev and Iskovskikh did was not only the classification of finite subgroups of $\operatorname{Cr}_2(\mathbb{C})$ up to isomorphism, but also the description of many conjugacy classes, which is clearly a much more delicate problem. For instance, it was established by Beauville and Bayle [BB00] that the conjugacy classes of involutions in $\operatorname{Cr}_2(\mathbb{C})$ are in one-to-one correspondence with the isomorphism classes of their fixed curves. In particular, an involution is *linearizable*, i.e. conjugate in $\operatorname{Cr}_2(\mathbb{C})$ to a linear one, if and only if its fixed curve is rational (notice that in this case it is crucial that the base field is algebraically closed: as shown in a recent work [CMYZ24], the aforementioned correspondence fails over a non-closed field, such as \mathbb{R}).

Nevertheless, despite numerous specific results (see, for example [DI09, Bla09]), the following basic question remained open: which finite subgroups of $\operatorname{Cr}_2(\mathbb{C})$ are linearizable? The goal of this paper is to provide a complete answer to this question. Before presenting our results, let us place the problem in a broader context and describe some existing approaches to solving it.

1.2. The Bogomolov–Prokhorov invariant. Consider the tower of the Cremona groups

$$\operatorname{Cr}_1(\mathbf{k}) \subset \operatorname{Cr}_2(\mathbf{k}) \subset \operatorname{Cr}_3(\mathbf{k}) \subset \operatorname{Cr}_4(\mathbf{k}) \subset \ldots,$$

where $\operatorname{Cr}_i(\mathbf{k})$ denotes the group of birational self-maps of $\mathbb{P}^i_{\mathbf{k}}$, and each embedding $\operatorname{Cr}_i(\mathbf{k}) \subset \operatorname{Cr}_{i+1}(\mathbf{k})$ is induced by adding a new variable. Two subgroups $G_1 \subset \operatorname{Cr}_n(\mathbf{k})$ and $G_2 \subset \operatorname{Cr}_m(\mathbf{k})$ are called *stably conjugate* if they are conjugate in some larger Cremona group $\operatorname{Cr}_N(\mathbf{k}) \supset \operatorname{Cr}_n(\mathbf{k}), \operatorname{Cr}_m(\mathbf{k})$, where $N \ge n, m$.

As will be recalled in Section 2.1, any finite subgroup $G \subset \operatorname{Cr}_n(\mathbf{k})$ is induced by a *biregular* action on some smooth projective rational variety X. The process of replacing the initial rational action of G by a regular one on X is called a *regularization* of G, and then one speaks about G-variety, i.e. the pair (X, G), where $G \subset \operatorname{Aut}(X)$. We say that two G-varieties (X, G) and (Y, G) are stably birational if there exists a G-birational map

$$X \times \mathbb{P}^n \dashrightarrow Y \times \mathbb{P}^m$$
,

where actions on \mathbb{P}^n and \mathbb{P}^m are trivial (so, the case G = id corresponds to classical notion of stable birationality). If (X, G) is stably birational to (\mathbb{P}^N, G) , then we say that G is *stably linearizable*. The following question can be viewed as an analogue of the famous Zariski cancellation problem in our setting:

Question 1.1. Let $G \subset Cr_2(\mathbf{k})$ be a stably linearizable finite subgroup. Is G linearizable?

As was observed by F. Bogomolov and Yu. Prokhorov [BP13], for a smooth projective G-variety X, the cohomology group $\mathrm{H}^1(G, \mathrm{Pic}(X))$ is a G-birational invariant (in the context of rationality questions over non-closed fields this observation goes back to Yu. Manin). Moreover, one has the following.

Theorem ([BP13, Corollary 2.5.1]). If (X, G) and (Y, G) are projective, smooth, and stably birational, then

$$\mathrm{H}^{1}(G, \mathrm{Pic}(X)) \simeq \mathrm{H}^{1}(G, \mathrm{Pic}(Y)).$$

Corollary ([BP13, Corollary 2.5.2]). If (X, G) is stably linearizable, then $H^1(H, Pic(X)) = 0$ for any subgroup $H \subset G$.

If the latter condition holds, the G-variety X is said to be H^1 -trivial. It turns out that in some cases, the invariant $H^1(G, Pic(X))$ can be computed in terms of G-fixed locus: **Theorem 1.2** ([BP13, Theorem 1.1, Corollary 1.2]). Let a finite cyclic group G of prime order p act on a non-singular projective rational surface X. Assume that G fixes (point-wise) a curve of genus g > 0. Then

$$\mathrm{H}^1(G, \mathrm{Pic}(X)) \simeq (\mathbb{Z}/p\mathbb{Z})^{2g}.$$

Moreover, the following are equivalent:

- (1) G is H^1 -trivial;
- (2) (X, G) is linearizable;
- (3) (X,G) is stably linearizable.

In particular, de Jonquières, Bertini and Geiser involutions are not stably linearizable: they all fix a curve of positive genera.

In [Pro15], Prokhorov further applies the invariant introduced by Bogomolov and himself and indicates where to look for negative answers to Question 1.1. He shows, for example, that a del Pezzo surface S acted on by a finite group G with $\operatorname{Pic}(S)^G \simeq \mathbb{Z}$, is H¹-trivial if and only if (i) either $K_S^2 \ge 5$, (ii) or S is the quartic

$$x_1^2 + \omega_3 x_2^2 + \omega_3^2 x_3^2 + x_4^2 = x_1^2 + \omega_3^2 x_2^2 + \omega_3 x_3^2 + x_5^2 = 0$$

in \mathbb{P}^4 , acted on by the group $G \simeq C_3 \rtimes C_4$ generated by two automorphisms

$$(x_1, x_2, x_3, x_4, x_5) \mapsto (x_2, x_3, x_1, \omega_3 x_4, \omega_3^2 x_5), \quad (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_3, x_2, -x_5, x_4) = (x_1, x_2, x_3, x_4, x_5) \mapsto (x_1, x_2, x_3, x_5) \mapsto (x_2, x_3, x_5) \mapsto (x_1, x_2, x_3, x_5) \mapsto (x_1, x_2, x_5) \mapsto ($$

where $\omega_3 = \exp(2\pi i/3)$. Our Main Theorem below shows that only a few groups in these two cases (i) and (ii) are linearizable. Constructions of *stable* linearizability in the remaining nonlinearizable cases are sporadic and remain an open problem even in dimension 2. For instance, as of the time of writing, it is unknown whether the group $G \simeq C_3 \rtimes C_4$ mentioned above is stably linearizable. Some of these constructions are mentioned below, see Remarks 4.5, 4.24, and 4.25.

1.3. The Burnside formalism. Another very recent technique for distinguishing birational actions of finite groups is the *Burnside group formalism*, introduced in the work [KT22a] of A. Kresch and Yu. Tschinkel and generalizing the *birational symbols groups* of M. Kontsevich, V. Pestun and Yu. Tschinkel [KPT23]. Here, we will limit ourselves to a rough sketch of this approach.

In [KT22a], the symbols group $\operatorname{Symb}_n(G)$ was defined as the free abelian group generated by symbols $(H, R \subset \mathbf{K}, \beta)$. Here, $H \subset G$ is an abelian subgroup, $R \subset C_G(H)/H$ is a subgroup, where $C_G(H)$ denotes the centralizer of H, \mathbf{K}/\mathbf{k} is a finitely generated extension of transcendence degree $d \leq n$ faithfully acted on by R, and $\beta = (b_1, \ldots, b_{n-d})$ is a sequence of nontrivial characters of H, generating $\operatorname{Hom}(H, \mathbf{k}^*)$. The quotient of $\operatorname{Symb}_n(G)$ by some involved conjugation and blow-up relations [KT22a, Section 4] gives the equivariant Burnside group $\operatorname{Burn}_n(G)$.

5

Now, let (X, G) be a smooth *n*-dimensional projective variety with a generically free action of G. By [RY00, Theorem 3.2] or [HKT21, Section 7.2], the action of G can be always brought to a standard form via equivariant blow-ups. This means that there is a G-invariant simple normal crossing divisor $\Delta \subset X$ such that G acts freely on $X \setminus \Delta$, and for every $g \in G$ and every irreducible component D of Δ one has either g(D) = D or $g(D) \cap D = \emptyset$. Let $\{D_j\}_{j\in\mathcal{J}}$ be the set of irreducible components with non-trivial (and hence cyclic) stabilizers $H_j \subset G$, considered up to conjugation in G. For each $j \in \mathcal{J}$, the elements of G which do not move D_j to an other component of Δ , give rise to a subgroup $R_j \subset C_G(H_j)/H_j$. Consider the subset $\mathcal{I} \subset \mathcal{J}$ corresponding to those divisors, together with the respective R_j -action, that cannot be obtained via equivariant blow-ups of any standard model of any G-variety (such divisors were called *incompressible*). Finally, let b_j be the character of H_j in the normal bundle to D_j . The formal sum

$$[X \circlearrowright G] = \sum_{i \in \mathcal{I}} (H_i, R_i \subset \mathbf{k}(D_i), b_i), \qquad (1.3)$$

viewed as an element of $\operatorname{Burn}_n(G)$, turns out to be a well-defined *G*-birational invariant [KT22a, Theorem 5.1].

There exist different versions of symbols groups and corresponding Burnside-type obstructions, see e.g. [TYZ24, Section 3] for an overview. The first applications of this formalism yield non-linearizable cyclic actions on certain cubic fourfolds [HKT21, 6]; a new proof [HKT21, 7.6] of non-linearizability of minimal D₆-action on the sextic del Pezzo surface (this is an Iskovskikh's example 4.22); non-conjugacy¹ of certain intransitive and imprimitive subgroups of PGL₃(\mathbb{C}) and PGL₄(\mathbb{C}) in Cr₂(\mathbb{C}) and Cr₃(\mathbb{C}), respectively (by showing that the corresponding actions give different classes in Burn₂ and Burn₃), see [KT22b, Sections 10-11] and [TYZ24, Sections 7-8]. Furthermore, there are examples of non-linearizable actions on some 3-dimensional quadrics [TYZ24, Section 9], and classification (obtained through combinations of various techniques) of non-linearizable actions on some prominent threefolds, such as the Segre cubic, the Burkhardt quartic, and some singular cubic threefolds [CTZ23, CTZ24, CMTZ24].

Some other equivariant birational invariants were recently proposed by L. Esser (the dual complex [Ess24]), T. Ciurca, S. Tanimoto and Yu. Tschinkel (an equivariant version of the formalism of intermediate Jacobian torsor obstructions [CTT24, CTZ24]), and by J. Blanc, I. Cheltsov, A. Duncan and Yu. Prokhorov (the Amitsur subgroup [BCDP23]). We refer the reader to these works for details.

1.4. Classification of linearizable actions. In this paper, we use an extremely powerful method from birational geometry known as the *Sarkisov program*, which, in principle, has no

 $^{^{1}}$ Note that this cannot be approached via the Bogomolov–Prokhorov invariant, which vanishes for linear actions.

limitations and allows to answer the question of the birationality of two Mori fibre spaces, and, in particular, their (equivariant) linearizability. The advantage of this method is that, in dimension 2, it is highly explicit (algorithmic); its application in higher dimensions is much more technically involved. In any case, it has enabled us to prove that the "majority" of subgroups of the plane Cremona group are non-linearizable, while the linearizable ones belong to the following compact list. We use the standard language of G-varieties, which is reminded in Section 2, to formulate our result.

Main Theorem. Let \mathbf{k} be an algebraically closed field of characteristic zero, and $G \subset \operatorname{Cr}_2(\mathbf{k})$ be a finite subgroup. Consider a regularization of G on a two-dimensional G-Mori fibre space S over the base B. Then G is linearizable if and only if the pair (S, G) is one of the following²:

K_S^2	Surface S	$Group \ G \subset \operatorname{Aut}(S)$	Reference			
	G-conic bundles (over $B \simeq \mathbb{P}^1$)					
$K_{S}^{2} = 8$	A Hirzebruch surface \mathbb{F}_n with n odd	— any	Theorem 6.1			
$K_{S}^{2} = 8$	A Hirzebruch surface \mathbb{F}_n with $n > 0$ even	$\begin{array}{c} - \ acts \ cyclically \ on \ B \\ - \ acts \ as \ D_{2m+1} \ on \ B \end{array}$	Theorem 6.1			
$K_{S}^{2} = 8$	The quadric surface $\mathbb{F}_0 \simeq \mathbb{P}^1 \times \mathbb{P}^1$	$\begin{array}{c} - \operatorname{C}_{n} \times_{Q} \operatorname{C}_{m} \\ - \operatorname{C}_{n} \times_{Q} \operatorname{D}_{2m+1} \\ - \operatorname{D}_{2n+1} \times_{Q} \operatorname{D}_{2m+1} \\ is \ dihedral \end{array}$	Theorem 6.14			
	G-del Pezzo surfaces (over $B = pt$)					
$K_{S}^{2} = 5$	The unique quintic del Pezzo surface	$-C_5$ $-D_5$	Proposition 4.4			
$K_{S}^{2} = 6$	The unique sextic del Pezzo surface	$-\mathrm{C}_{6}$ $-\mathrm{S}_{3}$	Proposition 4.23			
$K_{S}^{2} = 8$	The quadric surface $\mathbb{P}^1 \times \mathbb{P}^1$	$-(\mathbf{C}_n\times_Q\mathbf{C}_n)_{\bullet}\mathbf{C}_2$	Proposition 6.10			
$K_{S}^{2} = 9$	The projective plane \mathbb{P}^2	— Blichfeldt's list	Section 3.2			

This paper is organized as follows. In Section 2, we recall some key facts about rational G-surfaces and their minimal models, following the classical works of V. Iskovskikh. We demonstrate how the initial problem reduces to the linearization of finite groups acting on some very specific models. The main technical tool of the paper — the Sarkisov program — is also introduced. This paper employs a significant amount of (elementary) finite group theory, so relevant facts are isolated in Section 3. In Section 4, we derive part of the main theorem

 $^{^2\}mathrm{We}$ refer to Notations 3.1 for the group-theoretic notations.

related to G-del Pezzo surfaces. Although Section 5 is not used later on, it aims to fill a gap in the literature by explicitly describing, in matrix terms, finite groups acting on smooth two-dimensional quadrics (equivalently, finite subgroups of the projective orthogonal group PO(4)). In Section 6, we examine the linearizability of finite groups acting on Hirzebruch surfaces, which finishes the proof of our main result. Finally, in the Appendix we provide the supporting Magma code for Section 5 (note that this code is for the reader's convenience only and essentially is not used in any proof).

Acknowledgement. The authors thank Ivan Cheltsov for helpful discussions. The third author is grateful to the CNRS for its support through the grant PEPS JC/JC 2024.

2. Rational G-surfaces

Throughout the paper, we work over an algebraically closed field \mathbf{k} of characteristic zero. We first recall the classical approach to classification of finite subgroups in $Cr_2(\mathbf{k})$; we refer to [DI09, Section 3] for more details and proofs.

2.1. *G*-Minimal model program. Let *G* be a finite group. A *G*-surface is a triple (S, G, ι) , where *S* is a smooth projective surface over **k** and $\iota: G \hookrightarrow \operatorname{Aut}(S)$ is a monomorphism. A *G*-morphism of *G*-surfaces $(S_1, G, \iota_1) \to (S_2, G, \iota_2)$ is a morphism $f: S_1 \to S_2$ such that $f \circ \iota_1(G) = \iota_2(G) \circ f$. Similarly, one defines *G*-rational maps and *G*-birational maps.

Let X be an algebraic variety over \mathbf{k} and $G \subset Bir(X)$ be a finite group acting birationally and faithfully on X. Then one can resolve indeterminacies of the action of G by the following theorem.

Theorem 2.1 (e.g. [dFE02, Theorem 1.4]). Let X be an algebraic variety and $G \subset Bir(X)$ be a finite group acting birationally and faithfully on X. Then there exists a G-equivariant birational map $\phi: Y \dashrightarrow X$, where Y is a smooth algebraic variety acted on biregularly by G.

In this paper we are interested in the case $X = \mathbb{P}^2$, so the *G*-surface (Y, G, ι) is rational. Hence, after running the *G*-Minimal model program we find that (Y, G, ι) is *G*-birational to one of the following *G*-Mori fibre spaces:

- (i) a *G*-del Pezzo surface, i.e. a *G*-surface (S, G, ι) , such that S is a del Pezzo surface and $\operatorname{Pic}(S)^G \simeq \mathbb{Z}$;
- (ii) a *G*-conic bundle, i.e. a conic bundle $\pi: S \to \mathbb{P}^1$ such that, both S and \mathbb{P}^1 are acted on by G, $-K_S$ is π -ample and $\operatorname{Pic}(S)^G \simeq \mathbb{Z}^2$.


Conversely, given a rational G-surface S, any birational map $\psi: S \dashrightarrow \mathbb{P}^2$ yields an injective homomorphism

$$i_{\psi} \colon G \hookrightarrow \operatorname{Cr}_2(\mathbf{k}), \quad g \mapsto \psi \circ g \circ \psi^{-1}.$$

We say that G (or the *G*-surface *S*) is *linearizable* if there is a birational map $\psi: S \dashrightarrow \mathbb{P}^2$ such that $i_{\psi}(G) \subset \operatorname{Aut}(\mathbb{P}^2) \simeq \operatorname{PGL}_3(\mathbf{k})$. If *S'* is another rational *G*-surface with a birational map $\psi': S' \dashrightarrow \mathbb{P}^2$, then the subgroups $i_{\psi}(G)$ and $i_{\psi'}(G)$ are conjugate if and only if *G*-surfaces *S* and *S'* are *G*-birationally equivalent. In other words, a birational equivalence class of *G*-surfaces defines a conjugacy class of subgroups of $\operatorname{Cr}_2(\mathbf{k})$ isomorphic to *G*. To sum up, there is a natural bijection between the conjugacy classes of finite subgroups $G \subset \operatorname{Cr}_2(\mathbf{k})$ and *G*-birational equivalence classes of rational *G*-Mori fibre spaces of dimension 2.


2.2. Sarkisov program and birational rigidity. According to the equivariant version of the Sarkisov program (see e.g. [DI09, Section 7]), every G-birational map between two G-Mori fibre spaces can be decomposed into a sequence of G-isomorphisms and some "elementary" G-birational maps, called Sarkisov G-links. These links come in four types.

Type I.




where S is a G-del Pezzo surface, η is the blow-up of a G-orbit on S, and $\pi: T \to \mathbb{P}^1$ is a G-conic bundle.

Type II.




where η and η' are blow-ups of *G*-orbits on *S* and *S'*, of lengths *d* and *d'*, correspondently. In this case *S* and *S'* are *G*-del Pezzo surfaces if B = pt, or *S* and *S'* are *G*-conic bundles if $B \simeq \mathbb{P}^1$.

Type III.



This link is the inverse to the link of type I.





This link is the choice of a conic bundle structure on a G-conic bundle T, which has exactly two such structures. Note that in general such link is not represented by a biregular automorphism of T, which exchanges π and π' .


The complete classification of Sarkisov G-links between 2-dimensional G-Mori fibre spaces was obtained by V. Iskovskikh, see [Isk96, Theorem 2.6] and [DI09, Propositions 7.12, 7.13]. We will use extensively this classification in what follows.

Convention. From now on, when talking about Sarkisov links of some specific types, we always use the notation from the diagrams above. For example, for a Sarkisov link of type II starting at a G-del Pezzo surface S, η always denote the blow-up of S.

Some G-del Pezzo surfaces and G-conic bundles have essentially unique structure of a G-Mori fibre space. To be more precise, one introduces the following notions.

Definition 2.2. A *G*-del Pezzo surface *S* is called *G*-birationally rigid if for any *G*-birational map $\chi: S \dashrightarrow S'$ to a *G*-Mori fibre space *S'*, the surfaces *S* and *S'* are *G*-isomorphic (not necessarily via χ). Furthermore, *S* is called *G*-birationally superrigid, if every such map χ is already *G*-isomorphism.

Similarly, a G-conic bundle $\pi: S \to \mathbb{P}^1$ is G-birationally rigid, if for any G-birational map $\chi: S \dashrightarrow S'$ to the total space of another G-conic bundle $\pi': S' \to \mathbb{P}^1$, the latter is square G-birational to the former (not necessarily via χ), i.e. there is a diagram of G-maps



where δ is an isomorphism, and φ is birational. If this diagram holds already for $\varphi = \chi$, then $\pi: S \to \mathbb{P}^1$ is called *G*-birationally superrigid.

Here are two classic examples of G-birationally (super)rigid Mori fibre spaces.

Theorem 2.3 (Manin–Segre). Let S be a G-del Pezzo surface. If $K_S^2 \leq 3$, then S is Gbirationally rigid. If $K_S^2 = 1$, then S is G-birationally superrigid.

A modern proof of this theorem follows from the Noether–Fano method (or, formally, from the classification of Sarkisov G-links [Isk96, Theorem 2.6]).

The birational self-maps of G-conic bundles with at least 8 singular fibres are also easy to describe: they admit only fibrewise birational transformations.

Theorem 2.4 ([Isk69, Theorem 1.6], [CMYZ24, Theorem 2.10]). Let $\pi: S \to \mathbb{P}^1$ be a *G*-conic bundle such that $K_S^2 \leq 0$. Then the following two assertions hold:

- (i) S is not G-birational to a smooth (weak) del Pezzo surface;
- (ii) Moreover, such S is a G-birationally superrigid G-conic bundle.

2.3. *G*-minimal surfaces. So far we have reduced the problem of classification of finite subgroups in $\operatorname{Cr}_2(\mathbf{k})$ to classification of *G*-Mori fibre spaces up to *G*-birational equivalence. Note that *G*-conic bundles $\pi: S \to B$ do not have to be *G*-minimal in the absolute sense, i.e. it is not necessarily true that evey *G*-birational morphism $S \to T$ is *G*-isomorphism (a trivial example is the blow-up $\mathbb{F}_1 \to \mathbb{P}^2$ of a *G*-fixed point on \mathbb{P}^2). However, we have precise description of all such cases. All results below are essentially due to V. Iskovskikh.

Theorem 2.5 ([Isk80, Theorems 4 and 5]). Let $\pi: S \to \mathbb{P}^1$ be a *G*-conic bundle.

- (1) If S is not G-minimal, then S is a del Pezzo surface (in particular, $K_S^2 \ge 1$).
- (2) Assume that $1 \leq K_S^2 \leq 8$. Then S is G-minimal if and only if $K_S^2 \in \{1, 2, 4, 8\}$.
- (3) If $K_S^2 \in \{1, 2, 4\}$ then S is a del Pezzo surface if and only if there are exactly two G-conic bundle structures on S.

Lemma 2.6 (see [Isk80, Theorem 5]). Let S be a rational G-surface with $K_S^2 = 7$. Then S is not a G-Mori fibre space. In other words, S is neither a G-del Pezzo surface, nor a G-conic bundle.

For a G-conic bundle $\pi: S \to \mathbb{P}^1$, Noether's formula implies that $K_S^2 = 8 - c$, where c is the number of singular fibres of π . In particular, $K_S^2 \leq 8$. Besides, Theorem 2.4 says that G-conic bundles with $c \geq 8$ are G-birationally superrigid and hence are not G-birational to \mathbb{P}^2 . For our purposes we may assume that S is also G-minimal (in the absolute sense), hence $K_S^2 \in \{1, 2, 4, 8\}$ by Theorem 2.5.

Proposition 2.7. Let $\pi: S \to \mathbb{P}^1$ be a *G*-conic bundle. If $K_S^2 \in \{1, 2, 4\}$, then *G* is not linearizable.

Proof. Indeed, S is G-minimal by Theorem 2.5. In particular, it does not admit Sarkisov links of type III, i.e. contractions to G-del Pezzo surfaces. Hence, all Sarkisov G-links starting from S are of type II (elementary transformations) or IV and do not change K_S^2 . Therefore, G is not linearizable.

Summary. So, it remains to investigate the linearizability in the following two cases:

- S is a G-del Pezzo surface, where we can assume that $K_S^2 \ge 4$ by Theorem 2.3. If $K_S^2 = 8$, we can assume that S is not the blow-up of \mathbb{P}^2 in a G-fixed point, because such S is not G-minimal; for the same reason, we skip the case $K_S^2 = 7$, see Lemma 2.6.

— S is a G-conic bundle $\pi: S \to \mathbb{P}^1$ with no singular fibres, i.e. S is a Hirzebruch surface \mathbb{F}_n acted on by a finite group G.

3. Group theory

This Section is entirely devoted to auxiliary results from the theory of finite groups. These results are either elementary (nevertheless, we provide proofs for the reader's convenience) or pertain to the classical representation theory.

Notation 3.1. Throughout our paper, we use the following standard notations:

- $-\omega_n = \exp(2\pi i/n)$, where $n \in \mathbb{N}$;
- C_n is a cyclic group of order n;
- $V_4 \simeq C_2 \times C_2$ is Klein's Vierergruppe;
- Q₈ is the quaternion group;
- D_n is the dihedral group of order 2n;
- S_n is the permutation group of degree n;
- A_n is the alternating group of degree n;
- $\operatorname{Hol}(G) = G \rtimes \operatorname{Aut}(G)$ is the holomorph of a group G;
- $F_5 = Hol(C_5)$ is the Frobenius group of order 20;
- D(A) is the generalized dihedral group over an abelian group A;
- $-A_{\bullet}B$ is an extension (not necessarily split) of B with a normal subgroup A;
- $A \times_Q B$ is a fibred product of A and B over their common homomorphic image Q; see Section 3.3 for more details;
- $G \wr S_n$ is the wreath product, i.e. the semi-direct product $G^n \rtimes S_n$, where S_n acts on G^n by permuting the factors;
- $C_G(H)$ is the centralizer of $H \subset G$ in G;
- $N_G(H)$ is the normalizer of $H \subset G$ in G;
- Z(G) is the centre of a group G;
- $\operatorname{Inn}(G)$ is the inner automorphism group of a group G.

3.1. Klein's classification. First, we recall the following classical result of Felix Klein.

Proposition 3.2 ([Kle19]). If **k** is an algebraically closed field of characteristic zero, then every finite subgroup of $PGL_2(\mathbf{k})$ is isomorphic to C_n , D_n (where $n \ge 1$), A_4 , S_4 or A_5 . Moreover, there is only one conjugacy class for each of these groups.

In what follows, we will also need the following more general result which describes algebraic subgroups of $PGL_2(\mathbf{k})$.

Theorem 3.3 ([Kap57, p. 31], [NvdPT08, Theorem 1]). Up to conjugation, every algebraic subgroup of $PGL_2(\mathbf{k})$ is one of the following:

(1) $PGL_2(\mathbf{k})$.

- (2) A finite subgroup from the Klein's list: C_n, D_n, A_4, S_4, A_5 .
- (3) The image under the natural projection π : $SL_2(\mathbf{k}) \rightarrow PSL_2(\mathbf{k}) \simeq PGL_2(\mathbf{k})$ of the Borel (i.e. maximal solvable) subgroup

$$\mathbf{B} = \left\{ \begin{bmatrix} a & b \\ 0 & a^{-1} \end{bmatrix} : a \in \mathbf{k}^*, b \in \mathbf{k} \right\} \subset \mathrm{SL}_2(\mathbf{k}).$$

(4) The image under the natural projection π : $SL_2(\mathbf{k}) \rightarrow PSL_2(\mathbf{k}) \simeq PGL_2(\mathbf{k})$ of the infinite dihedral subgroup

$$\mathbf{D}_{\infty} = \left\{ \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix} : a \in \mathbf{k}^* \right\} \cup \left\{ \begin{bmatrix} 0 & -b \\ b^{-1} & 0 \end{bmatrix} : b \in \mathbf{k}^* \right\}.$$

Remark 3.4. In the case (3), let $\overline{B} = \pi(B)$, and consider the unipotent (abelian) subgroup

$$\overline{\mathbf{U}} = \left\{ \overline{\begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix}} : b \in \mathbf{k} \right\} \subset \overline{\mathbf{B}} = \left\{ \overline{\begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix}} : a \in \mathbf{k}^*, b \in \mathbf{k} \right\}$$

The elements of \overline{B} can be viewed as affine automorphisms of \mathbb{P}^1 , locally given by $z \mapsto az + b$. The map which sends every such automorphism to $a \in \mathbf{k}^*$ is a well-defined group homomorphism, which induces a short exact sequence

$$1 \longrightarrow \overline{\mathbf{U}} \longrightarrow \overline{\mathbf{B}} \longrightarrow \mathbf{k}^* \longrightarrow 1.$$
(3.5)

In the case (4), letting $\overline{\mathbb{D}}_{\infty} = \pi(\mathbb{D}_{\infty})$, we observe that the diagonal matrices diag $\{a, a^{-1}\}$, where $a \in \mathbf{k}^*$, constitute the connected component of the identity $\mathbb{D}_{\infty}^{\circ}$ and are mapped by π to a group isomorphic to \mathbf{k}^* . The other matrices are mapped to involutions. Furthermore, one has $\overline{\mathbb{D}}_{\infty} \simeq \mathbf{k}^* \rtimes \mathbb{C}_2$, where the action is by inversion: $t \mapsto t^{-1}$, i.e. $\overline{\mathbb{D}}_{\infty} \simeq \operatorname{Hol}(\mathbf{k}^*)$.

Let us fix the following presentations [Hup67, §19]:

- $C_n \simeq \langle r \mid r^n = \mathrm{id} \rangle.$
- $D_n \simeq \langle r, b \mid r^n = b^2 = (rb)^2 = \mathrm{id} \rangle.$
- $A_4 \simeq \langle a, b, c \mid a^2 = b^2 = c^3 = id$, $cac^{-1} = ab = ba$, $cbc^{-1} = a\rangle$. Note that $\langle a, b \rangle \simeq V_4$ is the derived subgroup of A_4 and one has $A_4 \simeq \langle a, b \rangle \rtimes \langle c \rangle$.
- $S_4 \simeq \langle a, b, c, d \mid a^2 = b^2 = c^3 = d^2 = \mathrm{id}, \ cac^{-1} = dad = ab = ba, \ cbc^{-1} = a, \ bd = db, \ dcd = c^{-1} \rangle.$ Here, one finds a unique copy of $A_4 \simeq \langle a, b, c \rangle$ inside, which is the derived subgroup of S_4 , and one has $S_4 \simeq \langle a, b, c \rangle \rtimes \langle d \rangle.$ $A_5 \simeq \langle e, f \mid e^5 = f^2 = (ef)^3 = \mathrm{id} \rangle.$

Notation 3.6. Let us fix the following identifications (see e.g. [Fab23, Theorem C]):

$$r \mapsto \mathbf{R}_n = \begin{bmatrix} 1 & 0 \\ 0 & \omega_n \end{bmatrix}, \ a \mapsto \mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \ b \mapsto \mathbf{B} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \ c \mapsto \mathbf{C} = \begin{bmatrix} i & -i \\ 1 & 1 \end{bmatrix},$$

$$d \mapsto \mathbf{D} = \begin{bmatrix} 1 & -i \\ i & -1 \end{bmatrix}, \ e \mapsto \mathbf{E} = \begin{bmatrix} \omega_5 & 0 \\ 0 & 1 \end{bmatrix}, \ f \mapsto \mathbf{F} = \begin{bmatrix} 1 & 1 - \omega_5 - \omega_5^{-1} \\ 1 & -1 \end{bmatrix}.$$

Then these identifications define embeddings of C_n , D_n , A_4 , S_4 and A_5 into $PGL_2(\mathbf{k})$.

The following description of orbits on \mathbb{P}^1 is classical:

Proposition 3.7 ([Spr77, 4.4]). Let G be a finite subgroup of $PGL_2(\mathbf{k})$. One has the following.

- (1) If $G \simeq C_n$ then it has 2 fixed points on \mathbb{P}^1 , namely [1:0] and [0:1], and any other point generates an orbit of length n.
- (2) If $G \simeq D_n$, then it has one orbit of length 2, namely [1:0] and [0:1], and one orbit of length n generated by [1:1]. Any other point generates an orbit of length 2n.
- (3) If $G \simeq A_4$ then it has two orbits of length 4 and one orbit of length 6. All other orbits are of length 12.
- (4) If $G \simeq S_4$ then it has one orbit of length 6, one orbit of length 8 and one orbit of length 12. All other orbits are of length 24.
- (5) If $G \simeq A_5$ then it has one orbit of length 12, one orbit of length 20, and one orbit of length 30. All other orbits are of length 60.

3.2. Blichfeldt's classification. Since we are interested in linearization of finite subgroups of $Cr_2(\mathbf{k})$, let us first recall the classification of finite subgroups of $PGL_3(\mathbf{k})$, which is essentially due to H. Blichfeldt.

Definition 3.8 ([Bli17]). We call a subgroup $\iota: G \hookrightarrow \operatorname{GL}_n(\mathbf{k})$ intransitive if the representation ι is reducible, and transitive otherwise. Further, a transitive group G is called imprimitive if there is a decomposition $\mathbf{k}^n = \bigoplus_{i=1}^m V_i$ into a direct sum of subspaces and G transitively acts on the set $\{V_i\}$. A transitive group G is called primitive if there is no such decomposition. Finally, we say that $G \subset \operatorname{PGL}_n(\mathbf{k})$ is (in)transitive or (im)primitive if its lift to $\operatorname{GL}_n(\mathbf{k})$ is such a group.

So, intransitive subgroups of $PGL_3(\mathbf{k})$ come in two types:

- I_1 The representation of G in $\operatorname{GL}_3(\mathbf{k})$ is a direct sum of three 1-dimensional representations. In other words, G fixes 3 non-collinear points on \mathbb{P}^2 and hence $G \simeq \operatorname{C}_n \times \operatorname{C}_m$, where $n, m \ge 1$, is a diagonal abelian group.
- I_2 The representation of G in $\operatorname{GL}_3(\mathbf{k})$ is a direct sum of 1-dimensional and 2-dimensional representations. In other words, G fixes a point $p \in \mathbb{P}^2$, and hence there is an embedding $G \hookrightarrow \operatorname{GL}(T_p \mathbb{P}^2) \simeq \operatorname{GL}_2(\mathbf{k})$. Obviously, every finite subgroup of $\operatorname{GL}_2(\mathbf{k})$ gives rise to an intransitive subgroup of $\operatorname{PGL}_3(\mathbf{k})$.

Assume that G is imprimitive. Then $\mathbf{k}^3 = V_1 \oplus V_2 \oplus V_3$, where $V_1 \simeq V_2 \simeq V_3 \simeq \mathbf{k}$ and G acts transitively on the set $\{V_1, V_2, V_3\}$. Hence G fits into the short exact sequence

 $1 \longrightarrow A \longrightarrow G \longrightarrow B \longrightarrow 1$

GAP	Order	Isomorphism class	Comments	
Primitive subgroups				
[36, 9]	36	$(\mathrm{C}_3\times\mathrm{C}_3)\rtimes\mathrm{C}_4$	subgroup of the Hessian group	
[72, 41]	72	$(\mathrm{C}_3\times\mathrm{C}_3)\rtimes\mathrm{Q}_8$	subgroup of the Hessian group	
[216, 153]	216	$(C_3 \times C_3) \rtimes SL_2(\mathbb{F}_3)$	the Hessian group	
[60, 5]	60	A_5	the simple icosahedral group	
[168, 42]	168	$\mathrm{PSL}_2(\mathbb{F}_7)$	the simple Klein group	
[360, 118]	360	A_6	the simple Valentiner group	

where $B \simeq C_3$ or $B \simeq S_3$, and $A \subset PGL_3(\mathbf{k})$ is an intransitive subgroup of type I_1 . Finally, there are 6 primitive subgroups of $PGL_3(\mathbf{k})$.

3.3. Goursat's lemma. Let Γ_1 and Γ_2 be two finite groups. Finite subgroups of the direct product $\Gamma_1 \times \Gamma_2$ can be determined using the classical *Goursat's lemma*. Recall that the *fibre product* of two groups G_1 and G_2 over a group Q is defined as

$$G_1 \times_Q G_2 = \{ (g_1, g_2) \in G_1 \times G_2 \colon \alpha(g_1) = \beta(g_2) \},\$$

where $\alpha: G_1 \to Q$ and $\beta: G_2 \to Q$ are surjective homomorphisms. Note that the data defining $G_1 \times_Q G_2$ is not only the groups G_1, G_2 and Q but also the homomorphisms α, β .

Lemma 3.9 (Goursat's lemma, [Gou89, p. 47]). Let Γ_1 and Γ_2 be two finite groups. There is a bijective correspondence between subgroups $G \subseteq \Gamma_1 \times \Gamma_2$ and 5-tuples $\{G_1, G_2, H_1, H_2, \varphi\}$, where G_1 is a subgroup of Γ_1 , H_1 is a normal subgroup of G_1 , G_2 is a subgroup of Γ_2 , H_2 is a normal subgroup of G_2 , and $\varphi: G_1/H_1 \xrightarrow{\sim} G_2/H_2$ is an isomorphism. Namely, the group corresponding to this 5-tuple is

$$G = \{ (g_1, g_2) \in G_1 \times G_2 \colon \varphi(g_1 H_1) = g_2 H_2 \}.$$

Conversely, let $G \subseteq \Gamma_1 \times \Gamma_2$ be a subgroup. Denote by $p_1 \colon \Gamma_1 \times \Gamma_2 \to \Gamma_1$ and $p_2 \colon \Gamma_1 \times \Gamma_2 \to \Gamma_2$ the natural projections, and set $G_1 = p_1(G)$, $G_2 = p_2(G)$. Set

 $H_1 = \ker p_2|_G = \{(g_1, \mathrm{id}) \in G, \ g_1 \in \Gamma_1\}, \ H_2 = \ker p_1|_G = \{(\mathrm{id}, g_2) \in G, \ g_2 \in \Gamma_2\},$

whose images by p_1 and p_2 define normal subgroups of G_1 and G_2 , respectively (denoted the same). Let $\pi_1: G_1 \to G_1/H_1$ and $\pi_2: G_2 \to G_2/H_2$ be the quotient homomorphisms. The map $\varphi: G_1/H_1 \to G_2/H_2$, $\varphi(g_1H_1) = g_2H_2$, where $g_2 \in \Gamma_2$ is any element such that $(g_1, g_2) \in G$, is an isomorphism. Furthermore, $G = G_1 \times_Q G_2$, where $Q = G_1/H_1$, $\alpha = \pi_1$, and $\beta = \varphi^{-1} \circ \pi_2$. **Lemma 3.10.** In the above notation, any subgroup $G \subseteq \Gamma_1 \times \Gamma_2$ fits into the short exact sequence

$$1 \longrightarrow H_1 \times H_2 \longrightarrow G \longrightarrow Q \longrightarrow 1$$

Proof. Indeed, the restriction of the homomorphism $\alpha \times \beta \colon G_1 \times G_2 \to Q \times Q$ to G has the kernel $H_1 \times H_2$ and the image is isomorphic to $\{(t,t) \in Q \times Q\} \simeq Q$.

Remark 3.11. We will often use explicit generators to describe a fibre product. They are given as follows. Let S be a subset of G_1 such that $\alpha(S) = Q$. Then the group $G_1 \times_Q G_2$ is generated by ker $\alpha \times \ker \beta$, and $\{(g_1, g_2), g_1 \in S, \alpha(g_1) = \beta(g_2)\}$.

3.4. Actions on sets. We will use several elementary lemmas about group actions on a set.

Lemma 3.12. Let A and B be two finite groups, and $G \subset A \times B$ be a subgroup such that the natural projection $G \to A$ is surjective. Assume that A acts on a set X and B acts on a set Y. Consider the induced action of G on $X \times Y$. Then the length of any orbit of G in $X \times Y$ is divisible by the length of some orbit of A on X.

Proof. Pick a point $(x_0, y_0) \in X \times Y$ and consider its *G*-orbit $\Sigma = \{(ax_0, by_0): (a, b) \in G\}$. Consider the projection $p_X: X \times Y \to X$. We claim that the set $\Sigma_X = p_X(\Sigma) \subset X$ is an *A*-orbit. Firstly, it is clearly *A*-invariant: taking any $x \in \Sigma_X$ there exists $y \in Y$ such that $(x, y) \in \Sigma$. By our assumption, for any $a \in A$ there exists $b \in B$ such that $(a, b) \in G$. Then $(ax, by) \in \Sigma$ and therefore $ax \in \Sigma_X$. Secondly, Σ_X is clearly a minimal *A*-invariant set. For if there is a proper *A*-invariant subset $\Sigma'_X \subsetneq \Sigma_X$, then taking two elements $x' \in \Sigma'_X$, $x \in \Sigma_X \setminus \Sigma'_X$ and their lifts $(x', y') \in \Sigma$, $(x, y) \in \Sigma$, there exists an element $(a, b) \in G$ such that (ax', by') = (x, y). In particular, $x = ax' \notin \Sigma'_X$; but this contradicts to Σ'_X being *A*-invariant.

It remains to show that for each $x_1 \in \Sigma_X$, the cardinality of the fibre $p_X^{-1}(x_1) \cap \Sigma$ is the same and equals to the cardinality of $p_X^{-1}(x_0) \cap \Sigma$. Indeed, there exists $a \in A$ such that $ax_0 = x_1$. Pick any lift $(a, b) \in G$ and consider the map of finite sets

$$p_X^{-1}(x_0) \cap \Sigma \to p_X^{-1}(x_1) \cap \Sigma, \ (x_0, y) \mapsto (ax_0, by) = (x_1, by),$$

which is obviously well defined and injective. The same holds for the map $p_X^{-1}(x_1) \cap \Sigma \to p_X^{-1}(x_0) \cap \Sigma$ which sends (x_1, y) to $(a^{-1}x_1, b^{-1}y) = (x_0, b^{-1}y)$. This finishes the proof. \Box

Lemma 3.13. Let G be a finite group and $H \subset G$ be a normal subgroup of finite index [G:H]. Assume that G acts on a set X and let Σ be an orbit of G. Write $\Sigma = \Sigma_1 \sqcup \ldots \sqcup \Sigma_n$ as the disjoint union of H-orbits Σ_i . Then all Σ_i have the same length and n divides [G:H]. In particular, the length of Σ is always divisible by the length of some orbit of H.

Proof. Indeed, the quotient group G/H acts on the space of orbits Σ/H , and this action is transitive. Therefore, by the Orbit-Stabilizer Theorem, [G:H] is divisible by the cardinality of Σ/H , i.e. by *n*. Consider any two *H*-orbits Σ_i and Σ_j , let $x_i \in \Sigma_i$ and $x_j \in \Sigma_j$, so that

 $\Sigma_i = Hx_i$ and $\Sigma_j = Hx_j$. By transitivity of G on Σ , there exists $g \in G$ such that $gx_i = x_j$. Then the map $\Sigma_i \to \Sigma_j, y \mapsto gy$ is a bijection.

3.5. Generalized dihedral groups. We remind the following elementary fact.

Lemma 3.14. Subgroups of the dihedral group $D_n = \langle r, s | r^n = s^2 = (sr)^2 = id \rangle$ are the following:

Cyclic: $\langle r^d \rangle \simeq C_{n/d}$, and $\langle r^k s \rangle$, where d divides n and $0 \leq k \leq n-1$. Dihedral: $\langle r^d, r^k s \rangle \simeq D_{n/d}$, where d < n divides n, and $0 \leq k \leq d-1$.

All cyclic subgroups $\langle r^d \rangle$ are normal, one has $D_n/\langle r^d \rangle \simeq D_d$, and these are all normal subgroups when n is odd. When n is even, there are two additional normal dihedral subgroups of index 2, namely $\langle r^2, s \rangle$ and $\langle r^2, rs \rangle$.

Let A be an abelian group. Recall that the generalized dihedral group D(A) is the semidirect product $A \rtimes C_2$, where C_2 acts by inverting the elements of A. In particular, for $A \simeq C_n$ one has $D(C_n) \simeq D_n$.

Lemma 3.15. Let $n, m \ge 3$ be odd integers. Then one has the following:

- (1) Any group $G = D_n \times_Q D_m$ is either $D_n \times D_m$, or a generalized dihedral group D(A), where A is a direct product of at most two cyclic groups.
- (2) Furthermore, $G \simeq D(A)$ admits a faithful 2-dimensional representation if and only if A is cyclic, so that G is isomorphic to a dihedral group.

Proof. Let us fix the presentations $D_n = \langle r_1, s_1 | r_1^n = s_1^2 = (s_1r_1)^2 = id \rangle$, $D_m = \langle r_2, s_2 | r_2^m = s_2^2 = (s_2r_2)^2 = id \rangle$. We then have the following possibilities for G:

- (i) $Q \simeq$ id. In this case, Lemma 3.10 implies that $G \simeq D_n \times D_m$.
- (ii) $Q \simeq C_2$. Since *n* and *m* are odd, Lemma 3.10 and Lemma 3.14 show that *G* is an extension of C_2 by $A = C_n \times C_m$. Let $\tau \in G$ be an element mapped to the generator of *Q*. It necessarily belongs to $D_n \times D_m \setminus A$ and, being in the fibre product, is of the form $(s_1r_1^k, s_2r_2^l)$. The conjugation by τ induces an inversion on *A*, hence the claim.
- (iii) $Q \simeq D_q$, where q divides both n and m. Then the group G fits into the short exact sequence

$$1 \longrightarrow \mathcal{C}_{n/q} \times \mathcal{C}_{m/q} \longrightarrow G \xrightarrow{\psi} \mathcal{D}_q \longrightarrow 1,$$

where the groups $C_{n/q}$ and $C_{m/q}$ are generated by r_1^q and r_2^q , respectively. Let $A = \psi^{-1}(C_q)$. This is an index 2 subgroup in G of odd order; therefore, all its elements are of the form (r_1^k, r_2^l) . Hence A is a subgroup of $C_n \times C_m \subset D_n \times D_m$. As in the previous case, we conclude that $G \simeq D(A)$.

To prove the second claim, it is enough to notice that Z(D(A)) = id for any non-trivial $A \subset C_n \times C_m$, since n and m are odd. In particular, if G admits a faithful 2-dimensional

representation, then G can be embedded in $PGL_2(\mathbf{k})$, hence A is a cyclic group. The converse is obvious.

4. G-del Pezzo surfaces

4.1. **Degree 4.** By Theorem 2.3, none of the groups G which act biregularly on a del Pezzo surface S with $\operatorname{Pic}(S)^G \simeq \mathbb{Z}$ and $K_S^2 \leq 3$ is linearizable. Let S be a G-del Pezzo surface of degree 4 and $\varphi \colon S \dashrightarrow S'$ be a G-birational map to another G-del Pezzo surface S'. It follows then from the classification of Sarkisov G-links that φ , if not an isomorphism, fits into the commutative diagram of G-maps

where π is the blow-up of a *G*-fixed point on *S*, which leads to a cubic surface *T* equipped with a structure of a *G*-conic bundle (so π is a link of type I); the maps χ_i are elementary transformations of *G*-conic bundles (links of type II), and π' is the blow-down of a *G*-orbit of (-1)-curves, where *S'* is again a *G*-del Pezzo surface of degree 4. We conclude that none of such groups *G* is linearizable.

4.2. **Degree 5.** Let S be a G-del Pezzo surface of degree 5. Recall that there is a single isomorphism class of del Pezzo surfaces of degree 5 over $\mathbf{k} = \overline{\mathbf{k}}$. Every Sarkisov G-link starting from S is of type II, where η blows up a G-orbit of length d, and one of the following holds:

- (1) $S \simeq S', d = 4, \chi$ is a birational Bertini involution;
- (2) $S \simeq S', d = 3, \chi$ is a birational Geiser involution;
- (3) $S' \simeq \mathbb{P}^1 \times \mathbb{P}^1, d = 2;$
- (4) $S' \simeq \mathbb{P}^2, d = 1.$

The following Proposition lists all possible groups G.

Proposition 4.2 ([DI09, Theorem 6.4]). Let S be a del Pezzo surface of degree 5 and $G \subset Aut(S)$ be a group such that $Pic(S)^G \simeq \mathbb{Z}$. Then G is isomorphic to one of the following five groups:

 $S_5, A_5, F_5, D_5, C_5,$

where $F_5 = \langle a, b \mid a^5 = b^4 = 1, bab^{-1} = a^3 \rangle$ is the Frobenius group³ of order 20.

Proposition 4.3 ([Che08, Example 6.3], [Che14, Theorem B.10]). The del Pezzo surface of degree 5 is A_5 - and S_5 -birationally superrigid.

Proof. None of these two groups has faithful 2-dimensional representations over \mathbb{C} , hence none of them has an orbit of size 1 or 2 on our surface: recall [BB73, Lemma 2.4] that if X

 3 GAP ID [20,3]

is an irreducible algebraic variety and $G \subset \operatorname{Aut}(X)$ is a finite group fixing a point $p \in X$ then the induced linear representation $G \hookrightarrow \operatorname{GL}(T_pX)$ is faithful. Furthermore, these groups have no subgroups of index 3 or 4, hence there are no equivariant birational Bertini or Geiser involutions on (with respect to these groups). The result follows. \Box

Proposition 4.4. Let S be a G-del Pezzo surface of degree 5. Then G is linearizable if and only if $G \simeq C_5$ or $G \simeq D_5$.

Proof. Proposition 4.3 implies that the groups A_5 and S_5 are not linearizable; in fact, S_5 is not a subgroup of $PGL_3(\mathbb{C})$. The Frobenius group F_5 is not a subgroup of $PGL_3(\mathbb{C})$ either: according to description given in Section 3.2, it is obviously not transitive, and moreover has no faithful 2-dimensional representations (in fact, by [Wol18, Theorem 1.1], there exists a unique F_5 -del Pezzo surface which is F_5 -birational to S, namely $\mathbb{P}^1 \times \mathbb{P}^1$).

We now show that the groups C_5 and D_5 are linearizable. Let G be any of these groups. It is enough to construct an explicit G-equivariant map $\mathbb{P}^2 \dashrightarrow S$, where S is a G-del Pezzo surface of degree 5, because there is a single G-isomorphism class of such surfaces. Indeed, there is a single conjugacy class in $\operatorname{Aut}(S) \simeq S_5$ of C_5 and D_5 , so any isomorphism $S \xrightarrow{\sim} S'$ to the del Pezzo surface S' of degree 5 can be made a G-isomorphism, after composing with a suitable automorphism of S'. Now, to construct an explicit G-birational map $\mathbb{P}^2 \dashrightarrow S$, one can take two linear automorphisms

$$r \colon [x:y:z] \mapsto [x:\omega_5 y:\omega_5^{-1} z], \quad s \colon [x:y:z] \mapsto [x:z:y],$$


and let $G_1 = \langle r \rangle \simeq C_5$, $G_2 = \langle r, s \rangle \simeq D_5$. The G_1 - and G_2 -orbit of the point [1:1:1] is a set of 5 points in general position on \mathbb{P}^2 , lying on the unique smooth conic $Q \subset \mathbb{P}^2$. By blowing up these points and contracting the strict transform of Q we get a G_1 -birational (respectively, G_2 -birational) map from \mathbb{P}^2 to S. \Box

Remark 4.5. The actions of A_5 and S_5 are stably linearizable. More precisely, by [Pro10, Proposition 4.7], Prokhorov shows that $S \times \mathbb{P}^1$ is S_5 -birational to the Segre cubic threefold $\sum_{i=0}^5 x_i = \sum_{i=0}^5 x_i^3 = 0$, which is obviously acted on by S_6 . Recall that, up to conjugation, the group S_6 has two subgroups isomorphic to S_5 , given by the standard and the non-standard embeddings (which differ by an outer automorphism of S_6). In the construction above, $S \times \mathbb{P}^1$ turns out to be S_5 -birational to the Segre cubic with a non-standard action of S_5 , which is linearizable. Recently, B. Hassett and Yu. Tschinkel gave another proof of this fact using the equivariant torsor formalism [HT23, §8.2].

4.3. **Degree 6.** Let S be a del Pezzo surface of degree $K_S^2 = 6$. Then S is the blow-up $\pi: S \to \mathbb{P}^2$ in three non-collinear points p_1, p_2, p_3 , which we may assume to be [1:0:0], [0:1:0] and [0:0:1], respectively. The surface S can be given as

$$\left\{ ([x_0:x_1:x_2], [y_0:y_1:y_2]) \in \mathbb{P}^2 \times \mathbb{P}^2 : \ x_0 y_0 = x_1 y_1 = x_2 y_2 \right\}.$$
(4.6)

The set of (-1)-curves on S consists of six curves: the exceptional divisors of blow-up $e_i = \pi^{-1}(p_i)$ and the strict transforms of the lines d_{ij} passing through p_i , p_j . In the anticanonical embedding $S \hookrightarrow \mathbb{P}^6$ these (-1)-curves form a hexagon Σ ; the configuration of lines is shown in the diagram (4.7) below.



Note that Σ is naturally acted on by Aut(S), so there is a homomorphism

$$\Phi: \operatorname{Aut}(S) \to \operatorname{Aut}(\Sigma) \simeq D_6 = \langle r, s \mid r^6 = s^2 = \operatorname{id}, srs^{-1} = r^{-1} \rangle,$$

where r is a rotation by $\pi/3$ and s is a reflection. The kernel of Φ is the maximal torus $\mathbb{T} \subset \mathrm{PGL}_3(\mathbf{k})$, isomorphic to $(\mathbf{k}^*)^3/\mathbf{k}^* \simeq (\mathbf{k}^*)^2$, and acts on S by

$$(\lambda_0, \lambda_1, \lambda_2) \cdot ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) = ([\lambda_0 x_0 : \lambda_1 x_1 : \lambda_2 x_2], [\lambda_0^{-1} y_0 : \lambda_1^{-1} y_1 : \lambda_2^{-1} y_2]).$$
(4.8)

The action of \mathbb{T} on $S \setminus \Sigma$ is faithful and transitive, and the automorphism group of Aut(S) fits into the short exact sequence

 $1 \longrightarrow \mathbb{T} \longrightarrow \operatorname{Aut}(S) \xrightarrow{\Phi} \mathcal{D}_6 \longrightarrow 1$

with $\Phi(\operatorname{Aut}(S)) \simeq D_6 \simeq S_3 \times C_2$, where C_2 is the lift of the standard quadratic involution, acting by

$$\iota: ([x_0:x_1:x_2], [y_0:y_1:y_2]) \mapsto ([y_0:y_1:y_2], [x_0:x_1:x_2]),$$
(4.9)

and S₃ acts naturally by permutations on each of the two triples x_0, x_1, x_2 and y_0, y_1, y_2 . In what follows, we will denote

$$\theta: ([x_0:x_1:x_2], [y_0:y_1:y_2]) \mapsto ([x_1:x_2:x_0], [y_1:y_2:y_0])$$
(4.10)

the generator of $C_3 \subset S_3$. Then D_6 is generated by the element $\rho = \theta \circ \iota$ of order 6 (rotation) and the element σ of order 2 (reflection), whose actions are given by

$$\rho: ([x_0:x_1:x_2], [y_0:y_1:y_2]) \mapsto ([y_1:y_2:y_0], [x_1:x_2:x_0]),$$
(4.11)

$$\sigma \colon ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([y_0 : y_2 : y_1], [x_0 : x_2 : x_1]).$$

$$(4.12)$$

Every Sarkisov link $\chi: S \dashrightarrow S'$ is of type II and starts with blowing up a G-orbit of length d. Then one of the following holds:

- (1) $S \simeq S', d = 5, \chi$ is a birational Bertini involution;
- (2) $S \simeq S', d = 4, \chi$ is a birational Geiser involution;
- (3) $d = 3, K_{S'}^2 = 6;$
- (4) $d = 2, K_{S'}^2 = 6;$
- (5) $d = 1, S' \simeq \mathbb{P}^1 \times \mathbb{P}^1.$

Remark 4.13. In the cases (3) and (4), the surfaces S' does not have to be *G*-isomorphic to *S*, see [Yas24, Example 3.9].

Lemma 4.14 ([Yas24, Lemma 3.7]). Let S be a G-del Pezzo surface of degree 6. Then G is of the form

$$T_{\bullet}\langle r \rangle \simeq T_{\bullet}C_6, \qquad T_{\bullet}\langle r^2, s \rangle \simeq T_{\bullet}S_3, \quad or \quad T_{\bullet}\langle r, s \rangle \simeq T_{\bullet}D_6,$$

where $T \simeq C_n \times C_m$ is a subgroup of \mathbb{T} .

Remark 4.15. The dihedral group $D_6 = \langle r, s \rangle$ contains two groups isomorphic to S_3 , but only for one of them the *G*-invariant Picard number is 1, namely for $\langle r^2, s \rangle$, which we denote S_3^{\min} . The group $\langle r^2, rs \rangle$ will be denoted S_3^{\min} . This is the quotient of D_6 by its centre $Z(D_6) \simeq C_2$.

The following lemma shows that we do not lose generality by choosing particular actions of the groups C_6 , S_3 and D_6 .

Lemma 4.16. Let S be a G-del Pezzo surface of degree 6, and G be isomorphic to C₆, S₃ or D₆. Then G is conjugate in Aut(S) to the groups $\langle \rho \rangle$, $\langle \rho^2, \sigma \rangle$ and $\langle \rho, \sigma \rangle$, respectively.

Proof. This is the content of [Pin24b, Propositions 5.6, 5.7, 5.8]. We give a short direct proof here for the largest of these groups $G \simeq D_6$. We may assume that it is generated by two elements $\overline{\rho}$ and $\overline{\sigma}$ whose images in Aut(Σ) are as in (4.11). The map $\overline{\rho}$ is given by

$$\overline{\rho}: ([x_0:x_1:x_2], [y_0:y_1:y_2]) \mapsto ([uy_1:vy_2:y_0], [u^{-1}x_1:v^{-1}x_2:x_0])$$
(4.17)

for some $u, v \in \mathbf{k}^*$. The map

$$\beta : ([x_0 : x_1 : x_2], [y_0 : y_1 : y_2]) \mapsto ([x_0 : v^{-1}x_1 : uv^{-1}x_2], [y_0 : vy_1, vu^{-1}y_2])$$

then conjugates $\overline{\rho}$ to ρ . After this conjugation, $\overline{\sigma}$ is given by

 $\overline{\sigma}: ([x_0:x_1:x_2], [y_0:y_1:y_2]) \mapsto ([ay_0:by_2:y_1], [a^{-1}x_0:b^{-1}x_2:x_1])$

for some $a, b \in \mathbf{k}^*$. The condition $\overline{\sigma}^2 = \text{id implies } b = 1$, while the condition $\overline{\sigma} \circ \rho \circ \overline{\sigma} = \rho^{-1}$ implies a = 1.

Lemma 4.18. Let S be a G-del Pezzo surface of degree 6. Then G fixes a point on S if and only if $G \cap \mathbb{T} = {\text{id}}$.

Proof. Assume that G fixes a point on S, but $G \cap \mathbb{T} \neq \{\text{id}\}$. Since \mathbb{T} can be identified with a subgroup of $\text{PGL}_3(\mathbf{k})$ which fixes 3 points on \mathbb{P}^2 , an element of \mathbb{T} fixing a point on $S \setminus \Sigma$ is necessarily trivial. Therefore, a fixed point $p \in S$ of G lies on Σ . But this implies $\operatorname{rk}\operatorname{Pic}(S)^G > 1$.

Conversely, suppose that $G \cap \mathbb{T} = \{\text{id}\}$. Then G is isomorphic to C₆, S₃ or D₆ by Lemma 4.14. It remains to apply Lemma 4.16 and notice that groups mentioned there fix the point $([1:1:1], [1:1:1]) \in S$.

Lemma 4.19. Let $\varphi: S_1 \dashrightarrow S_2$ be a *G*-birational map of *G*-del Pezzo surfaces of degree 6. Then *G* fixes a point on S_1 if and only if *G* fixes a point on S_2 .

Proof. It is enough to prove the necessity. Let $\varphi : (S_1, G, \iota_1) \dashrightarrow (S_2, G, \iota_2)$ be a *G*-birational map, and assume that $\iota_1(G)$ fixes a point on S_1 . By Lemma 4.18, the group $\iota_1(G)$ does not intersect the torus $\operatorname{Aut}^{\circ}(S_1)$ and hence is isomorphic to C_6 , S_3 or D_6 by Lemma 4.14. But this implies that $\iota_2(G)$ does not intersect the torus $\operatorname{Aut}^{\circ}(S_2)$: otherwise $\iota_2(G)/(\iota_2(G) \cap \operatorname{Aut}^{\circ}(S_2))$ is isomorphic to $\operatorname{C}_2, \operatorname{C}_3, \operatorname{C}_2^2, \operatorname{S}_3^{nmin}$ (see Remark 4.15), hence S_2 is not a *G*-del Pezzo surface. \Box

We now discuss (non)linearizability of three particular groups: $\langle \rho \rangle \simeq C_6$, $\langle \rho^2, \sigma \rangle \simeq S_3$ and $\langle \rho, \sigma \rangle \simeq D_6$.

Example 4.20. The cyclic group $G = \langle \rho \rangle \simeq C_6$ is linearizable. Indeed, the Sarkisov *G*-link centred at the *G*-fixed point ([1 : 1 : 1], [1 : 1 : 1]) $\in S$ leads to the *G*-del Pezzo surface $S' \simeq \mathbb{P}^1 \times \mathbb{P}^1$. Since *G* is cyclic, it fixes a point on S' (of course, this can be seen directly as in the next example), hence the stereographic projection from it linearizes the action of *G*.

Example 4.21. The symmetric group $G = \langle \rho^2, \sigma \rangle \simeq S_3$ is linearizable as well. Indeed, after the same *G*-link at the fixed point ([1 : 1 : 1], [1 : 1 : 1]), we arrive to $S' = \mathbb{P}^1 \times \mathbb{P}^1$ acted on by *G*. Recall that $\operatorname{Aut}(S') \simeq (\operatorname{PGL}_2(\mathbf{k}) \times \operatorname{PGL}_2(\mathbf{k})) \rtimes C_2$. Since $\operatorname{Pic}(S')^G = 1$, the group *G* maps non-trivially to C_2 factor of this semi-direct product, i.e. $G = \langle r, s \rangle$, where *r* is an automorphism of order 3 acting fibrewisely on *S'* and *s* is an involution which does not preserve the rulings of *S'*. It will be shown below in Proposition 6.10 that such a group fixes a point on *S'*, and the stereographic projection from this point linearizes the action of *G*.

Example 4.22 (*Iskovskikh's example*). Answering the question of V. Popov, V. Iskovskikh showed in [Isk08] that the group $G \simeq D_6$ generated by the automorphisms ρ and σ is not linearizable. The proof relied on the classical Sarkisov theory. Recently, another proof was given by B. Hassett, A. Kresch and Yu. Tschinkel [HKT21, Section 7.6] by using the Burnside group $Burn_2(G)$ and later by Yu. Tschinkel, K. Yang and Z. Zhang by using the combinatorial Burnside groups [TYZ22]. We refer to the Section 1.3 for more information.

We are ready to summarize the results of this section.

Proposition 4.23. Let S be a G-del Pezzo surface of degree 6. Then G is linearizable if and only if $G \simeq C_6$ or $G \simeq S_3$.

Proof. The sufficiency follows from Examples 4.20 and 4.21 and Lemma 4.16. To prove the necessity, we use the classification of G-links given above. Assume that G is linearizable. Since Bertini and Geiser involutions lead to a G-isomorphic surface, there exists a sequence of G-links $S \dashrightarrow S_1 \dashrightarrow S_1 \dashrightarrow \cdots \dashrightarrow S_n = S'$, where all S_i are G-del Pezzo surfaces of degree 6 and G fixes a point on S' — so that we construct a G-link to $\mathbb{P}^1 \times \mathbb{P}^1$. But then G fixes a point on S by Lemma 4.19. By Lemma 4.18, G does not intersect the torus $\operatorname{Aut}^{\circ}(S)$, and hence G maps isomorphically to a subgroup of D_6 . Now the result follows from Lemma 4.16 and Examples 4.20, 4.21 and 4.22.

Remark 4.24 (Cayley groups and stable linearizability). The (non)linearizability of D_6 and its subgroups was addressed in a different context by N. Lemire, V. Popov and Z. Reichstein in their seminal article [LPR06]. They called a connected linear algebraic group G over a field **K** a Cayley group if it admits a Cayley map, i.e. a G-equivariant birational isomorphism between the group variety G and its Lie algebra Lie(G). Let

$$T = \left\{ A = \begin{bmatrix} a_1 & 0 & 0 \\ 0 & a_2 & 0 \\ 0 & 0 & a_3 \end{bmatrix} : \det A = 1 \right\}, \quad \mathfrak{t} = \left\{ A = \begin{bmatrix} a_1 & 0 & 0 \\ 0 & a_2 & 0 \\ 0 & 0 & a_3 \end{bmatrix} : \operatorname{tr} A = 0 \right\}$$

be a maximal torus of $SL_3(\mathbb{C})$ and its Lie algebra. Both are obviously acted on by the group $W = S_3$ permuting a_1, a_2 and a_3 . Furthermore, T admits a natural compactification $\{x_0y_0z_0 = x_1y_1z_1\} \subset \operatorname{Proj} \mathbb{C}[x_0, x_1] \times \operatorname{Proj} \mathbb{C}[y_0, y_1] \times \operatorname{Proj} \mathbb{C}[z_0, z_1]$, which is a W-del Pezzo surface of degree 6. By Example 4.21, and as was also shown in [LPR06, 9.1], T and \mathfrak{t} are W-birational. By the Corollary of [LPR06, Lemma 3.5], this is equivalent to showing that $SL_3(\mathbf{k})$ is a Cayley group, because W is the Weyl group of $SL_3(\mathbb{C})$.

By the same principle, \mathbb{G}_2 is not Cayley: its Weyl group is $W = D_6$, the maximal torus and its Lie algebra are W-isomorphic to T and \mathfrak{t} , respectively. But the latter two are not W-birational, as showed in Iskovskikh's Example 4.22. Interestingly enough, by [LPR06, Proposition 9.1] $\mathbb{G}_2 \times \mathbb{G}_m^2$ is a Cayley group: the varieties $T \times \mathbb{A}^2$ and $\mathfrak{t} \times \mathbb{A}^2$ are D_6 -birational; in other words, D_6 is stably linearizable. It has recently been shown that one can replace \mathbb{A}^2 with \mathbb{A}^1 here [BvBT23, Proposition 12].

Remark 4.25. According to Proposition 4.23, any finite group G with $G \cap \mathbb{T} \neq \{id\}$ is not linearizable. In [HT23], B. Hassett and Yu. Tschinkel give an example of stably linearizable S₄-action on the sextic del Pezzo surface (here, one has $G \cap \mathbb{T} \simeq V_4$).

5. Finite groups acting on smooth quadric surfaces

Let $S = \mathbb{P}^1 \times \mathbb{P}^1$. Denote by σ the involution which permutes two rulings of S. The connected component $\operatorname{Aut}(S)^\circ$ of the identity is the subgroup of the automorphisms, which preserve both rulings. This subgroup is isomorphic to $\operatorname{PGL}_2(\mathbf{k}) \times \operatorname{PGL}_2(\mathbf{k})$. Then there is a

short exact sequence

$$1 \longrightarrow \mathrm{PGL}_2(\mathbf{k}) \times \mathrm{PGL}_2(\mathbf{k}) \longrightarrow \mathrm{Aut}(S) \xrightarrow{\psi} \langle \sigma \rangle \longrightarrow 1, \qquad (5.1)$$

which splits as the semidirect product $\operatorname{Aut}(S) \simeq (\operatorname{PGL}_2(\mathbf{k}) \times \operatorname{PGL}_2(\mathbf{k})) \rtimes \langle \sigma \rangle$, with $\langle \sigma \rangle \simeq C_2$ acting on $\operatorname{PGL}_2(\mathbf{k}) \times \operatorname{PGL}_2(\mathbf{k})$ by permuting two factors.

Proposition 5.2. Let $S = \mathbb{P}^1 \times \mathbb{P}^1$ and $G \subset \operatorname{Aut}(S)$.

(1) If $\operatorname{rk}\operatorname{Pic}(S)^G \simeq \mathbb{Z}$, then G fits into the short exact sequence

$$1 \longrightarrow H \times_Q H \longrightarrow G \longrightarrow \mathcal{C}_2 \longrightarrow 1, \tag{5.3}$$

where H is a finite subgroup of $PGL_2(\mathbf{k})$.

(2) If
$$\operatorname{rk}\operatorname{Pic}(S)^G \simeq \mathbb{Z}^2$$
, then $G \simeq H_1 \times_Q H_2$, where H_1 , H_2 are finite subgroups of $\operatorname{PGL}_2(\mathbf{k})$.

Proof. Let us restrict the exact sequence (5.1) to G. Clearly, $\psi(G) = \{\text{id}\}$ if and only if G preserves both rulings of S, which is equivalent to $\operatorname{Pic}(S)^G \simeq \mathbb{Z}^2$. If this is the case, then G is a subgroup of $\operatorname{PGL}_2(\mathbf{k}) \times \operatorname{PGL}_2(\mathbf{k})$ and the claim follows from Goursat's lemma 3.9 and Proposition 3.2.

Suppose that $\psi(G) = \langle \sigma \rangle$ and let $K = \operatorname{Ker} \psi|_G$. Let $p_1, p_2 \colon \operatorname{PGL}_2(\mathbf{k})^2 \to \operatorname{PGL}_2(\mathbf{k})$ be the projections onto the first and the second factors, respectively. By Goursat's lemma, we have $K = H_1 \times_Q H_2$, where $H_i = p_i(K)$; in particular, every element of K is of the form $(x, y) \mapsto (h_1 x, h_2 y)$, where (x, y) are local coordinates on $S, h_1 \in H_1, h_2 \in H_2$. The whole group G is the union of two cosets, K and τK , where $\tau(x, y) = (Ay, Bx)$ for some fixed automorphisms $A, B \in \operatorname{Aut}(\mathbb{P}^1)$. Since K is normal in G, then the conjugation by τ sends each automorphism $(h_1, h_2) \in K$ onto (Ah_2A^{-1}, Bh_1B^{-1}) . Therefore, H_1 is isomorphic to H_2 via the map $h_1 \mapsto Bh_1B^{-1}$. Indeed, it is clearly an injective homomorphism. To check surjectivity, we use that $\tau^2 = (AB, BA) \in K$, i.e. $AB \in H_1$. So, for any $h_2 \in H_2$ we have that $Ah_2A^{-1} \in H_1$ and hence $(AB)^{-1}Ah_2A^{-1}AB = B^{-1}h_2B \in H_1$ is sent to h_2 .

Notation 5.4. Everywhere below, the matrices R_n , A, B, C, D, E and F are defined as in Section 3.1, and I stands for the identity matrix. The elements of $Aut(S) \setminus Aut(S)^{\circ}$, i.e. those which do not preserve the rulings of S and which act as $(x, y) \mapsto (My, Nx)$ with $M, N \in PGL_2(\mathbf{k})$, will be denoted (M, N, σ) . Otherwise, we will write (M, N, id).

Every fibre product $G = H_1 \times_Q H_2$, where H_1, H_2 are subgroups of $\mathrm{PGL}_2(\mathbf{k})$, obviously acts on $S = \mathbb{P}^1 \times \mathbb{P}^1$ and forces $\mathrm{Pic}(S)^G \simeq \mathbb{Z}^2$. By contrast, not every group G fitting the exact sequence of the form (5.3) with $H \subset \mathrm{PGL}_2(\mathbf{k})$ actually embeds into $\mathrm{Aut}(S)$, as the following example shows.

Example 5.5. The group $G \simeq C_2^5$ obviously fits the exact sequence (5.3) with $H \simeq V_4$ and $Q = {id}$. However, it cannot be embedded⁴ in Aut($\mathbb{P}^1 \times \mathbb{P}^1$). Indeed, by Lemma 5.7

⁴In fact, this group cannot be embedded even into the Cremona group $Cr_2(\mathbf{k})$, see [Bea07].

below, the normalizer of $H = \langle A, B \rangle$ in $PGL_2(\mathbf{k})$ is $\langle A, B \rangle \rtimes \langle C, D \rangle \simeq V_4 \rtimes S_3 \simeq S_4$. Let $g = (M, N, \sigma) \in G$ be an element such that $\psi(g) = \sigma$. Then G is generated by $H \times H$ and g. Thus, by multiplying g by an element of $H \times H$, we may assume that $g = (T_1, T_2, \sigma)$, where $T_1, T_2 \in \langle C, D \rangle$. But $g^2 = (T_1T_2, T_2T_1, id) \in H \times H$, which forces $T_1 = T_2 \in \{I, D, CD, DC\}$, or $(T_1, T_2) \in \{(C, C^2), (C^2, C)\}$. In all these cases, the group G is isomorphic to $V_4 \wr C_2$.

So, our next goal is to fill the existing gap in the literature: we characterize completely those finite groups which admit a faithful action on $S = \mathbb{P}^1 \times \mathbb{P}^1$.

Theorem 5.6. Let $G \subset \operatorname{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$ be a finite subgroup. Then G is conjugate to a subgroup of one of the following groups:

GAP	Order	Isomorphism class	Generators
No id	48n	$D_n \times S_4$	$(\mathbf{R}_n, \mathbf{I}, \mathrm{id}), (\mathbf{B}, \mathbf{I}, \mathrm{id}), (\mathbf{I}, \mathbf{A}, \mathrm{id}), (\mathbf{I}, \mathbf{B}, \mathrm{id}), (\mathbf{I}, \mathbf{C}, \mathrm{id}), (\mathbf{I}, \mathbf{D}, \mathrm{id})$
No id	120n	$D_n \times A_5$	$(\mathbf{R}_n, \mathbf{I}, \mathrm{id}), (\mathbf{B}, \mathbf{I}, \mathrm{id}), (\mathbf{I}, \mathbf{E}, \mathrm{id}), (\mathbf{I}, \mathbf{F}, \mathrm{id})$
[1440, 5848]	1440	$S_4 \times A_5$	$(\mathrm{A},\mathrm{I},\mathrm{id}),(\mathrm{B},\mathrm{I},\mathrm{id}),(\mathrm{C},\mathrm{I},\mathrm{id}),(\mathrm{D},\mathrm{I},\mathrm{id}),(\mathrm{I},\mathrm{E},\mathrm{id}),(\mathrm{I},\mathrm{F},\mathrm{id})$
No id	$8n^2$	$D_n \wr C_2, n \ge 3$	$(\mathbf{R}_n, \mathbf{I}, \mathrm{id}), (\mathbf{B}, \mathbf{I}, \mathrm{id}), (\mathbf{I}, \mathbf{I}, \sigma)$
[1152, 157849]	1152	$S_4\wr C_2$	$(A, I, id), (B, I, id), (C, I, id), (D, I, id), (I, I, \sigma)$
No id	7200	$A_5\wr C_2$	$(E, I, id), (F, I, id), (I, I, \sigma)$

The proof of this theorem will be given at the end of the Section. We will begin with a complete description, in terms of matrix generators, of the subgroups of $\operatorname{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$ that satisfy the exact sequence (5.3) with H isomorphic to A_4 , S_4 , or A_5 .

Lemma 5.7. Let us consider V_4 , A_4 , S_4 and A_5 as subgroups of $PGL_2(\mathbf{k})$. Then their normalizers in $PGL_2(\mathbf{k})$ are the following:

- (1) $N_{PGL_2(\mathbf{k})}(V_4) \simeq S_4;$
- (2) $N_{PGL_2(\mathbf{k})}(A_4) \simeq S_4;$
- (3) $N_{PGL_2(\mathbf{k})}(S_4) \simeq S_4;$
- (4) $N_{PGL_2(\mathbf{k})}(A_5) \simeq A_5.$

Proof. Recall that all finite subgroups of $PGL_2(\mathbf{k})$ are unique up to conjugation. Since the group A_4 is normal in S_4 , then its normalizer contains S_4 . Similarly, S_4 contains a normal copy of V_4 , hence the normalizer of V_4 in $PGL_2(\mathbf{k})$ contains S_4 . Further, the normalizer of S_4 contains S_4 , while the normalizer of A_5 contains A_5 . Notice that the normalizer of a finite group in $PGL_2(\mathbf{k})$ is an algebraic group, hence we can use their classification provided by Theorem 3.3. In each case, the normalizer is clearly not the whole $PGL_2(\mathbf{k})$, and not the groups of type (3) or (4), as those are metabelian by Remark 3.4 and hence cannot contain a copy of S_4 or A_5 . The claim follows.

Next, we show how to conjugate fibred products $H \times_Q H$ to some "standard forms" in Aut(S). Recall that Aut(A₅) \simeq S₅ and one has Aut(A₅)/Inn(A₅) \simeq C₂. Any outer automorphism of A₅ is a conjugation by an odd permutation in S₅.

Proposition 5.8. Let H be one of the groups A_4 , S_4 or A_5 .

(1) Suppose that $H \simeq A_4$ or $H \simeq S_4$. Then every quotient Q = H/K of H defines uniquely the subgroup K and the fibre product $H \times_Q H$, which is conjugate in Aut(S) to the group

$$H \times_Q H = \{(h_1, h_2) \in H \times H : \overline{h}_1 = \overline{h}_2\} = \{(h, hk) : h \in H, k \in K\}.$$
 (5.9)

In particular, any action of H on S is conjugate to the diagonal one.

(2) Suppose that H ≃ A₅ and fix an outer automorphism ξ ∈ Aut(A₅) \ Inn(A₅). Then any fibre product H ×_Q H is conjugate in Aut(S) to one of the following groups:
(a) {(h₁, h₂) ∈ A₅ × A₅};

- (b) $\{(h,h): h \in A_5\}$, *i.e.* A_5 acts diagonally on S;
- (c) $\{(h,\xi(h)): h \in A_5\}.$

Proof. Let $K \subseteq H$ be a normal subgroup. The classification of normal subgroups of H implies that the isomorphism type of the quotient Q = H/K determines K uniquely. Vice versa, the isomorphism type of K determines the quotient Q. Therefore, by Goursat's Lemma 3.9, the fibred product $H \times_Q H$ is uniquely determined by H, Q and an automorphism $\delta \in \text{Aut}(Q)$, and in this case

$$H \times_Q H = \{(h_1, h_2) \in H \times H \colon \delta(\overline{h}_1) = \overline{h}_2\}$$

(1) Let us show that one can always obtain $\delta = \text{id}$ by conjugating this group in Aut(S). We may assume that $Q \neq \{\text{id}\}$. Then direct computations show that one has $H = K \rtimes Q$ for some complement Q to K in H; we then identify the quotient Q with a subgroup of Hand can write $H = q_1 K \sqcup \ldots \sqcup q_n K$, where $Q = \{q_1, \ldots, q_n\} \subset H$. More precisely, one of the following holds:

(i) $H = A_4$, $K = \{id\}$, $Q = A_4$ and $Aut(Q) \simeq S_4$. (ii) $H = A_4$, $K = V_4$, $Q = \langle (123) \rangle \simeq C_3$ and $Aut(Q) \simeq C_2$. (iii) $H = S_4$, $K = \{id\}$, $Q = S_4$ and $Aut(Q) = Inn(Q) \simeq S_4$. (iv) $H = S_4$, $K = V_4$, $Q = \langle (123), (12) \rangle \simeq S_3$ and $Aut(S_3) \simeq Inn(S_3) \simeq S_3$. (v) $H = S_4$, $K = A_4$, $Q = \langle (12) \rangle \simeq C_2$.

In particular we see that any automorphism $\delta \in \operatorname{Aut}(Q)$ is a conjugation $g \mapsto hgh^{-1}$ by an element $h \in S_4$ (note that in the case (ii) the only non-trivial automorphism $(123) \mapsto (123)^2 = (132)$ is the conjugation by (12)). Since every such h corresponds to an automorphism $\alpha_h \in \operatorname{Aut}(\mathbb{P}^1)$, the automorphism $(h_1, h_2) \mapsto (h_1, \alpha_h^{-1}h_2\alpha_h)$ conjugates $H \times_Q H$ to the fibre product with $\delta = \operatorname{id}$.

(2) Since A_5 is simple, then either $Q = \{id\}$ or $Q = A_5$. In the first case, we get the group (a). In the second case one has $H \times_Q H = \{(h, \delta(h)) \in A_5 \times A_5\}$ for $\delta \in Aut(A_5)$. If $\delta \in Inn(A_5)$, the same argument as in (1) shows that one can conjugate $H \times_Q H$ to the group (b). Otherwise $\delta = \gamma \circ \xi$ for some $\gamma \in Inn(A_5)$. Since γ corresponds to the conjugation by an automorphism of \mathbb{P}^1 , we can conjugate the whole group to (c).

Proposition 5.10. Assume that $H \simeq A_4$. Then, up to conjugation in Aut(S), there are the following cases for G, and only them.

GAP	Order	Isomorphism class	Generators
[24, 13]	24	$A_4 \times C_2$	$(\mathbf{A},\mathbf{A},\mathrm{id}),(\mathbf{B},\mathbf{B},\mathrm{id}),(\mathbf{C},\mathbf{C},\mathrm{id}),(\mathbf{I},\mathbf{I},\sigma)$
[24, 12]	24	S_4	$(\mathbf{A},\mathbf{A},\mathrm{id}),(\mathbf{B},\mathbf{B},\mathrm{id}),(\mathbf{C},\mathbf{C},\mathrm{id}),(\mathbf{D},\mathbf{D},\sigma)$
[96, 70]	96	$C_2^4 \rtimes C_6$	$(A, I, id), (B, I, id), (C, C, id), (I, I, \sigma)$
[96, 70]	96	$C_2^4 \rtimes C_6$	$(\mathbf{A},\mathbf{I},\mathrm{id}),(\mathbf{B},\mathbf{I},\mathrm{id}),(\mathbf{C},\mathbf{C},\mathrm{id}),(\mathbf{I},\mathbf{C},\sigma)$
[96, 227]	96	$C_2^2 \rtimes S_4$	$(\mathbf{A},\mathbf{I},\mathrm{id}),(\mathbf{B},\mathbf{I},\mathrm{id}),(\mathbf{C},\mathbf{C},\mathrm{id}),(\mathbf{D},\mathbf{D},\sigma)$
[288, 1025]	288	$A_4\wr C_2$	$(\mathbf{A},\mathbf{I},\mathrm{id}),(\mathbf{B},\mathbf{I},\mathrm{id}),(\mathbf{C},\mathbf{I},\mathrm{id}),(\mathbf{I},\mathbf{I},\sigma)$
[288, 1025]	288	$A_4\wr C_2$	$(\mathbf{A},\mathbf{I},\mathrm{id}),(\mathbf{B},\mathbf{I},\mathrm{id}),(\mathbf{C},\mathbf{I},\mathrm{id}),(\mathbf{D},\mathbf{D},\sigma)$

Proof. Any subgroup of $PGL_2(\mathbf{k})$ isomorphic to A_4 is conjugate to the group H generated by the matrices A, B, and C, see Section 3.1. We then apply Proposition 5.8. Note that $K = V_4 \triangleleft A_4$ is generated by A and B. Then, because of (5.9), we have one of the following possibilities for $H \times_Q H$:

- (1) $K = {id}: \langle (A, A), (B, B), (C, C) \rangle;$
- (2) $K = V_4$: $\langle (A, I), (B, I), (I, A), (I, B), (C, C) \rangle$
- (3) $K = A_4$: $\langle ; (A, I), (B, I), (C, I), (I, A), (I, B), (I, C) \rangle$.

It remains to identify an element $g \in G$ such that $\psi(g) = \sigma$. It is of the form $(M, N, \sigma) \in PGL_2(\mathbf{k})^2 \rtimes C_2$, and normalizes $H \times_Q H$. Since $(M, N, \sigma)^{-1} = (N^{-1}, M^{-1}, \sigma)$, then for any $P \in H$ we get that

$$(N^{-1}, M^{-1}, \sigma)(P, P, id)(M, N, \sigma) = (N^{-1}PN, M^{-1}PM, id)$$
 (5.11)

is an element of $H \times_Q H \subset H \times H$, which implies that M and N normalize H in PGL₂(**k**). By Lemma 5.7, the normalizer of H in PGL₂(**k**) is the group isomorphic to S₄ which contains H. Consider three cases.

Suppose that $H \times_Q H$ is of the form (1) among the three above possibilities. If $M \in H$, then, up to a multiplication by an element of $H \times_Q H$, we may assume that $g = (I, N, \sigma)$. Therefore, (5.11) implies $N^{-1}PN = P$ for all $P \in H$. But the centre of A_4 is trivial, hence N = I. If $M \notin H$, then (5.11) implies that $N^{-1}PN = M^{-1}PM$ for all $P \in H$, hence $MN^{-1} \in C_{S_4}(A_4) = \{id\}$, and we conclude that M = N, $g = (M, M, \sigma)$. Note that M = TD for $T \in H$ and D the matrix introduced in Section 3.1 (whose coset generates S_4/A_4). Multiplying further g by (T^{-1}, T^{-1}, id) , we achieve $g = (D, D\sigma)$.

Now suppose that $H \times_Q H$ is of the form (2). As before, if $M \in H$, then we may assume M = I. The condition (5.11) gives that $(N^{-1}PN, P)$ is an element of $H \times_Q H$ for any $P \in H$, hence $[N, P] \in K$ for all $P \in H$. This implies $N \in A_4$, hence N = T, N = TC or $N = TC^2$, where $T \in V_4$. Identifying $\langle (I, A), (I, B) \rangle$ with V_4 , we may further multiply g by an element of this group to get $g = (I, I, \sigma), g = (I, C, \sigma)$ or $g = (I, C^2, \sigma)$, respectively. But we can exclude $g = (I, C^2, \sigma)$, because is the same as $g = (I, C, \sigma)$, up to conjugation by (D, D, id), which normalizes $H \times_Q H$.

Now, if $M \notin H$, then (5.11) implies that $(N^{-1}PN, M^{-1}PM, id)$ is an element of $H \times_Q H$ for all $P \in H$, which is equivalent to $N^{-1}P^{-1}NM^{-1}PM$ being an element of K for all $P \in H$. But this holds if and only if $[P, MN^{-1}] \in NKN^{-1} = K$ for all $P \in H$. As above, we conclude that $MN^{-1} \in A_4$ and hence $g = (M, TM, \sigma)$, where $T \in A_4$. As above, we can multiply g by an element of $\langle (A, I), (B, I), (I, A), (I, B) \rangle$ to get $g = (M, M, \sigma), g = (M, CM, \sigma)$ or $g = (M, C^2M, \sigma)$. Since M = DT' for some $T' \in H$, then with a further multiplication by $(T'^{-1}, T'^{-1}, id) \in H \times_Q H$, H, we achive $g \in \{(D, D, \sigma), (D, CD, \sigma), (D, C^2D, \sigma)\}$. But we can exclude $g = (D, CD, \sigma)$ and $g = (D, C^2D, \sigma)$, because the squares of those elements are respectively (C^2, C, id) and (C, C^2, id) , which are not in $H \times_Q H = \langle (A, I), (B, I), (I, A), (I, B), (C, C) \rangle$.

Finally, assume that $H \times_Q H$ is of type (3). Note that $g^2 = (MN, NM, id) \in H \times H$, therefore $N = M^{-1}T$, where $T \in A_4$. Multiplying g by (I, T^{-1}, id) , we may assume $g = (M, M^{-1}, \sigma)$. Finally, since M = T' or M = T'D for $T' \in A_4$, multiplication by an element of $H \times H$ gives $g = (I, I, \sigma)$ or $g = (D, D, \sigma)$, respectively. \Box

Remark 5.12. Some generators of G, obtained during the proofs of Propositions 5.10, 5.13, may be redundant. This is taken into account in the *Generators* column of the corresponding tables. For example, as soon as G contains (I, I, σ) , the group G also contains $(I, I, \sigma)(M, N, id)(I, I, \sigma) = (N, M, id)$ for each $(M, N, id) \in G$. Similarly, suppose that we are in case (2) of the above proof, and $G \simeq C_2^2 \rtimes S_4$ is generated by (A, I, id), (B, I, id), (I, A, id), (I, B, id), (C, C, id) and (D, D, σ) . The relations in S_4 then yield (I, B, id) = (I, DBD, id) = $(D, D, \sigma)^{-1}(B, I, id)(D, D, \sigma)$, and $(I, A, id) = [(D, D, \sigma)^{-1}(A, I, id)(D, D, \sigma)] \cdot (I, B, id)^{-1}$. We leave other cases to the reader.

Proposition 5.13. Assume that $H \simeq S_4$. Then, up to conjugation in Aut(S), there are the following cases for G, and only them.

GAP	Order	Isomorphism class	Generators
[48, 48]	48	$S_4 \times C_2$	$(\mathbf{A},\mathbf{A},\mathrm{id}),(\mathbf{B},\mathbf{B},\mathrm{id}),(\mathbf{C},\mathbf{C},\mathrm{id}),(\mathbf{D},\mathbf{D},\mathrm{id}),(\mathbf{I},\mathbf{I},\sigma)$
[192, 955]	192	$C_2^4 \rtimes D_6$	$(\mathbf{A},\mathbf{I},\mathrm{id}),(\mathbf{B},\mathbf{I},\mathrm{id}),(\mathbf{C},\mathbf{C},\mathrm{id}),(\mathbf{D},\mathbf{D},\mathrm{id}),(\mathbf{I},\mathbf{I},\sigma)$
[576, 8654]	576	$\mathbf{A}_4^2 \rtimes \mathbf{C}_2^2$	$(\mathbf{A},\mathbf{I},\mathrm{id}),(\mathbf{B},\mathbf{I},\mathrm{id}),(\mathbf{C},\mathbf{I},\mathrm{id}),(\mathbf{D},\mathbf{D},\mathrm{id}),(\mathbf{I},\mathbf{I},\sigma)$

[576, 8652]	576	$A_4^2 \rtimes C_4$	$(A, I, id), (B, I, id), (C, I, id), (D, D, id), (I, D, \sigma)$
[1152, 157849]	1152	$S_4\wr C_2$	$(\mathbf{A},\mathbf{I},\mathrm{id}),(\mathbf{B},\mathbf{I},\mathrm{id}),(\mathbf{C},\mathbf{I},\mathrm{id}),(\mathbf{D},\mathbf{I},\mathrm{id}),(\mathbf{I},\mathbf{I},\sigma)$

Proof. First, we apply Proposition 5.8. Proper normal subgroups of S_4 are A_4 and V_4 , hence the only possibilities for $H \times_Q H$ are the following:

- (1) $K = \{ id \}$: $\langle (A, A), (B, B), (C, C), (D, D) \rangle$;
- (2) $K = V_4$: $\langle (A, I), (B, I), (I, A), (I, B), (C, C), (D, D) \rangle$;
- (3) $K = A_4$: $\langle (A, I), (B, I), (C, I), (I, A), (I, B), (I, C), (D, D) \rangle;$
- (4) $K = S_4$: $\langle (A, I), (B, I), (C, I), (D, I), (I, A), (I, B), (I, C), (I, D) \rangle$.

We now look for $g = (M, N, \sigma)$ such that G is generated by $H \times_Q H$ and h, i.e. $\psi(g) = \sigma$. By Lemma 5.7, the normalizer of H in PGL₂(**k**) is H itself. Therefore, $M, N \in H$. Up to a multiplication by an element of $H \times_Q H$, we may assume that M = I and hence $g = (I, N, \sigma)$. We again argue case by case.

If $H \times_Q H$ is of type (1), then the normalization condition (5.11) implies that $N \in Z(S_4) = \{id\}$. If $H \times_Q H$ is of type (2), then $[N, P] \in K$ for all $P \in H$, which implies $N \in V_4$. Multiplying g by an element of $\langle (I, A), (I, B) \rangle \simeq V_4$, we get $g = (I, I, \sigma)$. In the case (3), recall that the quotient of S_4 by $\langle A, B, C \rangle \simeq A_4$ is generated by the coset of D. Therefore, multiplying g by an element of $\langle (I, A), (I, B), (I, C) \rangle$, we achieve $g = (I, I, \sigma)$ or $g = (I, D, \sigma)$. Finally, in the case (4), we can always get $g = (I, I, \sigma)$.

Proposition 5.14. Assume that $H \simeq A_5$. Then, up to conjugation in Aut(S), there are the following cases for G, and only them.

GAP	Order	Isomorphism class	Generators
[120, 35]	120	$A_5 \times C_2$	$(\mathbf{E},\mathbf{E},\mathrm{id}),(\mathbf{F},\mathbf{F},\mathrm{id}),(\mathbf{I},\mathbf{I},\sigma)$
[120, 34]	120	S_5	$(\mathbf{E}, \xi(E), \mathrm{id}), (\mathbf{I}, \mathbf{I}, \sigma)$
No id	7200	$A_5\wr C_2$	$(I, E, id), (I, F, id), (E, I, id), (F, I, id), (I, I, \sigma)$

Proof. We again let $g = (M, N, \sigma) \in G$ be an element such that $\psi(g) = \sigma$. Since g normalizes $H \times_Q H$, we get that M and N normalize H in PGL₂(**k**), hence M, N \in H by Lemma 5.7. We now apply Proposition 5.8(2). If $H \times_Q H$ is of type (a), then one can multiply g by an element of $H \times_Q H$ to get $g = (I, I, \sigma)$. In the case (b), the normalization condition (5.11) implies $MN^{-1} \in C_{A_5}(A_5) = id$, hence M = N and we again can replace g = (M, M, id) by (I, I, σ) . Finally, assume that we are in case (c), i.e. $H \times_Q H = \{(P, \xi(P)) \colon P \in H)\}$. Multiplying g by $(M^{-1}, \xi(M^{-1}), id)$, we may assume M = I. Since for each $P \in H$, one has

$$(\mathbf{I}, \mathbf{N}, \sigma)^{-1}(\mathbf{P}, \xi(\mathbf{P}), \mathrm{id})(\mathbf{I}, \mathbf{N}, \sigma) = (\mathbf{N}^{-1}\xi(\mathbf{P})\mathbf{N}^{-1}, \mathbf{P}, \mathrm{id}) \in H \times_Q H_{\xi}$$

we get $P = \xi(N^{-1}\xi(P)N)$. Since ξ is of order 2, then we get that $N \in Z(A_5) = {id}$. \Box

Proof of Theorem 5.6. If $\operatorname{Pic}(S)^G \simeq \mathbb{Z}^2$, then G is a subgroup of $H_1 \times H_2$, where H_1 and H_2 are finite subgroups of $\operatorname{PGL}_2(\mathbf{k})$ acting fibrewisely on $\mathbb{P}^1 \times \mathbb{P}^1$. Clearly, we can consider the direct products of cyclic and dihedral groups as subgroups of a group of type $D_n \wr C_2$ from the Theorem, for a suitable n. Furthermore, $C_n \times A_4$ embeds into $D_n \times S_4$, while $C_n \times A_5$ embeds into $D_n \times A_5$.

Now suppose that $\operatorname{Pic}(S)^G \simeq \mathbb{Z}$ and hence G fits the exact sequence (5.3). If H is isomorphic to A_4, S_4 , or A_5 , then the result follows from Propositions 5.10, 5.13 and 5.14. Let $g = (M, N, \sigma) \in G$ be an element such that $\psi(g) = \sigma$. Then G is generated by $H \times_Q H$ and g. In particular, G is a subgroup of $\widehat{G} = \langle H \times H, g \rangle$.

Assume that H is cyclic. Then, up to conjugation in $\mathrm{PGL}_2(\mathbf{k})$, the group H is generated by R_n . Since g normalizes $H \times_Q H$, we deduce that M and N are diagonal or anti-diagonal. On the other hand, $g^2 \in H \times_Q H$, i.e. the matrices MN and NM are diagonal, hence either M and N are both diagonal, or both anti-diagonal. Let (u, v) be the affine coordinates on $\mathbb{P}^1 \times \mathbb{P}^1$, and let σ_0 be the automorphism $(u, v) \mapsto (v, u)$, i.e. $(\mathrm{I}, \mathrm{I}, \sigma)$. Then g is either of the form $g: (u, v) \mapsto (av, bu)$, or of the form $g: (u, v) \mapsto (av^{-1}, bu^{-1})$ for some $a, b \in \mathbf{k}^*$.

In the first case, the automorphism $\alpha : (u, v) \mapsto (u, a^{-1}v)$ commutes with $H \times H$ and conjugates g to $g' : (u, v) \mapsto (v, cu)$, where c = ab. The group $\alpha^{-1}\widehat{G}\alpha = \langle H \times H, g' \rangle$ contains an isomorphic copy of G. Since the automorphism $g'^2 : (u, v) \mapsto (cu, cv)$ belongs to $H \times H$, we get $c = \omega_n^k$ for some $k \in \mathbb{Z}$. Multiplying g' by an automorphism $(u, v) \mapsto (u, c^{-1}v)$, which belongs to $H \times H$, we find that $\alpha^{-1}\widehat{G}\alpha$ is generated by $H \times H$ and σ_0 ; in particular, it is isomorphic to $C_n \wr C_2$.

In the second case, the automorphism $\beta : (u, v) \mapsto (u, v^{-1})$ normalizes $H \times H$ and conjugates g to $g' : (u, v) \mapsto (av, b^{-1}u)$. Then $\beta^{-1}\widehat{G}\beta = \langle H \times H, g' \rangle$ contains a copy of G, and one repeats the argument from the previous case.

Assume that $H \simeq D_n$. Then H is conjugated in $PGL_2(\mathbf{k})$ to the group generated by \mathbb{R}_n and B. We may assume $n \ge 3$, since for $H \simeq V_4$ the group G is conjugate to a subgroup of $S_4 \wr \mathbb{C}_2$. Let $T \in H$ be any element, then there is $\mathbb{P} \in H$ so that $(T, \mathbb{P}, \mathrm{id}) \in H \times_Q H$. One has

$$(N^{-1}, M^{-1}, \sigma)(T, P, id)(M, N, \sigma) = (N^{-1}PN, M^{-1}TM, id) \in H \times H,$$
 (5.15)

and hence M normalizes H in $PGL_2(\mathbf{k})$. Now $C = \langle \mathbf{R}_n \rangle$ is a characteristic subgroup of H for $n \geq 3$, hence it is invariant under the conjugation by M, and therefore M is either diagonal or anti-diagonal. The same obviously holds for N. Now, by multiplying g by an element of $H \times H$, we may assume that M and N are either both diagonal, or both anti-diagonal. We finish the proof by applying the same process as in the cyclic case.

Remark 5.16. Our computations exhibit many non-conjugate embeddings of various isomorphic groups into $\operatorname{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$, e.g. of some semi-direct products in Propositions 5.10, 5.13.

6. LINEARIZATION OF FINITE GROUPS ACTING ON HIRZEBRUCH SURFACES

In the final section we study linearization of finite groups acting on Hirzebruch surfaces \mathbb{F}_n with $n \ge 0$. This section is divided into three parts. In the first part, we study Hirzebruch surfaces \mathbb{F}_n with $n \ge 1$. In the second and third parts, we investigate the linearization of groups acting on the quadric surface $\mathbb{F}_0 \simeq \mathbb{P}^1 \times \mathbb{P}^1$, according to the *G*-invariant Picard rank.

6.1. *G*-Hirzebruch surfaces. The goal of this section is to prove the following linearization criterion for Hirzebruch surfaces \mathbb{F}_n with $n \ge 1$.

Theorem 6.1. Let $n \ge 1$ be an integer and $G \subset \operatorname{Aut}(\mathbb{F}_n)$ be a finite group acting on \mathbb{F}_n , such that $\pi \colon \mathbb{F}_n \to \mathbb{P}^1$ is a *G*-conic bundle. Denote by \widehat{G} the image of *G* in the automorphism group of the base $\operatorname{Aut}(\mathbb{P}^1) \simeq \operatorname{PGL}_2(\mathbf{k})$. Then *G* is linearizable if and only if one of the following holds:

- (1) n is odd;
- (2) \widehat{G} is cyclic;
- (3) \widehat{G} is isomorphic to D_m with $m \in \mathbb{Z}$ odd.

Before proving this, we need some easy lemmas. First, recall the following classical definition.

Definition 6.2. Let $n \ge 0$ be a non-negative integer.

(1) An elementary transformation of the Hirzebruch surface \mathbb{F}_n is the following birational transformation. Let $\varphi: Y \to \mathbb{F}_n$ be the blow-up of a point p on a fibre F, \widetilde{F} be the strict transform of $F, \widetilde{\Sigma}_n$ be the strict transform of the (-n)-section $\Sigma_n \subset \mathbb{F}_n$ and Ebe the exceptional divisor. We have $(\widetilde{F})^2 = (\varphi^*F - E)^2 = F^2 - 1 = -1$. Then there is a morphism $\psi: Y \to Z$ blowing down \widetilde{F} . If $p \notin \Sigma_n$, then $\widetilde{\Sigma}_n^2 = \Sigma_n^2 = -n$ and $\widetilde{\Sigma}_n$ intersects \widetilde{F} transversely in exactly one point. Thus $\psi(\widetilde{\Sigma}_n)^2 = -n+1$ and $Z \simeq \mathbb{F}_{n-1}$. If $p \in \Sigma_n$, then $\widetilde{\Sigma}_n^2 = \Sigma_n^2 - 1 = -n-1$, $\widetilde{\Sigma}_n \cap \widetilde{F} = \emptyset$, so $\psi(\widetilde{\Sigma}_n)^2 = -n-1$ and $Z \simeq \mathbb{F}_{n+1}$. Therefore, one has the following diagram for an elementary transformation of \mathbb{F}_n .

$$\begin{array}{c}
 & Y \\
 & \psi \\
 & F_n - - - - - - \rightarrow Z = F_{n+1} \text{ or } F_{n-1} \\
 & \downarrow \\
 & P^1 = P^1
\end{array}$$
(6.3)

(2) Similarly, we define a *G*-elementary transformation $\mathbb{F}_n \dashrightarrow \mathbb{F}_m$, where *G* is a finite group acting on \mathbb{F}_n . In the diagram (6.3), the map φ is the blow-up of a *G*-orbit of length ℓ , and ψ is the contraction of the strict transforms of the fibres through the

blown-up points. We assume that no two points of the orbit lie in the same fibre; in what follows, we refer to this condition as to "conic bundle general position". If the blown-up orbit belongs to Σ_n , then $m = n + \ell$.

(3) More generally, given a G-conic bundle π: S → P¹, we define a G-elementary transformation (or a link of type II between G-conic bundles) χ: S --→ S', where π': S' → P¹ is another G-conic bundle, as the blow-up of a G-orbit on S, followed by the contraction of the strict transforms of the fibres through the blown-up points. Once again, we assume that no two points of the orbit lie in the same fibre. Note that χ does not change the number of singular fibres, i.e. one has K²_S = K²_{S'}.

Lemma 6.4. Any subgroup $G \subset \operatorname{Aut}(\mathbb{F}_n)$ preserves a curve on \mathbb{F}_n which has self-intersection n and is disjoint from Σ_n .

Proof. Recall that $S = \mathbb{F}_n$ has the unique section $\Sigma_n \subset \mathbb{F}_n$ of self-intersection -n. Let F be a fibre of S. The complete linear system $|\Sigma_n + nF|$ induces the birational morphism $\phi \colon S \to S' \subset \mathbb{P}^{n+1}$, which is the contraction of Σ_n onto the vertex of the cone S'. As Σ_n is preserved by G, this contraction is G-equivariant. Writing $\mathbb{P}^{n+1} = \mathbb{P}(L \oplus V)$ with L being 1-dimensional vector space corresponding to the unique singular (and hence G-fixed) point of S', we see that $\mathbb{P}(V)$ is a G-invariant hyperplane. It intersects S' in a G-invariant rational normal curve of degree n. Its preimage on S under ϕ is the required curve.

Now we will analyse G-elementary transformations of Hirzebruch surfaces in greater detail, based on the "arithmetic" of possible G-orbits; a similar idea was used e.g. in [Che14, Lemma B.15].

Lemma 6.5. Let $n \ge 1$ and consider a *G*-conic bundle $\pi \colon \mathbb{F}_n \to \mathbb{P}^1$. One has the following:

- (1) Let ℓ be the length of one of the orbits under the action of \widehat{G} on \mathbb{P}^1 . Then the conic bundle \mathbb{F}_n is G-birational to $\mathbb{F}_{|n+k\ell|}$, for any $k \in \mathbb{Z}$.
- (2) Let ℓ_i , i = 1, ..., r, be the lengths of orbits under the action of \widehat{G} on \mathbb{P}^1 , and let $d = \gcd\{\ell_i\}$. Then \mathbb{F}_n is birational to $\mathbb{F}_{|n+kd|}$ for any $k \in \mathbb{Z}$.

Proof. (1) The blow-up of \mathbb{F}_n at a *G*-orbit of length ℓ contained in Σ_n and contraction of the proper transforms of the fibres gives a *G*-birational map to $\mathbb{F}_{n+\ell}$. Moreover, the action of *G* on the $-(n + \ell)$ -curve of $\mathbb{F}_{n+\ell}$ is the same as the action of *G* on the (-n)-curve of \mathbb{F}_n . It proves the Lemma for $k \ge 0$. Let *C* be a *G*-invariant *n*-curve, given by Lemma 6.4. Notice that since *C* and Σ_n are both sections of the conic bundle, the action of *G* on *C* is the same as the action of *G* on Σ_n . The blow-up of \mathbb{F}_n at a *G*-orbit of length ℓ contained in *C* and contraction of the proper transforms of the fibres gives a *G*-birational map to $\mathbb{F}_{|n-\ell|}$. Moreover, the action of *G* on the $-|n - \ell|$ -curve of $\mathbb{F}_{|n-\ell|}$ is the same as the action of *G* on the (-n)-curve of \mathbb{F}_n . We proceed by induction. Suppose that constructed a sequence of *G*-elementary transformations $\mathbb{F}_n \to \mathbb{F}_{|n-k\ell|}$, where $k \ge 1$. If $n - k\ell \ge 0$, then an elementary transformation at a G-orbit of ℓ points, lying on the G-invariant n-curve as above, gives a map $\mathbb{F}_{n-k\ell} \longrightarrow \mathbb{F}_{|n-(k+1)\ell|}$. If $n-k\ell < 0$ then we use an elementary transformation at a G-orbit lying on $\Sigma_{|n-k\ell|}$ to get $\mathbb{F}_{|n-k\ell|} = \mathbb{F}_{k\ell-n} \dashrightarrow \mathbb{F}_{k\ell-n+\ell} = \mathbb{F}_{|n-(k+1)\ell|}$.

(2) By Bézout's identity, we have

$$kd = a_1\ell_1 + \ldots + a_s\ell_s - a_{s+1}\ell_{s+1} - \ldots - \ell_r a_r, \tag{6.6}$$

where $0 \leq s \leq r$ and $a_i \geq 0$ for all i = 1, ..., r. If $k \geq 0$, then, by performing G-elementary transformations as in (1), we get a sequence

$$\mathbb{F}_{n} \xrightarrow{\chi_{0}} \mathbb{F}_{n+a_{1}\ell_{1}} \xrightarrow{\chi_{1}} \mathbb{F}_{n+a_{1}\ell_{1}+a_{2}\ell_{2}} \xrightarrow{\chi_{2}} \dots \longrightarrow \mathbb{F}_{n+a_{1}\ell_{1}+\dots+a_{s}\ell_{s}} \xrightarrow{\chi_{s}} \mathbb{F}_{n+a_{1}\ell_{1}+\dots+a_{s}\ell_{s}-a_{s+1}\ell_{s+1}} \xrightarrow{\chi_{s+1}} \dots$$

$$(6.7)$$

which results in \mathbb{F}_{n+kd} . Suppose now that k < 0, and again write kd in the form (6.6). Perform the sequence of transformations (6.7) up to the s-th step. The map χ_s then leads to $\mathbb{F}_{|n+a_1\ell_1+\ldots+a_s\ell_s-a_{s+1}\ell_{s+1}|} = \mathbb{F}_{a_{s+1}\ell_{s+1}-n-a_1\ell_1-\ldots-a_s\ell_s}$. We can proceed by mapping

$$\mathbb{F}_{a_{s+1}\ell_{s+1}-n-a_{1}\ell_{1}-...-a_{s}\ell_{s}} \longrightarrow \mathbb{F}_{a_{s+2}\ell_{s+2}+a_{s+1}\ell_{s+1}-n-a_{1}\ell_{1}-...-a_{s}\ell_{s}} = \mathbb{F}_{|n+a_{1}\ell_{1}+...+a_{s}\ell_{s}-a_{s+1}\ell_{s+1}-a_{s+2}\ell_{s+2}|},$$
and so on until we get $\mathbb{F}_{|n+k,l|}$

and so on, until we get $\mathbb{F}_{|n+kd|}$.

Now, we deduce several corollaries.

Corollary 6.8. If \widehat{G} is cyclic or isomorphic to D_m with m odd, then \mathbb{F}_n is G-birational to \mathbb{F}_k , for any non-negative integer k. In particular, G is linearizable.

Proof. If \widehat{G} is cyclic, then it fixes a point on \mathbb{P}^1 . If \widehat{G} is isomorphic to D_m with m odd, then there is an orbit of length 2 and of odd length by Proposition 3.7. In both cases, by Lemma 6.5 (2), the surface \mathbb{F}_n is G-birational to $\mathbb{F}_{|n+s|}$ for any $s \in \mathbb{Z}$, hence the first claim. In particular, by taking s = 1 - n, we arrive at \mathbb{F}_1 where we can *G*-equivariantly contract the unique (-1)-curve to get \mathbb{P}^2 .

Corollary 6.9. The G-conic bundle \mathbb{F}_n is G-birational to any $\mathbb{F}_{|n+2k|}$, for any $k \in \mathbb{Z}$. So, if n is even, then \mathbb{F}_n is G-birational to the G-conic bundle $\mathbb{F}_0 \simeq \mathbb{P}^1 \times \mathbb{P}^1$. If n is odd, then \mathbb{F}_n is G-birational to \mathbb{F}_1 and hence G is linearizable in this case.

Proof. If \widehat{G} is cyclic, we conclude using Corollary 6.8. Otherwise, Proposition 3.7 implies that there is always a pair (or a triple) of orbits of \widehat{G} having 2 as the greatest common divisor of their lengths. By Lemma 6.5 (2), we conclude that \mathbb{F}_n is G-birational to $\mathbb{F}_{|n+2k|}$, for any $k \in \mathbb{Z}$.

6.2. Quadrics with invariant Picard group of rank 1. We will classify finite linearizable subgroups of Aut(S) according to the isomorphism class of H in the exact sequence (5.3).

Proposition 6.10. If H is cyclic, then G is linearizable.

Proof. We claim that G has a fixed point, so the stereographic projection from it linearizes the action of G. Indeed, the group $H \times_Q H$ is contained in $H \times H$, hence it has exactly 4 fixed points on S. We can choose the coordinates on S so that these points are

$$p_1 = ([1:0], [1:0]), \quad p_2 = ([1:0], [0:1]), \quad p_3 = ([0:1], [1:0]), \quad p_4 = ([0:1], [0:1]).$$

Let $g \in G$ be an element which is mapped to the generator of C_2 in the exact sequence (5.3), i.e. which does not preserve the rulings of S. Then G is generated by g and $H \times_Q H$. Since g normalizes $H \times_Q H$, it permutes the points p_1, p_2, p_3, p_4 . Since g swaps the rulings of S, it fixes p_1 and p_3 , or it fixes p_2 and p_4 , or it permutes p_1, p_2, p_3 and p_4 cyclically. In the first two cases, we are done. The third case is impossible, since g^2 would swap p_i with p_{i+2} , while it belongs to $H \times_Q H$, i.e. fixes all four points.

Remark 6.11. In the language of Sarkisov program, we linearize the action of G as follows. Let $\eta: T \to S = \mathbb{P}^1 \times \mathbb{P}^1$ be the blow-up of a G-fixed point $p \in S$. The surface T is a del Pezzo surface of degree 7 with three exceptional curves F_1, F_2 , and E, where E is the η -exceptional divisor, F_1 and F_2 are the preimages of fibres through p. One has $E \cdot F_1 = E \cdot F_2 = 1$, and $F_1 \cdot F_2 = 0$. Then there is a G-contraction $\eta': T \to \mathbb{P}^2$ of F_1 and F_2 onto a pair of points.

We are going to show that G is not linearizable in any of the other cases.

Observation. We use the Notations of Section 2.2. Every Sarkisov G-link starting from S is either of type I, where η blows up a G-orbit of length 2, or is of type II, and one of the following holds:

- (1) $S \simeq S', d = d' = 7, \chi$ is a birational Bertini involution;
- (2) $S \simeq S', d = d' = 6, \chi$ is a birational Geiser involution;
- (3) S' is a del Pezzo surface of degree 5, d = 5, d' = 2;
- (4) $S' \simeq \mathbb{P}^1 \times \mathbb{P}^1, d = d' = 4;$
- (5) S' is a del Pezzo surface of degree 6, d = 3, d' = 1;
- (6) $S' \simeq \mathbb{P}^2, d = 1, d' = 2.$

Birational Geiser and Bertini involutions lead to a *G*-isomorphic surface. By [Yas24, Proposition 4.3], links centred at orbits of length 4 also result in a surface which is *G*-isomorphic to *S*. Therefore, any *G*-link starting from *S* and leading to a non-isomorphic surface must be centred at an orbit of length $d \in \{1, 2, 3, 5\}$.

Proposition 6.12. Let $S = \mathbb{P}^1 \times \mathbb{P}^1$ and $G \subset \operatorname{Aut}(S)$ be a finite group such that $\operatorname{Pic}(S)^G \simeq \mathbb{Z}$. Assume that in the setting of the exact sequence (5.3) the group H is isomorphic to A_4, S_4 or A_5 . Then S is G-birationally rigid. In particular, G is not linearizable.

Proof. Note that an orbit of G is a disjoint union of orbits of $H \times_Q H$. Therefore, by Proposition 3.7 and Lemma 3.12, any orbit of G on S has length $2m \ge 4$. Hence S can admit only G-birational Geiser involutions and links at points of degree 4. As was observed before, they give a G-isomorphic surface.

Proposition 6.13. Let $S = \mathbb{P}^1 \times \mathbb{P}^1$ and $G \subset \operatorname{Aut}(S)$ be a finite group such that $\operatorname{Pic}(S)^G \simeq \mathbb{Z}$. Assume that in the setting of the exact sequence (5.3) the group H is dihedral D_n . Then G is not linearizable.

Proof. We may assume that there is a G-orbit $\Sigma \subset S$ in general position, i.e. its blow-up gives a del Pezzo surface. Then Lemma 3.13 implies that Σ is a disjoint union of at most 2 orbits of $H \times_Q H$. Suppose that $\Sigma = \Sigma_1 \sqcup \Sigma_2$, where $|\Sigma_1| = |\Sigma_2|$. By Proposition 3.7 and Lemma 3.12, these cardinalities are divisible by 2 or n. Then $|\Sigma|$ is an even number ≥ 4 , and therefore S is G-birationally rigid. So, we may assume that Σ is an orbit of $H \times_Q H$. By the same observations, we only need to investigate orbits of size 2, 3 and 5.

Suppose $|\Sigma| = 5$, i.e. $H \simeq D_5$. The link $S \dashrightarrow S'$ centred at Σ gives a *G*-del Pezzo surface S' of degree 5. Since Q can be isomorphic to {id}, C_2 or D_5 , we find that either Gis an extension of C_2 by D_5 , or G contains a copy of C_5^2 by Lemma 3.10. In the latter case, G cannot be embedded into $\operatorname{Aut}(S') \simeq S_5$, and neither $D_5 \times C_2$ can. Hence we may assume that $G \simeq F_5$. But in this case, G cannot be linearized by Proposition 4.4.

If $|\Sigma| = 3$, then $H \simeq S_3$, and the blow-up of Σ gives a del Pezzo surface T of degree 5. Since Q can be isomorphic to {id}, C_2 or S_3 , we find that either $G \simeq S_3 \times C_2$, or G contains a copy of C_3^2 by Lemma 3.10. In the latter case, G does not embed into $\operatorname{Aut}(T) \simeq S_5$, hence we may assume that we are in the former case. Then our Sarkisov link ends up on a G-del Pezzo surface of degree 6. Since $G \simeq S_3 \times C_2$, it cannot be linearized by Proposition 4.23.

Finally, case $|\Sigma| = 2$ corresponds to a link of type I leading to the del Pezzo surface T of degree 6 with a G-conic bundle structure. All Sarkisov G-links starting from a G-conic bundle with 2 singular fibres are either G-elementary transformations $T \dashrightarrow T'$, or links $T' \rightarrow S'$ of type III leading back to a G-del Pezzo surface S' of degree 8, so the composition of such links looks like in Diagram 4.1. In view of Propositions 6.10, 6.12 and the previous cases, it is enough to show that G does not fix a point on S', and hence cannot be further linearized. Let us denote cyclically the six (-1)-curves forming a hexagon on T by E_1, \ldots, E_6 . We may assume that E_1 and E_4 are the exceptional divisors over Σ , so in particular the action of G must swap E_1 and E_4 . They both form sections of the G-conic bundle $\pi: T \to \mathbb{P}^1$. On the other hand, $E_2 + E_3$ and $E_5 + E_6$ are the two singular fibres of π , whose irreducible components are swapped by G. By Theorem 2.5, the G-conic bundle $\pi': T' \to \mathbb{P}^1$ is again a del Pezzo surface of degree 6. Denote by D_1 and D_2 the singular fibres of π' . Besides their irreducible components, the surface T' has two more (-1)-curves L_1 and L_2 , which are disjoint sections of π' , swapped by G. Note that the points $E_2 \cap E_3$ and $E_5 \cap E_6$ are unique G-fixed points on T. Since a sequence of G-elementary transformations $T \dashrightarrow T'$ is an isomorphism away from G-orbit of curves of length > 1, the singular points of D_1 and D_2 are the unique G-fixed points on T'. Thus, the contraction $T' \to S'$ of L_1 and L_2 leaves no G-fixed points on S'.

6.3. Quadrics with invariant Picard group of rank 2. Suppose that $\operatorname{Pic}(S)^G \simeq \mathbb{Z}^2$. We endow $S = \mathbb{P}^1 \times \mathbb{P}^1$ with coordinates $([x_0 : x_1], [y_0 : y_1])$ and fix two projections π_1 and π_2 to $[x_0 : x_1]$ and $[y_0 : y_1]$, respectively. We often use the affine coordinates $x = x_1/x_0$ and $y = y_1/y_0$ on S, so that the points ([1 : 0], [1 : 0]) and ([0 : 1], [0 : 1]) correspond to (0, 0) and (∞, ∞) . We follow Notations 3.6 for the generators of finite subgroups of PGL₂(**k**).

By Proposition 5.2, the group G is the fibred product $H_1 \times_Q H_2$, where H_1 and H_2 are finite subgroups of PGL₂(**k**), acting on S fibrewisely (projections of G induced by π_1 and π_2 , respectively). The crucial observation, which follows from [DI09, Propositions 7.12, 7.13], is that G is linearizable if and only if there is a sequence of Sarkisov G-links

$$S = \mathbb{F}_0 \xrightarrow{\chi_0} S_1 \xrightarrow{\chi_1} S_2 \xrightarrow{\chi_2} S_3 \xrightarrow{\chi_3} \dots \xrightarrow{\chi_{N-1}} S_N = \mathbb{F}_1 \xrightarrow{\chi_N} \mathbb{P}^2,$$

where $S_i \simeq \mathbb{F}_{n_i}$ for all *i*, the links χ_i for $i = 0, \ldots, N - 1$ are either *G*-elementary transformations of Hirzebruch surfaces, or links of type IV (in which case $S_i \simeq S_{i+1} \simeq \mathbb{P}^1 \times \mathbb{P}^1$), and the last link χ_N is necessarily of type III, namely the equivariant blow-down of the unique (-1)-curve on \mathbb{F}_1 to the *G*-fixed point on \mathbb{P}^2 .

The main result of this subsection is the following.

Theorem 6.14. The group $G = H_1 \times_Q H_2$ is linearizable if and only if one of the following holds:

- (1) Both H_1 and H_2 are cyclic;
- (2) H_1 is cyclic, and $H_2 \simeq D_n$, where $n \ge 1$ is odd, or vice versa;
- (3) $H_1 \simeq D_n, H_2 \simeq D_m$, where $n, m \ge 3$ are odd, and G is isomorphic to a dihedral group.

We split the proof into several auxiliary statements.

Lemma 6.15. The group $G = C_n \times_Q C_m$ is linearizable for any $n, m \in \mathbb{Z}_{>0}$.


Proof. We may assume that G is a subgroup of $\langle \mathbf{R}_n \rangle \times \langle \mathbf{R}_m \rangle$, and hence fixes the point ([1:0], [1:0]); the stereographic projection from this point linearizes G.

Lemma 6.16. Let n and m be integers, with n odd. The group $G = D_n \times_Q C_m$ is linearizable.

Proof. The group C_m fixes a point on \mathbb{P}^1 , and hence G acts on the fibre F over this point. Therefore, the image of $H_1 \times_Q H_2$ in $\operatorname{Aut}(F) \simeq \operatorname{PGL}_2(\mathbf{k})$ is isomorphic to D_n with n odd. By Proposition 3.7, there is a G-orbit of size n on F. A G-elementary transformation centred at such orbit leads to the Hirzebruch surface \mathbb{F}_n and we are done by Corollary 6.9.

Lemma 6.17. If H_1 or H_2 is isomorphic to A_4, S_4, A_5 , or D_{2n} , then G is not linearizable.

Proof. We start with the following observation. Consider a G-elementary transformation



centred at a *G*-orbit $p_1, \ldots, p_{2k} \in \mathbb{F}_{2n}$ of length $2k \ge 2$. Let us show that *m* is even. Indeed, assume that *m* is odd. For $i = 1, \ldots, 2k$, let $F_i \subset \mathbb{F}_{2n}$ be the fibres through the blown-up points p_i , let $E_i = \eta^{-1}(p_i) \subset T$ be the exceptional divisors over these points, $\widetilde{F}_i \subset T$ be strict transforms of F_i , and $q_i = \eta'(F_i)$ be the resulting *G*-orbit on \mathbb{F}_m . Suppose that the curve $\Sigma_m \subset \mathbb{F}_m$ contains $t \ge 0$ points among q_i . Its strict transform $\widetilde{\Sigma}_m \subset T$ under η' then satisfies $\widetilde{\Sigma}_m^2 = -m - t$, it intersects exactly *t* curves \widetilde{F}_i on *T*, and 2k - t curves E_i on *T*. The η -images of these 2k - t curves are the blown-up points lying on $C = \eta(\widetilde{\Sigma}_m)$. Thus, $C^2 - (2k - t) = \widetilde{\Sigma}_m^2 = -m - t$, hence $C^2 = 2k - 2t - m$ is odd, which is impossible⁵ on \mathbb{F}_{2n} .

Now let H_1 and H_2 be as in the condition of the Lemma. It is sufficient to show that there is no sequence of *G*-elementary transformations (possibly alternating with links of type IV on \mathbb{F}_0) from $S = \mathbb{F}_0$ to \mathbb{F}_1 . By Proposition 3.7 and Lemma 3.12, the group $G = H_1 \times_Q H_2$ has only orbits of even length on *S*. Thus, the first *G*-elementary transformation brings us to some \mathbb{F}_{2n} . Since the lengths of the orbits is preserved under *G*-fibrewise transformations, a sequence of *G*-elementary transformations can only lead to Hirzebruch surfaces \mathbb{F}_N with *N* even.

It remains to study the linearizability of G when $H_1 \simeq D_n$ and $H_2 \simeq D_m$, with n and m odd. By using Notations 3.6, we can suppose that $D_n = \langle R_n, B \rangle$ and $D_m = \langle R_m, B \rangle$ as subgroups of PGL₂(**k**). Recall that possible fibre products $D_n \times_Q D_m$ were essentially described in Lemma 3.15.

Lemma 6.18. Let n, m be odd positive integers. If $G = D_n \times_Q D_m$ is not isomorphic to a dihedral group, then G is not linearizable.

Proof. If G is linearizable, then there is a sequence of Sarkisov G-links $S_1 = \mathbb{P}^1 \times \mathbb{P}^1 \dashrightarrow S_2 \dashrightarrow \cdots \dashrightarrow S_k = \mathbb{F}_1 \to \mathbb{P}^2$, where the last map is the G-equivariant blow-down of the unique (-1)-curve on \mathbb{F}_1 — a link of type III. The image of this curve is a G-fixed point $p \in \mathbb{P}^2$, and hence G admits a faithful representation in $\mathrm{GL}(T_p\mathbb{P}^2) \simeq \mathrm{GL}_2(\mathbf{k})$. We conclude by Lemma 3.15.

⁵Recall that $\operatorname{Pic}(\mathbb{F}_N)$ is generated by the class of a fibre F and the (-N)-section Σ_N (or a transversal fibre Σ_0 , such that $\Sigma_0 \cdot F = 1$, when N = 0). One has $F^2 = 0$, $F \cdot \Sigma_N = 1$, $\Sigma_N^2 = -N$. So, for a curve $C \sim aF + b\Sigma_N$ one has $C^2 = 2ab - Nb$.

Proposition 6.19 (Linearization via the Euclidean algorithm). Let n and m be odd positive integers. Suppose that the group $G = D_n \times_Q D_m$ is dihedral. Then G is linearizable.

Proof. For convenience, we set $D_1 = C_2$, so that $G = D_n \times_{D_d} D_m$, where $d \ge 1$ divides both n and m. By Lemma 3.10, there is a short exact sequence of groups

$$1 \longrightarrow \langle (\mathbf{R}_n^d, \mathrm{id}) \rangle \times \langle (\mathrm{id}, \mathbf{R}_m^d) \rangle \longrightarrow G \xrightarrow{\varrho} \mathbf{D}_d \longrightarrow 1.$$
(6.20)

Since G is dihedral, then ker ρ must be cyclic, so gcd(n/d, m/d) = 1 and ker ρ is generated by the element $(\mathbb{R}_n^d, \mathbb{R}_m^d)$. By Remark 3.11, the group G is generated by $(\mathbb{R}_n^d, \mathbb{R}_m^d)$ and $(\mathbb{B}, \mathbb{R}_m^v \mathbb{B})$ when d = 1, and by $(\mathbb{R}_n^d, \mathbb{R}_m^d)$, $(\mathbb{R}_n, \mathbb{R}_m^u)$ and $(\mathbb{B}, \mathbb{R}_m^v \mathbb{B})$ when⁶ d > 1; here, u, v are some positive integers. Let k = v/2 if v is even and k = (v + m)/2 if v is odd. By conjugating G in Aut(S) by the automorphism (id, \mathbb{R}_m^k), we may assume v = 0.

Let M = nm. Then

$$(\mathbf{R}_n^d, \mathbf{R}_m^d) = (\mathbf{R}_M^{dm}, \mathbf{R}_M^{dn}), \quad (\mathbf{R}_n, \mathbf{R}_m^u) = (\mathbf{R}_M^m, \mathbf{R}_M^{nu}).$$

Regardless of whether d = 1 or d > 1, the elements of G are all of the form $(\mathbb{R}_M^a, \mathbb{R}_M^b)$ or $(\mathbb{R}_M^a, \mathbb{R}_M^b)$ for some positive integers a, b. Since the latter ones are involutions, we can assume that the characteristic cyclic subgroup of G is generated by an element of the form $(\mathbb{R}_M^a, \mathbb{R}_M^b)$, i.e. by the map $(x, y) \mapsto (\omega_M^a x, \omega_M^b y)$.

Consider the following birational self-map of S and its inverse:

$$\varphi \colon (x,y) \vdash \to (x,x^{-1}y), \quad \varphi^{-1} \colon (x,y) \vdash \to (x,xy).$$

Given a biregular automorphism $g: (x, y) \mapsto (\alpha x, \beta y)$ of S for some $\alpha, \beta \in \mathbf{k}^*$, we have the following commutative diagram

where $\overline{g}: (x, y) \mapsto (\alpha x, \alpha^{-1}\beta y)$ is $\varphi \circ g \circ \varphi^{-1}$. The conjugation of the automorphism (B, B) by φ gives the same action. To sum up, φ is a birational equivalence between $(\mathbb{P}^1 \times \mathbb{P}^1, G)$ and $(\mathbb{P}^1 \times \mathbb{P}^1, \overline{G})$, where \overline{G} is generated by

$$(x,y)\mapsto (\omega_M^a x, \omega_M^{b-a} y), \ (x,y)\mapsto (x^{-1},y^{-1}).$$

The biregular automorphism $\sigma: (x, y) \mapsto (y, x)$ conjugates the map $(x, y) \mapsto (\alpha x, \beta y)$ to $(x, y) \mapsto (\beta x, \alpha y)$ and does not change (B, B). Therefore, up to conjugation by σ , we may assume $0 \leq a \leq b$. We now run the Euclidean algorithm for a and b: first, by iterating the conjugation by φ , we can replace $(\mathbf{R}_M^a, \mathbf{R}_M^b)$ by $(\mathbf{R}_M^a, \mathbf{R}_M^r)$, where r is the remainder of

⁶Note that the second case includes the sub-case n = m = d. As follows from Goursat's lemma, we have $G \simeq \{(g, \varphi(g)) : g \in D_n\}$, where $\varphi \in \operatorname{Aut}(D_n)$. It is well known that in the standard presentation of D_n , its automorphism group is generated by the maps $r \mapsto r^j$, $s \mapsto r^t s$, which agrees with the generators we chose.

division of b by a. By using σ , we replace $(\mathbf{R}_M^a, \mathbf{R}_M^r)$ by $(\mathbf{R}_M^r, \mathbf{R}_M^a)$, perform Euclidean division of a by r, and so on. The algorithm results in the generator $(\mathbf{R}_M^\ell, \mathrm{id})$, where $\ell = \gcd(a, b)$.

Hence, we birationally conjugated G to the dihedral group generated by $(\mathbb{R}_M^\ell, \mathrm{id})$ and (\mathbb{B}, \mathbb{B}) . The fibre over y = 1 is now faithfully acted on by G. Making an elementary transformation at the orbit of size N = |G|/2 in this fibre, we arrive to \mathbb{F}_N and hence can further linearize the action of G.

Proof of Theorem 6.14. Follows from Lemmas 6.15-6.18 and Proposition 6.19.

Appendix: Magma code

The following Magma code determines the GAP ID of finite subgroups of $\operatorname{Aut}(\mathbb{P}^1 \times \mathbb{P}^1) \simeq$ PO(4) defined by their matrix generators. The code is also available on the GitHub page of the first author [Pin24a].

```
K:=AlgebraicClosure(Rationals());
1
2 K4:=CartesianPower(K,4);
3 R<x>:=PolynomialRing(K);
4 i:=Roots(x^2+1)[1,1];
  w5:=Roots(x^{5}-1)[2,1];
5
6
  A := Matrix(K, 2, [1, 0, 0, -1]);
7
8 B:=Matrix(K,2,[0,1,1,0]);
9 C:=Matrix(K,2,[i,-i,1,1]);
10 D:=Matrix(K,2,[1,-i,i,-1]);
11 E:=Matrix(K,2,[w5,0,0,1]);
12 F:=Matrix(K,2,[1,1-w5-w5^-1,1,-1]);
13 I:=Matrix(K,2,[1,0,0,1]);
14
  // The following function takes two ordered sets of 4 points of P3
15
16 // in general position, and returns the unique matrix in PGL(4)
  // which takes the first set to the second one.
17
18
   AutP3:=function(points, images)
19
     points:=[<t:t in points[1]>,<t:t in points[2]>,<t:t in points[3]>,<t:t</pre>
20
      in points[4]>, <t:t in points[5]>];
     images:=[<t:t in images[1]>,<t:t in images[2]>,<t:t in images[3]>,<t:t</pre>
21
      in images[4]>,<t:t in images[5]>];
22
23
     A1<a,b,c,d>:=AffineSpace(K,4);
```

```
X1:=Scheme(A1,[a*points[2][1]+b*points[3][1]+c*points[4][1]+d*points
24
      [5] [1] -points [1] [1], a*points [2] [2] +b*points [3] [2] +c*points [4] [2] +d*
      points [5] [2] -points [1] [2], a*points [2] [3] +b*points [3] [3] +c*points [4] [3] +
      d*points [5] [3] -points [1] [3], a*points [2] [4]+b*points [3] [4]+c*points
      [4][4]+d*points[5][4]-points[1][4]]);
25
     sols1:=RationalPoints(X1);
26
     eltsM1:=[];
27
28
     for i in [1..4] do
29
       eltsM1 := eltsM1 cat ElementToSequence(sols1[1][i]*Vector(K,4,[points[i
30
      +1][1],points[i+1][2],points[i+1][3],points[i+1][4]]));
     end for:
31
       M1:=Transpose(Matrix(K,4,eltsM1));
32
33
     X2:=Scheme(A1,[a*images[2][1]+b*images[3][1]+c*images[4][1]+d*images
34
      [5] [1] - images [1] [1], a* images [2] [2] + b* images [3] [2] + c* images [4] [2] + d*
      images [5] [2] - images [1] [2], a*images [2] [3] + b*images [3] [3] + c*images [4] [3] +
      d*images [5] [3] - images [1] [3], a*images [2] [4] + b*images [3] [4] + c*images
      [4][4]+d*images[5][4]-images[1][4]]);
35
36
     sols2:=RationalPoints(X2);
     eltsM2:=[];
37
38
     for i in [1..4] do
39
       eltsM2 := eltsM2 cat ElementToSequence(sols2[1][i]*Vector(K,4,[images[i
40
      +1][1], images[i+1][2], images[i+1][3], images[i+1][4]]));
     end for;
41
42
     M2:=Transpose(Matrix(K,4,eltsM2));
43
     M := M2 * M1^{-1};
44
45
     return M;
   end function;
46
47
  // The following function takes a triple (M1,M2,s) in Aut(P1xP1),
48
   // and returns the matrix of PGL4 whose restriction on the quadric xt=yz
49
   // gives the corresponding automorphism after the Segre embedding.
50
51
   AutQuadSurf:=function(M1,M2,s)
52
     M1:=Matrix(K,2,ElementToSequence(M1));
53
     M2:=Matrix(K,2,ElementToSequence(M2));
54
     if s eq 0 then
55
```

56	g:=map <k4->K4 x:-><(ElementToSequence(M1*Matrix(K,2,1,[x[1],x[2]]))</k4->
	<pre>cat ElementToSequence(M2*Matrix(K,2,1,[x[3],x[4]])))[1],(</pre>
	<pre>ElementToSequence(M1*Matrix(K,2,1,[x[1],x[2]])) cat ElementToSequence(</pre>
	M2*Matrix(K,2,1,[x[3],x[4]])))[2],(ElementToSequence(M1*Matrix(K,2,1,[x
	<pre>[1],x[2]])) cat ElementToSequence(M2*Matrix(K,2,1,[x[3],x[4]])))[3],(</pre>
	<pre>ElementToSequence(M1*Matrix(K,2,1,[x[1],x[2]])) cat ElementToSequence(</pre>
	M2*Matrix(K,2,1,[x[3],x[4]])))[4]>>;
57	else
58	g:=map <k4->K4 x:-><(ElementToSequence(M1*Matrix(K,2,1,[x[3],x[4]]))</k4->
	<pre>cat ElementToSequence(M2*Matrix(K,2,1,[x[1],x[2]])))[1],(</pre>
	<pre>ElementToSequence(M1*Matrix(K,2,1,[x[3],x[4]])) cat ElementToSequence(</pre>
	M2*Matrix(K,2,1,[x[1],x[2]])))[2],(ElementToSequence(M1*Matrix(K,2,1,[x
	[3],x[4]])) cat ElementToSequence(M2*Matrix(K,2,1,[x[1],x[2]])))[3],(
	<pre>ElementToSequence(M1*Matrix(K,2,1,[x[3],x[4]])) cat ElementToSequence(</pre>
	M2*Matrix(K,2,1,[x[1],x[2]])))[4]>>;
59	end if;
60	
61	Phi:=map <k4->K4 x:-><x[1]*x[3],x[1]*x[4],x[2]*x[3],x[2]*x[4]>>;</x[1]*x[3],x[1]*x[4],x[2]*x[3],x[2]*x[4]></k4->
62	PhiInv:=map <k4->K4 x:-><x[1]+x[2],x[3]+x[4],x[1]+x[3],x[2]+x[4]>>;</x[1]+x[2],x[3]+x[4],x[1]+x[3],x[2]+x[4]></k4->
63	g1:=map <k4->K4 x:->Phi(g(PhiInv(x)))>;</k4->
64	
65	points:=[<2,8,1,4>,<1,1,0,0>,<1,0,1,0>,<0,1,0,1>,<0,0,2,1>];
66	<pre>images := [g1(points[1]),g1(points[2]),g1(points[3]),g1(points[4]),g1(</pre>
	<pre>points[5])];</pre>
67	<pre>return AutP3(points,images);</pre>
68	end function;
69	
70	// The following function takes a list of invertible matrices
71	// and their size, and returns the subgroup of PGL they generate.
72	
73	<pre>SGPGL:=function(matrices, dimension) gens:=[];</pre>
74	gens:=[]; for M in matrices do
$\frac{75}{76}$	<pre>gens:=Append(gens,M/Roots(x^dimension-R!Determinant(M))[1,1]);</pre>
70	end for:
78	G:=sub <gl(dimension,k) [gl(dimension,k) m: gens]="" in="" m="">;</gl(dimension,k) [gl(dimension,k) m:>
79	D:=[M: M in Center(G) IsScalar(M)];
80	GP:=quo <g d>;</g d>
81	<pre>return(GP);</pre>
82	end function;
83	
84	SGAutP1P1:=function(triples)

```
matrices:=[AutQuadSurf(t[1],t[2],t[3]):t in triples];
85
     return SGPGL(matrices,4);
86
   end function;
87
88
   // In what follows, IdentifyGroup is the built-in function
89
   // which returns the GAP ID of a finite group, if it exists.
90
   // Checking the generators given in Notation 3.6 for A4, S4, and A5:
91
92
   print "GAP ID of the subgroup of PGL(2) generated by A,B and C:",
93
       IdentifyGroup(SGPGL([A,B,C],2));
   print "GAP ID of the subgroup of PGL(2) generated by A,B,C and D:",
94
       IdentifyGroup(SGPGL([A,B,C,D],2));
   print "GAP ID of the subgroup of PGL(2) generated by E and F:",
95
       IdentifyGroup(SGPGL([E,F],2)),"\n";
96
   // Checking the GAP IDs in Proposition 5.10:
97
98
   print "GAP IDs of the subgroups of Aut(P^1xP^1) given in Proposition
99
       5.10:\n";
100
101 IdentifyGroup(SGAutP1P1([<A,A,0>,<B,B,0>,<C,C,0>,<I,I,1>]));
102 IdentifyGroup(SGAutP1P1([<A,A,0>,<B,B,0>,<C,C,0>,<D,D,1>]));
103 IdentifyGroup(SGAutP1P1([<A,I,0>,<B,I,0>,<C,C,0>,<I,I,1>]));
104 IdentifyGroup(SGAutP1P1([<A,I,0>,<B,I,0>,<C,C,0>,<I,C,1>]));
105 IdentifyGroup(SGAutP1P1([<A,I,0>,<B,I,0>,<C,C,0>,<D,D,1>]));
106 IdentifyGroup(SGAutP1P1([<A,I,0>,<B,I,0>,<C,I,0>,<I,I,1>]));
   IdentifyGroup(SGAutP1P1([<A,I,0>,<B,I,0>,<C,I,0>,<D,D,1>]));
107
108
   // Checking the GAP IDs in Proposition 5.13:
109
110
   print "\n GAP IDs of the subgroups of Aut(P^1xP^1) given in Proposition
111
       5.13:\n";
112
113 IdentifyGroup(SGAutP1P1([<A,A,0>,<B,B,0>,<C,C,0>,<D,D,0>,<I,I,1>]));
114 IdentifyGroup(SGAutP1P1([<A,I,0>,<B,I,0>,<C,C,0>,<D,D,0>,<I,I,1>]));
115 IdentifyGroup(SGAutP1P1([<A,I,0>,<B,I,0>,<C,I,0>,<D,D,0>,<I,I,1>]));
116 IdentifyGroup(SGAutP1P1([<A,I,0>,<B,I,0>,<C,I,0>,<D,D,0>,<I,D,1>]));
  IdentifyGroup(SGAutP1P1([<A,I,0>,<B,I,0>,<C,I,0>,<D,D,0>,<I,I,1>]));
117
118
119 // Printing a possible description of the last group in Propositon 5.13.
120 G:=SGAutP1P1([<A,I,O>,<B,I,O>,<C,I,O>,<D,I,O>,<I,I,1>]);
121 print "Last group. Order: ",Order(G)," description: ",GroupName(G);
```

References

- [BB00] L. Bayle & A. Beauville. Birational involutions of \mathbb{P}^2 . Asian J. Math., 4(1):11–17, 2000. 2
- [Bea07] A. Beauville. p-elementary subgroups of the Cremona group. J. Algebra, 314(2):553–564, 2007.
 23
- [BB04] A. Beauville & J. Blanc. On Cremona transformations of prime order. C. R., Math., Acad. Sci. Paris, 339(4):257–259, 2004. 2
- [BB73] A. Bialynicki-Birula. Some theorems on actions of algebraic groups. Ann. Math. (2), 98:480–497, 1973. 17
- [Bla06] J. Blanc. Finite abelian subgroups of the Cremona group of the plane. Genève: Univ. de Genève, Faculté des Sciences (Dissertation), 2006. 2
- [Bla09] J. Blanc. Linearisation of finite abelian subgroups of the Cremona group of the plane. Groups Geom. Dyn., 3(2):215–266, 2009. 3
- [BCDP23] J. Blanc, I. Cheltsov, A. Duncan & Y. Prokhorov. Finite quasisimple groups acting on rationally connected threefolds. *Math. Proc. Camb. Philos. Soc.*, 174(3):531–568, 2023. 5
- [Bli17] H. F. Blichfeldt. *Finite Collineation Groups*. University of Chicago Press, 1917. 13
- [BP13] F. Bogomolov & Y. Prokhorov. On stable conjugacy of finite subgroups of the plane Cremona group. I. Cent. Eur. J. Math., 11(12):2099–2105, 2013. 3, 4
- [BvBT23] C. Böhning, H.-C. G. von Bothmer & Y. Tschinkel. Equivariant birational geometry of cubic fourfolds and derived categories. Preprint, arXiv:2303.17678 [math.AG] (2023), 2023. 22
- [Che14] I. A. Cheltsov. Two local inequalities. Izv. Math., 78(2):375–426, 2014. 17, 31
- [Che08] I. Cheltsov. Log canonical thresholds of del Pezzo surfaces. Geom. Funct. Anal., 18(4):1118–1144, 2008. 17
- [CMYZ24] I. Cheltsov, F. Mangolte, E. Yasinsky & S. Zimmermann. Birational involutions of the real projective plane, J. Eur. Math. Soc. (JEMS), to appear., 2024. 3, 10
- [CMTZ24] I. Cheltsov, L. Marquand, Y. Tschinkel & Z. Zhang. Equivariant geometry of singular cubic threefolds, II. Preprint, arXiv:2405.02744 [math.AG] (2024), 2024. 5
- [CTZ23] I. Cheltsov, Y. Tschinkel & Z. Zhang. Equivariant geometry of the Segre cubic and the Burkhardt quartic. Preprint, arXiv:2308.15271 [math.AG] (2023), 2023. 5
- [CTZ24] I. Cheltsov, Y. Tschinkel & Z. Zhang. Equivariant geometry of singular cubic threefolds. Preprint, arXiv:2401.10974 [math.AG] (2024), 2024. 5
- [CTT24] T. Ciurca, S. Tanimoto & Y. Tschinkel. Intermediate Jacobians and linearizability. Preprint, arXiv:2403.06047 [math.AG] (2024), 2024. 5
- [dF04] T. de Fernex. On planar Cremona maps of prime order. Nagoya Math. J., 174:1–28, 2004. 2
- [dFE02] T. de Fernex & L. Ein. Resolution of indeterminacy of pairs. In Algebraic geometry. A volume in memory of Paolo Francia, pages 165–177. Berlin: de Gruyter, 2002. 7
- [DI09] I. V. Dolgachev & V. A. Iskovskikh. Finite subgroups of the plane Cremona group. In Algebra, arithmetic, and geometry. In honor of Yu. I. Manin on the occasion of his 70th birthday. Vol. I, pages 443–548. Boston, MA: Birkhäuser, 2009. 2, 3, 7, 8, 9, 17, 35
- [Ess24] L. Esser. The dual complex of a *G*-variety. *Math. Z.*, 308(3):8, 2024. Id/No 51. 5
- [Fab23] X. Faber. Finite *p*-irregular subgroups of $PGL_2(k)$. Matematica, 2(2):479–522, 2023. 12
- [Gou89] E. Goursat. Sur les substitutions orthogonales et les divisions régulières de l'espace. Ann. Sci. Éc. Norm. Supér. (3), 6:9–102, 1889. 14

- [HKT21] B. Hassett, A. Kresch & Y. Tschinkel. Symbols and equivariant birational geometry in small dimensions. In *Rationality of varieties. Proceedings of the conference, Island of Schiermonnikoog, The Netherlands, spring 2019*, pages 201–236. Cham: Birkhäuser, 2021. 5, 21
- [HT23] B. Hassett & Y. Tschinkel. Torsors and stable equivariant birational geometry. Nagoya Math. J., 250:275–297, 2023. 18, 22
- [Hup67] B. Huppert. Endliche Gruppen. I, volume 134 of Grundlehren Math. Wiss. Springer, Cham, 1967. 12
- [Isk69] V. A. Iskovskikh. Rational surfaces with pencil of rational curves. Math. USSR, Sb., 3:563–587, 1969. 10
- [Isk80] V. A. Iskovskikh. Minimal models of rational surfaces over arbitrary fields. Math. USSR, Izv., 14:17–39, 1980. 10
- [Isk96] V. A. Iskovskikh. Factorization of birational maps of rational surfaces from the viewpoint of Mori theory. Russ. Math. Surv., 51(4):585–652, 1996. 9
- [Isk08] V. A. Iskovskikh. Two non-conjugate embeddings of S₃ × Z₂ into the Cremona group. II. In Algebraic geometry in East Asia—Hanoi 2005. Proceedings of the 2nd international conference on algebraic geometry in East Asia, Hanoi, Vietnam, October 10−14, 2005, pages 251–267. Tokyo: Mathematical Society of Japan, 2008. 21
- [Kap57] I. Kaplansky. An introduction to differential algebra. Actualités Scientifiques et Industrielles. 1251. Publ. Inst. Math. Univ. Nancago. V. Paris: Hermann & Cie. 62 p. (1957)., 1957. 11
- [Kle19] F. Klein. Lectures on the icosahedron and the solution of equations of the fifth degree. Translated by George Gavin Morrice. With a new introduction and commentaries by Peter Slodowy. Translated by Lei Yang, volume 5 of CTM, Class. Top. Math. Beijing: Higher Education Press, reprint of the English translation of the 1884 German original edition edition, 2019. 11
- [KPT23] M. Kontsevich, V. Pestun & Y. Tschinkel. Equivariant birational geometry and modular symbols. J. Eur. Math. Soc. (JEMS), 25(1):153–202, 2023. 4
- [KT22a] A. Kresch & Y. Tschinkel. Equivariant birational types and Burnside volume. Ann. Sc. Norm. Super. Pisa, Cl. Sci. (5), 23(2):1013–1052, 2022. 4, 5
- [KT22b] A. Kresch & Y. Tschinkel. Equivariant Burnside groups and representation theory. Sel. Math., New Ser., 28(4):39, 2022. Id/No 81. 5
- [LPR06] N. Lemire, V. L. Popov & Z. Reichstein. Cayley groups. J. Am. Math. Soc., 19(4):921–967, 2006.
 22
- [NvdPT08] K. A. Nguyen, M. van der Put & J. Top. Algebraic subgroups of GL₂(ℂ). Indag. Math., New Ser., 19(2):287–297, 2008. 11
- [Pin24a] A. Pinardin. Antoine Pinardin's GitHub repository. https://github.com/antoinepinardin, 2024. Accessed: 2024-12-12. 38
- [Pin24b] A. Pinardin. G-solid rational surfaces. Eur. J. Math., 10(2):23, 2024. Id/No 33. 20
- [Pro15] Y. Prokhorov. On stable conjugacy of finite subgroups of the plane Cremona group. II. Mich. Math. J., 64(2):293–318, 2015. 4
- [Pro10] Y. G. Prokhorov. Fields of invariants of finite linear groups. In Cohomological and geometric approaches to rationality problems. New Perspectives, pages 245–273. Boston, MA: Birkhäuser, 2010. 18
- [RY00] Z. Reichstein & B. Youssin. Essential dimensions of algebraic groups and a resolution theorem for G-varieties. (With an appendix by János Kollár and Endre Szabó: Fixed points of group actions and rational maps). Can. J. Math., 52(5):1018–1056, 2000. 5
- [Spr77] T. A. Springer. Invariant theory, volume 585 of Lect. Notes Math. Springer, Cham, 1977. 13

- [TYZ22] Y. Tschinkel, K. Yang & Z. Zhang. Combinatorial Burnside groups. Res. Number Theory, 8(2):17, 2022. Id/No 33. 21
- [TYZ24] Y. Tschinkel, K. Yang & Z. Zhang. Equivariant birational geometry of linear actions. EMS Surv. Math. Sci., 11(2):235–276, 2024. 5
- [Wol18] J. Wolter. Equivariant birational geometry of quintic del Pezzo surface. Eur. J. Math., 4(3):1278– 1292, 2018. 18
- [Yas24] E. Yasinsky. On G-birational rigidity of del Pezzo surfaces. Épijournal de Géom. Algébr., EPIGA (to appear), 2024. 20, 33

SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH, EDINBURGH EH9 3JZ, UK *Email address*: antoine.pinardin@ed.ac.uk

London Institute for Mathematical Sciences, Royal Institution, 21 Albermarle St, London W1S 4BS, UK

Email address: ars@lims.ac.uk

L'Institut de Mathématiques de Bordeaux, Université de Bordeaux, 351 Cours de la Libération, 33405 Talence Cedex, France

Email address: egor.yasinsky@u-bordeaux.fr