

Non-Hermitian delocalization in 1D via emergent compactness

Liang-Hong Mo,^{1,2,3} Zhenyu Xiao,¹ Roderich Moessner,² and Hongzheng Zhao^{1,2,*}

¹*School of Physics, Peking University, 100871 Beijing, China.*

²*Max Planck Institute for the Physics of Complex Systems, Nöthnitzer Str. 38, 01187 Dresden, Germany.*

³*Department of Physics, Princeton University, Princeton, New Jersey 08544, USA*

(Dated: December 18, 2024)

Potential disorder in 1D leads to Anderson localization of the entire spectrum. Upon sacrificing hermiticity by adding non-reciprocal hopping, the non-Hermitian skin effect competes with localization. We find another route for delocalization, which involves *imaginary* potential disorder. While an entirely random potential generally still leads to localization, imposing minimal spatial structure to the disorder can protect delocalization: it endows the concomitant transfer matrix with an $SU(2)$ structure, whose compactness in turn translates into an infinite localization length. The fraction of delocalized states can be tuned by the choice of boundary conditions.

Introduction.— Anderson localization [1, 2] is an everlasting subject in condensed matter physics. The presence of quenched disorder significantly suppresses transport, especially in low dimensions. This stimulates decades of studies including, e.g., delocalization phase transitions with or without many-body interaction [3–5], and topological phases of matter in disordered systems [6–8].

Recently, localization in open systems has attracted considerable attention [9–31]. The interplay between non-Hermiticity and localization leads to a plethora of novel phenomena, unique in open quantum systems [32–35]. As shown in Hatano and Nelson’s pioneering work [9], in one-dimension (1D) asymmetric hopping can compete with quenched disorder, leading to delocalization, at variance with the scaling theory of localization for potential disorder [2]. Such delocalization originates from the non-trivial topology [15, 36], as elucidated by the effective field theory [21, 37]. Also, by introducing a deterministic quasi-periodic spatial structure delocalization may also appear [38–41], similar to their Hermitian counterpart [42–46].

Here, we address the intriguing question of whether 1D delocalized eigenstates can be realized by purely on-site energy gain and loss, without sacrificing reciprocity and spatial randomness. In general, a purely random complex potential localizes the entire spectrum, although prior work demonstrates the possibility of exotic transport dynamics in wave propagation [30, 47–50].

We provide an affirmative answer by constructing a concrete model with delocalized eigenstates and analytically tractable mobility edges. The key ingredient involves imposing a minimal structure on a binary potential, which we dub “dipolar disorder”. The corresponding transfer matrix generally belongs to the non-compact $SL(2, \mathbb{C})$ group [51, 52]; hence, a random product of such matrices leads to a positive Lyapunov exponent, i.e., a finite localization length [53]. However, the possible existence of an emergent compact $SU(2)$ structure, which we

identify, in the transfer matrix for the real-valued spectrum results in a zero Lyapunov exponent. It, in turn, leads to an infinite localization length, protecting delocalization against disorder. This $SU(2)$ structure can be revealed by a simple analytical procedure, cf. Fig. 1(a), allowing the exact determination of a mobility edge, for which the usual numerical approach faces difficulties, especially in the thermodynamic limit.

We numerically verify the theoretical prediction by diagonalizing the corresponding lattice model. A natural diagnostic of delocalization is the participation ratio of eigenstates, by which we obtain a localization-delocalization phase diagram at different energies, Fig. 1(b). The mobility edge precisely matches the analytical prediction. Crucially, the fraction of delocalized eigenstates depends on the boundary conditions, which are tunable by a twisted phase factor, cf. Fig. 1(c). For complex energy eigenvalues that go beyond this analytical framework, eigenstates are generally localized. However, for a given finite system size, we find strong numerical evidence showing that delocalization can persist as long as the imaginary component of the energy eigenvalue is sufficiently small.

The remainder of this account is organized as follows. First, we introduce the model with disordered imaginary potential and the dipolar structure. Then we elaborate on the emergent $SU(2)$ structure in the transfer matrix. By numerically calculating the participation ratio of each eigenstate, we demonstrate the existence of delocalization and the localization transition. Finally, we summarize and discuss open directions where the emergent compactness in the transfer matrix may play a crucial role.

The model.— The Schrödinger equation for 1D tight-binding Hamiltonian H with uniform and reciprocal nearest-neighbor hopping and on-site potential V_j reads

$$E\Psi_j = \Psi_{j+1} + \Psi_{j-1} + V_j\Psi_j, \quad j \in \mathbb{Z}, \quad (1)$$

where $\Psi_j \equiv \langle j|\Psi\rangle$ is the real-space wavefunction on site j and E defines the energy. It can be rewritten as

$$\begin{pmatrix} \Psi_{j+1} \\ \Psi_j \end{pmatrix} = T_j \begin{pmatrix} \Psi_j \\ \Psi_{j-1} \end{pmatrix}, \quad T_j = \begin{pmatrix} E - V_j & -1 \\ 1 & 0 \end{pmatrix}, \quad (2)$$

* hzhao@pku.edu.cn




FIG. 1. (a) Schematic for the emergent compact structure. The overall gray region denotes the non-compact $SL(2, \mathbb{C})$ group, which contains the compact $SU(2)$ structure (blue). The green region denotes a subset of $SL(2, \mathbb{C})$, inside which a matrix element as well as its conjugation can be simultaneously mapped to $SU(2)$ matrices, via a similarity transformation PW . The dipolar transfer matrices, \mathcal{M} and \mathcal{N} , belong to this subset and hence a random product of the two leads to zero Lyapunov exponent, i.e., an infinite localization length. (b) Participation ratio (PR) calculated for each eigenstate at different real energies E and disorder strength V . The emergent compactness leads to delocalized eigenstates (blue), which are separated from the localized states (red) by an exact mobility edge (black). This black line is obtained by analyzing the possible existence of a similarity transformation. In numerical simulation, we scan over all possible boundary conditions with a twisted angle K and sample different realizations of the disorder potential. (c) By tuning K , we can control the fraction of real energies (orange) in the entire spectrum and the delocalized eigenstates therein (blue). For periodic and anti-periodic boundary conditions $K=0, \pi$, the fraction of eigenstates is around 10^{-4} , a small but nonzero value. Yet, for other K values, most real energies are delocalized. For numerical simulation, we use $V = 0.3$, $2L = 1024$. We average over 100 random realizations to obtain both (b) and (c).

where T_j is the transfer matrix. For the entire chain with length N , the transfer matrix reads $T_{\text{tot}}^{(N)} = \prod_{j=1}^N T_j$. The corresponding Lyapunov exponent λ_L is defined as $\lambda_L \equiv \lim_{N \rightarrow \infty} \log \|T_{\text{tot}}^{(N)}\| / N$, where $\|\dots\|$ denotes the 2-norm [54]. For a given eigen-energy E , the localization length $\xi(E)$ can be obtained as the inverse of the Lyapunov exponent λ_L of $T_{\text{tot}}^{(N)}$ [55–57]. In the conventional Hermitian Anderson model where V_j is a real random variable, it has been shown that λ_L is positive for any energy E , therefore all eigenstates of the 1D Anderson model are localized.

Now we consider a non-Hermitian model with imaginary and binary potential, i.e., $V_j = \pm iV$, $V \in \mathbb{R}$. We focus on states with real energies and obtain two transfer matrices

$$T_{\pm} = \begin{pmatrix} E \pm iV & -1 \\ 1 & 0 \end{pmatrix}, \quad (3)$$

which satisfy $T_+ = T_-^*$ and belong to the group $SL(2, \mathbb{C})$. Its associated manifold is non-compact [51, 52]. According to Furstenberg’s theorem [58], in a non-compact manifold, a random product of matrices generally leads to a positive Lyapunov exponent. Therefore, just as the conventional Hermitian Anderson model, the onsite imaginary potential can localize all eigenstates, in accordance with previous predictions in Refs. [30, 47, 50].

Emergent compactness.— We impose a dipolar structure into the spatial disorder, such that the minimal building block of the disorder becomes either $\{iV, -iV\}$, or $\{-iV, iV\}$. Each block has a balanced energy gain (iV) and loss ($-iV$), resulting in a parity-time symme-

try within the block, also referred to as “local parity-time symmetry” [59]. By contrast, global parity-time symmetry is absent due to the spatial randomness of the blocks. The corresponding transfer matrices for two consecutive sites become $\mathcal{M} = T_+ T_-$, $\mathcal{N} = T_- T_+$, which satisfy $\mathcal{M} = \mathcal{N}^*$ and exhibit a real trace.

A peculiar feature appears in this dipolar construction: as schematically illustrated in Fig. 1(a), an emergent compactness arises because both \mathcal{M} and \mathcal{N} can simultaneously be transformed into $SU(2)$ matrices via a similarity transformation. As the corresponding manifold for the $SU(2)$ group is compact, the Lyapunov exponent for a series of random multiplication of \mathcal{M} and \mathcal{N} vanishes.

The crucial ingredient that allows for such a similarity transformation is simply:

$$\text{Tr}(\mathcal{M}\mathcal{N}) \leq 2. \quad (4)$$

This, together with the condition $\mathcal{M} = \mathcal{N}^*$, guarantees that the eigenvalues of both \mathcal{M} and \mathcal{N} satisfy the spectral requirements of the $SU(2)$ group. Numerically, one can efficiently scan over the entire parameter space and show that Eq. 4 leads to the blue region in Fig. 1(b), including the black line [60].

To construct the desired similarity transformation, we can first diagonalize the matrix \mathcal{M} via a normalized matrix P , which has the following general form $P = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. One can show that Eq. 4 ensures that the entire blue region satisfies $(b^*d - bd^*)(-a^*c + ac^*) \leq 0$. When the equal sign is taken, the matrix P is sufficient to simultaneously transform both \mathcal{M} and \mathcal{N} into two $SU(2)$ matrices. Oth-

erwise, one can construct a product of matrix P and W , where

$$W = \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{\frac{-a^*c+ac^*}{b^*d-bd^*}} \end{pmatrix}, \quad (5)$$

to achieve such a similar transformation. A detailed proof and a more general discussion is presented in the Appendix.

Consequently, inside the blue region in Fig. 1(b), the dipolar transfer matrices have an emergent compact $SU(2)$ structure, resulting in a vanishing Lyapunov exponent and a diverging localization length. In contrast, for parameters in the red region in Fig. 1(b), both \mathcal{M} and \mathcal{N} have an eigenvalue larger than one, thus they do not belong to any compact subgroup of $SL(2, \mathbb{C})$. Therefore, localization is expected.

Numerical results. — It is worth highlighting that, here the energy eigenvalue E is not a continuously tunable parameter. Rather, it depends on the specific spatial disorder realization as well as the boundary condition of the lattice. In the following, we show that a nonzero fraction of real eigenvalues, as well as delocalized eigenstates therein, can exist, and can be tunable by the twisted boundary condition.

For a system containing $2L$ sites (L dipoles), the spectrum can be obtained by solving the following bulk equation and boundary condition

$$\begin{pmatrix} \Psi_{2L+1} \\ \Psi_{2L} \end{pmatrix} = T_{\text{tot}}^{(2L)} \begin{pmatrix} \Psi_1 \\ \Psi_0 \end{pmatrix}, \quad \begin{pmatrix} \Psi_{2L+1} \\ \Psi_{2L} \end{pmatrix} = e^{iK} \begin{pmatrix} \Psi_1 \\ \Psi_0 \end{pmatrix}, \quad (6)$$

leading to

$$T_{\text{tot}}^{(2L)} \begin{pmatrix} \Psi_1 \\ \Psi_0 \end{pmatrix} = e^{iK} \begin{pmatrix} \Psi_1 \\ \Psi_0 \end{pmatrix}, \quad (7)$$

where we choose twisted boundary condition with a real number K [61]. $K=0$ corresponds to a periodic boundary condition.

If \mathcal{M} and \mathcal{N} can be simultaneously transformed into $SU(2)$ matrices, the eigenvalues of T_{tot} take the form of a $U(1)$ phase factor. Therefore, for a specific set of V and E , there exists a suitable boundary condition K such that the energy E exists in the spectrum.

To verify this, we numerically perform exact diagonalization of the Hamiltonian

$$\mathcal{H} = \sum_{j=0}^{2L-2} (\hat{c}_j^\dagger \hat{c}_{j+1} + h.c.) + \sum_{j=0}^{L-1} \left(V_j \hat{c}_{2j}^\dagger \hat{c}_{2j} - V_j \hat{c}_{2j+1}^\dagger \hat{c}_{2j+1} \right) + \left[\exp(iK) \hat{c}_0^\dagger \hat{c}_{2L-1} + h.c. \right], \quad (8)$$

and obtain the entire single-particle energy eigenvalues for different K . As expected, since the Hamiltonian is non-Hermitian, energy eigenvalues are mostly complex, see Fig. 3 (a). However, crucially, real eigenvalues also




FIG. 2. Scaling of the participation ratio (PR) for different system sizes. For eigenstates in the blue region of Fig. 1 (b), PR scales linearly with system size (black dashed line of slope 1 is a guide to the eye), confirming its delocalized nature. In contrast, for the eigenstates in the red region of Fig. 1 (b), PR is independent of system size, corresponding to localized states. We use the parameters $V = 0.5$ and $K = \pi/2$ for numerical simulation.

exist and their fraction depends on the boundary condition: As depicted in Fig. 1 (c), for periodic and anti-periodic boundary conditions ($K=0$ and π) this fraction (orange) is around 10^{-2} , and the fraction for delocalized eigenstates (blue) is around 10^{-4} , see details in the Supplementary Material (SM), Sec. SM 1. Whereas for other K values, a notable fraction of real eigenvalues appears, and most of the corresponding eigenstates are indeed delocalized. Note, these results can be efficiently obtained by counting the number of states located inside the black line in Fig. 1 (b).

To characterize the localization properties of the system, we calculate the participation ratio (PR) [38] for each normalized right eigenvector $|\Psi\rangle$,

$$\text{PR} = \left(\sum_{j=0}^{2L-1} |\langle j | \Psi \rangle|^4 \right)^{-1}. \quad (9)$$

For localized states, PR is of order $\mathcal{O}(1)$ and remains independent of the system size; for delocalized states, it becomes proportional to the system size, $\mathcal{O}(L)$. Fig. 1(b) presents the PR for all eigenstates with real eigenvalues, scanning over a wide range of the disorder strength V and different boundary conditions K . In the absence of disorder ($V=0$), all states are delocalized in space, exhibiting large values of PR (blue). For non-zero V , localized eigenstates appear with a notably smaller PR (red). Importantly, a mobility edge, separating the localized and delocalized regimes, is clearly visible and it precisely matches the analytical prediction (black line).

In Fig. 2, we depict the PR averaged over either the localized states (orange) or the delocalized states (blue)




FIG. 3. (a) PR for the entire complex spectrum. Our simulations use $V = 0.6$ and system size $L = 8192$. Most eigenstates with complex energies are localized, while a small fraction of energies near the real axis (bright region) exhibits large PR values, suggesting delocalization may persist. (b) System size scaling of PR, averaged over eigenstates within different imaginary energy windows. The real part of the energy is fixed within the range $[0.8, 1.2]$. For an eigenvalue sufficiently close to the real axis, PR exhibits linear scaling in L (the black dashed line of slope 1 is a guide to the eye). When deviating further away from the real axis, a power law scaling appears with a scaling exponent smaller than 1. A log-log scale is used here.

for different system sizes, where the linear system size dependence in PR confirms delocalization.

Complex spectrum.— The delocalized eigenstates and their localization transition have been well-established for real energies. For complex energies, the existence of the emergent $SU(2)$ structure is not guaranteed, nor is the existence of a delocalized eigenstate. In fact, most of the complex spectrum becomes localized, as shown in Fig. 3 (a), where PR for the entire spectrum is plotted.

For eigenvalues close to the real axis, matrix elements of T_{\pm} (Eq. 3) are still dominated by the real part of the energy. Therefore, despite the energy eigenvalues not being purely real, the resulting localization length can still be sufficiently large, even comparable to system sizes that are numerically accessible. Indeed, as shown in Fig. 3 (a), a bright region with large PR values appears, indicating the possible existence of delocalized eigenstates.

A systematic analysis of the system size dependence is presented in Fig. 3 (b). The real part of the energy is fixed within the range $\text{Re}(E) \in [0.8, 1.2]$, such that all eigenstates are delocalized if their energies are purely real. We average the PR over all eigenstates within different imaginary energy windows. For eigenstates with a small imaginary part, $\text{Im}(E) \in [10^{-3}, 10^{-2}]$ (orange), the averaged PR grows linearly with L , indicating that delocalization persists even for a complex eigenvalue. This can be a finite-size effect, but the localization length in practice is comparable to or larger than the current system size. Consider $\text{Im}(E) \in [10^{-2}, 10^{-1}]$ (light blue), where a crossover behavior appears, showing a power law scaling with a scaling exponent less than 1. In contrast,

for $\text{Im}(E) \in [0.1, 1]$ (dark blue), PR becomes independent of system size, implying localization.

Discussion.— We have presented a simple 1D non-Hermitian system with a purely imaginary disorder potential that leads to delocalized eigenstates. The crucial ingredient is an emergent compact $SU(2)$ structure in the concomitant transfer matrix. It leads to a zero Lyapunov exponent and hence an infinite localization length for real eigenvalues, as further supported by numerical simulation. For complex energies, delocalization persists as long as the imaginary component of the eigenvalue is sufficiently small. It would be worthwhile to perform further numerical investigations to clarify whether this is a finite-size phenomenon.

We also note that the spectrum exhibits an interesting D_2 symmetry, namely the mirror symmetry w.r.t. both real and imaginary axes. For the imaginary axis, the symmetry originates from a chiral symmetry, $PH^{\dagger}P^{-1} = -H$, where H is the single-particle Hamiltonian matrix of Eq. 8 and P denotes the chiral symmetry operator. This leads to the coexistence of the energies E and $-E^*$. Additionally, one can show that energy eigenvalues also appear in pairs, E and $-E$. This is more subtle and can be revealed by showing that the trace of the total transfer matrix only contains even powers of the energy E for twist boundary conditions. Details of the proof are illustrated in Sec. SM 2. The connection between this symmetry and delocalization is an intriguing subject for future study.

For disordered higher dimensional systems, a generic transfer matrix of large dimension normally appears, and the possible existence of a compact structure remains an open and interesting question.

While focusing on the eigenstate properties of the system, we emphasize that this emergent compact structure can also lead to interesting dynamical phenomena that may not exist in non-Hermitian systems with an entirely localized spectrum [30, 31, 62, 63].

There are various experimental platforms capable of realizing tunable control over the on-site gain and loss, such as photonic waveguides [64–67] and mechanical systems [68]. Therefore, the proposed dipolar potential structure, as well as the predicted delocalization should be readily accessible in these physical settings.

Finally, we emphasize that, due to the broad applicability of the transfer matrix method, the emergent compactness could have significant implications at a more general level, particularly in many-body systems. For instance, a similar transfer matrix method has been developed in the study of dual unitary and Gaussian quantum circuits [69, 70], as well as within the framework of time-dependent driven systems exhibiting conformal symmetry [71]. We anticipate that this emergent compactness will lead to more surprises and unexpected dynamical non-equilibrium many-body phenomena.

Acknowledgments.— We thank Kun Ding, Yumin Hu, Sen Mu, and Bo-Ting Chen for stimulating discussions. This work is supported by the National Natural Science

Foundation of China (Grant No. 12474214), by “The Fundamental Research Funds for the Central Universities, Peking University”, by “High-performance Comput-

ing Platform of Peking University” and by the Deutsche Forschungsgemeinschaft under the cluster of excellence ct.qmat (EXC 2147, project-id 390858490).

-
- [1] P. W. Anderson, Absence of Diffusion in Certain Random Lattices, *Phys. Rev.* **109**, 1492 (1958).
- [2] F. Evers and A. D. Mirlin, Anderson transitions, *Reviews of Modern Physics* **80**, 1355 (2008), [arXiv:0707.4378 \[cond-mat.mes-hall\]](#).
- [3] R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, *Annu. Rev. Condens. Matter Phys.* **6**, 15 (2015).
- [4] A. Smith, J. Knolle, D. L. Kovrizhin, and R. Moessner, Disorder-free localization, *Physical review letters* **118**, 266601 (2017).
- [5] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, *Reviews of Modern Physics* **91**, 021001 (2019).
- [6] J. Li, R.-L. Chu, J. K. Jain, and S.-Q. Shen, Topological anderson insulator, *Physical review letters* **102**, 136806 (2009).
- [7] P. Titum, E. Berg, M. S. Rudner, G. Refael, and N. H. Lindner, Anomalous floquet-anderson insulator as a nonadiabatic quantized charge pump, *Physical Review X* **6**, 021013 (2016).
- [8] S. Stützer, Y. Plotnik, Y. Lumer, P. Titum, N. H. Lindner, M. Segev, M. C. Rechtsman, and A. Szameit, Photonic topological anderson insulators, *Nature* **560**, 461 (2018).
- [9] N. Hatano and D. R. Nelson, Localization transitions in non-hermitian quantum mechanics, *Physical review letters* **77**, 570 (1996).
- [10] J. Feinberg and A. Zee, Non-hermitian random matrix theory: Method of hermitian reduction, *Nucl. Phys. B* **504**, 579 (1997).
- [11] J. Feinberg and A. Zee, Non-Hermitian localization and delocalization, *Phys. Rev. E* **59**, 6433 (1999).
- [12] A. Regensburger, M.-A. Miri, C. Bersch, J. Näger, G. Onishchukov, D. N. Christodoulides, and U. Peschel, Observation of defect states in pt-symmetric optical lattices, *Physical review letters* **110**, 223902 (2013).
- [13] B. Xu, T. Ohtsuki, and R. Shindou, Integer quantum magnon Hall plateau-plateau transition in a spin-ice model, *Phys. Rev. B* **94**, 220403(R) (2016).
- [14] K. Ding, G. Ma, M. Xiao, Z. Zhang, and C. T. Chan, Emergence, coalescence, and topological properties of multiple exceptional points and their experimental realization, *Physical Review X* **6**, 021007 (2016).
- [15] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, Topological Phases of Non-Hermitian Systems, *Phys. Rev. X* **8**, 031079 (2018).
- [16] R. Hamazaki, K. Kawabata, and M. Ueda, Non-hermitian many-body localization, *Physical review letters* **123**, 090603 (2019).
- [17] A. F. Tzortzakakis, K. G. Makris, and E. N. Economou, Non-Hermitian disorder in two-dimensional optical lattices, *Phys. Rev. B* **101**, 014202 (2020).
- [18] C. Wang and X. R. Wang, Level statistics of extended states in random non-Hermitian Hamiltonians, *Phys. Rev. B* **101**, 165114 (2020).
- [19] Y. Huang and B. I. Shklovskii, Anderson transition in three-dimensional systems with non-Hermitian disorder, *Phys. Rev. B* **101**, 014204 (2020).
- [20] Y. Huang and B. I. Shklovskii, Spectral rigidity of non-Hermitian symmetric random matrices near the Anderson transition, *Phys. Rev. B* **102**, 064212 (2020).
- [21] K. Kawabata and S. Ryu, Nonunitary Scaling Theory of Non-Hermitian Localization, *Phys. Rev. Lett.* **126**, 166801 (2021).
- [22] X. Luo, T. Ohtsuki, and R. Shindou, Universality Classes of the Anderson Transitions Driven by Non-Hermitian Disorder, *Phys. Rev. Lett.* **126**, 090402 (2021).
- [23] X. Luo, T. Ohtsuki, and R. Shindou, Transfer matrix study of the Anderson transition in non-Hermitian systems (2021), [arXiv:2103.05239 \[quant-ph\]](#).
- [24] X. Luo, Z. Xiao, K. Kawabata, T. Ohtsuki, and R. Shindou, Unifying the Anderson transitions in Hermitian and non-Hermitian systems, *Phys. Rev. Research* **4**, L022035 (2022).
- [25] W. Zhu and J. Gong, Hybrid skin-topological modes without asymmetric couplings, *Physical Review B* **106**, 035425 (2022).
- [26] Y. Li, C. Liang, C. Wang, C. Lu, and Y.-C. Liu, Gain-loss-induced hybrid skin-topological effect, *Physical Review Letters* **128**, 223903 (2022).
- [27] T. Zhang and Z. Cai, Emergent non-hermitian physics in a generalized lotka-volterra model, *Physical Review B* **108**, 104304 (2023).
- [28] P. Zhu, X.-Q. Sun, T. L. Hughes, and G. Bahl, Higher rank chirality and non-hermitian skin effect in a topolectrical circuit, *Nature communications* **14**, 720 (2023).
- [29] F. Thompson, Y. Huang, and A. Kamenev, Localization of Lindbladian fermions, *Phys. Rev. B* **109**, 174201 (2024).
- [30] B. Li, C. Chen, and Z. Wang, Universal non-hermitian transport in disordered systems, *arXiv preprint arXiv:2411.19905* (2024).
- [31] Z.-Y. Xing, S. Chen, and H. Hu, Universal spreading dynamics in quasiperiodic non-hermitian systems, *arXiv preprint arXiv:2412.01301* (2024).
- [32] S. Yao and Z. Wang, Edge states and topological invariants of non-hermitian systems, *Physical review letters* **121**, 086803 (2018).
- [33] K. Yokomizo and S. Murakami, Non-bloch band theory of non-hermitian systems, *Physical review letters* **123**, 066404 (2019).
- [34] N. Okuma, K. Kawabata, K. Shiozaki, and M. Sato, Topological origin of non-hermitian skin effects, *Physical review letters* **124**, 086801 (2020).
- [35] Y. Ashida, Z. Gong, and M. Ueda, Non-hermitian physics, *Advances in Physics* **69**, 249 (2020).
- [36] N. Okuma and M. Sato, Non-hermitian topological phenomena: A review, *Annual Review of Condensed Matter Physics* **14**, 83 (2023).
- [37] Z. Chen, K. Kawabata, A. Kulkarni, and S. Ryu,

- Field theory of non-hermitian disordered systems, arXiv preprint arXiv:2411.11878 (2024).
- [38] S. Longhi, Topological phase transition in non-hermitian quasicrystals, *Physical review letters* **122**, 237601 (2019).
- [39] Y. Liu, Q. Zhou, and S. Chen, Localization transition, spectrum structure, and winding numbers for one-dimensional non-hermitian quasicrystals, *Physical Review B* **104**, 024201 (2021).
- [40] L.-J. Zhai, G.-Y. Huang, and S. Yin, Cascade of the delocalization transition in a non-hermitian interpolating aubry-andré-fibonacci chain, *Physical Review B* **104**, 014202 (2021).
- [41] L.-M. Chen, Y. Zhou, S. A. Chen, and P. Ye, Quantum entanglement of non-hermitian quasicrystals, *Physical Review B* **105**, L121115 (2022).
- [42] S. Aubry and G. André, Analyticity breaking and anderson localization in incommensurate lattices, *Ann. Israel Phys. Soc* **3**, 18 (1980).
- [43] Y. Liu and R. Riklund, Electronic properties of perfect and nonperfect one-dimensional quasicrystals, *Physical Review B* **35**, 6034 (1987).
- [44] F. Axel and J. Peyrière, Spectrum and extended states in a harmonic chain with controlled disorder: effects of the thue-morse symmetry, *Journal of statistical physics* **57**, 1013 (1989).
- [45] J. Biddle, B. Wang, D. Priour Jr, and S. Das Sarma, Localization in one-dimensional incommensurate lattices beyond the aubry-andré model, *Physical Review A—Atomic, Molecular, and Optical Physics* **80**, 021603 (2009).
- [46] Y. Wang, X. Xia, L. Zhang, H. Yao, S. Chen, J. You, Q. Zhou, and X.-J. Liu, One-dimensional quasiperiodic mosaic lattice with exact mobility edges, *Physical Review Letters* **125**, 196604 (2020).
- [47] A. Basiri, Y. Bromberg, A. Yamilov, H. Cao, and T. Kottos, Light localization induced by a random imaginary refractive index, *Physical Review A* **90**, 043815 (2014).
- [48] I. Yusipov, T. Lapyteva, S. Denisov, and M. Ivanchenko, Localization in open quantum systems, *Physical review letters* **118**, 070402 (2017).
- [49] A. Tzortzakakis, K. Makris, A. Szameit, and E. Economou, Transport and spectral features in non-hermitian open systems, *Physical Review Research* **3**, 013208 (2021).
- [50] S. Weidemann, M. Kremer, S. Longhi, and A. Szameit, Coexistence of dynamical delocalization and spectral localization through stochastic dissipation, *Nature Photonics* **15**, 576 (2021).
- [51] B. C. Hall, *Lie groups, Lie algebras, and representations* (Springer, 2013).
- [52] P. Woit, *Quantum theory, groups and representations*, Vol. 4 (Springer, 2017).
- [53] F. Evers and A. D. Mirlin, Anderson transitions, *Reviews of Modern Physics* **80**, 1355 (2008).
- [54] The corresponding Lyapunov exponents λ_i ($i = 1, 2$; $\lambda_1 \geq \lambda_2$) are defined by the eigenvalues of $\lim_{N \rightarrow \infty} \log[(T_{\text{tot}}^{(N)})^\dagger T_{\text{tot}}^{(N)}]/2N$. The localization length is defined as $\xi(E) \equiv (\min_i |\lambda_i|)^{-1}$ for an eigenstate with energy E . For the 2×2 transfer matrix here, we always have $\lambda_2 = -\lambda_1$, and hence $\xi = \lambda_1^{-1}$. It is easy to prove that λ_1 is equivalent to the Lyapunov exponent defined through 2-norm.
- [55] R. Carmona, Exponential localization in one-dimensional disordered systems, *Duke Mathematical Journal* **49**, 191 (1982).
- [56] A. Comtet, C. Texier, and Y. Tourigny, Lyapunov exponents, one-dimensional anderson localization and products of random matrices, *Journal of Physics A: Mathematical and Theoretical* **46**, 254003 (2013).
- [57] K. Slevin and T. Ohtsuki, Critical exponent for the Anderson transition in the three-dimensional orthogonal universality class, *New J. Phys.* **16**, 015012 (2014).
- [58] H. Furstenberg, Noncommuting random products, *Transactions of the American Mathematical Society* **108**, 377 (1963).
- [59] O. Bendix, R. Fleischmann, T. Kottos, and B. Shapiro, Optical structures with local-symmetry, *Journal of Physics A: Mathematical and Theoretical* **43**, 265305 (2010).
- [60] The analytical expression for the mobility edge is $-4E^2 + E^4 + 2E^2V^2 + V^4 = 0$.
- [61] M. Kohmoto, Localization problem and mapping of one-dimensional wave equations in random and quasiperiodic media, *Physical Review B* **34**, 5043 (1986).
- [62] M. C. Zheng, D. N. Christodoulides, R. Fleischmann, and T. Kottos, Pt optical lattices and universality in beam dynamics, *Physical Review A—Atomic, Molecular, and Optical Physics* **82**, 010103 (2010).
- [63] A. Regensburger, C. Bersch, M.-A. Miri, G. Onishchukov, D. N. Christodoulides, and U. Peschel, Parity-time synthetic photonic lattices, *Nature* **488**, 167 (2012).
- [64] A. Guo, G. J. Salamo, D. Duchesne, R. Morandotti, M. Volatier-Ravat, V. Aimez, G. A. Siviloglou, and D. N. Christodoulides, Observation of pt-symmetry breaking in complex optical potentials, *Physical review letters* **103**, 093902 (2009).
- [65] L. Cong, J. Han, W. Zhang, and R. Singh, Temporal loss boundary engineered photonic cavity, *Nature Communications* **12**, 6940 (2021).
- [66] Q. Lin, T. Li, L. Xiao, K. Wang, W. Yi, and P. Xue, Topological phase transitions and mobility edges in non-hermitian quasicrystals, *Physical Review Letters* **129**, 113601 (2022).
- [67] A. Ma, T. Dai, J. Mao, Z. Fu, Y. Yang, X. Hu, Q. Gong, and J. Wang, Anisotropic quantum transport in a programmable photonic topological insulator, *Optica* **11**, 1533 (2024).
- [68] R. Süssstrunk and S. D. Huber, Classification of topological phonons in linear mechanical metamaterials, *Proceedings of the National Academy of Sciences* **113**, E4767 (2016).
- [69] P. W. Claeys and A. Lamacraft, Ergodic and nonergodic dual-unitary quantum circuits with arbitrary local hilbert space dimension, *Physical Review Letters* **126**, 100603 (2021).
- [70] E. Granet, C. Zhang, and H. Dreyer, Volume-law to area-law entanglement transition in a nonunitary periodic gaussian circuit, *Physical Review Letters* **130**, 230401 (2023).
- [71] R. Fan, Y. Gu, A. Vishwanath, and X. Wen, Emergent spatial structure and entanglement localization in floquet conformal field theory, *Physical Review X* **10**, 031036 (2020).

End Matter

Appendix.—

In the following, we prove that the condition $\text{tr}(\mathcal{M}\mathcal{N}) \in [-2, 2]$ leads to an emergent compact $\text{SU}(2)$ structure in the transfer matrix of our system.

We first consider two arbitrary $\text{SL}(2, \mathbb{C})$ matrices \mathcal{M}, \mathcal{N} and analyze the minimal conditions that allow for the existence of a similarity transformation to simultaneously transform them into two $\text{SU}(2)$ matrices. The general form of a $\text{SU}(2)$ matrix reads $\begin{pmatrix} \alpha & \beta \\ -\beta^* & \alpha^* \end{pmatrix}$. Then we apply this framework to our specific model being considered in the localization problem. These conditions read:

1. \mathcal{M}, \mathcal{N} are not triangular matrices and they satisfy $\mathcal{M} = \mathcal{N}^*$.
2. $\text{tr}(\mathcal{M}) \in [-2, 2]$.
3. $\text{tr}(\mathcal{M}\mathcal{N}) \leq 2$.

Proof. Condition 1 and 2 imply that the eigenvalues of \mathcal{M} and \mathcal{N} are $u + iv, u - iv$ with $u^2 + v^2 = 1$. Thus they can be expressed as

$$\mathcal{M} = P\Lambda P^{-1}, P = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \Lambda = \begin{pmatrix} u + iv & 0 \\ 0 & u - iv \end{pmatrix}, \quad (10)$$

$$\mathcal{N} = P^*\Lambda^*(P^*)^{-1}, \quad (11)$$

where each column of P is an eigenvector of \mathcal{M} . For simplicity, we consider P to be normalized and $\det(P) = 1$.

By construction, P diagonalizes \mathcal{M} into Λ , which is a $\text{SU}(2)$ matrix. Similarly, P transforms \mathcal{N} to

$$P^{-1}\mathcal{N}P = P^{-1}P^*\Lambda^*(P^*)^{-1}P, \quad (12)$$

which, in general, is not a $\text{SU}(2)$ matrix, and we have the explicit expression

$$P^{-1}P^* = \begin{pmatrix} -bc^* + a^*d & b^*d - bd^* \\ -a^*c + ac^* & -b^*c + ad^* \end{pmatrix}. \quad (13)$$

Note, its diagonal elements are conjugates of each other, an important feature of a generic $\text{SU}(2)$ matrix. However, its off-diagonal elements can break the $\text{SU}(2)$ structure. There are three possibilities:

Case 1: All off-diagonal elements are zero. $P^{-1}P^*$ reduces to the identity matrix, which is naturally a $\text{SU}(2)$ matrix;

Case 2: Only one of the off-diagonal elements is non-zero. If so, \mathcal{M} becomes a triangular matrix and hence Condition 1 is not satisfied.

Case 3: Both off-diagonal elements are non-zero. In this case, we need to introduce the one extra matrix

$$W = \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{\frac{-a^*c + ac^*}{b^*d - bd^*}} \end{pmatrix}. \quad (14)$$

By applying a similarity transformation W to Eq. 13, we have

$$U \equiv W^{-1}P^{-1}P^*W \quad (15)$$

$$= \begin{pmatrix} -bc^* + a^*d & (b^*d - bd^*)\sqrt{\frac{a^*c - ac^*}{-b^*d + bd^*}} \\ (b^*d - bd^*)\sqrt{\frac{a^*c - ac^*}{-b^*d + bd^*}} & -b^*c + ad^* \end{pmatrix}.$$

By comparing U with the general form of a $\text{SU}(2)$ matrix, one finds that if and only if $(b^*d - bd^*)(-a^*c + ac^*) < 0$, U becomes an $\text{SU}(2)$ matrix.

Therefore, by using the matrix PW one can transform \mathcal{N} as

$$(PW)^{-1}\mathcal{N}PW = U\Lambda^*U^{-1}, \quad (16)$$

which is now a product of three $\text{SU}(2)$ matrices.

In summary, based on the first two requirements as well as $(b^*d - bd^*)(-a^*c + ac^*) \leq 0$, \mathcal{M} and \mathcal{N} can be simultaneously similarly transformed into $\text{SU}(2)$ matrices.

In the following, we show that the condition $(b^*d - bd^*)(-a^*c + ac^*) \leq 0$ is equivalent to a simpler expression

$$\text{tr}(\mathcal{M}\mathcal{N}) \leq 2. \quad (17)$$

By using Eq. 10, we have

$$\text{tr}(\mathcal{M}\mathcal{N}) = 2u^2 + 2[(b^*c - ad^*)(bc^* - a^*d) + (b^*d - bd^*)(-a^*c + ac^*)]v^2. \quad (18)$$

Each term inside the four pairs of parentheses is nothing but the four elements of the matrix $P^{-1}P^*$. By noting that $\det(P^{-1}P^*) = 1$ and $u^2 + v^2 = 1$, we have

$$\begin{aligned} \text{tr}(\mathcal{M}\mathcal{N}) &= 2u^2 + 2[1 + 2(b^*d - bd^*)(-a^*c + ac^*)]v^2 \\ &= 2u^2 + 2v^2 + 4(b^*d - bd^*)(-a^*c + ac^*)v^2 \\ &= 2 + 4(b^*d - bd^*)(-a^*c + ac^*)v^2. \end{aligned} \quad (19)$$

Hence, $\text{tr}(\mathcal{M}\mathcal{N}) \leq 2$ implies $(b^*d - bd^*)(-a^*c + ac^*) \leq 0$. \square

In our disordered system, we consider the dipolar transfer matrices constructed by $\mathcal{M} = T_+T_-$, $\mathcal{N} = T_-T_+$ with

$$T_{\pm} = \begin{pmatrix} E \pm iV & -1 \\ 1 & 0 \end{pmatrix}, \quad (20)$$

leading to

$$\mathcal{M} = \begin{pmatrix} -1 + E^2 + V^2 & -E - iV \\ E - iV & -1 \end{pmatrix}, \quad (21)$$

$$\mathcal{N} = \begin{pmatrix} -1 + E^2 + V^2 & -E + iV \\ E + iV & -1 \end{pmatrix}, \quad (22)$$

which satisfies Condition 1 by construction.

Condition 3 leads to the parametric constraint, $-4E^2 + E^4 + 2E^2V^2 + V^4 \leq 0$, corresponding to the region inside the black line as shown in Fig. 1. It indeed naturally satisfies Condition 2, which corresponds to the region $E^2 + V^2 \in [0, 4]$. Therefore, for our protocol, we only require Condition 3 $\text{tr}(\mathcal{M}\mathcal{N}) \leq 2$.

Supplementary Material
for “Non-Hermitian delocalization in 1D via emergent compactness”

CONTENTS

SM 1. Fraction of delocalized states with real energy	8
SM 2. D_2 symmetry of the spectrum	8
A. Chiral symmetry	8
B. Symmetric energy pairs $\pm E$ in the spectrum	9

SM 1. FRACTION OF DELOCALIZED STATES WITH REAL ENERGY

The fraction of delocalized eigenstates depends on the specific spatial random realization. In Fig. 1 (c), the fraction is obtained by averaging over a large number of random realizations. This fraction quickly converges when we increase the size of the random ensemble, as shown in Fig. S1. In particular, for the periodic boundary condition $K = 0$ (blue), the fraction is around 10^{-4} , a very small but non-vanishing value.




FIG. S1. Fraction of the delocalized eigenstates with real energy. The fraction quickly converges for an increasing number of random realizations. For the periodic boundary condition, $K = 0$, this fraction is very small but non-zero. Here we use $V = 0.3$ and system size 1024, the same as Fig. 1(c) in the main text.

SM 2. D_2 SYMMETRY OF THE SPECTRUM

In this section, we show that the spectrum has a D_2 symmetry, namely, the mirror symmetry w.r.t. both real and imaginary axes.

A. Chiral symmetry

We first focus on the chiral symmetry, which ensures the coexistence of the energies E and $-E^*$ in the spectrum. Consider the single-particle Hamiltonian matrix H of Eq. 8. We now show that this Hamiltonian satisfies the following chiral symmetry

$$PH^\dagger P^{-1} = -H, \tag{S.1}$$

Proof. The explicit form of matrices \mathcal{M} and \mathcal{N} are

$$\mathcal{M} = \begin{pmatrix} -1 + E^2 + V^2 & -E - iV \\ E - iV & -1 \end{pmatrix}, \mathcal{N} = \begin{pmatrix} -1 + E^2 + V^2 & -E + iV \\ E + iV & -1 \end{pmatrix}. \quad (\text{S.8})$$

We expand them into Pauli matrices

$$\frac{-2 + E^2 + V^2}{2}\sigma_0 + \frac{E^2 + V^2}{2}\sigma_3 \pm iV\sigma_1 - iE\sigma_2 \equiv G_0\sigma_0 + G_3\sigma_3 \pm G_1\sigma_1 + G_2\sigma_2, \quad (\text{S.9})$$

where $-$ and $+$ correspond to \mathcal{M} and \mathcal{N} , respectively. Here $G_{i=0,1,2,3}$ denote the coefficient of the four matrices. Crucially, we note that G_0 and G_3 are polynomial functions of E^2 and even powers of $G_{i=1,2}$ are monomials of E^2 . For example, G_1^2 is V^2 , which is a monomial of zero power of E^2 . This observation turns out to be important for our argument, as further demonstrated below.

Now we consider a random product of \mathcal{M} and \mathcal{N} , e.g.,

$$\mathcal{M}_1\mathcal{M}_2\mathcal{N}_3\mathcal{M}_4 \dots \mathcal{N}_\alpha \dots \mathcal{M}_k, \quad (\text{S.10})$$

where we use the subindex $\alpha = 1, 2, 3, 4 \dots$ to track the position of each matrix. In terms of Pauli matrices, it becomes

$$\sum_{i_1=0}^3 C_{i_1}^1 \sigma_{i_1} \sum_{i_2=0}^3 C_{i_2}^2 \sigma_{i_2} \dots \sum_{i_\alpha=0}^3 C_{i_\alpha}^\alpha \sigma_{i_\alpha} \dots \sum_{i_k=0}^3 C_{i_k}^k \sigma_{i_k}. \quad (\text{S.11})$$

According to Eq. S.9, randomness only occurs in C_1^α which can take values of $\pm iV$, and $C_{0,2,3}^\alpha$ are fixed for all α . The expanded result of the polynomial multiplication is a linear combination of k -product of Pauli matrices, e.g.,

$$\sum_{\alpha=1}^k \sum_{i_\alpha=0}^3 C_{i_1}^1 \sigma_{i_1} C_{i_2}^2 \sigma_{i_2} \dots C_{i_\alpha}^\alpha \sigma_{i_\alpha} \dots C_{i_k}^k \sigma_{i_k}. \quad (\text{S.12})$$

We are going to prove that the summation of these 4^k terms, each being the k -product of Pauli strings, always has a trace that is a polynomial function of E^2 .

Utilizing the (anti-)commutation relations among Pauli matrices, each element in Eq. S.12 can be simplified as follows:

$$C_{i_1}^1 \sigma_{i_1} C_{i_2}^2 \sigma_{i_2} \dots C_{i_\alpha}^\alpha \sigma_{i_\alpha} \dots C_{i_k}^k \sigma_{i_k} = c_0 (G_0\sigma_0)^t (G_1\sigma_1)^p (G_2\sigma_2)^q (G_3\sigma_3)^l = c_0 G_0^t (G_1\sigma_1)^p (G_2\sigma_2)^q (G_3\sigma_3)^l, \quad (\text{S.13})$$

where t, p, q, l refer to the total number of different Pauli matrices corresponding to a given string $\{i_1, i_2, \dots, i_k\}$, satisfying $p + q + l = k - t$. The overall prefactor c_0 is either 1 or -1 , originating from the anti-commutation of two Pauli matrices as well as the randomness of the appearance of σ_1 in \mathcal{M} and \mathcal{N} .

A crucial observation is that any string $\{i_1, i_2, \dots, i_k\}$ containing at least one σ_2 and one σ_3 must have its unique counterpart in the expansion given by Eq. S.12. This counterpart is identified by locating the first pair of σ_2 and σ_3 from left to right in the sequence and swapping these two elements, regardless of whether they are adjacent. These two sequences share identical coefficients, but their permutation parities are opposite, resulting in mutual cancellation.

To illustrate, we can count the occurrences of σ_2 and σ_3 from left to right across the string and track their cumulative numbers. We define a number n as the position in the string where, for the first time, we encounter one σ_2 and σ_3 , such that the elements before n include either zero σ_2 or zero σ_3 .

If the n th term is σ_3 , then we swap it with the nearest σ_2 in the preceding $n - 1$ terms; if the n th term is σ_2 , then we swap it with the nearest σ_3 in the former $n - 1$ terms. After the swap, the resulting string will be a unique counterpart of the original.

This term naturally exists in the expansion (Eq. S.12) because of the distributive property in the polynomial multiplication.

For example, consider the string $\{i_1 = 1, i_2 = 2, i_3 = 1, i_4 = 3, i_5 = 2\}$ with $k = 5$. We proceed with the following steps:

- Counting each term, we notice:

$$\begin{aligned} C_{i_1=1}^1 \sigma_1 C_{i_2=2}^2 \sigma_2 & \quad (1 \sigma_2, 0 \sigma_3) \\ C_{i_1=1}^1 \sigma_1 C_{i_2=2}^2 \sigma_2 C_{i_3=1}^3 \sigma_1 & \quad (1 \sigma_2, 0 \sigma_3) \\ C_{i_1=1}^1 \sigma_1 C_{i_2=2}^2 \sigma_2 C_{i_3=1}^3 \sigma_1 C_{i_4=3}^4 \sigma_3 & \quad (1 \sigma_2, 1 \sigma_3) \end{aligned}$$

- In the fourth term, both a σ_2 and a σ_3 are present, meeting the criteria set forth above. Thus, we swap the σ_3 at the fourth position with the nearest σ_2 , which is at the second position.
- The resultant string after the swap is $\{i_1 = 1, i_2 = 3, i_3 = 1, i_4 = 2, i_5 = 2\}$ and its explicit form is $C_{i_1=1}^1 \sigma_1 C_{i_2=2}^2 \sigma_3 C_{i_3=1}^3 \sigma_1 C_{i_4=3}^4 \sigma_2 C_{i_5=2}^5 \sigma_2$, which is contained in the expansion of the $k = 5$ -terms polynomial multiplication in Eq. S.12. Also, it has the same coefficient, $C_{i_1=1}^1 C_{i_2=2}^2 C_{i_3=1}^3 C_{i_4=3}^4 C_{i_5=2}^5$, as the original string.

Now we demonstrate that swapping the nearest σ_2 and σ_3 in the sequence results in opposite permutation parities. The closest pair of σ_2 and σ_3 appears in two ways, either with an even or odd number of σ_1 matrices in between. In the even case, this simplifies to the swapping of the first adjacent pair of σ_2 and σ_3 , which naturally yields opposite permutation parities. For the other case, one only needs to consider the permutation parities of $\sigma_2\sigma_1\sigma_3$ and $\sigma_3\sigma_1\sigma_2$, which are also opposite.

Therefore, any string $\{i_1, i_2, \dots, i_k\}$ that contains at least one σ_2 and one σ_3 does not contribute to the trace of Eq. S.11. We only need to study Pauli strings without σ_2 and σ_3 appearing at the same time, i.e. either $q = 0$ or $l = 0$.

Case 1: $l = 0$

In this case, if both p and q are even, the trace is a polynomial of E^2 . Conversely, if either p or q is odd, the trace vanishes. Specifically:

- If only one of p or q is odd, then the trace of Eq. S.13 becomes proportional to either $\text{tr}(\sigma_1)$ or $\text{tr}(\sigma_2)$, which vanishes.
- If both p and q are odd, the trace is proportional to $\text{tr}(\sigma_3)$, which is also zero.

Case 2: $q = 0$

The analysis mirrors that of Case 1, where terms with even p and l survive, and their traces are polynomials of E^2 .

In summary, only terms with even powers of σ_1 and $\sigma_2(\sigma_3)$ contribute to the trace of Eq. S.12, thus the trace is a polynomial of E^2 . \square

In conclusion, we prove that the trace of any random multiplication of \mathcal{M} and \mathcal{N} is a polynomial of E^2 . In other words, E and $-E$ must coexist in the spectrum. This symmetry and the chiral symmetry result in the D_2 symmetry in the spectrum.