

PARTITIONS OF GRAPHS INTO SPECIAL BIPARTITE GRAPHS

LAJOS GYÖRFFY, ANDRÁS LONDON, GÁBOR V. NAGY, AND ANDRÁS PLUHÁR

ABSTRACT. We study the problem of partitioning the edge set of a graph into bipartite subgraphs under certain constraints defined by forbidden subgraphs. These constraints lead to both classical, such as partitioning into independent matchings and complete bipartite subgraphs, and novel partitioning problems. The theoretical framework we provide is motivated by clustering problems of real-world transaction graphs, which can naturally be formulated as edge partitioning problems under certain bipartite graph constraints.

1. INTRODUCTION AND RESULTS

A classical result of Graham and Pollak [7] shows that at least $n - 1$ complete bipartite graphs are needed to partition the edge set of the complete graph K_n (and it is easy to see that $n - 1$ complete bipartite graphs are enough). A natural generalization have led to the *biclique partitioning problem*, where the goal is to partition the edges of a graph G with the minimum number of bicliques. More generally, given a *host* graph G , partition its edges into subgraphs belonging to a given set, called the template class. Elements of the template class are called template graphs, or simply *templates* [11]. In the *covering* version of the problem, every edge of the host graph must belong to at least one (but not necessarily exactly one, as in the case of partitioning) template. It is easy to see that $\lceil \log_2 n \rceil$ bicliques are sufficient to cover K_n (cf. the aforementioned result of Graham and Pollak), showing that the gap between the partitioning and covering numbers for the same host graph can be significant. For a detailed overview, see [19].

In this study, we consider the graph partitioning problem under the constraint that the template class is defined by forbidden subgraphs. Specifically, we seek to partition the edge set of a given graph G into subgraphs that avoid certain forbidden configurations. We investigate how different sets of forbidden subgraphs impact the partitioning structure and the minimum number of subgraphs required to partition the edges of G .

Key words and phrases. Graph partition, Graham-Pollak, Ferrers graphs, nestedness, forbidden induced subgraphs.

The research leading to these results has received funding from the national project TKP2021-NVA-09. Project no. TKP2021-NVA-09 has been implemented with the support provided by the Ministry of Culture and Innovation of Hungary from the National Research, Development and Innovation Fund, financed under the TKP2021-NVA funding scheme.

Our main motivation for this theoretical framework stems from applications in clustering real-world transaction graphs, where similar constraints naturally arise.

Graph clustering and community detection play a crucial role in graph-based data mining and in the development of models on graphs; see for example [3, 17, 16, 18]. Motivated by applications such as social networks, these clustering methods aim to identify dense subgraphs and partition the graph into meaningful communities. While these methods often perform well in practice, defining a theoretically perfect clustering remains a challenging and unsolved problem, as discussed by Kleinberg [10].

Unlike clustering the nodes, edge partitioning, or edge-based clustering, allows each node to participate in multiple clusters, reflecting the fact that a single entity can belong to different groups depending on the context. This is particularly realistic in economic networks, where, for example, a company might interact with different industries through distinct transactions, such as being a supplier in one cluster and a customer in another. Generally, in the case of so-called *technological graphs* edge density based node clustering methods are not satisfactory. The background and examples are explained in [9, 14, 20], which motivated the development of completely different approaches to clustering. Here we only recall some facts that are essential for understanding. Certain social networks, and especially technological or transaction networks, tend to have fewer triangles and often exhibit tree-like structures [1], making disjoint dense clustering inherently inadequate. In addition, certain bipartite networks, such as pollination networks (of plant and pollinating animal species) or trade networks (of countries and imported/exported goods), often display special structures like *nestedness* [2, 20, 22]. In these networks, nodes can be ordered so that the neighborhood of each lower-ranked node contains that of any higher-ranked node. Ecological networks commonly exhibit this nested pattern, where specialists (lower-ranked) interacting with generalists (higher-ranked), which in turn interact with both specialists and other generalists.

In a former model, described in [14], the authors consider special *good (vertex) colorings* of a graph, where the edge structure between any two color classes is restricted. A particularly important case is when the bipartite subgraphs between color classes are induced $2K_2$ -free. Some of the special colorings with a minimum number of colors can be found in polynomial time, however, deciding cases like the $2K_2$ -free bipartite graphs is generally NP-complete.

A more application-driven approach is explored by Gera et al. [5, 6], where the goal is to find a covering of a graph such that the elements of the covering are not necessarily disjoint $2K_2$ -free bipartite graphs. In this paper, we propose a model that lies between the purely theoretical and the highly application-oriented approaches.

After the historical overview, we turn to the discussion of the edge partitioning results of this paper. We begin with formal definitions and notations. It is straightforward to check that induced $2K_2$ -free bipartite graphs are precisely the so-called Ferrers graphs.

Definition 1. A bipartite graph with color classes A and B is called a Ferrers graph if there exists an ordering a_1, \dots, a_n of the vertices in A such that $N(a_1) \subseteq N(a_2) \subseteq \dots \subseteq N(a_n)$ holds, where $N(a_i)$ denotes the neighborhood of a_i . (This condition implies the same “nestedness” for the neighborhoods of vertices of B as well.)

Ferrers graphs are named for the property that the 1’s in their bipartite adjacency matrix form a Ferrers diagram (of an integer partition) when the rows and columns are arranged in “degree-decreasing” order.

Definition 2. Fix a small graph H . An H -avoiding bipartite partition of a graph G is a set of bipartite graphs G_1, \dots, G_k such that $E(G) = \cup_{i=1}^k E(G_i)$, $E(G_i) \cap E(G_j) = \emptyset$ if $i \neq j$ and G_i does not contain H as an induced subgraph for all i . (The graphs G_1, \dots, G_k are also called template graphs.)

Notation 1. If we have more than one graph H (i.e. H_1, H_2, \dots), then we avoid a set of small graphs $\mathcal{H} = \{H_1, H_2, \dots\}$ among the graphs G_i and we write \mathcal{H} instead of H .

Of course, we want to have as few graphs as possible in an H -avoiding bipartite partition.

Notation 2. For a fixed graph H and G , we denote by $\chi'_H(G)$ the smallest integer k for which there exists an H -avoiding bipartite partition of G consisting of k graphs. We also use the notation $\chi'_H(n) := \chi'_H(K_n)$. For a set \mathcal{H} of forbidden subgraphs, the notations $\chi'_{\mathcal{H}}(G)$ and $\chi'_{\mathcal{H}}(n)$ are defined analogously.

Observation 1. For any fixed set \mathcal{H} of forbidden subgraphs, the sequence $(\chi'_{\mathcal{H}}(n))_{n=1}^{\infty}$ is monotonically increasing, that is, $\chi'_{\mathcal{H}}(n) \leq \chi'_{\mathcal{H}}(n+1)$ for all n .

Proof. Consider an (optimal) \mathcal{H} -avoiding bipartite partition G_1, \dots, G_t of K_{n+1} where $t = \chi'_{\mathcal{H}}(n+1)$. Let S be an n -element subset of $V(K_{n+1})$. Then the induced subgraphs $G_1|_S, \dots, G_t|_S$ clearly form an \mathcal{H} -avoiding bipartite partition of $K_{n+1}|_S \simeq K_n$ (empty graphs may appear here), proving that $\chi'_{\mathcal{H}}(n) \leq t = \chi'_{\mathcal{H}}(n+1)$. \square

At the end of this section in Figure 1 we list the small excluded subgraphs we will consider. We use K_n, P_n, C_n and S_n for the complete graph, path, cycle and star on n vertices, respectively. For graphs H_1 and H_2 on disjoint vertex sets, $H_1 + H_2$ denotes their disjoint union.

2. CLASSIFICATION OF SMALL EXCLUDED SUBGRAPHS

First we study the case $G = K_n$, to motivate more general problems and to show the difficulty of some examples even in this case.

Convention 1. In this section, we consider only bipartite partitions G_1, \dots, G_k in which no template graph has isolated vertices, and we also assume that none of the template graphs is empty (as we are interested in partitions with a minimal number of template graphs).




FIGURE 1. Small excluded subgraphs

Our most important case among the possible applications here will be the case $H = 2K_2$, but we will also consider other excluded small graphs H (for the sake of mathematical completeness, as seen by Král et al. in [12]). Some of the problems are related to other well-known problems, some are very easy or trivial, but some could be also hard. The values of $\chi'_H(n)$ come from the $[\log_2 n, \binom{n}{2}]$ interval.

2.1. Partitioning into bipartite graphs without restriction. It is a folklore fact that K_n cannot be partitioned into less than $\log_2 n$ (arbitrary) bipartite graphs, and $\lceil \log_2 n \rceil$ bipartite graphs are enough. (We also sketch the proof in the subsection ‘The proof of the lower bound in Theorem 9’.)

Observation 2. $\chi'_\emptyset(n) = \lceil \log_2 n \rceil$.

Observation 3. $\chi'_\mathcal{H}(n) \geq \lceil \log_2 n \rceil$, for any set \mathcal{H} of forbidden subgraphs.

2.2. P_3 -free bipartite graphs are matchings. If $H = P_3$, then the maximum degree in each template graph must be (at most) 1. Therefore, G_i is a matching for all $1 \leq i \leq k$. Therefore, the number of graphs in a minimal P_3 -avoiding bipartite partition of G is just the edge chromatic number of G , which is equal to $\Delta(G)$ or $\Delta(G) + 1$ by Vizing’s theorem (where $\Delta(G)$ denotes the maximum degree of G). The edge chromatic number of the complete graph $G = K_n$ is well known, which leads to the following.

Observation 4. $\chi'_{P_3}(n) = n - 1$, if n is even; and $\chi'_{P_3}(n) = n$, if n is odd.

Remark 1. The notation $\chi'_\mathcal{H}(G)$ comes from the fact, that $\chi'_{P_3}(G) = \chi'(G)$ for all graphs G , where $\chi'(G)$ is the chromatic index of the graph G .

2.3. $(K_2 + K_1)$ -free bipartite graphs are complete bipartite graphs. If $H = K_2 + K_1$ then there cannot be a single point independent of any other edge. Therefore, G_i is complete for all $1 \leq i \leq k$ and the number of graphs in a minimal H -avoiding bipartite partition of G gives the well-known Graham-Pollak theorem.

Theorem 5 (Graham–Pollak [7]). *There are at least $n - 1$ subgraphs in a complete bipartite partition of K_n (and $n - 1$ subgraphs are enough), that is, $\chi'_{K_2+K_1}(n) = n - 1$.*

Most of the proofs of this theorem use linear algebraic methods. No purely combinatorial proof is known (except a sophisticated one by Vishwanathan [21], which can be seen as a translation of the linear algebraic method). Even for weaker lower bounds, e.g. \sqrt{n} , we do not know any nice combinatorial proof.

2.4. $\{K_2 + K_1, P_3\}$ -free bipartite graphs are single edges. If $\mathcal{H} = \{K_2 + K_1, P_3\}$ then it is easy to see that the template graphs G_i are single edges for all $1 \leq i \leq k$. That is, the number of graphs in a minimal H -avoiding bipartite partition of G is just the number of edges $\binom{n}{2}$.

Observation 6. $\chi'_{\{K_2+K_1, P_3\}}(n) = \binom{n}{2}$.

We have listed all possible cases among forbidden graphs with three vertices. Now we consider all subsets \mathcal{H} of the set of 4-vertex excluded subgraphs shown in Figure 1.

2.5. P_4 -free bipartite graphs are disjoint complete graphs. If $H = P_4$, then it is straightforward to check that G_i is a vertex disjoint collection of complete bipartite graphs for all $1 \leq i \leq k$. If we divide the complete graph into two (almost) equal sets and draw all edges between the two color classes in G_1 , then there are $n^2/4$ edges and in G_2 we get two pieces of complete graphs with $n/2$ vertices where we can continue the division. (We do not require connectedness, so we can include more bipartite graphs in a G_i .) It shows that if n is a power of 2, then we can always divide the remaining graph into two parts, which yields $\log_2 n$ template graphs; and so $\chi'_{P_4}(n) = \log_2 n$ by Observation 3. In any other cases, we can bound $\chi'_{P_4}(n)$ by $\chi'_{P_4}(2^{\lceil \log_2 n \rceil})$ using the monotonicity of $\chi'_{P_4}(n)$.

Observation 7. $\chi'_{P_4}(n) = \lceil \log_2 n \rceil$.

If the graphs of the partition are required to be connected, we come to the assumption of the Graham-Pollak theorem.

2.6. C_4 -free bipartite graphs. If $H = C_4$, then G_i is C_4 -free bipartite graph for all $1 \leq i \leq k$. Since $|E(G_i)| \leq c_1 n^{3/2}$ (see Zarankiewicz[23]), we immediately get a lower bound $c_2 n^{1/2} \leq \chi'_{C_4}(n)$.

However, the upper bound (so is the problem) is still open. If we throw the point-line graph of large independent projective planes onto the complete graph K_n k times, then the probability that an edge is uncovered is about $(1 - c/\sqrt{n})^k$. If some edges are over covered (covered more than once), then we simply delete the surplus from the graphs after the first cover. Set $k = (3/c) \cdot \sqrt{n} \log n$. The Boole inequality readily gives the following, probably not sharp, upper bound: $\chi'_{C_4}(n) < \log n \cdot \sqrt{n}$.

$$\Pr(\text{edge } e \text{ is not covered}) \leq (1 - c/\sqrt{n})^k \leq e^{-kc/\sqrt{n}} = n^{-3}.$$

Then

$$\Pr(\text{some edge from } E(K_n) \text{ is not covered}) \leq \sum_1^{\binom{n}{2}} n^{-3} < n^{-1} < 1,$$

provided n is big enough.

Theorem 8. *For some positive constants a and b ,*

$$a\sqrt{n} \leq \chi'_{C_4}(n) \leq b\sqrt{n} \log n,$$

if n is large enough.

We think the next subsection is the most intriguing part of this section, which motivates that these types of questions can be difficult. Here the lower and upper bounds are very far apart, and the fact that there is no easy combinatorial proof of the Graham-Pollak theorem also shows the difficulty of this problem. However, it gives us a remarkable open problem.

2.7. $2K_2$ -free bipartite graphs are Ferrers graphs. If $H = 2K_2$, then each G_i is a Ferrers graph. The next result gives a non-trivial lower and upper bound on $\chi'_{2K_2}(n)$.

Theorem 9. $\lceil \log_2 n \rceil + \frac{1}{4}\sqrt{\lceil \log_2 n \rceil} - 1 < \chi'_{2K_2}(n) \leq 2 \lceil \sqrt{n} \rceil - 2$.

Theorem 9 is proved in the next section.

2.8. S_4 -free bipartite graphs. If $H = S_4$, then the maximal degree of each G_i is at most 2. Therefore, we need at least $\lceil (n-1)/2 \rceil$ template graphs. From the other side, Walecki [15] showed that K_n can be decomposed into $(n-1)/2$ Hamiltonian cycles for odd n , which yields the upper bound $(n-1)/2$. So the exact result is $\chi'_{S_4}(n) = \lceil (n-1)/2 \rceil$. (For even n , the upper bound follows from the inequality $\chi'_{S_4}(n) \leq \chi'_{S_4}(n+1)$.)

Observation 10. $\chi'_{S_4}(n) = \lceil (n-1)/2 \rceil$.

Remark 2. *The existence of a partition into Hamilton cycles in arbitrary graphs is an NP-complete problem. Even in 4-regular graphs.*

2.9. $\{2K_2, C_4\}$ -free bipartite graphs are double stars. If we avoid $2K_2$ and C_4 together, then our subgraphs G_i are connected and do not contain any induced cycle, because C_6, C_8, \dots contain induced $2K_2$. Hence, the template graphs are trees with $n-1$ edges and we get the lower bound $\lceil n/2 \rceil$.

From the other side, if we use double stars partitioning K_n , this is enough, as we can see in Figure 2 for even n . (The exact upper bound for odd n follows from the monotonicity of $\chi'_{\{2K_2, C_4\}}(n)$ again.)

Theorem 11. $\chi'_{\{2K_2, C_4\}}(n) = \lceil n/2 \rceil$.




FIGURE 2. Double star partition of K_6 .

2.10. $\{2K_2, C_4, P_4\}$ -free bipartite graphs are stars. If we also omit P_4 , then double stars are not allowed. What remains is a star decomposition which comes from the well-known solution of the Graham-Pollak theorem, and gives $n - 1$ as best result.

2.11. $\{C_4, P_4, S_4\}$ -free bipartite graphs are cherry orchards. If P_4, S_4, C_4 are the excluded subgraphs and the G_i graphs are bipartite, then the maximum degree is two and contains no cycles of any length in G_i . Therefore, each G_i consists of single edge and P_3 components, that is, the G_i graphs are a set of disjoint cherries (we will call them *cherry orchards*), for all i . One can see a single cherry orchard in Figure 3.




FIGURE 3. Cherry Orchard.

In a cherry orchard there are at most $\frac{2}{3}n$ edges. The lower bound to $\chi'_{\mathcal{H}}(n)$ comes from counting the edges: $\binom{n}{2} / \frac{2n}{3} \approx \frac{3}{4}n$.

To see the other side, we reach the $\frac{3}{4}n + o(n)$ upper bound in the following way. We divide the vertices of K_n into three (almost) equal parts, let us call them A, B and C , all containing $n/3$ vertices. The edges between each pair of sets can be divided into $n/3$ perfect matchings. We divide these $3 \cdot \left(\frac{n}{3}\right)^2$ edges into cherry orchards. First, we consider three cherry orchards, each with $\frac{2n}{3}$ edges. The first is the union of a perfect matching from (A, B) and (A, C) . The other two are the unions of perfect matchings from (B, A) and (B, C) ; and from (C, A) and (C, B) , respectively. Then we remove these three cherry orchards, and all degrees of vertices in the sets decreased by exactly four. We do the same in the next steps, where each step consists of three cherry orchards, until we run out of edges. With this procedure three cherry orchards decreases the degrees of vertices by four and we started with $\frac{2n}{3}$ degrees at each vertex

at the beginning. So with $\frac{2n/3}{4} \cdot 3 = \frac{n}{2}$ cherry orchards we covered all the edges between the parts.

Then we use induction on the sets A, B and C . Note, that this can be done in parallel. So, if $T(n)$ denotes the number of cherry orchards needed, then $T(n) \leq \frac{1}{2}n + T(\frac{n}{3}) = \frac{1}{2}n + \frac{1}{6}n + T(\frac{n}{9}) = (\frac{1}{2} + \frac{1}{6} + \frac{1}{18} + \dots) n = \frac{n}{2} (1 + \frac{1}{3} + \dots) = \frac{3}{4}n + o(n)$.

Theorem 12 (Cherry Orchard). *If $\mathcal{H} = \{P_4, C_4, S_4\}$, then $\chi'_{\mathcal{H}}(n) = \frac{3}{4}n + o(n)$.*

2.12. $\{C_4, P_4\}$ -free bipartite graphs are disjoint stars (star orchards). However if we do not require connectedness or maximum degree at most 2 (since we do not forbid $2K_2$ or S_4), then each template graph G_i can consist of arbitrarily many star components with arbitrary degrees. A trivial lower bound $\lceil \frac{n}{2} \rceil$ comes from the fact that the template graphs contain at most $n - 1$ edges.

Assume first that n is even. An upper bound can be derived from the former double star construction (see Figure 2): If we delete the central edge of each double star, then we have two disjoint stars in each graph G_i without any induced P_4 . The remaining extra edges form a matching which can be considered as a set of disjoint stars and the last template graph in the partition. This gives the upper bound $\frac{n}{2} + 1$. In fact, this is the exact value of $\chi'_{\{P_4, C_4\}}(n)$ for even n : Observe that at most one (star orchard) template graph can have $n - 1$ edges, i.e. at most one template graph can be a single star on n vertices, because the center of one such star cannot be contained in any other template graph. This means that the trivial lower bound $\frac{n}{2}$ cannot be sharp for even n , if $n \geq 4$, because each template would have exactly $n - 1$ edges in a star orchard partition of K_n with $\frac{n}{2}$ template graphs. So $\chi'_{\{P_4, C_4\}}(n) = \frac{n}{2} + 1$ for even $n \geq 4$.

It turns out that $\chi'_{\{P_4, C_4\}}(n) = \lceil \frac{n}{2} \rceil + 1$ holds for odd $n \geq 5$ as well. This case can be reduced to the case of even n . If n is odd, then a minimal star orchard partition of K_{n-1} can be extended to a star orchard partition of K_n with $\frac{n-1}{2} + 2 = \lceil \frac{n}{2} \rceil + 1$ template graphs by adding an n -vertex single star template. $\lceil \frac{n}{2} \rceil$ templates are not enough. This is because K_n cannot be partitioned into $\lceil \frac{n}{2} \rceil$ star orchards such that each template graph has at most $n - 2$ edges, as then the total number of edges in the templates would be at most

$$\lceil \frac{n}{2} \rceil \cdot (n - 2) = \frac{n + 1}{2} \cdot (n - 2) < \frac{n(n - 1)}{2} = E(K_n).$$

And for odd $n \geq 5$, K_n cannot be partitioned into $\lceil \frac{n}{2} \rceil$ star orchards such that one of them is a single n -vertex star, because the removal of the n -vertex star template would yield a star orchard partition of K_{n-1} with $\lceil \frac{n}{2} \rceil - 1 = \frac{n-1}{2}$ templates, which is impossible as seen in the previous paragraph.

So we obtained that

Theorem 13. *If $\mathcal{H} = \{P_4, C_4\}$, then $\chi'_{\mathcal{H}}(n) = \lceil \frac{n}{2} \rceil + 1$, for $n \geq 4$.*

The discussion of further examples will be less detailed. For example, we only include results where n meets some “natural” divisibility conditions. (The omitted cases are left to the reader as exercises.)

2.13. $\{P_4, 2K_2\}$ -free bipartite graphs, **Graham-Pollak again.**

Observation 14. *If $\mathcal{H} = \{P_4, 2K_2\}$, then we get back the Graham-Pollak theorem, hence $\chi'_{\mathcal{H}}(n) = n - 1$.*

2.14. $\{2K_2, S_4\}$ -free bipartite graphs are only C_4, P_4, P_3 or K_2 . The trivial lower bound can be attained by partitioning K_n into C_4 's (with divisibility conditions on n), see [13].

Observation 15. *If $\mathcal{H} = \{S_4, 2K_2\}$, then $\chi'_{\mathcal{H}}(n) = \frac{n(n-1)}{8}$, when $n \equiv 1 \pmod{8}$.*

2.15. $\{2K_2, S_4, P_4\}$ -free bipartite graphs are only C_4, P_3 or K_2 . An optimal partition is given by the same construction as above (for the same divisibility condition).

Observation 16. *If $\mathcal{H} = \{S_4, 2K_2, P_4\}$, then $\chi'_{\mathcal{H}}(n) = \frac{n(n-1)}{8}$, when $n \equiv 1 \pmod{8}$.*

2.16. $\{C_4, S_4\}$ -free bipartite graphs. In this case the maximum degree is at most 2 in each template (and there are no C_4 components). The Hamilton cycle decomposition of K_n is optimal here, as in case of S_4 -free partitioning.

Observation 17. *If $\mathcal{H} = \{C_4, S_4\}$, then $\chi'_{\mathcal{H}}(n) = \lceil \frac{n-1}{2} \rceil$, when $n > 4$.*

2.17. $\{P_4, S_4\}$ -free bipartite graphs are C_4 orchards (and their induced subgraphs). Danzinger et al. showed that if n is divisible by 4, then K_n can be partitioned into one perfect matching and $n/2 - 1$ graphs $(n/4)C_4$, where $(n/4)C_4$ denotes the vertex-disjoint union of $n/4$ cycles C_4 ; see Theorem 2.2 in [4]. This is a minimal $\{P_4, S_4\}$ -free bipartite partition of K_n with $n/2$ templates.

Observation 18. *If $\mathcal{H} = \{P_4, S_4\}$, then $\chi'_{\mathcal{H}}(n) = \frac{n}{2}$, when n is divisible by 4.*

2.18. $\{C_4, S_4, 2K_2\}$ -free bipartite graphs are only the paths P_4, P_3 or K_2 . In this case an optimal partition can be obtained from the Hamilton cycle decomposition of K_n , by cutting each Hamilton cycle into $n/3$ P_4 's (with divisibility conditions on n).

Observation 19. *If $\mathcal{H} = \{C_4, S_4, 2K_2\}$, then $\chi'_{\mathcal{H}}(n) = \frac{n(n-1)}{6}$, when $n \equiv 3 \pmod{6}$.*

2.19. $\{C_4, P_4, S_4, 2K_2\}$ -free bipartite graphs are cherries. It is a known corollary of Tutte's theorem on perfect matchings that every connected graph with an even number of edges can be decomposed into cherries P_3 . Hence the trivial lower bound is sharp.

Observation 20. *If $\mathcal{H} = \{P_4, C_4, S_4, 2K_2\}$, then $\chi'_{\mathcal{H}}(n) = \lceil \frac{n(n-1)}{4} \rceil$.*

3. THE PROOF OF THEOREM 9.

The proof of the upper bound. We construct a ($2K_2$ -free bipartite) partition of K_n consisting of at most $2\lceil\sqrt{n}\rceil - 2$ Ferrers graphs, for any $n \geq 1$.

First we assume that n is a square number, that is, $\lceil\sqrt{n}\rceil = \sqrt{n}$ holds. In this case we can also assume that the vertex set of K_n is the set $\mathcal{V} := \{1, \dots, \sqrt{n}\} \times \{1, \dots, \sqrt{n}\}$. We divide the edge set of K_n into two classes as follows. Let e be an edge of K_n between the vertices (x, y) and (x', y') , where $x \leq x'$. We say that e is a *descent* if $x < x'$ and $y \geq y'$. Otherwise we say that e is an *ascent*, i.e. e is an ascent if $x < x'$ and $y < y'$, or $x = x'$ (and $y \neq y'$). See also Figure 4. (If the vertices are regarded as points in the Cartesian plane and the edges are drawn as straight line segments, then the sign of the slope of an edge e determines whether e is a descent or an ascent.) Note that by our definition, “horizontal” edges are descents and “vertical” edges are ascents. By the *left endpoint* of a descent edge e we mean the (unique) endpoint of e which has smaller first coordinate; by the *lower endpoint* of an ascent edge f we mean the (unique) endpoint of f which has smaller second coordinate.




FIGURE 4. A descent edge e and an ascent edge f in K_{25}

After these preliminaries, we define $2\sqrt{n} - 2$ Ferrers graphs, $G_1, G_2, \dots, G_{\sqrt{n}-1}, G'_1, G'_2, \dots, G'_{\sqrt{n}-1}$, which admits a bipartite partition of K_n . For $i = 1, \dots, \sqrt{n} - 1$, let G_i be the bipartite graph with color classes $A_i = \{(x, y) \in \mathcal{V} : x = i\}$ and $B_i = \{(x, y) \in \mathcal{V} : x > i\}$ such that the vertices $a \in A_i$ and $b \in B_i$ are adjacent in G_i if and only if ab is a descent edge of K_n . For $j = 1, \dots, \sqrt{n} - 1$, let G'_j be the bipartite graph with color classes $A'_j = \{(x, y) \in \mathcal{V} : y = j\}$ and $B'_j = \{(x, y) \in \mathcal{V} : y > j\}$ such that the vertices $a \in A'_j$ and $b \in B'_j$ are adjacent in G'_j if and only if ab is an ascent edge of K_n .

First we check that these bipartite graphs are indeed Ferrers graphs. In a graph G_i , the neighborhood $N(i, y_0)$ of the vertex $(i, y_0) \in A_i$ is the set $\{(x, y) \in \mathcal{V} : x > i, y \leq y_0\}$; and so we actually have the nested property

$$N(i, 1) \subseteq N(i, 2) \subseteq N(i, 3) \subseteq \dots \subseteq N(i, \sqrt{n}).$$

FIGURE 5. Illustration of G_i and G'_j (for $n = 16$)

In a graph G'_j , the neighborhood $N(x_0, j)$ of the vertex $(x_0, j) \in A'_j$ is the set $\{(x, y) \in \mathcal{V} : x \geq x_0, y > j\}$; and so we have the nested property

$$N(1, j) \supseteq N(2, j) \supseteq N(3, j) \supseteq \cdots \supseteq N(\sqrt{n}, j)$$

again. Now all that remains is to show that the graphs $G_1, \dots, G_{\sqrt{n}-1}, G'_1, \dots, G'_{\sqrt{n}-1}$ partition K_n . Observe that for all $i = 1, \dots, \sqrt{n}-1$, the graph G_i contains all descent edges of K_n whose left endpoint has the first coordinate i , and for all $j = 1, \dots, \sqrt{n}-1$, the graph G'_j contains all ascent edges of K_n whose lower endpoint has the second coordinate j ; and all edges of K_n are contained in exactly one of these $2\sqrt{n} - 2$ edge classes. Thus we proved that $\chi'_{2K_2}(n) \leq 2\sqrt{n} - 2$ for all square numbers n .

If n is a non-square number, then consider the closest larger square number $n^* := (\lceil \sqrt{n} \rceil)^2$ (for which the previous case applies), and bound $\chi'_{2K_2}(n)$ using its monotonicity (see Observation 1):

$$\chi'_{2K_2}(n) \leq \chi'_{2K_2}(n^*) \leq 2\sqrt{n^*} - 2 = 2 \lceil \sqrt{n} \rceil - 2.$$

So the upper bound is proved for all positive integers n . \square

Before proving the lower bound, we present two required lemmas.

Lemma 21. *If a Ferrers graph F contains a set of m independent edges, then F has at least $m(m+1)/2$ edges.*

Proof. Let A and B be the two color classes of F . Now let $u_1v_1, u_2v_2, \dots, u_mv_m$ be m independent edges of F , where $u_1, \dots, u_m \in A$, $v_1, \dots, v_m \in B$. Since F is a Ferrers graph, we can assume without loss of generality that

$$N(u_1) \subseteq N(u_2) \subseteq \cdots \subseteq N(u_m).$$

This chain implies that u_iv_j is also an edge of F for all $1 \leq j \leq i \leq m$, cf. Figure 6. (The edge u_iv_i guarantees that $v_i \in N(u_i)$, for $i = 1, \dots, m$. Since $v_1 \in N(u_1) \subseteq N(u_2)$, thus u_2 is also adjacent to v_1 . Then $v_1, v_2 \in N(u_2) \subseteq N(u_3)$ implies that u_3 is adjacent to v_1 and v_2 ; and so on.) We have just found $1 + 2 + \cdots + m = m(m+1)/2$ distinct edges in F . \square

Lemma 22. *For fixed integers $k \geq 1$ and $d \geq 0$, assume that K_{2^k} , the complete graph on 2^k vertices, is partitioned into the bipartite graphs G_1, G_2, \dots, G_{k+d} . Then there*




FIGURE 6. Illustration of the proof of Lemma 21

exist (at least) k graphs in $\{G_1, \dots, G_{k+d}\}$ such that each contains a set of at least $\frac{2^{k-2}}{d+1}$ independent edges.

Proof. Let A_i and B_i be the two color classes of G_i , for $i = 1, \dots, k+d$. We can assume that $A_i \cup B_i = V(K_{2^k})$ by adding isolated vertices to the graph G_i if necessary. To each vertex v of K_{2^k} we assign a vector $\text{CL}(v) \in \{0, 1\}^{k+d}$, whose i^{th} coordinate is defined as 0, if $v \in A_i$, and it is defined as 1, if $v \in B_i$, for $i = 1, \dots, k+d$. We call the vector $\text{CL}(v)$ the *class-vector* of v . We mention some properties of class-vectors:

- (i) If $u \neq v$, then $\text{CL}(u) \neq \text{CL}(v)$.
- (ii) For fixed vertices $u \neq v$, let $S \subseteq \{1, \dots, k+d\}$ be the set of coordinates where $\text{CL}(u)$ and $\text{CL}(v)$ differ, and let G_i be the graph of the partition containing the edge uv (of K_{2^k}). Then the index i must be an element of S .

Both properties follow directly from the fact that if the edge $uv \in E(K_{2^k})$ is contained in the graph G_i of the partition, then $\text{CL}(u)$ and $\text{CL}(v)$ differ in coordinate i , because u and v are in different color classes of G_i . (In fact, property (i) is a corollary of (ii).)

The rest of the proof relies only on the information encoded in $\mathcal{C} := \{\text{CL}(v) : v \in V(K_{2^k})\}$, the set of class-vectors. Recall that $\mathcal{C} \subseteq \{0, 1\}^{k+d}$, and note that $|\mathcal{C}| = |V(K_{2^k})| = 2^k$ by property (i) above. From now on, we identify the vertices of K_{2^k} with their class-vectors, and so $V(K_{2^k}) = \mathcal{C}$ with a slight abuse of notation. We say that a graph G is *large* if it contains a set of at least $\frac{2^{k-2}}{d+1}$ independent edges; otherwise we say that G is *small*. We need to prove that there are at most d small graphs among G_1, \dots, G_{k+d} , i.e. this bipartite partition cannot have $d+1$ small graphs. To this end, we prove that there exists at least one large graph among arbitrary $d+1$ graphs from $\{G_1, \dots, G_{k+d}\}$. Without loss of generality, we show that at least one of the graphs G_1, \dots, G_{d+1} is large. We group the class-vectors in \mathcal{C} (vertices of K_{2^k}) by their last $k-1$ coordinates: For any $\mathbf{b} \in \{0, 1\}^{k-1}$, let $V_{\mathbf{b}} \subseteq \mathcal{C}$ be the set of those class-vectors (vertices) whose last $k-1$ coordinate is \mathbf{b} . ($V_{\mathbf{b}}$ might be the empty set.) The point is that for any fixed $\mathbf{b} \in \{0, 1\}^{k-1}$, in K_{2^k} , every edge between the vertices of $V_{\mathbf{b}}$ must be contained in one of the graphs G_1, \dots, G_{d+1} , by property (ii) of class-vectors. Since these edges form a complete graph $K_{\mathbf{b}}$ on vertex set $V_{\mathbf{b}}$, we can trivially find an independent edge set $M_{\mathbf{b}}$ of size $\lfloor \frac{|V_{\mathbf{b}}|}{2} \rfloor$ in $K_{\mathbf{b}}$. If we do this for all sets $V_{\mathbf{b}}$ (for all $\mathbf{b} \in \{0, 1\}^{k-1}$), and take the union of the resulting independent

edge sets $M_{\mathbf{b}}$, we yield an independent edge set M of size

$$\begin{aligned} |M| &= \sum_{\mathbf{b} \in \{0,1\}^{k-1}} \left\lfloor \frac{|V_{\mathbf{b}}|}{2} \right\rfloor \geq \sum_{\mathbf{b} \in \{0,1\}^{k-1}} \left(\frac{|V_{\mathbf{b}}|}{2} - \frac{1}{2} \right) = \\ &= \frac{1}{2} \left(\sum_{\mathbf{b} \in \{0,1\}^{k-1}} |V_{\mathbf{b}}| \right) - \frac{1}{2} \cdot 2^{k-1} = \frac{1}{2} |\mathcal{C}| - 2^{k-2} = \frac{1}{2} \cdot 2^k - 2^{k-2} = 2^{k-2} \end{aligned}$$

in K_{2^k} . Since every edge of M is contained in one of the graphs G_1, \dots, G_{d+1} , thus at least one of these $d+1$ graphs must contain at least $\frac{2^{k-2}}{d+1}$ independent edges (from M) by the pigeonhole principle, i.e. there is a large graph among G_1, \dots, G_{d+1} . This is what we wanted to prove, so the proof is now complete. \square

The proof of the lower bound in Theorem 9. First we assume that $n = 2^k$ for some positive integer k . In this case we have to prove that $\chi'_{2K_2}(2^k) > k + \frac{1}{4}\sqrt{k} - 1$. It is obvious that $\chi'_{2K_2}(2^k) \geq k$ because, as mentioned in Observation 3, it is a known (folklore) fact that K_{2^k} cannot be partitioned into less than k (arbitrary) bipartite graphs. (In the terminology of the proof of Lemma 22, a bipartite partition containing less than k graphs would result 2^k different class-vectors with less than k coordinates, which is a contradiction.) Now assume that the Ferrers graphs F_1, \dots, F_{k+d} partition K_{2^k} such that $k+d$ is minimal, i.e. $k+d = \chi'_{2K_2}(2^k)$. We already know that $d \geq 0$, and we want to prove that $d > \frac{1}{4}\sqrt{k} - 1$ also holds. We apply Lemma 22 first with $G_i := F_i$, for $i = 1, \dots, k+d$: This lemma guarantees that there exist k graphs F_{i_1}, \dots, F_{i_k} in our partition, each having at least $\frac{2^{k-2}}{d+1}$ independent edges. Then, by Lemma 21, each F_{i_s} has at least $\frac{1}{2} \cdot \frac{2^{k-2}}{d+1} \cdot \left(\frac{2^{k-2}}{d+1} + 1 \right)$ edges, for $s = 1, \dots, k$. We have found altogether at least $\frac{k}{2} \cdot \frac{2^{k-2}}{d+1} \cdot \left(\frac{2^{k-2}}{d+1} + 1 \right)$ distinct edges in K_{2^k} , which is bounded by $\frac{2^k(2^k-1)}{2}$, the total number of edges of K_{2^k} . The inequality

$$\frac{k}{2} \left(\frac{2^{k-2}}{d+1} \right)^2 < \frac{k}{2} \cdot \frac{2^{k-2}}{d+1} \cdot \left(\frac{2^{k-2}}{d+1} + 1 \right) \leq \frac{2^k(2^k-1)}{2} < \frac{1}{2} \cdot 4^k$$

can be transformed into

$$\frac{\sqrt{k}}{4} < d+1$$

by a simple calculation, which completes the proof for the case $n = 2^k$. (The verification of the case $n = 1$ is up to the reader.)

If n is not a power of 2, let k^* be the largest integer such that $2^{k^*} \leq n$, i.e. set $k^* := \lfloor \log_2 n \rfloor$. Then the previous case applies for $n^* := 2^{k^*}$, so by the monotonicity of $\chi'_{2K_2}(n)$ (see Observation 1) we have that

$$\chi'_{2K_2}(n) \geq \chi'_{2K_2}(2^{k^*}) > k^* + \frac{1}{4}\sqrt{k^*} - 1 = \lfloor \log_2 n \rfloor + \frac{1}{4}\sqrt{\lfloor \log_2 n \rfloor} - 1,$$

as stated. □

4. COMPLEXITY ISSUES

We address the problem of computing $\chi'_{\mathcal{H}}(G)$ for a fixed \mathcal{H} and arbitrary graph G .

Some cases:

Observation 23. *If $\mathcal{H} = \{P_3, K_2 + K_1\}$ then $\chi'_{\mathcal{H}}(G) = e(G)$.*

Observation 24. *$\mathcal{H} = \{P_3\}$ then the language $\mathcal{L} = \{G : \chi'_{\mathcal{H}}(G) = 3\}$ is NP-complete, see Holyer [8]. In fact $\chi'_{\mathcal{H}}(G) = \chi'(G)$ for all graph G , where $\chi'(G)$ is the chromatic index of the graph G .*

Theorem 25 (Cherry Orchard). *$\mathcal{H} = \{P_4, S_4, C_4\}$ then the language $\mathcal{L} = \{G : \chi'_{\mathcal{H}}(G) = 3\}$ is NP-complete.*




FIGURE 7. The gadget *bird*.

Proof. Recall that it is NP-complete to decide whether a cubic graph $\chi'(G) = 3$, see Holyer [8]. We construct a graph G^* such that $v(G^*) = v(G) + 9e(G)$, $e(G^*) = 14e(G)$ and $\chi'_{\mathcal{H}}(G^*) = 3$ iff $\chi'(G) = 3$. First let us substitute each edge of G with a gadget, let us call it a *bird*, see Figure 7. The edges x and y in the figure are called the *legs* of the bird. So, for each $e = (A, B) \in G$, the edge e is (removed and) replaced with a copy of the bird gadget such that the degree-1 end vertices of the legs are glued to the vertices A and B of G as indicated in Figure 7.

If $\chi'(G) = 3$, then the edge colors can be extended to an \mathcal{H} -avoiding (cherry-) coloring of G^* . Let the colors be 1, 2, 3 and assume that an edge $e = (A, B)$ got the color 1 in the original good coloring. Then in the new (cherry-)coloring both x and y are colored by 1 and let the other edges be colored by 1, 2 or 3, according to Figure 8, which is a good (cherry-)coloring of G^* .

To see the other direction, suppose that G^* has an \mathcal{H} -avoiding (cherry-)coloring with colors 1, 2, 3. If the two legs are monochromatic in each bird gadget, i.e. x and y have the same color for each e , then by removing the gadgets and undoing the substitution we get a good 3 edge-coloring of G .

On the other hand, in a well (cherry-)colored gadget the colors of x and y must be the same. To see this, suppose that x and y have different colors in a gadget. Then, by the pigeonhole principle, among the four edges starting from vertex S , there exists




FIGURE 8. The good (cherry-)coloring of the gadget.

a pair of edges (different from $\{x, y\}$) of the same color. Similarly, since the degree of P is six, the six incident edges can be grouped into three monochromatic pairs, which means that P must be the center of three cherries P_3 . Then the edge (S, Q) must have the same color as (S, A) or (S, B) , because (P, S) is contained in a cherry centered at P . This implies that the five edges incident to Q cannot be covered by cherries using 3 colors (the edges (S, Q) and (P, Q) are contained in cherries centered at S and P , respectively), which is a contradiction. \square

5. FURTHER REMARKS

This work can continue in a number of ways. One direction is to consider graphs other than K_n . Real-world graphs motivate the general case, but graphs like Q_d , the d -dimensional cube, can be interesting from a purely mathematical point of view. Developing exact or even good approximate algorithms to partition general graphs with special bipartite graphs are also interesting problems and valuable in terms of applications.

We consider the problem of finding the tight bounds for partitioning K_n with Ferrers graphs (cf. Theorem 9) as the most challenging next step.

REFERENCES

- [1] Adcock, A.B., Sullivan, B.D., Mahoney, M.W. (2013). Tree-like structure in large social and information networks. In *2013 IEEE 13th international conference on data mining*, pp. 1–10.
- [2] Bastolla, U., Fortuna, M.A., Pascual-García, A., Ferrera, A., Luque, B., Bascompte, J. (2009). The architecture of mutualistic networks minimizes competition and increases biodiversity. *Nature*, **458**(7241), pp. 1018–1020.
- [3] Bóta, A., Csizmadia, L. and Pluhár, A. (2010). Community detection and its use in Real Graphs. In *Proceedings of the 2010 Mini-Conference on Applied Theoretical Computer Science (MATCOS)*, pp. 95–99.
- [4] Danziger, P., Quattrocchi, G. and Stevens, B. (2009). The Hamilton–Waterloo problem for cycle sizes 3 and 4. *Journal of Combinatorial Designs*, **17**(4), pp. 342–352.
- [5] Gera, I., London, A. and Pluhár, A. (2022). Greedy algorithm for edge-based nested community detection. In *IEEE 2nd Conference on Information Technology and Data Science (CITDS)*, pp. 86–91.
- [6] Gera, I., London, A. (2023). Detecting and generating overlapping nested communities. *Applied Network Science*, **8**(1), p. 51.

- [7] Graham, R. L., Pollak, H. O. (1971). On the addressing problem for loop switching. *The Bell system technical journal*, **50(8)**, pp. 2495–2519.
- [8] Holyer, I. (1981). The NP-completeness of edge-coloring. *SIAM Journal on Computing*, **10(4)**, pp. 718–720.
- [9] Junttila, E. and Kaski, P. (2011). Segmented nestedness in binary data. In *Proceedings of the 2011 SIAM International Conference on Data Mining*, pp. 235–246.
- [10] Kleinberg, J. (2002). An impossibility theorem for clustering. In *Advances in Neural Information Processing Systems*, **15**.
- [11] Knauer, K., Ueckerdt, T. (2016). Three ways to cover a graph. *Discrete Mathematics*, **339(2)**, pp. 745–758.
- [12] Král, D., Kratochvíl, J., Tuza, Z. and Woeginger, G.J. (2001). Complexity of coloring graphs without forbidden induced subgraphs. In *27th International Workshop on Graph-Theoretic Concepts in Computer Science*, pp. 254–262.
- [13] Lavrov, M. (2018), 4-cycle decomposition of a complete graph, *Mathematics Stack Exchange*, <https://math.stackexchange.com/q/2979661> (version: 2018-11-01).
- [14] London, A., Martin, R. R. and Pluhár, A. (2022). Graph clustering via generalized colorings. *Theoretical Computer Science*, **918**, pp. 94–104.
- [15] Lucas, E. (1892), *Récréations Mathématiques* (Vol. 2.), Paris.
- [16] Newman, M.E. and Girvan, M. (2004). Finding and evaluating community structure in networks. *Physical review E*, **69(2)**, 026113.
- [17] Palla, G., Derényi, I., Farkas, I. and Vicsek, T. (2005). Uncovering the overlapping community structure of complex networks in nature and society. *Nature*, **435(9)**, pp. 814–818.
- [18] Schaeffer, S. E. (2007). Graph clustering. *Computer Science Review*, **1(1)**, pp. 27–64.
- [19] Schwartz, S. (2022). An overview of graph covering and partitioning. *Discrete Mathematics*, **345(8)**, 112884.
- [20] Uzzi, B. (1996). The sources and consequences of embeddedness for the economic performance of organizations: The network effect. *American Sociological Review*, **61**, pp. 674–698.
- [21] Vishwanathan, S. (2010). A counting proof of the Graham Pollak Theorem. *arXiv preprint arXiv:1007.1553*
- [22] Wright, D.H., Patterson, B.D., Mikkelsen, G.M., Cutler, A., Atmar, W. (1997). A comparative analysis of nested subset patterns of species composition. *Oecologia*, **113**, pp. 1–20.
- [23] Zarankiewicz, K. (1951). Problem p 101., *Colloq. Math.*, **2**, 301, pp. 5.

BOLYAI INSTITUTE, UNIVERSITY OF SZEGED AND JOHN VON NEUMANN UNIVERSITY, KECSKEMÉT
Email address: lgyorffy@math.u-szeged.hu

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF SZEGED
Email address: london@inf.u-szeged.hu

BOLYAI INSTITUTE, UNIVERSITY OF SZEGED AND JOHN VON NEUMANN UNIVERSITY, KECSKEMÉT
Email address: ngaba@math.u-szeged.hu

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF SZEGED
Email address: pluhar@inf.u-szeged.hu