Neutron Star Properties and Femtoscopic Constraints

I. Vidaña¹,V. Mantovani Sarti², J. Haidenbauer³, D. L. Mihaylov^{2,4}, L. Fabbietti²

¹Istituto Nazionale di Fisica Nucleare, Sezione di Catania, Dipartimento di Fisica "Ettore Majorana",

Università di Catania, Via Santa Sofia 64, I-95123 Catania, Italy

²Physik Department E62, Technische Universität München, Garching, Germany, EU

³ Institute for Advanced Simulation (IAS-4), Forschungszentrum Jülich, D-52425 Jülich, Germany and

⁴ Sofia University, Faculty of Physics, 5 J. Bourchier Blvd., Sofia, Bulgaria

We construct the equation of state of hypernuclear matter and study the structure of neutron stars employing a chiral hyperon-nucleon interaction of the Jülich–Bonn group tuned to femtoscopic Λp data of the ALICE collaboration, and $\Lambda\Lambda$ and ΞN interactions determined from Lattice QCD calculations by the HAL QCD collaboration that reproduce the femtoscopic $\Lambda\Lambda$ and Ξ^-p data. We employ the ab-initio microscopic Brueckner–Hartree–Fock theory extended to the strange baryon sector. A special focus is put on the uncertainties of the hyperon interactions and how they are effectively propagated to the composition, equation of state, and mass-radius relation of neutron stars. To such end, we consider the uncertainty due to the experimental error of the femtoscopic Λp data used to fix the chiral hyperon-nucleon interaction and the theoretical uncertainty, estimated from the residual cut-off dependence of this interaction. We find that the final maximum mass of a neutron star with hyperons is in the range $1.3-1.4~M_{\odot}$, in agreement with previous works. The hyperon puzzle, therefore, remains still an open issue if only two-body hyperon-nucleon and hyperon-hyperon interactions are considered.

I. INTRODUCTION

The presence of hyperons in finite and infinite nuclear systems allows us the study of baryon interactions from an enlarged perspective and, consequently, to extend our present knowledge of conventional nuclear physics to the SU(3)-flavor sector [1–3]. Particularly, a big interest has being put on the study of hyperonic matter (nuclear matter with nucleonic and hyperonic degrees of freedom) in connection with the physics of neutron star interiors [4]. Neutron star properties such as for instance their masses, thermal evolution, neutrino emissivity, or their gravitational instabilities are closely related to the underlying equation of state (EoS) of matter at high densities [5]. Therefore, despite the fact that hypernuclear matter is an idealized system, the theoretical determination of its EoS is an essential step towards the understanding of those properties, which can be affected by the presence of hyperons [6, 7]. However, this is a very challenging task since it requires the knowledge of two-, three- and, eventually, many-body interactions involving hyperons and nucleons, and the solution of the very complicated nuclear many-body problem [8].

Compared to the nucleon-nucleon (NN) interaction, the hyperon-nucleon (YN) and hyperon-hyperon (YY) ones are still poorly constrained, mainly due to the scarce amount of YN scattering data, and to the complete lack of them in the YY case. The reason are the experimental difficulties associated with the short lifetime of hyperons and the low-beam intensity fluxes. Three-baryon forces involving hyperons (YNN, YYN and YYY) [9–14] are even less constrained experimentally, although their role could be fundamental for the solution of the so-called hyperon puzzle, *i.e.*, the problem of the strong softening of the dense matter EoS due to the presence of hyperons which leads to maximum masses of neutron stars

not compatible with the recent observations of $\sim 2 M_{\odot}$ millisecond pulsars [15–17]. However, information on the two-body YN interactions, additional to that from scattering experiments and hypernuclei [1, 2], together with constraints on the YY ones and three-body forces has emerged recently thanks to the so-called femtoscopy technique by measuring the correlations (in momentum space) of YN pairs [18, 19], YY pairs [20] and YNN triads [21] produced in proton-proton (pp) collisions at the LHC by the ALICE collaboration. The comparison of the measured two- and three-body correlation functions with theoretical predictions permits one to test, improve and constrain the existing two- and three-body hyperon interactions.

The aim of the present work is to construct the EoS of hypernuclear matter and to study the structure of neutron stars employing the chiral $\Lambda N-\Sigma N$ interaction recently tuned to femtoscopic data of the ALICE collaboration [21, 22], and $\Lambda\Lambda$ and ΞN interactions determined from the lattice QCD simulations of the HAL QCD collaboration [23] which predict results in agreement with femtoscopy. In particular, we use the chiral YN interaction NLO19 of the Jülich-Bonn group [24], and variants of it recently established in a combined analysis of Λp scattering and correlation-function data by Mihaylov et al. [22]. Regarding the $\Lambda\Lambda$ and ΞN interactions we consider two specific sets of the HAL QCD potential that were employed in the analysis of data on the Ξ^-p correlation function by the ALICE Collaboration [18]. The EoS of hypernuclear matter is then constructed by using these interactions within the well known ab-initio microscopic Brueckner-Hartree-Fock (BHF) theory extended to the strange baryon sector [25–32]. The major novelty of this work is a detailed study of how the uncertainties on the YN and YY interactions that describe available current experimental data are effectively propagated to

the EoS and to the structure of a system as complex as a neutron star with strangeness content.

The paper is organized in the following way. Details on the YN and YY employed in this work are given in Sect. II. The theoretical (BHF) description of the hyperonic matter of the EoS is briefly reviewed in Sect. III. Results for the composition, the EoS and the structure of neutron stars are presented in Sect. IV with a special focus put on the uncertainties of the interactions and how they are effectively propagated to the EoS and the neutron star structure. Finally, a short summary and the main conclusions of this work are given in Sect. V.

II. YN AND YY INTERACTIONS

In this work we focus on the strangeness S=-1,-2 baryon-baryon interactions, considering explicitly the ΛN - ΣN and $\Lambda \Lambda$ - ΞN coupled channels, respectively. The novelty of this work lies in the fact that the interactions employed as input to determine the behavior of the hypernuclear matter EoS yield results in agreement with the recent Λp [19, 33], $\Lambda \Lambda$ [20] and $\Xi^- p$ [18, 34] correlation measurements performed by the ALICE collaboration.

The S = -1 YN interaction is described by the NLO19 potentials derived within SU(3) chiral effective field theory (χEFT) [24]. These potentials incorporate contributions up to next-to-leading order (NLO) in the chiral expansion, in particular they include contributions from one- and two-pseudoscalar-meson exchange diagrams, involving the Goldstone bosons π , K, η , and from fourbaryon contact terms (without and with two derivatives). where the latter encode the unresolved short-distance dynamics. The low energy constants (LECs) associated with these contact terms are free parameters, which need to be constrained by experimental data, and they have been so far established by a global fit to a set of 36 Λp and ΣN low energy scattering data points [24], available since the 1960s. SU(3) flavor symmetry has been imposed which reduces the number of independent contact terms or LECs, respectively [35].

Of specific interest for the present study are variants of those potentials which have been established in Ref. [22]. in the combined analysis of the above mentioned scattering data and the more recent Λp correlation function measured by ALICE [33]. In order to generate a slightly weaker ΛN interaction, suggested by those correlation data [22], the possible SU(3) symmetry breaking in the leading-order contact terms of the relevant ${}^{1}S_{0}$ and ${}^{3}S_{1}$ partial waves has been exploited, in line with χ EFT and the associated power counting [35, 36], so that the description of the cross sections in other YN reactions $(\Sigma^+ p, \Sigma^- p, \Sigma^- p \to \Lambda n)$ remains practically unchanged as compared to the original potential [24]. The allowed 1σ region in the singlet and triplet scattering length plane (f_0, f_1) obtained in [22] delivers an estimate on the current uncertainty arising from all the available experimental input. This uncertainty has been translated into variants of the NLO19 potential with cutoff 600 MeV, by an appropriate adjustment of the pertinent LECs. The yellow band in Fig. 2 in [22] indicates the resulting variation (uncertainty) in the Λ single-particle potential. Additionally, in that work the theoretical uncertainty [37] had been considered, represented by the grey band in Fig. 2 of Ref. [22].

For the interaction in the S=-2 sector we employ the $\Lambda\Lambda$ - ΞN potentials obtained by the HAL QCD collaboration from lattice QCD calculations [23]. The potentials are evaluated with (2+1)-flavour simulations close to the physical point. The systematic uncertainty on the potentials are provided by evaluating them for three different values of $t/a=11,12,13^1$, with t/a=12 considered by them as default. The statistical uncertainty of the calculations is evaluated using the jackknife variations. The coupling to the other S=-2 channels, $\Lambda\Sigma$ and $\Sigma\Sigma$, has been ignored in that work and is likewise neglected in the present study.

The analysis of the experimental $\Lambda\Lambda$ correlation in [20] significantly reduced the allowed space for the effective range parameters and showed a full compatibility with the scattering length and effective range predicted by the $\Lambda\Lambda$ potentials of the HAL QCD Collaboration. This agreement with the correlation data is shown in Fig. 3 of Ref. [20] with the red cross indicating the effective range parameters and the corresponding statistical uncertainties resulting from the default t/a = 12 potential parametrization in Table 2 of Ref. [23]. In the analysis of the $\Xi^- p$ correlation function, measured by the ALICE collaboration in two different systems, namely in pp [18] and p-Pb [34] collisions, the ΞN interaction of the HAL QCD Collaboration was used directly. In particular, the correlation measurement in pp collisions showed a good agreement with the HAL QCD potential, where the magenta band in panel (a) of Fig. 3 in Ref. [18] includes the statistical lattice uncertainties as well as t/a systematic variations between 11 and 13. Note that a detailed analysis of the $\Lambda\Lambda$ and Ξ^-p correlation functions based on the HAL QCD potentials has been also performed by Kamiya et al. [38], which confirmed the agreement between the theory predictions and experiment.

Finally, let us come to the pure nucleonic interaction included in the determination of the EoS. For the two-body NN part, we employ the well-known Argonne V18 potential [39]. Effects of three-nucleon forces (3NFs) are accounted for too, as explained in more detail in the next section.

 $^{^{1}}$ Here, t is the Euclidean time and a represents the lattice spacing used in the simulated grid.

$a \text{ (MeV fm}^3)$	$b \; (\text{MeV fm}^6)$	E_0 (MeV)	$K_0 \text{ (MeV)}$
-15.22	21.70	-16	160
-53.42	260.42	-16	270

TABLE I: Values of the coefficients a and b for the two extreme nucleonic EoS considered. The corresponding values of the binding energy E_0 and the nuclear incompressibility K_0 at saturation density are also given.

III. BHF APPROACH OF HYPERONIC MATTER

Our description of the hyperonic matter EoS starts with the construction of all the G-matrices that describe the interaction of two baryons immersed in a surrounding medium. They are obtained by solving the well known Bethe–Goldstone equation, written schematically as

$$G(\omega)_{B_1B_2;B_3B_4} = V_{B_1B_2;B_3B_4} + \frac{1}{\Omega} \sum_{B_iB_j} V_{B_1B_2;B_iB_j} \frac{Q_{B_iB_j}}{\omega - E_{B_i} - E_{B_j} + i\eta} \times G(\omega)_{B_iB_j;B_3B_4}$$
(1)

where the first (last) two subindices indicate the initial (final) two-baryon state compatible with a given value S of the strangeness, namely S=0 (NN), S=-1 (Λ N, Σ N) and S=-2 (Λ A, Ξ N); $V_{B_1B_2;B_3B_4}$ is the bare baryon-baryon interaction ($B_1B_2 \to B_3B_4$); Ω is the (large) volume enclosing the system; $Q_{B_iB_j}$ is the Pauli operator that takes into account the effect of the exclusion principle on the scattered baryons; and the starting energy ω corresponds to the sum of the nonrelativistic single-particle energies of the interacting baryons. We note that Eq. (1) is a coupled channel equation (see for instance Ref. [29] for a detailed discussion on the Bethe–Goldstone equation in coupled channels).

The single-particle energy of a baryon B_i with momentum \vec{k} is given by

$$E_{B_i}(\vec{k}) = M_{B_i} + \frac{\hbar^2 k^2}{2M_{B_i}} + Re[U_{B_i}(\vec{k})] .$$
 (2)

Here M_{B_i} denotes the rest mass of the baryon and the single-particle potential $U_{B_i}(\vec{k})$ represents the average mean field "felt" by the baryon owing to its interaction with the other baryons of the (hyper)nuclear medium. In the BHF theory $U_{B_i}(\vec{k})$ is calculated through the "onshell" G-matrix given by

$$U_{B_i}(\vec{k}) = \frac{1}{\Omega} \sum_{B_j} \sum_{\vec{k}'} n_{B_j}(|\vec{k}'|)$$

$$\times \langle \vec{k}\vec{k}'|G_{B_iB_i:B_jB_j}(\omega)|\vec{k}\vec{k}'\rangle_{\mathcal{A}}$$
(3)

where $n_{B_j}(|\vec{k}|) = \theta(k_{F_{B_j}} - |\vec{k}|)$ is the occupation number of the baryon species B_j with Fermi momentum $k_{F_{B_j}}$, the index \mathcal{A} indicates that the matrix elements are properly antisymmetrized when baryons B_i and B_j belong to the

same isomultiplet, and $\omega=E_{B_i}(\vec{k})+E_{B_j}(\vec{k}')$. We note here that in this work we have adopted the so-called continuous prescription when solving the Bethe–Goldstone equation, since as shown by the authors of Refs. [40, 41] the contribution to the energy per particle of nuclear matter from three-hole line diagrams (which account from three-body correlations) is minimized when this prescription is considered. Once a self-consistent solution of Eqs. (1-3) is achieved, the energy density of the system can be calculated in the BHF theory as

$$\varepsilon_{BHF} = \frac{1}{\Omega} \sum_{B_i} \sum_{\vec{k}} n_{B_i}(|\vec{k}|)$$

$$\times \left(M_{B_i} + \frac{\hbar^2 k^2}{2M_{B_i}} + \frac{1}{2} U_{B_i}(\vec{k}) \right) . \tag{4}$$

The inclusion of genuine 3NFs in the microscopic BHF approach is currently a task still far to be achieved because it would require the solution of a three-body Bethe-Faddeev equation in the nuclear medium. Therefore, the usual way of introducing them consist in adding an effective density-dependent two-nucleon force to the bare NN interaction when solving the Bethe-Goldstone equation. This effective force is obtained by averaging the genuine 3NF over the spatial, spin and isospin coordinates of one of the three nucleons (see e.g., Refs. [42– 44]). In this work, however, we have adopted a simpler alternative strategy which will allow us to explore the uncertainty on the neutron star properties associated to the softness/stiffness of the pure nucleonic part of the EoS. To such end, first, we construct, as described above, the energy density of hyperonic matter within the BHF approach using only two-body NN, YN and YY forces and then, we add to it the following phenomenological density- and isospin-dependent term that accounts for the effect of 3NFs

$$\varepsilon_{3NF} = (a\rho_N^2 + b\rho_N^3)(1+\beta^2)$$
 (5)

Here $\rho_N = \rho_n + \rho_p$ is the total nucleonic density and $\beta = (\rho_n - \rho_p)/(\rho_n + \rho_p)$ the isospin asymmetry parameter. The coefficients a and b are fitted to reproduce simultaneously the binding energy $E_0 = -16$ MeV of symmetric matter at nuclear saturation density $\rho_0 = 0.17$ fm⁻³ and the nuclear incompressibility K_0 , for which two extreme values compatible with current experimental uncertainties [45] have been considered: $K_0 = 160$ MeV and $K_0 = 270$ MeV corresponding, respectively, to a very soft and to a very stiff nucleonic EoS. The values of the coefficients a and b are given in Tab. I.





FIG. 1: (Color online). Composition of β -stable neutron star matter including nucleons, Λ , Σ^- and Ξ^- baryons. For the S=-1 YN interaction we use the NLO19 potential [24] and variants established in Ref. [22], while for the S=-2, the $\Lambda\Lambda$ and Ξ N potentials from the HAL QCD Collaboration are used, see text in Sect. IV. The upper (lower) panels (a) and (b) ((c) and (d)) show the results when a soft (stiff) nucleonic EoS with $K_0=160$ MeV ($K_0=270$ MeV) is considered. Left (right) panels (a) and (c) ((b) and (d)) show the composition assuming only S=-1 YN interactions (assuming also $\Lambda\Lambda$ and Ξ N interactions). The narrow darker colored bands represent the uncertainty from the femtoscopic determination of the Λp scattering lengths [22]. The wider faint bands show the overall theoretical uncertainty, estimated here from the residual cutoff dependence of the NLO19 interaction.

Once the total energy density $\varepsilon = \varepsilon_{BHF} + \varepsilon_{3NF} + \varepsilon_L$ (ε_L being the contribution of non interacting leptons) is known, the composition and the EoS of neutron star matter can be obtained from the requirement of equilibrium under weak β decay processes, $\mu_i = b_i \mu_n - q_i \mu_e$ (b_i and q_i denoting the baryon number and charge of species i) and electric charge neutrality, $\sum_i q_i \rho_i = 0$. The chemical potentials of the various species and the pressure are computed from the usual thermodynamical relations, $\mu_i = \partial \varepsilon / \partial \rho_i$ and $P = \rho^2 \partial (\varepsilon/\rho) / \partial \rho$. Finally, knowing the EoS, the equilibrium configurations of static non-rotating neutron stars are obtained by solving the well-known Tolman–Oppenheimer–Volkoff (TOV) equations [46, 47].

IV. NEUTRON STARS: EOS AND STRUCTURE

We start this section by showing in Fig. 1 the composition of β -stable neutron star matter as a function of the density ρ , expressed in units of ρ_0 . The upper ((a), (b)) and lower ((c), (d)) panels show, respectively, the content of matter when a soft $(K_0 = 160 \text{ MeV})$ or a stiffer $(K_0 = 270 \text{ MeV})$ nucleonic EoS is considered. The particle fractions when only S = -1 YN interactions are included in the calculation are displayed in the left ((a), (c)) panels of the figure, whereas those obtained when considering additionally the $S=-2~\Lambda\Lambda$ and ΞN interactions are shown in the right ((b), (d)) ones. The different colored bands reflect the uncertainties in the YN and YY interactions. A detailed discussion on how these uncertainties are effectively propagated to the composition, EoS and structure of neutron stars is given at the end of the section. Looking at the left and right panels we observe that, for a fixed nucleonic EoS, the onset den-






FIG. 2: (Color online). Particle chemical potentials as a function of the density (expressed in units of ρ_0) in β -stable neutron star matter including nucleons, Λ , Σ^- and Ξ^- baryons when a soft nucleonic EoS with $K_0 = 160$ MeV is considered (panel (a)) and when a scenario with a stiff nucleonic EoS with $K_0 = 270$ MeV is assumed (panel (b)). Results are shown using the chiral YN potential NLO19 with a cutoff of 600 MeV, and the $\Lambda\Lambda$ and Ξ N HAL QCD potentials for the sets of parameters given in Ref. [23] for t/a=12.

sity of the Λ is the same. This is due to the fact that, up to that density, matter is composed only of nucleons and leptons and, therefore, the S=-2 interactions play no role. We note also that the density at which the Σ^- appears is likewise not affected by these interactions simply because the $\Lambda\Sigma^-$ and $\Sigma^-\Sigma^-$ channels have not been included in the calculations. Comparing the upper ((a)-(b)) and lower panels ((c)-(d)) we see that the use of a stiffer nucleonic EoS leads to an earlier onset and a slightly larger population of Λ and Σ^- hyperons. Additionally, we should notice that whereas the Ξ^- hyperon appears around $4\rho_0$ when a softer nucleonic EoS is considered, it does not appear (or it appears at very large densities) when using a stiffer one.

To understand better all these features, we show in Fig. 2 the density dependence of the chemical potentials, μ_i , of the different baryon species obtained for the two nucleonic EoS together with the YN potential NLO19 with a 600 MeV cutoff, and the $\Lambda\Lambda$ and ΞN HAL QCD potentials for the sets of parameters given in Tabs. 2, 3 and 4 of Ref. [23] for t/a=12. Similar qualitative results have been obtained using other values of the cutoff and the parameters of the HAL QCD potentials. The earlier onset of the Λ and Σ^- hyperons when using a stiffer nucleonic EoS is now easy to understand since in this case the neutron (black line) and neutron plus electron (green line) chemical potentials increase more rapidly with the density (right panel), ensuring the chemical equilibrium condition for the appearance of these two hyperons is fulfilled at lower densities. Conversely, the softer nucleonic EoS scenario (left panel) leads to a later onset of such hyperons. The Ξ^- hyperon, which appears for the softer nucleonic EoS at around $4\rho_0$, does not appear below \sim $8\rho_0$ for the stiffer one. The reason is that in this case, due to the earlier appearance of the Σ^- , both the neutron and electron chemical potentials are smaller and, therefore, the Ξ^- onset condition is only fulfilled at very large densities ($\gtrsim 9\rho_0$).

The neutron star EoS and mass-radius relation corresponding to the particle compositions of the different cases just described are respectively shown in the upper ((a), (b)) and lower ((c), (d)) panels of Fig. 3. Results assuming only S = -1 YN potentials are displayed on the left ((a), (c)) panels whereas those including additionally the $\Lambda\Lambda$ and ΞN interactions in the S = -2 channels are presented on the right ((b), (d)) ones. In the lower panels (c) and (d), we also show the recently observed masses of three heavy millisecond pulsars PSR J1614-2230 (1.928 \pm 0.017 M_{\odot}) [15], PSR $J0348+0432 \ (2.01\pm0.04M_{\odot}) \ [16] \ and \ PSR \ J0740+6620$ $(2.14^{+0.10}_{-0.09}M_{\odot})$ [17] with their corresponding uncertainty bars, as well as constraints from the neutron star merger gravitational wave event GW170817 [48], and from the NICER mission [49–53].

Let us focus first on the EoS. As it is expected [6, 7], the appearance of hyperons leads to a softening of the EoS that is enhanced when the nucleonic contribution to the EoS is stiffer. This is because, in this case, the onset of the Λ and Σ^- hyperons is shifted to lower densities and their population is larger (see Fig. 1), as discussed before. A more quantitative estimate of such a softening effect can be obtained by comparing for instance the small dark blue bands in panel (a) or the small green ones in panel (b) of Fig. 3 for the soft and stiff nucleonic scenario. Particularly, in the high energy density region, one can observe a sizable reduction in the pressure of the order of about 20 - 25% when a stiff nucleonic EoS is used. As it is seen also in panel (b), an additional softening of the EoS is obtained when we take into account the effect of the $\Lambda\Lambda$ and ΞN interactions, due to their attractive character.

We now turn to describe the effects of the different




FIG. 3: (Color online). Panels (a) and (b): EoS of β -stable neutron star matter including nucleons, Λ , Σ^- and Ξ^- baryons. The EoS of nucleonic matter in β -equilibrium predicted for $K_0=160$ MeV (orange line) and $K_0=270$ MeV (red line) are shown for comparison. For the S=-1 YN interaction we use the NLO19 potential [24] and variants established in Ref. [22], while for the interactions in the S=-2 channels $\Lambda\Lambda$ and Ξ N the HAL QCD potentials described in Sect. IV are assumed. Panels (c) and (d): mass versus radius plots for the corresponding EoS shown in the upper panels. Recent observed masses of the millisecond pulsars PSR J1614-2230 [15], PSR J0348+0432 [16] and PSR J0740+6620 [17] with their corresponding error bars are also shown, as well as constraints from the gravitational wave event GW170817 [48], the NICER mission [49–53]. The reported astrophysical constraints for the GW170817 and NICER data are provided with 2σ uncertainty, while the Shapirodelay measurements for the heavy pulsars are reported with 1σ uncertainty.

EoS on the mass-radius relation of neutron stars. Results are shown in panels (c) and (d) of Fig. 3. We note first that the maximum mass of a neutron star with S = -1 and S = -2 hyperons in its interior is around $1.3-1.4~M_{\odot}$, as shown in panels (c) and (d) with the green and blue bands, respectively. We note also that such values seem to be quite insensitive to the nucleonic part of the EoS, washing out the rather large uncertainty in the maximum mass (from $\approx 2.3 \text{ M}_{\odot}$ to 1.8 M_{\odot}) achieved with a nucleons-only scenario. This insensitivity to the nucleonic EoS was already observed in Refs. [31, 32] where the Nijmegen YN and YY mesonexchange interaction models were used in combination with different NN potentials. As it was argued in these two works the reason for this should be traced to several strong compensation mechanisms caused by the appearance of hyperons that always leads to a soft EoS keeping the value of the maximum mass within $\sim 0.05 M_{\odot}$. Note,

however, that there still remain characteristic differences between the mass-radius relations that would eventually allow one to determine the EoS from observational constraints. In particular, there are significant differences for the corresponding radii achieved within the different EoS assumptions, which are linked to the maximum central baryon density. The overall stiffest EoS scenario with hyperons, achieved for the model with low nuclear incompressibility $K_0 = 160$ MeV, typically leads to radii of approximately 10 km, while larger values (≈ 12 km) are reached when the nuclear incompressibility is as well larger. Concerning the agreement with the reported measured masses and radii, we observe that the heavy pulsars data can be only reproduced by pure nucleonic EoSs. On the other hand, the gravitational wave constraints from GW170817, along with the NICER mass and radius measurements inferred for PSR J0030+0451, can be overall reproduced with the softer EoS with strange and multistrange hyperons. We note, however, that the latter shows a slight tension with the most recent NICER constraints on the PSR J0437-4715 millisecond pulsar.

Let us finally describe how the uncertainties in the hyperon interactions are effectively propagated to the composition, EoS and mass-radius relation of neutron stars. The variations within the S=-1 sector are generated by varying the LECs of the chiral YN potential NLO19 as described in [22]. There are two main sources of uncertainties, one related to the statistical errors of the AL-ICE data [54] and a theoretical one that is related to the truncation in the chiral expansion. Following Ref. [22], the data driven errors are considered by employing variants of the NLO19 YN potential with cutoff 600 MeV [24] whose LECs were readjusted in such a way that the scattering length $f_0 = 2.50$ fm is produced and variations of f_1 between the allowed values of 1.32 fm and 1.55 fm are achieved (see Table I in [22]). An essential feature of χ EFT is that, contrary to calculations with phenomenological models, an estimation for the theoretical uncertainty can be provided [37, 55]. In the present work it is estimated from the dependence of the results on the regulator cutoff. For that, the LECs are readjusted so that for all cutoff values considered in the original NLO19 potential (500-650 MeV) the scattering lengths of NLO19 (600) are obtained [24]. Note that the cutoff dependence provides only a lower bound for the uncertainty. For more elaborated ways to estimate the theoretical uncertainty in the context of nuclear matter calculations, that do not rely on cutoff variation, see [56].

Within the present work, the S = -2 sector includes the $\Lambda\Lambda$ and ΞN channels. The uncertainties for the ΞN interaction are inferred from the ALICE analysis on Ξ^-p correlations [18], in which the jackknife method was employed to obtain the statistical uncertainties from c.a. 24 different parameterizations of the lattice potential. To include the systematic uncertainty, the jackknife method has been applied to three different t/a = 11, 12, 13 values, resulting in a total of 71 different potentials. Out of these we have selected two specific sets of parameters that correspond to, respectively, the most and least attractive cases for the \(\Xi\)N interaction compatible with the femtoscopic $\Xi^- p$ data. Regarding the other three channels, $\Lambda\Lambda$, $\Lambda\Sigma$, and $\Sigma\Sigma$, in this work we take into account only $\Lambda\Lambda$ and its coupling to ΞN . Details on the potentials for $\Lambda\Lambda$ as well as for the transition $\Lambda\Lambda - \Xi N$ are provided in [23]. While the effect of these contributions to the final result are important, we have verified that their different allowed parameterizations lead to identical results within the precision of the present analysis. Thus, within the present work only the central values for t/a = 11 as listed in Tables 2 and 3 in [23] were considered.

The uncertainties on the composition, EoS and massradius relation, shown by the bands in Figs. 1 and 3 originate almost entirely from the interaction in the ΛN - ΣN system. These bands are constructed in the following way: For each one of the two pure nucleonic EoS, the wider bands are built by performing all possible calculations combining the NLO19 YN interactions with cutoff values 500-650 MeV, with two ΞN potentials corresponding to the two selected sets of parameters mentioned above, and the $\Lambda\Lambda$ and $\Lambda\Lambda-\Xi N$ ones for t/a=11 of Ref. [23]. The upper and lower extremes of the wider bands are then defined by the combination which gives, respectively, the stiffer and softer EoS. Similarly, the narrow bands are constructed by using the variants of the NLO19 YN interaction with cutoff 600 MeV, with LECs tuned to femtoscopic Λp data, combined with the same ΞN , $\Lambda\Lambda$ and $\Lambda\Lambda-\Xi N$ potentials. The extremes of the narrow bands are defined as those of the wider ones.

Not unexpectedly, the theoretical uncertainty due to the underlying $\Lambda N-\Sigma N$ interaction dominates the results. In fact, in analogous studies of (purely) nuclear matter properties based on NN interactions derived within χ EFT the theoretical uncertainty was found to be likewise large [56, 57]. Despite of much higher orders in the chiral expansion (N⁴LO) a reasonable convergence was only observed up to about twice the saturation density of nuclear matter [56]. Clearly, for improvements, among other things, a consistent inclusion of corresponding many-body forces is required [57]. We want to mention that there is an additional source of uncertainty due to the fact that the ΛN interaction in P- and higher partial waves is not constrained by empirical information. Judging from results in the literature, for example those in Ref. [58], it could be in the order of ± 3 MeV in the Λ single-particle potential at nuclear matter saturation density. We do not consider this uncertainty in the present calculation.

V. SUMMARY AND CONCLUSIONS

In this work we have employed the ab-initio microscopic Brueckner-Hartree-Fock theory extended to the strange baryon sector to construct the EoS of hypernuclear matter under β -equilibrium conditions, relevant for the study of neutron stars. To such end, we have used a chiral hyperon-nucleon interaction of the Jülich-Bonn group tuned to femtoscopic Λp data of the ALICE collaboration, and $\Lambda\Lambda$ and ΞN interactions determined from Lattice QCD calculations by the HAL QCD collaboration that reproduce the femtoscopic $\Lambda\Lambda$ and Ξ^-p data. To describe the pure nucleonic part of the EoS we have used the well-known Argonne V18 nucleon-nucleon potential. To account for the effect of three-nucleon forces we have added a phenomenological term to the total energy density of the system with parameters fitted to reproduce simultaneously the binding energy of symmetric nuclear matter at saturation and the nuclear incompressibility K_0 .

We have put a special focus on the interaction uncertainties and how they are effectively propagated to the composition, EoS and mass-radius relation of neutron stars. In particular, to explore the uncertainty on the neutron star properties associated to the softness/stiffness of the pure nucleonic part we have considered two extreme values of K_0 compatible with current heavy-ion experimental data. Regarding hyperons, we have considered the uncertainty due to the experimental error of the femtoscopic data used to fix the chiral hyperon-nucleon interaction and the theoretical uncertainty estimated from the residual cut-off dependence of this interaction, as well as the uncertainties on the $\Lambda\Lambda$ and ΞN interactions inferred from the analysis of correlation measurements. We have employed two specific sets of parameterizations of the ΞN potential from lattice QCD [59], out of the 71 explored in the ALICE analysis of $\Xi^- p$ correlations, that correspond, respectively, to the most and less attractive ΞN compatible with femtoscopic data. We have seen that the different parametrizations of the HAL QCD collaboration for the $\Lambda\Lambda$ potential and the transition $\Lambda\Lambda - \Xi N$ lead to identical results within the precision of the present analysis. Therefore, we have considered only one of the parameters sets listed in Tables 2 and 3 of Ref. [23].

Our results have shown, in agreement with previous works, that the appearance of hyperons in the interior of neutrons of neutron stars leads to a strong softening of the EoS and to a reduction of the maximum mass to values in the range $1.3-1.4\,M_{\odot}$, incompatible with current observations. In particular, the sizable range of maximum neutron star masses achieved within a pure nucleonic EoS, constrained to the current incompressibility limits, is significantly reduced once hyperons appear,

and the final maximum mass of a hyperonic star results quite independent of the nucleonic component of the EoS, as it was already observed in Refs. [31, 32]. Therefore, the hyperon puzzle remains still an open issue if only two-body hyperon-nucleon and hyperon-hyperon interactions are considered. Its solution requires a mechanism or mechanisms that could provide the additional repulsion that makes the EoS stiffer and, therefore, the maximum mass compatible with observations. One of such mechanisms could be the inclusion of hyperonic three-body forces [9–14] or, for instance, to consider a more exotic scenario where light QCD axions could appear in neutron stars [60]. Such possible mechanisms will be analyzed in future works.

Acknowledgments

This work was supported by ORIGINS cluster DFG under Germany's Excellence Strategy-EXC2094 - 390783311 and the DFG through the Grant SFB 1258 "Neutrinos and Dark Matter in Astro and Particle Physics". I.V. acknowledges the Excellent Cluster ORIGINS cluster DFG under Germany's Excellence Strategy-EXC2094 - 390783311 for its support during his visit at TUM in June and July 2024. V. M. S. was supported by the Deutsche Forschungsgemeinschaft (DFG) through the grant MA 8660/1-1.

- E. Botta, T. Bressani, and G. Garbarino, Eur. Phys. J. A 48, 41 (2012), arXiv:1203.5707 [nucl-ex].
- [2] A. Gal, E. V. Hungerford, and D. J. Millener, Rev. Mod. Phys. 88, 035004 (2016), arXiv:1605.00557 [nucl-th].
- [3] H. Lenske, M. Dhar, T. Gaitanos, and X. Cao, Prog. Part. Nucl. Phys. 98, 119 (2018).
- [4] L. Rezzolla, P. Pizzochero, D. I. Jones, N. Rea, and I. Vidaña, eds., The Physics and Astrophysics of Neutron Stars, Astrophysics and Space Science Library, Vol. 457 (Springer, 2018).
- [5] M. Baldo, Int. Rev. Nucl. Phys. 8, 1 (1999).
- [6] D. Chatterjee and I. Vidaña, Eur. Phys. J. A 52, 29 (2016), arXiv:1510.06306 [nucl-th].
- [7] I. Vidaña, Proc. Roy. Soc. Lond. A 474, 0145 (2018), arXiv:1803.00504 [nucl-th].
- [8] P. Ring and P. Schuck, The Nuclear Many-Body Problem, Physics and astronomy online library (Springer, 2004).
- [9] S. Petschauer, N. Kaiser, J. Haidenbauer, U.-G. Meißner, and W. Weise, Phys. Rev. C 93, 014001 (2016), arXiv:1511.02095 [nucl-th].
- [10] J. Haidenbauer, U. G. Meißner, N. Kaiser, and W. Weise, Eur. Phys. J. A 53, 121 (2017), arXiv:1612.03758 [nuclth].
- [11] M. Kohno, Phys. Rev. C 97, 035206 (2018), arXiv:1802.05388 [nucl-th].
- [12] D. Logoteta, I. Vidaña, and I. Bombaci, Eur. Phys. J. A 55, 207 (2019), arXiv:1906.11722 [nucl-th].
- [13] D. Gerstung et al., Eur. Phys. J. A 56, 175 (2020),

- arXiv:2001.10563 [nucl-th].
- [14] H. Tong, S. Elhatisari, and U.-G. Meißner, (2024), arXiv:2405.01887 [nucl-th].
- [15] P. Demorest, T. Pennucci, S. Ransom, M. Roberts, and J. Hessels, Nature 467, 1081 (2010), arXiv:1010.5788 [astro-ph.HE].
- [16] J. Antoniadis et~al., Science **340**, 6131 (2013), arXiv:1304.6875 [astro-ph.HE] .
- [17] H. T. Cromartie et al. (NANOGrav), Nature Astron. 4, 72 (2019), arXiv:1904.06759 [astro-ph.HE].
- [18] S. Acharya et al. (ALICE), Nature 588, 232 (2020), [Erratum: Nature 590, E13 (2021)], arXiv:2005.11495 [nuclex].
- [19] S. Acharya et al. (ALICE), Phys. Lett. B 833, 137272 (2022), arXiv:2104.04427 [nucl-ex].
- [20] S. Acharya et al. (ALICE), Phys. Lett. B 797, 134822 (2019), arXiv:1905.07209 [nucl-ex].
- [21] S. Acharya et al. (ALICE), Eur. Phys. J. A 59, 145 (2023), arXiv:2206.03344 [nucl-ex].
- [22] D. L. Mihaylov, J. Haidenbauer, and V. Mantovani Sarti, Phys. Lett. B 850, 138550 (2024), arXiv:2312.16970 [nucl-th].
- [23] K. Sasaki et al. (HAL QCD), Nucl. Phys. A 998, 121737 (2020), arXiv:1912.08630 [hep-lat].
- [24] J. Haidenbauer, U.-G. Meißner, and A. Nogga, Eur. Phys. J. A 56, 91 (2020), arXiv:1906.11681 [nucl-th].
- [25] H. J. Schulze, A. Lejeune, J. Cugnon, M. Baldo, and U. Lombardo, Phys. Lett. B 355, 21 (1995).

- [26] H. J. Schulze, M. Baldo, U. Lombardo, J. Cugnon, and A. Lejeune, Phys. Rev. C 57, 704 (1998).
- [27] M. Baldo, G. F. Burgio, and H. J. Schulze, Phys. Rev. C 58, 3688 (1998).
- [28] M. Baldo, G. F. Burgio, and H. J. Schulze, Phys. Rev. C 61, 055801 (2000), arXiv:nucl-th/9912066.
- [29] I. Vidaña, A. Polls, A. Ramos, M. Hjorth-Jensen, and V. G. J. Stoks, Phys. Rev. C 61, 025802 (2000), arXiv:nucl-th/9909019.
- [30] I. Vidaña, A. Polls, A. Ramos, L. Engvik, and M. Hjorth-Jensen, Phys. Rev. C 62, 035801 (2000), arXiv:nuclth/0004031.
- [31] H. J. Schulze, A. Polls, A. Ramos, and I. Vidaña, Phys. Rev. C 73, 058801 (2006).
- [32] H. J. Schulze and T. Rijken, Phys. Rev. C 84, 035801 (2011).
- [33] D. Mihaylov and J. González González, Eur. Phys. J. C 83, 590 (2023), arXiv:2305.08441 [hep-ph].
- [34] S. Acharya et al. (ALICE), Phys. Rev. Lett. 123, 112002 (2019), arXiv:1904.12198 [nucl-ex].
- [35] J. Haidenbauer, S. Petschauer, N. Kaiser, U.-G. Meißner, A. Nogga, and W. Weise, Nucl. Phys. A 915, 24 (2013), arXiv:1304.5339 [nucl-th].
- [36] S. Petschauer and N. Kaiser, Nucl. Phys. A 916, 1 (2013), arXiv:1305.3427 [nucl-th].
- [37] E. Epelbaum, H.-W. Hammer, and U.-G. Meißner, Rev. Mod. Phys. 81, 1773 (2009), arXiv:0811.1338 [nucl-th].
- [38] Y. Kamiya, K. Sasaki, T. Fukui, T. Hyodo, K. Morita, K. Ogata, A. Ohnishi, and T. Hatsuda, Phys. Rev. C 105, 014915 (2022), arXiv:2108.09644 [hep-ph].
- [39] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, Phys. Rev. C $\bf 51,$ 38 (1995), arXiv:nucl-th/9408016 .
- [40] H. Q. Song, M. Baldo, G. Giansiracusa, and U. Lombardo, Phys. Rev. Lett. 81, 1584 (1998).
- [41] M. Baldo, G. Giansiracusa, U. Lombardo, and H. Q. Song, Phys. Lett. B 473, 1 (2000).
- [42] B. A. Loiseau, Y. Nogami, and R. C. K., Nucl. Phys. A 165, 601 (1971), [Erratum: Nucl.Phys.A 176, 665–666 (1971)].

- [43] P. Grangé, M. Martzolff, Y. Nogami, D. W. L. Sprung, and C. K. Ross, Phys. Lett. B 60, 237 (1976).
- [44] M. Baldo and L. S. Ferreira, Phys. Rev. C 59, 682 (1999).
- [45] A. Le Fèvre, Y. Leifels, W. Reisdorf, J. Aichelin, and C. Hartnack, Nucl. Phys. A 945, 112 (2016), arXiv:1501.05246 [nucl-ex].
- [46] R. C. Tolman, Phys. Rev. **55**, 364 (1939).
- [47] J. R. Oppenheimer and G. M. Volkoff, Phys. Rev. 55, 374 (1939).
- [48] B. P. Abbott et al. (LIGO Scientific, Virgo), Phys. Rev. Lett. 119, 161101 (2017), arXiv:1710.05832 [gr-qc].
- [49] K. Gendreau and Z. Arzoumanian, Nature Astronomy 1, 895 (2017).
- [50] T. E. Riley et al., Astrophys. J. Lett. 887, L21 (2019), arXiv:1912.05702 [astro-ph.HE].
- [51] T. E. Riley et al., Astrophys. J. Lett. 918, L27 (2021), arXiv:2105.06980 [astro-ph.HE].
- [52] S. Vinciguerra et al., Astrophys. J. 961, 62 (2024), arXiv:2308.09469 [astro-ph.HE].
- [53] T. Salmi et al., arXiv:2406.14466 [astro-ph.HE] .
- [54] S. Acharya et al. (ALICE), Phys. Lett. B 811, 135849 (2020), arXiv:2004.08018 [nucl-ex].
- [55] E. Epelbaum, H. Krebs, and U.-G. Meißner, Eur. Phys. J. A 51, 53 (2015), arXiv:1412.0142 [nucl-th].
- [56] J. Hu, Y. Zhang, E. Epelbaum, U.-G. Meißner, and J. Meng, Phys. Rev. C 96, 034307 (2017), arXiv:1612.05433 [nucl-th].
- [57] F. Sammarruca, L. Coraggio, J. W. Holt, N. Itaco, R. Machleidt, and L. E. Marcucci, Phys. Rev. C 91, 054311 (2015), arXiv:1411.0136 [nucl-th].
- [58] S. Petschauer, J. Haidenbauer, N. Kaiser, U.-G. Meißner, and W. Weise, Eur. Phys. J. A 52, 15 (2016), arXiv:1507.08808 [nucl-th].
- [59] K. Sasaki, Private communication.
- [60] R. Balkin, J. Serra, K. Springmann, S. Stelzl, and A. Weiler, Phys. Rev. D 109, 095032 (2024), arXiv:2211.02661 [hep-ph].