

\mathcal{L} -LIE ALGEBROIDS OVER TOPOLOGICAL RINGED SPACES

MAINAK PODDAR AND ABHISHEK SARKAR

ABSTRACT. The notion of Lie algebroids over a topological ringed space provides a unified framework to study various geometric structures. This geometric concept is intimately connected with well-known algebraic structures, including Gerstenhaber algebras and Batalin–Vilkovisky algebras. We introduce more general concepts such as \mathcal{L} -Lie algebroids and \mathcal{A} -Gerstenhaber algebras, associated with a given Lie algebroid \mathcal{L} and Gerstenhaber algebra \mathcal{A} over a topological ringed space, respectively. Following this, we explore how several standard correspondences extend within this broader framework.

1. INTRODUCTION

The notion of Lie algebroids plays a prominent role in differential geometry and mathematical physics as they represent generalized infinitesimal symmetries of spaces, which are related to the corresponding global symmetries of spaces described by Lie groupoids [30]. Lie algebroids over a C^∞ -manifold generalize the notion of tangent bundle of a manifold and as well as the notion of Lie algebras [13,33]. The standard geometric notions (or structures) associated with a Lie algebroid such as exterior algebra of a Lie algebroid, Lie bialgebroid, Lie algebroid connections, Chevalley–Eilenberg–de Rham complex, etc., are defined for smooth vector bundles (see [30,44]). In recent years, some of these notions have been generalized in the context of complex geometry and algebraic geometry, using sheaf theoretic language (see [5,6,8,9,22,32,34,40]). These geometric notions are closely associated with some well known algebraic structures, such as Lie–Rinehart algebras, Gerstenhaber algebras (in short G-algebras), Batalin–Vilkovisky algebras (in short BV-algebras), universal enveloping algebra of a Lie–Rinehart algebra, etc., (see [19,21,25,29,33,44]). We consider sheaf theoretic (or algebro-geometric) analogues of these algebraic notions, and investigate some of the important relationships between them (see details in Section 2).

Unlike smooth vector bundles over a smooth manifold, the global sections of a holomorphic or algebraic vector bundle over a complex manifold or algebraic variety respectively, do not, in general, capture all the information about the bundle. Consequently, a sheaf-theoretic approach becomes essential in analytic and algebraic geometry, even in non-singular cases. To create a unified framework for studying geometric objects across diverse contexts, such as smooth, holomorphic, and algebraic settings, one relies on sheaves over locally ringed spaces (see [38,43]). When a space has singularities, the tangent bundle is replaced by the tangent sheaf, which consists of derivations acting on the structure sheaf of the space. Another important sheaf associated to these spaces is the logarithmic tangent sheaf, meaning the sheaf of logarithmic derivations corresponding to some divisor. Both of these notions share a common underlying structure as (quasi)coherent sheaves of Lie–Rinehart algebras, generally regarded as Lie algebroids (see [32,34,41]).

Recently, the notion of an L -Lie algebroid has found several applications in the literature. The notion was introduced in the smooth context by R. L. Klaasse who defined it is a Lie algebroid whose anchor factors through the anchor of a fixed Lie algebroid L (see [24]). He has used it in the study of log symplectic structures taking L to be a logarithmic tangent sheaf. The logarithmic Atiyah sheaf and the G -Atiyah sheaf employed by Biswas et al. [2–4] in the context of equivariant geometry are also examples of L -Lie algebroids. Note that when L is the tangential Lie algebroid, an L -Lie algebroid is simply a Lie algebroid.

The notion of an L -Lie algebroid may be informally thought of as a tethered or a rooted Lie algebroid. In this article, we formulate analogous rooted versions of the notions of differential graded algebras, Gerstenhaber algebras, strong differential Gerstenhaber algebras, BV algebras, Lie bialgebroids, etc. We always

AMS Mathematics Subject Classification (2020): 53D17, 32L10, 14F40, 17B66

Key words and phrases. Lie algebroids, tangent sheaf, Gerstenhaber algebras, BV-algebras, universal enveloping algebra, Lie algebroid (co)homology.

work in the setting of sheaves over locally ringed spaces. We generalize the relationships among the classical versions of these objects to their rooted avatars. Some of these are given a functorial interpretation.

Our basic object of study is an \mathcal{L} -Lie algebroid, where \mathcal{L} refers to a given Lie algebroid over a topological ringed space (see Section 3). This notion appears in various geometric settings, such as Poisson geometry [34], log symplectic structures [24], singular foliations [26], and triangular Lie bialgebroids [7,25]. Recall that, given a Lie algebroid \mathcal{L} , there is a sheaf of differential graded algebras $\Omega_{\mathcal{L}}^{\bullet}$, known as the Chevalley-Eilenberg-de Rham complex of \mathcal{L} [5,37]. Moreover, there is a sheaf of Gerstenhaber algebras $\mathcal{G}_{\mathcal{L}}$, given by the canonical Gerstenhaber algebra structure on the exterior algebra $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}$ endowed with the generalized Schouten-Nijenhuis bracket [9,38]. The smooth and algebraic counterparts of the above associations are described in [20,25,44]. We define more general notions, namely, $\Omega_{\mathcal{L}}^{\bullet}$ -differential graded algebras and $\mathcal{G}_{\mathcal{L}}$ -Gerstenhaber algebras for a given Lie algebroid \mathcal{L} over a topological ringed space (X, \mathcal{O}_X) . We establish analogous relationships among them (see Theorem 3.9 and Theorem 4.3).

There is a well known correspondence between Lie bialgebroid structures on a smooth vector bundle and strong differential Gerstenhaber algebra structures on its associated exterior algebra (see [20,25,44]). A strong differential Gerstenhaber algebra is a Gerstenhaber algebra together with a square zero differential of degree 1 which acts like a graded derivation with respect to both the exterior product and bracket operations. We introduce the notion of an \mathcal{L} -Lie bialgebroid and associate it with the notion of a strong differential $\mathcal{G}_{\mathcal{L}}$ -Gerstenhaber algebra (see Theorem 5.10). An \mathcal{L} -Lie bialgebroid is a pair of \mathcal{L} -Lie algebroids in duality with an additional compatibility condition as in the usual case.

For an \mathcal{O}_X -module \mathcal{E} the notion Atiyah algebroid $\mathcal{A}t(\mathcal{E})$ consists of first order differential operators on \mathcal{E} having scalar symbol ([5,22]). An \mathcal{L} -connection ∇ on an \mathcal{O}_X -module \mathcal{E} is an \mathcal{O}_X -linear morphism $\nabla : \mathcal{L} \rightarrow \mathcal{A}t(\mathcal{E})$ which satisfies the usual Leibnitz rule. We extend the notion of \mathcal{L} -connection and curvature of an \mathcal{O}_X -module \mathcal{E} to $(\mathcal{L}', \mathcal{L})$ -connections (and its curvature), for an \mathcal{L} -Lie algebroid \mathcal{L}' . For a smooth Lie algebroid L with a BV-algebra structure on the space of sections of the exterior bundle $\wedge^{\bullet} L$, P. Xu defined a notion of Lie algebroid homology of L with coefficient in line bundle $\wedge^n L$ (rank $L = n$) (see [44]). We consider a modified version of BV-algebra structure on $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}$ associated with a G -algebra structure on $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}'$ (see Theorem 6.7). We associate it with a \mathcal{L} -Lie algebroid structure on \mathcal{L}' with a flat $(\mathcal{L}', \mathcal{L})$ -connection on $\wedge_{\mathcal{O}_X}^n \mathcal{L}$, where $rank(\mathcal{L}) = n$. This defines a notion of homology for \mathcal{L} -Lie algebroids.

As a consequence of the BV-algebra structure on the exterior algebra of a smooth Lie algebroid, a duality theorem between its homology and cohomology was initially established by P. Xu ([44]) for a specific case and later generalized by J. Huebschmann ([20,21]) to the broader framework of Lie-Rinehart algebras. Building on these advancements, we extend this duality correspondence (see Theorem 8.9), from the context of finitely generated projective Lie-Rinehart algebras to the context of \mathcal{L} -Lie algebroids.

The universal enveloping algebroid of a Lie algebroid serves as an analogue of the sheaf of differential operators on a manifold (see [8,32,38,44]). It is equipped with a canonical filtered \mathbb{K}_X -algebra structure as well as an \mathcal{O}_X -bimodule structure (see [8,9,32]). This notion plays a pivotal role in exploring the homological algebra and representation theory of Lie algebroids (see [5,6,34,37]). A key ingredient in this is a version of the Poincaré-Birkhoff-Witt (PBW) theorem for Lie-Rinehart algebras, first established by G. Rinehart [36]. Additionally, a sheaf-theoretic analogue of this result for Lie algebroids has been developed (see [32,34,38]). We extend this framework to incorporate \mathcal{L} -Lie algebroids. Specifically, we generalize the PBW theorem to the setting of \mathcal{L} -Lie algebroids (see Theorem 7.7).

2. LIE ALGEBROIDS OVER (X, \mathcal{O}_X)

In this section, we recall some standard notions, namely topological ringed spaces, Lie-Rinehart algebras, Lie algebroids, Gerstenhaber algebras, Chevalley-Eilenberg-de Rham complex, Lie bialgebroids, BV-algebras and so on. Then, we recall some important relationships among them. In the later sections, we extend these notions in the algebro-geometric language and consider the analogous relationships in a more general setting.

2.1. Lie algebroids over topological ringed spaces. To define topological ringed spaces and Lie algebroids over them, we establish the following notation. Let \mathbb{K} denote either \mathbb{R} or \mathbb{C} (the real or complex number fields, respectively) or, an algebraically closed field of characteristic zero. The constant sheaf on a base space X , with stalks isomorphic to \mathbb{K} , is denoted by \mathbb{K}_X . The sheaf of \mathbb{K} -valued continuous functions on X , where \mathbb{K} is equipped with either the Euclidean or Zariski topology, is represented by C_X^0 .

Definition 2.1. *Let X be a topological space and \mathcal{O}_X be a \mathbb{K}_X -subalgebra of the sheaf of \mathbb{K} -algebras C_X^0 . Then the pair (X, \mathcal{O}_X) is said to be a topological ringed space.*

Remark 2.2. *An appropriate choice for the sheaf \mathcal{O}_X (known as the structure sheaf) over X provides various examples of topological ringed spaces (X, \mathcal{O}_X) coming from smooth manifolds, complex manifolds, analytic spaces and algebraic varieties. These topological ringed spaces are special kind of locally ringed spaces [43], we refer such spaces as special spaces.*

For a topological ringed space (X, \mathcal{O}_X) , the sheaf of derivations of \mathcal{O}_X is denoted by $Der_{\mathbb{K}_X}(\mathcal{O}_X)$. This is the subsheaf of the sheaf endomorphisms $End_{\mathbb{K}_X}(\mathcal{O}_X)$ on \mathcal{O}_X satisfies the Leibniz rule. More precisely, for an open set U of X , $Der_{\mathbb{K}_X}(\mathcal{O}_X)(U) := \{D : \mathcal{O}_U \rightarrow \mathcal{O}_U \text{ is a derivation}\} \subset End_{\mathbb{K}_X}(\mathcal{O}_U)$, where $\mathcal{O}_U = \mathcal{O}_X|_U$. That is, a section D of $Der_{\mathbb{K}_X}(\mathcal{O}_X)$ on U is a \mathbb{K} -linear derivation D of $\mathcal{O}_X(U)$ (or, $D \in Der_{\mathbb{K}}(\mathcal{O}_X(U))$) such that for open subsets W, V of U with $W \subset V$, satisfies the compatibility condition:

$$res_{VW}^{\mathcal{O}_X} \circ D_V = D_W \circ res_{VW}^{\mathcal{O}_X},$$

where $D_{V'} = D|_{V'}$, i.e. $D_{V'} = res_{UV'}^{End_{\mathbb{K}_X}(\mathcal{O}_X)}(D)$, for an open subset V' of U .

The sheaf $Der_{\mathbb{K}_X}(\mathcal{O}_X)$ over such a space (X, \mathcal{O}_X) has an canonical \mathcal{O}_X -module and \mathbb{K}_X -Lie algebra structure, which are compatible via a Leibniz rule [22,34,35]. For the special spaces, it is known as the tangent sheaf of X , and denoted by \mathcal{T}_X .

Remark 2.3. *A special topological ringed space (X, \mathcal{O}_X) is said to be smooth or nonsingular if its tangent sheaf $Der_{\mathbb{K}_X}(\mathcal{O}_X)$ is a locally free \mathcal{O}_X -module of finite rank.*

Recall that, a (\mathbb{K}, A) -Lie-Rinehart algebra L is a \mathbb{K} -Lie algebra L with an A -module structure (where A is a commutative unital \mathbb{K} -algebra) and an A -module homomorphism $\rho : L \rightarrow Der_{\mathbb{K}}(A)$ such that

- the map ρ is also a \mathbb{K} -Lie algebra homomorphism,
- $[x, ry] = r[x, y] + \rho(x)(r)y$ for all $x, y \in L, r \in A$.

The standard notion of Lie algebroids is defined on a smooth (real) vector bundle over a smooth manifold [30]. The space of global sections of a Lie algebroid forms a Lie-Rinehart algebra, capturing the entire structure [19]. In contrast, for holomorphic (or algebraic) vector bundles over complex manifolds (or algebraic varieties), the space of global sections does not capture the whole information. A sheaf-theoretic approach is necessary for analytic and algebraic geometry, applicable to both singular and non-singular cases.

Remark 2.4. *For a Lie algebroid L over a smooth manifold M , the space of sections $\Gamma(L|_U)$ of L on an open set $U \subset X$, forms an $(\mathbb{R}, C^\infty(U))$ -Lie-Rinehart algebra. In addition, the underlying $C^\infty(U)$ -module structure on $\Gamma(L|_U)$ is finitely generated (f.g.) projective. Moreover, a f.g. projective $(\mathbb{R}, C^\infty(M))$ -Lie-Rinehart algebra produces a smooth Lie algebroid over M . (Using an analogue of the Serre-Swan theorem, see [18]).*

We recall the definition of a Lie algebroid over a topological ringed space, which is a common generalization of the notion of tangent sheaf and a sheaf of Lie algebras (see [32,41]).

Definition 2.5. *(Lie algebroids over a topological ringed space) A Lie algebroid \mathcal{L} over a topological ringed space (X, \mathcal{O}_X) is a sheaf of \mathbb{K} -Lie algebras on X and an \mathcal{O}_X -module equipped with a homomorphism $\mathfrak{a} : (\mathcal{L}, [\cdot, \cdot]) \rightarrow (Der_{\mathbb{K}_X}(\mathcal{O}_X), [\cdot, \cdot]_c)$ of \mathcal{O}_X -modules and \mathbb{K}_X -Lie algebras, called the anchor map. The map \mathfrak{a} satisfies the Leibniz rule: $[D, fD'] = f[D, D'] + \mathfrak{a}(D)(f)D'$ for all $f \in \mathcal{O}_X$ and $D, D' \in \mathcal{L}$.*

Therefore, a Lie algebroid \mathcal{L} over (X, \mathcal{O}_X) can be viewed as a sheaf of $(\mathbb{K}_X, \mathcal{O}_X)$ -Lie-Rinehart algebras. We denote a Lie algebroid by $(\mathcal{L}, [\cdot, \cdot], \mathfrak{a})$ or simply by \mathcal{L} .

A morphism of Lie algebroids over a topological ringed space

$$\phi : (\mathcal{L}_1, [\cdot, \cdot]_1, \mathfrak{a}_1) \rightarrow (\mathcal{L}_2, [\cdot, \cdot]_2, \mathfrak{a}_2)$$

is a sheaf homomorphism of Lie-Rinehart algebras, i.e.

- $\phi : \mathcal{L}_1 \rightarrow \mathcal{L}_2$ is an \mathcal{O}_X -linear map,
- $\phi : (\mathcal{L}_1, [\cdot, \cdot]_1) \rightarrow (\mathcal{L}_2, [\cdot, \cdot]_2)$ is a \mathbb{K}_X -Lie algebra homomorphism,
- compatibility: $\mathfrak{a}_2 \circ \phi = \mathfrak{a}_1$.

Remark 2.6. Let (X, \mathcal{O}_X) be a special ringed space. There is a one-to-one correspondence between vector bundles over X (in the respective category) and locally free sheaves of \mathcal{O}_X -modules of finite rank.

In fact, the correspondence mentioned above is an equivalence of categories (see [34,35,43]).

Remark 2.7. For a locally free \mathcal{O}_X -module \mathcal{E} of finite rank n , the following properties hold:

- (1) For each $x \in X$, there exists an open neighborhood U_x of x in X such that $\mathcal{E}|_{U_x} \cong \mathcal{O}_X|_{U_x}^{\oplus n}$.
- (2) For each $x \in X$, the stalk \mathcal{E}_x is a free $\mathcal{O}_{X,x}$ -module of rank n .

Moreover, if \mathcal{E} is equipped with a Lie algebroid structure, this induces a Lie-Rinehart algebra structure on the space of sections of \mathcal{E} as well as on its stalks, retaining the above properties.

Using a standard approach, for a (locally free) Lie algebroid \mathcal{L} over (X, \mathcal{O}_X) , we mainly apply the algebraic properties of the $(\mathbb{K}, \mathcal{O}_X(U))$ -Lie-Rinehart algebra $\mathcal{L}(U)$ (which are free or projective module over $\mathcal{O}_X(U)$) at the local level to derive global results (see [5,6,8,9,22,32]).

Remark 2.8. A Lie algebroid over a smooth manifold (commonly referred to as a smooth Lie algebroid) X is equivalent to a locally free Lie algebroid of finite rank over the topological ringed space (X, C_X^∞) , where C_X^∞ denotes the sheaf of real valued smooth functions on X .

A Lie algebroid over a complex manifold X that is locally free of finite rank as an \mathcal{O}_X -module (\mathcal{O}_X is the sheaf of holomorphic functions on X) is the equivalent to a holomorphic Lie algebroid (see [34,39]).

Example 2.9. The standard Lie algebroid structure on the tangent sheaf of a real smooth manifold (or complex manifold) (X, \mathcal{O}_X) is described by the triple $(\text{Der}_{\mathbb{K}_X}(\mathcal{O}_X), [\cdot, \cdot]_c, id) =: \mathcal{T}_X$, where \mathcal{O}_X is the sheaf of smooth (or holomorphic) functions on X . The sheaf of smooth (or holomorphic) vector fields \mathfrak{X}_X on X is isomorphic to $\text{Der}_{\mathbb{K}_X}(\mathcal{O}_X)$, forming a locally free Lie algebroid of rank equal to the (real or complex) dimension of the manifold X .

Example 2.10. The sheaf of differential 1-forms (the cotangent sheaf) Ω_X^1 over a Poisson manifold X with Poisson bi-vector field $\pi \in H^0(X, \wedge^2 \mathcal{T}_X)$, has a canonical Lie algebroid structure, described as follows. The anchor map $\tilde{\pi} : \Omega_X^1 \rightarrow \mathcal{T}_X$ is defined by $\tilde{\pi}(\omega)(f) = \pi(\omega, df)$ for $f \in \mathcal{O}_X$, and the Lie bracket on Ω_X^1 is defined as $[\omega, \omega'] := L_{\tilde{\pi}(\omega)}\omega' - L_{\tilde{\pi}(\omega')}\omega - d(\pi(\omega, \omega'))$, for $f \in \mathcal{O}_X$ and $\omega, \omega' \in \Omega_X^1$ (see [14,34]).

Example 2.11. A (possibly singular) foliation \mathcal{F} on a real smooth manifold or a complex manifold (X, \mathcal{O}_X) is an \mathcal{O}_X -submodule of the Lie algebroid \mathcal{T}_X (or \mathfrak{X}_X), which is (i) stable under the Lie bracket and (ii) locally finitely generated. It provides a Stefan-Sussmann distribution or Nagano's distribution on X respectively, and forms a Lie algebroid over (X, \mathcal{O}_X) . These are coherent involutive subsheaves of the tangent sheaf. In particular, regular foliations arise from involutive subbundles of the tangent bundle (providing Frobenius distributions) are such examples. See [14,26,34] for more details.

Example 2.12. Let X be a complex manifold and \mathcal{O}_X denote the sheaf of holomorphic functions. The vanishing set (or the zero locus) $Y := V(\mathcal{I})$ of an ideal-sheaf $\mathcal{I} \subset \mathcal{O}_X$ is a subspace of X , which is not necessarily a submanifold and may have singularities. The sheaf of functions $\mathcal{O}_Y := \mathcal{O}_X/\mathcal{I}$ on Y serves as its structure sheaf, and the pair (Y, \mathcal{O}_Y) constitutes an analytic space (see in [34]).

Here, the tangent sheaf $\mathcal{T}_Y := (\text{Der}_{\mathbb{C}_Y}(\mathcal{O}_Y), [\cdot, \cdot]_c, id_Y)$ of Y is not necessarily a locally free Lie algebroid over (Y, \mathcal{O}_Y) (for example, if Y is a normal crossing divisor). Consider the sheaf of logarithmic derivations

(forms a generalized involutive distribution or singular foliation) [32,34] as

$$\mathcal{T}_X(-\log Y) := \{D \in \mathcal{T}_X \mid D(\mathcal{I}) \subset \mathcal{I}\} \hookrightarrow \mathcal{T}_X$$

(geometrically, it represents the sheaf of the vector fields on X whose flow preserves the divisor Y) with the canonical Lie algebroid structure. It is associated with \mathcal{T}_Y via the canonical map

$$\rho : \mathcal{T}_X(-\log Y) \rightarrow \mathcal{T}_Y$$

defined by $\rho(gD) = \bar{g}\tilde{D}$, for $g \in \mathcal{O}_X$, $D \in \mathcal{T}_X(-\log Y)$ as $\tilde{D}(\bar{f}) = D(\bar{f})$, where $\bar{h} = h + \mathcal{I} \in \mathcal{O}_Y$ for $h \in \mathcal{O}_X$.

There exist examples of Lie algebroids, such as free Lie algebroids [22] and sheaf of primitive elements [32], where the underlying module structure is not a coherent sheaf but rather a quasi-coherent sheaf.

2.2. Gerstenhaber algebras over a topological ringed space. In the study of deformations of associative algebras, M. Gerstenhaber discovered an algebraic structure ([16,17]), which is now known as the Gerstenhaber algebra. The notion of Gerstenhaber algebras also appears in the study of the space of multisections of a Lie algebroid over a smooth manifold [9,44]. These notions are appeared in the study of formality theory of Lie algebroids [7], generalizes the theory of deformation quantization (see [38]). We consider the notion of a sheaf of Gerstenhaber algebras as follows.

Definition 2.13. A sheaf of Gerstenhaber algebras on X or a Gerstenhaber algebra over (X, \mathcal{O}_X) consists of a triple $(\mathcal{A} := \bigoplus_{i \in \mathbb{Z}} \mathcal{A}^i, \wedge, [\cdot, \cdot])$ where each \mathcal{A}^i are \mathcal{O}_X -modules satisfying the following conditions:

- (\mathcal{A}, \wedge) is a graded commutative associative \mathcal{O}_X -algebra.
- $(\mathcal{A}[1], [\cdot, \cdot])$ is a graded \mathbb{K}_X -Lie algebra, where $\mathcal{A}[1] := \bigoplus_{i \in \mathbb{Z}} \mathcal{A}^{(i)}$ with $\mathcal{A}^{(i)} = \mathcal{A}^{i+1}$.
- For each $a \in \mathcal{A}^{(i)}$ the section $[a, \cdot]$ is a derivation with respect to \wedge with degree i , i.e. the operations \wedge and $[\cdot, \cdot]$ are compatible through the following Leibniz rule: for $a \in \mathcal{A}^i$, $b \in \mathcal{A}^j$ and $c \in \mathcal{A}$,

$$[a, b \wedge c] = [a, b] \wedge c + (-1)^{(i-1)j} b \wedge [a, c].$$

Remark 2.14. (Sheaves of) Gerstenhaber algebras are generalization (or graded analogue) of (sheaves of) Poisson algebras.

A morphism $\psi : (\mathcal{A} := \bigoplus_{i \in \mathbb{Z}} \mathcal{A}^i, \wedge, [\cdot, \cdot]) \rightarrow (\mathcal{A}' := \bigoplus_{i \in \mathbb{Z}} \mathcal{A}'^i, \wedge', [\cdot, \cdot]')$ of Gerstenhaber algebras over a topological ringed space (X, \mathcal{O}_X) is given as follows.

- The map $\psi : (\mathcal{A}, \wedge) \rightarrow (\mathcal{A}', \wedge')$ is a graded \mathcal{O}_X -algebra homomorphism, and
- the map $\psi : (\mathcal{A}[1], [\cdot, \cdot]) \rightarrow (\mathcal{A}'[1], [\cdot, \cdot]')$ is a graded \mathbb{K}_X -algebra homomorphism.

Remark 2.15. In most of the cases, we consider $\mathcal{A}^i = 0$ for $i < 0$ and $\mathcal{A}^0 = \mathcal{O}_X$ for a Gerstenhaber algebra $(\mathcal{A} := \bigoplus_{i \in \mathbb{Z}} \mathcal{A}^i, \wedge, [\cdot, \cdot])$ over (X, \mathcal{O}_X) . Notice that for such a Gerstenhaber algebra $(\mathcal{A} := \bigoplus_{i \geq 0} \mathcal{A}^i, \wedge, [\cdot, \cdot])$, the \mathcal{O}_X -module \mathcal{A}^1 canonically forms a Lie algebroid over (X, \mathcal{O}_X) , as described follows. The restriction of the G -bracket gives the \mathbb{K}_X -Lie algebra $(\mathcal{A}^1, [\cdot, \cdot]_{\mathcal{A}^1})$, and restricting the multiplication $\wedge : \mathcal{O}_X \otimes_{\mathbb{K}_X} \mathcal{A}^1 \rightarrow \mathcal{A}^1$ gives the \mathcal{O}_X -module structure. The anchor map is given by the assignment $D \mapsto [D, \cdot]$ for $D \in \mathcal{A}^1$.

This construction globalizes the algebraic counterpart discussed in [17,21].

Example 2.16. For a topological ringed space (X, \mathcal{O}_X) , the sheaf of multiderivations over \mathcal{O}_X is

$$\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{T}_X := \left(\bigoplus_{i=0}^{\infty} \wedge^i_{\mathcal{O}_X} \mathcal{T}_X, \wedge \right),$$

forms a sheaf of Gerstenhaber algebras. It is the sheafification of the presheaf $U \mapsto \wedge^{\bullet}_{\mathcal{O}_X(U)} \mathcal{T}_X(U)$ of exterior algebras for $\mathcal{T}_X(U)$ over $\mathcal{O}_X(U)$, where U varies in open sets of X . In particular, when X is a non-singular space, this is equivalent to the sheaf of multivector fields $\wedge^{\bullet}_{\mathcal{O}_X} \mathfrak{X}_X$ on X .

The Gerstenhaber structure is given by the Schouten-Nijenhuis bracket

$$[\cdot, \cdot] : \wedge^{\bullet}_{\mathcal{O}_X} \mathcal{T}_X[1] \otimes_{\mathcal{O}_X} \wedge^{\bullet}_{\mathcal{O}_X} \mathcal{T}_X[1] \rightarrow \wedge^{\bullet}_{\mathcal{O}_X} \mathcal{T}_X[1],$$

which is defined as:

$$[D_1 \wedge \cdots \wedge D_m, D_{m+1} \wedge \cdots \wedge D_n] := \sum_{i \leq m < j} (-1)^{i+j+m} [D_i, D_j]_c \wedge D_1 \wedge \cdots \wedge \hat{D}_i \cdots \wedge \hat{D}_j \cdots \wedge D_n,$$

where $D_i \in \mathcal{T}_X$ for $i = 1, \dots, n$, and $[\cdot, \cdot]_c$ denotes the commutator Lie bracket on \mathcal{T}_X .

Example 2.17. For a Lie algebroid \mathcal{L} over (X, \mathcal{O}_X) , the standard Gerstenhaber algebra associated to \mathcal{L} or the standard Gerstenhaber algebra structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}$ is given by the triple $\mathcal{G}_{\mathcal{L}} := (\bigoplus_{i=0}^\infty \wedge_{\mathcal{O}_X}^i \mathcal{L}, \wedge, [\cdot, \cdot]_{\mathcal{L}})$, where the graded \mathbb{K}_X -Lie bracket $[\cdot, \cdot]_{\mathcal{L}}$ is the generalization of the Schouten-Nijenhuis bracket in the context of Lie algebroids, replacing the commutator bracket $[\cdot, \cdot]_c$ of \mathcal{T}_X by the Lie bracket $[\cdot, \cdot]_{\mathcal{L}}$ of \mathcal{L} .

In the following we describe an algebro-geometric counterpart of the relationship between Lie algebroids and Gerstenhaber algebras (see [20,44] for the smooth and algebraic case).

Theorem 2.18. For an \mathcal{O}_X -module \mathcal{E} , there is a (sheaf of) Gerstenhaber algebra structure on the exterior algebra $\wedge_{\mathcal{O}_X}^\bullet \mathcal{E}$ if and only if \mathcal{E} has a Lie algebroid structure.

2.3. Chevalley-Eilenberg-de Rham complex. The Chevalley–Eilenberg–de Rham complex is commonly used to study the cohomology of Lie algebroids over smooth manifolds [30]. Additionally, its algebro-geometric counterparts are discussed in [5,6,34]. Here, we present a sheaf-theoretic generalization of this cochain complex, using the Lie-Rinehart cochain complex [36].

For a Lie algebroid \mathcal{L} over (X, \mathcal{O}_X) , first recall some of the key notions in the following. Denote ${}^p \text{Hom}_{\mathcal{O}_X(U)}(\wedge_{\mathcal{O}_X(U)}^\bullet \mathcal{L}(U), \mathcal{O}_X(U))$ (or in short ${}^p(\wedge_{\mathcal{O}_X(U)}^\bullet \mathcal{L}(U))^*$ by the module of $\mathcal{O}_X(U)$ -linear maps from $\wedge_{\mathcal{O}_X(U)}^\bullet \mathcal{L}(U)$ to $\mathcal{O}_X(U)$ which induces $\mathcal{O}_X(V)$ -module homomorphisms from $\wedge_{\mathcal{O}_X(V)}^\bullet \mathcal{L}(V)$ to $\mathcal{O}_X(V)$ for each open subset V in U , and are compatible with restriction maps of the presheaf $U \mapsto \wedge_{\mathcal{O}_X(U)}^\bullet \mathcal{L}(U)$ and of the sheaf \mathcal{O}_X . Thus, in the underlying process, we consider the presheaf ${}^p \wedge_{\mathcal{O}_X}^\bullet \mathcal{L}$ given by $U \mapsto \wedge_{\mathcal{O}_X(U)}^\bullet \mathcal{L}(U)$ together with all the presheaf homomorphisms from ${}^p \wedge_{\mathcal{O}_X}^\bullet \mathcal{L}$ to \mathcal{O}_X .

Consider the cochain complex of \mathcal{O}_X -modules associated to a Lie algebroid \mathcal{L}

$$(1) \quad \Omega_{\mathcal{L}}^\bullet := (\mathcal{H}om_{\mathcal{O}_X}(\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}, \mathcal{O}_X), d_{\mathcal{L}}).$$

It is the sheafification of the presheaf of cochain complexes (see [42] for such notions)

$$U \mapsto ({}^p \text{Hom}_{\mathcal{O}_X(U)}(\wedge_{\mathcal{O}_X(U)}^\bullet \mathcal{L}(U), \mathcal{O}_X(U)), d_{\mathcal{L}(U)}) =: \Omega_{\mathcal{L}(U)}^\bullet,$$

and the differential $d_{\mathcal{L}(U)} : {}^p(\wedge_{\mathcal{O}_X(U)}^\bullet \mathcal{L}(U))^* \rightarrow {}^p(\wedge_{\mathcal{O}_X(U)}^{\bullet+1} \mathcal{L}(U))^*$ is given by

$$\begin{aligned} d_{\mathcal{L}(U)}(\omega)(D_1 \wedge \cdots \wedge D_{k+1}) &= \sum_{i=1}^{k+1} (-1)^{i+1} \mathbf{a}(U)(D_i)(\omega(D_1 \wedge \cdots \wedge \hat{D}_i \wedge \cdots \wedge D_{k+1})) \\ &\quad + \sum_{i < j} (-1)^{i+j} \omega([D_i, D_j] \wedge D_1 \wedge \cdots \wedge \hat{D}_i \wedge \cdots \wedge \hat{D}_j \wedge \cdots \wedge D_{k+1}), \end{aligned}$$

where $D_1, \dots, D_{k+1} \in \mathcal{L}(U)$ and $\omega \in \Omega_{\mathcal{L}(U)}^k$, associated with an open subset U of X .

Equivalently one can consider, for an open set $U \subset X$, $\Omega_{\mathcal{L}}^\bullet(U) = (\text{Hom}_{\mathcal{O}_U}(\wedge_{\mathcal{O}_U}^\bullet \mathcal{L}_U, \mathcal{O}_U), d_{\mathcal{L}(U)})$, where a section $\omega \in \text{Hom}_{\mathcal{O}_U}(\wedge_{\mathcal{O}_U}^\bullet \mathcal{L}_U, \mathcal{O}_U)$ is a sheaf homomorphism (\mathcal{O}_U -linear map) compatible with restriction of the sheaf $V \mapsto \text{Hom}_{\mathcal{O}_V}(\wedge_{\mathcal{O}_V}^\bullet \mathcal{L}_V, \mathcal{O}_V)$ and the differential $V \mapsto d_{\mathcal{L}(V)}$, for open subset $V \subset U$.

Notice that, the differential $d_{\mathcal{L}}$ is a \mathbb{K}_X -linear map but not an \mathcal{O}_X -module homomorphism, satisfies the graded Leibniz rule: $d(\omega_1 \wedge \omega_2) = d\omega_1 \wedge \omega_2 + (-1)^k \omega_1 \wedge d\omega_2$, for $\omega \in \Omega_{\mathcal{L}(U)}^k$ and $\omega \in \Omega_{\mathcal{L}(U)}^l$, $k, l \in \mathbb{N} \cup \{0\}$.

Note that if \mathcal{L} is a locally free \mathcal{O}_X -modules of finite rank then we have

$$\mathcal{H}om_{\mathcal{O}_X}(\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}, \mathcal{O}_X) \cong \wedge_{\mathcal{O}_X}^\bullet \mathcal{L}^*.$$

This complex is equivalent to the standard complex associated with a smooth Lie algebroid [30] or a holomorphic Lie algebroid [34]. In particular, we get the Chevalley-Eilenberg complex when \mathcal{L} is a Lie algebroid over a point; and the de Rham complex Ω_X^\bullet when $\mathcal{L} = \mathcal{T}_X$ over a nonsingular space X . In general, the complex Ω_X^\bullet over (X, \mathcal{O}_X) produces *Zariski differentials*, not necessarily *Kähler differentials* (see [19]).

Remark 2.19. *In general, over a Noetherian separated scheme (algebraic varieties or analytic spaces) (X, \mathcal{O}_X) , one can consider the cochain complex of Kähler differentials $(\wedge^{\bullet}_{\mathcal{O}_X} \Omega_X^1, d_X)$ where Ω_X^1 is the \mathcal{O}_X -module of Kähler differentials 1-forms on X and the differential d_X is the extension of the canonical map $d_X^1 : \mathcal{O}_X \rightarrow \Omega_X^1$. But, notice that if Ω_X^1 is reflexive then $(\mathcal{T}_X)^* = (\Omega_X^1)^{**} \cong \Omega_X^1$ as \mathcal{O}_X -modules, recover the standard case for the Lie algebroid complex of $\mathcal{L} = \mathcal{T}_X$ discussed in [5,34].*

Next, we consider an algebro-geometric analogue of the correspondence between Lie algebroid structure on a smooth vector bundle and differential graded algebra structure on the exterior algebra of its dual bundle (see [25,44]).

Theorem 2.20. *Given a locally free \mathcal{O}_X -module \mathcal{E} of finite rank, \mathcal{E} has a Lie algebroid structure if and only if the exterior algebra of sheaf of multisections $\wedge^{\bullet} \mathcal{E}^* := \bigoplus_{k=0}^{\infty} \wedge^k \mathcal{E}^*$ of $\mathcal{E}^* = \mathcal{H}om_{\mathcal{O}_X}(\mathcal{E}, \mathcal{O}_X)$ forms a sheaf of differential graded algebras.*

First, we recall the notion of *hypercohomology* (see [11,42]) in the following, to consider an analogue of the Chevalley-Eilenberg-de Rham cohomology.

Let (X, \mathcal{O}_X) be a topological ringed space with \mathcal{O}_X -modules \mathcal{C}^i , for $i \geq 0$. Consider a cochain complex $(\mathcal{C}^{\bullet}, d)$ where $\mathcal{C}^{\bullet} := \bigoplus_i \mathcal{C}^i$ is a graded \mathcal{O}_X -module and $d : \mathcal{C}^{\bullet} \rightarrow \mathcal{C}^{\bullet+1}$ is the co-boundary map, i.e., d is a \mathbb{K}_X -linear map of degree 1 satisfying $d^2 = 0$. Similarly, for a chain complex, instead of a co-boundary map we have a boundary map $\partial : \mathcal{C}^{\bullet} \rightarrow \mathcal{C}^{\bullet-1}$, a \mathbb{K}_X -linear map of degree -1 satisfying $\partial^2 = 0$. To address the following topics of discussion consistently, we consider such (co)chain complexes where \mathcal{O}_X -linearity of the (co)boundary maps is not necessarily required.

Remark 2.21. *For an \mathcal{O}_X -module \mathcal{F} on X , there exists an injective resolution $\mathcal{F} \xrightarrow{\sim} \mathcal{C}^{\bullet}(\mathcal{F})$ (a quasi-isomorphism) of \mathcal{O}_X -modules, known as the flabby Godement resolution of \mathcal{F} . Thus, for a complex of \mathcal{O}_X -modules \mathcal{F}^{\bullet} , we consider the associated bicomplex of sheaves $\mathcal{C}^{\bullet}(\mathcal{F}^{\bullet}) = (\mathcal{C}^p(\mathcal{F}^q))$ ($p, q \in \mathbb{N} \cup \{0\}$) of injective resolutions. The original complex is embedded in the total complex $\mathcal{K}^{\bullet} = \text{tot}(\mathcal{C}^{\bullet}(\mathcal{F}^{\bullet}))$, and this embedding is a quasi-isomorphism. The cohomology of the associated complex of global sections $\mathcal{K}^{\bullet}(X) = \text{tot}(\mathcal{C}^{\bullet}(\mathcal{F}^{\bullet}))(X)$ is called the hypercohomology of \mathcal{F}^{\bullet} , and denoted by $\mathbb{H}^{\bullet}(X, \mathcal{F}^{\bullet})$.*

Denote the category of cochain complexes of sheaves of \mathcal{O}_X -modules over (X, \mathcal{O}_X) by $\text{Shv}(X)^{\bullet}$ and the category of \mathbb{K} -vector spaces by Vect . For $\mathcal{F}^{\bullet} \in \text{Shv}(X)^{\bullet}$, the k -th cohomology sheaf of the complex \mathcal{F}^{\bullet} is

$$\mathcal{H}^k(\mathcal{F}^{\bullet}) := \mathcal{K}er(\mathcal{F}^k \rightarrow \mathcal{F}^{k+1}) / \mathcal{I}m(\mathcal{F}^{k-1} \rightarrow \mathcal{F}^k)$$

(where $\mathcal{F}^{-1} := 0$) for $k \in \mathbb{N} \cup \{0\}$, considered as quotient of \mathbb{K}_X -vector spaces. Moreover, a map of complexes $\mathcal{F}^{\bullet} \rightarrow \mathcal{G}^{\bullet}$ is a quasi-isomorphism if the induced map on cohomology sheaves $\mathcal{H}^k(\mathcal{F}^{\bullet}) \rightarrow \mathcal{H}^k(\mathcal{G}^{\bullet})$ is an isomorphism for all k .

The k -th hypercohomology is a functor $\mathbb{H}^k(X, -) : \text{Shv}(X)^{\bullet} \rightarrow \text{Vect}$ for $k \in \mathbb{N} \cup \{0\}$ that satisfies the following two conditions:

- A quasi-isomorphism of complexes $f^{\bullet} : \mathcal{F}^{\bullet} \rightarrow \mathcal{G}^{\bullet}$ induces an isomorphism $\mathbb{H}^k(X, f^{\bullet})$.
- If \mathcal{I}^{\bullet} is a complex of injective sheaves then $\mathbb{H}^k(X, \mathcal{I}^{\bullet}) = H^k(\Gamma(X, \mathcal{I}^{\bullet}))$, where $\Gamma(X, \mathcal{I}^{\bullet})$ is the space of global sections of \mathcal{I}^{\bullet} .

Denote the graded vector space $\bigoplus_{n=0}^{\infty} \mathbb{H}^n(X, \mathcal{F}^{\bullet})$ as $\mathbb{H}^{\bullet}(X, \mathcal{F}^{\bullet})$.

The hypercohomology of the cochain complex (1) of \mathcal{O}_X -modules is called the Lie algebroid cohomology or Chevalley-Eilenberg-de Rham cohomology of \mathcal{L} with trivial coefficient and is denoted by $\mathbb{H}^{\bullet}(\mathcal{L}, \mathcal{O}_X)$.

Remark 2.22. *Later, we consider an analogue of the notion of Lie algebroid homology ([20,44]), which is obtained from a similar version of the above description of hypercohomology in the context of chain complexes of sheaves (see [42]), which we call the hyperhomology of the Lie algebroid.*

2.4. Lie bialgebroids over a topological ringed space. Lie bialgebroids over a smooth manifold, consists of two compatible Lie algebroids defined on dual vector bundles, is a standard notion appears in the study of Poisson geometry [25,30,31,44]. Here, we consider an analogue of it in the algebro-geometric settings.

Definition 2.23. A Lie bialgebroid over a topological ringed space (X, \mathcal{O}_X) is a pair $(\mathcal{L}, \mathcal{L}^*)$ of locally free Lie algebroids of finite rank in duality, where the Lie brackets satisfy a condition which can be expressed in terms of the differential $d_{\mathcal{L}^*} =: d_*$ on $\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{L}$ defined by the Lie algebroid structure of \mathcal{L}^* and the Gerstenhaber bracket $[\cdot, \cdot]$ on $\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{L}$ defined by the Lie algebroid structure on \mathcal{L} ,

$$d_*[D_1, D_2] = [d_*D_1, D_2] + [D_1, d_*D_2],$$

for all sections D_1, D_2 of \mathcal{L} .

In other words, a Lie bialgebroid over a topological ringed space (X, \mathcal{O}_X) is a dual pair $(\mathcal{L}, \mathcal{L}^*)$ of locally free Lie algebroids of finite rank such that the differential d_* , induced from the Lie algebroid structure on \mathcal{L}^* , is a derivation of the Lie bracket on \mathcal{L} .

Remark 2.24. The condition \mathcal{L}^* is a locally free \mathcal{O}_X -modules of finite rank implies

$$\mathrm{Hom}_{\mathcal{O}_X}(\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{L}^*, \mathcal{O}_X) \cong \wedge^{\bullet}_{\mathcal{O}_X} \mathcal{L}.$$

Example 2.25. For a Poisson manifold (X, π) , the dual pair $(\mathcal{T}_X, \Omega_X^1)$ forms a Lie bialgebroid, where the Lie bracket on the cotangent sheaf Ω_X^1 and the anchor map $\tilde{\pi} : \Omega_X^1 \rightarrow \mathcal{T}_X$ is given by the Poisson bivector field π on X (or, by the induced Poisson bracket $\{\cdot, \cdot\}$ on \mathcal{O}_X), see Example 2.10. More generally, there is an analogous notion, known as triangular Lie bialgebroid in the context of Lie algebroids [25].

Remark 2.26. Let $(\mathcal{L}, \mathcal{L}^*)$ be a pair of locally free Lie algebroids of finite rank in duality over a topological ringed space. Then $(\mathcal{L}, \mathcal{L}^*)$ is a Lie bialgebroid if and only if $(\mathcal{L}^*, \mathcal{L})$ is a Lie bialgebroid.

The notions Lie bialgebroids over a smooth manifold and strong differential Gerstenhaber algebras are interconnected [25, 44]. Here we define the notion strong differential Gerstenhaber algebras over a topological ringed space and extend such relationship.

Definition 2.27. A strong differential Gerstenhaber algebra over a topological ringed space (X, \mathcal{O}_X) is a Gerstenhaber algebra $\mathcal{A} := (\bigoplus_{i=0}^{\infty} \mathcal{A}^i, \wedge, [\cdot, \cdot])$ over (X, \mathcal{O}_X) equipped with a differential $\tilde{d} : \bigoplus_{i=0}^{\infty} \mathcal{A}^i \rightarrow \bigoplus_{i=0}^{\infty} \mathcal{A}^{i+1}$, which is a derivation of degree 1 with respect to \wedge satisfying $\tilde{d}^2 = 0$, and it is also a derivation of the graded Lie bracket $[\cdot, \cdot]$.

An algebro geometric analogue of the correspondence between these notions [25] is described as follows.

Theorem 2.28. The exterior algebra $\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{A}$ has a strong differential Gerstenhaber algebra structure over a topological ringed space (X, \mathcal{O}_X) for a locally free \mathcal{O}_X -module \mathcal{A} of finite rank if and only if the dual pair $(\mathcal{A}, \mathcal{A}^*)$ has a Lie bialgebroid over (X, \mathcal{O}_X) .

2.5. BV-algebras or exact Gerstenhaber algebras. The notion BV-algebras plays an important role in geometry and mathematical physics (see [20, 25, 44]). We consider a algebro-geometric analogue of it.

Definition 2.29. An exact Gerstenhaber algebra (or BV-algebra) over a topological ringed space (X, \mathcal{O}_X) is a Gerstenhaber algebra $(\mathcal{A} := \bigoplus_{i=0}^{\infty} \mathcal{A}^i, \wedge, [\cdot, \cdot])$ over (X, \mathcal{O}_X) equipped with an operator ∂ of degree -1 , square zero, and generating the Gerstenhaber bracket as

$$[a, b] = (-1)^i (\partial(a \wedge b) - \partial a \wedge b - (-1)^i a \wedge \partial b),$$

where $a \in \mathcal{A}^i \subset \mathcal{A}$ and $b \in \mathcal{A}$. The operator ∂ is called a generating operator.

We recall the notion of flat connections of a Lie algebroid using Atiyah algebroids [5, 22, 34, 37, 40].

For an \mathcal{O}_X -module \mathcal{E} , we form a Lie algebroid consisting of the sheaf of differential operators on \mathcal{E} of order ≤ 1 with scalar symbols, i.e.

$$\mathrm{At}(\mathcal{E}) := \{D \in \mathrm{End}_{\mathbb{K}_X}(\mathcal{E}) \mid D(fs) = fD(s) + \sigma_D(f)s \text{ for a unique } \sigma_D \in \mathcal{T}_X, \text{ where } f \in \mathcal{O}_X, s \in \mathcal{E}\},$$

(thus, here $\sigma_D(f) = [D, f]_c \in \mathcal{O}_X$ for $D \in \mathrm{At}(\mathcal{E})$ and $f \in \mathcal{O}_X$ holds) with the anchor map defined by

$$\sigma : \mathrm{At}(\mathcal{E}) \rightarrow \mathcal{T}_X \text{ where } D \mapsto \sigma_D$$

and the Lie bracket is commutator bracket. This Lie algebroid is the so-called Atiyah algebroid of the \mathcal{O}_X -module \mathcal{E} . It provides a short exact sequence (s.e.s.) of Lie algebroids over (X, \mathcal{O}_X) (an abelian Lie algebroid extension),

$$(2) \quad 0 \rightarrow \text{End}_{\mathcal{O}_X}(\mathcal{E}) \hookrightarrow \text{At}(\mathcal{E}) \xrightarrow{\sigma} \mathcal{T}_X \rightarrow 0.$$

In particular, when $\mathcal{E} = \mathcal{O}_X$ for a nonsingular space X , we have $\text{At}(\mathcal{O}_X) \cong \mathcal{O}_X \oplus \mathcal{T}_X$, equals to the sheaf differential operators on X of order ≤ 1 with scalar symbol.

Definition 2.30. Given Lie algebroid $(\mathcal{L}, [\cdot, \cdot], \mathfrak{a})$ on a topological ringed space (X, \mathcal{O}_X) , a \mathcal{L} -connection on an \mathcal{O}_X -module \mathcal{E} is defined by an \mathcal{O}_X -linear map

$$(3) \quad \begin{aligned} \nabla : \mathcal{L} &\rightarrow \text{At}(\mathcal{E}) \\ D &\mapsto \nabla_D \end{aligned}$$

satisfying the Leibniz rule $\nabla_D(f s) = f \nabla_D(s) + \mathfrak{a}(D)(f) s$ for sections $f \in \mathcal{O}_X$, $D \in \mathcal{L}$ and $s \in \mathcal{E}$ (see [5, 22, 34]). To ensure compatibility, we choose the condition $\sigma \circ \nabla = \mathfrak{a}$.

Remark 2.31. Equivalently, a \mathcal{L} -connection on \mathcal{E} is described by a \mathbb{K}_X -linear map, if in addition \mathcal{L} is a locally free \mathcal{O}_X -module of finite rank, as follows (see [34])

$$\nabla : \mathcal{E} \rightarrow \Omega_{\mathcal{L}}^1 \otimes_{\mathcal{O}_X} \mathcal{E}$$

satisfying the Leibniz rule $\nabla(f s) = f \nabla s + \check{\mathfrak{a}}(df) \otimes s$, where $\check{\mathfrak{a}} : \Omega_X^1 \rightarrow \Omega_{\mathcal{L}}^1$ is the dual of the anchor map.

Definition 2.32. A \mathcal{L} -connection on \mathcal{E} is said to be flat if the map (3) is a Lie algebroid homomorphism, i.e., the map (3) satisfies curvature zero condition: for $D, D' \in \mathcal{L}$

$$(4) \quad R_{\nabla}(D, D') := \nabla_{[D, D']} - [\nabla_D, \nabla_{D'}]_c = 0.$$

In that case, (\mathcal{E}, ∇) is said to be a representation of \mathcal{L} or an \mathcal{L} -module.

A connection $\nabla : \mathcal{T}_X \rightarrow \text{At}(\mathcal{E})$ that satisfies $\sigma \circ \nabla = \text{Id}_{\mathcal{T}_X}$ provides a splitting of the s.e.s. (2) as \mathcal{O}_X -modules. If its curvature is zero then the s.e.s. splits as Lie algebroids.

Next, we consider some special cases where X is a nonsingular space:

(i) For $\mathcal{L} = \mathcal{T}_X$ and \mathcal{E} a locally free \mathcal{O}_X -module of finite rank, a \mathcal{T}_X -connection on \mathcal{E} always exists if X either a smooth manifold, or a Stein manifold, or an affine variety (see [1]).

(ii) For $\mathcal{L} = \mathcal{T}_X(-\log Y)$, if a connection exists on an \mathcal{O}_X -module \mathcal{E} , it is called a logarithmic connection on \mathcal{E} . It forms a meromorphic connection with simple poles along the divisor Y (see [10, 34]). For instance, if X is a non-singular projective toric variety and Y is the boundary divisor, then a locally free \mathcal{O}_X -module \mathcal{E} admits a logarithmic connection if and only if it admits a torus equivariant structure (see [2, 3]).

An algebro-geometric analogue of the correspondence between these notions [44] is described as follows.

Theorem 2.33. Let \mathcal{L} be a locally free \mathcal{O}_X -module of rank n . Then there is an exact Gerstenhaber algebra (or BV-algebra) structure on the exterior algebra $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}$ if and only if there is a Lie algebroid structure on \mathcal{L} with a flat \mathcal{L} -connection on the \mathcal{O}_X -module $\wedge_{\mathcal{O}_X}^n \mathcal{L}$.

Remark 2.34. Let X be either a smooth manifold, or a Stein manifold, or an smooth affine variety, and \mathcal{O}_X be the associated structure sheaf (see [1, 15]). For a locally free Lie algebroid \mathcal{L} of rank n for some $n \in \mathbb{N}$, there always exists a flat \mathcal{L} -connection on $\wedge_{\mathcal{O}_X}^n \mathcal{L}$. Thus, there always exists a canonical BV-algebra structure on $\wedge_{\mathcal{O}_X}^n \mathcal{L}$ (see [44] for the smooth Lie algebroids case).

3. \mathcal{L} -LIE ALGEBROIDS

We define the notion of an \mathcal{L} -Lie algebroid for a given Lie algebroid \mathcal{L} over a topological ringed space (X, \mathcal{O}_X) . To generalize the relationship between Lie algebroids and sheaves of Gerstenhaber algebras (Theorem 2.18) in this context, we introduce the notion of an \mathcal{A} -Gerstenhaber algebra associated with a given Gerstenhaber algebra \mathcal{A} over (X, \mathcal{O}_X) .

Definition 3.1. Let $(\mathcal{L}, [\cdot, \cdot]_{\mathcal{L}}, \mathfrak{a})$ be a Lie algebroid over (X, \mathcal{O}_X) . A Lie algebroid $(\mathcal{L}', [\cdot, \cdot]_{\mathcal{L}'}, \mathfrak{a}')$ over (X, \mathcal{O}_X) is said to be a \mathcal{L} -Lie algebroid if there exists a Lie algebroid homomorphism

$$\phi : (\mathcal{L}', [\cdot, \cdot]_{\mathcal{L}'}, \mathfrak{a}') \rightarrow (\mathcal{L}, [\cdot, \cdot]_{\mathcal{L}}, \mathfrak{a}).$$

We denote this \mathcal{L} -Lie algebroid as (\mathcal{L}', ϕ) .

Example 3.2. (1) For a Lie algebroid \mathcal{L} , the pair (\mathcal{L}, id) forms a \mathcal{L} -Lie algebroid.

(2) Consider the tangent sheaf \mathcal{T}_X of a special ringed space (X, \mathcal{O}_X) (see Remark 2.2). Any Lie algebroid \mathcal{L} over (X, \mathcal{O}_X) is a \mathcal{T}_X -Lie algebroid. For example, the cotangent sheaf Ω_X^1 over a Poisson manifold X has a canonical \mathcal{T}_X -Lie algebroid structure.

(3) Any Lie subalgebroid \mathcal{L}' of a Lie algebroid \mathcal{L} naturally inherits the structure of an \mathcal{L} -Lie algebroid via the inclusion map $\iota : \mathcal{L}' \rightarrow \mathcal{L}$. Singular foliations offer such examples, including the sheaf of logarithmic derivations $\mathcal{T}_X(-\log Y) \subset \mathcal{T}_X$.

(4) For a triangular Lie bialgebroid $(\mathcal{L}^*, \mathcal{L}, \mathcal{P})$ where \mathcal{P} is the Poisson \mathcal{L} -bivector, can be viewed as a Lie algebroid homomorphism $\mathcal{P} : \mathcal{L}^* \rightarrow \mathcal{L}$ (see [25]). Therefore, $(\mathcal{L}^*, \mathcal{P})$ is a \mathcal{L} -Lie algebroid.

Example 3.3. Let G be a (smooth, holomorphic, or algebraic) Lie group acting on a G -manifold X , and let $\bar{\phi} : G' \rightarrow G$ be a Lie group homomorphism for another Lie group G' . This map $\bar{\phi}$ induces a natural G' -action on the G -manifold X . Denote the Lie algebra of a Lie group H by $\text{Lie}(H)$, and the associated constant sheaf by $\text{Lie}(H)_X$. Consider the associated transformation Lie algebroids $X \times \text{Lie}(G')$ and $X \times \text{Lie}(G)$ (see [14]) along with the canonical connecting map $\tilde{\phi} : X \times \text{Lie}(G') \rightarrow X \times \text{Lie}(G)$. Using the G -action on X , the map $\tilde{\phi}$ induces a Lie algebroid homomorphism over (X, \mathcal{O}_X) at the level of sheaves of sections, given by

$$\mathcal{O}_X \otimes_{\mathbb{K}_X} \text{Lie}(G')_X \rightarrow \mathcal{O}_X \otimes_{\mathbb{K}_X} \text{Lie}(G)_X.$$

Example 3.4. The notion of log Atiyah algebroid is appeared in the study of logarithmic connections on equivariant principal bundles (see [2, 12]) or on equivariant vector bundles [3]. Here, we recall the notion log Atiyah algebroid $\text{At}(\mathcal{E})(-\log Y)$ of an \mathcal{O}_X -module \mathcal{E} associated with an analytic subspace Y of a complex manifold X (or, a divisor Y of a smooth algebraic variety X), and describe it as a $\mathcal{T}_X(-\log Y)$ -Lie algebroid. First, consider the Atiyah algebroid $\text{At}(\mathcal{E})$ of \mathcal{E} and the Lie algebroid homomorphism $\iota : \mathcal{T}_X(-\log Y) \hookrightarrow \mathcal{T}_X$. The pull back of the Lie algebroid $\mathcal{T}_X(-\log Y)$ by the anchor map $\sigma : \text{At}(\mathcal{E}) \rightarrow \mathcal{T}_X$ is the log Atiyah algebroid of \mathcal{E} , i.e. $\text{At}(\mathcal{E})(-\log Y) = \sigma^{-1}(\mathcal{T}_X(-\log Y))$. Thus, $(\text{At}(\mathcal{E})(-\log Y), \bar{\sigma})$ is a $\mathcal{T}_X(-\log Y)$ -Lie algebroid, where $\bar{\sigma}$ is the restriction of the Lie algebroid homomorphism σ on $\text{At}(\mathcal{E})(-\log Y)$.

Remark 3.5. Consider the Atiyah algebroid $\text{At}(\mathcal{E})$ of an \mathcal{O}_X -module \mathcal{E} . For a given Lie algebroid \mathcal{L} , a flat \mathcal{L} -connection on \mathcal{E} is equivalent to have an $\text{At}(\mathcal{E})$ -Lie algebroid structure on \mathcal{L} (see Definitions 2.30, 2.32).

Example 3.6. Given a Lie algebroid $(\mathcal{L}, [\cdot, \cdot]_{\mathcal{L}}, \mathfrak{a})$ over (X, \mathcal{O}_X) and an element $[\omega] \in \mathbb{H}^2(\mathcal{L}, \mathcal{O}_X)$, we get a Lie algebroid extension of \mathcal{L} by \mathcal{O}_X (upto isomorphism) as follows (see [27, 39]).

Consider the \mathcal{O}_X -module $\mathcal{O}_X \oplus \mathcal{L}$ with the \mathbb{K}_X -Lie algebra structure given by

$$[(f, D), (f', D')]_{\omega} := (\mathfrak{a}(D)(f') - \mathfrak{a}(D')(f) + \omega(D, D'), [D, D']),$$

for $f, f' \in \mathcal{O}_X$, $D, D' \in \mathcal{L}$, together with the map

$$\mathfrak{a}_{\omega} : \mathcal{O}_X \oplus \mathcal{L} \rightarrow \mathcal{T}_X, (f, D) \mapsto \mathfrak{a}(D).$$

Thus, $\mathcal{L}_{\omega} := (\mathcal{O}_X \oplus \mathcal{L}, [\cdot, \cdot]_{\omega}, \mathfrak{a}_{\omega})$ forms a Lie algebroid over (X, \mathcal{O}_X) , provides the abelian Lie algebroid extension (an equivalence class)

$$\mathcal{O}_X \hookrightarrow \mathcal{L}_{\omega} \twoheadrightarrow \mathcal{L}.$$

Thus, $(\mathcal{L}_{\omega}, \text{pr}_2)$ is a \mathcal{L} -Lie algebroid for any 2-cocycle ω of the complex $\Omega_{\mathcal{L}}^{\bullet}$ (see (1)), where $\text{pr}_2 : \mathcal{O}_X \oplus \mathcal{L} \rightarrow \mathcal{L}$ denotes the projection onto the second component.

Definition 3.7. Let $\mathcal{A} := (\bigoplus_{i=0}^{\infty} \mathcal{A}^i, \wedge, [\cdot, \cdot])$ be a Gerstenhaber algebra over (X, \mathcal{O}_X) . A Gerstenhaber algebra $\mathcal{A}' := (\bigoplus_{i=0}^{\infty} \mathcal{A}'^i, \wedge', [\cdot, \cdot]')$ over (X, \mathcal{O}_X) is said to be a \mathcal{A} -Gerstenhaber algebra if there exist a Gerstenhaber algebra homomorphism $\psi : \mathcal{A}' \rightarrow \mathcal{A}$.

Let \mathcal{E} be an \mathcal{O}_X -module. Consider the exterior algebra $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{E} := (\bigoplus_{i=0}^{\infty} \wedge_{\mathcal{O}_X}^i \mathcal{E}, \wedge)$, a graded \mathcal{O}_X -algebra.

Example 3.8. (1) Let (\mathcal{L}', ϕ) be a \mathcal{L} -Lie algebroid. Then the induced map

$$\wedge_{\mathcal{O}_X}^{\bullet} \phi : \mathcal{G}_{\mathcal{L}'} := (\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}', [\cdot, \cdot]_{\mathcal{L}'}) \rightarrow \mathcal{G}_{\mathcal{L}} := (\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}, [\cdot, \cdot]_{\mathcal{L}})$$

provides a $\mathcal{G}_{\mathcal{L}}$ -Gerstenhaber algebra structure on $\mathcal{G}_{\mathcal{L}'}$ (see Example 2.17).

(2) A Gerstenhaber algebra over (X, \mathcal{O}_X) can be viewed as a $\mathcal{G}_{\mathcal{T}_X}$ -Gerstenhaber algebra.

In the following, we extend Theorem 2.18 to the framework of \mathcal{L} -Lie algebroids.

Theorem 3.9. For an \mathcal{O}_X -module \mathcal{E} and a Lie algebroid \mathcal{L} , the \mathcal{O}_X -algebra $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{E}$ has a $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}$ -Gerstenhaber algebra structure if and only if the \mathcal{O}_X -module \mathcal{E} has a \mathcal{L} -Lie algebroid structure.

Proof. Let the \mathcal{O}_X -algebra $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{E}$ has a $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}$ -Gerstenhaber algebra structure. Thus, $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{E}$ is equipped with a Gerstenhaber bracket

$$[\cdot, \cdot]^{\mathcal{E}} : \wedge_{\mathcal{O}_X}^i \mathcal{E} \otimes \wedge_{\mathcal{O}_X}^j \mathcal{E} \rightarrow \wedge_{\mathcal{O}_X}^{i+j-1} \mathcal{E},$$

for all $i, j \geq 0$ (note that $\wedge_{\mathcal{O}_X}^{-1} \mathcal{E} := \{0\}$ and $\wedge_{\mathcal{O}_X}^0 \mathcal{E} := \mathcal{O}_X$) and there is a Gerstenhaber algebra homomorphism

$$\psi : (\bigoplus_{i=0}^{\infty} \wedge_{\mathcal{O}_X}^i \mathcal{E}, \wedge, [\cdot, \cdot]^{\mathcal{E}}) \rightarrow (\bigoplus_{i=0}^{\infty} \wedge_{\mathcal{O}_X}^i \mathcal{L}, \wedge, [\cdot, \cdot]_{\mathcal{L}}).$$

Our claim is the triple $(\mathcal{E}, [\cdot, \cdot]_{\mathcal{E}} := [\cdot, \cdot]^{\mathcal{E}}|_{\mathcal{E}}, \mathfrak{a}' := \mathfrak{a} \circ \psi|_{\mathcal{E}})$ forms a \mathcal{L} -Lie algebroid. Since the homomorphism ψ is a degree zero map (graded map) and thus the restriction map

$$\phi := \psi|_{\mathcal{E}} : (\mathcal{E}, [\cdot, \cdot]_{\mathcal{E}}) \rightarrow (\mathcal{L}, [\cdot, \cdot]_{\mathcal{L}})$$

is an \mathcal{O}_X -module homomorphism and \mathbb{K}_X -Lie algebra homomorphism. For any two section $D, D' \in \mathcal{E}$ and a section $f \in \mathcal{O}_X$ we get the Leibniz identity

$$[D, f D']_{\mathcal{E}} = [D, f \wedge D']^{\mathcal{E}} = [D, f]^{\mathcal{E}} \wedge D' + f \wedge [D, D']^{\mathcal{E}} = \mathfrak{a}'(D)(f) D' + f [D, D']_{\mathcal{E}}$$

(since $\mathfrak{a}'(D)(f) = \mathfrak{a}(\phi(D))(f) = [\phi(D), f]_{\mathcal{L}} = [\psi(D), \psi(f)]_{\mathcal{L}} = \psi([D, f]^{\mathcal{E}}) = [D, f]_{\mathcal{E}}$ holds).

Let $(\mathcal{E}, \mathfrak{a}', [\cdot, \cdot]_{\mathcal{E}})$ be a \mathcal{L} -Lie algebroid. Thus, we have a Lie algebroid homomorphism

$$\phi : (\mathcal{E}, \mathfrak{a}', [\cdot, \cdot]_{\mathcal{E}}) \rightarrow (\mathcal{L}, \mathfrak{a}, [\cdot, \cdot]_{\mathcal{L}}).$$

We show that the homomorphism ϕ lifts or extends (uniquely) to a Gerstenhaber algebra homomorphism

$$\tilde{\phi} : (\bigoplus_{i=0}^{\infty} \wedge_{\mathcal{O}_X}^i \mathcal{E}, \wedge, [\cdot, \cdot]^{\mathcal{E}}) \rightarrow (\bigoplus_{i=0}^{\infty} \wedge_{\mathcal{O}_X}^i \mathcal{L}, \wedge, [\cdot, \cdot]_{\mathcal{L}})$$

between the canonical Gerstenhaber algebras associated with the Lie algebroids \mathcal{E} and \mathcal{L} respectively (see Example 2.17). Consider two homogeneous section $\tilde{D}, \tilde{D}' \in \wedge_{\mathcal{O}_X}^{\bullet} \mathcal{E}$, i.e. there is $f, g \in \mathcal{O}_X$ and $D_1, \dots, D_k, D'_1, \dots, D'_l \in \mathcal{E}$ such that $\tilde{D} = f D_1 \wedge \dots \wedge D_k$, $\tilde{D}' = g D'_1 \wedge \dots \wedge D'_l$. Therefore, the map $\tilde{\phi}$ is defined as follows.

$$\tilde{\phi}(\tilde{D} \wedge \tilde{D}') := f \phi(D_1) \wedge \dots \wedge \phi(D_k) \wedge g \phi(D'_1) \wedge \dots \wedge \phi(D'_l),$$

$$\tilde{\phi}([\tilde{D}, \tilde{D}']^{\mathcal{E}}) := [f \phi(D_1) \wedge \dots \wedge \phi(D_k), g \phi(D'_1) \wedge \dots \wedge \phi(D'_l)]_{\mathcal{L}}.$$

Hence, the map $\tilde{\phi}$ is a graded \mathcal{O}_X -algebra and graded \mathbb{K}_X -Lie algebra homomorphism. \square

Remark 3.10. The canonical correspondence between the category of Lie-Rinehart algebras and the category of Gerstenhaber algebras is established via standard adjoint functors (see [17, 21]). Its algebro-geometric counterpart is illustrated in Example 2.17 and Remark 2.14. Consequently, Theorem 3.9 can also be interpreted within this categorical framework.

4. $\Omega_{\mathcal{L}}^{\bullet}$ -DIFFERENTIAL GRADED ALGEBRAS

Lie algebroids gives rise to differential graded algebras, and conversely, a specific type of differential graded algebras can define Lie algebroids (see Theorem 2.20). To generalize this canonical correspondence in the context of \mathcal{L} -Lie algebroids, we introduce the concept of $\Omega_{\mathcal{L}}^{\bullet}$ -differential graded algebras. This notion extends

the framework of (sheaves of) differential graded algebras, enabling the study of \mathcal{L} -Lie algebroid structures on locally free \mathcal{O}_X -modules of finite rank, where (X, \mathcal{O}_X) is a special ringed space (see Remark 2.2).

Remark 4.1. For a locally free Lie algebroid \mathcal{L} of finite rank, we have the standard differential graded algebra $\Omega_{\mathcal{L}}^{\bullet} := (\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{L}^*, d_{\mathcal{L}})$ of the Chevalley-Eilenberg-de Rham complex.

Definition 4.2. Let $\mathcal{A}^{\bullet} := (\bigoplus_{i=0}^{\infty} \mathcal{A}^i, \wedge, d_{\mathcal{A}})$ be a differential graded algebra. It is said to be a $\Omega_{\mathcal{L}}^{\bullet}$ -differential graded algebra if for a given Lie algebroid \mathcal{L} , there is a differential graded algebra homomorphism

$$\psi : \Omega_{\mathcal{L}}^{\bullet} \rightarrow \mathcal{A}^{\bullet}$$

such that the differential $d_{\mathcal{A}}$ is given by the composition $\psi_1 \circ \mathfrak{a}^* \circ d_X$ of the maps

$$\mathcal{O}_X \xrightarrow{d_X} \Omega_X^1 \xrightarrow{\mathfrak{a}^*} \Omega_{\mathcal{L}}^1 \xrightarrow{\psi_1} \mathcal{A}^1,$$

where d_X represents the usual differential, \mathfrak{a}^* is the dual of \mathfrak{a} and ψ_1 is the restriction of ψ on $\Omega_{\mathcal{L}}^1 := \mathcal{L}^*$.

If the exterior algebra $\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{E}^*$ of the dual of an \mathcal{O}_X -module \mathcal{E} is equipped with a differential $d_{\mathcal{E}} : \wedge^{\bullet}_{\mathcal{O}_X} \mathcal{E}^* \rightarrow \wedge^{\bullet}_{\mathcal{O}_X} \mathcal{E}^{*+1}$, then the following result holds.

Theorem 4.3. Let \mathcal{L} be a locally free Lie algebroid of finite rank. A locally free \mathcal{O}_X -module \mathcal{E} of finite rank has a \mathcal{L} -Lie algebroid structure if and only if $\Omega_{\mathcal{E}}^{\bullet} := (\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{E}^*, d_{\mathcal{E}})$ has a $\Omega_{\mathcal{L}}^{\bullet}$ -differential graded algebra structure.

Proof. Let the \mathcal{O}_X -module \mathcal{E} has a \mathcal{L} -Lie algebroid structure given by a Lie algebroid homomorphism

$$\phi : \mathcal{E} \rightarrow \mathcal{L}.$$

We consider the Chevalley-Eilenberg-de Rham complex $\Omega_{\mathcal{E}}^{\bullet} := (\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{E}^*, d_{\mathcal{E}})$ associated to \mathcal{E} . Note that the differential $d_{\mathcal{E}}$ is given by the composition of the maps

$$\mathcal{O}_X \xrightarrow{d_X} \Omega_X^1 \xrightarrow{\mathfrak{a}^*} \Omega_{\mathcal{L}}^1 := \mathcal{L}^* \xrightarrow{\phi^*} \mathcal{E}^* =: \Omega_{\mathcal{E}}^1,$$

i.e. $d_{\mathcal{E}}$ is the canonical extension of the map $\phi^* \circ \mathfrak{a}^* \circ d_X$ where

$$d_{\mathcal{E}}f(D) = d_X f(\mathfrak{a}(\phi(D))) \text{ for all } D \in \mathcal{E} \text{ and } f \in \mathcal{O}_X.$$

We show that the induced map forms a homomorphism of the cochain complexes

$$\tilde{\phi}^* : \Omega_{\mathcal{L}}^{\bullet} \rightarrow \Omega_{\mathcal{E}}^{\bullet}$$

is defined by $\tilde{\phi}^*(\omega_1 \wedge \cdots \wedge \omega_k) = (\omega_1 \circ \phi) \wedge \cdots \wedge (\omega_k \circ \phi)$, for $\omega_1, \dots, \omega_k \in \mathcal{L}^*$, as follows. Consider $\tilde{\phi} := \wedge^k \phi$, thus for a section ω of $\Omega_{\mathcal{L}}^k := \wedge^k \Omega_{\mathcal{L}}^1$ and sections s_1, \dots, s_{k+1} of \mathcal{E} we get the following identities

$$\begin{aligned} & d_{\mathcal{E}}(\tilde{\phi}^*(\omega))(s_1 \wedge \cdots \wedge s_{k+1}) \\ &= d_{\mathcal{E}}(\omega \circ \tilde{\phi})(s_1 \wedge \cdots \wedge s_{k+1}) \\ &= \sum_{i=1}^{k+1} (-1)^{i+1} \mathfrak{a}_{\mathcal{E}}(s_i)((\omega \circ \tilde{\phi})(s_1 \wedge \cdots \wedge \hat{s}_i \wedge \cdots \wedge s_{k+1})) \\ &\quad + \sum_{i < j} (-1)^{i+j} (\omega \circ \tilde{\phi})([s_i, s_j] \wedge s_1 \wedge \cdots \wedge \hat{s}_i \wedge \cdots \wedge \hat{s}_j \wedge \cdots \wedge s_{k+1}) \\ &= \sum_{i=1}^{k+1} (-1)^{i+1} \mathfrak{a}_{\mathcal{L}}(\phi(s_i))((\omega(\phi(s_1) \wedge \cdots \wedge \hat{\phi}(s_i) \wedge \cdots \wedge \phi(s_{k+1}))) \\ &\quad + \sum_{i < j} (-1)^{i+j} \omega(\phi([s_i, s_j]) \wedge \phi(s_1) \wedge \cdots \wedge \hat{\phi}(s_i) \wedge \cdots \wedge \hat{\phi}(s_j) \wedge \cdots \wedge \phi(s_{k+1})) \\ &= d_{\mathcal{L}}(\omega)(\phi(s_1) \wedge \cdots \wedge \phi(s_{k+1})) \\ &= \tilde{\phi}^*(d_{\mathcal{L}}(\omega))(s_1 \wedge \cdots \wedge s_{k+1}), \end{aligned}$$

(this is an extension of the identity $\phi^*(d_{\mathcal{L}}(f))(s) = d_{\mathcal{L}}(f)(\phi(s))$ where $f \in \mathcal{O}_X$ and $s \in \mathcal{E}$), i.e. the required compatibility condition $d_{\mathcal{E}} \circ \tilde{\phi}^* = \tilde{\phi}^* \circ d_{\mathcal{L}}$ holds. By the construction the map $\tilde{\phi}^*$ is also a graded algebra map. Thus, it provides a $\Omega_{\mathcal{L}}^{\bullet}$ -differential graded algebra structure on $\Omega_{\mathcal{E}}^{\bullet}$.

Suppose, $\Omega_{\mathcal{E}}^{\bullet} := (\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{E}^*, d_{\mathcal{E}})$ is a $\Omega_{\mathcal{L}}^{\bullet}$ -differential graded algebra with connecting morphism $\psi : \Omega_{\mathcal{L}}^{\bullet} \rightarrow \Omega_{\mathcal{E}}^{\bullet}$. Consider the restriction map $\psi|_{\mathcal{L}^*} : \mathcal{L}^* \rightarrow \mathcal{E}^*$ and dualizing it we get a \mathcal{O}_X -module homomorphism $\phi : \mathcal{E} \rightarrow \mathcal{L}$, since both \mathcal{E} and \mathcal{L} are locally free \mathcal{O}_X -modules of finite rank (see Remark 2.7).

Define the \mathcal{O}_X -module homomorphism $\mathfrak{a}' : \mathcal{E} \rightarrow \mathcal{T}_X$ as

$$\mathfrak{a}'(D)(f) := d_{\mathcal{E}}f(D)$$

(well definedness of the map is given by the derivation property of the differential $d_{\mathcal{E}}$). Since $d_{\mathcal{E}} = \psi|_{\mathcal{L}^*} \circ \mathfrak{a}^* \circ d_X = (\mathfrak{a} \circ \phi)^* \circ d_X$ and thus $\mathfrak{a}'(D)(f) = (\mathfrak{a} \circ \phi)^*(d_X f)(D) = d_X f((\mathfrak{a} \circ \phi)(D)) = (\mathfrak{a} \circ \phi)(D)(f)$ for all $f \in \mathcal{O}_X$, $D \in \mathcal{E}$. Hence, the map \mathfrak{a}' should split as $\mathfrak{a} \circ \phi$.

Consider any two sections $D_1, D_2 \in \mathcal{E}$ and a section $\omega \in \mathcal{E}^*$, the Lie bracket $[\cdot, \cdot]_{\mathcal{E}}$ on \mathcal{E} is given by the following identity

$$\omega([D_1, D_2]_{\mathcal{E}}) := \mathfrak{a}'(D_1)(\omega(D_2)) - \mathfrak{a}'(D_2)(\omega(D_1)) - d_{\mathcal{E}}\omega(D_1 \wedge D_2).$$

The map \mathfrak{a}' is a \mathbb{K}_X -Lie algebra homomorphism as follows

$$\begin{aligned} \mathfrak{a}'([D_1, D_2]_{\mathcal{E}})(f) &= d_{\mathcal{E}}f([D_1, D_2]_{\mathcal{E}}) = \mathfrak{a}'(D_1)(d_{\mathcal{E}}f(D_2)) - \mathfrak{a}'(D_2)(d_{\mathcal{E}}f(D_1)) \\ &= \mathfrak{a}'(D_1)(\mathfrak{a}'(D_2)(f)) - \mathfrak{a}'(D_2)(\mathfrak{a}'(D_1)(f)) = [\mathfrak{a}'(D_1), \mathfrak{a}'(D_2)]_c(f), \end{aligned}$$

for all $D_1, D_2 \in \mathcal{E}$ and for all $f \in \mathcal{O}_X$.

To show the Leibniz identity for the map \mathfrak{a}' , we prove that for $f \in \mathcal{O}_X$ and $D_1, D_2 \in \mathcal{E}$ the following identity holds for all $\omega \in \mathcal{E}^*$

$$\omega([D_1, f D_2]_{\mathcal{E}}) = \omega(f [D_1, D_2]_{\mathcal{E}} + \mathfrak{a}'(D_1)(f) D_2).$$

Notice that we get the following identities

$$\begin{aligned} \omega([D_1, f D_2]_{\mathcal{E}}) &= \mathfrak{a}'(D_1)(f \omega(D_2)) - f \mathfrak{a}'(D_2)(\omega(D_1)) - f d_{\mathcal{E}}\omega(D_1 \wedge D_2) \\ &= \mathfrak{a}'(D_1)(f) \omega(D_2) + f \mathfrak{a}'(D_1)(\omega(D_2)) - f \mathfrak{a}'(D_2)(\omega(D_1)) - f d_{\mathcal{E}}\omega(D_1 \wedge D_2) \\ &= f (\mathfrak{a}'(D_1)(\omega(D_2)) - \mathfrak{a}'(D_2)(\omega(D_1)) - d_{\mathcal{E}}\omega(D_1 \wedge D_2)) + \mathfrak{a}'(D_1)(f) \omega(D_2) \\ &= f \omega([D_1, D_2]_{\mathcal{E}}) + \mathfrak{a}'(D_1)(f) \omega(D_2) = \omega(f [D_1, D_2]_{\mathcal{E}} + \mathfrak{a}'(D_1)(f) D_2). \end{aligned}$$

□

Remark 4.4. *The condition of locally free \mathcal{O}_X -modules of finite rank in the above result can be relaxed by considering both \mathcal{O}_X -modules \mathcal{E} and \mathcal{L} as reflexive sheaves. Under this assumption, in the second part of the proof, the restriction of the dga morphism $\psi : (\mathcal{H}om_{\mathcal{O}_X}(\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}, \mathcal{O}_X), d_{\mathcal{L}}) \rightarrow (\mathcal{H}om_{\mathcal{O}_X}(\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{E}, \mathcal{O}_X), d_{\mathcal{E}})$ to the grading 1 induces the same \mathcal{O}_X -module homomorphism $\phi : \mathcal{E} \rightarrow \mathcal{L}$, leveraging the reflexivity property. For the first part of the proof, we don't even need the reflexive condition.*

5. \mathcal{L} -LIE BIALGEBROIDS

There is a one to one correspondence between Lie bialgebroids and certain strong differential Gerstenhaber algebras (see Theorem 2.28). We extend this to a correspondence between two new notions, namely, \mathcal{L} -Lie bialgebroids and strong differential $\mathcal{G}_{\mathcal{L}}$ -Gerstenhaber algebras. In the following considerations, we take Lie algebroids that are locally free \mathcal{O}_X -modules of finite rank.

Definition 5.1. *A \mathcal{L} -Lie bialgebroid is a pair $(\mathcal{L}', \mathcal{L}'^*)$ of \mathcal{L} -Lie algebroids over (X, \mathcal{O}_X) that are in duality (with $\mathcal{L}'^* = (\mathcal{L}')^*$). The Lie bracket on \mathcal{L}' satisfies a compatibility condition expressed using the differential $d_{\mathcal{L}'^*} := d'_*$ on $\wedge_{\mathcal{O}_X}^{\bullet} \mathcal{L}'$ (since $(\mathcal{L}'^*)^* \cong \mathcal{L}'$), defined by the Lie algebroid structure of \mathcal{L}'^* . Specifically, d'_* acts as a derivation of the Lie bracket on \mathcal{L}' , meaning that for any sections D_1, D_2 of \mathcal{L}' , the following holds:*

$$d'_*[D_1, D_2] = [d'_*D_1, D_2] + [D_1, d'_*D_2].$$

Remark 5.2. A \mathcal{T}_X -Lie bialgebroid is same as a Lie bialgebroid over a non-singular space X .

Example 5.3. Let (X, π) be a Poisson manifold and $(\mathcal{O}_X, \{\cdot, \cdot\})$ be the associated Poisson algebra. Thus, there is a canonical Lie algebroid homomorphism $\tilde{\pi} : \Omega_X^1 \rightarrow \mathcal{T}_X$ induced from the Poisson structure on X . Consider a closed Poisson analytic subspace $Y := V(\mathcal{I})$ of X , i.e. the zero locus of an ideal sheaf \mathcal{I} of \mathcal{O}_X satisfying the condition $\{\mathcal{I}, \mathcal{O}_X\} \subset \mathcal{I}$. Thus, Y is an Ω_X^1 -invariant subspace, i.e. $\mathcal{I}m(\mathbf{a})$ preserves \mathcal{I} . In particular, when Y is a Poisson hypersurface (Y is of codimension 1), the anchor map \mathbf{a} induces a Lie algebroid homomorphism $\tilde{\pi} : \Omega_X^1(\log Y) \rightarrow \mathcal{T}_X(-\log Y)$, where $\Omega_X^1(\log Y)$ is the sheaf of meromorphic 1-forms with simple poles along Y . Therefore $(\Omega_X^1(\log Y), \mathcal{T}_X(-\log Y))$ forms a $\mathcal{T}_X(-\log Y)$ -Lie algebroid if in addition Y is free divisor. Moreover, the dual pair is forms a triangular Lie bialgeroid (see [34]).

Proposition 5.4. \mathcal{L} -Lie bialgebroid $(\mathcal{L}', \mathcal{L}'^*)$ is equivalent to \mathcal{L} -Lie bialgebroid $(\mathcal{L}'^*, \mathcal{L}')$.

Remark 5.5. Let $(\mathcal{L}, \mathcal{L}^*)$ be a Lie bialgebroid. Then the anchor map $\mathbf{a}' : \mathcal{L}^* \rightarrow \mathcal{T}_X$ induces the \mathcal{O}_X -module homomorphism $(\mathbf{a}')^* : \Omega_X^1 \rightarrow \mathcal{L}$, provides a Ω_X^\bullet -differential graded algebra structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}$ as an extension of the map $(\mathbf{a}')^* \circ d_X$.

Remark 5.6. Let $(\mathcal{L}, \mathcal{L}^*)$ be a Lie bialgebroid. Then there is a canonical strong differential Gerstenhaber algebra structure on $\mathcal{G}_{\mathcal{L}} := (\bigoplus_{i=0}^\infty \wedge_{\mathcal{O}_X}^i \mathcal{L}, \wedge, [\cdot, \cdot]^\mathcal{L})$ given through the differential $d_* : \wedge_{\mathcal{O}_X}^\bullet \mathcal{L} \rightarrow \wedge_{\mathcal{O}_X}^{\bullet+1} \mathcal{L}$, defined by the Lie algebroid structure of \mathcal{L}^* .

Definition 5.7. A strong differential $\mathcal{G}_{\mathcal{L}}$ -Gerstenhaber algebra is a $\mathcal{G}_{\mathcal{L}} := (\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}, [\cdot, \cdot]^\mathcal{L})$ -Gerstenhaber algebra $\mathcal{A} := (\bigoplus_{i=0}^\infty \mathcal{A}^i, \wedge, [\cdot, \cdot])$ equipped with a differential $\tilde{d} : \bigoplus_{i=0}^\infty \mathcal{A}^i \rightarrow \bigoplus_{i=0}^\infty \mathcal{A}^{i+1}$, which is a derivation of degree 1 with respect to \wedge satisfying $\tilde{d}^2 = 0$ and also it is a derivation of the graded Lie bracket $[\cdot, \cdot]$. In addition we need to have a differential graded algebra homomorphism $\psi : \Omega_{\mathcal{L}}^\bullet := (\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}^*, d_{\mathcal{L}}) \rightarrow (\mathcal{A}, \tilde{d})$.

Remark 5.8. A strong differential $\mathcal{G}_{\mathcal{T}_X}$ -Gerstenhaber algebra is same as sheaf of strong differential Gerstenhaber algebra.

Remark 5.9. Consider the standard case associated to $\mathcal{L} = \mathcal{T}_X$, where X is non singular, i.e. \mathcal{T}_X is a locally free \mathcal{O}_X -module of finite rank. For an locally free \mathcal{O}_X -module \mathcal{E} of finite rank, $\wedge_{\mathcal{O}_X}^\bullet \mathcal{E}$ has a strong differential Gerstenhaber algebra structure if and only if $(\mathcal{E}, \mathcal{E}^*)$ has a Lie bialgebroid structure.

Theorem 5.10. Let \mathcal{L}' be a locally free \mathcal{O}_X -module of finite rank. The pair $(\mathcal{L}', \mathcal{L}'^*)$ forms a \mathcal{L} -Lie bialgebroid if and only if there is a strong differential $\mathcal{G}_{\mathcal{L}}$ -Gerstenhaber algebra structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}'$.

Proof. Let $(\mathcal{L}', \mathcal{L}'^*)$ be a \mathcal{L} -Lie bialgebroids. The \mathcal{L} -Lie algebroid structure on \mathcal{L}' is given by a Lie algebroid homomorphism $\phi : \mathcal{L}' \rightarrow \mathcal{L}$, provides the canonical Gerstenhaber algebra homomorphism

$$\tilde{\phi} : \wedge_{\mathcal{O}_X}^\bullet \mathcal{L}' \rightarrow \wedge_{\mathcal{O}_X}^\bullet \mathcal{L},$$

and thus $(\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}', [\cdot, \cdot]^{\mathcal{L}'})$ forms a $\mathcal{G}_{\mathcal{L}}$ -Gerstenhaber algebra. The Chevalley-Eilenberg-de Rham complex $\Omega_{\mathcal{L}'^*}^\bullet$ of the Lie algebroid \mathcal{L}'^* provides the differential graded algebra $(\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}', d'_*)$ where the differential

$$d'_* : \wedge_{\mathcal{O}_X}^\bullet \mathcal{L}' \rightarrow \wedge_{\mathcal{O}_X}^{\bullet+1} \mathcal{L}'$$

is a derivation with respect to the standard Gerstenhaber bracket $[\cdot, \cdot]^{\mathcal{L}'}$. Now, consider the \mathcal{L} -Lie algebroid structure on \mathcal{L}'^* , is given by a Lie algebroid homomorphism $\psi : \mathcal{L}'^* \rightarrow \mathcal{L}$, provides the \mathcal{O}_X -module homomorphism $\psi^* : \mathcal{L}^* \rightarrow \mathcal{L}'$. Thus, we have the following sequence of \mathcal{O}_X -modules with \mathbb{K}_X -linear maps

$$\mathcal{O}_X \xrightarrow{d_X} \Omega_X^1 := \mathcal{T}_X^* \xrightarrow{\mathbf{a}^*} \Omega_{\mathcal{L}}^1 := \mathcal{L}^* \xrightarrow{\psi^*} \mathcal{L}'.$$

It provides the differential graded algebra homomorphism

$$\wedge^\bullet \psi^* : \Omega_{\mathcal{L}}^\bullet \rightarrow (\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}', d'_*).$$

Therefore, $(\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}', [\cdot, \cdot]^{\mathcal{L}'}, d'_*)$ forms a strong differential $\mathcal{G}_{\mathcal{L}}$ -Gerstenhaber algebras.

Let there is a strong differential $\mathcal{G}_{\mathcal{L}}$ -Gerstenhaber algebra structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}'$, i.e. there is a Gerstenhaber bracket $[\cdot, \cdot]$ and differential \tilde{d} on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}'$ satisfies the following properties. There exists a Gerstenhaber algebra

homomorphism

$$\bar{\phi} : (\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{L}', [\cdot, \cdot]) \rightarrow (\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{L}, [\cdot, \cdot]^{\mathcal{L}})$$

and a differential graded algebra homomorphism

$$\psi : (\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{L}^*, d_{\mathcal{L}}) \rightarrow (\wedge^{\bullet}_{\mathcal{O}_X} \mathcal{L}', \tilde{d}).$$

The homomorphism $\bar{\phi}$ provides a \mathcal{L} -Lie algebroid structure on \mathcal{L}' given by the restriction map

$$\phi := \bar{\phi}|_{\mathcal{L}'} : (\mathcal{L}', [\cdot, \cdot]_{\mathcal{L}'}) \rightarrow (\mathcal{L}, [\cdot, \cdot]_{\mathcal{L}})$$

and the homomorphism ψ provides a \mathcal{L} -Lie algebroid structure on \mathcal{L}'^* , given through dualizing the map $\psi_1 := \psi|_{\mathcal{L}'^*} : \mathcal{L}'^* \rightarrow \mathcal{L}'$ i.e. given by the map

$$\psi_1^* : \mathcal{L}'^* \rightarrow \mathcal{L}$$

The compatibility condition for the pair $(\mathcal{L}', \mathcal{L}'^*)$ of \mathcal{L} -Lie algebroids to become a \mathcal{L} -Lie bialgebroids is given by the derivation property of the differential map \tilde{d} with respect to the graded Lie bracket $[\cdot, \cdot]$. \square

Remark 5.11. *It is important to note that the dual pair of Lie algebroids considered here is not required to consist of vector bundles (or locally free sheaves of finite rank), as in [30,31,44]. However, to properly interpret the duality condition in the definition of a Lie bialgebroid, it is essential for the sheaves involved to be reflexive (see [34]). As a result, Theorem 5.10 can be extended to this more general setting.*

6. \mathcal{L} -LIE ALGEBROID HOMOLOGY

We extend the concepts of Lie algebroid connections and curvature and develop an analogue of Theorem 2.33 in a broader context. Subsequently, we generalize the notion of Lie algebroid homology [44] within this framework. These concepts serve as foundational tools for Section 8.

6.1. Generalized \mathcal{L} -connections. Let (\mathcal{L}', ϕ) be a \mathcal{L} -Lie algebroid. We consider such \mathcal{L}' -connection ∇' on an \mathcal{O}_X -module \mathcal{E} which splits by a \mathcal{L} -connection ∇ on \mathcal{E} as follows.

Definition 6.1. *A \mathcal{L}' -connection (\mathcal{E}, ∇') is said to be a $(\mathcal{L}', \mathcal{L})$ -connection if there exists a \mathcal{L} -connection (\mathcal{E}, ∇) such that $\nabla' = \nabla \circ \phi$.*

More generally, the above definition applies when considering an $\mathcal{I}m(\phi)$ -connection (\mathcal{E}, ∇) instead of an \mathcal{L} -connection, as $\mathcal{I}m(\phi)$ forms a Lie subalgebroid of \mathcal{L} .

Remark 6.2. (1) *A \mathcal{L} -connection (\mathcal{E}, ∇) provides the \mathcal{L}' -connection $(\mathcal{E}, \nabla \circ \phi)$.*

(2) *For a Lie algebroid \mathcal{L} over (X, \mathcal{O}_X) , a $(\mathcal{L}, \mathcal{L})$ -connection on an \mathcal{O}_X -module \mathcal{E} is same with a usual \mathcal{L} -connection on \mathcal{E} .*

Definition 6.3. *A $(\mathcal{L}', \mathcal{L})$ -connection (\mathcal{E}, ∇') is said to be a flat connection if the curvature of the connection ∇' is zero, i.e. for any $D_1, D_2 \in \mathcal{L}'$*

$$[\nabla'_{D_1}, \nabla'_{D_2}]_c = \nabla'_{[D_1, D_2]}.$$

Remark 6.4. (1) *For a flat \mathcal{L} -connection (\mathcal{E}, ∇) , the \mathcal{L}' -connection $(\mathcal{E}, \nabla \circ \phi)$ is also flat.*

(2) *Therefore, for a flat $(\mathcal{L}', \mathcal{L})$ -connection (\mathcal{E}, ∇') we get a Lie algebroid homomorphism*

$$\nabla' : \mathcal{L}' \rightarrow \mathcal{A}t(\mathcal{E}),$$

where $\mathcal{A}t(\mathcal{E})$ is the Atiyah algebroid of the \mathcal{O}_X -module \mathcal{E} . Thus, the \mathcal{O}_X -module homomorphism $\nabla : \mathcal{L} \rightarrow \text{End}_{\mathbb{K}_X}(\mathcal{E})$ satisfying the Leibniz rule, restricted on $\mathcal{I}m(\phi)$ is flat.

Example 6.5. *Let (Y, \mathcal{O}_Y) be a closed Poisson analytic hypersurface (or Poisson principal divisor) of a Poisson manifold (X, \mathcal{O}_X) . Then the canonical Lie algebroid homomorphism $\bar{\pi} : \Omega_X^1 \rightarrow \mathcal{T}_X$ induced by the Poisson bivector field π , extends to a Lie algebroid homomorphism (see Example 2.10 and Example 5.3)*

$$\bar{\pi} : \Omega_X^1(\log Y) \rightarrow \mathcal{T}_X(-\log Y).$$

Consider $\mathcal{L}' := \Omega_X^1(\log Y)$, $\mathcal{L} := \mathcal{T}_X(-\log Y)$. For any flat connection ∇ on an \mathcal{O}_X -module \mathcal{E} can be restricted to give a logarithmic connection (see Example 3.4)

$$\nabla' := \nabla|_{\mathcal{T}_X(-\log Y)} : \mathcal{T}_X(-\log Y) \rightarrow \mathcal{A}t(\mathcal{E})(-\log Y).$$

Thus, we get a flat $(\mathcal{L}', \mathcal{L})$ -connection $(\mathcal{E}, \nabla' \circ \bar{\pi})$.

Example 6.6. In the study of equivariant geometry (see [4]), for a holomorphic principal H -bundle E_H on a complex manifold X , where X is equipped with a holomorphic action of a connected complex Lie group G , a subbundle $At^G(E_H)$ of the holomorphic vector bundle $At(E_H) \oplus (X \times Lie(G))$ is constructed. The pair $(At^G(E_H), \beta \circ \alpha)$ forms a holomorphic Lie algebroid over X , where $\alpha : At^G(E_H) \rightarrow X \times Lie(G)$ is given by the composition map $At^G(E_H) \hookrightarrow At(E_H) \oplus (X \times Lie(G)) \xrightarrow{pr_2} X \times Lie(G)$ and $\beta : X \times Lie(G) \rightarrow TX$ is induced by the action of G on X . A G -connection on E_H is defined to be a holomorphic splitting of an appropriate short exact sequence, given by a section $\gamma : X \times Lie(G) \rightarrow At^G(E_H)$ (i.e. $\alpha \circ \gamma = id$). Consider the Lie algebroid formed by the sheaf of sections $\mathcal{A}t^G(E_H) := \Gamma_X(At^G(E_H))$ and $\mathcal{A}t(E_H) := \Gamma_X(At(E_H))$ over (X, \mathcal{O}_X) , with the morphism $\mathcal{A}t^G(E_H) \xrightarrow{pr_1} \mathcal{A}t(E_H)$. Consider $\mathcal{L} := \mathcal{O}_X \otimes_{\mathbb{C}_X} Lie(G)_X$, the sheaf of sections of the transformation Lie algebroid $X \times Lie(G)$ over X (see Example 3.3) together with the connection $\nabla := pr_1 \circ \gamma$ on $\mathcal{A}t(E_H)$. When E_H is G -equivariant, then it admits a flat G -connection ∇ . Now, consider another connected complex Lie group G' together with a Lie group homomorphism $\bar{\phi} : G' \rightarrow G$. We have an induced action of G' on E_H and a Lie algebroid homomorphism $\phi : \mathcal{L}' \rightarrow \mathcal{L}$ where $\mathcal{L}' := \mathcal{O}_X \otimes_{\mathbb{C}_X} Lie(G')_X$. Then, the above provides a flat $(\mathcal{L}', \mathcal{L})$ -connection on the \mathcal{O}_X -module corresponding to any vector bundle associated to E_H . Analogous notions can be constructed in the context of holomorphic (or smooth or algebraic) vector bundles, and more generally for sheaf theoretic settings (see Section 3.2, [23]).

We extend the Theorem 2.33 to our settings as follows.

Theorem 6.7. Let \mathcal{L} be a locally free \mathcal{O}_X -module of rank n (for some $n \in \mathbb{N}$) and $(\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}', [\cdot, \cdot]')$ a Gerstenhaber algebra for some \mathcal{O}_X -module \mathcal{L}' . Then there is an exact Gerstenhaber algebra (or BV-algebra) structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}$, say $(\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}, [\cdot, \cdot], \partial)$, together with a Gerstenhaber algebra homomorphism

$$\psi : (\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}', [\cdot, \cdot]') \rightarrow (\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}, [\cdot, \cdot])$$

if and only if there is a \mathcal{L} -Lie algebroid structure on \mathcal{L}' with a flat $(\mathcal{L}', \mathcal{L})$ -connection on the \mathcal{O}_X -module $\wedge_{\mathcal{O}_X}^n \mathcal{L}$.

Proof. Suppose $(\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}, [\cdot, \cdot], \partial)$ is an exact Gerstenhaber algebra together with a Gerstenhaber algebra homomorphism

$$\psi : (\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}', [\cdot, \cdot]') \rightarrow (\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}, [\cdot, \cdot]).$$

The Gerstenhaber algebra structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}$ provides a canonical Lie algebroid structure on \mathcal{L} . Moreover, the generating operator $\partial : \wedge_{\mathcal{O}_X}^\bullet \mathcal{L} \rightarrow \wedge_{\mathcal{O}_X}^{\bullet-1} \mathcal{L}$ provides a flat \mathcal{L} -connection ∇ on the locally free \mathcal{O}_X -module $\wedge_{\mathcal{O}_X}^n \mathcal{L}$ of rank 1, denote it as $(\wedge_{\mathcal{O}_X}^n \mathcal{L}, \nabla)$. This results are mentioned in Theorem 2.18 and Theorem 2.33. By Theorem 3.9, the Gerstenhaber algebra homomorphism ψ provides an \mathcal{L} -Lie algebroid structure on \mathcal{L}' through the connecting map $\psi|_{\mathcal{L}'}$. Therefore, we get the flat $(\mathcal{L}', \mathcal{L})$ -connection $(\wedge_{\mathcal{O}_X}^n \mathcal{L}, \nabla \circ \psi|_{\mathcal{L}'})$, i.e. we get a Lie algebroid homomorphism by the composition of the following maps

$$\mathcal{L}' \xrightarrow{\psi|_{\mathcal{L}'}} \mathcal{L} \xrightarrow{\nabla} \mathcal{A}t(\wedge_{\mathcal{O}_X}^n \mathcal{L}).$$

Let the locally free \mathcal{O}_X -module \mathcal{L} has a Lie algebroid structure and (\mathcal{L}', ϕ) be a \mathcal{L} -Lie algebroid. Suppose there is a flat $(\mathcal{L}', \mathcal{L})$ -connection ∇' on the \mathcal{O}_X -module $\wedge_{\mathcal{O}_X}^n \mathcal{L}$. Using Theorem 2.33 we get an exact gerstenhaber algebra structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}$ and by Theorem 2.18 we get a Gerstenhaber algebra structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}'$. Therefore, by Theorem 3.9, we get the canonical Gerstenhaber algebra homomorphism

$$\wedge^\bullet \phi : (\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}', [\cdot, \cdot]') \rightarrow (\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}, [\cdot, \cdot]).$$

□

Remark 6.8. Using Remark 2.34 and Theorem 6.7 we can conclude that there is a canonical flat $(\mathcal{L}', \mathcal{L})$ -connection on the \mathcal{O}_X -module $\wedge_{\mathcal{O}_X}^n \mathcal{L}$ and a canonical BV-algebra structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}$.

6.2. \mathcal{L} -Lie algebroid homology. Let \mathcal{L}' be a Lie subalgebroid of \mathcal{L} , where \mathcal{L} is a locally free \mathcal{O}_X -module of rank n for some $n \in \mathbb{N}$. Thus, (\mathcal{L}', ι) is a \mathcal{L} -Lie algebroid, where $\iota : \mathcal{L}' \rightarrow \mathcal{L}$ is the inclusion map.

Consider a flat $(\mathcal{L}', \mathcal{L})$ -connection on the \mathcal{O}_X -module $\wedge_{\mathcal{O}_X}^n \mathcal{L}$ of rank 1 (i.e. sheaf of sections of a line bundle over X). Let D be its corresponding generating operator, and

$$\partial = (-1)^{n-\bullet} D : \wedge_{\mathcal{O}_X}^\bullet \mathcal{L} \rightarrow \wedge_{\mathcal{O}_X}^{\bullet-1} \mathcal{L}$$

(later we see the importance for choosing the sign of ∂). Then $\partial^2 = 0$, and we obtain a chain complex of sheaves. Thus, the associated sheaves of homologies (see Remark 2.22)

$$\mathcal{H}_\bullet(\mathcal{L}, \nabla) := \text{Ker } \partial / \text{Im } \partial$$

is the sheafification of the presheaf $U \mapsto H_\bullet(\mathcal{L}(U), \nabla(U))$ of the associated homology groups on each open set U of X , considered as a graded \mathbb{K} -vector space. Applying the hyperhomology functor, we get the graded vector space $\mathbb{H}_\bullet(\mathcal{L}, \nabla)$ of hyperhomologies.

The \mathcal{L} -Lie algebroid (\mathcal{L}', ι) is said to be ∂ -invariant if the restriction map of ∂ on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}'$, say ∂' , is a map as follows

$$\partial' : \wedge_{\mathcal{O}_X}^\bullet \mathcal{L}' \rightarrow \wedge_{\mathcal{O}_X}^{\bullet-1} \mathcal{L}'.$$

Therefore, we get a graded vector space homomorphism between the associated homologies

$$\mathbb{H}_\bullet(\mathcal{L}', \nabla') \rightarrow \mathbb{H}_\bullet(\mathcal{L}, \nabla),$$

where $\nabla' = \nabla \circ \iota$.

7. $\mathcal{U}(\mathcal{O}_X, \mathcal{L})$ -UNIVERSAL ENVELOPING ALGEBROID

The universal enveloping algebra of a Lie-Rinehart algebra serves as a fundamental tool for exploring its homological algebra, analogous to the classical case of Lie algebras (see [19,20,27,36]). Its algebro-geometric counterpart is the universal enveloping algebroid, a concept that plays a central role in understanding the homological algebra of Lie algebroids (see [5,6,9,22,32,34]). A key ingredient in this is the Poincaré-Birkhoff-Witt (PBW) theorem, which has been established for Lie algebroids in a sheaf-theoretic framework (see [32]). We further generalize this theory to encompass the case of \mathcal{L} -Lie algebroids, which is useful in Section 8.

We begin by recalling the definition of the universal enveloping algebroid of a Lie algebroid.

7.1. Universal enveloping algebroid of a Lie algebroid. Let $(\mathcal{L}, [\cdot, \cdot], \mathfrak{a})$ be a Lie algebroid over (X, \mathcal{O}_X) . For each open set U of X , let $\mathcal{U}(\mathcal{O}_X(U), \mathcal{L}(U))$ be the universal enveloping algebra of the $(\mathbb{K}, \mathcal{O}_X(U))$ -Lie-Rinehart algebra $\mathcal{L}(U)$, or equivalently, of the Lie-Rinehart pair $(\mathcal{O}_X(U), \mathcal{L}(U))$ (see [19,33,36]). The sheafification of the canonical presheaf,

$$U \mapsto \mathcal{U}(\mathcal{O}_X(U), \mathcal{L}(U)),$$

is known as the universal enveloping algebroid of the Lie algebroid \mathcal{L} , and denoted by $\mathcal{U}(\mathcal{O}_X, \mathcal{L})$ (see [5,8,32,34,38] for details).

From the construction of $\mathcal{U}(\mathcal{O}_X, \mathcal{L})$, it is an associative \mathbb{K}_X -algebra and \mathcal{O}_X -bimodule. Moreover, there is a canonical \mathbb{K}_X -algebra monomorphism $\iota_X : \mathcal{O}_X \hookrightarrow \mathcal{U}(\mathcal{O}_X, \mathcal{L})$ and an \mathcal{O}_X -linear map $\iota_{\mathcal{L}} : \mathcal{L} \rightarrow \mathcal{U}(\mathcal{O}_X, \mathcal{L})$. The associative \mathbb{K}_X -algebra $\mathcal{U}(\mathcal{O}_X, \mathcal{L})$ is generated by \mathcal{O}_X and $\iota_{\mathcal{L}}(\mathcal{L})$ satisfy the following identities:

$$\bar{D} \bar{D}' - \bar{D}' \bar{D} = \overline{[D, D']}, \quad \bar{D} f - f \bar{D} = \mathfrak{a}(D)(f),$$

where $D, D' \in \mathcal{L}$, $f \in \mathcal{O}_X$, and $\bar{D} = \iota_{\mathcal{L}}(D)$ for all $D \in \mathcal{L}$. Hence, the map $\iota_{\mathcal{L}}$ can also be viewed as a \mathbb{K}_X -Lie algebra homomorphism.

The \mathbb{K}_X -algebra $\mathcal{U}(\mathcal{O}_X, \mathcal{L})$ has a natural filtration of \mathcal{O}_X -modules:

$$\mathcal{O}_X = \mathcal{U}_{(0)}(\mathcal{O}_X, \mathcal{L}) \subset \mathcal{U}_{(1)}(\mathcal{O}_X, \mathcal{L}) \subset \mathcal{U}_{(2)}(\mathcal{O}_X, \mathcal{L}) \subset \cdots,$$

where, $\mathcal{U}_{(n)}(\mathcal{O}_X, \mathcal{L})$ is spanned by the sections of \mathcal{O}_X and the powers $\iota_{\mathcal{L}}(\mathcal{L})^m$ for $m = 1, 2, \dots, n$. The direct sum of the quotient sheaves associated with each pair of consecutive sheaves appearing in the above filtration on $\mathcal{U}(\mathcal{O}_X, \mathcal{L})$ forms a sheaf of graded algebras denoted by $gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L}))$. Thus,

$$gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L})) = \bigoplus_{n \geq 0} \mathcal{U}_{(n)}(\mathcal{O}_X, \mathcal{L}) / \mathcal{U}_{(n-1)}(\mathcal{O}_X, \mathcal{L}).$$

Remark 7.1. *The universal enveloping algebroid $\mathcal{U}(\mathcal{O}_X, \mathcal{L})$ of a Lie algebroid \mathcal{L} over (X, \mathcal{O}_X) is characterized by the following universal property:*

Let \mathcal{A} be a sheaf of unital associative \mathbb{K}_X -algebra with sheaf homomorphisms $\phi : \mathcal{O}_X \rightarrow \mathcal{A}$ of \mathbb{K}_X -unital algebras and $\psi : (\mathcal{L}, [\cdot, \cdot]) \rightarrow (\mathcal{A}, [\cdot, \cdot]_c)$ of \mathcal{O}_X -linear \mathbb{K}_X -Lie algebras such that $\phi(f)\psi(D) = \psi(fD)$ and $[\psi(D), \phi(f)]_c = \phi(\mathfrak{a}(D(f)))$ holds for $f \in \mathcal{O}_X$ and $D \in \mathcal{L}$. Then, there exists a unique \mathcal{O}_X -linear homomorphism of unital \mathbb{K}_X -algebras $\tilde{\psi} : \mathcal{U}(\mathcal{O}_X, \mathcal{L}) \rightarrow \mathcal{A}$ such that $\tilde{\psi} \circ \iota = \phi$ and $\tilde{\psi} \circ \iota_{\mathcal{L}} = \psi$.

Example 7.2. *For a non-singular special space (X, \mathcal{O}_X) , the universal enveloping algebroid is isomorphic to the sheaf of differential operators \mathcal{D}_X on X (i.e. the sheaf of differential operators over \mathcal{O}_X , sometimes denoted as $\mathcal{D}iff(\mathcal{O}_X)$) [32, 38], i.e.*

$$\mathcal{U}(\mathcal{O}_X, \mathcal{T}_X) \cong \mathcal{D}iff(\mathcal{O}_X) =: \mathcal{D}_X.$$

Example 7.3. *Given a principal divisor Y in some complex manifold or smooth algebraic variety X , the sheaf of logarithmic derivations and the sheaf of logarithmic differential operators are denoted as $\mathcal{T}_X(-\log Y)$ and $\mathcal{D}_X(-\log Y)$, respectively [32]. In the case of a free divisor Y in X (i.e. $\mathcal{T}_X(-\log Y)$ is locally free \mathcal{O}_X -module [10, 32, 34]) we have (after sheafifying the local description given for the module of logarithmic derivations in [29]) the isomorphism*

$$\mathcal{U}(\mathcal{O}_X, \mathcal{T}_X(-\log Y)) \cong \mathcal{D}_X(-\log Y).$$

In the following we define an algebro-geometric analogue of the notion of algebra of differential operators on a module (see [28, 29, 38] for a local description).

In general, we consider the sheaf of differential operators on an \mathcal{O}_X -module \mathcal{E} , denote it by $\mathcal{D}iff(\mathcal{E})$ as follows. It forms a sheaf of almost commutative algebras, define inductively as follows

$$\mathcal{D}iff(\mathcal{E}) := \cup_{n \geq 0} \mathcal{D}iff^{\leq n}(\mathcal{E}),$$

where $\mathcal{D}iff^0(\mathcal{E}) := \mathcal{E}nd_{\mathcal{O}_X}(\mathcal{E})$, and for $n \geq 1$ consider the \mathcal{O}_X -modules

$$\mathcal{D}iff^{\leq n}(\mathcal{E}) := \{\tilde{D} \in \mathcal{E}nd_{\mathbb{K}_X}(\mathcal{E}) \mid [\tilde{D}, f]_c \in \mathcal{D}iff^{\leq n-1}(\mathcal{E}), f \in \mathcal{O}_X\}.$$

Note that, for any $\tilde{D} \in \mathcal{D}iff^{\leq n}(\mathcal{E})$, $[\tilde{D}, f]_c := \tilde{D} \circ f - f \circ \tilde{D} : \mathcal{E} \rightarrow \mathcal{E}$ is given by $s \mapsto \tilde{D}(f s) - f \tilde{D}(s)$, for $s \in \mathcal{E}$. Thus, there is the canonical map

$$\begin{aligned} \sigma : \mathcal{D}iff^{\leq 1}(\mathcal{E}) &\rightarrow \mathcal{D}er_{\mathbb{K}_X}(\mathcal{O}_X) \\ D &\mapsto [D, -]_c \end{aligned}$$

Remark 7.4. *It is easy to verify that for an \mathcal{O}_X -module \mathcal{E} , the notions $At(\mathcal{E})$ and $\mathcal{D}iff^{\leq 1}(\mathcal{E})$ are same. In addition, if there is a flat \mathcal{T}_X -connection on \mathcal{E} , the abelian Lie algebroid extension (2) splits.*

Theorem 7.5. (Generalized PBW theorem [32]) *If \mathcal{L} is a locally free Lie algebroid over (X, \mathcal{O}_X) then the graded algebra $gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L}))$ is isomorphic to the symmetric algebra $\mathcal{S}_{\mathcal{O}_X} \mathcal{L}$ as graded \mathcal{O}_X -algebras, given by the PBW map (also known as the symmetrization map)*

$$\begin{aligned} \psi_{\mathcal{L}} : \mathcal{S}_{\mathcal{O}_X} \mathcal{L} &\rightarrow gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L})) \\ f \otimes D_1 \otimes \cdots \otimes D_k &\mapsto \frac{1}{k!} f \sum_{\sigma \in S_k} \bar{D}_{\sigma(1)} \cdots \bar{D}_{\sigma(k)} \end{aligned}$$

where f is a section of \mathcal{O}_X and D_1, \dots, D_k are sections of \mathcal{L} .

Remark 7.6. *For a locally free Lie algebroid \mathcal{L} over (X, \mathcal{O}_X) , the homomorphism (as \mathbb{K}_X -algebras and \mathcal{O}_X -modules) $\iota_{\mathcal{L}} : \mathcal{L} \rightarrow \mathcal{U}(\mathcal{O}_X, \mathcal{L})$ become an embedding. As a consequence (using Remark 7.1), the category of representations of \mathcal{L} and category of left $\mathcal{U}(\mathcal{O}_X, \mathcal{L})$ -modules are equivalent (see [5, 34, 37]).*

For a \mathcal{L} -Lie algebroid (\mathcal{L}', ϕ) , we get the canonical map

$$\bar{\phi} := \mathcal{U}(\phi) : \mathcal{U}(\mathcal{O}_X, \mathcal{L}') \rightarrow \mathcal{U}(\mathcal{O}_X, \mathcal{L})$$

using the functorial property of $\mathcal{U}(\mathcal{O}_X, -)$. For the universal enveloping algebroids, we have a canonical \mathbb{K}_X -algebra and \mathcal{O}_X -bimodule structure, with a standard filtration. Considering their graded quotients, induces a canonical map

$$gr(\bar{\phi}) : gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L}')) \rightarrow gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L})).$$

Also, we get a canonical map between the associated sheaf of symmetric algebras, denote it by

$$\tilde{\phi} : Sym_{\mathcal{O}_X}(\mathcal{L}') \rightarrow Sym_{\mathcal{O}_X}(\mathcal{L}).$$

Theorem 7.7. *Let \mathcal{L} be a locally free Lie algebroid over (X, \mathcal{O}_X) and (\mathcal{L}', ϕ) a locally free \mathcal{L} -Lie algebroid. Then the PBW isomorphisms (of graded \mathcal{O}_X -algebras)*

$$\psi_{\mathcal{L}'} : gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L}')) \rightarrow Sym_{\mathcal{O}_X}(\mathcal{L}'), \quad \psi_{\mathcal{L}} : gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L})) \rightarrow Sym_{\mathcal{O}_X}(\mathcal{L}),$$

satisfies the compatibility condition $\tilde{\phi} \circ \psi_{\mathcal{L}'} = \psi_{\mathcal{L}} \circ gr(\bar{\phi})$.

Proof. The proof is straightforward by the following facts. We need to use the functorial property of $\mathcal{U}(\mathcal{O}_X, -)$ and of $Sym_{\mathcal{O}_X}(-)$, and then applying the generalized PBW theorem 7.5 on \mathcal{L}' and on \mathcal{L} . \square

Corollary 7.8. *For a locally free Lie algebroid $\tilde{\mathcal{L}}$ we have a canonical embedding $\iota_{\tilde{\mathcal{L}}} : \tilde{\mathcal{L}} \rightarrow \mathcal{U}(\mathcal{O}_X, \tilde{\mathcal{L}})$. Thus, for locally free Lie algebroids \mathcal{L}' and \mathcal{L} we have the following maps*

$$\mathcal{L}' \xrightarrow{\iota_{\mathcal{L}'}} \mathcal{U}(\mathcal{O}_X, \mathcal{L}') \xrightarrow{gr} gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L}')) \xrightarrow{\psi_{\mathcal{L}'}} Sym_{\mathcal{O}_X}(\mathcal{L}'),$$

and

$$\mathcal{L} \xrightarrow{\iota_{\mathcal{L}}} \mathcal{U}(\mathcal{O}_X, \mathcal{L}) \xrightarrow{gr} gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L})) \xrightarrow{\psi_{\mathcal{L}}} Sym_{\mathcal{O}_X}(\mathcal{L}).$$

Also, for \mathcal{L} -Lie algebroid (\mathcal{L}', ϕ) we have the following induced maps

$$gr(\mathcal{U}(\phi)) : gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L}')) \rightarrow gr(\mathcal{U}(\mathcal{O}_X, \mathcal{L})),$$

and

$$\tilde{\phi} : Sym_{\mathcal{O}_X}(\mathcal{L}') \rightarrow Sym_{\mathcal{O}_X}(\mathcal{L}).$$

Hence we get the following identity of maps as compatibility condition

$$\tilde{\phi} \circ \psi_{\mathcal{L}'} \circ gr \circ \iota_{\mathcal{L}'} = \psi_{\mathcal{L}} \circ gr \circ \iota_{\mathcal{L}} \circ \phi.$$

8. HOMOLOGY AND COHOMOLOGY CORRESPONDENCE

In this section, we extend the duality correspondence between Lie algebroid homology and cohomology [13,20,21,44] to the \mathcal{L} -Lie algebroid setting. The essential notions required for this extension are outlined in Remark 2.21, Section 6, and Section 7.

First, we prove a general result and then we apply it in our case.

Lemma 8.1. *Let $\theta : \mathcal{E}' \rightarrow \mathcal{E}$ be a \mathcal{O}_X -module monomorphism. Then it induces a Lie algebroid homomorphism*

$$\tilde{\theta} : At(\mathcal{E}) \rightarrow At(\mathcal{E}').$$

Proof. Consider the canonical map induced from θ in the following:

$$\bar{\theta} : \mathcal{E}nd_{\mathbb{K}_X}(\mathcal{E}) \rightarrow \mathcal{E}nd_{\mathbb{K}_X}(\mathcal{E}')$$

defined as $\bar{D} := \bar{\theta}(D)$ where $\theta(\bar{D}(s')) = D(\theta(s'))$, for $s' \in \mathcal{E}'$, $D \in \mathcal{E}nd_{\mathbb{K}_X}(\mathcal{E})$. The above map is well defined due to the condition of injectiveness of the map θ . It forms an \mathcal{O}_X -linear map which is also a \mathbb{K}_X -Lie algebra homomorphism as follows. To show $\bar{\theta}(fD) = \bar{f}\bar{D} = f\bar{D}$ for any $f \in \mathcal{O}_X$, $D \in \mathcal{E}nd_{\mathbb{K}_X}(\mathcal{E})$, notice that

$$\theta(\bar{f}\bar{D}(s')) = fD(\theta(s')) = f\theta(\bar{D})(s') = \theta(f\bar{D})(s'),$$

holds for all $s' \in \mathcal{E}$. On the other hand, to show $\bar{\theta}([D_1, D_2]_c) = [\bar{D}_1, \bar{D}_2]_c$ for any $D_1, D_2 \in \mathcal{E}$, notice that

$$\theta(\bar{\theta}([D_1, D_2]_c)(s')) = [D_1, D_2]_c(\theta(s')) = D_1(D_2(\theta(s'))) - D_2(D_1(\theta(s'))) = D_1(\theta(\bar{D}_2(s'))) - D_2(\theta(\bar{D}_1(s'))) = \theta(\bar{D}_1(\bar{D}_2(s'))) - \bar{D}_2(\bar{D}_1(s')),$$

for all $s' \in \mathcal{E}$, and then use the injectiveness of the map θ .

We want to show that the restriction of the map $\bar{\theta}$ on the \mathcal{O}_X -submodule and \mathbb{K}_X -Lie subalgebra $\mathcal{A}t(\mathcal{E})$ of $\mathcal{E}nd_{\mathbb{K}_X}(\mathcal{E})$ defines a Lie algebroid homomorphism

$$\bar{\theta} : \mathcal{A}t(\mathcal{E}) \rightarrow \mathcal{A}t(\mathcal{E}').$$

Let σ and σ' be the anchor maps for $\mathcal{A}t(\mathcal{E})$ and $\mathcal{A}t(\mathcal{E}')$ respectively, and $D \in \mathcal{A}t(\mathcal{E})$. Given any Lie algebroid homomorphism $\phi : \mathcal{A}t(\mathcal{E}) \rightarrow \mathcal{A}t(\mathcal{E}')$, we have $\sigma = \sigma' \circ \phi$. Thus, if $\bar{\theta}$ defines such a map then $\sigma'_{\bar{D}}(f) = (\sigma' \circ \bar{\theta})(D)(f) = \sigma(D)(f) = \sigma_D(f)$ holds for all $D \in \mathcal{A}t(\mathcal{E})$, $f \in \mathcal{O}_X$. Thus, our claim is $\bar{D} = \bar{\theta}(D) \in \mathcal{A}t(\mathcal{E}')$, i.e. \bar{D} satisfies the following Leibniz rule

$$\bar{D}(fs') = f\bar{D}(s') + \sigma_D(f)s',$$

for all $f \in \mathcal{O}_X$, $s' \in \mathcal{E}'$. Notice that, the following holds

$$\theta(\bar{D}(fs')) = D(\theta(fs')) = D(f\theta(s')) = fD(\theta(s')) + \sigma_D(f)\theta(s') = f\theta(\bar{D}(s')) + \sigma_D(f)\theta(s') = \theta(f\bar{D}(s') + \sigma_D(f)s').$$

The injectivity of the map θ provides the required condition for the well definedness of the restriction map $\bar{\theta}$. The well definedness of this map ensures that the map is a Lie algebroid homomorphism using definition of the map $\bar{\theta}$ and the notion of Atiyah algebroid. \square

Condition 8.2. Let (\mathcal{L}', ϕ) be a \mathcal{L} -Lie algebroid such that both \mathcal{L}' and \mathcal{L} are locally free \mathcal{O}_X -module of rank n for some $n \in \mathbb{N}$. Then the following results hold using Lemma 8.1.

Remark 8.3. In particular, if we consider $\mathcal{L}' = \mathcal{T}_X(-\log Y)$ and $\mathcal{L} = \mathcal{T}_X$ for some principal free divisor Y of a complex manifold (or smooth algebraic variety) X , the above conditions hold.

Theorem 8.4. Let ∇ be a flat \mathcal{L} -connection on $\wedge^n_{\mathcal{O}_X} \mathcal{L}$ and the map $\phi : \mathcal{L}' \rightarrow \mathcal{L}$ be an injective map. Then these induces a flat $(\mathcal{L}', \mathcal{L})$ -connection on $\wedge^n_{\mathcal{O}_X} \mathcal{L}'$, provides the graded vector space homomorphism

$$\mathbb{H}_\bullet(\phi) : \mathbb{H}_\bullet(\mathcal{L}', \nabla') \rightarrow \mathbb{H}_\bullet(\mathcal{L}, \nabla).$$

Proof. Consider the composition of the following Lie algebroid homomorphism

$$\mathcal{L}' \xrightarrow{\phi} \mathcal{L} \xrightarrow{\nabla} \mathcal{A}t(\wedge^n_{\mathcal{O}_X} \mathcal{L}) \xrightarrow{\theta} \mathcal{A}t(\wedge^n_{\mathcal{O}_X} \mathcal{L}'),$$

where $\theta := \widetilde{\wedge^n \phi}$ is the induced map associated with the map $\wedge^n \phi : \wedge^n_{\mathcal{O}_X} \mathcal{L}' \rightarrow \wedge^n_{\mathcal{O}_X} \mathcal{L}$ on the level of Atiyah algebroids. It provides the flat $(\mathcal{L}', \mathcal{L})$ -connection $\nabla' := \theta \circ \nabla \circ \phi$ on $\wedge^n_{\mathcal{O}_X} \mathcal{L}'$.

Let the flat \mathcal{L} -connection ∇ induces the exact generator $\partial : \wedge^\bullet_{\mathcal{O}_X} \mathcal{L} \rightarrow \wedge^{\bullet-1}_{\mathcal{O}_X} \mathcal{L}$ and the flat \mathcal{L}' -connection ∇' induces the exact generator $\partial' : \wedge^\bullet_{\mathcal{O}_X} \mathcal{L}' \rightarrow \wedge^{\bullet-1}_{\mathcal{O}_X} \mathcal{L}'$. We need to show that

$$\partial \circ \wedge^\bullet \phi = \wedge^{\bullet-1} \phi \circ \partial'.$$

One can verify that $\partial' = (\wedge^{\bullet-1} \phi)^{-1} \circ \partial \circ \wedge^\bullet \phi$ holds, using the isomorphism $\phi^{-1} : \mathcal{I}m(\mathcal{L}') \rightarrow \mathcal{L}'$.

The required graded vector space homomorphism $\mathbb{H}_\bullet(\phi)$ on the level of associated Lie algebroid homologies of \mathcal{L}' and \mathcal{L} with coefficient in $(\wedge^\bullet_{\mathcal{O}_X} \mathcal{L}', \nabla')$ and $(\wedge^\bullet_{\mathcal{O}_X} \mathcal{L}, \nabla)$ respectively, follows from the map ϕ . \square

Remark 8.5. Recall that for a flat \mathcal{L} -connection (or an \mathcal{L} -module) (\mathcal{E}, ∇) , we define an analogue of the complex (1), denoted by $\Omega^\bullet_{\mathcal{L}}(\mathcal{E})$, known as the Chevalley–Eilenberg–de Rham complex of \mathcal{L} with coefficient in \mathcal{E} , using the connection map ∇ in place of the anchor map \mathfrak{a} . The graded algebra of hypercohomologies associated with this complex is denoted by $\mathbb{H}^\bullet(\mathcal{L}, \mathcal{E})$ (see [5, 34, 37]). Notice that the standard case described in Section 2.3 is recovered when the \mathcal{L} -module $(\mathcal{O}_X, \mathfrak{a})$ is considered.

Remark 8.6. *Also, we have the induced graded vector space homomorphism (Condition 8.2)*

$$\mathbb{H}^\bullet(\phi) : \mathbb{H}^\bullet(\mathcal{L}, \wedge_{\mathcal{O}_X}^n \mathcal{L}) \rightarrow \mathbb{H}^\bullet(\mathcal{L}', \wedge_{\mathcal{O}_X}^n \mathcal{L}').$$

The map $\mathbb{H}^\bullet(\phi)$ is determined by the homomorphism of cochain complexes

$$\wedge_{\mathcal{O}_X}^\bullet \phi^* \otimes_{\mathcal{O}_X} \wedge_{\mathcal{O}_X}^n \phi^* : \wedge_{\mathcal{O}_X}^\bullet \mathcal{L}^* \otimes_{\mathcal{O}_X} \wedge_{\mathcal{O}_X}^n \mathcal{L} \rightarrow \wedge_{\mathcal{O}_X}^\bullet \mathcal{L}'^* \otimes_{\mathcal{O}_X} \wedge_{\mathcal{O}_X}^n \mathcal{L}',$$

where the dual map $\phi^* : \mathcal{L}^* \rightarrow \mathcal{L}'^*$ of ϕ can also be viewed as an \mathcal{O}_X -module homomorphism from \mathcal{L} to \mathcal{L}' , as both \mathcal{L} and \mathcal{L}' are locally free \mathcal{O}_X modules of finite rank.

Rinehart demonstrated that the cohomology of a (\mathbb{K}, A) -Lie-Rinehart algebra L with coefficient in an L -module E , which is projective over A , can be described using the derived functor Ext in the category of modules over the universal enveloping algebra $\mathcal{U}(A, L)$, as $H^\bullet(L, E) \cong Ext_{\mathcal{U}(A, L)}^\bullet(A, E)$ (see [19,36]).

Huebschmann employed a dual approach to this concept in an appropriate setting. Consider a (\mathbb{K}, A) -Lie-Rinehart algebra that is also a projective module of rank n (for some $n \in \mathbb{N}$) over R , together with a BV-algebra structure $(\wedge_A^\bullet L, \partial)$. In this context, Huebschmann introduced the notion of a flat right (A, L) -connection on R , which is equivalent to a flat left (A, L) -connection ∇ (induced from the generating operator ∂) on $\wedge_A^n L$. He has defined the notion of Lie-Rinehart homology $H_\bullet(L, A_\nabla)$ and expressed it by the derived functor Tor in the category of right $\mathcal{U}(A, L)$ -modules using the Rinehart resolution, where A_∇ can be viewed as $(\wedge_A^n L, \nabla)$. Explicitly, he showed that $H_\bullet(L, A_\nabla) \cong Tor_{\bullet}^{\mathcal{U}(A, L)}(A_\nabla, A)$ (see [20]).

In the following, we recall the duality theorem for Lie-Rinehart algebras (see Theorem 4, [20]). A very particular case of this theorem was first addressed in the context of smooth Lie algebroids by P. Xu ([44]). However, J. Huebschmann ([20,21]) later developed a more general formulation for Lie-Rinehart algebras, which does not directly follow from Xu's specific case. To extend these ideas to our setting, we use the description provided for Lie-Rinehart algebras as a local framework.

Theorem 8.7. *For a Lie-Rinehart algebra (A, L) where L is projective A -module of rank n for some $n \in \mathbb{N}$, the homology $H_\bullet(L, A_\nabla)$ of L with coefficient in the right (A, L) -module A , i.e., that of the BV-algebra $(\wedge_A^\bullet L, \partial)$ (where the exact generator ∂ is induced by the flat L -connection ∇) is naturally isomorphic (as graded vector spaces) to the cohomology $H^{n-\bullet}(L, \wedge_A^n L)$ of L with coefficients in the L -module $(\wedge_A^n L, \nabla)$.*

In particular, one can view the case of smooth context, by considering $A = C^\infty(M)$ for some smooth manifold M , and using the fact that finitely generated projective A -module is equivalent to a smooth vector bundle on M . That is, L is the space of global sections $\Gamma(\tilde{L})$ of some Lie algebroid \tilde{L} over M . Note that, in [44], the duality correspondence described with coefficient taking in the trivial line bundle $\wedge^n \tilde{L}$ (or in free A -module $\wedge_A^n L$ of rank 1). The case associated with non trivial line bundle as coefficient (twisted coefficients) is not described explicitly, but mentioned the possibility of that generality (see [13]).

It is well known assuming some description on Lie-Rinehart algebras, how to extend it in the context of Lie algebroids in the algebro-geometric settings (see [5,9,22,41]). Thus, we can follow the approach described in [20], and conclude that an analogue of the result Theorem 8.7 holds for locally free Lie algebroids of finite rank over (X, \mathcal{O}_X) context, as follows.

Remark 8.8. *Let \mathcal{L} be a Lie algebroid of rank n , having a BV-algebra structure on $\wedge_{\mathcal{O}_X}^\bullet \mathcal{L}$ with exact generator ∂ . Then the (hyper)homology of \mathcal{L} with coefficient in the right \mathcal{L} -module $\mathcal{O}_{X, \nabla}$ (this notion is equivalent to the left \mathcal{L} -module $(\wedge_{\mathcal{O}_X}^n \mathcal{L}, \nabla)$) where ∇ induces from ∂ , is naturally isomorphic to the (hyper)cohomology of \mathcal{L} with coefficient in the \mathcal{L} -module $(\wedge_{\mathcal{O}_X}^n \mathcal{L}, \nabla)$. Thus, there is the canonical (duality) isomorphism*

$$(5) \quad \tilde{\psi}_{\mathcal{L}} : \mathbb{H}_\bullet(\mathcal{L}, \nabla) \rightarrow \mathbb{H}^{n-\bullet}(\mathcal{L}, \wedge_{\mathcal{O}_X}^n \mathcal{L}).$$

Here, we show an analogue of the duality isomorphism (5) between Lie algebroid homology and Lie algebroid cohomology (see [20,21,44]), in the context of \mathcal{L} -Lie algebroids.

Theorem 8.9. *Let ∇ be a flat \mathcal{L} -connection on $\wedge_{\mathcal{O}_X}^n \mathcal{L}$ and the map $\phi : \mathcal{L}' \rightarrow \mathcal{L}$ be an injective map (satisfying Condition 8.2). Then, the canonical duality isomorphism*

$$\tilde{\psi}_{\mathcal{L}'} : \mathbb{H}_\bullet(\mathcal{L}', \nabla') \rightarrow \mathbb{H}^{n-\bullet}(\mathcal{L}', \wedge_{\mathcal{O}_X}^n \mathcal{L}')$$

factors through the duality isomorphism $\tilde{\psi}_{\mathcal{L}} : \mathbb{H}_{\bullet}(\mathcal{L}, \nabla) \cong \mathbb{H}^{n-\bullet}(\mathcal{L}, \wedge_{\mathcal{O}_X}^n \mathcal{L})$ as $\tilde{\psi}_{\mathcal{L}'} = \mathbb{H}^{n-\bullet}(\phi) \circ \tilde{\psi}_{\mathcal{L}} \circ \mathbb{H}_{\bullet}(\phi)$.

Proof. Let $\bar{\mathcal{L}}$ be a locally free Lie algebroid of rank n . If there is a flat $\bar{\mathcal{L}}$ -connection $\bar{\nabla}$ on $\wedge_{\mathcal{O}_X}^n \bar{\mathcal{L}}$, then we get the BV-algebra $(\wedge_{\mathcal{O}_X}^{\bullet} \bar{\mathcal{L}}, \bar{\delta})$ where $\bar{\delta} : \wedge_{\mathcal{O}_X}^{\bullet} \bar{\mathcal{L}} \rightarrow \wedge_{\mathcal{O}_X}^{\bullet-1} \bar{\mathcal{L}}$ is the BV-operator. An algebro-geometric analogue of the duality isomorphism for finitely generated projective Lie-Rinehart algebra (Theorem 8.7) is applicable for the Lie algebroid $\bar{\mathcal{L}}$ (Remark 8.8) as follows:

Using the locally free finite rank condition, there is an open cover $\{U_x \mid x \in X\}$ of X , for which we get a $\mathcal{O}_X|_{U_x}$ -module isomorphism $\bar{\mathcal{L}}|_{U_x} \cong \oplus^n \mathcal{O}_X|_{U_x}$. Consider the sheafification of the presheaf homomorphism

$$U \mapsto \bar{\psi}_{\bar{\mathcal{L}}(U)} : H_{\bullet}(\bar{\mathcal{L}}(U), \nabla_U) \rightarrow H^{n-\bullet}(\bar{\mathcal{L}}(U), \wedge_{\mathcal{O}_X(U)}^n \bar{\mathcal{L}}(U)),$$

which is an isomorphism as we have the local duality isomorphisms $\bar{\psi}_{\bar{\mathcal{L}}(U_x)}$ for each $x \in X$, denote it by $\bar{\psi}_{\bar{\mathcal{L}}}$. That is, the associated (co)homology sheaves are isomorphic as

$$\bar{\psi}_{\bar{\mathcal{L}}} : \mathcal{H}_{\bullet}(\bar{\mathcal{L}}, \nabla) \xrightarrow{\cong} \mathcal{H}^{n-\bullet}(\bar{\mathcal{L}}, \wedge_{\mathcal{O}_X}^n \bar{\mathcal{L}}).$$

Therefore, applying hyper(co)homology functor we get the graded vector space isomorphism

$$\tilde{\psi}_{\bar{\mathcal{L}}} := \mathbb{H}(\bar{\psi}_{\bar{\mathcal{L}}}) : \mathbb{H}_{\bullet}(\bar{\mathcal{L}}, \nabla) \xrightarrow{\cong} \mathbb{H}^{n-\bullet}(\bar{\mathcal{L}}, \wedge_{\mathcal{O}_X}^n \bar{\mathcal{L}}).$$

Thus, the above duality isomorphism holds for the Lie algebroids \mathcal{L} and \mathcal{L}' , denote these by $\psi_{\mathcal{L}}$ and $\psi'_{\mathcal{L}}$ respectively. Using the morphisms described in Theorem 8.4 and Remark 8.6 we get the required result. \square

Remarks on Lie Algebroids over Schemes. Our discussions apply to Lie algebroids over Noetherian separated schemes (or schemes of finite type) defined over a field of characteristic zero (see [5,6,22]). Moreover, these results can be extended to the broader context of Lie algebroids over ringed sites (see [8,9]), which requires adopting a more advanced abstraction perspective—such as the notions of formal geometry and coordinate spaces, rather than classical geometric approaches.

Acknowledgments. The second author expresses his heartfelt gratitude to Prof. Ashis Mandal for introducing him to several foundational concepts that have significantly shaped this work. He also extends his sincere thanks to Dr. Satyendra Kumar Mishra for his insightful suggestions and valuable conversations.

REFERENCES

- [1] M. F. Atiyah. Complex analytic connections in fibre bundles. *Trans. Amer. Math. Soc.*, 85:181–207, 1957.
- [2] Indranil Biswas, Arijit Dey, and Mainak Poddar. Equivariant principal bundles and logarithmic connections on toric varieties. *Pacific J. Math.*, 280(2):315–325, 2016.
- [3] Indranil Biswas, Vicente Muñoz, and Jonathan Sánchez. Equivariant vector bundles and logarithmic connections on toric varieties. *J. Algebra*, 384:227–241, 2013.
- [4] Indranil Biswas, S. Senthamarai Kannan, and D. S. Nagaraj. Equivariant principal bundles for G -actions and G -connections. *Complex Manifolds*, 2(1):178–185, 2015.
- [5] Ugo Bruzzo. Lie algebroid cohomology as a derived functor. *J. Algebra*, 483:245–261, 2017.
- [6] Ugo Bruzzo, Igor Mencattini, Vladimir N. Rubtsov, and Pietro Tortella. Nonabelian holomorphic Lie algebroid extensions. *Internat. J. Math.*, 26(5):1550040, 26, 2015.
- [7] Damien Calaque. Formality for Lie algebroids. *Comm. Math. Phys.*, 257(3):563–578, 2005.
- [8] Damien Calaque, Carlo A. Rossi, and Michel van den Bergh. Hochschild (co)homology for Lie algebroids. *Int. Math. Res. Not. IMRN*, (21):4098–4136, 2010.
- [9] Damien Calaque and Michel Van den Bergh. Hochschild cohomology and Atiyah classes. *Adv. Math.*, 224(5):1839–1889, 2010.
- [10] Francisco J. Castro-Jiménez, Luis Narváez-Macarro, and David Mond. Cohomology of the complement of a free divisor. *Trans. Amer. Math. Soc.*, 348(8):3037–3049, 1996.
- [11] V. I. Danilov. Cohomology of algebraic varieties. In *Current problems in mathematics. Fundamental directions, Vol. 35 (Russian)*, Itogi Nauki i Tekhniki, pages 5–130, 272. Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989.
- [12] Jyoti Dasgupta, Bivas Khan, and Mainak Poddar. Logarithmic connections on principal bundles over normal varieties. *arXiv:2211.03047v2 [math.AG]*, 2023.
- [13] Sam Evens, Jiang-Hua Lu, and Alan Weinstein. Transverse measures, the modular class and a cohomology pairing for Lie algebroids. *Quart. J. Math. Oxford Ser. (2)*, 50(200):417–436, 1999.

- [14] Rui Loja Fernandes. Lie algebroids, holonomy and characteristic classes. *Adv. Math.*, 170(1):119–179, 2002.
- [15] Franc Forstnerič. *Stein manifolds and holomorphic mappings*, volume 56 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer, Heidelberg, 2011. The homotopy principle in complex analysis.
- [16] Murray Gerstenhaber. The cohomology structure of an associative ring. *Ann. of Math. (2)*, 78:267–288, 1963.
- [17] Murray Gerstenhaber and Samuel D. Schack. Algebras, bialgebras, quantum groups, and algebraic deformations. In *Deformation theory and quantum groups with applications to mathematical physics (Amherst, MA, 1990)*, volume 134 of *Contemp. Math.*, pages 51–92. Amer. Math. Soc., Providence, RI, 1992.
- [18] Werner Greub, Stephen Halperin, and Ray Vanstone. *Connections, curvature, and cohomology*. Pure and Applied Mathematics, Vol. 47-III. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1976. Volume III: Cohomology of principal bundles and homogeneous spaces.
- [19] Johannes Huebschmann. Poisson cohomology and quantization. *J. Reine Angew. Math.*, 408:57–113, 1990.
- [20] Johannes Huebschmann. Lie-Rinehart algebras, Gerstenhaber algebras and Batalin-Vilkovisky algebras. *Ann. Inst. Fourier (Grenoble)*, 48(2):425–440, 1998.
- [21] Johannes Huebschmann. Duality for Lie-Rinehart algebras and the modular class. *J. Reine Angew. Math.*, 510:103–159, 1999.
- [22] Mikhail Kapranov. Free Lie algebroids and the space of paths. *Selecta Math. (N.S.)*, 13(2):277–319, 2007.
- [23] Bivas Khan and Mainak Poddar. G -connections on principal bundles over complete G -varieties. *J. Pure Appl. Algebra*, 229(1):Paper No. 107816, 24, 2025.
- [24] R. L. Klaasse. Geometric structures and Lie algebroids. *arXiv:1712.09560 [math.SG]*, Thesis (Ph.D.)-University of Utrecht., 2017.
- [25] Y. Kosmann-Schwarzbach. Exact Gerstenhaber algebras and Lie bialgebroids. *Acta Appl. Math.*, 41(1-3):153–165, 1995. Geometric and algebraic structures in differential equations.
- [26] Camille Laurent-Gengoux and Leonid Ryvkin. The neighborhood of a singular leaf. *J. Éc. polytech. Math.*, 8:1037–1064, 2021.
- [27] Helge Maakestad. Chern classes and Lie-Rinehart algebras. *Indag. Math. (N.S.)*, 18(4):589–599, 2007.
- [28] Helge Øystein Maakestad. Algebraic connections on projective modules with prescribed curvature. *J. Algebra*, 436:161–227, 2015.
- [29] Luis Narváez Macarro. Differential structures in commutative algebra. *Mini-course at the XXIII Brazilian Algebra Meeting, July 27 - August 1, Maringá, Brazil.*, 2014.
- [30] Kirill C. H. Mackenzie. *General theory of Lie groupoids and Lie algebroids*, volume 213 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2005.
- [31] Kirill C. H. Mackenzie and Ping Xu. Lie bialgebroids and Poisson groupoids. *Duke Math. J.*, 73(2):415–452, 1994.
- [32] Ashis Mandal and Abhishek Sarkar. On Lie algebroid over algebraic spaces. *Comm. Algebra*, 51(4):1594–1613, 2023.
- [33] I. Moerdijk and J. Mrčun. On the universal enveloping algebra of a Lie algebroid. *Proc. Amer. Math. Soc.*, 138(9):3135–3145, 2010.
- [34] Brent Pym. *Poisson Structures and Lie Algebroids in Complex Geometry*. ProQuest LLC, Ann Arbor, MI, 2013. Thesis (Ph.D.)-University of Toronto (Canada).
- [35] S. Ramanan. *Global calculus*, volume 65 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2005.
- [36] George S. Rinehart. Differential forms on general commutative algebras. *Trans. Amer. Math. Soc.*, 108:195–222, 1963.
- [37] A. Sarkar. Cohomology of Lie algebroid over algebraic spaces. *arXiv:2111.01735v8 [math.DG]*, 2024.
- [38] T. Schedler. Deformation of algebras in noncommutative geometry. *arXiv:1212.0914v3 [math.RA]*, 2019.
- [39] Pietro Tortella. Λ -modules and holomorphic Lie algebroid connections. *Cent. Eur. J. Math.*, 10(4):1422–1441, 2012.
- [40] Pietro Tortella. Representations of Atiyah algebroids and logarithmic connections. *Int. Math. Res. Not. IMRN*, (1):29–46, 2017.
- [41] J. Vilatero. On sheaves of lie-Rinehart algebras. *arXiv:2010.15463v2 [math.DG]*, 2021.
- [42] Charles Weibel. Cyclic homology for schemes. *Proc. Amer. Math. Soc.*, 124(6):1655–1662, 1996.
- [43] Charles A. Weibel. *The K-book*, volume 145 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2013. An introduction to algebraic K -theory.
- [44] Ping Xu. Gerstenhaber algebras and BV-algebras in Poisson geometry. *Comm. Math. Phys.*, 200(3):545–560, 1999.

Mainak Poddar, Department of Mathematics, Indian Institute of Science Education and Research Pune, Pune 411008, Maharashtra, India. e-mail: mainak@iiserpune.ac.in

Abhishek Sarkar Department of Mathematics, Indian Institute of Science Education and Research Pune, Pune 411008, Maharashtra, India. e-mail: abhisheksarkar49@gmail.com