

# An Integral over $(0, \pi)$ for the Distribution Function of a Sum of Independent Gamma Random Variables and for Quadratic Forms of Gaussian Variables

Thomas Royen

TH Bingen, University of Applied Sciences  
Berlinstrasse 109, D55411 Bingen, Germany

e-mail: [thomas.royen@t-online.de](mailto:thomas.royen@t-online.de)

## Abstract.

An integral over the interval  $(0, \pi)$  is given for the cumulative distribution function (cdf) of a sum of independent gamma random variables with different scale and shape parameters. The cdf of a positive definite quadratic form is obtained as a special case with identical shape parameters  $\alpha = \frac{1}{2}$ .

## 1. Introduction

There is a lot of literature on the distribution of quadratic forms, in particular with Gaussian random vectors, see e.g. the extensive chapter 4 in [3]. For all these series representations of the cdf a large number of coefficients has to be computed. It is the purpose of the underlying paper to avoid these computations and to use only the eigenvalues of a positive definite matrix.

Let  $X$  be an  $N_k(0, \Sigma)$ -random vector with a pos. def. covariance matrix  $\Sigma$ . For the quadratic form  $Q = X'CX$  with a pos. def. matrix  $C$  we have the equation

$$P\{Q \leq x\} = P\left\{\sum_{j=1}^k \lambda_j Z_j^2 \leq x\right\} \quad (1.1)$$

with the eigenvalues  $\lambda_j$  of  $\Sigma^{1/2}C\Sigma^{1/2}$  and an  $N_k(0, I_k)$ -random vector  $Z$ , where  $I_k$  denotes the  $k \times k$ -identity matrix. The determinant of a matrix  $M$  will be denoted by  $|M|$ , and  $\|M\|$  is the spectral norm of a symmetrical matrix.

The random variable  $\frac{1}{2}Z_j^2$  has a gamma-pdf  $g_\alpha(x) = \exp(-x)x^{\alpha-1} / \Gamma(\alpha)$  with  $\alpha = \frac{1}{2}$ . Therefore, the distribution of  $\frac{1}{2}Q$  is a special case of the distributions of sums of independent gamma random variables with pdfs  $\lambda_j^{-1}g_{\alpha_j}(\lambda_j^{-1}x)$ . The pdf  $f(x; \alpha_1, \dots, \alpha_k, \lambda_1, \dots, \lambda_k)$  of such a sum has the Laplace transform (Lt)

$$\prod_{j=1}^k (1 + \lambda_j t)^{-\alpha_j}. \quad (1.2)$$

For series expansions of the corresponding cdfs see [4] and for multivariate generalizations of these series see [5].

---

2020 Mathematics Subject Classification: 62E15, 60G50

Keywords and phrases: integral representations for probability distributions, sums of independent gamma random variables, distribution of quadratic forms of Gaussian variables.

## 2. The Integral Representations

For the intended integral representations we use a modified form of the Lt from (1.2). With

$$v = \frac{1}{2}(\lambda_{\max}^{-1} + \lambda_{\min}^{-1}), \quad z = (1 + v^{-1}t)^{-1}, \quad \text{and} \quad \alpha = \sum_{j=1}^k \alpha_j \quad (2.1)$$

we obtain

$$\begin{aligned} \prod_{j=1}^k (1 + \lambda_j t)^{-\alpha_j} &= \prod_{j=1}^k (1 + \lambda_j v t v^{-1})^{-\alpha_j} = \prod_{j=1}^k (1 + v^{-1}t + (\lambda_j v - 1) t v^{-1})^{-\alpha_j} = \\ &= z^\alpha \prod_{j=1}^k (1 + (\lambda_j v - 1)(1 - z))^{-\alpha_j} = \left( v^{-\alpha} \prod_{j=1}^k \lambda_j^{-\alpha_j} \right) z^\alpha \prod_{j=1}^k \left( 1 - (1 - \frac{1}{v \lambda_j}) z \right)^{-\alpha_j}, \end{aligned} \quad (2.2)$$

$$\text{and } \max |c_j| = \max \left| 1 - \frac{1}{v \lambda_j} \right| = \frac{\lambda_{\max} - \lambda_{\min}}{\lambda_{\max} + \lambda_{\min}} < 1.$$

To avoid series expansions with products of binomial series the orthogonality of the functions  $e^{in\varphi}$ ,  $n \in \mathbb{Z}$ , will be used for a product of a power series of  $y = r e^{i\varphi}$  and a power series of  $r^{-1} e^{-i\varphi}$ . For this goal the following function is defined for  $\alpha > 0$ ,  $x \geq 0$ ,  $y \in \mathbb{C}$ :

$$G_\alpha(x, y) := \sum_{n=0}^{\infty} G_{\alpha+n}(x) y^n \quad (2.3)$$

where  $G_\alpha(x)$  is the cdf with the gamma-pdf  $g_\alpha(x) = e^{-x} x^{\alpha-1} / \Gamma(\alpha)$ . Here we need only values  $|y| < 1$ .

Now the equation

$$G_\alpha(x, y) = (1 - y)^{-1} (G_\alpha(x) - y^{1-\alpha} e^{(y-1)x} G_\alpha(xy)). \quad (2.4)$$

is verified. Integration by parts shows  $G_\alpha(y) = \sum_{n=1}^{\infty} g_{\alpha+n}(y)$ , and it is

$$\begin{aligned} (1 - y) G_\alpha(x, y) &= G_\alpha(x) + \sum_{n=1}^{\infty} (G_{\alpha+n}(x) - G_{\alpha+n-1}(x)) y^n = G_\alpha(x) - \sum_{n=1}^{\infty} g_{\alpha+n}(x) y^n = \\ &= G_\alpha(x) - y^{1-\alpha} e^{(y-1)x} \sum_{n=1}^{\infty} g_{\alpha+n}(xy) = G_\alpha(x) - y^{1-\alpha} e^{(y-1)x} G_\alpha(xy). \end{aligned}$$

For  $y = 1$  we obtain  $G(x, 1) := \lim_{y \rightarrow 1} G(x, y) = x g_\alpha(x) + (1 + x - \alpha) G_\alpha(x)$ . Furthermore, a short calculation shows

$$G_\alpha(x, y) = (1 - y)^{-1} (G_{\alpha-1}(x) - y^{1-\alpha} e^{(y-1)x} G_{\alpha-1}(xy)), \quad \alpha \geq 1, \quad G_0 := 1. \quad (2.5)$$

In particular, it is  $G_{1+n}(y) = 1 - e^{-y} \sum_{k=0}^n \frac{y^k}{k!}$  and  $G_{1/2+n}(y) = \text{erf}(y^{1/2}) - e^{-y} \sum_{k=1}^n \frac{y^{k-1/2}}{\Gamma(k+1/2)}$ .

With binomial series we obtain for the Lt from (2.2)

$$\prod_{j=1}^k (1 + \lambda_j t)^{-\alpha_j} = \left( v^{-\alpha} \prod_{j=1}^k \lambda_j^{-\alpha_j} \right) \sum_{n=0}^{\infty} \left( \sum_{(n)} \prod_{j=1}^k \binom{\alpha_j + n_j - 1}{n_j} c_j^{n_j} \right) z^{\alpha+n}, \quad (2.6)$$

where  $\sum_{(n)}$  stands for  $\sum_{n_1 + \dots + n_k = n}$  and consequently for the corresponding cdf:

$$F(x; \alpha_1, \dots, \alpha_k, \lambda_1, \dots, \lambda_k) = \left( v^{-\alpha} \prod_{j=1}^k \lambda_j^{-\alpha_j} \right) \sum_{n=0}^{\infty} \left( \sum_{(n)} \prod_{j=1}^k \binom{\alpha_j + n_j - 1}{n_j} c_j^{n_j} \right) G_{\alpha+n}(vx). \quad (2.7)$$

Then with  $G_\alpha(x, y)$  from (2.4) or (2.5) it follows directly:

**Theorem.** Let  $X_1, \dots, X_k$  be independent gamma random variables with pdfs  $\lambda_j^{-1} g_{\alpha_j}(\lambda_j^{-1} x)$ . Then the cdf of  $\sum_{j=1}^k X_j$  is given by

$$F(x; \alpha_1, \dots, \alpha_k, \lambda_1, \dots, \lambda_k) = \left( v^{-\alpha} \prod_{j=1}^k \lambda_j^{-\alpha_j} \right) \frac{1}{\pi} \int_0^\pi \operatorname{Re} \left( \prod_{j=1}^k (1 - c_j r^{-1} e^{-i\varphi})^{-\alpha_j} \right) G_\alpha(vx, r e^{i\varphi}) d\varphi \quad (2.8)$$

with  $v$  from (2.1),  $c_j = 1 - \frac{1}{v\lambda_j}$  from (2.2) and any  $r \in \left( \frac{\lambda_{\max} - \lambda_{\min}}{\lambda_{\max} + \lambda_{\min}}, 1 \right)$ .

In particular, the cdf of the quadratic form  $\frac{1}{2} Q$  from (1.1) with an  $N_k(0, \Sigma)$ - random vector  $X$  is given by

$$F(x; \frac{1}{2}, \lambda_1, \dots, \lambda_k) = \left( v^{-k/2} |C\Sigma|^{-1/2} \right) \frac{1}{\pi} \int_0^\pi \operatorname{Re} \left( \prod_{j=1}^k (1 - c_j r^{-1} e^{-i\varphi})^{-1/2} \right) G_{k/2}(vx, r e^{i\varphi}) d\varphi. \quad (2.9)$$

For multivariate generalizations of formula (2.8) see [5], but the numerical evaluation of the corresponding multivariate integrals is frequently difficult because of a rather high total variation of the integrands. Here only such a formula is given for the cdf of a single  $p$ -variate gamma distribution with the Lt  $|I_p + \Sigma T|^{-\alpha}$ , where  $\Sigma$  is pos. def. with the eigenvalues  $\lambda_k$  and  $T := \operatorname{Diag}(t_1, \dots, t_p)$ ,  $t_k \geq 0$ . This distribution exists at least for  $2\alpha \in \mathbb{N} \cup ([\frac{p-1}{2}], \infty)$ , see [6], and for infinitely divisible multivariate gamma distributions for all  $\alpha > 0$ , see [1]. The corresponding cdf is

$$F(x_1, \dots, x_p; \alpha, \Sigma) = \left( v^{-\alpha p} |\Sigma|^{-\alpha} \right) \frac{1}{(2\pi)^p} \int_{(-\pi, \pi)^p} |I_p - CY^{-1}|^{-\alpha} \prod_{k=1}^p G_\alpha(vx_k, y_k) d\varphi_k \quad (2.10)$$

with  $Y = \operatorname{Diag}(y_1, \dots, y_p)$ ,  $y_k = r e^{i\varphi_k}$ ,  $C = I_p - (v\Sigma)^{-1}$ ,  $\|C\| = \frac{\lambda_{\max} - \lambda_{\min}}{\lambda_{\max} + \lambda_{\min}} < 1$  with the eigenvalues  $\lambda_k$  of

the pos. def. matrix  $\Sigma$  and any  $r \in \left( \frac{\lambda_{\max} - \lambda_{\min}}{\lambda_{\max} + \lambda_{\min}}, 1 \right)$ . However, there are numerical more favourable integral

representations for these multivariate gamma cdfs with  $p \leq 4$ , see [2]. The appendix of this paper contains presumably the most comprehensive collection of formulas for multivariate gamma distributions of this type.

## References

- [1] Bapat, R. B., Infinite divisibility of multivariate gamma distributions and M-matrices, Sankyā Series A **51**, (1989), 73-78.
- [2] Dickhaus, T. and Royen, T., A survey on multivariate chi-square distributions and their application in testing multiple hypotheses, Statistics **49**, (2015), 427-454.
- [3] Mathai, A. M. and Provost, S. B., Quadratic forms in random variables, Marcel Dekker, Inc., New York, Basel, Hongkong, 1992.
- [4] Moschopoulos, P. G., The distribution of the sum of independent gamma random variables, Ann. Inst. Statist. Math. **37**, (1985), 541-544.
- [5] Royen, T., Some representations for convolutions of multivariate gamma distribution, Far East J. Theor. Stat. **45**(1), (2013), 21-49.
- [6] Royen, T., A note on the existence of the multivariate gamma distribution, arXiv:1606.04747[math.PR].

