

Nonlocal Choquard equations involving critical Hardy-Littlewood-Sobolev exponent: the effect of the topology of the domain

Mohammed Ali Mohammed Alghamdi

*Department of Mathematics,
King Abdulaziz University, Jeddah 21589, Saudi Arabi
mamalgamdi@kau.edu.sa*

Hichem Chtioui

*Sfax University,
Faculty of Sciences of Sfax,
3018 Sfax, Tunisia.
Hichem.Chtioui@fss.rnu.tn*

Abstract We apply a topological method to prove existence of positive solutions for the nonlinear Choquard equation with upper critical exponent in the sense of Hardy-Littlewood-Sobolev inequality on bounded domains having nontrivial homology group.

Keywords: Choquard equation, Critical nonlinearity, Variational calculus, Singular Homology, Topological method.

MSC 2020. 35A16, 35J20, 35J60, 55N10

1 Introduction

During the last decades, the Choquard equation

$$-\Delta u + V(x)u = \left(\int_{\mathbb{R}^n} \frac{|u(y)|^q}{|x-y|^\mu} dy \right) |u|^{q-2} u \quad \text{in } \mathbb{R}^n, \quad (1.1)$$

where $n \geq 3$, $0 < \mu < n$ and $\frac{2n-\mu}{n} \leq q \leq \frac{2n-\mu}{n-2}$, has attracted the attention and the interest of a lot of researchers, since it appears in the modeling of various physical phenomena, such as quantum mechanics of a polaron at rest [27], Selfgravitatal collapse of a quantum mechanical wave function [26], modeling an electron trapped in its own hole as an approximation to Hartree-Fock theory of one component plasma [9] and so on.

Following standard variational theory, the solutions of the Choquard equation are the critical points of the Euler-Lagrange functional

$$I(u) = \frac{1}{2} \int_{\mathbb{R}^n} |\nabla u|^2 dx - \frac{1}{2q} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(y)|^q |u(x)|^q}{|x-y|^\mu} dx dy,$$

on $H^1(\mathbb{R}^n)$. Recall that by the celebrated Hardy-Littlewood-Sobolev inequality see [8] and [10], the second term of $I(u)$ is well defined on $H^1(\mathbb{R}^n)$, if and only if

$$2_\mu := \frac{2n - \mu}{n} \leq q \leq \frac{2n - \mu}{n - 2} := 2_\mu^*.$$

In this case, the functional I is continuously differentiable on $H^1(\mathbb{R}^n)$. The constants 2_μ and 2_μ^* are termed, respectively, as the lower critical exponent and the upper critical exponent in the sense of Hardy-Littlewood-Sobolev. It is remarkable that the Choquard equation is a nonlinear problem with superlinear nonlinearity, since $2_\mu > 1$, $\forall \mu \in (0, n)$, and it is a nonlocal problem. The nonlocal term which appears in the nonlinearity makes the problem particularly difficult. For $n = 3$, $q = 2$, $\mu = 1$ and V is a positive constant, Lieb [9] obtained the existence of a unique minimizing positive solution (up to translations). Under some restrictions on $V(x)$, Lions [25] proved the existence of a sequence of radially solutions. For $n \geq 3$, $\mu \in (0, n)$, $V = 1$, and suitable range of q , Moros-Van Schaftingen [21], [19] proved the existence of a nontrivial solution of (1.1), if and only if $2_\mu < q < 2_\mu^*$. Their result has obtained by using a Pohozaev type identity and minimizing arguments. Ma-Zhao [23] studied the positive and regularity of the minimizing solutions. For $q = 2_\mu$, the lower critical exponent and a nonconstant potential V , Moros-Van Schaftingen [20] and Cassani-Schaftingen-Zhang [5] studied the existence problem of positive solutions of equation (1.1). they provided sufficient and necessary conditions on the potential $V(x)$ to obtain minimizing solutions.

For the upper critical exponent $q = 2_\mu^*$ and $V = 0$, the Choquard equation reduces to

$$-\Delta u = \left(\int_{\mathbb{R}^n} \frac{|u(y)|^{2_\mu^*}}{|x - y|^\mu} dy \right) |u|^{2_\mu^* - 2} u \quad \text{in } \mathbb{R}^n. \quad (1.2)$$

Using moving plane techniques and analyzes of the associated integral system, Du-Yang [7] and Guo-Hu-Peng-Shuai [15] proved a uniqueness classification of the positive solutions of (1.2). Precisely, they proved that the positive solutions of (1.2) have to be of the form

$$\tilde{\delta}_{(a,\lambda)}(x) = (n(n-2))^{\frac{n-2}{4}} C_{n,\mu}^{\frac{2-n}{2(n-\mu+2)}} S^{\frac{(n-\mu)(2-n)}{4(n-\mu+2)}} \delta_{(a,\lambda)}(x), \quad x \in \mathbb{R}^n,$$

where

$$C_{n,\mu} = \pi^{\frac{\mu}{2}} \frac{\Gamma(\frac{n-\mu}{2})}{\Gamma(n - \frac{\mu}{2})} \left(\frac{\Gamma(n)}{\Gamma(\frac{n}{2})} \right)^{\frac{n-\mu}{n}}, \quad \Gamma(s) = \int_0^\infty \frac{e^{-t}}{t^{1-s}} dt, \quad s > 0,$$

is the Gamma function, S is the best constant of Sobolev and for $a \in \mathbb{R}^n$ and $\lambda > 0$,

$$\delta_{(a,\lambda)}(x) = \frac{\lambda^{\frac{n-2}{2}}}{(1 + \lambda^2 |x - a|^2)^{\frac{n-2}{2}}}.$$

In this paper we restrict our attention to problem (1.2) on bounded domains Ω of \mathbb{R}^n , $n \geq 3$. we are interested to study the existence problem of positive solutions of the following nonlocal Choquard equation with upper critical Hardy-Littlewood-Sobolev exponent

$$-\Delta u = \left(\int_{\Omega} \frac{|u(y)|^{2_\mu^*}}{|x - y|^\mu} dy \right) |u|^{2_\mu^* - 2} u \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega, \quad (1.3)$$

where $\mu \in (0, n)$ and $2_\mu^* = \frac{2n-\mu}{n-2}$.

Since 2_μ^* is critical, a lack of compactness occurs in the sense that the associated Euler Lagrange functional does not satisfy the Palais-Smale condition. This makes the problem of finding positive solutions more challenging. As the celebrate critical elliptic local equation

$$-\Delta u = |u|^{2^*-2} u \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega, \quad (1.4)$$

where $2^* = \frac{2n}{n-2}$ is the critical Sobolev exponent, a concentration compactness principle [24], [22] and Bahri-Coron representations [1], [2] play an important role in solving nonlinear problems of a non compact nature.

In [13], Goel-Radulescu-Sreenadh considred problem (1.3) where Ω is a bounded annular-type domain of \mathbb{R}^n , $n \geq 3$. They proved a global compactness Lemma and used a variational method to show existence of positive solutions when the inner hole of Ω is sufficiently small. Their result is motivated by the celebrate paper of Coron [17] for the local equation (1.4). In [12], Gao-Yang studied the Hartree type Brezis-Nivenberg problem; that is a perturbation of equation (1.3) by adding a linear term " λu , $\lambda \in \mathbb{R}$ " to the Choquard nonlinearity of (1.3). They proved existence results for a suitable range of λ and a non-existence result in the case of $\lambda \leq 0$ and Ω is star shaped domain with respect to the origin. In the papers [12], [11], [14] and [18], the method of Brezis-Nirenberg developed for the local elliptic equation, see [16], has been adopt to study nonlocal Choquard problems with upper critical nonlinearity.

Pushing more the resemblance of the local equation (1.4) and the nonlocal Choquard equation (1.3), we are led to investigate the effect of the domain's topology on the existence results of problem (1.3). In the pioneering paper [2], Bahri-Coron introduced a powerful topological argument which leads to prove the existence of positive solutions of equation (1.4) when Ω has a nontrivial homology group. In this paper we prove that the topological approach of Bahri-Coron can be successfully adopt to study the existence problem of positive solutions of the Choquard equation (1.3). Because of its nonlocal nature, additional difficulties arise in extending such a topological method to the present framework. This requires changes in the construction and novelties in the proof.

Let k be a non-negative integer. We denote $H_k(\Omega)$ the singular homology group of Ω of dimension k with \mathbb{Z}_2 -coefficients. We shall prove the following result.

Theorem 1.1 *Let $n \geq 3$ and let Ω be a bounded domain of \mathbb{R}^n . If there exists a positive integer k_0 such that $H_{k_0}(\Omega)$ is not null, then problem (1.3) has at least a positive solution.*

The proof of Theorem 1.1 will be the subject of Section 3 of this paper. We summarize the idea of the proof in the following three steps.

For a pair of topological spaces (X, Y) , $Y \subset X$, we denote $H_\ell(X, Y)$, $\ell \in \mathbb{N}$, the relative homology of (X, Y) .

Step 1. Under the assumption that $H_{k_0}(\Omega)$ is not trivial, we prove that there exist two sequences of topological pairs (B_p, B_{p-1}) and (E_p, F_p) , $p \in \mathbb{N} \setminus \{0\}$, a sequence of non-zero homology classes (ω_p) such that

$\omega_p \in H_\ell(B_p, B_{p-1})$, for any p and a sequence of relative homology homomorphisms

$$(\phi_p)_\ell: H_\ell(B_p, B_{p-1}) \rightarrow H_\ell(E_p, F_p)$$

such that the first homomorphism of the sequence satisfies

$$(\phi_1)_\ell(\omega_1) \neq 0. \quad (1.5)$$

Step 2. If we suppose that problem (1.3) has no positive solution, we prove that for any $p \in \mathbb{N} \setminus \{0\}$,

$$(\phi_p)_\ell(\omega_p) \neq 0 \implies (\phi_{p+1})_\ell(\omega_{p+1}) \neq 0. \quad (1.6)$$

Step 3. We prove the existence of a positive large integer p_0 such that

$$(\phi_{p_0})_\ell = 0 \text{ for any } \ell \in \mathbb{N}. \quad (1.7)$$

This concludes the proof.

In order to achieve the above steps, we state in the next Section the variational structure associated to problem (1.3) including some preparatory results. We first change the Euler-Lagrange functional I and work with a more convenient functional J . We study the behavior of sequences failing the Palais-Smale condition and Parametrize the neighborhoods of critical points at infinity [3]. Finally, we prove asymptotic estimates and expand the Euler-Lagrange functional J in these neighborhoods. Particularly, we determine exactly the levels of J where the Palais-Smale condition does not hold.

In the following we denote, $\langle u, v \rangle = \int_{\Omega} \nabla u \nabla v dx$ the inner product of $H_0^1(\Omega)$,

$|u|_{1,\Omega} = \left(\int_{\Omega} |\nabla u|^2 dx \right)^{\frac{1}{2}}$ the associated norm and the $\|u\|_{L^t} = \left(\int_{\Omega} |u|^t dx \right)^{\frac{1}{t}}$ norm of $L^t(\Omega)$.

2 Variational results and asymptotic expansions

We start this Section by recalling the Hardy-Littlewood-Sobolev inequality which play a fundamental role in defining the variational approach to critical problems with nonlocal Choquard nonlinearities.

Lemma 2.1 [8], [10]. *For $t_1, t_2 > 1$, such that $\frac{1}{t_1} + \frac{1}{t_2} = \frac{2n-\mu}{n}$, we have*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{f(y)g(x)}{|x-y|^\mu} dx dy \leq C \|f\|_{L^{t_1}} \|g\|_{L^{t_2}}, \quad \forall (f, g) \in L^{t_1}(\mathbb{R}^n) \times L^{t_2}(\mathbb{R}^n). \quad (2.1)$$

Here C is a positive constant independent of f and g .

Particularly, if $t_1 = t_2 = \frac{2n}{2n-\mu}$, then C equals to $C_{n,\mu}$; the positive constant introduced in the above Section. In this case, there is equality in (2.1), if and only if $f = g$ (up to a multiplicative constant) and

$$g(x) = A(\lambda^2 + |x-a|^2)^{\frac{\mu-2n}{2}},$$

where $A \in \mathbb{C}$, $\lambda \in \mathbb{R} \setminus \{0\}$ and $a \in \mathbb{R}^n$.

Let us observe that under the Hardy-Littlewood-Sobolev inequality, the integral

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(y)|^q |u(x)|^q}{|x-y|^\mu} dx dy \quad (2.2)$$

is well defined, if $|u|^q \in L^t(\mathbb{R}^n)$ such that $\frac{2}{t} = \frac{2n-\mu}{n}$. Thus, by Sobolev embedding, integral (2.2) is defined for $u \in H^1(\mathbb{R}^n)$, only if $\frac{2n-\mu}{n} \leq q \leq \frac{2n-\mu}{n-2}$.

In the special case of the upper critical Hardy-Littlewood-Sobolev exponent $2_\mu^* = \frac{2n-\mu}{n-2}$ and for an open set Ω of \mathbb{R}^n , $n \geq 3$, it is proved in [12] that

$$\|u\|_{HL} = \left(\int_{\Omega} \int_{\Omega} \frac{|u(y)|^{2_\mu^*} |u(x)|^{2_\mu^*}}{|x-y|^\mu} dx dy \right)^{\frac{1}{2 \cdot 2_\mu^*}}, \quad u \in L^{2_\mu^*}(\Omega),$$

is a norm on $L^{2_\mu^*}(\Omega)$, where $2_\mu^* = \frac{2n}{n-2}$. In this case the best constant $S_{HL}(\Omega)$ defined by

$$S_{HL}(\Omega) = \inf_{u \in H_0^1(\Omega) \setminus \{0\}} \frac{|u|_{1,\Omega}}{\|u\|_{HL}}, \quad (2.3)$$

is independent of Ω and it is never achieved except $\Omega = \mathbb{R}^n$, see [12]. In addition, the unique minimizers of $S_{HL}(\mathbb{R}^n)$ are unique and of the form $u = c\delta_{(a,\lambda)}$, where c is a positive constant and the parameters $(a, \lambda) \in \mathbb{R}^n \times \mathbb{R}^+$. Here

$$\delta_{(a,\lambda)} = \frac{\lambda^{\frac{n-2}{2}}}{(1 + \lambda^2 |x - a|^2)^{\frac{n-2}{2}}}, \quad x \in \mathbb{R}^n. \quad (2.4)$$

As a consequence of the Hardy-Littlewood-Sobolev inequality, the functional I defined by

$$I(u) = \frac{1}{2} |u|_{1\Omega}^2 - \frac{1}{2 \cdot 2_\mu^*} \|u\|_{HL}^{2 \cdot 2_\mu^*},$$

is well defined on $H_0^1(\Omega)$ and it is of class C^1 . It is straightforward to see that for any $u \in H_0^1(\Omega)$, we have

$$I'(u) = -\Delta u - \left(\int_{\Omega} \frac{|u(y)|^{2_\mu^*}}{|x-y|^\mu} dy \right) |u|^{2_\mu^*-2} u.$$

Therefore the solutions of problem (1.3) are the critical points of I . Since 2_μ^* is critical in the sense of Hardy-Littlewood-Sobolev inequality, the functional I does not satisfy the Palais-Smale condition. A description of sequences failing the Palais-Smale condition has been established in [13]. As a product, existence of a positive critical point of I is obtained under the assumption that Ω is a bounded annular-type domain with inner hole small enough. Let

$$\Sigma = \{u \in H_0^1(\Omega), |u|_{1\Omega} = 1\}.$$

On Σ , we define

$$J_1(u) = \sup_{\lambda \geq 0} I(\lambda u).$$

Lemma 2.2 *For any $u \in \Sigma$, there exists a unique $\lambda = \lambda(u) > 0$ such that $J_1(u) = I(\lambda(u)u)$.*

Proof. Let $u \in \Sigma$ and let $f(\lambda) = I(\lambda u)$, $\lambda \geq 0$. We have

$$f'(\lambda) = \lambda - \lambda^{2 \cdot 2_\mu^* - 1} \|u\|_{HL}^{2 \cdot 2_\mu^*}.$$

Therefore, f is an increasing function near 0, $f(0) = 0$ and $f(\lambda) \rightarrow -\infty$, as $\lambda \rightarrow +\infty$. Then there exists $\lambda = \lambda(u) > 0$ such that $f(\lambda(u)) = \max_{\lambda \geq 0} f(\lambda)$. Since $f'(\lambda(u)) = 0$, then $\lambda(u)$ is unique

and equals to $\|u\|_{HL}^{\frac{-2_\mu^*}{2_\mu^* - 1}}$. □

Let us observe that there is a correspondence between the critical points of J_1 and the critical points of I . Indeed,

Lemma 2.3 *Let $u \in \Sigma$. u is a critical point of J_1 if and only if $\lambda(u)u$ is a critical point of I .*

Proof. For any $u \in \Sigma$, we have by Lemma 2.2.

$$J_1'(u) = \lambda(u)I'(\lambda(u)u) + I'(\lambda(u)u)(u) \cdot \lambda'(u).$$

Using the fact that $f'(\lambda(u)) = 0$, where $f(\lambda)$ is defined in the proof of Lemma 2.2, we get $I'(\lambda(u)u)(u) = 0$ and hence $J_1'(u) = I'(\lambda(u)u)$, up to a positive multiplicative constant. The result follows. □

Note that by the result of Lemma 2.2, $J_1(u)$, $u \in \Sigma$, can be expressed by

$$J_1(u) = \left(\frac{1}{2} - \frac{1}{2 \cdot 2_\mu^*}\right) \frac{1}{\left(\int_{\Omega} \int_{\Omega} \frac{|u(y)|^{2_\mu^*} |u(x)|^{2_\mu^*}}{|x-y|^\mu} dx dy\right)^{\frac{1}{2_\mu^*-1}}},$$

since $\lambda(u) = \|u\|^{-\frac{2_\mu^*}{2_\mu^*-1}}$. Moreover, by Lemma 2.3, $J(u) := \left(\frac{1}{2} - \frac{1}{2 \cdot 2_\mu^*}\right)^{-1} J_1(u)^{2_\mu^*-1}$, can be also considered as an Euler-Lagrange functional associated to problem (1.3), in the sense that the solutions of (1.3) correspond to critical points of J . Then, to prove our result, it is more convenient for us to work with,

$$J(u) = \frac{1}{\int_{\Omega} \int_{\Omega} \frac{|u(y)|^{2_\mu^*} |u(x)|^{2_\mu^*}}{|x-y|^\mu} dx dy}, \quad u \in \Sigma.$$

By Hardy-Littlewood-Sobolev inequality and (2.3), the functional J is lower bounded on Σ and we have

$$\inf_{u \in \Sigma} J(u) = S_{HL}(\Omega)^{2 \cdot 2_\mu^*}.$$

Denote $\tilde{S}_{HL} = S_{HL}(\Omega)^{2 \cdot 2_\mu^*}$. Using the result of [12], \tilde{S}_{HL} is independent of Ω and it is never achieved except $\Omega = \mathbb{R}^n$. In this case, the minimizers functions are unique and of the form $c\delta_{(a,\lambda)}$, where $\delta_{(a,\lambda)}$ is defined in (2.4) and $c > 0$.

Let $a \in \mathbb{R}^n$ and $\lambda > 0$, we define

$$U_{(a,\lambda)}(x) = \gamma_0 \delta_{(a,\lambda)}(x), \quad x \in \mathbb{R}^n,$$

where

$$\gamma_0 = \left(\int_{\mathbb{R}^n} |\nabla \delta_{(a,\lambda)}|^2 dx\right)^{-\frac{1}{2}} = \left(n(n-2) \int_{\mathbb{R}^n} \frac{dx}{(1+|x|^2)^n}\right)^{-\frac{1}{2}}. \quad (2.5)$$

Lemma 2.4 For $a \in \mathbb{R}^n$ and $\lambda > 0$, $U_{(a,\lambda)}$ satisfies

$$-\Delta U_{(a,\lambda)} = \tilde{S}_{HL} \left(\int_{\mathbb{R}^n} \frac{U_{(a,\lambda)}^{2_\mu^*}(y)}{|x-y|^\mu} dy\right) U_{(a,\lambda)}^{2_\mu^*-1} \quad \text{in } \mathbb{R}^n.$$

Proof. From [7] and [12], we know that $\delta_{(a,\lambda)}$ satisfies

$$-\Delta \delta_{(a,\lambda)} = A_{HL} \left(\int_{\mathbb{R}^n} \frac{\delta_{(a,\lambda)}^{2_\mu^*}(y)}{|x-y|^\mu} dy\right) \delta_{(a,\lambda)}^{2_\mu^*-1} \quad \text{in } \mathbb{R}^n,$$

where, $A_{HL} = (n(n-2))^{\frac{n-\mu+2}{2}} C_{n,\mu}^{-1} S^{\frac{\mu-n}{2}}$. Therefore,

$$-\Delta U_{(a,\lambda)} = \gamma_0^{2-2 \cdot 2_\mu^*} A_{HL} \left(\int_{\mathbb{R}^n} \frac{U_{(a,\lambda)}^{2_\mu^*}(y)}{|x-y|^\mu} dy\right) U_{(a,\lambda)}^{2_\mu^*-1}.$$

Multiplying by $U_{(a,\lambda)}$ and integrate, we get

$$1 = \gamma_0^{2-2 \cdot 2_\mu^*} A_{HL} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a,\lambda)}(y)^{2_\mu^*} U_{(a,\lambda)}^{2_\mu^*}(x)}{|x-y|^\mu} dx dy,$$

and hence

$$\gamma_0^{2-22^*_\mu} A_{HL} = \frac{1}{\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a,\lambda)}(y)^{2^*_\mu} U_{(a,\lambda)}(x)^{2^*_\mu}}{|x-y|^\mu} dx dy} = \tilde{S}_{HL}.$$

□

For $a \in \Omega$, we denote

$$G(a, x) = \frac{\gamma_0}{|x-a|^{n-2}} - H(a, x),$$

where γ_0 is defined in (2.5) and

$$\begin{cases} \Delta_x H(a, x) = 0 & \text{in } \Omega, \\ H(a, x) = \frac{\gamma_0}{|x-a|^{n-2}} & \text{on } \partial\Omega. \end{cases}$$

Our aim in the remainder part of this Section is to establish an asymptotic expansion of the functional J on a suitable finite dimensional set in terms of the Green function G and its regular part H .

Denote $PU_{(a,\lambda)}$, $a \in \Omega$ and $\lambda > 0$, the projection of $U_{(a,\lambda)}$ on $H_0^1(\Omega)$. Namely, $PU_{(a,\lambda)}$ is the unique solution of

$$\begin{cases} -\Delta PU_{(a,\lambda)} = -\Delta U_{(a,\lambda)} & \text{in } \Omega, \\ PU_{(a,\lambda)} = 0 & \text{on } \partial\Omega. \end{cases}$$

We claim that

$$PU_{(a,\lambda)} = U_{(a,\lambda)} - \frac{H(a, \cdot)}{\lambda^{\frac{n-2}{2}}} + O\left(\frac{1}{\lambda^{\frac{n+2}{2}} d(a, \partial\Omega)^n}\right). \quad (2.6)$$

Indeed, denote $\theta_{(a,\lambda)} = U_{(a,\lambda)} - PU_{(a,\lambda)} - \frac{H(a, \cdot)}{\lambda^{\frac{n-2}{2}}}$. We have

$$\begin{cases} \Delta \theta_{(a,\lambda)} = 0 & \text{in } \Omega, \\ \theta_{(a,\lambda)}(x) = U_{(a,\lambda)} - \frac{\gamma_0}{\lambda^{\frac{n-2}{2}} |x-a|^{n-2}} & \text{on } \partial\Omega. \end{cases}$$

On the boundary $\partial\Omega$, we have

$$\theta_{(a,\lambda)}(x) = \frac{-\gamma_0}{\lambda^{\frac{n-2}{2}} |x-a|^{n-2}} \left(1 - \left(\frac{\lambda^2 |x-a|^2}{1 + \lambda^2 |x-a|^2}\right)^{\frac{n-2}{2}}\right),$$

and hence

$$|\theta_{(a,\lambda)}(x)| \leq \frac{\gamma_0}{\lambda^{\frac{n+2}{2}} |x-a|^n}, \quad \forall x \in \partial\Omega.$$

Claim (2.6) follows from the maximum principle Theorem.

Let M be a compact set included in Ω . We then have the following expansion of J .

Proposition 2.1 *Let $p \geq 1$ and let $a_1, \dots, a_p \in M$, $\alpha_1, \dots, \alpha_p \geq 0$ such that $\sum_{i=1}^p \alpha_i = 1$ and $\lambda > 0$. Denote $d_a = \min_{i \neq j} d(a_i, a_j)$. If λd_a is large enough, we have*

$$\begin{aligned} J\left(\frac{\sum_{i=1}^p \alpha_i PU_{(a_i,\lambda)}}{\left|\sum_{i=1}^p \alpha_i PU_{(a_i,\lambda)}\right|_{1\Omega}}\right) &= \tilde{S}_{HL} \frac{(\sum_{i=1}^p \alpha_i^2)^{2^*_\mu}}{\sum_{i=1}^p \alpha_i^{2^*_\mu}} \left[1 - 2^*_\mu n(n-2) \frac{\gamma_0 c_1}{\lambda^{n-2}} \left(\sum_{i=1}^p \left(\frac{\alpha_i^2}{\sum_{j=1}^p \alpha_j^2}\right)\right.\right. \\ &\quad \left.\left. - 2 \frac{\alpha_i^{22^*_\mu}}{\sum_{j=1}^p \alpha_j^{22^*_\mu}}\right) H(a_i, a_i) + \sum_{k \neq i} \left(\frac{2\alpha_i^{22^*_\mu-1} \alpha_k}{\sum_{j=1}^p \alpha_j^{22^*_\mu}} - \frac{\alpha_i \alpha_k}{\sum_{j=1}^p \alpha_j^2}\right) G(a_i, a_k)\right] + O\left(\frac{1}{(\lambda d_a)^{n-1}}\right). \end{aligned}$$

Here γ_0 is defined in (2.5) and $c_1 = \int_{\mathbb{R}^n} \frac{dx}{(1+|x|^2)^{\frac{n+2}{2}}}$.

Proof. Denote $u = \sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}$.

$$J\left(\frac{u}{|u|_{1\Omega}}\right) = \frac{|u|_{1\Omega}^{2 \cdot 2^*}}{\int_{\Omega} \int_{\Omega} \frac{u(y)^{2^*} u(x)^{2^*}}{|x-y|^\mu} dx dy} := \frac{N}{D}.$$

We expand the numerator as follows

$$N^{\frac{1}{2^*}} = \sum_{i=1}^p \alpha_i^2 |PU_{(a_i, \lambda)}|_{1\Omega}^2 + \sum_{i \neq j} \alpha_i \alpha_j \langle PU_{(a_i, \lambda)}, PU_{(a_j, \lambda)} \rangle.$$

Using estimate (2.6), for any $i = 1, \dots, p$, we have

$$\begin{aligned} |PU_{(a_i, \lambda)}|_{1\Omega}^2 &= \int_{\Omega} -\Delta U_{(a_i, \lambda)} PU_{(a_i, \lambda)} dx \\ &= \int_{\Omega} -\Delta U_{(a_i, \lambda)} \left(U_{(a_i, \lambda)} - \frac{H(a_i, x)}{\lambda^{\frac{n-2}{2}}} \right) dx + O\left(\frac{1}{\lambda^{\frac{n+2}{2}}} \int_{\mathbb{R}^n} U_{(a_i, \lambda)}^{2^*-1} dx \right). \end{aligned} \quad (2.7)$$

Observe that,

$$\begin{aligned} \int_{\Omega} -\Delta U_{(a_i, \lambda)} U_{(a_i, \lambda)} dx &= \tilde{S}_{HL} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a_i, \lambda)}^{2^*}(y) U_{(a_i, \lambda)}^{2^*}(x)}{|x-y|^\mu} dx dy + O\left(\frac{1}{\lambda^n}\right) \\ &= 1 + O\left(\frac{1}{\lambda^n}\right). \end{aligned} \quad (2.8)$$

By expanding $H(a_i, x)$ around a_i we have

$$\begin{aligned} \int_{\Omega} -\Delta U_{(a_i, \lambda)} \frac{H(a_i, x)}{\lambda^{\frac{n-2}{2}}} dx &= \tilde{S}_{HL} \frac{H(a_i, a_i)}{\lambda^{\frac{n-2}{2}}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a_i, \lambda)}^{2^*}(y) U_{(a_i, \lambda)}^{2^*-1}(x)}{|x-y|^\mu} dx dy + O\left(\frac{1}{\lambda^n}\right), \\ &= \frac{n(n-2)}{\gamma_0^{\frac{4}{n-2}}} \frac{H(a_i, a_i)}{\lambda^{\frac{n-2}{2}}} \int_{\mathbb{R}^n} U_{(a_i, \lambda)}^{2^*-1} dx + O\left(\frac{1}{\lambda^n}\right), \end{aligned}$$

since, $-\Delta U_{(a_i, \lambda)} = \gamma_0^{\frac{-4}{n-2}} n(n-2) U_{(a_i, \lambda)}^{2^*-1}$ in \mathbb{R}^n and therefore

$$\int_{\mathbb{R}^n} \frac{U_{(a_i, \lambda)}^{2^*}(y)}{|x-y|^\mu} dy = \frac{n(n-2)}{\tilde{S}_{HL} \gamma_0^{\frac{4}{n-2}}} U_{(a_i, \lambda)}^{2^*-2^*}(x). \quad (2.9)$$

Using the fact that

$$\int_{\mathbb{R}^n} U_{(a_i, \lambda)}^{2^*-1} dx = \frac{\gamma_0^{2^*-1}}{\lambda^{\frac{n-2}{2}}} \int_{\mathbb{R}^n} \frac{dx}{(1+|x|^2)^{\frac{n+2}{2}}},$$

we get

$$\int_{\Omega} -\Delta U_{(a_i, \lambda)} \frac{H(a_i, x)}{\lambda^{\frac{n-2}{2}}} dx = \gamma_0 n(n-2) c_1 \frac{H(a_i, a_i)}{\lambda^{n-2}} + O\left(\frac{1}{\lambda^n}\right). \quad (2.10)$$

From (2.8) and (2.10), equality (2.7) reduces to

$$|PU_{(a_i, \lambda)}|_{1\Omega}^2 = 1 - \gamma_0 n(n-2) c_1 \frac{H(a_i, a_i)}{\lambda^{n-2}} + O\left(\frac{1}{\lambda^n}\right). \quad (2.11)$$

Now, for $i \neq j$, we have

$$\begin{aligned} \langle PU_{(a_i, \lambda)}, PU_{(a_j, \lambda)} \rangle &= \int_{\Omega} -\Delta U_{(a_i, \lambda)} PU_{(a_j, \lambda)} dx \\ &= \tilde{S}_{HL} \int_{\Omega} \int_{\Omega} \frac{U_{(a_i, \lambda)}^{2^*_{\mu}}(y) U_{(a_i, \lambda)}^{2^*_{\mu}-1}(x) PU_{(a_j, \lambda)}(x)}{|x-y|^{\mu}} dx dy \end{aligned}$$

Using estimates (2.6) and (2.9), we get

$$\begin{aligned} \langle PU_{(a_i, \lambda)}, PU_{(a_j, \lambda)} \rangle &= \frac{n(n-2)}{\gamma_0^{\frac{4}{n-2}}} \left(\int_{\Omega} U_{(a_i, \lambda)}^{2^*_{\mu}-1} U_{(a_j, \lambda)} dx - \frac{1}{\lambda^{\frac{n-2}{2}}} \int_{\Omega} U_{(a_i, \lambda)}^{2^*_{\mu}-1} H(a_j, x) dx \right) + O\left(\frac{1}{\lambda^n}\right). \\ &= \frac{n(n-2)}{\gamma_0^{\frac{4}{n-2}}} \left(\gamma_0^{2^*_{\mu}} \int_{\Omega} \delta_{(a_i, \lambda)}^{2^*_{\mu}-1} \delta_{(a_j, \lambda)} dx - \frac{\gamma_0^{2^*_{\mu}-1}}{\lambda^{\frac{n-2}{2}}} \int_{\Omega} \delta_{(a_i, \lambda)}^{2^*_{\mu}-1} H(a_j, x) dx \right) + O\left(\frac{1}{\lambda^n}\right). \end{aligned}$$

From [3], we have

$$\int_{\Omega} \delta_{(a_i, \lambda)}^{2^*_{\mu}-1} \delta_{(a_j, \lambda)} dx = \frac{c_1}{(\lambda |a_i - a_j|)^{n-2}} + o\left(\frac{1}{(\lambda d_a)^{n-2}}\right),$$

and by expanding $H(a_j, x)$ around a_i , we have

$$\int_{\Omega} \delta_{(a_i, \lambda)}^{2^*_{\mu}-1} H(a_j, x) dx = c_1 \frac{H(a_i, a_j)}{\lambda^{\frac{n-2}{2}}} + o\left(\frac{1}{\lambda^{\frac{n-2}{2}}}\right).$$

Therefore,

$$\begin{aligned} \langle PU_{(a_i, \lambda)}, PU_{(a_j, \lambda)} \rangle &= n(n-2) \frac{\gamma_0 c_1}{\lambda^{n-2}} \left(\frac{\gamma_0}{|a_i - a_j|^{n-2}} - H(a_i, a_j) \right) + o\left(\frac{1}{(\lambda d_a)^{n-2}}\right), \\ &= n(n-2) \gamma_0 c_1 \frac{G(a_i, a_j)}{\lambda^{n-2}} + o\left(\frac{1}{(\lambda d_a)^{n-2}}\right). \end{aligned} \quad (2.12)$$

Estimates (2.11) and (2.12) yield

$$\begin{aligned} N^{\frac{1}{2^*_{\mu}}} &= \left(\sum_{i=1}^p \alpha_i^2 \right) \left(1 + \left(\sum_{i=1}^p \alpha_i^2 \right)^{-1} n(n-2) \frac{\gamma_0 c_1}{\lambda^{n-2}} \left(\sum_{i \neq j} \alpha_i \alpha_j G(a_i, a_j) \right. \right. \\ &\quad \left. \left. - \sum_{i=1}^p \alpha_i^2 H(a_i, a_i) \right) \right) + o\left(\frac{1}{(\lambda d_a)^{n-2}}\right), \end{aligned}$$

and therefore

$$\begin{aligned} N &= \left(\sum_{i=1}^p \alpha_i^2 \right)^{2^*_{\mu}} \left(1 + 2^*_{\mu} \left(\sum_{i=1}^p \alpha_i^2 \right)^{-1} n(n-2) \frac{\gamma_0 c_1}{\lambda^{n-2}} \left(\sum_{i \neq j} \alpha_i \alpha_j G(a_i, a_j) \right. \right. \\ &\quad \left. \left. - \sum_{i=1}^p \alpha_i^2 H(a_i, a_i) \right) \right) + o\left(\frac{1}{(\lambda d_a)^{n-2}}\right). \end{aligned} \quad (2.13)$$

We now turn to expand the denominator of $J(\frac{u}{|u|_{\Omega}})$. Let ρ be a small positive constant and denote $*$ the convolution product. We have

$$\begin{aligned}
D &= \int_{\Omega} \int_{\Omega} \frac{(\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda_i)}(y))^{2^*} (\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda_i)}(x))^{2^*}}{|x-y|^{\mu}} dx dy, \\
&= \int_{\Omega} \left(\frac{1}{|x|^{\mu}} * \left(\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}(x) \right)^{2^*} \right) \left(\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}(x) \right)^{2^*} dx, \\
&= \int_{\cup_{i=1}^p B(a_i, \rho)} \left(\frac{1}{|x|^{\mu}} * \left(\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}(x) \right)^{2^*} \right) \left(\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}(x) \right)^{2^*} dx \\
&+ \int_{\Omega \setminus \cup_{i=1}^p B(a_i, \rho)} \left(\frac{1}{|x|^{\mu}} * \left(\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}(x) \right)^{2^*} \right) \left(\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}(x) \right)^{2^*} dx \quad (2.14)
\end{aligned}$$

On $B(a_i, \rho)$, $i = 1, \dots, p$, we write

$$\sum_{k=1}^p \alpha_k PU_{(a_k, \lambda)} = \alpha_i U_{(a_i, \lambda)} + \sum_{k \neq i} \alpha_k PU_{(a_k, \lambda)} + \alpha_i (PU_{(a_i, \lambda)} - U_{(a_i, \lambda)}).$$

Therefore,

$$\begin{aligned}
&\int_{B(a_i, \rho)} \left(\frac{1}{|x|^{\mu}} * \left(\sum_{k=1}^p \alpha_k PU_{(a_k, \lambda)} \right)^{2^*} \right) \left(\sum_{k=1}^p \alpha_k PU_{(a_k, \lambda)} \right)^{2^*} dx \\
&= \int_{B(a_i, \rho)} \left(\frac{1}{|x|^{\mu}} * \left(\sum_{k=1}^p \alpha_k PU_{(a_i, \lambda)} \right)^{2^*} \right) (\alpha_i U_{(a_i, \lambda)})^{2^*} dx \\
&+ 2^* \int_{B(a_i, \rho)} \left(\frac{1}{|x|^{\mu}} * \left(\sum_{k=1}^p \alpha_k PU_{(a_k, \lambda)} \right)^{2^*} \right) (\alpha_i U_{(a_i, \lambda)})^{2^* - 1} \\
&\quad \left(\sum_{k \neq i} \alpha_k PU_{(a_k, \lambda)} + \alpha_i (PU_{(a_i, \lambda)} - U_{(a_i, \lambda)}) \right) dx + o\left(\frac{1}{(\lambda d_a)^{n-2}}\right), \\
&= \int_{B(a_i, \rho)} \left(\frac{1}{|x|^{\mu}} * (\alpha_i U_{(a_i, \lambda_i)})^{2^*} \right) (\alpha_i U_{(a_i, \lambda_i)})^{2^*} dx \\
&+ 2^* \int_{B(a_i, \rho)} \left(\frac{1}{|x|^{\mu}} * (\alpha_i U_{(a_i, \lambda)})^{2^* - 1} \left(\sum_{k \neq i} \alpha_k PU_{(a_k, \lambda)} \right. \right. \\
&\quad \left. \left. + \alpha_i (PU_{(a_i, \lambda)} - U_{(a_i, \lambda)}) \right) \right) (\alpha_i U_{(a_i, \lambda)})^{2^*} dx \\
&+ 2^* \int_{B(a_i, \rho)} \left(\frac{1}{|x|^{\mu}} * (\alpha_i U_{(a_i, \lambda)})^{2^*} (\alpha_i U_{(a_i, \lambda_i)})^{2^* - 1} \left(\sum_{k \neq i} \alpha_k PU_{(a_k, \lambda)} \right. \right. \\
&\quad \left. \left. + \alpha_i (PU_{(a_i, \lambda)} - U_{(a_i, \lambda)}) \right) \right) + o\left(\frac{1}{(\lambda d_a)^{n-2}}\right), \\
&= \alpha_i^{2 \cdot 2^*} \int_{\mathbb{R}^n} \left(\frac{1}{|x|^{\mu}} * U_{(a_i, \lambda)}^{2^*} \right) U_{(a_i, \lambda)}^{2^*} dx
\end{aligned}$$

$$\begin{aligned}
& + 2_\mu^* \int_{\mathbb{R}^n} \left(\frac{1}{|x|^\mu} * (\alpha_i U_{(a_i, \lambda)})^{2_\mu^* - 1} \left(\sum_{k \neq i} \alpha_k P U_{(a_k, \lambda)} \right) \right) (\alpha_i U_{(a_i, \lambda)})^{2_\mu^*} dx \\
& + 2_\mu^* \int_{\mathbb{R}^n} \left(\frac{1}{|x|^\mu} * (\alpha_i U_{(a_i, \lambda)})^{2_\mu^* - 1} \right) \left(\alpha_i (P U_{(a_i, \lambda)} - U_{(a_i, \lambda)}) \right) (\alpha_i U_{(a_i, \lambda)})^{2_\mu^*} dx \\
& + 2_\mu^* \int_{\mathbb{R}^n} \left(\frac{1}{|x|^\mu} * (\alpha_i U_{(a_i, \lambda)})^{2_\mu^*} \right) (\alpha_i U_{(a_i, \lambda)})^{2_\mu^* - 1} \left(\sum_{k \neq i} \alpha_k P U_{(a_k, \lambda)} \right) dx \\
& + 2_\mu^* \int_{\mathbb{R}^n} \left(\frac{1}{|x|^\mu} * (\alpha_i U_{(a_i, \lambda)})^{2_\mu^*} \right) (\alpha_i U_{(a_i, \lambda)})^{2_\mu^* - 1} \left(\alpha_i (P U_{(a_i, \lambda)} - U_{(a_i, \lambda)}) \right) dx + o\left(\frac{1}{(\lambda d_a)^{n-2}} \right).
\end{aligned}$$

Using the estimate of (2.6), we obtain that

$$\begin{aligned}
& \int_{B(a_i, \rho)} \left(\frac{1}{|x|^\mu} * \left(\sum_{k=1}^p \alpha_k P U_{(a_k, \lambda)} \right)^{2_\mu^*} \right) \left(\sum_{k=1}^p \alpha_k P U_{(a_k, \lambda)} \right)^{2_\mu^*} dx \\
& = \alpha_i^{2 \cdot 2_\mu^*} \tilde{S}_{HL}^{-1} + 2_\mu^* \sum_{k \neq i} \alpha_i^{2 \cdot 2_\mu^* - 1} \alpha_k \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a_i, \lambda)}^{2_\mu^* - 1}(y) P U_{(a_k, \lambda)}(y) U_{(a_i, \lambda)}^{2_\mu^*}(x)}{|x - y|^\mu} dx dy \\
& - 2_\mu^* \frac{\alpha_i^{2 \cdot 2_\mu^*}}{\lambda^{\frac{n-2}{2}}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a_i, \lambda)}^{2_\mu^* - 1}(y) H(a_i, y) U_{(a_i, \lambda)}^{2_\mu^*}(x)}{|x - y|^\mu} dx dy \\
& + 2_\mu^* \sum_{k \neq i} \alpha_i^{2 \cdot 2_\mu^* - 1} \alpha_k \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a_i, \lambda)}^{2_\mu^*}(y) U_{(a_i, \lambda)}^{2_\mu^* - 1}(x) P U_{(a_k, \lambda)}(x)}{|x - y|^\mu} dx dy \\
& - 2_\mu^* \frac{\alpha_i^{2 \cdot 2_\mu^*}}{\lambda^{\frac{n-2}{2}}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a_i, \lambda)}^{2_\mu^*}(y) U_{(a_i, \lambda)}^{2_\mu^* - 1}(x) H(a_i, x)}{|x - y|^\mu} dx dy + o\left(\frac{1}{(\lambda d_a)^{n-2}} \right), \\
& = \alpha_i^{2 \cdot 2_\mu^*} \tilde{S}_{HL}^{-1} + 2 \cdot 2_\mu^* \tilde{S}_{HL}^{-1} \sum_{k \neq i} \alpha_i^{2 \cdot 2_\mu^* - 1} \alpha_k \langle P U_{(a_i, \lambda)}, P U_{(a_k, \lambda)} \rangle \\
& - 2 \cdot 2_\mu^* \frac{\alpha_i^{2 \cdot 2_\mu^*}}{\lambda^{\frac{n-2}{2}}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a_i, \lambda)}^{2_\mu^*}(y) U_{(a_i, \lambda)}^{2_\mu^* - 1}(x) H(a_i, x)}{|x - y|^\mu} dx dy + o\left(\frac{1}{(\lambda d_a)^{n-2}} \right).
\end{aligned}$$

Observe that, by estimate (2.9), we have

$$\begin{aligned}
\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{U_{(a_i, \lambda)}^{2_\mu^*}(y) U_{(a_i, \lambda)}^{2_\mu^* - 1}(x) H(a_i, x)}{|x - y|^\mu} dx dy & = \tilde{S}_{HL}^{-1} \frac{n(n-2)}{\gamma_0^{\frac{4}{n-2}}} \int_{\mathbb{R}^n} U_{(a_i, \lambda)}^{2_\mu^* - 1}(x) H(a_i, x) dx \\
& = \tilde{S}_{HL}^{-1} n(n-2) c_1 \gamma_0 \frac{H(a_i, a_i)}{\lambda^{\frac{n-2}{2}}} + o\left(\frac{1}{\lambda^{\frac{n-2}{2}}} \right).
\end{aligned}$$

The above estimate with (2.12) yield

$$\begin{aligned}
& \int_{B(a_i, \rho)} \left(\frac{1}{|x|^\mu} * \left(\sum_{k=1}^p \alpha_k P U_{(a_k, \lambda)} \right)^{2_\mu^*} \right) \left(\sum_{k=1}^p \alpha_k P U_{(a_k, \lambda)} \right)^{2_\mu^*} dx \tag{2.15} \\
& = \tilde{S}_{HL}^{-1} \alpha_i^{2 \cdot 2_\mu^* - 2} \left[\alpha_i^2 + 2 \cdot 2_\mu^* n(n-2) \frac{c_1 \gamma_0}{\lambda^{n-2}} \left(\sum_{k \neq i} \alpha_i \alpha_k G(a_i, a_k) - \alpha_i^2 H(a_i, a_i) \right) \right] + o\left(\frac{1}{(\lambda d_a)^{n-2}} \right).
\end{aligned}$$

The remainder integral of (2.14) satisfies

$$\int_{\Omega \setminus \bigcup_{i=1}^p B(a_i, \rho)} \left(\frac{1}{|x|^\mu} * \left(\sum_{k=1}^p \alpha_i P U_{(a_i, \lambda)}(x) \right)^{2_\mu^*} \right) \left(\sum_{k=1}^p \alpha_i P U_{(a_i, \lambda)}(x) \right)^{2_\mu^*} dx = O\left(\frac{1}{(\lambda d_a)^n} \right). \tag{2.16}$$

Using (2.15) and (2.16), the expansion of (2.14) reduces to

$$\begin{aligned}
D &= \tilde{S}_{HL}^{-1} \left(\sum_{i=1}^p \alpha_i^{2 \cdot 2^*} \right) \left[1 + 2 \cdot 2^* n(n-2) \frac{c_1 \gamma_0}{\lambda^{n-2}} \left(\sum_{k \neq i} \frac{\alpha_i^{2 \cdot 2^* - 1} \alpha_k}{\sum_{j=1}^p \alpha_j^{2 \cdot 2^*}} G(a_i, a_k) \right. \right. \\
&\quad \left. \left. - \sum_{i=1}^p \frac{\alpha_i^{2 \cdot 2^*}}{\sum_{j=1}^p \alpha_j^{2 \cdot 2^*}} H(a_i, a_i) \right) \right] + o\left(\frac{1}{(\lambda d_a)^{n-2}}\right). \tag{2.17}
\end{aligned}$$

The expansion of Proposition 2.1 follows from (2.13) and (2.17). \square

Of course the above expansion is useful when $d_a = \min_{i \neq j} |a_i - a_j|$ is lower bounded by a positive constant independent of a . In this case, for any $\varepsilon > 0$, there exists $\bar{\lambda} = \bar{\lambda}(\varepsilon) > 0$ (independent of a , as long as a lies in a compact set of Ω) such that, for any $\lambda \geq \bar{\lambda}$,

$$J\left(\frac{\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}}{\left|\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}\right|_{1\Omega}}\right) \leq (p + \varepsilon)^{2^* - 1} \tilde{S}_{HL}, \tag{2.18}$$

since $\frac{(\sum_{i=1}^p \alpha_i^2)^{2^*}}{\sum_{i=1}^p \alpha_i^{2 \cdot 2^*}} \leq p^{2^* - 1}$.

If d_a is close to zero, or if a parameter α_i is close to zero for some index i , $i = 1, \dots, p$, we have

$$J\left(\frac{\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}}{\left|\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}\right|_{1\Omega}}\right) \leq p^{2^* - 1} \tilde{S}_{HL}. \tag{2.19}$$

This can be proved as the analogous assertion in [2].

3 Proof of the existence Theorem

To prove Theorem 1.1 we argue by contradiction. Therefore we suppose in what follows that problem (1.3) has no positive solution.

Let $-\partial J: \Sigma \rightarrow T\Sigma$ be the gradient of J . It is a continuous vector field, locally Lipschitz and bounded. For $u \in \Sigma$, we denote by $s \mapsto \eta(s, u)$, $s \geq 0$, the unique solution of the differential equation

$$\begin{cases} \dot{\eta} &= -\partial J(\eta) \\ \eta(0) &= u. \end{cases}$$

It is straightforward to see that $\eta(s, u)$ is defined for any $s \in [0, \infty)$, $J(\eta(s, u))$ is a decreasing function and $\partial J(\eta(s, u)) \rightarrow 0$, as $s \rightarrow \infty$. Moreover, if we denote by $\Sigma^+ = \{u \in \Sigma, u \geq 0\}$, then Σ^+ is invariant under the action of $(-\partial J)$, in the sense that, if $u \in \Sigma^+$ then $\eta(s, u) \in \Sigma^+$ for any $s \geq 0$. For any $p \in \mathbb{N} \setminus \{0\}$ and $\varepsilon > 0$, we define

$$\begin{aligned}
V(p, \varepsilon) &= \left\{ u \in \Sigma^+, \exists a_1, \dots, a_p \in \Omega, \exists \lambda_1, \dots, \lambda_p > 0 \text{ and } \alpha_1, \dots, \alpha_p > 0, \text{ s.t.}, \right. \\
&\quad \left| u - \sum_{i=1}^p \alpha_i PU_{(a_i, \lambda_i)} \right|_{1\Omega} < \varepsilon \text{ with } \lambda_i > \varepsilon^{-1}, \lambda_i d(a_i, \partial\Omega) > \varepsilon^{-1}, \forall i = 1, \dots, p \\
&\quad \text{and } \varepsilon_{ij} = \left(\frac{\lambda_i}{\lambda_j} + \frac{\lambda_j}{\lambda_i} + \lambda_i |a_i - a_j|^2 \right)^{\frac{2-n}{2}} < \varepsilon, \forall i \neq j \left. \right\}.
\end{aligned}$$

We have the following result

Proposition 3.1 *Assume that J has no positive critical point. Let $u \in \Sigma^+$. There exists a unique positive integer p which depends on u only, such that for any $\varepsilon > 0$ there exists $s_\varepsilon > 0$ such that for any $s > s_\varepsilon$, $\eta(s, u) \in V(p, \varepsilon)$. Moreover $J(\eta(s, u)) \rightarrow \ell$, as $s \rightarrow \infty$, with $\ell \leq p^{2^*_\mu - 1} \tilde{S}_{HL}$.*

Proof. Let $(s_k)_k$ be a positive sequence tending to ∞ . Denote $u_k = \eta(s_k, u)$, $u \in \Sigma^+$. Therefore, $(u_k)_k$ is a sequence of Σ^+ which satisfies

$$J(u_k) \rightarrow c \quad \text{and} \quad \partial J(u_k) \rightarrow 0.$$

Since we have supposed that J has no critical point in Σ^+ , by ([1], Proposition 1) and ([13], Lemma 4.5), Here exists $p \in \mathbb{N} \setminus \{0\}$ such that $u_k \in V(p, \varepsilon_k)$, $\forall k$, where $\varepsilon_k > 0$ and $\varepsilon_k \rightarrow 0$ as $k \rightarrow \infty$. Fix $\varepsilon > 0$. If the flow line $\eta(s, u)$ enter $V(p, \frac{\varepsilon}{2})$ at a time s_k and leave $V(p, \varepsilon)$ at a later time t_k , for $t \in [s_k, t_k]$, $\eta(s, u) \in V(p, \varepsilon) \setminus V(p, \frac{\varepsilon}{2})$. Therefore,

$$J(\eta(t_k, u)) - J(\eta(s_k, u)) = \int_{s_k}^{t_k} \widehat{J(\dot{\eta}(s, u))} ds \leq - \int_{s_k}^{t_k} |\partial J(\eta(s, u))|_{1\Omega}^2 ds \leq -\gamma(t_k - s_k), \quad (3.1)$$

where $\gamma = \gamma(\varepsilon)$ is the lower bound of $|\partial J(u)|_{1\Omega}$ on $V(p, \varepsilon) \setminus V(p, \frac{\varepsilon}{2})$. By ([1], Proposition 1) and ([13], Lemma 4.5), γ is positive. Now, let

$$0 < \beta = d(V(p, \varepsilon), V(p, \varepsilon)^c).$$

We have

$$\beta \leq d(\eta(s_k, u), \eta(t_k, u)) \leq \int_{s_k}^{t_k} |\dot{\eta}(s, u)|_{1\Omega} ds \leq M_0(t_k - s_k),$$

where $M_0 = \sup_{u \in V(p, \varepsilon)} |\partial J(u)|$ and hence $t_k - s_k \geq \frac{\beta}{M_0}$. This with (3.1) yield

$$J(\eta(t_k, u)) - J(\eta(s_k, u)) \leq -\frac{\gamma\beta}{M_0} \quad (3.2)$$

It follows from (3.2) that there is a finite number of such intervals $[s_k, t_k]$, since J is lower bounded. As a consequence, there exists $s_\varepsilon > 0$ such that $\eta(s, u) \in V(p, \varepsilon)$, $\forall s \geq s_\varepsilon$. The limit of $J(\eta(s, u))$ follows from an expansion like the one of Proposition 2.1. \square

In the following we denote by J_c , $c \in \mathbb{R}$, the set

$$J_c = \{u \in \Sigma^+, J(u) \leq c\}.$$

Let $p \in \mathbb{N} \setminus \{0\}$ and let ε_p be a constant such that

$$0 < \varepsilon_p < \tilde{S}_{HL}((p+1)^{2^*_\mu - 1} - p^{2^*_\mu - 1}).$$

Denote,

$$W_u(V(p, \varepsilon_p)) = \left\{ \eta(s, u), s \geq 0, u \in \bigcup_{1 \leq k \leq p} V(k, \varepsilon_k) \right\}.$$

We define,

$$E_p = J_{p^{2^*_\mu - 1} \tilde{S}_{HL} + \varepsilon_p} \cap W_u(V(p, \varepsilon_p)) \quad \text{and} \quad F_p = J_{p^{2^*_\mu - 1} \tilde{S}_{HL}} \cap W_u(V(p, \varepsilon_p)).$$

In this way,

$$F_{p-1} \subset E_{p-1} \subset F_p \subset E_p, \quad \forall p \geq 1,$$

with $F_0 = E_0 = \emptyset$.

For a pair of topological (X, Y) , $Y \subset X$, we denote $H_\ell(X, Y)$, $\ell \geq 0$, the relative homology of (X, Y) . We then have

Proposition 3.2 *Let $p \in \mathbb{N} \setminus \{0\}$. For any $\varepsilon \in (0, \varepsilon_p)$, there exists a continuous mapping*

$$R: (E_p, F_p) \rightarrow \left(J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap V(p, \varepsilon_p), J_{p^{2^*\mu-1}\tilde{S}_{HL}} \cap V(p, \varepsilon_p) \right),$$

which induces a sequence of isomorphisms

$$R_\ell: H_\ell(E_p, F_p) \rightarrow H_\ell \left(J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap V(p, \varepsilon_p), J_{p^{2^*\mu-1}\tilde{S}_{HL}} \cap V(p, \varepsilon_p) \right), \quad \ell \geq 0.$$

Proof. Let $0 < \varepsilon < \varepsilon_p$. Using the gradient flow $\eta(\cdot, \cdot)$ of the field $(-\partial J)$, the following retract by deformation holds,

$$J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon_p}} \simeq J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}},$$

since by Proposition 4.1, J has no critical value at infinity between the levels $p^{2^*\mu-1}\tilde{S}_{HL} + \varepsilon_p$ and $p^{2^*\mu-1}\tilde{S}_{HL} + \varepsilon$. Here \simeq denotes retract by deformation. It follows that

$$(E_p, F_p) \simeq \left(J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap W_u(V(p, \varepsilon_p)), F_p \right).$$

Let us denote by

$$r: (E_p, F_p) \rightarrow \left(J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap W_u(V(p, \varepsilon_p)), F_p \right)$$

the corresponding retract by deformation. Using the fact that

$$J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap W_u(V(p, \varepsilon_p)) \setminus F_p \subset V(p, \varepsilon_p),$$

we obtain that

$$J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap W_u(V(p, \varepsilon_p)) \setminus F_p = J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap V(p, \varepsilon_p) \setminus J_{p^{2^*\mu-1}\tilde{S}_{HL}} \cap V(p, \varepsilon_p),$$

and hence the following two pairs

$$\left(J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap W_u(V(p, \varepsilon_p)), F_p \right) \quad \text{and} \quad \left(J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap V(p, \varepsilon_p), J_{p^{2^*\mu-1}\tilde{S}_{HL}} \cap V(p, \varepsilon_p) \right),$$

are homotopically equivalent. Denote by

$$\hat{r}: \left(J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap W_u(V(p, \varepsilon_p)), F_p \right) \rightarrow \left(J_{p^{2^*\mu-1}\tilde{S}_{HL+\varepsilon}} \cap V(p, \varepsilon_p), J_{p^{2^*\mu-1}\tilde{S}_{HL}} \cap V(p, \varepsilon_p) \right),$$

the corresponding homotopy equivalence. Setting

$$R = \hat{r} \circ r.$$

R is an homotopy equivalence, since r and \hat{r} are. Hence the result follows. \square

We now introduce the following notations. For $p \in \mathbb{N} \setminus \{0\}$, we denote by

$$\Delta_{p-1} = \left\{ (\alpha_1, \dots, \alpha_p) \in [0, 1]^p, \sum_{i=1}^p \alpha_i = 1 \right\}$$

the standard p -simplex in \mathbb{R}^p . Let M be a fixed compact k -dimensional manifold in Ω , $1 \leq k \leq n-1$. We denote

$$B_p(M) = \left\{ \sum_{i=1}^p \alpha_i \delta_{a_i}, \alpha_i \in [0, 1], a_i \in M, \forall i = 1, \dots, p, \text{ and } \sum_{i=1}^p \alpha_i = 1 \right\},$$

where δ_{a_i} denotes the Dirac mass at a_i . We agree that $B_0(M) = \emptyset$. Lastly, denote ϱ_p the symmetric group of order p . We then have.

Proposition 3.3 For any $P \in \mathbb{N} \setminus \{0\}$, the homology $H_\ell(E_p, F_p)$, $\ell \geq 0$, of the pair (E_p, F_p) , is a module on the cohomology $H^\ell(\Omega^p/\varrho_p)$, $\ell \geq 0$ of Ω^p/ϱ_p . Moreover there exists a continuous mapping of topological pairs

$$\phi_p: (B_p(M), B_{p-1}(M)) \rightarrow (E_p, F_p),$$

such that the homology homomorphism

$$(\phi_p)_\ell: H_\ell(B_p(M), B_{p-1}(M)) \rightarrow (E_p, F_p), \quad \ell \geq 0,$$

is $H^\ell(\Omega^p/\varrho_p)$ -linear.

Proof. Let $p \in \mathbb{N} \setminus \{0\}$. For ε_p small enough and for $u \in V(p, \varepsilon_p)$, we optimize the approximation of u with respect to $\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda_i)}$. Namely, we set the following minimization problem

$$\min_{\alpha_i, a_i, \lambda_i} \left| u - \sum_{i=1}^p \alpha_i PU_{(a_i, \lambda_i)} \right|_{1\Omega}.$$

It is proved in [3] that this problem has a unique solution α_i, a_i, λ_i , $i = 1, \dots, p$ (modulo a permutation). It follows that there exists a continuous mapping

$$\chi: V(p, \varepsilon_p) \rightarrow \Omega^p/\varrho_p, \quad (3.3)$$

such that to any $u \in V(p, \varepsilon_p)$, we associate (a_1, \dots, a_p) ; the unique solution of the above problem of minimization.

Let $0 < \varepsilon < \varepsilon_p$ and χ_1 be the restriction of χ on $J_{p^{2^*}-1} \tilde{S}_{HL+\varepsilon} \cap V(p, \varepsilon_p)$. χ_1 is continuous and induces a cohomology homomorphism

$$(\chi_1)^\ell: H^\ell(\Omega^p/\varrho_p) \rightarrow H^\ell(J_{p^{2^*}-1} \tilde{S}_{HL+\varepsilon} \cap V(p, \varepsilon_p)), \quad \ell \geq 0.$$

From [4], we know that for any pair of topological spaces (X, Y) , the group of cohomology $H^\ell(X)$, $\ell \geq 0$, of X acts through the cap-product on the relative homology group $H_\ell(X, Y)$ of the pair (X, Y) and the action is linear. We apply this in our statement. We derive that

$$H_\ell \left(J_{p^{2^*}-1} \tilde{S}_{HL+\varepsilon} \cap V(p, \varepsilon_p), J_{p^{2^*}-1} \tilde{S}_{HL} \cap V(p, \varepsilon_p) \right), \quad \ell \geq 0,$$

is a module over $H_\ell \left(J_{p^{2^*}-1} \tilde{S}_{HL+\varepsilon} \cap V(p, \varepsilon_p) \right)$ and hence over $H^\ell(\Omega^p/\varrho_p)$ by $(\chi_1)^\ell$. Using the isomorphism R_ℓ of Proposition 3.2, we deduce that $H_\ell(E_p, F_p)$ is a module over $H^\ell(\Omega^p/\varrho_p)$. To construct the homology homomorphism $(\phi_p)_\ell$, $\ell \geq 0$, we first define on $M^p \times \Delta_{p-1}$ an equivalence relation such that the class of an element $(a_1, \dots, a_p, \alpha_1, \dots, \alpha_p) \in M^p \times \Delta_{p-1}$ equals to

$$\{(a_{\sigma(1)}, \dots, a_{\sigma(p)}, \alpha_{\sigma(1)}, \dots, \alpha_{\sigma(p)}), \sigma \in \varrho_p\}.$$

Denote $M^p \times_{\varrho_p} \Delta_{p-1}$, the related quotient space. We define

$$\begin{aligned} \pi_p: B_p(M) &\rightarrow M^p \times_{\varrho_p} \Delta_{p-1} \\ \sum_{i=1}^p \alpha_i \delta_i &\longmapsto (a_1, \dots, a_p, \alpha_1, \dots, \alpha_p). \end{aligned}$$

It is easy to verify that π_p is an homeomorphism. Let

$$S_p(M) = \{(a_1, \dots, a_p) \in M^p, \exists i \neq j, s.t., a_i = a_j\}.$$

π_p induces an homeomorphism of pairs that we denote again π_p ,

$$\pi_p: (B_p(M), B_{p-1}(M)) \rightarrow (M^p \times_{\varrho_p} \Delta_{p-1}, S_p(M) \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} \partial \Delta_{p-1}),$$

where $\partial \Delta_{p-1}$ denote the boundary of Δ_{p-1} . Let

$$(\pi_p)_\ell: H_\ell(B_p(M), B_{p-1}(M)) \rightarrow H_\ell(M^p \times_{\varrho_p} \Delta_{p-1}, S_p(M) \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} \partial \Delta_{p-1}), \ell \geq 0$$

be the homology isomorphism induced by π_p . Denote T_p a small ϱ_p -equivariant tubular open neighborhood of $S_p(M)$ in M^p such that $S_p(M)$ is a retract by deformation of T_p (see[4]) and denote

$$i_p: (M^p \times_{\varrho_p} \Delta_{p-1}, S_p(M) \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} \partial \Delta_{p-1}) \rightarrow (M^p \times_{\varrho_p} \Delta_{p-1}, \overline{T_p} \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} \partial \Delta_{p-1}),$$

the natural injection. It is an homotopy equivalence, since $\overline{T_p}$ retracts by deformation on $S_p(M)$. Therefore induces an homology isomorphism

$$(i_p)_\ell: H_\ell(M^p \times_{\varrho_p} \Delta_{p-1}, S_p(M) \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} \partial \Delta_{p-1}) \rightarrow H_\ell(M^p \times_{\varrho_p} \Delta_{p-1}, \overline{T_p} \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} \partial \Delta_{p-1}).$$

Let θ be a given small positive constant, We denote

$$\Delta_{p-1}^\theta = \left\{ (\alpha_i, \dots, \alpha_p) \in \Delta_{p-1}, \frac{\alpha_i}{\alpha_j} \in [1 - \theta, 1 + \theta], \forall i \neq j \right\}.$$

Using the fact that $(\Delta_{p-1}^\theta)^c$ retracts on $\partial \Delta_{p-1}$, there exists an homology isomorphism

$$(j_p)_\ell: H_\ell(M^p \times_{\varrho_p} \Delta_{p-1}, \overline{T_p} \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} \partial \Delta_{p-1}) \rightarrow H_\ell(M^p \times_{\varrho_p} \Delta_{p-1}, \overline{T_p} \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} (\Delta_{p-1}^\theta)^c),$$

$\ell \geq 0$. Let us denote

$$M_0^p = M^p \setminus T_p \tag{3.4}$$

By excision [6] of $T_p \times_{\varrho_p} \Delta_{p-1} \cup T_p \times_{\varrho_p} (\Delta_{p-1}^\theta)^c$, the two pairs

$$(M^p \times_{\varrho_p} \Delta_{p-1}, \overline{T_p} \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} (\partial \Delta_{p-1})^c) \text{ and } (M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial M_0^p \times_{\varrho_p} \Delta_{p-1} \cup M_0^p \times_{\varrho_p} (\Delta_{p-1}^\theta)^c)$$

have the same type of homotopy. Therefore

there exists a mapping

$$h_p: (M^p \times_{\varrho_p} \Delta_{p-1}, \overline{T_p} \times_{\varrho_p} \Delta_{p-1} \cup M^p \times_{\varrho_p} (\Delta_{p-1}^\theta)^c) \rightarrow (M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial M_0^p \times_{\varrho_p} \Delta_{p-1} \cup M_0^p \times_{\varrho_p} (\Delta_{p-1}^\theta)^c)$$

which defines an homotopy equivalence. Therefore, the corresponding homology homomorphism $(h_p)_\ell$, $\ell \geq 0$, is indeed an isomorphism.

Let for $\lambda > 0$, $f_{p,\lambda}$ denotes

$$f_{p,\lambda}: M^p \times_{\varrho_p} \Delta_{p-1} \longrightarrow \Sigma^+ \\ (a_1, \dots, a_p, \alpha_i, \dots, \alpha_p) \longmapsto \frac{\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}}{|\sum_{i=1}^p \alpha_i PU_{(a_i, \lambda)}|_{1\Omega}}.$$

Using the fact $d(a_i, a_j)$, $1 \leq i \neq j \leq p$ lower bounded by $d_1 > 0$, for every $(a_1, \dots, a_p) \in M_0^p$, where d_1 is the diameter of T_p , we can select $\bar{\lambda} = \bar{\lambda}(p)$ large enough, so that for any $\lambda \geq \bar{\lambda}$,

$$f_{p,\lambda}(M_0^p \times_{\varrho_p} \Delta_{p-1}) \subset \bigcup_{1 \leq k \leq p} V(k, \varepsilon_k) \subset W_u(V(p, \varepsilon_p)), \quad (3.5)$$

and

$$f_\lambda(\partial M_0^p \times_{\varrho_p} \Delta_{p-1} \cup_{\varrho_p} M_0^p \times (\Delta_{p-1}^\theta)^c) \subset \bigcup_{1 \leq k \leq p} V(k, \varepsilon_k) \subset W_u(V(p, \varepsilon_p)), \quad (3.6)$$

Moreover, by estimate (2.18), we have

$$f_{p,\lambda}(M_0^p \times \Delta_{p-1}) \subset J_{p^{2\mu^*-1}} \tilde{S}_{HL+\varepsilon_p}, \quad (3.7)$$

and by estimate (2.19), we have

$$f_{p,\lambda}(\partial M_0^p \times \Delta_{p-1} \cup_{\varrho_p} M_0^p \times (\partial \Delta_{p-1}^\theta)^c) \subset J_{p^{2\mu^*-1}} \tilde{S}_{HL}, \quad (3.8)$$

provided θ and d_1 are small. It follows from (3.5)–(3.8) that $f_{p,\lambda}$ maps the pair

$$(M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial M_0^p \times_{\varrho_p} \Delta_{p-1} \cup_{\varrho_p} M_0^p \times (\Delta_{p-1}^\theta)^c)$$

into (E_p, F_p) , we denote again

$$f_{p,\lambda}: (M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial M_0^p \times_{\varrho_p} \Delta_{p-1} \cup_{\varrho_p} M_0^p \times (\Delta_{p-1}^\theta)^c) \rightarrow (E_p, F_p)$$

the corresponding mapping. Passing to homology, we obtain the following homomorphism

$$(f_{p,\lambda})_\ell: H_\ell(M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial M_0^p \times_{\varrho_p} \Delta_{p-1} \cup_{\varrho_p} M_0^p \times (\Delta_{p-1}^\theta)^c) \rightarrow (E_p, F_p), \quad \ell \geq 0.$$

We now define the required mapping ϕ_p by:

$$\phi_p = f_{p,\lambda} \circ h_p \circ j_p \circ i_p \circ \pi_p.$$

By construction ϕ_p is continuous and hence induces an homology homomorphism $(\phi_p)_\ell$, $\ell \geq 0$, from $H_\ell(B_p(M), B_{p-1}(M))$ into $H_\ell(E_p, F_p)$.

Let us observe that $f_{p,\lambda}$ maps for $\theta > 0$ small enough and $\lambda > \bar{\lambda}$, the sets

$$M_0^p \times_{\varrho_p} \Delta_{p-1} \quad \text{into} \quad E_p \cap V(p, \varepsilon_p), \quad (3.9)$$

and

$$\partial(M_0^p \times_{\varrho_p} \Delta_{p-1}) = \partial M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta \cup_{\varrho_p} M_0^p \times \partial \Delta_{p-1}^\theta \quad \text{into} \quad F_p \cap V(p, \varepsilon_p). \quad (3.10)$$

Moreover, if we consider $M^p \times_{\varrho_p} \Delta_{p-1}$ as $B_p(M)$, through the homeomorphism π_p , we then have

$$(B_p(M), (M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta)^c) \simeq (B_p(M), B_{p-1}(M)),$$

since $(M_0^p)^c = \overline{T_p}$ retracts by deformation on $S_p(M)$ and $(\Delta_{p-1}^\theta)^c$ retracts on $\partial \Delta_{p-1}$. By excision

of $(M_0^p \times_{\varrho} \Delta_{p-1}^\theta)^c$, the two pairs $(M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta, \partial(M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta))$ and $(B_p(M), B_{p-1}(M))$ have the same type of homology. Let

$$q_p: (M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta, \partial(M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta)) \rightarrow (B_p(M), B_{p-1}(M))$$

be the associated homotopy equivalence.

Consider now the diagram of continuous maps,

$$\begin{array}{ccc} (B_p(M), B_{p-1}(M)) & \xrightarrow{\phi_p} & (E_p, F_p) \\ \uparrow q_p & & \uparrow \tilde{i}_p \\ (M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta, \partial(M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta)) & \xrightarrow{f_{p,\lambda}} & (E_p \cap V(p, \varepsilon_p), F_p \cap V(p, \varepsilon_p)) \\ \downarrow \hat{i}_p & & \downarrow \chi \\ M^p \times_{\varrho_p} \Delta_{p-1} & \xrightarrow{V_p} & \Omega^p / \varrho_p \end{array}$$

where V_p denotes the first projection, \tilde{i}_p and \hat{i}_p are the natural injection, χ is defined in (3.3), $f_{p,\lambda}$ is defined in (3.9) and (3.10) and q_p and ϕ_p are defined in the preceding steps.

Passing the above diagram to homology. Using the fact that $(\tilde{i}_p)_\ell$, $\ell \geq 0$, is an isomorphism and the homology isomorphism $(q_p)_\ell$ can be expressed by

$$(q_p)_\ell = (i)_\ell o (h_p)_\ell^{-1} o (j_p)_\ell^{-1} o (i_p)_\ell^{-1} o (\pi_p)_\ell^{-1}, \quad \ell \geq 0, \quad (3.11)$$

where

$$i: (M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta, \partial(M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta)) \rightarrow (M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial M_0^p \times_{\varrho_p} \Delta_{p-1} \cup M_0^p \times_{\varrho_p} (\Delta_{p-1}^\theta)^c),$$

denotes the natural injection, the obtained homological diagram is commutative and hence the homology homomorphism $(\phi_p)_\ell$, $\ell \geq 0$, is $H^\ell(\Omega^p / \varrho_p)$ -linear, in the sense that for any $\varphi^\ell \in H^\ell(\Omega^p / \varrho_p)$, and $[N_p, N_{p-1}]_\ell \in H_\ell(B_p(M), B_{p-1}(M))$, it holds

$$(\phi_p)_\ell(\varphi^\ell \cdot [N_p, N_{p-1}]_\ell) = \varphi^\ell \cdot (\phi_p)_\ell([N_p, N_{p-1}]_\ell). \quad (3.12)$$

Note that, it is proved that $H_\ell(E_p, F_p)$ is a module over $H^\ell(\Omega^p / \varrho_p)$ and by the cap-product, $H_\ell(B_p(M), B_{p-1}(M))$ is a module over $H^\ell(B_p(M))$ and thus over $H^\ell(\Omega^p / \varrho_p)$ via the first projection. the proof of the Proposition is thereby completed. \square

Let us observe that from (3.11), $(q_p)_\ell$ defines an isomorphism between

$$H_\ell(B_p(M), B_{p-1}(M)) \quad \text{and} \quad H_\ell(M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta, \partial(M_0^p \times_{\varrho_p} \Delta_{p-1}^\theta)), \quad \ell \geq 0.$$

Here M_0^p is defined in (3.4). We also note that $M_0^p \times_{\varrho_p} \Delta_{p-1}$ is a manifold of dimension $kp + p - 1$, ($k = \dim M$). It can be viewed as a singular $(k_p + p - 1)$ -chain with \mathbb{Z}_2 coefficients. Therefore the pair $(M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial(M_0^p \times_{\varrho_p} \Delta_{p-1}))$ defines a $(k_p + p - 1)$ -cycle in its self and thus gives arise to a non zero class denoted $[M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial(M_0^p \times_{\varrho_p} \Delta_{p-1})]$ in the relative homology of the pair $(M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial(M_0^p \times_{\varrho_p} \Delta_{p-1}))$. In the next, we denote

$$[B_p(M), B_{p-1}(M)] = (q_p)_\ell([M_0^p \times_{\varrho_p} \Delta_{p-1}, \partial(M_0^p \times_{\varrho_p} \Delta_{p-1})]). \quad (3.13)$$

Since $(q_p)_\ell$ is an isomorphism, then $[B_p(M), B_{p-1}(M)]$ defines a non zero class in the relative homology group $H_\ell(B_p(M), B_{p-1}(M))$ of the pair $(B_p(M), B_{p-1}(M))$, $\forall p \in \mathbb{N} \setminus \{0\}$. We have the following result.

Proposition 3.4 *Let ϕ_p , $p \in \mathbb{N} \setminus \{0\}$, be the continuous mapping defined in Proposition 3.3 and let $(\phi_p)_\ell$, $\ell \geq 0$, be the associated homology homomorphism. We then have*

$$(\phi_1)_\ell([B_1(M), B_0(M)]) \neq 0 \Rightarrow (\phi_p)_\ell([B_p(M), B_{p-1}(M)]) \neq 0 \forall p \in \mathbb{N} \setminus \{0\},$$

where $[B_p(M), B_{p-1}(M)]$ is defined in (3.13).

Proof. We argue by induction. For $p = 1$, we have

$$(\phi_1)_\ell([B_1(M), B_0(M)]) \neq 0.$$

Let $p \in \mathbb{N} \setminus \{0\}$. Assume that

$$(\phi_p)_\ell([B_p(M), B_{p-1}(M)]) \neq 0. \quad (3.14)$$

We claim that

$$(\phi_{p+1})_\ell([B_{p+1}(M), B_p(M)]) \neq 0. \quad (3.15)$$

For this, let

$$\partial_{p+1}: H_{\ell+1}(B_{p+1}(M), B_p(M)) \rightarrow H_\ell(B_p(M), B_{p-1}(M)), \ell \geq 0,$$

be the connecting homomorphism, see [6], and let $\varphi_{p+1} \in H^\ell(\Omega^{p+1}/\mathfrak{q}_{p+1})$ such that

$$\partial_{p+1}(\varphi_{p+1} \cdot [B_{p+1}(M), B_p(M)]) = [B_p(M), B_{p-1}(M)].$$

See ([2], Appendix C) for the existence of φ_{p+1} . By (3.14), we have

$$(\phi_p)_\ell(\partial_{p+1}(\varphi_{p+1} \cdot [B_{p+1}(M), B_p(M)])) \neq 0. \quad (3.16)$$

We now introduce the following relative homological diagram.

Since we have supposed that J has no critical points in Σ^+ , we derive from Proposition 3.1 that the natural injection $\bar{i}: E_p \hookrightarrow F_{p+1}$ is an homology equivalence. Thus it induces a relative homology isomorphism, denoted

$$(\bar{i})_\ell: H_\ell(E_{p+1}, E_p) \rightarrow H_\ell(E_{p+1}, F_{p+1}), \ell \geq 0.$$

Denote,

$$\delta_{p+1}: H_{\ell+1}(E_{p+1}, E_p) \rightarrow H_\ell(E_p, F_p), \ell \geq 0,$$

the connecting homomorphism of the triad (E_{p+1}, E_p, F_p) . the following diagram commutes.

$$\begin{array}{ccc} H_{\ell+1}(B_{p+1}(M), B_p(M)) & \xrightarrow{(\phi_{p+1})_\ell} & H_{\ell+1}(E_{p+1}, F_{p+1}) \\ \downarrow \partial_{p+1} & & \downarrow \overline{\partial_{p+1}} \\ H_\ell(B_p(M), B_{p-1}(M)) & \xrightarrow{(\phi_p)_\ell} & H_\ell(E_p, F_p) \end{array}$$

where $\overline{\partial_{p+1}} = \delta_{p+1} o(\bar{i})_{\ell+1}^{-1}$. We then have

$$(\phi_p)_\ell o \partial_{p+1} = \overline{\partial_{p+1}} o(\phi_{p+1})_\ell.$$

Thus from (3.16), we have

$$\overline{\partial_{p+1}}((\phi_{p+1})_\ell(\varphi_{p+1} \cdot [B_{p+1}(M), B_p(M)])) \neq 0$$

and therefore,

$$(\phi_{p+1})_\ell(\varphi_{p+1} \cdot [B_{p+1}(M), B_p(M)]) \neq 0$$

Using the fact that $(\phi_{p+1})_\ell$ is $H^\ell(\Omega^{p+1}/\rho)$ -linear, we get

$$\varphi_{p+1} \cdot (\phi_{p+1})_\ell([B_{p+1}(M), B_p(M)]) \neq 0,$$

and hence

$$(\phi_{p+1})_\ell([B_{p+1}(M), B_p(M)]) \neq 0.$$

Claim (3.15) is valid and the result follows. \square

We now state the following result

Proposition 3.5 *There exists a positive integer p_0 large enough such that for any $p \geq p_0$,*

$$\phi_p(B_{p+1}(M), B_{p-1}(M)) \subset (F_p, F_p).$$

Proof. Let $u = \sum_{i=1}^p \alpha_i \delta_{a_i} \in B_p(M)$.

$$\phi_p(u) = \frac{\sum_{i=1}^p \overline{\alpha_i} PU(\overline{a_i}, \lambda)}{\left| \sum_{i=1}^p \overline{\alpha_i} PU(\overline{a_i}, \lambda) \right|},$$

where $(\overline{a_1}, \dots, \overline{a_p}, \overline{\alpha_1}, \dots, \overline{\alpha_p}) = (h_p o j_p o i_p o \pi_p)(u)$. Here h_p, j_p, i_p and π_p are defined in the proof of Proposition 3.3. By (3.5), we have

$$\phi_p(u) \in W_u(V(p, \varepsilon_p)).$$

It remains to prove that $\phi_p(u) \in J_{p^{2\mu^*-1} \widetilde{S}_{HL}}$. For this we use the expansion of Proposition 2.1 and the fact that

$$\begin{aligned} H(a, a) &\leq C, \quad \forall a \in M, \\ G(a, b) &\geq \gamma > 0, \quad \forall a, b \in M, \quad \text{such that } d(a, b) \geq d_1, \end{aligned}$$

we get

$$J(\phi_p(u)) \geq P^{2\mu^*-1} \widetilde{S}_{HL} (1 + (\bar{c} - p\bar{\gamma}) \frac{1}{\lambda^{n-2}}),$$

where \bar{c} and $\bar{\gamma}$ are two position constants. Let p_0 such that $\bar{c} - p_0 \bar{\gamma} < 0$, we derive our result for any $p \geq p_0$. \square

Proof of Theorem 1.1 completed: Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be a bounded domain such that $H_{k_0}(\Omega) \neq 0$, $k_0 \geq 1$. there exists a singular k_0 -chain in Ω without boundary with \mathbb{Z}_2 -coefficients, (k_0 -cycle), which defines a non zero class in the homology group $H_\ell(\Omega)$, $\ell \geq 0$. following [6], the k_0 -cycle

can be viewed as a compact and closed k_0 -dimensional manifold M in Ω . Using the preceding notations, we prove that

$$(\phi_1)_\ell([B_1(M), B_0(M)]) \neq 0. \quad (3.17)$$

Indeed, if $(\phi_1)_\ell([B_1(M), B_0(M)]) = 0$, then using the mapping χ introduced in (3.3), we find that

$$(\chi)_\ell((\phi_1)_\ell([B_1(M), B_0(M)])) = [M, \partial M] = 0 \quad \text{in } H_\ell(\Omega).$$

This is absurd, since M has no boundary and defines a non zero class in $H_\ell(\Omega)$. The assertion (3.17) holds and hence claim (1.5) of the first step of the introduction is valid. Now if we suppose that J has no critical point in Σ^+ , it is proved in (3.14) and (3.15) that for any $p \in \mathbb{N} \setminus \{0\}$,

$$(\phi_p)_\ell([B_p(M), B_{p-1}(M)]) \neq 0 \Rightarrow (\phi_{p+1})_\ell([B_{p+1}(M), B_p(M)]) \neq 0.$$

Thus (1.6) follows. Lastly, it is proved in Proposition 3.5 that there exists a large positive integer p_0 such that the mapping ϕ_{p_0} is valued in the pair (F_{p_0}, F_{p_0}) . Therefore it is homologically trivial. This yields the assertion (1.7) and proves our Theorem.

References

- [1] Bahri, A., Coron, J.M., The scalar curvature problem on the standard three dimensional spheres, *J. Funct. Anal.* 95 (1991), 106–172.
- [2] Bahri, A., Coron, J.M., On a nonlinear Elliptic equation Involving the critical Sobolev Exponent: The effect of the topology on the domain, *Comm. Pure Appl. Math.* 41 (1988), 253–294.
- [3] Bahri, A., *Critical Point at Infinity in Some Variational Problems*, Pitman Res. Notes Math., vol. 182, Longman Sci. Tech., Harlow, 1989. 81.
- [4] Bredon, G., *Introduction to compact transformations Groups*, Academic Press, New York (1972).
- [5] Cassani, D., Scafington, J.V., Zhang, J., Groundstates for Choquard type equations with Hardy-Littlewood-Sobolev lower critical exponent, Available via <https://arxiv.org/pdf/1709.09448.pdf>.
- [6] Dold, A., *Lectures on Algebraic Topology*, Springer-Verlag Berlin and New York.
- [7] Du, L., Yang, M., Uniqueness and nondegeneracy of solutions for a critical nonlocal equation, *Discrete Contin. Dyn. Syst.* 39, 5847–5866 (2019)
- [8] Lieb, E., Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities, *Ann. of Math.* 118(1983), 349–374.
- [9] E. Lieb, Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation, *Studies in Appl. Math.*, 57(1976/77), 93–105.
- [10] Lieb, E., Loss, M., "Analysis". Graduate Studies in Mathematics, AMS, Providence, Rhode island, 2001.
- [11] Gao, F., Yang, M., On nonlocal Choquard equations with Hardy–Littlewood–Sobolev critical exponents, *J. Math. Anal. Appl.*, 448 (2017), 1006–1041.

- [12] Gao, F., Yang, M., The Brezis–Nirenberg type critical problem for the nonlinear Choquard equation, *Sci. China Math.* 61, 1219–1242 (2018).
- [13] Goel, D., Radulescu, V., Sreenadh, K., Coron problem for nonlocal equations involving Choquard nonlinearity, *Adv. Nonlinear Stud.* (2019), <https://doi.org/10.1515/ans-2019-2064>.
- [14] Goel D., Sreenadh K., Critical growth elliptic problems involving Hardy-Littlewood-Sobolev critical exponent in non-contractible domains, *Adv. Nonlinear Anal.* 9 (2020), no. 1, 803–835.
- [15] Guo, L., Hu, T., Peng, S., Shuai, W., Existence and uniqueness of solutions for Choquard equation involving Hardy–Littlewood–Sobolev critical exponent, *Calc. Var. Partial Differ. Equ.* 58, 128, 34 pp (2019),
- [16] Brézis, H., Nirenberg, L., Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Comm. Pure Appl. Math.*, 36 (1983), 437–477.
- [17] Coron, J.M., Topologie et cas limite des injections de Sobolev, *C. R. Acad. Sci. Paris Sér. I Math.* 299 (1984), no. 7, 209–212.
- [18] Squassina, M., Yang, M., Zhao, S., Local uniqueness of blow-up solutions for critical Hartree equations in bounded domain, *Calc. Var.* (2023) 62:217, <https://doi.org/10.1007/s00526-023-02551-1>.
- [19] Moroz, V., Schaftingen J.V., A guide to the Choquard equation, *J. Fixed Point Theory Appl.* 19 (2019), 773–813.
- [20] . Moroz, V., Schaftingen, J.V., Groundstates of nonlinear Choquard equations: Hardy–Littlewood–Sobolev critical exponent, *Commun. Contemp. Math.* 17, (2015).
- [21] Moroz V., Schaftingen J.V., Groundstates of nonlinear Choquard equations: existence, qualitative properties and decay asymptotics, *J. Funct. Anal.* 265 (2013), no. 2, 153–184,
- [22] Struwe, M., A global compactness result for elliptic boundary value problem involving limiting nonlinearities, *Math. Z.* 187, (1984), 511–517.
- [23] Ma, L., Zhao, L., Classification of positive solitary solutions of the nonlinear Choquard equation, *Ration. Mech. Anal.* 195(2010), 455–467.
- [24] Lions, P.L., The concentration compactness principle in the calculus of variations, (Part 1 and Part 2), *Riv. Mat. Iberoamericana* 1 (1985), 145–201, 45–121.
- [25] Lions, P.L., The Choquard equation and related questions, *Nonlinear Anal.* 4 (1980), 1063–1072.
- [26] Penrose, R., On gravity’s role in quantum state reduction, *Gen. Relativ. Gravitat.* 28 (1996), 581–600.
- [27] Pekar, S., *Untersuchung über die Elektronentheorie der Kristalle*, Akademie Verlag, Berlin, 1954.