

A partial proof of the Brouwer's conjecture

Slobodan Filipovski ^{*a}

^aFAMNIT, University of Primorska, Koper, Slovenia

Abstract

Let G be a simple graph with n vertices and m edges and let k be a natural number such that $k \leq n$. Brouwer conjectured that the sum of the k largest Laplacian eigenvalues of G is at most $m + \binom{k+1}{2}$. In this paper we prove that this conjecture is true for simple (m, n) -graphs where $n \leq m \leq \frac{\sqrt{3}-1}{4}(n-1)n$ and $k \in \left[\sqrt[3]{\frac{8m^2}{n-1} + 4mn + n^2}, n \right]$. Moreover, we prove that the conjecture is true for all simple (m, n) -graphs where $k(\leq n)$ is a natural number from the interval $\left[\sqrt{2n - 2m + 2\sqrt{2m^2 + mn(n-1)}}, 1 + \frac{8m^2}{n^2(n-1)} + \frac{4m}{n} \right]$.

1 Introduction

Let G be a simple graph with vertex set $V(G)$ and edge set $E(G)$ and let $|V(G)| = n$ and $|E(G)| = m$. The degree of a vertex $v \in V(G)$ denoted by $d(v)$, is the number of neighbours of v . The Laplacian matrix of G is the $n \times n$ matrix $L(G) = [l_{ij}]$ defined as $L(G) = D(G) - A(G)$, where $D(G)$ is the diagonal matrix of vertex degrees of the graph G , and $A(G)$ is the adjacency matrix of G . It is well known that $L(G)$ is a positive semidefinite matrix, consequently its eigenvalues are nonnegative real numbers. The eigenvalues of $L(G)$ are called the Laplacian eigenvalues of G and are denoted by $\mu_1(G) \geq \mu_2(G) \geq \dots \geq \mu_n(G)$. Since each row sum of $L(G)$ is 0, it holds $\mu_n(G) = 0$. In this paper, we study the sum $S_k(G) = \sum_{i=1}^k \mu_i(G)$ for $1 \leq k \leq n$. Brouwer [1] has conjectured the following upper bound for the sum $S_k(G)$.

Conjecture 1. [1] *Let G be a graph with vertices n and edges m . Then*

$$S_k(G) \leq m + \binom{k+1}{2}$$

for $k = 1, 2, \dots, n$.

*Corresponding author; slobodan.filipovski@famnit.upr.si

Based on a computer check, Brouwer [1] showed that the Conjecture 1 is true for all graphs with at most 10 vertices. There have been many partial proofs of this conjecture using specific methods from spectral graph theory. For $k = 1$, the conjecture follows from the well-known inequality $\mu_1(G) \leq |V(G)| = n$, see in [7]. The conjecture has been proved to be true for any graphs with $k = 2, n - 2, n - 3$, see [5, 8], trees [8], threshold graphs [9], unicyclic graphs [6], bicyclic graphs [6, 12], tricyclic graphs having no pendant vertices [10, 12], regular graphs [2, 11], split graphs [11], cographs [2, 11], planar graphs when $k \geq 11$ and bipartite graphs when $k \geq \sqrt{32n}$ [4]. Also the cases $k = n$ and $k = n - 1$ are straightforward. Another related upper bound was given by Zhou in [14]:

$$S_k(G) \leq \frac{2mk + \sqrt{mk(n-k-1)(n^2-n-2m)}}{n-1}$$

where $1 \leq k \leq n - 2$ and $m = |E(G)|$.

In this paper we prove that the Brouwer's conjecture is true for all simple (m, n) -graphs where $n \leq m \leq \frac{\sqrt{3}-1}{4}(n-1)n$ and $k \in \left[\sqrt[3]{\frac{8m^2}{n-1} + 4mn + n^2}, n \right]$. Moreover, we prove that this conjecture is true for all simple (m, n) -graphs where $k \leq n$ is a natural number from the interval $\left[\sqrt{2n-2m+2\sqrt{2m^2+mn(n-1)}}, 1 + \frac{8m^2}{n^2(n-1)} + \frac{4m}{n} \right]$.

2 A partial proof of the conjecture

Let J be all-ones matrix of size n and let $L' = J - L$. Since $(L')^T = J^T - L^T = J - L = L'$, we obtain that L' is a symmetric matrix. Let $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n$ be the eigenvalues of L , and let $\mu'_1 \geq \mu'_2 \geq \dots \geq \mu'_n$ be the eigenvalues of L' , in the decreasing order. Moreover, let $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ be the eigenvalues of J . It is well-known that $\lambda_1(J) = n$ and $\lambda_i(J) = 0$, for $i = 2, 3, \dots, n$. Applying the Weyl's inequality [13] we get the following result.

Lemma 2. a) $\mu_i + \mu'_{n-i+2} \leq 0$ for $i = 2, 3, \dots, n$.

b) $\mu_1 + \mu'_1 \geq n$.

Proof. a) From Weyl's inequality we get:

$$\mu_i(L) + \mu'_j(L') \leq \lambda_{i+j-n}(L + L') = \lambda_{i+j-n}(J). \quad (1)$$

Setting $j = n - i + 2$ in (1) we get the inequality:

$$\mu_i + \mu'_{n-i+2} \leq \lambda_2(J) = 0, \quad \text{for } i = 2, 3, \dots, n. \quad (2)$$

b) Since $\text{trace}(L) = \sum_{i=1}^n \mu_i$, $\text{trace}(L') = \sum_{i=1}^n \mu'_i$ and $\text{trace}(L) + \text{trace}(L') = \text{trace}(J) = n$ we have

$$\mu_1 + \mu'_1 + \sum_{i=2}^n (\mu_i + \mu'_{n-i+2}) = n.$$

From a) we get $\mu_1 + \mu'_1 \geq n$.

□

The *first Zagreb index* for a graph with a vertex set $\{v_1, v_2, \dots, v_n\}$ and vertex degrees $d_i = \deg(v_i)$ for $i = 1, \dots, n$ is defined by

$$M_1(G) = \sum_{i=1}^n d_i^2.$$

The following upper bound for M_1 , obtained by de Caen in [3], will be used in our proof.

Lemma 3. [3] *Let $G = (V, E)$ be a graph with n vertices and m edges. Then*

$$M_1(G) = \sum_{i=1}^n d_i^2 \leq \frac{2m^2}{n-1} + mn - 2m.$$

Moreover, if G is connected, then equality holds if and only if G is either a star $K_{1, n-1}$ or a complete graph K_n .

In the next theorem we give an upper bound for $S_k(G)$ in terms of m, n and k .

Theorem 4. *Let G be a simple graph on n vertices and m edges. Then*

$$S_k(G) = \sum_{i=1}^k \mu_i \leq n + \sqrt{(k-1) \left(\frac{2m^2}{n-1} + mn + \frac{n^2}{4} \right)}.$$

Proof. From $\text{trace}(L \cdot L^T) = \sum_{i=1}^n \sum_{j=1}^n l_{ij}^2$ and $L = L^T$ we obtain

$$\sum_{i=1}^n \mu_i^2 = \text{trace}(L^2) = \text{trace}(L \cdot L^T) = \sum_{i=1}^n \sum_{j=1}^n l_{ij}^2. \quad (3)$$

Analogously we get

$$\sum_{i=1}^n (\mu'_i)^2 = \text{trace}((L')^2) = \text{trace}(L' \cdot (L')^T) = \sum_{i=1}^n \sum_{j=1}^n (l'_{ij})^2. \quad (4)$$

Combining (3) and (4) we get:

$$\begin{aligned} \sum_{i=1}^n \mu_i^2 + \sum_{i=1}^n (\mu'_i)^2 &= \sum_{i=1}^n \sum_{j=1}^n l_{ij}^2 + \sum_{i=1}^n \sum_{j=1}^n (l'_{ij})^2 = \sum_{i=1}^n \sum_{j=1}^n (l_{ij}^2 + (1 - l_{ij})^2) \\ &= \sum_{i=1}^n \sum_{j=1}^n (2l_{ij}^2 - 2l_{ij} + 1) = \left(\sum_{i=1}^n 2d_i^2 - \sum_{i=1}^n 2d_i + n \right) + \left(\sum_{i=1}^n \sum_{j=1, j \neq i}^n (2l_{ij}^2 - 2l_{ij} + 1) \right) \\ &= (2M_1 - 4m + n) + ((4 + 4 + \dots + 4) + n^2 - n) \end{aligned}$$

$$= 2M_1 - 4m + n + 2m \cdot 4 + n^2 - n = 2M_1 + 4m + n^2.$$

Let $t = \frac{\mu_1}{n}$. Since $\mu_1 \leq n$ we get $0 < t \leq 1$. The inequality $\mu_1 + \mu'_1 \geq n$ implies

$$\mu_1^2 + (\mu'_1)^2 \geq n^2 t^2 + n^2 (1-t)^2 = n^2 (2t^2 - 2t + 1). \quad (5)$$

Moreover, from $\mu_i + \mu'_{n-i+2} \leq 0$, for $i = 2, 3, \dots, n$, and from $\mu_i \geq 0$ we easily conclude that $\mu_i^2 \leq (\mu'_{n-i+2})^2$. Therefore we have

$$\begin{aligned} 2M_1 + 4m + n^2 &= \sum_{i=1}^n \mu_i^2 + \sum_{i=1}^n (\mu'_i)^2 = (\mu_1^2 + (\mu'_1)^2) + \sum_{i=2}^n (\mu_i^2 + (\mu'_{n-i+2})^2) \\ &\geq (1 - 2t(1-t))n^2 + 2 \sum_{i=2}^n \mu_i^2. \end{aligned}$$

From the last inequality we get

$$\sum_{i=2}^k \mu_i^2 \leq \sum_{i=2}^n \mu_i^2 \leq \frac{2M_1 + 4m + n^2}{2} - \frac{(1 - 2t(1-t))n^2}{2}. \quad (6)$$

Now, based on the inequality (6) and applying the inequality between the arithmetic and quadratic means for the nonnegative eigenvalues $\mu_2, \mu_3, \dots, \mu_n$ we get

$$\sum_{i=2}^k \mu_i \leq \sqrt{(k-1)(\mu_2^2 + \dots + \mu_k^2)} \leq \sqrt{(k-1) \left(\frac{2M_1 + 4m + n^2}{2} - \frac{(1 - 2t(1-t))n^2}{2} \right)}. \quad (7)$$

Hence we get

$$S_k(G) = \sum_{i=1}^k \mu_i = \mu_1 + \sum_{i=2}^k \mu_i \leq nt + \sqrt{(k-1)(M_1 + 2m + n^2 t(1-t))}. \quad (8)$$

Setting $M_1 \leq \frac{2m^2}{n-1} - 2m + mn$ in (8) we obtain the following inequality

$$\sum_{i=1}^k \mu_i \leq nt + \sqrt{(k-1) \left(\frac{2m^2}{n-1} + mn + n^2 t(1-t) \right)}. \quad (9)$$

In the end, from $t \leq 1$ and $t(1-t) \leq \frac{1}{4}$ we have

$$S_k(G) = \sum_{i=1}^k \mu_i \leq n + \sqrt{(k-1) \left(\frac{2m^2}{n-1} + mn + \frac{n^2}{4} \right)}.$$

□

Theorem 5. *Let G be a simple graph on n vertices and m edges such that $n \leq m \leq \frac{\sqrt{3}-1}{4}(n-1)n$. The Brouwer's conjecture holds for all natural numbers k from the interval $\left[\sqrt[3]{\frac{8m^2}{n-1} + 4mn + n^2}, n \right]$.*

Proof. The bound in Theorem 4 and the assumption $n \leq m$ yield

$$S_k(G) \leq n + \sqrt{(k-1) \left(\frac{2m^2}{n-1} + mn + \frac{n^2}{4} \right)} < m + \sqrt{k \left(\frac{2m^2}{n-1} + mn + \frac{n^2}{4} \right)}. \quad (10)$$

Now, we are going to show that for $k \in \left[\sqrt[3]{\frac{8m^2}{n-1} + 4mn + n^2}, n \right]$, it holds

$$m + \sqrt{k \left(\frac{2m^2}{n-1} + mn + \frac{n^2}{4} \right)} \leq m + \binom{k+1}{2}.$$

Since $\frac{k^2}{2} < \binom{k+1}{2}$, it suffices to prove that

$$k \left(\frac{2m^2}{n-1} + mn + \frac{n^2}{4} \right) \leq \frac{k^4}{4},$$

which obviously holds for $k \geq \sqrt[3]{\frac{8m^2}{n-1} + 4mn + n^2}$.

In the end, we verify that for $m \leq \frac{\sqrt{3}-1}{4}(n-1)n$ it occurs $\sqrt[3]{\frac{8m^2}{n-1} + 4mn + n^2} < n$.

The inequality $\sqrt[3]{\frac{8m^2}{n-1} + 4mn + n^2} < n$ is equivalent to the quadratic inequality (in m)

$$8m^2 + 4mn(n-1) - n^2(n-1)^2 < 0. \quad (11)$$

Solving the inequality (11) we get $m \in \left(\frac{(-\sqrt{3}-1)}{4}n(n-1), \frac{(\sqrt{3}-1)}{4}n(n-1) \right)$.

□

Remark 6. We illustrate the above theorem for simple graphs on 100 vertices. From the condition $n \leq m \leq \frac{\sqrt{3}-1}{4}n(n-1)$ we have $100 \leq m \leq 1811$. Based on the restrictions for k , that is, $k \in \left[\sqrt[3]{\frac{8m^2}{n-1} + 4mn + n^2}, n \right]$, in the table below we consider particular examples and give the corresponding values of k for which the Brouwer's conjecture is true.

m	k
100	[38, 100]
200	[46, 100]
300	[52, 100]
400	[57, 100]
500	[62, 100]
600	[66, 100]
700	[70, 100]

Note that the Theorem 5 concerns the simple graphs with n vertices and with at most $\frac{(\sqrt{3}-1)}{4}n(n-1)$ edges. This is approximately $\frac{2}{5}$ of all connected graphs on n vertices. The next theorem extends the set of graphs for which the Brouwer's conjecture is true.

Theorem 7. Let G be a simple graph on n vertices and m edges such that

$$\sqrt{2n - 2m + 2\sqrt{2m^2 + mn(n-1)}} < 1 + \frac{8m^2}{n^2(n-1)} + \frac{4m}{n}.$$

The Brouwer's conjecture is true for any natural number $k \leq n$ from the interval

$$\left[\sqrt{2n - 2m + 2\sqrt{2m^2 + mn(n-1)}}, 1 + \frac{8m^2}{n^2(n-1)} + \frac{4m}{n} \right].$$

Proof. Let $g(t) = nt + \sqrt{(k-1)\left(\frac{2m^2}{n-1} + mn + n^2t(1-t)\right)}$ where $0 \leq t \leq 1$. Then $g'(t) = n + \frac{n^2(k-1)(1-2t)}{2\sqrt{(k-1)\left(\frac{2m^2}{n-1} + mn + n^2t(1-t)\right)}}$. If $0 \leq t \leq \frac{1}{2}$, then it is clear that $g'(t) > 0$.

Now, let $t > \frac{1}{2}$. We obtain

$$g'(t) > 0 \Leftrightarrow n > \frac{n^2(k-1)(2t-1)}{2\sqrt{(k-1)\left(\frac{2m^2}{n-1} + mn + n^2t(1-t)\right)}}. \quad (12)$$

The inequality (12) is equivalent to

$$\frac{8m^2}{n-1} + 4mn + 4n^2t(1-t) \geq n^2(k-1)(2t-1)^2. \quad (13)$$

We easily observe that the inequality in (13) holds since $2t-1 \leq 1$, $n^2t(1-t) \geq 0$ and $k \leq 1 + \frac{8m^2}{n^2(n-1)} + \frac{4m}{n}$.

Since $g'(t) > 0$, we get that $g(t)$ is an increasing function on the interval $[0, 1]$. Recalling the inequality (9) we have

$$S_k(G) \leq g(t) \leq g(1) = n + \sqrt{(k-1)\left(\frac{2m^2}{n-1} + mn\right)}.$$

Let us suppose that $k \geq \sqrt{2n - 2m + 2\sqrt{2m^2 + mn(n-1)}}$. We get $2m + k^2 \geq 2n + 2\sqrt{2m^2 + mn(n-1)}$. Hence

$$\begin{aligned} m + \binom{k+1}{2} &> m + \frac{k^2}{2} \geq n + \sqrt{2m^2 + mn(n-1)} = n + \sqrt{(n-1)\left(\frac{2m^2}{n-1} + mn\right)} \\ &\geq n + \sqrt{(k-1)\left(\frac{2m^2}{n-1} + mn\right)} = g(1) \geq S_k(G). \end{aligned}$$

□

Remark 8. We illustrate the above theorem for simple graphs on 100 vertices. From the condition $\sqrt{2n - 2m + 2\sqrt{2m^2 + mn(n-1)}} < 1 + \frac{8m^2}{n^2(n-1)} + \frac{4m}{n}$ we get $m \geq 1468$. Based on the restrictions for k , in the table below we consider particular examples and give the corresponding values of k for which the Brouwer's conjecture is true.

m	k
1500	[78, 79]
1600	[79, 85]
1700	[80, 92]
1800	[81, 99]
1900	[82, 100]
2000	[83, 100]
2100	[84, 100]

Acknowledgement: This work is supported in part by the Slovenian Research Agency (research program P1-0285 and research projects N1-0210, J1-3001, J1-3002, J1-3003, and J1-4414)

References

- [1] A. E. Brouwer and W. H. Haemers, *Spectra of Graphs*, Springer, New York, 2012.
- [2] J. Berndsen, *Three Problems in Algebraic Combinatorics*, Master's thesis, Eindhoven University of Technology, 2012.
- [3] D. de Caen, An upper bound on the sum of squares of degrees in a graph, *Discr. Math.* 185 (1998) 245–248.
- [4] J. N. Cooper, Constraints on Brouwer's Laplacian spectrum conjecture, *Linear Algebra Appl.* 615 (2021), 11–27.
- [5] X. Chen, On Brouwer's conjecture for the sum of k largest Laplacian eigenvalues of graphs, *Linear Algebra Appl.* 578 (2019), 402–410.
- [6] Z. Du and B. Zhou, Upper bounds for the sum of Laplacian eigenvalues of graphs, *Linear Algebra Appl.* 436 (2012), 3672–3683.
- [7] C. Godsil, G. Royle, *Algebraic Graph Theory*, Springer-Verlag, New York, 2001.
- [8] W. H. Haemers, A. Mohammadian, and B. Tayfeh-Rezaie, On the sum of Laplacian eigenvalues of graphs, *Linear Algebra Appl.* 432 (2010).
- [9] C. Helmberg and V. Trevisan, Spectral threshold dominance, Brouwer's conjecture and maximality of Laplacian energy, *Linear Algebra Appl.* 512 (2017), 18–31.
- [10] P. Kumar, S. Merajuddin, and S. Pirzada, Computing the sum of k largest Laplacian eigenvalues of tricyclic graphs, *Discrete Math. Lett.* 11 (2023), 14–18.
- [11] Mayank, *On Variants of the Grone-Merris Conjecture*, Master's thesis, Eindhoven University of Technology, 2010.

- [12] S. Wang, Y. Huang, and B. Liu, On a conjecture for the sum of Laplacian eigenvalues, *Math. Comput. Model.* 56 (2012), 60–68.
- [13] H. Weyl, Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung) bib *Mathematische Annalen* 71, no. 4 (1912): 441–479.
- [14] B. Zhou, On Laplacian Eigenvalues of a Graph, *Zeitschrift fur Naturforschung* A, 59, 2014.