

RELATIVE p -CLASS GROUPS AND p -SELMER GROUPS

DEBAJYOTI DE, DIPRAMIT MAJUMDAR, AND SUDIPA MONDAL

ABSTRACT. Let E be an elliptic curve with j -invariant 0 or 1728 and let \tilde{E} be a k^{th} twist of E . We show that for any prime p of good reduction of \tilde{E} , a degree k relative p -class group and the root number of \tilde{E} determines the dimension of the p -Selmer group of \tilde{E} . As a consequence, we construct families of large rank p -class group. We also relate congruent number and cube sum problem with relative p -class group.

1. INTRODUCTION

The Selmer group of an elliptic curve and class group of certain number field can sometime be embedded inside a Galois cohomology group and thus one expects that Selmer group of elliptic curve and certain class group should be closely related and this relationship has been explored by various mathematicians [Cas91, Sch96, Č15, Li19, PS21, JMS22]. In this article, we consider CM elliptic curves E with $j(E) \in \{0, 1728\}$. For the elliptic curves with $j(E) = 1728$, we show that the variation of rank of p -Selmer group of E in a quadratic (resp. quartic) twist family is completely determined by the root number and the variation of rank of certain component of p -class group of a family of relative quadratic (resp. quartic) extensions of a number field K_E (resp. F_E). More precisely, let E be an elliptic curve with $j(E) = 1728$ and let \tilde{E} be a quadratic (resp. quartic) twist of E . Let p be an odd prime such that both E and \tilde{E} have good reduction at p . Let $K = \mathbb{Q}(i)$ and \mathfrak{p} be a prime in K lying over p . For $C \in \{E, \tilde{E}\}$, let $L_{C,\mathfrak{p}} := K(C[\mathfrak{p}])$ be the \mathfrak{p} division field of C , then $L_{C,\mathfrak{p}}$ is Galois over K with cyclic Galois group G of order $N_{K/\mathbb{Q}}(\mathfrak{p}) - 1$. Let $K_{C,\mathfrak{p}}$ (resp. $F_{C,\mathfrak{p}}$) be the subfield of $L_{C,\mathfrak{p}}$ containing K such that $\text{Gal}(L_{C,\mathfrak{p}}/K_{C,\mathfrak{p}}) \cong \frac{\mathbb{Z}}{2\mathbb{Z}}$ (resp. $\text{Gal}(L_{C,\mathfrak{p}}/F_{C,\mathfrak{p}}) \cong \frac{\mathbb{Z}}{4\mathbb{Z}}$). Let χ_C denotes the character of G which gives the action of G on $C[\mathfrak{p}]$, and let $r_{\mathfrak{p}}(C)$ denotes the $k_{\mathfrak{p}} := \frac{\mathcal{O}_K}{\mathfrak{p}}$ -dimension of the χ_C component of the relative quadratic (resp. quartic) class group $\text{Cl}(L_{C,\mathfrak{p}}/K_{C,\mathfrak{p}})[p]$ (resp. $\text{Cl}(L_{C,\mathfrak{p}}/F_{C,\mathfrak{p}})[p]$). Finally, let $s_p(C)$ denotes the \mathbb{F}_p -dimension of p -Selmer group of C (over \mathbb{Q}), and $\omega(C)$ denotes the root number of C (over \mathbb{Q}).

Theorem A (Theorem 2.3, Theorem 2.4, Proposition 2.7). *Let \tilde{E} be a quadratic (resp. quartic) twist of an elliptic curve E with $j(E) = 1728$. With notation as above, $s_p(\tilde{E})$, the rank of p -Selmer group of \tilde{E} is completely determined by the root number $\omega(\tilde{E})$ and $r_{\mathfrak{p}}(\tilde{E})$, the $k_{\mathfrak{p}} := \frac{\mathcal{O}_K}{\mathfrak{p}}$ -dimension of the $\chi_{\tilde{E}}$ component of the relative p -class group $\text{Cl}(L_{\tilde{E},\mathfrak{p}}/K_{\tilde{E},\mathfrak{p}})[p]$ (resp. $\text{Cl}(L_{\tilde{E},\mathfrak{p}}/F_{\tilde{E},\mathfrak{p}})[p]$).*

Further, $K_{E,\mathfrak{p}} \cong K_{\tilde{E},\mathfrak{p}}$ (resp. $F_{E,\mathfrak{p}} \cong F_{\tilde{E},\mathfrak{p}}$) and the field (up to isomorphism) $K_{E,\mathfrak{p}}$ (resp. $F_{E,\mathfrak{p}}$) is independent of the choice of the prime \mathfrak{p} lying above of p .

Similarly, let E be an elliptic curve with $j(E) = 0$ and let \tilde{E} be a quadratic (resp. cubic, resp. sextic) twist of E . Let p be an odd prime such that both E and \tilde{E} have good reduction at p . Let $K = \mathbb{Q}(\zeta_3)$ and \mathfrak{p} be a prime in K lying over p . For $C \in \{E, \tilde{E}\}$, let $L_{C,\mathfrak{p}} := K(C[\mathfrak{p}])$

2020 *Mathematics Subject Classification*. Primary: 11R29, 11G05, Secondary: 11G15, 11R23 .

Key words and phrases. Ideal class groups, Selmer group, Elliptic curves, Division Field.

be the \mathfrak{p} division field of C , then $L_{C,\mathfrak{p}}$ is Galois over K with cyclic Galois group G of order $N_{K/\mathbb{Q}}(\mathfrak{p}) - 1$. Let $K'_{C,\mathfrak{p}}$ (resp. $K_{C,\mathfrak{p}}$, resp. $F_{C,\mathfrak{p}}$) be the subfield of $L_{C,\mathfrak{p}}$ containing K such that $\text{Gal}(L_{C,\mathfrak{p}}/K'_{C,\mathfrak{p}}) \cong \frac{\mathbb{Z}}{2\mathbb{Z}}$ (resp. $\text{Gal}(L_{C,\mathfrak{p}}/K_{C,\mathfrak{p}}) \cong \frac{\mathbb{Z}}{3\mathbb{Z}}$, resp. $\text{Gal}(L_{C,\mathfrak{p}}/F_{C,\mathfrak{p}}) \cong \frac{\mathbb{Z}}{6\mathbb{Z}}$). Let χ_C denotes the character of G which gives the action of G on $C[\mathfrak{p}]$ and let $r_{\mathfrak{p}}(C)$ denotes the $k_{\mathfrak{p}} := \frac{\mathcal{O}_K}{\mathfrak{p}}$ -dimension of the χ_C component of the relative quadratic (resp. cubic, resp. degree six) class group $\text{Cl}(L_{C,\mathfrak{p}}/K'_{C,\mathfrak{p}})[p]$ (resp. $\text{Cl}(L_{C,\mathfrak{p}}/K_{C,\mathfrak{p}})[p]$, resp. $\text{Cl}(L_{C,\mathfrak{p}}/F_{C,\mathfrak{p}})[p]$).

Theorem B (Theorem 3.2, Theorem 3.3, Proposition 3.6). *Let \tilde{E} be a quadratic (resp. cubic, resp. sextic) twist of an elliptic curve E with $j(E) = 0$. With notation as above, the rank of p -Selmer group of \tilde{E} is completely determined by the root number $\omega(\tilde{E})$ and $r_{\mathfrak{p}}(\tilde{E})$, the $k_{\mathfrak{p}} := \frac{\mathcal{O}_K}{\mathfrak{p}}$ -dimension of the $\chi_{\tilde{E}}$ component of the relative p -class group $\text{Cl}(L_{\tilde{E},\mathfrak{p}}/K'_{\tilde{E},\mathfrak{p}})[p]$ (resp. $\text{Cl}(L_{\tilde{E},\mathfrak{p}}/K_{\tilde{E},\mathfrak{p}})[p]$, resp. $\text{Cl}(L_{\tilde{E},\mathfrak{p}}/F_{\tilde{E},\mathfrak{p}})[p]$).*

Further, $K'_{E,\mathfrak{p}} \cong K'_{\tilde{E},\mathfrak{p}}$ (resp. $K_{E,\mathfrak{p}} \cong K_{\tilde{E},\mathfrak{p}}$, resp. $F_{E,\mathfrak{p}} \cong F_{\tilde{E},\mathfrak{p}}$) and the field (up to isomorphism) $K'_{E,\mathfrak{p}}$ (resp. $K_{E,\mathfrak{p}}$, resp. $F_{E,\mathfrak{p}}$) is independent of the choice of the prime \mathfrak{p} lying above of p .

In [Rub87a] the author showed that for an elliptic curve with $j(E) \in \{0, 1728\}$, the p -Selmer rank of E can be determined from the root number of E and the structure of the p -class group of the π -division field of E . Our results can be thought of as an refinement of the result of [Rub87a] which is obtained by studying the structure of π -division field of E .

In the first set of applications of our result, we give a reformulation of rank 0 p -converse theorems in terms of relative p -class group.

Proposition A (Corollary 2.5, Corollary 3.4). *Let E be an elliptic curve with $j(E) = 1728$ (resp. $j(E) = 0$) and $\omega(E) = +1$. The following statements are equivalent:*

- (1) $\text{rk}_{\mathbb{Q}} E = 0$ and $\text{III}(E/\mathbb{Q})$ is finite.
- (2) there exists an odd prime p such that E has good reduction at p and for a prime \mathfrak{p} in $\mathbb{Z}[i]$ (resp. \mathfrak{p} in $\mathbb{Z}[\zeta_3]$) lying over p , such that $\text{Cl}(L_{E,\mathfrak{p}}/F_{E,\mathfrak{p}})[p](\chi_E) = 0$.
- (3) $L(E, 1) \neq 0$.

Similarly, rank 1 p -converse theorem for these curves can be restated as follows:

Proposition B (Corollary 2.6, Corollary 3.5). *Let E be an elliptic curve with $j(E) = 1728$ (resp. $j(E) = 0$) and $\omega(E) = -1$. The following statements are equivalent:*

- (1) $\text{rk}_{\mathbb{Q}} E = 1$ and $\text{III}(E/\mathbb{Q})$ is finite.
- (2) there exists an odd prime p such that E has good reduction at p , $p = \mathfrak{p}\bar{\mathfrak{p}}$ in $\mathbb{Z}[i]$ (resp. in $\mathbb{Z}[\zeta_3]$), such that $\text{Cl}(L_{E,\mathfrak{p}}/F_{E,\mathfrak{p}})[p](\chi_E)$ is a cyclic group.
- (3) $\text{ord}_{s=1} L(E, s) = 1$.

As an application to the congruent number problem, we obtain

Corollary A (Corollary 2.8). *Let $n \equiv 1, 2, 3 \pmod{8}$ be a square free natural number which is a congruent number. For all odd primes p co-prime to n , the class number of the relative quadratic extension $L_{E,\mathfrak{p}}/K_{E,\mathfrak{p}}$ is divisible by p , here E is the elliptic curve $y^2 = x^3 - n^2x$.*

Similarly, as an application to the cube sum problem, we obtain

Corollary B (Corollary 3.7). *Let $n > 2$ be a cube-free integer for which the root number of $E : y^2 = x^3 - 432n^2$ is $+1$. If there are two rational numbers a and b such that $n = a^3 + b^3$, then for any prime p co-prime to $6n$, the class number of the relative cubic extension $L_{E,\mathfrak{p}}/K_{E,\mathfrak{p}}$ is divisible by p .*

Finally, we provide some modest evidence towards the conjecture on unboundedness of n -rank of class groups by using families of elliptic curves with large rank.

Conjecture 1.1. [BG18, Conjecture 1.2] *Let $n > 1$ be an integer and F be a number field. Then $\text{rk Cl}(L)[n]$ is unbounded when L varies in the extensions of F of degree d .*

The conjecture is widely open if d and n are co-prime. For $2 \leq d \leq (p-1)$, and if $\zeta_p \in F$, [BG18, Theorem 1.7] shows that there are infinitely many non-isomorphic number fields L such that $[L : F] = d$ and $\text{rk Cl}(L)[p] \geq 3 + \text{rk Cl}(F)[p]$. Towards this, for $d = 4$, we prove

Theorem C (Theorem 4.2). (1) *Let $E : y^2 = x^3 - x$, $p \equiv 3 \pmod{4}$ be a prime and let F denote the number field $F_{E,p}$. There are infinitely many non-isomorphic number fields L such that $[L : F] = 4$ and $\text{rk Cl}(L)[p] \geq 12 + \text{rk Cl}(F)[p]$.*
(2) *Let $E : y^2 = x^3 - x$, $p \equiv 1 \pmod{4}$ be a prime, \mathfrak{p} a prime in $\mathbb{Z}[i]$ lying above p and let F denote the number field $F_{E,\mathfrak{p}}$. There are infinitely many non-isomorphic number fields L such that $[L : F] = 4$ and $\text{rk Cl}(L)[p] \geq 6 + \text{rk Cl}(F)[p]$.*

We remark that if $\dim_{\mathbb{F}_p} \text{Sel}^p(E/\mathbb{Q})$ is unbounded as E varies over the CM elliptic curves with $j(E) = 1728$, we'll obtain that the Conjecture 1.1 is true for $d = 4$ and $F \cong F_{E,\mathfrak{p}}$. Similarly, for $d = 6$, we show

Theorem D (Theorem 4.6). (1) *Let $E : y^2 = x^3 + 1$, $p \equiv 2 \pmod{3}$ be a prime and let F denote the number field $F_{E,p}$. There are infinitely many non-isomorphic number fields L such that $[L : F] = 6$ and $\text{rk Cl}(L)[p] \geq 12 + \text{rk Cl}(F)[p]$.*
(2) *Let $E : y^2 = x^3 + 1$, $p \equiv 1 \pmod{3}$ be a prime, \mathfrak{p} a prime in $\mathbb{Z}[\zeta_3]$ lying above p and let F denote the number field $F_{E,\mathfrak{p}}$. There are infinitely many non-isomorphic number fields L such that $[L : F] = 6$ and $\text{rk Cl}(L)[p] \geq 6 + \text{rk Cl}(F)[p]$.*

Again, we remark that if $\dim_{\mathbb{F}_p} \text{Sel}^p(E/\mathbb{Q})$ is unbounded as E varies over the CM elliptic curves with $j(E) = 0$, we'll obtain that the Conjecture 1.1 is true for $d = 6$ and $F \cong F_{E,\mathfrak{p}}$.

The paper is structured as follows: In Section 2, we study the p -Selmer group of elliptic curves with j -invariant 1728. In Section 3, we study the p -Selmer group of elliptic curves with j -invariant 0. In Section 4, we construct class group with large p -rank. In the final Section 5, we use SAGE to compute class groups and use that information to determine the rank of various elliptic curves with j -invariant 0 and 1728.

Acknowledgements: We thank Ashay Burungale and B. Sury for discussions.

2. p -SELMER GROUP OF $E_D : Y^2 = X^3 - DX$

Let $E := E_1$ denote the elliptic curve $E : Y^2 = X^3 - X$. For a nonzero integer $D \in \mathbb{Z}$, the elliptic curve $E_D : Y^2 = X^3 - DX$ is the quartic twist of E by D . The elliptic curve E_D has CM by $\mathbb{Z}[i]$ given as $i(x, y) = (-x, iy)$ and for a prime $q \nmid 2D$, E_D has good ordinary reduction at q if and only if $q \equiv 1 \pmod{4}$. By $\omega(E_D)$, we denote the global root number of E_D .

The following lemmas provide an explicit description of n -division polynomial of E_D which will be useful for our study.

Lemma 2.1. *For every integer $n \geq 1$, there exists a homogeneous polynomial $f_n(Z, W) = c_n Z^{s_n} + \dots + d_n W^{s_n} \in \mathbb{Z}[Z, W]$, where $(c_n, d_n, s_n) = \begin{cases} (n, (-1)^{\frac{n-1}{2}}, \frac{n^2-1}{4}) & \text{if } n \text{ is odd,} \\ (\frac{n}{2}, \frac{n}{2}, \frac{n^2-4}{4}) & \text{if } n \text{ is even,} \end{cases}$*

such that

$$\psi_n = \begin{cases} f_n(x^2, D) & \text{if } n \text{ is odd,} \\ 2yf_n(x^2, D) & \text{if } n \text{ is even.} \end{cases}$$

Here $\psi_n \in \mathbb{Z}[x, y]$ is the n^{th} division polynomial for $E_D : y^2 = x^3 - Dx$.

Proof. Note that $f_1(Z, W) = f_2(Z, W) = 1$, $f_3(Z, W) = 3Z^2 - 6ZW - W^2$, $f_4(Z, W) = 2(Z^3 - 5WZ^2 - 5W^2Z + W^3)$. We prove the result by induction on n and using the recursive relations that define ψ_n .

Assume that the statement is true for $1 \leq t \leq n-1$. Let $n = 4m+1$, then $\psi_n = \psi_{2m+2}\psi_{2m}^3 - \psi_{2m-1}\psi_{2m+1}^3$, which implies that $\psi_n = 16y^4 f_{2m+2}(x^2, D)f_{2m}^3(x^2, D) - f_{2m-1}(x^2, D)f_{2m+1}^3(x^2, D)$. We define $f_{4m+1}(Z, W) = 16Z(Z-W)^2 f_{2m+2}(Z, W)f_{2m}^3(Z, W) - f_{2m-1}(Z, W)f_{2m+1}^3(Z, W)$ and using the induction on degree of $f_t(Z, W)$, one checks that $f_{4m+1}(Z, W)$ is a homogeneous polynomial in Z and W of degree $\frac{(4m+1)^2-1}{4}$ and we have $\psi_{4m+1} = f_{4m+1}(x^2, D)$. Similarly, one can show that $f_{4m+3}(Z, W) = f_{2m+3}(Z, W)f_{2m+1}^3(Z, W) - 16Z(Z-W)^2 f_{2m}(Z, W)f_{2m+2}^3(Z, W)$ and $f_{2m}(Z, W) = f_m(Z, W)[f_{m+2}(Z, W)f_{m-1}^2(Z, W) - f_{m-2}(Z, W)f_{m+1}^2(Z, W)]$. Also the formulas of s_n and d_n can be proved using induction. Finally, since

$$\psi_n = \begin{cases} nx^{\frac{n^2-1}{2}} + \dots, & \text{if } n \text{ is odd,} \\ y(nx^{\frac{n^2-4}{2}} + \dots), & \text{if } n \text{ is even,} \end{cases}$$

it follows that $c_n = n$ if n is odd and $c_n = \frac{n}{2}$ if n is even. \square

Lemma 2.2. For every integer $n \geq 1$, there exists a homogeneous polynomial $g_n(Z, W) = Z^{s_n} + \dots \in \mathbb{Z}[Z, W]$, where $s_n = \begin{cases} \frac{n^2-1}{4} & \text{if } n \text{ is odd,} \\ \frac{n^2}{4} & \text{if } n \text{ is even,} \end{cases}$

such that

$$\phi_n = \begin{cases} xg_n(x^2, D)^2 & \text{if } n \text{ is odd,} \\ g_n(x^2, D)^2 & \text{if } n \text{ is even.} \end{cases}$$

Here $\phi_n = x\psi_n^2 - \psi_{n-1}\psi_{n+1} \in \mathbb{Z}[x]$ and ψ_n is the n^{th} division polynomial for $E_D : y^2 = x^3 - Dx$.

Proof. Using induction as in the proof of the preceding lemma, one can show that there exists homogeneous polynomial $\tilde{g}_n(Z, W)$ of degree $2s_n$ such that

$$\phi_n = \begin{cases} x\tilde{g}_n(x^2, D) & \text{if } n \text{ is odd,} \\ \tilde{g}_n(x^2, D) & \text{if } n \text{ is even.} \end{cases}$$

Now from the point addition formula of elliptic curve, observe that $\frac{\phi_{2n}(P)}{\psi_{2n}(P)^2} = x(2nP) = \frac{\phi_2(nP)}{\psi_2(nP)^2}$ and hence $\phi_{2n}(P) = \phi_2(nP) \left(\frac{\psi_{2n}(P)}{\psi_2(nP)} \right)^2$. For the elliptic curve E_D , $\phi_2(P) = (x(P)^2 + D)^2$ and hence

$$\phi_{2n}(P) = \left(\frac{(x(nP)^2 + D)\psi_{2n}(P)}{\psi_2(nP)} \right)^2 = [\phi_n^2(P) + D\psi_n^4(P)]^2,$$

equivalently, $\tilde{g}_{2n}(x^2, D) = g_{2n}(x^2, D)^2 \in \mathbb{Z}[x]$.

On the other hand, if n is odd, then from the formula $4y\omega_n = \psi_{n+2}\psi_{n-1}^2 - \psi_{n-2}\psi_{n+1}^2$, it follows that $\omega_n = y\tilde{f}_n(x^2, D)$, where $\tilde{f}_n(Z, W) \in \mathbb{Z}[Z, W]$ is a homogeneous polynomial. Now consider the polynomial $h_n(Z, W) := Z\tilde{g}_n(Z, W)^2 - Wf_n(Z, W)^4 \in \mathbb{Z}[Z, W]$, then $\tilde{g}_n(x^2, D)h_n(x^2, D) = (x^2 - D)\tilde{f}_n(x^2, D)^2$ as we have $\omega_n^2 = \phi_n^3 - D\phi_n\psi_n^4 = \phi_n(\phi_n^2 - D\psi_n^4)$. Equivalently, in $\mathbb{Z}[X]$, we have the polynomial identity $\tilde{g}_n(X^2, D)h_n(X^2, D) = (X^2 - D)\tilde{f}_n(X^2, D)^2$. Since ϕ_a and

ψ_a are co-prime, it follows that $\tilde{g}_n(X^2, D)$ and $h_n(X^2, D)$ have no common root in $\overline{\mathbb{Q}}$. Now suppose that $(X \pm \sqrt{D})$ is a factor of $\tilde{g}_n(X^2, D) \in \overline{\mathbb{Q}}[X]$, but $(X \mp \sqrt{D})$ is not a factor of $\tilde{g}_n(X^2, D) \in \overline{\mathbb{Q}}[X]$. Since, for the polynomial $(x^2 - D)\tilde{f}_n(x^2, D)^2 \in \overline{\mathbb{Q}}[X]$, multiplicity of the roots $\pm\sqrt{D}$ are odd and for all other roots, the multiplicity is even, it follows that in the above mentioned scenario, $\deg(\tilde{g}_n(X^2, D))$ is odd, which is a contradiction. Now suppose that $(X^2 - D)$ is a factor of $\tilde{g}_n(X^2, D)$, and write $\tilde{g}_n(X^2, D) = (X^2 - D)p_n(X^2) \in \mathbb{Z}[X]$. Then it follows that $p_n(X^2) = q_n(X^2)^2$ where $q_n(X) \mid \tilde{f}_n(X, D) \in \mathbb{Z}[X]$, which in turn implies that $\deg(\tilde{g}_n(X^2, D)) \equiv 2 \pmod{4}$, contradiction. Thus, $\tilde{g}_n(X^2, D) = g_n(X^2)^2 \in \mathbb{Z}[X]$, where $g_n(X) \mid \tilde{f}_n(X, D) \in \mathbb{Z}[X]$. \square

Let us first fix a prime $p \equiv 3 \pmod{4}$, and for an integer D which is co-prime to p , let $s_p(E_D) := \dim_{\mathbb{F}_p} \text{Sel}^p(E_D/\mathbb{Q})$. Let $K = \mathbb{Q}(i)$ and $L_{D,p} := K(E_D[p])$ be the p -division field of E_D over K , then we have $G := \text{Gal}(L_{D,p}/K) \cong \frac{\mathbb{Z}}{(p^2-1)\mathbb{Z}}$. The following theorem shows that the variation of p -Selmer rank of quartic twist by D of E_1 is entirely controlled by the relative p -class group of a cyclic quartic extension of a fixed number field $F_{1,p}$ and root number of E_D .

Theorem 2.3. *Let $p \equiv 3 \pmod{4}$ be a prime and D be an integer co-prime to p . Let $K = \mathbb{Q}(i)$, $L_{D,p} := K(E_D[p])$ be the p -division field of E_D and $F_{D,p}$ be the field $K \subset F_{D,p} \subset L_{D,p}$ with $G = \text{Gal}(L_{D,p}/F_{D,p}) \cong \frac{\mathbb{Z}}{4\mathbb{Z}}$. Let χ_D denotes the character which gives the action of $\text{Gal}(L_{D,p}/K)$ on $E_D[p] \cong \mathbb{F}_{p^2}$ and $r_p(E_D) = \dim_{\mathbb{F}_{p^2}} \text{Cl}(L_{D,p}/F_{D,p})[p](\chi_D)$. Then we have,*

$$r_p(E_D) \leq s_p(E_D) \leq 1 + r_p(E_D),$$

and hence $\omega(E_D)$ and $r_p(E_D)$ uniquely determines $s_p(E_D)$. Moreover, we have $F_{D,p} \cong F_{1,p} \cong \frac{K[X]}{(f_p(X,1))}$ and $L_{D,p} \cong \frac{K[X]}{(f_p(X^4, D))}$.

Proof. From [Rub87a, Theorem 1], we see that $r_p(E_D) \leq s_p(E_D) \leq 1 + r_p(E_D)$, where $r_p(E_D) = \dim_{\mathbb{F}_{p^2}} \text{Hom}(\text{Cl}(L_{D,p}), E_D[p])^G = \dim_{\mathbb{F}_{p^2}} \text{Cl}(L_{D,p})[p](\chi_D)$. Now as $p \nmid [L_{D,p} : K]$, we have $\text{Cl}(L_{D,p})[p] \cong \text{Cl}(L_{D,p}/F_{D,p})[p] \times \text{Cl}(F_{D,p})[p]$ and as $\text{Gal}(L_{D,p}/F_{D,p})$ acts trivially on $\text{Cl}(F_{D,p})[p]$, it follows that $r_p(E_D) = \dim_{\mathbb{F}_{p^2}} \text{Cl}(L_{D,p}/F_{D,p})[p](\chi_D)$. From p -parity conjecture, we know that $s_p(E_D) \equiv \omega(E_D) \pmod{2}$, and hence $\omega(E_D)$ and $r_p(E_D)$ uniquely determines $s_p(E_D)$.

Let $P_D := (x(P_D), y(P_D))$ be a non-trivial element of $E_D[p]$, then $x(P_D)$ is a root of $\psi_p = f_p(X^2, D)$ (as in Lemma 2.1). Hence $x(P_D)^2$ is a root of $f_p(X, D)$ which is a polynomial of degree $\frac{p^2-1}{4}$ and, hence $[K(x(P_D)^2) : K] \leq \frac{p^2-1}{4}$. Since $[K(x(P_D)) : K(x(P_D)^2)] \leq 2$, $[K(y(P_D)) : K(x(P_D))] \leq 2$ and $[K(E_D[p]) : K] = p^2 - 1$, it follows that all the inequalities must be equality and, hence $f_p(X, D)$ is irreducible in $K[X]$ and, $K(x(P_D)^2) \cong \frac{K[X]}{(f_p(X, D))} := F_{D,p}$. Moreover, as $f_p(X, D)$ is a homogeneous polynomial, it follows that for every $D \geq 1$, $\frac{x(P_D)^2}{D}$ is a root of $f_p(X, 1)$ and, hence $F_{D,p} := K(x(P_D)^2) = K(\frac{x(P_D)^2}{D}) \cong K(x(P_1)^2) := F_{1,p}$ is independent of D .

Next, observe that as $[K(x(P_D)) : K] = \frac{p^2-1}{2}$. Since $x(P_D)$ is a root of $f_p(X^2, D)$, it follows that $f_p(X^2, D)$ is irreducible in $K[X]$ and, we have $K(x(P_D)) \cong \frac{\mathbb{Q}(i)[X]}{(f_p(X^2, D))} := K_{D,p}$. Further note that $L_{D,p}$ is a quadratic extension of $K_{D,p}$.

Finally, as $(p-1)P_D = -P_D$, it follows that $\frac{\phi_{p-1}(P_D)}{\psi_{p-1}(P_D)^2} = x((p-1)P_D) = x(-P_D) = x(P_D)$ or equivalently,

$$\frac{g_{p-1}(x(P_D)^2, D)^2}{4y(P_D)^2 f_{p-1}(x(P_D)^2, D)^2} = x(P_D).$$

As a consequence we see that $L_{D,p} := K(E_D[p]) \cong K\left(\sqrt{x(P_D)}\right) \cong \frac{K[X]}{(f_p(X^4, D))}$. \square

Next we consider the case $p \equiv 1 \pmod{4}$ and let $p\mathcal{O}_K = \mathfrak{p}\bar{\mathfrak{p}}$, where \mathfrak{p} and $\bar{\mathfrak{p}}$ are the primes in \mathcal{O}_K lying above p . Let $\pi = a + ib$ be the element in $\mathbb{Z}[i]$ such that $\pi \equiv 1 \pmod{2 + 2i}$ and, $\mathfrak{p} = (\pi)$ and, for every natural number D co-prime to p , we define $\pi_D = \left(\frac{D}{\pi}\right)_4 \pi$, where $(\cdot)_4$ denotes the 4th power residue symbol. With this notation, we have $\psi_D(\mathfrak{p}) = \pi_D$ (resp. $\psi_D(\bar{\mathfrak{p}}) = \bar{\pi}_D$), where ψ_D is the Grössencharacter associated to E_D . From now on, fix a prime \mathfrak{p} in \mathcal{O}_K lying above p . Let $L_{D,\mathfrak{p}} := K(E_D[\pi_D])$ and $G := \text{Gal}(L_{D,\mathfrak{p}}/K) \cong \frac{\mathbb{Z}}{(p-1)\mathbb{Z}}$. Finally, define $f_{D,p}(X) = 4X(X-D)g_{|a|}(X,D)^2 f_{|b|}(X,D)^2 + g_{|b|}(X,D)^2 f_{|a|}(X,D)^2 = pX^{\frac{p-1}{2}} + \dots + D^{\frac{p-1}{2}}$, where $\pi = a + ib \equiv 1 \pmod{2 + 2i}$, and the polynomials f_n, g_n are as in Lemma 2.1,2.2 respectively. The following theorem shows that the variation of p -Selmer rank of quartic twist by D of E_1 is entirely controlled by the relative p -class group of a cyclic quartic extension of a fixed number field $F_{1,p}$ and the root number of E_D .

Theorem 2.4. *Let $p \equiv 1 \pmod{4}$ be a prime with $(p, D) = 1$. Let $K = \mathbb{Q}(i)$ and \mathfrak{p} be a prime in \mathcal{O}_K lying above p . Let $L_{D,\mathfrak{p}} := K(E_D[\pi_D])$ and $F_{D,\mathfrak{p}}$ be the unique subfield of $L_{D,\mathfrak{p}}$ which contains K and satisfies $\text{Gal}(L_{D,\mathfrak{p}}/F_{D,\mathfrak{p}}) \cong \frac{\mathbb{Z}}{4\mathbb{Z}}$. Let χ_D be the character which gives the action of G on $E_D[\pi_D] \cong \mathbb{F}_p$ and $r_{\mathfrak{p}}(E_D) := \dim_{\mathbb{F}_p} \text{Cl}(L_{D,\mathfrak{p}}/F_{D,\mathfrak{p}})[p](\chi_D)$. Then we have*

$$r_{\mathfrak{p}}(E_D) \leq s_p(E_D) := \dim_{\mathbb{F}_p} \text{Sel}^p(E_D/\mathbb{Q}) \leq 1 + r_{\mathfrak{p}}(E_D),$$

and hence $\omega(E_D)$ and $r_{\mathfrak{p}}(E_D)$ uniquely determines $s_p(E_D)$. Further, we have $F_{D,\mathfrak{p}} \cong \frac{\mathbb{Q}[X]}{(f_{1,p}(X))}$ and $L_{D,\mathfrak{p}} \cong \frac{\mathbb{Q}[X]}{(f_{D,p}(X^4))}$.

Proof. From [Rub87a, Theorem 1], we see that $r_{\mathfrak{p}}(E_D) \leq s_p(E_D) \leq 1 + r_{\mathfrak{p}}(E_D)$, where $r_{\mathfrak{p}}(E_D) = \dim_{\mathbb{F}_p} \text{Hom}(\text{Cl}(L_{D,\mathfrak{p}}), E_D[\pi_D])^G = \dim_{\mathbb{F}_p} \text{Cl}(L_{D,\mathfrak{p}})[p](\chi_D)$. Now as $p \nmid [L_{D,\mathfrak{p}} : K]$, we have $\text{Cl}(L_{D,\mathfrak{p}})[p] \cong \text{Cl}(L_{D,\mathfrak{p}}/F_{D,\mathfrak{p}})[p] \times \text{Cl}(F_{D,\mathfrak{p}})[p]$ and, since $\text{Gal}(L_{D,\mathfrak{p}}/F_{D,\mathfrak{p}})$ acts trivially on $\text{Cl}(F_{D,\mathfrak{p}})[p]$, it follows that $r_{\mathfrak{p}}(E_D) = \dim_{\mathbb{F}_p} \text{Cl}(L_{D,\mathfrak{p}}/F_{D,\mathfrak{p}})[p](\chi_D)$. From p -parity conjecture, we know that $s_p(E_D) \equiv \omega(E_D) \pmod{2}$, and hence $\omega(E_D)$ and $r_{\mathfrak{p}}(E_D)$ uniquely determine $s_p(E_D)$.

Now let $P_D = (x(P_D), y(P_D))$ be a generator of $E_D[\pi_D]$. Note that $\pi_D = a_D + ib_D = i^k(a + ib)$ and hence $(|a_D|, |b_D|)$ is either $(|a|, |b|)$ or $(|b|, |a|)$. Since $\pi_D P_D = 0$, it follows that $x(|a_D|P_D) = -x(|b_D|P_D)$ and hence from Lemma 2.1 and Lemma 2.2, it follows that

$$\frac{x(P_D)g_{|a|}(x(P_D)^2, D)^2}{f_{|a|}(x(P_D)^2, D)^2} = -\frac{g_{|b|}(x(P_D)^2, D)^2}{4y(P_D)^2 f_{|b|}(x(P_D)^2, D)^2},$$

and hence $K(E_D[\pi_D]) = K(x(P_D), y(P_D)) \cong K(\sqrt{x(P_D)})$. Again from $\pi_D P_D = 0$, it follows that $x(P_D)^2$ is a root of $f_{D,p}(X)$, which is a homogeneous polynomial in D and X and hence it follows that $K(x(P_D)^2)$ is independent of D (upto isomorphism). Since $[K(\sqrt{x(P_D)}) : K(x(P_D)^2)] \leq 4$, it follows that $F_{D,\pi_D} \subset K(x(P_D)^2)$. We will show that $F_{D,\pi_D} = K(x(P_D)^2)$. Note that $f_{D,p}(X^4)$ can not be irreducible over K , as otherwise, $[K(\sqrt{x(P_D)}) : K] = 2(p-1)$ which is a contradiction. Consequently, $f_{D,p}(X^4)$ has at-least one irreducible factor of degree $(p-1)$ over K .

By [CW77, Lemma 4], we know that the conductor of $L_{D,\mathfrak{p}}$ over K is $\mathfrak{p}f_D$, where f_D is the conductor of Größencharacter ψ_D of E_D , in-particular, $\bar{\mathfrak{p}}$ is unramified in $L_{D,\mathfrak{p}}/K$. Further by [CW77, Lemma 5], we also know that \mathfrak{p} is totally ramified in $L_{D,\mathfrak{p}}/K$.

Similarly, writing $Q_D = (x(Q_D), y(Q_D))$ as a generator of $E_D[\bar{\pi}_D]$, we see that $L_{D,\bar{\mathfrak{p}}} := K(E_D[\bar{\pi}_D]) \cong K(\sqrt{x(Q_D)})$, \mathfrak{p} is unramified in $L_{D,\bar{\mathfrak{p}}}/K$ and $\bar{\mathfrak{p}}$ is totally ramified in $L_{D,\bar{\mathfrak{p}}}/K$. In-particular, we see that $L_{D,\mathfrak{p}} \cap L_{D,\bar{\mathfrak{p}}} = K$. Further, as $L_{D,\mathfrak{p}} \cong L_{D,\bar{\mathfrak{p}}}$, we see that for any isomorphism $\theta : L_{D,\mathfrak{p}} \rightarrow L_{D,\bar{\mathfrak{p}}}$, we have $\theta|_K$ is the non-trivial automorphism of K . Now if $f_{D,p}(X^4)$ has a unique irreducible factor, say $h(X)$, of degree $(p-1)$ over K , then both

$\sqrt{x(P_D)}$ and $\sqrt{x(Q_D)}$ are roots of $h(X)$ and there exists an isomorphism $\theta : L_{D,\mathfrak{p}} \rightarrow L_{D,\bar{\mathfrak{p}}}$ with $\theta(\sqrt{x(P_D)}) = \sqrt{x(Q_D)}$. Consequently, $h(X) = \overline{h(X)}$, which contradicts the fact $\theta|_K$ is the complex conjugation. Hence, $f_{D,\mathfrak{p}}(X^4) = h_1(X)h_2(X) \in K[X]$, where both $h_1(X)$ and $h_2(X)$ are irreducible polynomials of degree $p-1$ and $\sqrt{x(P_D)}$ is a root of $h_1(X)$ (say) and $\sqrt{x(Q_D)}$ is a root of $h_2(X)$, and hence $L_{D,\mathfrak{p}} \cong \frac{K[X]}{(h_1(X))}$ and $L_{D,\bar{\mathfrak{p}}} \cong \frac{K[X]}{(h_2(X))}$. Since $L_{D,\mathfrak{p}} \cong L_{D,\bar{\mathfrak{p}}}$ which induces complex conjugation when restricted to K , it follows that $h_2(X) = \overline{h_1(X)}$ and consequently, $f_{D,\mathfrak{p}}(X^4) = h_1(X)\overline{h_1(X)} \in K[X]$, which in-turn implies that $f_{D,\mathfrak{p}}(X^4)$ is irreducible of degree $2(p-1)$ over \mathbb{Q} and $L_{D,\mathfrak{p}} \cong \frac{\mathbb{Q}[X]}{(f_{D,\mathfrak{p}}(X^4))}$. Since, $f_{D,\mathfrak{p}}(X^4)$ is irreducible over \mathbb{Q} , it follows that $f_{D,\mathfrak{p}}(X)$ is irreducible over \mathbb{Q} . Again by degree argument we can say that $f_{D,\mathfrak{p}}(X)$ can not be irreducible over K . Using the above argument, it follows that we have an isomorphism $K(x(P_D)^2) \cong K(x(Q_D)^2)$ which induces complex conjugation on K , which in-turn implies that $f_{D,\mathfrak{p}}(X) = h_3(X)\overline{h_3(X)} \in K[X]$ and consequently, $F_{D,\mathfrak{p}} \cong \frac{\mathbb{Q}[X]}{(f_{D,\mathfrak{p}}(X))} \cong \frac{\mathbb{Q}[X]}{(f_{1,\mathfrak{p}}(X))}$. \square

As a consequence of Theorem 2.3 and Theorem 2.4, we obtain

Corollary 2.5. *Suppose that $\omega(E_D) = +1$. Then, $L(E_D, 1) \neq 0 \iff \text{rk}_{\mathbb{Q}} E_D = 0$ and $\text{III}(E_D/\mathbb{Q})$ is finite \iff for all but finitely many primes \mathfrak{p} in K with $\mathfrak{p} \nmid 2D$, $r_{\mathfrak{p}}(E_D) = 0 \iff$ there exists a prime \mathfrak{p} in K with $\mathfrak{p} \nmid 2D$ with $r_{\mathfrak{p}}(E_D) = 0$.*

Proof. First suppose that $L(E_D, 1) \neq 0$, then by the work of Coates-Wiles [CW77] and Rubin [Rub87b], it follows that $\text{rk}_{\mathbb{Q}} E_D = 0$ and $\text{III}(E_D/\mathbb{Q})$ is finite, which in-turn implies that for all but finitely many p , we have $\text{Sel}^p(E_D/\mathbb{Q}) = 0$. Consequently by either Theorem 2.3 or Theorem 2.4, for all but finitely many \mathfrak{p} , we have $r_{\mathfrak{p}}(E_D) = 0$. Now if $r_{\mathfrak{p}}(E_D) = 0$, then by either Theorem 2.3 or Theorem 2.4, we have $s_{\mathfrak{p}}(E_D) = 0$ and, by p -converse theorem of Rubin [Rub91], it follows that $L(E_D, 1) \neq 0$. \square

In a similar fashion, we prove:

Corollary 2.6. *Suppose that $\omega(E_D) = -1$. Then, $L(E_D, 1) = 0$ and $L'(E_D, 1) \neq 0 \iff \text{rk}_{\mathbb{Q}} E_D = 1$ and $\text{III}(E_D/\mathbb{Q})$ is finite \iff for all but finitely many primes \mathfrak{p} in K with $\mathfrak{p} \nmid 2D$ with $r_{\mathfrak{p}}(E_D) \leq 1 \iff$ there exists a prime $\mathfrak{p} \nmid 2D$ in K lying over $p \equiv 1 \pmod{4}$ with $r_{\mathfrak{p}}(E_D) \leq 1$.*

Proof. First suppose that $L(E_D, s)$ has simple zero at $s = 1$, then by the work of Gross-Zagier [GZ86] and Kolyvagin [Kol90], it follows that $\text{rk}_{\mathbb{Q}} E_D = 1$ and $\text{III}(E_D/\mathbb{Q})$ is finite, which in-turn implies that for all but finitely many p , we have $\dim_{F_p} \text{Sel}^p(E_D/\mathbb{Q}) = 1$. Consequently by either Theorem 2.3 or Theorem 2.4, for all but finitely many \mathfrak{p} , we have $r_{\mathfrak{p}}(E_D) \leq 1$. Now if for a prime $\mathfrak{p} \nmid 2D$ in K lying over $p \equiv 1 \pmod{4}$ we have $r_{\mathfrak{p}}(E_D) \leq 1$, then by Theorem 2.4, we have $s_{\mathfrak{p}}(E_D) = 1$ and by p -converse theorem of Burungale-Tian [BT20], it follows that $\text{ord}_{s=1} L(E_D, s) = 1$. \square

Observe that for every D and a prime $\mathfrak{p} \nmid 2D$ in K , we have a chain of sub-fields $K \subset F_{1,\mathfrak{p}} \subsetneq K_{D,\mathfrak{p}} \subsetneq L_{D,\mathfrak{p}}$. Note that we can explicitly describe $K_{D,\mathfrak{p}}$ as $K_{D,\mathfrak{p}} \cong \frac{K[X]}{(f_p(X^2, D))}$ if $p \equiv 3 \pmod{4}$, and $K_{D,\mathfrak{p}} \cong \frac{\mathbb{Q}[X]}{(f_{D,\mathfrak{p}}(X^2))}$ if $\mathfrak{p} \mid p$ with $p \equiv 1 \pmod{4}$. Further note that if $D_1 = D\alpha^4$ for some $\alpha \in K$, $L_{D_1,\mathfrak{p}} = L_{D,\mathfrak{p}}$ and if $D_2 = D\beta^2$ for some $\beta \in K$, then $K_{D_2,\mathfrak{p}} = K_{D,\mathfrak{p}}$. Consequently the following result shows that for quadratic twists of E_D , the variation of p -Selmer rank is controlled by a relative quadratic extension and the root number.

Proposition 2.7. *Let $\mathfrak{p} \nmid 2D$ be a prime in K lying over p . Let M be an integer co-prime to pD and $E_D^M : Y^2 = X^3 - DM^2X$ denote the quadratic twist of E_D by M . Then we have,*

$r_{\mathfrak{p}}(E_D^M) \leq s_p(E_D^M) \leq 1 + r_{\mathfrak{p}}(E_D^M)$, where $r_{\mathfrak{p}}(E_D^M) = \dim_{\mathcal{O}_K/\mathfrak{p}} Cl(L_{DM^2, \mathfrak{p}}/K_{DM^2, \mathfrak{p}})[p](\chi_{DM^2})$. Moreover, we have $K_{DM^2, \mathfrak{p}} \cong K_{D, \mathfrak{p}}$.

In-particular, for congruent number elliptic curves we obtain:

Corollary 2.8. *Let $n \equiv 1, 2, 3 \pmod{8}$ be a square-free integer. Suppose there exists a prime $\mathfrak{p} \nmid 2n$ such that $r_{\mathfrak{p}}(E_1^n) = 0$, then n is not a congruent number.*

In particular, if a square-free integer $n \equiv 1, 2, 3 \pmod{8}$ is a congruent number, then for all primes $\mathfrak{p} \nmid 2n$, we have $p \mid |Cl(L_{n^2, \mathfrak{p}}/K_{1, \mathfrak{p}})|$.

Proof. Since $n \equiv 1, 2, 3 \pmod{8}$, it follows that $\omega(E_1^n) = +1$ and, hence $r_{\mathfrak{p}}(E_1^n) = 0$ implies that $s_p(E_1^n) = 0$, which in-turn implies that $\text{rk}_{\mathbb{Q}} E_1^n = 0$, that is n is not a congruent number. Note that, $n \equiv 1, 2, 3 \pmod{8}$ is a congruent number which implies that $s_p(E_1^n) \geq 2$ or, equivalently $r_p(E_1^n) \geq 1$, which in-turn implies that $p \mid |Cl(L_{n^2, \mathfrak{p}}/K_{1, \mathfrak{p}})|$. \square

Given a non-negative integer k , by $D(k)$ we denote the integer $D(k) = (8k+5)(8k+6)(8k+7)$ and by $D(k)_{sf}$ we denote the square-free part of $D(k)$. One checks that for every $k \geq 0$, $D(k)_{sf}$ is a congruent number and, $D(k)_{sf} \equiv 2 \pmod{8}$. Now for a prime $p > 5$ and $p \equiv 1 \pmod{4}$, one obtains that the infinite set $S_p = \{D(pk)_{sf}\}_{k \geq 0}$ consists of congruent numbers which are co-prime to p . Similarly the infinite set $S_5 = \{D(5k+2)_{sf}, D(5k+4)_{sf}\}_{k \geq 0}$ consists of congruent numbers which are co-prime to 5.

Next observe that, for every integer s , the integer $f(s) = (2s+1)^4 + 24(2s+1)^2 + 16$ is a congruent number, and congruent to 1 modulo 8. For a prime $p \equiv 3 \pmod{4}$, the infinite set $T_p = \{f(ps)_{sf}\}_{s \geq 0}$ consists of congruent numbers which are co-prime to p . Consequently, we obtain the following:

Corollary 2.9. *Given any prime $p \geq 3$, there are infinitely many quadratic extensions $L_{\mathfrak{p}}$ of $K_{1, \mathfrak{p}} := K_{\mathfrak{p}}$ such that p divides the relative class number $L_{\mathfrak{p}}/K_{\mathfrak{p}}$.*

In a similar fashion to Corollary 2.8, we obtain:

Corollary 2.10. *Let $n \equiv 5, 6, 7 \pmod{8}$ be a square-free integer. Suppose that there exists a prime $p \equiv 1 \pmod{4}$ which is co-prime to n such that $r_{\mathfrak{p}}(E_1^n) \leq 1$, then n is a congruent number.*

3. p -SELMER GROUP OF $E_A : Y^2 = X^3 + A$

Let $E := E_1$ denote the elliptic curve $E : Y^2 = X^3 + 1$. For a nonzero integer $A \in \mathbb{Q}^{\times}/(\mathbb{Q}^{\times})^6$, the elliptic curve $E_A : Y^2 = X^3 + A$ is the sextic twist of E by A . The elliptic curve E_A has CM by $\mathbb{Z}[\zeta_3]$, given as $\zeta_3(x, y) = (\zeta_3 x, y)$, and E_A has good reduction at p for $p \nmid 6A$. Hence by Deuring's theorem, for primes $p \nmid 6A$, E_A has good ordinary reduction if and only if $p \equiv 1 \pmod{3}$. By $\omega(E_A)$, we denote the global root number of E_A .

Following lemma provides description of the n -th division polynomial for the elliptic curve E_A .

Lemma 3.1. *For every integer $n \geq 1$, there exists a homogeneous polynomial $f_n(Z, W) = c_n Z^{d_n} + \dots \in \mathbb{Z}[Z, W]$, where $c_n = \begin{cases} n & \text{if } n \text{ is odd,} \\ \frac{n}{2} & \text{if } n \text{ is even,} \end{cases}$ and the degree d_n is given by*

$$d_n = \begin{cases} \frac{n^2-1}{6} & \text{if } n \equiv 1, 5 \pmod{6}, \\ \frac{n^2-3}{6} & \text{if } n \equiv 3 \pmod{6}, \\ \frac{n^2-4}{6} & \text{if } n \equiv 2, 4 \pmod{6}, \\ \frac{n^2-6}{6} & \text{if } n \equiv 0 \pmod{6}, \end{cases}$$

such that

$$\psi_n = \begin{cases} f_n(y^2, A) = f_n(x^3 + A, A) & \text{if } n \equiv 1, 5 \pmod{6}, \\ xf_n(y^2, A) = xf_n(x^3 + A, A) & \text{if } n \equiv 3 \pmod{6}, \\ 2yf_n(y^2, A) = 2yf_n(x^3 + A, A) & \text{if } n \equiv 2, 4 \pmod{6}, \\ 2xyf_n(y^2, A) = 2xyf_n(x^3 + A, A) & \text{if } n \equiv 0 \pmod{6}. \end{cases}$$

Here ψ_n is the n^{th} division polynomial for $E_A : y^2 = x^3 + A$.

Proof. Note $f_1(Z, W) = f_2(Z, W) = 1$, $f_3(Z, W) = 3(Z + 3W)$, $f_4(Z, W) = 2(Z^2 + 18ZW - 27W^2)$. One can prove the result by induction on n and using the recursive relations that define ψ_n . \square

Let $p \equiv 5 \pmod{6}$ be a prime, and for an integer A co-prime to p , define $s_p(E_A) := \dim_{\mathbb{F}_p} \text{Sel}^p(E_A/\mathbb{Q})$. Let $K = \mathbb{Q}(\zeta_3)$ and $L_{A,p} := K(E_A[p])$ be the p -division field of E_A over K , which is of degree $12d_p = 2(p^2 - 1)$. Also we have $G := \text{Gal}(L_{A,p}/K) \cong \frac{\mathbb{Z}}{(p^2-1)\mathbb{Z}}$.

Theorem 3.2. *Let $p \equiv 5 \pmod{6}$ be a prime and A be an integer such that $(p, A) = 1$. Let $F_{A,p}$ be the subfield such that $K := \mathbb{Q}(\zeta_3) \subset F_{A,p} \subset L_{A,p} := K(E_A[p])$ with $\text{Gal}(L_{A,p}/F_{A,p}) \cong \mathbb{Z}/6\mathbb{Z}$. Let χ_A denote the character that gives the action of $G = \text{Gal}(L_{A,p}/K)$ on $E_A[p] \cong \mathbb{F}_{p^2}$ and $r_p(E_A) = \dim_{\mathbb{F}_{p^2}} \text{Cl}(L_{A,p}/F_{A,p})[p](\chi_A)$. Then we have,*

$$r_p(E_A) \leq s_p(E_A) \leq 1 + r_p(E_A),$$

and hence $\omega(E_A)$ and $r_p(E_A)$ uniquely determine $s_p(E_A)$.

Furthermore, let $K_{A,p}$ (resp. $K'_{A,p}$) be the subfield of $L_{A,p}$ such that $F_{A,p} \subset K_{A,p} \subset L_{A,p}$ (resp. $F_{A,p} \subset K'_{A,p} \subset L_{A,p}$) with $\text{Gal}(L_{A,p}/K_{A,p}) \cong \mathbb{Z}/3\mathbb{Z}$ (resp. $\text{Gal}(L_{A,p}/K'_{A,p}) \cong \mathbb{Z}/2\mathbb{Z}$), then

$$K_{A,p} \cong \frac{K[Y]}{(f_p(Y^2, A))}, K'_{A,p} \cong \frac{K[X]}{(f_p(X^3 + A, A))} \text{ and } F_{A,p} = K_{A,p} \cap K'_{A,p} \cong \frac{K[Z]}{(f_p(Z, 1))} \cong F_{1,p}.$$

Here $f_p(Z, W) \in \mathbb{Z}[Z, W]$ is the polynomial defined in Lemma 3.1.

Proof. The proof of the first part is same as in Theorem 2.3.

Let $P_A := (x(P_A), y(P_A)) \in E_A[p]$ be a nontrivial element. We note that $y(P_A)^2$ is a root of the polynomial $f_p(Y, A)$ which is a polynomial in Y of degree $\frac{p^2-1}{6}$. Using an argument similar to the proof of Theorem 2.3, we have $[K(y(P_A)^2) : K] = \frac{p^2-1}{6}$, which proves the irreducibility of $f_p(Y, A)$ over K . Similarly, we also conclude that $f_p(X^3 + A, A)$ is irreducible over K .

Since $f_p(Y, A)$ is a homogeneous polynomial with $y(P_A)^2$ being a root, we have that $\frac{y(P_A)^2}{A}$ is a root of $f_p(Y, 1)$, and hence $F_{A,p} := K(y(P_A)^2) \cong \frac{K[Z]}{(f_p(Z, 1))} \cong F_{1,p}$. Since $[K_{A,p} : K] = \frac{p^2-1}{3}$, and $y(P_A)$ is a root of $f_p(Y^2, A)$, we deduce that $f_p(Y^2, A)$ is irreducible, and hence $K_{A,p} \cong \frac{K[Y]}{(f_p(Y^2, A))} := K(y(P_A))$. Similarly, since $x(P_A)$ is a root of $f_p(X^3 + A, A)$ and $[K(x(P_A)) : K] = \frac{p^2-1}{2}$, we have $K'_{A,p} \cong \frac{K[X]}{(f_p(X^3 + A, A))} := K(x(P_A))$ and $F_{A,p} = K_{A,p} \cap K'_{A,p}$. \square

Next we consider the case $p \equiv 1 \pmod{6}$ and let $p\mathcal{O}_K = \mathfrak{p}\bar{\mathfrak{p}}$ where \mathfrak{p} and $\bar{\mathfrak{p}}$ are the primes in \mathcal{O}_K with $\mathfrak{p} \nmid 6A$ lying above p . We can write $\mathfrak{p} = (\pi)$ where $\pi \in \mathbb{Z}[\zeta_3]$ such that $\pi \equiv 2 \pmod{3}$. We define $\pi_A = -\left(\frac{4A}{\pi}\right)_6 \pi$, where $\left(\frac{\cdot}{\cdot}\right)_6$ denotes the 6th power residue symbol. Now the Grössencharacter associated to E_A is given by $\psi_{E_A/K}(\mathfrak{p}) = \pi_A$ (resp. $\psi_{E_A/K}(\bar{\mathfrak{p}}) = \bar{\pi}_A$). Writing $p = a^2 + ab + b^2$ with $a, b \in \mathbb{N}$ and a odd, we see that $\pi = \zeta_6^k(a - b\zeta_3^i)$ for a particular

choice of $0 \leq k \leq 5$ and $1 \leq i \leq 2$, hence $\pi_A = \zeta_6^t(a - b\zeta_3^i)$ for some $0 \leq t \leq 5$. Define

$$\widetilde{h}_{A,p} = \begin{cases} \frac{\psi_{2b}\psi_a^4 - \psi_{2a}\psi_b^4}{\psi_a\psi_b} = \frac{\psi_{2b}}{\psi_b}\psi_a^3 - \frac{\psi_{2a}}{\psi_2\psi_a}\left(\frac{\psi_b}{\psi_2}\right)^3(\psi_2^2)^2 & \text{if } b - a \text{ is odd,} \\ \frac{\psi_{2b}\psi_a^4 - \psi_{2a}\psi_b^4}{\psi_a\psi_b\psi_2} = \frac{\psi_{2b}}{\psi_2\psi_b}\psi_a^3 - \frac{\psi_{2a}}{\psi_2\psi_a}(\psi_b)^3 & \text{if } b - a \text{ is even,} \end{cases}$$

here ψ_n denotes the n -th division polynomial for the elliptic curve E_A . From Lemma 3.1, it follows that $\widetilde{h}_{A,p}$ is a polynomial with integer co-efficient in Y of degree $t_p = a^2 + b^2 - 1$ (resp. $t_p = a^2 + b^2 - 2$) if b is even (resp. odd) and moreover, $\widetilde{h}_{A,p}(Y) = h_{A,p}(Y^2) \in \mathbb{Z}[Y^2]$. In-particular, we have $\frac{2(p-1)}{3} \leq t_p < p-1$, and the equality holds if and only if $|a-b| \in \{1, 2\}$.

Theorem 3.3. *Let $p \equiv 1 \pmod{6}$ be a prime, co-prime to A . Let $K = \mathbb{Q}(\zeta_3)$ and \mathfrak{p} be a prime in \mathcal{O}_K lying above p . Let $L_{A,\mathfrak{p}} := K(E_A[\pi_A])$ and $F_{A,\mathfrak{p}}$ be the unique subfield of $L_{A,\mathfrak{p}}$ which contains K and satisfies $\text{Gal}(L_{A,\mathfrak{p}}/F_{A,\mathfrak{p}}) \cong \frac{\mathbb{Z}}{6\mathbb{Z}}$. Let χ_A be the character which gives the action of $G := \text{Gal}(L_{A,\mathfrak{p}}/K) \cong \frac{\mathbb{Z}}{(p-1)\mathbb{Z}}$ on $E_A[\pi_A] \cong \mathbb{F}_p$ and $r_{\mathfrak{p}}(E_A) := \dim_{\mathbb{F}_p} \text{Cl}(L_{A,\mathfrak{p}}/F_{A,\mathfrak{p}})[p](\chi_A)$. Then we have*

$$r_{\mathfrak{p}}(E_A) \leq s_{\mathfrak{p}}(E_A) := \dim_{F_{\mathfrak{p}}} \text{Sel}^p(E_A/\mathbb{Q}) \leq 1 + r_{\mathfrak{p}}(E_A),$$

and hence $\omega(E_A)$ and $r_{\mathfrak{p}}(E_A)$ uniquely determine $s_{\mathfrak{p}}(E_A)$.

Furthermore, let $K_{A,\mathfrak{p}}$ (resp. $K'_{A,\mathfrak{p}}$) be the subfield of $L_{A,\mathfrak{p}}$ such that $F_{A,\mathfrak{p}} \subset K_{A,\mathfrak{p}} \subset L_{A,\mathfrak{p}}$ (resp. $F_{A,\mathfrak{p}} \subset K'_{A,\mathfrak{p}} \subset L_{A,\mathfrak{p}}$) with $\text{Gal}(L_{A,\mathfrak{p}}/K_{A,\mathfrak{p}}) \cong \mathbb{Z}/3\mathbb{Z}$ (resp. $\text{Gal}(L_{A,\mathfrak{p}}/K'_{A,\mathfrak{p}}) \cong \mathbb{Z}/2\mathbb{Z}$), then

$$K_{A,\mathfrak{p}} \cong \frac{\mathbb{Q}[Y]}{(f_{A,p}(Y^2))}, K'_{A,\mathfrak{p}} \cong \frac{\mathbb{Q}[X]}{(f_{A,p}(X^3 + A))} \text{ and } F_{A,\mathfrak{p}} = K_{A,\mathfrak{p}} \cap K'_{A,\mathfrak{p}} \cong \frac{\mathbb{Q}[Y]}{(f_{1,p}(Y))} \cong F_{1,\mathfrak{p}},$$

where $f_{A,p}(Y)$ is the irreducible factor (over \mathbb{Q}) of degree $\frac{p-1}{3}$ of the polynomial $h_{A,p}(Y) \in \mathbb{Z}[Y]$.

Proof. The proof of the first part is same as in Theorem 2.3.

Now let $P_A = (x(P_A), y(P_A))$ be a generator of $E_A[\pi_A]$. Since $\pi_A P_A = 0$, it follows that $y(aP_A) = y(bP_A)$ and hence from Lemma 3.1, it follows that $y(P_A)$ is a root of the polynomial $\widetilde{h}_{A,p}(Y) = h_{A,p}(Y^2)$.

Note that $[K(x(P_A), y(P_A)) : K(y(P_A))] \leq 3$ and since $y(P_A)$ is root of a polynomial of degree $t_p < p-1$, it follows that $K_{A,\mathfrak{p}} = K(y(P_A))$ and $h_{A,p}(Y^2)$ has an irreducible factor of degree $\frac{p-1}{3}$ over K . Thus one of the following three situations hold: (a) $h_{A,p}(Y^2)$ has an unique irreducible factor of degree $\frac{p-1}{3}$ over \mathbb{Q} , (b) $h_{A,p}(Y^2)$ has two distinct irreducible factors of degree $\frac{p-1}{3}$ over \mathbb{Q} , or (c) $h_{A,p}(Y^2)$ has an unique irreducible factor of degree $\frac{2(p-1)}{3}$ over \mathbb{Q} which factors as $h(Y)\overline{h(Y)}$ over K . Considering $Q_A = (x(Q_A), y(Q_A))$ a generator of $E_A[\overline{\pi}_A]$ as in Theorem 2.4, and proceeding as in the proof of Theorem 2.4, we conclude that $h_{A,p}(Y^2)$ has an unique irreducible factor $g_{A,p}(Y)$ of degree $\frac{2(p-1)}{3}$ over \mathbb{Q} which factors as $g_{A,p}(Y) = h(Y)\overline{h(Y)}$ over K .

Since the minimal polynomial of $y(P_A)$ has degree $\frac{2(p-1)}{3}$ over \mathbb{Q} , it follows that the minimal polynomial of $y(P_A)^2$ (over \mathbb{Q}) has degree either $\frac{2(p-1)}{3}$ or $\frac{p-1}{3}$. Since $y(P_A)^2$ is a root of $h_{A,p}(Y)$ which has degree less than $\frac{p-1}{2}$, it follows that the minimal polynomial of $y(P_A)^2$ (over \mathbb{Q}) has degree $\frac{p-1}{3}$. Consequently, we see that $h_{A,p}(Y)$ has an irreducible factor $f_{A,p}(Y)$ of degree $\frac{p-1}{3}$ (which is necessarily unique) and $g_{A,p}(Y) = f_{A,p}(Y^2)$.

Since $h_{A,p}(Y)$ is a homogeneous polynomial in Y and A , it follows that $f_{A,p}(Y)$ is a homogeneous polynomial in Y and A . Consequently, $F_{1,\mathfrak{p}} = K(y(P_1)^2) \cong \frac{\mathbb{Q}[Y]}{(f_{1,p}(Y))} = K\left(\frac{y(P_A)^2}{A}\right) = K(y(P_A)^2) := F_{A,\mathfrak{p}}$ and $K_{A,\mathfrak{p}} = K(y(P_A)) \cong \frac{\mathbb{Q}[Y]}{(f_{A,p}(Y^2))}$. Finally, since $y(P_A)$ is a root of

$f_{A,p}(Y^2)$, it follows that $x(P_A)$ is a root of $f_{A,p}(X^3 + A)$ and we have $K'_{A,p} = K(x(P_A)) \cong \frac{\mathbb{Q}[X]}{(f_{A,p}(X^3+A))}$, $F_{A,p} = K_{A,p} \cap K'_{A,p}$. \square

As a consequence of Theorem 3.2 and 3.3, we derive the following results. The proof follows the same argument as in corollaries 2.5 and 2.6.

Corollary 3.4. *Suppose that $\omega(E_A) = +1$. Then $L(E_A, 1) \neq 0 \iff \text{rk}_{\mathbb{Q}} E_A = 0$ and $\text{III}(E_A/\mathbb{Q})$ is finite \iff for all but finitely many primes \mathfrak{p} in K with $\mathfrak{p} \nmid 6A$, $r_{\mathfrak{p}}(E_A) = 0 \iff$ there exists a prime \mathfrak{p} in K with $\mathfrak{p} \nmid 6A$ with $r_{\mathfrak{p}}(E_A) = 0$.*

Corollary 3.5. *Suppose that $\omega(E_A) = -1$. Then, $L(E_A, 1) = 0$ and $L'(E_A, 1) \neq 0 \iff \text{rk}_{\mathbb{Q}} E_A = 1$ and $\text{III}(E_A/\mathbb{Q})$ is finite \iff for all but finitely many primes \mathfrak{p} in K with $\mathfrak{p} \nmid 6A$ with $r_{\mathfrak{p}}(E_A) \leq 1 \iff$ there exists a prime $\mathfrak{p} \nmid 6A$ in K lying over $p \equiv 1 \pmod{6}$ with $r_{\mathfrak{p}}(E_A) \leq 1$.*

Observe that for every A and a prime $\mathfrak{p} \nmid 6A$ in K , we have the following fields diagram:




FIGURE 1

Further note that if $A_1 = A\alpha^4$ for some $\alpha \in K$, $L_{A_1,p} \cong L_{A,p}$, if $A_2 = A\beta^2$ for some $\beta \in K$, then $K_{A_2,p} \cong K_{A,p}$ and if $A_3 = A\gamma^3$ for some $\gamma \in K$, then $K'_{A_3,p} \cong K'_{A,p}$. Consequently the following result shows that for quadratic twists (resp. cubic twists) of E_A , the variation of p -Selmer rank is controlled by a relative quadratic extension (resp. a relative cubic extension) and the root number.

Proposition 3.6. *Let $\mathfrak{p} \nmid 6A$ be a prime in K lying over p .*

- (1) *Let D be a square free integer co-prime to $6A$ and $E_A^D : Y^2 = X^3 + AD^3$ denote the quadratic twist of E_A by D . Then we have, $r_{\mathfrak{p}}(E_A^D) \leq s_{\mathfrak{p}}(E_A^D) \leq 1 + r_{\mathfrak{p}}(E_A^D)$ where $r_{\mathfrak{p}}(E_A^D) = \dim_{\mathcal{O}_K/\mathfrak{p}} \text{Cl}(L_{AD^3,p}/K'_{AD^3,p})[p](\chi_{AD^3})$ with $K'_{AD^3,p} \cong K'_{A,p}$.*
- (2) *Let M be a cube free integer co-prime to $6A$ and $E_A^M : Y^2 = X^3 + AM^2$ denote the cubic twist of E_A by M . Then we have, $r_{\mathfrak{p}}(E_A^M) \leq s_{\mathfrak{p}}(E_A^M) \leq 1 + r_{\mathfrak{p}}(E_A^M)$ where $r_{\mathfrak{p}}(E_A^M) = \dim_{\mathcal{O}_K/\mathfrak{p}} \text{Cl}(L_{AM^2,p}/K_{AM^2,p})[p](\chi_{AM^2})$ with $K_{AM^2,p} \cong K_{A,p}$.*

We give an application of the above result for the rational cube sum problem. We say that a natural number n is a cube sum (resp. not a cube sum) if the equation $X^3 + Y^3 = n$ has (resp. does not have) a rational solution. It is well known that a cube free natural number $n > 2$ is a cube sum iff $\text{rk}_{\mathbb{Q}} E_{-432n^2} > 0$. Using Proposition 3.6(2), we relate cube sum problem to relative cubic class group.

Corollary 3.7. *Suppose $\omega(E_{-432}^n) = 1$ and there exists a prime $\mathfrak{p} \nmid 6n$ such that $r_{\mathfrak{p}}(E_{-432}^n) = 0$, then n is not a rational cube sum.*

In particular, if a cube free integer $n > 2$ is a cube sum for which $\omega(E_{-432}^n) = 1$, then for all primes $\mathfrak{p} \nmid 6n$, we have $p \mid |Cl(L_{-432n^2, \mathfrak{p}}/K_{-432, \mathfrak{p}})|$.

In particular since there are infinitely many primes $\ell \equiv 1 \pmod{9}$ which are cube sum [JMS23], we obtain

Corollary 3.8. *Given any prime $p \geq 5$, there are infinitely many cyclic cubic extensions L_p of $K_{-432, p} \cong K_{1, p} := K_p$ such that p divides the relative class number L_p/K_p .*

Corollary 3.9. *If $\omega(E_{-432}^n) = -1$ and there exists a prime $p \equiv 1 \pmod{6}$, which is co-prime to n such that $r_{\mathfrak{p}}(E_{-432}^n) \leq 1$, then n is a rational cube sum.*

4. CONSTRUCTION OF RELATIVE CLASS GROUPS WITH LARGE p -RANK

In this section we use known families of CM elliptic curves with large rank to construct number fields with large rank p -class group using Theorem 2.3, Theorem 2.4, Theorem 3.2 and Theorem 3.3. At first, following the construction of Kihara [Kih04], we show that given any odd prime p , there are infinitely many elliptic curves E with $j(E) = 1728$ for which p -Selmer rank of E is at least 7.

Proposition 4.1. *Let p be an odd prime. There exists infinitely many (non-isomorphic over \mathbb{Q}) elliptic curves $E_D : Y^2 = X^3 - DX$ with the following properties:*

- (1) E_D has good reduction at p ;
- (2) $\omega(E_D) = -1$;
- (3) $\text{rk}_{\mathbb{Q}} E_D \geq 6$.

Proof. We recall the construction of Kihara [Kih04, Theorem 1] of rank 6 (at-least) elliptic curves over $\mathbb{Q}(r, s, t)$. Define

$$\begin{aligned} u &= \frac{rs(2t^4 - r^4 - s^4)}{t(r^4 - s^4)}, \\ a &= \frac{r^4 + s^4 - t^4 - u^4}{2(r^2s^2 - t^2u^2)}, \\ b &= \frac{(s^2t^2 - r^2u^2)(r^2t^2 - s^2u^2)}{r^2s^2 - t^2u^2}, \end{aligned}$$

then the elliptic curve $\mathcal{E} : Y^2 = X^3 + b(a+1)(a-1)X$ has rank at-least 6 over $\mathbb{Q}(r, s, t)$. A simplification in SAGE yields that $b(a+1)(a-1) = -k(r, s, t)$, where

$$\begin{aligned} k(r, s, t) &= 2^{-8}(r^4 - s^4)^{-6}r^{-4}s^{-4}t^{-12}(r^4 + s^4 + 2r^2t^2)(r^4 + s^4 - 2r^2t^2)(r^4 + s^4 + 2s^2t^2) \\ &\quad \times (r^4 + s^4 - 2s^2t^2)[(r^4 + s^4)^2 + 4r^2s^2t^2(r^2 + s^2 + t^2)] \\ &\quad \times [(r^4 + s^4)^2 + 4r^2s^2t^2(r^2 - s^2 - t^2)][(r^4 + s^4)^2 - 4r^2s^2t^2(r^2 + s^2 - t^2)] \\ &\quad \times [(r^4 + s^4)^2 - 4r^2s^2t^2(r^2 - s^2 + t^2)]. \end{aligned}$$

Setting $D(t) := k(2p, 1, t)$, from the specialization theorem one obtains that for all but finitely many $t \in \mathbb{Q}$, $\text{rk}_{\mathbb{Q}} E_{D(t)} \geq 6$. Further from Faltings theorem [Fal83], it follows that for a fixed $t_0 \in \mathbb{Q}$, $E_{D(t)} \cong E_{D(t_0)}$ for finitely many $t \in \mathbb{Q}$. Thus to complete the proof, it is enough to show that there are infinitely many $t \in \mathbb{Q}$ for which $\nu_p(D(t)) \equiv 0 \pmod{4}$ and $\omega(E_{D(t)}) = -1$. Let us denote by $c = 16p^4 + 1$ and ℓ_i denote a prime $\ell_i \equiv 3 \pmod{4}$ and $\nu_{\ell_i}(c-2) \equiv 1 \pmod{2}$. Since $c-2 \equiv 3 \pmod{4}$, it follows that the number of distinct primes ℓ_i with the

above mentioned properties is odd, say $\ell_1, \dots, \ell_{2k+1}$ and let us define $d = 3\ell_1 \cdots \ell_{2k+1}$. A simple calculation yields that

$$\begin{aligned} D(4cdpt) &\doteq (c-2)^2(1+8\alpha cp^2t^2)(1-8\alpha cp^2t^2)(1+2\alpha ct^2)(1-2\alpha ct^2) \\ &\quad \times [1+16\alpha(4p^2+1)p^2t^2+16\alpha^2c^2p^2t^4][1-16\alpha(4p^2+1)p^2t^2+16\alpha^2c^2p^2t^4] \\ &\quad \times [1+16\alpha(4p^2-1)p^2t^2-16\alpha^2c^2p^2t^4][1-16\alpha(4p^2-1)p^2t^2-16\alpha^2c^2p^2t^4] \end{aligned}$$

where $\alpha = 16p^2d^2$ and by $q_1 \doteq q_2$ we understand that $\frac{q_1}{q_2} \in (\mathbb{Q}^*)^4$. Writing $t = \frac{m+n}{n}$ and by denoting $D(m, n) = D(4cdp\frac{m+n}{n})$, we see that

$$D(m, n) \doteq (c-2)^2 f(m, n),$$

where

$$\begin{aligned} f(x, y) &= (y^2 + 8\alpha cp^2(x+y)^2)(y^2 - 8\alpha cp^2(x+y)^2)(y^2 + 2\alpha c(x+y)^2)(y^2 - 2\alpha c(x+y)^2) \\ &\times [y^4 + 16\alpha(4p^2+1)p^2y^2(x+y)^2 + 16\alpha^2c^2p^2(x+y)^4][y^4 - 16\alpha(4p^2+1)p^2y^2(x+y)^2 + 16\alpha^2c^2p^2(x+y)^4] \\ &\times [y^4 + 16\alpha(4p^2-1)p^2y^2(x+y)^2 - 16\alpha^2c^2p^2(x+y)^4][y^4 - 16\alpha(4p^2-1)p^2y^2(x+y)^2 - 16\alpha^2c^2p^2(x+y)^4]. \end{aligned}$$

Since $f(x, y)$ is a homogeneous polynomial in 2-variables such that each of the irreducible factors has degree at most 4, from [Gre92], it follows that there are infinitely many $(m, n) \in \mathbb{N}^2$ such that $f(m, n)$ is a square-free integer. Note that the fact $f(m, n)$ is square-free necessarily implies that $(n, 2pcdm) = 1$, which in-turn implies that $p \nmid (c-2)^2 f(m, n)$, thus $E_{D(m, n)}$ has good reduction at p .

Finally, we use the formula of root number of E_D , given in [Liv95], to show that for $(m, n) \in \mathbb{N}^2$ for which $f(m, n)$ is square-free, we have $E_{D(m, n)}$ has root number -1 . One can check that if $x > 0$ and $y > 0$, then $f(x, y) > 0$ and hence, we have $\omega_\infty(E_{D(m, n)}) = \text{sgn}(-(c-2)^2 f(m, n)) = -1$. Further, as $(c-2)^2 f(m, n) \equiv 1 \pmod{16}$, it follows that $\omega_2(E_{D(m, n)}) = -1$. The set of primes $\ell \geq 3$ such that $\ell \equiv 3 \pmod{4}$ and $\nu_\ell(D(m, n)) \equiv 2 \pmod{4}$ is precisely $\{\ell_1, \dots, \ell_{2k+1}\}$ and for each ℓ_i , we have $\omega_{\ell_i}(E_{D(m, n)}) = -1$. For all other primes q , we have $\omega_q(E_{D(m, n)}) = 1$, consequently, we obtain $\omega(E_{D(m, n)}) = -1$. \square

Theorem 4.2. (1) *Let p be a prime such that $p \equiv 3 \pmod{4}$. There are infinitely many number fields $L_{n, p}$ of degree 4 over $F_{n, p} \cong F_{1, p}$, such that $\text{rkCl}(L_{n, p})[p] \geq 12 + \text{rkCl}(F_{1, p})[p]$, here $F_{n, p}$ is the field defined in Theorem 2.3.*

(2) *Let p be a prime such that $p \equiv 1 \pmod{4}$. There are infinitely many number fields $L_{n, p}$ of degree 4 over $F_{n, p} \cong F_{1, p}$, such that $\text{rkCl}(L_{n, p})[p] \geq 6 + \text{rkCl}(F_{1, p})[p]$, here $F_{n, p}$ is the field defined in Theorem 2.4.*

Proof. Given an odd prime p , let S_p be the set of fourth-power free integers D such that $p \nmid D$, $\text{rk}_{\mathbb{Q}} E_D \geq 6$ and $\omega(E_D) = -1$. The set S_p is an infinite set by Proposition 4.1.

For $n \in S_p$, define $L_{n, p} := K(E_n[\mathfrak{p}])$, where \mathfrak{p} is a prime in $K := \mathbb{Q}(i)$ lying over p . Since $L_{n, p}$ is a cyclic extension of degree $(N_{K/\mathbb{Q}}(\mathfrak{p}) - 1)$ of K , we have

$$\text{Disc}_{L_{n, p}/\mathbb{Q}} = (\text{Disc}_{K/\mathbb{Q}})^{[L_{n, p}:K]} N_{K/\mathbb{Q}}(\text{Disc}_{L_{n, p}/K}),$$

and from the conductor-discriminant formula, we have $\text{Disc}_{L_{n, p}/K} = \prod_{i=1}^{N_{K/\mathbb{Q}}(\mathfrak{p})-1} \text{cond}(\chi^i)$. It

follows that the primes that ramify in $L_{n, p}/\mathbb{Q}$ are precisely the primes dividing $2np$, since $\text{cond}(\chi) = f\mathfrak{p}$, where f is the conductor of the Hecke character of K associated to E_n (note that conductor of E_n is $|\text{Disc}_{K/\mathbb{Q}}|N_{K/\mathbb{Q}}(f)$). Hence the number fields $L_{n, p}$ are non-isomorphic for $n \in S_p$.

Since for $n \in S_p$, we have $\omega(E_n) = -1$ and hence $s_p(E_n) = \dim_{\mathbb{F}_p} \text{Sel}_p(E_n/\mathbb{Q})$ is odd, and

moreover as $\text{rk}_{\mathbb{Q}}E_n \geq 6$, we see that $s_p(E_n) \geq 7$. Consequently from Theorem 2.3 and Theorem 2.4, it follows that $r_p(E_n) \geq s_p(E_n) - 1 \geq 6$ and hence $Cl(L_{n,p}/F_{n,p})[p]$ contains at least 6-copies of $E_n[p]$. The result follows from the fact that $E_n[p] \cong \mathbb{Z}/p\mathbb{Z}$ for $p \equiv 1 \pmod{4}$ and $E_n[p] \cong \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z}$ for $p \equiv 3 \pmod{4}$. \square

In the following proposition, we make a quantitative estimate of the lower bound of number of number fields $L_{n,p}$ which satisfies the condition of Theorem 4.2. For simplicity of notation, for a number field L , by \mathcal{D}_L , we denote the quantity $\mathcal{D}_L = |\text{Disc}_{L/\mathbb{Q}}|^{\frac{1}{[L:\mathbb{Q}]}}$. For a prime $p \equiv 1 \pmod{4}$, let

$$\mathcal{N}_p(X) = \#\{L \supset F_p \cong F_{1,p} \mid 0 < \mathcal{D}_L < X, \text{Gal}(L/F_p) \cong \frac{\mathbb{Z}}{4\mathbb{Z}}, \text{rk}_p Cl(L/F_p)[p] \geq 6\}$$

and, for a prime $p \equiv 3 \pmod{4}$, let

$$\mathcal{N}_p(X) = \#\{L \supset F_p \cong F_{1,p} \mid 0 < \mathcal{D}_L < X, \text{Gal}(L/F_p) \cong \frac{\mathbb{Z}}{4\mathbb{Z}}, \text{rk}_p Cl(L/F_p)[p] \geq 12\}.$$

Proposition 4.3. *With the notation as above, for any odd prime p , we have*

$$\mathcal{N}_p(X) \gg c_p X^{\frac{1}{12}} \quad \text{where } c_p \neq 0 \text{ and } X \gg 0.$$

Proof. From [Gre92, Theorem (i)], we see that the number of square-free values $f(a, b) = n$ with $|n| = |f(a, b)| \leq X$ is at least $dX^{\frac{1}{12}}$, for a non-zero constant d . Since $\text{Disc}_{L_{n,p}/K} = N_{K/\mathbb{Q}}(\mathfrak{p})^{-1} \prod_{i=1}^n \text{cond}(\chi^i)$, $\text{cond}(\chi) = f\mathfrak{p}$, and, the conductor of $E_n = 4N_{K/\mathbb{Q}}(f) \leq 2^6 n^2$, it follows that $|\text{Disc}_{L_{n,p}/\mathbb{Q}}| \leq (4N_{K/\mathbb{Q}}(f)N_{K/\mathbb{Q}}(\mathfrak{p}))^{N_{K/\mathbb{Q}}(\mathfrak{p})-1} \leq (8n\sqrt{N_{K/\mathbb{Q}}(\mathfrak{p})})^{2(N_{K/\mathbb{Q}}(\mathfrak{p})-1)}$. Consequently, we have that $\mathcal{D}_{L_{n,p}} \leq 8|n|\sqrt{N_{K/\mathbb{Q}}(\mathfrak{p})}$ and the result follows. \square

Remark 4.4. *For a number field F , let $H_0 = F$ and inductively define H_n as the p -Hilbert class field of H_{n-1} and $H_\infty = \cup_{n \geq 0} H_n$, the maximal unramified p -extension of F . By work of [Gv64, Bos92, Haj97], it is known that if $\text{rk}Cl(F)[p] \geq 2 + 2\sqrt{1 + \text{rk}(\mathcal{O}_F^*)}$, then $\Gamma := \text{Gal}(H_\infty/F)$ is infinite, and moreover, Γ is not p -adic analytic and $\text{rk}Cl(H_n)[p]$ is unbounded as n tends to infinity. Unfortunately, it is not known if there are infinitely many number fields F of (fixed) degree d (which is co-prime to p) for which the maximal unramified p -extension is infinite.*

By applying Theorem 4.2(2) (resp. Theorem 4.2(1)) to $p = 3$ (resp. $p = 5$), we see that there are infinitely many number fields F of degree 16 (resp. of degree 8) containing $\mathbb{Q}(\zeta_{12})$ (resp. containing $\mathbb{Q}(i)$) whose maximal unramified 3-extension (resp. maximal unramified 5-extension) H_∞ is an infinite extension of F . Further, in this situation, $\Gamma = \text{Gal}(H_\infty/F)$ is not 3-adic (resp. 5-adic) analytic group and $\lim_{n \rightarrow \infty} \text{rk}Cl(H_n)[3] = \infty$ (resp. $\lim_{n \rightarrow \infty} \text{rk}Cl(H_n)[5] = \infty$).

We can obtain similar results using the large rank Mordell curves constructed by Mestre.

Proposition 4.5. *Let $p \geq 5$ be an odd prime. There exists infinitely many (non-isomorphic over \mathbb{Q}) elliptic curves $E_A : Y^2 = X^3 + A$ with the following properties:*

- (1) E_A has good reduction at p ;
- (2) $\text{rk}_{\mathbb{Q}}E_A \geq 7$.

Proof. We recall the construction of Mestre [Mes92, Theorem 2] of rank at-least 7 elliptic curves over $\mathbb{Q}(v)$ given by

$$\mathcal{E} : Y^2 = X^3 + (v^6 - 1)^2 Q(v^3)^2 P(v^3) := X^3 + A(v),$$

where $Q(v^3)$ and $P(v^3)$ are irreducible polynomials in $\mathbb{Z}[v]$ with $Q(v) = v^9 - v^8 - 8v^7 - 16v^6 + 194v^5 - 26v^4 + 336v^3 + 40v^2 - 11v + 3$ and $P(v) = 3^2 7^2 v^{156} + 2^6 5^3 v^{155} + \dots + 3^{12} 5^2$.

By the specialization theorem of Silverman, we have that for all but finitely many $v_0 \in \mathbb{Q}$, $\text{rk}_{\mathbb{Q}} E_{A(v_0)} \geq 7$. Further, from Faltings theorem, one sees that given a $v_0 \in \mathbb{Q}$, we have $E_{A(v)} \cong E_{A(v_0)}$ for finitely many $v \in \mathbb{Q}$.

For $p = 5$, we choose $v = \frac{1}{5k}$ as k varies over the set of natural numbers, which gives us the required family of elliptic curves. For $p > 5$, we choose $v = pk$ as k varies over the set of natural numbers and obtain the required family of elliptic curves. \square

As a consequence of Proposition 4.5, one proves (in a similar fashion to that of Theorem 4.2):

Theorem 4.6. (1) *Let p be an odd prime such that $p \equiv 5 \pmod{6}$. There are infinitely many number fields $L_{n,p}$ of degree 6 over $F_{n,p} \cong F_{1,p}$, such that $\text{rk} Cl(L_{n,p})[p] \geq 12 + \text{rk} Cl(F_{1,p})[p]$, where $F_{n,p}$ is the field defined in Theorem 3.2.*
(2) *Let p be an odd prime such that $p \equiv 1 \pmod{6}$. There are infinitely many number fields $L_{n,p}$ of degree 6 over $F_{n,p} \cong F_{1,p}$, such that $\text{rk} Cl(L_{n,p})[p] \geq 6 + \text{rk} Cl(F_{1,p})[p]$, where $F_{n,p}$ is the field defined in Theorem 3.3.*

Remark 4.7. *Assuming ABC conjecture, it is possible to give a quantitative lower bound for the number of number fields which satisfy the conditions of Theorem 4.6. More precisely, for a prime $p \equiv 1 \pmod{6}$, define $\mathcal{M}_p(X) = \#\{L \supset F_p \cong F_{1,p} \mid 0 < \mathcal{D}_L < X, \text{Gal}(L/F_p) \cong \frac{\mathbb{Z}}{6\mathbb{Z}}, \text{rk}_p Cl(L/F_p)[p] \geq 6\}$ and for a prime $p \equiv 5 \pmod{6}$, define $\mathcal{M}_p(X) = \#\{L \supset F_p \cong F_{1,p} \mid 0 < \mathcal{D}_L < X, \text{Gal}(L/F_p) \cong \frac{\mathbb{Z}}{6\mathbb{Z}}, \text{rk}_p Cl(L/F_p)[p] \geq 12\}$. Then using [Poo03, Theorem 3.5] and following the proof of Proposition 4.3, one can show that for $X \gg 0$, we have $\mathcal{M}_p(X) \gg c'_p X^{\frac{1}{534}}$ for some non-zero constant c'_p .*

We end the section by proving a conditional result on unboundedness of class group in degree 4 and 6 extensions.

Proposition 4.8. (1) *Fix an odd prime p and assume that $\dim_{\mathbb{F}_p} \text{Sel}^p(E_D/\mathbb{Q})$ is unbounded as D varies in the set of integers co-prime to p , where E_D denotes the elliptic curve $y^2 = x^3 - Dx$. Let F be a number field containing $F_{E_{1,p}}$ of degree prime to p . Then $\text{rk} Cl(L)[p]$ is unbounded as L varies in the extensions of F of degree 4.*
(2) *Fix a prime $p \geq 5$ and assume that $\dim_{\mathbb{F}_p} \text{Sel}^p(E_A/\mathbb{Q})$ is unbounded as A varies in the set of integers co-prime to p , where E_A denotes the elliptic curve $y^2 = x^3 + A$. Let F be a number field containing $F_{E_{1,p}}$ of degree prime to p . Then $\text{rk} Cl(L)[p]$ is unbounded as L varies in the extensions of F of degree 6.*

Proof. We'll prove the result for degree 4 extension. The proof for degree 6 extension is similar. By our assumption $s_p(E_D)$ is unbounded and hence by Theorem 2.3-2.4, $r_p(E_D)$ is unbounded, and hence $\text{rk} Cl(L_{E_D,p})[p]$ is unbounded. This proves the result for $F = F_{E_{1,p}}$. If $F \supset F_{E_{1,p}}$, then for each D , let L_D be a field containing $FL_{E_D,p}$ such that $[L_D : F] = 4$. Since $[FL_{E_D,p} : L_{E_D,p}]$ is co-prime to p , the norm map $N_{FL_{E_D,p}/L_{E_D,p}} : Cl(FL_{E_D,p})_p \rightarrow Cl(L_{E_D,p})_p$ is surjective, unboundedness of $\text{rk} Cl(L_D)[p]$ follows. \square

5. NUMERICAL COMPUTATION

For a prime p and an integer D co-prime to p , using SAGE [S⁺24], we compute $h_p(D) := \dim_{\mathbb{F}_p} Cl(L_{D,p}/F_{D,p})[p]$, here \mathfrak{p} is a prime in $\mathbb{Z}[i]$ lying over p . Since $h_p(D) \geq r_{\mathfrak{p}}(E_D)$, it follows that if $h_p(D) = 0$ (resp. $h_p(D) \leq 1$), then $r_{\mathfrak{p}}(E_D) = 0$ (resp. $r_{\mathfrak{p}}(E_D) \leq 1$), and hence the rank (resp. rank bound) of E_D is computed using Corollary 2.5 and Corollary 2.6 (resp.

Theorem 2.3 and Theorem 2.4). If the computation of $h_p(D)$ was done assuming GRH in SAGE, we have marked the corresponding entry in the table by * and if the resulting rank (or rank bound) depends on $h_p(D)$ which was computed assuming GRH, we put a * on the rank to specify that it depends on GRH.

Table for $E_D : Y^2 = X^3 - DX$ with $-50 \leq D \leq 50$

D	$\omega(E_D)$	$h_3(D)$	$h_5(D)$	$h_{13}(D)$	$\text{rk}_{\mathbb{Q}} E_D$	$\omega(E_{-D})$	$h_3(-D)$	$h_5(-D)$	$h_{13}(-D)$	$\text{rk}_{\mathbb{Q}} E_{-D}$
1	1	0	0	0	0	1	0	0	0	0
2	-1	0	0	0	1	1	0	0	0	0
3	1	NA	0	0	0	-1	NA	0	0	1
5	-1	0	NA	0	1	-1	2	NA	0	1
6	-1	NA	0	0	1	1	NA	0	0	0
7	-1	0	0	0	1	1	2	0	0	0
9	1	NA	0	0	0	-1	NA	0	0	1
10	-1	2	NA	0	1	1	0	NA	0	0
11	1	0	0	0	0	1	2	0	0	0
13	1	2	0	NA	0	-1	0	0	NA	1
14	-1	0	0	0	1	1	2	1	1	≤ 2
15	-1	NA	NA	0	1	-1	NA	NA	0	1
17	1	2	1	1*	≤ 2	1	0	0	0	0
18	1	NA	0	0	0	-1	NA	1	0	1
19	1	0	0	0	0	-1	0	0	0	1
21	-1	NA	0	0*	1	-1	NA	0	0	1
22	-1	0	0	0	1	1	0	0	0	0
23	-1	4	0	0*	1	1	0	0	0	0
25	-1	0	NA	0	1	1	0	NA	0	0
26	-1	2	0	NA	1	1	2	0	NA	0
27	1	NA	0	0	0	1	NA	0	0	0
29	1	0	0	0	0	-1	0*	0	0	1
30	-1	NA	NA	0*	1*	1	NA	NA	0	0
31	-1	0	0	0	1	-1	2*	0	0	1
33	1	NA	0	0	0	1	NA	1	1	≤ 2
34	-1	4*	0	1*	1	1	2*	1	1*	≤ 2
35	1	2*	NA	0	0	-1	0*	NA	0*	1*
37	-1	0*	1	0	1	-1	2*	0	0*	1
38	-1	2*	0	0	1	1	2*	0	0*	0
39	-1	NA	0	NA	1	1	NA	1	NA	≤ 2
41	-1	0*	0	0	1	1	0*	0	0*	0
42	-1	NA	0	0*	1	1	NA	0	0*	0
43	1	2*	0	0*	0	1	0*	0	0*	0
45	-1	NA	NA	0*	1*	1	NA	NA	0*	0*
46	-1	0*	0	1*	1	1	2*	1	1*	≤ 2
47	-1	0*	0	0*	1	-1	0*	0	0*	1
49	-1	0*	0	1*	1	-1	0*	0	0*	1
50	-1	0*	NA	0*	1*	1	2*	NA	0*	0*

Using SAGE, we compute $h_p(A) := \dim_{\mathbb{F}_p} Cl(L_{A,p}/F_{A,p})[p]$ (provided $p \nmid A$). Since $h_p(A) \geq r_p(E_A)$, the rank (resp. rank bound) of E_A is computed using Corollary 3.4 and Corollary 3.5 (resp. Theorem 3.2 and Theorem 3.3). Outcomes dependent on GRH are marked with *.

Table for $E_A : Y^2 = X^3 + A$ with $-50 \leq A \leq 50$

A	$\omega(E_A)$	$h_7(A)$	$h_{13}(A)$	$\text{rk}_{\mathbb{Q}} E_A$	$\omega(E_{-A})$	$h_7(-A)$	$h_{13}(-A)$	$\text{rk}_{\mathbb{Q}} E_{-A}$
1	1	0	0	0	1	0	0	0
2	-1	0	0*	1	-1	0	0*	1
3	-1	1	0*	1	1	0	0*	0
4	1	0	0	0	-1	0	0*	1
5	-1	0	0*	1	1	0	0*	0
6	1	0	0*	0	1	0	0*	0
7	1	NA	0*	0*	-1	NA	0*	1*
8	-1	0	0*	1	1	0	0*	0
9	-1	0	0*	1	1	0	0*	0
10	-1	0	0*	1	1	0	0*	0
11	-1	0	0*	1	1	1*	1*	$\leq 2^*$
12	-1	1	0*	1	1	0	0*	0
13	1	0	NA	0	-1	0*	NA	1*
14	1	NA	0*	0*	1	NA	0*	0*
15	1	1	1*	≤ 2	-1	0*	0*	1*
16	1	0	0	0	1	0	0*	0
17	1	1	1*	≤ 2	1	0*	0*	0*
18	-1	0	0*	1	-1	0*	0*	1*
19	-1	0*	0*	1*	-1	0*	0*	1*
20	1	0	0*	0	-1	0*	0*	1*
21	1	NA	0*	0*	-1	NA	0*	1*
22	-1	0	0*	1	-1	0*	0*	1*
23	1	1*	0*	0*	-1	1*	0*	1*
24	1	1	1*	$\leq 2^*$	1	0*	0*	0*
25	1	0	0*	0	-1	0*	0*	1*
26	-1	1*	NA	1*	1	1*	NA	$\leq 2^*$
28	-1	NA	0*	1*	-1	NA	0*	1*
29	1	0*	0*	0*	-1	0*	0*	1*
30	-1	0*	0*	1*	-1	0*	1*	1*
31	-1	0*	0*	1*	1	0*	0*	0*
32	1	0	0*	0*	1	0	0*	0*
33	-1	0*	1*	1*	1	0*	0*	0*
34	1	0*	0*	0*	1	0*	0*	0*
35	-1	NA	0*	1*	-1	NA	0*	1*
36	-1	0	0*	1	1	0	0*	0
37	1	1*	1*	$\leq 2^*$	1	0*	0*	0*
38	-1	0*	0*	1*	-1	0*	0*	1*
39	-1	0*	NA	1*	1	1*	NA	$\leq 2^*$
40	-1	0*	0*	1*	-1	0*	0*	1*
41	-1	0*	0*	1*	1	0*	0*	0*
42	1	NA	0*	0*	1	NA	0*	0*
43	1	2*	1*	$\leq 2^*$	-1	1*	0*	1*
44	-1	0*	0*	1*	-1	0*	0*	1*
45	1	0*	0*	0*	-1	0*	0*	1*
46	-1	0*	0*	1*	1	0*	0*	0*
47	-1	0*	0*	1*	1	1*	1*	$\leq 2^*$
48	-1	0	0*	1	-1	0*	0*	1*
49	1	NA	0*	0*	-1	NA	0*	1*
50	-1	0*	1*	1*	-1	0*	0*	1*

In the following table we compute rank of congruent number elliptic curve using the p -rank of relative quadratic class group $Cl(L_{n^2,p}/K_{n^2,p})$, which we denote by $h_p(n)$. As before, * denotes that the outcome is correct provided GRH is true.

Table for $E_{n^2} : Y^2 = X^3 - n^2X$ with $n < 50$

n	$\omega(E_{n^2})$	$h_3(n)$	$h_5(n)$	$h_{13}(n)$	$rk_{\mathbb{Q}} E_{n^2}$
1	1	0	0	0	0
2	1	0	0	0	0
3	1	NA	0	0	0
5	-1	0	NA	0	1
6	-1	NA	0	0	1
7	-1	0	0	1	1
10	1	0	NA	0	0
11	1	0	0	0	0
13	-1	0	0	NA	1
14	-1	0	0	0	1
15	-1	NA	NA	0	1
17	1	0	0	0	0
19	1	0	0	0	0
21	-1	NA	1	0	1
22	-1	0	0	1	1
23	-1	0	0	0	1

n	$\omega(E_{n^2})$	$h_3(n)$	$h_5(n)$	$h_{13}(n)$	$rk_{\mathbb{Q}} E_{n^2}$
26	1	2	0	NA	0
29	-1	2	0	0	1
30	-1	NA	NA	0	1
31	-1	2	0	0	1
33	1	NA	0	0	0
34	1	2	1	1	≤ 2
35	1	0	NA	0	0
37	-1	0	0	0	1
38	-1	2	1	0	1
39	-1	NA	0	NA	1
41	1	2	1	1	≤ 2
42	1	NA	0	0	0
43	1	2	0	0*	0
46	-1	2	0	0*	1
47	-1	2	0	0*	1

In the following table we compute rank of cube sum elliptic curve using the p -rank of relative cubic class group $Cl(L_{-432n^2,p}/K_{-432n^2,p})$, which we denote by $h_p(n)$. As before, * denotes that the outcome is correct provided GRH is true.

Table for $E_{-432n^2} : Y^2 = X^3 - 432n^2$ with $n \leq 50$

n	$\omega(E_{-432n^2})$	$h_7(n)$	$h_{13}(n)$	$rk_{\mathbb{Q}} E_{-432n^2}$
1	1	0	0	0
2	1	0	0	0
3	1	0	0*	0
4	1	0	0	0
5	1	0	0*	0
6	-1	0	0*	1
7	-1	NA	0*	1*
9	-1	0	0*	1
10	1	0	0*	0
11	1	0	0*	0
12	-1	0	0*	1
13	-1	0	NA	1
14	1	NA	0*	0*
15	-1	1	0*	1
17	-1	0	0*	1
18	1	0	0*	0
19	1	1	1*	≤ 2
20	-1	0	0*	1
21	1	NA	0*	0*
22	-1	0	0*	1
23	1	0	0*	0
25	1	0	0*	0

n	$\omega(E_{-432n^2})$	$h_7(n)$	$h_{13}(n)$	$rk_{\mathbb{Q}} E_{-432n^2}$
26	-1	0	NA	1
28	-1	NA	0*	1*
29	1	0	0*	0
30	1	1	1*	≤ 2
31	-1	0	0*	1
33	-1	0	0*	1
34	-1	0	0*	1
35	-1	NA	0*	1*
36	1	0	0*	0
37	1	1	1*	≤ 2
38	1	0	0*	0
39	1	0	NA	0
41	1	0	0*	0
42	-1	NA	0*	1*
43	-1	0	0*	1
44	1	0	0*	0
45	1	0	0*	0
46	1	0	0*	0
47	1	0	0*	0
49	-1	NA	0*	1*
50	-1	0	0*	1

REFERENCES

- [BG18] Y. Bilu and J. Gillibert, *Chevalley-weil theorem and subgroups of class groups*, Israel Journal of Mathematics **226** (2018), no. 2, 927–956.
- [Bos92] Nigel Boston, *Some cases of the Fontaine-Mazur conjecture*, J. Number Theory **42** (1992), no. 3, 285–291.
- [BT20] A. A. Burungale and Y. Tian, *p -converse to a theorem of Gross-Zagier*, Kolyvagin and Rubin, Invent. Math. **220** (2020), no. 1, 211–253.
- [Cas91] J. W. S. Cassels, *Lectures on elliptic curves*, London Mathematical Society Student Texts, vol. 24, Cambridge University Press, Cambridge, 1991.
- [CW77] J. Coates and A. Wiles, *On the conjecture of Birch and Swinnerton-Dyer*, Invent. Math. **39** (1977), no. 3, 223–251.
- [Fal83] G. Faltings, *Endlichkeitssätze für abelsche varietäten über zahlkörpern*, Inventiones Mathematicae **73** (1983), no. 3, 349–366.
- [Gre92] G. Greaves, *Power-free values of binary forms*, Quarterly Journal of Mathematics, Oxford Series (2) **43** (1992), no. 169, 45–65.
- [Gv64] E. S. Golod and I. R. Šafarevič, *On the class field tower*, Izv. Akad. Nauk SSSR Ser. Mat. **28** (1964), 261–272.
- [GZ86] B. H. Gross and D. B. Zagier, *Heegner points and derivatives of L -series*, Invent. Math. **84** (1986), no. 2, 225–320.
- [Haj97] Farshid Hajir, *On the growth of p -class groups in p -class field towers*, J. Algebra **188** (1997), no. 1, 256–271.
- [JMS22] Somnath Jha, Dipramit Majumdar, and Pratiksha Shingavekar, *3-Selmer group, ideal class groups and cube sum problem*, 2022, arXiv preprint arXiv:2207.12487.
- [JMS23] Somnath Jha, Dipramit Majumdar, and B. Sury, *Binary cubic forms and rational cube sum problem*, 2023, arXiv preprint arXiv:2301.06970.
- [Kih04] S. Kihara, *On the rank of the elliptic curve $y^2 = x^3 + kx$. ii*, Proc. Japan Acad. Ser. A Math. Sci. **80** (2004), no. 4, 24–25.
- [Kol90] V. A. Kolyvagin, *Euler systems*, The Grothendieck Festschrift, Vol. II, Progr. Math., vol. 87, Birkhäuser Boston, Boston, MA, 1990, pp. 435–483.
- [Li19] Chao Li, *2-Selmer groups, 2-class groups and rational points on elliptic curves*, Trans. Amer. Math. Soc. **371** (2019), no. 7, 4631–4653.
- [Liv95] E. Liverance, *A formula for the root number of a family of elliptic curves*, Journal of Number Theory **51** (1995), no. 2, 288–305.
- [Mes92] J.-F. Mestre, *Rang de courbes elliptiques d’invariant donné*, C. R. Acad. Sci. Paris Sér. I Math. **314** (1992), no. 12, 919–922.
- [Poo03] Bjorn Poonen, *Squarefree values of multivariable polynomials*, Duke Mathematical Journal **118** (2003), no. 2, 353–373.
- [PS21] Dipendra Prasad and Sudhanshu Shekhar, *Relating the Tate-Shafarevich group of an elliptic curve with the class group*, Pacific J. Math. **312** (2021), no. 1, 203–218.
- [Rub87a] K. Rubin, *Descents on elliptic curves with complex multiplication*, Séminaire de Théorie des Nombres, Paris 1985–86, Progr. Math., vol. 71, Birkhäuser Boston, Boston, MA, 1987, pp. 165–173.
- [Rub87b] ———, *Tate-Shafarevich groups and L -functions of elliptic curves with complex multiplication*, Invent. Math. **89** (1987), no. 3, 527–559.
- [Rub91] ———, *The “main conjectures” of Iwasawa theory for imaginary quadratic fields*, Invent. Math. **103** (1991), no. 1, 25–68.
- [S+24] W. A. Stein et al., *Sage Mathematics Software (Version 9.3)*, The Sage Group, 2024, <http://www.sagemath.org>.
- [Sch96] Edward F. Schaefer, *Class groups and Selmer groups*, J. Number Theory **56** (1996), no. 1, 79–114.
- [Č15] K. Česnavičius, *Selmer groups and class groups*, Compos. Math. **151** (2015), no. 3, 416–434.

(De) DEPARTMENT OF MATHEMATICS, IIT MADRAS, INDIA
Email address: debajyotide20@gmail.com

(Majumdar) DEPARTMENT OF MATHEMATICS, IIT MADRAS, INDIA
Email address: dipramit@iitm.ac.in

(Mondal) DEPARTMENT OF MATHEMATICS, IIT MADRAS, INDIA
Email address: sudipa.mondal123@gmail.com