Interplay of damage and repair in the control of epithelial tissue integrity in response to cyclic loading

Eleni Papafilippou¹, Lucia Baldauf², Guillaume Charras^{2,3,4}, Alexandre J. Kabla¹, and Aessandra Bonfanti⁵

¹Department of Engineering, University of Cambridge, Cambridge, UK ²London Centre for Nanotechnology, University College London, London, UK ³Institute for the Physics of Living Systems, University College London, London, UK ⁴Department of Cell and Developmental Biology, University College London, London, UK ⁵Department of Civil and Environmental Engineering, Politecnico di Milano, Milan, Italy

Keywords: epithelial tissues, remodelling, repair, fatigue, cyclic stretch

Abstract

Epithelial tissues are continuously exposed to cyclic stretch *in vivo*. Physiological stretching has been found to regulate soft tissue function at the molecular, cellular, and tissue scales, allowing tissues to preserve their homeostasis and adapt to challenges. In contrast, dysregulated or pathological stretching can induce damage and tissue fragilisation. Many mechanisms have been described for the repair of epithelial tissues across a range of time-scales. In this review, we present the timescales of (i) physiological cyclic loading regimes, (ii) strain-regulated remodelling and damage accumulation, and (iii) repair mechanisms in epithelial tissues. We discuss how fatigue in biological tissues differs from synthetic materials, in that damage can be partially or fully reversed by repair mechanisms acting on timescales shorter than cyclic loading. We highlight that timescales are critical to understanding the interplay between damage and repair in tissues that experience cyclic loading, opening up new avenues for exploring soft tissue homeostasis.

Introduction

During an average lifetime, the human heart epithelium undergoes more than 2.5 billion contraction and expansion cycles [1], far surpassing the durability of most synthetic polymers. Epithelial tissues are twodimensional sheets which line the internal structures and cavities of the body and are often monolayered, but can also be stratified or pseudo-stratified. They possess remarkable resilience to deformations and function as physical barriers. In dynamic mechanical environments, epithelia are challenged by intrinsic and extrinsic stresses, to which they respond over different length- and time-scales, from protein recruitment, to signalling and metabolic adaptation. The mechanical durability of epithelial monolayers raises important questions about the molecular and biophysical mechanisms underlying their resilience and is therefore an emerging research focus within mechanobiology. Here, we review the homeostatic and remodelling processes that preserve epithelial tissue integrity, focussing on the intrinsic timescales of epithelial remodelling and repair mechanisms. We propose that the integrity of epithelial tissues that undergo cyclic deformation requires a subtle balance between the timescales of damage and self-healing mechanisms and the period of the strain cycles.

1 Epithelia commonly experience cyclic loading

Cyclic loading is ubiquitous in normal physiology and manifests as periodic stretch and relaxation cycles in many organs. For instance, the heart undergoes oscillatory stress during contraction cycles and the intestinal lumen undergoes rhythmic deformation during peristalsis. Similarly, the bladder and lungs undergo periodic



Figure 1: a) Stress (kPa) and b) strain (%) amplitudes over frequency (Hz) for healthy (left column) and diseased (right column) epithelial tissues within different organs. Supporting data with exact values and references can be found in Table 1. Some icons have been adapted from NIAID NIH BIOART Source (bioart.niaid.nih.gov/bioart).

filling and voiding during physiological fluid flow. Although cyclic loading is essential for barrier function, excessive or abnormal loading can lead to tissue fatigue, micro-damage, and rupture. While some damage is a direct consequence of abnormal loading, other damage is indirect, arising through perturbation of processes normally regulated by cyclic loading. Indeed, cyclic deformation during normal breathing (25% strain) regulates molecular processes involved in cytoskeletal regulation and barrier function [2], while pathological strains (>37% in mechanical ventilation) can lead to altered junctional protein expression and paracellular gap formation, compromising barrier function [3]. Excessive or dysregulated cyclic stretch, characterised by perturbed stretch amplitude and frequency, has also been linked to pathologies such as heart valve prolapse, acute lung injury, and inflammatory bowel disease [4, 5, 6]. Despite its importance, the relationship between specific cyclic loading regimes and tissue integrity has not been systematically explored.

To provide an overview of common physiological and pathological cyclic deformations, we summarise data from studies which report cyclic loading regimes either in terms of stress amplitude vs frequency or strain amplitude vs frequency (Figure 1, Table 1). Stress measurements in epithelial tissues are often challenging to interpret due to the difficulty in isolating the contribution of the epithelium from that of coupled tissues, such as the stroma (comprising extracellular matrix and stromal cells) or underlying muscle layers. As a result, stress experienced by the epithelium alone is rarely reported. In contrast, strain offers a more reliable metric, as all coupled layers of the tissue deform cohesively to the same extent under mechanical loading. Cyclic strain characterisation is therefore a valuable metric for studying strain-controlled remodelling and self-healing in epithelia.

Tissue	Stress Amplitude	Frequency	Strain Amplitude	Icon
lung	1kPa [7]	0.2Hz [8]	30%[9] [MV 84% [5]]	A CONTRACTOR
aorta	100kPa [10, 11]	1Hz [12]	10-15% [1]	Ť
heart	-	1Hz [12]	16% [13] [ischemia 12% [13]]	
carotid	15kPa [14]	1Hz [12]	$\frac{10\% \ [15]}{[AS \ 8\% [15]]}$	1 ml
capillaries	35kPa [16] [thrombosis 29kPa [17]]	1Hz [12]	25% [18]	- TOP
heart valve	600kPa AV, 100kPa PV [1 9]	1Hz [12]	10% [20] [MVP 12% [4]]	
colon	50kPa RPC [6], 130kPa GMC [6] [IBS-D GMC 250kPa [6]] [IBS-C GMC <65kPa [6]]	0.02Hz slow RPC [6] 0.1Hz fast RPC [6] 10^{-4} Hz GMC [6] [IBS-D GMC > 10^{-4} Hz [6]] [IBS-C GMC < $5 \cdot 10^{-5}$ Hz [6]]	50% [21] [obstruction >50% [21]]	Ą
skin - walking	5.5kPa [22]	2Hz [23]	10% [24]	Ļ
skin - running	-	3Hz [23]	43% [24]	
skin - knee flexion	-	2Hz [23]	45% [25]	
stomach	30kPa [26]	0.05Hz [27] [FP 0.03Hz [28]]	30% [29] [FP 48%[29]]	25
small intestines	2.4kPa [30]	0.2Hz [31]	20% [31]	
bladder	51kPa [32]	10^{-4} Hz [33]	90% [32]	
uterus	1kPa [34] [endo. 3kPa [34]]	0.015Hz [34] [endo. 0.03Hz [34]]	$16\% \ [35]$	
ureter	3kPa [36]	0.05Hz [37]	$35\% \ [38]$	
esophageal	25kPa [39] [dysphagia 60kPa [39]]	0.1Hz [40] [achalasia 0.2Hz [40]]	$50\% \ [41, \ 42]$	

Table 1: Stress and strain cyclic loading parameters for organs lined by epithelia in health and disease.

Nomenclature: MV=Mechanical ventilation, AS=Ankylosing Spondylitis, AV=aortic valve, PV=pulmonary valve, MVP=Mitral valve prolapse, RPC=Rhythmic phasic contractions, GMC=giant migrating contractions, FP=functional dyspepsia, endo.=endometriosis.

2 Mechanical response of epithelia to strain

In response to deformation, the stress that epithelia experience depends on their rheology, which has been shown to be nonlinear and highly dependent on the magnitude and rate of strain. All levels from single molecular crosslinkers [43], to individual filaments [44] and filament networks [45] contribute unique time-dependent behaviours (Figure 2). Non-linear behaviours such as strain stiffening may represent an emergent adaptation of the tissue to resist sudden and transient increases in loading, such as shocks. Notably, epithelia can stiffen under strain, depending not only on the magnitude, but also the rate of deformation [46]**. If dysregulated loading is sustained, cells within the epithelium adapt by regulating both the cytoskeleton and junctional proteins, which act together to resist mechanical stresses and preserve barrier function. Adaptation can take place through post-translational modifications, protein recruitment (second to minute time-scales), or transcriptional regulation (longer time-scales) [47, 48] (reviewed in [49]**). Depending on the time-scale of cyclic loading distinct biophysical phenomena may contribute to damage and repair.

2.1 The dynamic actin cytoskeleton makes tissues resilient

The actin cytoskeleton, connected across cells via adherens junctions (AJs), forms a supracellular network that plays a key role in tissue mechanics. In AJs, E-cadherin binds to neighbouring cells through its extracellular domain and interfaces with actin through binding its intracellular domain to beta- and alpha-catenin. Both the actin cytoskeleton and AJs respond to and are regulated by mechanical tension. Actin networks exhibit a linear tension-strain response under small deformations and are regulated by mechanical forces [50]. At longer timescales, actin networks behave as viscous fluids due to turnover of crosslinkers and actin filaments $(\leq 1 \min, [51])$ (Figure 2ai), leading to stress relaxation in epithelial tissues [52]. In immediate response to load, several AJ proteins exhibit catch-bonding characteristics, where molecular bonds become more stable under a moderate load (Figure 2aii). E-cadherin trans-dimers, which bind adjacent cells to one another, are most stable under loads of around 30 pN [53]. The bond between actin filaments and alpha-catenin is stabilized 20-fold under 10 pN molecular load [54]. Similar behaviour also emerges in other intercellular adhesion proteins associated with the actin cytoskeleton: PDZ-domains from nectin-1 and JAM-A were recently shown to form catch bonds with a fadin [55]*. Interestingly, recent structural evidence suggests that nectin rather than E-cadherin may be the primary junctional protein involved in force transmission through the supracellular actin cytoskeleton in mature gut epithelia [56]**. Catch-bonding enables cells to rapidly strengthen intercellular junctions by extending bond lifetimes and forming new links, buffering transient tension. Vinculin recruitment reinforces cadherin-catenin complexes for added stability. Under prolonged loading, additional complexes are recruited, re-distributing the load to reduce each bond's lifetime and restore the proportion of bound links to homeostatic levels (Figure 2aii).

2.2 Intermediate filaments act as shock absorbers and provide mechanical memory

Intermediate filaments (IFs) and the desmosomes which connect them between neighbouring cells form a second supracellular network within tissues. Although evidence is conflicting on whether desmosomal proteins bear load while the epithelium is at rest [57, 58], the emergent properties of the keratin network connected by desmosomes dominate the mechanical response at large deformations $[46]^{**}$. At rest, bundles of keratin filaments appear wavy. When the tissue is stretched they gradually straighten and become progressively load bearing under large deformation [59], leading to a multi-stage strain stiffening response (Figure 2aiii). To capture this, recent modelling studies incorporate nonlinear components for the rate-dependent strain-stiffening of supracellular cytoskeletal networks, such as a 'slack' representing the waviness or 'degree of entanglement' of keratin bundles [46], acting as an emergent topological safety net [60]. Under sustained stretch, individual filaments can slide past one another [61]**, leading to intra-bundle flow and energy dissipation over tens of minutes. This mechanism may confer a long-term mechanical memory of the epithelium's deformation history. Mechanotransductory responses associated with keratin IFs are less well studied. Some pathways increase interfacing of the keratin network with the actin cytoskeleton via the cytolinker plectin in response to tension [62][63]**. Tensin 4 is recruited to keratin fibres subjected to tension and remains localised for minutes after deformation release [64], suggesting a role in adapting to cyclical loads with periods shorter than minutes. At longer timescales, keratins may indirectly regulate transcriptional responses by shielding the nucleus from deformation [65]. Adaptations of desmosomal junctions to stress may also contribute to



Figure 2: Response to deformation across length- and time-scales in epithelial monolayers. For each row, the left column represents the organisation before stretch, the middle column immediately after application of stretch, and the right column after prolonged stretch. a) Mechanisms of plasticity in cellular and molecular components in response to a step loading in strain. i) Actin filaments become stretched but may adopt a long-term new steady state with an increased number of actin filaments, ii) Intercellular junction proteins, such as alpha-catenin and E-cadherin, behave as catch bonds, becoming stabilised upon loading. Alpha-catenin unfolds to recruit vinculin and redistribute the applied load to the cadherin-catenin force chain. At longer time-scales, protein recruitment will return cadherin-catenin complexes to their homeostatic load, iii) Keratin filament bundles become loaded in response to stretch but dissipate stress by inter-filament siding in prolonged loading. b) Time-scales associated with emergent mechanisms and active feedback for cytoskeletal reorganisation, turnover, and protein synthesis. t_{lifetime} refers to the functional stabilization time-scale for the mechanisms of plasticity

mechanical memory, but the extent and timescale of mechanoresponsive desmosome remodelling remain less well-characterized [66].

2.3 Non-physiological strain leads to damage accumulation in living materials

Damage in tissues can result either from a single catastrophic high-amplitude deformation or from accumulation of defects due to repeated cycles of supraphysiological strain. We can apply the concept of fatigue from material engineering, where controlled cyclic loading is used to probe a material's endurance limits, energy dissipation, and failure mechanisms. In classic engineering materials, fatigue leads to cumulative damage - a progressive and irreversible process where repeated stress cycles gradually degrade the material's structure [67]. Even when individual cycles do not cause immediate macroscopic damage, microscopic lesions accumulate incrementally until failure. Living tissues, by contrast, possess repair mechanisms that will be discussed in Section 3.

In soft tissues, damage accumulation can either directly result from rupture of intercellular bonds and ECM adhesions, or indirectly arise from active responses of cells to strain. Cyclic stretch testing in mouse alveolar epithelial layers reveals that intercellular gaps form under hyperinflation (80% change in lung capacity), leading to a loss of epithelial barrier integrity. Intercellular junctions are destabilised due to downregulation of p120-catenin which induces paracellular gap formation [68]. Similarly, large amplitude uniaxial cyclic stretch (70%) of the oesophageal mucosa demonstrated hysteresis, permanent deformation, stress softening, and occasionally rupture [69]. Finally, mutations in keratin 5 or keratin 14 in epidermolysis bullosa simplex, have been linked to weakening of the epithelial sheet integrity upon shear stress [70].

Geometry is emerging as a critical factor in destabilising epithelia: the orientation of forces acting on junctional complexes is likely a major determinant in how junctions respond to load. For instance, tensile stress on AJs increases both E-cadherin and vinculin recruitment, whereas shear stress leads to loss of E-cadherin from AJs in the Drosophila germband $[71][72]^{**}$. Such geometry-dependent stability of junctions may explain why disease-related changes in junctional actin organisation lead to tissue fragility. For instance, pathological expression of the actin-bundling protein fascin-1 modifies the geometry of AJs by changing the curvature and alignment of junctional interfaces. This reduces junction stability and enhances cellular protrusive activity, leading to elongated shapes and polarization, which can induce cancer cell migration and invasion $[73]^*$.

Further detailed studies of the structures of intercellular contacts in epithelia are essential for improved understanding of how different cycling regimes affect epithelial stability. Such studies are becoming increasingly feasible thanks to recent advancements in the imaging of intercellular junctions (reviewed in $[74]^*$).

3 Repair mechanisms in epithelial tissues

Living materials are particularly intriguing when considering fatigue: damage can often be repaired through healing and remodelling processes. In this section, we examine mechanisms of repair at the molecular- and cellular-scale.

3.1 Re-binding of intercellular linkers

Repair at the molecular scale can either take the form of (i) gap or micro-lesion closure (when two surfaces have fully separated) or (ii) return to homeostatic intercellular bond density, where no significant intracellular gap has formed but the number of links connecting two cells has decreased due to force application.

A well-established model to study the biophysics of intercellular adhesion was developed by Bell [75]. A cell junction is defined as two surfaces connected by reversible links which can be bound or unbound. Each link has a constant binding probability k_{on} , and an unbinding probability that depends on force, $k_{off} = k_{off,0}e^{\frac{f}{t_0}}$. Rupture occurs when all linkers connecting the two surfaces are unbound. This model enables the exploration of critical loading conditions — such as strain ramps applied at varying rates — that result in junction failure. Previous studies have shown that the outcomes depend on the interplay between the binding and unbinding rates, the rate at which the load is applied, and the magnitude of the load itself [76, 46, 77]. Above a critical constant load, links are rapidly lost and failure ensues. However, the impact of time-varying loads has not been comprehensively explored, especially when loads only transiently surpass the critical load.

An interesting illustration of such a transient load comes from the *Drosophila* amnioserosa, which displays pulsative forces with a 200 s periodicity. When embryos are extrinsically loaded [78] or when myosin becomes over activated [79], gaps appear between cell surfaces. Myosin brings delaminated surfaces back together within 200-400 s [78]. Figure 3 illustrates how periods of low tension may allow for repair and self-healing in living materials. Thus, repair mechanisms may allow tissues to reach high tensions non-destructively for a limited period of time in the context of cyclic loading.

A key assumption of Bell's model is that force is distributed globally across all bound links. Subsequent extensions have implemented local load sharing to study crack initiation and critical length for junction failure [77]. While most models assume a rigid cell membrane, Li et al. accounted for viscoelastic cellular deformation [80]. Their findings suggest that viscosity can favour repair by prolonging the time it takes to separate membranes, thereby enhancing the likelihood of ligand-receptor pair rebinding. Once surfaces separate beyond the reach of intercellular linkers, cellular protrusions are needed to reconnect them. Consistent with this idea, recent work suggests that protrusive forces constantly act at junctions to maintain close membrane contact [81]*.

In cells, E-cadherins and desmosomal cadherins both assemble in distinct clusters whose size is force regulated [82]. Forces below 10 pN are predicted to create large numbers of small clusters (less than 5 E-cadherins each), while high forces result in a higher probability of forming few but large clusters [83]*. How this regulation takes place and why adhesions are organised in clusters remains unclear. One intriguing suggestion is that nanoclusters are the result of interdigitated microspikes extended by each cell to increase the area of intercellular contact. From a biophysical perspective, arranging links into clusters may enable junctions to be more adaptable and repair more efficiently. This raises further questions such as how forces are distributed among and within clusters, and how this may contribute to junction remodelling.

3.2 Wound healing through tissue fluidisation

Once a cellular or multicellular wound has appeared, different mechanisms are necessary to close the gap. These involve lamellipodial protrusion, to invade the gap, or the creation of a multicellular actin purse string that brings cell membranes into contact to regenerate new junctions. Closure of multicellular gaps necessitates reorganisation of the tissue involving cell intercalations. For example, in response to a multicellular wound, the *Drosophilla* wing disk epithelium first undergoes an initial fast closure phase (~20 min) attributed to myosin II-mediated purse string contraction which closes the gap at a rate of ~15 μ m²/min [84], followed by a slow phase of edge cell intercalations lasting up to four hours that necessitates unjamming of the epithelium[85]. Additionally, neighbour exchange in epithelial cells is reminiscent of a brittle to ductile transition as a result of topological defects which accelerates wound repair. During convergent extension, the Drosophila germ band undergoes 50% tissue strain and cell intercalation occurs over 25-30 min [86]. During this time, the original junction shrinks to a four-way contact point within 10 min and it takes another 10 min to build a new junction, completing neighbour exchange. How cells preserve tissue integrity during this process remains an area of active investigation.

Intriguingly, cyclic loading may favour intercalation by transiently decreasing the load on junctions. Indeed, experiments and modelling indicate that local heterogeneity in tension drives neighbour exchanges [87]. This suggests that physiological cyclic loading may promote remodelling and neighbour exchanges because of fluctuating tension at intercellular contacts. In support of this, unidirectional cyclic stretch was found to increase the frequency of cell rearrangements and induce elongation of epithelial cell colonies in the direction of stretch [88]. Mathematical modelling suggested that alterations in cell shape along with directional expansion of cell-cell interfaces can induce cell intercalation through mechanisms dependent on myosin II activity [89]*. Overall, by comparing the frequency and duration of external cyclic mechanical stimulation with the intrinsic timescales of cell migration, proliferation and intercalation, we can extract valuable insights about which tissue repair mechanisms may be at play in the different epithelial tissues subjected to cyclic loading.

Figure 3: Comparison of damage accumulation signatures for synthetic and living materials under fixed and cyclic loads (conceived in the context of intercellular junction dynamics but extends to other repair mechanisms). Under constant stress, damage accumulation in synthetic materials is a monotonic function (purple) until failure (black, dashed), similarly to living materials (green). Under cyclic stress, synthetic materials accumulate damage, but living materials may repair part or all of the accumulated damage. We speculate that the damage and repair mechanisms depend on the relationship between the relaxation portion of the cycle ($t_{relaxation}$) and the time-scale of the repair mechanism (t_{repair}). The linear repair functional form is a naive representation and may vary depending on the material, repair mechanism and imposed loading waveform, which haven't been comprehensively explored.

4 Conclusion

To grasp the conditions required for the stability of epithelial tissues under physiological cyclic deformations, it is crucial to consider the rich variety of damage and repair mechanisms in epithelial tissues, and the broad range of timescales at which they operate. We might speculate in particular that any damage accumulated during the stretch phases would be transient and repaired in the relaxed phases of the cycles.

Therefore, studying the impact of cyclic loading provides a powerful framework to dissect the interplay between damage accumulation and adaptive repair processes. Experimental and theoretical model systems provide a rich test-bed for these ideas. Refining our understanding of the timescales of damage, repair and remodelling mechanisms in epithelial tissues will be important to address questions about living materials' homeostasis and self-healing.

Acknowledgements

E.P. was supported by the Engineering and Physical Sciences Research Council Centre for Doctoral Training in Sensor Technologies for a Healthy and Sustainable Future [EP/S023046/1], L.B. was supported by an EMBO Postdoctoral Fellowship (grant no. ALTF 890-2023), L.B. and G.C. were supported by an sLoLa grant from the British Biotechnology and Biological Sciences Research council (BBSRC, grant no. BB/V019015/1) to G.C. and A.B. was supported by the European Research Executive Agency via the MSCA Postdoctoral Fellowships (BIOFRAC, Project number: 10110537).

Disclosure

The authors declare no conflict of interest.

References

- Jay D. Humphrey. Cardiovascular Solid Mechanics: Cells, Tissues, and Organs. Springer, New York, 2002.
- [2] R Christopher Geiger, Christopher D Kaufman, Ai P Lam, GR Scott Budinger, and David A Dean. Tubulin acetylation and histone deacetylase 6 activity in the lung under cyclic load. *American journal* of respiratory cell and molecular biology, 40(1):76–82, 2009.
- [3] Nurit Davidovich, Brian C DiPaolo, Gladys G Lawrence, Peter Chhour, Nadir Yehya, and Susan S Margulies. Cyclic stretch-induced oxidative stress increases pulmonary alveolar epithelial permeability. American journal of respiratory cell and molecular biology, 49(1):156–164, 2013.
- [4] K Carlos El-Tallawi, Peng Zhang, Robert Azencott, Jiwen He, Elizabeth L Herrera, Jiaqiong Xu, Mohammed Chamsi-Pasha, Jessen Jacob, Gerald M Lawrie, and William A Zoghbi. Valve strain quantitation in normal mitral valves and mitral prolapse with variable degrees of regurgitation. *Cardiovascular Imaging*, 14(6):1099–1109, 2021.
- [5] Michaela Kollisch-Singule, Bryanna Emr, Bradford Smith, Shreyas Roy, Sumeet Jain, Joshua Satalin, Kathy Snyder, Penny Andrews, Nader Habashi, Jason Bates, et al. Mechanical breath profile of airway pressure release ventilation: the effect on alveolar recruitment and microstrain in acute lung injury. JAMA surgery, 149(11):1138–1145, 2014.
- [6] Satish K. Sarna. Colonic Motility: From Bench Side to Bedside. Morgan & Claypool Life Sciences, San Rafael, CA, 2010.
- [7] Christophe Faisy, Francisco M Pinto, Morgan Le Guen, Emmanuel Naline, Stanislas Grassin Delyle, Edouard Sage, Maria-Luz Candenas, and Philippe Devillier. Airway response to acute mechanical stress in a human bronchial model of stretch. *Critical care*, 15:1–11, 2011.
- [8] J Brady Scott and Ramandeep Kaur. Monitoring breathing frequency, pattern, and effort. *Respiratory Care*, 65(6):793–806, 2020.
- [9] Jana Grune, Arata Tabuchi, and Wolfgang M Kuebler. Alveolar dynamics during mechanical ventilation in the healthy and injured lung. *Intensive care medicine experimental*, 7(Suppl 1):34, 2019.
- [10] Travis D. Homan, Stephen J. Bordes, and Erica Cichowski. Physiology, pulse pressure. StatPearls, 2023. Last updated: 10 July 2023.
- [11] John B West, Koichi Tsukimoto, Odile Mathieu-Costello, and Renato Prediletto. Stress failure in pulmonary capillaries. *Journal of Applied Physiology*, 70(4):1731–1742, 1991.
- [12] Anders Paul Persson, Alexandra Måneheim, Johan Economou Lundeberg, Artur Fedorowski, Jeff S Healey, Johan Sundström, Gunnar Engström, and Linda SB Johnson. Reference ranges for ambulatory heart rate measurements in a middle-aged population. *Heart*, 110(12):831–837, 2024.
- [13] Kimberly Howard-Quijano, Melissa McCabe, Alexander Cheng, Wei Zhou, Kentaro Yamakawa, Einat Mazor, Jennifer C Scovotti, and Aman Mahajan. Left ventricular endocardial and epicardial strain changes with apical myocardial ischemia in an open-chest porcine model. *Physiological Reports*, 4(24):e13042, 2016.
- [14] Effat Soleimani, Manijhe Mokhtari-Dizaji, Nasser Fatouraee, and Hazhir Saberi. Stress distribution analysis in healthy and stenosed carotid artery models reconstructed from in vivo ultrasonography. Ultrasonography, 40(3):428–441, Jul 2021.
- [15] Helena Forsblad-d'Elia, Lucy Law, Karin Bengtsson, Johan Smeds, Maria Ketonen, Björn Sundström, Lotta Ljung, Mats Geijer, Stefan Söderberg, and Per Lindqvist. Biomechanical properties of common carotid arteries assessed by circumferential 2d strain and β stiffness index in patients with ankylosing spondylitis. The Journal of rheumatology, 48(3):352–360, 2021.
- [16] Patrick J. Neligan. Fluid and electrolyte balance. Anaesthesia & Intensive Care Medicine, 25(2):107– 111, 2024.

- [17] E Seem and E Stranden. Transcapillary filtration in lower limbs with deep venous thrombosis; the role of the capillary filtration coefficient. Scandinavian journal of clinical and laboratory investigation, 50(3):331–336, 1990.
- [18] Louis Henquell, Paul L. LaCelle, and Carl R. Honig. Capillary diameter in rat heart in situ: Relation to erythrocyte deformability, o2 transport, and transmural o2 gradients. *Microvascular Research*, 12(3):259–274, 1976.
- [19] VM Parfeev, IV Grushetskii, and EV Smurova. Mechanical properties of elastomers for artificial leaflet heart valves. *Mechanics of Composite Materials*, 19(1):92–99, 1983.
- [20] Ailsa. J. Dalgliesh, Mojtaba Parvizi, Christopher Noble, and Leigh G. Griffiths. Effect of cyclic deformation on xenogeneic heart valve biomaterials. *PLoS One*, 14(6):e0214656, Jun 2019.
- [21] Tracy Jaffe and William M Thompson. Large-bowel obstruction in the adult: classic radiographic and ct findings, etiology, and mimics. *Radiology*, 275(3):651–663, 2015.
- [22] Hossein Joodaki and Matthew B. Panzer. Skin mechanical properties and modeling: A review. Journal of Engineering in Medicine, 232(4):323–343, 2018.
- [23] Alex Rowlands, Michelle R Stone, and Roger G Eston. Influence of speed and step frequency during walking and running on motion sensor output. *Medicine and science in sports and exercise*, 39(4):716, 2007.
- [24] João Manuel Pereira Figueiredo. Correlation of leg impedance and skin deformation during gait. Master's thesis, Universidade de Lisboa (Portugal), 2019.
- [25] Ashley M. Wessendorf and Dava J. Newman. Dynamic understanding of human-skin movement and strain-field analysis. *IEEE Transactions on Biomedical Engineering*, 59(12):3432–3438, 2012.
- [26] Keshavamurthy Indireshkumar, James G Brasseur, Henryk Faas, Geoffrey S Hebbard, Patrik Kunz, John Dent, Christine Feinle, Meijng Li, Peter Boesiger, Michael Fried, et al. Relative contributions of "pressure pump" and "peristaltic pump" to gastric emptying. *American Journal of Physiology-Gastrointestinal and Liver Physiology*, 278(4):G604–G616, 2000.
- [27] Ryohei Ebara, Shota Ishida, Takumi Miyagawa, and Yutaka Imai. Effects of peristaltic amplitude and frequency on gastric emptying and mixing: a simulation study. *Journal of the Royal Society Interface*, 20(198):20220780, Jan 2023.
- [28] Yusuf Kayar, Ahmet Danahoğlu, AA Kafee, Şükrü Okkesim, and Hakan Şentürk. Gastric myoelectrical activity abnormalities of electrogastrography in patients with functional dyspepsia. Turk J Gastroenterol, 27(5):415–420, 2016.
- [29] Adil E Bharucha, Armando Manduca, David S Lake, Jeff Fidler, Phillip Edwards, Roger C Grimm, Alan R Zinsmeister, and Stephen J Riederer. Gastric motor disturbances in patients with idiopathic rapid gastric emptying. *Neurogastroenterology & Motility*, 23(7):617–e252, 2011.
- [30] T Schmidt, N Hackelsberger, R Widmer, C Meisel, A Pfeiffer, and H Kaess. Ambulatory 24-hour jejunal motility in diarrhea-predominant irritable bowel syndrome. *Scandinavian journal of gastroenterology*, 31(6):581–589, 1996.
- [31] Jianhu Zhang, Cheri R Owen, Matthew A Sanders, Jerrold R Turner, and Marc D Basson. The motogenic effects of cyclic mechanical strain on intestinal epithelial monolayer wound closure are matrix dependent. *Gastroenterology*, 131(4):1179–1189, 2006.
- [32] Aron Parekh, Alexander D. Cigan, Silvia Wognum, Rebecca L. Heise, Michael B. Chancellor, and Michael S. Sacks. Ex vivo deformations of the urinary bladder wall during whole bladder filling: Contributions of extracellular matrix and smooth muscle. *Journal of biomechanics*, 43(9):1708–1716, 2010.
- [33] Jean F. Wyman, Caitlin H. Cain, Cynthia N. Epperson, Cynthia M. Fitzgerald, Sheila Gahagan, Diane K. Newman, Kyle Rudser, Amy L. Smith, Christine P. Vaughan, Siobhan Sutcliffe, and Prevention

of Lower Urinary Tract Symptoms (PLUS) Research Consortium. Urination frequency ranges in healthy women. Nursing Research, 71(5):341–352, 2022.

- [34] Carlo Bulletti, Dominique De Ziegler, Valeria Polli, Elena Del Ferro, Simone Palini, and Carlo Flamigni. Characteristics of uterine contractility during menses in women with mild to moderate endometriosis. *Fertility and sterility*, 77(6):1156–1161, 2002.
- [35] Osnat Eytan, Ilan Halevi, Joseph Har-Toov, Igal Wolman, David Elad, and Ariel J Jaffa. Characteristics of uterine peristalsis in spontaneous and induced cycles. *Fertility and sterility*, 76(2):337–341, 2001.
- [36] H Roshani, NF Dabhoiwala, T Dijkhuis, and WH Lamers. Intraluminal pressure changes in vivo in the middle and distal pig ureter during propagation of a peristaltic wave. Urology, 59(2):298–302, 2002.
- [37] Duane R Hickling, Tung-Tien Sun, and Xue-Ru Wu. Anatomy and physiology of the urinary tract: Relation to host defense and microbial infection. *Microbiology Spectrum*, 3(4):UTI-0016-2012, 2015.
- [38] Laxmikant G Keni, Satish Shenoy, KN Chethan, Padmaraj Hegde, K Prakashini, Masaaki Tamagawa, and Mohammad Zuber. Analyzing single peristaltic wave behavior in ureter with variable diameters: A ureterdynamic study. CFD Letters, 16(4):54–68, 2024.
- [39] Morris Traube, Riyad Albibi, and Richard W. McCallum. High-Amplitude Peristaltic Esophageal Contractions Associated With Chest Pain. JAMA, 250(19):2655–2659, 11 1983.
- [40] Dustin A Carlson, Wenjun Kou, and John E Pandolfino. The rhythm and rate of distension-induced esophageal contractility: a physiomarker of esophageal function. *Neurogastroenterology & Motility*, 32(5):e13794, 2020.
- [41] Richard E Leibbrandt, Phil G Dinning, Marcello Costa, Charles Cock, Lukasz Wiklendt, Guangsong Wang, Jan Tack, Dirk Van Beckevoort, Nathalie Rommel, and Taher I Omari. Characterization of esophageal physiology using mechanical state analysis. *Frontiers in systems neuroscience*, 10:10, 2016.
- [42] Joohee Lee, James Huprich, Christine Kujath, Karthik Ravi, Felicity Enders, Thomas C Smyrk, David A Katzka, Nicholas J Talley, and Jeffrey A Alexander. Esophageal diameter is decreased in some patients with eosinophilic esophagitis and might increase with topical corticosteroid therapy. *Clinical gastroenterology and hepatology*, 10(5):481–486, 2012.
- [43] Hyungsuk Lee, Benjamin Pelz, Jorge M Ferrer, Taeyoon Kim, Matthew J Lang, and Roger D Kamm. Cytoskeletal deformation at high strains and the role of cross-link unfolding or unbinding. *Cellular and Molecular Bioengineering*, 2:28–38, 2009.
- [44] Johanna Block, Hannes Witt, Andrea Candelli, Erwin JG Peterman, Gijs JL Wuite, Andreas Janshoff, and Sarah Köster. Nonlinear loading-rate-dependent force response of individual vimentin intermediate filaments to applied strain. *Physical review letters*, 118(4):048101, 2017.
- [45] Zanetta Kechagia, Matthias Eibauer, and Ohad Medalia. Structural determinants of intermediate filament mechanics. *Current Opinion in Cell Biology*, 89:102375, 2024.
- [46] **Julia Duque, Alessandra Bonfanti, Jonathan Fouchard, Lucia Baldauf, Sara R Azenha, Emma Ferber, Andrew Harris, Elias H Barriga, Alexandre J Kabla, and Guillaume Charras. Rupture strength of living cell monolayers. *Nature Materials*, pages 1–12, 2024.

This study presents the strain-rate controlled stiffening of epithelial monolayers under large deformations, studied through mechanical measurements, live imaging and computational models. The authors discuss the contribution of adhesive bond kinetics and history of deformation on the measured tissue strength, and compare the onset of fracture across wild-type and fragilized monolayers.

[47] Diana Pinheiro and Yohanns Bellaiche. Mechanical force-driven adherens junction remodeling and epithelial dynamics. *Developmental cell*, 47(1):3–19, 2018.

- [48] Sirio Dupont and Sara A Wickström. Mechanical regulation of chromatin and transcription. Nature Reviews Genetics, 23(10):624–643, 2022.
- [49] **Otger Campàs, Ivar Noordstra, and Alpha S Yap. Adherens junctions as molecular regulators of emergent tissue mechanics. Nature Reviews Molecular Cell Biology, 25(4):252–269, 2024.

This review examines the role of adherens junctions (AJs) in tissue mechanics. It highlights the molecular and cellular characteristics of AJs which influence emergent tissue mechanics, such as adhesion strength and force transmission. It also discusses the dynamic nature of AJs in response to mechanical cues, which is essential for processes like tissue morphogenesis and homeostasis.

- [50] Hugo Wioland, Antoine Jegou, and Guillaume Romet-Lemonne. Torsional stress generated by adf/cofilin on cross-linked actin filaments boosts their severing. *Proceedings of the National Academy of Sciences*, 116(7):2595–2602, 2019.
- [51] Marco Fritzsche, Alexandre Lewalle, Tom Duke, Karsten Kruse, and Guillaume Charras. Analysis of turnover dynamics of the submembranous actin cortex. *Molecular biology of the cell*, 24(6):757–767, 2013.
- [52] Nargess Khalilgharibi, Jonathan Fouchard, Nina Asadipour, Ricardo Barrientos, Maria Duda, Alessandra Bonfanti, Amina Yonis, Andrew Harris, Payman Mosaffa, Yasuyuki Fujita, et al. Stress relaxation in epithelial monolayers is controlled by the actomyosin cortex. *Nature physics*, 15(8):839–847, 2019.
- [53] Sabyasachi Rakshit, Yunxiang Zhang, Kristine Manibog, Omer Shafraz, and Sanjeevi Sivasankar. Ideal, catch, and slip bonds in cadherin adhesion. *Proceedings of the National Academy of Sciences*, 109(46):18815–18820, 2012.
- [54] Craig D Buckley, Jiongyi Tan, Karen L Anderson, Dorit Hanein, Niels Volkmann, William I Weis, W James Nelson, and Alexander R Dunn. The minimal cadherin-catenin complex binds to actin filaments under force. *Science*, 346(6209):1254211, 2014.
- [55] *Vipul T Vachharajani, Matthew P DeJong, and Alexander R Dunn. Pdz domains from the junctional proteins afadin and zo-1 act as mechanosensors. *bioRxiv*, pages 2023–09, 2023.

This study demonstrates the presence of molecular catch bonds which strengthen under tension between the PDZ domain of afadin and the intercellular domains of the adhesion molecules nectin-1 and JAM-A. The authors suggest that the mechanosensitivity of PDZ may allow the dynamic remodelling of junctional complexes.

[56] **Pierre Mangeol, Dominique Massey-Harroche, Michael Sebbagh, Fabrice Richard, André Le Bivic, and Pierre-François Lenne. The zonula adherens matura redefines the apical junction of intestinal epithelia. *Proceedings of the National Academy of Sciences*, 121(9):e2316722121, 2024.

This study uncovers a novel mechanical control module within epithelia that modulates barrier function by regulating cell packing and fluid permeability. By quantifying transitions between leak-tight and more permeable states, the authors reveal that the *zonula adherens matura* dual-belt organisation regulates epithelial mechanics.

- [57] Andrew J Price, Anna-Lena Cost, Hanna Ungewiß, Jens Waschke, Alexander R Dunn, and Carsten Grashoff. Mechanical loading of desmosomes depends on the magnitude and orientation of external stress. *Nature communications*, 9(1):5284, 2018.
- [58] Sindora R Baddam, Paul T Arsenovic, Vani Narayanan, Nicole R Duggan, Carl R Mayer, Shaston T Newman, Dahlia A Abutaleb, Abhinav Mohan, Andrew P Kowalczyk, and Daniel E Conway. The desmosomal cadherin desmoglein-2 experiences mechanical tension as demonstrated by a fret-based tension biosensor expressed in living cells. *Cells*, 7(7):66, 2018.

- [59] Andrew R Harris, Loic Peter, Julien Bellis, Buzz Baum, Alexandre J Kabla, and Guillaume T Charras. Characterizing the mechanics of cultured cell monolayers. *Proceedings of the National Academy of Sciences*, 109(41):16449–16454, 2012.
- [60] Marco Pensalfini, Tom Golde, Xavier Trepat, and Marino Arroyo. Nonaffine mechanics of entangled networks inspired by intermediate filaments. *Physical review letters*, 131(5):058101, 2023.
- [61] **Charlotta Lorenz, Johanna Forsting, Robert W Style, Stefan Klumpp, and Sarah Köster. Keratin filament mechanics and energy dissipation are determined by metal-like plasticity. *Matter*, 6(6):2019– 2033, 2023.

The authors mechanically characterise the different energy dissipation mechanisms of keratin and vimentin intermediate filaments. They find that the structural differences between the two types of filaments lead keratin filaments to elongate while maintaining their stiffness (similar to solid metals) and vimentin filaments to soften while maintaining their length (like double-network gels).

- [62] Magdalena Prechova, Zuzana Adamova, Anna-Lena Schweizer, Miloslava Maninova, Andreas Bauer, Delf Kah, Samuel M Meier-Menches, Gerhard Wiche, Ben Fabry, and Martin Gregor. Plectin-mediated cytoskeletal crosstalk controls cell tension and cohesion in epithelial sheets. *Journal of Cell Biology*, 221(3):e202105146, 2022.
- [63] **Wily G Ruiz, Dennis R Clayton, Tanmay Parakala-Jain, Marianela G Dalghi, Jonathan Franks, and Gerard Apodaca. The rat bladder umbrella cell keratin network: Organization, dependence on the plectin cytolinker, and responses to bladder filling. *Molecular Biology of the Cell*, 35(11):ar139, 2024.

This study examines how the structural organisation of the keratin intermediate filament network depends on the mechanosensitive cytolinker plectin. The findings elucidate the cellular mechanisms responsible for barrier function and integrity during the filling and voiding cycles in the bladder.

- [64] Joleen S Cheah, Kyle A Jacobs, Volkmar Heinrich, Su Hao Lo, and Soichiro Yamada. Force-induced recruitment of cten along keratin network in epithelial cells. *Proceedings of the National Academy of Sciences*, 116(40):19799–19801, 2019.
- [65] Ana C Laly, Kristina Sliogeryte, Oscar J Pundel, Rosie Ross, Michael C Keeling, Deepa Avisetti, Ahmad Waseem, Núria Gavara, and John T Connelly. The keratin network of intermediate filaments regulates keratinocyte rigidity sensing and nuclear mechanotransduction. *Science Advances*, 7(5):eabd6187, 2021.
- [66] Joshua A Broussard, Avinash Jaiganesh, Hoda Zarkoob, Daniel E Conway, Alexander R Dunn, Horacio D Espinosa, Paul A Janmey, and Kathleen J Green. Scaling up single-cell mechanics to multicellular tissues-the role of the intermediate filament-desmosome network. *Journal of cell science*, 133(6):jcs228031, 2020.
- [67] Yung-Li Lee and Jian Pan. Fatigue Testing and Analysis: Theory and Practice, pages 57–76. Elsevier, 2005.
- [68] Yuelan Wang, Richard D Minshall, David E Schwartz, and Guochang Hu. Cyclic stretch induces alveolar epithelial barrier dysfunction via calpain-mediated degradation of p120-catenin. American Journal of Physiology-Lung Cellular and Molecular Physiology, 301(2):L197–L206, 2011.
- [69] Ciara Durcan, Mokarram Hossain, Grégory Chagnon, Djordje Perić, Georges Karam, Lara Bsiesy, and Edouard Girard. Experimental investigations of the human oesophagus: anisotropic properties of the embalmed mucosa–submucosa layer under large deformation. *Biomechanics and Modeling in Mechanobi*ology, 21(6):1685–1702, 2022.
- [70] Melanie Homberg, Lena Ramms, Nicole Schwarz, Georg Dreissen, Rudolf E Leube, Rudolf Merkel, Bernd Hoffmann, and Thomas M Magin. Distinct impact of two keratin mutations causing epider-

molysis bullosa simplex on keratinocyte adhesion and stiffness. *Journal of Investigative Dermatology*, 135(10):2437–2445, 2015.

- [71] Girish R Kale, Xingbo Yang, Jean-Marc Philippe, Madhav Mani, Pierre-François Lenne, and Thomas Lecuit. Distinct contributions of tensile and shear stress on e-cadherin levels during morphogenesis. *Nature communications*, 9(1):5021, 2018.
- [72] **Kenji Nishizawa, Shao-Zhen Lin, Claire Chardès, Jean-François Rupprecht, and Pierre-François Lenne. Two-point optical manipulation reveals mechanosensitive remodeling of cell-cell contacts in vivo. Proceedings of the National Academy of Sciences, 120(13):e2212389120, 2023.

This study investigates how cells sense and respond to mechanical cues via stress fibres and focal adhesions. Using optical manipulation, the authors can probe the mechanics of cell-cell contacts by exreting localised contractile and extensile stresses. They find that stress fibres and focal adhesions exhibit dynamic mechanosensitive responses and restructuring.

[73] *Amin Esmaeilniakooshkghazi, Eric Pham, Sudeep P George, Afzal Ahrorov, Fabian R Villagomez, Michael Byington, Srijita Mukhopadhyay, Srinivas Patnaik, Jacinta C Conrad, Monali Naik, et al. In colon cancer cells fascin1 regulates adherens junction remodeling. *The FASEB Journal*, 37(3):e22786, 2023.

This study identifies a novel role for fascin1 in adherens junction (AJ) remodelling in colon cancer cells, showing that it destabilizes AJs by remodelling junctional actin and actomyosin contractility. This drives collective plasticity which enhances tumor cell migration, growth and dissemination offering new insights into the mechanics of colorectal carcinoma progression.

[74] *Vera Janssen and Stephan Huveneers. Cell-cell junctions in focus-imaging junctional architectures and dynamics at high resolution. *Journal of Cell Science*, 137(20), 2024.

This review presents recent advances in microscopy tools, such as biosensors and optogenetic probes, along with new insights from super-resolution and volume electron microscopy on the nanoscale organization and membrane morphology of tight junctions, adherens junctions and desmosomes.

- [75] George I Bell. Models for the specific adhesion of cells to cells: a theoretical framework for adhesion mediated by reversible bonds between cell surface molecules. *Science*, 200(4342):618–627, 1978.
- [76] Thorsten Erdmann and Ulrich S Schwarz. Stochastic dynamics of adhesion clusters under shared constant force and with rebinding. *The Journal of chemical physics*, 121(18):8997–9017, 2004.
- [77] Yuval Mulla, Giorgio Oliveri, Johannes TB Overvelde, and Gijsje H Koenderink. Crack initiation in viscoelastic materials. *Physical review letters*, 120(26):268002, 2018.
- [78] Angughali Sumi, Peran Hayes, Arturo D'Angelo, Julien Colombelli, Guillaume Salbreux, Kai Dierkes, and Jérôme Solon. Adherens junction length during tissue contraction is controlled by the mechanosensitive activity of actomyosin and junctional recycling. *Developmental cell*, 47(4):453–463, 2018.
- [79] J Duque and N Gorfinkiel. Integration of actomyosin contractility with cell-cell adhesion during dorsal closure. dev. camb. engl. 143, 4676–4686, 2016.
- [80] Long Li, Wenyan Zhang, and Jizeng Wang. A viscoelastic–stochastic model of the effects of cytoskeleton remodelling on cell adhesion. *Royal Society open science*, 3(10):160539, 2016.
- [81] *Yosuke Senju, Toiba Mushtaq, Helena Vihinen, Aki Manninen, Juha Saarikangas, Katharina Ven, Ulrike Engel, Markku Varjosalo, Eija Jokitalo, and Pekka Lappalainen. Actin-rich lamellipodia-like protrusions contribute to the integrity of epithelial cell–cell junctions. *Journal of Biological Chemistry*, 299(5), 2023.

This study explores the role of actin dynamics in maintaining the integrity of epithelial adherens junctions (AJs). It supports that lamelipodia-like protrusions at cell-cell junctions are critical for reinforcing and stabilising epithelial junctions. The findings provide valuable insights into mechanisms underlying cell migration and diseases such as cancer and inlammatory disorders.

[82] *Sergey M Troyanovsky. Adherens junction: the ensemble of specialized cadherin clusters. *Trends in cell biology*, 33(5):374–387, 2023.

This review explores the phenomenon of clustering due to cis interactions of cadherins located in the same cell. The authors suggest that cadherin clusters are tuned by mechanical forces and reinforce weak and unstable cadherin trans dimers. The authors conclude that the lifetime, strength and plasticity of adherens junctions should be examined in the context of molecular clustering.

- [83] Yang Chen, Julia Brasch, Oliver J Harrison, and Tamara C Bidone. Computational model of e-cadherin clustering under force. *Biophysical Journal*, 120(22):4944–4954, 2021.
- [84] Anna B Kobb, Teresa Zulueta-Coarasa, and Rodrigo Fernandez-Gonzalez. Tension regulates myosin dynamics during drosophila embryonic wound repair. *Journal of Cell Science*, 130(4):689–696, 2017.
- [85] Robert J Tetley, Michael F Staddon, Davide Heller, Andreas Hoppe, Shiladitya Banerjee, and Yanlan Mao. Tissue fluidity promotes epithelial wound healing. *Nature physics*, 15(11):1195–1203, 2019.
- [86] Sérgio de Matos Simões, Avantika Mainieri, and Jennifer A Zallen. Rho gtpase and shroom direct planar polarized actomyosin contractility during convergent extension. *Journal of Cell Biology*, 204(4):575–589, 2014.
- [87] Scott Curran, Charlotte Strandkvist, Jasper Bathmann, Marc de Gennes, Alexandre Kabla, Guillaume Salbreux, and Buzz Baum. Myosin ii controls junction fluctuations to guide epithelial tissue ordering. *Developmental cell*, 43(4):480–492, 2017.
- [88] Jordi Comelles, Soumya Ss, Linjie Lu, Emilie Le Maout, Sudakar Anvitha, Guillaume Salbreux, Frank Jülicher, Mandar M Inamdar, and Daniel Riveline. Epithelial colonies in vitro elongate through collective effects. *Elife*, 10:e57730, 2021.
- [89] *Jui-Chien Lien and Yu-li Wang. Cyclic stretching combined with cell-cell adhesion is sufficient for inducing cell intercalation. *Biophysical Journal*, 122(15):3146–3158, 2023.

This study combines experimental and mathematical modelling approaches to assess the effect of uniaxial cyclic stretch on epithelial cell intercalation. The key results suggest that cyclic stretch can induce cell intercalation by inducing cell shape changes and reorientation in the presence of dynamic intercellulat adhesions.