SCATTERING THEORY FOR THE DEFOCUSING 3D NLS IN THE EXTERIOR OF A STRICTLY CONVEX OBSTACLE

XUAN LIU, YILIN SONG, AND JIQIANG ZHENG

ABSTRACT. In this paper, we investigate the global well-posedness and scattering theory for the defocusing nonlinear Schrödinger equation $iu_t + \Delta_\Omega u = |u|^\alpha u$ in the exterior domain Ω of a smooth, compact and strictly convex obstacle in \mathbb{R}^3 . It is conjectured that in Euclidean space, if the solution has a prior bound in the critical Sobolev space, that is, $u \in L^\infty_t(I; \dot{H}^{s_c}_x(\mathbb{R}^3))$ with $s_c := \frac{3}{2} - \frac{2}{\alpha} \in (0, \frac{3}{2})$, then u is global and scatters. In this paper, assuming that this conjecture holds, we prove that if u is a solution to the nonlinear Schrödinger equation in exterior domain Ω with Dirichlet boundary condition and satisfies $u \in L^\infty_t(I; \dot{H}^{s_c}_D(\Omega))$ with $s_c \in \left[\frac{1}{2}, \frac{3}{2}\right)$, then u is global and scatters.

The main ingredients in our proof are the linear profile decompositions in the critical space $\dot{H}_D^{sc}(\Omega)$ and embed it to the nonlinear equation. Inspired by Killip-Visan-Zhang [Amer. J. Math. 138 (2016)], we overcome the difficulty caused by the breakdown of the scaling and translation invariance. This allows us to utilize the concentration-compactness/rigidity argument of Kenig and Merle [Invent. Math. 166 (2006)]. To preclude the minimal counterexamples, we established the long-time Stricharrtz estimates and the frequency-localized Morawetz estimates.

Contents

| 1. Introduction | 1 |
|---|----|
| 1.1. Outline of the proof of Theorem 1.2 | 4 |
| 2. Preliminaries | 7 |
| 2.1. Notation and useful lemmas | 7 |
| 2.2. Littlewood-Paley theory on exterior domains | 9 |
| 2.3. Strichartz estimates, local well-posedness, and the stability result | 10 |
| 2.4. Convergence results | 13 |
| 3. Linear profile decomposition | 14 |
| 4. Embedding of nonlinear profiles | 25 |
| 5. Reduction to Almost Periodic Solutions | 35 |
| 6. The cases $1 < s_c < \frac{3}{2}$ and $s_c = \frac{1}{2}$. | 39 |
| 7. The case $\frac{1}{2} < s_c < 1$. | 41 |
| 7.1. Long time Strichartz estimates | 42 |
| 7.2. A Frequency-Localized Lin-Strauss Morawetz Inequality | 43 |
| 7.3. Rule out the minimal conuterexample | 47 |
| References | 48 |

1. Introduction

We study the defocusing nonlinear Schrödinger equation in the exterior domain Ω of a smooth compact, strictly convex obstacle in \mathbb{R}^3 with Dirichlet boundary condition:

$$\begin{cases} iu_t + \Delta_{\Omega} u = |u|^{\alpha} u, \\ u(0, x) = u_0(x), \\ u(t, x)|_{x \in \partial\Omega} = 0, \end{cases}$$

$$(1.1)$$

where u is a complex-valued function defined in $\mathbb{R} \times \Omega$ and $-\Delta_{\Omega}$ denotes the Dirichlet Laplacian on Ω . The Dirichlet-Laplacian is the unique self-adjoint operator on $L^2(\Omega)$ corresponding to the following

quadratic form

$$Q: H^1_0(\Omega) \to [0,\infty) \quad \text{with} \quad Q(f) := \int_{\Omega} \overline{\nabla f(x)} \cdot \nabla f(x) \, dx.$$

We take initial data $u_0 \in \dot{H}^s_D(\Omega)$, where for $s \geq 0$, the homogeneous Sobolev space is defined by the functional calculus as the completion of $C_c^{\infty}(\Omega)$ with respect to the norm

$$||f||_{\dot{H}^{s}_{\mathcal{D}}(\Omega)} := ||(-\Delta_{\Omega})^{s/2}f||_{L^{2}(\Omega)}.$$

It is easy to find that the solution u to equation (1.1) with sufficient smooth conditions possesses the mass and energy conservation laws:

$$\begin{split} M_{\Omega}[u(t)] &:= \int_{\Omega} |u(t,x)|^2 dx = M_{\Omega}[u_0], \\ E_{\Omega}[u(t)] &:= \frac{1}{2} \int_{\Omega} |\nabla u(t,x)|^2 dx + \frac{1}{\alpha+2} \int_{\Omega} |u(t,x)|^{\alpha+2} dx = E_{\Omega}[u_0]. \end{split}$$

When posed on the whole Euclidean space \mathbb{R}^3 , the Cauchy problem (1.1) is scale-invariant. More precisely, the scaling transformation

$$u(t,x) \longmapsto \lambda^{\frac{2}{\alpha}} u(\lambda x, \lambda^2 t)$$
 for $\lambda > 0$,

leaves the class of solutions to $\text{NLS}_{\mathbb{R}^3}$ invariant. This transformation also identifies the critical space $\dot{H}_x^{s_c}$, where the critical regularity s_c is given by $s_c := \frac{3}{2} - \frac{2}{\alpha}$. We call (1.1) mass-critical if $s_c = 0$, energy-critical if $s_c = 1$, inter-critical if $0 < s_c < 1$ and energy-supercritical if $s_c > 1$ respectively. Although the obstacle in the domain alters certain aspects of the equation, it does not affect the problem's inherent dimensionality. Therefore, (1.1) maintains the same criticality and is classified as $\dot{H}_D^{s_c}(\Omega)$ critical.

Throughout this paper, we restrict ourselves to the following notion of solution.

Definition 1.1 (Solution). A function $u: I \times \Omega \to \mathbb{C}$ on a non-empty interval $I \ni 0$ is called a solution to (1.1) if it satisfies $u \in C_t \dot{H}^{s_c}_D(K \times \Omega) \cap L^{\frac{5\alpha}{2}}_{t,x}(K \times \Omega)$ for every compact subset $K \subset I$ and obeys the Duhamel formula

$$u(t) = e^{it\Delta_{\Omega}}u_0 - i\int_0^t e^{i(t-s)\Delta_{\Omega}}(|u|^{\alpha}u)(s) ds$$

for each $t \in I$. We refer to the interval I as the lifespan of u. We say that u is a maximal-lifespan solution if the solution cannot be extended to any strictly larger interval. We say that u is a global solution if $I = \mathbb{R}$.

The assumption that the solution lies in the space $L_{t,x}^{\frac{5\alpha}{2}}(I \times \Omega)$ locally in time is natural since by the Strichartz estimate (see Proposition 2.15 below), the linear flow always lies in this space. Also, if a solution u to (1.1) is global, with $\|u\|_{L_{t,x}^{\frac{5\alpha}{2}}(I \times \Omega)} < \infty$, then it *scatters*; that is, there exist unique $u_{\pm} \in \dot{H}_D^{s_c}(\Omega)$ such that

$$\lim_{t \to +\infty} \|u(t) - e^{it\Delta_{\Omega}} u_{\pm}\|_{\dot{H}^{s_c}_{D}(\Omega)} = 0.$$

The study of NLS in exterior domains was initiated in [5]. The authors proved a local existence result for the 3d sub-cubic (i.e., $\alpha < 3$) NLS $_{\Omega}$ equation, assuming that the obstacle is non-trapping. Subsequently, Anton [1] extended these result to the cubic nonlinearity, while Planchon-Vega [48] extended it to the energy-subcritical NLS $_{\Omega}$ equation in dimension d=3. Later, Planchon and Ivanovici [23] established the small data scattering theory for the energy-critical NLS $_{\Omega}$ equation in dimension d=3. For NLS outside a smooth, compact, strictly convex obstacle Ω in \mathbb{R}^3 , Killip-Visan-Zhang [35] proved that for arbitrarily large initial data, the corresponding solutions to the defocusing energy-critical equation scatter in the energy space. For related results in the focusing case, see e.g. [15, 37, 59, 60].

In this paper, we investigate the $\dot{H}^{s_c}_D(\Omega)$ critical global well-posedness and scattering theory for the defocusing nonlinear Schrödinger equation (1.1) in the exterior domain Ω of a smooth, compact and strictly convex obstacle in \mathbb{R}^3 . To put the problem in context, let us first recall some earlier results for the equivalent problem posed in the whole Euclidean space \mathbb{R}^d . The study of global well-posedness and scattering theory for nonlinear Schrödinger equations

$$iu_t + \Delta u = \pm |u|^{\alpha} u, \qquad (t, x) \in \mathbb{R} \times \mathbb{R}^d$$
 (1.2)

in \dot{H}^{s_c} has seen significant advancements in recent years. Due to the presence of conserved quantities at the critical regularity, the mass- and energy-critical equations have been the most widely studied. For the defocusing energy-critical NLS, it is now known that arbitrary data in \dot{H}^1_x lead to solutions that are global and scatter. This was proven first for radial initial data by Bourgain [3], Grillakis [18], and Tao [52] and later for arbitrary data by Colliander- Keel-Staffilani-Takaoka-Tao, [7], Ryckman-Visan [49] and Visan [56,57] (For results in the focusing case, see [12,24,29]). For the mass-critical NLS, it has also been established that arbitrary data in L^2_x lead to solutions that are global and scatter. This was proven through the use of minimal counterexamples, first for radial data in dimensions $d \geq 2$ (see [28,34,55]), and later for arbitrary data in all dimensions by Dodson [8–11].

Killip-Visan [32] and Visan [57] revisited the defocusing energy-critical problem in dimensions $d \in \{3,4\}$ from the perspective of minimal counterexamples, utilizing techniques developed by Dodson [8] in the mass-critical setting. In particular, they established a "long-time Strichartz estimate" for almost periodic solutions, which serves to rule out the existence of frequency-cascade solutions. Additionally, they derived a frequency-localized interaction Morawetz inequality (which may in turn be used to preclude the existence of soliton-like solutions).

Unlike the energy- and mass-critical problems, for any other $s_c \neq 0, 1$, there are no conserved quantities that control the growth in time of the \dot{H}^{s_c} norm of the solutions. It is conjectured that, assuming some a priori control of a critical norm, global well-posedness and scattering hold for any $s_c > 0$ and in any spatial dimension:

Conjecture 1.1. Let $d \geq 1$, $\alpha \geq \frac{4}{d}$, and $s_c = \frac{d}{2} - \frac{2}{\alpha}$. Assume $u : I \times \mathbb{R}^d \to \mathbb{C}$ is a maximal-lifespan solution to (1.2) such that

$$u \in L_t^{\infty} \dot{H}_x^{s_c}(I \times \mathbb{R}^d),$$

then u is global and scatters as $t \to \pm \infty$.

The first work dealing with Conjecture 1.1 is attributed to Kenig and Merle [25] at the case $d=3, s_c=\frac{1}{2}$ by using their concentration-compactness method developed in [24] and the scaling-critical Lin-Strauss Morawetz inequality. Subsequently, Murphy [46] extended the methods of [25] to higher dimensions, resolving Conjecture 1.1 for $d\geq 3$ and $s_c=\frac{1}{2}$. In the inter-critical case $(0< s_c<1)$, Murphy [45, 47] developed a long-time Strichartz estimate in the spirit of [8] and proved Conjecture 1.1 for the general data in the case

$$\begin{cases} \frac{1}{2} \le s_c \le \frac{3}{4}, & d = 3\\ \frac{1}{2} \le s_c < 1, & d = 4\\ \frac{1}{2} < s_c < 1, & d = 5; \end{cases}$$

and for the radial data in the case $d=3, s_c\in(0,\frac{1}{2})\cup(\frac{3}{4},1)$. Later, Gao-Miao-Yang [16] resolved Conjecture 1.1 for radial initial data in the case $d\geq 4, 0< s_c<\frac{1}{2}$; Gao-Zhao [17] resolved Conjecture 1.1 for general initial data in the case $d\geq 5, \frac{1}{2}< s_c<1$. See also [58] for earlier partial results regarding these cases. Recently, Yu [61] resolved Conjecture 1.1 in the case $d=2, s_c=\frac{1}{2}$, by first developing a long-time Strichartz estimate in the spirit of [10] and then utilizing the interaction Morawetz estimate from Planchon-Vega [48] to exclude the minimal counterexamples. See Table 1.

In the energy-supercritical case $(s_c>1)$, Killip and Visan [29] were the first to resolve Conjecture 1.1 for $d\geq 5$ under certain conditions on s_c . Subsequently, Murphy [47] addressed the conjecture for radial initial data in the case d=3 and $s_c\in(1,\frac32)$. By developing long-time Strichartz estimates for the energy-supercritical regime, Miao-Murphy-Zheng [44] and Dodson-Miao-Murphy-Zheng [13] resolved the Conjecture 1.1 for general initial data when d=4 and $1< s_c\leq \frac32$. For the case d=4 and $\frac32< s_c<2$ with radial initial data, see the work of Lu and Zheng [43]. More recently, Zhao [63] and Li-Li [42] resolved the Conjecture 1.1 in the case $d\geq 5$ and $1< s_c<\frac d2$. For $d\geq 8$, their results also required α to be an even number. See Table 2.

Analogous to Conjecture 1.1, it is conjectured that for the Schrödinger equation in the exterior domain Ω of a smooth, compact, strictly convex obstacle in \mathbb{R}^3 :

Conjecture 1.2. Let $\alpha > \frac{4}{3}$ and $s_c = \frac{3}{2} - \frac{2}{\alpha}$. Assume $u : I \times \Omega \to \mathbb{C}$ is a maximal-lifespan solution to (1.1) such that

$$u \in L_t^{\infty} \dot{H}_D^{s_c}(I \times \Omega), \tag{1.3}$$

then u is global and scatters as $t \to \pm \infty$.

| | $0 < s_c < \frac{1}{2}$ | $s_c = \frac{1}{2}$ | $\frac{1}{2} < s_c < 1$ |
|-----------|----------------------------|---------------------|---|
| d = 1 | no results | \ | \ |
| d=2 | no results | Yu [61] | no results |
| d=3 | radial, Murphy [47] | Kenig-Merle [25] | $\frac{1}{2} < s_c \le \frac{3}{4}, \text{Murphy [45]}$ radial, $\frac{3}{4} < s_c < 1, \text{Murphy [47]}$ |
| $d \ge 4$ | radial, Gao-Miao-Yang [16] | Murphy [46] | Gao-Zhao [17], Murphy [45], Xie-Fang [58] |

Table 1. Results for Conjecture 1.1 in the sub-critical case: $0 < s_c < 1$

Table 2. Results for Conjecture 1.1 in the super-critical case: $1 < s_c < \frac{d}{2}$

| d=3 | $1 < s_c < \frac{3}{2}$, radial, Murphy [47] |
|-----------|--|
| d=4 | $1 < s_c < \frac{3}{2}$, Miao-Murphy-Zheng [44]; $s_c = \frac{3}{2}$, Dodson-Miao-Murphy-Zheng [13]; $\frac{3}{2} < s_c < 2$, radial, Lu-Zheng [43] |
| $d \ge 5$ | $1 < s_c < \frac{d}{2}$, and assume α is even when $d \ge 8$, Killip-Visan [29], Zhao [63], Li-Li [42] |

Killip-Visan-Zhang [35] first resolved Conjecture 1.2 in the case d=3 and $s_c=1$. Since this corresponds to the energy-critical setting, the energy conservation law eliminates the need for the assumption (1.3); it suffices to require the initial data to belong to $\dot{H}_D^1(\Omega)$. In this paper, under the assumption that Conjecture 1.1 holds in Euclidean space, we resolve Conjecture 1.2 in the case d=3 and $\frac{1}{2} \leq s_c < \frac{3}{2}$. Our main result is as follows:

Theorem 1.2. Let $s_c \in [\frac{1}{2}, \frac{3}{2})$. Assume that Conjection 1.1 holds. Then Conjection 1.2 holds.

Remark 1.3. In Section 4, we will embed the solutions in the limit geometries into Ω via the stability theorem 2.19. To achieve this, we assume that Conjecture 1.1 holds true, so that the solutions in the limit geometries satisfy uniform spacetime bounds; then the solutions to NLS_{Ω} will inherit these spacetime bounds. These solutions to NLS_{Ω} will appear again as nonlinear profiles in Proposition 5.1.

Remark 1.4. In [25, 45, 46], Kenig-Merle and Murphy proved that Conjecture 1.1 holds for $s_c \in [\frac{1}{2}, \frac{3}{4}]$; and for $s_c \in (\frac{3}{4}, 1) \cup (1, \frac{3}{2})$, Murphy [47] resolved the conjecture 1.1 for the case of radial initial data. Therefore, in Theorem 1.2, we only need to assume that Conjecture 1.1 holds for non-radial initial data when $s_c \in (\frac{3}{4}, 1) \cup (1, \frac{3}{2})$.

1.1. Outline of the proof of Theorem 1.2. We proceed by contradiction and assume that Theorem 1.2 is false. Observing that Theorem 2.16 guarantees the global existence and scattering for sufficiently small initial data. From that we deduce the existence of a critical threshold size. Below this threshold, the theorem holds, but above it, solutions with arbitrarily large scattering size can be found. By employing a limiting argument, we establish the existence of minimal counterexamples, which are blowup solutions precisely at the critical threshold. Due to their minimality, these solutions exhibit compactness properties that ultimately conflict with the dispersive nature of the equation. Consequently, we can exclude their existence and conclude that Theorem 1.2 holds.

A key characteristic of these minimal counterexamples is their almost periodicity modulo the symmetries of the equation. We briefly discuss this property and its immediate implications; for a detailed analysis, the reader is referred to [33].

Definition 1.5. Let $s_c > 0$. A solution $u: I \times \Omega \to \mathbb{C}$ to (1.1) is called almost periodic if (1.3) holds and there exist function $C: \mathbb{R}^+ \to \mathbb{R}^+$ such that for all $t \in I$ and a

$$\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u(t,x)\|_{L_{x}^{2}(\Omega\cap\{x:|x|>C(\eta)\})}+\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}P_{>C(\eta)}^{\Omega}u(t,x)\|_{L_{x}^{2}(\Omega)}<\eta,$$

where $P^{\Omega}_{>N}$ denotes the Littlewood-Paley projections adapted to the Dirichlet Laplacian on Ω (c.f. (2.4)). We call C the compactness modulus function.

Remark 1.6. Using the equivalence of norms in Lemma 2.13, it is straightforward to deduce that when $\{u(t): t \in I\}$ is precompact in $\dot{H}^{s_c}_D(\Omega)$, then $u: I \times \Omega \to \mathbb{C}$ is almost periodic and there exist functions

 $C, c: \mathbb{R}^+ \to \mathbb{R}^+$ such that for all $t \in I$ and all $\eta > 0$,

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}} P_{< c(\eta)}^{\Omega} u(t, x)\|_{L_x^2(\Omega)} + \|(-\Delta_{\Omega})^{\frac{s_c}{2}} P_{> C(\eta)}^{\Omega} u(t, x)\|_{L_x^2(\Omega)} < \eta.$$

$$(1.4)$$

To proceed, we require the following result, which relates the interval length of an almost periodic solution to its Strichartz norms. This result can be established by adapting the proof of [33, Lemma 5.21] (the only difference being that we need to use the chain rule (2.1) instead of the chain rule in Euclidean space).

Lemma 1.7. Let $s_c \in [\frac{1}{2}, \frac{3}{2})$, and suppose $u : I \times \Omega \to \mathbb{C}$ is an almost periodic solution to (1.1). Then $|I| \lesssim_u \|(-\Delta_{\Omega})^{\frac{s_c}{2}} u\|_{L^2_+ L^6_{\alpha}(I \times \Omega)}^2 \lesssim_u 1 + |I|.$

With these preliminaries established, we can now describe the first major step in the proof of Theorem 1.2.

Theorem 1.8 (Reduction to almost periodic solutions). Suppose that Theorem 1.2 fails for some $s_c \in$ $[\frac{1}{2},\frac{3}{2})$. Then there exists a global solution $u:\mathbb{R}\times\Omega\to\mathbb{C}$ to (1.1) such that $u\in L^{\infty}_t\dot{H}^{s_c}_D(\mathbb{R}\times\Omega)$, whose orbit $\{u(t): t \in \mathbb{R}\}$ is precompact in $\dot{H}_D^{s_c}(\Omega)$ and there exists R > 0 such that

$$\int_{\Omega \cap \{|x| \le R\}} |u(t,x)|^{\frac{3\alpha}{2}} dx \gtrsim 1 \quad uniformly \ for \quad t \in \mathbb{R}.$$
 (1.5)

Remark 1.9. Indeed, Theorem 1.8 is valid for all $s_c \in (0, \frac{3}{2})$. The restriction $s_c \ge \frac{1}{2}$ in Theorem 1.2 arises from the limitations imposed by the indices in Theorem 2.6, which make it challenging to exclude almost periodic solutions in Theorem 1.8 when $s_c \in (0, \frac{1}{2})$. See Remark 7.6 for more details.

The reduction to almost periodic solutions is now widely regarded as a standard technique in the study of dispersive equations at critical regularity. Keraani [27] was the first to prove the existence of minimal blowup solutions, while Kenig-Merle [24] were the first to use them to establish a global well-posedness result. Since then, this technique has proven to be extremely useful; see [25,28-30,32-34,44-47] for many more examples of this technique in action (and note that this is by no means an exhaustive list). For a good introduction to these methods, see [33].

The proof of Theorem 1.8 requires three main ingredients. First, one needs the linear and nonlinear profile decomposition. For the linear profile decomposition, the case $s_c = 1$ was established in [35] and we will follow the argument presented in that paper. The profile decomposition is the key to prove the existence of minimal blow-up solutions, which will be discussed below. The tool that will be used to prove the linear profile decomposition is the inverse Strichartz inequality. This inequality indicates that the solution with non-trivial space-time bounds must concentrate at least one bubble. Applying the inverse Strichartz inequality repeatedly, it can be shown that the linear solution is concentrating at multiple bubbles with the remainder term after passing a subsequence.

Once we have obtained the linear profile decomposition, the next step is to construct the nonlinear profiles. The nonlinear profiles are the solution to NLS_{Ω} with initial data corresponding to each linear profiles. Due to the presence of the boundary, suitable scaling and space translations lead to the study of NLS in different geometries, which is the significant distinction between our setting and the Euclidean setting. The main difficulty is that we cannot know if a profile with certain initial data is also contained in the exterior domain. Moreover, it can lives at any scale and in any possible location. To deal with this, we will follow the argument established in [35] which will associated the each profiles to different limiting cases. Moreover, we will consider the following three scenarios arising from scaling and spatial translation of Ω . We denote it by the rescaled domain $\Omega_n = \lambda_n^{-1}(\Omega - \{x_n\})$ for the first two cases and $\Omega_n = \lambda_n^{-1} R_n^{-1} (\Omega - \{x_n^*\})$ where $x_n^* \in \partial \Omega$ such that $|x_n - x_n^*| = \operatorname{dist}(x_n, \Omega^c)$ and $R_n \in \operatorname{SO}(3)$ satisfies $R_n e_3 = \frac{x_n - x_n^*}{|x_n - x_n^*|}$.

- (1) When $\lambda_n \to \infty$, the rescaled domain Ω_n will approximate \mathbb{R}^3 , (2) When $\frac{\operatorname{dist}(x_n,\Omega^c)}{\lambda_n} \to \infty$, the domain Ω_n^c are retreating to infinity, (3) When $\lambda_n \to 0$ and $\frac{\operatorname{dist}(x_n,\Omega^c)}{\lambda_n} = K > 0$, the domain Ω_n will be approximated by the half-space.

The second ingredient is a stability result for the nonlinear equation (see e.g. Theorem 2.19 below). The third ingredient is a decoupling statement for nonlinear profiles. The last two ingredients are closely related, in the sense that the decoupling must hold in a space that is dictated by the stability theory. Most precisely, this means that the decoupling must hold in a space with s_c derivatives. Keraani [26] showed how to prove such a decoupling statement in the context of the mass- and energy-critical NLS; however, these arguments rely on pointwise estimates to bound the difference of nonlinearities and hence fail to be directly applicable in the presence of fractional derivatives. In [29], Killip and Visan devised a strategy that is applicable in the energy-supercritical setting, while Murphy [45] developed a strategy tailored to the energy-subcritical setting. In particular, by employing a Strichartz square function that provides estimates equivalent to those of $|\nabla|^{s_c}$, they can reduce the problem to a framework where Keraani's arguments can be directly applied. In this paper, we adopt the strategies presented in [29, 45]. Specifically, by appropriately selecting the parameters and applying the equivalence theorem (Theorem 2.6), we reduce the proof of the decoupling for nonlinear profiles to the cases addressed in [29, 45].

We therefore have all of the ingredients necessary for Theorem 1.8. Putting them together via the usual arguments, we can deduce that Theorem 1.8 holds. Hence to prove Theorem 1.2, it suffices to preclude the existence of the solutions described in Theorem 1.8. To this end, we will use versions of the Lin-Strauss Morawetz inequality

$$\int \int_{I \times \Omega} \frac{|u(t,x)|^{\alpha+2}}{|x|} dx dt \lesssim |||\nabla|^{1/2} u||_{L_t^{\infty} L_x^2(I \times \Omega)}^2, \tag{1.6}$$

which will be used in Section 6 to preclude the existence of the almost periodic solutions in Theorem 1.8 when $s_c = \frac{1}{2}$. However, when $s_c > \frac{1}{2}$, we cannot use the estimate (1.6) directly, as the solutions we consider only belong to $\dot{H}_D^{s_c}(\Omega)$ (and hence the right-hand side of (1.6) need not be finite).

For $s_c > 1/2$, one needs to suppress the low frequencies of solutions in order to access the estimate (1.6). For 3D radial energy-critical NLS [3], Bourgain accomplished this by proving a space-localized version of (1.6) (see also [18,55]). In Section 6, we adopt the this approach to preclude the existence of the almost periodic solutions in Theorem 1.8 when $1 < s_c < 3/2$. As one of the error terms resulting from space-localization requires control of the solution at the level of \dot{H}_D^1 , a different approach is needed to handle the range $\frac{1}{2} < s_c < 1$. To this end, in Section 7, we prove a version of (1.6) localized to high frequencies. This will also be used to preclude the existence of the almost periodic solutions in Theorem 1.8 when $\frac{1}{2} < s_c < 1$.

The remainder of the paper is organized as follows:

In Section 2, we record some notation and the materials that is needed for the analysis: equivalence of Sobolev spaces and the product rule for the Dirichlet Laplacian; Littlewood-Paley theory and Bernstein inequalities; Strichartz estimates; local and stability theories for (1.1); local smoothing; the convergence of functions of the Dirichlet Laplacian as the underlying domains converge; and the behavior of the linear propagator under domain convergence.

In Section 3, we first prove the refined and inverse Strichartz inequalities (Proposition 3.1 and Proposition 3.2). These results demonstrate that linear evolutions with non-trivial spacetime norms must exhibit a bubble of concentration. This is subsequently utilized to derive the linear profile decomposition for the linear propagator $e^{it\Delta_{\Omega}}$ in $\dot{H}_{D}^{s_{c}}(\Omega)$ (see Theorem 3.5 below).

In Section 4, we show that nonlinear solutions in the limiting geometries can be embedded into Ω . Since nonlinear solutions in the limiting geometries admit global spacetime bounds (Here we need to assume that Conjecture 1.1 holds true), we deduce that solutions to NLS_{Ω} , whose characteristic length scale and location conform closely with one of these limiting cases, inherit these spacetime bounds. These solutions to NLS_{Ω} will reappear as nonlinear profiles in Section 5.

In Section 5, we establish the existence of almost periodic solutions (Theorem 1.8), whose proof is primarily reduced to demonstrating the Palais-Smale condition (Proposition 5.1). To this end, we employ the profile decomposition developed in Section 4, the stability theorem (Theorem 2.19) presented in Section 2, as well as the arguments from [29, 45] to prove the decoupling for nonlinear profiles.

In Section 6, we rule out the almost periodic solutions as in Theorem 1.8 for the cases $1 < s_c < \frac{3}{2}$ and $s_c = \frac{1}{2}$. The proof is based on a space-localized Lin-Strauss Morawetz inequality as in the work of Bourgain [3].

In Section 7, we rule out solutions as in Theorem 1.8 with $\frac{1}{2} < s_c < 1$. The primary technical tool employed is the long-time Strichartz estimate (Proposition 7.1), which was originally developed by Dodson [8] for the mass-critical NLS. Then we establish a frequency-localized Lin-Strauss Morawetz inequality (Proposition 7.3) to rule out almost periodic solutions as in Theorem 1.8. In this process, we

truncate the solution to high frequencies and utilize Proposition 7.1 to control the error terms arising from the frequency projection.

2. Preliminaries

2.1. Notation and useful lemmas. We express $X \lesssim Y$ or $Y \gtrsim X$ to denote that $X \leq CY$ for some absolute constant C > 0, which might change from line to line. If the implicit constant relies on additional variables, this will be shown with subscripts. We employ O(Y) to represent any quantity X such that $|X| \lesssim Y$. The notation $X \sim Y$ implies that $X \lesssim Y \lesssim X$. The term o(1) is used to describe a quantity that converges to zero. We will also use s+ or s-, which means that there exists a small positive number ε such that it is equal to $s+\varepsilon$ or $s-\varepsilon$ respectively.

Throughout this paper, we let $s_c = \frac{3}{2} - \frac{2}{\alpha} \in (0, \frac{3}{2})$. Further restrictions on the range of s_c are imposed only in Section 6 and Section 7. Ω will stand for the exterior domain of a smooth, compact, strictly convex obstacle in \mathbb{R}^3 . Without loss of generality, we assume $0 \in \Omega^c$. The notation diam := diam(Ω^c) is used to denote the diameter of the obstacle, and $d(x) := \operatorname{dist}(x, \Omega^c)$ denotes the distance from a point $x \in \mathbb{R}^3$ to the obstacle.

We first state the Hardy inequality on the exterior domain.

Lemma 2.1 (Hardy's inequality, [36]). Let $d \ge 3$, $1 and <math>0 < s < \min\{1 + \frac{1}{p}, \frac{3}{p}\}$, then for any $f \in C_c^{\infty}(\Omega)$, we have

$$\left\| \frac{f(x)}{d(x)} \right\|_{L^p(\Omega)} \lesssim \left\| (-\Delta_{\Omega})^{\frac{s}{2}} f \right\|_{L^p(\Omega)},$$

where $d(x) = dist(x, \Omega^c)$.

We will use the following refined version of Fatou's lemma due to Brezis and Lieb.

Lemma 2.2 (Refined Fatou, [4]). Let $0 and assume that <math>\{f_n\} \subset L^p(\mathbb{R}^d)$ with $\limsup_{n \to \infty} \|f_n\|_p < \infty$. If $f_n \to f$ almost everywhere, then

$$\int_{\mathbb{R}^d} ||f_n|^p - |f_n - f|^p - |f|^p| \, dx \to 0 \quad as \quad n \to \infty.$$

In particular, $||f_n||_{L^p}^p - ||f_n - f||_{L^p}^p \to ||f||_{L^p}^p$.

The following fractional difference estimate will be used in the proof of Lemma 2.18.

Lemma 2.3 (Derivatives of differences, [29]). Let $F(u) = |u|^p u$ with p > 0 and let 0 < s < 1. Then for $1 < q, q_1, q_2 < \infty$ such that $\frac{1}{q} = \frac{1}{q_1} + \frac{p}{q_2}$, we have

$$\|\nabla|^{s}[F(u+v)-F(u)]\|_{L^{q}(\mathbb{R}^{d})} \lesssim \|\nabla|^{s}u\|_{L^{q_{1}}(\mathbb{R}^{d})}^{p}\|v\|_{L^{q_{2}}(\mathbb{R}^{d})} + \|\nabla|^{s}v\|_{L^{q_{1}}(\mathbb{R}^{d})}^{p}\|u+v\|_{L^{q_{2}}(\mathbb{R}^{d})}.$$

We will also use the following heat kernel estimate due to Q. S. Zhang [62].

Lemma 2.4 (Heat kernel estimate [62]). Let Ω denote the exterior of a smooth, compact, convex obstacle in \mathbb{R}^d for $d \geq 3$. Then there exists c > 0 such that

$$|e^{t\Delta_{\Omega}}(x,y)| \lesssim \left(\frac{d(x)}{\sqrt{t} \wedge diam} \wedge 1\right) \left(\frac{d(y)}{\sqrt{t} \wedge diam} \wedge 1\right) e^{-\frac{c|x-y|^2}{t}} t^{-\frac{d}{2}},$$

uniformly for $x, y \in \Omega$ and $t \geq 0$; recall that $A \wedge B = \min\{A, B\}$. Moreover, the reverse inequality holds after suitable modification of c and the implicit constant.

There is a natural family of Sobolev spaces associated with powers of the Dirichlet Laplacian. Our notation for these is as follows.

Definition 2.5. For $s \geq 0$ and $1 , let <math>\dot{H}_D^{s,p}(\Omega)$ and $H_D^{s,p}(\Omega)$ denote the completions of $C_c^{\infty}(\Omega)$ under the norms

$$||f||_{\dot{H}^{s,p}_D(\Omega)} := ||(-\Delta_{\Omega})^{s/2}f||_{L^p} \quad and \quad ||f||_{H^{s,p}_D(\Omega)} := ||(1-\Delta_{\Omega})^{s/2}f||_{L^p}.$$

When p=2 we write $\dot{H}^{s}_{D}(\Omega)$ and $H^{s}_{D}(\Omega)$ for $\dot{H}^{s,2}_{D}(\Omega)$ and $H^{s,2}_{D}(\Omega)$, respectively.

The following result from [37] establishes a connection between Sobolev spaces defined with respect to the Dirichlet Laplacian and those defined through conventional Fourier multipliers. The constraints on regularity s are important, as shown by counterexamples in [37].

Theorem 2.6 (Equivalence of Sobolev spaces, [37]). Let $d \geq 3$ and let Ω denote the complement of a compact convex body $\Omega^c \subset \mathbb{R}^d$ with smooth boundary. Let $1 . If <math>0 \leq s < \min\left\{1 + \frac{1}{p}, \frac{d}{p}\right\}$, then

$$\|(-\Delta_{\mathbb{R}^d})^{s/2}f\|_{L^p} \sim_{d,p,s} \|(-\Delta_{\Omega})^{s/2}f\|_{L^p}$$
 for all $f \in C_c^{\infty}(\Omega)$.

This result allows us to transfer the L^p -product rule for fractional derivatives and the chain rule directly from the Euclidean setting, provided we respect the restrictions on s and p.

Lemma 2.7. For all $f, g \in C_c^{\infty}(\Omega)$, we have

$$\|(-\Delta_{\Omega})^{s/2}(fg)\|_{L^{p}(\Omega)} \lesssim \|(-\Delta_{\Omega})^{s/2}f\|_{L^{p_{1}}(\Omega)}\|g\|_{L^{p_{2}}(\Omega)} + \|f\|_{L^{q_{1}}(\Omega)}\|(-\Delta_{\Omega})^{s/2}g\|_{L^{q_{2}}(\Omega)}$$

with the exponents satisfying $1 < p, p_1, q_2 < \infty, 1 < p_2, q_1 \le \infty$,

$$\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{q_1} + \frac{1}{q_2}, \quad and \quad 0 < s < \min\left\{1 + \frac{1}{p_1}, 1 + \frac{1}{q_2}, \frac{3}{p_1}, \frac{3}{q_2}\right\}.$$

Lemma 2.8. Suppose $G \in C^2(\mathbb{C})$ and $1 < p, p_1, p_2 < \infty$ are such that $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$. Then for all $0 < s < \min\left\{2, \frac{3}{p_2}\right\}$,

$$\|(-\Delta_{\Omega})^{\frac{s}{2}}G(u)\|_{L^{p}(\Omega)} \lesssim \|G'(u)\|_{L^{p_1}(\Omega)}\|(-\Delta_{\Omega})^{\frac{s}{2}}u\|_{L^{p_2}(\Omega)}.$$

In particular, in Section 7, we will use the following fractional chain rule:

Corollary 2.9. Given $u \in L_t^{\infty} \dot{H}_D^{s_c}(I \times \Omega) \cap L_t^2 \dot{H}_D^{s_c,6}(I \times \Omega)$,

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}}(|u|^{\alpha}u)\|_{L_t^2 L_x^{\frac{6}{5}}(I \times \Omega)} \lesssim \|(-\Delta_{\Omega})^{\frac{s_c}{2}}u\|_{L_t^{\infty} L_x^2}^{\alpha}\|(-\Delta_{\Omega})^{\frac{s_c}{2}}u\|_{L_t^2 L_x^6(I \times \Omega)}. \tag{2.1}$$

Proof. Using the equivalence theorem 2.6, the chain rule in Euclidean space, and applying the equivalence theorem 2.6 again, we obtain

$$\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}(|u|^{\alpha}u)\|_{L_{t}^{2}L_{x}^{\frac{6}{2}}(I\times\Omega)} \lesssim \|u\|_{L_{t}^{2\alpha}L_{x}^{3\alpha}(I\times\Omega)}^{\alpha}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u\|_{L_{t}^{\infty}L_{x}^{2}(I\times\Omega)}. \tag{2.2}$$

Moreover, by Sobolev embedding and H?lder's inequality, we have

$$\|u\|_{L_{t}^{2\alpha}L_{x}^{3\alpha}(I\times\Omega)}^{\alpha} \lesssim \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u\|_{L_{t}^{2\alpha}L_{x}^{\frac{6\alpha}{3\alpha-2}}(I\times\Omega)}^{\alpha} \lesssim \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u\|_{L_{t}^{\infty}L_{x}^{2}(I\times\Omega)}^{\alpha-1}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u\|_{L_{t}^{2}L_{x}^{6}(I\times\Omega)}. \tag{2.3}$$

Combining (2.2) and (2.3), we obtain the desired inequality (2.1).

We will also use the local smoothing estimate. The particular version we need is [35, Lemma 2.13].

Lemma 2.10. Let $u = e^{it\Delta_{\Omega}}u_0$. Then

$$\int_{\mathbb{R}} \int_{\Omega} |\nabla u(t,x)|^2 \langle R^{-1}(x-z) \rangle^{-3} dx dt \lesssim R \|u_0\|_{L^2(\Omega)} \|\nabla u_0\|_{L^2(\Omega)},$$

uniformly for $z \in \mathbb{R}^3$ and R > 0.

A direct consequence of the local smoothing estimate is the following result, which will be used to prove Lemma 5.2.

Corollary 2.11. Given $w_0 \in \dot{H}_D^{s_c}(\Omega)$,

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}}e^{it\Delta_{\Omega}}w_0\|_{L^2_{t,x}([\tau-T,\tau+T]\times\{|x-z|\leq R\})}\lesssim T^{\frac{2(5\alpha-4)}{10\alpha(s_c+2)}}R^{\frac{15\alpha-4}{10\alpha(s_c+2)}}\|e^{it\Delta_{\Omega}}w_0\|_{L^2_{t}(\mathbb{R}\times\Omega)}^{\frac{1}{2(s_c+2)}}\|w_0\|_{\dot{H}^{s_c}_D(\Omega)}^{1-\frac{1}{2(s_c+2)}},$$

uniformly in w_0 and the parameters $R, T > 0, \tau \in \mathbb{R}$, and $z \in \mathbb{R}^3$.

Proof. Replacing w_0 by $e^{i\tau\Delta_{\Omega}}w_0$, we see that it suffices to treat the case $\tau=0$.

Given N > 0, using the Hölder, Bernstein, and Strichartz inequalities, as well as the equivalence of Sobolev spaces, we have

$$\begin{split} & \||\nabla|^{s_c} e^{it\Delta_{\Omega}} P^{\Omega}_{< N} w_0\|_{L^2_{t,x}([-T,T] \times \{|x-z| \leq R\})} \\ & \lesssim T^{\frac{5\alpha-4}{10\alpha}} R^{\frac{3(5\alpha+4)}{40\alpha}} \||\nabla|^{s_c} e^{it\Delta_{\Omega}} P^{\Omega}_{< N} w_0\|_{L^{\frac{5\alpha}{2}}_t L^{\frac{40\alpha}{15\alpha-4}}_x} \\ & \lesssim T^{\frac{5\alpha-4}{10\alpha}} R^{\frac{3(5\alpha+4)}{40\alpha}} N^{\frac{s_c}{4}} \||\nabla|^{\frac{3}{4}s_c} e^{it\Delta_{\Omega}} P^{\Omega}_{< N} w_0\|_{L^{\frac{5\alpha}{2}}_t L^{\frac{40\alpha}{15\alpha-4}}_x} \\ & \lesssim T^{\frac{5\alpha-4}{10\alpha}} R^{\frac{3(5\alpha+4)}{40\alpha}} N^{\frac{s_c}{4}} \|e^{it\Delta_{\Omega}} P^{\Omega}_{< N} w_0\|_{L^{\frac{5\alpha}{2}}_t}^{\frac{1}{4}} \||\nabla|^{s_c} e^{it\Delta_{\Omega}} P^{\Omega}_{\leq N} w_0\|_{L^{\frac{5\alpha}{2}}_t L^{\frac{30\alpha}{15\alpha-8}}_x}^{\frac{30\alpha}{15\alpha-8}} \\ & \lesssim T^{\frac{5\alpha-4}{10\alpha}} R^{\frac{3(5\alpha+4)}{40\alpha}} N^{\frac{s_c}{4}} \|e^{it\Delta_{\Omega}} P^{\Omega}_{< N} w_0\|_{L^{\frac{1}{4}}_{t,x}}^{\frac{1}{4}} \|w_0\|_{\dot{H}^{s_c}(\Omega)}^{\frac{3}{4}}. \end{split}$$

We estimate the high frequencies using Lemma 2.10 and the Bernstein inequality:

$$\begin{split} \||\nabla|^{s_c} e^{it\Delta_{\Omega}} P_{\geq N}^{\Omega} w_0\|_{L^2_{t,x}([-T,T] \times \{|x-z| \leq R\})}^2 \\ &\lesssim R \|P_{\geq N}^{\Omega} |\nabla|^{s_c-1} w_0\|_{L^2_x} \||\nabla|^{s_c} P_{\geq N}^{\Omega} w_0\|_{L^2_x} \lesssim RN^{-1} \|w_0\|_{\dot{H}^{s_c}_D(\Omega)}^2. \end{split}$$

The desired estimate in Corollary 2.11 now follows by optimizing in the choice of N.

2.2. Littlewood-Paley theory on exterior domains. Fix $\phi : [0, \infty) \to [0, 1]$, a smooth non-negative function obeying

$$\phi(\lambda) = 1$$
 for $0 \le \lambda \le 1$, and $\phi(\lambda) = 0$ for $\lambda \ge 2$.

For each dyadic number $N \in 2^{\mathbb{Z}}$, we then define

$$\phi_N(\lambda) := \phi(\lambda/N), \quad \psi_N(\lambda) := \phi_N(\lambda) - \phi_{N/2}(\lambda);$$

notice that $\{\psi_N(\lambda)\}_{N\in\mathbb{Z}^{\mathbb{Z}}}$ forms a partition of unity for $(0,\infty)$.

With these functions in place, we can now introduce the Littlewood-Paley projections adapted to the Dirichlet Laplacian on Ω and defined via the functional calculus for self-adjoint operators:

$$P_{\leq N}^{\Omega} := \phi_N(\sqrt{-\Delta_{\Omega}}), \quad P_N^{\Omega} := \psi_N(\sqrt{-\Delta_{\Omega}}), \quad P_{>N}^{\Omega} := I - P_{\leq N}^{\Omega}. \tag{2.4}$$

For brevity we will often write $f_N := P_N^{\Omega} f$ and similarly for the other projections.

We will write $P_N^{\mathbb{R}^3}$, and so forth, to represent the analogous operators associated to the usual Laplacian in the full Euclidean space. We will also need the analogous operators on the halfspace $\mathbb{H} = \{x \in \mathbb{R}^3 : x \cdot e_3 > 0\}$ where $e_3 = (0, 0, 1)$, which we denote by $P_N^{\mathbb{H}}$, and so forth.

Just like their Euclidean counterparts, the following two basic estimates are well-known.

Lemma 2.12 (Bernstein estimates, [37]). For any $f \in C_c^{\infty}(\Omega)$, we have

$$\|P_{\leq N}^{\Omega} f\|_{L^{p}(\Omega)} + \|P_{N}^{\Omega} f\|_{L^{p}(\Omega)} \lesssim \|f\|_{L^{p}(\Omega)} \quad \text{for } 1
$$\|P_{\leq N}^{\Omega} f\|_{L^{q}(\Omega)} + \|P_{N}^{\Omega} f\|_{L^{q}(\Omega)} \lesssim N^{3\left(\frac{1}{p} - \frac{1}{q}\right)} \|f\|_{L^{p}(\Omega)} \quad \text{for } 1 \leq p < q \leq \infty,$$

$$N^{s} \|P_{N}^{\Omega} f\|_{L^{p}(\Omega)} \sim \|(-\Delta_{\Omega})^{s/2} P_{N}^{\Omega} f\|_{L^{p}(\Omega)} \quad \text{for } 1$$$$

Here, the implicit constants depend only on p, q, and s.

Lemma 2.13 (Square function estimate, [37]). Fix $1 . For all <math>f \in C_c^{\infty}(\Omega)$,

$$||f||_{L^p(\Omega)} \sim \left\| \left(\sum_{N \in 2^{\mathbb{Z}}} |P_N^{\Omega} f|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\Omega)}.$$

2.3. Strichartz estimates, local well-posedness, and the stability result. Strichartz estimates for domains exterior to a compact, smooth, strictly convex obstacle were proved by Ivanovici [21] with the exception of the endpoint $L_t^2 L_x^6$, see also [2]. Subsequently, Ivanovici and Lebeau [22] proved the dispersive estimate for d=3.

Lemma 2.14 (Dispersive estimate, [22]).

$$||e^{it\Delta_{\Omega}}f||_{L_{x}^{\infty}(\Omega)} \lesssim |t|^{-\frac{3}{2}}||f||_{L_{x}^{1}(\Omega)}.$$
 (2.5)

For $d \geq 4$, Ivanovici and Lebeau [22] also demonstrated through the construction of explicit counterexamples that the dispersive estimate no longer holds, even for the exterior of the unit ball. However, for d = 5, 7, Li-Xu-Zhang [41] established the dispersive estimates for solutions with radially symmetric initial data outside the unit ball.

Combining the dispersive estimate (2.5) with the Theorem of Keel-Tao [38], we obtain the following Strichartz estimates:

Proposition 2.15 (Strichartz estimates [2,21,22]). Let $q, \tilde{q} \geq 2$, and $2 \leq r, \tilde{r} \leq \infty$ satisfying

$$\frac{2}{q} + \frac{3}{r} = \frac{2}{\tilde{q}} + \frac{3}{\tilde{r}} = \frac{3}{2}.$$

Then, the solution u to $(i\partial_t + \Delta_\Omega)u = F$ on an interval $I \ni 0$ satisfies

$$||u||_{L_t^q L_x^r(I \times \Omega)} \lesssim ||u_0||_{L_x^2(\Omega)} + ||F||_{L_t^{\tilde{q}'} L_x^{\tilde{r}'}(I \times \Omega)}. \tag{2.3}$$

By the Strichartz estimate and the standard contraction mapping principle, we can establish the following local well-posedness result.

Theorem 2.16. Let $\Omega \subset \mathbb{R}^3$ be the exterior of a smooth compact strictly convex obstacle. There exists $\eta > 0$ such that if $u_0 \in \dot{H}^{s_c}_D(\Omega)$ obeys

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}}e^{it\Delta_{\Omega}}u_0\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)} \leq \eta \tag{2.6}$$

for some time interval $I \ni 0$, then there is a unique strong solution to (1.1) on the time interval I; moreover,

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}}u\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)} \lesssim \eta.$$

Remark 2.17.

- (1) If u_0 has small $\dot{H}_D^{s_c}(\Omega)$ norm, then Proposition 2.15 guarantees that (2.6) holds with $I = \mathbb{R}$. Thus global well-posedness for small data is a corollary of this theorem.
- (2) For large initial data u_0 , the existence of some small open interval $I \ni 0$ for which (2.6) holds follows from combining the monotone convergence theorem with Proposition 2.15. In this way, we obtain local well-posedness for all data in $\dot{H}_{D}^{c}(\Omega)$.
- (3) The argument below holds equally well for initial data prescribed as $t \to \pm \infty$, thus proving the existence of wave operators.

Proof. Throughout the proof, all space-time norms will be on $I \times \Omega$. Consider the map

$$\Phi: u \mapsto e^{it\Delta_{\Omega}}u_0 - i\int_0^t e^{i(t-s)\Delta_{\Omega}}(|u|^{\alpha}u)(s)ds.$$

We will show this is a contraction on the ball

$$B := \left\{ u \in L_t^{\infty} \dot{H}_D^{s_c} \cap L_t^{\frac{5\alpha}{2}} \dot{H}_D^{s_c, \frac{30\alpha}{15\alpha - 8}} : \| (-\Delta_{\Omega})^{\frac{s_c}{2}} u \|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}}} \le 2\eta, \right.$$

and
$$||u||_{L_t^{\infty}\dot{H}_D^{s_c}} \le 2||u_0||_{\dot{H}_D^{s_c}}, ||u||_{L_x^{\frac{5\alpha}{2}}L_x^{\frac{5\alpha}{2}}} \le 2C\eta$$

under the metric given by

$$d(u,v) := \|u - v\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}}.$$

To see that Φ maps the ball B to itself, we use the Strichartz inequality followed by Lemma 2.7, (2.6), Sobolev embedding, and then Theorem 2.6:

$$\begin{split} & \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_{c}}{2}} \Phi(u) \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}} \\ & \leq \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_{c}}{2}} e^{it\Delta_{\Omega}} u_{0} \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}} + C \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_{c}}{2}} \left(|u|^{\alpha} u \right) \right\|_{L_{t}^{\frac{5\alpha}{2(\alpha + 1)}} L_{x}^{\frac{30\alpha}{27\alpha - 8}}} \\ & \leq \eta + C \| u \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} \| \left(-\Delta_{\Omega} \right)^{\frac{s_{c}}{2}} u \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}} \leq \eta + C \| \left(-\Delta_{\Omega} \right)^{\frac{s_{c}}{2}} u \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}} \\ & \leq \eta + C (2\eta)^{\alpha + 1} \leq 2\eta, \end{split}$$

provided η is chosen sufficiently small.

Similar estimates give

$$\|\Phi(u)\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{5\alpha}{2}}} \leq C\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}\Phi(u)\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{16\alpha-8}}} \leq 2C\eta,$$

and

$$\begin{split} \|\Phi(u)\|_{L_{t}^{\infty}\dot{H}_{D}^{s_{c}}(\Omega)} &\leq \|u_{0}\|_{\dot{H}_{D}^{s_{c}}(\Omega)} + C\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}(|u|^{\alpha}u)\|_{L_{t}^{\frac{5\alpha}{2(\alpha+1)}}L_{x}^{\frac{30\alpha}{27\alpha-8}}} \\ &\leq \|u_{0}\|_{\dot{H}_{D}^{s_{c}}(\Omega)} + C\|u\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}} \\ &\leq \|u_{0}\|_{\dot{H}_{D}^{s_{c}}(\Omega)} + C(2\eta)^{\alpha+1} \leq 2\|u_{0}\|_{\dot{H}_{D}^{s_{c}}(\Omega)}, \end{split}$$

provided η is chosen small enough. This shows that Φ maps the ball B to itself.

Finally, to prove that Φ is a contraction, we argue as above:

$$\begin{split} d(\Phi(u),\Phi(v)) &\leq C \||u|^{\alpha}u - |v|^{\alpha}v\|_{L_{t}^{\frac{5\alpha}{2(\alpha+1)}}L_{x}^{\frac{30\alpha}{27\alpha-8}}} \\ &\leq Cd(u,v) \left(\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}}^{\alpha} + \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}v\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}}^{\alpha} \right) \\ &\leq 2Cd(u,v)(2\eta)^{\alpha} \leq \frac{1}{2}d(u,v), \end{split}$$

provided η is chosen small enough.

Below, we present the stability theorem for the Schrödinger equation in the exterior domain. Its proof relies on the following nonlinear estimate.

Lemma 2.18. For any $u, v \in L_t^{\frac{5\alpha}{2}} \dot{H}_D^{s_c, \frac{30\alpha}{15\alpha - 8}}(I \times \Omega)$, the following inequality holds:

$$\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}} (|u+v|^{\alpha}(u+v) - |u|^{\alpha}u) \|_{L_{t}^{\frac{5\alpha}{2(\alpha+1)}} L_{x}^{\frac{30\alpha}{2(\alpha+1)}}}$$

$$\lesssim \left(\|u\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} + \|v\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} \right) (\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}}} + \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}v\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}}})^{2},$$

$$(2.7)$$

where all the space-time integrals are over $I \times \Omega$. Note that since $s_c > 0$, we have $\alpha > \frac{4}{3}$.

Proof. We first consider the case $s_c < 1$. Applying Lemma 2.3 and the equivalence theorem 2.6, we obtain

$$\begin{split} & \| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} \left(|u+v|^{\alpha} (u+v) - |u|^{\alpha} u \right) \|_{L_{t}^{\frac{5\alpha}{2(\alpha+1)}} L_{x}^{\frac{30\alpha}{27\alpha-8}}} \\ & \lesssim \|v\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} \| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} u \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{16\alpha-8}}} + \|u+v\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} \| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{16\alpha-8}}}, \end{split}$$

which together with Sobolev embedding yields (2.7).

Next, we turn to the case $s_c > 1$. Writing $F(u) = |u|^{\alpha} u$, we have

$$|\nabla|^{s_c} (|u+v|^{\alpha}(u+v) - |u|^{\alpha}u) = |\nabla|^{s_c-1} [F'(u+v) - F'(u)] \nabla u + |\nabla|^{s_c-1} [F'(u+v) \nabla v].$$

Using the fractional differentiation rule and Sobolev embedding, we obtain

$$\begin{aligned} & \| |\nabla|^{s_{c}-1} [F'(u+v)\nabla v] \|_{L_{t}^{\frac{5\alpha}{2(\alpha+1)}} L_{x}^{\frac{30\alpha}{2'\alpha-8}}} \\ & \lesssim \| |\nabla|^{s_{c}-1} F'(u+v) \|_{L_{t}^{\frac{5}{2}} L_{x}^{\frac{5\alpha}{2(\alpha-1)}}} \| \nabla v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{15\alpha}{5\alpha+6}}} \\ & + \| u+v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}} \| |\nabla|^{s_{c}} v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}}} \\ & \lesssim \| u+v \|_{L_{t}^{\frac{\alpha-1}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} \| |\nabla|^{s_{c}} (u+v) \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}}} \| |\nabla|^{s_{c}} v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}}}. \end{aligned} \tag{2.8}$$

Similarly, using the fractional differentiation rule, Sobolev embedding, and Lemma 2.3, we have

$$\begin{split} & \| |\nabla|^{s_{c}-1} [(F'(u+v)-F'(u)) \, \nabla u] \|_{L_{t}^{\frac{5\alpha}{2}(\alpha+1)} L_{x}^{\frac{30\alpha}{27\alpha-8}}} \\ & \lesssim \| |\nabla|^{s_{c}-1} \, (F'(u+v)-F'(u)) \, \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{17\alpha-20}}} \| \nabla u \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{15\alpha}{3\alpha+6}}} + \| F'(u+v)-F'(u) \|_{L_{t}^{\frac{5}{2}} L_{x}^{\frac{5}{2}}} \| \nabla|^{s_{c}} u | \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}}} \\ & \lesssim \left(\| |\nabla|^{s_{c}-1} u \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{3\alpha-8}}} \| v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} + \| |\nabla|^{s_{c}-1} v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{3\alpha-8}}} \| u + v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} \right) \| |\nabla|^{s_{c}} u \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{3\alpha-8}}} \\ & + \left(\| u + v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} + \| v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} \right) \| v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} \| |\nabla^{s_{c}} u | \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{3\alpha-8}}} \\ & \lesssim \left(\| u \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} + \| v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}}^{\frac{5\alpha}{2}} \right) (\| |\nabla|^{s_{c}} u \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{2}}}^{\frac{30\alpha}{15\alpha-8}} + \| |\nabla|^{s_{c}} v \|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}}}^{\frac{30\alpha}{2}} \right)^{2}. \end{split} \tag{2.9}$$

Combining (2.8) and (2.9), and using the equivalence theorem 2.6, we obtain (2.7).

Now, we are in position to give the stability result for the Schrödinger equation (1.1).

Theorem 2.19 (Stability result). Let Ω be the exterior of a smooth compact strictly convex obstacle in \mathbb{R}^3 . Let I a compact time interval and let \tilde{u} be an approximate solution to (1.1) on $I \times \Omega$ in the sense that

$$i\tilde{u}_t = -\Delta_{\Omega}\tilde{u} + |\tilde{u}|^{\alpha}\tilde{u} + e \tag{2.10}$$

for some function e. Assume that

$$\|\tilde{u}\|_{L^{\infty}_t \dot{H}^{s_c}_D(I \times \Omega)} \leq E \quad and \quad \|\tilde{u}\|_{L^{\frac{5\alpha}{2}}_t L^{\frac{5\alpha}{2}}_x(I \times \Omega)} \leq L$$

for some positive constants E and L. Assume also the smallness conditions

$$\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}e^{i(t-t_{0})\Delta_{\Omega}}(u_{0}-\tilde{u}(t_{0}))\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)} \leq \varepsilon,$$

$$\|e\|_{\dot{N}^{s_{c}}((I\times\Omega))} := \inf\left\{\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}e\|_{L_{t}^{q'}L_{x}^{r'}(I\times\Omega)} : \frac{2}{q} + \frac{3}{r} = \frac{3}{2}\right\} \leq \varepsilon.$$
(2.11)

for some $0 < \varepsilon < \varepsilon_1 = \varepsilon_1(E, L)$. Then, there exists a unique strong solution $u : I \times \Omega \to \mathbb{C}$ to (1.1) with initial data u_0 at time $t = t_0$ satisfying

$$\begin{split} \|(-\Delta_{\Omega})^{\frac{s_c}{2}}(u-\tilde{u})\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)} &\leq C(E,L)\varepsilon, \\ \|(-\Delta_{\Omega})^{\frac{s_c}{2}}u\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)} &\leq C(E,L). \end{split}$$

Proof. We provide only a brief outline of the proof; the standard proof can be found in [7,49,53].

Define $w = u - \widetilde{u}$ so that $(i\partial_t + \Delta_\Omega)w = |u|^\alpha u - |\widetilde{u}|^\alpha \widetilde{u} - e$. It then follows from Lemma 2.18, Strichartz estimate, and (2.11) that

$$\begin{split} \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}w\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)} &\lesssim \varepsilon + \left(\|\widetilde{u}\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{5\alpha}{2}}(I\times\Omega)}^{\alpha-1} + \|w\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{5\alpha}{2}}(I\times\Omega)}^{\alpha-1}\right) \\ &\times \left(\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}\widetilde{u}\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)} + \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}w\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)}^{\frac{5\alpha}{2}}\right)^{2}. \end{split}$$

We first note that there exists $\delta > 0$ such that, under the additional assumption

$$\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}\widetilde{u}\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)} \leq \delta, \tag{2.12}$$

we can use the continuity method to obtain

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}}w\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{30\alpha}{15\alpha-8}}(I\times\Omega)} \lesssim \varepsilon. \tag{2.13}$$

For the general case, we divide the interval I into a finite number of smaller intervals I_j , $1 \le j \le n$, such that on each subinterval I_j , the $L_t^{\frac{5\alpha}{2}}L_x^{\frac{5\alpha}{2}}$ norm of \widetilde{u} is sufficiently small. Then using equation (2.10), the Strichartz estimate, and the continuity method on each subinterval I_j , we know that (2.12) holds on each I_j , thus obtaining that (2.13) holds on each I_j . Summing the estimates over all I_j , we obtain the desired estimate in Theorem 2.19.

2.4. Convergence results. The region Ω is not invariant under scaling or translation; indeed, under suitable choices of such operations, the obstacle may shrink to a point, march off to infinity, or even expand to fill a halfspace. In this subsection, we record from [35] some statements about the behavior of functions of the Dirichlet Laplacian under such circumstances and the convergence of propagators in Strichartz spaces. These will be used in the proof of linear profile decomposition (Proposition 3.5).

Throughout this subsection, we write

$$G_{\mathcal{O}}(x, y; \lambda) := (-\Delta_{\mathcal{O}} - \lambda)^{-1} (x, y)$$

for the Green's function of the Dirichlet Laplacian in a general open set \mathcal{O} .

Definition 2.20 ([35]). Given a sequence $\{\mathcal{O}_n\}_n$ of open subsets of \mathbb{R}^3 , we define

$$\widetilde{\lim} \mathcal{O}_n := \left\{ x \in \mathbb{R}^3 : \liminf_{n \to \infty} \operatorname{dist} (x, \mathcal{O}_n^c) > 0 \right\}.$$

Writing $\tilde{O} = \widetilde{\lim} \mathcal{O}_n$, we say $\mathcal{O}_n \to \mathcal{O}$ if the following two conditions hold: the symmetric difference $\mathcal{O} \triangle \tilde{O}$ is a finite set and

$$G_{\mathcal{O}_n}(x, y; \lambda) \to G_{\mathcal{O}}(x, y; \lambda)$$
 (2.14)

for all $\lambda \in (-2, -1)$, all $x \in \mathcal{O}$, and uniformly for y in compact subsets of $\mathcal{O} \setminus \{x\}$.

Remark 2.21. We restrict λ to the interval (-2, -1) in (2.14) for simplicity and because it allows us to invoke the maximum principle when verifying this hypothesis. Indeed, Killip-Visan-Zhang [35, Lemma 3.4] proved that this convergence actually holds for all $\lambda \in \mathbb{C} \setminus [0, \infty)$.

Given sequences of scaling and translation parameters $N_n \in 2^{\mathbb{Z}}$ and $x_n \in \Omega$, we would like to consider the domains $\Omega_n := N_n (\Omega - \{x_n\})$. When $\Omega_n \to \Omega_{\infty}$ in the sense of Definition 2.20, Killip, Visan and Zhang [35] used the maximum principle to prove the convergence of the corresponding Green's functions. Then, by applying the Helffer-Sjöstrand formula and using the convergence of the Green's functions, they obtain the following two convergence results:

Proposition 2.22. Assume $\Omega_n \to \Omega_\infty$ in the sense of Definition 2.20 and let $\Theta \in C_0^\infty((0,\infty))$. Then,

$$\|(\Theta(-\Delta_{\Omega_n}) - \Theta(-\Delta_{\Omega_{\infty}})) \,\delta_y\|_{\dot{H}^{-s_c}(\mathbb{R}^3)} \to 0 \quad \text{when} \quad n \to \infty,$$
(2.15)

uniformly for y in compact subsets of $\lim \Omega_n$. Moreover, for any fixed $t \in \mathbb{R}$ and $h \in C_0^{\infty}(\lim \Omega_n)$, we have

$$\lim_{n \to \infty} \left\| e^{it\Delta_{\Omega_n}} h - e^{it\Delta_{\Omega_\infty}} h \right\|_{\dot{H}^{-s_c}(\mathbb{R}^3)} = 0.$$

Proposition 2.23. Let $\Omega_n \to \Omega_\infty$ in the sense of Definition 2.20. Then we have

$$\left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} f - \left(-\Delta_{\Omega_{\infty}} \right)^{\frac{s_c}{2}} f \right\|_{L^2(\mathbb{R}^3)} \to 0$$

for all $f \in C_0^{\infty}(\widetilde{\lim} \Omega_n)$.

Remark 2.24. Killip, Visan and Zhang [35] proved Proposition 2.22 and Proposition 2.23 for the case when $s_c = 1$. Using their results and interpolation, we can easily extend this to the general case where $s_c \in (0, \frac{3}{2})$.

Next, we state the convergence of the Schrödinger propagators within the Strichartz norms. We rescale and translate the domain Ω to $\Omega_n = N_n * (\{\Omega\} - x_n)$ which depends on the parameters $N_n \in 2^{\mathbb{Z}}$ and $x_n \in \Omega$ conforming to one of the following three scenarios (recall that $d(x_n) := \operatorname{dist}(x_n, \Omega^c)$):

$$\begin{cases} \text{(i) } N_n \to 0 \quad \text{and} \quad -N_n x_n \to x_\infty \in \mathbb{R}^3, \\ \text{(ii) } N_n d(x_n) \to \infty, \\ \text{(iii) } N_n \to \infty \quad \text{and} \quad N_n d(x_n) \to d_\infty > 0. \end{cases}$$

Indeed, in the linear profile decomposition, there are four cases needed to be discussed (see Theorem 3.5 below). The first case will not be included in these three scenarios since there is no change of geometry in that case. In Case (i) and (ii), $\Omega_n \to \mathbb{R}^3$ while in Case (iii), $\Omega_n \to \mathbb{H}$.

After these preparation, we can state the convergence of linear Schrödinger propagators. See Theorem 4.1 and Corollary 4.2 in Killip-Visan-Zhang [35].

Theorem 2.25. Let Ω_n be as above and let Ω_{∞} be such that $\Omega_n \to \Omega_{\infty}$. Then, for all $\phi \in C_0^{\infty}(\widetilde{\lim} \Omega_n)$, $\lim_{n \to \infty} \left\| e^{it\Delta_{\Omega_n}} \phi - e^{it\Delta_{\Omega_{\infty}}} \phi \right\|_{L^{\frac{5\alpha}{2}}(\mathbb{R} \times \mathbb{R}^3)} = 0.$

3. Linear profile decomposition

In this section, we prove a linear profile decomposition for the Schrödinger propagator $e^{it\Delta_{\Omega}}$ for initial data $u_0 \in \dot{H}^{s_c}_D(\Omega)$ with $s_c \in (0, \frac{3}{2})$. For $s_c = 1$, Killip-Visan-Zhang [35] prove the linear profile decomposition for $e^{it\Delta_{\Omega}}$ by observing that the profiles can live in different limiting geometries and use the linear profile decomposition for $e^{it\Delta_{\mathbb{R}^d}}$ as a black-box. In this section, we use the linear profile decomposition for $e^{it\Delta_{\mathbb{R}^d}}$ in $\dot{H}^{s_c}(\mathbb{R}^d)$ as a black-box (see e.g. [50]), and extend the result of Killip-Visan-Zhang [35] to the general $\dot{H}^{s_c}_D(\Omega)$ setting.

Throughout this section, we denote $\Theta: \mathbb{R}^3 \to [0,1]$ the smooth function by

$$\Theta(x) = \begin{cases} 0, & |x| \leqslant \frac{1}{4}, \\ 1, & |x| \geqslant \frac{1}{2}. \end{cases}$$

We start with a refined Strichartz estimates.

Proposition 3.1 (Refined Strichartz estimate). Let $s_c \in (0, \frac{3}{2})$ and $f \in \dot{H}_D^{s_c}(\Omega)$. Then we have

$$\|e^{it\Delta_{\Omega}}f\|_{L_{t,x}^{q_0}(\mathbb{R}\times\Omega)} \lesssim \|f\|_{\dot{H}_{D}^{q_0}}^{\frac{2}{q_0}} \sup_{N\in2^{\mathbb{Z}}} \|e^{it\Delta_{\Omega}}P_{N}^{\Omega}f\|_{L_{t,x}^{q_0}(\mathbb{R}\times\Omega)}^{1-\frac{2}{q_0}}, \tag{3.1}$$

where $q_0 := \frac{10}{3-2s_c} = \frac{5\alpha}{2}$.

Proof. Throughout the proof, all space-time norms are taken over $\mathbb{R} \times \Omega$ and we set $u(t) = e^{it\Delta\Omega} f$.

We divide the proof of Proposition 3.1 into two cases.

Case One. First suppose $s_c > \frac{1}{4}$, so that $q_0 = \frac{10}{3-2s_c} > 4$. By the square function estimate (Lemma 2.13), Bernstein inequality and Strichartz estimates, we have

$$\begin{split} \|u\|_{L_{t,x}^{q_0}}^{q_0} &\lesssim \iint \left[\sum_{N} |u_N|^2\right]^{\frac{q_0}{2}} dx \, dt \lesssim \sum_{N_1 \leq N_2} \iint \left[\sum_{N} |u_N|^2\right]^{\frac{q_0}{2} - 2} |u_{N_1}|^2 |u_{N_2}|^2 \, dx \, dt \\ &\lesssim \|u\|_{L_{t,x}^{q_0}}^{q_0 - 4} \sum_{N_1 \leq N_2} \|u_{N_1}\|_{L_{t}^{q_0} L_{x}^{q_0 +}} \|u_{N_2}\|_{L_{t}^{q_0} L_{x}^{q_0 -}} \prod_{j=1}^{2} \|u_{N_j}\|_{L_{t,x}^{q_0}} \\ &\lesssim \|f\|_{\dot{H}_{D}^{s_c}}^{q_0 - 4} \sup_{N} \|u_N\|_{L_{t,x}^{q_0}}^2 \sum_{N_1 \leq N_2} \left(\frac{N_1}{N_2}\right)^{0 +} \prod_{j=1}^{2} \|u_{N_j}\|_{L_{t}^{q_0} \dot{H}_{x}^{s_c, r_0}} \\ &\lesssim \|f\|_{\dot{H}_{D}^{s_c}}^{q_0 - 4} \sup_{N} \|u_N\|_{L_{t,x}^{q_0}}^2 \sum_{N_1 \leq N_2} \left(\frac{N_1}{N_2}\right)^{0 +} \|f_{N_1}\|_{\dot{H}_{x}^{s_c}} \|f_{N_2}\|_{\dot{H}_{x}^{s_c}} \\ &\lesssim \|f\|_{\dot{H}_{D}^{s_c}}^{q_0 - 2} \sup_{N} \|e^{it\Delta_{\Omega}} f_N\|_{L_{t,x}^{q_0}}^2, \end{split}$$

where $r_0 = \frac{9+4s_c}{10}$ such that (q_0, r_0) is admissible pair. Therefore, we complete the proof of the first case.

Case Two. Suppose $\frac{1}{4} \leqslant s_c < \frac{3}{2}$, so that $2 < q_0 \le 4$. Arguing similar to the first case, we observe that

$$\begin{split} \|u\|_{L^{q_0}_{t,x}}^{q_0} &\lesssim \iint \left[\sum_N |u_N|^2\right]^{\frac{q_0}{2}} dx \, dt \lesssim \iint \left[\sum_N |u_N|^{\frac{q_0}{2}}\right]^2 dx \, dt \\ &\lesssim \sum_{N_1 \leq N_2} \iint |u_{N_1}|^{\frac{q_0}{2}} |u_{N_2}|^{\frac{q_0}{2}} \, dx \, dt \\ &\lesssim \sum_{N_1 \leq N_2} \|u_{N_1}\|_{L^{q_0}_t L^{q_0+}_x} \|u_{N_2}\|_{L^{q_0}_t L^{q_0-}_x} \prod_{j=1}^2 \|u_{N_j}\|_{L^{q_0}_{t,x}}^{\frac{q_0}{2}-1} \\ &\lesssim \sup_N \|e^{it\Delta_\Omega} f_N\|_{L^{q_0}_{t,x}}^{q_0-2} \|f\|_{L^{q_0}_t}^2, \end{split}$$

giving the desired result in this case.

The above refined Strichartz estimates show that the linear solution with nontrivial space-time norms must concentrate in an annulus. The following inverse Strichartz inequality further shows that the linear solution contains at least one bubble near some space-time point.

Proposition 3.2 (Inverse Strichartz estimate). Let $\{f_n\} \in \dot{H}_D^{s_c}(\Omega)$. Assume that

$$\lim_{n \to \infty} \|f_n\|_{\dot{H}^{s_c}_D(\Omega)} = A < \infty, \quad and \quad \lim_{n \to \infty} \left\| e^{it\Delta_{\Omega}} f_n \right\|_{L^{q_0}_{t,x}(\mathbb{R} \times \Omega)} = \varepsilon > 0. \tag{3.2}$$

Then, there exists a subsequence in n, $\{\phi_n\} \in \dot{H}^{s_c}_D(\Omega)$, $\{N_n\} \subset 2^{\mathbb{Z}}$, $\{(t_n, x_n)\} \subset \mathbb{R} \times \Omega$ conforming to one of the four scenarios below such that

$$\liminf_{n \to \infty} \|\phi_n\|_{\dot{H}_D^{s_c}(\Omega)} \gtrsim \varepsilon^{\frac{15}{s_c(4s_c+4)}} A^{\frac{4s_c^2 + 4s_c - 15}{2s_c(2s_c+2)}},\tag{3.3}$$

$$\liminf_{n \to \infty} \left\{ \|f_n\|_{\dot{H}^{s_c}_D(\Omega)}^2 - \|f_n - \phi_n\|_{\dot{H}^{s_c}_D(\Omega)}^2 - \|\phi_n\|_{\dot{H}^{s_c}_D(\Omega)}^2 \right\} \gtrsim \varepsilon^{\frac{15}{s_c(2s_c + 2)}} A^{\frac{4s_c^2 + 4s_c - 15}{s_c(2s_c + 2)}}, \tag{3.4}$$

$$\liminf_{n \to \infty} \left\{ \|e^{it\Delta_{\Omega}} f_n\|_{L^{q_0}_{t,x}(\mathbb{R} \times \Omega)}^{q_0} - \|e^{it\Delta_{\Omega}} (f_n - \phi_n)\|_{L^{q_0}_{t,x}(\mathbb{R} \times \Omega)}^{q_0} \right\} \gtrsim \varepsilon^{\frac{75}{2s_c(s_c+1)}} A^{\frac{20s_c^2 + 20s_c - 75}{2s_c(s_c+1)}}.$$
(3.5)

The four scenarios are given below:

- Case 1. $N_n \equiv N_\infty \in 2^{\mathbb{Z}}$ and $x_n \to x_\infty \in \Omega$. In this case, we take $\phi \in \dot{H}_D^{s_c}(\Omega)$ and the subsequence such that $e^{it_n\Delta_\Omega}f_n \rightharpoonup \phi$ weakly in $\dot{H}_D^{s_c}(\Omega)$ and we denote $\phi_n = e^{-it_n\Delta_\Omega}\phi$.
- Case 2. $N_n \to 0$ and $-N_n x_n \to x_\infty \in \mathbb{R}^3$. In this case, we can find $\tilde{\phi} \in \dot{H}^{s_c}(\mathbb{R}^3)$ and the subsequence such that

$$g_n(x) = N_n^{s_c - \frac{3}{2}} (e^{it_n \Delta_{\Omega}} f_n) (N_n^{-1} x + x_n) \rightharpoonup \tilde{\phi}(x)$$

weakly convergence in $\dot{H}^{s_c}(\mathbb{R}^3)$. We set

$$\phi_n(x) := N_n^{\frac{3}{2} - s_c} e^{-it_n \Delta_{\Omega}} [(\chi_n \tilde{\phi})(N_n(x - x_n))],$$

where $\chi_n(x) = \chi(N_n^{-1}x + x_n)$ and $\chi(x) = \Theta(\frac{d(x)}{\operatorname{diam}(\Omega^c)})$.

• Case 3. $N_n d(x_n) \to \infty$. In this case, we take $\tilde{\phi} \in \dot{H}^{s_c}(\mathbb{R}^3)$ and the subsequence such that

$$g_n(x) = N_n^{s_c - \frac{3}{2}} (e^{it_n \Delta_{\Omega}} f_n) (N_n^{-1} x + x_n) \rightharpoonup \tilde{\phi}(x)$$

weakly in $\dot{H}^{s_c}(\mathbb{R}^3)$. We set

$$\phi_n(x) := N_n^{\frac{3}{2} - s_c} e^{-it_n \Delta_{\Omega}} [(\chi_n \tilde{\phi})(N_n(x - x_n))],$$

where $\chi_n(x) = 1 - \Theta\left(\frac{|x|}{N_n d(x_n)}\right)$.

• Case 4. $N_n \to \infty$ and $N_n d(x_n) \to d_\infty > 0$. In this case, we find $\tilde{\phi} \in \dot{H}^{s_c}_D(\mathbb{H})$ and the subsequence such that

$$g_n(x) = N_n^{s_c - \frac{3}{2}} (e^{it_n \Delta_{\Omega}} f_n) (N_n^{-1} R_n x + x_n^*) \rightharpoonup \tilde{\phi}(x)$$

weakly in $\dot{H}^{s_c}(\mathbb{R}^3)$. We set

$$\phi_n(x) = N_n^{\frac{3}{2} - s_c} e^{-it_n \Delta_{\Omega}} [\tilde{\phi}(N_n R_n^{-1}(\cdot - x_n^*))],$$

where $R_n \in SO(3)$ satisfies $R_n e_3 = \frac{x_n - x_n^*}{|x_n - x_n^*|}$ and $x_n^* \in \partial\Omega$ such that $d(x_n) = |x_n - x_n^*|$.

Proof. Using the refined Strichartz estimate (3.1) and (3.2), we see that for each n, there exists N_n such that

$$\left\| e^{it\Delta_{\Omega}} P_{N_n}^{\Omega} f_n \right\|_{L_{+\infty}^{q_0}(\mathbb{R} \times \Omega)} \gtrsim \left\| e^{it\Delta_{\Omega}} f_n \right\|_{L_{+\infty}^{q_0}(\mathbb{R} \times \Omega)}^{\frac{q_0}{q_0 - 2}} \left\| f_n \right\|_{\dot{H}_{-\infty}^{r_c}(\Omega)}^{\frac{2}{q_0 - 2}} \gtrsim \varepsilon^{\frac{q_0}{q_0 - 2}} A^{-\frac{2}{q_0 - 2}}.$$

By Strichartz, Bernstein and (3.2), we obtain

$$\|e^{it\Delta_{\Omega}}P_{N_n}^{\Omega}f_n\|_{L_{t,n}^{q_0}(\mathbb{R}\times\Omega)} \lesssim N_n^{-s_c}A.$$

Combining the above two estimates and using Hölder's inequality, we obtain

$$\varepsilon^{\frac{q_0}{q_0-2}} A^{-\frac{2}{q_0-2}} \lesssim \|e^{it\Delta_{\Omega}} P_{N_n}^{\Omega} f_n\|_{L_{t,x}^{q_0}(\mathbb{R}\times\Omega)} \lesssim \|e^{it\Delta_{\Omega}} P_{N_n}^{\Omega} f_n\|_{L_{t,x}^{\frac{10}{3}}(\mathbb{R}\times\Omega)}^{1-\frac{2s_c}{3}} \|e^{it\Delta_{\Omega}} P_{N_n}^{\Omega} f_n\|_{L_{t,x}^{\infty}(\mathbb{R}\times\Omega)}^{\frac{2s_c}{3}} \\
\lesssim N_n^{-s_c(1-\frac{2}{3}s_c)} A^{1-\frac{2s_c}{3}} \|e^{it\Delta_{\Omega}} P_{N_n}^{\Omega} f_n\|_{L_{\infty}^{\infty}(\mathbb{R}\times\Omega)}^{\frac{2s_c}{3}},$$

which implies

$$\left\| e^{it\Delta_{\Omega}} P_{N_n}^{\Omega} f_n \right\|_{L^{\infty}_{t,x}(\mathbb{R} \times \Omega)} \gtrsim N_n^{\frac{3}{2} - s_c} \varepsilon^{\frac{15}{s_c(4s_c + 4)}} A^{\frac{4s_c^2 + 4s_c - 15}{2s_c(2s_c + 2)}}.$$

Thus there exist $x_n \in \mathbb{R}$ and $t_n \in \mathbb{R}$ such that

$$\left| \left(e^{it_n \Delta_{\Omega}} P_{N_n}^{\Omega} f_n \right) (x_n) \right| \gtrsim N_n^{\frac{3}{2} - s_c} \varepsilon^{\frac{15}{s_c (4s_c + 4)}} A^{\frac{4s_c^2 + 4s_c - 15}{2s_c (2s_c + 2)}}.$$

$$(3.6)$$

Note that the four cases in Proposition 3.2 are completely determined by the behavior of x_n and N_n . We first claim that

$$N_n d(x_n) \gtrsim \varepsilon^{\frac{15}{s_c(4s_c+4)}} A^{-\frac{15}{2s_c(2s_c+2)}}.$$
 (3.7)

Indeed, using the heat kernel bound (Lemma 2.4), we have

$$\int_{\Omega} |e^{t\Delta_{\Omega}/N_n^2}(x_n, y)|^2 dy \lesssim N_n^6 \int_{\Omega} \left| (N_n d(x_n))(N_n (d(x_n) + N_n |x_n - y|))e^{-cN_n^2 |x_n - y|^2} \right|^2 dy$$
$$\lesssim (N_n d(x_n))^2 (N_n (d(x_n) + 1))^2 N_n^3.$$

Writting

$$(e^{it_n\Delta_{\Omega}}P_{N_n}^{\Omega}f_n)(x_n) = \int_{\Omega} [e^{\Delta_{\Omega}/N_n^2}(x_n, y)P_{\leq 2N_n}^{\Omega}e^{-\Delta_{\Omega}/N_n^2}e^{it_n\Delta_{\Omega}}P_{N_n}^{\Omega}f_n](y)dy,$$

using (3.6), and Cauchy-Schwartz gives

$$N_n^{\frac{3}{2}-s_c} \varepsilon^{\frac{15}{s_c(4s_c+4)}} A^{\frac{4s_c^2+4s_c-15}{2s_c(2s_c+2)}} \lesssim (N_n d(x_n))(N_n d(x_n)+1) N_n^{\frac{3}{2}} \|P_{\leq 2N_n}^{\Omega} e^{-\Delta_{\Omega}/N_n^2} e^{it_n \Delta_{\Omega}} P_{N_n}^{\Omega} f_n\|_{L^2(\Omega)}$$

$$\lesssim (N_n d(x_n))(N_n d(x_n)+1) N_n^{\frac{3}{2}-s_c} A.$$

Then claim (3.7) follows.

Due to (3.7) and passing the subsequence, we only need to consider the following four cases:

Case 1. $N_n \sim 1$ and $N_n d(x_n) \sim 1$,

Case 2. $N_n \to 0$ and $N_n d(x_n) \lesssim 1$,

Case 3. $N_n d(x_n) \to \infty$ as $n \to \infty$,

Case 4. $N_n \to \infty$ and $N_n d(x_n) \sim 1$.

We will treat these cases in order.

Case 1. After passing through the subsequence, we may assume that

$$N_n \equiv N_\infty \in 2^{\mathbb{Z}}$$
 and $x_n \to x_\infty \in \Omega$.

Let

$$g_n(x) := N_n^{s_c - \frac{3}{2}} (e^{it_n \Delta_{\Omega}} f_n) (N_n^{-1} x + x_n).$$

Since f_n is supported in Ω , g_n is supported in $\Omega_n := N_n(\Omega - \{x_n\})$. Moreover, we have

$$||g_n||_{\dot{H}_D^{s_c}(\Omega_n)} = ||f_n||_{\dot{H}_D^{s_c}(\Omega)} \lesssim A.$$

Passing to a further subsequence, we find a $\tilde{\phi}$ such that $g_n \rightharpoonup \tilde{\phi}$ in $\dot{H}^{s_c}(\mathbb{R}^3)$ as $n \to \infty$. Rescaling this weak convergence, we have

$$e^{it_n \Delta_{\Omega}} f_n(x) \rightharpoonup \phi(x) := N_{\infty}^{\frac{3}{2} - s_c} \tilde{\phi}(N_{\infty}(x - x_{\infty})) \text{ in } \dot{H}_D^{s_c}(\Omega). \tag{3.8}$$

Since $\dot{H}^{s_c}_D(\Omega)$ is a weakly closed subset of $\dot{H}^{s_c}(\mathbb{R}^3)$, $\phi \in \dot{H}^{s_c}_D(\Omega)$.

We next prove that ϕ is non-trivial. To this end, let $h:=P_{N_{\infty}}^{\Omega}\delta_{x_{\infty}}$. Then the Bernstein inequality implies

$$\left\| \left(-\Delta_{\Omega} \right)^{-\frac{s_c}{2}} h \right\|_{L^2(\Omega)} = \left\| \left(-\Delta_{\Omega} \right)^{-\frac{s_c}{2}} P_{N_{\infty}}^{\Omega} \delta_{x_{\infty}} \right\|_{L^2(\Omega)} \lesssim N_{\infty}^{\frac{3}{2} - s_c}, \tag{3.9}$$

which reveals that $h \in \dot{H}_{D}^{-s_c}(\Omega)$. On the other hand, we note that

$$\langle \phi, h \rangle = \lim_{n \to \infty} \langle e^{it_n \Delta_{\Omega}} f_n, h \rangle = \lim_{n \to \infty} \langle e^{it_n \Delta_{\Omega}} f_n, P_{N_{\infty}}^{\Omega} \delta_{x_{\infty}} \rangle$$

$$= \lim_{n \to \infty} \left(e^{it_n \Delta_{\Omega}} P_{N_n}^{\Omega} f_n \right) (x_n) + \lim_{n \to \infty} \left\langle e^{it_n \Delta_{\Omega}} f_n, P_{N_{\infty}}^{\Omega} \left(\delta_{x_{\infty}} - \delta_{x_n} \right) \right\rangle. \tag{3.10}$$

We first claim that the second limit in (3.10) vanishes. In fact, since $d(x_n) \sim 1$, we have the following by using the Bernstein inequality,

$$\|P_{N_{\infty}}^{\Omega}e^{it_{n}\Delta_{\Omega}}f_{n}\|_{L_{\infty}^{\infty}} \lesssim N_{\infty}^{\frac{3}{2}-s_{c}}A \quad \text{ and } \quad \|\Delta P_{N_{\infty}}^{\Omega}e^{it_{n}\Delta_{\Omega}}f_{n}\|_{L_{\infty}^{\infty}} \lesssim N_{\infty}^{\frac{3}{2}+s_{c}}A.$$

Thus for sufficiently large n, it follows from the fundamental theorem of calculus, and the basic elliptic estimate

$$\|\nabla v\|_{L^{\infty}(|x| \le R)} \lesssim R^{-1} \|v\|_{L^{\infty}(|x| \le 2R)} + R\|\Delta v\|_{L^{\infty}(|x| \le 2R)},\tag{3.11}$$

that

$$\left| \left\langle e^{it_n \Delta_{\Omega}} f_n, P_{N_{\infty}}^{\Omega} \left(\delta_{x_{\infty}} - \delta_{x_n} \right) \right\rangle \right| \lesssim \left| x_{\infty} - x_n \right| \left\| \nabla P_{N_{\infty}}^{\Omega} e^{it_n \Delta_{\Omega}} f_n \right\|_{L^{\infty}(|x| \leq R)}$$

$$\lesssim \left(\frac{N_{\infty}^{\frac{3}{2} - s_c}}{d(x_n)} + N_{\infty}^{\frac{3}{2} + s_c} d(x_n) \right) A |x_{\infty} - x_n|,$$
(3.12)

which converges to zero as $n \to \infty$.

Therefore, it follows from (3.6), (3.9), (3.10), and (3.12) that

$$N_{\infty}^{\frac{3}{2}-s_{c}} \varepsilon^{\frac{15}{s_{c}(4s_{c}+4)}} A^{\frac{4s_{c}^{2}+4s_{c}-15}{2s_{c}(2s_{c}+2)}} \lesssim |\langle \phi, h \rangle| \lesssim \|\phi\|_{\dot{H}_{D}^{s_{c}}(\Omega)} \|h\|_{\dot{H}_{D}^{-s_{c}}(\Omega)} \lesssim N_{\infty}^{\frac{3}{2}-s_{c}} \|\phi\|_{\dot{H}_{D}^{s_{c}}(\Omega)}, \tag{3.13}$$

which gives (3.3).

Next, since $\dot{H}_D^{s_c}(\Omega)$ is a Hilbert space, (3.4) is a direct consequence of (3.3) and (3.8).

It remains to prove decoupling for the $L_x^{q_0}$ norm in (3.5). We note that

$$(i\partial_t)^{\frac{s_c}{2}}e^{it\Delta_{\Omega}} = (-\Delta_{\Omega})^{\frac{s_c}{2}}e^{it\Delta_{\Omega}}.$$

Thus, applying Hölder's inequality on the arbitrary compact domain $K \subset \mathbb{R} \times \mathbb{R}^3$, we have

$$\left\|e^{it\Delta_{\Omega}}e^{it_n\Delta_{\Omega}}f_n\right\|_{H^{\frac{s_c}{2}}_{+,2}(K)} \lesssim \left\|\langle -\Delta_{\Omega} \rangle^{\frac{s_c}{2}}e^{i(t+t_n)\Delta_{\Omega}}f_n\right\|_{L^2_{t,x}(K)} \lesssim_K A.$$

By Rellich-Kondrachov compactness theorem and using the diagonal argument, we get after passing the subsequence,

$$e^{it\Delta_{\Omega}}e^{it_n\Delta_{\Omega}}f_n \to e^{it\Delta_{\Omega}}\phi$$
 strongly in $L^2_{t,x}(K)$,

and

$$e^{it\Delta_{\Omega}}e^{it_n\Delta_{\Omega}}f_n \to e^{it\Delta_{\Omega}}\phi(x)$$
 a.e. in $\mathbb{R} \times \mathbb{R}^3$.

By refined Fatou Lemma 2.2, and a change of variables, we have

$$\lim_{n \to \infty} \left(\|e^{it\Delta_{\Omega}} f_n\|_{L^{q_0}_{t,x}(\mathbb{R} \times \Omega)}^{q_0} - \|e^{it\Delta_{\Omega}} (f_n - \phi_n)\|_{L^{q_0}_{t,x}(\mathbb{R} \times \Omega)}^{q_0} \right) = \|e^{it\Delta_{\Omega}} \phi\|_{L^{q_0}_{t,x}(\mathbb{R} \times \Omega)}^{q_0},$$

from which (3.5) will follow once we prove

$$\|e^{it\Delta_{\Omega}}\phi\|_{L^{q_0}_{t,x}(\mathbb{R}\times\Omega)} \gtrsim \varepsilon^{\frac{15}{s_c(4s_c+4)}} A^{\frac{4s_c^2+4s_c-15}{2s_c(2s_c+2)}}.$$
 (3.14)

We now prove (3.14). By the Mikhlin multiplier theorem, we have the uniform estimate for $|t| \leq N_{\infty}^{-2}$,

$$\left\|e^{it\Delta_{\Omega}}P_{\leq 2N_{\infty}}^{\Omega}\right\|_{L_{x}^{q'_{0}}\to L_{x}^{q'_{0}}}\lesssim 1, \quad \text{with} \quad q'_{0}=\frac{10}{2s_{c}+7}.$$

Combining this with the Bernstein inequality, we get

$$\|e^{it\Delta_{\Omega}}h\|_{L_{x}^{q'_{0}}}\lesssim \left\|e^{it\Delta_{\Omega}}P_{\leq 2N_{\infty}}^{\Omega}\right\|_{L_{x}^{q'_{0}}\to L_{x}^{q'_{0}}}\left\|P_{N_{\infty}}^{\Omega}\delta_{\infty}\right\|_{L_{x}^{q'_{0}}}\lesssim N_{\infty}^{\frac{9-6s_{c}}{10}}.$$

This together with (3.13) yields

$$N_{\infty}^{\frac{3}{2}-s_c} \varepsilon^{\frac{15}{s_c(4s_c+4)}} A^{\frac{4s_c^2+4s_c-15}{2s_c(2s_c+2)}} \lesssim |\langle \phi, h \rangle| = |\langle e^{it\Delta_{\Omega}} \phi, e^{it\Delta_{\Omega}} h \rangle| \lesssim N_{\infty}^{\frac{9-6s_c}{10}} \|e^{it\Delta_{\Omega}} \phi\|_{L_{\infty}^{q_0}(\mathbb{R} \times \Omega)},$$

uniformly for $|t| \leq N_{\infty}^{-2}$. Integrating in t then justifies (3.14).

Case 2. As $N_n \to 0$, the condition $N_n d(x_n) \lesssim 1$ indicates that the sequence $\{N_n x_n\}_{n \geq 1}$ is bounded. Thus, up to a subsequence, we suppose that $-N_n x_n \to x_\infty \in \mathbb{R}^3$ as $n \to \infty$. Similar to Case 1, we set $\Omega_n := N_n(\Omega - \{x_n\})$. Since $N_n \to 0$, we note that in this case, the rescaled obstacles Ω_n^c shrink to x_∞ when $n \to \infty$.

Noticing that f_n is bounded in $\dot{H}^{s_c}_D(\Omega)$, then the sequence g_n is bounded in $\dot{H}^{s_c}_D(\Omega_n) \subset \dot{H}^{s_c}(\mathbb{R}^3)$. Passing a subsequence, we find $\tilde{\phi}$ such that $g_n \rightharpoonup \tilde{\phi}$ in $\dot{H}^{s_c}(\mathbb{R}^3)$ when $n \to \infty$.

Next we claim that

$$\chi_n \tilde{\phi} \to \tilde{\phi}$$
, or equivalently, $(1 - \chi(N_n^{-1}x + x_n)) \tilde{\phi}(x) \to 0$ in $\dot{H}^{s_c}(\mathbb{R}^3)$. (3.15)

In fact, let

$$B_n := \left\{ x \in \mathbb{R}^3 : \operatorname{dist}(x, \Omega_n^c) \le \operatorname{diam}(\Omega_n^c) \right\}.$$

Then B_n contains supp $(1 - \chi_n)$ and supp $(\nabla \chi_n)$. Since $N_n \to 0$, the measure of B_n shrinks to zero as $n \to \infty$. (3.15) then follows from Hölder, Sobolev inequalities, fractional chain rule and the monotone convergence theorem.

With (3.15) in place, the proofs of (3.3) and (3.4) now follow their Case 1 counterparts very closely. First, we prove (3.3). We set $h := P_1^{\mathbb{R}^3} \delta_0$. Then

$$\left\langle \tilde{\phi}, h \right\rangle = \lim_{n \to \infty} \left\langle g_n, h \right\rangle = \lim_{n \to \infty} \left\langle g_n, P_1^{\Omega_n} \delta_0 \right\rangle + \lim_{n \to \infty} \left\langle g_n, \left(P_1^{\mathbb{R}^3} - P_1^{\Omega_n} \right) \delta_0 \right\rangle.$$

By Proposition 2.22 and the uniform boundedness of $||g_n||_{\dot{H}^{s_c}(\mathbb{R}^3)}$, the second term vanishes. Hence, by the definition of g_n and a change of variables, we have

$$\left| \left\langle \tilde{\phi}, h \right\rangle \right| = \left| \lim_{n \to \infty} \left\langle g_n, P_1^{\Omega_n} \delta_0 \right\rangle \right| = \left| \lim_{n \to \infty} \left\langle f_n, N_n^{s_c + \frac{3}{2}} \left(P_1^{\Omega_n} \delta_0 \right) \left(N_n (x - x_n) \right) \right\rangle \right|$$

$$= \left| \lim_{n \to \infty} \left\langle f_n, N_n^{s_c - \frac{3}{2}} P_{N_n}^{\Omega} \delta_{x_n} \right\rangle \right| \gtrsim \varepsilon^{\frac{15}{s_c (4s_c + 4)}} A^{\frac{4s_c^2 + 4s_c - 15}{2s_c (2s_c + 2)}}, \tag{3.16}$$

where the last inequality follows from (3.6). Thus, arguing as (3.13), we get

$$\|\tilde{\phi}\|_{\dot{H}^{s_c}(\mathbb{R}^3)} \gtrsim \varepsilon^{\frac{15}{s_c(4s_c+4)}} A^{\frac{4s_c^2+4s_c-15}{2s_c(2s_c+2)}},$$

which together with (3.15) yields (3.3).

To prove the decoupling estimate in $\dot{H}_{D}^{s_{c}}(\Omega)$, we write

$$||f_{n}||_{\dot{H}_{D}^{s_{c}}(\Omega)}^{2} - ||f_{n} - \phi_{n}||_{\dot{H}_{D}^{s_{c}}(\Omega)}^{2} = 2\langle f_{n}, \phi_{n} \rangle_{\dot{H}_{D}^{s_{c}}(\Omega)}^{2} - ||\phi_{n}||_{\dot{H}_{D}^{s_{c}}(\Omega)}^{2}$$

$$= 2 \left\langle N_{n}^{s_{c} - \frac{3}{2}} f_{n}(N_{n}^{-1}x + x_{n}), \tilde{\phi}(x)\chi(x) \right\rangle_{\dot{H}_{D}^{s_{c}}(\Omega_{n})}^{2} - ||\chi_{n}\tilde{\phi}||_{\dot{H}_{D}^{s_{c}}(\Omega_{n})}^{2}$$

$$= 2 \left\langle g_{n}, \tilde{\phi} \right\rangle_{\dot{H}^{s_{c}}(\mathbb{R}^{3})}^{2} - 2 \left\langle g_{n}, \tilde{\phi}(1 - \chi_{n}) \right\rangle_{\dot{H}^{s_{c}}(\mathbb{R}^{3})}^{2} - ||\chi_{n}\tilde{\phi}||_{\dot{H}_{D}^{s_{c}}(\Omega_{n})}^{2}.$$

From the weak convergence of g_n to $\tilde{\phi}$, (3.15), and (3.3), we conclude

$$\lim_{n \to \infty} \left(\|f_n\|_{\dot{H}^{s_c}_D(\Omega)}^2 - \|f_n - \phi_n\|_{\dot{H}^{s_c}_D(\Omega)}^2 \right) = \left\|\tilde{\phi}\right\|_{\dot{H}^{s_c}(\mathbb{R}^3)}^2 \gtrsim \varepsilon^{\frac{15}{s_c(2s_c + 2)}} A^{\frac{4s_c^2 + 4s_c - 15}{s_c(2s_c + 2)}}.$$

This verifies (3.4).

We now prove the decoupling for $L_{t,x}^{q_0}(\mathbb{R}\times\Omega)$ norm by showing

$$\lim_{n \to \infty} \inf \left(\|e^{it\Delta_{\Omega}} f_n\|_{L^{q_0}_{t,x}}^{q_0} - \|e^{it\Delta_{\Omega}} (f_n - \phi_n)\|_{L^{q_0}_{t,x}}^{q_0} \right) = \|e^{it\Delta_{\Omega}} \tilde{\phi}\|_{L^{q_0}_{t,x}}^{q_0}.$$
(3.17)

Notice that (3.5) then follows from the lower bound

$$\|e^{it\Delta_{\Omega}}\tilde{\phi}\|_{L_{x}^{q_{0}}}^{q_{0}} \gtrsim \left(\varepsilon^{\frac{15}{s_{c}(4s_{c}+4)}} A^{\frac{4s_{c}^{2}+4s_{c}-15}{2s_{c}(2s_{c}+2)}}\right)^{q_{0}}.$$
 (3.18)

The proof of (3.18) is almost the same as in Case 1 and we omit the details.

It remains to prove (3.17). We need two simple observations: the first one is

$$e^{it\Delta_{\Omega_n}}(g_n - \chi_n\tilde{\phi}) \to 0$$
 a.e. in $\mathbb{R} \times \mathbb{R}^3$, (3.19)

while the second one is

$$\left\| e^{it\Delta_{\Omega_n}} \chi_n \tilde{\phi} - e^{it\Delta_{\mathbb{R}^3}} \tilde{\phi} \right\|_{L^{q_0}_{*,n}(\mathbb{R} \times \mathbb{R}^3)} \to 0. \tag{3.20}$$

For (3.19), using the definition of $\tilde{\phi}$ together with (3.15), we deduce

$$g_n - \chi_n \tilde{\phi} \rightharpoonup 0$$
 weakly in $\dot{H}^{s_c}(\mathbb{R}^3)$.

Thus, by Lemma 3.4, $e^{it\Delta_{\Omega}}[g_n - \chi_n\widetilde{\phi}] \to 0$ weakly in $\dot{H}^{s_c}(\mathbb{R}^3)$ for each $t \in \mathbb{R}$. By the same argument as in Case 1, using the fact that $(i\partial_t)^{s_c/2}e^{it\Delta_{\Omega_n}} = (-\Delta_{\Omega})^{s_c/2}e^{it\Delta_{\Omega_n}}$ and passing to a subsequence, we obtain (3.19). For (3.20), we can simply use (3.15), Strichartz inequality and Theorem 2.25 to deduce the desired result.

With (3.19) and (3.20) in hand, passing to a subsequence if necessary, we obtain

$$e^{it\Delta_{\Omega_n}}g_n - e^{it\Delta_{\mathbb{R}^3}}\tilde{\phi} \to 0$$
 a.e. in $\mathbb{R} \times \mathbb{R}^3$.

Thus, by the refined Fatou lemma 2.2, we obtain

$$\liminf_{n \to \infty} \left(\|e^{it\Delta_{\Omega_n}} g_n\|_{L^{q_0}_{t,x}}^{q_0} - \|e^{it\Delta_{\Omega_n}} g_n - e^{it\Delta_{\mathbb{R}^3}} \tilde{\phi}\|_{L^{q_0}_{t,x}}^{q_0} \right) = \|e^{it\Delta_{\mathbb{R}^3}} \tilde{\phi}\|_{L^{q_0}_{t,x}}^{q_0}.$$

Combining this with (3.20), (3.15) and using a rescaling argument, we obtain (3.17).

Case 3. The proof of this case is parallel to that of Case 2. The differing geometry of the two cases enters only in the use of Proposition 2.22 and the analogue of estimate (3.15). As these first two inputs have already been proven in all cases, we only need to show

$$\chi_n \tilde{\phi} \to \tilde{\phi}$$
, or equivalently, $\Theta\left(\frac{|x|}{\operatorname{dist}(0,\Omega_n^c)}\right) \tilde{\phi}(x) \to 0 \text{ in } \dot{H}^{s_c}(\mathbb{R}^3).$ (3.21)

To do this, we let $B_n := \{x \in \mathbb{R}^3 : |x| \ge \frac{1}{4} \operatorname{dist}(0, \Omega_n^c)\}$. Then by Hölder's inequality and Sobolev embedding, we have

$$\left\|\Theta\left(\frac{|x|}{\operatorname{dist}\left(0,\Omega_{n}^{c}\right)}\right)\tilde{\phi}(x)\right\|_{\dot{H}^{s_{c}}(\mathbb{R}^{3})} \lesssim \left\|\left(-\Delta\right)^{\frac{s_{c}}{2}}\tilde{\phi}\right\|_{L^{2}(B_{n})} + \left\|\tilde{\phi}\right\|_{L^{\frac{6}{3-2s_{c}}}(B_{n})}.$$

which converges to 0 by the monotone convergence theorem.

Case 4. Passing to a subsequence, we may assume that $N_n d(x_n) \to d_\infty > 0$. By weak sequential compactness of balls in $\dot{H}^{s_c}(\mathbb{R}^3)$, we can find a subsequence and a $\tilde{\phi} \in \dot{H}^{s_c}(\mathbb{R}^3)$ such that $g_n \rightharpoonup \tilde{\phi}$ in $\dot{H}^{s_c}(\mathbb{R}^3)$. Using the following useful characterization of Sobolev spaces,

$$\dot{H}_D^{s_c}(\mathbb{H}) = \left\{ g \in \dot{H}^{s_c}(\mathbb{R}^3) : \int_{\mathbb{R}^3} g(x)\psi(x)dx = 0 \text{ for all } \psi \in C_c^{\infty}(-\mathbb{H}) \right\},$$

we find $\tilde{\phi} \in \dot{H}_D^{s_c}(\mathbb{H})$ by noticing that for any compact set K in the halfspace, $K \subset \Omega_n^c$ for n sufficiently large, where $\Omega_n := N_n R_n^{-1} (\Omega - \{x_n^*\}) \supset \operatorname{supp}(g_n)$. As $\tilde{\phi} \in \dot{H}_D^{s_c}(\mathbb{H})$, we have, as is easily seen from

$$x \in \mathbb{H} \iff N_n^{-1}R_nx + x_n^* \in \mathbb{H}_n := \left\{ y : (x_n - x_n^*)(y - x_n^*) > 0 \right\} \subset \Omega;$$

indeed, $\partial \mathbb{H}_n$ is the tangent plane to $\partial \Omega$ at x_n^* . This inclusion further shows that

$$\|\tilde{\phi}\|_{\dot{H}_{D}^{s_{c}}(\mathbb{H})} = \|\phi_{n}\|_{\dot{H}_{D}^{s_{c}}(\mathbb{H}_{n})} = \|\phi_{n}\|_{\dot{H}_{D}^{s_{c}}(\Omega)}. \tag{3.22}$$

To prove (3.3), it remains to find a lower bound on $\|\tilde{\phi}\|_{\dot{H}^{s_c}_D(\mathbb{H})}$. To this end, we let $h := P_1^{\mathbb{H}} \delta_{d_{\infty} e_3}$. Using the Bernstein inequality, we have

$$\left\| \left(-\Delta_{\mathbb{H}} \right)^{-\frac{s_c}{2}} h \right\|_{L^2(\Omega)} \lesssim 1. \tag{3.23}$$

In other words, $h \in \dot{H}_D^{-s_c}(\mathbb{H})$. Now, we set $\tilde{x}_n := N_n R_n^{-1}(x_n - x_n^*)$; by hypothesis $\tilde{x}_n \to d_\infty e_3$. It follows that from Proposition 2.22 that

$$\langle \tilde{\phi}, h \rangle = \lim_{n \to \infty} \left(\langle g_n, P_1^{\Omega_n} \delta_{\tilde{x}_n} \rangle + \langle g_n, (P_1^{\mathbb{H}} - P_1^{\Omega_n}) \delta_{d_{\infty} e_3} \rangle + \langle g_n, P_1^{\Omega_n} (\delta_{d_{\infty} e_3} - \delta_{\tilde{x}_n}) \rangle \right)$$

$$= \lim_{n \to \infty} \left(N_n^{s_c - \frac{3}{2}} (e^{it_n \Delta_{\Omega}} P_{N_n}^{\Omega} f_n)(x_n) + \langle g_n, P_1^{\Omega_n} (\delta_{d_{\infty} e_3} - \delta_{\tilde{x}_n}) \rangle \right).$$

Arguing as in the treatment of (3.12) and applying (3.11) to $v(x) = \left(P_1^{\Omega_n} g_n\right)(x + \tilde{x}_n)$ with $R = \frac{1}{2}N_n d(x_n)$, we obtain

$$\left| \left\langle g_n, P_1^{\Omega_n} \left(\delta_{d_\infty e_3} - \delta_{\tilde{x}_n} \right) \right\rangle \right| \lesssim A \left(d_\infty^{-1} + d_\infty \right) |d_\infty e_3 - \tilde{x}_n| \xrightarrow{n \to \infty} 0.$$

In conclusion, we have

$$\left|\left<\tilde{\phi},h\right>\right|\gtrsim \varepsilon^{\frac{15}{s_c(2s_c+2)}}A^{\frac{4s_c^2+4s_c-15}{s_c(2s_c+2)}},$$

which together with (3.22) and (3.23) gives (3.3).

Finally, by the same argument as that used in Case 2, we obtain (3.4). This completes the proof of Proposition 3.2.

To show the linear profile decomposition for the Schrödinger flow $e^{it\Delta_{\Omega}}$, we also need the following two weak convergence results.

Lemma 3.3 (Weak convergence). Suppose that $\Omega_n \equiv \Omega$ or $\{\Omega_n\}$ conforms to one of the last three scenarios considered in Proposition 3.2. Let $f \in C_0^{\infty}(\widetilde{\lim} \Omega_n)$ and let $\{(t_n, x_n)\}_{n \geqslant 1} \subset \mathbb{R} \times \mathbb{R}^3$. Then we have

$$e^{it_n \Delta_{\Omega_n}} f(x + x_n) \rightharpoonup 0 \tag{3.24}$$

weakly in $\dot{H}^{s_c}(\mathbb{R}^3)$ when $n \to \infty$ with $|t_n| \to \infty$ or $|x_n| \to \infty$.

Proof. Killip-Visan-Zhang [35, Lemma 5.4] proved that $\{e^{it_n\Delta_{\Omega_n}}f(x+x_n)\}_{n=1}^{\infty}$ converges weakly to zero as $n\to\infty$ in $\dot{H}^1(\mathbb{R}^3)$. Noting that $\{e^{it_n\Delta_{\Omega_n}}f(x+x_n)\}_{n=1}^{\infty}$ is also bounded in $\dot{H}^{s_c}(\mathbb{R}^3)$, it therefore converges to zero in $\dot{H}^{s_c}(\mathbb{R}^3)$ as well.

Lemma 3.4 (Weak convergence). Assume $\Omega_n \equiv \Omega$ or $\{\Omega_n\}$ conforms to one of the last three scenarios considered in Proposition 3.2. Let $f_n \in \dot{H}^{s_c}_D(\Omega_n)$ be such that $f_n \rightharpoonup 0$ weakly in $\dot{H}^{s_c}(\mathbb{R}^3)$ and let $t_n \rightarrow t_\infty \in \mathbb{R}$. Then

$$e^{it_n\Delta_{\Omega_n}}f_n \rightharpoonup 0$$
 weakly in $\dot{H}^{s_c}(\mathbb{R}^3)$.

Proof. Given any $\phi \in C_c^{\infty}(\mathbb{R}^3)$,

$$\left| \left\langle \left(e^{it_n \Delta_{\Omega_n}} - e^{it_\infty \Delta_{\Omega_n}} \right) f_n, \phi \right\rangle_{\dot{H}^{s_c}(\mathbb{R}^3)} \right| \lesssim |t_n - t_\infty|^{\frac{s_c}{2}} \| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} f_n \|_{L^2} \| \phi \|_{\dot{H}^{2s_c}},$$

which converges to zero as $n \to \infty$. To obtain the last inequality above, we have used the spectral theorem together with the elementary inequality $|e^{it_n\lambda} - e^{it_\infty\lambda}| \lesssim |t_n - t_\infty|^{s_c/2} \lambda^{s_c/2}$ for $\lambda \ge 0$. Thus, we are left to prove

$$\int_{\mathbb{R}^3} |\nabla|^{s_c} \left[e^{it_\infty \Delta_{\Omega_n}} f_n \right](x) |\nabla|^{s_c} \bar{\phi}(x) dx = \int_{\mathbb{R}^3} e^{it_\infty \Delta_{\Omega_n}} f_n(x) (-\Delta)^{s_c} \bar{\phi}(x) dx \to 0 \quad \text{as} \quad n \to \infty$$

for all $\phi \in C_0^{\infty}(\mathbb{R}^3)$. As $\{e^{it_{\infty}\Delta_{\Omega_n}}f_n\}_{n=1}^{\infty}$ is uniformly bounded in $\dot{H}^{s_c}(\mathbb{R}^3)$, it suffices to show (using the fact that the measure of $\Omega_n\Delta(\widetilde{\lim}\Omega_n)$ converges to zero)

$$\int_{\mathbb{R}^3} e^{it_\infty \Delta_{\Omega_n}} f_n(x) \bar{\phi}(x) dx \to 0 \quad \text{as} \quad n \to \infty$$
(3.25)

for all $\phi \in C_c^{\infty}(\widetilde{\lim}\Omega_n)$. To prove (3.25), we write

$$\langle e^{it_{\infty}\Delta_{\Omega_n}} f_n, \phi \rangle = \langle f_n, [e^{-it_{\infty}\Delta_{\Omega_n}} - e^{-it_{\infty}\Delta_{\Omega_{\infty}}}] \phi \rangle + \langle f_n, e^{-it_{\infty}\Delta_{\Omega_{\infty}}} \phi \rangle,$$

where Ω_{∞} denotes the limit of Ω_n . The first term converges to zero by Proposition 2.22. As $f_n \to 0$ in $\dot{H}^{s_c}(\mathbb{R}^3)$, to see that the second term converges to zero, we merely need to prove that $e^{-it_{\infty}\Delta_{\Omega_{\infty}}}\phi \in \dot{H}^{-s_c}(\mathbb{R}^3)$ for all $\phi \in C_0^{\infty}(\widetilde{\lim}\Omega_n)$. This in fact follows from the Mikhlin multiplier theorem and Bernstein's inequality:

$$\begin{split} \|e^{-it_{\infty}\Delta_{\Omega_{\infty}}}\phi\|_{\dot{H}^{-s_{c}}(\mathbb{R}^{3})} &\lesssim \|e^{-it_{\infty}\Delta_{\Omega_{\infty}}}P_{\leq 1}^{\Omega_{\infty}}\phi\|_{L^{\frac{6}{2s_{c}+3}}(\mathbb{R}^{3})} + \sum_{N\geq 1} \|e^{-it_{\infty}\Delta_{\Omega_{\infty}}}P_{N}^{\Omega_{\infty}}\phi\|_{L^{\frac{6}{2s_{c}+3}}(\mathbb{R}^{3})} \\ &\lesssim \|\phi\|_{L^{\frac{6}{2s_{c}+3}}(\mathbb{R}^{3})} + \|(-\Delta_{\Omega_{\infty}})^{2}\phi\|_{L^{\frac{6}{2s_{c}+3}}(\mathbb{R}^{3})} \lesssim_{\phi} 1. \end{split}$$

This completes the proof of the lemma.

Now, we are in position to give the linear profile decomposition for the Schrödinger propagator $e^{it\Delta_{\Omega}}$ in $\dot{H}_{D}^{s_{c}}(\Omega)$. Indeed, this follows from the application of Proposition 3.1 and 3.2.

Theorem 3.5 $(\dot{H}_{D}^{s_{c}}(\Omega))$ linear profile decomposition). Let $\{f_{n}\}_{n\geq 1}$ be a bounded sequence in $\dot{H}_{D}^{s_{c}}(\Omega)$. Passing a subsequence, there exist $J^{*} \in \{0,1,\cdots,\infty\}$, $\{\phi_{n}^{j}\}_{j=1}^{J^{*}} \subset \dot{H}_{D}^{s_{c}}(\Omega)$, $\{\lambda_{n}^{j}\}_{j=1}^{J^{*}} \subset (0,\infty)$ and $\{(t_{n},x_{n})\}_{j=1}^{J^{*}} \subset \mathbb{R} \times \Omega$ such that for each j, one of the following four cases hold

• Case 1. $\lambda_n^j \equiv \lambda_\infty^j$, $x_n^j = x_\infty^j$ and there exists a $\phi^j \in \dot{H}_D^{s_c}(\Omega)$ such that

$$\phi_n^j = e^{it_n^j(\lambda_n^j)^2 \Delta_\Omega} \phi^j.$$

We define $[G_n^j f](x) := (\lambda_n^j)^{s_c - \frac{3}{2}} f(\frac{x - x_n^j}{\lambda_n^j})$ and $\Omega_n^j = (\lambda_n^j)^{-1} (\Omega - \{x_n^j\}).$

• Case 2. $\lambda_n^j \to \infty$, $-\frac{x_n^j}{\lambda_n^j} \to x_\infty^j \in \mathbb{R}^3$. There exists a $\phi^j \in \dot{H}^{s_c}(\mathbb{R}^3)$ such that

$$\phi_n^j(x) = G_n^j \left(e^{it_n^j \Delta_{\Omega_n^j}} (\chi_n^j \phi^j) \right)(x) \quad \text{with} \quad [G_n^j f](x) := (\lambda_n^j)^{s_c - \frac{3}{2}} f\left(\frac{x - x_n^j}{\lambda^j} \right),$$

$$\Omega_n^j = (\lambda_n^j)^{-1} (\Omega - \{x_n^j\}), \qquad \chi_n^j(x) = \chi(\lambda_n^j x + x_n^j) \quad and \quad \chi(x) = \Theta\left(\frac{d(x)}{\operatorname{diam}(\Omega^c)}\right).$$

• Case 3. $\frac{d(x_n^j)}{\lambda_n^j} \to \infty$ and there exists a $\phi^j \in \dot{H}^{s_c}(\mathbb{R}^3)$ such that

$$\phi_n^j(x) := G_n^j \left(e^{it_n^j \Delta_{\Omega_n^j}} (\chi_n^j \phi^j) \right)(x) \quad \text{with} \quad [G_n^j f](x) := (\lambda_n^j)^{s_c - \frac{3}{2}} f\left(\frac{x - x_n^j}{\lambda_n^j} \right),$$

where

$$\Omega_n^j = (\lambda_n^j)^{-1} (\Omega - \{x_n^j\}), \quad and \quad \chi_n^j(x) := 1 - \Theta(\frac{\lambda_n^j |x|}{d(x_n^j)}).$$
 (3.26)

• Case 4. $\lambda_n^j \to 0$, $\frac{d(x_n^j)}{\lambda_n^j} \to \infty$ and there exists a $\phi^j \in \dot{H}^{s_c}(\mathbb{H})$ such that

$$\phi_n^j(x) := G_n^j \Big(e^{it_n^j \Delta_{\Omega_n^j}} \phi^j \Big)(x) \quad \text{with} \quad [G_n^j f](x) = (\lambda_n^j)^{s_c - \frac{3}{2}} f\Big(\frac{(R_n^j)^{-1} (x - (x_n^j)^*)}{\lambda_n^j} \Big),$$

 $\Omega_n^j = (\lambda_n^j)^{-1}(R_n^j)(\Omega - \{(x_n^j)^*\}), \ (x_n^j)^* \in \partial\Omega \ is \ chosen \ by \ d(x_n) = |x_n^j - (x_n^j)^*| \ and \ R_n^j \in SO(3)$ satisfies $R_n^j e_3 = \frac{x_n^j - (x_n^j)^*}{|x_n^j - (x_n^j)^*|}.$

Moreover, for any finite $0 \le J \le J^*$, we have the profile decomposition

$$f_n = \sum_{j=1}^{J} \phi_n^j + W_n^J,$$

where

• For all n and all $J \ge 1$, we have $W_n^J \in \dot{H}_D^{s_c}(\Omega)$, with

$$\lim_{J\to J^*}\limsup_{n\to\infty}\left\|e^{it\Delta_\Omega}W_n^J\right\|_{L^{q_0}_{t,x}(\mathbb{R}\times\Omega)}=0. \tag{3.27}$$

• For any $J \geq 1$, we have the following decoupling properties of $\dot{H}_D^{s_c}(\Omega)$ norms:

$$\lim_{n \to \infty} \left(\|f_n\|_{\dot{H}_D^{s_c}(\Omega)}^2 - \sum_{j=1}^J \|\phi_n^j\|_{\dot{H}_D^{s_c}(\Omega)}^2 - \|W_n^J\|_{\dot{H}_D^{s_c}(\Omega)}^2 \right) = 0. \tag{3.28}$$

• For any $1 \le J \le J^*$, we have

$$e^{it_n^J \Delta_{\Omega_n^J}} (G_n^J)^{-1} W_n^J \rightharpoonup 0 \quad weakly \ in \quad \dot{H}_D^{s_c}(\mathbb{R}^3).$$
 (3.29)

• For all $j \neq k$, we have the following asymptotic orthogonality property

$$\lim_{n \to \infty} \frac{\lambda_n^j}{\lambda_n^k} + \frac{\lambda_n^k}{\lambda_n^j} + \frac{|x_n^j - x_n^k|^2}{\lambda_n^j \lambda_n^k} + \frac{|t_n^j (\lambda_n^j)^2 - t_n^k (\lambda_n^k)^2|}{\lambda_n^j \lambda_n^k} = \infty.$$

$$(3.30)$$

Lastly, we may additionally assume that either $t_n^j \equiv 0$ or $|t_n^j| \to \infty$ for each j.

Proof. We will use induction argument to finish the proof and extract one bubble at each time. In the first step, we set $W_n^0 := f_n$. Suppose we have a decomposition up to level $J \geq 0$ satisfying (3.28) and (3.29). Passing to a subsequence if necessary, we denote

$$A_J := \lim_{n \to \infty} \big\| W_n^J \big\|_{\dot{H}^{s_c}_D(\Omega)} \quad \text{and} \quad \varepsilon_J := \lim_{n \to \infty} \big\| e^{it\Delta_\Omega} W_n^J \big\|_{L^{q_0}_{t,x}(\mathbb{R} \times \Omega)} \,.$$

If $\varepsilon_J = 0$, then we can stop the induction and let $J^* = J$. If not, we apply the inverse Strichartz inequality (cf.Proposition 3.2) to W_n^J . Passing to a subsequence in n, we find $\{\phi_n^{J+1}\}\subseteq \dot{H}_D^{s_c}(\Omega), \{\lambda_n^{J+1}\}\subseteq 2^{\mathbb{Z}}$, and $\{x_n^{J+1}\}\subseteq \Omega$, which conform to one of the four cases listed in the theorem. Then we rename the parameters given by Proposition 3.2 as follows: $\lambda_n^{J+1}:=N_n^{-1}$ and $t_n^{J+1}:=-N_n^2t_n$. The profiles $\tilde{\phi}^{J+1}$ are defined as weak limits in the following way

$$\tilde{\phi}^{J+1} = w \lim_{n \to \infty} (G_n^{J+1})^{-1} [e^{-it_n^{J+1} (\lambda_n^{J+1})^2 \Delta_{\Omega}} W_n^J] = w \lim_{n \to \infty} e^{-it_n^{J+1} \Delta_{\Omega_n^{J+1}}} [\left(G_n^{J+1}\right)^{-1} W_n^J],$$

where G_n^{J+1} is defined in the statement of the theorem. In Cases 2, 3, 4, we define $\phi^{J+1} := \tilde{\phi}^{J+1}$, while in Case 1,

$$\phi^{J+1}(x) := G_{\infty}^{J+1} \tilde{\phi}^{J+1}(x) := \left(\lambda_{\infty}^{J+1}\right)^{s_c - \frac{3}{2}} \tilde{\phi}^{J+1} \left(\frac{x - x_{\infty}^{J+1}}{\lambda_{\infty}^{J+1}}\right).$$

Finally, ϕ_n^{J+1} is defined as in the statement of the theorem.

For case 1, we can rewrite ϕ_n^{J+1} as

$$\phi_n^{J+1} = e^{it_n^{J+1}(\lambda_n^{J+1})^2 \Delta_{\Omega}} \tilde{\phi}^{J+1} = G_{\infty}^{J+1} e^{it_n^{J+1} \Delta_{\Omega_{\infty}^{J+1}}} \tilde{\phi}^{J+1},$$

where $\Omega_{\infty}^{J+1} := (\lambda_{\infty}^{J+1})^{-1} (\Omega - \{x_{\infty}^{J+1}\})$. We notice that in all four cases, we have

$$\lim_{n \to \infty} \left\| e^{-it_n^{J+1} \Delta_{\Omega_n^{J+1}}} \left(G_n^{J+1} \right)^{-1} \phi_n^{J+1} - \tilde{\phi}^{J+1} \right\|_{\dot{H}^{s_c}(\mathbb{R}^3)} = 0; \tag{3.31}$$

see also (3.15) and (3.21) for Cases 2 and Case 3. Next, we define $W_n^{J+1}:=W_n^J-\phi_n^{J+1}$. By (3.31) and the construction of $\tilde{\phi}^{J+1}$ in each case, we have

$$e^{-it_n^{J+1}\Delta_{\Omega_n^{J+1}}} \left(G_n^{J+1}\right)^{-1} W_n^{J+1} \rightharpoonup 0 \quad \text{ in } \dot{H}^{s_c}(\mathbb{R}^3) \quad \text{as} \quad n \to \infty,$$

which proves (3.29) at the level J+1. Moreover, (3.4) implies that

$$\lim_{n \to \infty} \left(\left\| W_n^J \right\|_{\dot{H}^{s_c}_D(\Omega)}^2 - \left\| \phi_n^{J+1} \right\|_{\dot{H}^{s_c}_D(\Omega)}^2 - \left\| W_n^{J+1} \right\|_{\dot{H}^{s_c}_D(\Omega)}^2 \right) = 0.$$

This together with the inductive hypothesis gives (3.28) at the level J+1.

From Proposition 3.2, passing to a further subsequence, we obtain

$$A_{J+1}^{2} = \lim_{n \to \infty} \|W_{n}^{J+1}\|_{\dot{H}_{D}^{s_{c}}(\Omega)}^{2} \leqslant A_{J}^{2} \left(1 - C\left(\frac{\varepsilon_{J}}{A_{J}}\right)^{\frac{15}{s_{c}(2s_{c}+2)}}\right) \leq A_{J}^{2},$$

$$\varepsilon_{J+1}^{q_{0}} = \lim_{n \to \infty} \|e^{it\Delta_{\Omega}}W_{n}^{J+1}\|_{L_{t,x}^{q_{0}}(\mathbb{R}\times\Omega)}^{q_{0}} \leq \varepsilon_{J}^{\frac{10}{3-2s_{c}}} \left(1 - C\left(\frac{\varepsilon_{J}}{A_{J}}\right)^{\frac{75}{s_{c}(2s_{c}+2)(3-2s_{c})}}\right).$$
(3.32)

If $\varepsilon_{J+1} = 0$, we can stop and set $J^* = J+1$; moreover (3.27) is automatic. If $\varepsilon_{J+1} > 0$, then we continue the induction process. If this process does not terminate in finitely many steps, then we set $J^* = \infty$; in this case, (3.32) implies $\varepsilon_J \xrightarrow{J \to \infty} 0$ and so (3.27) follows.

Next we verify the asymptotic orthogonality condition (3.30) by a contradiction argument. Assume (3.30) fails to be true for some pair (j, k). Without loss of generality, we suppose that j < k and (3.30) holds for all pairs (j, l) with j < l < k. Passing to a subsequence, we let

$$\frac{\lambda_n^j}{\lambda_n^k} \to \lambda_0 \in (0, \infty), \quad \frac{x_n^j - x_n^k}{\sqrt{\lambda_n^j \lambda_n^k}} \to x_0, \quad \text{and} \quad \frac{t_n^j (\lambda_n^j)^2 - t_n^k (\lambda_n^k)^2}{\lambda_n^j \lambda_n^k} \to t_0 \quad \text{as} \quad n \to \infty.$$
 (3.33)

From the inductive relation

$$W_n^{k-1} = W_n^j - \sum_{l=j+1}^{k-1} \phi_n^l$$

and the definition of $\tilde{\phi}^k$, we obtain

$$\begin{split} \tilde{\phi}^k &= w \lim_{n \to \infty} e^{-it_n^k \Delta_{\Omega_n^k}} \left[\left(G_n^k \right)^{-1} W_n^{k-1} \right] \\ &= w \lim_{n \to \infty} e^{-it_n^k \Delta_{\Omega_n^k}} \left[\left(G_n^k \right)^{-1} W_n^j \right] - \sum_{l=j+1}^{k-1} w \lim_{n \to \infty} e^{-it_n^k \Delta_{\Omega_n^k}} \left[\left(G_n^k \right)^{-1} \phi_n^l \right] \\ &=: A_1 + A_2. \end{split}$$

Next, we claim that the weak limits in A_1 and A_2 are zero, which would be a contradiction to $\tilde{\phi}^k \neq 0$. Rewriting A_1 as follows:

$$\begin{split} e^{-it_{n}^{k}\Delta_{\Omega_{n}^{k}}} \left[\left(G_{n}^{k} \right)^{-1} W_{n}^{j} \right] &= e^{-it_{n}^{k}\Delta_{\Omega_{n}^{k}}} \left(G_{n}^{k} \right)^{-1} G_{n}^{j} e^{it_{n}^{j}\Delta_{\Omega_{n}^{j}}} \left[e^{-it_{n}^{j}\Delta_{\Omega_{n}^{j}}} \left(G_{n}^{j} \right)^{-1} W_{n}^{j} \right] \\ &= \left(G_{n}^{k} \right)^{-1} G_{n}^{j} e^{i \left(t_{n}^{j} - t_{n}^{k} \frac{(\lambda_{n}^{k})^{2}}{(\lambda_{n}^{j})^{2}} \right) \Delta_{\Omega_{n}^{j}}} \left[e^{-it_{n}^{j}\Delta_{\Omega_{n}^{j}}} \left(G_{n}^{j} \right)^{-1} W_{n}^{j} \right]. \end{split}$$

Note that by (3.33),

$$t_n^j - t_n^k \frac{(\lambda_n^k)^2}{(\lambda_n^j)^2} = \frac{t_n^j (\lambda_n^j)^2 - t_n^k (\lambda_n^k)^2}{\lambda_n^j \lambda_n^k} \cdot \frac{\lambda_n^k}{\lambda_n^j} \to \frac{t_0}{\lambda_0}.$$
(3.34)

Using this together with (3.29), Lemma 3.4, and the fact that the adjoints of the unitary operators $(G_n^k)^{-1}G_n^j$ converge strongly, we obtain $A_1=0$.

To finish the proof of (3.30), it suffices to show $A_2 = 0$. For all j < l < k, we write

$$e^{-it_n^k\Delta_{\Omega_n^k}}\left[\left(G_n^k\right)^{-1}\phi_n^l\right] = \left(G_n^k\right)^{-1}G_n^j e^{i\left(t_n^j - t_n^k\frac{\left(\lambda_n^k\right)^2}{\left(\lambda_n^j\right)^2}\right)\Delta_{\Omega_n^j}}\left[e^{-it_n^j\Delta_{\Omega_n^j}}(G_n^j)^{-1}\phi_n^l\right].$$

By (3.34) and Lemma 3.4, it suffices to show

$$e^{-it_n^j\Delta_{\Omega_n^j}}\left[\left(G_n^j\right)^{-1}\phi_n^l\right] \rightharpoonup 0 \quad \text{weakly in} \quad \dot{H}^{s_c}(\mathbb{R}^3).$$

By density, this reduces to prove the following: for all $\phi \in C_0^{\infty}\left(\widetilde{\lim} \Omega_n^l\right)$,

$$I_n := e^{-it_n^j \Delta_{\Omega_n^j}} (G_n^j)^{-1} G_n^l e^{it_n^l \Delta_{\Omega_n^l}} \phi \rightharpoonup 0 \quad \text{weakly in} \quad \dot{H}^{s_c}(\mathbb{R}^3) \quad \text{as} \quad n \to \infty.$$
 (3.35)

Depending on which cases j and l fall into, we can rewrite I_n as follows:

• Case (a): If both j and l conform to Case 1, 2, or 3, then

$$I_n = \left(\frac{\lambda_n^j}{\lambda_n^l}\right)^{\frac{3}{2} - s_c} \left[e^{i\left(t_n^l - t_n^j \left(\frac{\lambda_n^j}{\lambda_n^l}\right)^2\right) \Delta_{\Omega_n^l}} \phi \right] \left(\frac{\lambda_n^j x + x_n^j - x_n^l}{\lambda_n^l}\right).$$

• Case (b): If j conforms to Case 1, 2, or 3 and l to Case 4, then

$$I_n = \left(\frac{\lambda_n^j}{\lambda_n^l}\right)^{\frac{3}{2} - s_c} \left[e^{i\left(t_n^l - t_n^j \left(\frac{\lambda_n^j}{\lambda_n^l}\right)^2\right) \Delta_{\Omega_n^l}} \phi \right] \left(\frac{(R_n^l)^{-1} (\lambda_n^j x + x_n^j - (x_n^l)^*)}{\lambda_n^l}\right).$$

 \bullet Case (c): If j conforms to Case 4 and l to Case 1, 2, or 3, then

$$I_n = \left(\frac{\lambda_n^j}{\lambda_n^l}\right)^{\frac{3}{2} - s_c} \left[e^{i\left(t_n^l - t_n^j \left(\frac{\lambda_n^j}{\lambda_n^l}\right)^2\right) \Delta_{\Omega_n^l}} \phi \right] \left(\frac{R_n^j \lambda_n^j x + (x_n^j)^* - x_n^l}{\lambda_n^l}\right).$$

• Case (d): If both j and l conform to Case 4, then

$$I_n = \left(\frac{\lambda_n^j}{\lambda_n^l}\right)^{\frac{3}{2} - s_c} \left[e^{i\left(t_n^l - t_n^j \left(\frac{\lambda_n^j}{\lambda_n^l}\right)^2\right) \Delta_{\Omega_n^l}} \phi \right] \left(\frac{(R_n^l)^{-1} (R_n^j \lambda_n^j x + (x_n^j)^* - (x_n^l)^*)}{\lambda_n^l}\right).$$

We first prove (3.35) when the scaling parameters are not comparable, that is,

$$\lim_{n \to \infty} \left(\frac{\lambda_n^j}{\lambda_n^l} + \frac{\lambda_n^l}{\lambda_n^j} \right) = \infty. \tag{3.36}$$

In this case, we shall treat all the four cases simultaneously. By the Cauchy-Schwarz inequality and (3.36), for all $\psi \in C_c^{\infty}(\mathbb{R}^3)$, we have

$$\begin{split} \left| \langle I_n, \psi \rangle_{\dot{H}^{s_c}(\mathbb{R}^3)} \right| &\lesssim \min \left(\| (-\Delta)^{s_c} I_n \|_{L^2(\mathbb{R}^3)} \| \psi \|_{L^2(\mathbb{R}^3)}, \| I_n \|_{L^2(\mathbb{R}^3)} \| (-\Delta)^{s_c} \psi \|_{L^2(\mathbb{R}^3)} \right) \\ &\lesssim \min \left(\left(\frac{\lambda_n^j}{\lambda_n^l} \right)^{s_c} \| (-\Delta)^{s_c} \phi \|_{L^2(\mathbb{R}^3)} \| \psi \|_{L^2(\mathbb{R}^3)}, \left(\frac{\lambda_n^l}{\lambda_n^j} \right)^{s_c} \| \phi \|_{L^2(\mathbb{R}^3)} \| (-\Delta)^{s_c} \psi \|_{L^2(\mathbb{R}^3)} \right), \end{split}$$

which converges to zero when $n \to \infty$. Thus, in this case $A_2 = 0$ and we get the desired contradiction. Hence, we may assume

$$\lim_{n \to \infty} \frac{\lambda_n^j}{\lambda_n^l} = \lambda_0 \in (0, \infty).$$

Arguing like the previous case, we assume the time parameters diverge, i.e.

$$\lim_{n \to \infty} \frac{|t_n^j(\lambda_n^j)^2 - t_n^l(\lambda_n^l)^2|}{\lambda_n^j \lambda_n^l} = \infty.$$
(3.37)

Then, we get

$$|t_n^l - t_n^j \frac{(\lambda_n^j)^2}{(\lambda_n^l)^2}| = \frac{|t_n^l (\lambda_n^l)^2 - t_n^j (\lambda_n^j)^2|}{\lambda_n^j \lambda_n^l} \cdot \frac{\lambda_n^j}{\lambda_n^l} \to \infty$$

as $n \to \infty$.

First, we consider Case (a). By (3.37) and Lemma 3.3, we have that

$$\lambda_0^{\frac{3}{2}-s_c} \left(e^{i(t_n^l - t_n^j (\frac{\lambda_n^j}{\lambda_n^l})^2) \Delta_{\Omega_n^l}} \phi \right) (\lambda_0 x + (\lambda_n^l)^{-1} (x_n^j - x_n^l)) \rightharpoonup 0 \quad \text{weakly in} \quad \dot{H}^{s_c}(\mathbb{R}^3),$$

which implies (3.35). In Cases (b), (c), and (d), the proof proceeds similarly since SO(3) is a compact group; indeed, passing to a subsequence we may assume that $R_n^j \to R_0$ and $R_n^l \to R_1$, which places us in the same situation as in Case (a).

Finally, we treat the case that

$$\frac{\lambda_n^j}{\lambda_n^l} \to \lambda_0, \quad \frac{t_n^l(\lambda_n^l)^2 - t_n^j(\lambda_n^j)^2}{\lambda_n^j \lambda_n^l} \to t_0 \quad \text{but} \quad \frac{|x_n^j - x_n^l|^2}{\lambda_n^j \lambda_n^l} \to \infty.$$
 (3.38)

Then we also have $t_n^l - t_n^j (\lambda_n^j)^2 / (\lambda_n^l)^2 \to \lambda_0 t_0$. Thus for Case (a), it sufficies to show that

$$\lambda_0^{\frac{3}{2}-s_c} e^{it_0 \lambda_0 \Delta_{\Omega_n^l}} \phi(\lambda_0 x + y_n) \rightharpoonup 0 \quad \text{weakly in} \quad \dot{H}^{s_c}(\mathbb{R}^3), \tag{3.39}$$

where

$$y_n := \frac{x_n^j - x_n^l}{\lambda_n^l} = \frac{x_n^j - x_n^l}{(\lambda_n^l \lambda_n^j)^{\frac{1}{2}}} \cdot \sqrt{\frac{\lambda_n^j}{\lambda_n^l}} \to \infty \quad \text{as} \quad n \to \infty.$$

The desired weak convergence (3.39) follows from the weak convergence Lemma 3.3.

Since SO(3) is a compact group, in Case (b) we can proceed similarly if we can show

$$\frac{|x_n^j - (x_n^l)^*|}{\lambda_n^l} \to \infty \quad \text{as} \quad n \to \infty.$$
 (3.40)

then (3.35) follows similar to Case (a). In fact, (3.40) follows from the triangle inequality

$$\frac{|x_n^j-(x_n^l)^*|}{\lambda_n^l} \geq \frac{|x_n^j-x_n^l|}{\lambda_n^l} - \frac{|x_n^l-(x_n^l)^*|}{\lambda_n^l} \geq \frac{|x_n^j-x_n^l|}{\lambda_n^l} - 2d_\infty^l \to \infty.$$

Case (c) is symmetric to Case (b), thus we have Case (c) immediately.

Now, we handle case (d). For sufficiently large n, we have

$$\frac{\left|(x_n^j)^* - (x_n^l)^*\right|}{\lambda_n^l} \ge \frac{\left|x_n^j - x_n^l\right|}{\lambda_n^l} - \frac{\left|x_n^j - (x_n^j)^*\right|}{\lambda_n^l} - \frac{\left|x_n^l - (x_n^l)^*\right|}{\lambda_n^l}$$

$$\ge \frac{\left|x_n^j - x_n^l\right|}{\sqrt{\lambda_n^l \lambda_n^j}} \cdot \sqrt{\frac{\lambda_n^j}{\lambda_n^l}} - \frac{d(x_n^j)\lambda_n^j}{\lambda_n^j \lambda_n^l} - \frac{d(x_n^l)}{\lambda_n^l}$$

$$\ge \frac{1}{2}\sqrt{\lambda_0} \frac{\left|x_n^j - x_n^l\right|}{\sqrt{\lambda_n^l \lambda_n^j}} - 2\lambda_0 d_\infty^j - 2d_\infty^l \to \infty \quad \text{as} \quad n \to \infty.$$

The desired weak convergence follows again from Lemma 3.3.

4. Embedding of nonlinear profiles

In Section 5, we will use the linear profile decomposition obtained in the previous section to prove Theorem 1.8. The most critical step is to derive a Palais-Smale condition for minimizing sequences of blowup solutions to (1.1). This essentially reduces to proving a nonlinear profile decomposition for solutions to NLS_{Ω} . To achieve this, we must address the possibility that the nonlinear profiles we extract are solutions to the \dot{H}^{s_c} -critical equation in different limiting geometries. In this section, we will see how to embed these nonlinear profiles corresponding to different limiting geometries back inside Ω , following the arguments presented in [35]. Since nonlinear solutions in the limiting geometries admit global spacetime bounds, we deduce that solutions to NLS_{Ω} , corresponding to Cases 2, 3, and 4 in Theorem 3.5 respectively, inherit these spacetime bounds. These solutions to NLS_{Ω} reappear as nonlinear profiles in Proposition 5.1. This section contains three theorems, Theorems 4.1, 4.3, and 4.4, corresponding to Cases 2, 3, and 4 in Theorem 3.5, respectively.

As in the previous section, we denote $\Theta: \mathbb{R}^3 \to [0,1]$ the smooth function such that

$$\Theta(x) = \begin{cases} 0, & |x| \le \frac{1}{4}, \\ 1, & |x| \ge \frac{1}{2}. \end{cases}$$

Our first result in this section consider the scenario when the rescaled obstacles Ω_n^c are shrinking to a point (i.e. Case 2 in Theorem 3.5).

Theorem 4.1 (Embedding nonlinear profiles for shrinking obstacles). Let $\{\lambda_n\} \subset \mathbb{Z}^{\mathbb{Z}}$ be such that $\lambda_n \to \infty$. Let $\{t_n\} \subset \mathbb{R}$ be such that $t_n \equiv 0$ or $|t_n| \to \infty$. Suppose that $\{x_n\} \subset \Omega$ satisfies $-\lambda_n^{-1}x_n \to x_\infty \in \mathbb{R}^3$. Let $\phi \in \dot{H}^{s_c}(\mathbb{R}^3)$ and

$$\phi_n(x) := \lambda_n^{s_c - \frac{3}{2}} e^{it_n \lambda_n^2 \Delta_{\Omega}} \left[(\chi_n \phi) \left(\frac{x - x_n}{\lambda_n} \right) \right],$$

where $\chi_n(x) = \chi(\lambda_n x + x_n)$ with $\chi(x) = \Theta(\frac{d(x)}{diam \Omega^c})$. Then for n sufficiently large, there exists a global solution v_n to (1.1) with initial data $v_n(0) = \phi_n$ such that

$$||v_n||_{L_{t,\sigma}^{\frac{5\alpha}{2}}(\mathbb{R}\times\Omega)}\lesssim 1,$$

with the implicit constant depending only on $\|\phi\|_{\dot{H}^{s_c}}$. Moreover, for any $\varepsilon > 0$, there exists $N_{\varepsilon} \in \mathbb{N}$ and $\psi_{\varepsilon} \in C_0^{\infty}(\mathbb{R} \times \mathbb{R}^3)$ such that for all $n \geq N_{\varepsilon}$

$$\left\| (-\Delta_{\Omega})^{\frac{s_c}{2}} \left[v_n(t - \lambda_n^2 t_n, x + x_n) - \lambda_n^{s_c - \frac{3}{2}} \psi_{\varepsilon}(\lambda_n^{-2} t, \lambda_n^{-1} x) \right] \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \mathbb{R}^3)} < \varepsilon. \tag{4.1}$$

Proof. Our proof follows the idea of [35, Theorem 6.1]. For the first step, we will construct the global solution to \dot{H}^{s_c} -critical NLS in the limiting geometry of Ω_n .

Step 1: Constructing the global solution to $NLS_{\mathbb{R}^3}$.

Let $\theta = \frac{1}{100(\alpha+1)}$. The construction of the global solution on \mathbb{R}^3 depends on the choice of time parameter t_n . If $t_n \equiv 0$, let w_n and w_∞ be the solutions to $\mathrm{NLS}_{\mathbb{R}^3}$ with initial data $w_n(0) = \phi_{\leq \lambda_n^{\theta}}$ and $w_\infty(0) = \phi$.

Otherwise, if $t_n \to \pm \infty$, let w_n be the solutions to $NLS_{\mathbb{R}^3}$ such that

$$\|w_n(t) - e^{it\Delta}\phi_{\leq \lambda_n^{\theta}}\|_{\dot{H}^{s_c}(\mathbb{R}^3)} \to 0$$
, as $t \to \pm \infty$.

Similarly, we denote w_{∞} by the solution to $\mathrm{NLS}_{\mathbb{R}^3}$ such that

$$\|w_{\infty}(t) - e^{it\Delta}\phi\|_{\dot{H}^{s_c}(\mathbb{R}^3)} \to 0, \quad \text{as} \quad t \to \pm \infty.$$
 (4.2)

By [45] and assumption made in Theorem 1.2, both $w_n(t)$ and $w_{\infty}(t)$ are global solutions and satisfy

$$||w_n||_{L_{\star}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}(\mathbb{R} \times \mathbb{R}^3)} + ||w_{\infty}||_{L_{\star}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}(\mathbb{R} \times \mathbb{R}^3)} \lesssim 1.$$
 (4.3)

Moreover, by the perturbation theory in [45],

$$\lim_{n \to \infty} \|w_n(t) - w_{\infty}(t)\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}(\mathbb{R} \times \mathbb{R}^3)} = 0.$$
 (4.4)

From the Bernstein inequality, we have

$$\|\phi_{\leq \lambda_n^{\theta}}\|_{\dot{H}^s(\mathbb{R}^3)} \lesssim \lambda_n^{\theta(s-s_c)}, \quad \text{for any} \quad s \geqslant s_c.$$

The persistence of regularity yields that

$$\||\nabla|^s w_n\|_{\dot{S}^{s_c}(\mathbb{R}\times\mathbb{R}^3)} \lesssim \lambda_n^{\theta s}$$
 for any $s \geqslant 0$,

which together with the Gagliardo-Nirenberg inequality

$$\|f\|_{L^{\infty}(\mathbb{R}^{3})} \lesssim \|f\|_{\dot{H}^{s_{c}}(\mathbb{R}^{3})}^{\frac{1}{2}} \|f\|_{\dot{H}^{3-s_{c}}(\mathbb{R}^{3})}^{\frac{1}{2}}$$

implies that

$$\||\nabla|^s w_n\|_{L^{\infty}(\mathbb{R}\times\mathbb{R}^3)} \lesssim \lambda_n^{\theta(s+\frac{3}{2}-s_c)}, \quad \text{for all} \quad s \ge 0.$$
 (4.5)

Finally, using the structure of the $NLS_{\mathbb{R}^3}$, we have

$$\|\partial_t w_n\|_{L^{\infty}_{t,x}(\mathbb{R}\times\mathbb{R}^3)} \lesssim \|\Delta w_n\|_{L^{\infty}_{t,x}(\mathbb{R}\times\mathbb{R}^3)} + \|w_n\|_{L^{\infty}_{t,x}(\mathbb{R}\times\mathbb{R}^3)}^{\alpha+1} \lesssim \lambda_n^{\theta(\frac{7}{2}-s_c)}.$$
(4.6)

Step 2. Constructing the approximate solution to (1.1).

As discussed in Case 2 of Proposition 3.2, we let $\Omega_n = \lambda_n^{-1}(\Omega - \{x_n\})$. One may want to embed $w_n(t)$ to Ω_n by taking $\tilde{v}_n(t) = \chi_n w_n(t)$ directly. However, this is not a approximation of (1.1). Instead, we take

$$z_n(t) := i \int_0^t e^{i(t-\tau)\Delta_{\Omega_n}} (\Delta_{\Omega_n} \chi_n) w_n(\tau, -\lambda_n^{-1} x_n) d\tau.$$

This can allow us to control the reflected waves near the boundary. Moreover, we have the following properties.

Lemma 4.2. For all T > 0, we have

$$\lim_{n \to \infty} \| (-\Delta_{\Omega})^{\frac{s_c}{2}} z_n \|_{L_{+}^{\frac{5\alpha}{2}} L_{\pi}^{\frac{20\alpha}{15\alpha - 8}}([-T, T] \times \Omega_n)} = 0, \tag{4.7}$$

$$\left\| (-\Delta_{\Omega_n})^{\frac{s}{2}} z_n \right\|_{L_t^{\infty} L_x^2([-T,T] \times \Omega_n)} \lesssim \lambda_n^{s - \frac{3}{2} + \theta(\frac{7}{2} - s_c)} (T + \lambda_n^{-2\theta}) \quad \text{for all} \quad 0 \le s < \frac{3}{2}.$$
 (4.8)

Proof. Integrating by parts, we write

$$z_n(t) = -\int_0^t \left(e^{it\Delta_{\Omega_n}} \partial_{\tau} e^{-i\tau\Delta_{\Omega_n}} \chi_n \right) w_n(\tau, -\lambda_n^{-1} x_n) d\tau$$
$$= -\chi_n w_n(t, -\lambda_n^{-1} x_n) + e^{it\Delta_{\Omega_n}} \left(\chi_n w_n(0, -\lambda_n^{-1} x_n) \right)$$
$$+ \int_0^t \left(e^{i(t-\tau)\Delta_{\Omega_n}} \chi_n \right) \partial_{\tau} w_n(\tau, -\lambda_n^{-1} x_n) d\tau.$$

By the Strichartz estimate, the equivalence of Sobolev norms, (4.5) and (4.6), we have

$$\begin{split} & \left\| (-\Delta_{\Omega_n})^{\frac{s}{2}} z_n \right\|_{L_t^{\infty} L_x^2([-T,T] \times \Omega_n)} \\ & \lesssim \left\| (-\Delta)^{\frac{s}{2}} \chi_n w_n(t, -\lambda_n^{-1} x_n) \right\|_{L_t^{\infty} L_x^2([-T,T] \times \Omega_n)} + \left\| (-\Delta_{\Omega_n})^{\frac{s}{2}} \chi_n w_n(0, -\lambda_n^{-1} x_n) \right\|_{L^2([-T,T] \times \Omega_n)} \\ & + \left\| (-\Delta)^{\frac{s}{2}} \chi_n \partial_t w_n(t, -\lambda_n^{-1} x_n) \right\|_{L_t^1 L_x^2([-T,T] \times \Omega_n)} \\ & \lesssim \lambda_n^{s - \frac{3}{2} + \theta(\frac{3}{2} - s_c)} + T \lambda_n^{s - \frac{3}{2} + \theta(\frac{7}{2} - s_c)}. \end{split}$$

This proves (4.8). By a similar argument, we can prove (4.7). This completes the proof of lemma 4.2.

We are now in a position to construct the approximate solution

$$\tilde{v}_n(t,x) := \begin{cases} \lambda_n^{s_c - \frac{3}{2}} (\chi_n w_n + z_n) (\lambda_n^{-2} t, \lambda_n^{-1} (x - x_n)), & |t| \leqslant \lambda_n^2 T, \\ e^{i(t - \lambda_n^2 T) \Delta_{\Omega}} \tilde{v}_n(\lambda_n^2 T, x), & t > \lambda_n^2 T, \\ e^{i(t + \lambda_n^2 T) \Delta_{\Omega}} \tilde{v}_n(-\lambda_n^2 T, x), & t < -\lambda_n^2 T, \end{cases}$$

where T > 0 is a parameter to be taken later. We first note that \tilde{v}_n has finite scattering norm. In fact, it follows from Lemma 4.2, the Strichartz estimate and change of variables that

$$\|\tilde{v}_{n}\|_{L_{t,x}^{\frac{5\alpha}{2}}(\mathbb{R}\times\Omega)} \lesssim \|\chi_{n}w_{n} + z_{n}\|_{L_{t,x}^{\frac{5\alpha}{2}}([-T,T]\times\Omega)} + \|(\chi_{n}w_{n} + z_{n})(\pm T)\|_{\dot{H}_{D}^{s_{c}}(\Omega_{n})}$$

$$\lesssim \|w_{n}\|_{L_{t,x}^{\frac{5\alpha}{2}}(\mathbb{R}\times\mathbb{R}^{3})} + \|z_{n}\|_{L_{t,x}^{\frac{5\alpha}{2}}([-T,T]\times\Omega)} + \|\chi_{n}\|_{L_{x}^{\infty}(\Omega_{n})} \||\nabla|^{s_{c}}w_{n}\|_{L_{t}^{\infty}L_{x}^{2}(\mathbb{R}\times\mathbb{R}^{3})}$$

$$+ \||\nabla|^{s_{c}}\chi_{n}\|_{L_{t}^{\frac{3}{2}}} \|w_{n}\|_{L_{t}^{\infty}L_{x}^{\frac{6}{3-2s_{c}}}(\mathbb{R}\times\mathbb{R}^{3})} + \|(-\Delta_{\Omega_{n}})^{\frac{s_{c}}{2}}z_{n}\|_{L_{t}^{\infty}L_{x}^{2}([-T,T]\times\Omega)}$$

$$\lesssim 1 + \|z_{n}\|_{L_{t}^{\frac{5\alpha}{2}}([-T,T]\times\Omega)} + + \|(-\Delta_{\Omega_{n}})^{\frac{s_{c}}{2}}z_{n}\|_{L_{t}^{\infty}L_{x}^{2}([-T,T]\times\Omega)} < +\infty.$$

$$(4.9)$$

Step 3. Asymptotic agreement of the initial data.

In this step, we show that

$$\lim_{T \to \infty} \limsup_{n \to \infty} \left\| e^{it\Delta_{\Omega}} \left(\tilde{v}_n(\lambda_n^2 t_n) - \phi_n \right) \right\|_{L_{\star}^{\frac{5\alpha}{2}} \dot{H}_{D}^{s_{\mathcal{C}}, \frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \Omega)} = 0. \tag{4.10}$$

We first consider the case when $t_n \equiv 0$. By using Hölder's inequality, the Strichartz estimate and change of variables, we obtain

$$\|e^{it\Delta_{\Omega}} (\tilde{v}_{n}(0) - \phi_{n})\|_{L_{t}^{\frac{5\alpha}{2}} \dot{H}_{D}^{s_{c}, \frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \Omega)} \lesssim \|\tilde{v}_{n}(0) - \phi_{n}\|_{\dot{H}_{D}^{s_{c}}(\Omega)} \lesssim \|\chi_{n} \phi_{\leq \lambda_{n}^{\theta}} - \chi_{n} \phi\|_{\dot{H}_{D}^{s_{c}}(\Omega)}$$

$$\lesssim \||\nabla|^{s_{c}} \chi_{n}\|_{L_{x}^{\frac{3}{s_{c}}}(\Omega)} \|\phi_{\leq \lambda_{n}^{\theta}} - \phi\|_{L_{x}^{\frac{6}{3-2s_{c}}}(\Omega)} + \|\chi_{n}\|_{L_{x}^{\infty}(\Omega)} \||\nabla|^{s_{c}} (\phi_{\leq \lambda_{n}^{\theta}} - \phi)\|_{L_{x}^{2}(\Omega)} \to 0, \quad \text{as} \quad n \to \infty.$$

It is sufficient to prove the case that $|t_n| \to \infty$. By symmetry, it remains to show the case that $t_n \to +\infty$; the case $t_n \to -\infty$ can be handled in a similar way. Since T > 0 is fixed, for n sufficiently large, we have that $t_n > T$; so that

$$\tilde{v}_n(\lambda_n^2 t_n, x) = e^{i(t_n - T)\lambda_n^2 \Delta_{\Omega}} \tilde{v}_n(\lambda_n^2 T, x) = e^{i(t_n - T)\lambda_n^2 \Delta_{\Omega}} \left[\lambda_n^{s_c - \frac{3}{2}} (\chi_n w_n + z_n) \left(T, \frac{x - x_n}{\lambda_n} \right) \right].$$

Then by change of variables, Hölder's inequality and the Strichartz estimate, we have

$$\begin{split} & \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_c}{2}} e^{it\Delta_{\Omega}} \left[\tilde{v}_n(\lambda_n^2 t_n) - \phi_n \right] \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \Omega)} \\ &= \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} \left[e^{i(t-T)\Delta_{\Omega_n}} (\chi_n w_n + z_n)(T) - e^{it\Delta_{\Omega_n}} (\chi_n \phi) \right] \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \Omega_n)} \\ &\lesssim \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} z_n(T) \right\|_{L_x^2} + \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} \left(\chi_n(w_n - w_{\infty})(T) \right) \right\|_{L_x^2} \\ &+ \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} \left[e^{i(t-T)\Delta_{\Omega_n}} (\chi_n w_{\infty})(T) - e^{it\Delta_{\Omega_n}} (\chi_n \phi) \right] \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \Omega_n)}. \end{split}$$

Using (4.4) and (4.8), we have

$$\begin{split} & \left\| (-\Delta_{\Omega_n})^{\frac{s_c}{2}} z_n(T) \right\|_{L_x^2} + \left\| (-\Delta_{\Omega_n})^{\frac{s_c}{2}} \left(\chi_n(w_n - w_\infty)(T) \right) \right\|_{L_x^2} \\ & \lesssim \lambda_n^{s_c - \frac{3}{2} + \theta(\frac{7}{2} - s_c)} (T + \lambda_n^{-2\theta}) + \left\| (-\Delta_{\Omega_n})^{\frac{s_c}{2}} \right) \chi_n \right\|_{L_x^{\frac{3}{s_c}}} \|w_n - w_\infty\|_{L_t^{\infty} L_x^{\frac{6}{3 - 2s_c}}} \\ & + \|\chi_n\|_{L^{\infty}} \| (-\Delta_{\Omega_n})^{\frac{s_c}{2}} (w_n - w_\infty) \|_{L_t^{\infty} L_x^2} \to 0 \quad \text{as} \quad n \to \infty. \end{split}$$

Thus, we are left to verify that

$$\lim_{T \to \infty} \limsup_{n \to \infty} \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} \left[e^{i(t-T)\Delta_{\Omega_n}} (\chi_n w_\infty)(T) - e^{it\Delta_{\Omega_n}} (\chi_n \phi) \right] \right\|_{L^{\frac{5\alpha}{2}}_{x} L^{\frac{30\alpha}{75\alpha-8}}_{x}(\mathbb{R} \times \Omega_n)} = 0.$$

By the triangle inequality and the Strichartz estimate,

$$\begin{split} & \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} e^{i(t-T)\Delta_{\Omega_n}} \left(\chi_n w_\infty(T) \right) - e^{it\Delta_{\Omega_n}} \left(\chi_n \phi \right) \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}} (\mathbb{R} \times \Omega_n)} \\ & \lesssim \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} \left(\chi_n w_\infty(T) \right) - \chi_n (-\Delta)^{\frac{s_c}{2}} w_\infty(T) \right\|_{L_x^2} \\ & + \left\| \left[e^{i(t-T)\Delta_{\Omega_n}} - e^{i(t-T)\Delta} \right] \left[\chi_n (-\Delta)^{\frac{s_c}{2}} w_\infty(T) \right] \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}} (\mathbb{R} \times \Omega_n)} \\ & + \left\| e^{-iT\Delta} \left[\chi_n (-\Delta)^{\frac{s_c}{2}} w_\infty(T) \right] - \chi_n (-\Delta)^{\frac{s_c}{2}} \phi \right\|_{L_x^2} \\ & + \left\| \left[e^{it\Delta_{\Omega_n}} - e^{it\Delta} \right] \left[\chi_n (-\Delta)^{\frac{s_c}{2}} \phi \right] \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}} (\mathbb{R} \times \Omega_n)} \\ & + \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} \left(\chi_n \phi \right) - \chi_n (-\Delta)^{\frac{s_c}{2}} \phi \right\|_{L_x^2} \end{split}$$

The fact that I_2 and I_4 converge to zero as $n \to \infty$ follows from Theorem 2.25 and the density in L^2_x of C_c^{∞} functions supported in \mathbb{R}^3 minus a point. Next, we estimate I_1 , I_3 and I_5 . First, by triangle inequality, Proposition 2.23 and the monotone convergence theorem, for any $f \in \dot{H}^{s_c}(\mathbb{R}^3)$, we have

$$\begin{split} & \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} (\chi_n f) - \chi_n (-\Delta)^{\frac{s_c}{2}} f \right\|_{L_x^2} \\ & \lesssim \left\| (1 - \chi_n) (-\Delta)^{\frac{s_c}{2}} f \right\|_{L_x^2} + \left\| (-\Delta)^{\frac{s_c}{2}} \left((1 - \chi_n) f \right) \right\|_{L_x^2} \\ & + \left\| (-\Delta_{\Omega_n})^{\frac{s_c}{2}} (\chi_n f) - (-\Delta)^{\frac{s_c}{2}} (\chi_n f) \right\|_{L_x^2} \to 0 \quad \text{as} \quad n \to \infty. \end{split}$$

This completes the proof of I_5 and so the proof of I_1 .

Finally, for the term I_3 , we use (4.2) and the monotone convergence theorem to obtain

$$I_{3} \lesssim \left\| (1 - \chi_{n})(-\Delta)^{\frac{s_{c}}{2}} w_{\infty}(T) \right\|_{L_{x}^{2}} + \left\| (1 - \chi_{n})(-\Delta)^{\frac{s_{c}}{2}} \right\|_{L_{x}^{2}} + \left\| e^{-iT\Delta}(-\Delta)^{\frac{s_{c}}{2}} w_{\infty}(T) - (-\Delta)^{\frac{s_{c}}{2}} \phi \right\|_{L_{x}^{2}} \to 0,$$

by first taking $n \to \infty$ and then $T \to \infty$.

Step 4. We show that \tilde{v}_n is an approximate solution to (1.1) in the sense

$$i\partial_t \tilde{v}_n + \Delta_{\Omega} \tilde{v}_n = |\tilde{v}_n|^{\alpha} \tilde{v}_n + e_n,$$

where e_n satisfies the smallness condition

$$\lim_{T \to \infty} \limsup_{n \to \infty} \|e_n\|_{\dot{N}^{s_c}(\mathbb{R} \times \Omega)} = 0. \tag{4.11}$$

First, we consider the large time scale $t > \lambda_n^2 T$. By symmetry, we can treat the case $t < -\lambda_n^2 T$ in a similar way. Using the equivalence of Sobolev spaces, Strichartz estimates and Hölder's inequality, we

have

$$\begin{split} & \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_c}{2}} e_n \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{27\alpha - 8}} (\{t > \lambda_n^2 T\} \times \Omega)} \lesssim & \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_c}{2}} \left(\left| \tilde{v}_n \right|^{\alpha} \tilde{v}_n \right) \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{27\alpha - 8}} (\{t > \lambda_n^2 T\} \times \Omega)} \\ & \lesssim & \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_c}{2}} \tilde{v}_n \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}} (\{t > \lambda_n^2 T\} \times \Omega)} \|\tilde{v}_n\|_{L_t^{\frac{5\alpha}{2}} (\{t > \lambda_n^2 T\} \times \Omega)}^{\alpha} \\ & \lesssim & \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_c}{2}} \left[\chi_n w_n(T) + z_n(T) \right] \right\|_{L_x^2} \|\tilde{v}_n\|_{L_t^{\frac{5\alpha}{2}} (\{t > \lambda_n^2 T\} \times \Omega)}^{\alpha} \\ & \lesssim & \left(1 + \lambda_n^{s_c - \frac{3}{2} + \theta(\frac{7}{2} - s_c)} (T + \lambda_n^{-2\theta}) \right) \|\tilde{v}_n\|_{L_t^{\frac{5\alpha}{2}} (\{t > \lambda_n^2 T\} \times \Omega)}^{\alpha} . \end{split}$$

Therefore, to establish (4.11), it suffices to prove that

$$\lim_{T \to \infty} \limsup_{n \to \infty} \left\| e^{i(t - \lambda_n^2 T)\Delta_{\Omega}} \tilde{v}_n(\lambda_n^2 T) \right\|_{L_{t,T}^{\frac{5\alpha}{2}}(\{t > \lambda_n^2 T\} \times \Omega)} = 0. \tag{4.12}$$

We now prove (4.12). By the spacetime bounds (4.3), the global solution w_{∞} scatters. Let ϕ_{+} denote the forward asymptotic state, that is,

$$\|w_{\infty} - e^{it\Delta}\phi_{+}\|_{\dot{H}^{s_{c}}(\mathbb{R}^{3})} \to 0, \quad \text{as} \quad t \to \pm \infty.$$
 (4.13)

It then follows from Strichartz estimate, Hölder's inequality and change of variables that

$$\begin{split} & \left\| e^{i(t-\lambda_n^2T)\Delta_\Omega} \tilde{v}_n(\lambda_n^2T) \right\|_{L_{t,x}^{\frac{5\alpha}{2}}(\{t>\lambda_n^2T\}\times\Omega)} \lesssim \left\| e^{it\Delta_{\Omega_n}} \left(\chi_n w_n(T) + z_n(T)\right) \right\|_{L_{t,x}^{\frac{5\alpha}{2}}([0,\infty)\times\Omega_n)} \\ & \lesssim \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} z_n(T) \right\|_{L_x^2} + \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} \left[\chi_n(w_n(T) - w_\infty(T)) \right] \right\|_{L_x^2} \\ & + \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} \left[\chi_n(w_\infty(T) - e^{iT\Delta}w_+) \right] \right\|_{L_x^2} + \left\| e^{it\Delta_{\Omega_n}} \left[\chi_n e^{iT\Delta}w_+ \right] \right\|_{L_{t,x}^{\frac{5\alpha}{2}}([0,\infty)\times\Omega_n)} \\ & \lesssim \lambda_n^{s_c - \frac{3}{2} + \theta(\frac{7}{2} - s_c)} (T + \lambda_n^{-2\theta}) + \left\| w_n(T) - w_\infty(T) \right\|_{\dot{H}^{s_c}} + \left\| w_\infty(T) - e^{iT\Delta}w_+ \right\|_{\dot{H}^{s_c}} \\ & + \left\| \left[e^{it\Delta_{\Omega_n}} - e^{it\Delta} \right] \left[\chi_n e^{iT\Delta}w_+ \right] \right\|_{L_{t,x}^{\frac{5\alpha}{2}}([0,\infty)\times\mathbb{R}^3)} + \left\| \left(-\Delta \right)^{\frac{s_c}{2}} \left[(1 - \chi_n) e^{iT\Delta}w_+ \right] \right\|_{L_x^2} \\ & + \left\| e^{it\Delta}w_+ \right\|_{L_{t,x}^{\frac{5\alpha}{2}}((T,\infty)\times\mathbb{R}^3)}, \end{split}$$

which converges to zero by first letting $n \to \infty$ and then $T \to \infty$ by (4.8), (4.13), Theorem 2.25, and the monotone convergence theorem.

Now, we consider the case that $|t_n| \leq \lambda_n^2 T$. For these values of time, by the direct calculus we have

$$\begin{split} e_n(t,x) &= [(i\partial_t + \Delta_\Omega) \tilde{v}_n - |\tilde{v}_n|^\alpha \tilde{v}_n](t,x) \\ &= -\lambda_n^{s_c - \frac{7}{2}} [\Delta \chi_n] (\lambda_n^{-1}(x-x_n)) w_n (\lambda_n^{-2}t, -\lambda_n^{-1}x_n) + \lambda_n^{s_c - \frac{7}{2}} [\Delta \chi_n w_n] (\lambda_n^{-2}t, \lambda_n^{-1}(x-x_n)) \\ &+ 2\lambda_n^{s_c - \frac{7}{2}} (\nabla \chi_n \cdot \nabla w_n) (\lambda_n^{-2}t, \lambda_n^{-1}(x-x_n)) \\ &+ \lambda_n^{s_c - \frac{7}{2}} [\chi_n |w_n|^\alpha w_n - |\chi_n w_n + z_n|^\alpha (\chi_n w_n + z_n)] (\lambda_n^{-2}t, \lambda_n^{-1}(x-x_n)). \end{split}$$

By a change of variables and the equivalence of Sobolev norms Theorem 2.6, we obtain

$$\begin{split} \left\| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} e_{n} \right\|_{\dot{N}^{s_{c}}(\mathbb{R} \times \Omega)} &\lesssim \left\| (-\Delta)^{\frac{s_{c}}{2}} [\Delta \chi_{n}(w_{n}(t, x) - w_{n}(t, \lambda_{n}^{-1} x_{n}))] \right\|_{L_{t}^{2} L_{x}^{\frac{6}{5}}([-T, T] \times \Omega_{n})} \\ &+ \left\| (-\Delta)^{\frac{s_{c}}{2}} \left(\nabla \chi_{n} \nabla w_{n} \right) \right\|_{L_{t}^{2} L_{x}^{\frac{6}{5}}([-T, T] \times \Omega_{n})} \\ &+ \left\| (-\Delta)^{\frac{s_{c}}{2}} \left[(\chi_{n} - \chi_{n}^{\alpha+1}) |w_{n}|^{\alpha} w_{n} \right\|_{L_{t}^{2} L_{x}^{\frac{6}{5}}([-T, T] \times \Omega_{n})} \\ &+ \left\| (-\Delta)^{s_{c}} [|\chi_{n} w_{n} + z_{n}|^{\alpha} (\chi_{n} w_{n} z_{n}) - |\chi_{n} w_{n}|^{\alpha} \chi_{n} w_{n}] \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}([-T, T] \times \Omega_{n})} \\ &\stackrel{\triangle}{=} J_{1} + J_{2} + J_{3} + J_{4}. \end{split}$$

Using Hölder, the fundamental theorem of calculus, and (4.5), we estimate

$$J_{1} \lesssim T^{\frac{1}{2}} \| (-\Delta)^{\frac{s_{c}}{2}} (w_{n}(t,x) - w_{n}(t,-\lambda_{n}^{-1}x_{n})) \|_{L_{t,x}^{\infty}} \| \Delta \chi_{n} \|_{L_{\frac{6}{5}}^{\frac{6}{5}}}$$

$$+ T^{\frac{1}{2}} \| w_{n} - w_{n}(t,-\lambda_{n}^{-1}x_{n}) \|_{L_{t,x}^{\infty}(\mathbb{R} \times \text{supp}\Delta \chi_{n})} \| (-\Delta)^{\frac{s_{c}}{2}} (\Delta \chi_{n}) \|_{L_{x}^{\frac{6}{5}}}$$

$$\leq T^{\frac{1}{2}} \lambda_{n}^{-\frac{1}{2} + \frac{3}{2}\theta} + T^{\frac{1}{2}} \lambda_{n}^{-1 + \theta(\frac{5}{2} - s_{c})} \lambda_{n}^{s_{c} - \frac{1}{2}} \to 0 \quad \text{as} \quad n \to \infty.$$

By a similar argument, we can show that $J_2 \to 0$ as $n \to \infty$ and we omit the details. Next, we turn our attention to J_3 . By Lemma 2.7, Hölder's inequality and (4.5), we have

$$J_{3} \lesssim \left\| |\nabla|^{s_{c}} \chi_{n} \right\|_{L_{x}^{\frac{6}{5}}} \|w_{n}\|_{L_{x}^{\infty}L_{x}^{\infty}}^{\alpha+1} + \left\| \chi_{n} - \chi_{n}^{\alpha+1} \right\|_{L_{x}^{\frac{6}{5}}} \|w_{n}\|_{L_{x}^{\infty}L_{x}^{\infty}}^{\alpha} \left\| |\nabla|^{s_{c}} w_{n} \right\|_{L_{x}^{\infty}L_{x}^{\infty}} \lesssim \lambda_{n}^{s_{c} - \frac{5}{2} + \theta(\alpha+1)(\frac{3}{2} - s_{c})} + \lambda_{n}^{-\frac{5}{2} + \theta\alpha(\frac{3}{2} - s_{c}) + \frac{3}{2}\theta} \to 0 \quad \text{as} \quad n \to \infty.$$

Finally, we consider J_4 . By Lemma 2.18,

$$J_{4} \lesssim \left(\|\chi_{n} w_{n}\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}([-T,T] \times \Omega_{n})}^{\frac{\alpha-1}{2}} + \|z_{n}\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{5\alpha}{2}}([-T,T] \times \Omega_{n})}^{\frac{\alpha-1}{2}} \right) \times \left(\||\nabla|^{s_{c}} (\chi_{n} w_{n})\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{2}}([-T,T] \times \Omega_{n})}^{\frac{30\alpha}{2} L_{x}^{\frac{30\alpha}{2}} L_{x}^{\frac{30\alpha}{2}}([-T,T] \times \Omega_{n})} + \||\nabla|^{s_{c}} z_{n}\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{2}}([-T,T] \times \Omega_{n})}^{\frac{30\alpha}{2}} \right)^{2}.$$

$$(4.14)$$

Using the fractional product rule and (4.3), we have

$$\begin{aligned} & \| |\nabla|^{s_c}(\chi_n w_n) \|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha-8}}([-T,T] \times \Omega_n)} \lesssim \| |\nabla|^{s_c} \chi_n \|_{L_x^{\frac{20\alpha}{15\alpha-8}}} \| w_n \|_{L_t^{\infty} L_x^{\infty}} + \| \chi_n \|_{L_x^{\frac{30\alpha}{15\alpha-8}}} \| |\nabla|^{s_c} w_n \|_{L_t^{\infty} L_x^{\infty}} \\ & \lesssim T^{\frac{2}{5\alpha}} \lambda_n^{s_c - \frac{15\alpha-8}{30\alpha} \times 3 + \theta(\frac{3}{2} - s_c)} + T^{\frac{2}{5\alpha}} \lambda_n^{-\frac{15\alpha-8}{30\alpha} \times 3 + \frac{3}{2}\theta} = T^{\frac{2}{5\alpha}} \lambda_n^{\frac{3(2s_c - 3)}{10} + \theta(\frac{3}{2} - s_c)} + T^{\frac{2}{5\alpha}} \lambda_n^{-\frac{3}{2} + \frac{4}{5\alpha} + \frac{3}{2}\theta}, \end{aligned}$$

which converges to 0 as $n \to \infty$. This together with (4.3), Lemma 4.2 and (4.14) gives $J_4 \to 0$ as $n \to \infty$. This completes the proof of (4.11).

Step 5. Constructing v_n and approximation by C_c^{∞} functions.

By (4.9), (4.10), and invoking the stability Theorem 2.19, for sufficiently large n and T, we obtain a global solution v_n to (1.1) with initial data $v_n(0) = \phi_n$. Moreover, it has finite scattering norm and satisfies

$$\lim_{T \to \infty} \limsup_{n \to \infty} \left\| v_n(t - \lambda_n^2 t_n) - \tilde{v}_n(t) \right\|_{L^{\frac{5\alpha}{2}} L^{\frac{5\alpha}{2}}_x(\mathbb{R} \times \Omega)} = 0. \tag{4.15}$$

Therefore, to prove Theorem 4.1, it sufficies to prove the approximation (4.1).

This follows from a standard argument; see, for example, [33,35]. Here, we only provide a brief outline of the proof. First, by the density argument, we can take $\psi_{\varepsilon} \in C_0^{\infty}(\mathbb{R} \times \mathbb{R}^3)$ such that

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}}(w_{\infty} - \psi_{\varepsilon})\|_{L_{\tau}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \mathbb{R}^3)} < \varepsilon.$$

$$(4.16)$$

Then, using the change of variables and the triangle inequality, we obtain

$$\left\| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} \left[v_{n}(t - \lambda_{n}^{2}t_{n}, x + x_{n}) - \lambda_{n}^{s_{c} - \frac{3}{2}} \psi_{\varepsilon}(\lambda_{n}^{-2}t, \lambda_{n}^{-1}x) \right] \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \mathbb{R}^{3})}$$

$$\lesssim \left\| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} (w_{\infty} - \psi_{\varepsilon}) \right\|_{\dot{X}^{s_{c}}(\mathbb{R} \times \mathbb{R}^{3})} + \left\| v_{n}(t - \lambda_{n}^{2}t_{n}) - \tilde{v}_{n}(t) \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \mathbb{R}^{3})}$$

$$(4.17)$$

$$+ \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_c}{2}} \left[\tilde{v}_n(t, x) - \lambda_n^{s_c - \frac{3}{2}} w_{\infty}(\lambda_n^{-2} t, \lambda_n^{-1} (x - x_n)) \right] \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \mathbb{R}^3)}. \tag{4.18}$$

Clearly, by (4.15) and (4.16), we have (4.17) $\lesssim \varepsilon$. For (4.18), note that by (4.4), for sufficiently large n, w_n approximates w_∞ and $\chi_n(x) \to 1$. As \widetilde{v}_n is constructed through w_n , χ_n , and z_n , we can use Lemma 4.2, the triangle inequality, the Strichartz estimate, and Theorem 2.25 to show that for sufficiently large n, (4.18) is also small, which yields (4.1).

Next, we concerns the scenario when the rescaled obstacles Ω_n^c (where $\Omega_n = \lambda_n^{-1} (\Omega - \{x_n\})$) are retreating to infinity, which corresponds to Case 3 of Theorem 3.5.

Theorem 4.3 (Embedding of nonlinear profiles for retreating obstacles). Let $\{t_n\} \subset \mathbb{R}$ be such that $t_n \equiv 0$ or $|t_n| \to +\infty$. Let $\{x_n\} \subset \Omega$ and $\{\lambda_n\} \subset 2^{\mathbb{Z}}$ satisfy that $\frac{d(x_n)}{\lambda_n} \to \infty$. Suppose that $\phi \in \dot{H}^{s_c}(\mathbb{R}^3)$

$$\phi_n(x) = \lambda_n^{s_c - \frac{3}{2}} e^{i\lambda_n^2 t_n \Delta_{\Omega}} \left[(\chi_n \phi) \left(\frac{x - x_n}{\lambda_n} \right) \right]$$

with $\chi_n(x) = 1 - \Theta(\lambda_n |x|/d(x_n))$. Then for sufficiently large n, there exists a global solution v_n to (1.1) with initial data $v_n(0) = \phi_n$, which satisfies

$$||v_n||_{L^{\frac{5\alpha}{2}}_{+^2}(\mathbb{R}\times\Omega)} \lesssim ||\phi||_{\dot{H}^{s_c}} 1. \tag{4.19}$$

Furthermore, for every $\varepsilon > 0$, there exist $N_{\varepsilon} > 0$ and $\psi_{\varepsilon} \in C_0^{\infty}(\mathbb{R} \times \mathbb{R}^3)$ such that for $n \geq N_{\varepsilon}$, we get

$$\left\| (-\Delta_{\Omega})^{\frac{s_c}{2}} \left[v_n(t - \lambda_n^2 t_n, x + x_n) - \lambda_n^{s_c - \frac{3}{2}} \psi_{\varepsilon}(\lambda_n^{-2} t, \lambda_n^{-1} x) \right] \right\|_{L_{\star}^{\frac{5\alpha}{2}} L_{\tau}^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \mathbb{R}^3)} < \varepsilon. \tag{4.20}$$

Proof. Similar to the proof of Theorem 4.1, we also divide the proof of Theorem 4.3 into five steps. For the sake of simpleness, we denote $-\Delta_{\mathbb{R}^3} = -\Delta$.

Step 1. Constructing the global solution to ${\rm NLS}_{\mathbb{R}^3}.$

Let $\theta = \frac{1}{100(\alpha+1)}$. As in the proof of Theorem 4.1, if $t_n \equiv 0$, we let w_n and w_∞ be solutions to $NLS_{\mathbb{R}^3}$ with initial data $w_n(0) = P_{\leq d(x_n)^{\theta} \lambda_n^{-\theta}} \phi$ and $w_\infty(0) = \phi$. If $t_n \to \pm \infty$, we let w_n and w_∞ be solutions to $NLS_{\mathbb{R}^3}$ with

$$\begin{cases} \|w_n(t) - e^{it\Delta} P_{\leq d(x_n)^{\theta} \lambda_n^{-\theta}} \phi \|_{\dot{H}^{s_c}(\mathbb{R}^3)} \to 0, \\ \|w_{\infty}(t) - e^{it\Delta} \phi \|_{\dot{H}^{s_c}(\mathbb{R}^3)} \to 0. \end{cases}$$

By the assumption made in Theorem 1.2, we have that w_n and w_∞ are global solutions with uniformly bounded Strichartz norms. Moreover, arguing as in the proof of Theorem 4.1 and invoking Theorem 2.19, we see that w_n and w_∞ satisfy the following properties:

$$\begin{cases}
\|w_n\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{5\alpha}{2}}(\mathbb{R}\times\mathbb{R}^3)} + \|w_\infty\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{5\alpha}{2}}(\mathbb{R}\times\mathbb{R}^3)} \lesssim 1, \\
\||\nabla|^{s_c}(w_n - w_\infty)\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R}\times\mathbb{R}^3)} \to 0 \quad \text{as} \quad t \to \pm\infty, \\
\||\nabla|^s w_n\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{5\alpha}{2}}(\mathbb{R}\times\mathbb{R}^3)} \lesssim \left(\frac{d(x_n)}{\lambda_n}\right)^{\theta s}, \quad \text{for all} \quad s \ge 0.
\end{cases} \tag{4.21}$$

Step 2. Constructing the approximate solution to (1.1).

Fix T > 0 to be chosen later. We define

$$\tilde{v}_n(t,x) \stackrel{\triangle}{=} \begin{cases} \lambda_n^{s_c - \frac{3}{2}} (\chi_n w_n) (\lambda_n^{-2} t, \lambda_n^{-1} (x - x_n)), & |t| \leq \lambda_n^2 T, \\ e^{i(t - \lambda_n^2 T) \Delta_{\Omega}} \tilde{v}_n (\lambda_n^2 T, x), & t > \lambda_n^2 T, \\ e^{i(t + \lambda_n^2 T) \Delta_{\Omega}} \tilde{v}_n (-\lambda_n^2 T, x), & t < -\lambda_n^2 T. \end{cases}$$

Similar to (4.9), using (4.21), it is easy to see that \tilde{v}_n has finite scattering norm. Step 3. Agreement of the initial data:

$$\lim_{T \to \infty} \limsup_{n \to \infty} \left\| e^{it\Delta_{\Omega}} \left(\tilde{v}_n(\lambda_n^2 t_n) - \phi_n \right) \right\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \Omega)} = 0. \tag{4.22}$$

By the same argument as used in the proof of Step 3 in Theorem 4.1, we can prove (4.22) in the cases of $t_n \equiv 0$ and $|t_n| \to \infty$ by applying a change of variables, the Strichartz estimate, and using (4.21).

Step 4. Showing that \tilde{v}_n is the approximate solution to (1.1) in the sense that

$$\lim_{T \to \infty} \limsup_{n \to \infty} \left\| (i\partial_t + \Delta_{\Omega})\tilde{v}_n - |\tilde{v}_n|^{\alpha} \tilde{v}_n \right\|_{\dot{N}^{s_c}(\mathbb{R} \times \Omega)} = 0. \tag{4.23}$$

Similar to (4.12), it sufficies to prove

$$\lim_{T \to \infty} \limsup_{n \to \infty} \left\| e^{i(t - \lambda_n^2 T)\Delta_{\Omega}} \tilde{v}_n(\lambda_n^2 T) \right\|_{L_{\star}^{\frac{5\alpha}{2}}(\{t > \lambda_n^2 T\} \times \Omega)} = 0. \tag{4.24}$$

Let w_+ be the asymptotic state of w_{∞} . Then by Strichartz estimates and the change of variables, we get

$$\begin{aligned} & \left\| e^{i(t-\lambda_{n}^{2}T)\Delta_{\Omega}} \tilde{v}_{n}(\lambda_{n}^{2}T) \right\|_{L_{t,x}^{\frac{5\alpha}{2}}(\{t>\lambda_{n}^{2}T\}\times\Omega)} = \left\| e^{it\Delta_{\Omega_{n}}} (\chi_{n}w_{n}(T)) \right\|_{L_{t,x}^{\frac{5\alpha}{2}}((0,\infty)\times\Omega)} \\ & \lesssim \left\| e^{it\Delta_{\Omega_{n}}} [\chi_{n}e^{iT\Delta}w_{+}] \right\|_{L_{t,x}^{\frac{5\alpha}{2}}((0,\infty)\times\Omega_{n})} + \left\| \chi_{n} [w_{\infty}(T) - e^{iT\Delta}w_{+}] \right\|_{\dot{H}^{s_{c}}(\mathbb{R}^{3})} + \left\| \chi_{n} [w_{\infty}(T) - w_{n}(T)] \right\|_{\dot{H}^{s_{c}}(\mathbb{R}^{3})} \\ & \lesssim \left\| \left(e^{it\Delta_{\Omega_{n}}} - e^{it\Delta} \right) [\chi_{n}e^{iT\Delta}w_{+}] \right\|_{L_{t,x}^{\frac{5\alpha}{2}}((0,\infty)\times\mathbb{R}^{3})} + \left\| (1 - \chi_{n})e^{iT\Delta}w_{+} \right\|_{\dot{H}^{s_{c}}(\mathbb{R}^{3})} \\ & + \left\| e^{it\Delta}w_{+} \right\|_{L_{t,x}^{\frac{5\alpha}{2}}((T,\infty)\times\mathbb{R}^{3})} + \left\| w_{\infty}(T) - e^{iT\Delta}w_{+} \right\|_{\dot{H}^{s_{c}}(\mathbb{R}^{3})} + \left\| w_{\infty}(T) - w_{n}(T) \right\|_{\dot{H}^{s_{c}}(\mathbb{R}^{3})}, \end{aligned}$$

which converges to zero by first letting $n \to \infty$ and then $T \to \infty$ in view of Theorem 2.25, (4.21) and the monotone convergence theorem.

Finally, we consider the intermediate time scale $|t| \leq \lambda_n^2 T$. We compute

$$[(i\partial_{t} + \Delta_{\Omega})\tilde{v}_{n} - |\tilde{v}_{n}|^{\alpha}\tilde{v}_{n}](t,x) = \lambda_{n}^{s_{c} - \frac{7}{2}}[\Delta\chi_{n}w_{n}](\lambda_{n}^{-2}t, \lambda_{n}^{-1}(x - x_{n}))$$

$$+ 2\lambda_{n}^{s_{c} - \frac{7}{2}}(\nabla\chi_{n} \cdot \nabla w_{n})(\lambda_{n}^{-2}t, \lambda_{n}^{-1}(x - x_{n}))$$

$$+ \lambda_{n}^{s_{c} - \frac{7}{2}}[(\chi_{n} - \chi_{n}^{\alpha+1})|w_{n}|^{\alpha}w_{n}](\lambda_{n}^{-2}t, \lambda_{n}^{-1}(x - x_{n})).$$

Note that the cut-off function $\chi_n \sim 1_{|x| \sim \frac{d(x_n)}{\lambda_n}}$ and $\frac{d(x_n)}{\lambda_n} \to \infty$ as $n \to \infty$. Therefore, we can modified the proof in step 4 of Theorem 4.3 with minor change to obtain (4.23).

Step 5. Constructing v_n and approximation by C_c^{∞} functions.

By (4.22), (4.23) and invoking the stability Theorem 2.19, for sufficiently large n we obtain a global solution v_n to (1.1) with initial data $v_n(0) = \phi_n$. Moreover, it satisfies

$$\|v_n\|_{L^{\frac{5\alpha}{2}}_{t,x}(\mathbb{R}\times\Omega)} \lesssim 1, \quad \text{and} \quad \lim_{T\to\infty} \limsup_{n\to\infty} \|v_n(t-\lambda_n^2t_n) - \tilde{v}_n(t)\|_{\dot{H}^{s_c}_D(\Omega)} = 0.$$

Finially, by the same argument as that used to derive (4.1), we can obtain the convergence (4.20) and omit the details. This completes the proof of Theorem 4.3.

At last, we treat the case that the obstacle expands to fill the half-space, i.e. Case 4 in Theorem 3.5.

Theorem 4.4 (Embedding the nonlinear profiles: the half-space case). Let $\{t_n\} \subset \mathbb{R}$ be such that $t_n \equiv 0$ and $|t_n| \to \infty$. Let $\{\lambda_n\} \subset 2^{\mathbb{Z}}$ and $\{x_n\} \subset \Omega$ be such that

$$\lambda_n \to 0$$
, and $\frac{d(x_n)}{\lambda_n} \to d_\infty > 0$.

Let $x_n^* \in \partial \Omega$ be such that $|x_n - x_n^*| = d(x_n)$ and $R_n \in SO(3)$ be such that $R_n e_3 = \frac{x_n - x_n^*}{|x_n - x_n^*|}$. Finally, let $\phi \in \dot{H}_D^{s_c}(\mathbb{H})$, we define

$$\phi_n(x) = \lambda_n^{s_c - \frac{3}{2}} e^{i\lambda_n^2 t_n \Delta_{\Omega}} \phi\left(\frac{R_n^{-1}(x_n - x_n^*)}{\lambda_n}\right).$$

Then for n sufficiently large, there exists a global solution v_n to (1.1) with initial data $v_n(0) = \phi_n$, which also satisfies

$$||v_n||_{L_{t,x}^{\frac{5\alpha}{2}}(\mathbb{R}\times\Omega)}\lesssim 1.$$

Furthermore, for every $\varepsilon > 0$, there exists $N_{\varepsilon} \in \mathbb{N}$ and $\psi_{\varepsilon} \in C_0^{\infty}(\mathbb{R} \times \mathbb{H})$ so that for every $n \geq N_{\varepsilon}$, we have

$$\left\| \left(-\Delta_{\Omega} \right)^{\frac{s_c}{2}} \left[v_n(t - \lambda_n^2 t_n, R_n x + x_n^*) - \lambda_n^{\frac{s_c - \frac{3}{2}}{2}} \psi_{\varepsilon}(\lambda_n^{-2} t, \lambda_n^{-1} x) \right] \right\|_{L^{\frac{5\alpha}{2}} L_{\pi^{\frac{30\alpha}{5\alpha - 8}}(\mathbb{R} \times \mathbb{R}^3)}^{\frac{30\alpha}{5\alpha - 8}} < \varepsilon. \tag{4.25}$$

Proof. Again, we divide the proof of this theorem into five main steps.

Step 1. Construction of the global solution to $NLS_{\mathbb{R}^3}$.

Let $\theta \ll 1$. For $t_n \equiv 0$, let U_n and U_{∞} be solutions to NLS_H with initial data $U_n(0) = \phi_{\lambda_n^{-\theta}}$ and $U_{\infty}(0) = \phi$. If $|t_n| \to +\infty$, we set U_n and U_{∞} be solutions to NLS_{R3} satisfying

$$\|U_n(t) - e^{it\Delta_{\mathbb{H}}} \phi_{\leq \lambda_n^{-\theta}}\|_{\dot{H}^{s_c}_{D}(\mathbb{H})} \to 0 \quad \text{and} \quad \|U_{\infty}(t) - e^{it\Delta_{\mathbb{H}}} \phi\|_{\dot{H}^{s_c}_{D}(\mathbb{H})} \to 0, \quad \text{as} \quad t \to \pm \infty.$$
 (4.26)

In all cases, the assumption made in Theorem 1.2 implies that

$$||U_n||_{L_{\star}^{\frac{5\alpha}{2}}L_{x}^{\frac{5\alpha}{2}}(\mathbb{R}\times\mathbb{H})} + ||U_{\infty}||_{L_{\star}^{\frac{5\alpha}{2}}L_{x}^{\frac{5\alpha}{2}}(\mathbb{R}\times\mathbb{H})} \lesssim 1.$$

Indeed, the solution to $NLS_{\mathbb{H}}$ can be extended to the solution of $NLS_{\mathbb{R}^3}$ by reflecting along the boundary $\partial \mathbb{H}$. Moreover, arguing as in the proof of the previous two embedding theorems, and using the stability theorem and the persistence of regularity, we have

$$\begin{cases}
\lim_{n \to \infty} \|U_n - U_\infty\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{5\alpha}{2}}(\mathbb{R} \times \mathbb{H})} = 0, \\
\|(-\Delta_{\mathbb{H}})^{\frac{s}{2}} U_n\|_{L_t^{\infty} L_x^2(\mathbb{R} \times \mathbb{H})} \lesssim \lambda_n^{\theta(s-1)}.
\end{cases}$$
(4.27)

Step 2. Construction of the approximate solution to (1.1).

Let $\Omega_n := \lambda_n^{-1} R_n^{-1} (\Omega - \{x_n^*\})$ and let T > 0 to be chosen later. On the intermediate time scale that $|t| < \lambda_n^2 T$, we embed U_n by using a boundary straightening diffeomorphism Ψ_n of a neighborhood of zero in Ω_n of size $L_n := \lambda_n^{-2\theta}$ into a corresponding neighborhood in \mathbb{H} .

To do this, we define a smooth function ψ_n on the set $|x^{\perp}| \leq L_n$ so that $x^{\perp} \mapsto (x^{\perp}, -\psi_n(x^{\perp}))$ traces out $\partial \Omega_n$. Here and below we write $x \in \mathbb{R}^3$ as $x = (x^{\perp}, x_3)$. By our choice of R_n , $\partial \Omega_n$ has unit normal e_3 at zero. Moreover, $\partial \Omega_n$ has curvatures that are $O(\lambda_n)$. Thus, ψ_n satisfies the following:

$$\begin{cases}
\psi_n(0) = 0, \quad \nabla \psi_n(0) = 0, \quad |\nabla \psi_n(x^{\perp})| \lesssim \lambda_n^{1-2\theta}, \\
|\partial^{\alpha} \psi_n(x^{\perp})| \lesssim \lambda_n^{|\alpha|-1} \quad \text{for all} \quad |\alpha| \ge 2.
\end{cases}$$
(4.28)

We now define the map $\Psi_n: \Omega_n \cap \{|x^{\perp}| \leq L_n\} \to \mathbb{H}$ and a cutoff $\chi_n: \mathbb{R}^3 \to [0,1]$ via

$$\Psi_n(x) := (x^{\perp}, x_3 + \psi_n(x^{\perp}))$$
 and $\chi_n(x) := 1 - \Theta(\frac{x}{L_n})$.

Note that on the domain of Ψ_n , which contains supp χ_n , we have:

$$|\det(\partial \Psi_n)| \sim 1 \quad \text{and} \quad |\partial \Psi_n| \lesssim 1.$$
 (4.29)

Now, we are in position to define the approximate solution. Let $\tilde{U}_n := \chi_n U_n$ and define

$$\tilde{v}_n(t,x) := \begin{cases} \lambda_n^{s_c - \frac{3}{2}} [\tilde{U}_n(\lambda_n^{-2}t) \circ \Psi_n] (\lambda_n^{-1}R_n^{-1}(x - x_n^*)), & |t| \leq \lambda_n^2 T, \\ e^{i(t - \lambda_n^2 T)\Delta_\Omega} \tilde{v}_n(\lambda_n^2 T, x), & t > \lambda_n^2 T, \\ e^{i(t + \lambda_n^2 T)\Delta_\Omega} \tilde{v}_n(-\lambda_n^2 T, x), & t < -\lambda_n^2 T. \end{cases}$$

We first prove that \tilde{v}_n has finite scattering size. Indeed, by the Strichartz inequality, a change of variables, and (4.29),

$$\|\tilde{v}_n\|_{L_{t,x}^{\frac{5\alpha}{2}}(\mathbb{R}\times\Omega)} \lesssim \|\tilde{U}_n \circ \Psi_n\|_{L_{t,x}^{\frac{5\alpha}{2}}(\mathbb{R}\times\Omega_n)} + \|\tilde{U}_n(\pm T) \circ \Psi_n\|_{\dot{H}_D^{s_c}(\Omega_n)}$$

$$\lesssim \|\tilde{U}_n\|_{L_{t,x}^{\frac{5\alpha}{2}}(\mathbb{R}\times\mathbb{H})} + \|\tilde{U}_n(\pm T)\|_{\dot{H}_D^{s_c}(\mathbb{H})} \lesssim 1. \tag{4.30}$$

Step 3. Asymptotic agreement with the initial data:

$$\lim_{T \to \infty} \limsup_{n \to \infty} \|(-\Delta_{\Omega})^{\frac{s_c}{2}} e^{it\Delta_{\Omega}} [\tilde{v}_n(\lambda_n^2 t_n) - \phi_n] \|_{L_{\star}^{\frac{5\alpha}{2}} L_{\tau}^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \Omega)} = 0. \tag{4.31}$$

First, we consider the case that $t_n \equiv 0$. By Strichartz and a change of variables,

$$\begin{split} & \left\| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} e^{it\Delta_{\Omega}} (\tilde{v}_{n}(0) - \phi_{n}) \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}} (\mathbb{R} \times \Omega)} \lesssim \left\| (\chi_{n} \phi_{\leq \lambda_{n}^{-\theta}}) \circ \Psi_{n} - \phi \right\|_{\dot{H}_{D}^{s_{c}}(\Omega_{n})} \\ & \lesssim \left\| (-\Delta)^{\frac{s_{c}}{2}} \left((\chi_{n} \phi_{> \lambda_{n}^{-\theta}}) \circ \Psi_{n} \right) \right\|_{L_{x}^{2}} + \left\| (-\Delta)^{\frac{s_{c}}{2}} [(\chi_{n} \phi) \circ \Psi_{n} - \chi_{n} \phi] \right\|_{L_{x}^{2}} + \left\| (-\Delta)^{\frac{s_{c}}{2}} ((1 - \chi_{n}) \phi) \right\|_{L_{x}^{2}}. \end{split}$$

As $\lambda_n \to 0$, we get $\|\phi_{>\lambda_n^{-\theta}}\|_{\dot{H}^{s_c}} \to 0$ as $n \to \infty$; thus, using (4.29) we see that the first term converges to 0. For the second term, we note that $\Psi_n(x) \to x$ in C^1 ; thus, approximating ϕ by $C_0^{\infty}(\mathbb{H})$ functions we see that the second term converges to 0. Finally, the last term converges to 0 by the dominated convergence theorem and $L_n = \lambda_n^{-2\theta} \to \infty$.

It remains to prove (4.31) when $t_n \to +\infty$; the case when $t_n \to -\infty$ can be treated similarly. Note that as T > 0 is fixed, for n sufficiently large, we have that $t_n > T$; so that

$$\tilde{v}_n(\lambda_n^2 t_n, x) = e^{i(t_n - T)\lambda_n^2 \Delta_\Omega} \left[\lambda_n^{s_c - \frac{3}{2}} (\tilde{U}_n(T) \circ \Psi_n) (\lambda_n^{-1} R_n^{-1} (x - x_n^*))\right].$$

A change of variables then yields that

$$\begin{aligned} & \left\| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} e^{it\Delta_{\Omega}} (\tilde{v}_{n}(\lambda_{n}^{2}t_{n}) - \phi_{n}) \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha - 8}}(\mathbb{R} \times \Omega)} \\ & \lesssim & \left\| (-\Delta_{\Omega_{n}})^{\frac{s_{c}}{2}} (\tilde{U}_{n}(T) \circ \Psi_{n} - U_{\infty}(T)) \right\|_{L_{x}^{2}} \end{aligned}$$

$$(4.32)$$

$$+ \left\| \left(-\Delta_{\Omega_n} \right)^{\frac{s_c}{2}} \left(e^{i(t-T)\Delta_{\Omega_n}} U_{\infty}(T) - e^{it\Delta_{\Omega_n}} \phi \right) \right\|_{L_{\star}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}}(\mathbb{R} \times \Omega_n)}. \tag{4.33}$$

By the triangle inequality,

$$(4.32) \lesssim \|(-\Delta_{\Omega_n})^{\frac{s_c}{2}} \left((\chi_n U_{\infty}(T)) \circ \Psi_n - U_{\infty}(T) \right) \|_{L_x^2} + \|(-\Delta_{\Omega_n})^{\frac{s_c}{2}} \left((\chi_n (U_n(T) - U_{\infty}(T))) \circ \Psi_n \right) \|_{L_x^2},$$

which converges to zero as $n \to \infty$ by the fact that $\Psi_n(x) \to x$ in C^1 and (4.27). For the second term, by the Strichartz estimate, Proposition 2.23, Theorem 2.25, and (4.26), we see that

$$\begin{aligned} (4.33) &\lesssim \left\| e^{i(t-T)\Delta_{\Omega_{n}}} (-\Delta_{\mathbb{H}})^{\frac{s_{c}}{2}} U_{\infty}(T) - e^{it\Delta_{\Omega_{n}}} (-\Delta_{\mathbb{H}})^{\frac{s_{c}}{2}} \phi \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}} (\mathbb{R} \times \Omega_{n})} \\ &+ \left\| \left((-\Delta_{\Omega_{n}})^{\frac{s_{c}}{2}} - (-\Delta_{\mathbb{H}})^{\frac{s_{c}}{2}} \right) U_{\infty}(T) \right\|_{L_{x}^{2}} + \left\| \left((-\Delta_{\Omega_{n}})^{\frac{s_{c}}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}} (\mathbb{R} \times \Omega_{n}) \right) \right\|_{L_{x}^{2}} \\ &\lesssim \left\| \left(e^{i(t-T)\Delta_{\Omega_{n}}} - e^{i(t-T)\Delta_{\mathbb{H}}} \right) \left(-\Delta_{\mathbb{H}} \right)^{\frac{s_{c}}{2}} U_{\infty}(T) \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}} (\mathbb{R} \times \Omega_{n})} \\ &+ \left\| \left(e^{it\Delta_{\Omega_{n}}} - e^{it\Delta_{\mathbb{H}}} \right) \left(-\Delta_{\mathbb{H}} \right)^{\frac{s_{c}}{2}} \phi \right\|_{L_{t}^{\frac{5\alpha}{2}} L_{x}^{\frac{30\alpha}{15\alpha-8}} (\mathbb{R} \times \Omega_{n})} \\ &+ \left\| e^{-iT\Delta_{\mathbb{H}}} U_{\infty}(T) - \phi \right\|_{\dot{H}_{D}^{c}(\mathbb{H})} + o(1), \end{aligned}$$

and that this converges to zero by first taking $n \to \infty$ and then $T \to \infty$.

Step 4. Proving that \tilde{v}_n is approximate solution to (1.1) in the following sense

$$\lim_{T \to \infty} \limsup_{n \to \infty} \left\| (i\partial_t + \Delta_{\Omega})\tilde{v}_n - |\tilde{v}_n|^{\alpha} \tilde{v}_n \right\|_{\dot{N}^{s_c}(\mathbb{R} \times \Omega)} = 0.$$
(4.34)

We first control the contribution of $|t| \ge \lambda_n^2 T$. By the same argument as that used in step 4 of Theorem 4.1, this reduces to proving

$$\lim_{T \to \infty} \limsup_{n \to \infty} \|e^{i(t-\lambda_n^2 T)\Delta_{\Omega}} \tilde{v}_n(\lambda_n^2 T)\|_{L_{t,x}^{\frac{5\alpha}{2}}(\{t > \lambda_n^2 T\} \times \Omega)} = 0. \tag{4.35}$$

Let U_+ denote the scattering state of U_{∞} in forward-time direction. By the Strichartz estimate, Theorem 2.25 and the monotone convergence theorem, we have that

$$\begin{split} & \left\| e^{i(t-\lambda_n^2T)\Delta_{\Omega}} \tilde{v}_n(\lambda_n^2T) \right\|_{L_{t,x}^{\frac{5\alpha}{2}}((\lambda_n^2T,\infty)\times\Omega)} = \left\| e^{i(t-T)\Delta_{\Omega_n}} (\tilde{U}_n(T)\circ\Psi_n) \right\|_{L_{t,x}^{\frac{5\alpha}{2}}((T,\infty)\times\Omega_n)} \\ & \lesssim \left\| \left(e^{i(t-T)\Delta_{\Omega_n}} - e^{i(t-T)\Delta_{\mathbb{H}}} \right) (e^{iT\Delta_{\mathbb{H}}} U_+) \right\|_{L_{t,x}^{\frac{5\alpha}{2}}((0,\infty)\times\Omega_n)} + \left\| e^{it\Delta_{\mathbb{H}}} U_+ \right\|_{L_{t,x}^{\frac{5\alpha}{2}}((T,\infty)\times\mathbb{R}^3_+)} + o(1), \end{split}$$

and that this converges to zero by Theorem 2.25 and the monotone convergence theorem by first taking $n \to \infty$ and then $T \to \infty$.

Next, we handle the middle time interval $\{|t| \leq \lambda_n^2 T\}$. By the direct computation, we have

$$\Delta(\widetilde{U}_n\circ\Psi_n)=(\partial_k\widetilde{U}_n\circ\Psi_n)\Delta\Psi_n^k+(\partial_{kl}\widetilde{U}_n\circ\Psi_n)\partial_j\Psi_n^l\partial_j\Psi_n^k,$$

where Ψ_n^k denotes the kth component of Ψ_n and repeated indices are summed. Recall that $\Psi_n(x) = x + (0, \psi_n(x^{\perp}))$, thus we have

$$\Delta \Psi_n^k = O(\partial^2 \psi_n), \quad \partial_j \Psi_n^l = \delta_{jl} + O(\partial \psi_n),$$
$$\partial_j \Psi_n^l \partial_j \Psi_n^k = \delta_{jl} \delta_{jk} + O(\partial \psi_n) + O((\partial \psi_n)^2),$$

where we use O to denote a collection of similar terms. Therefore,

$$\begin{split} &(\partial_k \widetilde{U}_n \circ \Psi_n) \Delta \Psi_n^k = O\big((\partial \widetilde{U}_n \circ \Psi_n)(\partial^2 \psi_n)\big), \\ &(\partial_{kl} \widetilde{U}_n \circ \Psi_n) \partial_j \Psi_n^l \partial_j \Psi_n^k = \Delta \widetilde{U}_n \circ \Psi_n + O\big(\big(\partial^2 \widetilde{U}_n \circ \Psi_n\big)\big(\partial \psi_n + (\partial \psi_n)^2\big)\big) \end{split}$$

and so

$$(i\partial_t + \Delta_{\Omega_n})(\widetilde{U}_n \circ \Psi_n) - (|\widetilde{U}_n|^{\alpha}\widetilde{U}_n) \circ \Psi_n = [(i\partial_t + \Delta_{\mathbb{H}})\widetilde{U}_n - |\widetilde{U}_n|^4\widetilde{U}_n] \circ \Psi_n + O((\partial\widetilde{U}_n \circ \Psi_n)(\partial^2\psi_n)) + O((\partial^2\widetilde{U}_n \circ \Psi_n)(\partial\psi_n + (\partial\psi_n)^2)).$$

By a change of variables and (4.29), we get

$$\begin{split} & \left\| \left(-\Delta_{\Omega} \right)^{\frac{s_{c}}{2}} \left((i\partial_{t} + \Delta_{\Omega}) \tilde{v}_{n} - |\tilde{v}_{n}|^{\alpha} \tilde{v}_{n} \right) \right\|_{L_{t}^{1} L_{x}^{2} (\{|t| \leq \lambda_{n}^{2} T\} \times \Omega)} \\ &= \left\| \left(-\Delta_{\Omega_{n}} \right)^{\frac{s_{c}}{2}} \left((i\partial_{t} + \Delta_{\Omega_{n}}) (\tilde{U}_{n} \circ \Psi_{n}) - (|\tilde{U}_{n}|^{\alpha} \tilde{U}_{n}) \circ \Psi_{n} \right) \right\|_{L_{t}^{1} L_{x}^{2} (\{|t| \leq \lambda_{n}^{2} T\} \times \Omega_{n})} \\ &\lesssim \left\| \left(-\Delta_{\Omega_{n}} \right)^{\frac{s_{c}}{2}} \left(((i\partial_{t} + \Delta_{\mathbb{H}}) \tilde{U}_{n} - |\tilde{U}_{n}|^{\alpha} \tilde{U}_{n}) \circ \Psi_{n} \right) \right\|_{L_{t}^{1} L_{x}^{2} ([-T,T] \times \Omega_{n})} \\ &+ \left\| \left(-\Delta_{\Omega_{n}} \right)^{\frac{s_{c}}{2}} \left((\partial \tilde{U}_{n} \circ \Psi_{n}) \partial^{2} \psi_{n} \right) \right\|_{L_{t}^{1} L_{x}^{2} ([-T,T] \times \Omega_{n})} \\ &+ \left\| \left(-\Delta_{\Omega_{n}} \right)^{\frac{s_{c}}{2}} \left((\partial^{2} \tilde{U}_{n} \circ \Psi_{n}) \left(\partial \psi_{n} + (\partial \psi_{n})^{2} \right) \right) \right\|_{L_{t}^{1} L_{x}^{2} ([-T,T] \times \Omega_{n})} \\ &\lesssim \left\| \left(-\Delta \right)^{\frac{s_{c}}{2}} \left((i\partial_{t} + \Delta_{\mathbb{H}}) \tilde{U}_{n} - |\tilde{U}_{n}|^{\alpha} \tilde{U}_{n} \right) \right\|_{L_{t}^{1} L_{x}^{2} ([-T,T] \times \mathbb{H})} \end{aligned} \tag{4.36}$$

$$+ \left\| (-\Delta)^{\frac{sc}{2}} \left((\partial \tilde{U}_n \circ \Psi_n) \partial^2 \psi_n \right) \right\|_{L^1_t L^2_x([-T,T] \times \Omega_n)} \tag{4.37}$$

$$+ \left\| \left(-\Delta \right)^{\frac{s_c}{2}} \left(\left(\partial^2 \tilde{U}_n \circ \Psi_n \right) \left(\partial \psi_n + \left(\partial \psi_n \right)^2 \right) \right) \right\|_{L^1_t L^2_x([-T,T] \times \Omega_n)}. \tag{4.38}$$

By direct computation

$$(i\partial_t + \Delta_{\mathbb{H}})\tilde{U}_n - |\tilde{U}_n|^{\alpha}\tilde{U}_n = (\chi_n - \chi_n^{\alpha+1})|U_n|^4U_n + 2\nabla\chi_n \cdot \nabla w_n + \Delta\chi_n w_n. \tag{4.39}$$

For fixed T>0, using fractional product rule, (4.27), (4.28), (4.29) and $\lambda_n\to 0$, it is easy to see that (4.37), (4.38) and the $\dot{N}^{s_c}(\mathbb{R}\times\mathbb{H})$ norm of the last two terms in (4.39) converges to 0 as $n\to\infty$.

Therefore, the proof of (4.34) reduces to show that the $\dot{N}^{s_c}(\mathbb{R} \times \mathbb{H})$ norm of the first term in (4.39) converges to 0 as $n \to \infty$. To this end, we estimate

$$\begin{split} &\|(\chi_n-\chi_n^{\alpha+1})|U_n|^{\alpha+1}U_n\|_{\dot{N}^{s_c}([-T,T]\times\mathbb{H})}\\ &\lesssim \|(\chi_n-\chi_n^{\alpha+1})|U_n|^{\alpha+1}|\nabla|^{s_c}U_n\|_{L_t^{\frac{5\alpha}{2(\alpha+1)}}L_x^{\frac{30\alpha}{27\alpha-8}}([-T,T]\times\mathbb{H})} + \||U_n|^{\alpha+1}|\nabla|^{s_c}\chi_n\|_{L_t^{\frac{5\alpha}{2(\alpha+1)}}L_x^{\frac{30\alpha}{27\alpha-8}}([-T,T]\times\mathbb{H})}\\ &\lesssim \|U_n1_{|x|\sim L_n}\|_{L_{t,x}^{\frac{5\alpha}{2}}}^{\alpha}\||\nabla|^{s_c}U_n\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{30\alpha}{15\alpha-8}}} + \|U_n1_{|x|\sim L_n}\|_{L_{t,x}^{\frac{5\alpha}{2}}}^{\alpha}\||\nabla|^{s_c}U_n\|_{L_t^{\frac{5\alpha}{2}}L_x^{\frac{30\alpha}{15\alpha-8}}} \||\nabla|^{s_c}\chi_n\|_{L_x^{\frac{3c}{2}}}\\ &\lesssim \|1_{|x|\sim L_n}U_\infty\|_{L_{t,x}^{\frac{5\alpha}{2}}}^{\alpha} + \|U_\infty-U_n\|_{L_{t,x}^{\frac{5\alpha}{2}}}^{\alpha} \to 0 \quad \text{as} \quad n\to\infty. \end{split}$$

This completes the proof of (4.34).

Step 5. Constructing v_n and approximating by compactly supported functions.

Similar to Theorem 4.1 and Theorem 4.3, using (4.30), (4.31), (4.34) and the stability theorem 2.19, for n large enough we obtain a global solution v_n to (1.1) with initial data $v_n(0) = \phi_n$, which satisfies (4.19). Moreover, the similar argument used in Theorem 4.1 and Theorem 4.3 also gives (4.20) and we omit the details.

5. Reduction to Almost Periodic Solutions

The goal of this section is to prove Theorem 1.8. The proof relies on establishing a Palais-Smale condition for minimizing sequences of blowup solutions to (1.1), enabling us to conclude that the failure of Theorem 1.2 would necessitate the existence of minimal counterexamples exhibiting the properties described in Theorem 1.8. We follow the approach outlined in [30, Section 3]. By now, this general strategy has become fairly standard; see, for example, [24, 25, 33, 54] for analogous results in different scenarios. Thus, we will focus on presenting the essential steps, offering detailed explanations only where new complications arise in our setting.

Throughout this section, we use the notation

$$S_I(u) := \int_I \int_{\Omega} |u(t,x)|^{\frac{5\alpha}{2}} dx dt.$$
 (5.1)

Assume Theorem 1.2 fails for some $s_c \in [\frac{1}{2}, \frac{3}{2})$. We define the function $L: [0, \infty) \to [0, \infty)$ as

$$L(E) := \sup\{S_I(u) : u : I \times \Omega \to \mathbb{C} \text{ solving (1.1) with } \sup_{t \in I} \|u(t)\|_{\dot{H}^{s_c}_D(\Omega)}^2 \le E\}.$$

It is important to note that L is non-decreasing, and Theorem 2.16 provides the bound

$$L(E) \lesssim E^{\frac{5\alpha}{4}}$$
 for E sufficiently small. (5.2)

This leads to the existence of a unique critical value $E_c \in (0, \infty]$ such that $L(E) < \infty$ for $E < E_c$ and $L(E) = \infty$ for $E > E_c$. The failure of Theorem indicates that $0 < E_c < \infty$.

A crucial element in establishing Theorem 1.8 is verifying a Palais–Smale condition. Once the following proposition is established, the derivation of Theorem follows a standard route (see [30]).

Proposition 5.1 (Palais–Smale condition modulo symmetries). Let $u_n: I_n \times \Omega \to \mathbb{C}$ be a sequence of solutions to (1.1) such that

$$\limsup_{n \to \infty} \sup_{t \in I_n} \|u_n(t)\|_{\dot{H}^{s_c}_D(\Omega)}^2 = E_c,$$

and suppose $t_n \in I_n$ are such that

$$\lim_{n \to \infty} S_{[t_n, \sup I_n]}(u_n) = \lim_{n \to \infty} S_{[\inf I_n, t_n]}(u_n) = \infty.$$

$$(5.3)$$

Then the sequence $u_n(t_n)$ has a subsequence that converges strongly in $\dot{H}_D^{s_c}(\Omega)$.

We now sketch the proof of this proposition, following the argument as it appears in [30]. As in that setting, the main ingredients will be a linear profile decomposition (Theorem 3.5 in our case) and a stability result (Theorem 2.19 in our case).

We begin by translating so that each $t_n = 0$, and apply the linear profile decomposition Theorem 3.5 to write

$$u_n(0) = \sum_{i=1}^{J} \phi_n^j + w_n^J \tag{5.4}$$

with the properties stated in Theorem 3.5.

Next, we construct the nonlinear profiles. If j conforms to Case 1, we invoke Theorem 2.16 and define $v^j: I^j \times \mathbb{R}^d \to \mathbb{C}$ as the maximal-lifespan solution to (1.1) satisfying

$$\begin{cases} v^j(0) := \phi^j & \text{if } t_n^j \equiv 0, \\ v^j \text{ scatters to } \phi^j \text{ as } t \to \pm \infty & \text{if } t_n^j \to \pm \infty. \end{cases}$$

We then define the nonlinear profiles $v_n^j(t,x) := v^j(t+t_n^j(\lambda_n^j)^2,x)$. Then v_n^j is also a solution to (1.1) on the time interval $I_n^j := I^j - \{t_n^j(\lambda_n^j)^2\}$. In particular, for n sufficiently large, we have $0 \in I_n^j$ and

$$\lim_{n \to \infty} \|v_n^j(0) - \phi_n^j\|_{\dot{H}^{s_c}_D(\Omega)} = 0.$$

When j conforms to Cases 2, 3, or 4, we apply the nonlinear embedding theorems of the previous section to construct the nonlinear profiles. More precisely, let v_n^j be the global solutions to (1.1) constructed in Theorems 4.1, 4.3, or 4.4, respectively.

The $\dot{H}_{D}^{s_{c}}(\Omega)$ decoupling of the profiles ϕ^{j} in (3.28) together with the definition of E_{c} ensures that for sufficiently large j, the profiles v_{n}^{j} are global and scatter, say $j \geq J_{0}$; indeed, for large enough j, we are in the small-data regime. Then we want to show that there exists some $1 \leq j_{0} < J_{0}$ such that

$$\lim_{n \to \infty} \sup_{[0, \sup I_n^{j_0})} (v_n^{j_0}) = \infty.$$
 (5.5)

Once a 'bad' nonlinear profile like (5.5) is obtained, we can prove that only one such profile exists. To see this, one needs to adapt the argument from [30, Lemma 3.3] to see that the $\dot{H}_{D}^{s_{c}}(\Omega)$ decoupling persists in time. Then, the 'critical' nature of E_{c} can be used to rule out the possibility of multiple profiles. Thus there is a single profile in the decomposition (5.4) (i.e. $J^{*}=1$) and we can write

$$u_n(0) = \phi_n + w_n \quad \text{with} \quad \lim_{n \to \infty} ||w_n||_{\dot{H}_D^1(\Omega)} = 0.$$
 (5.6)

If ϕ_n conforms to Cases 2, 3, or 4, then by Theorems 4.1, 4.3, or 4.4, there are global solutions v_n to (1.1) with data $v_n(0) = \phi_n$ that admit a uniform space-time bound. By Theorem 2.19, this space-time bound extends to the solutions u_n for n large enough, which contradicts (5.3). Therefore, ϕ_n must conform to Case 1 and (5.6) becomes

$$u_n(0) = e^{it_n \lambda_n^2 \Delta_{\Omega}} \phi + w_n \quad \text{with} \quad \lim_{n \to \infty} ||w_n||_{\dot{H}_D^{s_c}(\Omega)} = 0$$

and $t_n \equiv 0$ or $t_n \to \pm \infty$. If $t_n \equiv 0$, then we obtain the desired compactness. Thus, we only need to preclude that $t_n \to \pm \infty$.

Let us suppose $t_n \to \infty$; the case $t_n \to -\infty$ can be treated symmetrically. In this case, the Strichartz inequality and the monotone convergence theorem yield

$$S_{\geq 0}\left(e^{it\Delta_{\Omega}}u_n(0)\right) = S_{\geq 0}\left(e^{i(t+t_n\lambda_n^2)\Delta_{\Omega}}\phi + e^{it\Delta_{\Omega}}w_n\right) \longrightarrow 0 \text{ as } n \to \infty.$$

By the small data theory, this implies that $S_{\geq 0}(u_n) \to 0$, which contradicts (5.3).

We now prove the existence of at least one bad profile by assuming, for contradiction, that no bad nonlinear profiles exist. In this scenario, we have:

$$\sum_{j\geq 1} S_{[0,\infty)}(v_n^j) \lesssim_{E_c} 1. \tag{5.7}$$

Indeed, for sufficiently large n, we remain in the small-data regime. Then by the small-data local well posedness, $S_{[0,\infty)}(v_n^j) \lesssim \|v_n^j\|_{\dot{H}^{s_c}_D(\Omega)}$, which together with the decoupling (3.28) allows us to bound the tail by E_c .

Next, we use (5.7) and the stability result (Theorem 2.19) to constrain the scattering size of u_n , contradicting (5.3). To proceed, we define the approximations

$$u_n^J(t) = \sum_{j=1}^J v_n^j(t) + e^{it\Delta} w_n^J.$$
 (5.8)

By the construction of v_n^j , it is easy to verify that

$$\lim_{n \to \infty} \sup_{n \to \infty} \|u_n(0) - u_n^J(0)\|_{\dot{H}_D^{s_c}(\Omega)} = 0.$$
 (5.9)

Furthermore, we claim:

$$\lim_{J \to \infty} \limsup_{n \to \infty} S_{[0,\infty)}(u_n^J) \lesssim_{E_c} 1.$$
(5.10)

To justify (5.10), observe that by (3.27) and (5.7), it suffices to prove

$$\lim_{J \to \infty} \limsup_{n \to \infty} \left| S_{[0,\infty)} \left(\sum_{j=1}^{J} v_n^j \right) - \sum_{j=1}^{J} S_{[0,\infty)}(v_n^j) \right| = 0.$$
 (5.11)

Note that

$$\left| \left| \sum_{j=1}^J v_n^j \right|^{\frac{5\alpha}{2}} - \sum_{j=1}^J \left| v_n^j \right|^{\frac{5\alpha}{2}} \right| \lesssim_J \sum_{j \neq k} \left| v_n^j \right|^{\frac{5\alpha}{2} - 1} \left| v_n^k \right|.$$

It follows from Hölder's inequality that

$$LHS(5.11) \lesssim_{J} \sum_{j \neq k} \|v_n^j\|_{L_t^{\frac{5\alpha}{2}} L_x^{\frac{5\alpha}{2}}([0,\infty) \times \Omega)}^{\frac{5\alpha}{2}} \|v_n^j v_n^k\|_{L_t^{\frac{5\alpha}{4}} L_x^{\frac{5\alpha}{4}}([0,\infty) \times \Omega)}.$$
 (5.12)

Following Keraani's argument [26, Lemma 2.7], with $j \neq k$, we can first use (4.1), (4.20) and (4.25) to approximate v^j and v^k by compactly supported functions in $\mathbb{R} \times \mathbb{R}^3$, then using the asymptotic orthogonality (3.30) to demonstrate

$$\lim_{n \to \infty} \sup \left(\|v_n^j v_n^k\|_{L_t^{\frac{5\alpha}{4}} L_x^{\frac{5\alpha}{4}}([0,\infty) \times \Omega)} + \|v_n^j (-\Delta_{\Omega})^{\frac{s_c}{2}} v_n^k\|_{L_t^{\frac{5\alpha}{4}} L_x^{\frac{15\alpha}{15\alpha - 8}}([0,\infty) \times \Omega)} \right) = 0.$$
 (5.13)

Combining this with (5.12), we see that (5.11) (and hence (5.10)) is valid.

With (5.9) and (5.10) in place, proving that u_n^J asymptotically solves (1.1) reduces to showing:

$$\lim_{J \to \infty} \limsup_{n \to \infty} \|(i\partial_t + \Delta)u_n^J - |u_n^J|^\alpha u_n^J\|_{\dot{N}^{s_c}([0,\infty) \times \Omega)} = 0.$$
 (5.14)

Once this is established, we can apply the stability Theorem 2.19 to bound the scattering size of u_n , contradicting (5.3). This completes the proof of proposition 5.1.

It sufficies to prove (5.14), which relys on demonstrating:

Lemma 5.2 (Decoupling of nonlinear profiles). Let $F(u) = |u|^{\alpha}u$. Then

$$\lim_{J \to \infty} \limsup_{n \to \infty} \|F(\sum_{j=1}^{J} v_n^j) - \sum_{j=1}^{J} F(v_n^j)\|_{\dot{N}^{s_c}([0,\infty) \times \Omega)} = 0, \tag{5.15}$$

$$\lim_{J \to \infty} \limsup_{n \to \infty} \|F(u_n^J - e^{it\Delta} w_n^J) - F(u_n^J)\|_{\dot{N}^{s_c}([0,\infty) \times \Omega)} = 0.$$
 (5.16)

In the energy-critical setting, i.e., $s_c = 1$, one can instead use the pointwise estimate

$$\left| \nabla \left(F \left(\sum_{j=1}^{J} v_n^j \right) - \sum_{j=1}^{J} F(v_n^j) \right) \right| \lesssim_J \sum_{j \neq k} |\nabla v_n^j| |v_n^k|^{\alpha}$$
 (5.17)

and (5.13) to prove (5.15) and (5.16); the key is to exhibit terms that all contain some v_n^j paired against some v_n^k for $j \neq k$. In the case $s_c = 0$, there are also pointwise estimates similar to (5.17). However, when $s_c \neq 0, 1$, a new difficulty arises as the nonlocal operator $|\nabla|^{s_c}$ does not respect pointwise estimates in the spirit of (5.17).

To address this issue, in the subcritical case ($s_c < 1$), Murphy [45] employs the Littlewood-Paley square function estimates, which hold for all s > 0 and $1 < r < \infty$:

$$\|(\sum N^{2s}|f_N(x)|^2)^{1/2}\|_{L_x^r(\mathbb{R}^d)} \sim \|(\sum N^{2s}|f_{>N}(x)|^2)^{1/2}\|_{L_x^r(\mathbb{R}^d)} \sim \||\nabla|^s f\|_{L_x^r(\mathbb{R}^d)}, \tag{5.18}$$

to work at the level of individual frequencies. By utilizing maximal function and vector maximal function estimates, he adapts the standard arguments to this context. In the supercritical case $(s_c > 1)$, Killip and Visan [29] employed the following equivalence (see, e.g., [51]):

$$\||\nabla|^s f\|_{L^q_x} \sim \|\mathcal{D}_s(f)\|_{L^q_x},$$
 (5.19)

where the operator \mathcal{D}_s is defined as

$$\mathcal{D}_s(f)(x) := \left(\int_0^\infty \left| \int_{|y| < 1} \frac{|f(x + ry) - f(x)|}{r^{1 + 2s}} \, dy \right|^2 dr \right)^{1/2},$$

which behaves like $|\nabla|^s$ under symmetries. They then used the following pointwise inequality:

$$\mathcal{D}_s(w \cdot [F'(u+v) - F'(u)]) \lesssim \mathcal{D}_s(w)|v|^{\alpha} + M(|w|)M(|v|)[\mathcal{D}_s(u+v) + \mathcal{D}_s(u)],$$

where M denotes the Hardy-Littlewood maximal function. By combining this inequality with various permutations of the techniques discussed above, they adapted the standard arguments to this context.

In this paper, we follow the arguments in [29,45] and sketch the proof of Lemma 5.2.

Proof of (5.15). By induction, it suffices to treat the case of two summands. To simplify notation, we write $f = v_n^j$ and $g = v_n^k$ for some $j \neq k$, and are left to show

$$|||f+g|^{\alpha}(f+g)-|f|^{\alpha}f-|g|^{\alpha}g||_{\dot{N}^{s_c}([0,\infty)\times\Omega)}\to 0$$
 as $n\to\infty$.

We first rewrite

$$|f+g|^{\alpha}(f+g) - |f|^{\alpha}f - |g|^{\alpha}g = (|f+g|^{\alpha} - |f|^{\alpha})f + (|f+g|^{\alpha} - |g|^{\alpha})g.$$

By symmetry, it suffices to treat

$$\|\left(|f+g|^{\alpha}-|f|^{\alpha}\right)f\|_{\dot{N}^{s_{c}}([0,\infty)\times\Omega)}.$$
(5.20)

We then utilize Theorem 2.6 and the Littlewood-Paley square function estimates (5.18) to reduce (5.20) to handling

$$\left\| \left(\sum_{N} \left| |\nabla|^{s_c} P_N \left(\left(|f + g|^{\alpha} - |f|^{\alpha} \right) f \right) \right|^2 \right)^{\frac{1}{2}} \right\|_{L^{\frac{5\alpha}{2(\alpha+1)}} L^{\frac{30\alpha}{2'\alpha-8}}}. \tag{5.21}$$

Then the key step is to perform a decomposition such that all resulting terms to estimate have f paired against g inside a single integrand. For such terms, the asymptotic orthogonality (5.13) can be used.

Following the arguments in [45], we decompose (5.21) into terms such that each term contains pairings of f and g. For instance, one of the terms is

$$\|\left(\sum_{N} |N^{s_c} f_{>N} M(g|f|^{\alpha-1})|^2\right)^{1/2}\|_{L_{t}^{\frac{5\alpha}{2(\alpha+1)}} L_{x}^{\frac{30\alpha}{27\alpha-8}}}.$$
(5.22)

Using Hölder's inequality and maximal function estimates, this term can be controlled as

$$\|(\sum_{N}|N^{s_c}f_{>N}|^2)^{1/2}\|_{L_{t}^{\frac{5\alpha}{2}}L_{x}^{\frac{30\alpha}{15\alpha-8}}}\||g||f|^{\alpha-1}\|_{L_{t,x}^{\frac{d+2}{2}}}.$$

By (5.18), the first term is bounded by $|||\nabla|^{s_c}v_n^j||_{L_{t,x}^{\frac{2(d+2)}{d}}}$, which is further bounded by the construction of v_n^j . The second term vanishes as $n\to\infty$ due to the asymptotic orthogonality of parameters (5.13). The other terms similar to (5.22) can be handled similarly, thereby completing the proof of (5.15).

Proof of (5.16). For this term, we will need to make use of (3.27) instead of (5.13). The proof is, in fact, similar to that of (5.15). By employing the same arguments as in the proof of (5.13), we obtain terms involving either $e^{it\Delta}w_n^J$ or $|\nabla|^{s_c}e^{it\Delta}w_n^J$. The terms in which $e^{it\Delta}w_n^J$ appears without derivatives will be relatively straightforward to handle, as (3.27) can be applied directly. However, the terms involving $|\nabla|^{s_c}e^{it\Delta}w_n^J$ require more careful analysis. Indeed, we need to first apply the local smoothing estimate provided in Corollary 2.11 and then use (3.27) to show that these terms vanish as $n \to \infty$.

We now apply the Palais-Smale condition in Proposition 5.1 to prove Theorem 1.8.

Proof of Theorem 1.8. If Theorem 1.2 fails to be true, using a standard argument (see e.g. [33, Theorem 5.2]), we can employ the Palais-Smale condition to construct a minimal counterexample $u: I \times \Omega \to \mathbb{C}$ that satisfies

$$S_{>0}(u) = S_{<0}(u) = \infty, \tag{5.23}$$

and whose orbit $\{u(t): t \in I\}$ is precompact in $\dot{H}_D^{s_c}(\Omega)$. Moreover, due to the compact modulation parameter $N(t) \equiv 1$, we deduce that the maximal lifespan interval $I = \mathbb{R}$ (see e.g. [33, Corollary 5.19]).

Finally, we prove the lower bound (1.5). We argue by contradiction. Suppose there exist sequences $R_n \to \infty$ and $\{t_n\} \subset \mathbb{R}$ such that

$$\int_{\Omega \cap \{|x| \le R_n\}} |u(t_n, x)|^{\frac{3}{2}\alpha} dx \to 0.$$

Passing to a subsequence, we find $u(t_n) \to \phi$ in $\dot{H}_D^{s_c}(\Omega)$ for some non-zero $\phi \in \dot{H}_D^{s_c}(\Omega)$. Note that if ϕ were zero, then the solution u would have a $\dot{H}_D^{s_c}(\Omega)$ norm less than the small data threshold, which would contradict (5.23). By Sobolev embedding, $u(t_n) \to \phi$ in $L^{\frac{3}{2}\alpha}$, and since $R_n \to \infty$,

$$\int_{\Omega} |\phi(x)|^{\frac{3}{2}\alpha} dx = \lim_{n \to \infty} \int_{\Omega \cap \{|x| \le R_n\}} |\phi(x)|^{\frac{3}{2}\alpha} dx = \lim_{n \to \infty} \int_{\Omega \cap \{|x| \le R_n\}} |u(t_n, x)|^{\frac{3}{2}\alpha} dx = 0.$$

This contradicts the fact that $\phi \neq 0$, thereby completing the proof of Theorem 1.8.

6. The cases
$$1 < s_c < \frac{3}{2}$$
 and $s_c = \frac{1}{2}$.

In this section, we rule out the existence of almost periodic solutions as in Theorem 1.8 in the cases $1 < s_c < 3/2$ and $s_c = \frac{1}{2}$. We employ a space-localized Morawetz inequality as in the work of Bourgain [3] on the radial energy-critical NLS. See also [18,52].

Lemma 6.1 (Morawetz inequality). Let $1 < s_c < \frac{3}{2}$ and let u be a solution to (1.1) on the time interval I. Then for any $A \ge 1$ with $A|I|^{1/2} \ge diam(\Omega^c)$ we have

$$\int_{I} \int_{|x| \le A|I|^{1/2}, x \in \Omega} \frac{|u(t, x)|^{\alpha + 2}}{|x|} dx dt \lesssim A|I|^{s_c - 1/2}, \tag{6.1}$$

where the implicit constant depends only on $\|u\|_{\dot{H}^{s_c}_{\mathcal{D}}(\Omega)}$.

Proof. Let $\phi(x)$ be a smooth, radial bump function such that $\phi(x) = 1$ for $|x| \le 1$ and $\phi(x) = 0$ for |x| > 2. We set $R \ge \operatorname{diam}(\Omega^c)$ and denote $a(x) := |x| \phi\left(\frac{x}{R}\right)$. Then, for $|x| \le R$ we have

$$\partial_j \partial_k a(x)$$
 is positive definite, $\nabla a(x) = \frac{x}{|x|}$, and $\Delta \Delta a(x) < 0$. (6.2)

For |x| > R, we have the following rough bounds:

$$|\partial_k a(x)| \lesssim 1, \quad |\partial_j \partial_k a(x)| \lesssim \frac{1}{R}, \quad \text{and} \quad |\Delta \Delta a(x)| \lesssim \frac{1}{R^3}.$$
 (6.3)

By the direct calculus, we have the following identity

$$2\partial_t \operatorname{Im}(\bar{u}\partial_j u) = -4\partial_k \operatorname{Re}(\partial_k u \partial_j \bar{u}) + \partial_j \Delta(|u|^2) - \frac{2\alpha}{\alpha + 2} \partial_j (|u|^{\alpha + 2}). \tag{6.4}$$

Multiplying by $\partial_j a$ in both sides and integrating over Ω , we obtain

$$2\partial_{t}\operatorname{Im}\int_{\Omega}\bar{u}\partial_{j}u\partial_{j}a\,dx$$

$$= -4\operatorname{Re}\int_{\Omega}\partial_{k}(\partial_{k}u\partial_{j}\bar{u})\partial_{j}a\,dx + \int_{\Omega}\partial_{j}\Delta(|u|^{2})\partial_{j}a\,dx - \frac{2\alpha}{\alpha+2}\int_{\Omega}\partial_{j}(|u|^{\alpha+2})\partial_{j}a\,dx. \tag{6.5}$$

Now, we give the upper bound of the left-hand side of (6.5) which follows immediately from Hölder and the Sobolev embedding:

$$2\partial_{t} \operatorname{Im} \int_{\Omega} \bar{u} \partial_{j} u \partial_{j} a \, dx \lesssim \|u\|_{L_{x}^{\frac{6}{3-2s_{c}}}(\Omega)} \|\nabla u\|_{L_{x}^{\frac{6}{5-2s_{c}}}(\Omega)} \|\nabla a\|_{L_{x}^{\frac{3}{2s_{c}-1}}(\Omega)} \lesssim R^{2s_{c}-1} \|u\|_{\dot{H}_{D}^{s_{c}}(\Omega)}. \tag{6.6}$$

Next, we find a lower bound on right-hand side of (6.5). By using the Gauss theorem, we get

$$-4\operatorname{Re}\int_{\Omega}\partial_{k}(\partial_{k}u\partial_{j}\bar{u})\partial_{j}a\,dx = 4\operatorname{Re}\int_{\partial\Omega}\partial_{k}u\partial_{j}a\partial_{j}\bar{u}\vec{n}_{k}\,d\sigma(x) + 4\operatorname{Re}\int_{\Omega}\partial_{k}u\partial_{j}\bar{u}\partial_{k}\partial_{j}a\,dx,$$

where \vec{n} denotes the outer normal vector to Ω^c . We write $\partial_j \bar{u} \vec{n}_j = \nabla \bar{u} \cdot \vec{n} = \bar{u}_n$ and $\partial_j a n_j = \nabla a \cdot \vec{n} = a_n$. Moreover, from the Dirichlet boundary condition, the tangential derivative of u vanishes on the boundary:

$$\nabla u = (\nabla u \cdot \vec{n})\vec{n} = u_n \vec{n}, \text{ and } \partial_i \overline{u}_i \partial_i a = u_n a_n \vec{n}$$

Combining the analysis above, we obtain

$$-4\operatorname{Re}\int_{\Omega}\partial_{k}(\partial_{k}u\partial_{j}\bar{u})\partial_{j}a\,dx \geq 4\int_{\partial\Omega}a_{n}|u_{n}|^{2}\,d\sigma(x) + 4\int_{|x|\geq R}\partial_{k}u\partial_{j}\bar{u}\partial_{k}\partial_{j}a\,dx$$

$$\geq 4\int_{\partial\Omega}a_{n}|u_{n}|^{2}\,d\sigma(x) - \|\Delta a\|_{L_{x}^{\frac{3}{2(s_{c}-1)}}(\{x:|x|>R\})}\|\nabla u\|_{L_{x}^{\frac{6}{5-2s_{c}}}(\Omega)}^{2}$$

$$\geq 4\int_{\partial\Omega}a_{n}|u_{n}|^{2}\,d\sigma(x) - CR^{2s_{c}-3}\|u\|_{\dot{H}_{D}^{s_{c}}(\Omega)}^{2}.$$

We can bound the second term on RHS of (6.5) in a similar way:

$$\int_{\Omega} \partial_{j} \Delta(|u|^{2}) \partial_{j} a \, dx = \int_{\Omega} \partial_{j} (\Delta(|u|^{2}) \partial_{j} a) dx - \int_{\Omega} \Delta(|u|^{2}) \Delta a \, dx$$

$$= -\int_{\partial\Omega} \Delta(|u|^{2}) \partial_{j} a \vec{n}_{j} \, d\sigma(x) - \int_{\Omega} |u|^{2} \Delta \Delta a \, dx$$

$$= -2 \int_{\partial\Omega} |\nabla u|^{2} a_{n} \, d\sigma(x) - \int_{|x| \leq R} |u|^{2} \Delta^{2} a \, dx - \int_{|x| \geq R} |u|^{2} \Delta^{2} a \, dx$$

$$\geq -2 \int_{\partial\Omega} |u_{n}|^{2} a_{n} \, d\sigma(x) - ||u||^{2}_{L_{x}^{\frac{6}{3-2s_{c}}}(\Omega)} ||\Delta^{2} a||_{L_{x}^{\frac{3}{2s_{c}}}(\{x:|x|>R\})}$$

$$\geq -2 \int_{\partial\Omega} |u_{n}|^{2} a_{n} \, d\sigma(x) - CR^{3-2s_{c}} ||u||_{\dot{H}_{D}^{s}(\Omega)}^{2s_{c}}. \tag{6.7}$$

Finally, it remains to estimate the third term on right-hand side of (6.5) by using (6.2) and (6.3):

$$-\frac{2\alpha}{\alpha+2} \int_{\Omega} \partial_{j} (|u|^{\alpha+2}) \partial_{j} a \, dx = \frac{2\alpha}{\alpha+2} \int_{\Omega} |u|^{\alpha+2} \Delta a \, dx$$

$$= \frac{4\alpha}{\alpha+2} \int_{|x| \leq R} \frac{|u|^{\alpha+2}}{|x|} \, dx - \frac{4\alpha}{\alpha+2} \int_{\Omega \cap \{x:|x| > R\}} \Delta a |u|^{\alpha+2} dx$$

$$\geq \frac{4\alpha}{\alpha+2} \int_{|x| \leq R} \frac{|u|^{\alpha+2}}{|x|} \, dx - \|\Delta a\|_{L_{x}^{\frac{3}{2s_{c}-1}}(\{x:|x| > R\})} \|u\|_{L_{x}^{\frac{6}{3-2s_{c}}}(\Omega)}^{\alpha+2}$$

$$\geq \frac{4\alpha}{\alpha+2} \int_{|x| \leq R} \frac{|u|^{\alpha+2}}{|x|} \, dx - CR^{2s_{c}-3} \|u\|_{\dot{H}_{D}^{sc}(\Omega)}^{\alpha+2}. \tag{6.8}$$

Putting these together and using the fact that $a_n \geq 0$ on $\partial \Omega$, we have

$$LHS(6.5) \gtrsim \int_{|x| \le R} \frac{|u|^{\alpha+2}}{|x|} dx - R^{2s_c - 3} (\|u\|_{\dot{H}_D^{s_c}(\Omega)}^2 + \|u\|_{\dot{H}_D^{s_c}(\Omega)}^{\alpha+2}). \tag{6.9}$$

Integrating (6.5) over I and using the upper bound for the left-hand side of (6.5) and the lower bound for the right-hand side of (6.5), we finally deduce

$$\int_{I} \int_{|x| < R, x \in \Omega} \frac{|u|^{\alpha + 2}}{|x|} dx dt \lesssim R^{2s_c - 1} + R^{2s_c - 3} |I|.$$

Taking $R = A|I|^{1/2}$ yields (6.1). This completes the proof of the lemma.

In the proof of Lemma 6.1, by taking $R \to +\infty$ and using the same argument as in [6, Lemma 2.3] to control the upper bound of the Morawetz action, we can obtain the following non-spatially localized Lin-Strauss Morawetz inequality.

Lemma 6.2 (Morawetz inequality). Let $s_c = \frac{1}{2}$ and let u be a solution to (1.1) on the time interval I. Then we have

$$\int_{I} \int_{\Omega} \frac{|u(t,x)|^{\alpha+2}}{|x|} dx dt \lesssim ||u||_{L_{t}^{\infty}\dot{H}_{D}^{s_{c}}(\Omega)}^{2}.$$
(6.10)

We now use Lemma 6.1 and Lemma 6.2 to prove the following.

Theorem 6.3. There are no almost periodic solutions u to (1.1) as in Theorem 1.8 with $1 < s_c < 3/2$.

Proof. By contradiction, we suppose that there exists a minimal blow-up solution u whose the orbit is precompact in Hilbert space $\dot{H}_{D}^{s_c}(\Omega)$ and satisfies (1.5). Integrating over I with length $|I| \geq 1$, we obtain

$$|I| \lesssim R \int_I \int_{\Omega \cap \{|x| \leq R\}} \frac{|u(t,x)|^{\alpha+2}}{|x|} \, dx \, dt \lesssim R \int_I \int_{\Omega \cap \{|x| \leq R|I|^{1/2}\}} \frac{|u(t,x)|^{\alpha+2}}{|x|} \, dx \, dt.$$

On the other hand, for $R|I|^{1/2} \ge 1$, the Morawetz inequality (Lemma 6.1) and Lemma 6.2 yield that

$$|I| \lesssim \int_{I} \int_{\Omega \cap I|x| \leq R|I|^{1/2}} \frac{|u(t,x)|^{\alpha+2}}{|x|} dx dt \lesssim R|I|^{s_c-1/2},$$

with the implicit constant depending only on $\|u\|_{\dot{H}^{s_c}_D(\Omega)}$. Choosing I sufficiently large depending on R and $\|u\|_{\dot{H}^{s_c}_D(\Omega)}$, we get a contradiction, which completes the proof of Theorem 6.3.

7. The case
$$\frac{1}{2} < s_c < 1$$
.

In this section, we rule out the almost periodic solutions in the case $1/2 < s_c < 1$. The main technical tool we use is a long-time Strichartz estimate, which will be proved in subsection 7.1. In subsection 7.2, we prove a frequency-localized Lin-Strauss Morawetz inequality, which we use in subsection 7.3 to rule out the almost periodic solutions.

7.1. Long time Strichartz estimates. In this subsection, we prove a long-time Strichartz estimates adapted to the Lin-Strauss Morawetz inequality. Such estimates were first developed by Dodson [8] and have been proven to be a powerful tool for ruling out the existence of a minimal counterexample, see [32, 57] for the energy-critical setting, [45, 47, 61] for the inter-critical setting, and [13, 43, 44, 47] for the super-critical setting. In this paper, we establish, for the first time, a long-time Strichartz estimate in the setting of the exterior domain Schrödinger equation. This long time Strichartz estimate will be a crucial technical tool in subsection 7.2 to control the error terms arising from frequency truncation in the Lin-Strauss Morawetz inequality.

Throughout Section 7, we use the following notation:

$$A_I(N) := \|(-\Delta_{\Omega})^{\frac{s_c}{2}} P_{\leq N}^{\Omega} u\|_{L_t^2 L_x^6(I \times \Omega)}.$$

The main result of this subsection is the following.

Proposition 7.1 (Long time Strichartz estimate). Let $u : \mathbb{R} \times \Omega \to \mathbb{C}$ be an almost periodic solution to (1.1) with $1/2 < s_c < 1$. Then for any N > 0, we have

$$A_I(N) \lesssim_u 1 + N^{s_c - 1/2} |I|^{1/2}.$$
 (7.1)

Moreover, for any $\varepsilon > 0$, there exists $N_0 = N_0(\varepsilon) > 0$ so that for any $N \leq N_0$,

$$A_I(N) \lesssim_u \varepsilon (1 + N^{s_c - 1/2} |I|^{1/2}).$$
 (7.2)

We prove Proposition 7.1 by induction. The inductive step relies on the following.

Lemma 7.2. Let $\eta > 0$, u and I be as above. For any N > 0, we have

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}} P_{\leq N}^{\Omega} F(u)\|_{L_t^2 L_x^{6/5}(I \times \Omega)} \lesssim_u C_{\eta} \|u_{\leq N/\eta}\|_{L_t^{\infty} \dot{H}_D^{s_c}(\Omega)} N^{s_c - 1/2} |I|^{1/2} + \sum_{M > N/\eta} \left(\frac{N}{M}\right)^{s_c} A_I(M).$$

Proof. We fix $0 < \eta < 1$ and decompose the nonlinearity as follows:

$$F(u) = F(u_{\leq N/\eta}) + [F(u) - F(u_{\leq N/\eta})].$$

Using the fractional chain rule (2.1), Hölder and Sobolev embedding, we estimate

$$\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}P_{\leq N}^{\Omega}F(u_{\leq N/\eta})\|_{L_{t}^{2}L_{x}^{6/5}(I\times\Omega)}$$

$$\lesssim \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{\leq N/\eta}\|_{L_{t}^{\infty}L_{x}^{2}(I\times\Omega)}^{\alpha}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{\leq N/\eta}\|_{L_{t}^{2}L_{x}^{6}(I\times\Omega)}$$

$$\lesssim \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}P_{\leq c(\eta)}^{\Omega}u_{\leq N/\eta}\|_{L_{t}^{\infty}L_{x}^{2}(I\times\Omega)}^{\alpha}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{\leq N/\eta}\|_{L_{t}^{2}L_{x}^{6}(I\times\Omega)}$$

$$+ \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}P_{>c(\eta)}^{\Omega}u_{\leq N/\eta}\|_{L_{x}^{\infty}L_{x}^{2}(I\times\Omega)}^{\alpha}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{\leq N/\eta}\|_{L_{t}^{2}L_{x}^{6}(I\times\Omega)}.$$

$$(7.3)$$

For the first term, we use (1.4) to get

$$(7.3) \lesssim \eta^{\alpha} \| (-\Delta_{\Omega})^{\frac{s_c}{2}} u_{\leq N/\eta} \|_{L_t^2 L_x^6(I \times \Omega)} \lesssim \eta^{s_c} A_I(N/\eta).$$
 (7.5)

For the next term, we note that we only need to consider the case $c(\eta) < N/\eta$, in which case we have $1 \lesssim_{\eta} N^{s_c-1/2}$. Using Bernstein and Lemma 1.7, we estimate

$$(7.4) \lesssim_{u} C_{\eta} N^{s_{c}-1/2} \| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} u_{\leq N/\eta} \|_{L_{t}^{\infty} L_{x}^{2}(I \times \Omega)} \lesssim_{u} C_{\eta} N^{s_{c}-1/2} \| u_{\leq N/\eta} \|_{L_{t}^{\infty} \dot{H}_{D}^{s_{c}}(I \times \Omega)}.$$
 (7.6)

Combining (7.5) and (7.6), we obtain

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}} P_{\leq N}^{\Omega} F(u_{\leq N/\eta})\|_{L_t^2 L_x^{6/5}(I \times \Omega)} \lesssim_u \eta^{s_c} A_I(N/\eta) + C_{\eta} \|u_{\leq N/\eta}\|_{L_t^{\infty} \dot{H}_D^{s_c}(I \times \Omega)} N^{s_c - 1/2} |I|^{\frac{1}{2}}.$$
 (7.7)

Next, we use Bernstein, Hölder and Sobolev embedding to estimate

$$\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}P_{\leq N}^{\Omega}\left(F(u) - F(u_{\leq N/\eta})\right)\|_{L_{t}^{2}L_{x}^{6/5}(I\times\Omega)}$$

$$\lesssim N^{s_{c}}\|u\|_{L_{t}^{\infty}L_{x}^{3\alpha/2}(I\times\Omega)}^{\alpha}\sum_{M>N/\eta}\|u_{M}\|_{L_{t}^{2}L_{x}^{6}(I\times\Omega)}$$

$$\lesssim N^{s_{c}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u\|_{L_{t}^{\infty}L_{x}^{2}(I\times\Omega)}^{\alpha}\sum_{M>N/\eta}M^{-s_{c}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{M}\|_{L_{t}^{2}L_{x}^{6}(I\times\Omega)}$$

$$\lesssim u\sum_{M>N/\eta}\left(\frac{N}{M}\right)^{s_{c}}A_{I}(M),$$

which together with (7.7) yields the desired estimate in Lemma 7.2.

We now turn to the proof of Proposition 7.1.

Proof of Proposition 7.1. We proceed by induction. For the base case, we take N > 1, so that by Lemma 1.7, we have

$$A_I(N) \lesssim_u 1 + |I|^{1/2} \le C_u \left[1 + N^{s_c - 1/2} |I|^{1/2} \right].$$
 (7.8)

This inequality clearly remains true if we replace C_u by any larger constant.

We now suppose that (7.8) holds at frequencies $\geq 2N$; we will use Lemma 7.2 to show that it holds at frequency N.

Applying Strichartz and Lemma 7.2 gives

$$A_{I}(N) \leq \tilde{C}_{u} \left[\inf_{t \in I} \|u_{\leq N}(t)\|_{\dot{H}_{D}^{s_{c}}(\Omega)} + C_{\eta} \|u_{\leq N/\eta}\|_{L_{t}^{\infty} \dot{H}_{D}^{s_{c}}(I \times \Omega)} N^{s_{c}-1/2} |I|^{1/2} \right.$$

$$\left. + \sum_{M \geq N/\eta} \left(\frac{N}{M} \right)^{s_{c}} A_{I}(M) \right]$$

$$\leq \tilde{C}_{u} \left[1 + C_{\eta} N^{s_{c}-1/2} |I|^{1/2} + \sum_{M \geq N/\eta} \left(\frac{N}{M} \right)^{s_{c}} A_{I}(M) \right]. \tag{7.9}$$

By the inductive hypothesis, we have

$$\begin{split} A_I(N) &\leq \tilde{C}_u \left[1 + C_\eta N^{s_c - 1/2} |I|^{1/2} + \sum_{M \geq N/\eta} \left(\frac{N}{M} \right)^{s_c} \left(C_u + C_u M^{s_c - 1/2} |I|^{1/2} \right) \right] \\ &\leq \tilde{C}_u \left[1 + C_\eta N^{s_c - 1/2} |I|^{1/2} \right] + C_u \tilde{C}_u \left[\eta^{s_c} + \eta^{1/2} N^{s_c - 1/2} |I|^{1/2} \right]. \end{split}$$

Choosing η sufficiently small depending on \tilde{C}_{u} , we obtain

$$A_I(N) \le \tilde{C}_u(1 + C_{\eta}N^{s_c - 1/2}|I|^{1/2}) + \frac{1}{2}C_u(1 + N^{s_c - 1/2}|I|^{1/2}).$$

Finally, choosing C_u possibly even larger to guarantee $C_u \geq 2(1 + C_\eta)\tilde{C}_u$, we deduce from the above inequality that

$$A_I(N) \le C_u(1 + N^{s_c - 1/2}|I|^{1/2}).$$

This completes the proof of (7.1) by induction.

With (7.1) in place, we can prove (7.2) by continuing from (7.9). In fact, for any $\eta > 0$ sufficiently small, the almost periodicity implies that

$$\lim_{N\to 0} \|u_{\leq N/\eta}\|_{L^\infty_t \dot{H}^{s_c}_D(I\times\Omega)} = 0.$$

This completes the proof of Proposition 7.1.

7.2. A Frequency-Localized Lin-Strauss Morawetz Inequality.

Proposition 7.3 (Frequency-localized Morawetz). Let $u : \mathbb{R} \times \Omega \to \mathbb{C}$ be an almost periodic solution to (1.1) with $1/2 < s_c < 1$. Then for any $\eta > 0$, there exists $N_0 = N_0(\eta) \in (0,1)$ such that for $N < N_0$ and $I \subset \mathbb{R}$, we have

$$\int_{I} \int_{\Omega} \frac{|u_{>N}(t,x)|^{\alpha+2}}{|x|} dx dt \lesssim_{u} \eta \left(N^{1-2s_{c}} + |I|\right).$$

To prove Proposition 7.3, we start by truncating the low frequencies of the solution and focus on $u_{>N}$ for some N > 0. Since $u_{>N}$ is not a solution to (1.1), we need to estimate the additional error terms brought by this frequency cut-off. To achieve this, we take N sufficiently small to maintain the most part of the solution and utilize the estimates proved in Proposition 7.1.

Lemma 7.4 (High and low frequency control). Let u and I be as above. With all spacetime norms over $I \times \Omega$, we have the following:

(i)Let (q,r) be an admissible pair. For any N>0 and $0 \le s < 1/2$, we have

$$\|(-\Delta_{\Omega})^{\frac{s}{2}}u_{>N}\|_{L_{t}^{q}L_{x}^{r}} \lesssim_{u} N^{s-s_{c}}(1+N^{2s_{c}-1}|I|)^{1/q}.$$
(7.10)

(ii) For any $\eta > 0$ and $0 < s < s_c$, there exists a parameter $N_1 = N_1(\eta)$ such that for $N < N_1$, we have

$$\|(-\Delta_{\Omega})^{\frac{s}{2}}u_{>N}\|_{L_{t}^{\infty}L_{x}^{2}} \lesssim_{u} \eta N^{s-s_{c}}.$$
(7.11)

(iii) For any $\eta > 0$, there exists a parameter $N_2 = N_2(\eta)$ such that for $N < N_2$, we have

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}} u_{< N}\|_{L^2_{-L^6}} \lesssim_u \eta (1 + N^{2s_c - 1} |I|)^{1/2}. \tag{7.12}$$

Proof. For (7.10), we first apply the interpolation, (1.3) and (7.1) to derive

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}}u_{< N}\|_{L_t^q L_x^r} \lesssim \|u\|_{L_t^{\infty}\dot{H}_x^{s_c}(\Omega)}^{1-\frac{2}{q}}\|(-\Delta_{\Omega})^{\frac{s_c}{2}}u_{< N}\|_{L_t^2 L_x^6}^{\frac{2}{q}} \lesssim (1+N^{2s_c-1}|I|)^{1/q}.$$

Then by Bernstein,

$$||u_{>N}||_{L_t^q L_x^r} \lesssim \sum_{M>N} M^{s-s_c} ||(-\Delta_{\Omega})^{\frac{s_c}{2}} u_M||_{L_t^q L_x^r}$$

$$\lesssim_u \sum_{M>N} M^{s-s_c} (1 + M^{2s_c-1} |I|)^{1/q} \lesssim_u N^{s-s_c} (1 + N^{2s_c-1} |I|)^{1/q}.$$

For (7.11), we use the property of the almost-periodic solution (1.4) and Bernstein's inequality to estimate

$$\|(-\Delta_{\Omega})^{\frac{s}{2}}u_{>N}\|_{L_{t}^{\infty}L_{x}^{2}} \lesssim c(\eta)^{s-s_{c}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{>c(\eta)}\|_{L_{t}^{\infty}L_{x}^{2}} + N^{s-s_{c}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{N\leq \cdot \leq c(\eta)}\|_{L_{t}^{\infty}L_{x}^{2}} \leq c(\eta)^{s-s_{c}} + \eta N^{s-s_{c}}.$$

Taking $N_1(\eta) = \eta^{1/(s_c - s)} c(\eta)$, we obtain the desired estimate (7.11).

For the last inequality
$$(7.12)$$
, it is equal to (7.2) .

We now proceed to prove Proposition 7.3.

Proof of Proposition 7.3. Throughout the proof, we take the norms over $I \times \Omega$ and thus we omit it for short.

Assume that $0 < \eta \ll 1$ and taking

$$N < \min\{N_1(\eta), \eta^2 N_2(\eta^{2s_c})\},$$

where N_1 and N_2 are as in Lemma 7.4. As a consequence, (7.10) implies that

$$||u_{>N/\eta^2}||_{L^2_+L^6_\sigma} \lesssim_u \eta N^{-s_c} (1 + N^{2s_c - 1}|I|)^{1/2}. \tag{7.13}$$

Moreover, using the fact $N/\eta^2 < N_2(\eta^{2s_c})$ we can apply (7.12) to show

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}} u_{< N/\eta^2}\|_{L^2_x L^6_x} \lesssim_u \eta (1 + N^{2s_c - 1} |I|)^{1/2}. \tag{7.14}$$

Then we define the Morawetz action by the following

$$\operatorname{Mor}(t) = 2\operatorname{Im} \int_{\Omega} \frac{x}{|x|} \cdot \nabla u_{>N}(t,x) \overline{u_{>N}(t,x)} dx.$$

Since $(i\partial_t + \Delta_\Omega)u_{>N} = P_{>N}^\Omega(F(u))$ and by the direct calculus, we have

$$\partial_t \operatorname{Mor}(t) = -4\operatorname{Re} \int_{\Omega} \partial_k (\partial_k u_{>N} \partial_j \overline{u}_{>N}) \frac{x_j}{|x|} dx + \int_{\Omega} \partial_j \Delta(|u|^2) \frac{x_j}{|x|} dx + 2 \int_{\Omega} \frac{x}{|x|} \cdot \{P_{>N}^{\Omega}(F(u)), u_{>N}\}_p dx,$$

where the momentum bracket $\{\cdot,\cdot\}_p$ is defined by $\{f,g\}_p := \operatorname{Re}(f\nabla \overline{g} - g\nabla \overline{f})$. Using the same argument as that used to derive (6.7), (6.8), and the fact that $\partial_{jk}|x|$ is positive definite, we have

$$-4\operatorname{Re}\int_{\Omega}\partial_{k}(\partial_{k}u_{>N}\partial_{j}\overline{u}_{>N})\frac{x_{j}}{|x|}dx+\int_{\Omega}\partial_{j}\Delta(|u|^{2})\frac{x_{j}}{|x|}dx\geq2\int_{\partial\Omega}|\nabla u_{>N}\cdot\vec{n}|^{2}\frac{x}{|x|}\cdot\vec{n}d\sigma(x)>0,$$

where \vec{n} denotes the outer normal to Ω^c . Hence

$$\partial_t \operatorname{Mor}(t) \ge 2 \int_{\Omega} \frac{x}{|x|} \cdot \{P_{>N}^{\Omega}(F(u)), u_{>N}\}_p dx.$$

The fundamental theorem of calculus yields that

$$\int_{I \times \Omega} \frac{x}{|x|} \cdot \{P_{>N}^{\Omega}(F(u)), u_{>N}\}_p dx \lesssim \|\text{Mor}(t)\|_{L_t^{\infty}(I)}. \tag{7.15}$$

By standard computation, one hase

$$\{F(u), u\}_p = -\frac{\alpha}{\alpha + 2} \nabla(|u|^{\alpha+2}).$$

Thus, we can write the truncate momentum bracket to the following

$$\begin{split} &\{P_{>N}^{\Omega}(F(u)), u_{>N}\}_{p} \\ &= \{F(u), u\}_{p} - \{F(u_{\leq N}), u_{\leq N}\}_{p} - \{F(u) - F(u_{\leq N}), u_{\leq N}\}_{p} - \{P_{\leq N}^{\Omega}(F(u)), u_{>N}\}_{p} \\ &= -\frac{\alpha}{\alpha + 2} \nabla(|u|^{\alpha + 2} - |u_{\leq N}|^{\alpha + 2}) - \{F(u) - F(u_{\leq N}), u_{\leq N}\}_{p} - \{P_{\leq N}^{\Omega}(F(u)), u_{>N}\}_{p} \\ &:= I + II + III. \end{split}$$

Integrating by parts, the contribution of I in the integral (7.15) can be bounded by a multiple of:

$$\int_{I} \int_{\Omega} \frac{|u_{>N}(t,x)|^{\alpha+2}}{|x|} dxdt \tag{7.16}$$

and to the right-hand side of (7.15) a multiple of:

$$\left\| \frac{1}{|x|} (u_{\leq N})^{\alpha+1} u_{>N} \right\|_{L_t^1 L_x^1} + \left\| \frac{1}{|x|} u_{\leq N} (u_{>N})^{\alpha+1} \right\|_{L_t^1 L_x^1} := I_1 + I_2. \tag{7.17}$$

For the second term II, we will use the divergence theorem to find that

$$II \lesssim \left\| \frac{1}{|x|} u_{\leq N} [F(u) - F(u_{\leq N})] \right\|_{L_t^1 L_x^1} + \left\| \nabla u_{\leq N} [F(u) - F(u_{\leq N})] \right\|_{L_t^1 L_x^1} := II_1 + II_2. \tag{7.18}$$

Finally, for the last term III, we use integrating by parts when the derivative acts on $u_{>N}$. Then we get that

$$III \lesssim \left\| \frac{1}{|x|} u_{>N} P_{\leq N}^{\Omega}(F(u)) \right\|_{L_{t}^{1} L_{x}^{1}} + \left\| u_{>N} \nabla P_{\leq N}^{\Omega}(F(u)) \right\|_{L_{t}^{1} L_{x}^{1}}. \tag{7.19}$$

Thus, building upon (7.15), we conclude that it remains to show:

$$\|\operatorname{Mor}\|_{L^{\infty}(I)} \lesssim_{u} \eta N^{1-2s_{c}},\tag{7.20}$$

and that the error terms (7.17) through (7.19) are enough to obtain the needed estimate, which can be controlled by $\eta(N^{1-2s_c} + |I|)$.

Next, we begin to prove (7.20). Making use of the Bernstein estimate, the Hardy inequality and (7.11) to obtain

$$\|\operatorname{Mor}\|_{L_{t}^{\infty}(I)} \lesssim \||\nabla|^{-1/2} \nabla u_{>N}\|_{L_{t}^{\infty} L_{x}^{2}} \||\nabla|^{1/2} \left(\frac{x}{|x|} u_{>N}\right)\|_{L_{t}^{\infty} L_{x}^{2}} \lesssim \||\nabla|^{1/2} u_{>N}\|_{L_{t}^{\infty} L_{x}^{2}}^{2} \lesssim \|(-\Delta_{\Omega})^{\frac{1}{4}} u_{>N}\|_{L_{t}^{\infty} L_{x}^{2}}^{2} \lesssim u \, \eta N^{1-2s_{c}}.$$

We next turn to give the estimate of the error terms (7.17) through (7.19). To estimate I_1 , we first note that by interpolation, (1.3) and (7.12),

$$\|(-\Delta_{\Omega})^{\frac{s_c}{2}}u_{\leq N}\|_{L_{t}^{2(\alpha+1)}L_{x}^{\frac{6(\alpha+1)}{3\alpha+1}}}^{\alpha+1}\lesssim \|(-\Delta_{\Omega})^{\frac{s_c}{2}}u\|_{L_{t}^{\infty}L_{x}^{2}}^{\alpha}\|(-\Delta_{\Omega})^{\frac{s_c}{2}}u_{\leq N}\|_{L_{t}^{2}L_{x}^{6}}\lesssim \eta(1+N^{2s_c-1}|I|)^{\frac{1}{2}}.$$

It then follows from Hölder, Hardy's inequality, Sobolev embedding, Bernstein and (7.10) that

$$\begin{split} I_{1} &\lesssim \left\| \frac{1}{|x|^{\alpha+1}} u_{\leq N} \right\|_{L_{t}^{2(\alpha+1)} L_{x}^{\frac{6(\alpha+1)}{5}}}^{\alpha+1} \|u_{>N}\|_{L_{t}^{2} L_{x}^{6}} \lesssim \|(-\Delta_{\Omega})^{\frac{1}{2(\alpha+1)}} u_{\leq N} \|_{L_{t}^{2(\alpha+1)} L_{x}^{\frac{6(\alpha+1)}{5}}}^{\alpha+1} \|u_{>N}\|_{L_{t}^{2} L_{x}^{6}} \\ &\lesssim \|(-\Delta_{\Omega})^{\frac{3\alpha-2}{4(\alpha+1)}} u_{\leq N} \|_{L_{t}^{2(\alpha+1)} L_{x}^{\frac{6(\alpha+1)}{3\alpha+1}}}^{\alpha+1} \|u_{>N}\|_{L_{t}^{2} L_{x}^{6}} \lesssim N^{1-s_{c}} \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}} u_{\leq N} \|_{L_{t}^{2(\alpha+1)} L_{x}^{\frac{6(\alpha+1)}{3\alpha+1}}}^{\alpha+1} \|u_{>N}\|_{L_{t}^{2} L_{x}^{6}} \\ &\lesssim_{u} \eta N^{1-2s_{c}} (1+N^{2s_{c}-1}|I|). \end{split}$$

For I_2 , we divide it into two cases. If $|u_{\leq N}| \lesssim |u_{>N}|$, then it is very similar to the term I_1 , which we have already treated. Thus, it suffices to consider the case $|u_{\leq N}| \ll |u_{>N}|$, which can be absorbed into the left-hand side of (7.15), if we can show

$$\left\| \frac{1}{|x|} |u_{>N}|^{\alpha+2} \right\|_{L^1_{t,x}} < \infty. \tag{7.21}$$

To prove (7.21), we apply Hardy's inequality and Sobolev embedding to deduce

$$\left\| \frac{1}{|x|} |u_{>N}|^{\alpha+2} \right\|_{L^1_{t,x}} \lesssim \left\| |x|^{-\frac{1}{\alpha+2}} u_{>N} \right\|_{L^{\alpha+2}_{t,x}}^{\alpha+2} \lesssim \left\| (-\Delta_{\Omega})^{\frac{1}{2(\alpha+1)}} u_{>N} \right\|_{L^{\alpha+2}_{t,x}}^{\alpha+2} \lesssim \left\| (-\Delta_{\Omega})^{\frac{3\alpha-2}{4(\alpha+2)}} u_{>N} \right\|_{L^{\alpha+2}_{t}L^{\frac{6(\alpha+2)}{3\alpha+2}}_{x}}^{\alpha+2}.$$

Moreover, by the Bernstein inequality and Lemma 1.7, we have

$$\left\| (-\Delta_{\Omega})^{\frac{3\alpha-2}{4(\alpha+2)}} u_{>N} \right\|_{L_{t}^{\alpha+2} L_{x}^{\frac{6(\alpha+2)}{3\alpha+2}}}^{\alpha+2} \lesssim_{u} N^{1-2s_{c}} \left\| (-\Delta_{\Omega})^{\frac{s_{c}}{2}} u \right\|_{L_{t}^{\alpha+2} L_{x}^{\frac{6(\alpha+2)}{3\alpha+2}}}^{\alpha+2} \lesssim_{u} N^{1-2s_{c}} \left(1+|I|\right) < \infty.$$

Next, we turn to II_1 . By Minkowski's inequality, we have

$$\left\| \frac{1}{|x|} u_{\leq N} [F(u) - F(u_{\leq N})] \right\|_{L^{1}_{t,x}} \lesssim \left\| \frac{1}{|x|} (u_{\leq N})^{\alpha+1} u_{> N} \right\|_{L^{1}_{t,x}} + \left\| \frac{1}{|x|} u_{\leq N} (u_{> N})^{\alpha+1} \right\|_{L^{1}_{t,x}},$$

which is exactly (7.17). Thus II_1 is done. For II_2 , we estimate

$$II_2 \lesssim \|\nabla u_{\leq N}|u_{>N}|^{\alpha}u_{>N}\|_{L^1_tL^1_x} + \|\nabla u_{\leq N}|u_{< N}|^{\alpha}u_{>N}\|_{L^1_tL^1_x}.$$

For the first term, we use Hölder's inequality, Sobolev embedding, Bernstein's estimate, (1.4) and (7.10) to show

$$\begin{split} &\|\nabla u_{\leq N}|u_{>N}|^{\alpha}u_{>N}\|_{L^{1}_{t}L^{1}_{x}} \lesssim \|\nabla u_{\leq N}\|_{L^{\infty}_{t}L^{\frac{3}{1+\varepsilon}}_{x}} \||u_{>N}|^{\alpha}u_{>N}\|_{L^{1}_{t}L^{\frac{3}{2-\varepsilon}}_{x}} \\ &\lesssim N^{1-s_{c}+3(\frac{1}{2}-\frac{1+\varepsilon}{3})} \|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{\leq N}\|_{L^{\infty}_{t}L^{2}_{x}} \|u_{>N}\|_{L^{\infty}_{t}L^{\frac{3\alpha}{2}}_{x}}^{\alpha-1} \|u_{>N}\|_{L^{2}_{t}L^{\frac{6\alpha}{2-\varepsilon\alpha}}_{x}}^{2} \\ &\lesssim \eta N^{1-s_{c}+3(\frac{1}{2}-\frac{1+\varepsilon}{3})} \|(-\Delta_{\Omega})^{\frac{s}{2}}u_{>N}\|_{L^{2}_{t}L^{6}_{x}}^{2} \lesssim \eta N^{1-s_{c}+3(\frac{1}{2}-\frac{1+\varepsilon}{3})} N^{2(s-s_{c})} (1+N^{2s_{c}-1}|I|) \\ &\lesssim \eta N^{1-2s_{c}} (1+N^{2s_{c}-1}|I|), \end{split}$$

where $\varepsilon > 0$ is a fixed sufficiently small constant and $s := \frac{1}{2} - \frac{2-\varepsilon\alpha}{2\alpha}$. For the second term, we use Hölder, Sobolev embedding, Bernstein, (7.10) and (7.12) to estimate

$$\begin{split} &\|\nabla u_{\leq N}|u_{< N}|^{\alpha}u_{> N}\|_{L_{t}^{1}L_{x}^{1}} \\ &\lesssim \|\nabla u_{\leq N}\|_{L_{t}^{\infty}L_{x}^{2}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{< N}\|_{L_{t}^{4}L_{x}^{3}}^{\alpha}\|u_{> N}\|_{L_{t}^{\frac{4-\alpha}{4-\alpha}}L_{x}^{\frac{6}{\alpha-1}}} \\ &\lesssim N^{1-s_{c}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{\leq N}\|_{L_{t}^{\infty}L_{x}^{2}}^{\frac{2}{\alpha+1}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{< N}\|_{L_{t}^{2}L_{x}^{6}}^{\frac{4-\alpha}{\alpha-1}}\|u_{> N}\|_{L_{t}^{2}L_{x}^{6}}^{\frac{2}{\alpha-1}}\|u_{> N}\|_{L_{t}^{\infty}L_{x}^{2}}^{\frac{2}{\alpha-1}} \\ &\lesssim N^{1-s_{c}}\eta^{\alpha}(1+N^{2s_{c}-1}|I|)^{\frac{\alpha}{4}}N^{-s_{c}}(1+N^{2s_{c}-1}|I|)^{\frac{4-\alpha}{4}}N^{-s_{c}(\frac{\alpha}{2}-1)} \\ &\lesssim \eta N^{1-2s_{c}}(1+N^{2s_{c}-1}|I|). \end{split}$$

Thus II_2 is acceptable.

Finally, we turn to III. By Hardy's inequality and Theorem 2.6,

$$(7.19) \lesssim \|u_{>N}\|_{L_{t}^{2}L_{x}^{6}} \left\| \frac{1}{|x|} P_{\leq N}^{\Omega}(F(u)) \right\|_{L_{t}^{2}L_{x}^{6/5}} + \|u_{>N}\|_{L_{t}^{2}L_{x}^{6}} \|\nabla P_{\leq N}^{\Omega}(F(u))\|_{L_{t}^{2}L_{x}^{6/5}}$$
$$\lesssim \|u_{>N}\|_{L_{t}^{2}L_{x}^{6}} \|(-\Delta_{\Omega})^{\frac{1}{2}} P_{\leq N}^{\Omega}(F(u))\|_{L_{t}^{2}L_{x}^{6/5}}.$$

Thus, by (7.10) it remains to prove

$$\|(-\Delta_{\Omega})^{\frac{1}{2}} P_{\leq N}^{\Omega}(F(u))\|_{L_{*}^{2} L_{x}^{6/5}} \lesssim_{u} \eta N^{1-s_{c}} (1+N^{2s_{c}-1}|I|)^{1/2}.$$

To this end, we use Hölder's inequality, Bernstein's estimate, the fractional chain rule (2.1), (1.3), (7.13), and (7.14) to estimate

$$\begin{split} &\|(-\Delta_{\Omega})^{\frac{1}{2}}P_{\leq N}^{\Omega}(F(u))\|_{L_{t}^{2}L_{x}^{6/5}} \\ &\lesssim N\|F(u) - F(u_{\leq N/\eta^{2}})\|_{L_{t}^{2}L_{x}^{6/5}} + N^{1-s_{c}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}F(u_{\leq N/\eta^{2}})\|_{L_{t}^{2}L_{x}^{6/5}} \\ &\lesssim N\|u\|_{L_{t}^{\infty}L_{x}^{3\alpha/2}}^{\alpha}\|u_{>N/\eta^{2}}\|_{L_{t}^{2}L_{x}^{6}} + N^{1-s_{c}}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u\|_{L_{t}^{\infty}L_{x}^{2}}^{\alpha}\|(-\Delta_{\Omega})^{\frac{s_{c}}{2}}u_{\leq N/\eta^{2}}\|_{L_{t}^{2}L_{x}^{6}} \\ &\lesssim u \, \eta N^{1-s_{c}}(1+N^{2s_{c}-1}|I|)^{1/2}. \end{split}$$

$$(7.22)$$

This completes the proof of Proposition 7.3.

7.3. Rule out the minimal conuterexample. In this section, we preclude the existence of the almost periodic solutions to (1.1) with $1/2 < s_c < 1$. We will rely on the frequency-localized Lin-Strauss Morawetz inequality established in Subsection 7.2.

Theorem 7.5. There are no almost periodic solutions to (1.1) as in Theorem 1.8 with $1/2 < s_c < 1$.

Proof. Suppose u were such a solution. Let $\eta > 0$ and let $I \subset [0, \infty)$ be a compact time interval. By Proposition 7.3, we find that for N sufficiently small, we have

$$\int_{I} \int_{\Omega} \frac{|u_{>N}(t,x)|^{\alpha+2}}{|x|} dx dt \lesssim_{u} \eta(N^{1-2s_{c}} + |I|).$$
 (7.23)

Next, we prove the lower bound for the LHS(7.23). As the orbit $\{u(t): t \in \mathbb{R}\}$ is precompact in $\dot{H}_D^{s_c}(\Omega)$ and $\dot{H}_D^{s_c}(\Omega) \hookrightarrow L^{\frac{3\alpha}{2}}(\Omega)$, we see that for any $\varepsilon > 0$, there exists $c(\varepsilon) > 0$ such that

$$\int_{\Omega} |u_{< c(\varepsilon)}(t,x)|^{\frac{3\alpha}{2}} dx < \varepsilon \quad \text{uniformly for} \quad t \in \mathbb{R}.$$

Combining this with (1.5), we obtain that for N sufficiently small,

$$\int_{\Omega \cap \{|x| < R\}} |u_{>N}(t,x)|^{\frac{3\alpha}{2}} dx \gtrsim 1 \quad \text{uniformly for} \quad t \in \mathbb{R}.$$

Moreover, by Hölder, we have that for N sufficiently small,

$$1 \lesssim \int_{\Omega \cap \{|x| \leq R\}} |u_{>N}(t,x)|^{\frac{3\alpha}{2}} dx \lesssim \left(\int_{\Omega \cap \{|x| \leq R\}} |u_{>N}(t,x)|^{\alpha+2} dx \right)^{\frac{3\alpha}{2(\alpha+2)}} R^{\frac{3(4-\alpha)}{2(\alpha+2)}}.$$

Therefore, we obtain, for N sufficiently small,

$$\int_{I} \int_{\Omega} \frac{|u_{>N}(t,x)|^{\alpha+2}}{|x|} dx dt \ge \frac{1}{R} \int_{I} \int_{\Omega \cap \{x: |x| < R\}} |u_{>N}(t,x)|^{\alpha+2} dx dt \gtrsim R^{-\frac{4}{\alpha}} |I|.$$
 (7.24)

Combining (7.23) and (7.24), choosing $\eta = \eta(R)$ sufficiently small, we deduce $|I| \lesssim_u N^{1-2s_c}$ uniformly in I. We obtain a contradiction by taking |I| sufficiently large. This finishes the proof of Theorem 7.5. \square

Remark 7.6. Finally, we discuss the possibility of extending Theorem 1.2 to the case $0 < s_c < \frac{1}{2}$. In this case, we also want to employ a frequency-localized Lin-Strauss Morawetz inequality to preclude almost periodic solutions. To achieve this, we truncate the high frequencies of the solution and work with $u_{< N}$ for some N > 0. Similar to Lemma 7.4, we first establish a long-time Strichartz estimate and then prove (see e.g. [47, Lemma 5.7]):

$$\|(-\Delta_{\Omega})^{\frac{s}{2}} u_{\leq N}\|_{L_{t}^{2} L_{x}^{6}(I \times \Omega)} \lesssim N^{s-s_{c}} (1 + N^{2s_{c}-1} |I|)^{\frac{1}{2}}, \quad for \quad s > \frac{1}{2}.$$
 (7.25)

However, when using the Morawetz inequality to exclude almost periodic solutions, what we actually require is the estimate $||\nabla|^s u_{\leq N}||_{L^2_t L^6_x(I \times \Omega)}$, $s > \frac{1}{2}$. Notably, since $s > \frac{1}{2}$, p = 6 does not satisfy the index relationships in Theorem 2.6, we cannot derive the required estimate from (7.25). Therefore, in the case $0 < s_c < \frac{1}{2}$, we still lack the necessary estimates to exclude almost periodic solutions.

References

- [1] R. Anton, Global existence for defocusing cubic NLS and Gross-Pitaevskii equations in three-dimensional exterior domains, J. Math. Pures Appl. (9) 89 (4) (2008), 335-354.
- [2] M. D. Blair, H. F. Smith, C. D. Sogge, Strichartz estimates and the nonlinear Schrödinger equation on manifolds with boundary, Math. Ann. 354 (4) (2012), 1397-1430.
- [3] J. Bourgain, Global wellposedness of defocusing critical nonlinear Schrödinger equation in the radial case, J. Amer. Math. Soc. 12 (1999), 145-171.
- [4] H. Brézis, E. Lieb, A relation between pointwise convergence of functions and convergence of functionals, Proc. Amer. Math. Soc. 88 (1983), no. 3, 486–490.
- [5] N. Burq, P. Gérard, N. Tzvetkov, On nonlinear Schrödinger equations in exterior domains, Ann. Inst. Henri Poincaré, Anal. Non Linéaire 21 (3) (2004), 295-318.
- [6] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, Global existence and scattering for rough solutions of a nonlinear Schrödinger equation on ℝ³, Comm. Pure Appl. Math. 57 (2004), 987–1014.
- [7] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in \mathbb{R}^3 , Ann. of Math. 167 (2008), 767–865.
- [8] B. Dodson, Global well-posedness and scattering for the defocusing, L²-critical nonlinear Schrödinger equation when d≥ 3, J. Amer. Math. Soc. 25 (2012), 429–463.
- [9] B. Dodson, Global well-posedness and scattering for the mass critical nonlinear Schrödinger equation with mass below the mass of the ground state, Adv. Math. 285 (2015), 1589–1618.
- [10] B. Dodson, Global well-posedness and scattering for the defocusing L²-critical nonlinear Schrödinger equation when d = 2, Duke Math. J. 165 (2016), 3435–3516.
- [11] B. Dodson, Global well-posedness and scattering for the defocusing L²-critical nonlinear Schrödinger equation when d = 1, Am. J. Math. 138 (2016), 531–569.
- [12] B. Dodson, Global well-posedness and scattering for the focusing, cubic Schrödinger equation in dimension d=4, Ann. Sci. Ec. Norm. Supér. (4), **52** (2019), no. 1, 139–180.
- [13] B. Dodson, C. X. Miao, J. Murphy, J. Zheng, The defocusing quintic NLS in four space dimensions, Ann. Inst. H. Poincaré Anal. Non Linéaire 34 (2017), 759–787.
- [14] B. Dodson, J. Murphy, A new proof of scattering below the ground state for the non-radial focusing NLS. Math. Res. Lett., 25 (2018), 1805-1825.
- [15] T. Duyckaerts, O. Landoulsi, S. Roudenko, Threshold solutions in the focusing 3D cubic NLS equation outside a strictly convex obstacle, Journal of Functional Analysis, 282 (2022), 109326.
- [16] C. Gao, C. Miao, J. Yang, The Interctitical Defocusing Nonlinear Schrödinger Equations with Radial Initial Data in Dimensions Four and Higher, Anal. Theory Appl. 35 (2019), 205–234.
- [17] C. Gao, Z. Zhao, On scattering for the defocusing high dimensional inter-critical NLS, J. Differential Equations 267 (2019), 6198–6215.
- [18] M. Grillakis, On nonlinear Schrödinger equations, Comm. Part. Diff. Eqs. 25 (2000), 1827-1844.
- [19] A. Hassell, A. Sikora, Riesz transforms in one dimension, Indiana Univ. Math. J. 58 (2009), no. 2, 823–852.
- [20] O. Ivanovici, Precised smoothing effect in the exterior of balls, Asymptot. Anal. 53 (4) (2007), 189-208.
- [21] O. Ivanovici, On the Schrödinger equation outside strictly convex obstacles, Anal. PDE 3 (3) (2010), 261–293.
- [22] O. Ivanovici, G. Lebeau, Dispersion for the wave and the Schrödinger equations outside strictly convex obstacles and counterexamples, Comptes Rendus Math. 355 (2017), 774–779.
- [23] O. Ivanovici, F. Planchon, On the energy critical Schrödinger equation in 3D non-trapping domains, Ann. Inst. Henri Poincaré, Anal. Non Linéaire 27 (5) (2010), 1153–1177.
- [24] C. E. Kenig, F. Merle, Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case, Invent. Math. 166 (2006), no. 3, 645-675.
- [25] C. E. Kenig, F. Merle, Scattering for $\dot{H}^{1/2}$ bounded solutions to the cubic, defocusing NLS in 3 dimensions, Trans. Am. Math. Soc. **362** (2010), 1937–1962.
- [26] S. Keraani, On the defect of compactness for the Strichartz estimates of the Schrödinger equations, J. Differential Equations 175 (2001), no. 2, 353–392.
- [27] S. Keraani, On the blow up phenomenon of the critical nonlinear Schr ödinger equation, J. Funct. Anal., 235 (2006), 171-192.
- [28] R. Killip, T. Tao, M. Visan, The cubic nonlinear Schrödinger equation in two dimensions with radial data, J. Eur. Math. Soc. 11 (2009), 1203–1258.
- [30] R. Killip, M. Visan, The focusing energy-critical nonlinear Schrödinger equation in dimensions five and higher, Amer. J. Math. 132 (2010), 361–424.
- [31] R. Killip, M. Visan, The radial defocusing energy-supercritical nonlinear wave equation in all space dimensions, Proc. Amer. Math. Soc. 139 (2011), 1805–1817.
- [32] R. Killip, M. Visan, Global well-posedness and scattering for the defocusing quintic NLS in three dimensions, Anal. PDE 5 (2012), 855–885.
- [33] R. Killip, M. Visan, Nonlinear Schrödinger equations at critical regularity, in: Evolution Equations, in: Clay Math. Proc., vol. 17, Amer. Math. Soc., Providence, RI, 2013, pp. 325–437.
- [34] R. Killip, M. Visan, X. Zhang, The mass-critical nonlinear Schrödinger equation with radial data in dimensions three and higher, Anal. PDE 1 (2008), 229–266.

- [35] R. Killip, M. Visan, X. Zhang, Quintic NLS in the exterior of a strictly convex obstacle, Amer. J. Math. 138 (2016), no. 5, 1193–1346.
- [36] R. Killip, M. Visan, X. Zhang, The focusing cubic NLS on exterior domains in three dimensions, Appl. Math. Res. Express. AMRX 2016, no. 1, 146–180.
- [37] R. Killip, M. Visan, X. Zhang, Riesz transforms outside a convex obstacle, Int. Math. Res. Not. IMRN 2016, no. 19, 5875-5921.
- [38] M. Keel, T. Tao, Endpoint Strichartz estimates, Amer. J. Math. 120 (1998), no. 5, 955–980.
- [39] O. Landoulsi, Construction of a solitary wave solution of the nonlinear focusing Schrödinger equation outside a strictly convex obstacle in the L²-supercritical case, Discrete Contin. Dyn. Syst., Ser. A 41 (2) (2021), 701–746.
- [40] D. Li, H. Smith, X. Zhang, Global well-posedness and scattering for defocusing energy-critical NLS in the exterior of balls with radial data, Math. Res. Lett. 19 (2012), no. 1, 213–232.
- [41] D. Li, G. Xu, X. Zhang, On the dispersive estimate for the Dirichlet Schrödinger propagator and applications to energy critical NLS, Canad. J. Math. 66 (2014), no. 5, 1110–1142.
- [42] J. Li, K. Li, The Defocusing Energy-supercritical Nonlinear Schrödinger Equation in High Dimensions, SIAM J. Math. Anal., 54 (2022), no.3, 3253-3274.
- [43] C. Lu, J. Zheng, The radial defocusing energy-supercritical NLS in dimension four, J. Differ. Equ. 262 (2017), 4390–4414.
- [44] C. Miao, J. Murphy, J. Zheng, The defocusing energy-supercritical NLS in four space dimensions, J. Funct. Anal. 267 (2014), 1662–1724.
- [45] J. Murphy, Inter-critical NLS: Critical H's-bounds imply scattering, SIAM J. Math. Anal. 46 (2014), 939–997.
- [46] J. Murphy, The defocusing $\dot{H}^{1/2}$ -critical NLS in high dimensions, Discrete Contin. Dyn. Syst. Ser. A **34** (2014), 733–748.
- [47] J. Murphy, The radial defocusing nonlinear Schrödinger equation in three space dimensions, Comm. Partial Differential Equations 40 (2015), 265–308.
- [48] F. Planchon, L. Vega, Bilinear virial identities and applications, Ann. Sci. Éc. Norm. Supér. (4) 42 (2009), 261–290.
- [49] E. Ryckman, M. Visan, Global well-posedness and scattering for the defocusing energy critical nonlinear Schrödinger equation in ℝ¹⁺⁴, Am. J. Math. 129 (2007), 1–60.
- [50] S. Shao, Maximizers for the Strichartz and the Sobolev-Strichartz inequalities for the Schrödinger equation, Electron. J. Differential Equations 2009, No. 3, 13 pp.
- [51] R. S. Strichartz, Multipliers on fractional Sobolev spaces, J. Math. Mech., 16, (1967), 1031–1060.
- [52] T. Tao, Global well-posedness and scattering for the higher-dimensional energy-critical non-linear Schrödinger equation for radial data, New York J. Math. 11 (2005), 57–80.
- [53] T. Tao, M. Visan, Stability of energy-critical nonlinear Schrödinger equations in high dimensions, Electron. J. Differ. Equ. 118 (2005), 1–28.
- [54] T. Tao, M. Visan, X. Zhang, Minimal-mass blowup solutions of the mass-critical NLS, Forum Math. 20 (2008), 881– 010
- [55] T. Tao, M. Visan, X. Zhang, Global well-posedness and scattering for the mass-critical nonlinear Schrödinger equation for radial data in high dimensions, Duke Math. J. 140 (2007), 165–202.
- [56] M. Visan, The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions, Duke Math. J. 138 (2007), 281–374.
- [57] M. Visan, Global well-posedness and scattering for the defocusing cubic nonlinear Schrödinger equation in four dimensions, Int. Math. Res. Not. IMRN 2012 (2012), 1037–1067.
- [58] J. Xie, D. Fang, Global well-posedness and scattering for the defocusing H

 s-critical NLS, Chin. Ann. Math. 34B (2013), 801–842.
- [59] C. Xu, T. Zhao, J. Zheng, Scattering for 3D cubic focusing NLS on the domain outside a convex obstacle revisited. Acta. Math. Sin.-English Ser. 38 (2022), 1054-1068.
- $[60] \ \text{K. Yang, } \textit{The focusing NLS on exterior domains in three dimensions.} \ \text{Commun. Pure Appl. Anal.} \ \textbf{6} \ (2017), \ 2269-2297.$
- [61] X. Yu, Global well-posedness and scattering for the defocusing $\dot{H}^{1/2}$ -critical nonlinear Schrödinger equation in \mathbb{R}^2 , Anal. PDE 14 (2021), 1037–1067.
- [62] Q. S. Zhang, The global behavior of heat kernels in exterior domains, J. Funct. Anal. 200 (2003), no. 1, 160–176.
- [63] T. F. Zhao, The Defocusing Energy-supercritical NLS in Higher Dimensions, Acta Mathematica Sinica, English Series, 33 (2017), 911–925.

School of Mathematics, Hangzhou Normal University, Hangzhou, 311121, China Email address: liuxuan95@hznu.edu.cn

Yilin Song

The Graduate School of China Academy of Engineering Physics, Beijing 100088, P. R. China $Email\ address$: songyilin21@gscaep.ac.cn

JIQIANG ZHENG

Institute of Applied Physics and Computational Mathematics, Beijing, 100088, China.

NATIONAL KEY LABORATORY OF COMPUTATIONAL PHYSICS, BEIJING 100088, CHINA

 $Email\ address: {\tt zheng_jiqiang@iapcm.ac.cn, zhengjiqiang@gmail.com}$