

Integral Motivic Cohomology of BSO_4

Alexander Port

December 19, 2024

Abstract

Motivic cohomology is a powerful tool in algebraic geometry with associated realization maps giving important information about the relations between cohomological invariants of schemes and their classifying spaces. The problem of computing general cohomological invariants of these classifying spaces is ongoing. Most relevant to this paper is (1) Totaro's construction of the Chow ring of a classifying space in general and his use of this to study symmetric groups in [Tot99], (2) Guillot's similar examination for the Lie groups G_2 and $Spin(7)$ in [Gui07], (3) Field's computation of the Chow ring of $BSO(2n, \mathbb{C})$ in [Fie12], and (4) Yagita's work on the \mathbb{Z}_2 -motivic cohomology of BSO_4 and BG_2 in [Yag10].

The work presented in this paper covers the computation of the motivic cohomology of BSO_4 with integral coefficients. The primary approach draws on methods laid out by Guillot and Yagita ([Gui07], [Yag10]). These results lay the groundwork for future work, most immediately the analogous computation for BG_2 ([Por21]).

1 Notation and Conventions

- Maps:

- The short exact sequence $0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}_2 \rightarrow 0$ induces the long exact sequence

$$\dots \rightarrow H^*(BSO_4; \mathbb{Z}) \xrightarrow{\times 2} H^*(BSO_4; \mathbb{Z}) \xrightarrow{\mu} H^*(BSO_4; \mathbb{Z}_2) \xrightarrow{\tilde{\beta}} H^{*+1}(BSO_4; \mathbb{Z}) \rightarrow \dots$$

- The short exact sequence $0 \rightarrow \mathbb{Z}_2 \xrightarrow{\times 2} \mathbb{Z}_4 \rightarrow \mathbb{Z}_2 \rightarrow 0$ induces a similar long exact sequence

$$\dots \rightarrow H^*(BSO_4; \mathbb{Z}_2) \xrightarrow{\times 2} H^*(BSO_4; \mathbb{Z}_4) \rightarrow H^*(BSO_4; \mathbb{Z}_2) \xrightarrow{\beta} H^{*+1}(BSO_4; \mathbb{Z}_2) \rightarrow \dots$$

- Note that μ is the map induced by reducing coefficients mod 2, $\tilde{\beta}$ is the *integral Bockstein homomorphism*, and β is the *Bockstein homomorphism* (where $\beta = \mu \circ \tilde{\beta}$).

- The Milnor operations $Q_i : H^{*,*'}(X; \mathbb{Z}_p) \rightarrow H^{*+2p^i-1, *'+p^i-1}(X; \mathbb{Z}_p)$ are defined inductively as $Q_0 = \beta$ and $Q_{i+1} = [Q_i, P^{p^i}]$ where $P^k : H^{*,*'}(X; \mathbb{Z}_p) \rightarrow H^{*+2k(p-1), *'+k(p-1)}(X; \mathbb{Z}_p)$ is a reduced p^{th} -power operation.

- Generators:

- In the context of ordinary cohomology and Chow rings, the subscript of a generator is usually its degree. For motivic cohomology, a subscript of n means the degree is $(2n, n)$. The one exception to this is $H^{*,*'}((\mathbb{G}_m)^n; \mathbb{Z}) = H^{*,*'}(\text{Spec}(\mathbb{C}); \mathbb{Z})[t_1, \dots, t_n]$ where $\deg(t_i) = (2, 1)$ for all i ([Fie12], [Gui07]).

- Let $d_m \in H^{2m}(BSO_n; \mathbb{Z})$ be the m^{th} Chern class of the standard representation of SO_n ([Gui07]).
- Let ζ be the universal n -plane bundle over BSO_n , let $w_m \in H^m(BSO_n; \mathbb{Z}_2)$ be the m^{th} Stiefel-Whitney class of ζ , and let $p_m \in H^{4m}(BSO_n; \mathbb{Z})$ be the m^{th} Pontryagin class of the complexification $\zeta \otimes \mathbb{C}$ ([Bro82]).
- τ refers to a generator of $H^{*,*'}(Spec(\mathbb{C}); \mathbb{Z}_p)$ with $deg(\tau) = (0, 1)$ and corresponds to a choice of generator of $H^{0,1}(Spec(\mathbb{C}); \mathbb{Z}_p) \cong \mathbb{Z}_p$.

- Shorthand:

- We use the shorthand $Q(n)$ to refer to the exterior algebra $\Lambda_{\mathbb{Z}_2}(Q_0, \dots, Q_n)$ ([Yag10]).
- If $I = \{i_1, \dots, i_l\}$ we denote the products $w(2I) = w_{2i_1} \dots w_{2i_l}$ and $p(I) = p_{i_1} \dots p_{i_l}$.
- If S is a set and R is a commutative ring then we use $R\{S\}$ to denote the free R -module with S as its basis.
- If A is an abelian group then ${}_l A = \{a \in A : la = 0_A\}$ is the l -torsion subgroup of A .

2 Known Computations

2.1 Classical BSO_4 Computations

The computational strategy used for computing $H^{*,*'}(BSO_4; \mathbb{Z})$ is to take lifts of classical generators (against the realization map $t : H^{*,*'}(BSO_4; A) \rightarrow H^*(BSO_4; A)$ for $A = \mathbb{Z}$ or \mathbb{Z}_p) and to search for additional generators using the universal coefficient theorem. As such, the following represents a summary of the needed classical results regarding the cohomology and Chow ring of BSO_4 :

Theorem 2.1. ([Bro82] 1.3) *The mod 2 cohomology ring of BSO_n is given by*

$$H^*(BSO_n; \mathbb{Z}_2) = \mathbb{Z}_2[w_2, w_3, \dots, w_{2n}]$$

Theorem 2.2. ([Bro82] 1.5) *The integral cohomology ring of BSO_n is $H^*(BSO_n; \mathbb{Z}) = \bar{R}_n / \bar{I}_n$ where*

$$\bar{R}_n = \mathbb{Z}[p_1, \dots, p_{\lfloor \frac{n-1}{2} \rfloor}, X_n, \tilde{\beta}w(2I) : 1 \leq i_1 < \dots < i_l \leq \lfloor \frac{n-1}{2} \rfloor]$$

and \bar{I}_n is the ideal generated by the following relations: (1) $2\tilde{\beta}w(I) = 0$, (2) $\tilde{\beta}w(2I)\tilde{\beta}w(2J)$ is equal to

$$\sum_{k \in I} \tilde{\beta}w_{2k} p((I - \{k\}) \cap J) \tilde{\beta}w(2((I - \{k\}) \cup J - (I - \{k\}) \cap J))$$

(3) if $n = 2k + 1$ is odd then $X_n = \tilde{\beta}w_{2k}$, and (4) if $n = 2k$ is even then $X_n^2 = p_k$. Furthermore $\mu(p_i) = w_{2i}^2$ for all i .

Theorem 2.3. ([Fie12] 1) *The Chow ring of BSO_n is given by*

$$CH^*(BSO_n) = \mathbb{Z}[d_2, \dots, d_{2n}, y_n] / (2d_{\text{odd}}, y_n d_{\text{odd}}, y_n^2 + (-1)^n 2^{2n-2} d_{2n})$$

where y_n is the class which maps to 2^{n-1} times the Euler class of ζ_n under the cycle class map.

Note that when $n = 2k$ is even, the above class X_n is the Euler class $e = e(\zeta \otimes \mathbb{C})$ as it satisfies the needed property of $e \smile e = p_k$. As such, we will simply replace X_n with $\sqrt{p_k}$ moving forward.

Corollary 2.4. ([Por21] 5.3.4) *Taking $n = 4$ with the above theorems yields*

$$\begin{aligned} H^*(BSO_4; \mathbb{Z}_2) &= \mathbb{Z}_2[w_2, w_3, w_4] \\ H^*(BSO_4; \mathbb{Z}) &= \mathbb{Z}[\tilde{\beta}w_2, p_1, \sqrt{p_2}]/(2\tilde{\beta}w_2) \\ CH^*(BSO_4) &= \mathbb{Z}[d_2, d_3, d_4, y_2]/(2d_3, y_2d_3, y_2^2 - 4d_4) \end{aligned} \quad (1)$$

2.2 Re-framing Yagita's $H^{*,*'}(BSO_4; \mathbb{Z}_2)$

With this in place we can now move into the motivic setting. Recalling that $H^{2n,n}(X; A) = CH^n(X) \otimes A$, we can take d_2, d_3, d_4 to be motivic cohomology classes where

$$\deg(d_2) = (4, 2) \quad \deg(d_3) = (6, 3) \quad \deg(d_4) = (8, 4) \quad (2)$$

However, to avoid confusion we denote the motivic Stiefel-Whitney classes as w_i^{alg} (i.e. the ‘‘algebraic’’ lifts of their classical counterparts). As seen by Corollary 2.2 of [HN18]:

$$\deg(w_{2i}^{alg}) = (2i, i+1) \quad \deg(w_{2i+1}^{alg}) = (2i+1, i+1) \quad (3)$$

With these generators and conventions in place we can consider Yagita's original statement of the mod 2 motivic cohomology of BSO_4 :

Theorem 2.5. ([Yag10] 9.4) *The motivic cohomology $H^{*,*'}(BSO_4; \mathbb{Z}_2)$ is given by*

$$\begin{aligned} H^{*,*'}(BSO_4; \mathbb{Z}_2) &= \mathbb{Z}_2[\mu(d_2), \mu(d_4)]\{y_{0,2}\} \oplus \mathbb{Z}_2[\tau, \mu(d_2)] \otimes (\mathbb{Z}_2[\mu(d_4)]\{1\} \oplus \mathbb{Z}_2[\mu(d_3)] \otimes Q(1)\{w_2^{alg}\} \\ &\quad \oplus \mathbb{Z}_2[\mu(d_4)] \otimes (\mathbb{Z}_2[\mu(d_3)]Q(2) - \mathbb{Z}_2\{1\})\{a\}) \end{aligned}$$

where (1) a is a virtual element corresponding to element that in $H^0(BSO_4; H^3(B\mathbb{Z}_2; \mathbb{Z}_2))$ and is taken to have degree $(3, 3)$, and (2) the degrees of the generators are given by $\deg(\tau) = (0, 1)$, $\deg(y_{0,2}) = (4, 2)$, and $\deg(\mu(d_i)) = (2i, i)$. Furthermore, realizations involving a are given as follows:

$$t_2(\mu(d_2)a) = w_2w_3w_4 \quad t_2(Q_0a) = w_4 \quad t_2(Q_1a) = w_2w_4 \quad t_2(Q_2a) = w_2w_4^2$$

The following theorem will re-frame Yagita's above generators in a different context which will be useful in the following section.

Theorem 2.6. ([Por21] 7.3.3) *The motivic cohomology ring $H^{*,*'}(BSO_4; \mathbb{Z}_2)$ is given by*

$$\mathbb{Z}_2[\tau, y_{0,2}, w_2^{alg}, w_3^{alg}, w_4^{alg}, \tau^{-2}(w_2^{alg})^2, \tau^{-1}(w_3^{alg})^2, \tau^{-2}(w_4^{alg})^2, \tau^{-1}w_2^{alg}w_3^{alg}, \tau^{-1}w_2^{alg}w_4^{alg}, \tau^{-1}w_3^{alg}w_4^{alg}]/I$$

where degrees are given by

$$\deg(\tau) = (0, 1) \quad \deg(y_{0,2}) = (4, 2) \quad \deg(w_2^{alg}) = (2, 2) \quad \deg(w_3^{alg}) = (3, 2) \quad \deg(w_4^{alg}) = (4, 3)$$

and I is the ideal generated by the following relations:

$$\begin{aligned} \tau y_{0,2} &= 0 & y_{0,2}^2 &= 0 & y_{0,2}w_2^{alg} &= 0 & y_{0,2}w_3^{alg} &= 0 & y_{0,2}w_4^{alg} &= 0 \\ y_{0,2}\tau^{-1}(w_3^{alg})^2 &= 0 & y_{0,2}\tau^{-1}w_2^{alg}w_3^{alg} &= 0 & y_{0,2}\tau^{-1}w_2^{alg}w_4^{alg} &= 0 & y_{0,2}\tau^{-1}w_3^{alg}w_4^{alg} &= 0 \end{aligned}$$

Finally, Harada and Nakada expand on Yagita's result and show that $\text{Ker}(t_2) = \mathbb{Z}_2[\mu(d_2), \mu(d_4)]\{y_{0,2}\}$ ([HN18] 0.1); translating this into the new notation we get that

$$\text{Ker}(t_2) = \mathbb{Z}_2[\tau^{-2}(w_2^{alg})^2, \tau^{-2}(w_4^{alg})^2]\{y_{0,2}\} \quad (4)$$

3 Integral Motivic Cohomology of BSO_4

The goal here is to compute the integral motivic cohomology of BSO_4 using the universal coefficient theorem (UCT) as presented on pg. 27 of "Lecture Notes on Motivic Cohomology" by Carlo Mazza, Vladimir Voevodsky and Charles Weibel as well its classical analog ([MVW06], see [nLab] universal coefficient theorem for classical version used here). Here we obtain a short exact sequence on cohomology:

$$0 \rightarrow H^*(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2 \xrightarrow{\mu_C} H^*(BSO_4; \mathbb{Z}_2) \xrightarrow{\tilde{\beta}_C} {}_2H^{*+1}(BSO_4; \mathbb{Z}) \rightarrow 0$$

These maps μ_C and $\tilde{\beta}_C$ are closely related to μ and $\tilde{\beta}$ in that $\mu(x) = \mu_C(x \otimes 1)$ for all $x \in H^*(BSO_4; \mathbb{Z})$ and $\tilde{\beta}_C$ is the same as β with the only different being the codomain. With a slight abuse of notation with μ , β and $\tilde{\beta}$ being reused, the motivic version of these constructions can be summarized as the following ([MVW06] pg. 27):

$$\begin{array}{ccccccc} & H^{*,*'}(BSO_4; \mathbb{Z}) & & & H^{*+1,*'}(BSO_4; \mathbb{Z}) & & \\ & \downarrow \otimes 1 & \searrow \mu & & \nearrow \tilde{\beta} & & \\ 0 & \longrightarrow & H^{*,*'}(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2 & \xrightarrow{\mu_M} & H^{*,*'}(BSO_4; \mathbb{Z}_2) & \xrightarrow{\tilde{\beta}_M} & {}_2H^{*+1,*'}(BSO_4; \mathbb{Z}) \longrightarrow 0 \\ & & & & \searrow \beta & & \uparrow \mu \\ & & & & & & H^{*+1,*'}(BSO_4; \mathbb{Z}_2) \end{array}$$

Note that the subscript C 's and M 's are meant to distinguish between "classical" and "motivic". All together, these sequences form the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^{*,*'}(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2 & \xrightarrow{\mu_M} & H^{*,*'}(BSO_4; \mathbb{Z}_2) & \xrightarrow{\tilde{\beta}_M} & {}_2H^{*+1,*'}(BSO_4; \mathbb{Z}) \longrightarrow 0 \\ & & \downarrow t_1 & \dashrightarrow & \downarrow t_2 & \dashrightarrow & \downarrow t_3 \\ 0 & \longrightarrow & H^*(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2 & \xrightarrow{\mu_C} & H^*(BSO_4; \mathbb{Z}_2) & \xrightarrow{\tilde{\beta}_C} & {}_2H^{*+1}(BSO_4; \mathbb{Z}) \longrightarrow 0 \end{array}$$

The maps t_i are variants of the realization map $t^{m,n} : H^{m,n}(BSO_4; \mathbb{Z}) \rightarrow H^m(BSO_4; \mathbb{Z})$ where $t = \bigoplus_{m,n} t^{m,n}$.

There is an important conclusion to draw here that is relevant in the proof of the main theorem of this section (see 3.1 below). The map μ_C is injective, so $\varphi \in \text{Ker}(\mu_C \circ t_1)$ iff $\varphi \in \text{Ker}(t_1)$. Similarly, μ_M is injective so $\varphi \in \text{Ker}(t_2 \circ \mu_M)$ iff $\mu_M(\varphi) \in \text{Ker}(t_2)$. But $\mu_C \circ t_1 = t_2 \circ \mu_M$ by commutativity and therefore

$$\varphi \in \text{Ker}(t_1) \iff \mu_M(\varphi) \in \text{Ker}(t_2) \quad (5)$$

Theorem 3.1. *Let the d_i 's be Chern classes of the standard representation of SO_4 ([Gui07]). Let the w_{2i} 's be the even Stiefel-Whitney classes of the inclusion $G_2 \subset SO_7$ where $\text{deg}(w_{2i}) = (2i, i+1)$ ([Yag10]). Let*

the class y_2 be a lift of twice the classical Euler class of universal 4-plane bundle over BSO_4 ([Fie12]). Take $\tilde{\beta} : H^{*,*'}(BSO_4; \mathbb{Z}_2) \rightarrow H^{*+1,*'}(BSO_4; \mathbb{Z})$ to be the integral Bockstein homomorphism. Finally, let τ be a generator of $H^{0,1}(Spec(\mathbb{C}); \mathbb{Z}_2) \cong \mathbb{Z}_2$. Recalling that $H^*(BSO_4; \mathbb{Z}) = \mathbb{Z}[p_1, \sqrt{p_2}, \tilde{\beta}w_2]/(2\tilde{\beta}w_2)$ for SO_4 defined over \mathbb{C} ([Bro82]), the integral motivic cohomology ring of BSO_4 is

$$H^{*,*'}(BSO_4; \mathbb{Z}) = H^{*,*'}(Spec(\mathbb{C}); \mathbb{Z})[y_2, d_2, d_3, d_4, \tilde{\beta}\tau^k w_2, \tilde{\beta}\tau^{k-1} w_2 w_4 : k \geq 0]/I$$

where I is the ideal generated by the following relations:

$$\begin{aligned} 2d_3 &= 0 & y_2 d_3 &= 0 & y_2^2 - 4d_4 &= 0 \\ 2\tilde{\beta}\tau^k w_2 &= 0 & 2\tilde{\beta}\tau^{k-1} w_2 w_4 &= 0 & y_2 \tilde{\beta}\tau^k w_2 &= 0 & y_2 \tilde{\beta}\tau^{k-1} w_2 w_4 &= 0 \\ (\tilde{\beta}\tau^{k_1} w_2)(\tilde{\beta}\tau^{k_2} w_2) &= (\tilde{\beta}\tau^{k_3} w_2)(\tilde{\beta}\tau^{k_4} w_2) & \iff & k_1 + k_2 &= k_3 + k_4 \\ (\tilde{\beta}\tau^{k_1-1} w_2 w_4)(\tilde{\beta}\tau^{k_2-1} w_2 w_4) &= (\tilde{\beta}\tau^{k_3-1} w_2 w_4)(\tilde{\beta}\tau^{k_4-1} w_2 w_4) & \iff & k_1 + k_2 &= k_3 + k_4 \\ (\tilde{\beta}\tau^{k_1} w_2)(\tilde{\beta}\tau^{k_2-1} w_2 w_4) &= (\tilde{\beta}\tau^{k_3} w_2)(\tilde{\beta}\tau^{k_4-1} w_2 w_4) & \iff & k_1 + k_2 &= k_3 + k_4 \\ (\tilde{\beta}\tau^{k-1} w_2 w_4)^2 &= d_4(\tilde{\beta}\tau^k w_2)^2 & (\tilde{\beta}\tau^k w_2)^3 &= d_3 \tilde{\beta}\tau^{3k+1} w_2 \end{aligned}$$

The degrees of the generators are as follows:

$$\begin{aligned} \deg(d_i) &= (2i, i) & \deg(y_2) &= (4, 2) \\ \deg(\tilde{\beta}\tau^k w_2) &= (3, 2+k) & \deg(\tilde{\beta}\tau^{k-1} w_2 w_4) &= (7, 4+k) \end{aligned}$$

Furthermore, the images of the generators under the realization map $t : H^{*,*'}(BSO_4; \mathbb{Z}) \rightarrow H^*(BSO_4; \mathbb{Z})$ are $t(d_2) = -p_1$, $t(d_3) = (\tilde{\beta}w_2)^2$, $t(d_4) = p_2$, $t(y_2) = 2\sqrt{p_2}$, $t(\tilde{\beta}\tau^k w_2) = \tilde{\beta}w_2$, and $t(\tilde{\beta}\tau^{k-1} w_2 w_4) = \sqrt{p_2} \tilde{\beta}w_2$.

Proof: Note that the integral lifts are sometimes in bold font for emphasis in diagrams.

The below consideration of only 2-torsion is sufficient by a transfer argument. We know that $SL_2 \times SL_2$ is a 2-fold cover of SO_4 and that $H^{*,*'}(BSL_2; \mathbb{Z}) = H^{*,*'}(Spec(\mathbb{C}); \mathbb{Z})[c_2]$ does not contain torsion outside of $H^{*,*'}(Spec(\mathbb{C}); \mathbb{Z})$. Therefore, the only torsion that can appear in the generators of $H^{*,*'}(BSO_4; \mathbb{Z})$ over $H^{*,*'}(Spec(\mathbb{C}); \mathbb{Z})$ is 2-torsion.

This method relies on the fact that $H^*(BSO_4; \mathbb{Z})$, $H^*(BSO_4; \mathbb{Z}_2)$, $H^{*,*'}(BSO_4; \mathbb{Z}_2)$, and the realization map $t_2 : H^{*,*'}(BSO_4; \mathbb{Z}_2) \rightarrow H^*(BSO_4; \mathbb{Z}_2)$. Together with the universal coefficient theorem and $CH^*(BSO_4)$, these allow us to compute $H^{*,*'}(BSO_4; \mathbb{Z})$ in its entirety.

$$\begin{array}{ccccccccc} 0 & \longrightarrow & H^{*,*'}(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2 & \xrightarrow{\mu_M} & H^{*,*'}(BSO_4; \mathbb{Z}_2) & \xrightarrow{\tilde{\beta}_M} & {}_2H^{*+1,*'}(BSO_4; \mathbb{Z}) & \longrightarrow & 0 \\ & & \downarrow t_1 & \searrow & \downarrow t_2 & \searrow & \downarrow t_3 & & \\ 0 & \longrightarrow & H^*(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2 & \xrightarrow{\mu_C} & H^*(BSO_4; \mathbb{Z}_2) & \xrightarrow{\tilde{\beta}_C} & {}_2H^{*+1}(BSO_4; \mathbb{Z}) & \longrightarrow & 0 \end{array}$$

To fill in the lower left entry, tensoring $H^*(BSO_4; \mathbb{Z})$ with \mathbb{Z}_2 gives $H^*(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2 \cong \mathbb{Z}_2[p_1, \sqrt{p_2}, \tilde{\beta}w_2]$.

Lemma 3.2. In the exact sequence $0 \rightarrow H^*(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2 \xrightarrow{\mu_C} H^*(BSO_4; \mathbb{Z}_2) \xrightarrow{\tilde{\beta}_C} {}_2H^{*+1}(BSO_4; \mathbb{Z}) \rightarrow 0$:

$$\begin{aligned} \mu_C(p_1^{k_1}(\sqrt{p_2})^{k_2}(\tilde{\beta}w_2)^{k_3} \otimes 1) &= w_2^{2k_1} w_3^{k_3} w_4^{k_2} & \tilde{\beta}_C(w_2^{2k_1} w_3^{k_3} w_4^{k_2}) &= 0 \\ \tilde{\beta}_C(w_2^{2k_1+1} w_3^{k_3} w_4^{k_2}) &= p_1^{k_1}(\sqrt{p_2})^{k_2}(\tilde{\beta}w_2)^{k_3+1} \end{aligned}$$

Proof of Lemma: Brown's Theorem 1.5 includes the statement that $\mu(p_i) = w_2^2$ for all i . This shows that

$\mu_C(p_1 \otimes 1) = w_2^2$ and $\mu_C(\sqrt{p_2} \otimes 1) = w_4$. Injectivity of μ_C along with the facts that $H^3(BSO_4; \mathbb{Z}) = \mathbb{Z}_2\{\tilde{\beta}w_2\}$ and $H^3(BSO_4; \mathbb{Z}_2) = \mathbb{Z}_2\{w_3\}$ gives that $\mu_C(\tilde{\beta}w_2 \otimes 1) = w_3$. The first line follows from exactness.

The approach for the second line is different because $\tilde{\beta}_C$ is a module homomorphism, not a ring homomorphism. Clearly $\tilde{\beta}_C(w_2) = \tilde{\beta}w_2$, as indicated by Brown's notation. More generally, we know that $Sq^1 = \beta = \mu \circ \tilde{\beta}_C$ is simply the Bockstein homomorphism and the first Steenrod squaring operation. On Stiefel-Whitney classes, β is characterized by the fact that $\beta(w_{2i}) = w_{2i+1}$ and $\beta(w_{2i+1}) = 0$ ([HN18]); this along with induction on the Cartan formula and the fact that $w_5 = 0$ gives the result:

$$\begin{aligned}\beta(w_2^{2k_1+1}) &= w_2\beta(w_2^{2k_1}) + \beta(w_2)w_2^{2k_1} = w_2^{2k_1}w_3 = \mu(p_1^{k_1}\tilde{\beta}w_2) \\ \beta(w_2^{2k_1+1}w_3^{k_3}) &= w_2^{2k_1+1}\beta(w_3^{k_3}) + \beta(w_2^{2k_1+1})w_3^{k_3} = w_2^{2k_1}w_3^{k_3+1} = \mu(p_1^{k_1}(\tilde{\beta}w_2)^{k_3+1}) \\ \beta(w_2^{2k_1+1}w_3^{k_3}w_4^{k_2}) &= w_2^{2k_1+1}w_3^{k_3}\beta(w_4^{k_2}) + \beta(w_2^{2k_1+1}w_3^{k_3})w_4^{k_2} = w_2^{2k_1}w_3^{k_3+1}w_4^{k_2} = \mu(p_1^{k_1}(\sqrt{p_2})^{k_2}(\tilde{\beta}w_2)^{k_3+1})\end{aligned}$$

□

The next step is to consider the elements of $CH^*(BSO_4)$ as lifts of classes in $H^*(BSO_4; \mathbb{Z})$; in particular, we need to contextualize d_2, d_3, d_4 and y_2 . It well-known that Pontryagin classes have lifts to the motivic setting via $t(d_{2i}) = (-1)^i p_i$ ([Bro82]), so the following diagrams are known from [Yag10]:

$$\begin{array}{ccc} d_2 & \xrightarrow{\mu} & \tau^{-2}(w_2^{alg})^2 \\ \downarrow t & & \downarrow t_2 \\ -p_1 & \xrightarrow{\mu} & w_2^2 \end{array} \qquad \begin{array}{ccc} d_3 & \xrightarrow{\mu} & \tau^{-1}(w_3^{alg})^2 \\ \downarrow t & & \downarrow t_2 \\ (\tilde{\beta}w_2)^2 & \xrightarrow{\mu} & w_3^2 \end{array}$$

$$\begin{array}{ccc} d_4 & \xrightarrow{\mu} & \tau^{-2}(w_4^{alg})^2 \\ \downarrow t & & \downarrow t_2 \\ p_2 & \xrightarrow{\mu} & w_4^2 \end{array} \qquad \begin{array}{ccc} y_2 & \xrightarrow{\mu} & y_{0,2} \\ \downarrow t & & \downarrow t_2 \\ 2\sqrt{p_2} & \xrightarrow{\mu} & 0 \end{array}$$

On the other hand, there is no lift e^{alg} of the Euler class $e = \sqrt{p_2}$. We know from Field's Theorem 1 that her class $y_2 \in CH^*(BSO_4)$ is mapped to $2\sqrt{p_2}$ under the realization map; clearly this implies that twice the Euler class does have a lift in $H^{*,*'}(BSO_4; \mathbb{Z})$. We run into trouble if we instead attempt to lift only $\sqrt{p_2}$, as seen in detail below:

Lemma 3.3. *There is no lift of $\sqrt{p_2} \in H^*(BSO_4; \mathbb{Z})$ to $H^{*,*'}(BSO_4; \mathbb{Z})$.*

Proof of Lemma: Suppose such of lift $e^{alg} \in H^{4,n}(BSO_4; \mathbb{Z})$ of $\sqrt{p_2}$ exists for some $n \geq 2$.

- Clearly $(\sqrt{p_2})^2 - p_2 = 0$; thus $((e^{alg})^2 - d_4) \otimes 1 \in Ker(t_1)$ and so $\mu_M(((e^{alg})^2 - d_4) \otimes 1) \in Ker(t_2)$.
- Simplifying gives $\mu_M(((e^{alg})^2 - d_4) \otimes 1) = \mu_M(e^{alg} \otimes 1)^2 - \mu(d_4)$.
- We know from Yagita's work that $H^{8,4}(BSO_4; \mathbb{Z}_2) = \mathbb{Z}_2\{\mu(d_2)^2, y_{0,2}\mu(d_2), \mu(d_4)\}$ and if $k \geq 5$ then $H^{8,k}(BSO_4; \mathbb{Z}_2) = \mathbb{Z}_2\{\tau^{k-5}w_4^{alg}\mu(d_2), \tau^{n-5}w_2^{alg}\mu(d_3), \tau^{n-4}\mu(d_2)^2, \tau^{n-4}\mu(d_4)\}$.
- Computations show that t_2 sends the following elements of $H^{*,*'}(BSO_4; \mathbb{Z}_2)$ to their counterparts in $H^*(BSO_4; \mathbb{Z}_2)$: $\tau \mapsto 1$, $\mu(d_2)^2 \mapsto w_2^4$, $y_{0,2}\mu(d_2) \mapsto 0$, $\mu(d_4) \mapsto w_4^2$, $w_4^{alg}\mu(d_2) \mapsto w_2^2w_4$, and

$w_2^{alg} \mu(d_3) \mapsto w_2 w_3^2$. Thus, $\mu_M(e^{alg} \otimes 1)^2 \in H^{8,2n}(BSO_4; \mathbb{Z}_2)$ must be equal to $\tau^{2n-4} \mu(d_4)$ so that $t_2(\mu_M(e^{alg} \otimes 1)^2) = w_4^2$.

- This implies that if e^{alg} exists then it must satisfy the property that $\mu_M(e^{alg} \otimes 1)$ is equal to $\tau^{n_3-2} x$ for some $x \in H^{4,2}(BSO_4; \mathbb{Z}_2)$ such that $x^2 = \mu(d_4)$. The problem is that no such element x exists because $H^{4,2}(BSO_4; \mathbb{Z}_2) = \mathbb{Z}_2\{\mu(d_2), y_{0,2}\}$ and neither of these two generators satisfy this property.
- Thus, the lift e^{alg} cannot exist.

Visually, the above argument can be summarized as follows:

$$\begin{array}{ccc}
& & w_4^{alg} \xrightarrow{\tilde{\beta}_M} 0 \\
& & \downarrow t_2 \\
\sqrt{p_2} \mapsto & \xrightarrow{\mu} & w_4 \xrightarrow{\tilde{\beta}_C} 0 \\
& & \downarrow t
\end{array}$$

□

The key to the remainder of this proof is that whenever ${}_2H^{m+1}(BSO_4; \mathbb{Z})$ is non-zero there will be elements of $H^{m+1}(BSO_4; \mathbb{Z})$ that can be lifted to non-zero elements of $H^{m+1,n}(BSO_4; \mathbb{Z})$ for some n . To see this, note that we can rewrite $H^*(BSO_4; \mathbb{Z})$ in the following way:

$$H^*(BSO_4; \mathbb{Z}) = \mathbb{Z}\{p_1^i(\sqrt{p_2})^j : i, j \in \mathbb{N}\} \oplus \mathbb{Z}_2\{p_1^i(\sqrt{p_2})^j(\tilde{\beta}w_2)^k : i, j \in \mathbb{N}, k \in \mathbb{Z}_{>0}\}$$

So ${}_2H^{m+1}(BSO_4; \mathbb{Z}) = \mathbb{Z}_2\{p_1^i(\sqrt{p_2})^j(\tilde{\beta}w_2)^k : i, j \in \mathbb{N}, k \in \mathbb{Z}_{>0}, 4i + 4j + 3k = m + 1\}$ is non-trivial precisely when $[\frac{1}{4}(m+2), \frac{1}{3}(m+1)] \cap \mathbb{Z} \neq \emptyset$; a quick calculation shows that this is equivalent to having $m+1 \in \{3, 6, 7\} \cup \mathbb{Z}_{\geq 9}$. Conversely, we now know that ${}_2H^{m+1}(BSO_4; \mathbb{Z}) = \{0\}$ only if $m+1 \in \{1, 2, 4, 5, 8\}$.

Consider the values of $0 \rightarrow H^m(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2 \xrightarrow{\mu_C} H^m(BSO_4; \mathbb{Z}_2) \xrightarrow{\tilde{\beta}_C} {}_2H^{m+1}(BSO_4; \mathbb{Z}) \rightarrow 0$ for the first several values of m , as given by the Lemma 3.2:

$$\begin{array}{l}
m = 0 : \quad 0 \longrightarrow \mathbb{Z}_2\{1 \otimes 1\} \longrightarrow \mathbb{Z}_2\{1\} \longrightarrow 0 \longrightarrow 0 \\
m = 1 : \quad 0 \longrightarrow 0 \longrightarrow 0 \longrightarrow 0 \longrightarrow 0 \\
m = 2 : \quad 0 \longrightarrow 0 \longrightarrow \mathbb{Z}_2\{w_2\} \longrightarrow \mathbb{Z}_2\{\tilde{\beta}w_2\} \longrightarrow 0 \\
m = 3 : \quad 0 \longrightarrow \mathbb{Z}_2\{\tilde{\beta}w_2 \otimes 1\} \longrightarrow \mathbb{Z}_2\{w_3\} \longrightarrow 0 \longrightarrow 0 \\
m = 4 : \quad 0 \longrightarrow \mathbb{Z}_2\{p_1 \otimes 1, \sqrt{p_2} \otimes 1\} \longrightarrow \mathbb{Z}_2\{w_2^2, w_4\} \longrightarrow 0 \longrightarrow 0 \\
m = 5 : \quad 0 \longrightarrow 0 \longrightarrow \mathbb{Z}_2\{w_2 w_3\} \longrightarrow \mathbb{Z}_2\{(\tilde{\beta}w_2)^2\} \longrightarrow 0 \\
m = 6 : \quad 0 \longrightarrow \mathbb{Z}_2\{(\tilde{\beta}w_2)^2 \otimes 1\} \longrightarrow \mathbb{Z}_2\{w_2^3, w_2 w_4, w_3^2\} \longrightarrow \mathbb{Z}_2\{p_1 \tilde{\beta}w_2, \sqrt{p_2} \tilde{\beta}w_2\} \longrightarrow 0
\end{array}$$

Keeping in mind that $\tilde{\beta}_C \circ t_2 = t_3 \circ \tilde{\beta}_M$:

- The $m = 2$ case shows that $\tilde{\beta}_C(w_2) = \tilde{\beta}w_2$. Because $t_2(\tau^k w_2^{alg}) = w_2$ for any choice of k , we have that $t_3 \circ \tilde{\beta}_M(\tau^k w_2^{alg}) = \tilde{\beta}w_2$. This means that $\tilde{\beta}_M(\tau^k w_2^{alg})$ is non-zero in ${}_2H^{3,2+i}(BSO_4; \mathbb{Z})$, i.e. $\tilde{\beta}_M(\tau^i w_2^{alg}) \neq 0$ but $2\tilde{\beta}_M(\tau^i w_2^{alg}) = 0$. In other words, $\tilde{\beta}_M(\tau^k w_2^{alg})$ is a lift of $\tilde{\beta}w_2$ for all k .

- Note that this demonstrates the distinction between the algebraic lifts w_i^{alg} and the topological lifts w_i^{top} in $H^{*,*'}(BSO_4; \mathbb{Z}_2)$. Both give rise to lifts in $H^{*,*'}(BSO_4; \mathbb{Z})$ and, in fact, so do infinitely many other classes:

$$\begin{array}{ccc}
w_2^{alg} & \xrightarrow{\tilde{\beta}_M} & \tilde{\beta}_M(w_2^{alg}) & \xrightarrow{\mu_M} & w_3^{alg} \\
\downarrow t_2 & & \downarrow t_3 & & \downarrow t_2 \\
w_2 & \xrightarrow{\tilde{\beta}_C} & \tilde{\beta}_C w_2 & \xrightarrow{\mu_C} & w_3
\end{array}
\qquad
\begin{array}{ccc}
\tau w_2^{alg} & \xrightarrow{\tilde{\beta}_M} & \tilde{\beta}_M(\tau w_2^{alg}) & \xrightarrow{\mu_M} & w_3^{top} \\
\downarrow t_2 & & \downarrow t_3 & & \downarrow t_2 \\
w_2 & \xrightarrow{\tilde{\beta}_C} & \tilde{\beta}_C w_2 & \xrightarrow{\mu_C} & w_3
\end{array}$$

- Similarly, the $m = 5$ case shows that $\tilde{\beta}_M(\tau^k w_2^{alg} w_3^{alg})$ is a lift of $(\tilde{\beta}_C w_2)^2$ for all k .
- On the other hand, the Cartan formula gives that $\beta(\tau^k w_3^{alg}) = 0$; it is well-known that $\beta(\tau^k) = 0$ because no integral analog of τ exists ([MVW06] 4.2) and we also know that $\beta(w_3^{alg}) = 0$ ([HN18]). This shows that $\beta(\tau^k w_3^{alg})$ is not a non-trivial lift of any element of $H^4(BSO_4; \mathbb{Z})$.
- The same reasoning applied to the $m = 4$ case shows that $\tilde{\beta}_M(\tau^k w_2^2)$ and $\tilde{\beta}_M(\tau^k w_4)$ are both 0 and thus are not considered lifts either.

More generally, all 2-torsion elements in $H^*(BSO_4; \mathbb{Z})$ will have lifts to 2-torsion elements in $H^{*,*'}(BSO_4; \mathbb{Z})$. As above, these lifts will be of the form $\tilde{\beta}(\tau^k x)$ for classes $x \in H^{*,*'}(BSO_4; \mathbb{Z}_2)$ such that $\tilde{\beta}_C \circ t_2(x)$ is 2-torsion.

Ultimately the result of the above argument is that the integral motivic cohomology of BSO_4 has infinitely many generators. For example, each $\tilde{\beta}_M(\tau^k w_2^{alg}) \in H^{3,2+k}(BSO_4; \mathbb{Z})$ is 2-torsion lift satisfying the property $\tau^k w_3^{alg} = \mu \circ \tilde{\beta}_M(\tau^k w_2^{alg})$. It is key to determine the relations between these $\tilde{\beta}_M(\tau^k w_2^{alg})$ for different k . One immediate relation mod 2 is given by

$$k \geq k' \implies \mu \circ \tilde{\beta}_M(\tau^k w_2^{alg}) = \tau^{k-k'} \mu \circ \tilde{\beta}_M(\tau^{k'} w_2^{alg}) \quad (6)$$

It would be convenient if there were a similar relation on the integral side of things, i.e. $k \geq k'$ implies $\tilde{\beta}_M(\tau^k w_2) = T^{k-k'} \tilde{\beta}_M(\tau^{k'} w_2)$ for some $T \in H^{0,1}(BSO_4; \mathbb{Z})$. Unfortunately that class would have to live in $H^{0,1}(Spec(\mathbb{C}); \mathbb{Z})$ and it is known that no such class exists ([MVW06] 4.2).

Clearly the integral case is more complicated than the mod 2 case due to the fact that no integral analog of τ exists. Before proceeding any further, the following lemma will help to greatly simplify arguments moving forward:

Lemma 3.4. *Generators of 2-torsion in $H^{*,*'}(BSO_4; \mathbb{Z})$ that are not 2-divisible are equal iff their mod 2 reductions in $H^{*,*'}(BSO_4; \mathbb{Z}_2)$ are equal.*

Proof of Lemma: Suppose $x, y \in H^{*,*'}(BSO_4; \mathbb{Z})$ are 2-torsion and are not 2-divisible. Clearly if $x = y$ then $\mu(x) = \mu(y)$, so it remains to go in the other direction. The elements x and y are not 2-divisible so $x \otimes 1$ and $y \otimes 1$ are both non-zero in $H^{*,*'}(BSO_4; \mathbb{Z}) \otimes \mathbb{Z}_2$.

Assume that $\mu(x) = \mu(y)$. This implies that $\mu_M(x \otimes 1) = \mu_M(y \otimes 1)$. Injectivity of μ_M gives that $x \otimes 1 = y \otimes 1$. Thus, some odd multiples of x and y must be equal, i.e. $(2i+1)x = (2j+1)y$ for some i and j . But x and y are 2-torsion and therefore $2ix$ and $2jy$ are zero; we conclude that $x = y$.

□

We have shown that there are infinitely many lifts of $\tilde{\beta}w_2$; as stated above, in principal any 2-torsion element of $H^*(BSO_4; \mathbb{Z})$ will have a lift in a similar form. Applying the same construction as done for $\tilde{\beta}w_2$ to other elements of Yagita's presentation yields all remaining lifts (where here we make a slight simplification to the notation of the lifts):

$$\begin{array}{ccccc}
\tau^k w_2^{alg} & \xrightarrow{\tilde{\beta}_M} & \tilde{\beta} \tau^k w_2 & \xrightarrow{\mu} & \tau^k w_3^{alg} \\
\downarrow t_2 & & \downarrow t & & \downarrow t_2 \\
w_2 & \xrightarrow{\tilde{\beta}_C} & \tilde{\beta} w_2 & \xrightarrow{\mu} & w_3 \\
\\
\tau^{k-1} w_2^{alg} w_3^{alg} & \xrightarrow{\tilde{\beta}_M} & \tilde{\beta} \tau^{k-1} w_2 w_3 & \xrightarrow{\mu} & \tau^{k-1} (w_3^{alg})^2 \\
\downarrow t_2 & & \downarrow t & & \downarrow t_2 \\
w_2 w_3 & \xrightarrow{\tilde{\beta}_C} & (\tilde{\beta} w_2)^2 & \xrightarrow{\mu} & w_3^2 \\
\\
\tau^{k-1} w_2^{alg} w_4^{alg} & \xrightarrow{\tilde{\beta}_M} & \tilde{\beta} \tau^{k-1} w_2 w_4 & \xrightarrow{\mu} & \tau^{k-1} w_3^{alg} w_4^{alg} \\
\downarrow t_2 & & \downarrow t & & \downarrow t_2 \\
w_2 w_4 & \xrightarrow{\tilde{\beta}_C} & \sqrt{p_2} \tilde{\beta} w_2 & \xrightarrow{\mu} & w_3 w_4 \\
\\
\tau^{k-1} w_2^{alg} w_3^{alg} w_4^{alg} & \xrightarrow{\tilde{\beta}_M} & \tilde{\beta} \tau^{k-1} w_2 w_3 w_4 & \xrightarrow{\mu} & \tau^{k-1} (w_3^{alg})^2 w_4^{alg} \\
\downarrow t_2 & & \downarrow t & & \downarrow t_2 \\
w_2 w_3 w_4 & \xrightarrow{\tilde{\beta}_C} & \sqrt{p_2} (\tilde{\beta} w_2)^2 & \xrightarrow{\mu} & w_3^2 w_4 \\
\\
\tau^{k-2} w_2^{alg} (w_4^{alg})^2 & \xrightarrow{\tilde{\beta}_M} & \tilde{\beta} \tau^{k-2} w_2 w_4^2 & \xrightarrow{\mu} & \tau^{k-2} w_3^{alg} (w_4^{alg})^2 \\
\downarrow t_2 & & \downarrow t & & \downarrow t_2 \\
w_2 w_4^2 & \xrightarrow{\tilde{\beta}_C} & p_2 \tilde{\beta} w_2 & \xrightarrow{\mu} & w_3 w_4^2 \\
\\
\tau^{k-3} w_2^{alg} w_3^{alg} (w_4^{alg})^2 & \xrightarrow{\tilde{\beta}_M} & \tilde{\beta} \tau^{k-3} w_2 w_3 w_4^2 & \xrightarrow{\mu} & \tau^{k-3} (w_3^{alg})^2 (w_4^{alg})^2 \\
\downarrow t_2 & & \downarrow t & & \downarrow t_2 \\
w_2 w_3 w_4^2 & \xrightarrow{\tilde{\beta}_C} & p_2 (\tilde{\beta} w_2)^2 & \xrightarrow{\mu} & w_3^2 w_4^2
\end{array}$$

One important conclusion to draw from these diagrams is that the generator d_3 is of this form:

$$d_3 = \tilde{\beta} \tau^{-1} w_2 w_3 \tag{7}$$

There are two other types lift that we have seen, namely those involving d_2 , d_4 , and y_2 . However, these types do not involve τ because multiple lifts can only exist for 2-torsion elements in this construction:

$$\begin{array}{ccc}
d_2^{k_1} d_4^{k_2} & \xrightarrow{\mu} & \mu(d_2)^{k_1} \mu(d_4)^{k_2} \\
\downarrow t & & \downarrow t_2 \\
(-p_1)^{k_1} (\sqrt{p_2})^{2k_2} & \xrightarrow{\mu} & w_2^{2k_1} w_4^{2k_2}
\end{array}$$

$$\begin{array}{ccc}
y_2 d_2^{k_1} d_4^{k_2} & \xrightarrow{\mu} & y_{0,2} \mu(d_2)^{k_1} \mu(d_4)^{k_2} \\
\downarrow t & & \downarrow t_2 \\
2(-p_1)^{k_1} (\sqrt{p_2})^{2k_2+1} & \xrightarrow{\mu} & 0
\end{array}$$

If we take “simple” to mean a product of only generators with no coefficients, we see that almost all simple elements of $H^*(BSO_4; \mathbb{Z})$ have lifts. There is only one type of simple element that remains to be considered, namely the case of the power of $\sqrt{p_2}$ being odd and there being no $\tilde{\beta}w_2$. The Euler class falls into this category and so the following lemma is a natural continuation of the logic from Lemma 3.3:

Lemma 3.5. *An element $p_1^{j_1} (\sqrt{p_2})^{j_2} (\tilde{\beta}w_2)^{j_3} \in H^*(BSO_4; \mathbb{Z})$ lifts $H^{*,*'}(BSO_4; \mathbb{Z})$ iff j_2 is even or $j_3 > 0$.*

Proof of Lemma: We’ve seen that if j_3 is greater than zero then there is a lift. In addition, we know that $(-p_1)^{k_1} (\sqrt{p_2})^{2k_2}$ lifts to $d_2^{k_1} d_4^{k_2}$. Thus, it suffices to show that if $j_2 = 2k_2 + 1$ and $k_3 = 0$ then there is no lift:

- Suppose a lift of $l^{alg} \in H^{4k_1+8k_2+4, n}(BSO_4; \mathbb{Z})$ of $p_1^{k_1} (\sqrt{p_2})^{2k_2+1}$ exists.
- Clearly $(p_1^{k_1} (\sqrt{p_2})^{2k_2+1}) = p_1^{2k_1} (\sqrt{p_2})^{4k_2+2}$ because p_1 and p_2 are both even degree.
- But $p_1^{2k_1} (\sqrt{p_2})^{4k_2+2} = p_1^{2k_1} p_2^{2k_2+1}$ lifts to $d_2^{2k_1} d_4^{2k_2+1}$ and so $((l^{alg})^2 - d_2^{2k_1} d_4^{2k_2+1}) \otimes 1$ is in $\text{Ker}(t_1)$.
- Thus, $\mu_M(((l^{alg})^2 - d_2^{2k_1} d_4^{2k_2+1}) \otimes 1) = \mu_M(l^{alg} \otimes 1)^2 - \mu(d_2)^{2k_1} \mu(d_4)^{2k_2+1}$ is in $\text{Ker}(t_2)$.
- This gives $\mu_M(l^{alg} \otimes 1)^2 = \tau^{2n-4k_1-8k_2-4} \mu(d_2)^{2k_1} \mu(d_4)^{2k_2+1}$ and there must be some $x \in H^{4,2}(BSO_4; \mathbb{Z}_2)$ such that $x^2 = \mu(d_4)$ where $\mu_M(l^{alg} \otimes 1)$ is equal to $\tau^{n-2k_1-4k_2-2} \mu(d_2)^{k_1} \mu(d_4)^{k_2} x$.
- No such x exists in $H^{4,2}(BSO_4; \mathbb{Z}_2) = \mathbb{Z}_2\{\mu(d_2), y_{0,2}\}$ and so l^{alg} does not exist.

□

The first objective is to remove any generators that can be rewritten completely in terms of other generators. All the generators from the first six families in Table 1 are 2-torsion but not 2-divisible and so Lemma 3.4 applies for finding integral relations. The driving idea for these relations comes from [Por21] 7.3.1; in particular, if two classes in $H^{*,*'}(BSO_4; \mathbb{Z}_2) - \text{Ker}(t_2)$ have the same realizations and degrees then they must be equal. This fact allows us to rewrite families 2, 4, 5 and 6:

$$\begin{aligned}
\tilde{\beta}\tau^{-1}w_2w_3 &= d_3 & \tilde{\beta}\tau^{k_1+k_2}w_2w_3 &= (\tilde{\beta}\tau^{k_1}w_2)(\tilde{\beta}\tau^{k_2}w_2) \\
\tilde{\beta}\tau^{k_1+k_2-1}w_2w_3w_4 &= (\tilde{\beta}\tau^{k_1}w_2)(\tilde{\beta}\tau^{k_2-1}w_2w_4) \\
\tilde{\beta}\tau^{k_1-2}w_2w_4^2 &= d_4\tilde{\beta}\tau^{k_1}w_2 \\
\tilde{\beta}\tau^{-3}w_2w_3w_4^2 &= d_3d_4 & \tilde{\beta}\tau^{k_1+k_2-2}w_2w_3w_4^2 &= d_4(\tilde{\beta}\tau^{k_1}w_2)(\tilde{\beta}\tau^{k_2}w_2)
\end{aligned} \tag{8}$$

As a more detailed example, we can consider why $\tilde{\beta}\tau^{-1}w_2w_3$ is equal to d_3 . We know that $t(\tilde{\beta}\tau^{-1}w_2w_3)$ and $t(d_3)$ are both equal to $(\tilde{\beta}w_2)^{-1}$ and both have degree $(6, 3)$. Our choice of notation gives $\mu(d_3) = \tau^{-1}(w_3^{alg})^2$, so Lemma 3.4 and [Por21] 7.3.1 give that $\mu(\tilde{\beta}\tau^{-1}w_2w_3) = \tau^{-1}(w_3^{alg})^2$ and therefore $\tilde{\beta}\tau^{-1}w_2w_3 = d_3$. In short, the only integral generators that appear are $y_2, d_2, d_3, d_4, \tilde{\beta}\tau^k w_2$ and $\tilde{\beta}\tau^{k-1} w_2 w_4$.

It remains to determine how these six types of generators interact with each other. First, recall the original relations from Field:

$$2d_3 = 0 \quad y_2d_3 = 0 \quad y_2^2 - 4d_4 = 0 \tag{9}$$

Family	Form	Coefficient	Power of p_1	Power of $\sqrt{p_2}$	Power of $\tilde{\beta}w_2$	Form of Lifts
1	$p_1^{k_2}(\tilde{\beta}w_2)^{2k_3+1}$	1	any	0	odd	$d_2^{k_2}d_3^{k_3}\tilde{\beta}\tau^{k_1}w_2$
2	$p_1^{k_2}(\tilde{\beta}w_2)^{2k_3+2}$	1	any	0	even > 0	$d_2^{k_2}d_3^{k_3}\tilde{\beta}\tau^{k_1-1}w_2w_3$
3	$p_1^{k_2}(\sqrt{p_2})^{2k_4+1}(\tilde{\beta}w_2)^{2k_3+1}$	1	any	odd	odd	$d_2^{k_2}d_3^{k_3}d_4^{k_4}\tilde{\beta}\tau^{k_1-1}w_2w_4$
4	$p_1^{k_2}(\sqrt{p_2})^{2k_4+1}(\tilde{\beta}w_2)^{2k_3+2}$	1	any	odd	even > 0	$d_2^{k_2}d_3^{k_3}d_4^{k_4}\tilde{\beta}\tau^{k_1-1}w_2w_3w_4$
5	$p_1^{k_2}(\sqrt{p_2})^{2k_4+2}(\tilde{\beta}w_2)^{2k_3+1}$	1	any	even > 0	odd	$d_2^{k_2}d_3^{k_3}d_4^{k_4}\tilde{\beta}\tau^{k_1-2}w_2w_4^2$
6	$p_1^{k_2}(\sqrt{p_2})^{2k_4+2}(\tilde{\beta}w_2)^{2k_3+2}$	1	any	even > 0	even > 0	$d_2^{k_2}d_3^{k_3}d_4^{k_4}\tilde{\beta}\tau^{k_1-3}w_2w_3w_4^2$
7	$\lambda p_1^{k_2}(\sqrt{p_2})^{2k_4}$	any	any	even	0	$\lambda(-d_2)^{k_2}d_4^{k_4}$
8	$2\lambda p_1^{k_2}(\sqrt{p_2})^{2k_4+1}$	even	any	odd	0	$\lambda y_2(-d_2)^{k_2}d_4^{k_4}$
9	$(2\lambda + 1)p_1^{k_1}(\sqrt{p_2})^{2k_2+1}$	odd	any	odd	0	N/A

Table 1: The elements of $H^{*,*'}(BSO_4; \mathbb{Z})$ fall into nine families. Eight of these have lifts as above but the ninth does not.

Next, the commutative diagram construction shows that the new classes are all 2-torsion:

$$2\tilde{\beta}\tau^k w_2 = 0 \quad 2\tilde{\beta}\tau^{k-1} w_2 w_4 = 0 \quad (10)$$

Lemma 3.4 along with $\mu(\tilde{\beta}\tau^k w_2) = \tau^k w_3^{alg}$ and $\mu(\tilde{\beta}\tau^k w_2 w_4) = \tau^k w_3^{alg} w_4^{alg}$ give the following:

$$\begin{aligned} (\tilde{\beta}\tau^{k_1} w_2)(\tilde{\beta}\tau^{k_2} w_2) &= (\tilde{\beta}\tau^{k_3} w_2)(\tilde{\beta}\tau^{k_4} w_2) \iff k_1 + k_2 = k_3 + k_4 \\ (\tilde{\beta}\tau^{k_1-1} w_2 w_4)(\tilde{\beta}\tau^{k_2-1} w_2 w_4) &= (\tilde{\beta}\tau^{k_3-1} w_2 w_4)(\tilde{\beta}\tau^{k_4-1} w_2 w_4) \iff k_1 + k_2 = k_3 + k_4 \\ (\tilde{\beta}\tau^{k_1} w_2)(\tilde{\beta}\tau^{k_2-1} w_2 w_4) &= (\tilde{\beta}\tau^{k_3} w_2)(\tilde{\beta}\tau^{k_4-1} w_2 w_4) \iff k_1 + k_2 = k_3 + k_4 \end{aligned} \quad (11)$$

Finally, d_2 does not interact with any other generators and interactions of new generators with y_2 are similar to those between y_2 and d_3 :

$$y_2 \tilde{\beta}\tau^k w_2 = 0 \quad y_2 \tilde{\beta}\tau^{k-1} w_2 w_4 = 0 \quad (12)$$

These final equations are shown in the same way one shows that $y_2 d_3 = 0$. Yagita's presentation shows that $y_{0,2} \in H^{*,*'}(BSO_4; \mathbb{Z}_2)$ only appears as part of $\mathbb{Z}_2[\mu(d_2), \mu(d_4)]\{y_{0,2}\}$. This implies that $y_{0,2}$ multiplied by any generator other than $\mu(d_2)$ or $\mu(d_4)$ is 0 ([HN18] Theorem 0.1). Keeping in mind that $2\tilde{\beta}w_2 = 0$, $t(y_2) = 2\sqrt{p_2}$, $t(d_3) = (\tilde{\beta}w_2)^2$, $t(\tilde{\beta}\tau^k w_2) = \tilde{\beta}w_2$ and $t(\tilde{\beta}\tau^k w_2) = \sqrt{p_2}\tilde{\beta}w_2$:

$$\begin{array}{ccc} \begin{array}{ccc} y_2 d_3 & \xrightarrow{\mu} & y_{0,2} \mu(d_3) \\ \downarrow t & & \downarrow t_2 \\ 0 & \xrightarrow{\mu} & 0 \end{array} & \begin{array}{ccc} y_2 \tilde{\beta}\tau^k w_2 & \xrightarrow{\mu} & \tau^k y_{0,2} w_3^{alg} \\ \downarrow t & & \downarrow t_2 \\ 0 & \xrightarrow{\mu} & 0 \end{array} & \begin{array}{ccc} y_2 \tilde{\beta}\tau^{k-1} w_2 w_4 & \xrightarrow{\mu} & \tau^{k-1} y_{0,2} w_2^{alg} w_4^{alg} \\ \downarrow t & & \downarrow t_2 \\ 0 & \xrightarrow{\mu} & 0 \end{array} \end{array}$$

Thus, the upper right entry of each of these four diagrams must be equal to 0. We can demonstrate that the upper left corners must all be 0 as well using $y_2 d_3$ as an example:

- $\mu_M(y_2 d_3 \otimes 1) = y_{0,2} \mu(d_3)$ and μ_M is injective, so $y_2 d_3 \otimes 1 = 0$
- y_2 and d_3 are both not 2-divisible, so therefore $y_2 d_3 = 0$

The claim that $y_2 \tilde{\beta}\tau^k w_2 = 0$ and $y_2 \tilde{\beta}\tau^{k-1} w_2 w_4 = 0$ follows from the same logic because $\tilde{\beta}\tau^k w_2$ and $\tilde{\beta}\tau^{k-1} w_2 w_4$ are not 2-divisible. □

4 Conclusion

The above represents a portion of the author's thesis work, the full version of which can be found at [Por21]. There are several additional results presented there, most notably the analogous computation for the integral motivic cohomology of BG_2 using the results presented in this paper. A future goal is to also make the BG_2 part of this work a standalone paper.

References

- [Bro82] Brown, Edgar H., Jr.: The cohomology of BSO_n and BO_n with integer coefficients. Proc. Amer. Math. Soc. 85 (1982), no. 2, 283–288.
- [Fie12] Field, R. E.: The Chow ring of the classifying space $BSO(2n, \mathbb{C})$. J. Algebra 350 (2012), 330–339.

- [Gui07] Guillot, Pierre: The Chow rings of G_2 and $Spin(7)$. *J. Reine Angew. Math.* 604 (2007), 137–158.
- [HN18] Harada, Masana; Nakada, Masayuki: The motivic cohomology of BSO_n . *Math. Proc. Cambridge Philos. Soc.* 164 (2018), no. 3, 461–471.
- [MVW06] Mazza, Carlo; Voevodsky, Vladimir; Weibel, Charles: Lecture notes on motivic cohomology. *Clay Mathematics Monographs*, 2. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2006. xiv+216 pp. ISBN: 978-0-8218-3847-1; 0-8218-3847-4.
- [nLab] nLab authors: retrieved January 13, 2021.
- [Por21] Port, Alexander: Motivic Cohomology of BG_2 . USC Digital Library of Thesis Submissions.
- [Tot99] Totaro, Burt: The Chow ring of a classifying space. *Algebraic K-theory (Seattle, WA, 1997)*, 249–281, *Proc. Sympos. Pure Math.*, 67, Amer. Math. Soc., Providence, RI, 1999.
- [Vez00] Vezzosi, Gabriele: On the Chow ring of the classifying stack of $PGL_{3,\mathbb{C}}$. *J. Reine Angew. Math.* 523 (2000), 1–54.
- [Yag10] Yagita, Nobuaki: Coniveau filtration of cohomology of groups. *Proc. Lond. Math. Soc.* (3) 101 (2010), no. 1, 179–206.