Modeling coskewness with zero correlation and correlation with zero coskewness

Carole Bernard, Jinghui Chen[†] and Steven Vanduffel[‡]

December 19, 2024

Abstract

This paper shows that one needs to be careful when making statements on potential links between correlation and coskewness. Specifically, we first show that, on the one hand, it is possible to observe any possible values of coskewness among symmetric random variables but zero pairwise correlations of these variables. On the other hand, it is also possible to have zero coskewness and any level of correlation. Second, we generalize this result to the case of arbitrary marginal distributions showing the absence of a general link between rank correlation and standardized rank coskewness.

Keywords: Coskewness, Correlation, Rank coskewness, Rank correlation, Copula, Marginal distribution.

^{*}Carole Bernard, Department of Accounting, Law and Finance, Grenoble Ecole de Management (GEM) and Department of Economics at Vrije Universiteit Brussel (VUB). (email: carole.bernard@grenoble-em.com).

[†]Corresponding author: Jinghui Chen, Department of Economics at Vrije Universiteit Brussel (VUB). (email: jinghui.chen@vub.be).

[‡]Steven Vanduffel, Department of Economics at Vrije Universiteit Brussel (VUB). (email: steven.vanduffel@vub.be).

1 Introduction

Let $X_i \sim F_i$, i = 1, 2, ..., d be random variables such that μ_i and σ_i are their respective means and standard deviations, and their second moments are finite. One of the essential characteristics of dependency of a random vector $\mathbf{X} = (X_1, X_2, ..., X_d)$ is the kth order standardized central mixed moments

$$\mathbb{E}\left(\left(\frac{X_1-\mu_1}{\sigma_1}\right)^{k_1}\left(\frac{X_2-\mu_2}{\sigma_2}\right)^{k_2}\cdots\left(\frac{X_d-\mu_d}{\sigma_d}\right)^{k_d}\right)$$

where k_i , i = 1, 2, ..., d, are non-negative integers such that $\sum_{i=1}^{d} k_i = k$. Specifically, the Pearson correlation coefficient (Pearson, 1895) is obtained when $k_1 = k_2 = 1$ (d = 2) and coskewness is obtained when $k_1 = k_2 = k_3 = 1$ (d = 3).

The correlation coefficient between X_i and X_j denoted as ρ_{ij} , i, j = 1, 2, ..., d, is given as

$$\rho_{ij} = \frac{\mathbb{E}((X_i - \mu_i)(X_j - \mu_j))}{\sigma_i \sigma_j},$$

and the correlation matrix is a d by d matrix. Jondeau and Rockinger (2006) define the d by d^2 coskewness matrix of a d-dimensional random vector \mathbf{X} , as a matrix that contains all coskewnesses. The coskewness of X_i , X_j and X_k , denoted by $S(X_i, X_j, X_k)$, $i, j, k = 1, 2, \ldots, d$, is given as

$$S(X_i, X_j, X_k) = \frac{\mathbb{E}((X_i - \mu_i)(X_j - \mu_j)(X_k - \mu_k))}{\sigma_i \sigma_j \sigma_k}.$$

The coskewness matrix is denoted by M_d , so that, for example, when d = 3,

where $s_{ijk} = S(X_i, X_j, X_k)$, i, j, k = 1, 2, 3. The coskewness matrix M_d is invariant w.r.t. location and scale parameters, i.e., a linear transformation of X_i , i = 1, 2, ..., d, does not affect M_d . However, the coskewness $S(X_i, X_j, X_k)$ generally depends on the marginal distributions and copula among the three variables; see Bernard et al. (2023).

It is well-known that correlation always takes values in [-1, 1]. However, such affirmation is not true for higher co-moments such as coskewness and cokurtosis. In particular, no universal range of values for coskewness works for all distributions; see Bernard et al. (2023). We thus use the notion of standardized rank coskewness, which is normalized and takes values in [-1, 1].

This paper studies whether a relationship exists between correlation and coskewness. At first glance, it is easy to think that the answer is affirmative because the mathematical for-

mulas of correlation and coskewness share some similarities. Moreover, correlations do not determine the dependence but at least impose some structure. For instance, the maximum and minimum correlation between two random variables are obtained by comonotonic and antimonotonic dependence, respectively. Hence, one could expect a link between the second cross and the third cross moment. For example, Beddock and Karehnke (2020) use a split bivariate normal model to illustrate that the coskewness is monotonic to the correlation; see their Table 3. However, such conclusion heavily depends on the model assumed (here the split bivariate normal), and the remaining of this paper is dedicated to showing that, in general, there is no link between correlation and coskewness and that such conclusions can only be made under specific model assumptions.

The paper is organized as follows. In Section 2, we present counterexamples based on three symmetrically distributed random variables. In Section 3, we generalize the result to the case of random variables with arbitrary marginal distributions. Section 4 provides some elements to justify statements that appear in previous literature on the link between coskewness and tail risk. The last section draws the conclusion.

2 Correlation and coskewness with symmetric marginals

In this section, we aim to show that, in general, there is no link between the coskewness and the correlation coefficient in the case of symmetric distributions. Let F_i , i = 1, 2, 3, be symmetric distributions, i.e., $X_i \sim F_i$. For symmetric case, we have explicit copulas to obtain the maximizing and minimizing coskewness (see Bernard et al., 2023). Moreover, the symmetric distribution appears as a benchmark in many applications in finance, such as optimal portfolio choice. More general distributions are discovered in Section 3.

The goal of Section 2 is to prove the following two propositions.

Proposition 2.1. Let (X_1, X_2, X_3) be a random vector with symmetric marginals. For any given value of coskewness, ranging between the minimum and maximum admissible values, there exists a dependence model such that the coskewness among the three variables attains this value, and such that the pairwise correlations are all equal to zero.

Proof. In Section 2.1, we construct such a model. \Box Proposition 2.2. Let (X_1, X_2, X_3) be a random vector with symmetric marginals. For every given set of correlations among the three variables, there exists a dependence model such that their coskewness is equal to zero.

Proof. In Section 2.2, we construct such a model. \Box

2.1 Arbitrary coskewness and zero correlation

We recall that the range of possible values for coskewness depends on the choice of marginal distributions. The following lemma recalls Theorems 3.1 and 3.2 of Bernard et al. (2023). We thus do not provide a proof.

Lemma 2.1 (Theorems 3.1 and 3.2 of Bernard et al. (2023)). Let $X_i \sim F_i$ in which the F_i are symmetric, i = 1, 2, 3, and $U \sim U[0, 1]$. The explicit bounds \underline{S} and \overline{S} for the coskewness of X_1 , X_2 and X_3 are

$$\underline{S} := -\mathbb{E}\left(G_1^{-1}(U)G_2^{-1}(U)G_3^{-1}(U)\right) \leq S(X_1,X_2,X_3) \leq \bar{S} := \mathbb{E}\left(G_1^{-1}(U)G_2^{-1}(U)G_3^{-1}(U)\right),$$

in which G_i is the distribution of $|(X_i - \mu_i)/\sigma_i|$. The maximum coskewness \bar{S} is attained for $\bar{S} = S(Y_1, Y_2, Y_3)$ in which $Y_i = F_i^{-1}(U_i)$ with U_i as in

$$U_1 = U,$$

$$U_2 = IJU + I(1-J)(1-U) + (1-I)JU + (1-I)(1-J)(1-U),$$

$$U_3 = IJU + I(1-J)(1-U) + (1-I)J(1-U) + (1-I)(1-J)U,$$
(2.1)

where $I = \mathbb{1}_{U > \frac{1}{2}}$, $J = \mathbb{1}_{V > \frac{1}{2}}$, and $V \stackrel{d}{=} U[0,1]$ is independent of U. The minimum coskewness \underline{S} is attained for $\underline{S} = S(H_1, H_2, H_3)$ in which $H_i = F_i^{-1}(U_i)$ with U_i as in

$$U_1 = U,$$

$$U_2 = IJU + I(1-J)(1-U) + (1-I)JU + (1-I)(1-J)(1-U),$$

$$U_3 = IJ(1-U) + I(1-J)U + (1-I)JU + (1-I)(1-J)(1-U).$$
(2.2)

When F_i , i = 1, 2, 3, are symmetric, the bounds can be computed explicitly; see Table 2 in Bernard et al. (2023).

We now construct a model in which the coskewness varies from \underline{S} to \bar{S} but where the pairwise correlations of these variables are always equal to zero. To do so, let us introduce a mixture copula C^{λ} for $\lambda \in [0,1]$ based on Lemma 2.1. We refer to Lindsay (1995) for a study on mixture models.

Definition 2.1 (Mixture Copula). Let $X_i \sim F_i$, i = 1, 2, 3, in which the F_i are symmetric, $U \stackrel{d}{=} V \sim U[0,1]$ such that $U \perp V$, $B \sim Bernoulli(\lambda)$ where B is independent of X_i , U and V, and $\lambda \in [0,1]$. Define two indicator functions $I = \mathbb{1}_{U > \frac{1}{2}}$ and $J = \mathbb{1}_{V > \frac{1}{2}}$. The dependence structure of $X_1 = F_1^{-1}(U_1)$, $X_2 = F_2^{-1}(U_2)$ and $X_3 = F_3^{-1}(U_3^{\lambda})$ is called a mixture copula C^{λ} when the trivariate random vector $(U_1, U_2, U_3^{\lambda})$ is given as

$$U_1 = U,$$

$$U_2 = IJU + I(1-J)(1-U) + (1-I)JU + (1-I)(1-J)(1-U),$$

$$U_3^{\lambda} = BU_3^M + (1-B)U_3^m,$$
(2.3)

where

$$U_3^M = IJU + I(1-J)(1-U) + (1-I)J(1-U) + (1-I)(1-J)U$$

and

$$U_3^m = IJ(1-U) + I(1-J)U + (1-I)JU + (1-I)(1-J)(1-U).$$

 U_j in (2.1) and U_j in (2.2) are the same for j = 1, 2, thus $U_j^m = U_j^M = U_j^\lambda = U_j$. Note that McNeil et al. (2022) use the same principle to mix U and 1 - U to study the property of Kendall's tau.

Proposition 2.3. Let (X_1, X_2, X_3) be a trivariate random vector with symmetric marginals F_i for i = 1, 2, 3, i.e. $X_i \sim F_i$ and having the mixture copula C^{λ} . The coskewness $S(X_1, X_2, X_3)$ can take any values from the minimum to the maximum by varying the parameter λ in the mixture copula C^{λ} .

Proof. Without loss of generality, we assume that X_i , i = 1, 2, 3, have zero means and unit variances. With the mixture copula C^{λ} , we have

$$\begin{split} X_1 &= F_1^{-1}(U), \\ X_2 &= F_2^{-1}(U_2), \\ X_3 &= BF_3^{-1}(U_3^M) + (1-B)F_3^{-1}(U_3^m). \end{split}$$

Then, the coskewness of X_1 , X_2 and X_3 is

$$\begin{split} S(X_1, X_2, X_3) = & \mathbb{E}\left(F_1^{-1}(U)F_2^{-1}(U_2)\left(BF_3^{-1}(U_3^M) + (1-B)F_3^{-1}(U_3^m)\right)\right) \\ = & \mathbb{E}\left(BF_1^{-1}(U)F_2^{-1}(U_2)F_3^{-1}(U_3^M)\right) + \mathbb{E}\left((1-B)F_1^{-1}(U)F_2^{-1}(U_2)F_3^{-1}(U_3^m)\right) \\ = & \lambda \mathbb{E}\left(F_1^{-1}(U)F_2^{-1}(U_2)F_3^{-1}(U_3^M)\right) + (1-\lambda)\mathbb{E}\left(F_1^{-1}(U)F_2^{-1}(U_2)F_3^{-1}(U_3^m)\right) \\ = & \lambda \bar{S} + (1-\lambda)\underline{S}. \end{split}$$

The third equation holds because B is independent of X_i , U and V.

The proof of Proposition 2.3 shows that the mixture copula C^{λ} leads to a coskewness that is a linear combination between the maximum coskewness and the minimum coskewness with weights driven by the parameter λ . We then consider a trivariate random vector with symmetric marginals and the mixture copula C^{λ} . Thus, the mixture random variables $X_j = F_j^{-1}(U_j)$ for j = 1, 2 and $X_3^{\lambda} = F_3^{-1}(U_3^{\lambda})$, in which U_j and U_3^{λ} are in (2.3). Let us denote this model as $(X_1, X_2, X_3^{\lambda})$. In Appendix A, we provide a numerical method to simulate the dependence structure C^{λ} and thus the model $(X_1, X_2, X_3^{\lambda})$.

We proceed by simulating the mixture copula $C^{\lambda} = (U_1, U_2, U_3^{\lambda})$ using Algorithm A.1




Figure 1: Effect of the parameter λ on coskewness in the case of three normal variables. The coskewness is obtained by simulation with number of simulations $n=10^5$. The approximate minimum $(\lambda=0)$ and maximum coskewness $(\lambda=1)$ are $-1.59\approx -\frac{2\sqrt{2\pi}}{\pi}$ and $1.59\approx \frac{2\sqrt{2\pi}}{\pi}$, respectively.

in Appendix A. Figure 1 illustrates that this model allows us to span all possible levels of coskewness. This result follows immediately by the construction of the mixture copula and by the continuity of coskewness with respect to the parameter λ . Moreover, the plot proves Proposition 2.3 numerically. Given the behaviour of coskewness as a linear function of λ , we can then use λ to represent the level of coskewness.

We can prove that the correlation coefficient is equal to zero in this mixture model. Hence, we obtain the following proposition.

Proposition 2.4. Let (X_1, X_2, X_3) be a trivariate random vector with symmetric marginals F_i for i = 1, 2, 3, i.e. $X_i \sim F_i$ and having the mixture copula C^{λ} . The pairwise correlation coefficients of the three variables are equal to zero, while their coskewness takes arbitrary values (depending on the value of λ) between the minimum and the maximum.

Proof. We only need to prove that correlations are equal to zero. Without loss of generality, we assume that all X_i are symmetrically distributed random variables with zero means and unit variances. Observe that for an indicator function

$$\mathbb{1}_{A}(\omega) = \begin{cases} 1, & \text{if } \omega \in A, \\ 0, & \text{otherwise,} \end{cases}$$

we have $\mathbb{1}_A f(x) + (1 - \mathbb{1}_A) f(y) = f(\mathbb{1}_A x + (1 - \mathbb{1}_A) y)$ for all functions. Note that I, J, B, 1 - I, 1 - J and 1 - B in dependence structure C^{λ} are all indicator functions as well as their products. Thus, under assumptions of symmetric marginals and dependence structure C^{λ} ,

we have

$$\begin{split} X_1 = & F_1^{-1}(U), \\ X_2 = & IJF_2^{-1}(U) + I(1-J)F_2^{-1}(1-U) + (1-I)JF_2^{-1}(U) + (1-I)(1-J)F_2^{-1}(1-U), \\ X_3 = & IJF_3^{-1}(BU + (1-B)(1-U)) + I(1-J)F_3^{-1}(B(1-U) + (1-B)U) \\ & + (1-I)JF_3^{-1}(B(1-U) + (1-B)U) + (1-I)(1-J)F_3^{-1}(BU + (1-B)(1-U)). \end{split}$$

Note that $\Phi^{-1}(U)$ in X_1 can be expanded as follow

$$X_1 = IJF_1^{-1}(U) + I(1-J)F_1^{-1}(U) + (1-I)JF_1^{-1}(U) + (1-I)(1-J)F_1^{-1}(U).$$

We now prove that ρ_{12} equals zero using $F_2^{-1}(U) = -F_2^{-1}(1-U)$. We obtain

$$\rho_{12} = \mathbb{E}(X_1 X_2) = \frac{1}{2} \left[\mathbb{E}(IF_1^{-1}(U)F_2^{-1}(U)) + \mathbb{E}(IF_1^{-1}(U)F_2^{-1}(1-U)) \right]$$

$$+ \mathbb{E}((1-I)F_1^{-1}(U)F_2^{-1}(U)) + \mathbb{E}((1-I)F_1^{-1}(U)F_2^{-1}(1-U)) \right]$$

$$= \frac{1}{2} \left[\mathbb{E}(IF_1^{-1}(U)F_2^{-1}(U)) - \mathbb{E}(IF_1^{-1}(U)F_2^{-1}(U)) \right]$$

$$+ \mathbb{E}((1-I)F_1^{-1}(U)F_2^{-1}(U)) - \mathbb{E}((1-I)F_1^{-1}(U)F_2^{-1}(U)) \right] = 0.$$

Similarly, we have $\rho_{13} = 0$ since

$$\mathbb{E}(IF_1^{-1}(U)F_3^{-1}(BU + (1-B)(1-U))) = -\mathbb{E}(IF_1^{-1}(U)F_3^{-1}(B(1-U) + (1-B)U))$$

and

$$\mathbb{E}((1-I)F_1^{-1}(U)F_3^{-1}(B(1-U)+(1-B)U)) = -\mathbb{E}((1-I)F_1^{-1}(U)F_3^{-1}(BU+(1-B)(1-U))).$$

The proof that $\rho_{23} = 0$ is similar and omitted.

Proposition 2.1 follows as a corollary of Proposition 2.4.

2.2 Arbitrary correlation and zero coskewness

Proposition 2.5. Let (X_1, X_2, X_3) be a trivariate Gaussian random vector. The coskewness $S(X_1, X_2, X_3)$ of X_1 , X_2 and X_3 , equals zero for any possible values of correlation ρ_{ij} of X_i and X_j , i, j = 1, 2, 3 and $i \neq j$.

Proof. We only need to prove that coskewness is equal to zero. It is well-known that the

trivariate Gaussian random vector (X_1, X_2, X_3) can be expressed as

$$X_{1} = \mu_{1} + \sigma_{1} Z_{1},$$

$$X_{2} = \mu_{2} + \sigma_{2} \left(\rho_{12} Z_{1} + a Z_{2} \right),$$

$$X_{3} = \mu_{3} + \sigma_{3} \left(\rho_{13} Z_{1} + \frac{\rho_{23} - \rho_{12} \rho_{13}}{a} Z_{2} + \frac{b}{a} Z_{3} \right),$$

$$(2.4)$$

where $a = \sqrt{1 - \rho_{12}^2}$, $b = \sqrt{1 - \rho_{12}^2 - \rho_{13}^2 - \rho_{23}^2 + 2\rho_{12}\rho_{13}\rho_{23}}$, and Z_1 , Z_2 and Z_3 are independent standard normally distributed random variables. The coskewness of X_1 , X_2 and X_3 is

$$S(X_{1}, X_{2}, X_{3}) = \mathbb{E}\left(Z_{1} \left(\rho_{12}Z_{1} + aZ_{2}\right) \left(\rho_{13}Z_{1} + \frac{\rho_{23} - \rho_{12}\rho_{13}}{a}Z_{2} + \frac{b}{a}Z_{3}\right)\right)$$

$$= \rho_{12}\rho_{13}\mathbb{E}Z_{1}^{3} + \left(\frac{\rho_{12}(\rho_{23} - \rho_{12}\rho_{13})}{a} + a\rho_{13}\right)\mathbb{E}(Z_{1}^{2}Z_{2})$$

$$+ \frac{b\rho_{12}}{a}\mathbb{E}(Z_{1}^{2}Z_{3}) + (\rho_{23} - \rho_{12}\rho_{13})\mathbb{E}(Z_{1}Z_{2}^{2}) + b\mathbb{E}(Z_{1}Z_{2}Z_{3})$$

$$= 0.$$

The last equation for $S(X_1, X_2, X_3)$ holds because $\mathbb{E}Z_1 = \mathbb{E}Z_2 = \mathbb{E}Z_3 = \mathbb{E}Z_1^3 = 0$.

Proposition 2.2 follows as a corollary of Proposition 2.5.

3 Rank correlation and rank coskewness

The range of possible coskewness generally depends on the choice of distributions. Thus in this section, we study the standardized rank coskewness (Bernard et al., 2023) as it always takes values in [-1, 1].

We first recall the definitions of the standardized rank coskewness from Bernard et al. (2023) and the rank correlation.

Definition 3.1 (Standardized Rank Coskewness). Let $X_i \sim F_i$, i = 1, 2, 3, such that F_i are strictly increasing and continuous. The standardized rank coskewness of X_1 , X_2 and X_3 denoted by $RS(X_1, X_2, X_3)$ is defined as $RS(X_1, X_2, X_3) = \frac{4\sqrt{3}}{9}S(F_1(X_1), F_2(X_2), F_3(X_3))$. Hence,

$$RS(X_1, X_2, X_3) = 32\mathbb{E}\left(\left(F_1(X_1) - \frac{1}{2}\right)\left(F_2(X_2) - \frac{1}{2}\right)\left(F_3(X_3) - \frac{1}{2}\right)\right).$$

Definition 3.2 (Rank Correlation). Let $X_i \sim F_i$, i = 1, 2, such that F_i are strictly increasing and continuous. The Spearman's correlation of X_1 and X_2 denoted by ρ_{12}^S is defined

as

$$\rho_{12}^S = 12\mathbb{E}\left(\left(F_1(X_1) - \frac{1}{2}\right)\left(F_2(X_2) - \frac{1}{2}\right)\right).$$

The goal of Section 3 is to prove the following two propositions.

Proposition 3.1. Let $X_i \sim F_i$, i = 1, 2, 3, such that F_i are strictly increasing and continuous. For any value in [-1, 1], one can construct a dependence model such that standardized rank coskewness among X_1 , X_2 and X_3 has that value. In contrast, the pairwise rank correlations among them are equal to zero.

Proof. In Section 3.1, we construct such a model.

Proposition 3.2. Let $X_i \sim F_i$, i = 1, 2, 3, such that F_i are strictly increasing and continuous. There exists a dependence model such that the pairwise rank correlations among X_1 , X_2 and X_3 can take any possible values in [-1, 1], but their standardized rank coskewness is zero.

Proof. In Section 3.2, we construct such a model.

3.1 Arbitrary rank coskewness and zero rank correlation

Proposition 3.3. Let (X_1, X_2, X_3) be a trivariate random vector with strictly increasing and continuous marginals F_i , i = 1, 2, 3, and the mixture copula C^{λ} . The standardized rank coskewness can take any possible values in [-1, 1], while the rank correlation coefficients ρ_{ij}^S of X_i and X_j , j = 1, 2, 3 and $j \neq i$, are equal to zero.

Proof. We only need to prove that the rank correlation coefficients are equal to zero. Lemma 2.1, in this case, still holds because $F_i(X_i)$ are distributed as standard uniform. Thus, we have

$$F_1(X_1) = U,$$

$$F_2(X_2) = IJU + I(1-J)(1-U) + (1-I)JU + (1-I)(1-J)(1-U),$$

$$F_3(X_3) = IJ(BU + (1-B)(1-U)) + I(1-J)(B(1-U) + (1-B)U) + (1-I)J(B(1-U) + (1-B)U) + (1-I)J(B(1-U) + (1-B)U) + (1-I)(B(1-U)).$$

We now prove that rank correlation is equal to zero. It is

$$\rho_{12}^{S} = 12\mathbb{E}\left(\left(F_{1}(X_{1}) - \frac{1}{2}\right)\left(F_{2}(X_{2}) - \frac{1}{2}\right)\right) \\
= 6\left[\mathbb{E}\left(IU^{2}\right) + \mathbb{E}\left(IU\left(1 - U\right)\right) + \mathbb{E}\left((1 - I)U^{2}\right) + \mathbb{E}\left((1 - I)U\left(1 - U\right)\right) - \frac{1}{2}\right] \\
= 6\left[\mathbb{E}\left(IU\right) + \mathbb{E}\left((1 - I)U\right) - \frac{1}{2}\right] = 0.$$

Similarly, we have

$$\begin{split} \rho_{13}^S = &12\mathbb{E}\left(\left(F_1(X_1) - \frac{1}{2}\right)\left(F_3(X_3) - \frac{1}{2}\right)\right) \\ = &6\left[\mathbb{E}\left(I(BU^2 + (1-B)(1-U)U) + I(B(1-U)U + (1-B)U^2) \right. \\ &\left. + (1-I)(B(1-U)U + (1-B)U^2) + (1-I)(BU^2 + (1-B)(1-U)U)\right) - \frac{1}{2}\right] \\ = &6\left[\mathbb{E}\left(BU^2 + (1-B)(1-U)U + B(1-U)U + (1-B)U^2\right) - \frac{1}{2}\right] \\ = &6\left[\mathbb{E}\left((1-B)U + BU\right) - \frac{1}{2}\right] = 0. \end{split}$$

 $\rho_{23}^S = 0$ can be similarly proven.

Proposition 3.1 follows as a corollary of Proposition 3.3.

3.2 Arbitrary rank correlation and zero rank coskewness

Proposition 3.4. Let (X_1, X_2, X_3) be a trivariate random vector with strictly increasing and continuous marginals F_i , i = 1, 2, 3, and Gaussian copula. The standardized rank coskewness $RS(X_1, X_2, X_3)$ of X_1 , X_2 and X_3 is equal to zero for any possible values of rank correlation ρ_{ij}^S of X_i and X_j , i, j = 1, 2, 3 and $i \neq j$.

Proof. Recall Equation (2.4), we have $F_1(X_1) = \Phi(H_1)$, $F_2(X_2) = \Phi(H_2)$ and $F_3(X_3) = \Phi(H_3)$, where

$$H_1 = Z_1,$$

$$H_2 = \rho_{12}Z_1 + \sqrt{1 - \rho_{12}^2}Z_2,$$

$$H_3 = \rho_{13}Z_1 + \frac{\rho_{23} - \rho_{12}\rho_{13}}{\sqrt{1 - \rho_{12}^2}}Z_2 + \frac{\sqrt{1 - \rho_{12}^2 - \rho_{13}^2 - \rho_{23}^2 + 2\rho_{12}\rho_{13}\rho_{23}}}{\sqrt{1 - \rho_{12}^2}}Z_3.$$

Pearson (1907) proves the relationship between the Pearson correlation and the Spearman

rank correlation under Gaussian copula, i.e. for i, j = 1, 2, 3 and $i \neq j$,

$$\rho_{ij} = 2\sin\left(\frac{\pi}{6}\rho_{ij}^S\right).$$

Thus,

$$\rho_{ij}^S = \frac{6}{\pi} \arcsin\left(\frac{\rho_{ij}}{2}\right) \in [-1, 1].$$

This implies $\mathbb{E}(F_i(X_i)F_j(X_j)) = \mathbb{E}(\Phi(H_i)\Phi(H_j)) = \frac{1}{2\pi}\arcsin\left(\frac{\rho_{ij}}{2}\right) + \frac{1}{4}$. The rank coskewness of X_1, X_2 and X_3 is

$$RS(X_{1}, X_{2}, X_{3}) = 32\mathbb{E}\left(\left(F_{1}(X_{1}) - \frac{1}{2}\right)\left(F_{2}(X_{2}) - \frac{1}{2}\right)\left(F_{3}(X_{3}) - \frac{1}{2}\right)\right)$$

$$= 32\left[\mathbb{E}(F_{1}(X_{1})F_{2}(X_{2})F_{3}(X_{3})) - \frac{1}{2}\mathbb{E}(F_{1}(X_{1})F_{2}(X_{2})) - \frac{1}{2}\mathbb{E}(F_{1}(X_{1})F_{3}(X_{3})) - \frac{1}{2}\mathbb{E}(F_{2}(X_{2})F_{3}(X_{3})) + \frac{1}{4}\right]$$

$$= 32\left[\mathbb{E}(F_{1}(X_{1})F_{2}(X_{2})F_{3}(X_{3})) - \frac{1}{4\pi}\arcsin\left(\frac{\rho_{12}}{2}\right) - \frac{1}{4\pi}\arcsin\left(\frac{\rho_{13}}{2}\right) - \frac{1}{4\pi}\arcsin\left(\frac{\rho_{23}}{2}\right) - \frac{1}{8}\right].$$

We assume f_{H_1,H_2,H_3} is the joint density function of (H_1,H_2,H_3) . Define X_1' , X_2' and X_3' as three independent standard normally distributed random variables such that they are independent of X_1 , X_2 and X_3 . Let $(Y_1,Y_2,Y_3) = \left(\frac{X_1'-H_1}{\sqrt{2}},\frac{X_2'-H_2}{\sqrt{2}},\frac{X_3'-H_3}{\sqrt{2}}\right)$. (Y_1,Y_2,Y_3) is also trivariate normal with zero means and unit variances, and the pairwise correlation coefficients are equal to $\frac{\rho_{ij}}{2}$. Then,

$$\mathbb{E}(F_{1}(X_{1})F_{2}(X_{2})F_{3}(X_{3})) = \mathbb{E}(\Phi(H_{1})\Phi(H_{2})\Phi(H_{3}))$$

$$= \int_{\mathbb{R}} \int_{\mathbb{R}} \int_{\mathbb{R}} \Phi(x_{1})\Phi(x_{2})\Phi(x_{3})f_{H_{1},H_{2},H_{3}}(x_{1},x_{2},x_{3})dx_{1}dx_{2}dx_{3}$$

$$= \int_{\mathbb{R}} \int_{\mathbb{R}} \int_{\mathbb{R}} \mathbb{P}(X'_{1} \leq x_{1}, X'_{2} \leq x_{2}, X'_{3} \leq x_{3})f_{H_{1},H_{2},H_{3}}(x_{1},x_{2},x_{3})dx_{1}dx_{2}dx_{3}$$

$$= \int_{\mathbb{R}} \int_{\mathbb{R}} \mathbb{P}(X'_{1} \leq x_{1}, X'_{2} \leq x_{2}, X'_{3} \leq x_{3}|H_{1} = x_{1}, H_{2} = x_{2}, H_{3} = x_{3})$$

$$f_{H_{1},H_{2},H_{3}}(x_{1},x_{2},x_{3})dx_{1}dx_{2}dx_{3}$$

$$= \mathbb{P}(X'_{1} \leq H_{1}, X'_{2} \leq H_{2}, X'_{3} \leq H_{3})$$

$$= \mathbb{P}\left(\frac{X'_{1} - H_{1}}{\sqrt{2}} \leq 0, \frac{X'_{2} - H_{2}}{\sqrt{2}} \leq 0, \frac{X'_{3} - H_{3}}{\sqrt{2}} \leq 0\right)$$

$$= \mathbb{P}\left(Y_{1} < 0, Y_{2} < 0, Y_{3} < 0\right).$$

Rose et al. (2002) prove that

$$\mathbb{P}\left(Y_1 \leq 0, Y_2 \leq 0, Y_3 \leq 0\right) = \frac{1}{4\pi} \left[\arcsin\left(\rho_{Y_1 Y_2}\right) + \arcsin\left(\rho_{Y_1 Y_3}\right) + \arcsin\left(\rho_{Y_2 Y_3}\right)\right] + \frac{1}{8}.$$
Therefore, $RS(X_1, X_2, X_3) = 0.$

Proposition 3.2 follows as a corollary of Proposition 3.4. Let us illustrate this feature with more examples of rank coskewness in the cases of strictly increasing and continuous marginals and various copulas.

Example 3.1. Rank coskewness and rank correlation for strictly increasing and continuous marginals under various dependence assumptions.

Assume that $X_i \sim F_i$ for i = 1, 2, 3 and $U \sim U[0, 1]$.

(1) With the comonotonic copula, we can know $F_i(X_i) = U$. The rank coskewness is

$$RS(X_1, X_2, X_3) = 32\mathbb{E}\left(\left(U - \frac{1}{2}\right)^3\right) = 0.$$

The rank correlations are $\rho_{12}^S = \rho_{13}^S = \rho_{23}^S = 1$.

(2) From Rüschendorf and Uckelmann (2002), the mixing copula is the dependence such that $\sum_{i=1}^{3} U_i = \frac{3}{2}$ where

$$U_{1} = U;$$

$$U_{2} = \begin{cases}
-2U + 1, & \text{if } 0 \le U \le \frac{1}{2}; \\
-2U + 2, & \text{if } \frac{1}{2} \le U \le 1;
\end{cases}$$

$$U_{3} = \begin{cases}
U + \frac{1}{2}, & \text{if } 0 \le U \le \frac{1}{2}; \\
U - \frac{1}{2}, & \text{if } \frac{1}{2} \le U \le 1.
\end{cases}$$

Under the mixing copula, we find that the rank coskewness of X_1 , X_2 and X_3 is given as

$$RS(X_1, X_2, X_3) = 32 \left[\mathbb{E}\left(\left(U - \frac{1}{2}\right) \left(-2U + \frac{1}{2}\right) U \mathbb{1}_{U \in [0, \frac{1}{2}]}\right) + \mathbb{E}\left(\left(U - \frac{1}{2}\right) \left(-2U + \frac{3}{2}\right) (U - 1) \mathbb{1}_{U \in [\frac{1}{2}, 1]}\right) \right]$$
$$= 0.$$

The rank correlations are $\rho_{12}^S=-1,\,\rho_{13}^S=1$ and $\rho_{23}^S=-1.$

(3) Under the independence copula, we have that $F_i(X_i) = U_i$, in which X_1 , X_2 and X_3 are independent. Moreover, the rank coskewness is

$$RS(X_1, X_2, X_3) = 32\mathbb{E}\left(U_1 - \frac{1}{2}\right)\mathbb{E}\left(U_2 - \frac{1}{2}\right)\mathbb{E}\left(U_3 - \frac{1}{2}\right) = 0.$$

The rank correlations are $\rho_{12}^S=\rho_{13}^S=\rho_{23}^S=0.$

These examples also support the proof for Proposition 3.2.

4 Coskewness and tail risk

Rather than using Pearson correlation as in the previous sections, we utilize the event conditional correlation coefficient to analyze the relationship between two random variables, X_i and X_j , i, j = 1, 2, 3, given a particular event \mathcal{A} ; see the same definition in Maugis (2014). This coefficient, denoted as $\rho_{ij|\mathcal{A}}$ and given as

$$\rho_{ij|\mathcal{A}} = \frac{\mathbb{E}((X_i - \mu_{X_i|\mathcal{A}})(X_j - \mu_{X_j|\mathcal{A}})|\mathcal{A})}{\sigma_{X_i|\mathcal{A}}\sigma_{X_i|\mathcal{A}}}$$
(4.1)

quantifies the degree of correlation between X and Y, conditioned on event A. Similarly, $\mu_{X_i|A}$ and $\sigma_{X_i|A}$ are the respective conditional mean and standard deviation of X_i .

One notable application of conditional correlation in risk management is the exceedance correlation, where event A is defined as exceeding a certain threshold, i.e., $\{X_i > \theta_1, X_j > \theta_1, X_j > \theta_1, X_j > \theta_2, X_j > \theta_3, X_j >$ θ_2 or $\{X_i \leq \theta_1, X_i \leq \theta_2\}$. Longin and Solnik (2001) first introduce the concept of exceedance correlation to study the dependence structure of international equity markets, while more recent studies, such as Sakurai and Kurosaki (2020), apply this concept to investigate the relationship between oil and the US stock market. In some cases, the exceedance correlation is calculated using the inverse of the cumulative distribution functions of X_i and X_j , denoted as $\theta_1 = F_i^{-1}(p)$ and $\theta_2 = F_j^{-1}(p)$, respectively, in which $p \in [0,1]$. For example, Garcia and Tsafack (2011) use this approach to test the co-movement trend between international equity and bond markets. However, it should be noted that the exceedance correlation is constantly equal to one under specific dependence structures, as described by Equations (2.1) and (2.2). Another interesting conditional correlation in finance is when the event A is the overall volatility of the market (Z) greater than a crisis volatility threshold (z), i.e., we consider $\rho_{ij|Z>z_c}$. X_i and X_j can be two asset returns in the conditional correlation. Banks are considerably interested in estimating $\rho_{ij|Z>z_c}$ efficiently. Kalkbrener and Packham (2015) conducted a study of $\rho_{ij|Z>z_c}$ on determining the appropriate amount of funds to allocate towards crisis management, while Kenett et al. (2015) researched efficient asset allocation during a crisis.

In this subsection, we investigate the relationship between the coskewness and the down-




Figure 2: The effect of coskewness on conditional correlations $\rho_{ij|\mathcal{A}}$. The random vector $(X_1, X_2, X_3^{\lambda})$ has normal marginals and the mixture copula C^{λ} . The event \mathcal{A} is $\{S < \mu_S \text{ where } S = \sum_{i=1}^3 X_i \text{ and } \mu_S = \mathbb{E}S\}$. The coskewness and downside risks are obtained by implementing the Algorithm A.1 with $n = 10^5$.

side risk, which is a type of conditional correlation when event \mathcal{A} is $\{S < \mu_S \text{ where } S = \sum_{i=1}^{3} X_i \text{ and } \mu_S = \mathbb{E}S\}$ in (4.1). Downside risk was first proposed by Bawa and Lindenberg (1977) as a measure of risk for developing a capital asset pricing model and has gained significant interest in portfolio optimization. We refer to Lettau et al. (2014) and Zhang et al. (2021) for further applications of downside risk in finance.

Ang et al. (2006) study the relationship between downside risk and coskewness and find that the risks differ. In this study, we aim to explore if there exists a theoretical connection between downside risk and coskewness risk. To do so, we use the same parameter settings as in Section 2 for Algorithm A.1 but adjust the last step to compute the conditional correlation.

Figure 2 illustrates the relationship between the coskewness of three normal random variables with the mixture copula C^{λ} and the pairwise downside risks, $\rho_{ij|S<\mu_S}$, i,j=1,2,3 and $i\neq j$. Our result shows that as the coskewness becomes more negative, the downside risk sharply increases. Moreover, the reduction rate of downside risk slows down as the coskewness increases. Overall, we find that the downside risk decreases as the coskewness increases, confirming the empirical findings of Ang et al. (2006) and Huang et al. (2012). They conclude that higher downside risk leads to higher average stock returns, while coskewness risk has the opposite effect. That is, higher coskewness is associated with lower downside risk.

5 Conclusion

In this paper, we provide some propositions and examples to illustrate that, in general, there is no link between coskewness and correlation. Under the assumption of some specific models, the coskewness does not affect the correlation, and vice versa. Specifically, the coskewness of three symmetrically distributed random variables takes any values between the maximum and minimum, but the pairwise correlations are equal to zero. Moreover, under the trivariate Gaussian model assumption, the pairwise correlations can reach all possible values, while coskewness equals zero. We generalize the result using the standardized rank coskewness and the rank correlation for all continuous and strictly increasing marginal distributions. Therefore, one needs to be careful when finding potential links between the coskewness and the correlation empirically and theoretically.

Appendix A. Simulation of the dependence structures C^{λ}

As the function Φ^{-1} is cumbersome to deal with, we propose the following algorithm to compute the coskewness and the pairwise correlation coefficients of mixture random variables. We set $\mu_i = 0$ and $\sigma_i = 1$ because the location and scale parameters do not affect the coskewness and correlation coefficient.

Algorithm A.1.

- 1. Set the mixture parameter $\lambda \in [0, 1]$.
- 2. Simulate $\mathbf{u} = (u_1, \dots, u_n)$, $\mathbf{v} = (v_1, \dots, v_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ where u_i , v_i and b_i $i = 1, \dots, n$, are respective n sampled values from random variables $U \sim U[0, 1]$, $V \sim U[0, 1]$ and $B \sim Bernoulli(\lambda)$.
- 3. Compute discrete maximizing and minimizing copulas $\mathbf{u}_{\mathbf{j}}^{\mathbf{M}} = (u_{1j}^{M}, \dots, u_{nj}^{M})$ and $\mathbf{u}_{\mathbf{j}}^{\mathbf{m}} = (u_{1j}^{m}, \dots, u_{nj}^{m})$, j = 1, 2, 3, using \mathbf{u} and \mathbf{v} in terms of copulas (2.1) and (2.2), respectively.
- 4. Compute discrete mixture copula $\mathbf{c}_{\mathbf{j}}^{\lambda} = (c_{1j}^{\lambda}, \dots, c_{nj}^{\lambda})$ where $c_{ij}^{\lambda} = b_i u_{ij}^M + (1 b_i) u_{ij}^M$.
- 5. Compute discrete mixture random variables $\mathbf{x_j} = (x_{1j}, \dots, x_{nj})$ where $x_{ij} = \Phi^{-1}(c_{ij}^{\lambda})$.
- 6. Compute $\bar{x}_j = \frac{1}{n} \sum_{i=1}^n x_{ij}$ and $s_j = \sqrt{\frac{1}{n} \sum_{i=1}^n [(x_{ij} \bar{x}_j)^2]}$. Then $\rho_{jk} = \frac{\frac{1}{n} \sum_{i=1}^n [(x_{ij} \bar{x}_j)(x_{ik} \bar{x}_k)]}{s_j s_k}$ for k = 1, 2, 3 and $k \neq j$, and $S(X_1, X_2, X_3) = \frac{\frac{1}{n} \sum_{i=1}^n [(x_{i1} \bar{x}_1)(x_{i2} \bar{x}_2)(x_{i3} \bar{x}_3)]}{s_1 s_2 s_3}$.

References

- Ang, A., J. Chen, and Y. Xing (2006). Downside risk. Review of Financial Studies 19(4), 1191–1239.
- Bawa, V. S. and E. B. Lindenberg (1977). Capital market equilibrium in a mean-lower partial moment framework. *Journal of Financial Economics* 5(2), 189–200.
- Beddock, A. and P. Karehnke (2020). Two skewed risks. Preprint.
- Bernard, C., J. Chen, L. Rüschendorf, and S. Vanduffel (2023). Coskewness under dependence uncertainty. Statistics & Probability Letters 199, 109853.
- Garcia, R. and G. Tsafack (2011). Dependence structure and extreme comovements in international equity and bond markets. *Journal of Banking & Finance* 35(8), 1954–1970.
- Huang, W., Q. Liu, S. G. Rhee, and F. Wu (2012). Extreme downside risk and expected stock returns. *Journal of Banking & Finance* 36(5), 1492–1502.
- Jondeau, E. and M. Rockinger (2006). Optimal portfolio allocation under higher moments. *European Financial Management* 12(1), 29–55.
- Kalkbrener, M. and N. Packham (2015). Correlation under stress in normal variance mixture models. Mathematical Finance 25(2), 426–456.
- Kenett, D. Y., X. Huang, I. Vodenska, S. Havlin, and H. E. Stanley (2015). Partial correlation analysis: Applications for financial markets. *Quantitative Finance* 15(4), 569–578.
- Lettau, M., M. Maggiori, and M. Weber (2014). Conditional risk premia in currency markets and other asset classes. *Journal of Financial Economics* 114(2), 197–225.
- Lindsay, B. G. (1995). Mixture models: theory, geometry, and applications. Institute of Mathematical Statistics.
- Longin, F. and B. Solnik (2001). Extreme correlation of international equity markets. *Journal of Finance* 56(2), 649-676.
- Maugis, P. (2014). Event conditional correlation: Or how non-linear linear dependence can be. arXiv preprint arXiv:1401.1130.
- McNeil, A. J., J. G. Nešlehová, and A. D. Smith (2022). On attainability of Kendall's tau matrices and concordance signatures. *Journal of Multivariate Analysis* 191, 105033.
- Pearson, K. (1895). Note on regression and inheritance in the case of two parents. *Proceedings of the Royal Society of London* 58(347-352), 240–242.
- Pearson, K. (1907). On further methods of determining correlation, Volume 16. Dulau and Company.
- Rose, C., M. D. Smith, et al. (2002). Mathematical statistics with Mathematica, Volume 1. Springer.
- Rüschendorf, L. and L. Uckelmann (2002). Variance minimization and random variables with constant sum. In *Distributions with given marginals and statistical modelling*, pp. 211–222. Springer.
- Sakurai, Y. and T. Kurosaki (2020). How has the relationship between oil and the us stock market changed after the Covid-19 crisis? *Finance Research Letters* 37, 101773.
- Zhang, W., Y. Li, X. Xiong, and P. Wang (2021). Downside risk and the cross-section of cryptocurrency returns. *Journal of Banking & Finance* 133, 106246.