

# Dense halves in balanced bipartitions of $K_4$ -free graphs

Yue Xu and Xiao-Dong Zhang\*

School of Mathematical Sciences, MOE-LSC, SHL-MAC,  
Shanghai Jiao Tong University, 800 Dongchuan Road, Shanghai, 200240, P.R.  
China

September 2024

## Abstract

Let  $G$  be a simple graph on even vertices. Denote  $D_{2,\infty}^b(G)$  by the minimal value of  $\max\{e(A), e(A^c)\}$  over all 2-balanced partition  $V(G) = A \cup A^c$  where  $|A| = |A^c|$ . In this paper we first provide a lower bound on  $D_{2,\infty}^b(G)$  by examining the graph  $2n$ -blow-up of  $I_7 \vee C_5$ , where  $I_7$  represents a 7-vertex independent set and  $C_5$  is a 5-cycle, which disproves Conjecture 2.6 in [2]. Furthermore, we present two upper bounds on  $D_{2,\infty}^b(G)$  for a  $K_4$ -free graph and a join of independent set and triangle-free graph.

## 1 Introduction

Let  $G$  be a simple graph with vertex set  $V(G)$  and edge set  $E(G)$ . We denote the number of vertices by  $|G| = |V(G)|$  and the number of edges by  $e(G) = |E(G)|$ . Given vertex-subset  $A$ , we denote  $e(G[A])$  the number of edges in  $G$  with both endpoints from  $A$ . If  $A$  and  $B$  are two disjoint vertex subsets, we denote  $e(G[A], G[B])$  the number of edges in  $G$  connecting vertices from  $A$  to vertices from  $B$ . When  $G$  is clear from context, we also write  $e(A)$  and  $e(A, B)$ . A vertex 2-partition of a graph  $G$  is a partition  $V(G) = A \cup A^c$  of its vertex subset into 2 classes. And we denote  $D_2(G) := \min_A (e(A) + e(A^c))$  where the minimum is taken over all 2-partitions. By definition,  $D_2(G)$  also represents the minimum number of edges which have to be removed to make  $G$  bipartite. Furthermore, 2-partition is balanced if  $|A| = |A^c|$ . And we denote

$$D_2^b(G) := \min_A (e(A) + e(A^c)); D_{2,\infty}^b(G) := \min_A (\max\{e(A), e(A^c)\})$$

where the minimum is taken over all 2-balanced-partitions  $V(G) = A \cup A^c$ . For a graph  $H$ , we define a graph  $G$  as  $H$ -free if  $G$  does not contain  $H$  as a subgraph.

---

\*Corresponding author: xiaodong@sjtu.edu.cn

Given two graphs  $G_1$  and  $G_2$ , the join of  $G_1$  and  $G_2$ , denoted by  $G_1 \vee G_2$ , is defined as a graph whose vertex set is  $V(G_1) \cup V(G_2)$ , and whose edge set is  $E(G_1) \cup E(G_2) \cup (v_1, v_2) \mid v_1 \in V(G_1), v_2 \in V(G_2)$ . For a graph  $G$  and an integer  $n$ , the  $n$ -blow-up of  $G$ , denoted by  $G^n$ , is defined as a graph obtained by replacing every vertex  $v$  of  $G$  with  $n$  vertices, where a copy of  $u$  is adjacent to a copy of  $v$  in the blow-up graph if and only if  $u$  is adjacent to  $v$  in  $G$ .

Erdős once offered a \$250 reward to the solution of the sparse half conjecture:

**Conjecture 1.1.** [5] *Let  $G$  be a triangle-free graph on  $n$  vertices. If  $n$  is even, then there exists a balanced 2-partition  $V(G) = A \cup A^c$  such that*

$$\min\{e(A), e(A^c)\} \leq \frac{n^2}{50}.$$

The most recent progress by Razborov[10] demonstrates that every triangle-free graph  $n$  has a subset of vertices of size  $n/2$  spanning at most  $(27/1024)n^2$  edges. Another related conjecture of Erdős stated that:

**Conjecture 1.2.** [4] *Let  $G$  be a triangle-free graph on  $n$  vertices.*

$$D_2(G) \leq \frac{n^2}{25}.$$

The most recent progress by Balogh, Clemen and Lidický[1] demonstrates that every triangle-free graph  $G$  on  $n$  vertices can be made bipartite by deleting at most  $n^2/23.5$  edges. Several years later, they also raised and solved 2 related problems: for every triangle-free graph on  $n$  vertices, where  $n$  is sufficiently large,  $D_2^b(G) \leq n^2/16$  and  $D_{2,\infty}^b(G) \leq n^2/18$ . In addition, [3],[12],[8] investigate the related 2-partition problem concerning a fixed number of graph edges and solve some of these problems.

For  $K_4$ -free graphs, significant work has been done on related problems. Sudakov [13] proved that every  $K_4$ -free graph  $G$  on  $n$  vertices can be made bipartite by deleting at most  $n^2/9$  edges. Recently, Reiher [11], building on the work of Liu and Ma [9], proved that every  $K_4$ -free graph contains a set of size  $n/2$  spanning at most  $n^2/18$ , thereby solving the conjecture proposed in [6]. Additionally, Balogh, Clemen and Lidický[2] recently inquired about the upper bounds of  $D_2^b(G)$  and  $D_{2,\infty}^b(G)$  for  $K_4$ -free graph, and proposed the following conjectures:

**Conjecture 1.3.** *Let  $G$  be a  $K_4$ -free graph on  $n$  vertices. If  $n$  is even, then*

$$D_2^b(G) \leq \frac{n^2}{9}.$$

**Conjecture 1.4.** *Let  $G$  be a  $K_4$ -free graph on  $n$  vertices. If  $n$  is even, then*

$$D_{2,\infty}^b(G) \leq \frac{n^2}{16}.$$

Reiher's work[11] mentioned in the previous paragraph implies that Conjectures 1.3 and 1.4 holds for regular graphs. Furthermore, Balogh, Clemen and Lidický[2] demonstrated that Conjectures 1.3 and 1.4 holds for 3-partite graph.

**Theorem 1.5.** *Let  $G$  be a 3-partite graph on  $n$  vertices. If  $n$  is even,*

$$D_{2,\infty}^b(G) \leq \frac{n^2}{16}.$$

*The equality holds when  $G$  is a complete 3-partite graph with class sizes  $n/2, n/4$  and  $n/4$ .*

The main result of this paper is to provide a family of counterexamples to Conjecture 1.4, as well as to establish an upper bound on  $D_{2,\infty}^b(G)$  for  $K_4$ -free graphs. Additionally, we present an upper bound on  $D_{2,\infty}^b(G)$  for the join of an independent set and a triangle-free graph.

**Theorem 1.6.** *Let  $H = I_7 \vee C_5$ , where  $I_7$  is a 7-vertex independent set and  $C_5$  is a 5-cycle.*

$$D_{2,\infty}^b(H^{2n}) = \frac{37}{24^2} |H^{2n}|^2 > \frac{|H^{2n}|^2}{16}. \quad (1)$$

*where  $H^{2n}$  denotes the  $2n$ -blow-up of  $H$ .*

**Remark 1.7.** *Theorem 1.6 disproves Conjecture 1.4.*

**Theorem 1.8.** *Let  $G$  be a  $K_4$ -free graph on  $n$  vertices. If  $n$  is even and greater than  $10^5$ ,*

$$D_{2,\infty}^b(G) < 0.074n^2. \quad (2)$$

**Theorem 1.9.** *Let  $G$  be the join of an independent set and a triangle-free set, where  $|G| = n$ . If  $n$  is even and large enough,*

$$D_{2,\infty}^b(G) \leq \frac{5}{72}n^2. \quad (3)$$

The organization of this paper is as follows. In section 2, we prove Theorem 1.6. In section 3, we present some preliminaries for Theorem 1.8 and Theorem 1.9. In section 4 we complete the proof of Theorem 1.8, by considering two cases regarding the independent number. In section 5, we prove Theorem 1.9.




Figure 1:  $H : I \vee C_5$

## 2 The lower bound for the maximum value of $D_{2,\infty}^b(G)$

In this section, we will prove Theorem 1.6 and provide a lower bound on the maximum value of  $D_{2,\infty}^b(G)$  for  $K_4$ -free  $G$ .

**Proposition 2.1.** *Let  $H = I_7 \vee C_5$ , where  $I_7$  is a 7-vertex independent set and  $C_5$  is a 5-cycle.*

$$D_{2,\infty}^b(H) = \frac{5}{72}|H|^2.$$

*Proof.* Let  $V(H) = A \cup A^c$  be the 2-balanced partition of  $H$ . Without loss of generality, let  $A$  be the vertex set that contains at least half of the vertices in  $C_5$ . According to the table below,  $e(A) \geq 10 = 5/72 \cdot |H|^2$ .

vertices from $C_5$	min edges from original $C_5$	edges from original $e(K, C_5)$	min $e(A)$
3	1	$3 \times 3 = 9$	10
4	3	$4 \times 2 = 8$	11
5	5	$5 \times 3 = 5$	10

When  $A$  contains  $V(C_5)$ ,  $\max\{e(A), e(A^c)\} = \max(10, 0) = 10$ . Hence,  $D_{2,\infty}^b(H) = 10 = 5|H|^2/72$ .  $\square$

We can apply the similar method to determine the value of  $D_{2,\infty}^b(H^{2n})$ , where  $H^{2n}$  represents the  $2n$ -blow-up of  $H$ .

**Lemma 2.2.** Let  $C_5^n$  be a graph with vertex set  $V(C_5^n) = A_1 \cup A_2 \cup A_3 \cup A_4 \cup A_5$  and edges  $xy \in C_5^n$  iff  $x \in A_i$  and  $y \in A_{i+1}$  for some  $i \in [5]$ , where  $A_6 := A_1$  and  $|A_i| = n$ . Let  $G_0$  be a subgraph of  $C_5^n$  with  $|V(G_0) \cap A_i| = a_i$  for  $i \in [5]$  and we denote the values  $b_1, b_2, b_3, b_4, b_5$  as the non-increasing rearrangement of  $a_1, a_2, a_3, a_4, a_5$ .

1. If  $T(b) = b_1b_4 + b_4b_3 + b_3b_2 + b_2b_5 + b_5b_1$ , then  $e(G_0) = a_1a_2 + a_2a_3 + a_3a_4 + a_4a_5 + a_5a_1 \geq T(b)$ .

2. If  $|V(G_0)| = pn + q$ , where  $p$  and  $q$  are non-negative integers with  $p < 5$  and  $q < n$ ,

$$\text{then } e(G) \geq f(n, p, q) = \begin{cases} 0 & p \leq 1 \\ qn & p = 2 \\ n^2 + 2qn & p = 3 \\ 3n^2 + 2qn & p = 4 \end{cases}. \text{ Equality holds if } b_i = \begin{cases} n & i \leq p \\ q & i = p + 1 \\ 0 & \text{other cases} \end{cases}.$$




Figure 2:  $C_5^n$  and its subgraph  $G_0$

*Proof.* Without loss of generality, let  $a_1$  denote the largest element among  $\{a_n\}$ , and assume that  $a_2 \geq a_5$ . Based on the arrangements of these elements in accordance with their magnitude, we obtain a total of  $\frac{4!}{2} = 12$  distinct permutations. Let  $S_a = a_1a_2 + a_1a_5 + a_2a_3 + a_3a_4 + a_4a_5$  and  $T(b) = b_1b_4 + b_1b_5 + b_2b_3 + b_2b_5 + b_3b_4$  where  $b = (b_1, b_2, b_3, b_4, b_5)$ .

Note that  $(x_1y_1 + x_2y_2) - (x_1y_2 + x_2y_1) = (x_1 - x_2)(y_1 - y_2)$ . Therefore, the following claim holds.

*Claim 2.1* If  $x_1 \geq x_2$  and  $y_1 \geq y_2$ , then  $x_1y_1 + x_2y_2 \geq x_1y_2 + x_2y_1$ .

Next, we will show that  $e(G_0) = S_a \geq T(b)$  always holds.

*Case 1:*  $b_2 = a_2, b_3 = a_5$ . By Claim 2.1,  $S_a \geq b_1b_2 + b_1b_3 + b_2b_5 + b_4b_5 + b_3b_4 = T(b) + (b_3 - b_5)(b_1 - b_2) + (b_1 - b_5)(b_2 - b_4) \geq T(b)$ .

*Case 2:*  $b_2 = a_2, b_4 = a_5$ . By Claim 2.1,  $S_a \geq b_1b_2 + b_1b_4 + b_2b_5 + b_3b_5 + b_3b_4 = T(b) + (b_2 - b_5)(b_1 - b_3) \geq T(b)$ .

*Case 3:*  $b_2 = a_2, b_5 = a_5$ . By Claim 2.1,  $S_a \geq b_1b_2 + b_1b_5 + b_2b_4 + b_3b_4 + b_3b_5 = T(b) + (b_1 - b_2)(b_3 - b_4) + (b_1 - b_5)(b_2 - b_3) \geq T(b)$ .

*Case 4:*  $b_3 = a_2, b_4 = a_5$ . By Claim 2.1,  $S_a \geq b_1b_3 + b_1b_4 + b_3b_5 + b_2b_5 + b_2b_4 = T(b) + (b_1 - b_3)(b_3 - b_5) + (b_4 - b_5)(b_2 - b_3) \geq T(b)$ .

*Case 5:*  $b_3 = a_2, b_5 = a_5$ . By Claim 2.1,  $S_a \geq b_1b_3 + b_1b_5 + b_3b_4 + b_2b_4 + b_2b_5 = T(b) + (b_1 - b_2)(b_3 - b_4) \geq T(b)$ .

*Case 6:*  $b_4 = a_2, b_5 = a_5$ . By Claim 2.1,  $S_a \geq T(b)$ . Equality holds if  $(b_2, b_3) = (a_3, a_2)$ .

Case 1 to 6 together show that  $S_a \geq T(b)$  under different permutations.

(2)

If  $i < j$ , let  $b^{ij} = (b_1^{ij}, b_2^{ij}, b_3^{ij}, b_4^{ij}, b_5^{ij})$ , where  $b_k^{ij} = \begin{cases} b_k + 1 & k = i \\ b_k - 1 & k = j \\ b_k & \text{other cases} \end{cases}$ . Since  $b_i \geq b_j$

for  $i < j$ ,  $T(b^{ij}) - T(b) \leq 0$  for  $i < j$ , according to the table given below.

$T(b^{ij}) - T(b)$	$i = 1$	$i = 2$	$i = 3$	$i = 4$
$j = 2$	$b_4 - b_3$			
$j = 3$	$b_5 - b_2$	$b_5 - b_4 + b_3 - b_2 - 1$		
$j = 4$	$b_5 - b_3 + b_4 - b_1 - 1$	$b_5 - b_1$	$b_2 - b_1 + b_4 - b_3 - 1$	
$j = 5$	$b_5 - b_2 + b_4 - b_1 - 1$	$b_5 - b_1 + b_3 - b_2 - 1$	$b_4 - b_1$	$b_3 - b_2$

Table 1:  $T(b^{ij}) - T(b)$

Let  $b^* = (b_1^*, b_2^*, b_3^*, b_4^*, b_5^*)$ , where  $b_i^* = \begin{cases} n & i \leq p \\ q & i = p + 1 \\ 0 & \text{other cases} \end{cases}$ . For any  $b$  satisfying  $b_i \geq b_j$

for  $i > j$  and  $\sum_{i=1}^5 b_i = pn + q$ , we can apply the following algorithm to get a new value  $\min(T(b)) = T(b^*)$  not less than  $T(b)$ .

---

**Algorithm 1** Algorithm of  $\min(T(b))$ 

---

**Input:**  $b, b^*$ **Output:**  $\min(T(b))$ 

- 1: declare  $T$  as function defined in Lemma 2.2
  - 2: **while**  $b \neq b^*$  **do**
  - 3:    $x = \min(i)$  such that  $b_i < n$ ,  $y = \max(j)$  such that  $b_j > 0$
  - 4:   **if**  $x < y$  **then**
  - 5:      $b_x = b_x + 1$ ;  $b_y = b_y - 1$ ;  $\min(T(b)) = T(b)$ ;
  - 6:   **end if**
  - 7: **end while**
  - 8: **return**  $\min(T(b))$
- 

Note that each operation in the while loop guarantees  $b_i \geq b_j$  for  $i > j$  and  $\sum_{i=1}^5 b_i = pn + q$ . In this case, according to Table 1,  $T(b)$  does not increase after each iteration. In addition,  $\|b - b^*\|$  decreases by 2 in each iteration, where  $\|\cdot\|$  represents the sum of the absolute values of each component of the vector. This implies that after  $\|b - b^*\|/2$  iterations, the while loop will terminate. At this time,  $\min(T(b)) = T(b^*)$ .

Therefore,  $e(G_0) \geq T(b^*) = f(n, p, q)$ . Equality holds if  $a_1 \geq a_4 \geq a_3 \geq a_2 \geq a_5$ , and  $\{a_n\}$  contains  $p$  elements equal to  $n$ , 1 element equal to  $q$ , and the rest are equal to 0.  $\square$

**Remark 2.3.** *Actually, if  $e(G_0) = T(b) = f(n, p, q)$ ,  $T(b)$  should not decrease after each iteration. This implies if  $b \neq b^*$ ,  $b$  can only be  $(b_1, b_2, 0, 0, 0)$  and  $(n, n, n, b_4, b_5)$  according to Table 1. Hence, if  $pn + q = 2n + q$ , the equality holds if and only if  $b = (n, n, q, 0, 0)$ .*

Next, we will show  $D_{2,\infty}^b(H^{2n}) = 37n^2$  to prove Theorem 1.6.

*Proof of Theorem 1.6.* Let  $V(H^{2n}) = A \cup A^c$  be the 2-balanced partition of  $H^{2n}$  such that  $\max\{e(A), e(A^c)\} = D_{2,\infty}^b(H^{2n})$ . Here,  $|A| = 12n \geq |C_5^{2n}|$ . If  $A \supset V(C_5^{2n})$ ,  $e(A) = 40n^2 > 37n^2$ . Otherwise, let  $|A \cap V(C_5^{2n})| = p \cdot 2n + q$ , where  $p$  and  $q$  are non-negative integers with  $p < 5$  and  $q < 2n$ . Without loss of generality, let  $|A \cap V(C_5^{2n})| \geq |V(C_5^{2n})|/2 = 5n$ . If this condition does not hold, we can let  $A = A^c$ , where  $V(H^{2n}) = A^c \cup (A^c)^c$  forms the same 2-balanced partition. Hence,  $p \geq 2$ , and when  $p = 2$ ,  $q > n$ . Therefore, by Lemma 2.7

$$\begin{aligned} e(A) &= (|A| - |A \cap V(C_5^{2n})|)|A \cap V(C_5^{2n})| + e(A \cap V(C_5^{2n})) \\ &\geq (2pn + q)(12n - 2pn - q) + f(2n, p, q) \\ &= \begin{cases} 32n^2 + q(6n - q) & p = 2, n \leq q < 2n \\ 40n^2 + 4nq - q^2 & p = 3, 0 \leq q < 2n \\ 44n^2 - q^2 & p = 4, 0 \leq q < 2n \end{cases} \end{aligned}$$

Hence,  $e(A) \geq 37n^2$ . Equality holds if and only if  $|A \cap V(C_5^{2n})|$  and  $e(A) = (2pn + q)(12n - 2pn - q) + f(n, p, q)$ . By Remark 2.3, this implies  $A$  contains all vertices in  $A_1$ ,  $A_4$  and half vertices in  $A_3$  in Figure 2, then  $e(A^c) = (|A^c| - |A^c \cap V(C_5^{2n})|)|A \cap V(C_5^{2n})| + e(A^c \cap V(C_5^{2n})) = 5n \cdot 7n + 2n^2 = 37n^2$ .

Therefore,  $D_{2,\infty}^b(G) \geq e(A) \geq 37n^2$ . Additionally,  $D_{2,\infty}^b(G) = 37n^2$  if and only if  $A$  contains  $7n$  vertices from  $|G[A] \cap C_5^{2n}| = K_{n,2n} \cup I_{2n}$ , where  $K_{n,2n}$  represents a complete bipartite with class sizes  $2n$  and  $n$ , and  $I_{2n}$  represents a  $2n$ -vertex independents.

□

### 3 Some technical lemmas

In this section, we will introduce the notations and lemmas that will be used later in this paper. Let  $\alpha(G)$  be the independent number of  $G$ . For  $v \in V(G)$ , let  $N_G(v)$  be the set of neighbors of  $v$  in  $G$  and  $d_G(v) = |N_G(v)|$ . If  $G$  is clear context, we also write  $N(v)$  and  $d(v)$ . We also omit floors and ceiling when they are not essential in our proofs.

The following 3 lemmas will be applied in this paper to estimate the upper bound. For completeness we include their proofs here.

**Lemma 3.1.** *Let  $G$  be a  $K_4$ -free graph on  $n$  vertices.*

1. *If  $Z$  is a maximal triangle-free induced graph of  $G$ , then*

$$e(G) \leq \min \left\{ \frac{|Z||G|}{2}, |Z|(|G| - \frac{3}{4}|Z|) \right\}. \quad (4)$$

2. *If  $Z_0$  is a triangle-free induced graph of  $G$  satisfying  $|Z_0| \geq \frac{2}{3}|G|$ ,*

$$e(G) \leq |Z_0|(|G| - \frac{3}{4}|Z_0|). \quad (5)$$

*Proof.* For any vertex  $v \in G$ , where  $G$  is  $K_4$ -free, its neighbor  $N(v)$  is triangle-free, so  $d(v) \leq |Z|$ , which implies  $e(G) = \frac{1}{2} \sum_{v \in V(G)} d(v) \leq \frac{|Z||G|}{2}$ . In addition, by Turán's Theorem

$$e(Z) \leq \frac{|Z|^2}{4}. \text{ Hence, } e(G) \leq e(Z) + e(Z, G \setminus Z) + 2e(G \setminus Z) \leq \frac{|Z|^2}{4} + \sum_{v \in V(G \setminus Z)} d(v) \leq |Z|(|G| - \frac{3}{4}|Z|).$$

Clearly, the function  $f(x) := x(|G| - \frac{3}{4}x)$  decreases with respect to  $x \geq \frac{2}{3}|G|$ . Hence,  $|Z_0| \geq \frac{2}{3}|G|$  implies inequality (5). □

**Lemma 3.2.** *Let  $G$  be a triangle-free graph on  $n$  vertices. If  $I_0$  is an independent set of  $G$  with  $|I_0| \geq \frac{|G|}{2}$ , then  $e(G) \leq |I_0|(|G| - |I_0|)$ .*



*Proof.* Let  $I$  be a maximal independent set of triangle-free graph  $G$ . For any vertex  $v \in G$ , its neighbor  $N(v)$  is an independent set, so  $d(v) \leq |I|$ , which implies  $e(G) \leq e(I) + 2e(I, G \setminus I) = \sum_{v \in V(G \setminus I)} d(v) \leq (|G| - |I|)|I|$ .

Clearly, the function  $f(x) := x(|G| - x)$  is decreasing with respect to  $x \geq \frac{n}{2}$ . Hence,  $|I_0| \geq |G|/2$  implies  $e(G) \leq |I_0|(|G| - |I_0|)$ .  $\square$

Moreover, if a  $K_4$ -free graph  $G$  contains a large independent set, we can provide an upper bound by Lemma 3.1.

**Lemma 3.3.** *Let  $G$  be a  $K_4$ -free graph on  $n$  vertices and  $I$  be an independent set of  $G$ . If  $Z$  is a largest triangle-free induced graph of  $G \setminus I$ , then*

$$e(G) \leq \min \left\{ \frac{|Z|(|G| + |I|)}{2}, |Z|(|G| - \frac{3}{4}|Z|) \right\}. \quad (6)$$

Moreover, if  $|I| > \frac{|G|}{3}$ , then

$$e(G) \leq \frac{1}{4}(|G| - |I|)(|G| + 3|I|) \quad (7)$$

*Proof.* For any vertex  $v \in I$ , its neighbor  $N(v)$  is triangle-free and  $N(v) \subset G \setminus I$ . Hence,  $d(v) = |N(v)| \leq |Z|$ , which implies  $e(V(I), V(G \setminus I)) \leq \sum_{v \in V(I)} d(v) \leq |Z||I|$ . By Lemma

2.1,  $e(G \setminus I) \leq \min \left\{ \frac{|Z|(|G| - |I|)}{2}, |Z|(|G| - |I| - \frac{3}{4}|Z|) \right\}$ . Therefore,  $e(G) = e(I) + e(G \setminus I) + e(G \setminus I) \leq \min \left\{ \frac{|Z|(|G| + |I|)}{2}, |Z|(|G| - \frac{3}{4}|Z|) \right\}$ .

Clearly, the function  $f(x) := x(|G| - \frac{3}{4}x)$  is decreasing with respect to  $x \geq \frac{2}{3}|G|$ . Hence,  $|Z| \leq |G \setminus I| < \frac{2}{3}|G|$  implies  $e(G) \leq |Z|(|G| - \frac{3}{4}|Z|) \leq (|G| - |I|)(\frac{1}{4}|G| + \frac{3}{4}|I|)$ .  $\square$

The following conclusions can help us find a proper balanced 2-partition.

**Proposition 3.4.** *Let  $G$  be a graph on  $n$  vertices and  $V(G) = B \cup C$ . If  $p \in \{0, 1, \dots, |V(B)|\}$ , there exists a vertex subset  $Y$  (where  $Y \subset B$ ) of size  $p$  such that*

$$e(Y, C) \leq \frac{p}{|B|}e(B, C)$$

This proposition can be found in the literature (e.g., [7]).

**Theorem 3.5.** [14] *Let  $G$  be a graph with  $n$  vertices and  $m$  edges. If  $n$  is even, then  $G$  admits a balanced 2-partition  $V(G) = A \cup A^c$  such that*

$$\max\{e(A), e(A^c)\} \leq \frac{m + \Delta(G) - \delta(G)}{4}$$

where  $\Delta(G)$  and  $\delta(G)$  denote the maximum and minimum degree of the graph  $G$ .

**Corollary 3.6.** *Let  $G$  be an  $n$ -vertex graph where  $n$  is an even number. If  $|E(G)| \leq (4m_0 - \epsilon)n^2$  with  $\epsilon > 0$  and  $n > 1/\epsilon$ , then there exists a balanced 2-partition  $A \cup A^c$  such that  $\max\{e(A), e(A^c)\} < m_0n^2$ .*

In this paper, we set  $m_0 = 0.074$ ,  $\epsilon = 10^{-5}$  in Section 4, and  $m_0 = \frac{5}{72}$ ,  $\epsilon = 0.001$  in Section 5.

We also need the following theorems to further deal with the triangle-free graphs.

**Theorem 3.7.** [1] *Let  $H$  be a triangle-free graph on  $n$  vertices. If  $|E(H)| \geq 0.3197\binom{n}{2}$  with  $n$  large enough,*

$$D_2(H) \leq \frac{n^2}{25}.$$

**Theorem 3.8.** [5] *For a  $K_3$ -free graph with  $n$  vertices and  $m$  edges,  $D_2(G) \leq m - \frac{4m^2}{n^2}$ .*

## 4 A new upper bound for $D_{2,\infty}^b(G)$

In this section we assume that the order of  $G$  is even and large enough. We will prove Theorem 1.8 by considering two cases based on the independent number of  $G$ .

**Lemma 4.1.** *Let  $G$  be a  $K_4$ -free graph on  $n$  vertices, where  $n > 10^5$ . If  $G$  contains an independent set  $I$  of size  $0.28n$ , then*

$$D_{2,\infty}^b(G) < 0.074n^2.$$

*Proof.* Let  $Z$  be a maximal triangle-free induced subgraph in  $G \setminus I$  and  $|Z| = zn$ .

**Case 1.1:**  $|Z| \leq 0.45n$ . By (6) of Lemma 3.3,

$$\begin{aligned} e(G) &\leq \frac{|Z|(|G| + |I|)}{2} \\ &= (0.45n \cdot 0.64n) = 0.288n^2 \end{aligned}$$

Therefore, by Corollary 3.6,  $D_{2,\infty}^b(G) < 0.074n^2$ .

**Case 1.2:**  $|Z| > 0.45n$ . Now, there exists a triangle-free induced graph  $Z_0 \in G \setminus I$  of size  $0.45n$ . Choose a vertex subset  $A$  of size  $n/2$  such that  $A \supset I$  and  $A^c \supset Z_0$ .

Since  $0.28n > \frac{|A|}{3}$ , by (7) of Lemma 3.3,  $e(A) \leq \frac{1}{4}(|A| - |I|)(|A| + 3|I|) = 0.0737n^2 < 0.074n^2$ . On the other hand, since the triangle-free graph  $Z_0 \subset A^c$  and  $|Z_0| \geq \frac{2}{3}|A^c|$ , by ,

$$e(A^c) \leq |Z_0|(|A^c| - \frac{3}{4}|Z_0|) = (0.45n \cdot (0.45n - \frac{3}{4} \cdot 0.45n)) = 0.073125n^2.$$

Therefore,  $\max\{e(A), e(A^c)\} < 0.074n^2$ .

□

According to Corollary 3.6, we only need to consider the graph whose edges are more than  $0.02959n^2$ . Hence, we introduce the following lemma.

**Lemma 4.2.** *Let  $G$  be a graph with  $n$  vertices and  $m$  edges. There exists an edge  $v_1v_2$  such that*

$$d(v_1) + d(v_2) \geq \frac{4m}{n}. \quad (8)$$

If  $G$  is  $K_4$ -free and  $m \geq \frac{n^2}{4}$ , the independent number of  $G$  satisfies  $\alpha(G) \geq \frac{4m}{n} - n$ .

*Proof.* By Cauchy-Schwarz inequality,

$$\sum_{ab \text{ is an edge}} (d(a) + d(b)) = \sum_{v \in V(G)} d(v)^2 \geq \frac{\left(\sum_{v \in V(G)} d(v)\right)^2}{n} = \frac{4m^2}{n}.$$

Therefore, there exists an edge  $v_1v_2$  such that  $d(v_1) + d(v_2) \geq \frac{\sum_{ab \text{ is an edge}} (d(a) + d(b))}{e(G)} = \frac{4m}{n}$ .

Furthermore, if  $G$  is  $K_4$ -free,  $N(v_1) \cap N(v_2)$  is an independent set. Therefore,  $|N(v_1) \cap N(v_2)| \geq |N(v_1)| + |N(v_2)| - |V(G)|$ , which implies  $\alpha(G) \geq \frac{4m}{n} - n$ . □

By Lemma 4.2,  $e(G) \geq 0.02959n^2$  implies  $\alpha(G) \geq \frac{4m}{n} - n \geq 0.1836n$  when  $G$  is  $K_4$ -free.

**Lemma 4.3.** *Let  $G$  be a  $K_4$ -free graph on  $n$  vertices, where  $n > 10^5$ . If the independent number of  $G$  satisfies  $\alpha(G) < 0.28n$  and  $e(G) \geq 0.02959n^2$ , then*

$$D_{2,\infty}^b(G) < 0.074n^2.$$

*Proof.* When  $e(G) > 0.02959n^2$ , Lemma 4.2 ensures that  $\alpha(G) > 0.18n$  and there exists an edge  $v_1v_2$  such that  $d(v_1) + d(v_2) \geq 1.18n$ . We denote  $N_1 = N(v_1)$ ,  $N_2 = N(v_2)$ . Let  $|N_1 \setminus N_2| = a_1n$ ,  $|N_2 \setminus N_1| = a_2n$  and  $|N_1 \cap N_2| = cn$ , then

$$\begin{cases} a_1 + a_2 + 2c \geq 1.18 \\ a_1 + a_2 + c \leq 1 \\ 0.18 < c < 0.28 \end{cases} \quad (9)$$

Without loss of generality, let  $a_1 \geq a_2$  which implies  $a_1 + c \geq \frac{1+0.18}{2} = 0.59$  and  $d(v_1) \geq d(v_2)$ . In addition, since  $N_1$  is triangle-free, for any  $v \in N_1$ ,  $N_{N_1}(v)$  is an independent set. Hence,  $d_{N_1}(v) < 0.28n$ . Similarly,  $d_{N_2}(v) < 0.28n$ .

**Case 3.1:**  $a_2 + c \leq 0.39$ .

This implies  $a_1 + c \geq 0.79$ . Note that  $e(N_1) = \frac{1}{2} \sum_{v \in N_1} d_{N_1}(v) \leq 0.14n \cdot |N_1|$ . In addition, by Turán's theorem, the  $K_4$ -free graph  $G - N_1$  contains at most  $\frac{1}{3}|V(G) \setminus N_1|^2$  edges. Therefore,

$$\begin{aligned} e(G) &\leq e(N_1) + e(N_1, V(G) \setminus N_1) + e(G \setminus N_1) \\ &\leq n^2(0.14(a_1 + c) + (a_1 + c)(1 - a_1 - c) + \frac{1}{3}(1 - a_1 - c)^2) \\ &= n^2 \left( -\frac{2}{3}(a_1 + c)^2 + \left(\frac{1}{3} + 0.14\right)(a_1 + c) + \frac{1}{3} \right) \\ &\leq n^2 \left( \left(-\frac{2}{3}\right) \cdot 0.79^2 + \left(\frac{1}{3} + 0.14\right) \cdot 0.79 + \frac{1}{3} \right) = 0.2912n^2. \end{aligned}$$

This implies  $D_{2,\infty}^b(G) \leq 0.074n^2$  by Corollary 3.6. The last inequality holds because the derivative is below 0 when  $d(v_1) \geq 0.79n$ .

**Case 3.2:**  $a_2 + c \geq 0.45$ .

By (9),  $a_1 + a_2 + c \geq 0.9$ . In addition,  $a_1 + c \geq a_2 + c \geq 0.45$ , so we can choose a vertex subset  $A$  of size  $n/2$  such that  $|A \cap N_1| = 0.45n$  and  $|A^c \cap N_1| = 0.45n$ . Therefore,  $G[A]$  and  $G[A^c]$  contains a triangle-free  $0.45n$ -vertex induced graph. By (5) of Lemma 3.1,  $\max\{e(A), e(A^c)\} \leq (0.45n \cdot (0.5n - 3/4 \cdot 0.45n)) < 0.074n^2$ .

**Case 3.3:**  $0.39 < a_2 + c < 0.45$ .

**Case 3.3.1:**  $0.18 < c \leq 0.23$ . Here,  $a_1 \geq 1.18 - a_2 - 2c > 0.5$ .

Since  $d_{N_1}(x) \leq 0.28n, d_{N_2}(x) \leq 0.28n$ ,  $e(N_1 \cap N_2, N_1 \setminus N_2) = \sum_{v \in V(N_1 \cap N_2)} d(v) \leq 0.28cn^2$ . Since  $n/2 - |N_2| = (0.5 - a_2 - c)n \in (0.05n, 0.11n) \subset (0, a_1]$ , by Proposition 3.4 (let  $B = N_1 \setminus N_2, C = N_1 \cap N_2$  and  $p = n/2 - |N_2|$ ), there exists a vertex subset  $Y$  of size  $(n/2 - |N_2|)$  such that  $N(v_2) \subset A \subset N(v_1) \cup N(v_2)$  and

$$e(Y, N_1 \cap N_2) \leq \frac{\frac{1}{2} - a_2 - c}{a_1} 0.28cn^2.$$

Let  $A = Y \cup (N_2 \setminus N_1)$ . Since  $N_1$  and  $N_2$  are triangle-free, by Turán's Theorem,

$$\begin{aligned}
e(A) &\leq e(N_1 \cap N_2, Y) + e(N_2 \setminus N_1, Y) + e(A \setminus N(v_2)) + e(N(v_2)) \\
&\leq n^2 \left( 0.28c \frac{\frac{1}{2} - a_2 - c}{a_1} + a_2 \left( \frac{1}{2} - a_2 - c \right) + \frac{\left( \frac{1}{2} - a_2 - c \right)^2}{4} + \frac{(a_2 + c)^2}{4} \right) \\
&\stackrel{t=a_2+c}{=} n^2 \left( \frac{0.28}{a_1} c \left( \frac{1}{2} - t \right) + (t - c) \left( \frac{1}{2} - t \right) + \frac{1}{4} \left( \frac{1}{2} - t \right)^2 + \frac{1}{4} t^2 \right) \\
&= n^2 \left( \left( \frac{0.28}{a_1} - 1 \right) c \left( \frac{1}{2} - t \right) - \frac{1}{2} t^2 + \frac{1}{4} t + \frac{1}{16} \right)
\end{aligned}$$

Since  $1 - c \geq a_1 \geq 1.18 - t - c$ ,  $\left( \frac{0.28}{a_1} - 1 \right) c \geq \left( \frac{0.28}{1.18-t-c} - 1 \right) c \stackrel{def}{=} g_t(c)$ . When  $t \in (0.39, 0.45]$  is fixed,  $\frac{dg_t(c)}{dc} = \frac{0.28(1.18-t) - (1.18-t-c)^2}{(1.18-t-c)^2} < 0$ . Hence, the function reaches the maximum when  $c = 0.18$ , which implies:

$$\frac{e(A)}{n^2} \leq 0.18 \left( \frac{0.28}{1-t} - 1 \right) \left( \frac{1}{2} - t \right) - \frac{1}{2} t^2 + \frac{1}{4} t + \frac{1}{16}.$$

Notice that  $f(t) := 0.18 \left( \frac{0.28}{1-t} - 1 \right) \left( \frac{1}{2} - t \right) - \frac{1}{2} t^2 + \frac{1}{4} t + \frac{1}{16}$  is decreasing with respect to  $t \in [0.39, 0.45]$ , because  $f'(t) = 0.43 - t - \frac{0.0252}{(t-1)^2} < 0$  when  $t \in [0.39, 0.45]$ . Hence,  $e(A) \leq n^2 f(0.39) < 0.0733n^2$ . On the other hand,  $|A^c \cap N_1| \geq (a_1 + a_2 + c - 0.5)n \geq (0.68 - c)n \geq 0.45n$ , so by (5) of Lemma 3.1,  $e(A^c) < 0.074n^2$ . Therefore,  $D_{2,\infty}^b(G) \leq 0.074n^2$ .

**Case 3.3.2:**  $0.23 < c \leq 0.28$ . To be precise, we are considering the following five cases:  $c \in (0.22 + 0.01i, 0.23 + 0.01i]$  for  $i \in [5]$ .

Since  $a_2 + c \geq 0.45$  and  $a_1 + a_2 + c \geq 1.18 - c$ ,  $0.5 - 0.01i - a_2 - c \in [0, a_1]$ . Hence, similarly to **Case 3.3.1**, by Proposition 3.4 there exists a vertex subset  $Y_1 \subset (N_1 \setminus N_2)$  with  $|Y_1| = (0.5 - 0.01i - a_2 - c)n$  such that

$$e(Y_1, N_1 \cap N_2) \leq \frac{0.5 - 0.01i - a_2 - c}{a_1} e(N_1 \setminus N_2, N_1 \cap N_2) \leq n^2 \frac{0.5 - 0.01i - a_2 - c}{a_1} 0.28cn^2.$$

Now, we can choose a vertex subset  $A$  of size  $n/2$  such that  $A \supset (N_2 \cup Y_1)$  and  $|A^c \cap (N_1 \setminus N_2)| \geq 0.45$ . This holds because, for each  $i$ ,  $a_1 + a_2 + c \geq 1.18 - c \geq 0.95 - 0.01i$  when  $c \in (0.22 + 0.01i, 0.23 + 0.01i]$ . Here, the condition  $|A^c \cap (N_1 \setminus N_2)| \geq 0.45n$  ensures that  $e(A^c) < 0.074n^2$  by (5) of Lemma 3.1. On the other hand,

$$\begin{aligned}
e(A) &= e(Y_1, N_1 \cap N_2) + e(Y_1, N_2 \setminus N_1) + e(Y_1) + e(N_2) \\
&\quad + e(Y_1 \cup N_2, A \setminus (Y_1 \cup N_2)) + e(A \setminus (Y_1 \cup N_2))
\end{aligned}$$

Let  $t = a_2 + c$ . Now,  $\frac{e(Y_1, N_1 \cap N_2)}{n^2} \leq 0.28c \cdot \frac{0.5 - 0.01i - a_2 - c}{a_1} \leq 0.28c \cdot \frac{0.5 - 0.01i - t}{1.18 - t - c}$ . Moreover,  $G(I_1)$  is triangle-free, so  $e(I_1) \leq \frac{1}{4}(0.5 - 0.01i - t)^2$ ; while  $G(N_2)$  is triangle-free graph and contains an independent set of size  $c > \frac{t}{2}$ , so  $e(I_2) \leq (t - c)c$  by Lemma 3.2. Therefore, we let

$$\begin{aligned} \frac{e(A)}{n^2} &\leq 0.28c \frac{0.5 - 0.01i - t}{1.18 - t - c} + (t - c)(0.5 - 0.01i - t) + (t - c)c + \frac{1}{4}(0.5 - 0.01i - t)^2 \\ &\quad + (0.5 - 0.01i) \cdot 0.01i + \frac{1}{3}(0.01i)^2 \stackrel{def}{=} F_i(c, t) \end{aligned}$$

where  $0.39 < t < 0.45$ ,  $c \in (0.22 + 0.01i, 0.23 + 0.01i]$  and  $i \in [5]$ . Hence,

$$\begin{aligned} \frac{dF_i(c, t)}{dc} &= (0.5 - 0.01i - t) \left( \frac{0.28}{1.18 - t - c} + \frac{0.28c}{(1.18 - t - c)^2} - 1 \right) + t - 2c \\ &< (0.5 - 0.01i - t) \left( \frac{0.28}{0.5 - 0.01i} + \frac{0.28(0.23 + 0.01i)}{(0.5 - 0.01i)^2} - 1 \right) + 0.45 - 0.44 - 0.02i \\ &\leq (0.5 - 0.01i - t) \left( \frac{0.28}{0.45} + \frac{0.28 \cdot 0.28}{0.45^2} - 1 \right) - 0.01 \\ &\leq 0.1 \cdot 0.01 - 0.01 < 0 \end{aligned}$$

Therefore,  $F_i(c, t) \leq F_i(0.22 + i, t)$ . Actually,

$$F_i(0.22 + i, t) = \begin{cases} -\frac{3t^2}{4} + \frac{141t}{200} - \frac{12077}{120000} + \frac{161t}{2500} - \frac{7889}{250000} & i = 1 \\ -\frac{3t^2}{4} + \frac{18t}{25} - \frac{791}{7500} + \frac{42t}{625} - \frac{504}{15625} & i = 2 \\ -\frac{3t^2}{4} + \frac{147t}{200} - \frac{883}{8000} + \frac{7t}{100} - \frac{329}{10000} & i = 3 \\ -\frac{3t^2}{4} + \frac{3t}{4} - \frac{3461}{30000} + \frac{91t}{1250} - \frac{2093}{62500} & i = 4 \\ -\frac{3t^2}{4} + \frac{153t}{200} - \frac{14453}{120000} + \frac{189t}{2500} - \frac{1701}{50000} & i = 5 \end{cases} \quad (10)$$

Below, we will show that  $G_i(t) := F_i(0.22 + i, t)(0.96 - 0.01i - t) - 0.0739(0.96 - 0.01i -$

$t) < 0$  which implies  $e(A) < 0.074n^2$ .

$$G_i(t) = \begin{cases} \left(-\frac{3}{4}t^2 + \frac{141}{200}t - \frac{12077}{120000}\right) \left(\frac{19}{20} - t\right) - \frac{161}{2500}t + \frac{7889}{250000} - 0.0739 \left(t - \frac{19}{20}\right) & i = 1 \\ \left(-\frac{3}{4}t^2 + \frac{18}{25}t - \frac{791}{7500}\right) \left(\frac{47}{50} - t\right) - \frac{42}{625}t + \frac{504}{15625} - 0.0739 \left(t - \frac{47}{50}\right) & i = 2 \\ \left(-\frac{3}{4}t^2 + \frac{147}{200}t - \frac{883}{8000}\right) \left(\frac{93}{100} - t\right) - \frac{7}{100}t + \frac{329}{10000} - 0.0739 \left(t - \frac{93}{100}\right) & i = 3 \\ \left(-\frac{3}{4}t^2 + \frac{3}{4}t - \frac{3461}{30000}\right) \left(\frac{23}{25} - t\right) - \frac{91}{1250}t + \frac{2093}{62500} - 0.0739 \left(t - \frac{23}{25}\right) & i = 4 \\ \left(-\frac{3}{4}t^2 + \frac{153}{200}t - \frac{14453}{120000}\right) \left(\frac{91}{100} - t\right) - \frac{189}{2500}t + \frac{1701}{50000} - 0.0739 \left(t - \frac{91}{100}\right) & i = 5 \end{cases} \quad (11)$$

$$< \begin{cases} 0.75t^3 - 1.4175t^2 + 0.77990t - 0.13416 & i = 1 \\ 0.75t^3 - 1.425t^2 + 0.78898t - 0.13625 & i = 2 \\ 0.75t^3 - 1.4325t^2 + 0.79783t - 0.13838 & i = 3 \\ 0.75t^3 - 1.44t^2 + 0.80648t - 0.14054 & i = 4 \\ 0.75t^3 - 1.4475t^2 + 0.81490t - 0.14273 & i = 5 \end{cases} \quad (12)$$

The figure of functions in (12) is given below.




Figure 3: Graph of (12)

Therefore, the  $A$  we choose ensures  $\max\{e(A), e(A^c)\} \leq 0.074n^2$ .  $\square$

*Proof of Theorem 1.8.* If  $e(G) \leq 0.02959n^2$ , then by Corollary 3.6,  $D_{2,\infty}^b(G) < 0.074n^2$ . If  $e(G) > 0.02959n^2$ , by Lemmas 4.1 and 4.3,  $D_{2,\infty}^b(G) < 0.074n^2$ .  $\square$

## 5 An upper bound for $I \vee H$

*Proof of Theorem 1.9.* Let  $G = H \vee I$ , where  $I$  is an independent set and  $H$  is a triangle-free graph  $H$ . We will prove the Theorem 1.9 by considering the four cases regarding the

order of the independents set.

**Case 1:**  $|I| > 0.62n$ . Now, there exists 2 disjoint vertex-subsets  $I_1$  and  $I_2$  with  $|I_1| = |I_2| = 0.31n$ . Choose a vertex subset  $A$  of size  $n/2$  such that  $A \supset I_1$  and  $A^c \supset I_2$ . Since  $|I_1| = |I_2| \geq \frac{1}{3}|A| = \frac{1}{3}|A^c|$ , by (7) of Lemma 3.3

$$\max\{e(A), e(A^c)\} \leq \frac{1}{4}(0.5n - 0.31n)(0.5n + 3 \cdot 0.31n) = 0.067925n^2 < \frac{5}{72}n^2$$

**Case 2**  $|I| \in (\frac{7}{12}n, 0.62n]$ . If  $e(H) \leq 0.16|H|^2$ , then  $e(G) < 0.271n^2$ , which implies  $D_{2,\infty}^b(G) < \frac{5}{72}n^2$  by Corollary 2.6.

If  $e(H) > 0.16|H|^2$ , we can delete  $\frac{|H|^2}{25}$  edges to make  $H$  bipartite by Theorem 3.7. This implies  $G$  can be made tripartite by deleting  $\frac{|H|^2}{25}$  edges. Moreover,  $|H| = n - |I| < \frac{5n}{12}$ . Therefore, by Theorem 1.4,

$$D_{2,\infty}(G) \leq \frac{|H|^2}{25} + \frac{n^2}{16} < \frac{5}{72}n^2.$$

**Case 3**  $|I| \in [0.5n, \frac{7}{12}n]$ . Let  $|I| = \alpha n$ . Here,  $\alpha \in [0.5, 7/12]$ . If  $\frac{e(H)}{n^2} < \frac{5}{72} - (1 - \alpha)(\alpha - \frac{1}{2})$ , then choose a vertex subset  $A$  of size  $\frac{n}{2}$  such that  $A^c \supset V(H)$ . Then,  $\max\{e(A), e(A^c)\} < \frac{5}{72}n^2$ .

If  $\frac{e(H)}{n^2} \geq \frac{5}{72} - (1 - \alpha)(\alpha - \frac{1}{2})$ , then by Theorem 3.8, we can delete  $(e(H) - \frac{4e(H)^2}{|H|^2})$  edges to make  $H$  bipartite. This implies  $G$  can be made tripartite by deleting  $e(H) - \frac{4e(H)^2}{|H|^2}$  edges. Let  $e_0 = e(H)/n^2$ . By Theorem 1.4,

$$\begin{aligned} \frac{D_{2,\infty}^b(G)}{n^2} &\leq \frac{1}{16} + \frac{e(H)}{n^2} - \frac{4e(H)^2}{|H|^2 n^2} \\ &= \frac{1}{16} + e_0 - \frac{4e_0^2}{(1 - \alpha)^2} \stackrel{def}{=} g_\alpha(e_0) \end{aligned}$$

where  $\alpha \in [0.5, 7/12]$  and  $e_0 \geq \frac{5}{72} - (1 - \alpha)(\alpha - \frac{1}{2})$ . Actually, for a fixed  $\alpha$ , the function  $g_\alpha(e_0)$  is decreasing with respect to  $[\frac{(1-\alpha)^2}{8}, +\infty)$ . Moreover, when  $\alpha \in [0.5, 7/12]$ ,  $\frac{5}{72} - (1 - \alpha)(\alpha - \frac{1}{2}) - \frac{(1-\alpha)^2}{8} = \frac{1}{72}(21\alpha - 16)(3\alpha - 2) > 0$ . Hence,  $g_\alpha(e_0) \leq g_\alpha(\frac{5}{72} - (1 - \alpha)(\alpha - \frac{1}{2}))$ . Let  $f(\alpha) := g_\alpha(\frac{5}{72} - (1 - \alpha)(\alpha - \frac{1}{2})) - \frac{1}{16} = \frac{5}{72} + (\alpha - 1)(\alpha - \frac{1}{2}) - \frac{25}{1296(1-\alpha)^2} - \frac{5}{9} \cdot \frac{\frac{1}{2} - \alpha}{1 - \alpha} - 4(\alpha - \frac{1}{2})^2$ . Note that when  $\alpha \in [0.5, 7/12]$ ,



$$f'(\alpha) = \frac{5}{18} \cdot \frac{1}{(\alpha-1)^2} \left( 1 - \frac{5}{36(1-\alpha)} \right) - 6\alpha + \frac{5}{2} \quad (13)$$

$$\leq \frac{5}{18} \cdot \frac{1}{(\alpha-1)^2} \cdot \frac{2}{3} - 6\alpha + \frac{5}{2} \quad (14)$$

$$= \frac{1}{(\alpha-1)^2} \left( \frac{5}{27} - (6\alpha - \frac{5}{2})(\alpha-1)^2 \right) < 0 \quad (15)$$

(15) holds because the function  $f_0(\alpha) = (6\alpha - 2.5)(\alpha - 1)^2$  is increasing with respect to  $(-\infty, 11/18]$ , which implies  $(6\alpha - \frac{5}{2})(\alpha - 1)^2 - \frac{5}{27} \leq (6 \cdot 7/12 - 5/2)(7/12 - 1)^2 < 0$ . Hence, if  $\alpha \in [0.5, 7/12]$ ,  $f(\alpha) \leq f(7/12) = 1/144$ .

Therefore,  $\frac{D_{2,\infty}^b(G)}{n^2} \leq \frac{1}{16} + f(\frac{7}{12}) \leq 5/72$ .

**Case 4**  $\alpha(G) < 0.5n$ . If  $e(H) \leq (\frac{5}{18} - 0.001)n^2 - |H|(n - |H|)$  (which implies  $e(G) \leq (\frac{5}{18} - 0.001)n^2$ ), then  $\frac{D_{2,\infty}^b(G)}{|G|^2} < \frac{5}{72}$  by Corollary 3.6.

If  $e(H) > (\frac{5}{18} - 0.001)n^2 - |H|(n - |H|)$ , then there exists a vertex  $v \in V(H)$  satisfying  $d_H(v) \geq \frac{2e(H)}{|H|} \geq \frac{0.553n^2}{|H|} + 2|H| - 2n$ . Since  $H$  is triangle-free,  $N_H(v)$  is an independent set with  $|N_H(v)| > 0.553n^2|H| + 2|H| - 2n$ . Hence, we can let  $I_0$  be the independent set of  $H$  with  $|I_0| = \frac{0.553n^2}{|H|} + 2|H| - 2n$ . By AM-GM inequality,  $|V(I \cup I_0)| \geq \frac{0.553n^2}{|H|} + |H| - n \geq (2\sqrt{0.553} - 1)n^2 \geq 0.486n$ .

If  $|V(I \cup I_0)| > \frac{n}{2}$ , we can choose a vertex subset  $A$  of size  $\frac{n}{2}$  such that  $I \subset A \subset I \cup I_0$ . Now,  $A$  is triangle-free, and  $A^c$  which is a subgraph of  $H$  is also triangle-free. Therefore, by Mantel's theorem,  $\frac{D_{2,\infty}^b(G)}{n^2} \leq \frac{1}{16}n^2 < \frac{5}{72}n^2$ .

If  $|V(I \cup I_0)| < \frac{n}{2}$ , we can choose a vertex subset  $A$  of size  $\frac{n}{2}$  such that  $A \supset I \cup I_0$ . Note that  $I \cup I_0$  is a triangle-free graph with  $|I \cup I_0| \geq 0.486n$ , so by (2) of Lemma 3.1  $e(A) \leq |I \cup I_0|(|A| - \frac{3}{4}|I \cup I_0|) = 0.065853n^2 < \frac{5}{72}n^2$ . On the other hand, since  $A^c$  is triangle-free,  $e(A) \leq \frac{(n/2)^2}{4} < \frac{5}{72}n^2$ . Therefore,  $D_{2,\infty}^b(G) < \frac{5}{72}n^2$  in this case.

□

## Acknowledgement

This work is partly supported by the National Natural Science Foundation of China (Nos.12371354, 12161141003) and Science and Technology Commission of Shanghai Municipality (No. 22JC1403600), National Key R&D Program of China under Grant No. 2022YFA1006400.

## References

- [1] J. Balogh, F. C. Clemen, and B. Lidický. Max cuts in triangle-free graphs. *Trends Math. (Extended Abstracts EuroComb 2021)*, 14:509–514, 2021.
- [2] J. Balogh, F. C. Clemen, and B. Lidický. 10 problems for partitions of triangle-free graphs. *Eur. J. Comb.*, page 103841, 2023.
- [3] B. Bollobás and A. D. Scott. Exact bounds for judicious partitions of graphs. *Combinatorica*, 19(4):473–486, 1999.
- [4] P. Erdős. Some old and new problems in various branches of combinatorics. *Discrete Math.*, 165(166):227–231, 1997.
- [5] P. Erdős, R. J. Faudree, J. Pach, and J. H. Spencer. How to make a graph bipartite. *J. Combin. Theory Ser. B*, 45(1):86–98, 1988.
- [6] P. Erdős, R. J. Faudree, C. C. Rousseau, and R. H. Schelp. A local density condition for triangles. *Discrete Math.*, 127(1-3):153–161, 1994.
- [7] M. Krivelevich. On the edge distribution in triangle-free graphs. *J. Combin. Theory Ser. B*, 63(2):245–260, 1995.
- [8] C. Lee, P.-S. Loh, and B. Sudakov. Bisections of graphs. *J. Combin. Theory Ser. B*, 103(5):599–629, 2013.
- [9] X. Liu and J. Ma. Sparse halves in  $K_4$ -free graphs. *J. Graph Theory*, 99(1):5–25, 2022.
- [10] A. A. Razborov. More about sparse halves in triangle-free graphs. *Mat. Sb.*, 213(1):109, jan 2022.
- [11] C. Reiher.  $K_4$ -free graphs have sparse halves. *Bull. Lond. Math. Soc.*, 55:1178–1195, 12 2022.
- [12] A. Scott. Judicious partitions and related problems. *London Math. Soc. Lecture Note Ser.*, 327:95–117, 2005.
- [13] B. Sudakov. Making a  $K_4$ -free graph bipartite. *Combinatorica*, 27(4):509–518, 2007.
- [14] B. Xu, J. Yan, and X. Yu. A note on balanced bipartitions. *Discrete Math.*, 310(20):2613–2617, 2010.