Γ-EXPANSION OF THE MEASURE-CURRENT LARGE DEVIATIONS RATE FUNCTIONAL OF NON-REVERSIBLE FINITE-STATE MARKOV CHAINS

S. KIM AND C. LANDIM

ABSTRACT. Consider a sequence of continuous-time Markov chains $(X_t^{(n)}:t\geq 0)$ evolving on a fixed finite state space V. Let I_n be the measure-current large deviations rate functional for $X_t^{(n)}$, as $t\to\infty$. Under a hypothesis on the jump rates, we prove that I_n can be written as $I_n=\mathbf{I}^{(0)}+\sum_{1\leq p\leq \mathfrak{q}}(1/\theta_n^{(p)})\mathbf{I}^{(p)}$ for some rate functionals $\mathbf{I}^{(p)}$. The weights $\theta_n^{(p)}$ correspond to the time-scales at which the sequence of Markov chains $X_t^{(n)}$ evolve among the metastable wells, and the rate functionals $\mathbf{I}^{(p)}$ characterise the asymptotic Markovian dynamics among these wells. This expansion provides therefore an alternative description of the metastable behavior of a sequence of Markovian dynamics.

1. Introduction

Let V be a finite set. Denote by $(X_t : t \ge 0)$ a V-valued, irreducible continuoustime Markov chain, whose jump rates are represented by R(x, y). The generator reads as

$$(\mathcal{L}f)(x) = \sum_{y \in V} R(x,y) \left\{ f(y) - f(x) \right\}, \quad f \colon V \to \mathbb{R}.$$

Let π be the unique stationary state. The so-called Matrix tree Theorem [15, Lemma 6.3.1] provides a representation of the measure π in terms of arborescences of the set V.

Denote by $\mathcal{P}(V)$ the space of probability measures on V endowed with the weak topology, and by L_t the empirical measure of the chain X_t defined as:

$$L_t := \frac{1}{t} \int_0^t \delta_{X_s} \, \mathrm{d}s \,, \tag{1.1}$$

where δ_x , $x \in V$, represents the Dirac measure concentrated at x. Thus, L_t is a random element of $\mathcal{P}(V)$ and $L_t(V_0)$, $V_0 \subset V$, stands for the average amount of time the process X_t stays at V_0 in the time interval [0, t].

As the Markov chain X_t is irreducible, by the ergodic theorem, for any starting point $x \in V$, as $t \to \infty$, the empirical measure L_t converges in probability to the stationary state π .

Denote by E the set of directed edges:

$$E := \{ (x, y) \in V \times V : y \neq x, R(x, y) > 0 \}.$$
 (1.2)

²⁰¹⁰ Mathematics Subject Classification. Primary 60K35, 60F10, 82C22.

Key words and phrases. Large deviations, empirical current, Γ-convergence, metastability.

and by $(Q_t: t \geq 0)$ the empirical flow defined by

$$Q_t(x,y) := \frac{1}{t} \sum_{0 \le s \le t} \mathbf{1} \{X_{s-} = x, X_s = y\}.$$

In words, $t Q_t(x, y)$ counts the number of times the process X jumped from x to y in the time interval [0, t]. By [7, Section 2.1], for any starting point $x \in V$, as $t \to \infty$, $Q_t(x, y)$ converges in probability to $\pi(x) R(x, y)$.

A function $J : E \to \mathbb{R}_+$ is called a *flow*. The set of flows defined on E is represented by \mathfrak{F}_E . We sometimes refer to a flow in \mathfrak{F}_E as an E-flow. The divergence of a flow J at a vertex $x \in V$, denoted by $(\operatorname{div} J)(x)$, is given by

$$(\operatorname{div} J)(x) = \sum_{y:(x,y)\in E} J(x,y) - \sum_{y:(y,x)\in E} J(y,x) .$$

A flow is said to be divergence-free if $(\operatorname{div} J)(x) = 0$ for all $x \in V$. Let $\mathfrak{F}_E^{\operatorname{div}}$ be the set of divergence-free E-flows.

Denote by $D(\mathbb{R}_+, W)$, W a finite set, the space of right-continuous functions $\mathfrak{f} \colon \mathbb{R}_+ \to W$ with left-limits endowed with the Skorohod topology and its associated Borel σ -algebra. Let \mathbf{P}_x , $x \in V$, be the probability measure on the path space $D(\mathbb{R}_+, V)$ induced by the Markov chain X_t starting from x. Expectation with respect to \mathbf{P}_x is represented by \mathbf{E}_x .

Let $\Phi \colon \mathbb{R}_+ \times \mathbb{R}_+ \to [0, +\infty]$ be the function defined by

$$\Phi(q,p) := \begin{cases}
p & \text{if } q = 0, \\
q \log \frac{q}{p} - (q - p) & \text{if } q \in (0, +\infty) \text{ and } p \in (0, +\infty), \\
+\infty & \text{if } q \in (0, +\infty) \text{ and } p = 0.
\end{cases} (1.3)$$

For p > 0, $\Phi(\cdot, p)$ is a nonnegative convex function which vanishes only at q = p. Actually, $\Phi(\cdot, p)$ is the large deviations rate functional of a Poisson process with parameter p.

Denote by $J_{\mu,R} \in \mathfrak{F}_E$, $\mu \in \mathfrak{P}(V)$, the flow defined by

$$J_{\mu B}(x,y) := \mu(x) R(x,y), \quad (x,y) \in E.$$
 (1.4)

Let $\Upsilon_{E,R} \colon \mathcal{P}(V) \times \mathfrak{F}_E \to [0,+\infty]$ be the functional defined by

$$\Upsilon_{E,R}(\mu, J) := \sum_{(x,y)\in E} \Phi(J(x,y), J_{\mu,R}(x,y)).$$
 (1.5)

Let $I: \mathcal{P}(V) \times \mathfrak{F}_E \to [0, +\infty]$ be the measure-current large deviations rate functional defined by

$$I(\mu, J) := \begin{cases} \Upsilon_{E,R}(\mu, J) & J \in \mathfrak{F}_E^{\text{div}}, \\ +\infty & \text{otherwise}. \end{cases}$$
 (1.6)

Bertini, Faggionato and Gabrielli [7] proved a large deviations principle for the pair (L_t, Q_t) . Mind that Condition 2.2 in [7] is trivially satisfied in the case where V is finite.

Theorem 1.1. For every closed set C of $\mathfrak{P}(V) \times E$, and every open set G of $\mathfrak{P}(V) \times E$,

$$\limsup_{t \to \infty} \sup_{x \in V} \frac{1}{t} \log \mathbf{P}_x \left[(L_t, Q_t) \in C \right] \le - \inf_{(\mu, J) \in C} I(\mu, J) ,$$

$$\liminf_{t \to \infty} \inf_{x \in V} \frac{1}{t} \log \mathbf{P}_x \left[(L_t, Q_t) \in G \right] \ge - \inf_{(\mu, J) \in G} I(\mu, J) .$$

Moreover, I is a convex rate function with compact level sets.

By [6, Theorem 1.6], the projection of the rate functional $I(\mu, J)$ on the first coordinate yields the Donsker-Varadhan large deviations functional [13] for the empirical measure:

$$\inf_{J \in \mathfrak{F}_E^{\text{div}}} I(\mu, J) = \sup_{u > 0} - \int \frac{\mathcal{L}u}{u} \, d\mu =: \mathfrak{I}(\mu) .$$

Γ-convergence. Suppose now that $(X_t^{(n)}: t \geq 0)$ is a sequence of V-valued, irreducible continuous-time Markov chains, whose jump rates are represented by $R_n(x,y)$. We add a subscript n to the previous notation to refer to the Markov chain $X^{(n)}$. In particular, the generator, the stationary state, and the rate functions are represented by \mathcal{L}_n , π_n , I_n , \mathcal{I}_n , respectively. We assume, however, that the set of directed edges E introduced in (1.2) does not depend on n: for all $n \geq 1$,

$$R_n(x,y) > 0$$
 if and only if $(x,y) \in E$. (1.7)

We assume furthermore that the jump rates satisfy condition (2.5).

In this article, we investigate the Γ -convergence of the measure-current large deviations rate functional I_n . Referring to [11] for an overview, we recall the definition of Γ -convergence. Fix a Polish space \mathcal{X} and a sequence $(U_n : n \in \mathbb{N})$ of functionals on \mathcal{X} , $U_n : \mathcal{X} \to [0, +\infty]$. The sequence $U_n \Gamma$ -converges to the functional $U : \mathcal{X} \to [0, +\infty]$, i.e. $U_n \xrightarrow{\Gamma} U$, if and only if the two following conditions are met:

- (i) Γ -liminf. The functional U is a Γ -liminf for the sequence U_n : For each $x \in \mathcal{X}$ and each sequence $x_n \to x$, we have that $\liminf_n U_n(x_n) \geq U(x)$.
- (ii) Γ -limsup. The functional U is a Γ -limsup for the sequence U_n : For each $x \in \mathcal{X}$ there exists a sequence $x_n \to x$ such that $\limsup_n U_n(x_n) \leq U(x)$.

The main result of the article provides a Γ -expansion of the rate functional I_n . It states that there exist sequences $(\theta_n^{(p)}: n \ge 1)$, $1 \le p \le \mathfrak{q}$, and large deviations rate functionals $\mathbf{I}^{(p)}: \mathcal{P}(V) \times \mathfrak{F}_E \to [0, +\infty], \ 0 \le p \le \mathfrak{q}$ such that $\theta_n^{(1)} \to \infty$, $\theta_n^{(p)}/\theta_n^{(p+1)} \to 0$, $1 \le p < \mathfrak{q}$, and $\theta_n^{(p)}I_n$ Γ -converges to $\mathbf{I}^{(p)}$. We summarize this result writing a Γ -expansion for I_n :

$$I_n = \mathbf{I}^{(0)} + \sum_{p=1}^{\mathfrak{q}} \frac{1}{\theta_n^{(p)}} \mathbf{I}^{(p)} .$$
 (1.8)

The rate function I_n thus encodes all the characteristics of the metastable behavior of the sequence of Markov chains $X_t^{(n)}$. The factors $\theta_n^{(p)}$ provide the time-scales at which it is observed and the rate functions $\mathbf{I}^{(p)}$ correspond to the generators describing the synthetic evolution.

Next result is a simple consequence of the large deviations principle stated in Theorem 1.1 and the Γ -convergence informally described in the previous paragraphs. (cf. Corollary 4.3 in [25]).

Corollary 1.2. Fix $0 \le p \le \mathfrak{q}$ and set $\theta_n^{(0)} = 1$. For every closed set C of $\mathfrak{P}(V) \times E$, and every open set G of $\mathfrak{P}(V) \times E$,

$$\limsup_{n\to\infty} \limsup_{t\to\infty} \frac{\theta_n^{(p)}}{t} \sup_{x\in V} \log \mathbf{P}_x^n \big[(L_t, Q_t) \in C \big] \leq -\inf_{(\mu, J)\in C} \mathbf{I}^{(p)}(\mu, J) ,$$

$$\liminf_{n\to\infty} \liminf_{t\to\infty} \frac{\theta_n^{(p)}}{t} \inf_{x\in V} \log \mathbf{P}_x^n \big[(L_t, Q_t) \in G \big] \ge - \inf_{(\mu, J)\in G} \mathbf{I}^{(p)}(\mu, J) .$$

2. Notation and results

In this section we state the main result of the article. This requires some notation. Denote by $\lambda_n(x)$, $x \in V$, the holding rates of the Markov chain $X_t^{(n)}$ and by $p_n(x,y)$, $x,y \in V$, the jump probabilities, so that $R_n(x,y) = \lambda_n(x) p_n(x,y)$. The generator reads therefore as

$$(\mathcal{L}_n f)(x) = \sum_{y \in V} R_n(x, y) \left\{ f(y) - f(x) \right\}, \quad f \colon V \to \mathbb{R}.$$

Let E' be a proper subset of E. Denote by $\mathfrak{F}_{E'}$ the set of flows in \mathfrak{F}_E such that J(x,y)=0 for all $(x,y)\in E\setminus E'$:

$$\mathfrak{F}_{E'} := \left\{ J \in \mathfrak{F}_E : J(x, y) = 0 \text{ for all } (x, y) \in E \setminus E' \right\}. \tag{2.1}$$

As before, denote by $\mathfrak{F}_{E'}^{\text{div}}$ the elements of $\mathfrak{F}_{E'}$ which are divergence-free.

The Γ -convergence. Assume that $\lim_n R_n(x,y)$ exists for all $(x,y) \in E$, and denote by $\mathbb{R}_0(x,y) \in [0,\infty)$ its limit:

$$\mathbb{R}_0(x,y) := \lim_n R_n(x,y) , \quad (x,y) \in E .$$
 (2.2)

Let \mathbb{E}_0 be the set of edges whose asymptotic rate is positive:

$$\mathbb{E}_0 := \{ (x, y) \in E : \mathbb{R}_0(x, y) > 0 \},$$

and assume that $\mathbb{E}_0 \neq \emptyset$. The jump rates $\mathbb{R}_0(x,y)$ induce a continuous-time Markov chain on V, denoted by $(\mathbb{X}_t : t \geq 0)$, which, of course, may be reducible. Denote by $\mathbb{L}^{(0)}$ its generator.

Denote by $\mathcal{V}_1, \dots, \mathcal{V}_n$, $n \geq 1$, the closed irreducible classes of \mathbb{X}_t , and let

$$S := \{1, \dots, \mathfrak{n}\}, \quad \mathcal{V} := \bigcup_{j \in S} \mathcal{V}_j, \quad \Delta := V \setminus \mathcal{V}.$$
 (2.3)

The set Δ may be empty and some of the sets \mathcal{V}_j may be singletons.

Let $\mathbf{I}^{(0)}: \mathcal{P}(V) \times \mathfrak{F}_E \to [0, +\infty]$ be the functional given by

$$\mathbf{I}^{(0)}(\mu, J) := \begin{cases} \Upsilon_{\mathbb{E}_0, \mathbb{R}_0}(\mu, J) & J \in \mathfrak{F}_{\mathbb{E}_0}^{\mathrm{div}}, \\ +\infty & \text{otherwise}, \end{cases}$$
 (2.4)

which is the measure-current large deviations rate functional of X_t .

Proposition 2.1. The functional I_n Γ -converges to $\mathbf{I}^{(0)}$.

We prove Proposition 2.1 in Section 3.

The main assumption. To examine the Γ -convergence of the measure-current large deviations rate functionals at longer time-scales, we introduce a natural hypothesis on the jump rates proposed in [2] and adopted in [8, 14, 21, 24].

For two sequences of positive real numbers $(\alpha_n : n \ge 1)$, $(\beta_n : n \ge 1)$, $\alpha_n \prec \beta_n$ or $\beta_n \succ \alpha_n$ means that $\lim_{n\to\infty} \alpha_n/\beta_n = 0$. Similarly, $\alpha_n \preceq \beta_n$ or $\beta_n \succeq \alpha_n$ indicates that either $\alpha_n \prec \beta_n$ or α_n/β_n converges to a positive real number $a \in (0, \infty)$.

Two sequences of positive real numbers $(\alpha_n : n \ge 1)$, $(\beta_n : n \ge 1)$ are said to be comparable if $\alpha_n \prec \beta_n$, $\beta_n \prec \alpha_n$ or $\alpha_n/\beta_n \to a \in (0, \infty)$. This condition excludes the possibility that $\liminf_n \alpha_n/\beta_n \ne \limsup_n \alpha_n/\beta_n$.

A set of sequences $(\alpha_n^{\mathfrak{u}}:n\geq 1)$, $\mathfrak{u}\in\mathfrak{R}$, of positive real numbers, indexed by some finite set \mathfrak{R} , is said to be comparable if for all $\mathfrak{u},\mathfrak{v}\in\mathfrak{R}$ the sequence $(\alpha_n^{\mathfrak{u}}:n\geq 1)$, $(\alpha_n^{\mathfrak{v}}:n\geq 1)$ are comparable.

Recall that we denote by E the set of directed edges (independent of n) with positive jump rates. Let $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$, and $\Sigma_m, m \geq 1$, be the set of functions $k \colon E \to \mathbb{Z}_+$ such that $\sum_{(x,y)\in E} k(x,y) = m$. We assume, hereafter, that for every $m \geq 1$ the set of sequences

$$\left(\prod_{(x,y)\in E} R_n(x,y)^{k(x,y)} : n \ge 1\right), \quad k \in \Sigma_m , \qquad (2.5)$$

is comparable. In Remark 2.4, we comment on this assumption.

Tree decomposition. If the Markov chain \mathbb{X}_t has only one closed irreducible class, the Γ-expansion of I_n has only one term, $\mathbf{I}^{(0)}$. Indeed, in this case, by Lemma 3.1, $\mathbf{I}^{(0)}(\mu, J) = 0$ implies that μ is the stationary state of the Markov chain \mathbb{X}_t and $J = J_{\mu,\mathbb{R}_0}$. In particular, as I_n converges to $\mathbf{I}^{(0)}$, for any sequence $\beta_n \to \infty$, $\beta_n I_n(\mu_n, J_n) \to \infty$ for any sequence (μ_n, J_n) converging to $(\mu, J) \neq (\pi, J_{\pi,\mathbb{R}_0})$ if π represents the stationary state of the Markov chain \mathbb{X}_t .

Assume therefore that there are more than one closed irreducible class, in other words, that the constant $\mathfrak n$ introduced in (2.3) is larger than or equal to 2: $\mathfrak n \geq 2$. Under this assumption, (1.7), and (2.5), [3,24] constructed a rooted tree which describes the behaviour of the Markov chain $X_t^{(n)}$ at all different time-scales. We recall the construction below.

The tree satisfies the following conditions:

- (a) Each vertex of the tree represents a subset of V;
- (b) Each generation forms a partition of V;
- (c) The children of each vertex form a partition of the parent;
- (d) The generation p+1 is strictly coarser than the generation p.

The tree is constructed by induction starting from the leaves to the root. It corresponds to a deterministic coalescence process. Denote by \mathfrak{q} the number of steps in the recursive construction of the tree. At each level $1 \leq p \leq \mathfrak{q}$, the procedure generates a partition $\{\mathcal{V}_1^{(p)},\ldots,\mathcal{V}_{\mathfrak{n}_p}^{(p)},\Delta_p\}$, a time-scale $\theta_n^{(p)}$ and a $\{1,\ldots,\mathfrak{n}_p\}$ -valued continuous-time Markov chain $\mathbb{X}_t^{(p)}$ which describes the evolution of the chain $X_{t\theta_n^{(p)}}^{(n)}$ among the subsets $\mathcal{V}_1^{(p)},\ldots,\mathcal{V}_{\mathfrak{n}_p}^{(p)}$, called hereafter wells.

among the subsets $\mathcal{V}_1^{(p)}, \dots, \mathcal{V}_{\mathfrak{n}_p}^{(p)}$, called hereafter wells. The leaves are the sets $\mathcal{V}_1, \dots, \mathcal{V}_{\mathfrak{n}}, \Delta$ introduced in (2.3). We proceed by induction. Let $S_1 = S$, $\mathfrak{n}_1 = \mathfrak{n}$, $\mathcal{V}_j^{(1)} = \mathcal{V}_j$, $j \in S_1$, $\Delta_1 = \Delta$, and assume that the recursion has produced the sets $\mathcal{V}_1^{(p)}, \dots, \mathcal{V}_{\mathfrak{n}_p}^{(p)}, \Delta_p$ for some $p \geq 1$, which forms a partition of V.

Denote by H_A , H_A^+ , $A \subset V$, the hitting and return time of A:

$$H_{\mathcal{A}} := \inf \{ t > 0 : X_t^{(n)} \in \mathcal{A} \}, \quad H_{\mathcal{A}}^+ := \inf \{ t > \tau_1 : X_t^{(n)} \in \mathcal{A} \}, \quad (2.6)$$

where τ_1 represents the time of the first jump of the chain $X_t^{(n)}$: $\tau_1 = \inf\{t > 0 : X_t^{(n)} \neq X_0^{(n)}\}.$

For two non-empty, disjoint subsets \mathcal{A} , \mathcal{B} of V, denote by $\operatorname{cap}_n(\mathcal{A}, \mathcal{B})$ the capacity between \mathcal{A} and \mathcal{B} :

$$\operatorname{cap}_{n}(\mathcal{A}, \mathcal{B}) := \sum_{x \in \mathcal{A}} \pi_{n}(x) \lambda_{n}(x) \mathbf{P}_{x}^{n} \left[H_{\mathcal{B}} < H_{\mathcal{A}}^{+} \right]. \tag{2.7}$$

Set $S_p = \{1, \dots, \mathfrak{n}_p\}$, and let $\theta_n^{(p)}$ be defined by

$$\frac{1}{\theta_n^{(p)}} := \sum_{i \in S_p} \frac{\text{cap}_n(\mathcal{V}_i^{(p)}, \check{\mathcal{V}}_i^{(p)})}{\pi_n(\mathcal{V}_i^{(p)})} , \quad \text{where } \check{\mathcal{V}}_i^{(p)} := \bigcup_{j \in S_p \setminus \{i\}} \mathcal{V}_j^{(p)} . \tag{2.8}$$

By [24, Assertion 8.B],

$$\theta_n^{(p-1)} \prec \theta_n^{(p)} \,. \tag{2.9}$$

The ratio $\pi_n(\mathcal{V}_i^{(p)})/\text{cap}_n(\mathcal{V}_i^{(p)}, \check{\mathcal{V}}_i^{(p)})$ represents the time it takes for the chain $X_t^{(n)}$, starting from a point in $\mathcal{V}_i^{(p)}$ to reach the set $\check{\mathcal{V}}_i^{(p)}$. Therefore, $\theta_n^{(p)}$ corresponds to the smallest time needed to observe such a jump.

Let $\Psi_p: V \to S_p \cup \{0\}$ be the projection which sends the points in $\mathcal{V}_j^{(p)}$ to j and the elements of Δ_p to 0:

$$\Psi_p \,:=\, \sum_{k\in S_n} k\; \chi_{{\boldsymbol{\mathcal{V}}}_k^{(p)}}\;.$$

In this formula and below, χ_A stands for the indicator function of the set A. Next theorem follows from the main result in [24] and [23].

Theorem 2.2. Assume that conditions (1.7), (2.5) are in force. Then, for each $j \in S_p$, $x \in \mathcal{V}_j^{(p)}$, under the measure \mathbf{P}_x^n , the finite-dimensional distributions of the sequence of $(S_p \cup \{0\})$ -valued processes $\Psi_p(X_{t\theta_n^{(p)}}^{(n)})$ converge to the finite-dimensional distributions of a S_p -valued Markov chain, represented by $\mathbb{X}_t^{(p)}$.

The process $\mathbb{X}_t^{(p)}$ describes therefore how the chain $X_t^{(n)}$ evolves among the wells $\mathcal{V}_j^{(p)}$ in the time-scale $\theta_n^{(p)}$. Note that the Markov chain $\mathbb{X}_t^{(p)}$ takes value in S_p , while the process $\Psi_p(X_{t\theta_s^{(p)}}^{(n)})$ may also be equal to 0.

Denote by $r^{(p)}(j,k)$ the jump rates of the S_p -valued continuous-time Markov chain $(\mathbb{X}_t^{(p)}:t\geq 0)$. By [24, Theorem 2.7], there exist $j, k\in S_p$ such that $r^{(p)}(j,k)>0$. Actually, by the proof of this result,

$$\sum_{k \neq j} r^{(p)}(j,k) > 0 \text{ for all } j \in S_p \text{ such that } \lim_{n \to \infty} \theta_n^{(p)} \frac{\text{cap}_n(\mathcal{V}_j^{(p)}, \check{\mathcal{V}}_j^{(p)})}{\pi_n(\mathcal{V}_j^{(p)})} > 0 . \quad (2.10)$$

Denote by $\mathfrak{R}_1^{(p)},\ldots,\mathfrak{R}_{\mathfrak{n}_{p+1}}^{(p)}$ the recurrent classes of the S_p -valued chain $\mathbb{X}_t^{(p)}$, and by \mathfrak{T}_p the transient states. Let $\mathfrak{R}^{(p)}=\cup_j\mathfrak{R}_j^{(p)}$, and observe that $\{\mathfrak{R}_1^{(p)},\ldots,\mathfrak{R}_{\mathfrak{n}_{p+1}}^{(p)},\mathfrak{T}_p\}$

forms a partition of the set S_p . This partition of S_p induces a new partition of the set V. Let

$$\mathcal{V}_{m}^{(p+1)} := \bigcup_{j \in \mathfrak{R}_{m}^{(p)}} \mathcal{V}_{j}^{(p)} , \quad \mathfrak{T}^{(p+1)} := \bigcup_{j \in \mathfrak{T}_{p}} \mathcal{V}_{j}^{(p)} , \quad m \in S_{p+1} := \{1, \dots, \mathfrak{n}_{p+1}\} ,$$

so that $V = \Delta_{p+1} \cup \mathcal{V}^{(p+1)}$, where

$$\mathcal{V}^{(p+1)} = \bigcup_{m \in S_{p+1}} \mathcal{V}_m^{(p+1)} , \quad \Delta_{p+1} := \Delta_p \cup \mathcal{T}^{(p+1)} . \tag{2.11}$$

The subsets $\mathcal{V}_1^{(p+1)}, \ldots, \mathcal{V}_{\mathfrak{n}_{p+1}}^{(p+1)}, \Delta_{p+1}$ of V are the result of the recursive procedure. We claim that conditions (a)–(d) hold at step p+1 if they are fulfilled up to step p in the induction argument.

The sets $\mathcal{V}_1^{(p+1)}, \ldots, \mathcal{V}_{\mathfrak{n}_{p+1}}^{(p+1)}, \Delta_{p+1}$ constitute a partition of V because the sets $\mathfrak{R}_1^{(p)}, \ldots, \mathfrak{R}_{\mathfrak{n}_{p+1}}^{(p)}, \mathfrak{T}_p$ form a partition of S_p , and the sets $\mathcal{V}_1^{(p)}, \ldots, \mathcal{V}_{\mathfrak{n}_p}^{(p)}, \Delta_p$ one of V. Conditions (a)–(c) are therefore satisfied.

To show that the partition obtained at step p+1 is strictly coarser than $\{\mathcal{V}_1^{(p)},\ldots,\mathcal{V}_{\mathfrak{n}_p}^{(p)},\Delta_p\}$, observe that, by (2.10), $r^{(p)}(j,k)>0$ for some $k\neq j\in S_p$. Hence, either j is a transient state for the process $\mathbb{X}_t^{(p)}$ or the closed recurrent class which contains j also contains k. In the first case $\Delta_p \subseteq \Delta_{p+1}$, and in the second one there exists $m\in S_{p+1}$ such that $\mathcal{V}_j^{(p)}\cup\mathcal{V}_k^{(p)}\subset\mathcal{V}_m^{(p+1)}$. Therefore, the new partition $\{\mathcal{V}_1^{(p+1)},\ldots,\mathcal{V}_{\mathfrak{n}_{p+1}}^{(p+1)},\Delta_{p+1}\}$ of V satisfies the condition (d).

The construction terminates when the S_p -valued Markov chain $\mathbb{X}_t^{(p)}$ has only one recurrent class so that $\mathfrak{n}_{p+1} = 1$. In this situation, the partition at step p+1 is $\mathcal{V}_1^{(p+1)}$, Δ_{p+1} .

This completes the construction of the rooted tree. Recall that we denote by \mathfrak{q} the number of steps of the scheme. As claimed at the beginning of the procedure, for each $1 \leq p \leq \mathfrak{q}$, we generated a time-scale $\theta_n^{(p)}$, a partition $\mathcal{P}_p = \{\mathcal{V}_1^{(p)}, \dots, \mathcal{V}_{\mathfrak{n}_p}^{(p)}, \Delta_p\}$, where $\mathcal{P}_1 = \{\mathcal{V}_1, \dots, \mathcal{V}_{\mathfrak{n}}, \Delta\}$, $\mathcal{P}_{\mathfrak{q}+1} = \{\mathcal{V}_1^{(\mathfrak{q}+1)}, \Delta_{\mathfrak{q}+1}\}$, and a S_p -valued continuous-time Markov chain $\mathbb{X}_t^{(p)}$.

The partitions $\mathcal{P}_1, \ldots, \mathcal{P}_{\mathfrak{q}+1}$ form a rooted tree whose root (0-th generation) is V, first generation is $\{\mathcal{V}_1^{(\mathfrak{q}+1)}, \Delta_{\mathfrak{q}+1}\}$ and last $((\mathfrak{q}+1)$ -th) generation is $\{\mathcal{V}_1, \ldots, \mathcal{V}_{\mathfrak{n}}, \Delta\}$. Note that the set $\mathcal{V}^{(p+1)}$ corresponds to the set of recurrent points for the chain $\mathbb{X}_t^{(p)}$. In contrast, the points in Δ_{p+1} are either transient for this chain or negligible in the sense that the chain $X_t^{(n)}$ remains a negligible amount of time on the set Δ_p in the time-scale $\theta_p^{(p)}$ (cf. [2,24]).

A set of measures. We construct in this subsection a set of probability measures $\pi_j^{(p)}$, $1 \leq p \leq \mathfrak{q} + 1$, $j \in S_p$, on V which describe the evolution of the chain $X_t^{(n)}$ and such that

the support of
$$\pi_j^{(p)}$$
 is the set $\mathcal{V}_j^{(p)}$. (2.12)

We proceed by induction. Let $\pi_j^{(1)}$, $j \in S_1$, be the stationary states of the Markov chain \mathbb{X}_t restricted to the closed irreducible classes $\mathcal{V}_j^{(1)} = \mathcal{V}_j$. Clearly, condition (2.12) is fulfilled.

Fix $1 \leq p \leq \mathfrak{q}$, and assume that the probability measures $\pi_j^{(p)}$, $j \in S_p$, have been defined and satisfy condition (2.12). Denote by $M_m^{(p)}(\cdot)$, $m \in S_{p+1}$, the stationary state of the Markov chain $\mathbb{X}_t^{(p)}$ restricted to the closed irreducible class $\mathfrak{R}_m^{(p)}$. The measure $M_m^{(p)}$ is understood as a measure on $S_p = \{1, \ldots, \mathfrak{n}_p\}$ which vanishes on the complement of $\mathfrak{R}_m^{(p)}$. Let $\pi_m^{(p+1)}$ be the probability measure on V given by

$$\pi_m^{(p+1)}(x) := \sum_{j \in \mathfrak{R}_m^{(p)}} M_m^{(p)}(j) \, \pi_j^{(p)}(x) \;, \quad x \in V \;. \tag{2.13}$$

Clearly, condition (2.12) holds, and the measure $\pi_m^{(p+1)}$, $1 \leq p \leq \mathfrak{q}$, $m \in S_{p+1}$, is a convex combination of the measures $\pi_j^{(p)}$, $j \in \mathfrak{R}_m^{(p)}$. Moreover, by [8, Theorem 3.1 and Proposition 3.2], for all $z \in \mathcal{V}_i^{(p)}$,

$$\lim_{n \to \infty} \frac{\pi_n(z)}{\pi_n(\mathcal{V}_i^{(p)})} = \pi_j^{(p)}(z) \in (0,1] , \quad \lim_{n \to \infty} \pi_n(\Delta_{\mathfrak{q}+1}) = 0 . \tag{2.14}$$

By (2.13), the measures $\pi_j^{(p)}$, $2 \leq p \leq \mathfrak{q} + 1$, $j \in S_p$, are convex combinations of the measures $\pi_k^{(1)}$, $k \in S_1$. By (2.14), for all $x \in \mathcal{V}^{(\mathfrak{q}+1)}$, $\lim_{n \to \infty} \pi_n(x)$ exists and belongs to (0,1]. By (2.14), and since by (1.c) $\Delta_p \subset \Delta_{p+1}$ for $1 \leq p \leq \mathfrak{q}$, $\lim_{n \to \infty} \pi_n(\Delta_p) = 0$ for all p.

The Γ -expansion. We are now in a position to state the main result of this article. Let $\mathbb{L}^{(p)}$, $1 \leq p \leq \mathfrak{q}$, be the generator of the S_p -valued Markov chain $\mathbb{X}_t^{(p)}$. Denote by $\mathbb{P}(S_p)$, $1 \leq p \leq \mathfrak{q}$, the set of probability measures on S_p . Let $\mathbb{I}^{(p)} : \mathbb{P}(S_p) \to [0, +\infty]$ be the Donsker-Varadhan large deviations rate functional of $\mathbb{X}_t^{(p)}$ given by

$$\mathbb{I}^{(p)}(\omega) := \sup_{\mathbf{h}} - \sum_{j \in S_p} \omega_j e^{-\mathbf{h}(j)} \left(\mathbb{L}^{(p)} e^{\mathbf{h}} \right) (j) , \qquad (2.15)$$

where the supremum is carried over all functions $\mathbf{h}: S_p \to \mathbb{R}$. Denote by $\mathbf{I}^{(p)}: \mathcal{P}(V) \times \mathfrak{F}_E \to [0, +\infty]$ the functional given by

$$\mathbf{I}^{(p)}(\mu, J) := \begin{cases} \mathbb{I}^{(p)}(\omega) & \text{if } \mu = \sum_{j \in S_p} \omega_j \, \pi_j^{(p)} & \text{for } \omega \in \mathcal{P}(S_p) \text{ and } J = J_{\mu, \mathbb{R}_0} ,\\ + \infty & \text{otherwise} . \end{cases}$$

$$(2.16)$$

The main result of the article reads as follows.

Theorem 2.3. For each $1 \le p \le \mathfrak{q}$, the functional $\theta_n^{(p)}I_n$ Γ -converges to $\mathbf{I}^{(p)}$.

We complete this section with some comments on condition (2.5).

Remark 2.4. The hypothesis (2.5) on the jump rates is taken from [2] and [24]. It is a natural condition in the investigation of the asymptotic behavior of sequences of Markov chains. Indeed, a first reasonable hypothesis to impose consists in assuming that the jump rates converge, as stated in (2.2).

In longer time-scales, the process is expected to remain long times in a set of wells and to perform very short excursions among the points which separate the wells. It is therefore also natural to require the jump rates of the trace process to converge. (We refer to [1, Section 6], [20] for the definition of the trace process.)

Fix $V_0 \subset V$, and denote by $R_n^{V_0}$ the jump rates of the trace in $V \setminus V_0$ of the Markov chain $X_t^{(n)}$. According to the displayed equation after Corollary 6.2 in [1], $R_n^{V_0}$ can be expressed as a sum of products of terms of the form

$$\frac{R_n(x_0, x_1) \cdots R_n(x_{m-1}, x_m)}{\sum_{b=1}^q R_n(y_0^b, y_1^b) \cdots R_n(y_{m-1}^b, y_m^b)}$$
(2.17)

for some $m \ge 1$. Here $x_i, y_j^b \in V$. Mind that the number of terms in each product, m, is always the same.

Condition (2.5) is precisely the one needed to guarantee that such expressions have a limit (which might be $+\infty$) and do not oscillate.

Finally, as observed in [2] (see Remark 2.2 in [8]), assumption (2.5) is fulfilled by all statistical mechanics models which evolve on a fixed finite state space and whose metastable behaviour has been derived. This includes the Ising model [5,10,27,28], the Potts model with or without a small external field [18,19,26], the Blume-Capel model [12,22], and conservative Kawasaki dynamics [4,9,16,17].

3. The first time-scale

In this section, we investigate the first time-scale. In detail, we prove Proposition 2.1 and identify the set of zeros of the functional $\mathbf{I}^{(0)}$.

A flow $J: E \to \mathbb{R}_+$ is called a *cycle* if there exists a set of distinct edges $(x_0, x_1), \ldots, (x_{n-1}, x_n)$ in E and a constant $a \neq 0$ such that $x_n = x_0$ and

$$J(x,y) = \begin{cases} a & \text{if } (x,y) = (x_j, x_{j+1}) \text{ for some } 0 \le j < n, \\ 0 & \text{otherwise}. \end{cases}$$

Clearly, every cycle is divergence-free and every divergence-free flow can be expressed as a finite sum of cycles. To prove the latter argument, it suffices to observe the following. Given any $J \in \mathfrak{F}_E^{\mathrm{div}}$, take $(x,y) \in E$ such that

$$J(x,y) \, = \, \min \, \left\{ \, J(z,w) : (z,w) \in E, \ \, J(z,w) > 0 \, \right\} \, , \tag{3.1} \label{eq:3.1}$$

and find a set of distinct edges $(x_0, x_1), \ldots, (x_{n-1}, x_n)$ with $(x_0, x_1) = (x, y)$ and $x_n = x_0$ such that $J(x_j, x_{j+1}) > 0$ for each $0 \le j < n$, which is possible since J is divergence-free. Define a cycle J_0 which has value J(x, y) along the n edges. Then, by the minimality in $(3.1), J - J_0$ is again a divergence-free flow which has strictly less number of edges with a positive value. Iterating this procedure, which ends in finite steps, we obtain the desired decomposition as a sum of cycles.

Proof of Proposition 2.1. We first consider the Γ – \limsup . Fix a pair (μ, J) in $\mathfrak{P}(V) \times \mathfrak{F}_E$. We may assume that $\mathbf{I}^{(0)}(\mu, J) < \infty$, otherwise there is nothing to prove. Under this restriction, J belongs to $\mathfrak{F}_{\mathbb{E}_0}^{\mathrm{div}}$, thus to $\mathfrak{F}_E^{\mathrm{div}}$.

Let (μ_n, J_n) be the sequence constant equal to (μ, J) . Since J belongs to $\mathfrak{F}_E^{\text{div}}$, by (1.5) and (1.6),

$$I_n(\mu, J) = \Upsilon_{E,R_n}(\mu, J) = \sum_{(x,y)\in E} \Phi(J(x,y), J_{\mu,R_n}(x,y)).$$

It remains to show that

$$\lim_{n \to \infty} \Phi(J(x,y), J_{\mu,R_n}(x,y)) = \Phi(J(x,y), J_{\mu,\mathbb{R}_0}(x,y)) \quad \text{for all } (x,y) \in E.$$
(3.2)

Suppose that J(x,y)=0. By (1.3), $\Phi(0,\cdot)$ is continuous. Thus, by (2.2) and the definition (1.4) of the flow $J_{\mu,R_n}(x,y)$, (3.2) holds for edges $(x,y) \in E$ such that J(x,y)=0. On the other hand, if J(x,y)>0, since $\mathbf{I}^{(0)}(\mu,J)<\infty$, by (1.3), $\mu(x)\mathbb{R}_0(x,y)=J_{\mu,\mathbb{R}_0}(x,y)>0$. Thus, by (2.2), (3.2) also holds in this case. This completes the proof of the Γ -limsup. Note that we proved that

$$\lim_{n\to\infty} I_n(\mu,J) = \mathbf{I}^{(0)}(\mu,J) .$$

We turn to the Γ -lim inf. Fix a pair (μ, J) in $\mathcal{P}(V) \times \mathfrak{F}_E$ and a sequence (μ_n, J_n) in $\mathcal{P}(V) \times \mathfrak{F}_E$ converging to (μ, J) . If J is not divergence-free, for n sufficiently large J_n is not divergence-free either and for those n's $I_n(\mu_n, J_n) = \mathbf{I}^{(0)}(\mu, J) = \infty$.

Assume that J is divergence-free and that J(x,y) > 0 for some $(x,y) \notin \mathbb{E}_0$. In this case $\mathbf{I}^{(0)}(\mu, J) = \infty$ (because $J_{\mu,\mathbb{R}_0}(x,y) = \mu(x) \mathbb{R}_0(x,y) = 0$ so that, in view of (1.3), $\Phi(J(x,y), J_{\mu,\mathbb{R}_0}(x,y)) = +\infty$). On the other hand, since Φ is positive, $J_n(x,y) \to J(x,y) > 0$ and $J_{\mu_n,R_n}(x,y) \le R_n(x,y) \to 0$,

$$\liminf_{n\to\infty} I_n(\mu_n, J_n) \geq \liminf_{n\to\infty} \Phi(J_n(x, y), J_{\mu_n, R_n}(x, y)) = \infty = \mathbf{I}^{(0)}(\mu, J).$$

The previous arguments show that we may restrict our attention to divergencefree flows that vanish on edges which do not belong to \mathbb{E}_0 , that is to flows in $\mathfrak{F}_{\mathbb{E}_0}^{\text{div}}$. Assume that J belongs to this set. Since Φ is positive,

$$\liminf_{n \to \infty} I_n(\mu_n, J_n) \geq \liminf_{n \to \infty} \sum_{(x,y) \in \mathbb{E}_0} \Phi(J_n(x,y), J_{\mu_n, R_n}(x,y)).$$

Fix $(x,y) \in \mathbb{E}_0$. We consider three cases. If $\mu(x) > 0$, $\Phi(J_n(x,y), J_{\mu_n,R_n}(x,y))$ converges to $\Phi(J(x,y), J_{\mu,\mathbb{R}_0}(x,y))$. If $\mu(x) = 0$ and J(x,y) = 0, then, as Φ is positive,

$$\liminf_{n\to\infty}\Phi\big(\,J_n(x,y)\,,\,J_{\mu_n,R_n}(x,y)\,\big)\,\geq\,0\,=\,\Phi\big(\,J(x,y)\,,\,J_{\mu,\mathbb{R}_0}(x,y)\,\big)\;.$$

Finally, if $\mu(x) = 0$ and J(x,y) > 0, then $\liminf_{n \to \infty} \Phi(J_n(x,y), J_{\mu_n,R_n}(x,y)) = \infty = \Phi(J(x,y), J_{\mu,R_n}(x,y))$. This completes the proof of the proposition.

The functional $\mathbf{I}^{(0)}$. The set of divergence-free flows in $\mathfrak{F}^{\mathrm{div}}_{\mathbb{E}_0}$ has a simple structure. Denote by $\mathbb{E}_{0,j}$, $1 \leq j \leq \mathfrak{n}$, the set of directed edges in \mathbb{E}_0 whose endpoints belong to \mathcal{V}_j :

$$\mathbb{E}_{0,j} := \left\{ (x,y) \in \mathbb{E}_0 : x, y \in \mathcal{V}_j \right\}.$$

We say that y is equivalent to x, $y \sim x$ if y = x or if there exist sequences $x = x_0, \ldots, x_\ell = y$ and $y = y_0, \ldots, y_m = x$ such that $\mathbb{R}_0(x_i, x_{i+1}) > 0$, $\mathbb{R}_0(y_j, y_{j+1}) > 0$ for all $0 \le i < \ell$, $0 \le j < m$. Denote by $\mathcal{C}_1, \ldots, \mathcal{C}_m$ the equivalent classes which are not one of the sets \mathcal{V}_j , $1 \le j \le \mathfrak{n}$. Then, denote by $\mathbb{E}_{0,k}^{\mathrm{Tr}}$, $1 \le k \le \mathfrak{m}$, the set of directed edges in \mathbb{E}_0 whose endpoints belong to \mathcal{C}_k :

$$\mathbb{E}_{0,k}^{\mathrm{Tr}} := \left\{ (x,y) \in \mathbb{E}_0 : x, y \in \mathcal{C}_k \right\}.$$

We claim that any \mathbb{E}_0 -divergence-free flow is a sum of $\mathbb{E}_{0,j}$ - or $\mathbb{E}_{0,k}^{\mathrm{Tr}}$ -divergence-free flows (that is, flows whose edges belong to $\mathbb{E}_{0,j}$ or $\mathbb{E}_{0,k}^{\mathrm{Tr}}$, respectively). To see this, recall that $J \in \mathfrak{F}_{\mathbb{E}_0}^{\mathrm{div}}$ implies that

$$\sum_{y:(x,y)\in\mathbb{E}_0}J(x,y)=\sum_{y:(y,x)\in\mathbb{E}_0}J(y,x)\quad\text{for all }x\in V\;.$$

Adding this for all $x \in \mathcal{V}_j$ for fixed j, all edges in $\mathbb{E}_{0,j}$ cancel out with each other and we obtain that

$$\sum_{x \in \mathcal{V}_j} \sum_{y \notin \mathcal{V}_j : (x,y) \in \mathbb{E}_0} J(x,y) = \sum_{x \in \mathcal{V}_j} \sum_{y \notin \mathcal{V}_j : (y,x) \in \mathbb{E}_0} J(y,x) .$$

Since V_j is an irreducible component, the left-hand side is zero. Thus, we must have J(y,x)=0 for all $(y,x)\in\mathbb{E}_0$ such that $x\in\mathcal{V}_j$ and $y\notin\mathcal{V}_j$. Applying the same idea to the collection \mathcal{C}_k , we also obtain that J(y,x)=0 for all $(y,x)\in\mathbb{E}_0$ such that $x\in\mathcal{C}_k$ and $y\notin\mathcal{C}_k$. These two observations imply that each $J\in\mathfrak{F}_{\mathbb{E}_0}^{\mathrm{div}}$ is allowed to be positive only on the edges in $\mathbb{E}_{0,j}$ or $\mathbb{E}_{0,k}^{\mathrm{Tr}}$, as desired.

Now, we are ready to characterize the zeros of the functional $\mathbf{I}^{(0)}$. Recall from the definition after (2.12) that $\pi_j^{(1)}$, $1 \leq j \leq \mathfrak{n}$ denotes the invariant probability measure on \mathcal{V}_j for the \mathbb{R}_0 -chain.

Lemma 3.1. Fix a probability measure μ in $\mathfrak{P}(V)$ and a flow J in $\mathfrak{F}_{\mathbb{E}_0}^{\mathrm{div}}$. Then, $\mathbf{I}^{(0)}(\mu, J) = 0$ if and only if there exist weights ω_j , $1 \leq j \leq \mathfrak{n}$, such that $\omega_j \geq 0$, $\sum_j \omega_j = 1$, $\mu = \sum_j \omega_j \pi_j^{(1)}$, and $J = J_{\mu,\mathbb{R}_0}$.

Proof. First, assume that $\mathbf{I}^{(0)}(\mu, J) = 0$ and recall formula (2.4). Since Φ is positive, $\Phi(J(x,y), J_{\mu,\mathbb{R}_0}(x,y)) = 0$ for all $(x,y) \in \mathbb{E}_0$. As $\Phi(q,p) = 0$ if, and only if, q = p, $J(x,y) = J_{\mu,\mathbb{R}_0}(x,y)$ for all $(x,y) \in \mathbb{E}_0$. As J is a divergence-free $\mathbb{E}_{0,j}$ -flow when restricted to each \mathcal{V}_j , so is J_{μ,\mathbb{R}_0} . This implies that μ is a stationary state for the \mathcal{V} -valued (cf. (2.3)) Markov chain with jump rates \mathbb{R}_0 . Moreover, since each \mathcal{C}_k is a transient collection, there exists at least one $x \in \mathcal{C}_k$ such that $\mathbb{R}_0(x,y) > 0$ for some $y \notin \mathcal{C}_k$. As \mathcal{C}_k is a transient class and $x \in \mathcal{C}_k$, $y \notin \mathcal{C}_k$, and $J \in \mathfrak{F}_{\mathbb{E}_0}^{\mathrm{div}}$, J(x,y) has to vanish. Since J(x,y) = 0, this implies that $\mu(x) = 0$, which then implies that $\mu = 0$ on the whole \mathcal{C}_k . In summary, μ is a convex combination of only the ergodic measures on \mathcal{V} , that are, $\pi_j^{(1)}$, $1 \le j \le \mathfrak{n}$, as claimed.

Finally, suppose that $\mu = \sum_{j} \omega_{j} \pi_{j}^{(1)}$ for some probability measure ω on $S_{1} = \{1, \ldots, \mathfrak{n}\}$ and $J = J_{\mu, \mathbb{R}_{0}}$. By (2.4),

$$\mathbf{I}^{(0)}(\mu, J) \, = \, \sum_{(x,y) \in \mathbb{E}_0} \Phi \big(\, J(x,y) \, , \, J_{\mu,\mathbb{R}_0}(x,y) \, \big) = \sum_{(x,y) \in \mathbb{E}_0} \mu(x) \, \mathbb{R}_0(x,y) \, ,$$

where the second equality follows from (1.3) and (1.5). Since μ vanishes outside \mathcal{V} and \mathbb{R}_0 is zero going out from \mathcal{V} , the last summation equals

$$\sum_{j=1}^{\mathfrak{n}} \sum_{(x,y) \in \mathbb{E}_{0,j}} \mu(x) \, \mathbb{R}_0(x,y) \; .$$

The summation in $(x, y) \in \mathbb{E}_{0,j}$ is zero for all j since μ is stationary in each \mathcal{V}_j with respect to \mathbb{X}_t . This concludes the proof of Lemma 3.1.

4. Longer time-scales

Recall from (2.15) that $\mathbb{I}^{(p)}$, $1 \leq p \leq \mathfrak{q}$ denotes the Donsker-Varadhan large deviations rate functional of $\mathbb{X}_t^{(p)}$. Then (2.16) implies that, for all $\mu \in \mathcal{P}(V)$,

$$\mathbb{I}^{(p)}(\mu) = \inf_{J \in \mathfrak{F}_E} \mathbf{I}^{(p)}(\mu, J) . \tag{4.1}$$

We start with a lemma which states that the functionals $\mathbf{I}^{(p)}$, $1 \leq p \leq \mathfrak{q}$, form a hierarchical structure of zeros.

Lemma 4.1. For each $1 \le p \le \mathfrak{q}$, $\mathbf{I}^{(p)}(\mu, J) < \infty$ if and only if $\mathbf{I}^{(p-1)}(\mu, J) = 0$.

Proof. Suppose first that $\mathbf{I}^{(p)}(\mu, J) < \infty$, so that by (2.16), $\mu = \sum_{j=1}^{\mathfrak{n}_p} \omega_j \, \pi_j^{(p)}$ and $J = J_{\mu,\mathbb{R}_0}$. Then, $\mathbb{I}^{(p)}(\mu) = \mathbf{I}^{(p)}(\mu, J_{\mu,\mathbb{R}_0}) < \infty$, thus by [21, Lemma 5.1 and Eq. (5.1)], we have $\mathbf{I}^{(p-1)}(\mu, J_{\mu,\mathbb{R}_0}) = \mathbb{I}^{(p-1)}(\mu) = 0$ for $p \geq 2$. If p = 1, we also have $\mathbf{I}^{(0)}(\mu, J_{\mu,\mathbb{R}_0}) = 0$ by Lemma 3.1.

Next, suppose that $\mathbf{I}^{(p-1)}(\mu, J) = 0$. If p = 1, then by Lemma 3.1 we have $\mu = \sum_{j=1}^{\mathfrak{n}} \omega_j \, \pi_j^{(1)}$ and $J = J_{\mu,\mathbb{R}_0}$, so that $\mathbf{I}^{(1)}(\mu, J) = \mathbb{I}^{(1)}(\mu) < \infty$ by (2.16) and [21, Eq. (2.18)]. If $p \geq 2$, then by (2.16), $\mu = \sum_{j=1}^{\mathfrak{n}_{p-1}} \omega_j \, \pi_j^{(p-1)}$, $J = J_{\mu,\mathbb{R}_0}$, and moreover $\mathbb{I}^{(p-1)}(\mu) = 0$. Via [21, Lemma 5.1], this implies that $\mathbb{I}^{(p)}(\mu) < \infty$, thus again by (2.16), $\mathbf{I}^{(p)}(\mu, J) = \mathbb{I}^{(p)}(\mu) < \infty$. This finishes the verification of Lemma 4.1.

Now, we provide the proof of Theorem 2.3.

Proof of Theorem 2.3. We proceed by induction. Suppose that the statement holds for p-1, where the 0-th step holds by Proposition 2.1.

Denote by $\mathfrak{I}_n: \mathfrak{P}(V) \to [0,+\infty]$ the Donsker-Varadhan large deviations rate functional of the original process $X_t^{(n)}$. Recall from [21, Theorem 2.5] that $\theta_n^{(p)} \mathfrak{I}_n$ Γ -converges to $\mathbb{I}^{(p)}$.

First, we consider the Γ – \limsup . Fix $(\mu, J) \in \mathcal{P}(V) \times \mathfrak{F}_E$. We may assume that $\mu = \sum_{j=1}^{\mathfrak{n}_p} \omega_j \, \pi_j^{(p)}$ and $J = J_{\mu,\mathbb{R}_0}$ since otherwise there is nothing to prove. By the Γ -convergence $\theta_n^{(p)} \mathfrak{I}_n \to \mathbb{I}^{(p)}$, there exists a sequence of probability measures $(\nu_n : n \geq 1)$ on V that converges to μ and satisfies

$$\limsup_{n \to \infty} \, \theta_n^{(p)} \, \mathfrak{I}_n(\nu_n) \, \leq \, \mathbb{I}^{(p)}(\mu) \, \, .$$

By [6, Theorem 1.6], there exists a unique flow $J_n^* \in \mathfrak{F}_E$ such that

$$\mathfrak{I}_n(\nu_n) = I_n(\nu_n, J_n^*) \,. \tag{4.2}$$

Then by (2.16), we have

$$\limsup_{n \to \infty} \theta_n^{(p)} I_n(\nu_n, J_n^*) \le \mathbb{I}^{(p)}(\mu) = \mathbf{I}^{(p)}(\mu, J_{\mu, \mathbb{R}_0}).$$

To conclude the Γ – \limsup part, it remains to prove that $\lim_{n\to\infty} J_n^* = J_{\mu,\mathbb{R}_0}$. By [6, Eq. (2.9)], there exists a function $g_n \colon V \to \mathbb{R}$ such that

$$\log \frac{J_n^*(x,y)}{J_{\nu_n,R_n}(x,y)} = g_n(y) - g_n(x) \quad \text{for each } (x,y) \in E.$$
 (4.3)

Then by the definition of I_n , (4.2), and [6, Eq. (2.17)], we can calculate

$$\mathfrak{I}_{n}(\nu_{n}) = \sum_{(x,y)\in E} \Upsilon_{E,R_{n}}(\nu_{n}, J_{n}^{*}) = \sum_{(x,y)\in E} \nu_{n}(x) R_{n}(x,y) \left(1 - e^{g_{n}(y) - g_{n}(x)}\right).$$

Thus, by [21, Lemma A.3], \mathcal{I}_n and g_n satisfy

$$J_n(\nu_n) = \sum_{(x,y)\in E} \nu_n(x) R_n(x,y) \left((g_n(y) - g_n(x)) e^{g_n(y) - g_n(x)} - e^{g_n(y) - g_n(x)} + 1 \right)$$

where the right-hand side is a sum of nonnegative terms. Since $\nu_n \to \mu$ which is a zero of $\mathbb{I}^{(0)}$ by Lemma 3.1, we have $\lim_n \mathfrak{I}_n(\nu_n) = \mathbb{I}^{(0)}(\mu) = 0$. Hence, for each

 $(x,y) \in E$, the summand in the right-hand side of (4.4) vanishes as $n \to \infty$. Finally, since $|1-e^a| \le ae^a - a + 1$ for all $a \in \mathbb{R}$, we obtain that

$$\lim_{n \to \infty} \nu_n(x) R_n(x,y) | 1 - e^{g_n(y) - g_n(x)} | = 0.$$

Via (4.3) and the fact that $\nu_n(x) R_n(x,y) \to \mu(x) \mathbb{R}_0(x,y)$, the displayed identity

concludes the proof of $\lim_n J_n^* = J_{\mu,\mathbb{R}_0}$, thus of the $\Gamma - \limsup$ part. Next, we consider the $\Gamma - \liminf$. Fix $(\mu, J) \in \mathcal{P}(V) \times \mathfrak{F}_E$ and an arbitrary sequence $(\mu_n, J_n) \in \mathcal{P}(V) \times \mathfrak{F}_E$, $n \geq 1$, that converges to (μ, J) . We have

$$\liminf_{n \to \infty} \theta_n^{(p)} \, \mathfrak{I}_n(\mu_n) \, \ge \, \mathbb{I}^{(p)}(\mu) \, . \tag{4.5}$$

If $\mathbf{I}^{(p)}(\mu, J) < \infty$, then we easily obtain that

$$\liminf_{n\to\infty} \theta_n^{(p)} I_n(\mu_n, J_n) \ge \liminf_{n\to\infty} \theta_n^{(p)} \mathfrak{I}_n(\mu_n) \ge \mathbb{I}^{(p)}(\mu) = \mathbf{I}^{(p)}(\mu, J) ,$$

where the first inequality holds by (4.1), the second inequality holds by (4.5), and the equality holds by (2.16). Finally, suppose that $\mathbf{I}^{(p)}(\mu, J) = \infty$. Then by Lemma 4.1, $\mathbf{I}^{(p-1)}(\mu, J) > 0$. By the induction hypothesis at level p-1,

$$\liminf_{n \to \infty} \theta_n^{(p-1)} I_n(\mu_n, J_n) \ge \mathbf{I}^{(p-1)}(\mu, J) > 0.$$

Since $\theta_n^{(p-1)} \prec \theta_n^{(p)}$ by (2.9), we conclude that

$$\liminf_{n \to \infty} \theta_n^{(p)} I_n(\mu_n, J_n) = \infty = \mathbf{I}^{(p)}(\mu, J) ,$$

finishing the proof of Theorem 2.3.

Acknowledgement. C. L. has been partially supported by FAPERJ CNE E-26/201.117/2021, by CNPq Bolsa de Produtividade em Pesquisa PQ 305779/2022-2. S. K. has been supported by the KIAS Individual Grant (HP095101) at the Korea Institute for Advanced Study. S. K. would like to thank IMPA (Rio de Janeiro) for the warm hospitality during his stay in July 2024.

References

- [1] J. Beltrán, C. Landim: Tunneling and metastability of continuous time Markov chains. J. Stat. Phys. **140** 1065–1114 (2010).
- [2] J. Beltrán, C. Landim; Metastability of reversible finite state Markov processes. Stoch. Proc. Appl. **121** 1633–1677 (2011).
- [3] J. Beltrán, C. Landim; Tunneling and metastability of continuous time Markov chains II. J. Stat. Phys. 149, 598-618 (2012).
- [4] J. Beltrán, C. Landim; Tunneling of the Kawasaki dynamics at low temperatures in two dimensions. Ann. Inst. H. Poincaré, Probab. Statist. 51, 59-88 (2015).
- [5] G. Ben Arous, R. Cerf, Metastability of the three-dimensional Ising model on a torus at very low temperature, Electron. J. Probab. 1 (1996) Research Paper 10.
- [6] L. Bertini, A. Faggionato, D. Gabrielli: From level 2.5 to level 2 large deviations for continuous time Markov chains. Markov Processes and Related Fields 20, 545–562 (2014).
- [7] L. Bertini, A. Faggionato, D. Gabrielli: Large deviations of the empirical flow for continuous time Markov chains. Annales de l'IHP Probabilités et statistiques 51, 867-900 (2015).
- [8] L. Bertini, D. Gabrielli, C. Landim: Metastable expansion of finite state Markov chains level two large deviations rate functions. Ann. Appl. Probab. 34, 3820–3869 (2024).
- A. Bovier, F. den Hollander, F.R. Nardi: Sharp asymptotics for Kawasaki dynamics on a finite box with open boundary. Probab. Theory Relat. Fields 135, 265-310 (2006).
- A. Bovier, F. Manzo: Metastability in Glauber dynamics in the low-temperature limit: beyond exponential asymptotics, J. Stat. Phys. 107, 757–779 (2002).
- [11] A. Braides; \(\Gamma\)-Convergence for beginners. Oxford University Press, Oxford, 2002.

- [12] E. Cirillo, E. Olivieri: Metastability and nucleation for the Blume-Capel model. Different mechanisms of transition. J. Stat. Phys. 83, 473-554 (1996).
- [13] M. D. Donsker, S. R. S. Varadhan: Asymptotic evaluation of certain Markov proces expectations for large time, I, Comm. Pure Appl. Math., 28, 1-47 (1975).
- [14] M. Freidlin, L. Koralov: Metastable Distributions of Markov Chains with Rare Transitions. J. Stat. Phys. 167, 1355–1375 (2017).
- [15] M. I. Freidlin, A. D. Wentzell: Random perturbations of dynamical systems. Second edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 260. Springer-Verlag, New York, 1998.
- [16] A. Gaudillière, F. Den Hollander, F.R. Nardi, E. Olivieri, E. Scoppola: Ideal gas approximation for a two-dimensional rarefied gas under Kawasaki dynamics, Stochastic Process. Appl. 119, 737–774 (2009).
- [17] F. den Hollander, F. Nardi, A. Troiani: Metastability for Kawasaki dynamics at low temperature with two types of particles, Electron. J. Probab. 17: 1–26 (2012).
- [18] S. Kim, I. Seo: Approximation method to metastability: an application to non-reversible, two-dimensional Ising and Potts models without external fields (2022). arXiv:2212.13746
- [19] S. Kim, I. Seo: Energy landscape and metastability of stochastic Ising and Potts models on three-dimensional lattices without external fields, Electron. J. Probab. 29: 1–70 (2024).
- [20] C. Landim: Metastable Markov chains. Probability Surveys 16, 143–227 (2019). DOI: 10.1214/18-PS310
- [21] C. Landim: Metastability from the large deviations point of view: A Γ-expansion of the level two large deviations rate functional of non-reversible finite-state Markov chains. Stoch. Proc. Appl. 165, 275–315 (2023).
- [22] C. Landim, P. Lemire; Metastability of the two-dimensional Blume-Capel model with zero chemical potential and small magnetic field. J. Stat. Phys. 164, 346–376 (2016).
- [23] C. Landim, M. Loulakis, M. Mourragui: Metastable Markov chains: from the convergence of the trace to the convergence of the finite-dimensional distributions. Electron. J. Probab. 23, paper no. 95 (2018).
- [24] C. Landim, T. Xu; Metastability of finite state Markov chains: a recursive procedure to identify slow variables for model reduction. ALEA Lat. Am. J. Probab. Math. Stat. 13, 725-751 (2016).
- [25] M. Mariani: A Γ-convergence approach to large deviations. Ann. Sc. Norm. Super. Pisa Cl. Sci. 18, 951–976 (2018).
- [26] F. R. Nardi, A. Zocca: Tunneling behavior of Ising and Potts models in the low-temperature regime. Stochastic Process. Appl. 129, 4556–4575 (2019).
- [27] E. J. Neves, R. H. Schonmann: Critical droplets and metastability for a Glauber dynamics at very low temperatures. Comm. Math. Phys. 137, 209–230 (1991).
- [28] E. J. Neves, R. H. Schonmann: Behavior of droplets for a class of Glauber dynamics at very low temperature. Probab. Theory Related Fields 91, 331–354 (1992).

SEONWOO KIM

Korea Institute for Advanced Study

85 Hoegi-ro,

Dongdaemun-gu, Seoul 02455, South Korea.

Email address: seonwookim@kias.re.kr

Claudio Landim

IMPA

ESTRADA DONA CASTORINA 110,

J. Botanico, 22460 Rio de Janeiro, Brazil

and

CNRS UMR 6085, Université de Rouen,

AVENUE DE L'UNIVERSITÉ, BP.12, TECHNOPÔLE DU MADRILLET,

F76801 SAINT-ÉTIENNE-DU-ROUVRAY, FRANCE.

Email address: landim@impa.br