On stochastic control problems with higher-order moments

Yike Wang^{*}, Jingzhen Liu[†], Alain Bensoussan[‡], Ka-Fai Cedric Yiu[§], Jiaqin Wei.[¶]

Abstract

In this paper, we focus on a class of time-inconsistent stochastic control problems, where the objective function includes the mean and several higher-order central moments of the terminal value of state. To tackle the time-inconsistency, we seek both the closed-loop and the open-loop Nash equilibrium controls as time-consistent solutions. We establish a partial differential equation (PDE) system for deriving a closed-loop Nash equilibrium control, which does not include the equilibrium value function and is different from the extended Hamilton-Jacobi-Bellman (HJB) equations as in Björk et al. (Finance Stoch. 21: 331-360, 2017). We show that our PDE system is equivalent to the extended HJB equations that seems difficult to be solved for our higher-order moment problems. In deriving an open-loop Nash equilibrium control, due to the non-separable higher-order moments in the objective function, we make some moment estimates in addition to the standard perturbation argument for developing a maximum principle. Then, the problem is reduced to solving a flow of forward-backward stochastic differential equations. In particular, we investigate linear controlled dynamics and some objective functions affine in the mean. The closed-loop and the open-loop Nash equilibrium controls are identical, which are independent of the state value, random path and the preference on the odd-order central moments. By sending the highest order of central moments to infinity, we obtain the time-consistent solutions to some control problems whose objective functions include some penalty functions for deviation.

Keywords: stochastic control, higher-order moment, time-consistent, closed-loop Nash equilibrium control, open-loop Nash equilibrium control

AMS2010 classification: Primary: 93E20, 91G80; Secondary: 91B08, 49N90

1 Introduction

Stochastic control theories have been widely studied in the past few decades. It is a quite important mathematical technic in the field of finance and economics, when a dynamic decision is supposed to be made in a multi-period problem. Dynamic programming and Bellman principle of optimality, see [\(Yong & Zhou](#page-30-0), [1999](#page-30-0)), represent the property of optimal control in a large class of dynamic optimization problems: An optimal control based on the current information is still optimal in future. However, there are plenty of "non-standard" problems, for which the dynamic programming method cannot be directly used. In those problems, an optimal control for some initial epochs may no longer be optimal for some future epochs in the sense of global maximization/minimization, which may induce the decision makers

[∗] School of Finance, Chongqing Technology and Business University, Chongqing 400067, China.

[†]Corresponding author. China Institute for Actuarial Science, Central University of Finance and Economics, Beijing 100081, China.

[†]Naveen Jindal School of Management, University of Texas at Dallas, Dallas, TX 75083.

[§]Department of Applied Mathematics, The Hong Kong Polytechnic University, Hunghom, Kowloon, Hong Kong, China. ¶Key Laboratory of Advanced Theory and Application in Statistics and Data Science–MOE, School of Statistics, East China Normal University, Shanghai 200241, China.

to change their approach and to modify their previous optimal plan at every instant. This phenomenon is known as "time-inconsistency".

The studies on time-inconsistent control problems can be traced back to [\(Strotz](#page-29-0), [1955;](#page-29-0) [Pollak,](#page-29-1) [1968;](#page-29-1) [Peleg & Yaari,](#page-29-2) [1973;](#page-29-2) [Goldman](#page-28-0), [1980;](#page-28-0) [Laibson,](#page-29-3) [1997,](#page-29-3) [1998](#page-29-4)). In the early years, the research on timeinconsistent control problems can be almost classified into two main categories:

- the problems whose objective function includes non-exponential discounting factors, like $J(t, u) =$ $\mathbb{E}[\int_t^T Q(t,s)H(s,X_s^u,u_s)ds + Q(t,T)F(X_T^u)|\mathcal{F}_t];$
- and the problems whose objective function includes non-linear functions of the conditional expectation of state, like $J(t, u) = G(\mathbb{E}[X_T^u | \mathcal{F}_t]).$

Here u and X^u respectively denote the control process and its corresponding state process. Later, the non-exponential discounting problem was extended to the initial-state dependence problem, while the problem with non-linear functions of expectation was extended to the mean-field type control problem. On the one hand, the initial-state dependence means that some functions for evaluating the future control, such as the integrand in the objective function, explicitly rely on the current time or the current value of state. See, e.g., [\(Björk, Khapko, & Murgoci,](#page-28-1) [2017](#page-28-1), (6.1)). This initial-state dependence in linearquadratic (LQ) problems and utility maximization problems has attracted much attention. The related literature includes, e.g., [\(Yong,](#page-30-1) [2011](#page-30-1), [2012;](#page-30-2) [Ekeland, Mbodji, & Pirvu](#page-28-2), [2012](#page-28-2); [Zhao, Shen, & Wei](#page-30-3), [2014;](#page-30-3) [Liu, Lin, Yiu, & Wei,](#page-29-5) [2020](#page-29-5); [Zhang, Purcal, & Wei,](#page-30-4) [2021\)](#page-30-4). On the other hand, in many mean-field type control problems, the conditional expectation $\mathbb{E}[X_T^u|\mathcal{F}_t]$ also appears in the controlled dynamic equation of X^u . More generally, researchers incorporate the distribution of X_s^u into the controlled dynamic equation (known as McKean-Vlasov equation) to make theoretical extensions. In terms of application, the meanfield type control problems, sometimes combined with LQ problems, have been deeply investigated for several years, see, e.g., [\(Hu, Jin, & Zhou,](#page-29-6) [2012,](#page-29-6) [2017;](#page-29-7) [Bensoussan, Frehse, & Yam,](#page-28-3) [2013;](#page-28-3) [Yong,](#page-30-5) [2017\)](#page-30-5) and references therein. As the most celebrated application, the mean-variance dynamic portfolio selection, which can be traced back to [\(Markowitz,](#page-29-8) [1952](#page-29-8)), was pioneered by [\(Li & Ng,](#page-29-9) [2000](#page-29-9); [Zhou & Li](#page-30-6), [2000\)](#page-30-6) and developed by a huge body of literature including [\(Basak & Chabakauri,](#page-28-4) [2010;](#page-28-4) [Czichowsky,](#page-28-5) [2013;](#page-28-5) [Björk, Murgoci, & Zhou,](#page-28-6) [2014\)](#page-28-6), etc.

A straightforward extension of mean-variance dynamic portfolio selection is to incorporate higherorder (central) moments into the objective function. Given that the distribution of the terminal wealth can be determined by all central/original moments if its moment generating function exists, an objective function including central/origin moments as many as possible might be regarded as an approximate quantification for the preferences on the distribution.

Moreover, objective functions with higher-order central moments can also arise from improving the classical single-period mean-variance model pioneered by [\(Markowitz](#page-29-8), [1952\)](#page-29-8). As is well known, in the classical model, the investors aim to maximize the expected return when the variance does not exceed a given level, or to minimize the variance when the expected return does not fall below a given level. By Pareto efficiency or the effective frontier theory (see also [\(Yong & Zhou,](#page-30-0) [1999,](#page-30-0) Section 6.8) or [\(Zhou & Li,](#page-30-6) [2000\)](#page-30-6)), the primal problem can be embedded into a class of unconstrained optimization problems with "mean-variance utility", where the utility is increasing with the mean and decreasing with the variance. In the same manner, for the maximizer of mean-variance-skewness utility, which is increasing with respect to the mean and decreasing with respect to the variance and the skewness, the investor cannot further improve the expected return without increasing the variance and the skewness. Notably, when the mean and the variance are fixed, a larger skewness usually indicates a smaller mode or median, which is not preferred by some investors. This coincides with the negative risk premium of skewness found by [\(Chabi-Yo,](#page-28-7) [2012](#page-28-7)) in the multi-period capital asset pricing model with higher-order moments. However, a positive skewness usually arises from a long right tail, which may be preferred by some investors. This implies that it is also rational to assume that the mean-variance-skewness utility is increasing with the skewness. The above discussion on benefit and drawback of positive skewness is still valid for other odd-order central moments, since they can also be treated as measures of the symmetry of a return's distribution.

Furthermore, we have the following informal expansion for the expected utility of the random return X (see also [\(Samuelson,](#page-29-10) [1970;](#page-29-10) [Jean](#page-29-11), [1971](#page-29-11); [Scott & Horvath,](#page-29-12) [1980\)](#page-29-12)),

$$
\mathbb{E}[\Psi(X)] = \Psi(0) + \Psi'(0)\mathbb{E}[X] + \sum_{j=2}^{n} \frac{1}{j!} \Psi^{(j)}(\mathbb{E}[X])\mathbb{E}[(X - \mathbb{E}[X])^{j}] + \bar{R},
$$

where E denotes the expectation operator, Ψ is a utility function, and \bar{R} is the residual. The mean and the higher-order central moments are included in the expansion. From stochastic dominance theories (see, e.g., [\(Ekern,](#page-28-8) [1980\)](#page-28-8)), $(-1)^{j+1}\Psi^{(j)}$ for every j is supposed to be positive, which provides appropriate assumptions for the monotonicity of the utility with respect to the central moments.

Besides, in the objective function, the combination of higher-order central moments can act as a penalty function for the deviation $X - \mathbb{E}[X]$, or can generate a utility function of the random return X net of the endogenous reference level $\mathbb{E}[X]$. For example, for the objective function

$$
J = \mathbb{E}[X] - \sum_{j=2}^{\infty} \frac{(-c)^j}{j!} \mathbb{E}[(X - \mathbb{E}[X])^j] = \mathbb{E}[X] - \mathbb{E}[e^{-c(X - \mathbb{E}[X])}] + 1, \quad c > 0,
$$
 (1)

the function $S(x) = \exp(-cx) - 1$ provides a relatively small reward (resp. large penalty) for positive (resp. negative) deviation from the expected return. Also, the abovementioned objective function J can be regarded as a weighted average of the mean $\mathbb{E}[X]$ for a risk-neutral preference on X, and the constant absolute risk aversion (CARA) utility function of $X - \mathbb{E}[X]$, whose CARA coefficient is c. The inspiration about endogenous reference level can be also found in the study on, e.g., consumer's habit formation in behavioral finance (see also [\(Liu et al.,](#page-29-5) [2020\)](#page-29-5)). The related results for the abovementioned objective function can be found in Section [5.3.1](#page-23-0) in this paper.

In the last few decades, there have been several studies devoted to the portfolio selection problems with higher-order moments. Researchers proceeded with static (namely, single-period) mean-varianceskewness problem. [\(Konno & Suzuki,](#page-29-13) [1995](#page-29-13)) found that the "efficient frontier" for the mean-varianceskewness problem is non-concave (see Figure 1 for its 3D image and Figure 2 for its 2D section, therein), which may lead to a significant increase in the complexity of further analysis. At first, the 3D efficient frontier has a flat area and many kinks, where the solution switches from a corner point to an interior point or vice versa. When drawing a "capital market surface" as an analog to capital market line, which should be a ruled surface tangent to the 3D efficient frontier, one can find that the intersection of the 3D efficient frontier and the capital market surface is sometimes discontinuous. Furthermore, it is difficult to derive an explicit solution for mean-variance-skewness utility maximization over the capital market surface, since the indifference surface of utility is not necessarily convex. [\(De Athayde & Flôres,](#page-28-9) [2004](#page-28-9)) considered the variance minimization problem with mean-skewness constraints to determine the efficient frontier. The result is not always the same as that of the skewness maximization problem with mean-variance constraints or the mean maximization problem with variance-skewness constraints. This also indicates the non-concavity of the efficient frontier. To handle the non-concavity, many researchers have tried to obtain the optimal solution by specifying the objective function, by considering some multi-object problems, or by designing some numerical algorithms. See, e.g., [\(Lai](#page-29-14), [1991;](#page-29-14) [Konno, Shirakawa, & Yamazaki](#page-29-15), [1993;](#page-29-15) [Joro & Na,](#page-29-16) [2006;](#page-29-16) [Briec, Kerstens, & Jokung,](#page-28-10) [2007;](#page-28-10) [Harvey, Liechty, Liechty, & Müller](#page-28-11), [2010;](#page-28-11) [Boissaux & Schiltz](#page-28-12), [2013;](#page-28-12) [Landsman, Makov, & Shushi,](#page-29-17) [2020;](#page-29-17) [Mehlawat, Gupta, & Khan,](#page-29-18) [2021;](#page-29-18) [Zhen & Chen,](#page-30-7) [2022\)](#page-30-7).

For dynamic maximization problems in multi-period or continuous-time models, the non-concavity of efficient frontier, or the non-convexity of optimization problem, makes it difficult to derive the value function from the associated Hamilton-Bellman-Jacobi (HJB) equation. The stochastic maximum principle has also been considered (see [\(Boissaux & Schiltz](#page-28-13), [2010\)](#page-28-13)), but it is still quite hard to derive an explicit expression of the maximizer/minimizer. We should note that unlike the systematic skewness and co-skewness of yield rates of all assets emphasized in [\(Kraus & Litzenberger](#page-29-19), [1976\)](#page-29-19), it is the total skewness of terminal wealth that the objective function of a dynamic optimization problem includes. In fact, the geometric Brownian motion, or the Brownian drift-diffusion model in general, has already provided the multivariate distribution of the long-term yield rates of those assets.

Rather than a maximizer/minimizer for the objective function, we consider time-consistent solutions for our higher-order moment problems. Notably, the model studied in this paper is fairly general, although it arises from the portfolio problems. On the one hand, following the definition as in [\(Björk & Murgoci,](#page-28-14) [2014;](#page-28-14) [Björk et al.](#page-28-1), [2017;](#page-28-1) [Wang & Zheng,](#page-30-8) [2021\)](#page-30-8), we try to seek an optimal feedback control only against some spike variations in form of feedback function, which is known as the closed-loop Nash equilibrium control, or namely, the Nash equilibrium control law. The problem is reduced to solving an extended HJB system, which contains a Bellman equation for the equilibrium value function and some partial differential equations (PDEs) known as the Feynman-Kac formula for original moments. On the other hand, referring to [\(Hu et al.](#page-29-6), [2012,](#page-29-6) [2017\)](#page-29-7), we try to seek an open-loop Nash equilibrium control, which is optimal against some spike variations in form of stochastic process. A standard perturbation argument from [\(Yong & Zhou](#page-30-0), [1999,](#page-30-0) Chapter 3) is adopted to develop a sufficient maximum principle. And then the solution can be derived from a flow of forward-backward stochastic differential equations (FBSDEs).

The main contributions of this paper are as follows. Firstly, we establish a PDE system for deriving a closed-loop Nash equilibrium control. The PDE system does not include the so-called equilibrium value function and is different from the well-known extended HJB equations as in [\(Björk et al.,](#page-28-1) [2017\)](#page-28-1). After using a sequence of multipliers to connect the PDEs arising from the Feynman-Kac formula for original moments, we establish a verification theorem to show the sufficiency of a feedback control to the optimality. Our innovative methodology is similar to that in [\(Wang & Zheng,](#page-30-8) [2021\)](#page-30-8), where a generalized equilibrium value function is introduced which reduces to the classical equilibrium value function in a diagonal manner. However, using the system given by [\(Wang & Zheng,](#page-30-8) [2021\)](#page-30-8) is not appropriate to solve our problem, where the terminal value of the state and its mean and higher-order moments are not separable. For our approach, when handling the problems whose objective function can be re-expressed as a linear combination of mean and higher-order central moments, it is not necessary to solve any Bellman equation or PDE to arrive at the time-consistent solution. Secondly, we derive a sufficient maximum principle with a flow of FBSDEs in seeking open-loop Nash equilibrium controls. For a linear control model, the maximum principle can be further simplified. Notably, compared with the classical cases, e.g., in [\(Peng,](#page-29-20) [1990;](#page-29-20) [Yong & Zhou](#page-30-0), [1999](#page-30-0); [Buckdahn, Djehiche, & Li,](#page-28-15) [2011;](#page-28-15) [Hu et al.,](#page-29-6) [2012\)](#page-29-6), the estimation of more expansion terms needs to be investigated in the perturbation argument, due to the non-separable higher-order moments in the objective function. Thirdly, we consider a particular case with a linear control model, where the objective function is affine in the mean. Both categories of time-consistent solutions, closed-loop and open-loop, are derived. It is interesting to find that they are identical, and they are independent of both the state and random path. Moreover, the preference on odd-order central moments does not affect the solution. Finally, we consider several limiting cases where

the objective functions are linear combinations of mean and central moments and the highest order of the central moments tends to infinity. The closed-form solution in each case is derived from solving an algebraic equation rather than a differential equation. Moreover, by a heuristic approach with the use of Fourier cosine transform, we obtain a candidate time-consistent solution for a fairly general even penalty function.

The rest of this paper is organized as follows. In Section [2,](#page-4-0) we introduce the problem with higherorder central moments in a general formulation. In Section [3,](#page-7-0) an extended HJB system with a verification theorem for closed-loop Nash equilibrium control is studied. In Section [4,](#page-9-0) we derive a sufficient maximum principle for open-loop Nash equilibrium control, and we reduce the problem to solving a flow of FBSDEs. In Section [5,](#page-15-0) we apply our results to solve several particular linear control problems, and show their state/path-independent solutions. In Section [6,](#page-27-0) we make some concluding remarks.

2 Model and problem formulation

Let T denote a fixed finite time-horizon, $(\Omega, \mathcal{F}, \mathbb{P})$ denote a complete probability space, on which a one-dimensional standard Brownian motion $W := \{W_t\}_{t\in[0,T]}$ is defined, and $\mathbb E$ be the expectation operator. Let $\mathbb{F} := {\{\mathcal{F}_t\}}_{t \in [0,T]}$ be the right-continuous, completed natural filtration generated by W, and $\mathbb{E}_t[\cdot] := \mathbb{E}[\cdot | \mathcal{F}_t]$ for short. For the sake of brevity, in this paper we usually omit the statement $\mathbb{P}\text{-a.s.}$ (namely, almost surely with respect to \mathbb{P}) for equalities and inequalities involving conditional expectations or Itô integrals, unless otherwise noted. We suppose that the dynamics of the real-valued state process ${X_t}_{t\in[0,T]}$ evolve as the following fairly general controlled stochastic differential equation (SDE),

$$
dX_s = b(s, X_s, u_s)ds + \sigma(s, X_s, u_s)dW_s, \quad X_0 = x_0,
$$
\n(2)

and the primal object is to maximize

$$
J(0, x_0, u) = \mathbb{E}\Big[\Psi\big(0, x_0, X_T, \mathbb{E}[X_T], \mathbb{E}\big[(X_T - \mathbb{E}[X_T])^2\big], \dots, \mathbb{E}\big[(X_T - \mathbb{E}[X_T])^n\big]\big)\Big]
$$

for a fixed integer $n > 2$. When the time goes to t and the state moves to x, or namely, for the initial pair (t, x) , it is supposed to maximize the objective function

$$
J(t, x, u) := \mathbb{E}_t \Big[\Psi(t, x, X_T^{t, x, u}, \mathbb{E}_t [X_T^{t, x, u}], \mathbb{E}_t \big[(X_T^{t, x, u} - \mathbb{E}_t [X_T^{t, x, u}])^2 \big],
$$

$$
\dots, \mathbb{E}_t \big[(X_T^{t, x, u} - \mathbb{E}_t [X_T^{t, x, u}])^n \big] \big] \Big],
$$
 (3)

where $\{X_s^{t,x,u}\}_{s\in[t,T]}$ is given by the following controlled SDE:

$$
dX_s^{t,x,u} = b(s, X_s^{t,x,u}, u_s)ds + \sigma(s, X_s^{t,x,u}, u_s)dW_s, \quad X_t^{t,x,u} = x.
$$
 (4)

To reduce the complexity of derivation, the following basic assumptions are adopted:

- 1. $\{u_t\}_{t\in[0,T]}$ is valued in \mathcal{R} , a subspace of some Euclidean space.
- 2. The initial condition $X_0 = x_0 \in \mathbb{R}$ and the functions $b, \sigma : [0, T] \times \mathbb{R} \times \mathcal{R} \to \mathbb{R}$ and $\Psi : [0, T] \times \mathbb{R} \times \mathbb{R}$ $\mathbb{R} \times \mathbb{R}^n \to \mathbb{R}$ are given a priori.
- 3. $b(t, x, u)$ and $\sigma(t, x, u)$ are uniformly continuous, and twice differentiable in x. Moreover, (b, σ) and their partial derivatives $(b_x, b_{xx}, \sigma_x, \sigma_{xx})$ are Lipschitz continuous in (x, u) , for which the Lipschitz

continuity parameters are uniformly bounded for all $t \in [0, T]$; and $|b(\cdot, 0, \cdot)|$ and $|\sigma(\cdot, 0, \cdot)|$ are uniformly bounded.

4. $\Psi(t, x, y, \vec{z})$ is continuously differentiable in $\vec{z} := (z_1, \ldots, z_n)$ and twice differentiable in y. Moreover, the functions $(\Psi, \Psi_y, \Psi_{yy}, \Psi_{z_1}, \dots, \Psi_{z_n})$ are Lipschitz continuous in (y, \vec{z}) , for which the Lipschitz continuity parameters are uniformly bounded for all (t, x) ; and $|\Psi(\cdot, \cdot, 0, \vec{0})|$ is uniformly bounded.

Definition 2.1. A stochastic process $u = \{u_t\}_{t\in[0,T]}$ is named an "admissible control", if it is R-valued and $\mathbb{F}\text{-}adapted$, and the corresponding controlled SDE [\(2\)](#page-4-1) admits a unique $\mathbb{F}\text{-}adapted$ strong solution such that $\mathbb{E}[|X_T|^{2n-2+\epsilon}] < \infty$ for some $\epsilon > 0$. Furthermore, a deterministic function $\hat{u}: [0,T] \times \mathbb{R} \to \mathcal{R}$ is named an "admissible feedback control", if u given by $u_t = \hat{u}(t, X_t^{0,x_0,u})$ is an admissible control.

Remark 2.2. The additional condition $\mathbb{E}[|X_T|^{2n-2+\epsilon}] < \infty$ is employed for the case $\Psi_{yz_n} \neq 0$ in Section [4.](#page-9-0) If $\Psi_{yz_m} \equiv 0$ for all $m \geq n/2$, then $\mathbb{E}[|X_T|^{2n-2+\epsilon}] < \infty$ can be weakened to $\mathbb{E}[|X_T|^n] < \infty$.

Let \mathcal{U}_0 denote the set of all the admissible controls and $\mathcal U$ denote the set of all the admissible feedback controls. With a slight abuse of notation, corresponding to $\hat{u} \in \mathcal{U}$, we let $\{X_s^{t,x,\hat{u}}\}_{s \in [t,T]}$ and $J(t,x,\hat{u})$ be the state process and the objective function, respectively. In addition, for the sake of brevity, we let $z_0 := 1, \partial_x := \frac{\partial}{\partial x}, f_{\vec{z}} := (f_{z_1}, \ldots, f_{z_n})$ for any f continuously differentiable in $\vec{z}, f_{\vec{z}}\vec{z}$ denote the Hessian matrix of f with respect to \vec{z} for any f twice differentiable in \vec{z} , and $\langle \cdot, \cdot \rangle$ denote the scalar product (or namely, inner product) of two row vectors; and employ the infinitesimal operator \mathcal{D}_{ζ} by

$$
\mathcal{D}_{\zeta}f(t,x) := f_t(t,x) + f_x(t,x)b(t,x,\zeta) + \frac{1}{2}f_{xx}(t,x)\big(\sigma(t,x,\zeta)\big)^2
$$

for $\zeta \in \mathcal{R}$ and any bivariate function f continuously differentiable in the first argument and twice differentiable in the second argument.

It is well-known that the dynamic maximization problems, maximizing [\(3\)](#page-4-2) subject to [\(4\)](#page-4-3) over all admissible controls u (or all $\hat{u} \in \mathcal{U}$) for all initial pairs (t, x) , are usually time-inconsistent, due to the fairly general initial-pair-dependent function $\Psi(t, x, \cdot, \cdot)$ and the central moments in the objective function. In other words, any maximizer u for some initial pair (t, x) may not realize the maximization of [\(3\)](#page-4-2) for the initial pair $(s, X_s^{t,x,u})$. To overcome the time-inconsistency, researchers formulate the problem as a game between the agent and her/his future selves, pioneered by [\(Strotz,](#page-29-0) [1955\)](#page-29-0). A subgame perfect Nash equilibrium point for this game is regarded as a time-consistent solution, since any deviation from it at any time instant will be worse off and each of the future selves has no incentive to change the equilibrium strategy. In this paper, we take this game-theoretic perspective and seek a closed-loop Nash equilibrium control (see also [\(Björk et al.,](#page-28-1) [2017](#page-28-1))) and an open-loop Nash equilibrium control (see also [\(Hu et al.,](#page-29-6) [2012,](#page-29-6) [2017\)](#page-29-7)), which are defined as follows.

Definition 2.3. $\tilde{u} \in \mathcal{U}$ is a closed-loop Nash equilibrium control (CNEC), if

$$
0 \le \liminf_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \big(J(t, \tilde{X}_t, \tilde{u}) - J(t, \tilde{X}_t, \tilde{u}^{t, \varepsilon, \zeta}) \big), \quad \mathbb{P}-a.s., \ a.e. \ t \in [0, T), \tag{5}
$$

for $\tilde{X} := X^{0,x_0,\tilde{u}}$ and any constant $\zeta \in \mathcal{R}$ satisfying $\tilde{u}^{t,\varepsilon,\zeta} \in \mathcal{U}$, where $\tilde{u}^{t,\varepsilon,\zeta}$ is a spike variation of \tilde{u} given by $\tilde{u}^{t,\varepsilon,\zeta}(s,\cdot) = \tilde{u}(s,\cdot)1_{\{s\notin[t,t+\varepsilon)\}} + \zeta1_{\{s\in[t,t+\varepsilon)\}}$. Moreover, corresponding to any fixed CNEC \tilde{u} , $V(t, x) := J(t, x, \tilde{u})$ is called an equilibrium value function (EVF).

Remark 2.4. Our Definition [2.3](#page-5-0) for CNEC is a bit weaker than that in [\(Björk et al.](#page-28-1), [2017](#page-28-1)), while [\(Björk et al.](#page-28-1), [2017\)](#page-28-1) used an arbitrary function to make the perturbation (namely, $\zeta = \zeta(s, y)$). In fact, our modification is due to the use of Dynkin's formula for, e.g., [\(16\)](#page-8-0), where the feedback control should

be continuous in time. In other words, replacing the constant ζ by the function $\zeta(t,x)$ continuous in t, as well as replacing \mathcal{D}_{ζ} by $\mathcal{D}_{\zeta(t,x)}$, does not affect the validness of results in this work. Apart from that, our Definition [2.3](#page-5-0) for CNEC is also much weaker than the strong equilibrium strategy in (Huang \mathcal{B} Zhou, [2021](#page-28-16); He \mathcal{C} Jiang, 2021). Although the weak equilibrium strategy might correspond to a stationary point rather than a maximum as mentioned in $(Björk \text{ et al., } 2017,$ $(Björk \text{ et al., } 2017,$ $(Björk \text{ et al., } 2017,$ Remark 3.5), the coming sufficient condition guarantees that $J(t, \bar{X}_t, \bar{u}^{t,\varepsilon,\zeta}) - J(t, \bar{X}_t, \bar{u})$, even if positive, is at most $o(\varepsilon)$ (see the proof for Theorem [3.2\)](#page-7-1). Notably, (He $\mathcal B$ Jiang, [2021](#page-28-16)) has shown that many classical time-inconsistent control problems, including the mean-variance problem in $(Basak \& Chabakauri, 2010)$ $(Basak \& Chabakauri, 2010)$, have no proper strong equilibrium strategy, if the perturbation ζ could be a smooth function. Therefore, we do not intend to find the strong equilibrium strategy for our problem, but leave this work to our future research.

Definition 2.5. $\bar{u} \in \mathcal{U}_0$ is an open-loop Nash equilibrium control (ONEC), if

$$
0 \le \liminf_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \big(J(t, \bar{X}_t, \bar{u}) - J(t, \bar{X}_t, \bar{u}^{t, \varepsilon, \zeta}) \big), \quad \mathbb{P}-a.s., \ a.e. \ t \in [0, T), \tag{6}
$$

 $for \ \bar{X} := X^{0,x_0,\bar{u}}$ and any \mathcal{F}_t -measurable random variable ζ satisfying $\bar{u}^{t,\varepsilon,\zeta} \in \mathcal{U}_0$, where $\bar{u}^{t,\varepsilon,\zeta}$ is a spike variation of \bar{u} given by $\bar{u}_s^{t,\varepsilon,\zeta} = \bar{u}_s + \zeta 1_{\{s \in [t,t+\varepsilon)\}}.$

Remark 2.6. Distinct from the substitution used in the spike variation for CNEC, the spike variation for ONEC uses the addition of a perturbation. This modification helps limit the differences, such as $\sigma(s,\bar{X}_s,\bar{u}_s+\zeta)-\sigma(s,\bar{X}_s,\bar{u}_s)$ in the proof for Lemma [4.1,](#page-10-0) by a multiple of $|\zeta|$, given that the admissibility of u does not ensure the boundedness of every u_s . In other words, if we had adopted some adequate assumptions for the growth or boundedness of (b, σ) and the integrability of u, the spike variation for ONEC could have been generalized as $\bar{u}_s^{t,\epsilon,\zeta} = \bar{u}_s \mathbb{1}_{\{s \notin [t,t+\epsilon)\}} + \zeta_s \mathbb{1}_{\{s \in [t,t+\epsilon)\}}$. See, e.g., [\(Peng,](#page-29-20) [1990](#page-29-20); [Buckdahn et al.](#page-28-15), [2011](#page-28-15); [Alia,](#page-28-17) [2019\)](#page-28-17).

Remark 2.7. There is a recent work (Liang, Xia, $\&$ Yuan, [2023\)](#page-29-22) that characterized open-loop equilibria for dynamic portfolio problem with a fairly general objective function $g(\mathbb{P}_{X_T}^t)$ named non-linear law-dependent preference, where $\mathbb{P}_{X_T}^t$ denotes the regular conditional distribution of X_T given \mathcal{F}_t . The problem therein reduces to maximizing an expected utility (resp. a mean-variance objective function), when the Fréchet derivative $\nabla g(\mathbb{P}_{X_T}^t, \cdot)$ is a $\mathbb{P}_{X_T}^t$ -independent utility function (resp. an affine function of $X_T - \mathbb{E}_t[X_T]$). So many celebrated studies are covered in [\(Liang et al.](#page-29-22), [2023](#page-29-22)). However, [\(Liang et al.,](#page-29-22) [2023](#page-29-22), Assumption 3.1) imposed the concavity assumption on $\nabla q(\mu, \cdot)$, which is not valid for our prob-lems. For instance, to apply the results in [\(Liang et al.](#page-29-22), [2023](#page-29-22)) to $\Psi(\mathbb{E}_{t}[(X_T - \mathbb{E}_{t}[X_T])^3])$, we shall $let g(\mu) = \Psi(\int_{\mathbb{R}} x^3 \mu(dx) - 3 \int_{\mathbb{R}} x^2 \mu(dx) \times \int_{\mathbb{R}} x \mu(dx) + 2(\int_{\mathbb{R}} x \mu(dx))^3$. Then, $\nabla g(\mu, x) = \Psi' \times (x^3 3x^2\int_{\mathbb{R}}y\mu(dy)-3x\int_{\mathbb{R}}y^2\mu(dy)+6x(\int_{\mathbb{R}}y\mu(dy))^2),$ which is not necessarily concave in x. For another example, for $g(\mathbb{P}_{X_T}^t) = \kappa \mathbb{E}_t[X_T] - \mathbb{E}_t[S(X_T - \mathbb{E}_t[X_T])]$ with continuously differentiable $S : \mathbb{R} \to \mathbb{R}$, as the unified form of the limiting cases we investigate in Section [5.3,](#page-22-0) we shall let $g(\mu) = \int_{\mathbb{R}} (\kappa x - S(x - \int_{\mathbb{R}} y \mu(dy))) \mu(dx)$. Then, $\nabla g(\mu, x) = \kappa x - S(x - \int_{\mathbb{R}} y\mu(dy)) + x \int_{\mathbb{R}} S'(z - \int_{\mathbb{R}} y\mu(dy))\mu(dz)$, which is concave in x if and only if S is concave. In particular, $\nabla g(\mu, \cdot)$ is concave for the mean-exponential utility [\(1\)](#page-2-0) (see also Sec-tion [5.3.1\)](#page-23-0), while it is not concave for $S(x) = (1 - \cos(cx))/c$ (see Section [5.3.3\)](#page-25-0). Due to the presence of non-concavity in our objective functions as well as the generality of our model, we employ the spike variation method for which the second-order variational equation and its adjoint equation are involved (see (20) and (25)), rather than refer to the spirit of convex variation method for which only the first adjoint duality analysis is considered as in [\(Liang et al.,](#page-29-22) [2023](#page-29-22)). Conversely, by our Theorem 4.5 , some verification results in [\(Liang et al.](#page-29-22), [2023](#page-29-22)) are recovered. Interested readers can refer to Example [4.6](#page-13-2) for detailed discussions.

3 Sufficient condition for CNEC: extended HJB-PDE

In this section, we introduce an extended HJB system with several integrality and smoothness conditions for deriving a CNEC. For any $\hat{u} \in \mathcal{U}$, let $m_j^{\hat{u}}(t,x) := \mathbb{E}_t[(X_T^{t,x,\hat{u}})^j]$ and $\vec{m}^{\hat{u}} := (m_1^{\hat{u}}, \dots, m_n^{\hat{u}})$. Without loss of generality, we consider the following fairly general objective function with higher-order original moments,

$$
J(t, x, \hat{u}) = \mathbb{E}_t \big[\Phi(t, x, X_T^{t, x, \hat{u}}, \vec{m}^{\hat{u}}(t, x)) \big],\tag{7}
$$

where Φ has the same properties as of Ψ , e.g. $\Phi(t, x, y, \vec{z})$ is continuously differentiable in \vec{z} and twice differentiable in y . Obviously, in terms of our problem with higher-order central moments,

$$
\Phi(t, x, y, \vec{z}) = \Psi\bigg(t, x, y, z_1, z_2 - z_1^2, z_3 - 3z_1z_2 + 2z_1^3, \dots, \sum_{j=0}^n \binom{n}{j} (-1)^j z_1^j z_{n-j}\bigg). \tag{8}
$$

In particular, we have

$$
\Phi(t, x, y, z, z^2, \dots, z^n) = \Psi(t, x, y, z, 0, \dots, 0).
$$

For the (\mathbb{F}, \mathbb{P}) -martingales $\{m_j^{\hat{u}}(v, X_v^{t,x,\hat{u}}) = \mathbb{E}_v[(X_T^{t,x,\hat{u}})^j]\}_{v \in [t,T]}$ and $\{\mathbb{E}_v[\Phi(s, y, X_T^{t,x,\hat{u}}, \vec{z})]\}_{v \in [t,T]}$ for any $(t, x), (s, y, \vec{z})$ and appropriate \hat{u} , Feynman-Kac formula gives their PDE representation (see, e.g., [\(Yong & Zhou](#page-30-0), [1999](#page-30-0), Theorem 7.4.1, p. 373)). As an analogue, for any $\tilde{u} \in \mathcal{U}$, applying Itô's rule to $m^{(j)}(v, X_v^{t,x,\tilde{u}})$ and $U^{s,y,\vec{z}}(v, X_v^{t,x,\tilde{u}})$ produces the following lemma. The proof is straightforward, so we omit it due to the page limit.

Lemma 3.1. Fix $\tilde{u} \in \mathcal{U}$ and $(s, y, \vec{z}) \in [0, T) \times \mathbb{R} \times \mathbb{R}^n$. Assume that $U^{s,y,\vec{z}}$ and $\vec{m} := (m^{(1)}, \ldots, m^{(n)})$ are the classical solutions of the following PDEs on $[0, T] \times \mathbb{R}$:

$$
0 = \mathcal{D}_{\tilde{u}(t,x)} U^{s,y,\vec{z}}(t,x), \qquad s.t. \qquad U^{s,y,\vec{z}}(T,x) = \Phi(s,y,x,\vec{z}); \tag{9}
$$

$$
0 = \mathcal{D}_{\tilde{u}(t,x)} m^{(j)}(t,x), \qquad s.t. \qquad m^{(j)}(T,x) = x^j, \quad \forall j = 1,2,\ldots,n,
$$
 (10)

respectively; and for any $(t, x) \in [0, T) \times \mathbb{R}$,

$$
\mathbb{E}\bigg[\int_{t}^{T}\big|U_{x}^{s,y,\vec{z}}(v,X_{v}^{t,x,\tilde{u}})\sigma\big(v,X_{v}^{t,x,\tilde{u}},\tilde{u}(v,X_{v}^{t,x,\tilde{u}})\big)\big|^{2}dv\bigg]<\infty;\tag{11}
$$

$$
\mathbb{E}\bigg[\int_t^T \big|m_x^{(j)}(v, X_v^{t,x,\tilde{u}})\sigma\big(v, X_v^{t,x,\tilde{u}}, \tilde{u}(v, X_v^{t,x,\tilde{u}})\big)\big|^2 dv\bigg] < \infty, \quad \forall j = 1, 2, \dots, n. \tag{12}
$$

Then, $U^{s,y,\vec{z}}(t,x) = \mathbb{E}_t[\Phi(s,y,X_T^{t,x,\tilde{u}},\vec{z})]$ and $\vec{m}(t,x) = \vec{m}^{\tilde{u}}(t,x)$. Furthermore, if \tilde{u} is a CNEC, then $V(t,x) = U^{t,x,\vec{m}(t,x)}(t,x)$ is an EVF.

Similarly, for fixed $\tilde{u} \in \mathcal{U}$ and $(s, y) \in [0, T) \times \mathbb{R}$, if $\vec{\lambda}^{s, y}$ is the classical solution of

$$
0 = \mathcal{D}_{\tilde{u}(t,x)} \vec{\lambda}^{s,y}(t,x), \quad s.t. \quad \vec{\lambda}^{s,y}(T,x) = \Phi_{\tilde{z}}(s,y,x,\vec{m}(s,y)), \tag{13}
$$

$$
\mathbb{E}\bigg[\int_{t}^{T} \langle \vec{\lambda}^{s,y}_{x}(v, X^{t,x,\tilde{u}}_{v}), \vec{\lambda}^{s,y}_{x}(v, X^{t,x,\tilde{u}}_{v}) \rangle \big|\sigma\big(v, X^{t,x,\tilde{u}}_{v}, \tilde{u}(v, X^{t,x,\tilde{u}}_{v})\big)\big|^{2} dv\bigg] < \infty,
$$
\n(14)

then $\vec{\lambda}^{s,y}(t,x) = \mathbb{E}_t[\Phi_{\vec{z}}(s, y, X_T^{t,x,\tilde{u}}, \vec{m}(s,y))].$ For the sake of notational brevity, we omit the statement of the dependence of \tilde{u} for $(\vec{m}, U^{s,y,\vec{z}}, \vec{\lambda}^{s,y})$ in what follows, unless otherwise mentioned. As a main result, the following theorem shows that a CNEC can be derived from a system including the abovementioned PDEs and square-integrability conditions with an optimality condition.

Theorem 3.2. For a fixed $\tilde{u} \in \mathcal{U}$ and any $(s, y, \vec{z}) \in [0, T) \times \mathbb{R} \times \mathbb{R}^n$, assume that $(\vec{m}, U^{s,y,\vec{z}}, \vec{\lambda}^{s,y})$ are

the classical solution of the PDEs (9) , (10) and (13) with the optimality condition

$$
\tilde{u}(t,x) \in \underset{\zeta \in \mathcal{R}}{\arg \max} \{ \mathcal{D}_{\zeta} U^{t,x,\vec{m}(t,x)}(t,x) + \langle \vec{\lambda}^{t,x}(t,x), \mathcal{D}_{\zeta} \vec{m}(t,x) \rangle \}
$$
\n(15)

and the square-integrability conditions (11) , (12) and (14) . Then, \tilde{u} is a CNEC.

Proof. Write $X^{t, \tilde{X}_t, \zeta} := X^{t, \tilde{X}_t, \hat{u}(\cdot, \cdot)}$ with $\hat{u}(\cdot, \cdot) \equiv \zeta \in \mathcal{R}$ for short. This slight abuse of notation is used throughout the rest of this paper, unless otherwise stated. Based on the previous results for $(\vec{m}, U^{s,y,\vec{z}}, \vec{\lambda}^{s,y})$, let us proceed with

$$
m_j^{\tilde{u}^{t,\varepsilon,\zeta}}(t,x) = \mathbb{E}_t[m_j^{\tilde{u}}(t+\varepsilon, X_{t+\varepsilon}^{t,x,\zeta})] = m_j^{\tilde{u}}(t,x) + \mathcal{D}_{\zeta}m_j^{\tilde{u}}(t,x)\varepsilon + o(\varepsilon),
$$
\n(16)

where the first equality follows from the definition of $m_j^{\hat{u}}(\cdot,\cdot)$ and the tower property (or namely, iterated conditioning) and the last equality is a straightforward application of Dynkin's formula. Applying the tower property and Dynkin's formula again yields

$$
\mathbb{E}_t[\Phi(s,y,X_T^{t+\varepsilon,X_{t+\varepsilon}^{t,x,\zeta},\tilde{u}},\tilde{z})] = U^{s,y,\tilde{z}}(t,x) + \mathcal{D}_{\zeta}U^{s,y,\tilde{z}}(t,x)\varepsilon + o(\varepsilon).
$$

Then, given the smoothness of $\Phi(t, x, \cdot, \cdot)$, we have

$$
J(t, x, \tilde{u}^{t, \varepsilon, \zeta}) = \mathbb{E}_t \left[\Phi\big(t, x, X_T^{t + \varepsilon, X_{t + \varepsilon}^{t, x, \zeta}, \tilde{u}}, \mathbb{E}_t[\vec{m}^{\tilde{u}}(t + \varepsilon, X_{t + \varepsilon}^{t, x, \zeta})] \big) \right]
$$

= $J(t, x, \tilde{u}) + \left(\mathcal{D}_{\zeta} U^{t, x, \vec{m}(t, x)}(t, x) + \langle \vec{\lambda}^{t, x}(t, x), \mathcal{D}_{\zeta} \vec{m}(t, x) \rangle \right) \varepsilon + o(\varepsilon)$
 $\leq J(t, x, \tilde{u}) + o(\varepsilon),$

where the last inequality follows from (15) with the linear combination of the PDEs in (9) and (10) , that is, $0 = \mathcal{D}_{\tilde{u}(t,x)}U^{t,x,\tilde{m}(t,x)}(t,x) + \langle \vec{\lambda}^{t,x}(t,x), \mathcal{D}_{\tilde{u}(t,x)}\vec{m}(t,x)\rangle$. Consequently, $\liminf_{\varepsilon \downarrow 0} \frac{1}{\varepsilon}[J(t,x,\tilde{u}) J(t, x, \tilde{u}^{t,\varepsilon,\zeta})] \geq 0$, and hence, \tilde{u} is a CNEC.

In addition to the previous smoothness condition for (Ψ, Φ) , we now assume that $\Phi = \Phi(t, x, y, \vec{z})$ is differentiable in t and twice differentiable in (x, \bar{z}) . Then, we are able to establish a Bellman equation for the EVF.

Theorem 3.3 (Verification theorem). Assume that the following properties hold:

- 1. For a fixed $\tilde{u} \in \mathcal{U}$ and any $(s, y, \vec{z}) \in [0, T) \times \mathbb{R} \times \mathbb{R}^n$, $(\vec{m}, U^{s,y,\vec{z}}, \vec{\lambda}^{s,y})$ fulfill the PDEs [\(9\)](#page-7-2), [\(10\)](#page-7-3) and (13) with the square-integrability condition (11) , (12) and (14) .
- 2. $U^{s,y,\vec{z}}(t,x)$ is continuously differentiable in s and twice differentiable in (y,\vec{z}) .
- 3. For $\tilde{U}(t,x) := U^{t,x,\vec{m}(t,x)}(t,x)$, V is a classical solution of

$$
0 = \max_{\zeta \in \mathcal{R}} \left\{ \mathcal{D}_{\zeta} V(t, x) - \mathcal{D}_{\zeta} \tilde{U}(t, x) + \mathcal{D}_{\zeta} U^{t, x, \vec{m}(t, x)}(t, x) + \langle \vec{\lambda}^{t, x}(t, x), \mathcal{D}_{\zeta} \vec{m}(t, x) \rangle \right\},\tag{17}
$$

whereby $\tilde{u}(t, x)$ realizes the maximum, with the terminal condition

$$
V(T, x) = \Phi(T, x, x, x, x^2, \dots, x^n) \equiv \Psi(T, x, x, x, 0, \dots, 0).
$$
 (18)

4. For any $(t, x) \in [0, T) \times \mathbb{R}$ and $f = V_x, \tilde{U}_x$,

$$
\mathbb{E}\bigg[\int_t^T \Big|f(v,X_v^{t,x,\tilde{u}})\sigma\big(v,X_v^{t,x,\tilde{u}},\tilde{u}(v,X_v^{t,x,\tilde{u}})\big)\Big|^2 dv\bigg] < \infty.
$$

Then, \tilde{u} is a CNEC, and $V(t, x) = J(t, x, \tilde{u})$; that is, V is an EVF.

Proof. Plugging the maximizer $\tilde{u}(t, x)$ with the PDEs in [\(9\)](#page-7-2) and [\(10\)](#page-7-3) into the right-hand side of [\(17\)](#page-8-2) yields $\mathcal{D}_{\tilde{u}(t,x)}V(t,x) = \mathcal{D}_{\tilde{u}(t,x)}\tilde{U}(t,x)$. Given the square-integrability conditions for (V_x, \tilde{U}_x) , and that $V(T, x) = \tilde{U}(T, x)$ follows from [\(18\)](#page-8-3) and the terminal conditions in [\(9\)](#page-7-2) and [\(10\)](#page-7-3), we can show $V = \tilde{U}$ by applying Itô's rule to $V(v, X_v^{t,x,\tilde{u}}) - \tilde{U}(v, X_v^{t,x,\tilde{u}})$. Consequently, [\(17\)](#page-8-2) gives [\(15\)](#page-8-1), and then we conclude that \tilde{u} is a CNEC according to Theorem [3.2.](#page-7-1) Moreover, due to Lemma [3.1,](#page-7-8) $V(t, x) = \tilde{U}(t, x) = J(t, x, \tilde{u})$ is an EVF.

Collecting (9) , (10) , (13) , (17) and (18) , we obtain an extended HJB system, which can be regarded as an extension of that in [\(Björk et al.,](#page-28-1) [2017,](#page-28-1) Definition 4.1). Thanks to Theorem [3.2,](#page-7-1) it is not necessary to solve the Bellman equation [\(17\)](#page-8-2) for deriving a CNEC in our approach. To end this section, we consider the special case with $\Phi(\cdot, \cdot, y, \vec{z}) \equiv \tilde{\Phi}(y, \vec{z})$; that is, $\Phi(t, x, y, \vec{z})$ is independent of the initial pair (t, x) . The results are used to investigate some particular problems in Section [5.](#page-15-0) In this case, with $\vec{m}(t,x) = \vec{m}^{\tilde{u}}(t,x)$ corresponding to the CNEC \tilde{u} , the row vector $\vec{\lambda}^{s,y}(t,x) = \mathbb{E}_t[\tilde{\Phi}_{\vec{z}}(X_T^{t,x,\tilde{u}}, \vec{m}(s,y))]$ and the matrix $\mu^{s,y}(t,x) := \mathbb{E}_t[\tilde{\Phi}_{\vec{z}\vec{z}}(X_T^{t,x,\tilde{u}}, \vec{m}(s,y))],$ plugging the result of applying the chain rule to $\mathcal{D}_{\zeta}\tilde{U}(t,x)$ into the right-hand side of [\(17\)](#page-8-2) yields the Bellman equation

$$
0 = \max_{\zeta \in \mathcal{R}} \left\{ \mathcal{D}_{\zeta} V(t, x) - \langle \vec{\lambda}_x^{t, x}(t, x), \vec{m}_x(t, x) \rangle \big(\sigma(t, x, \zeta) \big)^2 + \frac{1}{2} \langle \vec{m}_x(t, x) \mu^{t, x}(t, x), \vec{m}_x(t, x) \rangle \big(\sigma(t, x, \zeta) \big)^2 \right\},\tag{19}
$$

whereby $\tilde{u}(t, x)$ realizes the maximum on the right-hand side.

4 Sufficient condition for ONEC: maximum principle

In this section, we refer to the spike variation method in [\(Yong & Zhou](#page-30-0), [1999](#page-30-0), pp. 124–140) to develop a sufficient maximum principle for our problem. To verify that $\bar{u} \in \mathcal{U}_0$ is an ONEC, it suffices to prove that $J(t, \bar{X}_t, \bar{u}) - J(t, \bar{X}_t, \bar{u}^{t,\epsilon,\zeta}) \ge o(\epsilon)$ for any t and any spike variation $\bar{u}^{t,\epsilon,\zeta} \in \mathcal{U}_0$ of \bar{u} . We fix $t \in [0, T)$ and suppose that the values of the \mathcal{F}_t -measurable random variables (\bar{X}_t, ζ) are given conditioned on \mathcal{F}_t , and then write $X^{\varepsilon} := X^{t, \bar{X}_t, \bar{u}^{t,\varepsilon,\zeta}}$ for short. In addition, for the sake of brevity, we write $\bar{f}(s) := f(s, \bar{X}_s, \bar{u}_s)$ and $\delta f(s) := f(s, \bar{X}_s, \bar{u}_s + \zeta) - \bar{f}(s)$ for $f = b, b_x, b_{xx}, \sigma, \sigma_x, \sigma_{xx}$. To derive an expansion of $J(t, \bar{X}_t, \bar{u}) - J(t, \bar{X}_t, \bar{u}^{t,\epsilon,\zeta})$ with respect to ϵ , let us proceed with the first-order and the second-order variational equations

$$
\begin{cases} dy_s^{\varepsilon} = \bar{b}_x(s)y_s^{\varepsilon}ds + (\bar{\sigma}_x(s)y_s^{\varepsilon} + 1_{\{s \in [t, t + \varepsilon)\}}\delta\sigma(s))dW_s, \quad y_t^{\varepsilon} = 0; \\ dz_s^{\varepsilon} = (\bar{b}_x(s)z_s^{\varepsilon} + \frac{1}{2}\bar{b}_{xx}(s)(y_s^{\varepsilon})^2 + 1_{\{s \in [t, t + \varepsilon)\}}\delta b(s)\Big)ds \\ \qquad + (\bar{\sigma}_x(s)z_s^{\varepsilon} + \frac{1}{2}\bar{\sigma}_{xx}(s)(y_s^{\varepsilon})^2 + 1_{\{s \in [t, t + \varepsilon)\}}\delta\sigma_x(s)y_s^{\varepsilon}\Big)dW_s, \quad z_t^{\varepsilon} = 0. \end{cases} \tag{20}
$$

Following the same line of proof as for [\(Yong & Zhou](#page-30-0), [1999,](#page-30-0) Theorem 4.4, p. 128), we can show that for any positive integer k ,

$$
\begin{cases}\n\sup_{s\in[t,T]} \mathbb{E}_t[|X_s^{\varepsilon} - \bar{X}_s|^{2k}] = O(\varepsilon^k), & \sup_{s\in[t,T]} \mathbb{E}_t[|y_s^{\varepsilon}|^{2k}] = O(\varepsilon^k), \\
\sup_{s\in[t,T]} \mathbb{E}_t[|z_s^{\varepsilon}|^{2k}] = O(\varepsilon^{2k}), & \sup_{s\in[t,T]} \mathbb{E}_t[|X_s^{\varepsilon} - \bar{X}_s - y_s^{\varepsilon}|^{2k}] = O(\varepsilon^{2k}), \\
\sup_{s\in[t,T]} \mathbb{E}_t[|X_s^{\varepsilon} - \bar{X}_s - y_s^{\varepsilon} - z_s^{\varepsilon}|^{2k}] = o(\varepsilon^{2k}),\n\end{cases}
$$
\n(21)

and hence, for $i = 0, 1, \ldots, n$,

$$
\mathbb{E}_t[(X_T^{\varepsilon})^i] = \mathbb{E}_t[(\bar{X}_T)^i] + i\mathbb{E}_t[(\bar{X}_T)^{i-1}(y_T^{\varepsilon} + z_T^{\varepsilon})] + {i \choose 2} \mathbb{E}_t[(\bar{X}_T)^{i-2}(y_T^{\varepsilon})^2] + o(\varepsilon).
$$
 (22)

From [\(21\)](#page-10-1), we have $\sup_{s\in[t,T]} |\mathbb{E}_t[y_s^{\varepsilon}]|^2 \leq \sup_{s\in[t,T]} \mathbb{E}_t[|y_s^{\varepsilon}|^2] = O(\varepsilon)$, which implies that $\mathbb{E}_t[y_s^{\varepsilon}]$ and $\sqrt{\varepsilon}$ might be infinitesimals of the same order. As a consequence, the terms in the form of $\mathbb{E}_t[A_y \mathbb{F}_T] \mathbb{E}_t[B_y \mathbb{F}_T]$ in the expansion of $J(t, \bar{X}_t, \bar{u}^{t,\epsilon,\zeta}) - J(t, \bar{X}_t, \bar{u})$ would seriously interfere with the establishment of adjoint equations. Fortunately, the following lemma ensures that the abovementioned dilemma does not occur.

Lemma 4.1. Fix $\tau \in (t, T]$. Suppose that ξ is an (ε, ζ) -independent \mathcal{F}_{τ} -measurable random variable, and $\mathbb{E}[\left|\xi\right|^{2+\varrho}] < \infty$ for some $\varrho > 0$. Then, $\left|\mathbb{E}_t[\xi y_\tau^{\varepsilon}] \right|^2 = o(\varepsilon)$, and $\sup_{\tau \in [t,T]} \left|\mathbb{E}_t[y_\tau^{\varepsilon}] \right|^2 = o(\varepsilon)$.

Proof. Inspired by the proof for [\(Buckdahn et al.](#page-28-15), [2011](#page-28-15), Proposition 3.1), we introduce the factor

$$
\mathcal{E}_s := \exp\left(\frac{1}{2}\int_t^s \left(\bar{\sigma}_x(v)\right)^2 dv - \int_t^s \bar{\sigma}_x(v) dW_v - \int_t^s \bar{b}_x(v) dv\right),
$$

which solves the SDE $d\mathcal{E}_s = \mathcal{E}_s(\bar{\sigma}_x(s))^2 ds - \mathcal{E}_s(\bar{b}_x(s)ds + \bar{\sigma}_x(s)dW_s)$ and results in

$$
\mathcal{E}_{\tau} y_{\tau}^{\varepsilon} = \int_{t}^{\tau} 1_{\{s \in [t, t + \varepsilon)\}} \mathcal{E}_{s} \delta \sigma(s) \big(dW_{s} - \bar{\sigma}_{x}(s) ds \big) = \int_{t}^{\tau \wedge \varepsilon} \mathcal{E}_{s} \delta \sigma(s) \big(dW_{s} - \bar{\sigma}_{x}(s) ds \big).
$$

By Doob's maximal inequality, for any $p > 1$, we obtain

$$
\mathbb{E}\left[\sup_{s\in[t,T]}|\mathcal{E}_s|^p\right] \leq e^{pT\sup|b_x|+pT\sup|\sigma_x|^2}\mathbb{E}\left[\sup_{s\in[t,T]}|e^{\int_t^s \bar{b}_x(v)dv - \int_t^s (\bar{\sigma}_x(v))^2 dv}\mathcal{E}_s|^p\right]
$$

$$
\leq \left(\frac{p}{p-1}\right)^p e^{pT\sup|b_x|+\frac{1}{2}(p^2+p)T\sup|\sigma_x|^2}.
$$

In the same manner, we can show that

$$
\label{eq:3.16} \begin{aligned} &\mathbb{E}\bigg[\sup_{s\in[t,T]}|\mathcal{E}_s|^{-p}\bigg]\leq \Big(\frac{p}{p-1}\Big)^p e^{pT\sup|\bar{b}_x|+\frac{1}{2}(p^2-p)T\sup|\bar{\sigma}_x|^2},\quad \forall p>1;\\ &\mathbb{E}\bigg[\sup_{s\in[t,T]} \Big|\frac{\mathcal{E}_s}{\mathcal{E}_\tau}\Big|^p\bigg]\leq \Big(\frac{|p|}{|p|-1}\Big)^{|p|} e^{|p|T\sup|\bar{b}_x|+\frac{1}{2}(|p|^2+|p|)T\sup|\bar{\sigma}_x|^2},\quad \forall \tau\in[t,T],\ |p|>1. \end{aligned}
$$

Applying the martingale representation theorem to $\{\mathbb{E}_s[\xi/\mathcal{E}_\tau]\}_{s\in[t,\tau]}$, we conclude that there exists a square-integrable F-predictable process $\{\gamma_{s,\tau}\}_{s\in[t,\tau]}$ such that $\xi/\mathcal{E}_{\tau} = \mathbb{E}_{t}[\xi/\mathcal{E}_{\tau}] + \int_{t}^{\tau} \gamma_{s,\tau} dW_{s}$. Consequently,

$$
\mathbb{E}_t[\xi y_\tau^\varepsilon] = \mathbb{E}_t\bigg[\frac{\xi}{\mathcal{E}_\tau}(\mathcal{E}_\tau y_\tau^\varepsilon)\bigg] = \mathbb{E}_t\bigg[\int_t^{\tau \wedge \varepsilon} \gamma_{s,\tau} \mathcal{E}_s \delta \sigma(s) ds - \int_t^{\tau \wedge \varepsilon} \xi \frac{\mathcal{E}_s}{\mathcal{E}_\tau} \delta \sigma(s) \bar{\sigma}_x(s) ds\bigg],
$$

for which we have the following moment estimates with some appropriate constants (K_1, K_2, K_3, K_4)

independent of τ :

 $\overline{}$ $\overline{}$ $\overline{}$ \mid

$$
\left| \mathbb{E}_{t} \left[\int_{t}^{\tau \wedge \varepsilon} \gamma_{s,\tau} \mathcal{E}_{s} \delta \sigma(s) ds \right] \right|^{2} \leq K_{1} \mathbb{E}_{t} \left[\int_{t}^{\tau \wedge \varepsilon} |\mathcal{E}_{s}|^{2} ds \right] \mathbb{E}_{t} \left[\int_{t}^{\tau \wedge \varepsilon} |\gamma_{s,\tau}|^{2} ds \right] \n\leq K_{2} \varepsilon \mathbb{E}_{t} \left[\int_{t}^{\tau \wedge \varepsilon} |\gamma_{s,\tau}|^{2} ds \right],
$$
\n
$$
\mathbb{E}_{t} \left[\int_{t}^{\tau \wedge \varepsilon} \xi \frac{\mathcal{E}_{s}}{\mathcal{E}_{\tau}} \delta \sigma(s) \bar{\sigma}_{x}(s) ds \right] \right|^{2} \leq K_{3} \mathbb{E}_{t} \left[\int_{t}^{\tau \wedge \varepsilon} \left| \frac{\mathcal{E}_{s}}{\mathcal{E}_{\tau}} \right|^{2} ds \right] \mathbb{E}_{t} \left[\int_{t}^{\tau \wedge \varepsilon} \xi^{2} ds \right] \leq K_{4} \varepsilon^{2}.
$$

Hence, $|\mathbb{E}_{t}[\xi y_{\tau}^{\varepsilon}]|^{2} \leq 2(K_{2} \vee K_{4})(\varepsilon \mathbb{E}_{t}[\int_{t}^{\tau \wedge \varepsilon} |\gamma_{s,\tau}|^{2} ds] + \varepsilon^{2}) = o(\varepsilon)$. Furthermore, setting $\xi = \bar{b}_{x}(\tau)$ yields $|\mathbb{E}_t[y_\tau^\varepsilon \overline{b}_x(\tau)]|^2 \leq K_5(\varepsilon \mathbb{E}_t[\int_t^{\tau \wedge \varepsilon} |\gamma_{s,\tau}|^2 ds] + \varepsilon^2)$ for some constant K_5 independent of τ . Consequently,

$$
\int_t^T |\mathbb{E}_t[y_s^{\varepsilon} \bar{b}_x(s)]|^2 ds \le K_5 \bigg(\varepsilon \int_t^T \int_t^{\tau \wedge \varepsilon} \mathbb{E}_t[|\gamma_{s,\tau}|^2] ds d\tau + T \varepsilon^2 \bigg) = o(\varepsilon),
$$

where the last equality follows from the dominated convergence theorem with

$$
\int_t^\tau \mathbb{E}_t[|\gamma_{s,\tau}|^2]ds = \mathbb{E}_t\bigg[\Big|\frac{\bar{b}_x(\tau)}{\mathcal{E}_\tau} - \mathbb{E}_t\Big[\frac{\bar{b}_x(\tau)}{\mathcal{E}_\tau}\Big]\Big|^2\bigg] \le 4(\sup|b_x|)^2 \mathbb{E}_t\bigg[\sup_{s\in[t,T]}|\mathcal{E}_s|^{-2}\bigg].
$$

Finally, by [\(20\)](#page-9-1) and Cauchy-Schwarz inequality, we obtain

$$
\sup_{\tau \in [t,T]} |\mathbb{E}_t[y_\tau^{\varepsilon}]|^2 = \sup_{\tau \in [t,T]} \left| \int_t^\tau \mathbb{E}_t[y_s^{\varepsilon} \overline{b}_x(s)] ds \right|^2 \le T \int_t^T |\mathbb{E}_t[y_s^{\varepsilon} \overline{b}_x(s)]|^2 ds = o(\varepsilon),
$$

and thus, this proof is complete. \Box

Remark 4.2. In comparison, [\(Buckdahn et al.](#page-28-15), [2011](#page-28-15), Lemma 3.2) for proving Proposition 3.1 therein has used \mathbb{L}^p -integrability for all $p \geq 1$, which is much stronger than our condition. Lemma [4.1](#page-10-0) shows that the condition slightly stronger than square-integrability is sufficient to derive the first-order expansion of $J(t, \bar{X}_t, \bar{u}^{t,\epsilon,\zeta}) - J(t, \bar{X}_t, \bar{u})$ for our problem. By Lemma [4.1,](#page-10-0) one can further obtain the estimates

$$
\begin{aligned} |\mathbb{E}_t[\xi y_\tau^\varepsilon]\mathbb{E}_t[z_\tau^\varepsilon]|&\leq \frac{1}{2}(\mathbb{E}_t[\xi y_\tau^\varepsilon])^2+\frac{1}{2}\mathbb{E}_t[|z_\tau^\varepsilon|^2]=o(\varepsilon),\\ |\mathbb{E}_t[y_\tau^\varepsilon]\mathbb{E}_t[\xi z_\tau^\varepsilon]|&\leq \frac{1}{2}(\mathbb{E}_t[y_\tau^\varepsilon])^2+\frac{1}{2}\mathbb{E}_t[\xi^2]\mathbb{E}_t[|z_\tau^\varepsilon|^2]=o(\varepsilon),\quad etc. \end{aligned}
$$

As a result, many terms in the expansion of $J(t, \bar{X}_t, \bar{u}^{t,\epsilon,\zeta}) - J(t, \bar{X}_t, \bar{u})$ can be shown to be higher order infinitesimals of ε .

For the sake of brevity, we let $M_j^u(t,x) := \mathbb{E}_t[(X_T^{t,x,u} - \mathbb{E}_t[X_T^{t,x,u}])^j]$ for the *j*-th conditional central moment, and $\vec{M}^u(t,x) := (\mathbb{E}_t[X_T^{t,x,u}], M_2^u(t,x), \dots, M_n^u(t,x))$. In addition, for $f = \Psi, \Psi_y, \Psi_{yy}, \Psi_{yz_j}, \Psi_{z_j},$ $\Psi_{z_iz_j}$, we write $\bar{f} := f(t, \bar{X}_t, \bar{X}_T, \vec{M}^{\bar{u}}(t, \bar{X}_t)).$

Lemma 4.3. For an arbitrarily fixed $\bar{u} \in \mathcal{U}_0$, any $t \in [0, T)$, a sufficiently small $\varepsilon \in (0, T - t]$ and an \mathcal{F}_t -measurable random variable ζ such that $\bar{u}^{t,\varepsilon,\zeta} \in \mathcal{U}_0$, we have

$$
J(t, \bar{X}_t, \bar{u}^{t,\varepsilon,\zeta}) = J(t, \bar{X}_t, \bar{u}) + \mathbb{E}_t \Big[\bar{\Psi}_y(y_T^{\varepsilon} + z_T^{\varepsilon}) + \frac{1}{2} \bar{\Psi}_{yy}(y_T^{\varepsilon})^2 \Big] + \mathbb{E}_t [\bar{\Psi}_{z_1}] \mathbb{E}_t [y_T^{\varepsilon} + z_T^{\varepsilon}] + \sum_{j=2}^n j \mathbb{E}_t [\bar{\Psi}_{z_j}] \mathbb{E}_t \big[\big((\bar{X}_T - \mathbb{E}_t [\bar{X}_T])^{j-1} - M_{j-1}^{\bar{u}}(t, x) \big) (y_T^{\varepsilon} + z_T^{\varepsilon}) \big] + \sum_{j=2}^n \binom{j}{2} \mathbb{E}_t [\bar{\Psi}_{z_j}] \mathbb{E}_t \big[(\bar{X}_T - \mathbb{E}_t [\bar{X}_T])^{j-2} (y_T^{\varepsilon})^2 \big] + o(\varepsilon).
$$
 (23)

Proof. It follows from [\(22\)](#page-10-2) and Lemma [4.1](#page-10-0) that

$$
\begin{split} & (\mathbb{E}_{t}[X_{T}^{\varepsilon}])^{j-i}\mathbb{E}_{t}[(X_{T}^{\varepsilon})^{i}] - (\mathbb{E}_{t}[\bar{X}_{T}])^{j-i}\mathbb{E}_{t}[(\bar{X}_{T})^{i}] \\ &= (j-i)1_{\{i \leq j-1\}}(\mathbb{E}_{t}[\bar{X}_{T}])^{j-i-1}\mathbb{E}_{t}[(\bar{X}_{T})^{i}]\mathbb{E}_{t}[y_{T}^{\varepsilon} + z_{T}^{\varepsilon}] \\ &+ 1_{\{i \geq 1\}}i(\mathbb{E}_{t}[\bar{X}_{T}])^{j-i}\mathbb{E}_{t}[(\bar{X}_{T})^{i-1}(y_{T}^{\varepsilon} + z_{T}^{\varepsilon})] \\ &+ 1_{\{i \geq 2\}}\binom{i}{2}(\mathbb{E}_{t}[\bar{X}_{T}])^{j-i}\mathbb{E}_{t}[(\bar{X}_{T})^{i-2}(y_{T}^{\varepsilon})^{2}] + o(\varepsilon), \end{split}
$$

which leads to

$$
M_j^{\bar{u}^{t,\varepsilon,\zeta}}(t,\bar{X}_t) = M_j^{\bar{u}}(t,\bar{X}_t) + j \mathbb{E}_t \left[\left((\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^{j-1} - M_{j-1}^{\bar{u}}(t,\bar{X}_t) \right) (y_T^{\varepsilon} + z_T^{\varepsilon}) \right] + \binom{j}{2} \mathbb{E}_t \left[(\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^{j-2} (y_T^{\varepsilon})^2 \right] + o(\varepsilon), \quad j \ge 2.
$$
 (24)

On the one hand, in the same line of proof as for [\(Yong & Zhou,](#page-30-0) [1999,](#page-30-0) (4.31), p. 128), one can obtain

$$
\mathbb{E}_{t} \left[\Psi(t, \bar{X}_{t}, X_{T}^{\varepsilon}, \vec{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_{t})) \right] - \mathbb{E}_{t} \left[\Psi(t, \bar{X}_{t}, \bar{X}_{T}, \vec{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_{t})) \right]
$$
\n
$$
= \mathbb{E}_{t} \left[\Psi_{y}(t, \bar{X}_{t}, \bar{X}_{T}, \vec{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_{t})) (y_{T}^{\varepsilon} + z_{T}^{\varepsilon}) \right]
$$
\n
$$
+ \frac{1}{2} \mathbb{E}_{t} \left[\Psi_{yy}(t, \bar{X}_{t}, \bar{X}_{T}, \vec{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_{t})) (y_{T}^{\varepsilon})^{2} \right] + o(\varepsilon).
$$

By (21) , (24) , Lemma [4.1](#page-10-0) and the Lipschitz continuity of (Φ, Φ_y) , we obtain

$$
\mathbb{E}_t[\Psi_y(t, \bar{X}_t, \bar{X}_T, \bar{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_t))y_T^{\varepsilon}] - \mathbb{E}_t[\bar{\Psi}_y y_T^{\varepsilon}] = o(\varepsilon),
$$

\n
$$
\mathbb{E}_t[\Psi_y(t, \bar{X}_t, \bar{X}_T, \bar{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_t))z_T^{\varepsilon}] - \mathbb{E}_t[\bar{\Psi}_y z_T^{\varepsilon}] = o(\varepsilon),
$$

\n
$$
\mathbb{E}_t[\Psi_{yy}(t, \bar{X}_t, \bar{X}_T, \bar{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_t))(y_T^{\varepsilon})^2] - \mathbb{E}_t[\bar{\Psi}_{yy}(y_T^{\varepsilon})^2] = o(\varepsilon).
$$

On the other hand, by Taylor's expansion with the use of [\(21\)](#page-10-1), [\(24\)](#page-12-0) and Lemma [4.1,](#page-10-0) we can show that

$$
\mathbb{E}_{t} \left[\Psi(t, \bar{X}_{t}, \bar{X}_{T}, \bar{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_{t})) \right] - \mathbb{E}_{t} [\bar{\Psi}] - \mathbb{E}_{t} [\bar{\Psi}_{z_{1}}] \mathbb{E}_{t} [y_{T}^{\varepsilon} + z_{T}^{\varepsilon}]
$$
\n
$$
= \sum_{j=2}^{n} j \mathbb{E}_{t} [\bar{\Psi}_{z_{j}}] \mathbb{E}_{t} \left[\left((\bar{X}_{T} - \mathbb{E}_{t} [\bar{X}_{T}])^{j-1} - M_{j-1}^{\bar{u}}(t, \bar{X}_{t}) \right) (y_{T}^{\varepsilon} + z_{T}^{\varepsilon}) \right]
$$
\n
$$
+ \sum_{j=2}^{n} \frac{j(j-1)}{2} \mathbb{E}_{t} [\bar{\Psi}_{z_{j}}] \mathbb{E}_{t} \left[(\bar{X}_{T} - \mathbb{E}_{t} [\bar{X}_{T}])^{j-2} (y_{T}^{\varepsilon})^{2} \right] + o(\varepsilon).
$$

Therefore, plugging the abovementioned expansions into the decomposition

$$
J(t, \bar{X}_t, \bar{u}^{t,\varepsilon,\zeta}) - J(t, \bar{X}_t, \bar{u})
$$

=
$$
\left(\mathbb{E}_t \big[\Psi(t, \bar{X}_t, X_T^{\varepsilon}, \bar{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_t)) \big] - \mathbb{E}_t \big[\Psi(t, \bar{X}_t, \bar{X}_T, \bar{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_t)) \big] \right)
$$

$$
+ \left(\mathbb{E}_t \big[\Psi(t, \bar{X}_t, \bar{X}_T, \bar{M}^{\bar{u}^{t,\varepsilon,\zeta}}(t, \bar{X}_t)) \big] - \mathbb{E}_t[\bar{\Psi}] \right)
$$

immediately yields the desired result. $\hfill \square$

Now we are able to state the maximum principles for our control problems.

Theorem 4.4. Suppose that corresponding to $\bar{u} \in \mathcal{U}_0$, the following FBSDE admits a square-integrable

solution $(\bar{X}, Y^t, \mathcal{Y}^t, \mathcal{Z}^t, \mathcal{Z}^t)$ for a.e. $t \in [0, T)$:

$$
\begin{cases}\nd\bar{X}_s = \bar{b}(s)ds + \bar{\sigma}(s)dW_s, \ \forall s \in [0, T], \quad \bar{X}_0 = x_0; \\
dY_s^t = -(Y_s^t \bar{b}_x(s) + Y_s^t \bar{\sigma}_x(s))ds + Y_s^t dW_s, \ \forall s \in [t, T], \\
Y_T^t = \bar{\Psi}_y + \mathbb{E}_t[\bar{\Psi}_{z_1}] + \sum_{j=2}^n j \mathbb{E}_t[\bar{\Psi}_{z_j}]((\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^{j-1} - M_{j-1}^{\bar{u}}(t, \bar{X}_t)); \\
dZ_s^t = -(2Z_s^t \bar{b}_x(s) + Z_s^t |\bar{\sigma}_x(s)|^2 + 2Z_s^t \bar{\sigma}_x(s) + Y_s^t \bar{b}_{xx}(s) + Y_s^t \bar{\sigma}_{xx}(s))ds \\
&+ Z_s^t dW_s, \ \forall s \in [t, T], \\
Z_T^t = \bar{\Psi}_{yy} + \sum_{j=2}^n j(j-1) \mathbb{E}_t[\bar{\Psi}_{z_j}](\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^{j-2},\n\end{cases} \tag{25}
$$

such that

$$
\limsup_{s\downarrow t}\mathbb{E}_t\bigg[\frac{1}{2}Z_s^t\big(|\sigma(s,\bar{X}_s,\bar{u}_s+\zeta)|^2-|\bar{\sigma}(s)|^2\big)+Y_s^t\delta b(s)+\big(\mathcal{Y}_s^t-Z_s^t\bar{\sigma}(s)\big)\delta\sigma(s)\bigg]\leq 0
$$

for any \mathcal{F}_t -measurable random variable ζ satisfying $\bar{u}^{t,\varepsilon,\zeta} \in \mathcal{U}_0$. Then, \bar{u} is an ONEC.

Proof. According to the FBSDE [\(25\)](#page-13-0), [\(23\)](#page-11-0) can be re-expressed as

$$
J(t, \bar{X}_t, \bar{u}^{t,\varepsilon,\zeta}) - J(t, \bar{X}_t, \bar{u}) = \mathbb{E}_t[Y_T^t(y_T^{\varepsilon} + z_T^{\varepsilon})] + \frac{1}{2}\mathbb{E}_t[Z_T^t(y_T^{\varepsilon})^2] + o(\varepsilon).
$$

In the same line of duality analysis as in [\(Yong & Zhou](#page-30-0), [1999,](#page-30-0) Section 3.4.3, pp. 134-137), we obtain

$$
J(t, \bar{X}_t, \bar{u}^{t,\varepsilon,\zeta}) - J(t, \bar{X}_t, \bar{u})
$$

\n
$$
= \mathbb{E}_t \bigg[\int_t^{t+\varepsilon} \left(Y_s^t \delta b(s) + Y_s^t \delta \sigma(s) + \frac{1}{2} Z_s^t |\delta \sigma(s)|^2 \right) ds \bigg] + o(\varepsilon)
$$

\n
$$
= \mathbb{E}_t \bigg[\int_t^{t+\varepsilon} \left(\frac{1}{2} Z_s^t (|\sigma(s, \bar{X}_s, \bar{u}_s + \zeta)|^2 - |\bar{\sigma}(s)|^2) + Y_s^t \delta b(s) + (\mathcal{Y}_s^t - Z_s^t \bar{\sigma}(s)) \delta \sigma(s) \right) ds \bigg] + o(\varepsilon) \leq o(\varepsilon),
$$
\n(26)

which implies that \bar{u} is an ONEC.

Theorem 4.5. Suppose that $b(t, x, u)$ and $\sigma(t, x, u)$ are both affine in u. Then, $\bar{u} \in \mathcal{U}_0$ is an ONEC, if for a.e. $t \in [0, T)$, the FBSDE [\(25\)](#page-13-0) admits a square-integrable solution $(\bar{X}, Y^t, \mathcal{Y}^t, \mathcal{Z}^t, \mathcal{Z}^t)$ such that

$$
\lim_{s\downarrow t} \mathbb{E}_t[Y_s^t b_u(s, \bar{X}_s, u) + \mathcal{Y}_s^t \sigma_u(s, \bar{X}_s, u)] = 0, \quad \lim_{s\downarrow t} \mathbb{E}_t[Z_s^t] \le 0, \quad \mathbb{P}-a.s.
$$
\n(27)

Proof. Given that $\lim_{s \downarrow t} \mathbb{E}_t[Z_s^t] \leq 0$, from the first equality in [\(26\)](#page-13-3) we have

$$
\lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \big(J(t, \bar{X}_t, \bar{u}^{t,\varepsilon,\zeta}) - J(t, \bar{X}_t, \bar{u}) \big) \le \lim_{\varepsilon \downarrow 0} \frac{\zeta}{\varepsilon} \int_t^{t+\varepsilon} \mathbb{E}_t[Y_s^t b_u(s, \bar{X}_s, u) + \mathcal{Y}_s^t \sigma_u(s, \bar{X}_s, u)] ds.
$$

Due to the square-integrability of (Y^t, Y^t) and the Lipschitz continuity of $b(t, x, \cdot)$ and $\sigma(t, x, \cdot)$, we obtain $\mathbb{E}_t \int_t^T |Y_s^t b_u(s, \bar{X}_s, u) + \mathcal{Y}_s^t \sigma_u(s, \bar{X}_s, u)| ds < \infty.$ Hence,

$$
\lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \big(J(t, \bar{X}_t, \bar{u}^{t, \varepsilon, \zeta}) - J(t, \bar{X}_t, \bar{u}) \big) \le \zeta \lim_{s \downarrow t} \mathbb{E}_t \big[Y_s^t b_u(s, \bar{X}_s, u) + \mathcal{Y}_s^t \sigma_u(s, \bar{X}_s, u) \big] = 0,
$$

which implies that \bar{u} is an ONEC.

Example 4.6. As mentioned in Remark [2.7,](#page-6-0) we now consider $\Psi(\vec{z}) = g(p^t(\cdot, \vec{z}))$, where $p^t(\cdot, \vec{z})$ is a \vec{z} -indexed regular conditional law given \mathcal{F}_t under \mathbb{P} . This setting captures some non-linear law-dependent preferences as in [\(Liang et al.](#page-29-22), [2023\)](#page-29-22), as $p^t(\cdot, \vec{M}^u(t,x))$ could be the conditional law of $X_T^{t,x,u}$ characterized by its conditional expectation and higher-order central moments. Let $\mathcal{B}(\mathbb{R})$ be the Borel σ -algebra generated by all open intervals in \mathbb{R}, \mathcal{P} be the Banach space of all finite signed measures on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$, the non-empty convex set $\mathcal{P}_0 \subseteq \mathcal{P}$ be the domain of the fairly general functional g, and $\mathcal{L}(\mathcal{S}_1, \mathcal{S}_2)$ be the collection of all continuous linear functions from S_1 to S_2 . In addition to the basic assumptions for Ψ in Section [2,](#page-4-0) we shall introduces the following assumptions that can be viewed as continuous Gâteaux differentiability:

1. There exists $\nabla g : \mathcal{P}_0 \times \mathbb{R} \to \mathbb{R}$ such that for any $p, q \in \mathcal{P}_0$,

$$
\lim_{\varepsilon \downarrow 0} \frac{g(p + \varepsilon (q - p)) - g(p)}{\varepsilon} =: dg(p, q - p) = \int_{\mathbb{R}} \nabla g(p, x) \big(q(dx) - p(dx) \big),
$$

 $\int_{\mathbb{R}} |\nabla g(p,x)| q(dx) < \infty$, and $p \mapsto dg(p, \cdot) \in \mathcal{L}(\mathcal{P}, \mathbb{R})$ is continuous.

2. For any $p \in \mathcal{P}_0$, there exists $p_{\vec{z}} : \mathcal{B}(\mathbb{R}) \times \mathbb{R}^n \to \mathbb{R}^n$ such that for any $\vec{z}, \vec{h} \in \mathbb{R}^n$,

$$
\lim_{\varepsilon \downarrow 0} \frac{p(B, \vec{z} + \varepsilon \vec{h}) - p(B, \vec{z})}{\varepsilon} = \langle p_{\vec{z}}(B, \vec{z}), \vec{h} \rangle, \quad \forall B \in \mathcal{B}(\mathbb{R}),
$$

and $\vec{z} \mapsto \int_{\mathbb{R}} \nabla g(q,x) \langle p_{\vec{z}}(dx,\vec{z}), \cdot \rangle \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^n)$ is continuous for any $q \in \mathcal{P}_0$.

Notably, ∇g is not unique due to $\int_{\mathbb{R}} (q(dx) - p(dx)) = 0$ for probability measures (p, q) . By the chain rule, $\Psi_{\vec{z}}(\vec{z}) = \int_{\mathbb{R}} \nabla g(p^t(\cdot, \vec{z}), x) p_{\vec{z}}^t(dx)$ follows. Then, according to Theorems [4.4](#page-12-1) and [4.5,](#page-13-1) the ONEC for $\Psi(\vec{M}^u(t,x)) = g(p^t(\cdot, \vec{M}^u(t,x)))$ can be characterized by [\(25\)](#page-13-0) with

$$
\begin{cases}\nY_T^t = \int_{\mathbb{R}} \nabla g\Big(p^t(\cdot, \vec{M}^{\bar{u}}(t,x)), x\Big) \langle p_{\bar{z}}^t(dx, \vec{M}^{\bar{u}}(t,x)), \vec{Q}(\bar{X}_T, \vec{M}^{\bar{u}}(t,x)) \rangle, \\
Z_T^t = \int_{\mathbb{R}} \nabla g\Big(p^t(\cdot, \vec{M}^{\bar{u}}(t,x)), x\Big) \langle p_{\bar{z}}^t(dx, \vec{M}^{\bar{u}}(t,x)), \vec{Q}_x(\bar{X}_T, \vec{M}^{\bar{u}}(t,x)) \rangle,\n\end{cases} \tag{28}
$$

where $\vec{Q}(x, \vec{z}) := (Q_1(x, \vec{z}), \dots, Q_n(x, \vec{z}))$ for $Q_1(x, \vec{z}) := 1$, $Q_2(x, \vec{z}) := 2(x - z_1)$ and other $Q_j(x, \vec{z}) :=$ $j(x-z_1)^{j-1} - jz_{j-1}$, and $\vec{Q}_x(x,\vec{z}) := \partial_x \vec{Q}(x,\vec{z})$. On the other hand, we write $\vec{M}_X^t := (\mathbb{E}_t[X], \mathbb{E}_t[(X - \vec{Q}_x(X_t), \vec{Q}_x(X_t), \vec{Q}_x(X_t))$ $\mathbb{E}_t[X])^2], \ldots, \mathbb{E}_t[(X - \mathbb{E}_t[X])^n]$ analogous to $\vec{M}^u(t, X_t)$, and then obtain

$$
\lim_{\varepsilon \to 0} \frac{1}{\varepsilon} \left\| \vec{M}^t_{X + \varepsilon \Delta} - \vec{M}^t_X - \mathbb{E}_t \left[\int_X^{X + \varepsilon \Delta} \vec{Q}(v, \vec{M}^t_X) \rangle dv \right] \right\| = 0,
$$

where $\|\cdot\|$ is the Euclidean norm, from

$$
\mathbb{E}_t[(X+\varepsilon\Delta-\mathbb{E}_t[X+\varepsilon\Delta])^j] - \mathbb{E}_t[(X-\mathbb{E}_t[X])^j] = \mathbb{E}_t\bigg[\int_X^{X+\varepsilon\Delta} Q_j(v, \vec{M}_X)dv\bigg] + o(\varepsilon).
$$

Consequently, for an arbitrarily fixed $c \in \mathbb{R}$,

$$
g(p^t(\cdot, \vec{M}^t_{X+\varepsilon\Delta})) - g(p^t(\cdot, \vec{M}^t_X))
$$

\n
$$
= \mathbb{E}_t \bigg[\int_X^{X+\varepsilon\Delta} dv \int_{\mathbb{R}} \nabla g(p^t(\cdot, \vec{M}^t_X), x) \langle p^t_{\vec{z}}(dx, \vec{M}^t_X), \vec{Q}(v, \vec{M}^t_X) \rangle \bigg] + o(\varepsilon)
$$

\n
$$
= \int_{\mathbb{R}} \bigg(\int_c^s \int_{\mathbb{R}} \nabla g(p^t(\cdot, \vec{M}^t_X), x) \langle p^t_{\vec{z}}(dx, \vec{M}^t_X), \vec{Q}(v, \vec{M}^t_X) \rangle dv \bigg) (\mathbb{P}^t_{X+\varepsilon\Delta}(ds) - \mathbb{P}^t_X(ds))
$$

 $+ o(\varepsilon)$

where \mathbb{P}_{X}^{t} denotes the regular conditional law of X given \mathcal{F}_{t} under \mathbb{P} . This implies that if fixing p^{t} and writing $G(\mathbb{P}_{X}^{t}) = g(p^{t}(\cdot, \vec{M}_{X}^{t}))$, then one can choose

$$
\nabla G(\mathbb{P}_X^t, x) = \int_c^x \int_{\mathbb{R}} \nabla g(p^t(\cdot, \vec{M}_X^t), s) \langle p_{\vec{z}}^t(ds, \vec{M}_X^t), \vec{Q}(v, \vec{M}_X^t) \rangle dv, \quad \forall c \in \mathbb{R},
$$

so that [\(28\)](#page-14-0) can be re-expressed as

$$
Y^t_T=\partial_x \nabla G(\mathbb{P}^t_{\bar{X}_T},x)|_{x=\bar{X}_T},\quad Z^t_T=\partial_x^2 \nabla G(\mathbb{P}^t_{\bar{X}_T},x)|_{x=\bar{X}_T}.
$$

Furthermore, if $p^t(\cdot, \vec{M}_{\bar{X}_T}^t) = \mathbb{P}_{\bar{X}_T}^t$, then $G \equiv g$, and the corresponding verification result in [\(Liang et al.,](#page-29-22) [2023](#page-29-22)) gets recovered to some extent. Notably, taking advantage of the second-order adjoint equation for Z^t , we do not need to impose the concavity condition on $\nabla g(\mathbb{P}^t_{\bar{X}_T},\cdot)$ as in [\(Liang et al.,](#page-29-22) [2023](#page-29-22), Assumption 3.1). For instance, if the functions (b, σ) are affine in each of (x, u) as in [\(Liang et al.,](#page-29-22) [2023](#page-29-22)), then it follows from [\(25\)](#page-13-0) that

$$
Z_s^t = \mathbb{E}_s \left[Z_T^t e^{2 \int_s^T \bar{\sigma}_x(v) dW_v - \int_s^T |\bar{\sigma}_x(v)|^2 dv + 2 \int_s^T \bar{b}_x(v) dv \right],
$$

which automatically satisfies that $\lim_{s\downarrow t} \mathbb{E}_t[Z_s^t] \leq 0$ under the concavity assumption of $\nabla g(\mathbb{P}_{X_T}^t, \cdot)$.

5 Particular cases with linear controlled SDE

Let $\mathcal{R} = \mathbb{R}$, $b(t, x, u) = A_t x + B_t u + C_t$ and $\sigma(t, x, u) = D_t u + F_t$, where $A, B, C, D, F : [0, T] \to \mathbb{R}$ are deterministic continuous functions with $|D_t| > 0$ for every t, and the constant $\kappa \geq 0$. Suppose that $\Psi(t, x, X_T, \vec{z})$ is independent of x and X_T and affine in z_1 , and write it as $\kappa z_1 + \psi(t, z_2, \dots, z_n)$. That is, the objective function is

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] + \psi(t, M_2^u(t, x), \dots, M_n^u(t, x)),
$$
\n(29)

and the dynamics of $X^{t,x,u}$ evolve as the linear controlled SDE:

$$
dX_s^{t,x,u} = (A_s X_s^{t,x,u} + B_s u_s + C_s)ds + (D_s u_s + F_s)dW_s, \quad X_t^{t,x,u} = x.
$$
 (30)

In addition, for the sake of brevity, we let $\vec{M}_{-1}^u(t,x) := (M_2^u(t,x), \ldots, M_n^u(t,x)), \alpha_0(\cdot) = 1, \alpha_1(\cdot) = 0,$ $\vec{\alpha}(y) := (\alpha_2(y), \dots, \alpha_n(y))$ for $\alpha_j(y) := 1_{\{j \in 2\mathbb{N}\}}(j-1)!!y^{j/2}, y_t^{\beta} := \int_t^T |D_s \beta_s|^2 ds,$

$$
\theta_t := \int_t^T \Big|\frac{B_s}{D_s}\Big|^2 ds,\quad and \quad \Theta(t,x) := xe^{\int_t^T A_v dv} + \int_t^T e^{\int_s^T A_v dv} \Big(C_s - \frac{B_s}{D_s} F_s\Big) ds.
$$

Remark 5.1. The objective function [\(29\)](#page-15-1) arises from the theory of efficient frontier. See also [\(Li & Ng,](#page-29-9) [2000](#page-29-9); [Zhou & Li](#page-30-6), [2000\)](#page-30-6). Generally, the investors try to make trade-offs between the mean and the combination of higher-order central moments, which respectively stand for reward and risk. To see an economic insight, one can let the state X be the household/individual wealth, and u be the investment amount on share, with the interest rate A, the volatility rate D, the market price of risk B/D , the income rate C and the white noise coefficient F . Notably, in the coming Theorem [5.2,](#page-16-0) the CNEC is not only independent of the wealth level X , but also independent of the income rate C .

5.1 Closed-loop Nash equilibrium control

According to [\(8\)](#page-7-9),

$$
\Phi(t, \cdot, \cdot, \vec{z}) \equiv \phi(t, \vec{z}) := \kappa z_1 + \psi\bigg(t, z_2 - z_1^2, \dots, \sum_{j=0}^n \binom{n}{j} (-1)^j z_1^j z_{n-j}\bigg). \tag{31}
$$

Then, we conclude that $U^{s,\cdot,\vec{z}}(\cdot,\cdot) \equiv \phi(s,\vec{z})$ and $\vec{\lambda}^{s,y}(\cdot,\cdot) \equiv \phi_{\vec{z}}(s,\vec{m}^{\tilde{u}}(s,y))$ for Theorem [3.2.](#page-7-1) Therefore, to determine a CNEC, it suffices to find a solution for

$$
\begin{cases}\n\mathcal{D}_{\tilde{u}(t,x)}\vec{m}(t,x) = 0, \quad s.t. \quad \vec{m}(T,x) = (x, x^2, \dots, x^n); \\
\tilde{u}(t,x) = -\frac{B_t}{|D_t|^2} \frac{\langle \phi_{\tilde{z}}(t, \vec{m}(t,x)), \vec{m}_x(t,x) \rangle}{\langle \phi_{\tilde{z}}(t, \vec{m}(t,x)), \vec{m}_{xx}(t,x) \rangle} - \frac{F_t}{D_t}; \\
0 > \langle \phi_{\tilde{z}}(t, \vec{m}(t,x)), \vec{m}_{xx}(t,x) \rangle.\n\end{cases}
$$
\n(32)

Theorem 5.2. Let $\{\beta_t\}_{t\in[0,T]}$ fulfill the integral equation

$$
\frac{\kappa B_t}{|D_t|^2} + \beta_t \sum_{1 \le j \le \frac{n}{2}} 2j(2j-1)\alpha_{2j-2}(y_t^{\beta})\psi_{z_{2j}}(t, \vec{\alpha}(y_t^{\beta})) = 0,
$$
\n(33)

and satisfy the condition

$$
\sum_{1 \le j \le \frac{n}{2}} j(2j-1)\alpha_{2j-2}(y_t^{\beta})\psi_{z_{2j}}(t, \vec{\alpha}(y_t^{\beta})) < 0.
$$
\n(34)

Then, $(\tilde{u}, \tilde{m}^{\tilde{u}})$ determined by

$$
\tilde{u}(t,x) = \beta_t e^{-\int_t^T A_v dv} - \frac{F_t}{D_t}
$$
\n(35)

solves the PDE system (32) , and hence, \tilde{u} is a CNEC for the control problem with the objective function [\(29\)](#page-15-1) and the controlled SDE [\(30\)](#page-15-2).

Proof. Plugging [\(35\)](#page-16-2) into the SDE in [\(30\)](#page-15-2) yields

$$
dX_s^{t,x,\tilde{u}} = \left(A_s X_s^{t,x,\tilde{u}} + B_s \beta_s e^{-\int_s^T A_v dv} + C_s - \frac{B_s}{D_s} F_s\right) ds + D_s \beta_s e^{-\int_s^T A_v dv} dW_s,
$$

which leads to $X_T^{t,x,\tilde{u}} = \Theta(t,x) + \int_t^T B_s \beta_s ds + \int_t^T D_s \beta_s dW_s$. Since $\partial_x X_T^{t,x,\tilde{u}} = \exp(\int_t^T A_v dv)$ is deterministic, then

$$
\begin{cases}\n\partial_x m_j^{\tilde{u}}(t,x) = j \mathbb{E}_t[(X_T^{t,x,\tilde{u}})^{j-1}] \partial_x X_T^{t,x,\tilde{u}} = j m_{j-1}^{\tilde{u}}(t,x) e^{\int_t^T A_v dv}, \\
\partial_x^2 m_j^{\tilde{u}}(t,x) = j \partial_x m_{j-1}^{\tilde{u}}(t,x) e^{\int_t^T A_v dv} = j(j-1) m_{j-2}^{\tilde{u}}(t,x) e^{\int_t^T A_v dv}.\n\end{cases}
$$
\n(36)

Since $X_T^{t,x,\tilde{u}} - \mathbb{E}_t[X_T^{t,x,\tilde{u}}] \sim N(0, y_t^{\beta}),$ we can show that

$$
\sum_{j=0}^{i} {i \choose j} (-1)^{i-j} (m_1^{\tilde{u}}(t, x))^{i-j} m_j^{\tilde{u}}(t, x) \equiv M_i^{\tilde{u}}(t, x) = \alpha_i(y_t^{\beta}), \quad \forall i \in \mathbb{N}.
$$
 (37)

Consequently, $\phi(t, \vec{m}^{\tilde{u}}(t, x)) = \kappa m_1^{\tilde{u}}(t, x) + \psi(t, \vec{\alpha}(y_t^{\beta}))$ follows from [\(29\)](#page-15-1), and hence,

$$
\langle \phi_{\vec{z}}(t, \vec{m}^{\tilde{u}}(t, x)), \vec{m}^{\tilde{u}}_{x}(t, x) \rangle = \partial_x \phi(t, \vec{m}^{\tilde{u}}(t, x)) = \kappa \partial_x m_1^{\tilde{u}}(t, x) = \kappa e^{\int_t^T A_v dv},
$$

while it follows from (31) , (36) and (37) that

$$
\langle \phi_{\vec{z}}(t, \vec{m}^{\tilde{u}}(t, x)), \vec{m}^{\tilde{u}}_{xx}(t, x) \rangle = \sum_{j=2}^{n} \sum_{i=j}^{n} {i \choose j} \left(-m^{\tilde{u}}_{1}(t, x) \right)^{i-j} \psi_{z_{i}}(t, \vec{\alpha}(y^{\beta}_{t})) \partial_{x}^{2} m_{j}^{\tilde{u}}(t, x)
$$

$$
= e^{2 \int_{t}^{T} A_{v} dv} \sum_{i=2}^{n} i(i-1) \alpha_{i-2}(y^{\beta}_{t}) \psi_{z_{i}}(t, \vec{\alpha}(y^{\beta}_{t}))
$$

$$
= e^{2 \int_{t}^{T} A_{v} dv} \sum_{1 \leq j \leq \frac{n}{2}} 2j(2j-1) \alpha_{2j-2}(y^{\beta}_{t}) \psi_{z_{2j}}(t, \vec{\alpha}(y^{\beta}_{t})).
$$

Therefore, $\langle \phi_{\vec{z}}(t, \vec{m}(t, x)), \vec{m}_{xx}^{\tilde{u}}(t, x) \rangle < 0$ immediately arises from the condition [\(34\)](#page-16-6). On the other hand, it follows from the equation [\(33\)](#page-16-7) that

$$
-\frac{B_t}{|D_t|^2}\frac{\langle \phi_{\vec{z}}(t,\vec{m}(t,x)),\vec{m}_x(t,x)\rangle}{\langle \phi_{\vec{z}}(t,\vec{m}(t,x)),\vec{m}_{xx}(t,x)\rangle}=\beta_t e^{-\int_t^T A_v dv}.
$$

Hence, $(\tilde{u}, \tilde{m}^{\tilde{u}})$ determined by [\(35\)](#page-16-2) solves the PDE system [\(32\)](#page-16-1).

Remark 5.3. In view of [\(33\)](#page-16-7) and [\(34\)](#page-16-6), one can find that corresponding to the CNEC \tilde{u} given by [\(35\)](#page-16-2), $X_T^{t,x,\tilde{u}}$ is normally distributed, and hence all of its $(2j+1)$ -th conditional central moment $\alpha_i(y_t^{\beta})$ vanish. Consequently, \tilde{u} does not vary as the preferences on those odd-order central moments change. In other words, if \tilde{u} given by [\(33\)](#page-16-7)–[\(35\)](#page-16-2) is a CNEC for the objective function [\(29\)](#page-15-1), then it is also a CNEC for

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] + \psi(t, M_2^u(t, x), 0, M_4^u(t, x), 0, \dots, 1_{\{n \in 2\mathbb{N}\}} M_n^u(t, x)),
$$

Furthermore, for any odd polynomial function $P(t, \cdot)$, \tilde{u} is also a CNEC for

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] + \psi(t, \vec{M}_{-1}^u(t, x)) + \mathbb{E}_t[P(t, X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])].
$$

Intuitively speaking, the independence of the preferences on the odd-order central moments for the CNEC \tilde{u} meets the risk attitude that a positive odd-order central moment has both benefits (e.g., a long right tail) and drawbacks (e.g., a small mode or median), as we have mentioned in Section [1.](#page-0-0) As far as results are concerned, the CNEC does not only sacrifice those benefits, but also avoids those drawbacks, so that agents with different risk attitudes towards odd-order central moments can get the same time-consistent strategy. However, this explanation strongly relies on the artificial structure of our objective function [\(29\)](#page-15-1), and is likely to fail for more general problems. For instance, if we only change the constant κ therein to some function $\kappa(x)$, then the proof of Theorem [5.2](#page-16-0) no longer holds, and one may obtain an x-dependent $\tilde{u}(t, x)$ distinct from [\(35\)](#page-16-2). Thus, this leads to a loss of the independence of the preferences on the odd-order central moments for the CNEC \tilde{u} . Summing up, for the sake of rigor, we prefer to think of this independence as a mathematical consequence, rather than a necessary result for some intuitive characteristics.

If $\kappa = 0$, then $\beta \equiv 0$ solves [\(33\)](#page-16-7). Moreover, if $\psi_{z_2}(t, \vec{0}) < 0$ arising from [\(34\)](#page-16-6) holds for every t, then $\tilde{u}(t, x) = -F_t/D_t$ gives a CNEC such that the corresponding X_T is deterministic. The case with $\kappa > 0$ is slightly more complicated. The following theorem provides a sufficient condition for the existence and uniqueness of solution for [\(33\)](#page-16-7) with $\kappa > 0$.

Theorem 5.4. Suppose that

$$
f(t,y) := -\bigg(\sum_{1 \le j \le \frac{n}{2}} 2j(2j-1)\alpha_{2j-2}(y)\psi_{z_{2j}}(t, \vec{\alpha}(y))\bigg)^{-1}
$$

is strictly positive and uniformly bounded over $[0, T] \times [0, \infty)$. Then, [\(33\)](#page-16-7) with $\kappa > 0$ admits a unique classical solution β .

Proof. [\(33\)](#page-16-7) can be re-expressed as $|D_t|^2 \beta_t = \kappa B_t f(t, y_t^{\beta})$. If β solves it, then

$$
0 \le \beta_t = \kappa B_t |D_t|^{-2} f(t, y_t^{\beta}) \le \kappa B_t |D_t|^{-2} \sup_{y \ge 0} f(t, y) < \infty,
$$

$$
0 \le y_t^{\beta} \le \int_0^T |D_s \beta_s|^2 ds \le T \kappa^2 (\inf D)^{-2} (\sup |B|^2) (\sup |f|^2) < \infty,
$$

and $y = y^{\beta}$ solves $y'_t + |\kappa B_t|^2 |D_t|^{-2} |f(t, y_t)|^2 = 0$. Conversely, we conclude that $|f(t, \cdot)|^2$ is Lipschitz continuous on $[0, T\kappa^2(\inf D)^{-2}(\sup |B|^2)(\sup |f|^2)],$ since

$$
\left| \left(f(t,y) \right)^2 - \left(f(t,z) \right)^2 \right| = |f(t,y)f(t,z)||f(t,y) + f(t,z)| \left| \frac{1}{f(t,y)} - \frac{1}{f(t,z)} \right|, \quad \forall y, z \ge 0.
$$

Then, due to the existence and uniqueness theorem for first-order ordinary differential equation, there exists a unique $\{y_t\}_{t\in[0,T]}$ such that $|D_t|^2 y_t' + |\kappa B_t f(t, y_t)|^2 = 0$ with $y_T = 0$. Hence, $\beta_t = \kappa B_t |D_t|^{-2} f(t, y_t)$ gives the unique solution to [\(33\)](#page-16-7).

Apart from the result given by Theorem [5.2,](#page-16-0) according to Lemma [3.1](#page-7-8) and Theorem [3.3,](#page-8-4) seeking a CNEC can be reduced to solving the following Bellman equation:

$$
0 = \max_{\zeta \in \mathbb{R}} \left\{ V_t(t, x) + V_x(t, x)(A_t x + B_t u + C_t) + \frac{1}{2} V_{xx}(t, x)(D_t \zeta + F_t)^2 - \phi_t(t, \vec{m}^{\tilde{u}}(t, x)) - \frac{1}{2} (D_t \zeta + F_t)^2 \langle \vec{m}^{\tilde{u}}_x(t, x) \phi_{\vec{z}\vec{z}}(t, \vec{m}^{\tilde{u}}(t, x)), \vec{m}^{\tilde{u}}_x(t, x) \rangle \right\},
$$
(38)

subject to the terminal condition $V(T, x) = \kappa x + \psi(T, \vec{0}).$

Theorem 5.5. Suppose that the integral equation [\(33\)](#page-16-7) admits the solution β such that [\(34\)](#page-16-6) holds, $\int_t^T B_s \beta_s ds < \infty$ and $y_t^{\beta} \equiv \int_t^T |D_s \beta_s|^2 ds < \infty$. Then,

$$
V(t,x) = \kappa \Theta(t,x) + \kappa \int_{t}^{T} B_{s} \beta_{s} ds + \psi(t, \vec{\alpha}(y_{t}^{\beta}))
$$
\n(39)

fulfills (38) , and (35) realizes the maximum on the right-hand side of (38) .

Proof. For twice differentiable functions $H(\vec{z})$ and $h(\vec{z})$ with

$$
H(\vec{z}) = h\bigg(z_1, z_2 - z_1^2, z_3 - 3z_2z_1 + 2z_1^3, \dots, \sum_{j=0}^n \binom{n}{j} (-1)^j z_1^j z_{n-j}\bigg),
$$

by straightforward calculation, one can obtain

$$
\sum_{i=1}^{n} \sum_{j=1}^{n} ij H_{z_i z_j}(\vec{z}) z_{i-1} z_{j-1} = h_{z_1 z_1} - \sum_{k=2}^{n} k(k-1) h_{z_k} \sum_{i=0}^{k-2} {k-2 \choose i} (-1)^{k-2-i} z_1^{k-2-i} z_i,
$$

where $h_{z_1z_1} = h_{z_1z_1}(z_1, z_2 - z_1^2, \ldots, \sum_{j=0}^n {n \choose j} (-1)^j z_1^j z_{n-j})$ and so on for brevity. Applying this differentiation result to (31) , with the first identity in (36) , we have

$$
\left\langle \vec{m}_x^{\tilde{u}}(t,x) \phi_{\vec{z}\vec{z}}\big(t, \vec{m}^{\tilde{u}}(t,x)\big), \vec{m}_x^{\tilde{u}}(t,x) \right\rangle
$$

$$
= e^{2\int_t^T A_v dv} \sum_{i=1}^n \sum_{j=1}^n ij \phi_{z_i z_j}(t, \vec{m}^{\tilde{u}}(t, x)) m_{i-1}^{\tilde{u}}(t, x) m_{j-1}^{\tilde{u}}(t, x)
$$

\n
$$
= -e^{2\int_t^T A_v dv} \sum_{k=2}^n k(k-1) M_{k-2}^{\tilde{u}}(t, x) \psi_{z_k}(t, \vec{M}_{-1}^{\tilde{u}}(t, x))
$$

\n
$$
= -e^{2\int_t^T A_v dv} \sum_{1 \le j \le \frac{n}{2}} 2j(2j-1) \alpha_{2j-2}(y_t^{\beta}) \psi_{z_{2j}}(t, \vec{\alpha}(y_t^{\beta})).
$$

Then, plugging the above result with $\phi_t(t, \vec{m}^{\tilde{u}}(t,x)) = \psi_t(t, \vec{M}^{\tilde{u}}_{-1}(t,x)) = \psi_t(t, \vec{\alpha}(y_t^{\beta})), \frac{1}{2}\kappa B_t \beta_t =$ $\partial_t \psi(t, \vec{\alpha}(y_t^{\beta})) - \psi_t(t, \vec{\alpha}(y_t^{\beta}))$ arising from [\(33\)](#page-16-7) and the EVF candidate [\(39\)](#page-18-1) into the Bellman equation [\(38\)](#page-18-0), yields

$$
0 = \max_{\zeta \in \mathbb{R}} \left\{ e^{\int_t^T A_v dv} \left(\zeta + \frac{F_t}{D_t} \right) \kappa B_t - \frac{1}{2} \kappa B_t \beta_t + \frac{1}{2} e^{2 \int_t^T A_v dv} \left(\zeta + \frac{F_t}{D_t} \right)^2 |D_t|^2 \sum_{1 \le j \le \frac{n}{2}} 2j(2j - 1) \alpha_{2j - 2}(y_t^{\beta}) \psi_{z_{2j}}(t, \vec{\alpha}(y_t^{\beta})) \right\}.
$$
 (40)

Notably, the rest of this proof is to show the validness of [\(40\)](#page-19-0) and that $\tilde{u}(t, x)$ given by [\(35\)](#page-16-2) realizes the maximum on the right-hand side of (40) . Since (34) holds, the maximization condition for (40) can be re-expressed as

$$
0 = \frac{\kappa B_t}{|D_t|^2} + \left(\zeta^* + \frac{F_t}{D_t}\right) e^{\int_t^T A_v dv} \sum_{1 \le j \le \frac{n}{2}} 2j(2j-1)\alpha_{2j-2}(y_t^{\beta}) \psi_{z_{2j}}(t, \vec{\alpha}(y_t^{\beta})),
$$

where ζ^* represents the maximizer. In comparison with [\(33\)](#page-16-7), $\zeta^* = \tilde{u}(t, x)$ follows. Then, substituting the maximizer $\zeta^* = \tilde{u}(t, x)$ back into [\(40\)](#page-19-0), by using [\(33\)](#page-16-7) again, one can find that the right-hand side of [\(40\)](#page-19-0) vanishes. So we complete this proof.

Example 5.6. For the following objective functions arising from mean-variance-skewness problems with arbitrarily fixed $\kappa_2 > 0$ and κ_3 (which degenerates to a mean-variance objective function when $\kappa_3 = 0$),

$$
J_3(t, x, u) = \mathbb{E}_t[X_T^{t, x, u}] + \sum_{j=2}^3 (-1)^{j+1} \frac{\kappa_j}{j!} \mathbb{E}_t \big[(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])^j \big],
$$

Theorem [5.2](#page-16-0) provides a (κ_3, x) -independent CNEC, as the following:

$$
\tilde{u}_{MV}(t,x) = \frac{B_t}{\kappa_2 |D_t|^2} e^{-\int_t^T A_v dv} - \frac{F_t}{D_t}
$$

,

which due to Remark [5.3](#page-17-0) is also a CNEC for

$$
J_{MVS}(t, x, u) = \mathbb{E}_t[X_T^{t, x, u}] - \frac{\kappa_2}{2} \mathbb{E}_t \left[(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])^2 \right] + \frac{\kappa_3}{6} \frac{\mathbb{E}_t \left[(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])^3 \right]}{\left(\mathbb{E}_t \left[(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])^2 \right] \right)^{\frac{3}{2}}}.
$$

For the objective function related to mean-variance-skewness-kurtosis problems,

$$
J_4(t, x, u) = \mathbb{E}_t[X_T^{t, x, u}] + \sum_{j=2}^4 (-1)^{j+1} \frac{\kappa_j}{j!} \mathbb{E}_t \left[(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])^j \right],
$$

where $\kappa_2, \kappa_4 \geq 0$ with $|\kappa_2| + |\kappa_4| > 0$, the CNEC given by Theorem [5.2](#page-16-0) is

$$
\tilde{u}_4(t,x) = \left(\kappa_2^3 + \frac{3}{2}\kappa_4\theta_t\right)^{-\frac{1}{3}} \frac{B_t}{|D_t|^2} e^{-\int_t^T A_v dv} - \frac{F_t}{D_t},
$$

which is also (κ_3, x) -independent. In fact, according to Theorem [5.2](#page-16-0) and the definition of (θ_t, y_t^{β}) , we are supposed to solve $\theta'_t = y'_t(\kappa_2 + \frac{1}{2}\kappa_4y_t)^2$ for y_t . Integrating the both sides over [t, T] with rearranging the terms yields $\kappa_2 + \frac{1}{2}\kappa_4 y_t = (\kappa_2^3 + \frac{3}{2}\kappa_4 \theta_t)^{1/3}$, which leads to the desired result by differentiating the both sides w.r.t. t. More generally, for the objective function

$$
J_n(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] + \sum_{j=2}^n (-1)^{j+1} \frac{\kappa_j}{j!} \mathbb{E}_t\left[(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])^j \right],\tag{41}
$$

as a linear combination of mean and central moments, where all $\kappa_{2j} \geq 0$ and at least one $\kappa_{2j} > 0$, Theorem [5.2](#page-16-0) suggests that seeking a CNEC can be reduced to solving

$$
0 = \left| \frac{\kappa B_t}{D_t} \right|^2 + y'_t \left| \sum_{1 \le j \le \frac{n}{2}} \kappa_{2j} \frac{(y_t)^{j-1}}{(2j-2)!!} \right|^2
$$

as well as the integral equation $y_t = y_t^{\beta}$. Integrating the both sides of the above differential equation over $[t, T]$ yields the algebraic equation

$$
\kappa^2 \theta_t = \int_0^{y_t} \bigg| \sum_{0 \le j \le \frac{n}{2} - 1} \kappa_{2j+2} \frac{z^j}{(2j)!!} \bigg|^2 dz \equiv 2 \int_0^{\frac{1}{2} y_t} \bigg| \sum_{0 \le j \le \frac{n}{2} - 1} \kappa_{2j+2} \frac{z^j}{j!} \bigg|^2 dz. \tag{42}
$$

If $\kappa = 0$, then [\(42\)](#page-20-0) has the unique solution $y \equiv 0$, which leads to $\beta \equiv 0$. Otherwise, the unique solution y_t of [\(42\)](#page-20-0) is decreasing and continuously differentiable. Then, [\(35\)](#page-16-2) with $\beta_t = (D_t)^{-1} \sqrt{-y_t}$ gives the CNEC.

Example 5.7. Interested readers may focus on the mean-variance-standardized moments objective function

$$
J_{MVSM}(t,x,u) = \mathbb{E}_t[X_T^{t,x,u}] - \frac{\kappa_2}{2} \mathbb{E}_t \left[(X_T^{t,x,u} - \mathbb{E}_t[X_T^{t,x,u}])^2 \right] + \sum_{j=3}^n (-1)^{j+1} \frac{\kappa_j}{j!} \frac{\mathbb{E}_t \left[(X_T^{t,x,u} - \mathbb{E}_t[X_T^{t,x,u}])^j \right]}{\left(\mathbb{E}_t \left[(X_T^{t,x,u} - \mathbb{E}_t[X_T^{t,x,u}])^2 \right] \right)^{\frac{j}{2}}};
$$

that is, all higher-order central moments except the variance in (41) are replaced by standardized moments of the corresponding order. For this situation, we let

$$
\psi(t, z_2, \dots, z_n) = -\frac{\kappa_2}{2} z_2 + \sum_{j=3}^n (-1)^{j+1} \frac{\kappa_j}{j!} \frac{z_j}{|z_2|^{\frac{j}{2}}}.
$$

Consequently, the left-hand side of [\(33\)](#page-16-7) equals to $B_t|D_t|^{-2} - \kappa_2\beta_t$. Therefore, \tilde{u}_{MV} in the last Example [5.6](#page-19-1) is also a CNEC for J_{MVSM} , if $\kappa_2 > 0$.

5.2 Open-loop Nash equilibrium control

Now we show that the state-independent CNEC given by [\(35\)](#page-16-2) is exactly a path-independent ONEC. According to Theorem [4.5,](#page-13-1) seeking an ONEC can be reduced to solving the linear FBSDE

$$
\begin{cases}\nd\bar{X}_s = (A_s\bar{X}_s + B_s\bar{u}_s + C_s)ds + (D_s\bar{u}_s + F_s)dW_s, \ \forall s \in [0, T], \ \bar{X}_0 = x_0; \\
dY_s^t = -A_sY_s^tds + \mathcal{Y}_s^tdW_s, \ \forall s \in [t, T], \\
Y_T^t = \kappa + \sum_{j=2}^n j\psi_{z_j}(t, \vec{M}_{-1}^{\bar{u}}(t, \bar{X}_t))\big((\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^{j-1} - M_{j-1}^{\bar{u}}(t, \bar{X}_t)\big); \\
dZ_s^t = -2A_sZ_s^tds + \mathcal{Z}_s^tdW_s, \ \forall s \in [t, T], \\
Z_T^t = \sum_{j=2}^n j(j-1)\psi_{z_j}(t, \vec{M}_{-1}^{\bar{u}}(t, \bar{X}_t))(\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^{j-2},\n\end{cases} \tag{43}
$$

with the following optimality condition:

$$
\lim_{s \downarrow t} \mathbb{E}_t[B_s Y_s^t + D_s \mathcal{Y}_s^t] = 0, \quad \lim_{s \downarrow t} \mathbb{E}_t[Z_s^t] \le 0, \quad \mathbb{P}-a.s., \ a.e. \ t \in [0, T). \tag{44}
$$

Lemma 5.8. Suppose that $\bar{u}_t = \beta_t \exp(-\int_t^T A_v dv) - F_t/D_t$, where β is an arbitrarily fixed deterministic continuous function. Then, [\(43\)](#page-21-0) admits the unique solution

$$
\begin{cases}\n\bar{X}_s = x_0 e^{\int_0^s A_v dv} + \int_0^s e^{\int_r^s A_v dv} \Big(C_\tau - \frac{B_\tau}{D_\tau} F_\tau \Big) d\tau \\
+ e^{-\int_s^T A_v dv} \Big(\int_0^s B_\tau \beta_\tau d\tau + \int_0^s D_\tau \beta_\tau dW_\tau \Big), \\
Y_s^t = e^{\int_s^T A_v dv} \Big(\kappa + \sum_{j=2}^n j \psi_{z_j} (t, \vec{\alpha}(y_t^\beta)) \Big(\mathbb{E}_s \Big[\Big(\int_t^T D_\tau \beta_\tau dW_\tau \Big)^{j-1} \Big] - \alpha_{j-1}(y_t^\beta) \Big) \Big), \\
\mathcal{Y}_s^t = e^{\int_s^T A_v dv} D_s \beta_s \sum_{j=2}^n j(j-1) \psi_{z_j} (t, \vec{\alpha}(y_t^\beta)) \mathbb{E}_s \Big[\Big(\int_t^T D_\tau \beta_\tau dW_\tau \Big)^{j-2} \Big], \\
Z_s^t = e^{2 \int_s^T A_v dv} \sum_{j=2}^n j(j-1) \psi_{z_j} (t, \vec{\alpha}(y_t^\beta)) \mathbb{E}_s \Big[\Big(\int_t^T D_\tau \beta_\tau dW_\tau \Big)^{j-2} \Big], \\
Z_s^t = e^{2 \int_s^T A_v dv} D_s \beta_s \sum_{j=3}^n j(j-1) (j-2) \psi_{z_j} (t, \vec{\alpha}(y_t^\beta)) \mathbb{E}_s \Big[\Big(\int_t^T D_\tau \beta_\tau dW_\tau \Big)^{j-3} \Big].\n\end{cases}
$$

Proof. The explicit expression of \bar{X} arises from the SDE

$$
d\bar{X}_s = \left(A_s \bar{X}_s + B_s \beta_s e^{-\int_s^T A_v dv} + C_s - \frac{B_s}{D_s} F_s\right) ds + D_s \beta_s e^{-\int_s^T A_v dv} dW_s.
$$

Then, we have $\bar{X}_T = \mathbb{E}_t[\bar{X}_T] + \int_t^T D_s \beta_s dW_s$, and hence $\vec{M}_{-1}^{\bar{u}}(t, \bar{X}_t) = \vec{\alpha}(y_t^{\beta})$. See also the proof for Theorem [5.2.](#page-16-0) Consequently, by the Lipschitz continuity of $\psi(t, \cdot)$, we conclude that Y_T^t and Z_T^t are squareintegrable. Due to [\(Yong & Zhou,](#page-30-0) [1999,](#page-30-0) Theorem 7.2.2, p. 349), each backward stochastic differential equation (BSDE) in [\(43\)](#page-21-0) admits a unique square-integrable F-predictable solution. Furthermore, from the BSDEs in [\(43\)](#page-21-0) one can obtain

$$
Y^t_s = e^{\int_s^T A_v dv} \mathbb{E}_s[Y^t_T], \qquad Z^t_s = e^{2\int_s^T A_v dv} \mathbb{E}_s[Z^t_T],
$$

which lead to the desired explicit expression of (Y^t, Z^t) . In order to derive the explicit expression of \mathcal{Y}^t

from $Y_s^t \exp(-\int_s^T A_v dv) = \mathbb{E}_s[Y_T^t]$, we differentiate the both sides of

$$
\sum_{j=0}^{\infty} \frac{z^j}{j!} \mathbb{E}_s \left[(\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^j \right] = \mathbb{E}_s \left[e^{z \int_t^T D_v \beta_v dW_v} \right] = e^{z \int_t^s D_v \beta_v dW_v + \frac{1}{2} z^2 y_s^{\beta}}.
$$

with respect to s to show that $d\mathbb{E}_s[(\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^j] = j\mathbb{E}_s[(\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^{j-1}]D_s\beta_s dW_s$. Consequently,

$$
d\left(Y_s^t e^{-\int_s^T A_v dv}\right) = \sum_{j=2}^n j(j-1)\psi_{z_j}\left(t, \vec{\alpha}(y_t^{\beta})\right) \mathbb{E}_s\left[\left(\bar{X}_T - \mathbb{E}_t[\bar{X}_T]\right)^{j-2}\right] D_s \beta_s dW_s.
$$

By comparing $d(Y_s^t \exp(-\int_s^T A_v dv)) = \exp(-\int_s^T A_v dv) \mathcal{Y}_s^t dW_s$ from [\(43\)](#page-21-0) with the abovementioned result, we obtain the desired expression of \mathcal{Y}^t according to the existence and uniqueness of \mathcal{Y}^t . In the same manner, comparing

$$
d\Big(Z_s^t e^{-2\int_s^T A_v dv}\Big) = \sum_{j=3}^n j(j-1)(j-2)\psi_{z_j}(t, \vec{\alpha}(y_t^{\beta})) \mathbb{E}_s[(\bar{X}_T - \mathbb{E}_t[\bar{X}_T])^{j-3}]D_s\beta_s dW_s
$$

with $d(Z_s^t \exp(-2\int_s^T A_v dv)) = \exp(-2\int_s^T A_v dv) Z_s^t dW_s$ from [\(43\)](#page-21-0) yields the desired expression of \mathcal{Z}^t . **Theorem 5.9.** Let β fulfill [\(33\)](#page-16-7) and [\(34\)](#page-16-6). Then, for the objective function [\(29\)](#page-15-1) and the controlled SDE [\(30\)](#page-15-2), $\{\beta_t \exp(-\int_t^T A_v dv) - F_t/D_t\}_{t \in [0,T]}$ is an ONEC.

Proof. From Lemma [5.8,](#page-21-1) we obtain $\lim_{s\downarrow t} \mathbb{E}_t[B_s Y_s^t + D_s \mathcal{Y}_s^t] = B_t Y_t^t + D_t \mathcal{Y}_t^t$ and $\lim_{s\downarrow t} \mathbb{E}_t[Z_s^t] = Z_t^t$, where

$$
\begin{cases} Y_t^t = \kappa e^{\int_t^T A_v dv}, \\ \mathcal{Y}_t^t = e^{\int_t^T A_v dv} D_t \beta_t \sum_{1 \le j \le \frac{n}{2}} 2j(2j-1) \alpha_{2j-2}(y_t^{\beta}) \psi_{z_{2j}}(t, \vec{\alpha}(y_t^{\beta})), \\ Z_t^t = e^{2 \int_t^T A_v dv} \sum_{1 \le j \le \frac{n}{2}} 2j(2j-1) \alpha_{2j-2}(y_t^{\beta}) \psi_{z_{2j}}(t, \vec{\alpha}(y_t^{\beta})). \end{cases}
$$

 $D_t \mathcal{Y}_t^t = -B_t Y_t^t$ follows from [\(33\)](#page-16-7), while $Z_t^t < 0$ follows from [\(34\)](#page-16-6). Thus, for β given by (33) and (34), the optimality condition [\(44\)](#page-21-2) holds. According to Theorem [4.5,](#page-13-1) \bar{u} is an ONEC.

5.3 Limiting cases

In the previous subsections, we have shown that the univariate time function given by (33) – (35) is a CNEC and an ONEC for the control problem with the objective function [\(29\)](#page-15-1) and the controlled SDE [\(30\)](#page-15-2). In particular, for the objective function [\(41\)](#page-20-1) as a linear combination of mean and central moments that researchers may be more concerned about, deriving the state-independent CNEC or the path-independent ONEC is reduced to solving the algebraic equation [\(42\)](#page-20-0). However, the celebrated Abel-Ruffini Theorem and Galois Theory indicate that there is no solution in radicals to [\(42\)](#page-20-0) of finite degree $n \geq 6$ with arbitrary coefficients κ_{2j} . In this subsection, we let n tend to infinity. As a result, we provide a heuristic approach to find state-independent CNEC, as well as path-independent ONEC, for several artificial objective functions. Notably, the theories in Sections [3](#page-7-0) and [4](#page-9-0) are only suitable for the problem with finitely many higher-order moments, so we also provide verification procedures by applying those theories to $\Psi(t, x, y, \vec{z}) \equiv \Phi(y, z_1)$ (i.e. without any higher-order moment) for the sake of rigor.

5.3.1 Limiting case I: mean-exponential utility function

Let us consider the objective function

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] + \sum_{j=2}^{\infty} \frac{(-c)^{j-1}}{j!} \mathbb{E}_t[(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])^j]
$$
(45)

for fixed $\kappa > 0$ and $c > 0$, or equivalently,

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] - \frac{1}{c} \mathbb{E}_t \left[e^{-c(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])} \right] + \frac{1}{c}.
$$
 (46)

Notably, [\(46\)](#page-23-1) can be regarded as a weighted average of the expectation $\mathbb{E}_t[X_T^{t,x,u}]$ and the expected exponential utility for $X_T^{t,x,u} - \mathbb{E}_t[X_T^{t,x,u}]$, for which c represents the CARA coefficient. By Theorems [5.2](#page-16-0) and [5.9,](#page-22-1) to determine a CNEC and an ONEC for [\(45\)](#page-23-2), it suffices to solve $\kappa^2 \theta_t = \exp(c^2 y_t) - 1$ arising from [\(41\)](#page-20-1) and [\(42\)](#page-20-0). As a result, $y_t = c^{-2} \ln(1 + \kappa^2 \theta_t)$, and hence, $\beta_t = \kappa c^{-1} B_t |D_t|^{-2} (1 + \kappa^2 \theta_t)^{-1/2}$ solves $y_t^{\beta} = y_t$. Moreover, [\(34\)](#page-16-6) for $n = \infty$ holds, since

$$
\sum_{j=1}^{\infty} j(2j-1)\alpha_{2j-2}(y_t^{\beta})\psi_{z_{2j}}(t,\vec{\alpha}(y_t^{\beta})) = -\frac{c}{2} \sum_{j=1}^{\infty} \frac{(c^2 y_t^{\beta})^{j-1}}{(2j-2)!!} = -\frac{c}{2} e^{\frac{1}{2}c^2 y_t^{\beta}} < 0.
$$

Hence, for the CNEC \tilde{u} , the ONEC \bar{u} and the EVF, we have

$$
\tilde{u}(t,x) = \bar{u}_t = \frac{\kappa B_t e^{-\int_t^T A_v dv}}{c|D_t|^2 \sqrt{1 + \kappa^2 \theta_t}} - \frac{F_t}{D_t},\tag{47}
$$

$$
V(t,x) = J(t,x,\tilde{u}) = \kappa \Theta(t,x) + \frac{\sqrt{1 + \kappa^2 \theta_t} - 1}{c}.
$$
\n(48)

The rest of this subsection is to show that \tilde{u} (resp. \bar{u}) is exactly a CNEC (resp. an ONEC) for [\(46\)](#page-23-1). We may refer to the total of the following steps as a "verification procedure". On the one hand, we let $\Phi(t, x, y, \vec{z}) \equiv \Phi(y, z_1) = \kappa y - c^{-1} e^{-c(y - z_1)} + c^{-1}$ for [\(46\)](#page-23-1). According to [\(19\)](#page-9-2), the Bellman equation for deriving a CNEC is

$$
0 = \max_{\zeta \in \mathbb{R}} \left\{ V_t(t, x) + V_x(t, x)(A_t x + B_t \zeta + C_t) + \frac{1}{2} V_{xx}(t, x)(D_t \zeta + F_t)^2 - \left(\lambda_x^{t, x}(t, x) \partial_x m_1^{\tilde{u}}(t, x) + \frac{1}{2} \mu^{t, x}(t, x) \left(\partial_x m_1^{\tilde{u}}(t, x) \right)^2 \right) (D_t \zeta + F_t)^2 \right\},
$$
\n(49)

where

$$
\begin{cases}\n\lambda^{s,y}(t,x) = \mathbb{E}_t \big[\Phi_{z_1} \big(X_T^{t,x,\tilde{u}}, m_1^{\tilde{u}}(s,y) \big) \big] = -\mathbb{E}_t \big[e^{-c(X_T^{t,x,\tilde{u}} - m_1^{\tilde{u}}(s,y))} \big], \\
\mu^{s,y}(t,x) = \mathbb{E}_t \big[\Phi_{z_1 z_1} \big(X_T^{t,x,\tilde{u}}, m_1^{\tilde{u}}(s,y) \big) \big] = c \lambda^{s,y}(t,x).\n\end{cases} \tag{50}
$$

Now we show that V given by [\(48\)](#page-23-3) fulfills [\(49\)](#page-23-4), whereby \tilde{u} given by [\(47\)](#page-23-5) realizes the maximum. Corresponding to \tilde{u} given by [\(47\)](#page-23-5), we have

$$
\lambda_x^{s,y}(t,x) = ce^{\int_t^T A_v dv} \mathbb{E}_t \left[e^{-c(X_T^{t,x,\tilde{u}} - m_1^{\tilde{u}}(s,y))} \right] = -ce^{\int_t^T A_v dv} \lambda^{s,y}(t,x)
$$

and $\lambda^{t,x}(t,x) = -\mathbb{E}_t[\exp(-c\int_t^T D_s\beta_s dW_s)] = -\sqrt{1+\kappa^2\theta_t}$. It follows from [\(48\)](#page-23-3) with the abovementioned

results that the right-hand side of [\(49\)](#page-23-4) is equal to

$$
\max_{\zeta \in \mathbb{R}} \left\{ -\frac{\kappa^2 |B_t|^2}{2c|D_t|^2 \sqrt{1 + \kappa^2 \theta_t}} + \kappa e^{\int_t^T A_v dv} \frac{B_t}{D_t} (D_t \zeta + F_t) -\frac{c}{2} \sqrt{1 + \kappa^2 \theta_t} e^{2 \int_t^T A_v dv} (D_t \zeta + F_t)^2 \right\} = 0,
$$

for which $\zeta^* = \tilde{u}(t, x)$ given by [\(47\)](#page-23-5) satisfies the first-order derivative condition:

$$
0 = \kappa e^{\int_t^T A_v dv} B_t - c\sqrt{1 + \kappa^2 \theta_t} e^{2 \int_t^T A_v dv} D_t (D_t \zeta^* + F_t).
$$

So we are done. On the other hand, to show that \bar{u} given by [\(47\)](#page-23-5) is an ONEC for [\(46\)](#page-23-1), it suffices to show the validity of [\(44\)](#page-21-2) with $(Y^t, \mathcal{Y}^t, Z^t)$ given by the BSDEs

$$
\begin{cases}\ndY_s^t = -A_s Y_s^t ds + Y_s^t dW_s, & \forall s \in [t, T], \\
Y_T^t = \Psi_y(\bar{X}_T, \mathbb{E}_t[\bar{X}_T]) + \mathbb{E}_t[\Psi_{z_1}(\bar{X}_T, \mathbb{E}_t[\bar{X}_T])] = \kappa + e^{-c \int_t^T D_v \beta_v dW_v} - e^{\frac{1}{2}c^2 y_t^{\beta}}; \\
dZ_s^t = -2A_s Z_s^t ds + Z_s^t dW_s, & \forall s \in [t, T], \\
Z_T^t = \Psi_{yy}(\bar{X}_T, \mathbb{E}_t[\bar{X}_T]) = -ce^{-c \int_t^T D_v \beta_v dW_v},\n\end{cases}
$$

according to Theorem [4.5](#page-13-1) with $\bar{X}_T - \mathbb{E}_t[\bar{X}_T] = \int_t^T D_s \beta_s dW_s$. From

$$
\begin{cases} Y_s^t e^{-\int_s^T A_v dv} = \kappa + \mathbb{E}_s \big[e^{-c \int_t^T D_v \beta_v dW_v} \big] - e^{\frac{1}{2}c^2 y_t^{\beta}}, \\ d\big(Y_s^t e^{-\int_s^T A_v dv}\big) = e^{-\int_s^T A_v dv} \mathcal{Y}_s^t dW_s, \end{cases}
$$

we obtain $\mathcal{Y}_{s}^{t} = -c \exp(\int_{s}^{T} A_{v} dv + \frac{1}{2} c_{v}^{2} y_{t}^{\beta} - c \int_{t}^{s} D_{v} \beta_{v} dW_{v}) D_{s} \beta_{s}$. In the same manner, we can show that $Z_s^t = -c \exp(2 \int_s^T A_v dv) \mathbb{E}_s[\exp(-c \int_t^T D_v \beta_v dW_v)].$ Consequently,

$$
\lim_{s\downarrow t}\mathbb{E}_t[B_sY_s^t+D_s\mathcal{Y}_s^t]=B_tY_t^t+D_t\mathcal{Y}_t^t=0,\quad \lim_{s\downarrow t}\mathbb{E}_t[Z_s^t]=Z_t^t<0.
$$

Therefore, [\(44\)](#page-21-2) holds, and hence \bar{u} given by [\(47\)](#page-23-5) is an ONEC for [\(46\)](#page-23-1).

5.3.2 Limiting case II: hyperbolic cosine penalty function

For arbitrarily fixed $\kappa \geq 0$ and $c > 0$, we consider the objective function

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] - \frac{1}{c} \mathbb{E}_t \left[\cosh \left(c(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}]) \right) \right] + \frac{1}{c},\tag{51}
$$

where the hyperbolic cosine function is the penalty function for the deviation $X_T^{t,x,u} - \mathbb{E}_t[X_T^{t,x,u}]$. Roughly speaking, [\(51\)](#page-24-0) is an analogue to the conventional mean-variance objective function, but it replaces the quadratic function for variance with a hyperbolic cosine function. Notably, [\(51\)](#page-24-0) can be re-expressed as

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] - \sum_{j=1}^{\infty} \frac{c^{2j-1}}{(2j)!} \mathbb{E}_t\left[(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])^{2j} \right],\tag{52}
$$

which is the result of removing all odd-order central moments from [\(45\)](#page-23-2). This implies that \tilde{u} (resp. \bar{u}) given by (47) is a CNEC (resp. an ONEC) for (52) , and V given by (48) is the EVF. Interested readers can follow our verification procedure as in the previous Section [5.3.1](#page-23-0) to see that \tilde{u} (resp. \bar{u}) is exactly a

CNEC (resp. an ONEC) for [\(51\)](#page-24-0). Here we omit the statement due to the page limit.

5.3.3 Limiting case III: cosine penalty function

In this subsection, we employ an irrational penalty function $S(x) = (1 - \cos(cx))/c$ with $c > 0$; that is,

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] + \frac{1}{c} \mathbb{E}_t \left[\cos \left(c(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}]) \right) \right] - \frac{1}{c}.
$$
 (53)

Some results in the following are parallel to those in Section [5.3.2,](#page-24-2) because of the link between $cos(\cdot)$ and cosh(\cdot). In addition, for technic reasons we assume that $\kappa^2 \theta_0 < 1$, so that the marginal reward from $\mathbb{E}_t[X_T^{t,x,u}]$ and the penalty for the deviation $X_T^{t,x,u} - \mathbb{E}_t[X_T^{t,x,u}]$ are both bounded. Let us rewrite [\(53\)](#page-25-1) as

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] - \sum_{j=1}^{\infty} \frac{(-1)^{j-1} c^{2j-1}}{(2j)!} \mathbb{E}_t \left[(X_T^{t, x, u} - \mathbb{E}_t [X_T^{t, x, u}])^{2j} \right].
$$
 (54)

Following the same line as deriving [\(47\)](#page-23-5) and [\(48\)](#page-23-3), we obtain $y_t^{\beta} = -c^{-2} \ln(1 - \kappa^2 \theta_t)$, which solves $\kappa^2 \theta_t = 1 - \exp(-c^2 y_t)$. Then, $\beta_t = \kappa c^{-1} B_t |D_t|^{-2} (1 - \kappa^2 \theta_t)^{-1/2}$ and

$$
\sum_{j=1}^{\infty} j(2j-1)\alpha_{2j-2}(y_t^{\beta})\psi_{z_{2j}}(t,\vec{\alpha}(y_t^{\beta})) = -\frac{c}{2} \sum_{j=1}^{\infty} \frac{(-c^2 y_t^{\beta})^{j-1}}{(2j-2)!!} = -\frac{c}{2} e^{-\frac{1}{2}c^2 y_t^{\beta}} < 0.
$$

Therefore, for the CNEC \tilde{u} , the ONEC \bar{u} and the EVF V, we have

$$
\tilde{u}(t,x) = \bar{u}_t = \frac{\kappa B_t e^{-\int_t^T A_v dv}}{c|D_t|^2 \sqrt{1 - \kappa^2 \theta_t}} - \frac{F_t}{D_t}, \quad V(t,x) = \kappa \Theta(t,x) + \frac{1 - \sqrt{1 - \kappa^2 \theta_t}}{c}.
$$

Here we omit the statement for the verification procedure due to the page limit, and encourage interested readers to try it on their own. In fact, the verification procedure for a more general case will be provided in the coming Section [5.3.4.](#page-25-2)

5.3.4 Limiting case IV: ambiguous cosine penalty function

Let H be an $\mathbb F$ -independent random variable whose all original moments of positive integer order exist, and $S(x) = 1 - \mathbb{E}[\cos(Hx)]$ be the penalty function for

$$
J(t, x, u) = \kappa \mathbb{E}_t[X_T^{t, x, u}] + \mathbb{E}_t \left[\cos \left(H(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}]) \right) \right] - 1
$$

$$
= \kappa \mathbb{E}_t[X_T^{t, x, u}] - \sum_{j=1}^{\infty} \frac{(-1)^{j-1}}{(2j)!} \mathbb{E}[H^{2j}] \mathbb{E}_t \left[(X_T^{t, x, u} - \mathbb{E}_t[X_T^{t, x, u}])^{2j} \right].
$$

(55)

In fact, when the agent cannot specify an exact penalty function $S_h(x) = 1 - \cos(hx)$ because she/he cannot have complete information about the principal or the environment, it might be an appropriate choice to try the expected penalty function $S(x) = \mathbb{E}[S_H(x)] = 1 - \mathbb{E}[\cos(Hx)]$ with a given distribution of H. Since

$$
\sum_{j=1}^{\infty} j(2j-1)\alpha_{2j-2}(y_t^{\beta})\psi_{z_{2j}}(t,\vec{\alpha}(y_t^{\beta})) = -\frac{1}{2}\sum_{j=1}^{\infty} \frac{(-y_t^{\beta})^{j-1}}{(2j-2)!!} \mathbb{E}[H^{2j}] = -\frac{1}{2} \mathbb{E}[H^2 e^{-\frac{1}{2}H^2 y_t^{\beta}}],
$$

the condition [\(34\)](#page-16-6) for $n = \infty$ holds. Then, for a CNEC/ONEC candidate in the form of [\(35\)](#page-16-2), due to [\(41\)](#page-20-1) and [\(42\)](#page-20-0), it is supposed to solve $\kappa^2 \theta_t = \int_0^{y_t} \left(\mathbb{E} \left[H^2 e^{-\frac{1}{2} H^2 z} \right] \right)^2 dz$ for the unique solution y_t^{β} . As a result, $\beta_t = \kappa B_t |D_t|^{-2} / \mathbb{E}[H^2 \exp(-H^2 y_t^{\beta}/2)].$

Now we are going to show that for [\(55\)](#page-25-3),

$$
\tilde{u}(t,x) = \bar{u}_t = e^{-\int_t^T A_v dv} \frac{\kappa B_t}{|D_t|^2} \left(\mathbb{E} \left[H^2 e^{-\frac{1}{2} H^2 y_t^{\beta}} \right] \right)^{-1} - \frac{F_t}{D_t}
$$
\n(56)

exactly gives a CNEC and an ONEC, and the corresponding EVF is

$$
V(t,x) = \kappa \Theta(t,x) + 1 - \mathbb{E} \left[e^{-\frac{1}{2}H^2 y_t^{\beta}} \right].
$$
\n(57)

On the one hand, let $\Phi(y, z_1) = \kappa y - S(y - z_1)$ and

$$
\begin{cases}\n\lambda^{s,y}(t,x) = \mathbb{E}_t \big[\Phi_{z_1} (X_T^{t,x,\tilde{u}}, m_1^{\tilde{u}}(s,y)) \big] = \mathbb{E}_t \big[S' (X_T^{t,x,\tilde{u}} - m_1^{\tilde{u}}(s,y)) \big], \\
\mu^{s,y}(t,x) = \mathbb{E}_t \big[\Phi_{z_1 z_1} (X_T^{t,x,\tilde{u}}, m_1^{\tilde{u}}(s,y)) \big] = -\mathbb{E}_t \big[S'' (X_T^{t,x,\tilde{u}} - m_1^{\tilde{u}}(s,y)) \big],\n\end{cases}
$$

which leads to $\lambda_x^{s,y}(t,x) = -\exp(\int_t^T A_v dv) \mu^{s,y}(t,x)$ and

$$
\mu^{t,x}(t,x) = -\mathbb{E}_t \left[H^2 \cos \left(H \int_t^T D_v \beta_v dW_v \right) \right] = -\mathbb{E} \left[H^2 e^{-\frac{1}{2} H^2 y_t^{\beta}} \right].
$$

Then, for (57) , the right-hand side of (49) is equal to

$$
\max_{\zeta \in \mathbb{R}} \left\{ -\frac{\kappa^2 |B_t|^2}{2|D_t|^2 \mathbb{E} \left[H^2 e^{-\frac{1}{2} H^2 y_t^{\beta}} \right]} + \kappa e^{\int_t^T A_v dv} \frac{B_t}{D_t} (D_t \zeta + F_t) -\frac{1}{2} \mathbb{E} \left[H^2 e^{-\frac{1}{2} H^2 y_t^{\beta}} \right] e^{2 \int_t^T A_v dv} (D_t \zeta + F_t)^2 \right\} = 0,
$$

whereby [\(56\)](#page-26-1) realizes the maximum. Hence, we obtain the desired conclusion for (\tilde{u}, V) . On the other hand, since the following BSDE arising from [\(43\)](#page-21-0),

$$
\begin{cases} dY_s^t = -A_s Y_s^t ds + \mathcal{Y}_s^t dW_s, \quad \forall s \in [t, T], \\ Y_T^t = \kappa - \mathbb{E}_T \left[H \sin \left(H \int_t^T D_v \beta_v dW_v \right) \right], \end{cases}
$$

admits the solution

$$
\begin{cases}\nY_s^t e^{-\int_s^T A_v dv} = \kappa - \mathbb{E}_s \left[H \sin \left(H \int_t^s D_v \beta_v dW_v \right) e^{-\frac{1}{2} H^2 y_s^{\beta}} \right], \\
\mathcal{Y}_s^t e^{-\int_s^T A_v dv} = - \mathbb{E}_s \left[H^2 \cos \left(H \int_t^s D_v \beta_v dW_v \right) e^{-\frac{1}{2} h^2 y_s^{\beta}} \right] D_s \beta_s,\n\end{cases}
$$

we obtain $\lim_{s\downarrow t} \mathbb{E}_t[B_s Y_s^t + D_s Y_s^t] = B_t Y_t^t + D_t Y_t^t = 0$. Moreover, given that $dZ_s^t = -2A_s Z_s^t ds + Z_s^t dW_s$ and $Z_T^t = -\mathbb{E}_T[H^2\cos(H\int_t^T D_v\beta_v dW_v)]$ from [\(43\)](#page-21-0), we obtain $\lim_{s\downarrow t} Z_s^t = Z_t^t = -\mathbb{E}_t[H^2\exp(-\frac{1}{2}H^2y_t^{\beta})]$ 0. Hence, we obtain the desired conclusion for \bar{u} . So we are done.

Remark 5.10. Let i be the imaginary unit for (generalized) Fourier transform. For a fairly general even penalty function $S(x)$, its Fourier (cosine) transform

$$
\hat{S}(h) := \int_{-\infty}^{+\infty} S(x)e^{-ihx}dx = \int_{-\infty}^{+\infty} S(x)\cos(hx)dx
$$

is also an even real-valued function. By inverse Fourier (cosine) transform,

$$
S(x) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \hat{S}(h) \cos(hx) dh = \sum_{j=0}^{\infty} \frac{(-1)^j x^{2j}}{2\pi (2j)!} \int_{-\infty}^{+\infty} \hat{S}(h) h^{2j} dh.
$$

This is a generalization of the form $S(x) = 1 - \mathbb{E}[\cos(Hx)]$. Roughly speaking, one can treat $\hat{S}/2\pi$ as a "probability density function", or replace the probability measure with $\hat{S}(h)dh/2\pi$ in our previous derivation. For example, for $S(x) = 1 - \exp(-\frac{1}{2}x^2)$,

$$
\frac{1}{2\pi}\hat{S}(h) = \delta(h) - \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ihx - \frac{x^2}{2}} dx = \delta(h) - \frac{1}{\sqrt{2\pi}} e^{-\frac{h^2}{2}},
$$

where $\delta(\cdot)$ is Dirac delta function. In terms of determining β for the CNEC/ONEC in the form of [\(35\)](#page-16-2), according to (42) , it is supposed to solve

$$
\kappa^{2} \theta_{t} = \int_{0}^{y_{t}^{\beta}} \bigg(\int_{-\infty}^{+\infty} \frac{\hat{S}(h)}{2\pi} h^{2} e^{-\frac{1}{2}h^{2}z} dh \bigg)^{2} dz.
$$

As a result, if $\int_{-\infty}^{+\infty} \hat{S}(h)h^2 \exp(-h^2 y_t^{\beta}/2) dh > 0$, interested readers can arrive at the CNEC/ONEC and its corresponding EVF, as the following:

$$
\tilde{u}(t,x) = \bar{u}_t = e^{-\int_t^T A_v dv} \frac{\kappa B_t}{|D_t|^2} \left(\int_{-\infty}^{+\infty} \frac{\hat{S}(h)}{2\pi} h^2 e^{-\frac{1}{2}h^2 y_t^{\beta}} dh \right)^{-1} - \frac{F_t}{D_t},
$$

$$
V(t,x) = \kappa \Theta(t,x) + 1 - \int_{-\infty}^{+\infty} \frac{\hat{S}(h)}{2\pi} e^{-\frac{1}{2}h^2 y_t^{\beta}} dh.
$$

Notably, any real function S defined on an interval symmetric about the origin has the decomposition $S(x) = (S(x) + S(-x))/2 + (S(x) - S(-x))/2$, where the first part is even and the second part is odd. According to Remark [5.3,](#page-17-0) the state-independent CNEC and the path-independent ONEC for a fairly qeneral penalty function $S(x)$ might be the same as those for $(S(x) + S(-x))/2$, for which the above Fourier cosine transform method can be used.

6 Concluding remark

We have studied the time-consistent stochastic control problems with higher-order central moments of the terminal value of state process. On the one hand, seeking a closed-loop Nash equilibrium control is reduced to solving a PDE system, where the so-called equilibrium value function as well as its Bellman equation is not necessary. On the other hand, referring to a standard perturbation argument for spike variation, we derive a sufficient maximum principle with a flow of FBSDEs for open-loop Nash equilibrium control. After the necessary theory is established, we consider the linear control problems, where the objective functional is affine in the mean. In many cases, the closed-loop and the open-loop Nash equilibrium controls are identical and independent of the state, random path and the preference on odd-order central moments.

Acknowledgments

The research of the first author is supported by National Natural Science Foundation of China under Grant 12401611 and CTBU Research Projects Grant 2355010. The research of the second author is supported by Major Program of the Key Research Institute on Humanities and Social Science of China Ministry of Education under Grant 22JJD790091. The third author acknowledges the funding support from NSF-DMS Grant 2204795. The fourth author acknowledges the funding support from the Research Grants Council of the Hong Kong Special Administrative Region Grant PolyU15223419.

References

- Alia, I. (2019). A non-exponential discounting time-inconsistent stochastic optimal control problem for jump-diffusion. Math. Control Relat. Fields, 9(3), 541-570. doi: 10.3934/mcrf.2019025
- Basak, S., & Chabakauri, G. (2010). Dynamic mean-variance asset allocation. Rev. Financ. Stud., 23(8), 2970-3016. doi: 10.1093/rfs/hhq028
- Bensoussan, A., Frehse, J., & Yam, P. (2013). Mean field games and mean field type control theory. New York: Springer-Verlag. doi: 10.1007/978-1-4614-8508-7
- Björk, T., Khapko, M., & Murgoci, A. (2017). On time-inconsistent stochastic control in continuous time. Finance Stoch., 21(2), 331-360. doi: 10.1007/s00780-017-0327-5
- Björk, T., & Murgoci, A. (2014). A theory of Markovian time-inconsistent stochastic control in discrete time. Finance Stoch., 18(3), 545-592. doi: 10.1007/s00780-014-0234-y
- Björk, T., Murgoci, A., & Zhou, X. Y. (2014). Mean variance portfolio optimization with state dependent risk aversion. *Math. Finance*, $24(1)$, 1-24. doi: 10.1111/j.1467-9965.2011.00515.x
- Boissaux, M., & Schiltz, J. (2010). An optimal control approach to portfolio optimization with conditioning information. Luxembourg School of Finance Working Paper, 57(19), 2519-2526. doi: 10.1080/00268976.2010.508052
- Boissaux, M., & Schiltz, J. (2013). Conditioned higher-moment portfolio: Optimization using optimal control. in: Terraza v., razafitombo h. (eds) understanding investment funds. London: Palgrave Macmillan. doi: 10.1057/9781137273611_6
- Briec, W., Kerstens, K., & Jokung, O. (2007). Mean-variance-skewness portfolio performance gauging: A general shortage function and dual approach. Management Sci., 53(1), 135-149. doi: 10.1287/ mnsc.1060.0596
- Buckdahn, R., Djehiche, B., & Li, J. (2011). A general stochastic maximum principle for SDEs of mean-field type. Appl. Math. Optim., 64(2), 197-216. doi: 10.1007/s00245-011-9136-y
- Chabi-Yo, F. (2012). Pricing kernels with stochastic skewness and volatility risk. Management Sci., 58(3), 624-640. doi: 10.1287/mnsc.1110.1424
- Czichowsky, C. (2013). Time-consistent mean-variance portfolio selection in discrete and continuous time. Finance Stoch., 17(2), 227-271. doi: 10.1007/s00780-012-0189-9
- De Athayde, G. M., & Flôres, R. G. (2004). Finding a maximum skewness portfolio - a general solution to three-moments portfolio choice. J. Econom. Dynam. Control, 28(7), 1335-1352. doi: 10.1016/ s0165-1889(02)00084-2
- Ekeland, I., Mbodji, O., & Pirvu, T. A. (2012). Time-consistent portfolio management. SIAM J. Finan. Math., 3(1), 1-32. doi: 10.1137/100810034
- Ekern, S. (1980). Increasing Nth degree risk. Econom. Lett., 6(4), 329-333. doi: 10.1016/0165-1765(80) 90005-1
- Goldman, S. M. (1980). Consistent plans. Rev. Econ. Stud., 47(3), 533-537. doi: 10.2307/2297304
- Harvey, C. R., Liechty, J. C., Liechty, M. W., & Müller, P. (2010). Portfolio selection with higher moments. Quant. Finance, 10(5), 469-485. doi: 10.1080/14697681003756877
- He, X. D., & Jiang, Z. L. (2021). On the equilibrium strategies for time-inconsistent problems in continuous time. SIAM J. Control Optim., 59(5), 3860-3886. doi: 10.1137/20M1382106
- Hu, Y., Jin, H. Q., & Zhou, X. Y. (2012). Time-inconsistent stochastic linear-quadratic control. SIAM J. Control Optim., 50(3), 1548-1572. doi: 10.1137/110853960
- Hu, Y., Jin, H. Q., & Zhou, X. Y. (2017). Time-inconsistent stochastic linear-quadratic control: Characterization and uniqueness of equilibrium. $SIAM$ J. Control Optim., $52(2)$, 1261-1279. doi: 10.1137/15M1019040
- Huang, Y. J., & Zhou, Z. (2021). Strong and weak equilibria for time-inconsistent stochastic control in continuous time. Math. Oper. Res., 46(2), 428-451. doi: 10.1287/moor.2020.1066
- Jean, W. H. (1971). The extension of portfolio analysis to three or more parameters. J. Financ. Quant. Anal., $6(1)$, 505-515. doi: 10.2307/2330125
- Joro, T., & Na, P. (2006). Portfolio performance evaluation in a mean-variance-skewness framework. European J. Oper. Res., 175(1), 446-461. doi: 10.1016/j.ejor.2005.05.006
- Konno, H., Shirakawa, H., & Yamazaki, H. (1993). A mean-absolute deviation-skewness portfolio optimization model. Ann. Oper. Res., 45(1), 205-220. doi: 10.1007/bf02282050
- Konno, H., & Suzuki, K. (1995). A mean-variance-skewness portfolio optimization model. J. Oper. Res. Soc. Japan, 38(2), 173-187. doi: 10.15807/jorsj.38.173
- Kraus, A., & Litzenberger, R. H. (1976). Skewness preference and the valuation of risk assets. J. Finance, $31(4)$, 1085-1100. doi: $10.1111/j.1540-6261.1976.$ tb01961.x
- Lai, T. Y. (1991). Portfolio selection with skewness: A multiple-objective approach. Rev. Quant. Finance Account., 1(3), 293-305. doi: 10.1007/bf02408382
- Laibson, D. (1997). Golden eggs and hyperbolic discounting. Quart. J. of Econom., 112(2), 443-477. doi: 10.1162/003355397555253
- Laibson, D. (1998). Life-cycle consumption and hyperbolic discount functions. European Econom. Rev., 42(3-5), 861-871. doi: 10.1016/S0014-2921(97)00132-3
- Landsman, Z., Makov, U., & Shushi, T. (2020). Analytic solution to the portfolio optimization problem in a mean-variance-skewness model. European J. Finance, 26(2-3), 165-178. doi: 10.1080/1351847X .2019.1618363
- Li, D., & Ng, W. (2000). Optimal dynamic portfolio selection: Multi-period mean-variance formulation. Math. Finance, 10(3), 387-406. doi: 10.1111/1467-9965.00100
- Liang, Z. X., Xia, J. M., & Yuan, F. Y. (2023). Dynamic portfolio selection for nonlinear law-dependent preferences. arXiv:2311.06745v2, 1-36. doi: 10.48550/arXiv.2311.06745
- Liu, J. Z., Lin, L. Y., Yiu, K. F. C., & Wei, J. Q. (2020). Non-exponential discounting portfolio management with habit formation. *Math. Control Relat. Fields*, 10(4), 761-783. doi: 10.3934/ mcrf.2020019
- Markowitz, H. (1952). Portfolio selection. J. Finance, 7(1), 77-91. doi: 10.2307/2975974
- Mehlawat, M. K., Gupta, P., & Khan, A. Z. (2021). Portfolio optimization using higher moments in an uncertain random environment. *Inform. Sci.*, 567, 348-374. doi: 10.1016/j.ins.2021.03.019
- Peleg, B., & Yaari, M. E. (1973). On the existence of a consistent course of action when tastes are changing. Rev. Econ. Stud., 40(3), 391-401. doi: 10.2307/2296458
- Peng, S. G. (1990). A general stochastic maximum principle for optimal control problems. SIAM J. Control Optim., 28(4), 966-979. doi: 10.1137/0328054
- Pollak, R. A. (1968). Consistent planning. Rev. Econ. Stud., 35(2), 201-208. doi: 10.2307/2296548
- Samuelson, P. A. (1970). The fundamental approximation theorem of portfolio analysis in terms of means, variances and higher moments. Rev. Econ. Stud., 37(4), 537-542. doi: 10.2307/2296483
- Scott, R. C., & Horvath, P. A. (1980). On the direction of preference for moments of higher order than the variance. J. Finance, 35(4), 915-919. doi: 10.1111/j.1540-6261.1980.tb03509.x
- Strotz, R. H. (1955). Myopia and inconsistency in dynamic utility maximization. Rev. Econ. Stud., 23(3), 165-180. doi: 10.1007/978-1-349-15492-0_10
- Wang, T. X., & Zheng, H. (2021). Closed-loop equilibrium strategies for general time-inconsistent optimal control problems. SIAM J. Control Optim., 59(5), 3152-3178. doi: 10.1137/20M1377242
- Yong, J. M. (2011). A deterministic linear quadratic time-inconsistent optimal control problem. Math. Control Relat. Fields, 1(1), 83-118. doi: 10.3934/mcrf.2011.1.83
- Yong, J. M. (2012). Time-inconsistent optimal control problems and the equilibrium HJB equation. Math. Control Relat. Fields, 2(3), 271-329. doi: 10.3934/mcrf.2012.2.271
- Yong, J. M. (2017). Linear-quadratic optimal control problems for mean-field stochastic differential equations – time-consistent solutions. Trans. Amer. Math. Soc., $369(8)$, 5467-5523. doi: 10.1090/ tran/6502
- Yong, J. M., & Zhou, X. Y. (1999). Stochastic controls: Hamiltonian systems and HJB equations (Vol. 43). New York: Springer-Verlag. doi: 10.1007/978-1-4612-1466-3
- Zhang, J. H., Purcal, S., & Wei, J. Q. (2021). Optimal life insurance and annuity demand under hyperbolic discounting when bequests are luxury goods. Insurance Math. Econom., 101, 80-90. doi: 10.1016/j.insmatheco.2020.07.003
- Zhao, Q., Shen, Y., & Wei, J. Q. (2014). Consumption-investment strategies with non-exponential discounting and logarithmic utility. European J. Oper. Res., 238(3), 824-835. doi: 10.1016/j.ejor .2014.04.034
- Zhen, F., & Chen, J. N. (2022). A closed-form mean-variance-skewness portfolio strategy. Finance Res. Lett., 47(2), 102933. doi: 10.1016/j.frl.2022.102933
- Zhou, X. Y., & Li, D. (2000). Continuous-time mean-variance portfolio selection: a stochastic LQ framework. Appl. Math. Optim., 42(1), 19-33. doi: 10.1007/s002450010003