On the Maximum and Minimum of a Multivariate Poisson Distribution

Zheng Liu¹, Feifan Shi¹, Jing Yao¹, Yang Yang¹

¹ Center for Financial Engineering, Soochow University, China

15th June 2024

Abstract

In this paper, we investigate the cumulative distribution functions (CDFs) of the maximum and minimum of multivariate Poisson distributions with three dependence structures, namely, the common shock, comonotonic shock and thinning-dependence models. In particular, we formulate the definition of a thinning-dependent multivariate Poisson distribution based on Wang and Yuen (2005). We derive explicit CDFs of the maximum and minimum of the multivariate Poisson random vectors and conduct asymptotic analyses on them. Our results reveal the substantial difference between the three dependence structures for multivariate Poisson distribution and may suggest an alternative method for studying the dependence for other multivariate distributions. We further provide numerical examples demonstrating obtained results.

Keywords: Maximum; Minimum; Multivariate Poisson distribution; Common shock structure; Comonotonic shock structure; Thinning-dependence structure.

1 Introduction

Studies on the distribution of the maximum (minimum) of a d-dimension random vector $\mathbf{X}=(X_1,\cdots,X_d)$ have been garnering considerable ongoing attention in the literature. Such a distribution of the maximum (minimum) serves as a tailor-made tool for modeling extreme events in diverse fields. For instance, it is widely employed in hydrology (Salas et al., 2018), biostatistics (Cheng et al., 2008), as well as insurance and finance applications (Embrechts et al., 2013). Therefore, the investigation of the distribution of $M^{\mathbf{X}}=\max{(X_1,\cdots,X_d)}$ or equivalently that of $m^{\mathbf{X}}=\min{(X_1,\cdots,X_d)}$ holds particular importance and many scholars have delved into this topic in the literature

For the classic normal distribution, in the case of d=2 (i.e., the bivariate normal distribution), Basu and Ghosh (1978) investigated the distribution of M^X , while Cain (1994) obtained its first two moments using the moment-generating function. Building upon this work, Ker (2001) examined how the mean and variance of M^X behave with changes in the mean, variance, and covariance of X. When considering $d \geq 3$ (i.e., the multivariate normal distribution case), Nadarajah et al. (2019) derived explicit expressions for the probability density function (PDF) and moments of M^X . Furthermore, as a variant of the normal distribution, the log-normal distribution is frequently utilized for modeling stochastic price/loss in finance and insurance. Its maximum and minimum hold considerable significance in applications, and these distributions have been studied by Lien (1986, 2005).

For other distributions, Jamalizadeh and Balakrishnan (2010) investigated the distributions of order statistics and linear combinations of order statistics derived from an elliptical distribution. They showed that these distributions can be expressed as mixtures of unified skew-elliptical distributions. Hakamipour et al. (2011) derived the distributional properties of the minimum and maximum when X has a bivariate Pareto distribution, and this result was extended to the multivariate case by Nadarajah and Woo (2013). Unlike previous studies,

^{*}Corresponding author, e-mail: j.yao@suda.edu.cn

Arellano-Valle and Genton (2008) derived a more general expression for the maximum distribution applicable to absolutely continuous random vectors, and they specifically investigated the scenario where the random vector follows an elliptically contoured distribution.

However, as mentioned above, the existing literature focuses on continuous random vectors, leaving a research gap concerning multivariate discrete distributions. The primary reasons for this gap could be two-fold. On one hand, discrete random vectors do not have PDF, rendering many abovementioned methods, such as Arellano-Valle and Genton (2008)'s general expression, inapplicable. For discrete distributions, we are limited to study the cumulative distribution function (CDF) and probability mass function (PMF). On the other hand, the probability that multiple variables within a random vector reach maximum or minimum values simultaneously is strictly larger than 0, i.e.

$$\mathbb{P}(M^{\mathbf{X}} = X_i = X_j, \forall i \neq j) > 0, \quad \text{(or } \mathbb{P}(m^{\mathbf{X}} = X_i = X_j, \forall i \neq j) > 0, \text{)},$$

which makes studying the maximum (minimum) of multivariate discrete random vectors even more challenging. In the continuous case, these probabilities are all equal to zero.

Despite these challenges, addressing this research gap is crucial due to the widespread applications and significance of discrete distributions in various fields. Among discrete distributions, the Poisson distribution, as a fundamental cornerstone in probability theory, plays a pivotal role in mathematical statistics, financial risk management, and biology (see Bailey, 1995; Chua and Goh, 2005; Inouye et al., 2017; Lindskog and McNeil, 2003). In this regard, we focus on the distribution of maximum and minimum in the context of multivariate Poisson distributions in this paper. Particularly, the common shock structure (Yuen et al., 2015) is one of the most prevalent and effectively depicts the dependence on many real-world events. For instance, within the insurance industry, the claims of different types of insurance business are often dependent. A natural disaster may trigger various types of insurance claims, including but not limited to medical, death, and property claims, etc. Such systematic impact of the natural disaster can be captured by a common shock in generating multivariate claims. Furthermore, Schulz et al. (2021) pointed out that the common shock structure inherently imposes constraints on pairwise covariances, requiring them to be positive and identical, which may not accurately represent some real-world scenarios. To overcome this limitation, Schulz et al. (2021) extended the common shock structure to a comonotonic shock structure. This extension provides enhanced flexibility in capturing the underlying dependence structure. By contrast, the thinning-dependence structure, introduced by Wang and Yuen (2005) to characterize continuoustime risk models in credit risk theory and the insurance business, incorporates a number of common background risks that may cause common claims of X_i , i = 1, ..., d. Likewise, in the insurance industry example mentioned earlier, the thinning-dependence structure not only captures the impact of background common risk sources (e.g. a natural disaster) but also gives the probabilities to different types of insurance claims (X_i) triggered by the same background risk sources, offering another flexible dependence with a more interpretive structure.

Although the three dependence structures share a similar idea by correlating marginal Poisson random variables via certain common factors, they distinguish from each other in how the common factors contribute to the marginal random variables $X_i, i=1,\ldots,d$. The common shock structure utilizes a unique random variable Y_0 to depict the impact of the common claim. This seems intuitive and straightforward, but also imposes specific constraints. The comonotonic shock is a natural extension to common shock. It replaces the unique random shock variable Y_0 in marginal random variables by a comonotonic random vector $\mathbf{Z} = (Z_1, \ldots, Z_d)$, providing more flexible "shocks" in the dependent structures. Nevertheless, both common shock and comonotonic shock structures incorporate the common factor as "shocks" directly into the system. Specifically, they add the "shocks" (the shock variable Y_0 or the shock vector \mathbf{Z}) into another independent random vector $\mathbf{Y} = (Y_1, \ldots, Y_d)$, which stands for the idiosyncratic

$$(Z_1,\ldots,Z_d) = \left(F_1^{-1}(U),\ldots,F_d^{-1}(U)\right),\,$$

where for each $j \in \{1, ..., d\}$, F_j denotes the distribution of Z_j . More details can be found in Schulz (2019). Moreover, in this paper, for a real function $F, F^{-1}(u) = \inf\{t \in \mathbb{R} : F(t) \geq u\}$.

¹A random vector (Z_1, \ldots, Z_d) is comonotonic if and only if each component can be represented in terms of a common underlying uniform random variable on the unit interval. That is, for a uniform random variable $U \sim \mathcal{U}(0, 1)$, the comonotonic vector can be written as

marginal shocks. On the contrary, simultaneous claims (shocks) in the thinning-dependence structure are triggered by commonly shared background random variables (risk sources), tying marginal Poisson random variables up via an explanatory probabilistic mechanism. We present more details on the three dependence structures in Section 2.

In this paper, we explore the maximum and minimum of the multivariate Poisson distribution with dependence. Our contributions to the existing literature can be summarized in threefold. Firstly, to the best of our knowledge, we are the first to investigate the distribution of maximum and minimum of the multivariate Poisson distribution with dependence. More specifically, we explicitly derive the CDFs of the M^X and m^X of the multivariate Poisson distribution under the three aforementioned dependent structures, namely, the common shock, comonotonic shock, and thinning-dependence structures. In doing so, we theoretically formulate a multivariate Poisson distribution with thinning-dependence structure, which broadens the scope of this dependence structure from stochastic processes to multivariate probability distributions, offering a pedagogical contribution.

Secondly, we perform asymptotic analysis of all derived CDFs and obtain several interesting results that underpin the substantial differences between the three dependence structures. For example, when the dimension $d \to \infty$, we find that the asymptotic behaviors of derived CDFs under the common shock and comonotonic shock structures are similar, whereas the one under the thinning-dependence structure exhibits evident discrepancy. Specifically, the asymptotic behavior of M^X aligns with the commonly shared background risk sources (more precisely, the convolution of background random variables) under the thinning-dependence structure. By contrast, under the common shock and comonotonic shock structures, the CDFs of M^{X} are asymptotically determined by the idiosyncratic random vector Y (Y_0 for common shock structure), which is independent of common shocks. Furthermore, despite the tight connection, the asymptotic analysis on d for comonotonic structure is mathematically much more challenging than common shock structure because the dimension of the shock vector Z increases along d, indicating the number of comonotonic shock variables Z_i also approach infinity. As such, the comonotonic structure differs significantly from the common shock structure and requires additional assumptions on the comonotonic shock vector Z to obtain similar asymptotic results. Otherwise, the asymptotic results may no longer hold. To illustrate this point, we provide a counterexample in Remark 4.3 where the assumptions on Z are not valid. Such finding reveals the intrinsic difference between the common shock and comonotonic shock structures, and the difference is more obvious as d is high. Moreover, we also consider the asymptotic scenarios when certain key parameters dominate the dependence structures. For instance, it is intuitive to expect that when the rate of the common shock variable Y_0 dominates the other Poisson rates, the distribution of M^X and m^X should considerably depend on the distribution of Y_0 . Our results validate this analytically and quantify the convergency rate between the CDF of Y_0 and those of M^X and m^X , which is in line with intuitive expectations. Overall, our asymptotic analyses, to some extent, provide an alternative new way to study similar dependence structures for other multivariate distributions from an asymptotic standpoint.

Thirdly, we present numerical examples to demonstrate the influence of different dependent structures on the M^X and m^X of multivariate Poisson distributions. To highlight this impact, it is essential to ensure consistent settings for a relatively "fair" comparison of the three structures. To this end, we first consider a setting that marginal Poisson rates. As for internal correlations, note that ensuring a setting of pairwisely identical correlations is very difficult, especially for large d, because it requires equal $(d \times d - 1)/2$ correlation coefficients across all three structures with identical marginal Poisson rates. Hence, instead of pairwise correlation, we consider the same average correlation coefficient for the three structures, which can independently describe the overall correlation of a multivariate random vector. Based on the above settings, we compare their CDFs in numerical examples under two scenarios: one where the common shock and thinning-dependence structures differ significantly and another where they are very similar. Our results show that when the structures are similar, the CDF values are close. However, when the structural differences are significant, the difference in CDF is pronounced. These findings strongly demonstrate that the construction of dependent structures plays a crucial role in determining the maximum and minimum values. Compared with Karlis (2003); Schulz et al. (2021) and Wang and Yuen (2005), our numerical results further analyze these three dependent structures from the perspective of their maximum (minimum) values, offering an additional viewpoint to discuss the similarities and differences between them.

The rest of paper is organized as follows. Section 2 outlines the definition of the multivariate Poisson-

distributed random variable under common shock, comonotonic shock, and thinning-dependence structure. The CDFs of M^X and m^X are obtained in Section 3. In Section 4, we show the results of our asymptotic analysis under the three structures. Section 5 presents the numerical illustration and Section 6 concludes the paper with the discussions.

2 Multivariate Poisson distribution with dependence

In this section, we utilize three distinct dependence structures: common shock, comonotonic shock, and thinning-dependence structures, to construct multivariate Poisson distributions. Specifically, we provide the mathematical definitions of these three multivariate Poisson distributions. Throughout this paper, $\mathcal{P}(\theta)$ denotes a univariate Poisson distribution with rate θ , $G_{\theta}(\cdot)$ represents its CDF, and $\bar{G}_{\theta}(\cdot)$ is the corresponding survival function.

2.1 Common shock structure

In the literature, several methods for constructing the common shock structure have been proposed, such as those by Mahamunulu (1967) and Kawamura (1976), which consider multiple random variables as the common shock components. Although these approaches are intuitive and flexible, deriving the explicit expression of the distribution becomes complex, as noted by Karlis and Meligkotsidou (2007) and Inouye et al. (2017). This complexity poses challenges for studying the distribution of the maximum or minimum. Therefore, in this paper, following the approach of Karlis (2003), we consider only the single common shock variable case. The definition of the multivariate common shock Poisson distribution is given as follows:

Definition 2.1 A random vector $(X_1, ..., X_d)$ is said to have a multivariate common shock Poisson distribution with parameters $(\theta_0, \theta_1, ..., \theta_d) \in (0, \infty)^{d+1}$ if it can be expressed as follow:

$$X_i = Y_0 + Y_i, \quad i = 1, 2, \dots, d,$$
 (1)

where $Y_i \sim \mathcal{P}(\theta_i)$, i = 0, 1, ..., d are mutually independent Poisson random variables.

The multivariate Poisson model with the common shock structure is applied in various fields, including market research and epidemiology (see Karlis, 2003). However, this structure also imposes strong limitations. Specifically, it imposes constraints on the correlation between any two variables in a multivariate common shock Poisson vector. For arbitrary pair of Poisson random variables with rates θ_i and θ_j , Griffiths et al. (1979) demonstrated that their correlation falls in the interval $[\rho_{\min}, \rho_{\max}]$ where

$$\rho_{\min} = \frac{1}{\sqrt{\theta_{i}\theta_{j}}} \left[-\theta_{i}\theta_{j} - \sum_{m \in \mathbb{N}} \sum_{n \in \mathbb{N}} \min \left\{ 0, G_{\theta_{i}}\left(m\right) + G_{\theta_{j}}\left(n\right) - 1 \right\} \right],$$

$$\rho_{\max} = \frac{1}{\sqrt{\theta_{i}\theta_{j}}} \left[-\theta_{i}\theta_{j} + \sum_{m \in \mathbb{N}} \sum_{n \in \mathbb{N}} \min \left\{ \overline{G}_{\theta_{i}}\left(m\right), \overline{G}_{\theta_{j}}\left(n\right) \right\} \right].$$

Nevertheless, within the common shock structure, the correlation coefficient will not reach ρ_{max} unless the marginal Poisson rates are identical.

2.2 Comonotonic shock structure

To establish a more flexible dependence structure, Schulz et al. (2021) substituted a comonotonic vector Z for the single common shock variable Y_0 , allowing each X_i , $i=1,\ldots,d$, to have its own comonotonic shock variable Z_i . It is important to highlight that since Z is comonotonic, all comonotonic shock variables Z_i are controlled by the same uniform random variable U, essentially reflecting the influence of the common factor. This structure enhances the flexibility of dependent structures while maintaining intuitiveness and interpretability. Its mathematical definition is as follows:

Definition 2.2 A random vector $(X_1, ..., X_d)$ is said to follow the multivariate comonotonic shock Poisson distribution with parameters $\Lambda_d = (\lambda_1, ..., \lambda_d) \in (0, \infty)^d$ and $\theta \in [0, 1]$, if it can be expressed in

$$(X_1,\ldots,X_d)=(Y_1,\ldots,Y_d)+(Z_1,\ldots,Z_d),$$

in terms of two independent random vectors (Y_1, \ldots, Y_d) and (Z_1, \ldots, Z_d) such that

- (i) Y_1, \ldots, Y_d are mutually independent and, for all $j \in \{1, \ldots, d\}$, $Y_j \sim \mathcal{P}((1 \theta) \lambda_j)$;
- (ii) Z_1, \ldots, Z_d are comonotonic and, for all $j \in \{1, \ldots, d\}$, $Z_j \sim \mathcal{P}(\theta \lambda_j)$.

Notably, the marginals are mutually independent when $\theta=0$, and (X_1,\ldots,X_d) is comonotonic when $\theta=1$. Moreover, when $\lambda_1=\lambda_2=\cdots=\lambda_d$, the multivariate comonotonic shock Poisson distribution reduces to the multivariate common shock Poisson distribution. In contrast to the common shock structure, the introduction of the comonotonic shock vector enhances the flexibility of this dependent structure. For instance, the correlation coefficient of any pair (X_i,X_j) , $i\neq j$, can attain ρ_{\max} without requiring the marginal Poisson rates to be equal.

2.3 Thinning-dependence structure

Since Wang and Yuen (2005) introduced correlated Poisson processes with a thinning-dependence structure, most studies have applied this framework exclusively to describe the correlation of stochastic processes. In this paper, we extend the thinning-dependence structure to random vectors and propose a multivariate thinning-dependence Poisson distribution, defined as follows.

Definition 2.3 Assuming

$$X = (X_1, \dots, X_d), \quad X_j = \sum_{k=1}^l X_j^k, \quad j = 1, \dots, d,$$
 (2)

and Y_1, Y_2, \ldots, Y_l are independent Poisson random variables with parameters $\theta_1, \ldots, \theta_l$, respectively. Then, $X = (X_1, \ldots, X_d)$ is said to follow the multivariate thinning-dependence Poisson distribution, if, (i) the random variables $X_j^k | Y_k = y_k, j = 1, \ldots, d, k = 1, \ldots, l$ follow the binomial distribution with parameters

 y_k and p_j^k , that is,

 $\mathbb{P}\left(X_j^k = z | Y_k = y_k\right) = \binom{y_k}{z} (p_j^k)^z \left(1 - p_j^k\right)^{y_k - z},\tag{3}$

where $z \leq y_k$;

(ii) X_1^k, \ldots, X_d^k are conditionally independent given $Y_k = y_k, k = 1, \ldots, l$;

(iii) the l random vectors $(Y_1, X_1^1, \dots, X_d^1), \dots, (Y_l, X_1^l, \dots, X_d^l)$ are independent.

Remark 2.1 For $j=1,\ldots,d,$ $k=1,\ldots,l$, the random variables X_j and X_j^k are all Poisson-distributed. Specifically, we have $X_j^k \sim \mathcal{P}(p_j^k \theta_k)$ and $X_j \sim \mathcal{P}(\sum_{k=1}^l p_j^k \theta_k)$, for $j=1,\ldots,d,$ $k=1,\ldots,l$. This follows from Definition 2.3, where

$$\mathbb{P}\left(X_j^k = z\right) = \sum_{y_k = z}^{\infty} \mathbb{P}\left(X_j^k = z | Y_k = y_k\right) \mathbb{P}(Y_k = y_k) = e^{-(p_j^k \theta_k)} \frac{(p_j^k \theta_k)^z}{z!},$$

implying $X_j^k \sim \mathcal{P}(p_j^k \theta_k)$. Moreover, due to the independence of X_j^1, \dots, X_j^l mentioned in (iii) of Definition 2.3, we have $X_j \sim \mathcal{P}(\sum_{k=1}^l p_j^k \theta_k)$.

Remark 2.2 We provide an explanation of credit risk to Definition 2.3 to further illustrate the idea behind the dependence structure. Consider d firms with a solo systematic background risk (e.g., the default risk of major clients who have business with all firms). Let Y_1 and $X_j^1, j=1,\ldots,d$, denote the default times of a common client and the firms, respectively. For each firm, the systematic default of Y_1 may trigger the j-th firm's default with probability p_j^1 . Hence, it is straightforward to see $X_j^1|Y_1=y_1$ indicates y_1 trials of systematic default event, which follows a binomial distribution, and these trials are mutually independent when the solo background risk is known.

Remark 2.3 The thinning-dependence structure may reduce to common shock structure. Specifically, consider the case where l = d + 1, and

$$\begin{bmatrix} p_1^1 & p_1^2 & \cdots & p_1^d & p_1^{d+1} \\ p_2^1 & p_2^2 & \cdots & p_2^d & p_2^{d+1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ p_d^1 & p_d^2 & \cdots & p_d^d & p_d^{d+1} \end{bmatrix} = \begin{bmatrix} 1 & 0 & \cdots & 0 & 1 \\ 0 & 1 & \cdots & 0 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 1 \end{bmatrix},$$

where p_j^k , $j=1,\ldots,d, k=1,\ldots,d+1$ are the probabilities given in Definition 2.3. In this scenario, background random variable $Y^{d+1}\sim \mathcal{P}(\theta_{d+1})$ affects X_j^{d+1} , $j=1,\ldots,d$ with probability 1, implying that the Y^{d+1} reduces to the common shock variable. Meanwhile, for $k=1,\ldots,d$, the background random variable $Y_k\sim \mathcal{P}(\theta_k)$ affects only X_k^k with probability 1 and does not affect X_j^k , for $j\neq k$, implying $X_k^k=Y^k$, $X_j^k=0$. Thus, according on Definition 2.3, for $j=1,\ldots,d$, we have $X_j=\sum_{k=1}^{d+1}X_j^k=Y_j+Y_{d+1}$, indicating that the multivariate thinning-dependence Poisson distribution reduces to the multivariate common shock Poisson distribution under this parameter setting.

3 CDFs of maximum and minimum of multivariate Poisson distribution

In this section, we explicitly derive CDFs for the maximum and minimum of the multivariate Poisson distribution under the common shock, comonotonic shock, and thinning-dependence structures.

Before presenting the aforementioned results, we first introduce the CDF ², and survival function of the univariate Poisson random variable in the following lemma.

Lemma 3.1 Let a random variable $X \sim \mathcal{P}(\lambda)$, then the CDF and survival function of X are

$$G_{\lambda}(x) = \frac{\Gamma(x+1,\lambda)}{\Gamma(x+1)}, \quad \bar{G}_{\lambda}(x) = \frac{\gamma(x+1,\lambda)}{\Gamma(x+1)},$$

respectively, where

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt, \quad \Gamma(z, x) = \int_x^\infty t^{z-1} e^{-t} dt, \quad \gamma(z, x) = \int_0^x t^{z-1} e^{-t} dt,$$

are the gamma function, upper incomplete gamma function, and lower incomplete gamma function, respectively.

3.1 Common shock structure

For the multivariate common shock Poisson distribution defined in Definition 2.1, we can obtain the CDFs of the maximum and minimum by the conditional technique in the following theorem.

Theorem 3.1 Suppose $X = (X_1, ..., X_d)$ follows the multivariate common shock Poisson distribution defined in Definition 2.1, then the CDFs of $M^X = \max(X_1, ..., X_d)$ and $m^X = \min(X_1, ..., X_d)$ are

$$F_{Mx}\left(x\right) = \sum_{y=0}^{x} \frac{\theta_{0}^{y} e^{-\theta_{0}} \prod_{j=1}^{d} \Gamma\left(x-y+1,\theta_{j}\right)}{\Gamma\left(y+1\right) \Gamma^{d}\left(x-y+1,\right)}, \quad F_{mx}\left(x\right) = \sum_{y=0}^{x} \frac{\theta_{0}^{y} e^{-\theta_{0}} \left[\Gamma^{d}\left(x-y+1\right) - \prod_{j=1}^{d} \gamma\left(x-y+1,\theta_{j}\right)\right]}{\Gamma\left(y+1\right) \Gamma^{d}\left(x-y+1\right)},$$

where $x \in \mathbb{N}$.

Proof For the CDF of M^X , conditioning on $Y_0 = y_0$, we have,

$$F_{MX}(x) = \sum_{y=0}^{x} \left[\mathbb{P}(Y_0 = y) \prod_{j=1}^{d} \mathbb{P}(Y_j \le x - y) \right] = \sum_{y=0}^{x} \frac{\theta_0^y e^{-\theta_0} \prod_{j=1}^{d} \Gamma(x - y + 1, \theta_j)}{\Gamma(y + 1) \Gamma^d(x - y + 1)},$$

²Notably, the distributions studied in this paper are discrete, all defined on non-negative integers. Let $\hat{F}(x)$ denote the CDF when $x \in \mathbb{N}$. When x > 0 and $\notin \mathbb{N}$, $F(x) = \hat{F}(\lfloor x \rfloor)$, where $\lfloor \cdot \rfloor$ represents rounding down. Therefore, for convenience, we only need to focus on the case where the variable x assumes non-negative integer values in Sections 3 and 4 unless otherwise specified.

where the last equality is from Lemma 3.1. Similarly, for the CDF of m^X , we have,

$$F_{m\mathbf{x}}\left(x\right) = \sum_{y=0}^{x} \frac{\theta_{0}^{y} e^{-\theta_{0}} \left[\Gamma^{d}\left(x-y+1\right) - \prod_{j=1}^{d} \gamma\left(x-y+1,\theta_{j}\right)\right]}{\Gamma\left(y+1\right) \Gamma^{d}\left(x-y+1\right)}.$$

3.2 Comonotonic shock structure

Similar to the common shock structure, we can utilize the conditional technique to explicitly derive the CDFs of M^X and m^X again. However, replacing X's single common shock variable with a d-dimensional components shock vector adds additional complexity to the derivation of the distributions of M^X and m^X , as detailed in the following theorem.

Theorem 3.2 Suppose $X = (X_1, ..., X_d)$ follows the multivariate comonotonic shock Poisson distribution defined in Definition 2.2, then the CDFs of $M^X = \max(X_1, ..., X_d)$ and $m^X = \min(X_1, ..., X_d)$ are

$$F_{MX}(x) = \sum_{z_1=0}^{x} \cdots \sum_{z_d=0}^{x} \prod_{j=1}^{d} g_{(1-\theta)\lambda_j}(x-z_j) \min \{G_{\theta\lambda_1}(z_1), \dots, G_{\theta\lambda_d}(z_d)\},$$
(4)

or equivalently,

$$F_{M}x(x) = \sum_{z_{1}=0}^{x} \cdots \sum_{z_{d}=0}^{x} \prod_{j=1}^{d} G_{(1-\theta)\lambda_{j}}(x-z_{j}) C_{\Lambda,\theta}(z_{1},\ldots,z_{d}),$$
 (5)

and

$$F_{mx}(x) = \sum_{z_1=0}^{\infty} \cdots \sum_{z_d=0}^{\infty} \prod_{j=1}^{d} g_{(1-\theta)\lambda_j}(z_j) \max \{G_{\theta\lambda_1}(x-z_1), \dots, G_{\theta\lambda_d}(x-z_d)\},$$
 (6)

or equivalently,

$$F_{m\mathbf{x}}(x) = 1 - \sum_{z_1=0}^{\infty} \cdots \sum_{z_d=0}^{\infty} \prod_{j=1}^{d} \overline{G}_{(1-\theta)\lambda_j}(x-z_j) C_{\Lambda,\theta}(z_1,\ldots,z_d),$$

$$(7)$$

where $x \in \mathbb{N}$, $g_{\lambda}(x)$ denotes the PMF of the Poisson distribution with parameter λ ,

$$C_{\Lambda,\theta}\left(z_{1},\ldots,z_{d}\right)=\left[\min\left\{G_{\theta\lambda_{1}}\left(z_{1}\right),\ldots,G_{\theta\lambda_{d}}\left(z_{d}\right)\right\}-\max\left\{G_{\theta\lambda_{1}}\left(z_{1}-1\right),\ldots,G_{\theta\lambda_{d}}\left(z_{d}-1\right)\right\}\right]_{+},$$
 and $\left[\cdot\right]_{+}=\max\{0,\cdot\}$.

Proof Firstly, utilizing the conditional technique on Z or Y, the CDF of M^X can be written as the following two forms:

$$F_{MX}(x) = \sum_{z_1=0}^{x} \cdots \sum_{z_d=0}^{x} \mathbb{P}(Z_1 = z_1, \dots, Z_d = z_d) \mathbb{P}(Y_1 \le x - z_1, \dots, Y_d \le x - z_d)$$

$$= \sum_{z_1=0}^{x} \cdots \sum_{z_d=0}^{x} \mathbb{P}(Y_1 = x - z_1, \dots, Y_d = x - z_d) \mathbb{P}(Z_1 \le z_1, \dots, Z_d \le z_d).$$
(8)

The key point of above formula is to calculate $\mathbb{P}(Z_1 = z_1, \dots, Z_d = z_d)$ and $\mathbb{P}(Z_1 \leq z_1, \dots, Z_d \leq z_d)$. To obtain (4), we calculate the first one,

$$C_{\Lambda,\theta}(z_{1},...,z_{d}) = \mathbb{P}(Z_{1} = z_{1},...,Z_{d} = z_{d})$$

$$= \mathbb{P}(U \in (G_{\theta\lambda_{1}}(z_{1} - 1), G_{\theta\lambda_{1}}(z_{1})],...,U \in (G_{\theta\lambda_{d}}(z_{d} - 1), G_{\theta\lambda_{d}}(z_{d})])$$

$$= \mathbb{P}(\max\{G_{\theta\lambda_{1}}(z_{1} - 1),...,G_{\theta\lambda_{d}}(z_{d} - 1)\} < U \leq \min\{G_{\theta\lambda_{1}}(z_{1}),...,G_{\theta\lambda_{d}}(z_{d})\})$$

$$= \left[\min \left\{G_{\theta \lambda_1}\left(z_1\right), \dots, G_{\theta \lambda_d}\left(z_d\right)\right\} - \max \left\{G_{\theta \lambda_1}\left(z_1 - 1\right), \dots, G_{\theta \lambda_d}\left(z_d - 1\right)\right\}\right]_+,$$

where U is a uniform random variable and $U \sim \mathcal{U}(0, 1)$.

To obtain (5), we calculate the second one,

$$\mathbb{P}\left(Z_{1} \leq z_{1}, \ldots, Z_{d} \leq z_{d}\right) = \min\left\{G_{\theta\lambda_{1}}\left(z_{1}\right), \ldots, G_{\theta\lambda_{d}}\left(z_{d}\right)\right\}.$$

Secondly, for the CDF of m^X , we have,

$$F_{m}\mathbf{x}(x) = 1 - \sum_{z_{1}=0}^{\infty} \cdots \sum_{z_{d}=0}^{\infty} \mathbb{P}(Z_{1} = z_{1}, \dots, Z_{d} = z_{d}) \, \mathbb{P}(Y_{1} \ge x - z_{1} + 1, \dots, Y_{d} \ge x - z_{d} + 1)$$

$$= 1 - \sum_{z_{1}=0}^{\infty} \cdots \sum_{z_{d}=0}^{\infty} \mathbb{P}(Y_{1} = z_{1}, \dots, Y_{d} = z_{d}) \, \mathbb{P}(Z_{1} \ge x - z_{1} + 1, \dots, z_{d} \ge x - z_{d} + 1). \tag{9}$$

Meanwhile, we have

$$\mathbb{P}(Z_1 \geq z_1, \dots, Z_d \geq z_d) = 1 - \max\{G_{\theta \lambda_1}(z_1 - 1), \dots, G_{\theta \lambda_d}(z_d - 1)\},\$$

and

$$1 = \sum_{z_1=0}^{\infty} \cdots \sum_{z_d=0}^{\infty} \prod_{j=1}^{d} g_{(1-\theta)\lambda_j}(z_j) = \sum_{z_1=0}^{\infty} g_{(1-\theta)\lambda_1}(z_1) \cdots \sum_{z_d=0}^{\infty} g_{(1-\theta)\lambda_d}(z_d).$$

Therefore, we can derive two forms of expression for $F_{mx}(x)$: For (6),

$$F_{m}x(x) = 1 - \sum_{z_{1}=0}^{\infty} \cdots \sum_{z_{d}=0}^{\infty} \mathbb{P}\left(Y_{1} = z_{1}, \dots, Y_{d} = z_{d}\right) \mathbb{P}\left(Z_{1} \geq x - z_{1} + 1, \dots, z_{d} \geq x - z_{d} + 1\right)$$
$$= \sum_{z_{1}=0}^{\infty} \cdots \sum_{z_{d}=0}^{\infty} \prod_{j=1}^{d} g_{(1-\theta)\lambda_{j}}(z_{j}) \max \left\{G_{\theta\lambda_{1}}(x - z_{1}), \dots, G_{\theta\lambda_{d}}(x - z_{d})\right\}.$$

For (7),

$$F_{m}\mathbf{x}(x) = 1 - \sum_{z_{1}=0}^{\infty} \cdots \sum_{z_{d}=0}^{\infty} \mathbb{P}(Z_{1} = z, \dots, Z_{d} = z_{d}) \, \mathbb{P}(Y_{1} \ge x - z_{1} + 1, \dots, Y_{d} \ge x - z_{d} + 1)$$

$$= 1 - \sum_{z_{1}=0}^{\infty} \cdots \sum_{z_{d}=0}^{\infty} \prod_{i=1}^{d} \overline{G}_{(1-\theta)\lambda_{j}}(x - z_{j}) \, C_{\Lambda,\theta}(z_{1}, \dots, z_{d}).$$

Notably, in (8) and (9), we equivalently express the events $\{Y_1+Z_1 \leq x_1,\ldots,Y_d+Z_d \leq x_d\}$ and $\{Y_1+Z_1 \geq x_1+1,\ldots,Y_d+Z_d \geq x_d+1\}$ in two different forms, respectively. Consequently, we analytically derive two distinct CDFs for M^X , specifically (4) and (5). Similarly, for m^X , we derive (6) and (7). Although these expressions are equivalent, one may be more convenient for specific asymptotic analyses. For example, we use (5) in Proposition 4.5, while Propositions 4.4 and 4.6 utilize (4).

3.3 Thinning-dependence structure

As illustrated in Definition 2.3, for a multivariate thinning-dependence Poisson distributed random vector X, the components of marginal random variable X_j are controlled by background random variable Y_k (i.e., $X_j^k|Y_k=y_k\sim \text{Binomial}(p_j^k,y_k)$). This mechanism links all the marginal random variables through l background random variables Y_1,\ldots,Y_l indirectly. Therefore, it is necessary to first investigate the specific mechanism by which the background random variables influence the marginal random variables, i.e., the conditional marginal random variable $X_j|Y_1=y_1,\cdots,Y_l=y_l$. These results are presented in the following lemmas. The proof of Lemma 3.2 is straightforward and thus omitted here.

Lemma 3.2 Let $X = (X_1, ..., X_d)$ follows the multivariate thinning-dependence Poisson distribution, then $X_j, j = 1, ..., d$ are mutually independent given $Y_1 = y_1, ..., Y_l = y_l$.

 $X_j|Y_1=y_1,\cdots,Y_l=y_l$ may involve the sum of independent, non-identically distributed binomial random variables, complicating the derivation of the CDFs for M^X and m^X under the thinning-dependence structure. Hence, we first provide the explicit PMF of these conditional marginal random variables in the following lemma.

Lemma 3.3 Let $X = (X_1, ..., X_d)$ follows the multivariate thinning-dependence Poisson distribution defined in Definition 2.3. Then, for any j = 1, ..., d, when $s \le y_1 + \cdots + y_l$, we have,

$$P\left(X_{j} = s | Y_{1} = y_{1}, \cdots, Y_{l} = y_{l}\right) = \sum_{x_{j}^{1} = \max\left(0, s - \sum_{i=2}^{l} y_{i}\right)}^{\min\left(s, y_{1}\right)} \left\{ P\left(X_{j}^{1} = x_{j}^{1}\right) \sum_{x_{j}^{2} = \max\left(0, s - x_{j}^{1} - \sum_{i=3}^{l} y_{i}\right)}^{\min\left(s - x_{j}^{1}, y_{2}\right)} P\left(X_{j}^{2} = x_{j}^{2}\right) \right\}$$

$$\dots \left[\sum_{x_{j}^{k} = \max\left(0, s - \sum_{i=1}^{k-1} x_{j}^{i} - \sum_{i=k+1}^{l} y_{i}\right)}^{P\left(X_{j}^{k} = x_{j}^{k}\right)} P\left(X_{j}^{k} = x_{j}^{k}\right) \right]$$

$$\dots \left(\sum_{x_{j}^{l-1} = \max\left(0, s - \sum_{i=1}^{l-2} x_{j}^{i}, y_{l-1}\right)}^{\min\left(s - \sum_{i=1}^{l-2} x_{j}^{i}, y_{l-1}\right)} P\left(X_{j}^{l-1} = x_{j}^{l-1}\right) P\left(X_{j}^{l} = s - \sum_{i=1}^{l-1} x_{j}^{i}\right) \right] .$$

$$(10)$$

Proof From Definition 2.3, we can deduce that, given $Y_1 = y_1, \ldots, Y_l = y_l$, the random variables $X_j^k \sim \text{Binomial}(y_k, p_j^k)$, $k = 1, \ldots, l$, are independent. Thus, for any $j = 1, \ldots, d$, $X_j | Y_1 = y_1, \ldots, Y_l = y_l$ is the sum of d independent non-identically distributed binomial random variables, and its PMF is given by Eisinga et al. (2013).

Using the conditional marginal distribution and employing the conditional technique once more, we can derive the following theorem.

Theorem 3.3 Suppose $X = (X_1, ..., X_d)$ follows the multivariate thinning-dependence Poisson distribution defined in Definition 2.3, then the CDFs of $M^X = \max(X_1, ..., X_d)$ and $m^X = \min(X_1, ..., X_d)$ are

$$F_{MX}(x) = \sum_{y_1 + \dots + y_l \le x} \prod_{k=1}^{l} \frac{\theta_k^{y_k} e^{-\theta_k}}{\Gamma(y_k + 1)} + \sum_{y_1 + \dots + y_l \ge x + 1} \left\{ \left[\prod_{j=1}^{d} \sum_{s=0}^{x} h_j(s, y_1, \dots, y_l) \right] \left[\prod_{k=1}^{l} \frac{\theta_k^{y_k} e^{-\theta_k}}{\Gamma(y_k + 1)} \right] \right\}$$
(11)

and

$$F_{mx}(x) = \sum_{y_1 + \dots + y_l \le x} \prod_{k=1}^{l} \frac{\theta_k^{y_k} e^{-\theta_k}}{\Gamma(y_k + 1)} + \sum_{y_1 + \dots + y_l \ge x + 1} \left\{ \left[1 - \prod_{j=1}^{d} \left(1 - \sum_{s=0}^{x} h_j(s, y_1, \dots, y_l) \right) \right] \left[\prod_{k=1}^{l} \frac{\theta_k^{y_k} e^{-\theta_k}}{\Gamma(y_k + 1)} \right] \right\},$$
(12)

where $h_j(s, y_1, ..., y_l) = P(X_j = s | Y_1 = y_1, ..., Y_l = y_l)$, as explicitly given in Lemma 3.3.

Proof For $F_{MX}(x)$, we have,

$$\begin{split} F_{M^{\mathbf{X}}}\left(x\right) &= \mathbb{P}\left(X_{1} \leq x, \cdots, X_{d} \leq x\right) \\ &= \sum_{y_{k}=0, k=1, \dots, l}^{\infty} \mathbb{P}\left(Y_{1} = y_{1}, \cdots, Y_{l} = y_{l}\right) \mathbb{P}\left(X_{1} \leq x, \dots, X_{d} \leq x | Y_{1} = y_{1}, \dots, Y_{l} = y_{l}\right) \\ &= \sum_{y_{1}+\dots+y_{l} \leq x} \prod_{k=1}^{l} \mathbb{P}\left(Y_{k} = y_{k}\right) + \sum_{y_{1}+\dots+y_{l} \geq x+1} \prod_{j=1}^{d} \mathbb{P}\left(X_{j} \leq x | Y_{1} = y_{1}, \dots, Y_{l} = y_{l}\right) \prod_{k=1}^{l} \mathbb{P}\left(Y_{k} = y_{k}\right) \\ &= \sum_{y_{1}+\dots+y_{l} \leq x} \prod_{k=1}^{l} \frac{\theta_{k}^{y_{k}} e^{-\theta_{k}}}{\Gamma\left(y_{k}+1\right)} + \sum_{y_{1}+\dots+y_{l} \geq x+1} \left\{ \left[\prod_{j=1}^{d} \sum_{s=0}^{x} h_{j}\left(s, y_{1}, \dots, y_{m}\right)\right] \left[\prod_{k=1}^{l} \frac{\theta_{k}^{y_{k}} e^{-\theta_{k}}}{\Gamma\left(y_{k}+1\right)}\right] \right\}, \end{split}$$

For $F_{m} \mathbf{x}(x)$, we have,

$$F_{m\mathbf{x}}(x) = 1 - \mathbb{P}(X_1 > x, \dots, X_d > x)$$

$$= \sum_{y_1+\dots+y_l\geq x+1} \left\{ \left[1 - \prod_{j=1}^d \left(1 - \sum_{s=0}^x h_j\left(s,y_1,\dots,y_l\right)\right)\right] \left[\prod_{k=1}^l \frac{\theta_k^{y_k}e^{-\theta_k}}{\Gamma\left(y_k+1\right)}\right] \right\} + \sum_{y_1+\dots+y_l\leq x} \prod_{k=1}^l \frac{\theta_k^{y_k}e^{-\theta_k}}{\Gamma\left(y_k+1\right)}.$$

When the number of background random variables $l \ge 2$, the conditional marginal distribution (10) involves the sum of independent, non-identically distributed binomial random variables, making the explicit expressions (11) and (12) more complex. By contrast, in the case of l = 1, an elegant formula employing the regularized incomplete beta function is presented in the following corollary.

Corollary 3.1 Suppose $X = (X_1, ..., X_d)$ follows the multivariate thinning-dependence Poisson distribution defined in Definition 2.3 with l = 1, then the CDFs of $M^X = \max(X_1, ..., X_d)$ and $m^X = \min(X_1, ..., X_d)$ are

$$F_{MX}(x) = \frac{\Gamma(x+1,\theta_1)}{\Gamma(x+1)} + \sum_{y=x+1}^{\infty} \frac{\theta_1^y e^{-\theta_1}}{\Gamma(y+1)} \prod_{j=1}^d I_{1-p_j}(y-x,x+1),$$

and

$$F_{m\mathbf{x}}\left(x\right) = \frac{\Gamma\left(x+1,\theta_{1}\right)}{\Gamma\left(x+1\right)} + \sum_{y=x+1}^{\infty} \frac{\theta_{1}^{y}e^{-\theta_{1}}}{\Gamma\left(y+1\right)} \left[1 - \prod_{j=1}^{d} I_{p_{j}}\left(x+1,y-x\right)\right],$$

where $I_x\left(a,b\right) = \frac{\int_0^x t^{a-1}(1-t)^{b-1}dt}{\int_0^1 t^{a-1}(1-t)^{b-1}dt}$ is regularized incomplete beta function.

4 Asymptotic analysis

In this section, we conduct asymptotic analyses for the CDFs of the maximum and minimum of the aforementioned three multivariate Poisson distributions. Specifically, we work out two types of asymptotic results; one investigates the derived CDFs' asymptotic behaviors when the dimension $d \to \infty$, and the other examines scenarios where certain parameters (excluding d) dominate the dependence structure. We indeed obtain some interesting results that reveal the essential difference between the three dependence structures. Compared to the existing literature, our results offer an alternative approach to analyzing dependent structures from the perspective of the distribution of maximum (minimum) values and their asymptotic properties. For instance, the gap in the asymptotic results for dimension d between the common shock and comonotonic shock structures reveals a crucial difference in extending the common shock variable to a comonotonic shock vector, and we provide a counterexample to highlight this point. These findings supplement the work of Schulz et al. (2021) and further showcase the theoretical extension from common shock to comonotonic shocks.

We first define the asymptotic equivalence as follows.

Definition 4.1 Given functions f(x) and g(x), we define a binary relation

$$f(x) \sim g(x) \ (as \ x \to x_0, \ x_0 \in \mathbb{R}) \ if and only if \lim_{x \to x_0} \frac{f(x)}{g(x)} = 1.$$

Remark 4.1 For the sake of clarity, we further introduce following notations. For arbitrary random vector $\boldsymbol{\xi} = (\xi_1, \dots, \xi_d)$, denote

$$M_{-j}^{\xi} = \max(\xi_1, \dots, \xi_{j-1}, \xi_{j+1}, \dots, \xi_d), \quad m_{-j}^{\xi} = \min(\xi_1, \dots, \xi_{j-1}, \xi_{j+1}, \dots, \xi_d).$$

4.1 Common shock model

In this subsection, we will discuss the asymptotic analysis for the Poisson rates of Y_j , j = 1, ..., d, and Y_0 for the multivariate common shock Poisson distribution, as presented in Proposition 4.1 and Proposition 4.2, respectively. Moreover, we consider the asymptotic result for the dimension of X in Proposition 4.3.

Proposition 4.1 Let $X = (X_1, ..., X_d)$ follows the multivariate common shock Poisson distribution defined in Definition 2.1, then

$$F_{M}\mathbf{x}\left(x\right)\sim F_{X_{i}}\left(x\right)F_{M_{-i}^{\mathbf{Y}}}\left(x\right),\quad F_{m}\mathbf{x}\left(x\right)\sim F_{m_{-i}^{\mathbf{X}}}(x),\ as\ \theta_{i}\rightarrow\infty.$$

Proof Without loss of generality, we assume i=d, and the results for other cases can be similarly derived. Firstly, we consider the maximum distribution $F_{M^X}(x)$. From the Definition 2.1, the marginal distribution of X is

$$F_{X_i}(x) = \sum_{y=0}^{x} \frac{e^{-\theta_0} \theta_0^y \Gamma(x-y+1, \theta_i)}{\Gamma(y+1) \Gamma(x-y+1)}.$$

Therefore,

$$\lim_{\theta_{d}\rightarrow+\infty}\frac{F_{M^{X}}\left(x\right)}{F_{X_{d}}\left(x\right)}=\lim_{\theta_{d}\rightarrow+\infty}\frac{\sum_{y=0}^{x}\frac{\theta_{0}^{y}e^{-\theta_{0}}\prod_{j=1}^{d}\Gamma\left(x-y+1,\theta_{j}\right)}{\Gamma\left(y+1\right)\Gamma^{d}\left(x-y+1\right)}}{\sum_{y=0}^{x}\frac{e^{-\theta_{0}}\theta_{0}^{y}\Gamma\left(x-y+1,\theta_{d}\right)}{\Gamma\left(y+1\right)\Gamma\left(x-y+1\right)}}.$$

It is noteworthy that, for any given $z = 0, \dots, x$, we have,

$$\lim_{\theta_{d} \to +\infty} \frac{\frac{\theta_{0}^{z} e^{-\theta_{0}} \prod_{j=1}^{d} \Gamma(x-z+1,\theta_{j})}{\Gamma(z+1)\Gamma^{d}(x-z+1)}}{\sum_{y=0}^{x} \frac{e^{-\theta_{0}} \theta_{0}^{y} \Gamma(x-y+1,\theta_{d})}{\Gamma(y+1)\Gamma(x-y+1)}} = \frac{\theta_{0}^{z} e^{-\theta_{0}} \Gamma(x-y+1) \prod_{j=1}^{d-1} \Gamma(x-z+1,\theta_{j})}{\Gamma(z+1)\Gamma^{d}(x-z+1)}}{\sum_{y=0}^{x} \frac{\theta_{0}^{y} e^{-\theta_{0}}}{\Gamma(y+1)} \lim_{\theta_{d} \to +\infty} \frac{\Gamma(x-y+1,\theta_{d})}{\Gamma(x-z+1,\theta_{d})}},$$
(13)

and the limit part of the right side of (13) is

$$\lim_{\theta_d \to +\infty} \frac{\Gamma(x-y+1,\theta_d)}{\Gamma(x-z+1,\theta_d)} = \lim_{\theta_d \to \infty} \frac{\int_{\theta_d}^{+\infty} t^{x-y} e^{-t} dt}{\int_{\theta_d}^{+\infty} t^{x-z} e^{-t} dt} = \begin{cases} 0, & \text{if } z < y, \\ 1, & \text{if } z = y, \\ +\infty, & \text{if } z > y. \end{cases}$$

Hence,

$$\lim_{\theta_d \to +\infty} \frac{F_{M^{\mathbf{X}}}(x)}{F_{X_d}(x)} = \prod_{j=1}^{d-1} \frac{\Gamma(x+1,\theta_j)}{\Gamma(x+1)} = F_{m_{-d}^{\mathbf{X}}}(x).$$

Secondly, for $F_{mx}(x)$, we can similarly deduce that

$$\lim_{\theta_{d} \to \infty} \frac{F_{m} \mathbf{x} (x)}{F_{m} \mathbf{x}_{-d} (x)} = \lim_{\theta_{d} \to \infty} \frac{\sum_{y=0}^{x} \frac{\theta_{0}^{y} e^{-\theta_{0}} \left[\Gamma^{d} (x-y+1) - \prod_{j=1}^{d} \gamma (x-y+1,\theta_{j}) \right]}{\Gamma(y+1) \Gamma^{d} (x-y+1)}}{\sum_{y=0}^{x} \frac{\theta_{0}^{y} e^{-\theta_{0}} \left[\Gamma^{d-1} (x-y+1) - \prod_{j=1}^{d-1} \gamma (x-y+1,\theta_{j}) \right]}{\Gamma(y+1) \Gamma^{d-1} (x-y+1)}} = 1.$$

The following proposition analyzes the asymptotic equivalence of $F_{M}x(x)$ and $F_{m}x(x)$ as the rate of the common shock variable Y_0 approaches infinity.

Proposition 4.2 Let $X = (X_1, ..., X_d)$ follows the multivariate common shock Poisson distribution defined in Definition 2.1, then

$$F_{M\mathbf{x}}(x) \sim e^{-\sum_{j=1}^{d} \theta_{j}} F_{Y_{0}}(x), \quad F_{m\mathbf{x}}(x) \sim \left[1 - \prod_{j=1}^{d} \left(1 - e^{-\theta_{j}}\right)\right] F_{Y_{0}}(x), \text{ as } \theta_{0} \to \infty.$$

Proof For $F_{Mx}(x)$, we have

$$\lim_{\theta_{0}\to+\infty} \frac{F_{M}x(x)}{e^{-\sum_{j=1}^{d}\theta_{j}}F_{Y_{0}}(x)} = \lim_{\theta_{0}\to+\infty} \sum_{y=0}^{x} \frac{\theta_{0}^{y}e^{-\theta_{0}}\prod_{j=1}^{d}\Gamma(x-y+1,\theta_{j})}{\Gamma(y+1)\Gamma^{d}(x-y+1)} \frac{\Gamma(x+1)}{e^{-\sum_{j=1}^{d}\theta_{j}}\Gamma(x+1,\theta_{0})}$$

$$= \sum_{y=0}^{x} \lim_{\theta_{0}\to+\infty} \frac{\theta_{0}^{y}e^{-\theta_{0}}}{\Gamma(x+1,\theta_{0})} e^{\sum_{j=1}^{d}\theta_{j}} \frac{\Gamma(x+1)\prod_{j=1}^{d}\Gamma(x-y+1,\theta_{j})}{\Gamma(y+1)\Gamma^{d}(x-y+1)}.$$
(14)

The limit part of the right side of (14) is

$$\lim_{\theta_0 \to +\infty} \frac{\theta_0^y e^{-\theta_0}}{\Gamma(x+1, \theta_0)} = \lim_{\theta_0 \to +\infty} \frac{\theta_0^y e^{-\theta_0}}{\int_{\theta_0}^{+\infty} t^x e^{-t} dt} = \begin{cases} 0, & \text{if } y < x, \\ 1, & \text{if } y = x. \end{cases}$$
(15)

Therefore,

$$\lim_{\theta_0 \to +\infty} \frac{F_M(x)}{e^{-\sum_{j=1}^d \theta_j} F_{Y_0}(x)} = e^{\sum_{j=1}^d \theta_j} \frac{\Gamma(x+1) \prod_{j=1}^d \Gamma(1,\theta_j)}{\Gamma(x+1) \Gamma^d(1)} = 1.$$

The proof of $F_{mx}(x)$ can be obtained in a similar way.

Before presenting the asymptotic analysis results concerning the dimension of X, we first provide the following lemma, which can be directly derived from the properties of the Gamma function.

Lemma 4.1 Sequence $a_n = \frac{\Gamma(n)}{\Gamma(n,k)}$ is strictly monotonically decreasing.

Proposition 4.3 Let $X = (X_1, ..., X_d)$ follows the multivariate common shock Poisson distribution defined in Definition 2.1, then

$$F_{Mx}(x) \sim F_{Y_0}(0)F_{MY}(x)$$
, $F_{mx}(x) \sim F_{Y_0}(x)$, as $d \to \infty$.

Proof On one hand, for $F_{M}x(x)$, our objective is to prove

$$\lim_{d \to \infty} \frac{F_{M} \mathbf{x} (x)}{F_{Y_0}(0) F_{M} \mathbf{y} (x)} = \lim_{d \to \infty} \frac{F_{M} \mathbf{x} (x)}{e^{-\theta_0} F_{M} \mathbf{y} (x)} = 1.$$
 (16)

Notably, Y_j , j = 1, ..., d are mutually independent, thus,

$$F_{MY}(x) = \mathbb{P}\left(M^Y \le x\right) = \prod_{j=1}^d F_{Y_j}(x) = \frac{\prod_{j=1}^d \Gamma\left(x+1, \theta_j\right)}{\Gamma^d\left(x+1\right)}.$$

Therefore, we have,

$$\frac{F_{MX}\left(x\right)}{e^{-\theta_{0}}F_{MY}\left(x\right)} = 1 + \sum_{y=1}^{x} \frac{\theta_{0}^{y}}{\Gamma\left(y+1\right)} \prod_{j=1}^{d} \frac{\Gamma\left(x+1\right)\Gamma\left(x-y+1,\theta_{j}\right)}{\Gamma\left(x-y+1\right)\Gamma\left(x+1,\theta_{j}\right)}.$$

Moreover, from Lemma 4.1, for $y=1,\ldots,x$, we have, $\frac{\Gamma(x+1)\Gamma(x-y+1,\theta_j)}{\Gamma(x-y+1)\Gamma(x+1,\theta_j)}<1$, which implies that $\lim_{d\to\infty}\frac{F_{MX}(x)}{e^{-\theta_0}F_{MY}(x)}=1$.

The proof of $F_{mx}(x)$ follows a similar approach to that of $F_{Mx}(x)$.

4.2 Comonotonic shock model

In this subsection, we discuss the asymptotic analysis of the derived CDFs for the multivariate comonotonic shock Poisson distribution. We focus on the parameters λ_j , $j=1,\ldots,d$, as presented in Proposition 4.4, and the parameter θ in Propositions 4.5 and 4.6. Furthermore, we also consider the asymptotic result of the dimension of X in Proposition 4.7.

Proposition 4.4 Let $X = (X_1, ..., X_d)$ follows the multivariate comonotonic shock Poisson distribution defined in Definition 2.2, then

$$F_{M}\mathbf{x}\left(x\right) \sim F_{X_{i}}\left(x\right)F_{M}\mathbf{y}_{.}\left(x\right), \quad F_{m}\mathbf{x}\left(x\right) \sim F_{m}\mathbf{x}_{.}\left(x\right), \text{ as } \lambda_{i} \to \infty.$$

Proof We first consider the marginal distribution of X, it can be written as:

$$F_{X_i}(x) = \mathbb{P}\left(X_i \le x\right) = \sum_{z_i=0}^{x} g_{(1-\theta)\lambda_i}(x-z_i) G_{\theta\lambda_i}(z_i).$$

Notably, for any given z_1, \ldots, z_d and $z_i = 0, 1, \ldots, x_d$

$$\min \{G_{\theta \lambda_1}(z_1), \dots, G_{\theta \lambda_d}(z_d)\} \sim G_{\theta \lambda_i}(z_i), \quad \text{as } \lambda_i \to \infty,$$

hence, for the distribution of the maximum, $F_{Mx}(x)$, we have:

$$\lim_{\lambda_{i} \to \infty} \frac{F_{Mx}(x)}{F_{X_{i}}(x)} = \lim_{\lambda_{i} \to \infty} \frac{\sum_{z_{1}=0}^{x} \cdots \sum_{z_{d}=0}^{x} \min \{G_{\theta\lambda_{1}}(z_{1}), \dots, G_{\theta\lambda_{d}}(z_{d})\} \prod_{j=1}^{d} g_{(1-\theta)\lambda_{j}}(x-z_{j})}{\sum_{z_{i}=0}^{x} g_{(1-\theta)\lambda_{i}}(x-z_{i}) G_{\theta\lambda_{i}}(z_{i})}$$

$$= \lim_{\lambda_{i} \to \infty} \frac{\sum_{z_{1}=0}^{x} g_{(1-\theta)\lambda_{1}}(x-z_{1}) \cdots \sum_{z_{i}=0}^{x} g_{(1-\theta)\lambda_{i}}(x-z_{i}) G_{\theta\lambda_{i}}(z_{i}) \cdots \sum_{z_{d}=0}^{x} g_{(1-\theta)\lambda_{d}}(x-z_{d})}{\sum_{z_{i}=0}^{x} g_{(1-\theta)\lambda_{i}}(x-z_{i}) G_{\theta\lambda_{i}}(z_{i})}$$

$$= \prod_{j=1, j \neq i}^{d} G_{(1-\theta)\lambda_{j}}(x).$$

Similarly, the asymptotic analysis of F_{mx} can be proved using the same method.

Proposition 4.5 Let $X = (X_1, ..., X_d)$ follows the multivariate comonotonic shock Poisson distribution defined in Definition 2.2, then

$$F_{MX}(x) \sim F_{MY}(x)$$
, $F_{mX}(x) \sim F_{mY}(x)$, as $\theta \to 0$.

Proof Firstly, according to (i) in Definition 2.2, the CDFs of M^Y and m^Y are given by:

$$F_{MY}(x) = \prod_{j=1}^{d} G_{(1-\theta)\lambda_{j}}(x), \quad F_{mY}(x) = 1 - \prod_{j=1}^{d} \overline{G}_{(1-\theta)\lambda_{j}}(x).$$
 (17)

Meanwhile, we have:

$$\lim_{\theta \to 0} C_{\Lambda,\theta} (z_1, \dots, z_d) = \begin{cases} 1, & z_1 = \dots = z_d = 0, \\ 0, & \text{otherwise.} \end{cases}$$
 (18)

Thus, combining (18) with (5), (7) and (17), we have

$$\lim_{\theta \to 0} \frac{F_{MX}\left(x\right)}{F_{MY}\left(x\right)} = \lim_{\theta \to 0} \sum_{z_{1}=0}^{x} \cdots \sum_{z_{d}=0}^{x} \prod_{j=1}^{d} \frac{G_{\left(1-\theta\right)\lambda_{j}}\left(x-z_{j}\right)}{G_{\left(1-\theta\right)\lambda_{j}}\left(x\right)} C_{\Lambda,\theta}\left(z_{1},\ldots,z_{d}\right) = \prod_{j=1}^{d} \frac{G_{\left(1-\theta\right)\lambda_{j}}\left(x\right)}{G_{\left(1-\theta\right)\lambda_{j}}\left(x\right)} = 1,$$

and

$$\lim_{\theta \to 0} \frac{F_{m^{\mathbf{X}}}\left(x\right)}{F_{m^{\mathbf{Y}}}\left(x\right)} = \lim_{\theta \to 0} \frac{1 - \sum_{z_{1}=0}^{\infty} \cdots \sum_{z_{d}=0}^{\infty} \prod_{j=1}^{d} \overline{G}_{(1-\theta)\lambda_{j}}\left(x - z_{j}\right) C_{\Lambda,\theta}\left(z_{1}, \dots, z_{d}\right)}{1 - \prod_{j=1}^{d} \overline{G}_{(1-\theta)\lambda_{j}}\left(x\right)} = 1.$$

Proposition 4.6 Let $X = (X_1, ..., X_d)$ follows the multivariate comonotonic shock Poisson distribution defined in Definition 2.2, then

$$F_{MX}(x) \sim F_{MZ}(x)$$
, $F_{mX}(x) \sim F_{mZ}(x)$, as $\theta \to 1$.

Proof Firstly, according to (ii) in Definition 2.2, the CDFs of $M^{\mathbb{Z}}$ and $m^{\mathbb{Z}}$ are:

$$F_{Mz}(x) = \min \{G_{\theta\lambda_1}(x), \dots, G_{\theta\lambda_d}(x)\}, \quad F_{mz}(x) = \max \{G_{\theta\lambda_1}(x), \dots, G_{\theta\lambda_d}(x)\}.$$
 (19)

Meanwhile, we have

$$\lim_{\theta \to 1} \prod_{j=1}^{d} g_{(1-\theta)\lambda_j} \left(x - z_j \right) = \begin{cases} 1, & z_1 = \dots = z_d = x, \\ 0, & \text{otherwise.} \end{cases}$$
 (20)

Thus, combining (20) with (4) and (19), we have

$$\lim_{\theta \to 1} \frac{F_{Mx}(x)}{F_{Mz}(x)} = \lim_{\theta \to 1} \frac{\sum_{z_1=0}^{x} \cdots \sum_{z_d=0}^{x} \prod_{j=1}^{d} g_{(1-\theta)\lambda_j}(x-z_j) \min \{G_{\theta\lambda_1}(z_1), \dots, G_{\theta\lambda_d}(z_d)\}}{\min \{G_{\theta\lambda_1}(x), \dots, G_{\theta\lambda_d}(x)\}}$$

$$= \frac{\min \{G_{\lambda_1}(x), \dots, G_{\lambda_d}(z_d)\}}{\min \{G_{\lambda_1}(x), \dots, G_{\lambda_d}(z_d)\}} = 1.$$

and the proof of $F_{mx}(x)$ follows a similar approach to that of $F_{Mx}(x)$.

Next, we give the asymptotic analysis concerning the dimension d. We first introduce some notations. For a multivariate comonotonic shock Poisson distribution $X=(X_1,\ldots,X_d)$, we denote that $G_{x,d}:=\{G_{\theta\lambda_1}(x),\ldots,G_{\theta\lambda_d}(x)\}$ for $x=0,1,\ldots$ and $\Lambda=\lim_{d\to\infty}\Lambda_d$. Thus, we denote the bonded infinite set of real numbers $G^{(1)}:=\lim_{d\to\infty}G_{0,d}=\{G_{\theta\lambda_1}(0),G_{\theta\lambda_2}(0),\ldots\}$, its infimum is denoted as $m^{(1)}=\inf G^{(1)}$, and we denote $m^{(0)}:=0$. For $k=2,3,\ldots$, we denote that, $G^{(k)}:=G^{(k-1)}\setminus\{m^{(k-1)}\},m^{(k)}:=\inf G^{(k)}$. Similarly, we denote the bonded infinite set of real numbers $\tilde{G}_x^{(1)}:=\lim_{d\to\infty}G_{x,d}=\{G_{\theta\lambda_1}(x),G_{\theta\lambda_2}(x),\ldots\}$ and its supremum and infimum are denoted as $\tilde{M}_x^{(1)}=\sup \tilde{G}_x^{(1)}$ and $\tilde{m}_x^{(1)}=\inf \tilde{G}_x^{(1)}$. For $k=2,3,\ldots$, we denote $\tilde{G}_x^{(k)}=\tilde{G}_x^{(k-1)}\setminus\{\tilde{M}_x^{(k-1)}\},\tilde{M}_x^{(k)}=\sup G_x^{(k)}$.

Despite the structural similarities between comonotonic shock and common shock, the asymptotic analysis of d in comonotonic shock is intricate. Unlike the common shock with only one shock variable Y_0 , the number of shock variables in comonotonic shock tends to infinity as $d \to \infty$. This expansion poses significant challenges to the asymptotic analysis. To address this problem, we introduce the following assumptions.

Assumption 4.1 There exists $k \in \mathbb{N}_+$, such that $G^{(1)} \supseteq G^{(2)} \supseteq \cdots \supseteq G^{(k)} = G^{(k+1)}$.

Assumption 4.2 There exists $k \in \mathbb{N}_+$, such that $\tilde{G}^{(1)} \supseteq \tilde{G}^{(2)} \supseteq \cdots \supseteq \tilde{G}^{(k)} = \tilde{G}^{(k+1)}$.

Proposition 4.7 Let $X = (X_1, ..., X_d)$ follows the multivariate comonotonic shock Poisson distribution defined in Definition 2.2. Then if Assumption 4.1 holds, we have

$$F_{MX}(x) \sim cF_{MY}(x)$$
, as $d \to \infty$.

If Assumption 4.2 holds, we have

$$F_{m\mathbf{x}}(x) \sim \tilde{M}_{x}^{(k)} + \tilde{c}(x), \quad as \ d \to \infty,$$

where c and $\tilde{c}(x)$ are given in (23) and (24), respectively.

Proof Since in Theorem 3.2, the number of summation operations in (4)-(7) is the same as the dimension of the multivariate Poisson distribution, making it difficult to perform asymptotic analysis of d directly using them. To this end, we derive $F_{M} \mathbf{x}(x)$ and $F_{m} \mathbf{x}(x)$ by conditioning on the underlying uniform random variable U that generates the comonotonic shock random vector:

$$F_{MX}(x) = \int_{0}^{1} \prod_{j=1}^{d} G_{(1-\theta)\lambda_{j}}\left(x - G_{\theta\lambda_{j}}^{-1}(u)\right) du, \quad F_{mX}(x) = \int_{0}^{1} \left(1 - \prod_{j=1}^{d} \bar{G}_{(1-\theta)\lambda_{j}}\left(x - G_{\theta\lambda_{j}}^{-1}(u)\right)\right) du. \tag{21}$$

Notably, although the explicitness of (21) is lower than that of (4)-(7), the former are more suitable for the asymptotic analysis of d.

From Assumption 4.1, let us rearrange the elements in $G^{(1)}$, with the first k-1 elements from $G_{\theta\lambda_1}(0)=m^{(1)}$ to $G_{\theta\lambda_{k-1}}(0)=m^{(k-1)}$ in sequence. Notice that for $u\in(m^{(i)},m^{(i+1)}), i=0,1,\ldots,k-1$, we have, $G_{\theta\lambda_l}^{-1}(u)\geq 1, l=1,2\ldots,i$, and $G_{\theta\lambda_l}^{-1}(u)=0, l=i+1$. Hence, we have,

$$\lim_{d \to \infty} \frac{F_{MX}(x)}{F_{MY}(x)} = \int_0^1 \prod_{j=1}^{\infty} \frac{G_{(1-\theta)\lambda_j}\left(x - G_{\theta\lambda_j}^{-1}(u)\right)}{G_{(1-\theta)\lambda_j}\left(x\right)} du = c + \int_{m^{(k)}}^1 \prod_{j=1}^{\infty} \frac{G_{(1-\theta)\lambda_j}\left(x - G_{\theta\lambda_j}^{-1}(u)\right)}{G_{(1-\theta)\lambda_j}\left(x\right)} du, \quad (22)$$

where

$$c = \int_{m^{(0)}}^{m^{(1)}} 1 du + \int_{m^{(1)}}^{m^{(2)}} \frac{G_{(1-\theta)\lambda_1}(x - G_{\theta\lambda_1^{-1}}(u))}{G_{(1-\theta)\lambda_1}(x)} du + \dots + \int_{m^{(k-1)}}^{m^{(k)}} \prod_{j=1}^{k-1} \frac{G_{(1-\theta)\lambda_j}\left(x - G_{\theta\lambda_j}^{-1}(u)\right)}{G_{(1-\theta)\lambda_j}\left(x\right)} du$$
 (23)

is a constant.

By the supremum property, for $u\in(m^{(k)},1)$, there exists $u_1:=G_{\theta\lambda_{a_1}}(0)\in G^{(k)}\subset G^{(1)}$, such that $G_{\theta\lambda_{a_1}}(0)\in(m^{(k)},u)$. Similarly, there exists $u_2:=G_{\theta\lambda_{a_2}}(0)\in G^{(k)}\subset G^{(1)}$, such that $G_{\theta\lambda_{a_2}}(0)\in(m^{(k)},u_1)$.

Repeat this operation, we can find a sequence $\{a_n\}$ such that no two of $\{u_n\} = \{G_{\theta\lambda_{a_n}}\}$ are same and $G_{\theta\lambda_{a_n}}^{-1}(u) \ge 1$ for $n = 1, \ldots$ Thus, we have,

$$\int_{m^{(k)}}^{1} \prod_{j=1}^{\infty} \frac{G_{(1-\theta)\lambda_{j}}\left(x-G_{\theta\lambda_{j}}^{-1}(u)\right)}{G_{(1-\theta)\lambda_{j}}\left(x\right)} du = \int_{m^{(k)}}^{1} \prod_{n=1}^{\infty} \frac{G_{(1-\theta)\lambda_{a_{n}}}\left(x-G_{\theta\lambda_{a_{n}}}^{-1}(u)\right)}{G_{(1-\theta)\lambda_{a_{n}}}\left(x\right)} \prod_{\text{Others } \lambda_{j} \in \Lambda} \frac{G_{(1-\theta)\lambda_{j}}\left(x-G_{\theta\lambda_{j}}^{-1}(u)\right)}{G_{(1-\theta)\lambda_{j}}\left(x\right)} du = 0.$$

Similarly, for $F_{mx}(x)$, we have,

$$\begin{split} &\lim_{d \to \infty} \int_0^1 \left(1 - \prod_{j=1}^d \bar{G}_{(1-\theta)\lambda_j} \left(x - G_{\theta\lambda_j}^{-1}(u) \right) \right) du \\ &= \int_0^{\tilde{m}_x^{(1)}} 1 du + \int_{\tilde{m}_x^{(1)}}^{\tilde{M}_x^{(1)}} \left(1 - \prod_{j=1}^\infty \bar{G}_{(1-\theta)\lambda_j} \left(x - G_{\theta\lambda_j}^{-1}(u) \right) \right) du + \int_{\tilde{M}_x^{(1)}}^1 \left(1 - \prod_{j=1}^\infty 1 \right) du \\ &= \tilde{m}_x^{(1)} + \int_{\tilde{m}_x^{(1)}}^{\tilde{M}_x^{(1)}} \left(1 - \prod_{j=1}^\infty \bar{G}_{(1-\theta)\lambda_j} \left(x - G_{\theta\lambda_j}^{-1}(u) \right) \right) du. \end{split}$$

From Assumption 4.2, let us rearrange the elements in $\tilde{G}_x^{(1)}$, with the first k-1 elements from $\tilde{M}_x^{(1)} := G_{\theta \lambda_1}(x)$ to $\tilde{M}_x^{(k-1)} := G_{\theta \lambda_{k-1}}(x)$ in sequence. Similar to (22), we have,

$$\int_{\tilde{m}_{x}^{(1)}}^{\tilde{M}_{x}^{(1)}} \left(1 - \prod_{j=1}^{\infty} \bar{G}_{(1-\theta)\lambda_{j}} \left(x - G_{\theta\lambda_{j}}^{-1}(u) \right) \right) du = \int_{\tilde{m}_{x}^{(1)}}^{\tilde{M}_{x}^{(k)}} \left(1 - \prod_{j=1}^{\infty} \bar{G}_{(1-\theta)\lambda_{j}} \left(x - G_{\theta\lambda_{j}}^{-1}(u) \right) \right) du + \tilde{c}(x),$$

where $\tilde{c}(x) = 0$ for k = 1, and when $k = 1, 2, \dots, \tilde{c}(x)$ is given by

$$\tilde{c}(x) = \int_{\tilde{M}_{x}^{(k)}}^{\tilde{M}_{x}^{(k-1)}} \left(1 - \prod_{j=1}^{k-1} \bar{G}_{(1-\theta)\lambda_{j}} \left(x - G_{\theta\lambda_{j}}^{-1}(u) \right) \right) du + \dots + \int_{\tilde{M}_{x}^{(2)}}^{\tilde{M}_{x}^{(1)}} \left(1 - \bar{G}_{(1-\theta)\lambda_{1}} \left(x - G_{\theta\lambda_{1}}^{-1}(u) \right) \right) du$$
(24)

is a constant given x.

By the supremum property, we can find a sequence $\{b_n\}$ such that $G_{\theta\lambda_{b_n}}^{-1}(u) \leq x$ for $u \in (\tilde{m}_x^{(1)}, \tilde{M}_x^{(k)})$ and $n = 1, \ldots$. Thus, we have,

$$\begin{split} &\int_{\tilde{m}_x^{(1)}}^{\tilde{M}_x^{(k)}} \left(1 - \prod_{j=1}^{\infty} \bar{G}_{(1-\theta)\lambda_j} \left(x - G_{\theta\lambda_j}^{-1}(u)\right)\right) du \\ &= \int_{\tilde{m}_x^{(1)}}^{\tilde{M}_x^{(k)}} \left(1 - \prod_{n=1}^{\infty} \bar{G}_{(1-\theta)\lambda_{b_n}} \left(x - G_{\theta\lambda_{b_n}}^{-1}(u)\right) \prod_{\text{Others } \lambda_j \in \Lambda} \bar{G}_{(1-\theta)\lambda_j} \left(x - G_{\theta\lambda_j}^{-1}(u)\right)\right) du = \tilde{M}_x^{(k)} - \tilde{m}_x^{(1)}. \, \Box \end{split}$$

Remark 4.2 If k=1 satisfies Assumption 4.1, we observe $F_{M^X}(x) \sim m^{(1)} F_{M^Y}(x)$ as $d \to \infty$. Notably, $m^{(1)}$ represents the infimum of the CDF values of each comonotonic shock variable, Z_i , at x=0. This result is similar to that under the common shock model (Proposition 4.3). In both dependent structures, the CDFs of M^X are asymptotically equivalent to $F_Y(x)$ multiplied by a constant related to the value of the CDF of the "shock variables" at 0. Similarly, if k=1 satisfies Assumption 4.2, we have $F_{m^X}(x) \sim \tilde{M}_x^{(1)}$ as $d \to \infty$, also closely resembling the results under the common shock model.

Remark 4.3 If Assumption 4.1 or Assumption 4.2 is not valid, the asymptotic analysis of d may not be realistic for $F_{M} \times (x)$ or $F_{m} \times (x)$. Considering Assumption 4.1 as an instance, let us denote that

$$G_1^{(1)} := \{ \frac{1}{2} - \frac{1}{n}, n = 2, \dots \} \cup G^{(1)'}, \quad \inf G^{(1)'} = \frac{1}{2} \notin G^{(1)'},$$

$$G_2^{(1)} := \{ \frac{2}{3} + \frac{1}{12n}, n = 1, 2, \dots \}.$$

where $\sup G^{(1)'} \le 1$ and $\inf G^{(1)'} = \frac{1}{2} \notin G^{(1)'}$.

If $G^{(1)}=G_1^{(1)}$, we can divide the integration interval in (21) into two parts. The first part consists of countable non-empty intervals $(m^{(i)},m^{(i+1)}], i=0,1,\ldots$, and the second part is $(\frac{1}{2},1]$. Notably, the total interval length of the first part is $\sum_{i=0}^{\infty}m^{(i+1)}-m^{(i)}=\frac{1}{2}$. Thus, we have:

$$\lim_{d \to \infty} \frac{F_{Mx}(x)}{F_{My}(x)} = \int_{m^{(0)}}^{m^{(1)}} 1 du + \int_{m^{(1)}}^{m^{(2)}} \frac{G_{(1-\theta)\lambda_1}(x - G_{\theta\lambda_1^{-1}}(u))}{G_{(1-\theta)\lambda_1}(x)} du + \dots + \int_{\frac{1}{2}}^{1} \prod_{j=1}^{\infty} \frac{G_{(1-\theta)\lambda_j}\left(x - G_{\theta\lambda_j^{-1}}(u)\right)}{G_{(1-\theta)\lambda_j}(x)} du$$

$$= \int_{m^{(0)}}^{m^{(1)}} 1 du + \int_{m^{(1)}}^{m^{(2)}} \frac{G_{(1-\theta)\lambda_1}(x - G_{\theta\lambda_1^{-1}}(u))}{G_{(1-\theta)\lambda_1}(x)} du + \dots$$

$$< \int_{m^{(0)}}^{m^{(1)}} 1 du + \int_{m^{(0)}}^{m^{(1)}} 1 du + \dots + \int_{m^{(0)}}^{m^{(1)}} 1 du + \dots$$

$$< \frac{1}{2}.$$

However, if $G^{(1)} = G_2^{(1)}$, we have:

$$\lim_{d\to\infty}\frac{F_{M^{\mathbf{X}}}(x)}{F_{M^{\mathbf{Y}}(x)}} = \int_{m^{(0)}}^{m^{(1)}}1du + \int_{m^{(1)}}^{m^{(2)}}\frac{G_{(1-\theta)\lambda_1}(x-G_{\theta\lambda_1^{-1}}(u))}{G_{(1-\theta)\lambda_1}(x)}du + \dots > \int_{m^{(0)}}^{m^{(1)}}1du = \frac{2}{3}$$

The above two remarks illustrate the differences and connections between the structurally similar common shock and comonotonic shock structures, highlighting that while the extension of comonotonic shock is straightforward, it leads to significantly different probabilistic outcomes, particularly in the asymptotic analysis of d on M^X and m^X . And the difference becomes more evident when the dimension is high.

4.3 Thinning-dependence structure

In this subsection, we discuss the asymptotic analysis for parameters p_i^k , for i = 1, ..., d, k = 1, ..., l for the multivariate thinning-dependence Poisson distribution as presented in Proposition 4.8, and 4.9. Meanwhile, we also consider the asymptotic result on the dimension of X in Proposition 4.10.

Proposition 4.8 Let $X = (X_1, ..., X_d)$ follows the multivariate thinning-dependence Poisson distribution defined in Definition 2.3, then

$$F_{M\mathbf{x}}(x) \sim F_{X_i}(x), \quad F_{m\mathbf{x}}(x) \sim F_{m\mathbf{x}_i}(x), \text{ as } p_i^k \to 1, k = 1, \dots, l.$$

Proof For any $i = 1, \ldots, d$, $k = 1, \ldots, l$, and $x_i^k = 0, 1, \ldots, y_k$, we have,

$$\lim_{p_i^k \to 1} \mathbb{P}(X_i^k = x_i^k | Y_k = y_k) = \begin{cases} 1, & \text{if } x_i^k = y_k, \\ 0, & \text{if } x_i^k < y_k. \end{cases}$$
 (25)

Recall Theorem 3.3. For both $F_{Mx}(x)$ and $F_{mx}(x)$, p_i^k , $k=1,\ldots,l$ appear only in $\sum_{s=0}^x h_j(s,y_1,\ldots,y_l)$. This quantity can be expressed as:

$$\sum_{s=0}^{x} h_{j}(s, y_{1}, \dots, y_{l}) = \mathbb{P}\left(\sum_{k=1}^{l} X_{i}^{k} \leq x | Y_{1} = y_{1}, \dots, Y_{l} = y_{l}\right)$$

$$= \sum_{x_{i}^{1} + \dots + x_{i}^{l} \leq x} \mathbb{P}\left(X_{i}^{1} = x_{i}^{1}, \dots, X_{i}^{l} = x_{i}^{l} | Y_{1} = y_{1}, \dots, Y_{l} = y_{l}\right)$$

$$= \sum_{x_{i}^{1} + \dots + x_{i}^{l} \leq x} \mathbb{P}\left(X_{i}^{1} = x_{i}^{1} | Y_{1} = y_{1}\right) \dots \mathbb{P}\left(X_{i}^{l} = x_{i}^{l} | Y_{l} = y_{l}\right),$$
(26)

where the last equality is from Lemma 3.2. Thus, from (25), as $p_i^k \to 1, k = 1, ..., l$, (26) can be rewritten as follow:

$$\lim_{p_i^k \to 1, k=1, \dots, l} \sum_{x_i^1 + \dots + x_i^l \le x} \mathbb{P}\left(X_i^1 = x_i^1 | Y_1 = y_1\right) \dots \mathbb{P}\left(X_i^l = x_i^l | Y_l = y_l\right) = \begin{cases} 1, & \text{if } x_i^1 = y_1, \dots, x_i^l = y_l, \\ 0, & \text{otherwise.} \end{cases}$$

Therefore, we have,

$$\lim_{p_i^k \to 1, k=1, \dots, l} \frac{F_{MX}(x)}{F_{X_i}(x)} = \frac{\sum_{y_1 + \dots + y_l \le x} \prod_{k=1}^l \frac{\theta_k^{y_k} e^{-\theta_k}}{\Gamma(y_k + 1)}}{\sum_{y_1 + \dots + y_l \le x} \prod_{k=1}^l \frac{\theta_k^{y_k} e^{-\theta_k}}{\Gamma(y_k + 1)}} = 1.$$

The asymptotic result of $F_{mx}\left(x\right)$ can be obtained in a similar manner

Proposition 4.9 Let $X = (X_1, ..., X_d)$ follows the multivariate thinning-dependence Poisson distribution defined in Definition 2.3, then

$$F_{M\mathbf{x}}(x) \sim F_{M\mathbf{x}_i}(x)$$
, $F_{m\mathbf{x}}(x) \sim 1$, as $p_i^k \to 0$, $k = 1, \dots, l$.

Proof For any $i=1,\ldots,d,\,k=1,\ldots,l,$ and $x_i^k=0,1,\ldots,y_k,$ we have,

$$\lim_{p_i^k \to 0} \mathbb{P}(X_i^k = x_i^k | Y_k = y_k) = \begin{cases} 1, & \text{if } x_i^k = 0, \\ 0, & \text{if } x_i^k = 1, \dots, y_k. \end{cases}$$

The remaining proof is in the same line with Proposition 4.8.

From Definition 2.3, we can directly deduce that $M^X \leq Y_1 + \cdots + Y_l$, and the lower bound of $m^X \geq 0$; see also Remark 2.2. When there are more marginal risks, i.e. the dimension d is high, more trials are involved in the system, and it is intuitive to expect the upper/lower bound is reached with a higher chance. Consequently, we prove the following proposition of the asymptotic results for M^X and m^X when $d \to \infty$, which is indeed consistent with intuitions.

Proposition 4.10 Let $X = (X_1, ..., X_d)$ follows the multivariate thinning-dependence Poisson distribution defined in Definition 2.3, then

$$F_{M}\mathbf{x}(x) \sim F_{Y_1} * F_{Y_2} * \cdots * F_{Y_l}(x), \quad F_{m}\mathbf{x}(x) \sim 1, \text{ as } d \to \infty.$$

where * denotes the convolution operator.

Proof Notably, for $j=1,\ldots,d$, $\mathbb{P}\left(X_j\leq x|Y_1=y_1,\ldots,Y_l=y_l\right)$ and $1-\mathbb{P}\left(X_j\leq x|Y_1=y_1,\ldots,Y_l=y_l\right)$ are strictly positive and less than 1 when $y_1+\cdots+y_l\geq x+1$. Thus, in this scenario,

$$\lim_{d \to \infty} \prod_{j=1}^{d} \mathbb{P}(X_j \le x | Y_1 = y_1, \dots, Y_l = y_l) = 0, \quad \lim_{d \to \infty} \prod_{j=1}^{d} 1 - \mathbb{P}(X_j \le x | Y_1 = y_1, \dots, Y_l = y_l) = 0.$$

Therefore, when $d \to \infty$, (11) and (12) can be written as:

$$\lim_{d \to \infty} F_{M} x (x) = \sum_{y_1 + \dots + y_l \le x} \prod_{k=1}^l \frac{\theta_k^{y_k} e^{-\theta_k}}{\Gamma(y_k + 1)} = F_{Y_1 + \dots + Y_l}(x),$$

and,

$$\lim_{d\to\infty}F_{m}\mathbf{x}\left(x\right)=\sum_{y_{1}+\cdots+y_{l}\leq x}\prod_{k=1}^{l}\mathbb{P}\left(Y_{k}=y_{k}\right)+\sum_{y_{1}+\cdots+y_{l}\geq x+1}\prod_{k=1}^{l}\mathbb{P}\left(Y_{k}=y_{k}\right)=\prod_{k=1}^{l}\sum_{y_{k}=0}^{\infty}\mathbb{P}\left(Y_{k}=y_{k}\right)=1.\square$$

Remark 4.4 The asymptotic results for d under the thinning-dependence structure is significantly different, compared to those under the common shock and comonotonic shock structures. Specifically, as $d \to \infty$, M^X 's behavior is determined by the common background factors under the thinning-dependence structure, whereas under the common shock and comonotonic shock structures, M^X behaves consistently with the idiosyncratic factors.

Combining Remarks 4.2 to 4.4, we find that the differences in the asymptotic analysis of d under the three structures correspond precisely to the intrinsic differences in how these structures are constructed, particularly in the way the common factor is introduced. This not only sheds new light on analyzing dependent structures but also plausibly suggests that similar results may be obtained when utilizing these structures to construct other distributions.

5 Numerical results

In this section, we present numerical results of the CDFs of the multivariate Poisson distribution and its asymptotic analysis under the common shock, comonotonic shock, and thinning-dependence structures. Due to space limitation, the numerical examples focus solely on the distribution of M^X .

5.1 Numerical result of CDFs

To compare the effects of different dependent structures on the maximum (minimum) of multivariate Poisson distributions, we ensure that the marginal distributions of the three multivariate Poisson distributions are identical in our numerical example, and that the correlations within each distribution are also consistent. Intuitively, the most direct way to ensure the same correlation is to equalize the correlation coefficients of the three multivariate Poisson distributions. However, this is often difficult to achieve while maintaining equal marginal distributions. Therefore, we use the average correlation coefficient to characterize the internal correlation within each multivariate distribution in this paper. To illustrate the results, we take the 3-dimensional case as an example. Specifically, for a 3-dimensional random vector $\mathbf{X} = (X_1, X_2, X_3)$, the average correlation coefficient $\bar{\rho}$ is given by

$$\bar{\rho} = \frac{\text{Var}(X_1 + X_2 + X_3) - \text{Var}(X_1) - \text{Var}(X_2) - \text{Var}(X_3)}{2\left(\sqrt{\text{Var}(X_1)\text{Var}(X_2)} + \sqrt{\text{Var}(X_2)\text{Var}(X_3)} + \sqrt{\text{Var}(X_1)\text{Var}(X_3)}\right)}.$$

Hence, the average correlation coefficient of the multivariate common shock, comonotonic shock, thinning-dependence Poisson distribution, denoted as ρ_1 , ρ_2 and ρ_3 , are given by

$$\bar{\rho}_{1} = \frac{3\theta_{0}}{\sqrt{(\theta_{0} + \theta_{1})(\theta_{0} + \theta_{2})} + \sqrt{(\theta_{0} + \theta_{1})(\theta_{0} + \theta_{3})} + \sqrt{(\theta_{0} + \theta_{2})(\theta_{0} + \theta_{3})}},$$

$$\bar{\rho}_{2} = \frac{m_{\lambda_{1},\lambda_{2}}(\theta) + m_{\lambda_{1},\lambda_{3}}(\theta) + m_{\lambda_{2},\lambda_{3}}(\theta)}{\sqrt{\lambda_{1}\lambda_{2}} + \sqrt{\lambda_{1}\lambda_{3}} + \sqrt{\lambda_{2}\lambda_{3}}},$$

$$\bar{\rho}_{3} = \frac{\sum_{k=1}^{l} p_{1}^{k} p_{2}^{k} + \sum_{k=1}^{l} p_{1}^{k} p_{3}^{k} + \sum_{k=1}^{l} p_{2}^{k} p_{3}^{k}}{\sqrt{\sum_{k=1}^{l} p_{1}^{k} \sum_{k=1}^{l} p_{2}^{k}} + \sqrt{\sum_{k=1}^{l} p_{1}^{k} \sum_{k=1}^{l} p_{3}^{k}} + \sqrt{\sum_{k=1}^{l} p_{2}^{k} \sum_{k=1}^{l} p_{3}^{k}}},$$

$$(27)$$

where
$$m_{\lambda_{j},\lambda_{k}}\left(\theta\right)=\operatorname{cov}\left(X_{j},X_{k}\right)=\sum_{m=0}^{\infty}\sum_{n=0}^{\infty}\min\left\{\overline{G}_{\theta\lambda_{j}}\left(m\right),\overline{G}_{\theta\lambda_{k}}\left(m\right)\right\}-\theta^{2}\lambda_{j}\lambda_{k}.$$

In Table 1, we compare the CDFs of M^X for three types of multivariate Poisson. In the case of l=1, for the multivariate common shock Poisson distribution, we set $\theta_0=3.3804, \theta_1=2.6196, \theta_2=3.6196, \theta_3=4.6196$. For the multivariate common shock Poisson distribution, we set $\theta_0=3.3804, \theta_1=2.6196, \theta_2=3.6196, \theta_3=4.6196$; for the multivariate comonotonic shock Poisson distribution, we set $\lambda_1=6, \lambda_2=7, \lambda_3=8, \theta=0.5$; for the multivariate thinning-dependence Poisson distribution, we set $l=1, \theta_1=15.3829, p_1=0.39, p_2=0.4551, p_3=0.5201$. This setting ensures equal marginal distributions and similar average correlation coefficients, with $\bar{\rho}_1\approx\bar{\rho}_2\approx\bar{\rho}_3\approx0.48$. In the case of l=2, for the multivariate common shock Poisson distribution, we set $\theta_0=8.2, \theta_1=\theta_2=0.8, \theta_3=1.8$; for the multivariate comonotonic shock Poisson distribution, we set $\lambda_1=\lambda_2=9, \lambda_3=10, \theta=0.5$; For the multivariate thinning-dependence Poisson distribution, we set $\theta_1=8.2, \theta_2=19.69, p_1^1=p_2^1=p_3^1=1, p_1^2=p_2^2=0.0406, p_3^2=0.0914$. This setup also ensures equal marginal distributions and similar average correlation coefficients, with $\bar{\rho}_1\approx\bar{\rho}_2\approx\bar{\rho}_3\approx0.9$.

According to (2), when l=1, each X_i in the thinning-dependence structure is derived from only one X_i^1 , meaning that all X_i are only dependent on the same Y_1 . This points out a significant difference between the thinning-dependence structure and the other two structures. However, when l=2, not only all X_i have two cumulative sources, but also, if $p_i^1=1$ simultaneously, the first cumulative source X_i^1 of X_i becomes a common Y_1 . In this case, the thinning-dependence structure closely resembles the other two structures. The numerical results in Table 1 also confirm this point. When l=1, the CDF of the thinning-dependence structure is greater than that of the common shock and comonotonic shock structures, indicating that the M^X in the former structure, which adds the common factors indirectly, is indeed larger than in the common shock and comonotonic shock structures, which directly add common factors. Moreover, the gap in CDF values between the thinning-dependence

Table 1: Numerical results of the multivariate Poisson distribution.

x	l=2			l=1			
	Common	Comonotonic	Thinning	Common	Comonotonic	Thinning	
1	1.58×10^{-5}	1.50×10^{-5}	1.77×10^{-5}	6.70×10^{-7}	8.89×10^{-7}	7.17×10^{-5}	
2	1.17×10^{-3}	1.17×10^{-3}	1.27×10^{-3}	3.01×10^{-5}	3.85×10^{-5}	7.71×10^{-4}	
3	5.37×10^{-3}	5.48×10^{-3}	5.68×10^{-3}	4.57×10^{-4}	5.65×10^{-4}	4.41×10^{-3}	
4	1.77×10^{-2}	1.83×10^{-2}	1.85×10^{-2}	3.42×10^{-3}	4.08×10^{-3}	1.66×10^{-2}	
5	4.57×10^{-2}	4.72×10^{-2}	4.72×10^{-2}	1.55×10^{-2}	1.78×10^{-2}	4.66×10^{-2}	
6	0.0973	0.1004	0.0997	0.0488	0.0540	0.1039	
7	0.1771	0.1821	0.1802	0.1162	0.1247	0.1937	
8	0.2833	0.2903	0.2870	0.2229	0.2336	0.3123	
9	0.4076	0.4158	0.4114	0.3610	0.3718	0.4480	
10	0.5371	0.5456	0.5406	0.5119	0.5208	0.5846	
11	0.6587	0.6662	0.6616	0.6542	0.6604	0.7073	
12	0.7626	0.7683	0.7649	0.7724	0.7762	0.8069	
13	0.8442	0.8481	0.8458	0.8605	0.8625	0.8805	
14	0.9034	0.9058	0.9045	0.9200	0.9209	0.9305	
15	0.9433	0.9447	0.9440	0.9569	0.9573	0.9620	

structure and the common (comonotonic) shock structures is significantly larger when l=2, especially for smaller values of x. These findings highlight the significant impact of different dependence structures on the maximum of the multivariate Poisson distribution.

5.2 Numerical result of asymptotic analysis

In this subsection, we first provide numerical examples for the dimension d: For the multivariate common shock Poisson distribution, we set $\theta_0=5, \theta_1=\cdots=\theta_d=5$; For the multivariate comonotonic shock Poisson distribution, we set $\lambda_j=8-2/j, j=1,2,\ldots,d, \theta=0.5$; For the multivariate thinning-dependence Poisson distribution, we set $l=1, \theta_1=8, p_j=0.7, j=1,2,\ldots,d$. The asymptotic analysis results are presented in Table 2. Secondly, we provide numerical examples for the asymptotic analysis of other parameters. For the multivariate common shock Poisson distribution, in Proposition 4.1, we set $d=10, \theta_j=6$ for $j=1,\ldots,10$ and $j\neq i$; in Proposition 4.2, we set $d=2, \theta_1=\theta_2=2$. For the multivariate comonotonic shock Poisson distribution, in Proposition 4.4, we set $d=3, \theta=1/2, \lambda_1=\lambda_2=20$; in Proposition 4.5, we set $d=3, \lambda_1=\lambda_2=\lambda_3=10$; in Proposition 4.6, we set $d=3, \lambda_1=\lambda_2=\lambda_3=5$. For the multivariate thinning-dependence Poisson distribution, in Proposition 4.8, we set $d=1, d=10, \theta_1=10, \theta_1=10,$

6 Discussion and conclusion

In this paper, we comprehensively investigate the distribution of the maximum and minimum of multivariate Poisson distributions, which is the cornerstone of the discrete distributions. To the best of our knowledge, there is very

Table 2: Numerical results for the asymptotic analysis of dimension d, illustrating the ratio of the asymptotic results (AR) to its corresponding maximum distribution (FM) when x=7.

Dependent Structure	Proposition	d = 10	d = 20	d = 50	d = 70	d = 200
Common	Proposition 4.3	0.35464	0.71591	0.99193	0.99938	1
Comonotonic	Proposition 4.7	0.15952	0.35401	0.80749	0.92822	0.99389
Thinning	Proposition 4.10	0.83133	0.91117	0.98441	0.99520	1

Table 3: Numerical results for the asymptotic analysis of other parameters are presented, illustrating the ratio between the FM and its AR when x = 7.

	Proposition 4.1	θ_i	10	100	200	1000	5000
Common		FM/AR	0.08830	0.71232	0.84105	0.96542	0.99296
Common	Proposition 4.2	θ_0	10	50	500	5000	8000
		AR/FM	0.07647	0.42064	0.89728	0.98890	0.99304
	Proposition 4.4	λ_i	10	20	40	50	100
		FM/AR	0.00229	0.06308	0.91665	1	1
Comonotonic	Proposition 4.5	θ	0.66	0.46	0.31	0.11	0.01
Comonotonic		FM/AR	0.08487	0.09044	0.14666	0.46898	0.99210
	Proposition 4.6	θ	0.0005	0.3005	0.5005	0.9005	0.99999
		FM/AR	0.65094	0.69323	0.72714	0.88655	0.99998
	Proposition 4.8	p_i^k	0.009	0.309	0.509	0.709	0.959
Thinning		FM/AR	0.56523	0.57256	0.64028	0.80086	0.99029
Timiling	Proposition 4.9	p_i^k	0.951	0.751	0.551	0.251	0.001
		FM/AR	0.46755	0.77821	0.95398	0.99972	1

limited literature on the maximum distribution for discrete random vectors. Our work not only derives explicit maximum and minimum distribution functions for three types of multivariate Poisson distributions but also conducts asymptotic analyses for all derived CDFs. These results address a research gap in the literature, providing valuable insights into relevant studies on this topic. In particular, our asymptotic analysis may indicate a new way to study the dependence structures for other multivariate distributions.

As a matter of fact, the Poisson distribution, being a classical discrete distribution, has a wide range of applications, such as in credit risk (Liang and Wang, 2012), reinsurance (Yuen et al., 2015), and price modeling (Holỳ and Tomanová, 2022). In future research, we also anticipate utilizing our findings on the distribution of the maximum and minimum of multivariate Poisson distributions to model such practical problems, providing new perspectives and solutions to these important areas.

Funding

This work is supported by the National Natural Science Foundation of China [grant number 12371474]; the Postgraduate Research & Practice Innovation Program of Jiangsu Province [grant number KYCX24_3353].

References

- Reinaldo B Arellano-Valle and Marc G Genton. On the exact distribution of the maximum of absolutely continuous dependent random variables. *Statistics & Probability Letters*, 78(1):27–35, 2008.
- Norman TJ Bailey. Statistical methods in biology. Cambridge university press, 1995.
- AP Basu and JK Ghosh. Identifiability of the multinormal and other distributions under competing risks model. *Journal of Multivariate Analysis*, 8(3):413–429, 1978.
- Michael Cain. The moment-generating function of the minimum of bivariate normal random variables. *The American Statistician*, 48(2):124–125, 1994.
- Bin Cheng, Shein-Chung Chow, David Burt, and Dennis Cosmatos. Statistical assessment of qt/qtc prolongation based on maximum of correlated normal random variables. *Journal of Biopharmaceutical Statistics*, 18(3): 494–501, 2008.
- DK Chua and Yang Miang Goh. Poisson model of construction incident occurrence. *Journal of Construction Engineering and Management*, 131(6):715–722, 2005.
- Rob Eisinga, Manfred Te Grotenhuis, and Ben Pelzer. Saddlepoint approximations for the sum of independent non-identically distributed binomial random variables. *Statistica Neerlandica*, 67(2):190–201, 2013.
- Paul Embrechts, Claudia Klüppelberg, and Thomas Mikosch. *Modelling extremal events: for insurance and finance*, volume 33. Springer Science & Business Media, 2013.
- RC Griffiths, RK Milne, and R Wood. Aspects of correlation in bivariate poisson distributions and processes. *Australian Journal of Statistics*, 21(3):238–255, 1979.
- Nooshin Hakamipour, Adel Mohammadpour, and Saralees Nadarajah. Extremes of a bivariate pareto distribution. *Information Technology and Control*, 40(1):83–87, 2011.
- Vladimír Holỳ and Petra Tomanová. Modeling price clustering in high-frequency prices. *Quantitative Finance*, 22 (9):1649–1663, 2022.
- David I Inouye, Eunho Yang, Genevera I Allen, and Pradeep Ravikumar. A review of multivariate distributions for count data derived from the poisson distribution. *Wiley Interdisciplinary Reviews: Computational Statistics*, 9 (3):e1398, 2017.
- Ahad Jamalizadeh and N Balakrishnan. Distributions of order statistics and linear combinations of order statistics from an elliptical distribution as mixtures of unified skew-elliptical distributions. *Journal of Multivariate Analysis*, 101(6):1412–1427, 2010.
- Dimitris Karlis. An em algorithm for multivariate poisson distribution and related models. *Journal of Applied Statistics*, 30(1):63–77, 2003.
- Dimitris Karlis and Loukia Meligkotsidou. Finite mixtures of multivariate poisson distributions with application. *Journal of statistical Planning and Inference*, 137(6):1942–1960, 2007.
- Kazutomo Kawamura. The structure of trivariate poisson distribution. In *Kodai Mathematical Seminar Reports*, volume 28, pages 1–8. Department of Mathematics, Tokyo Institute of Technology, 1976.
- Alan P Ker. On the maximum of bivariate normal random variables. Extremes, 4:185–190, 2001.
- Xue Liang and Guojing Wang. On a reduced form credit risk model with common shock and regime switching. *Insurance: Mathematics and Economics*, 51(3):567–575, 2012.

- Da-Hsiang Donald Lien. Moments of ordered bivariate log-normal distributions. *Economics Letters*, 20(1):45–47, 1986.
- Donald Lien. On the minimum and maximum of bivariate lognormal random variables. Extremes, 8:79–83, 2005.
- Filip Lindskog and Alexander J McNeil. Common poisson shock models: applications to insurance and credit risk modelling. *ASTIN Bulletin: The Journal of the IAA*, 33(2):209–238, 2003.
- DM Mahamunulu. A note on regression in the multivariate poisson distribution. *Journal of the American Statistical Association*, 62(317):251–258, 1967.
- Saralees Nadarajah and Jungsoo Woo. On the maximum and minimum of multivariate pareto random variables. *Information Technology and Control*, 42(3):242–246, 2013.
- Saralees Nadarajah, Emmanuel Afuecheta, and Stephen Chan. On the distribution of maximum of multivariate normal random vectors. *Communications in Statistics-Theory and Methods*, 48(10):2425–2445, 2019.
- JD Salas, J Obeysekera, and RM Vogel. Techniques for assessing water infrastructure for nonstationary extreme events: a review. *Hydrological Sciences Journal*, 63(3):325–352, 2018.
- Juliana Schulz. *Multivariate Poisson models based on comonotonic and counter-monotonic shocks*. McGill University (Canada), 2019.
- Juliana Schulz, Christian Genest, and Mhamed Mesfioui. A multivariate poisson model based on comonotonic shocks. *International Statistical Review*, 89(2):323–348, 2021.
- Guojing Wang and Kam C Yuen. On a correlated aggregate claims model with thinning-dependence structure. *Insurance: Mathematics and Economics*, 36(3):456–468, 2005.
- Kam Chuen Yuen, Zhibin Liang, and Ming Zhou. Optimal proportional reinsurance with common shock dependence. *Insurance: Mathematics and Economics*, 64:1–13, 2015.