A Model-Based Clustering Approach for Bounded Data Using Transformation-Based Gaussian Mixture Models

Luca Scrucca 🗅

Department of Statistical Sciences, University of Bologna

December 19, 2024

Abstract

The clustering of bounded data presents unique challenges in statistical analysis due to the constraints imposed on the data values. This paper introduces a novel method for model-based clustering specifically designed for bounded data. Building on the transformation-based approach to Gaussian mixture density estimation introduced by Scrucca (2019), we extend this framework to develop a probabilistic clustering algorithm for data with bounded support that allows for accurate clustering while respecting the natural bounds of the variables. In our proposal, a flexible range-power transformation is employed to map the data from its bounded domain to the unrestricted real space, hence enabling the estimation of Gaussian mixture models in the transformed space. This approach leads to improved cluster recovery and interpretation, especially for complex distributions within bounded domains. The performance of the proposed method is evaluated through real-world data applications involving both fully and partially bounded data, in both univariate and multivariate settings. The results demonstrate the effectiveness and advantages of our approach over traditional and advanced model-based clustering techniques that employ distributions with bounded support.

Keywords: model-based clustering, Bounded data, Gaussian mixture models, Data transformation, Expectation-Maximization algorithm.

Contents

1	Introduction	3
2	Methods	4
	2.1 Finite mixture modeling for clustering	4
	2.2 A transformation-based approach for Gaussian mixtures clustering	5
	2.3 Maximum likelihood estimation via the EM algorithm	7
	2.4 Clustering and classification uncertainty	8
3	Applications	10
	3.1 Enzyme data	10
	3.2 Wholesale customer segmentation	11
	3.3 Human Development Index	16
4	Conclusion	18

1 Introduction

Clustering is a fundamental task in data analysis, aiming to identify natural groupings or patterns within a dataset and thus facilitate the discovery of underlying structures and relationships among data points. While numerous model-based clustering methods exist for unconstrained data, either having symmetric or asymmetric distributions, the clustering of bounded data presents unique challenges that require specialized approaches. Data with bounded support frequently arise in various fields, including economics, biology, environmental science, and social sciences. Examples include percentages, proportions, non-negative variables, e.g. arising from physical measurements, and any metric naturally limited to a fixed range. Traditional clustering methods often struggle when applied to bounded data due to their inability to adequately account for the constraints imposed by the data bounds, leading to inaccurate inference and suboptimal clustering results.

Model-based clustering, which assumes that data are generated from a mixture of probability distributions, provides a principled and flexible framework for clustering. Gaussian mixture models (GMMs) assume the data are generated from a mixture of Gaussian distributions, with each component representing a distinct cluster. They are widely used due to their flexibility to model complex data structures and their adaptability to a variety of clustering problems.

However, the direct application of GMMs to bounded data is problematic due to the inherent constraints on data values. In this context, alternative model-based approaches that account for natural bounds in the data can be adopted by assuming bounded distributions for the component densities.

For instance, in the case of positive, right-skewed data, a mixture of gamma distributions (John, 1970) provides a natural alternative. Bagnato and Punzo (2013) and Young et al. (2019) proposed mixture of univariate unimodal gamma densities, while Wiper et al. (2001) discussed a similar model in a Bayesian framework, although also restricted to univariate data. Another potential family of distributions is introduced in Karlis and Santourian (2009), who employed a mixture of normal inverse-Gaussian distributions. This model extends naturally to the multivariate case, allowing for skewness, fat tails and, as by-product, can handle variables bounded at zero.

For data bounded at both the lower and upper limits, Bagnato and Punzo (2013) proposed the mixture of univariate unimodal beta densities. Building on this, Dean and Nugent (2013) developed a model-based clustering procedure for data confined within the unit hypercube using a mixture of univariate unimodal beta distributions. However, their approach in the multivariate case is constrained by the assumption of conditional independence, whereby variables are considered conditionally independent given component membership.

A different approach was introduced in speech processing, where a bounded Gaussian mixture model (BGMM; Hedelin and Skoglund, 2000) was employed and later extended to a bounded generalized Gaussian mixture model (BGGMM; Lindblom and Samuelsson, 2003). The BGGMM encompasses several models, including the standard GMM and BGMM, as special cases. These methods rely on truncated GMMs, where the unbounded Gaussian densities are multiplied by an indicator function that equal 1 if the component densities lie within the bounded data region and 0 otherwise; then, a subsequent normalization step ensures a proper marginal density distribution.

A common alternative to modeling bounded distributions directly is the use of data transformation techniques. Transforming bounded data onto an unbounded scale enables the application of standard statistical methods, then followed by an inverse transformation to express the results obtained in the original data space. For instance, transformations such as the Box-Cox (Lo and Gottardo, 2012) and Manly (Zhu and Melnykov, 2018) transformations have been employed to manage skewness in component distributions.

More recently, Gallaugher et al. (2020) compared the performance of Gaussian mixtures in handling skewed data or outliers with that of mixtures of skewed distributions, such as the variancegamma distribution (McNicholas et al., 2017, VG;) and the generalized hyperbolic distribution (Browne and McNicholas, 2015, GH). They also examined mixtures incorporating transformations aimed at achieving near-normality, including the power transformation (Yeo and Johnson, 2000) and the Manly transformation (Manly, 1976). Their results indicate that no single method consistently outperforms the others, with the optimal choice depending on the specific characteristics and context of the data being analyzed.

In this paper, we propose a novel model-based clustering method specifically designed for bounded data. Our approach builds upon the range-power transformation framework for Gaussian mixtures proposed by Scrucca (2019). The method involves mapping bounded data into an unbounded space, where standard Gaussian mixture models can be applied, followed by an inverse transformation to recover the clustering results in the original bounded space. This approach has proven effective in real data application, offering a general and flexible framework for clustering bounded data.

In Section 2, we first review the model-based approach to clustering, with particular emphasis on the Gaussian mixture model (GMM). The proposed approach is then presented, introducing the range-power transformation and its integration within the finite mixture model framework. We discuss maximum likelihood estimation via the EM algorithm, as well as model selection and methods for assessing clustering and classification uncertainty in this specific context. Section 3 applies the method to real-data examples, covering both fully and partially bounded data in univariate and multivariate contexts. We compare our method against both the standard GMM and more advanced model-based clustering techniques that incorporate distributions with bounded support specific to the analyzed features. Finally, Section 4 summarizes the main contributions of this work, discusses its strengths and limitations, and outlines potential directions for future research.

2 Methods

2.1 Finite mixture modeling for clustering

Consider a multivariate dataset $\mathcal{D} = \{x_i\}_{i=1}^n$ of *n* observations, where each observation x_i is drawn from a *d*-dimensional random vector x with unbounded support $\mathcal{S}_{\mathcal{X}} \equiv \mathbb{R}^d$.

In model-based clustering we aim to partition the observations into G distinct groups or clusters by employing a finite mixture model (FMM; McLachlan and Peel, 2000; McLachlan et al., 2019). Within this framework, each mixture component is directly associated with a cluster, effectively representing a distinct grouping of the data. In its general form, a FMM can be expressed as:

$$f(\boldsymbol{x}; \boldsymbol{\Psi}) = \sum_{k=1}^{G} \pi_k f_k(\boldsymbol{x}; \boldsymbol{\theta}_k), \tag{1}$$

where π_k represents the mixing proportions or weights, subject to the constraints $\pi_k > 0$ and $\sum_{k=1}^{G} \pi_k = 1$, and $f_k(\boldsymbol{x}; \boldsymbol{\theta}_k)$ denotes the multivariate density function of the kth component with parameters vector $\boldsymbol{\theta}_k$ $(k = 1, \ldots, G)$. If the density function $f_k()$ is usually assumed to be known, the parameters of the FMM, $\boldsymbol{\Psi} = (\pi_1, \ldots, \pi_{G-1}, \boldsymbol{\theta}_1, \ldots, \boldsymbol{\theta}_G)$, are unknown and need to be estimated from the data.

One of the earlier, and still the most popular, model for continuous data is the Gaussian mixture model (GMM), which is obtained from (1) assuming a multivariate Gaussian distribution for each component density (Fraley and Raftery, 2002; Bouveyron et al., 2019; Gormley et al., 2023; Scrucca et al., 2023). A *G*-component GMM can be expressed as

$$f(\boldsymbol{x}; \boldsymbol{\Psi}) = \sum_{k=1}^{G} \pi_k \phi(\boldsymbol{x}; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k),$$
(2)

where $\phi(\boldsymbol{x}; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)$ denotes the multivariate Gaussian density function with mean $\boldsymbol{\mu}_k$ and covariance matrix $\boldsymbol{\Sigma}_k$ for the k-th mixture component.

Parsimonious covariances decomposition can be obtained by imposing constraints on the geometric characteristics, such as volume, shape, and orientation, of corresponding ellipsoids. This is achieved using the following covariance matrices eigen-decomposition (Banfield and Raftery, 1993; Celeux and Govaert, 1995):

$$\boldsymbol{\Sigma}_k = \lambda_k \boldsymbol{U}_k \boldsymbol{\Delta}_k \boldsymbol{U}_k^{\mathsf{T}}$$

where $\lambda_k = |\mathbf{\Sigma}_k|^{1/d}$ is a scalar controlling the volume, $\mathbf{\Delta}_k$ is a diagonal matrix controlling the shape, such that $|\mathbf{\Delta}_k| = 1$ and with the normalized eigenvalues of $\mathbf{\Sigma}_k$ in decreasing order, and \mathbf{U}_k is an orthogonal matrix of eigenvectors of $\mathbf{\Sigma}_k$ that controls the orientation. Further details on the resulting 14 different models can be found in Scrucca et al. (2023, Sec. 2.2.1).

Maximum likelihood estimation of GMM parameters, $\Psi = (\pi_1, \ldots, \pi_{G-1}, \mu_1, \ldots, \mu_G, \Sigma_1, \ldots, \Sigma_G)$, is commonly carried out using the Expectation-Maximization (EM) algorithm (Dempster et al., 1977). Let z_i denote the membership multinomial latent variable, with $z_{ik} = 1$ if observation *i* belongs to cluster *k*, and $z_{ik} = 0$ otherwise. The EM algorithm iteratively maximizes the completedata log-likelihood:

$$\ell_C(\boldsymbol{\Psi}) = \sum_{i=1}^n \sum_{k=1}^G z_{ik} \left\{ \log \pi_k + \log \phi(\boldsymbol{x}_i; \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \right\}$$

by alternating between two steps, the Expectation (E) step and the Maximization (M) step. For a comprehensive treatment of estimation using the EM algorithm and its properties, refer to McLachlan and Krishnan (2008). Details on the M-step for the different covariance parameterizations can be found in Celeux and Govaert (1995) and (Scrucca et al., 2023, Sec. 2.2.2).

2.2 A transformation-based approach for Gaussian mixtures clustering

A direct application of Gaussian mixtures, as formulated in (2), may lead to inaccurate density estimates and erroneous clustering assignments when some or all the features of the dataset are bounded. To overcome these limitations, we propose employing the transformation-based approach introduced by Scrucca (2019). The core idea of this approach is to map the bounded features to an unbounded space, where standard GMMs can be applied more effectively, and then transform the results back to the original bounded space, where clustering assignments can be straightforwardly obtained.

Let \boldsymbol{x} represent a random vector from a distribution with bounded support $S_{\mathcal{X}} \subset \mathbb{R}^d$. Consider the family of continuous monotonic transformations, denoted by $\boldsymbol{y} = t(\boldsymbol{x}; \boldsymbol{\lambda})$, where $\boldsymbol{\lambda} = [\lambda_1, \lambda_2, \dots, \lambda_d]^{\top} \in \boldsymbol{\Lambda}$ represents the vector of transformation parameters, mapping the bounded

support of data to an unbounded space, $S_{\mathcal{Y}} \equiv \mathbb{R}^d$. Note that if one or more features do not require transformation, the corresponding λ parameters can be set to 1 and kept fixed during the fitting process.

On the transformed space, the data can be modeled using a standard GMM:

$$h(\boldsymbol{y};\boldsymbol{\Psi}) = \sum_{k=1}^{G} \pi_k \phi(\boldsymbol{y};\boldsymbol{\mu}_k,\boldsymbol{\Sigma}_k),$$

where Ψ refers here to the parameters in the transformed scale, so μ_k and Σ_k are, respectively, the mean vector and the covariance matrix of transformed features. Then, by applying the change-of-variable theorem, the density on the original scale can be recovered as:

$$f(\boldsymbol{x};\boldsymbol{\Psi},\boldsymbol{\lambda}) = h(t(\boldsymbol{x};\boldsymbol{\lambda})) \times |\boldsymbol{J}(t(\boldsymbol{x};\boldsymbol{\lambda}))|,$$
(3)

where $J(t(\boldsymbol{x};\boldsymbol{\lambda}))$ is the Jacobian of the transformation. Since we consider coordinate-independent transformations

$$t(\boldsymbol{x};\boldsymbol{\lambda}) = [t(x_1;\lambda_1), t(x_2;\lambda_2), \dots, t(x_d;\lambda_d)]^{\top},$$

the Jacobian simplifies to a product of first derivatives:

$$\boldsymbol{J}(t(\boldsymbol{x};\boldsymbol{\lambda})) = \prod_{j=1}^{d} t'(x_j;\lambda_j).$$
(4)

The transformation $\boldsymbol{y} = t(\boldsymbol{x}; \boldsymbol{\lambda})$ adopted is based on the *range-power transformation* introduced in Scrucca (2019), which is suitable for both partially and completely bounded data. For a data feature $x_j \in (l_j, +\infty)$, where l_j represents its lower bound, the range-power transformation is defined as:

$$t(x_j;\lambda_j) = \begin{cases} \frac{(x_j - l_j)^{\lambda_j} - 1}{\lambda_j} & \text{if } \lambda_j \neq 0\\ \log(x_j - l_j) & \text{if } \lambda_j = 0, \end{cases}$$
(5)

with continuous first derivative equal to

$$t'(x_j;\lambda_j) = \frac{\partial t(x_j;\lambda_j)}{\partial x_j} = (x_j - l_j)^{\lambda_j - 1}$$
(6)

for any $\lambda_j \in \Lambda$ $(j = 1, \ldots, d)$.

If a variable $x_j \in (l_j, u_j)$, where l_j and u_j represent the lower and upper bounds, respectively, the range-power transformation is defined as:

$$t(x_j;\lambda_j) = \begin{cases} \frac{\left(\frac{x_j - l_j}{u_j - x_j}\right)^{\lambda_j} - 1}{\lambda_j} & \text{if } \lambda_j \neq 0\\ \log\left(\frac{x_j - l_j}{u_j - x_j}\right) & \text{if } \lambda_j = 0, \end{cases}$$
(7)

with continuous first derivative given by

$$t'(x_j;\lambda_j) = \frac{\partial t(x_j;\lambda_j)}{\partial x_j} = \begin{cases} \left(\frac{x_j - l_j}{u_j - x_j}\right)^{\lambda_j - 1} \frac{u_j - l_j}{(u_j - x_j)^2} & \text{if } \lambda_j \neq 0\\ \frac{1}{x_j - l_j} + \frac{1}{u_j - x_j} & \text{if } \lambda_j = 0. \end{cases}$$
(8)

Equations (6) and (8) can be used to compute the Jacobian in (4). Together with the rangepower transformations in equations (5) and (7)), these allow for the estimation of the mixture density in (3). However, both the unknown mixture parameters Ψ and the transformation parameters λ need to be estimated.

2.3 Maximum likelihood estimation via the EM algorithm

Recalling the density in (3), the log-likelihood of the observed data can be expressed as:

$$\ell(\boldsymbol{\Psi},\boldsymbol{\lambda}) = \sum_{i=1}^{n} \log \left(\sum_{k=1}^{G} \pi_k \phi(t(\boldsymbol{x}_i;\boldsymbol{\lambda});\boldsymbol{\mu}_k,\boldsymbol{\Sigma}_k) \times |\boldsymbol{J}(t(\boldsymbol{x}_i;\boldsymbol{\lambda}))| \right).$$

Maximum likelihood estimation can be pursued via the EM algorithm by maximizing the completedata log-likelihood:

$$\ell_C(\boldsymbol{\Psi}, \boldsymbol{\lambda}) = \sum_{i=1}^n \sum_{k=1}^G z_{ik} \left\{ \log \pi_k + \log \phi(t(\boldsymbol{x}_i; \boldsymbol{\lambda}); \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) + \log |\boldsymbol{J}(t(\boldsymbol{x}_i; \boldsymbol{\lambda}))| \right\},$$

where z_i is the latent variable for cluster membership described in Section 2.1.

At iteration m of the EM algorithm, the conditional expectation of the complete-data loglikelihood given the observed data and the current parameter values can be expressed as:

$$Q(\boldsymbol{\Psi},\boldsymbol{\lambda};\boldsymbol{\Psi}^{(m)},\boldsymbol{\lambda}^{(m)}) = \sum_{i=1}^{n} \sum_{k=1}^{G} \widehat{z}_{ik}^{(m+1)} \left\{ \log \pi_k + \log \phi(t(\boldsymbol{x}_i;\boldsymbol{\lambda});\boldsymbol{\mu}_k,\boldsymbol{\Sigma}_k) + \log |\boldsymbol{J}(t(\boldsymbol{x}_i;\boldsymbol{\lambda}))| \right\},$$

where $\hat{z}_{ik}^{(m+1)} = \mathbb{E}(I(\boldsymbol{z}_i = k) | t(\boldsymbol{x}_i, \hat{\boldsymbol{\lambda}}^{(m)}), \boldsymbol{\Psi}^{(m)})$. Therefore, in the E-step the posterior probabilities are updated using

$$\widehat{z}_{ik}^{(m+1)} = \frac{\widehat{\pi}_k^{(m)} \phi\left(t(\boldsymbol{x}_i; \widehat{\boldsymbol{\lambda}}^{(m)}); \widehat{\boldsymbol{\mu}}_k^{(m)}, \widehat{\boldsymbol{\Sigma}}_k^{(m)}\right)}{\sum_{g=1}^G \widehat{\pi}_g^{(m)} \phi\left(t(\boldsymbol{x}_i; \widehat{\boldsymbol{\lambda}}^{(m)}); \widehat{\boldsymbol{\mu}}_g^{(m)}, \widehat{\boldsymbol{\Sigma}}_g^{(m)}\right)}.$$

In the M-step, the parameters (Ψ, λ) are updated by maximizing the *Q*-function, given the previous values of the parameters and the updated posterior probabilities. Following the Expectation-Conditional-Maximization (ECM) algorithm introduced by Meng and Rubin (1993), this maximization is carried out in two steps. In the first step, the updated value $\hat{\lambda}^{(m+1)}$ is computed by maximizing the Q-function with respect to λ . Since no closed-form solution is available, this requires numerical optimization, such as a Newton-type algorithm or a related method. The remaining parameters are then obtained as in standard EM algorithm but accounting for the updated transformation parameters $\hat{\lambda}^{(m+1)}$, i.e.

$$\widehat{\pi}_{k}^{(m+1)} = \frac{\sum_{i=1}^{n} \widehat{z}_{ik}^{(m+1)}}{n} \quad \text{and} \quad \widehat{\mu}_{k}^{(m+1)} = \frac{\sum_{i=1}^{n} \widehat{z}_{ik}^{(m+1)} t(\boldsymbol{x}_{i}; \widehat{\boldsymbol{\lambda}}^{(m+1)})}{\sum_{i=1}^{n} \widehat{z}_{ik}^{(m+1)}}.$$

The update formula for the covariance matrices depends on the assumed eigen-decomposition model. In the most general case of unconstrained covariance matrices, i.e. the VVV model in mclust nomenclature (see Scrucca et al., 2023, Table 2.1), we have

$$\widehat{\boldsymbol{\Sigma}}_{k}^{(m+1)} = \frac{\sum_{i=1}^{n} \widehat{z}_{ik}^{(m+1)} \left(t(\boldsymbol{x}_{i}; \widehat{\boldsymbol{\lambda}}^{(m+1)}) - \widehat{\boldsymbol{\mu}}_{k}^{(m+1)} \right) \left(t(\boldsymbol{x}_{i}; \widehat{\boldsymbol{\lambda}}^{(m+1)}) - \widehat{\boldsymbol{\mu}}_{k}^{(m+1)} \right)^{\top}}{\sum_{i=1}^{n} \widehat{z}_{ik}^{(m+1)}}.$$

Initialization of the above EM algorithm can be obtained by first estimating the optimal marginal transformations, and then using the final classification from a k-means algorithm on the range-power transformed variables. Alternatively, partitions obtained from model-based agglomerative hierarchical clustering (MBAHC; Scrucca and Raftery, 2015) can be used. This initial partition of data points is used to start the algorithm from the M-step. Finally, the EM algorithm is stopped when the log-likelihood improvement falls below a specified tolerance value or a maximum number of iterations is reached.

Model selection in finite mixture modeling is typically carried out using information criteria, such as the Bayesian Information Criterion (BIC; Schwarz, 1978), or the Integrated Completedata Likelihood criterion (ICL; Biernacki et al., 2000). Both criteria penalize model complexity, favoring more parsimonious models unless the additional parameters are justified by a significant improvement in the likelihood. Additionally, ICL introduces an extra penalization for clustering overlap, promoting models that produce well-separated clusters.

2.4 Clustering and classification uncertainty

In model-based clustering, assigning a data point to one of the identified clusters is straightforward using the maximum a posteriori (MAP) principle. According to MAP, each observation is assigned to the cluster that has the highest posterior probability.

Consider a partition of the data $\mathcal{D} = \{\boldsymbol{x}_i\}_{i=1}^n$ into G clusters, denoted as $\mathcal{C} = \{C_1, C_2, \ldots, C_G\}$, where $C_k \cap C_g = \emptyset$ (for $k \neq g$) and $\bigcup_{k=1}^G C_k = \mathcal{D}$. The MAP procedure assigns an observation \boldsymbol{x}_i to a cluster $C_{\hat{k}}$ according to the rule:

$$\boldsymbol{x}_i \in C_{\widehat{k}}$$
 with $\widehat{k} = \underset{k \in \{1, \dots, G\}}{\operatorname{arg\,max}} \widehat{z}_{ik},$

where \hat{z}_{ik} represents the posterior probability of observation x_i belonging to cluster k, i.e.

$$\widehat{z}_{ik} = \frac{\widehat{\pi}_k \phi\left(t(\boldsymbol{x}_i; \widehat{\boldsymbol{\lambda}}); \widehat{\boldsymbol{\mu}}_k, \widehat{\boldsymbol{\Sigma}}_k\right)}{\sum_{g=1}^G \widehat{\pi}_k \phi\left(t(\boldsymbol{x}_i; \widehat{\boldsymbol{\lambda}}); \widehat{\boldsymbol{\mu}}_g, \widehat{\boldsymbol{\Sigma}}_g\right)}.$$

Hard classification based on the MAP principle assumes the most likely cluster is the correct one. However, once each data point is assigned to its most probable cluster, it becomes essential to evaluate the uncertainty of these assignments. Assessing clustering uncertainty is thus crucial, as some data points may not clearly belong to a single cluster, especially when they lie near cluster boundaries. This also allows to distinguish between well-separated clusters and those that exhibit substantial overlap.

To measure this uncertainty, we can examine the distribution of posterior probabilities across all clusters rather than focusing solely on the maximum probability. The uncertainty for each data point is captured by the following index:

$$u_i = 1 - \max_k \widehat{z}_{ik}.$$

This score ranges from 0 to (G-1)/G, with values close to 0 indicating low classification uncertainty, while values near the upper bound suggest greater uncertainty.

An alternative measure of uncertainty can be derived from the entropy. For each data point, we can compute

$$e_i = -\sum_{k=1}^G \widehat{z}_{ik} \log(\widehat{z}_{ik}),$$

which yields values in the range $[0, \log(G)]$, with higher entropy values indicating greater uncertainty in the assignment.

An overall index of classification uncertainty can be obtained by summing the entropy values for all data points:

$$E = \sum_{i=1}^{n} e_i = -\sum_{i=1}^{n} \sum_{k=1}^{G} \hat{z}_{ik} \log(\hat{z}_{ik}).$$

This classification entropy measures the uncertainty in clustering, taking the value zero when each data points is assigned to its respective cluster with probability 1. As uncertainty increases, the entropy value grows accordingly. For easier interpretation, a normalized version of classification entropy, ranging between 0 and 1, can be calculated by taking the ratio of its value to the maximum possible value. This leads to the *Normalized Entropy Criterion* (NEC) proposed by Celeux and Soromenho (1996):

$$NEC = \frac{E}{n\log(G)},$$

for G > 1, and with the implicit assumption that NEC = 0 when G = 1.

As a final note, we caution against aiming for the smallest possible value of NEC when assessing clustering results. A very small NEC suggests well-separated clusters, with low uncertainty in the MAP assignments. On the contrary, an increase in NEC may indicate significant overlap among clusters. While this could be a sign of poor clustering, often it reflects the inherent nature of the data, where the clusters are not well separated. Therefore, while NEC is a useful measure of the overall uncertainty in a clustering solution, it should not be used as the sole criterion for evaluating the quality of the resulting partition.

3 Applications

3.1 Enzyme data

The enzyme data set, originally described by Bechtel et al. (1993), consists of enzymatic activity measurements in the blood for an enzyme involved in the metabolism of carcinogenic substances, from a sample of n = 245 unrelated individuals. The primary interest lies in identifying subgroups of slow or fast metabolizers, which serve as markers of genetic polymorphism in the general population. This benchmark data set for mixture models is notable for containing at least two components, one of which exhibits clear skewness. Richardson and Green (1997) employed a Bayesian GMM estimated using reversible jump MCMC to analyze the distribution of enzymatic activity. Their analysis suggests that a model with 3 to 5 mixture components is plausible, with the three-component solution being preferred, particularly in line with a simple underlying genetic model. In contrast, Karlis and Santourian (2009) applied a Normal Inverse Gaussian Mixture (NIGM) and identified two distinct, non-overlapping clusters.



Figure 1: (a) Histogram of the enzyme data and (b) plot of component densities, scaled by the corresponding mixing probabilities, estimated using the Gaussian Mixture Model for Bounded data (GMMB).

Figure 1a presents the histogram of the enzyme data, while Figure 1b shows the estimated densities from the selected two-component GMMB model with unconstrained variances and a range-power transformation parameter in (5) equals to $\hat{\lambda} = 0.3666$.

This model, identified as the best fit according to both the BIC and ICL criteria, distinguishes between two clusters: one consisting of 152 observations characterized by a skewed distribution and enzymatic activity levels below 0.5, and a second, comprising 93 observations, forming a nearly symmetric group with enzymatic activity levels above 0.5. These results align with those reported by Karlis and Santourian (2009) and have been reproduced here using the function MGHM() from the **MixtureMissing** R package (Tong and Tortora, 2024).

Although the GMMB model provides a better fit, as indicated by higher BIC and ICL values, and is more parsimonious with fewer parameters to estimate, it reflects a slightly larger classification uncertainty than the NIGM model, as measured by NEC. This is primarily due to those observations near the classification boundary at an enzymatic level of 0.5. In any case, both models represent a clear improvement over the standard GMM, as shown in Table 1.

Table 1: Model comparison for the clustering of the enzyme data using Gaussian Mixture Model (GMM), mixture of Normal Inverse Gaussian (NIGM) distributions, and Gaussian Mixture Model for Bounded data (GMMB), with the number of components selected according to the BIC criterion.

Model	log-likelihood	df	BIC	ICL	NEC
GMM(V,2)	-54.6401	5	-136.7865	-148.9526	0.1109
$\mathrm{NIGM}(2)$	-41.4723	9	-132.4558	-132.6530	0.0052
GMMB(V,2)	-46.1870	6	-125.3815	-129.6447	0.0208

3.2 Wholesale customer segmentation

Customer segmentation aims to divide customers into distinct groups based on common characteristics, such as purchasing behavior. This partition helps businesses to tailor their marketing efforts, products, and services to specific customer segments, potentially maximizing customer satisfaction and profitability.

The wholesale customers dataset, freely available on the UCI Machine Learning Repository (Cardoso, 2013), is a widely-used dataset to explore clustering and segmentation techniques. The dataset includes 440 observations, each representing a customer of a wholesale distributor. For each client, annual spending, in monetary units, is recorded for six product categories: fresh, milk, grocery, frozen, detergents paper, and delicatessen. Two additional categorical variables are available, corresponding to the region and customer channel. Note that, unlike other authors who have analyzed this dataset, such as Punzo and Tortora (2021), we conduct our analysis using the variables in their original scale.

Since annual expenditure cannot take negative values, variables used in the segmentation process are naturally bounded at zero. Standard clustering techniques often assume unbounded data, which is clearly not appropriate in the current context and, by failing to capture the true underlying structure, can lead to suboptimal segmentation.

Table 2 reports the results obtained by fitting the standard GMM for unbounded features, the GMMB model proposed in this paper, and some models discussed in Punzo and Tortora (2021), namely the *Mixture of Contaminated Normal* distributions (MCNM) and the *Mixture of Multiple Scaled Contaminated Normal* distributions (MSCNM). For all models, the number of mixing components is fixed at G = 2. The MCNM model is fitted using the function CNmixt() in the **ContaminatedMixt** R package (Punzo et al., 2018, 2023), while the MSCNM is fitted using the function mscn() in the **MSclust** R package (Tortora et al., 2024). Models are compared using the BIC, ICL, NEC and the adjusted Rand index (Hubert and Arabie, 1985, ARI;).

Both the MCNM and MSCNM models provide significant improvements over the standard GMM, offering superior fit as evidenced by higher BIC values, and better classification performance with lower error rates (ER) and higher adjusted Rand index (ARI). However, both models are outperformed by the GMMB model, which not only achieves a higher BIC, indicating a better overall fit, but also approximately halves the ER and doubles the ARI, further emphasizing its superior classification capabilities.

Both the MCNM and MSCNM models show significant improvements over the standard GMM, offering a superior fit, as indicated by higher BIC values, and enhanced classification performance, with higher adjusted Rand index (ARI) values. However, these models also exhibit increased classification uncertainty, as reflected by higher NEC values, compared to the sharper but less accurate classifications of the GMM. Nonetheless, both models are outperformed by the GMMB model, which not only achieves a higher BIC, indicating an improved overall fit, but also doubles the ARI, further emphasizing its superior classification capabilities. As for the MCNM and MSCNM models, the classification uncertainty of GMMB is higher than that of GMM, reflecting a more pronounced overlap among mixture components in GMMB model.

Table 2: Model comparison for the clustering of the wholesale dataset, using Gaussian Mixture Model (GMM), Mixture of Contaminated Normals (MCNM), Mixture of Multiple Scaled Contaminated Normals (MSCNM), and Gaussian Mixture Model for Bounded data (GMMB), all fitted using G = 2 mixture components.

Model	log-likelihood	df	BIC	ICL	NEC	ARI
GMM(VVV,2)	-25069.70	55	-50474.18	-50495.52	0.0796	0.1028
MCNM(VVV,2)	-24614.60	59	-49588.32	-49626.70	0.1439	0.3808
MSCNM(2)	-24498.57	79	-49477.99	-49491.15	0.1290	0.3642
GMMB(VVE,2)	-23909.79	46	-48099.57	-52003.48	0.1539	0.6585

Figure 2 shows the range-power transformations in (5) for each variable in the wholesale dataset, as estimated by the GMMB model. All transformations exhibit a log-type shape, with λ values ranging from 0.05 to 0.3. These transformations effectively reduce the pronounced positive skewness in the marginal distributions, as illustrated in the corresponding marginal histograms.

Figure 3 presents the scatterplot matrix of the features from the wholesale dataset, with points representing wholesale customers marked by their assigned cluster. The distributions within each segment exhibit noticeable skewness due to the zero-lower bound constraint of the features, a characteristic effectively captured by the GMMB model used for the segmentation.

Finally, Figure 4 shows the average latent profiles for the two clusters, with the marginal means represented by the dashed line. This graph clearly highlights the key features that distinguish the two segments of wholesale distributor customers, with the "delicatessen" and "frozen" product categories not playing a significant role in the segmentation



Figure 2: Plots displaying the selected range-power transformation $t(X; \hat{\lambda})$ as a function of the original feature X for variables in the wholesale dataset, with marginal histograms showing the distribution before and after the applied transformation.

Figure 3: Scatterplot matrix of variables in the wholesale dataset, with wholesale customers marked by their cluster membership.

Figure 4: Latent profiles plot displaying the estimated means by cluster (solid lines) from the fitted GMMB model, and the marginal means (dashed line) for each feature in the wholesale dataset.

3.3 Human Development Index

The Human Development Index (HDI) is a composite index provided by the United Nations Development Programme (UNDP) and designed to measure the level of human development in world countries (Herre and Arriagada, 2023).

The HDI combines three key dimensions of human development: *health*, measured by life expectancy at birth, *education*, measured by expected years of schooling (for children of school entering age) and average years of schooling (for adults aged 25 and older), and *standard of living*, measured by Gross National Income (GNI) per capita. These three indicators are first normalized to have a minimum of 0 and a maximum of 1, then are combined by calculating the geometric mean. The resulting HDI value for each country represents an overall score in the range [0, 1], with higher values indicating a better level of human development.

Figure 5: (a) Histogram of the Human Development Index (HDI) distribution for the year 2022. (b) Plot of component densities, scaled by the corresponding mixing probabilities, estimated using the Gaussian Mixture Model for Bounded data (GMMB).

Figure 5a shows the distribution of the Human Development Index (HDI) for the year 2022, revealing a clearly multimodal pattern. HDI values span a range from approximately 0.4 to just under 1, reflecting significant variation across countries.

According to the BIC criterion, the best-fitting GMMB model comprises three components with equal variance and a range-power transformation parameter in (7) equals to $\hat{\lambda} = -0.12$. The corresponding component densities (scaled by their mixing probabilities) are shown in Figure 5b. This model indicates the presence of three distinct clusters of countries, which can be broadly categorized as low, medium, and high human development clusters.

Figure 6 presents a world map with countries colored according to their respective cluster memberships. The high-HDI cluster includes mainly developed countries, such as those in North America and Europe, along with Japan, Australia, New Zealand, and the United Arab Emirates. The low-HDI cluster consists mostly of underdeveloped countries, primarily in Africa, with additional members in Asia (Afghanistan, Pakistan, Yemen, Nepal, Myanmar, Laos, Cambodia) and Central America (Guatemala, Haiti, Honduras). The remaining countries, mainly from Central and South America, Eastern Europe, and parts of Asia, fall into the medium-HDI cluster.

Figure 6: World map showing countries clustered by Human Development Index (HDI) using the Gaussian Mixture Model for Bounded data (GMMB).

The selected GMMB model can be compared to the standard Gaussian Mixture Model (GMM) and a Mixture of Beta distributions (BMM; Bagnato and Punzo, 2013; Dean and Nugent, 2013), with the number of components again selected using the BIC criterion. The results in Table 3 indicate that the GMMB model achieves a better fit, as evidenced by a higher BIC, along with a slightly reduced classification uncertainty, as shown by a lower NEC. Overall, the GMMB approach offers a reasonable and interpretable clustering solution with lowered uncertainty, further highlighting its effectiveness in this context.

Table 3: Model comparison for the clustering of the Human Development Index (HDI), using Gaussian Mixture Model (GMM), Beta Mixture Model (BMM), and Gaussian Mixture Model for Bounded data (GMMB), with the number of components selected according to the BIC criterion.

Model	log-likelihood	df	BIC	ICL	NEC
$\overline{\mathrm{GMM}(\mathrm{V},3)}$	96.6144	8	152.5775	127.8651	0.1660
BMM(3)	97.8045	8	154.9578	128.7258	0.1609
GMMB(E,3)	97.8727	7	160.1756	133.8794	0.1575

4 Conclusion

This study aimed to develop a model-based clustering approach for bounded data, extending earlier work of Scrucca (2019) on Gaussian mixture density estimation. The primary research question focused on overcoming the unique challenges posed by bounded data, particularly by transforming bounded features into an unbounded space to facilitate the application of GMMs. Through the proposed range-power transformation, our method provides a flexible and interpretable clustering framework that preserves data bounds, enhancing the accuracy and applicability of GMMs for real-world bounded data scenarios.

The analyzed datasets demonstrated that this transformation-based approach offers a unified and flexible framework capable of handling diverse bounded data structures while ensuring model parsimony and interpretability. Compared to traditional GMMs and other model-based approaches using distributions with limited support, our method showed improved clustering partitions and interpretability, highlighting its relevance for practical applications involving the analysis of bounded data.

Despite the advantages mentioned above, some limitations remain. The proposed transformationbased approach requires selecting a set of appropriate transformation parameters. However, the use of a coordinate-wise approach may prove suboptimal if cluster-dependent data transformations are needed, a topic warranting further exploration in future research.

Additionally, the suitability of the proposal for handling heavy-tailed or clusters could be investigated. This would allow to extend the transformation-based approach also in case of unbounded but non-spherical cluster distributions, hence offering an alternative approach to the several existing methods for non-Gaussian clusters (McNicholas, 2016).

Code and Data Availability All the analyses have been conducted in R (R Core Team, 2024) using the mclust and mclustAddons packages (Fraley et al., 2024; Scrucca, 2024). Source code and datasets to reproduce the analyses are available in a GitHub repository at https://github.com/luca-scr/MclustBounded.

References

- Bagnato, L. and Punzo, A. (2013). Finite mixtures of unimodal beta and gamma densities and the k-bumps algorithm. Computational Statistics, 28(4):1571–1597.
- Banfield, J. and Raftery, A. E. (1993). Model-based Gaussian and non-Gaussian clustering. *Bio-metrics*, 49:803–821.
- Bechtel, Y. C., Bonaiti-Pellie, C., Poisson, N., Magnette, J., and Bechtel, P. R. (1993). A population and family study N-acetyltransferase using caffeine urinary metabolites. *Clinical Pharmacology* & Therapeutics, 54(2):134–141.
- Biernacki, C., Celeux, G., and Govaert, G. (2000). Assessing a mixture model for clustering with the integrated completed likelihood. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 22(7):719–725.
- Bouveyron, C., Celeux, G., Murphy, T. B., and Raftery, A. E. (2019). *Model-Based Clustering and Classification for Data Science: With Applications in R.* Cambridge University Press.
- Browne, R. P. and McNicholas, P. D. (2015). A mixture of generalized hyperbolic distributions. Canadian Journal of Statistics, 43(2):176–198.
- Cardoso, M. (2013). Wholesale customers. UCI Machine Learning Repository.
- Celeux, G. and Govaert, G. (1995). Gaussian parsimonious clustering models. *Pattern Recognition*, 28:781–793.
- Celeux, G. and Soromenho, G. (1996). An entropy criterion for assessing the number of clusters in a mixture model. *Journal of Classification*, 13(2):195–212.
- Dean, N. and Nugent, R. (2013). Clustering student skill set profiles in a unit hypercube using mixtures of multivariate betas. Advances in Data Analysis and Classification, 7(3):339–357.
- Dempster, A. P., Laird, N. M., and Rubin, D. B. (1977). Maximum likelihood from incomplete data via the EM algorithm (with discussion). *Journal of the Royal Statistical Society: Series B* (Statistical Methodology), 39:1–38.
- Fraley, C. and Raftery, A. E. (2002). Model-based clustering, discriminant analysis, and density estimation. *Journal of the American Statistical Association*, 97(458):611–631.
- Fraley, C., Raftery, A. E., and Scrucca, L. (2024). mclust: Gaussian Mixture Modelling for Model-Based Clustering, Classification, and Density Estimation. R package version 6.1.1.
- Gallaugher, M. P. B., McNicholas, P. D., Melnykov, V., and Zhu, X. (2020). Skewed distributions or transformations? modelling skewness for a cluster analysis.
- Gormley, I. C., Murphy, T. B., and Raftery, A. E. (2023). Model-based clustering. Annual Review of Statistics and Its Application, 10:573–595.
- Hedelin, P. and Skoglund, J. (2000). Vector quantization based on Gaussian mixture models. *IEEE Transactions on Speech and Audio Processing*, 8(4):385–401.

- Herre, B. and Arriagada, P. (2023). The human development index and related indices: what they are and what we can learn from them. *Our World in Data*.
- Hubert, L. and Arabie, P. (1985). Comparing partitions. Journal of Classification, 2:193–218.
- John, S. (1970). On identifying the population of origin of each observation in a mixture of observations from two gamma populations. *Technometrics*, 12(3):565–568.
- Karlis, D. and Santourian, A. (2009). Model-based clustering with non-elliptically contoured distributions. *Statistics and Computing*, 19(1):73–83.
- Lindblom, J. and Samuelsson, J. (2003). Bounded support Gaussian mixture modeling of speech spectra. *IEEE Transactions on Speech and Audio Processing*, 11(1):88–99.
- Lo, K. and Gottardo, R. (2012). Flexible mixture modeling via the multivariate t distribution with the box-cox transformation: an alternative to the skew-t distribution. *Statistics and computing*, 22(1):33–52.
- Manly, B. F. J. (1976). Exponential data transformations. The Statistician, 25(1):37–42.
- McLachlan, G. J. and Krishnan, T. (2008). *The EM Algorithm and Extensions*. Wiley-Interscience, Hoboken, New Jersey, 2nd edition.
- McLachlan, G. J., Lee, S. X., and Rathnayake, S. I. (2019). Finite mixture models. Annual Review of Statistics and Its Application, 6(1):355–378.
- McLachlan, G. J. and Peel, D. (2000). Finite Mixture Models. Wiley, New York.
- McNicholas, P. D. (2016). Mixture Model-Based Classification. Chapman & Hall/CRC.
- McNicholas, S. M., McNicholas, P. D., and Browne, R. P. (2017). A mixture of variance-gamma factor analyzers. In Ahmed, S. E., editor, *Big and Complex Data Analysis: Methodologies and Applications*, pages 369–385, Cham. Springer International Publishing.
- Meng, X.-L. and Rubin, D. B. (1993). Maximum likelihood estimation via the ECM algorithm: A general framework. *Biometrika*, 80(2):267–278.
- Punzo, A., Mazza, A., and McNicholas, P. D. (2018). ContaminatedMixt: An R package for fitting parsimonious mixtures of multivariate contaminated normal distributions. *Journal of Statistical Software*, 85(10):1–25.
- Punzo, A., Mazza, A., and McNicholas, P. D. (2023). ContaminatedMixt: Clustering and Classification with the Contaminated Normal. R package version 1.3.8.
- Punzo, A. and Tortora, C. (2021). Multiple scaled contaminated normal distribution and its application in clustering. *Statistical Modelling*, 21(4):332–358.
- R Core Team (2024). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Richardson, S. and Green, P. J. (1997). On bayesian analysis of mixtures with an unknown number of components (with discussion). Journal of the Royal Statistical Society: Series B (Statistical Methodology), 59(4):731–792.

- Schwarz, G. (1978). Estimating the dimension of a model. The Annals of Statistics, 6(2):461–464.
- Scrucca, L. (2019). A transformation-based approach to Gaussian mixture density estimation for bounded data. *Biometrical Journal*, 61(4):873–888.
- Scrucca, L. (2024). mclustAddons: Addons for the 'mclust' Package. R package version 0.9.1.
- Scrucca, L., Fraley, C., Murphy, T. B., and Raftery, A. E. (2023). Model-Based Clustering, Classification, and Density Estimation Using mclust in R. Chapman & Hall/CRC, Boca Raton, FL.
- Scrucca, L. and Raftery, A. E. (2015). Improved initialisation of model-based clustering using Gaussian hierarchical partitions. Advances in Data Analysis and Classification, 4(9):447–460.
- Tong, H. and Tortora, C. (2024). MixtureMissing: Robust and Flexible Model-Based Clustering for Data Sets with Missing Values at Random. R package version 3.0.2.
- Tortora, C., Punzo, A., and Tran, L. (2024). MSclust: Multiple-Scaled Clustering. R package version 1.0.4.
- Wiper, M., Insua, D. R., and Ruggeri, F. (2001). Mixtures of gamma distributions with applications. Journal of Computational and Graphical Statistics, 10(3):440–454.
- Yeo, I.-K. and Johnson, R. A. (2000). A new family of power transformations to improve normality or symmetry. *Biometrika*, 87(4):954–959.
- Young, D. S., Chen, X., Hewage, D. C., and Nilo-Poyanco, R. (2019). Finite mixture-of-gamma distributions: estimation, inference, and model-based clustering. Advances in Data Analysis and Classification, 13(4):1053–1082.
- Zhu, X. and Melnykov, V. (2018). Manly transformation in finite mixture modeling. *Computational Statistics & Data Analysis*, 121:190–208.