GELFAND-TSETLIN CRYSTALS OF KOSTANT-KUMAR MODULES

MRIGENDRA SINGH KUSHWAHA

ABSTRACT. We give Gelfand-Tsetlin crystals for the Kostant-Kumar modules for the finite simple Lie algebra of type A. Kostant-Kumar modules are cyclic submodules of the tensor product of two irreducible highest weight modules of a symmetrizable Kac-Moody Lie algebras. In this case (type A), we also provide a polytopal model for Kostant-Kumar modules in terms of *BiKogan faces*.

1. INTRODUCTION

1.1. Let \mathfrak{g} be a symmetrizable Kac-Moody Lie algebra, \mathfrak{h} be a Cartan subalgebra, \mathfrak{b} be the Borel subalgebra containing \mathfrak{h} , \mathfrak{b}^- be the opposite Borel subalgebra to \mathfrak{b} , P be the weight lattice, P^+ be the set of dominant integral weights and W be the Weyl group of \mathfrak{g} . Let $V(\lambda)$ denote the irreducible highest weight module of \mathfrak{g} for $\lambda \in P^+$. Given $\lambda, \mu \in P^+$ and $w \in W$, the Kostant-Kumar (KK) module $K(\lambda, w, \mu)$ is a cyclic submodule of $V(\lambda) \otimes V(\mu)$ generated by the vector $v_{\lambda} \otimes v_{w\mu}$ (see [8]), that is

$$K(\lambda, w, \mu) := \mathfrak{Ug}(v_{\lambda} \otimes v_{w\mu}), \tag{1.1}$$

where v_{λ} is the highest weight vector of $V(\lambda)$, $v_{w\mu}$ be a non zero weight vector of weight $w\mu$ in $V(\mu)$ and \mathfrak{Ug} be the universal enveloping algebra of \mathfrak{g} . The following results are well-known (see [8]):

(1) $K(\lambda, 1, \mu) \cong V(\lambda + \mu)$ where 1 denote the identity element of W,

(2) $K(\lambda, w_0, \mu) = V(\lambda) \otimes V(\mu)$ where w_0 is the longest element of W,

(3) $K(\lambda, w, \mu) \subseteq K(\lambda, w', \mu)$ if $w \leq w'$ in the Bruhat order in W.

The item (3) shows that KK submodules form a filtration of $V(\lambda) \otimes V(\mu)$ with respect to the Bruhat order in W. Further results on KK-modules can be found in the following references [6–9, 12].

In our paper [8], we gave a path model for the KK-modules in terms of Lakshmibai-Seshadri (LS) paths provided \mathfrak{g} is either symmetric or of finite type, which is a generalization of the Littelmann's path model for the tensor product of two irreducible highest weight modules of \mathfrak{g} [10, 11].

In this paper, we consider $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$ the set of $n \times n$ complex matrices with trace zero and provide a polytopal model for the KK-modules in terms of pairs of Gelfand-Tsetlin patterns (see theorem 3.2.1) which is a very simple description. It is a generalization of well known polytopal model in terms of pairs of Gelfand-Tsetlin patterns for the tensor product of two irreducible highest weight modules of \mathfrak{g} . The crucial ingredient for the above polytopal model

²⁰¹⁰ Mathematics Subject Classification. 17B67, 17B10, 05E10.

Key words and phrases. Gelfand-Tsetlin polytope, Demazure crystals, Kogan faces, BiKogan faces.



FIGURE 2.1. Gelfand-Tsetlin array for n = 5.

of KK-modules is an association of permutation $\mathfrak{p}(P,Q)$ to a given pair of GT patterns (P,Q). The description of $\mathfrak{p}(P,Q)$ is visually simple and a procedure for it is stated in section 3.3.

1.2. We recall that for $w \in W$ and $\mu \in P^+$, the *Demazure module* $V_w(\mu)$ is the cyclic $\mathfrak{U}\mathfrak{b}$ -submodule and the *opposite Demazure module* $V^w(\mu)$ is the cyclic $\mathfrak{U}\mathfrak{b}^-$ -submodule of $V(\mu)$ generated by a nonzero vector $v_{w\mu}$ of weight $w\mu$, where $\mathfrak{U}\mathfrak{b}$ and $\mathfrak{U}\mathfrak{b}^-$ are the universal enveloping algebra of \mathfrak{b} and \mathfrak{b}^- respectively.

In section 5.6, we define BiKogan faces of a product of two Gelfand-Tsetlin polytopes, in the same spirit of Kogan faces and dual Kogan faces (see [2], [5]) of a Gelfand-Tsetlin polytope. A certain union of Kogan faces (resp. dual Kogan faces) gives a polytopal model for Demazure crystals (resp. opposite Demazure crystals). In the same spirit, our polytopal model for the KK-modules is described as a certain union of BiKogan faces (see Theorem 5.6.2).

1.3. Acknowledgement. I thank Sankaran Viswanath for his insightful suggestions and comments on the draft of the paper. I also extend my thanks to K. N. Raghavan for many useful discussions.

2. Gelfand-Tsetlin pattern

2.1. Let $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$. In this case, the Weyl group W can be identified with the permutation group S_n of the set $\{1, 2, \ldots, n\}$ and the set of dominant integral weights P^+ with the set of integer partitions with *n*-parts such that the n^{th} -part is zero. A sequence $\mu = (\mu_1, \mu_2, \ldots, \mu_n) \in \mathbb{Z}_{\geq 0}^n$ is called an integer partition if $\mu_i \geq \mu_{i+1}$ for all $i = 1, 2, \ldots, n-1$. There are many combinatorial models for the irreducible representation $V(\mu)$ indexed

by $\mu \in P^+$. The set of integral Gelfand-Tsetlin (GT) patterns of shape μ is one of the combinatorial models of $V(\mu)$, introduced by Gelfand and Tsetlin [3].

2.2. A GT-pattern of size n is a triangular array $A = (a_{ij})_{n \ge i \ge j \ge 1} \in \mathbb{R}^{n(n+1)/2}$ of real numbers (see Figure 2.1) satisfying the following ("North-East" and "South-East") inequalities for all $1 \le j < i \le n$: $NE_{ij}(A) = a_{ij} - a_{i-1,j} \ge 0$ (marked by red) and $SE_{ij}(A) = a_{i-1,j} - a_{i,j+1} \ge 0$ (marked by blue). For a given integer partition $\mu = (\mu_1, \mu_2, \dots, \mu_n)$ with n-parts, let $GT(\mu)$ denote the set of all GT-patterns of size n with fixed bottom row $a_{ni} = \mu_i$

for $1 \le i \le n$. Note that $GT(\mu)$ is a polytope and it is called the GT-polytope of shape μ . The *weight* of a GT-pattern A of size n is defined as follows:

$$wt(A) := (p_1, p_2, \dots, p_n)$$
 where $p_i = \sum_{j=1}^i a_{ij} - \sum_{j=1}^{i-1} a_{i-1,j}$.

For example, let $\mu = (5, 4, 2, 1, 0)$ and the following be a GT pattern of shape μ ,



the weight of the above GT-pattern is (3, 2, 4, 2, 1).

2.3. The character of $V(\mu)$ defined by $\operatorname{Char} V(\mu) = \sum_{\beta \in P} d_{\mu}(\beta) \exp^{\beta}$ where $d_{\mu}(\beta) = \dim V(\mu)_{\beta}$ is the dimension of the β -weight space. Let $\operatorname{GT}_{\mathbb{Z}}(\mu)$ denote the set of all integral points in the polytope $\operatorname{GT}(\mu)$. For a given $\mu \in P^+$, the set of integral GT-patterns $\operatorname{GT}_{\mathbb{Z}}(\mu)$ of shape μ is a combinatorial model for the irreducible module $V(\mu)$ and $\operatorname{Char} V(\mu) = \sum_{P \in \operatorname{GT}_{\mathbb{Z}}(\mu)} \exp^{wt(P)}$. By work of Littelmann [10] [11] it is well known that

$$\operatorname{GT}_{\mathbb{Z}}(\lambda) \times \operatorname{GT}_{\mathbb{Z}}(\mu) = \{ (P, Q) \mid P \in \operatorname{GT}_{\mathbb{Z}}(\lambda), \ Q \in \operatorname{GT}_{\mathbb{Z}}(\mu) \}$$

is a combinatorial model for the tensor product $V(\lambda) \otimes V(\mu)$ in the sense that:

Char
$$V(\lambda) \otimes V(\mu) = \sum_{(P,Q) \in \operatorname{GT}_{\mathbb{Z}}(\lambda) \times \operatorname{GT}_{\mathbb{Z}}(\mu)} \exp^{wt(P) + wt(Q)}$$
. (2.1)

For given $\lambda, \mu \in P^+$ and a permutation $w \in S_n$, we determine a subset $\operatorname{GT}_{\mathbb{Z}}(\lambda, w, \mu)$ of $\operatorname{GT}_{\mathbb{Z}}(\lambda) \times \operatorname{GT}_{\mathbb{Z}}(\mu)$ and we show that $\operatorname{GT}_{\mathbb{Z}}(\lambda, w, \mu)$ is a combinatorial model for the KK-module $K(\lambda, w, \mu)$ (see Theorem 3.2.1).

3. Gelfand-Tsetlin Crystal for the KK-module

3.1. We recall here a binary operation * on the permutation group S_n by Deodhar [1] (more generally on the Weyl group). Let $w \in S_n$ and s be a simple reflection in S_n then $w*s := \max\{w, ws\}$, where $\max\{w, ws\} = w$ if $w \ge ws$ and $\max\{w, ws\} = ws$ if $ws \ge w$ in the Bruhat order. The binary operation * is an associative operation, then for any $w, w' \in S_n$ the permutation w*w' is well defined and $w*w' = \max I(w)I(w')$ [8, §2 Remark 2.9], where I(w) is the Bruhat interval of w, that is $I(w) = \{u \in S_n \mid u \le w\}$.

Definition 3.1.1 (Demazure product). Let $w = s_{i_1}s_{i_2}\cdots s_{i_k}$ be a word which may not be reduced in the simple reflections s_{i_j} for $1 \le j \le k$. The Demazure product of w is defined as follows:

$$*(w) := s_{i_1} * s_{i_2} * \cdots * s_{i_k}.$$

For example $*(s_1s_3s_1s_2s_2) = s_3s_1s_2$.



FIGURE 3.1. The blue edges $a_{i-1,j} \longrightarrow a_{i,j+1}$ labelled by s_j and the red edges $a_{ij} \longrightarrow a_{i-1,j}$ are labelled by s_{i-j} .

Lemma 3.1.2. Let $w = s_{i_1}s_{i_2}\cdots s_{i_k}$ be a word which may not be reduced then the set $\{*(u) \mid u \text{ is a subword of } w\}$ has a unique maximal element and the unique maximal element is *(w).

Proof. Let $u = s_{i_{j_1}} s_{i_{j_2}} \cdots s_{i_{j_t}}$ be a subword of w where $j_s \in \{1, 2, \ldots, k\}$ for $s = 1, 2, \ldots, t$. We know that $*(w) = \max I(s_{i_1})I(s_{i_2})\cdots I(s_{i_k})$, similarly $*(u) = \max I(s_{i_{j_1}})I(s_{i_{j_2}})\cdots I(s_{i_{j_t}})$. The above expression implies that $*(u) \leq *(w)$.

Lemma 3.1.3. Let $w = s_{i_1}s_{i_2}\cdots s_{i_k}$ be a word which may not be reduced, where s_{i_j} are simple reflections for $1 \le j \le k$. Consider $w' = s_{i_1}\cdots s_{i_k}$ and $w'' = s_{i_{t+1}}\cdots s_{i_k}$ for some t such that 1 < t < k. Then, *(w) = *(w') * (*(w'')).

The above Lemma follows since * is an associative operation.

3.2. Now we will state the main theorem. For that, we need the following subset of GT polytope. For a given pair $(P, Q) \in \operatorname{GT}(\lambda) \times \operatorname{GT}(\mu)$ we associate a permutation $\mathfrak{p}(P, Q) \in S_n$ below in equation (3.3). Next we define a subset $\operatorname{GT}(\lambda, w, \mu)$ of $\operatorname{GT}(\lambda) \times \operatorname{GT}(\mu)$ for a given permutation w as follows:

$$\operatorname{GT}(\lambda, w, \mu) := \{ (P, Q) \in \operatorname{GT}(\lambda) \times \operatorname{GT}(\mu) \mid \mathfrak{p}(P, Q) \le w \}$$
(3.1)

where \leq is the Bruhat order in S_n . We denote by $\operatorname{GT}_{\mathbb{Z}}(\lambda, w, \mu)$ the set of all integral points in $\operatorname{GT}(\lambda, w, \mu)$. Now we state our main theorem which provides a Gelfand-Tsetlin crystal for KK-modules.

Theorem 3.2.1. For a given $\lambda, \mu \in P^+$ and $w \in S_n$, the set $GT_{\mathbb{Z}}(\lambda, w, \mu)$ is a Gelfand-Tsetlin model for the KK-module $K(\lambda, w, \mu)$ in the sense that

Char
$$K(\lambda, w, \mu) = \sum_{(P,Q)\in \operatorname{GT}_{\mathbb{Z}}(\lambda, w, \mu)} \exp^{wt(P) + wt(Q)},$$
 (3.2)

and $GT_{\mathbb{Z}}(\lambda, w, \mu)$ is e_i, f_i invariant where e_i and f_i are raising and lowering crystal operators defined below in the section 5.

We will prove theorem 3.2.1 at the end of the section 6.

3.3. Association of a permutation to a GT pair (P,Q).

Let $(P, Q) \in \operatorname{GT}(\lambda) \times \operatorname{GT}(\mu)$ for given partitions λ, μ with *n*-parts. Let $P = (a_{ij})_{n \geq i \geq j \geq 1}$ be a GT-pattern of size *n* as shown in Figure 2.1. Consider the labeling of South-East edges of *P* (marked by blue) $a_{i-1,j} \to a_{i,j+1}$ by the simple reflection s_j as shown in the left in Figure 3.1 for $n \geq i > j \geq 1$. Let $Q = (a_{ij})_{n \geq i \geq j \geq 1}$ be a GT-pattern of size *n* as shown in Figure 2.1. Consider the labeling of North-East edges of *Q* (marked by red) $a_{ij} \to a_{i-1,j}$ by the simple reflection s_{i-j} as shown in the right in Figure 3.1 for $n \geq i > j \geq 1$.

Having done this, we give a procedure to associate a word w(P,Q) in the following two steps. Simultaneously, we will do an example given in Figure 3.2 for a better understanding of the procedure. Consider the left GT-pattern is P and the right GT-pattern is Q in Figure 3.2.

Procedure 1. (1) First read only those labels in P for which $a_{i-1,j} = a_{i,j+1}$ for $n \ge i > j \ge 1$, from bottom to top and left to right, write them from left to right and denote this word by $\mathfrak{f}(P)$.

In the example of Figure 3.2, we read simple reflections along the blue path which moves from bottom to top and the resulting word is $f(P) = s_1 s_2 s_4 s_2 s_1 s_1$.

(2) Next, read only those labels in Q for which $a_{i,j} = a_{i-1,j}$ for $n \ge i > j \ge 1$, from top to bottom and left to right, write them from left to right and denote this word by $\mathfrak{i}(Q)$.

In the example of Figure 3.2, we read simple reflections along the red path which moves from top to bottom and the resulting word is $i(Q) = s_1 s_3 s_2 s_4 s_2$.

The associated word to the pair (P,Q) is $w(P,Q) := \mathfrak{f}(P)\mathfrak{i}(Q)$. In our example of Figure 3.2 we have $w(P,Q) = s_1s_2s_4s_2s_1s_1 \ s_1s_3s_2s_4s_2$.

The permutation associated to the pair (P, Q) is:

$$\mathfrak{p}(P,Q) := \ast(w(P,Q)) \cdot w_0 \tag{3.3}$$

where $*(\cdot)$ is the Demazure product (see 3.1.1) and w_0 is the longest length element in S_n . In our example $*(w(P,Q)) = s_1 s_4 s_2 s_1 s_3 s_4 s_2$ and $\mathfrak{p}(P,Q) = s_2 s_1 s_3$.



FIGURE 3.2. Associated word is $w(P,Q) = s_1 s_2 s_4 s_2 s_1 s_1 s_3 s_2 s_4 s_2$.

The following is an immediate Corollary of theorem 3.2.1.

Corollary 3.3.1. For given $\lambda, \mu \in P^+$, let $\operatorname{GT}_{\mathbb{Z}}(\lambda + \mu)$ be the collection of all pairs (P, Q)in $\operatorname{GT}_{\mathbb{Z}}(\lambda) \times \operatorname{GT}_{\mathbb{Z}}(\mu)$ such that the word w(P, Q) contains w_0 as a subword then,

char
$$V(\lambda + \mu) = \sum_{(P,Q)\in \operatorname{GT}_{\mathbb{Z}}(\lambda+\mu)} \exp^{wt(P)+wt(Q)}$$
.

4. Lakshmibai-Seshadri path model for the KK-modules

4.1. We recall the Lakshmibai-Seshadri (LS) path model from [8]. LS path introduced by Littelmann [10], it is a piecewise linear path π : $[0,1] \to P_{\mathbb{R}}$ such that $\pi(0) = 0$ and $\pi(1) \in P$, where $P_{\mathbb{R}}$ is the real vector space $P \otimes_{\mathbb{Z}} \mathbb{R}$. Let $\lambda \in P^+$, an LS path π of shape λ is given by a sequence

$$\pi = (w_1 > w_2 > \dots > w_r; 0 < a_1 < a_2 < \dots < a_r = 1)$$

which satisfies "chain conditions", where $w_i \in W/W_{\lambda}$ and W_{λ} is the stabilizer of λ . The cosets w_1 and w_r are called the initial direction and final direction of π respectively. Let P_{λ} denote the set of all LS paths of shape λ .

Let $\pi \otimes \pi'$ denote the concatenation of two LS paths π and π' and

$$P_{\lambda} \circledast P_{\mu} = \{ \pi \circledast \pi' \mid \pi \in P_{\lambda}, \ \pi' \in P_{\mu} \}.$$

Note that we can identify $P_{\lambda} \otimes P_{\mu}$ with $P_{\lambda} \times P_{\mu}$ as a set by assuming $\pi \otimes \pi' = (\pi, \pi')$. We associate a Weyl group element $\mathfrak{w}(\pi \otimes \pi')$ to each element $\pi \otimes \pi' \in P_{\lambda} \otimes P_{\mu}$ as follows (see [8], section 3.1):

$$\mathfrak{w}(\pi \circledast \pi') := \min\{W_{\lambda} \ I(\tau^{-1}) \ \varphi \ W_{\mu}\}$$

$$(4.1)$$

where τ is any representative of the final direction of π and φ is any representative of the initial direction of π' .

Definition 4.1.1 (KK set [8]). For given $\lambda, \mu \in P^+$ and $w \in W$, the KK set defined as follows:

$$P(\lambda, w, \mu) = \{ \pi \circledast \pi' \in P_{\lambda} \circledast P_{\mu} \mid \mathfrak{w}(\pi \circledast \pi') \le w \}.$$

$$(4.2)$$

The following theorem shows that the KK set $P(\lambda, w, \mu)$ is a path model for the KK-module $K(\lambda, w, \mu)$.

Theorem 4.1.2 ([8]). Let \mathfrak{g} be a symmetrizable Kac-Moody Lie algebra which is either symmetric or of finite type, then the KK set $P(\lambda, w, \mu)$ is a path model for the KK-module $K(\lambda, w, \mu)$ in the sense that

char
$$K(\lambda, w, \mu) = \sum_{\pi \circledast \pi' \in P(\lambda, w, \mu)} \exp^{\pi(1) + \pi'(1)},$$
 (4.3)

and $K(\lambda, w, \mu)$ is crystal operators e_i, f_i invariant.

4.2. We will simplify the definition of the Weyl group element $\mathfrak{w}(\pi \otimes \pi')$. First, we recall some corollaries and lemmas from [8] which are needed to prove this simplification.

Lemma 4.2.1 ([8], Corollary 2.5). Let K be a subset of Weyl group W that admits a unique minimal element say u. Then, for any two elements σ_1 and σ_2 in W, the set $I(\sigma_1)KI(\sigma_2)$ has a unique minimal element and this element is the unique minimal element in $I(\sigma_1)uI(\sigma_2)$.

Lemma 4.2.2 ([8], Remark 2.4). Let σ and w be elements of W. The set $\{I(\sigma)w\}$ has a unique maximal element and

$$max \{ I(\sigma)w \} = \sigma * w.$$

Lemma 4.2.3. Let K be a subset of W which admits a unique minimum, then

$$min \{u \mid u \in K\} = (max \{uw_0 \mid u \in K\}) \cdot w_0$$

where w_0 is the longest element in W.

The above lemma follows since $u \leq v$ if and only if $vw_0 \leq uw_0$ in the Bruhat order.

Lemma 4.2.4 ([8], Remark 2.9). Let w and w' be two element in W then, I(w)I(w') = I(w * w'). Moreover, if $\ell(ww') = \ell(w) + \ell(w')$ then I(w)I(w') = I(ww'), where $\ell(\cdot)$ is the length function on W.

The following proposition is a simplification of $\mathfrak{w}(\pi \otimes \pi')$ defined in equation (4.1).

Proposition 4.2.5. For given an element $\pi \otimes \pi' \in P_{\lambda} \otimes P_{\mu}$, the Weyl group element associated in the equation (4.1) is:

$$\mathfrak{w}(\pi \circledast \pi') = (\widehat{\tau}^{-1} * \widetilde{\varphi} w_0) \cdot w_0 \tag{4.4}$$

where $\hat{\tau}$ is the maximal representative of the final direction of π and $\tilde{\varphi}$ is the minimal representative of the initial direction of π' .

Proof. For given $\pi \otimes \pi' \in P_{\lambda} \otimes P_{\mu}$ the permutation $\mathfrak{w}(\pi \otimes \pi')$ defined in equation (4.1) is

$$\mathfrak{w}(\pi \otimes \pi') := \min\{W_{\lambda} \ I(\tau^{-1}) \ \varphi \ W_{\mu}\}$$

$$(4.5)$$

where τ is any representative of the final direction of π and φ is any representative of the initial direction of π' . Let $w_{0\lambda}$ be the longest element in the stabilizer W_{λ} . Then equation (4.5) can be written as follows:

$$\mathfrak{w}(\pi \otimes \pi') := \min\{I(w_{0\lambda}) \ I(\tau^{-1}) \ \varphi \ W_{\mu}\}.$$
(4.6)

By lemma 4.2.1 we can write equation (4.6) as

$$\min\{I(w_{0\lambda}) \ I(\tau^{-1}) \ \varphi \ W_{\mu}\} = \min\{I(w_{0\lambda}) \ \min\{\ I(\tau^{-1}) \ \min\{\varphi \ W_{\mu}\}\}\}$$
(4.7)

$$= \min\{I(w_{0\lambda}) \min\{I(\tau^{-1}) \widetilde{\varphi}\}\}$$
(4.8)

$$= \min\{I(w_{0\lambda}) \ I(\tau^{-1}) \ \widetilde{\varphi}\}, \tag{4.9}$$

since $\min\{\varphi W_{\mu}\} = \widetilde{\varphi}$. By lemma 4.2.4, equation (4.9) can be written as follows,

$$\min\{I(w_{0\lambda}) \ I(\tau^{-1}) \ \widetilde{\varphi}\} = \min\{I(w_{0\lambda} * \tau^{-1}) \ \widetilde{\varphi}\}.$$
(4.10)

We can consider τ is the minimal representative of coset τW_{λ} then $\hat{\tau} = \tau w_{0\lambda}$ is the maximal coset representative of τW_{λ} . Then, equation (4.10) is:

$$\min\{I(w_{0\lambda} * \tau^{-1}) \ \widetilde{\varphi}\} = \min\{I(w_{0\lambda}\tau^{-1}) \ \widetilde{\varphi}\} = \min\{I((\tau w_{0\lambda})^{-1}) \ \widetilde{\varphi}\} = \min\{I(\widehat{\tau}^{-1}) \ \widetilde{\varphi}\}.$$

By lemma 4.2.3 $\min\{I(\hat{\tau}^{-1}) \ \widetilde{\varphi}\} = (\max\{I(\hat{\tau}^{-1}) \ \widetilde{\varphi}w_0\}) \cdot w_0$, and by lemma 4.2.2 we have $\max\{I(\hat{\tau}^{-1}) \ \widetilde{\varphi}w_0\} = \hat{\tau}^{-1} * \widetilde{\varphi}w_0.$

5. Demazure crystals and Kogan faces

5.1. We will recall the definition of *crystal* and *crystal isomorphism* from [4]. Let \mathfrak{g} be a symmetrizable Kac-Moody Lie algebra, \mathfrak{h} be a Cartan subalgebra, and \mathfrak{h}^* be the dual subspace of \mathfrak{h} . Denote by $\langle \cdot, \cdot \rangle : \mathfrak{h}^* \times \mathfrak{h} \to \mathbb{C}$ the canonical pairing and by P the weight lattice of \mathfrak{g} . Let $\{\alpha_i \mid i \in I\} \subset \mathfrak{h}^*$ be the set of simple roots and $\{\alpha_i^{\vee} \mid i \in I\} \subset \mathfrak{h}$ be the set of simple coroots, where $I = \{1, 2, \ldots, n-1\}$ is an indexing of the vertices of the Dynkin diagram.

Definition 5.1.1 (Crystal [4]). A crystal is a set \mathfrak{B} equipped with the following maps:

- $wt: \mathfrak{B} \to P$.
- $\psi_i : \mathfrak{B} \to \mathbb{Z} \cup \{-\infty\} \text{ and } \varphi_i : \mathfrak{B} \to \mathbb{Z} \cup \{-\infty\} \text{ for } i \in I.$
- $e_i: \mathfrak{B} \to \mathfrak{B} \cup \{0\}$ and $f_i: \mathfrak{B} \to \mathfrak{B} \cup \{0\}$ for $i \in I$.

where $-\infty$ and 0 are additional elements that are not contained in \mathbb{Z} and \mathfrak{B} , respectively, such that the following holds: for $i \in I$ and $b \in \mathfrak{B}$,

 $\begin{array}{l} (1) \ \varphi_i(b) = \psi_i(b) + \langle wt(b), \alpha_i^{\vee} \rangle. \\ (2) \ wt(e_i(b)) = wt(b) + \alpha_i, \ \psi_i(e_i(b)) = \psi_i(b) - 1 \ and \ \varphi_i(e_i(b)) = \varphi_i(b) + 1, \ if \ e_i(b) \in \mathfrak{B}. \\ (3) \ wt(f_i(b)) = wt(b) - \alpha_i, \ \psi_i(f_i(b)) = \psi_i(b) + 1 \ and \ \varphi_i(f_i(b)) = \varphi_i(b) - 1, \ if \ f_i(b) \in \mathfrak{B}. \\ (4) \ e_i(b) = b' \ for \ b' \in \mathfrak{B}, \ if \ and \ only \ if \ f_i(b') = b \ for \ b \in \mathfrak{B}. \\ (5) \ e_i(b) = f_i(b) = 0, \ if \ \varphi_i(b) = -\infty. \end{array}$

Definition 5.1.2 (Crystal Isomorphism [4]). Let \mathfrak{B}_1 and \mathfrak{B}_2 be two crystals. A map ϕ : $\mathfrak{B}_1 \cup \{0\} \to \mathfrak{B}_2 \cup \{0\}$ is called crystal isomorphism if ϕ is bijective and satisfies the followings:

- (1) $\phi(0) = 0$.
- (2) $wt(b) = wt(\phi(b)), \psi_i(b) = \psi_i(\phi(b)) \text{ and } \varphi_i(b) = \varphi_i(\phi(b)) \text{ for all } i \in I \text{ and } b \in \mathfrak{B}_1 \text{ such that } \phi(b) \in \mathfrak{B}_2.$

(3)
$$\phi(e_i(b)) = e_i(\phi(b))$$
 and $\phi(f_i(b)) = f_i(\phi(b))$ for all $b \in \mathfrak{B}_1$.

where we set $e_i(\phi(b)) = f_i(\phi(0)) = 0$, if $\phi(b) = 0$.

Littelmann has shown that the set of LS paths P_{μ} of shape μ for $\mu \in P^+$ is a crystal. For the definition of crystal operators e_i and f_i for $i \in I$ on LS paths we give references of Littelmann [10] and [11] which he calls the *root operators*.

5.2. Crystal operators on GT patterns. For a given integer partition $\mu = (\mu_1, \mu_2, \dots, \mu_n)$ the set of integral GT-patterns $\operatorname{GT}_{\mathbb{Z}}(\mu)$ of shape μ is also a crystal. We will recall the definition of crystal operators e_i and f_i for $i \in I = \{1, 2, \dots, n-1\}$ on $\operatorname{GT}_{\mathbb{Z}}(\mu)$.

Let P be the following GT pattern of shape μ .



where $a_{n,i} = \mu_i$ for i = 1, 2, ..., n.

For a fixed $i \in I$ we will recall the definition of e_i and f_i . Define the integers d_t for $1 \leq t \leq i+1$ inductively as follows, set $d_1 := a_{i,1} - a_{i+1,1}$ and

$$d_t := d_{t-1} + a_{i,t-1} + a_{i,t} - a_{i-1,t-1} - a_{i+1,t} \text{ for } 1 < t \le i+1,$$

where assume that $a_{k,k+1} = 0$ for $k \in \{i - 1, i\}$. Note that $d_1 \leq 0$ since $a_{i,1} \leq a_{i+1,1}$. Let $d = \min\{d_1, d_2, \ldots, d_{i+1}\}, m$ be the smallest integer such that $d_m = d$ and M be the largest integer such that $d_M = d$.

The action of e_i on P defined as follows:

- (1) if $d_m = 0$ then, $e_i(P) = 0$.
- (2) if $d_m < 0$ then, $e_i(P) = P'$ where P' have all entries same as P except the entry a_{im} will change to $a_{im} + 1$.

The action of f_i on P defined as follows,

- (1) if M = i + 1 then, $f_i(P) = 0$.
- (2) if M < i + 1 then, $f_i(P) = P'$ where P' have all entries same as P except the entry a_{iM} will change to $a_{iM} 1$.

The following proposition 5.2.1 is a well-known fact and a variation of proof can be seen in [11].

Proposition 5.2.1. There exists a crystal isomorphism $\phi : \operatorname{GT}_{\mathbb{Z}}(\mu) \to P_{\mu}$.

5.3. For an element $G \in \operatorname{GT}_{\mathbb{Z}}(\mu)$, We define the initial and final direction of G as the same as LS path $\phi(G)$. There exists a unique element G^0_{μ} in $\operatorname{GT}_{\mathbb{Z}}(\mu)$ such that $e_i(G^0_{\mu}) = 0$ for all $i \in I$ and a unique element G^*_{μ} in $\operatorname{GT}_{\mathbb{Z}}(\mu)$ such that $f_i(G^*_{\mu}) = 0$ for all $i \in I$. The element G^0_{μ} is called the highest-weight element and G^*_{μ} is called the lowest-weight element of $\operatorname{GT}_{\mathbb{Z}}(\mu)$.

Definition 5.3.1. Given $w \in S_n$, fix a reduced decomposition $w = s_{i_1}s_{i_2}\cdots s_{i_k}$. The set

$$Dem(\mu, w) := \{ f_{i_1}^{m_1} f_{i_2}^{m_2} \cdots f_{i_k}^{m_k} (G^0_\mu) \mid m_j \ge 0; \ 1 \le j \le k \} \subseteq GT_{\mathbb{Z}}(\mu)$$

is called a Demazure crystal.

Definition 5.3.2. Given $w \in S_n$, fix a reduced decomposition $ww_0 = s_{j_1}s_{j_2}\cdots s_{i_t}$. The set

$$Dem(\mu, w)^{op} := \{ e_{j_1}^{m_1} e_{j_2}^{m_2} \cdots e_{j_t}^{m_t} G_{\mu}^* \mid m_s \ge 0; \quad 1 \le s \le t \} \subseteq GT_{\mathbb{Z}}(\mu)$$

is called an opposite Demazure crystal.

The following two propositions determine the minimal representative of the initial direction and the maximal representative of the final direction of a given $G \in \operatorname{GT}_{\mathbb{Z}}(\mu)$ in terms of Demazure and opposite Demazure crystals respectively.

Proposition 5.3.3. Let $G \in GT_{\mathbb{Z}}(\mu)$ and $S = \{w \in S_n \mid G \in Dem(\mu, w)\}$. Then the set S has a unique minimal element and **min** S is the minimal representative of the initial direction of G.

Proof. From work of Littelmann [10, §5.2], Demazure crystals can be defined in terms of initial direction as follows:

$$Dem(\mu, w) = \{T \in GT_{\mathbb{Z}}(\mu) \mid init(T) \le wS_{\mu}\}$$
(5.1)

where $\operatorname{init}(T)$ denote the initial direction of T. Let $\widetilde{\varphi}$ be the minimal representative of the initial direction of G. From the above expression (5.1) of Demazure crystal, clearly $\widetilde{\varphi} \in S$ and for any $w \in S$ we have $\operatorname{init}(G) = \widetilde{\varphi}S_{\mu} \leq wS_{\mu}$ which implies that $\widetilde{\varphi} \leq w$.

Proposition 5.3.4. Let $G \in GT_{\mathbb{Z}}(\mu)$ and $\overline{S} = \{w \in S_n \mid G \in Dem(\mu, w)^{op}\}$. Then the set \overline{S} has a unique maximal element and $\max \overline{S}$ is the maximal representative of the final direction of G.

Proof. Similar to Demazure crystals, opposite Demazure crystals can be defined in terms of final direction as follows:

$$\operatorname{Dem}(\mu, w)^{op} = \{T \in \operatorname{GT}_{\mathbb{Z}}(\mu) \mid \operatorname{fin}(T) \ge w S_{\mu}\}$$
(5.2)

where fin(T) denote the final direction of T. Let $\hat{\tau}$ be the maximal representative of the final direction of G. From the above expression (5.2) of opposite Demazure crystal, clearly $\hat{\tau} \in \overline{S}$ and for any $w \in \overline{S}$ we have fin(G) = $\hat{\tau}S_{\mu} \geq wS_{\mu}$ which implies that $\hat{\tau} \geq w$.

Next, we will introduce Kogan faces and dual Kogan faces which establish a connection between Demazure Crystals and opposite Demazure crystals, for more details see [2].

5.4. Kogan faces of GT polytope. Let $\mu \in P^+$ and $GT(\mu)$ be the GT polytope of shape μ defined in the section 2. Fix a subset $F \subseteq \{(i, j) : n \ge i > j \ge 1\}$. Consider the face of $GT(\mu)$ obtained by setting North-East inequalities $NE_{ij} = a_{ij} - a_{i-1,j} = 0$ for $(i, j) \in F$ and leaving all other inequalities untouched. We call this the Kogan face $K(\mu, F)$. To each pair i > j, associate the simple transposition $s_{i-j} \in S_n$. We list the elements of F in lexicographically increasing order: (i, j) precedes $(i', j') \Leftrightarrow$ either i < i', or i = i' and j < j'. Denote the product of the corresponding s_{i-j} in this order by $\sigma(F)$. If len $\sigma(F) = |F|$, i.e., this word is reduced, we say that $K(\mu, F)$ is reduced Kogan face and set [2, Definition 5.1]:

$$\varpi(F) = w_0 \,\sigma(F) \,w_0.$$

For $w \in S_n$, let $K(\mu, w) := \bigcup K(\mu, F)$, the union over reduced Kogan face $K(\mu, F)$ for which $\varpi(F) = w$. Denote the set of integral points in $K(\mu, w)$ by $K_{\mathbb{Z}}(\mu, w)$.

We state the following proposition by Fujita which establishes a connection between Kogan faces and Demazure crystals.

Proposition 5.4.1 ([2], Corollary 5.19). $K_{\mathbb{Z}}(\mu, w_0 w) = Dem(\mu, w)$.

5.5. Dual Kogan faces of GT polytope. Fix a subset $F \subseteq \{(i, j) : n \ge i > j \ge 1\}$. Consider the face of $GT(\mu)$ obtained by setting South-East inequalities $SE_{ij} = a_{i-1,j} - a_{i,j+1} = 0$ for $(i, j) \in F$ and leaving all other inequalities untouched. We call this the *dual* Kogan face $\overline{K}(\mu, F)$. To each pair $(i, j) \in \{(i, j) : n \ge i > j \ge 1\}$, associate the simple transposition $s_j \in S_n$. We consider a total order on pairs (i, j) defined by (i, j) precedes $(i', j') \Leftrightarrow$ either i < i', or i = i' and j > j'. We list the elements of F in the increasing order relative to this total order. Denote the product of the corresponding s_j in this order by $\overline{\sigma}(F)$. If len $\overline{\sigma}(F) = |F|$, i.e., this word is reduced, we say that $\overline{K}(\mu, F)$ is a reduced dual Kogan face and set [2, Definition 5.1]:

$$\overline{\varpi}(F) = w_0 \,\overline{\sigma}(F) \, w_0.$$

For $w \in S_n$, let $\overline{\mathrm{K}}(\mu, w) := \bigcup \overline{\mathrm{K}}(\mu, F)$, the union over reduced dual Kogan faces $\overline{\mathrm{K}}(\mu, F)$ for which $\overline{\varpi}(F) = w$. Denote the set of integral points in $\overline{\mathrm{K}}(\mu, w)$ by $\overline{\mathrm{K}}_{\mathbb{Z}}(\mu, w)$.

We state the following proposition by Fujita which establishes a connection between dual Kogan faces and opposite Demazure crystals.

Proposition 5.5.1 ([2], Corollary 5.2). $\overline{\mathrm{K}}_{\mathbb{Z}}(\mu, w_0 w w_0) = \mathrm{Dem}(\mu, w)^{op}$.

5.6. BiKogan faces. Let λ, μ be two elements of P^+ . Consider the GT polytopes $\operatorname{GT}(\lambda)$ and $\operatorname{GT}(\mu)$, with a typical element of the former given by $(a_{ij})_{n \geq i \geq j \geq 1}$ and that of the latter given by $(b_{ij})_{n \geq i \geq j \geq 1}$ as in figure 2.1. Fix subsets F, F' of the set $\{(i, j) : n \geq i > j \geq 1\}$. Consider the face of the Cartesian product $\operatorname{GT}(\lambda) \times \operatorname{GT}(\mu)$ of GT polytopes obtained by setting $a_{i-1,j} - a_{i,j+1} = 0$ for all $(i, j) \in F$ and $b_{ij} - b_{i-1,j} = 0$ for all $(i, j) \in F'$ and leaving all other inequalities untouched. Observe that the former equalities concern SE differences, while the latter are NE differences. We call this face a *BiKogan face* and denote it by $K_{\lambda,\mu}(F, F')$.

Define an element $\varpi(F, F') \in S_n$ associated to the pair (F, F') as follows: $\varpi(F, F') = \overline{\sigma}(F)^{-1} \cdot \sigma(F')$. We say the BiKogan face $K_{\lambda,\mu}(F, F')$ is reduced if $\operatorname{len}(\varpi(F, F')) = |F| + |F'|$. For a given $w \in S_n$, let

$$\mathbf{K}_{\lambda,\mu}(w) := \cup \mathbf{K}_{\lambda,\mu}(F, F'),$$

the union being taken over all reduced BiKogan faces for which $\varpi(F, F') = w$.

Lemma 5.6.1. Let $K_{\lambda,\mu}(F, F')$ be reduced for a given pair (F, F'). If $(P, Q) \in K_{\lambda,\mu}(F, F')$ then $\varpi(F, F') \leq *(w(P, Q))$.

Proof. Let $(P,Q) \in K_{\lambda,\mu}(F,F')$ and $P = (a_{ij})_{n \ge i \ge j \ge 1}$, $Q = (b_{ij})_{n \ge i \ge j \ge 1}$. Then F is a subset of $\{(i,j) \mid a_{i-1,j} = a_{i,j+1}\}$ and F' is a subset of $\{(i,j) \mid b_{ij} = b_{i-1,j}\}$. From the item (1) and (2) of Procedure 1 observe that $\sigma(F')$ and $\overline{\sigma}(F)^{-1}$ are subwords of $\mathfrak{i}(Q)$ and $\mathfrak{f}(P)$ respectively. Then we have $\varpi(F,F') = \overline{\sigma}(F)^{-1}\sigma(F)$ is a subword of the word $\mathfrak{f}(P)\mathfrak{i}(Q) = w(P,Q)$. Since $K_{\lambda,\mu}(F,F')$ is reduced BiKogan face then $\varpi(F,F') \le *(w(P,Q))$.

We state the following theorem which gives another description of $GT(\lambda, w, \mu)$ in terms of BiKogan faces.

Theorem 5.6.2. Let λ, μ be two elements in P^+ and w be a permutation in S_n then $K_{\lambda,\mu}(ww_0) = GT(\lambda, w, \mu)$. In particular, the set of integral points in $K_{\lambda,\mu}(ww_0)$ is a combinatorial model for the KK-module $K(\lambda, w, \mu)$.

Proof. Let $(P,Q) \in K_{\lambda,\mu}(ww_0)$. Then there exist subsets F, F' of the set $\{(i,j) \mid n \geq i > j \geq 1\}$ such that the BiKogan face $K_{\lambda,\mu}(F,F')$ is reduced, $\varpi(F,F') = ww_0$ and $(P,Q) \in K_{\lambda,\mu}(F,F')$. From the lemma 5.6.1, $\varpi(F,F') \leq *(w(P,Q))$. Hence $\mathfrak{p}(P,Q) = *(w(P,Q))w_0 \leq w$ which implies that $(P,Q) \in \mathrm{GT}(\lambda, w, \mu)$.

For the converse, let $(P,Q) \in \operatorname{GT}(\lambda, w, \mu)$ then $\mathfrak{p}(P,Q) = *(w(P,Q))w_0 \leq w$. Since $*(w(P,Q)) \geq ww_0$, then we can choose subsets F, F' of the set $\{(i,j) \mid n \geq i > j \geq 1\}$ such that the BiKogan face $K_{\lambda,\mu}(F,F')$ is reduced, $(P,Q) \in K_{\lambda,\mu}(F,F')$ and $\varpi(F,F') = ww_0$. Hence $(P,Q) \in K_{\lambda,\mu}(ww_0)$.

6. Proof of the theorem 3.2.1

6.1. We state the following two lemmas 6.1.1 and 6.1.2 which are the main ingredients to prove theorem 3.2.1. For a given $P \in \operatorname{GT}_{\mathbb{Z}}(\mu)$, these lemmas provide a procedure to determine the minimal representative of the initial direction and the maximal representative of the final direction of $\phi(P)$ respectively.

Lemma 6.1.1. Let i(P) be the word associated in item (2) of the procedure 1 then, the minimal representative of the initial direction of the LS path $\phi(P)$ is $*(i(P))w_0$.

Lemma 6.1.2. Let $\mathfrak{f}(P)$ be the word associated in item (1) of the procedure 1 then, the maximal representative of the final direction of the LS path $\phi(P)$ is $(*(\mathfrak{f}(P)))^{-1}$.

We will give a proof of lemma 6.1.1 and 6.1.2 later in this section.

The following two corollaries are immediate results of the above lemmas which gives another characterization of the Demazure crystals and opposite Demazure crystals.

Corollary 6.1.3. With the notation in the lemma 6.1.1

$$\operatorname{Dem}(\mu, w) = \{ P \in \operatorname{GT}_{\mathbb{Z}}(\mu) \mid \ast(\mathfrak{i}(P))w_0 \le w \}.$$

Corollary 6.1.4. With the notation in the lemma 6.1.2

 $Dem(\mu, w)^{op} = \{ P \in GT_{\mathbb{Z}}(\mu) \mid w_0(*(\mathfrak{f}(P)))^{-1} \le w_0 w \}.$

6.2. The next proposition will give a connection between the LS path model and the GT model for the KK-modules. We will use it to prove theorem 3.2.1.

Proposition 6.2.1. The following diagram commutes



In other words

$$\mathfrak{p}(P,Q) = \mathfrak{w}(\phi(P) \otimes \phi(Q)) \text{ for all } (P,Q) \in \mathrm{GT}_{\mathbb{Z}}(\lambda) \times \mathrm{GT}_{\mathbb{Z}}(\mu)$$
(6.1)

where $\phi \times \phi$ is the crystal isomorphism given in the proposition 5.2.1, \mathfrak{p} is defined in the equation (3.3) and \mathfrak{w} is defined in the equation (4.1).

Proof. For a given pair $(P,Q) \in \operatorname{GT}_{\mathbb{Z}}(\lambda) \times \operatorname{GT}_{\mathbb{Z}}(\mu)$, by lemma 4.2.5 we know that

$$\mathfrak{w}(\phi(P) \circledast \phi(Q)) = (\widehat{\tau}^{-1} * \widetilde{\varphi} w_0) \cdot w_0 \tag{6.2}$$

where $\hat{\tau}$ is the maximal representative of the final direction of $\phi(P)$ and $\tilde{\varphi}$ is the minimal representative of the initial direction of $\phi(Q)$. The association of Weyl group element to the

pair (P, Q) is given by the equation (3.3) is

$$\mathfrak{p}(P,Q) = \ast(w(P,Q)) \cdot w_0. \tag{6.3}$$

From the equations (6.2) and (6.3), it is enough to prove that

$$\widehat{\tau}^{-1} * \widetilde{\varphi} w_0 = *(w(P,Q)). \tag{6.4}$$

We have $w(P,Q) = \mathfrak{f}(P)\mathfrak{i}(Q)$ in the procedure 1 and by lemma 3.1.3

$$*(w(P,Q)) = *(\mathfrak{f}(P)\mathfrak{i}(Q)) = *\mathfrak{f}(P) * (*\mathfrak{i}(Q)).$$

Finally by lemma 6.1.2 and lemma 6.1.1 we have

$$*\mathfrak{f}(P)*(*\mathfrak{i}(Q))=\widehat{\tau}^{-1}*\widetilde{\varphi}w_0.$$

Finally, we can prove the theorem 3.2.1.

Proof of theorem 3.2.1. It is clear from the above equation (6.1) that

$$GT_{\mathbb{Z}}(\lambda, w, \mu) = \{ (P, Q) \in GT_{\mathbb{Z}}(\lambda) \times GT_{\mathbb{Z}}(\mu) \mid \phi(P) \circledast \phi(Q) \in P(\lambda, w, \mu) \}$$
(6.5)

Combining theorem 4.1.2, equation (6.5) and $\phi \times \phi : \operatorname{GT}_{\mathbb{Z}}(\lambda) \times \operatorname{GT}_{\mathbb{Z}}(\mu) \mapsto P_{\lambda} \otimes P_{\mu}$ is a crystal isomorphism proves the theorem 3.2.1.

6.3. Next, we will prove lemmas 6.1.1 and lemma 6.1.2.

Proof of lemma 6.1.1. Let $P = (a_{ij})_{n \ge i \ge j \ge 1}$ be an integral GT-pattern of shape μ . Define the set $\mathfrak{F}_P := \{(i, j) | a_{ij} - a_{i-1,j} = 0 \text{ for } n \ge i > j \ge 1\}$. We associate a word $w(\mathfrak{F}_P)$ corresponding to \mathfrak{F}_P as follows: write s_{i-j} from left to right for all $(i, j) \in \mathfrak{F}_P$ in the total order defined in the section 5.4. Note that $\mathfrak{i}(P) = w(\mathfrak{F}_P)$.

From the proposition 5.4.1, for any $F \subseteq \mathfrak{F}_P$ such that $K(\mu, F)$ is a reduced Kogan face then $P \in K_{\mathbb{Z}}(\mu, F)$ and hence $P \in \text{Dem}(\mu, \sigma(F)w_0)$. Further, if $P \in \text{Dem}(\mu, w)$ then there exist a subset $F \subseteq \mathfrak{F}_P$ such that $K(\mu, F)$ is reduced, $w = \sigma(F)w_0$ and $P \in K_{\mathbb{Z}}(\mu, F)$.

There exist a subset $F^0 \subseteq \mathfrak{F}_P$ such that $\mathrm{K}(\mu, F^0)$ is reduced, $\sigma(F^0) = *(w(\mathfrak{F}_P))$ and $P \in \mathrm{Dem}(\mu, \sigma(F^0)w_0)$. By lemma 3.1.2 $\sigma(F^0) = *(w(\mathfrak{F}_P))$ is the unique maximal element of the set

 $\{\sigma(F) \mid F \subseteq \mathfrak{F}_P \text{ and } \mathrm{K}(\mu, F) \text{ is reduced}\}.$

Hence by above arguments $\sigma(F^0)w_0$ is the unique minimal permutation of the set

$$\{w \mid P \in \operatorname{Dem}(\mu, w)\}.$$

Now we appeal to the proposition 5.3.3 to conclude that $\sigma(F^0)w_0 = *(w(\mathfrak{F}_P))w_0$ is the minimal representative of the initial direction of $\phi(P)$.

Proof of lemma 6.1.2. The proof of this lemma is exactly similar to the proof of the above lemma 6.1.1. However, we will record the proof below for completeness. Le $P = (a_{ij})_{n \ge i > j \ge 1}$ be an integral GT-pattern of shape μ . Define the set $\mathfrak{H}_P := \{(i, j) | a_{i-1,j} - a_{i,j+1} = 0 \text{ for } n \ge i > j \ge 1\}$. We associate a word $w(\mathfrak{H}_P)$ corresponding to \mathfrak{H}_P as follows: write s_j from left to right for all $(i, j) \in \mathfrak{H}_P$ in the total order defined in the section 5.5. Note that $\operatorname{rev}(\mathfrak{f}(P)) = w(\mathfrak{H}_P)$ where $\operatorname{rev}(\mathfrak{f}(P))$ is the word obtained by writing the word $\mathfrak{f}(P)$ in the reverse order, then $(*(\mathfrak{f}(P)))^{-1} = *(w(\mathfrak{H}_P)).$

From the proposition 5.5.1, for any $F \subseteq \mathfrak{H}_P$ such that $\overline{\mathrm{K}}(\mu, F)$ is a reduced dual Kogan face then $P \in \overline{\mathrm{K}}_{\mathbb{Z}}(\mu, F)$ and hence $P \in \mathrm{Dem}(\mu, \overline{\sigma}(F))^{op}$. Further, if $P \in \mathrm{Dem}(\mu, w)^{op}$ then there exist a subset $F \subseteq \mathfrak{H}_P$ such that $\overline{\mathrm{K}}(\mu, F)$ is reduced, $w = \overline{\sigma}(F)$ and $P \in \overline{\mathrm{K}}_{\mathbb{Z}}(\mu, F)$.

There exist a subset $\overline{F'} \subseteq \mathfrak{H}_P$ such that $\overline{K}(\mu, \overline{F'})$ is reduced, $\overline{\sigma}(\overline{F'}) = *(w(\mathfrak{H}_P))$ and $P \in \text{Dem}(\mu, \overline{\sigma}(\overline{F'}))$. By lemma 3.1.2 $\overline{\sigma}(\overline{F'}) = *(w(\mathfrak{H}_P))$ is the unique maximal element of the set

 $\{\overline{\sigma}(F) \mid F \subseteq \mathfrak{H}_P \text{ and } \overline{\mathrm{K}}(\mu, F) \text{ is reduced}\}.$

Hence by above arguments $\overline{\sigma}(\overline{F'})$ is the unique maximal permutation of the set

$$\{w \mid P \in \operatorname{Dem}(\mu, w)^{op}\}.$$

Now we appeal to the proposition 5.3.4 to conclude that $\overline{\sigma}(\overline{F'}) = *(w(\mathfrak{H}_P))$ is the maximal representative of the final direction of $\phi(P)$.

7. Data Availibility Statement

This manuscript has no associated data.

References

- Deodhar, Vinay V., A splitting criterion for the Bruhat orderings on Coxeter groups, Communications in Algebra, 15, (1987), pp. 1889–1894, ISSN: 0092-7872, URL https://doi.org/10.1080/00927878708823511.
- Fujita, Naoki, Schubert calculus from polyhedral parametrizations of Demazure crystals, Advances in Mathematics, 397, paper number. 108201, 2022, https://doi.org/10.1016/j.aim.2022.108201.
- [3] Gelfand, I. M. and Tsetlin, M. L., Finite-dimensional representations of the group of unimodular matrices, Dokl. Akad. Nauk SSSR, 71, pages:825–828, (1950)
- [4] Kashiwara, Masaki, The crystal base and Littlemann's refined Demazure character formula, Duke Mathematical Journal, 71, pages:839–858, (1993)
- [5] Mikhail Kogan, Schubert Geometry of Flag Varieties and Gelfand-Cetlin Theory, MIT, (2000)
- [6] Kumar, Shrawan, Proof of the Parthasarathy-Ranga Rao-Varadarajan conjecture, Inventiones Mathematicae, 93, (1988), PAGES:117–130
- [7] Kumar, Shrawan, A refinement of the PRV conjecture, Inventiones Mathematicae, 97, (1989), PAGES:305-311, https://doi.org/10.1007/BF01389044.
- [8] M S Kushwaha and K N Raghavan and S Viswanath, A study of Kostant-Kumar modules via Littelmann paths, Advances in Mathematics, 381, 107614, (2021)
- [9] M S Kushwaha and K N Raghavan and S Viswanath, The saturation problem for refined Littlewood-Richardson coefficients, Séminaire Lotharingien de Combinatoire, 85B.52, 12 pp, (2021)
- [10] Littelmann, Peter, A Littlewood-Richardson rule for symmetrizable Kac-Moody algebras, Inventiones Mathematicae, 116, (1994), PAGES:329–346, https://doi.org/10.1007/BF01231564.
- [11] Littelmann, Peter, Paths and root operators in representation theory, Annals of Mathematics. Second Series, 142, (1995), PAGES:499-525
- [12] Mathieu, Olivier, Construction du groupe de Kac-Moody et applications, Comptes Rendus des Séances de l'Académie des Sciences. Série I. Mathématique, **306**, (1988), PAGES:227–230.

DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCE, UNIVERSITY OF DELHI, NEW DELHI - 110007

Email address: mrigendra1540gmail.com, mskushwaha@maths.du.ac.in