

Milnor-Witt motivic decomposition of Stiefel varieties

Keyao Peng

1 Introduction

This paper extends the research presented in [4], which investigated the cohomology groups $H_\eta^*(V_k(\mathbb{A}^n))$ with η -inverted coefficients by employing the \mathbb{A}^1 -Leray spectral sequence [1]. For foundational results and notations relevant to this study, we refer the reader to [4].

In this work, we will initially compute the integral MW-motivic cohomology groups $H_{\text{MW}}^{p,q}(V_k(\mathbb{A}^n))$ associated with Stiefel varieties.

Theorem 1.1 (Theorem 3.4). *Let $i \in N_{n,k}$. The generators α_i are elements of $H_{\text{MW}}^{2i+1,i+1}(V_k(\mathbb{A}^n))$, and for $J \in F_{n,k}$, the generators β_J belong to $H_{\text{MW}}^{d(J)}(V_k(\mathbb{A}^n))$. For any subset $I \subset N_{n,k}$, define*

$$b_I = \prod_{J \in F_{n,k}, J \subset I} \beta_J \prod_{i \in I_T} \alpha_i \in H_{\text{MW}}^{d(I)}(V_k(\mathbb{A}^n)).$$

Then, we establish the following isomorphism of graded $H_{\text{MW}}^{*,*}(K)$ -modules:

$$H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n)) \cong \bigoplus_{I \subset N_{n,k}} (H_{\text{MW}})_I b_I,$$

where

$$(H_{\text{MW}})_I = \begin{cases} H_{\text{MW}}^{*,*}(K), & \text{if } I = I_F, \\ H_M^{*,*}(K), & \text{if } \text{Max}(I_T) \text{ is even,} \\ {}^n H_M^{*,*}(K), & \text{if } \text{Max}(I_T) \text{ is odd.} \end{cases}$$

It is important to note that our computations focus solely on the additive structures; the ring structures are considerably more complex.

Building upon these computations, we will proceed to establish the following MW-motive decomposition:

Theorem 1.2 (Theorem 4.2). *Let $C_\eta(S)$ denote the cone of $\tilde{M}(S) \xrightarrow{\eta} \tilde{M}(S)(1)[1]$ and define the following:*

$$\begin{aligned} \widetilde{\mathbf{HS}}_{2k} &:= \tilde{M}(\mathbb{A}^{2k} \setminus \{0\}) \cong \tilde{M}(S) \oplus \tilde{M}(S)(2k)[4k-1], \\ \widetilde{\mathbf{HS}}_{2k+1} &:= \tilde{M}(S) \oplus C_\eta(S)(2k)[4k] \oplus \tilde{M}(S)(4k+1)[8k]. \end{aligned}$$

Then, for any $i, j \in \mathbb{N}$, we obtain an MW-motivic decomposition of the following forms:

$$\begin{aligned} \tilde{M}(V_{2j}(\mathbb{A}^{2i})) &\cong \widetilde{\mathbf{HS}}_{2i} \otimes \widetilde{\mathbf{HS}}_{2i-1} \otimes \widetilde{\mathbf{HS}}_{2i-3} \otimes \dots \otimes \widetilde{\mathbf{HS}}_{2i-1-2(j-2)} \otimes \tilde{M}(\mathbb{A}^{2i+1-2j} \setminus \{0\}), \\ \tilde{M}(V_{2j+1}(\mathbb{A}^{2i})) &\cong \widetilde{\mathbf{HS}}_{2i} \otimes \widetilde{\mathbf{HS}}_{2i-1} \otimes \widetilde{\mathbf{HS}}_{2i-3} \otimes \dots \otimes \widetilde{\mathbf{HS}}_{2i-1-2(j-1)}, \\ \tilde{M}(V_{2j}(\mathbb{A}^{2i+1})) &\cong \widetilde{\mathbf{HS}}_{2i+1} \otimes \widetilde{\mathbf{HS}}_{2i-1} \otimes \dots \otimes \widetilde{\mathbf{HS}}_{2i+1-2(j-1)}, \\ \tilde{M}(V_{2j+1}(\mathbb{A}^{2i+1})) &\cong \widetilde{\mathbf{HS}}_{2i+1} \otimes \widetilde{\mathbf{HS}}_{2i-1} \otimes \dots \otimes \widetilde{\mathbf{HS}}_{2i+1-2(j-1)} \otimes \tilde{M}(\mathbb{A}^{2i+1-2j} \setminus \{0\}). \end{aligned}$$

Notation

Let $C_\eta(S)$ denote the cone of $\tilde{M}(S) \xrightarrow{\eta} \tilde{M}(S)(1)[1]$. We sometimes use the index of degree $(p, \{q\}) := (p+q, q)$. A cohomology theory $\mathbb{E} \in \mathcal{SH}(K)$ is termed **coconnected** if $\mathbb{E}^{p,\{q\}}(K) = 0$ for $p > 0$.

2 The \mathbb{A}^1 -Thom-Gysin Sequence

We will follow the computation from [2] to determine the total cohomology group $H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n), \mathbb{Z})$ with integral coefficients as a module. The key element in this determination is the \mathbb{A}^1 -Thom-Gysin sequence, which directly results from the \mathbb{A}^1 -Leray spectral sequence.

Theorem 2.1 ([1, Thm 5.2.8]). *Let $f : X \rightarrow B$ be a Zariski locally trivial smooth fibration over a field K , where the fiber F of f is an \mathbb{A}^1 -homology sphere (i.e., $\Sigma^r F \cong \Sigma^p \mathbb{G}_m^{\wedge q}$). Let M be a homotopy module with a ring structure over K , and assume that $H_\delta^i(f_* M_X)$ is K -simple over B . Then there exists an Euler class $e(f) \in H^{p-r+1}(B, M_q)$, such that the Thom-Gysin sequence is given by:*

$$\cdots \rightarrow H^i(B, M_j) \rightarrow H^i(X, M_j) \rightarrow H^{i+r-p}(B, M_{j-q}) \xrightarrow{\cup e(f)} H^{i+1}(B, M_j) \rightarrow \cdots$$

where the connecting morphism is given by the product with $e(f)$.

In fact, we can generalize this to an Atiyah-Hirzebruch type spectral sequence by replacing M with a spectrum $\mathbb{E} \in \mathcal{SH}(S)$, where S is a Noetherian base scheme over the field K .

Proposition 2.2 (\mathbb{A}^1 -Thom-Gysin sequence). *Let $\mathbb{E} \in \mathcal{SH}(S)$ be a ring spectrum, and let $f : V \rightarrow B$ be a rank- n vector bundle that is \mathbb{E} -oriented. Let $X := V \setminus B$ denote the complement of the zero section, which constitutes a sphere bundle. Then, we acquire the Thom-Gysin exact sequence:*

$$\cdots \rightarrow \mathbb{E}^{i,j}(B) \rightarrow \mathbb{E}^{i,j}(X) \xrightarrow{r} \mathbb{E}^{i-(2n-1),j-n}(B) \xrightarrow{\cup e(f)} \mathbb{E}^{i+1,j}(B) \rightarrow \cdots$$

where $e(f) \in \mathbb{E}^{2n,n}(B)$ represents the Euler class of the vector bundle.

Proof. Since $f : V \rightarrow B$ is oriented with respect to \mathbb{E} , we have the Thom isomorphism $\phi : \mathbb{E}^{i-2n,j-n}(B) \xrightarrow{\cong} \mathbb{E}^{i,j}(\text{Th}(V))$, where $\text{Th}(V) := V/X$ is the Thom space. Let $\tau = \phi(1) \in \mathbb{E}^{2n,n}(\text{Th}(V))$ denote the Thom class. Furthermore, there exists a homotopy equivalence $f^* : \mathbb{E}^{i,j}(B) \xrightarrow{\cong} \mathbb{E}^{i,j}(V)$. Let the morphism $\text{Th}(f)^* : \mathbb{E}^{i,j}(B) \xrightarrow{\cong} \mathbb{E}^{i,j}(\text{Th}(V))$ be induced by the projection. We note that for the Thom isomorphism, $\phi(a) = \text{Th}(f)^*(a) \cup \tau$.

Now, consider the Gysin sequences associated with the cofiber sequence $X \rightarrow V \xrightarrow{q} \text{Th}(V)$:

$$\begin{array}{ccccccc} \mathbb{E}^{i,j}(V) & \longrightarrow & \mathbb{E}^{i,j}(X) & \longrightarrow & \mathbb{E}^{i+1,j}(\text{Th}(V)) & \xrightarrow{q^*} & \mathbb{E}^{i+1,j}(V) \\ \uparrow f^* & & \parallel & & \uparrow \phi & & \uparrow f^* \\ \mathbb{E}^{i,j}(B) & \longrightarrow & \mathbb{E}^{i,j}(X) & \longrightarrow & \mathbb{E}^{i-(2n-1),j-n}(B) & \xrightarrow{\partial} & \mathbb{E}^{i+1,j}(B) \end{array}$$

We only need to determine the morphism $\partial := (f^*)^{-1}q^*\phi$. By the properties of the Thom isomorphism, we have

$$\partial(a) = (f^*)^{-1}q^*(\text{Th}(f)^*(a) \cup \tau) = (f^*)^{-1}(f^*(a) \cup q^*(\tau)) = a \cup (f^*)^{-1}q^*(\tau).$$

By the definition of the Euler class, $e(f) := (f^*)^{-1}q^*(\tau)$, we can conclude our proof. \square

Pursuing the diagram further, we can derive the following corollary:

Corollary 2.3. *We maintain the previously established notation. Let $g : \mathbb{A}^n \setminus \{0\} \rightarrow X$ be a fiber embedding induced by the base point $b : pt \rightarrow B$. Then the pullback $g^* : \mathbb{E}^{i,j}(X) \rightarrow \mathbb{E}^{i,j}(\mathbb{A}^n \setminus \{0\})$ is defined as follows:*

$$\mathbb{E}^{i,j}(X) \xrightarrow{r} \mathbb{E}^{i-(2n-1),j-n}(B) \xrightarrow{b^*} \mathbb{E}^{i-(2n-1),j-n}(pt) \xrightarrow{\cup \theta} \mathbb{E}^{i,j}(\mathbb{A}^n \setminus \{0\}),$$

where $\theta \in \mathbb{E}^{2n-1,n}(\mathbb{A}^n \setminus \{0\})$ serves as a generator.

3 MW-Motivic Cohomology Groups of $V_k(\mathbb{A}^n)$

We apply this theorem to the fiber sequences of Stiefel varieties over the field K :

$$\mathbb{A}^{n-k} \setminus \{0\} \rightarrow V_{k+1}(\mathbb{A}^n) \xrightarrow{f_{n,k}} V_k(\mathbb{A}^n).$$

The fiber $\mathbb{A}^{n-k} \setminus \{0\} \cong \Sigma^{n-k-1} \mathbb{G}_m^{\wedge n-k}$ is precisely an \mathbb{A}^1 -sphere and corresponds to a H_{MW} -oriented bundle, which is, in fact, GL -oriented as shown in [4, §3.6, §3.7]. The Thom-Gysin sequence for MW motivic cohomology $H_{MW}^{i,j}(-) := H_{MW}^{i,j}(-, \mathbb{Z})$ is expressed as follows:

$$\dots \xrightarrow{\cup e(f_{n,k})} H_{MW}^{i,j}(V_k(\mathbb{A}^n)) \rightarrow H_{MW}^{i,j}(V_{k+1}(\mathbb{A}^n)) \rightarrow H_{MW}^{i-(2(n-k)-1), j-(n-k)}(V_k(\mathbb{A}^n)) \xrightarrow{\cup e(f_{n,k})} H_{MW}^{i+1,j}(V_k(\mathbb{A}^n)) \rightarrow \dots$$

Consequently, we derive the short exact sequence:

$$0 \rightarrow \text{Coker}(\cup e(f_{n,k})) \rightarrow H_{MW}^{i,j} \rightarrow \text{Ker}(\cup e(f_{n,k})) \rightarrow 0. \quad (1)$$

To carry out calculations, it suffices to understand the Euler class $e(f_{n,k}) \in H_{MW}^{2(n-k), n-k}(V_k(\mathbb{A}^n))$.

We begin by reviewing some facts about the cone $C_\eta(K) \in \widetilde{DM}(K)$, which is associated with the morphism $\widetilde{M}(K)(1)[1] \xrightarrow{\eta} \widetilde{M}(K) \rightarrow C_\eta(K)$.

Proposition 3.1 ([6, Prop 2.2]). *The following exact sequence splits:*

$$H_{MW}^{i-2, j-1}(K) \rightarrow H_{MW}^{i,j}(C_\eta(K)) \rightarrow H_{MW}^{i,j}(K) \xrightarrow{\eta} H_{MW}^{i,j}(C_\eta(K)) = \begin{cases} H_M^{i-2, j-1}(K) \oplus H_M^{i,j}(K), & \text{if } i \neq j \text{ or } i = j < 0, \\ H_M^{i-2, j-1}(K) \oplus 2K_j^M(K), & \text{if } i = j \geq 0 \end{cases}$$

Here, $H_M^{i,j}(-) := H_M^{i,j}(-, \mathbb{Z})$ denotes the motivic cohomology as defined by Voevodsky [3].

We also note the following exact sequence of abelian groups:

$$0 \rightarrow 2K_j^M(K) \cong \eta K_j^{MW}(K) \rightarrow K_j^{MW}(K) \xrightarrow{\eta} K_{j-1}^{MW}(K) \rightarrow K_{j-1}^M(K) \rightarrow 0.$$

This motivates us to define a modified cohomology group:

$$\eta H_M^{i,j}(K) = \begin{cases} H_M^{i,j}(K), & \text{if } i \neq j, \\ 2K_j^M(K), & \text{if } i = j. \end{cases}$$

Consequently, we have the following exact sequence of the graded cohomology group $H_{MW}^{*,*}(K) = \bigoplus_{i,j} H_{MW}^{i,j}(K)$:

$$0 \rightarrow \eta H_M^{*,*}(K) \rightarrow H_{MW}^{*,*}(K) \xrightarrow{\eta} H_M^{*-1, *-1}(K) \rightarrow H_M^{*-1, *-1}(K) \rightarrow 0.$$

Additionally, we have $H_{MW}^{*,*}(C_\eta(K)) = H_M^{*,*}(K)\theta_{2,1} \oplus \eta H_M^{*,*}(K)$.

To compute the cohomology of $V_2(\mathbb{A}^n)$, we utilize the following theorem:

Proposition 3.2 ([4, Cor 3.9]). *The following motivic decompositions are valid in $\widetilde{DM}(S)$:*

$$\widetilde{M}(V_2(\mathbb{A}^n)) \cong \widetilde{M}(\mathbb{A}^n \setminus \{0\}) \otimes \widetilde{M}(\mathbb{A}^{n-1} \setminus \{0\})$$

for n even, and

$$\widetilde{M}(V_2(\mathbb{A}^n)) \cong \widetilde{HS}_n$$

for n odd.

Corollary 3.3. *We have an isomorphism of graded $H_{MW}^{*,*}(K)$ -modules:*

$$H_{MW}^{*,*}(V_2(\mathbb{A}^n)) = \begin{cases} H_{MW}^{*,*}(K)[\beta_{n-1}, \beta_{n-2}]/(\beta_{n-1}^2, \beta_{n-2}^2), & \text{if } n \text{ is even,} \\ H_{MW}^{*,*}(K)\beta_{n-1, n-2} \oplus H_M^{*,*}(K)\alpha_{n-1} \oplus \eta H_M^{*,*}(K)\alpha_{n-2} \oplus H_{MW}^{*,*}(K), & \text{if } n \text{ is odd,} \end{cases}$$

where $\deg(\beta_i) = \deg(\alpha_i) = (2i+1, i+1) = (i, \{i+1\})$, meaning that $\alpha_i \in H_{MW}^{2i+1, i+1}(V_2(\mathbb{A}^n))$, and $\deg(\beta_{n-1, n-2}) = (4n-4, 2n-1) = (2n-3, \{2n-1\})$.

Now, we can state our main result for $V_k(\mathbb{A}^n)$. Let $N_{n,k} = \{t \in \mathbb{Z} \mid n-k \leq t \leq n-1\}$. We define the *free pairs* as follows:

$$F_{n,k} = \{\{n-1\} \mid \text{if } n \text{ is even}\} \sqcup \{\{i, i-1\} \subset N_{n,k} \mid \text{if } i \text{ is even}\} \sqcup \{\{n-k\} \mid \text{if } n-k \text{ is even}\}.$$

Additionally, we define the *free part* and the *torsion part* of a subset $I \subset N_{n,k}$ as follows:

$$I_F = \bigsqcup_{J \in F_{n,k}, J \subset I} J, \quad I_T = I \setminus I_F.$$

Furthermore, we define the degree for a subset I as $d(I) = (\sum_{i \in I} (2i+1), \sum_{i \in I} (i+1))$.

Theorem 3.4. *Let $i \in N_{n,k}$. The generators α_i are elements of $H_{\text{MW}}^{2i+1, i+1}(V_k(\mathbb{A}^n))$, and for $J \in F_{n,k}$, the generators β_J belong to $H_{\text{MW}}^{d(J)}(V_k(\mathbb{A}^n))$. For any subset $I \subset N_{n,k}$, define*

$$b_I = \prod_{J \in F_{n,k}, J \subset I} \beta_J \prod_{i \in I_T} \alpha_i \in H_{\text{MW}}^{d(I)}(V_k(\mathbb{A}^n)).$$

Then, we establish the following isomorphism of graded $H_{\text{MW}}^{*,*}(K)$ -modules:

$$H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n)) \cong \bigoplus_{I \subset N_{n,k}} (H_{\text{MW}})_{I} b_I,$$

where

$$(H_{\text{MW}})_{I} = \begin{cases} H_{\text{MW}}^{*,*}(K), & \text{if } I = I_F, \\ H_{\text{M}}^{*,*}(K), & \text{if } \text{Max}(I_T) \text{ is even,} \\ \eta H_{\text{M}}^{*,*}(K), & \text{if } \text{Max}(I_T) \text{ is odd.} \end{cases}$$

We can compute the Euler class as a corollary:

Corollary 3.5. *For $k \geq 2$, we have*

$$H_{\text{MW}}^{2(n-k), n-k}(V_k(\mathbb{A}^n)) = \begin{cases} K_{-1}^{\text{MW}} \beta_{n-k}, & \text{if } n-k \text{ is even,} \\ 0, & \text{if } n-k \text{ is odd.} \end{cases}$$

Consequently, the Euler class satisfies $e(f_{n,k}) = \eta \beta_{n-k}$ if $n-k$ is even, and $e(f_{n,k}) = 0$ if $n-k$ is odd.

Proof. The first part of the statement follows from comparing the degrees of generators in Theorem 3.4. Specifically, since b_{n-k} has the lowest degree $(n-k, \{n-k+1\})$ and H_{MW} is coconnected (i.e., $H_{\text{MW}}^{p, \{q\}}(K) = 0$ for $p > 0$), we conclude that

$$H_{\text{MW}}^{2(n-k), n-k}(V_k(\mathbb{A}^n)) = (H_{\text{MW}})_{n-k}^{-1, -1} b_{n-k}.$$

For $n-k$ even, we have $b_{n-k} = \beta_{n-k}$ and $(H_{\text{MW}})_{n-k}^{-1, -1} = K_{-1}^{\text{MW}}$. Conversely, for $n-k$ odd, it follows that $(H_{\text{MW}})_{n-k}^{-1, -1} = \eta H_{\text{M}}^{-1, -1}(K) = 0$.

The known values for the Euler classes in the case $k=1$ are provided by [4, Prop 3.8]. Specifically, in these situations, it is stated that $e(f_{n,1}) = n_\epsilon \eta \beta_{n-1} \in H_{\text{MW}}^{2(n-1), n-1}(\mathbb{A}^n \setminus \{0\})$. This implies that $e(f_{n,1}) = 0$ if n is even, and $e(f_{n,1}) = \eta \beta_{n-1}$ if n is odd, where $\beta_{n-1} \in H_{\text{MW}}^{2n-1, n}(\mathbb{A}^n \setminus \{0\})$ is a generator.

We can consider the following diagram:

$$\begin{array}{ccccc} \mathbb{A}^{n-k} \setminus \{0\} & \longrightarrow & V_2(\mathbb{A}^{n-k+1}) & \xrightarrow{f_{n-k+1,1}} & \mathbb{A}^{n-k+1} \setminus \{0\} \\ & & \downarrow & \lrcorner & \downarrow g \\ & & \mathbb{A}^{n-k} \setminus \{0\} & \longrightarrow & V_{k+1}(\mathbb{A}^n) & \xrightarrow{f_{n,k}} & V_k(\mathbb{A}^n) \\ & & & & \downarrow & & \downarrow \\ & & & & V_{k-1}(\mathbb{A}^n) & \longleftarrow & V_{k-1}(\mathbb{A}^n) \end{array}$$

Utilizing Corollary 2.3, we observe that when $n-k$ is even, the pullback

$$g^* : H_{\text{MW}}^{2(n-k), n-k}(V_k(\mathbb{A}^n)) \rightarrow H_{\text{MW}}^{2(n-k), n-k}(\mathbb{A}^{n-k+1} \setminus \{0\})$$

is an isomorphism. Furthermore, the Euler class is compatible with the pullback, expressed as $g^*(e(f_{n,k})) = e(f_{n-k+1,1})$. Consequently, we derive that $e(f_{n,k}) = \eta\beta_{n-k}$. \square

Although we do not provide the multiplicative structure of the graded cohomology group, we can state a useful fact about it.

Corollary 3.6. *Assume $n - k$ is even, and let $\beta_{n-k} \in H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n))$ be a generator with $n > k$. Then we have*

$$\eta\beta_{n-k}^2 = 0.$$

Proof. By induction and by comparing the degree, we find that the degree of β_{n-k}^2 is given by $\deg(\beta_{n-k}^2) = (2(n-k), \{2(n-k+1)\})$. Again, exploiting the coconnectedness of H_{MW} , we conclude that $\beta_{n-k}^2 \in (H_{\text{MW}})_{2(n-k)}^{1,1} b_{2(n-k)} = K_1^M \alpha_{2(n-k)}$ if $2(n-k) < n$, or that $\beta_{n-k}^2 = 0$ if $2(n-k) \geq n$.

In both scenarios, we establish that $\eta\beta_{n-k}^2 = 0$. \square

Remark 1. *According to the computations in motivic cohomology [5], we obtain the following result:*

$$\beta_{n-k}^2 = \begin{cases} 0, & \text{if } 2(n-k) \geq n, \\ [-1]_{K^M} \alpha_{2(n-k)}, & \text{if } 2(n-k) < n. \end{cases}$$

We prove our main theorem by induction.

Proposition 3.7. *Assuming that Theorem 3.4 holds for $V_k(\mathbb{A}^n)$, then the exact sequence (1) splits. Furthermore, we have the following isomorphism of $H_{\text{MW}}^{*,*}(K)$ -modules:*

$$H_{\text{MW}}^{*,*}(V_{k+1}(\mathbb{A}^n)) \cong \begin{cases} H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n)) \oplus H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n))\beta_{n-k-1}, & \text{if } n-k \text{ is odd,} \\ H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)) \oplus \text{Coker}_\eta(H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)))\alpha_{n-k} \\ \oplus \text{Ker}_\eta(H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)))\alpha_{n-k-1} \oplus H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n))\beta_{n-k, n-k-1}, & \text{if } n-k \text{ is even.} \end{cases}$$

Here, $\text{Coker}_\eta(H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)))$ and $\text{Ker}_\eta(H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)))$ denote the cokernel and kernel of η , respectively:

$$0 \rightarrow \text{Ker}_\eta(H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n))) \rightarrow H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)) \xrightarrow{\eta} H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)) \rightarrow \text{Coker}_\eta(H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n))) \rightarrow 0.$$

Proof. Since the boundary morphism is evident from Corollaries 3.5 and 3.6, it suffices to demonstrate that the exact sequence (1) splits.

When $n - k$ is odd, the result is straightforward. In this case, $H_{\text{MW}}^{*,*}(V_{k+1}(\mathbb{A}^n))$ serves as a $H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n))$ -module, and $H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n))$ itself is a free $H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n))$ -module. We denote the generators by 1 and β_{n-k-1} .

When $n - k$ is even, under our assumption, we have the isomorphism

$$H_{\text{MW}}^{*,*}(V_k(\mathbb{A}^n)) \cong H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)) \oplus H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n))\beta_{n-k}.$$

From Corollary 3.5, we can identify the Euler class as $e(f_{n,k}) = \eta\beta_{n-k}$. Moreover, invoking Corollary 3.6, we can rewrite the exact sequence (1) as:

$$\begin{aligned} 0 &\rightarrow H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)) \oplus \text{Coker}_\eta(H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)))\alpha_{n-k} \\ &\rightarrow H_{\text{MW}}^{*,*}(V_{k+1}(\mathbb{A}^n)) \\ &\rightarrow \text{Ker}_\eta(H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n)))\alpha_{n-k-1} \oplus H_{\text{MW}}^{*,*}(V_{k-1}(\mathbb{A}^n))\beta_{n-k, n-k-1} \rightarrow 0. \end{aligned}$$

To demonstrate that this sequence splits, it is sufficient to consider the η -torsion part. This is evident when comparing it with the exact sequence of $H_M^{*,*}(V_{k+1}(\mathbb{A}^n))$ presented in [5]. \square

Proof of Theorem 3.4. We observe that the following relations hold:

$$\begin{aligned} \text{Coker}_\eta(H_{\text{MW}}^{*,*}(K)) &= H_M^{*,*}(K), \quad \text{Ker}_\eta(H_{\text{MW}}^{*,*}(K)) = {}^\eta H_M^{*,*}(K), \\ \text{Coker}_\eta(H_M^{*,*}(K)) &= \text{Ker}_\eta(H_M^{*,*}(K)) = H_M^{*,*}(K), \\ \text{Coker}_\eta({}^\eta H_M^{*,*}(K)) &= \text{Ker}_\eta({}^\eta H_M^{*,*}(K)) = {}^\eta H_M^{*,*}(K). \end{aligned}$$

These observations provide the necessary conditions to establish the existence of a direct summand in Theorem 3.4. \square

4 Decomposition of MW-motive of $V_k(\mathbb{A}^n)$

It is noteworthy that throughout the proof, we relied solely on the coconnectedness of \mathbf{H}_{MW} as described in Corollaries 3.5 and 3.6. Consequently, the same computation applies to all coconnected MW-motives $\mathbb{E} \in \widetilde{\text{DM}}(K)$. In particular, this assertion holds for $C_\eta = C_\eta(K)$.

Theorem 4.1. *Let $\mathbb{E} \in \widetilde{\text{DM}}(K)$ be a coconnected MW-motive, and let ${}^\eta\mathbb{E}$ and \mathbb{E}/η denote the kernel and cokernel of the morphism η . The sequence is defined as follows:*

$$0 \rightarrow {}^\eta\mathbb{E}^{*,*}(K) \rightarrow \mathbb{E}^{*,*}(K) \xrightarrow{\eta} \mathbb{E}^{*,*}(K) \rightarrow (\mathbb{E}/\eta)^{*,*}(K) \rightarrow 0$$

We use the same notation as in Theorem 3.4. Then, we establish an isomorphism of graded $\mathbb{E}^{*,*}(K)$ -modules:

$$\mathbb{E}^{*,*}(V_k(\mathbb{A}^n)) = \bigoplus_{I \subset N_{n,k}} \mathbb{E}_I b_I$$

where

$$\mathbb{E}_I = \begin{cases} \mathbb{E}^{*,*}(K), & \text{if } I = I_F, \\ (\mathbb{E}/\eta)^{*,*}(K), & \text{if } \text{Max}(I_T) \text{ is even,} \\ {}^\eta\mathbb{E}^{*,*}(K), & \text{if } \text{Max}(I_T) \text{ is odd.} \end{cases}$$

We now proceed to prove the conjecture stated in [4, Conj 1.4]. Recall that $C_\eta(S)$ represent the cone of the morphism $\widetilde{\text{M}}(S) \xrightarrow{\eta} \widetilde{\text{M}}(S)(1)[1]$, and

$$\begin{aligned} \widetilde{\text{HS}}_{2k} &:= \widetilde{\text{M}}(\mathbb{A}^{2k} \setminus \{0\}) \cong \widetilde{\text{M}}(S) \oplus \widetilde{\text{M}}(S)(2k)[4k-1], \\ \widetilde{\text{HS}}_{2k+1} &:= \widetilde{\text{M}}(S) \oplus C_\eta(S)(2k)[4k] \oplus \widetilde{\text{M}}(S)(4k+1)[8k]. \end{aligned}$$

Theorem 4.2. *For any $i, j \in \mathbb{N}$, we obtain an MW-motivic decomposition of the following forms:*

$$\begin{aligned} \widetilde{\text{M}}(V_{2j}(\mathbb{A}^{2i})) &\cong \widetilde{\text{HS}}_{2i} \otimes \widetilde{\text{HS}}_{2i-1} \otimes \widetilde{\text{HS}}_{2i-3} \otimes \dots \otimes \widetilde{\text{HS}}_{2i-1-2(j-2)} \otimes \widetilde{\text{M}}(\mathbb{A}^{2i+1-2j} \setminus \{0\}) \\ \widetilde{\text{M}}(V_{2j+1}(\mathbb{A}^{2i})) &\cong \widetilde{\text{HS}}_{2i} \otimes \widetilde{\text{HS}}_{2i-1} \otimes \widetilde{\text{HS}}_{2i-3} \otimes \dots \otimes \widetilde{\text{HS}}_{2i-1-2(j-1)} \\ \widetilde{\text{M}}(V_{2j}(\mathbb{A}^{2i+1})) &\cong \widetilde{\text{HS}}_{2i+1} \otimes \widetilde{\text{HS}}_{2i-1} \otimes \dots \otimes \widetilde{\text{HS}}_{2i+1-2(j-1)} \\ \widetilde{\text{M}}(V_{2j+1}(\mathbb{A}^{2i+1})) &\cong \widetilde{\text{HS}}_{2i+1} \otimes \widetilde{\text{HS}}_{2i-1} \otimes \dots \otimes \widetilde{\text{HS}}_{2i+1-2(j-1)} \otimes \widetilde{\text{M}}(\mathbb{A}^{2i+1-2j} \setminus \{0\}). \end{aligned}$$

Proof. We will only prove the case when $S = \text{Spec } K$.

Let $Y_{k,n}$ denote the right-hand side of the isomorphism in the theorem. By definition, we have the following cofiber sequences for $Y_{k,n}$:

$$Y_{k-1,n-1} \rightarrow Y_{k,n} \rightarrow \widetilde{\mathbb{Z}}(n)[2n-1] \otimes Y_{k-1,n-1} \xrightarrow{0}$$

for n even, and

$$Y_{k-1,n-1} \rightarrow Y_{k,n} \rightarrow \widetilde{\mathbb{Z}}(n)[2n-1] \otimes Y_{k-1,n-1} \xrightarrow{\beta}$$

for n odd, where β is defined as follows:

$$\widetilde{\mathbb{Z}}(n)[2n-1] \otimes (Y_{k-2,n-2} \oplus Y_{k-2,n-2}(n-1)[2n-3]) \xrightarrow{\begin{bmatrix} 0 & \eta \\ 0 & 0 \end{bmatrix}} Y_{k-2,n-2}[1] \oplus Y_{k-2,n-2}(n-1)[2n-2].$$

From Theorem 4.1, we observe that for coconnected \mathbb{E} , the following holds:

$$\text{Hom}_{\widetilde{\text{DM}}(K)}(\widetilde{\text{M}}(V_k(\mathbb{A}^n)), \mathbb{E}(q)[p]) \cong \text{Hom}_{\widetilde{\text{DM}}(K)}(Y_{k,n}, \mathbb{E}(q)[p]).$$

It is noteworthy that $Y_{k,n}$ is a direct sum of $\widetilde{\mathbb{Z}}(q)[p]$ and $C_\eta(q)[p]$. In particular, we have

$$\mathrm{Hom}_{\widetilde{\mathrm{DM}}(K)}(\widetilde{M}(V_k(\mathbb{A}^n)), Y_{k,n}) \cong \mathrm{Hom}_{\widetilde{\mathrm{DM}}(K)}(Y_{k,n}, Y_{k,n}).$$

Let $\iota_{k,n} : \widetilde{M}(V_k(\mathbb{A}^n)) \rightarrow Y_{k,n}$ denote the morphism corresponding to the identity of $Y_{k,n}$. We now need to prove inductively that $\iota_{k,n}$ is an isomorphism. For $k = 1$, this is evident, and for $k = 2$, the result follows from [4, Cor 3.9].

For $V_k(\mathbb{A}^n)$, we have the following cofiber sequence in $\mathcal{H}(K)$ (see [4, §3.2]):

$$V_{k-1}(\mathbb{A}^{n-1})_+ \rightarrow V_k(\mathbb{A}^n)_+ \rightarrow \mathbb{A}^n \setminus \{0\} \wedge V_{k-1}(\mathbb{A}^{n-1})_+.$$

When n is even, one can construct the splitting projection using that of $Y_{k,n}$:

$$\mathrm{Hom}_{\widetilde{\mathrm{DM}}(K)}(\widetilde{M}(V_k(\mathbb{A}^n)), \widetilde{M}(V_{k-1}(\mathbb{A}^{n-1}))) \cong \mathrm{Hom}_{\widetilde{\mathrm{DM}}(K)}(\widetilde{M}(V_k(\mathbb{A}^n)), Y_{k-1,n-1}) \cong \mathrm{Hom}_{\widetilde{\mathrm{DM}}(K)}(Y_{k,n}, Y_{k-1,n-1}).$$

Thus, it follows that

$$\widetilde{M}(V_k(\mathbb{A}^n)) \cong \widetilde{M}(V_{k-1}(\mathbb{A}^{n-1})) \oplus \widetilde{M}(V_{k-1}(\mathbb{A}^{n-1}))(n)[2n-1] \cong Y_{k-1,n-1} \oplus Y_{k-1,n-1}(n)[2n-1] \cong Y_{k,n}.$$

For n odd, by employing the same argument used in the proof of [4, Prop 3.13], we arrive at the following cofiber sequences in $\widetilde{\mathrm{DM}}(K)$:

The equations presented elucidate the relationships between various mathematical structures in the study. Specifically, we have the following mappings:

$$\widetilde{M}(V_{k-2}(\mathbb{A}^{n-2})) \rightarrow \widetilde{M}(V_k(\mathbb{A}^n)) \rightarrow Q_{k,n}$$

and

$$C_\eta \otimes \widetilde{M}(V_{k-2}(\mathbb{A}^{n-2}))(n-1)[2n-2] \rightarrow Q_{k,n} \rightarrow \widetilde{Z}(n)[2n-1] \otimes \widetilde{M}(V_{k-2}(\mathbb{A}^{n-2}))(n-1)[2n-3]$$

In a similar vein, we can construct the splitting projections, leading us ultimately to the isomorphisms:

$$\begin{aligned} \widetilde{M}(V_k(\mathbb{A}^n)) &\cong \widetilde{M}(V_{k-2}(\mathbb{A}^{n-2})) \oplus \left(C_\eta \otimes \widetilde{M}(V_{k-2}(\mathbb{A}^{n-2}))(n-1)[2n-2] \right) \oplus \widetilde{M}(V_{k-2}(\mathbb{A}^{n-2}))(2n-1)[4n-4] \\ &\cong Y_{k-2,n-2} \oplus (C_\eta \otimes Y_{k-2,n-2}(n-1)[2n-2]) \oplus Y_{k-2,n-2}(2n-1)[4n-4] \cong Y_{k,n} \end{aligned}$$

□

References

- [1] Aravind Asok, Frédéric Déglise, and Jan Nagel. The homotopy lera spectral sequence. *Motivic Homotopy Theory and Refined Enumerative Geometry*, 2018.
- [2] Martin Čadek, Mamoru Mimura, and Jiří Vanžura. The cohomology rings of real stiefel manifolds with integer coefficients. *Journal of Mathematics of Kyoto University*, 43(2):411–428, 2003.
- [3] Carlo Mazza, Vladimir Voevodsky, and Charles A Weibel. *Lecture notes on motivic cohomology*, volume 2. American Mathematical Soc., 2006.
- [4] Keyao Peng. Milnor-witt motivic cohomology and linear algebraic groups. *Advances in Mathematics*, 458:109973, 2024.
- [5] Ben Williams. The motivic cohomology of stiefel varieties. *Journal of K-theory*, 10(1):141–163, 2012.
- [6] Nanjun Yang. Split milnor-witt motives and its applications to fiber bundles. *Cambridge Journal of Mathematics*, 10(2):935–1004, 2022.