GRADED FROBENIUS ALGEBRAS

JONATHAN CLIVIO

Abstract. We construct a PROP which encodes 2D-TQFTs with a grading. This defines a graded Frobenius algebra as algebras over this PROP. We also give a description of graded Frobenius algebras in terms of maps and relations. This structure naturally arises as the cohomology of manifolds, loop homology and Hochschild homology of Frobenius algebras. In addition, we give a comprehensive description of the signs that arise in suspending algebras over PROPs.

1. INTRODUCTION

The goal of this paper is to clarify what is meant by a graded Frobenius algebra. To this end, we introduce a PROP that describes a Frobenius algebra with multiplication in a fixed degree c and comultiplication of a fixed degree d.

There is a need for such a definition, because the straight-forward attempt of considering Frobenius algebras in graded abelian groups runs into problems. Defining a Frobenius algebra in graded abelian groups is unproblematic if all maps are degree-preserving i.e. they are maps in degree 0. But this attempt fails to describe some non-trivial examples if maps of different degrees are considered: for example, we consider a graded abelian group A with a unital multiplication $\mu: A \otimes A \rightarrow A$ and a counital comultiplication $\nu: A \to A \otimes A$ on it satisfying the usual Frobenius relation

(1)
$$
(\mu \otimes id) \circ (id \otimes \nu) = \nu \circ \mu = (id \otimes \mu) \circ (\nu \otimes id).
$$

We now assume that μ and ν are of some degree $|\mu| = c$ and $|\nu| = d$. If $c + d$ is odd and we work over a field where $1 \neq -1$, the only example that is described is the trivial algebra $A = 0$. We prove this in Proposition [3.6.](#page-11-0)

Another attempt to describe examples of Frobenius algebras with maps in non-zero degree is to suspend a Frobenius algebra. Suppose A is a Frobenius algebra with multiplication μ and ν in degree 0. Suspending A gives an algebra ΣA with a multiplication $\Sigma \mu$ in degree -1 and a comultiplication $\Sigma \nu$ in degree 1. Suspension and desuspension of algebras therefore leaves the sum of the degree of the multiplication and the degree of the comultiplication invariant. Thus an example of A with a multiplication μ in degree 0 and a comultiplication in degree $d \neq 0$ cannot be described as the suspension of an algebra with only operations in degree 0. Moreover, the canonical map

(2)
$$
\Sigma: \text{Hom}(A^{\otimes k}, A^{\otimes l}) \to \text{Hom}((\Sigma A)^{\otimes k}, (\Sigma A)^{\otimes l})
$$

is only functorial and symmetric monoidal up to sign. We compute these signs in Lemma [5.10.](#page-23-0) These signs also illustrate why Frobenius algebras in graded abelian groups are not the natural concept to consider. For example if μ and ν satisfy the Frobenius relation as in [\(1\)](#page-0-0), the signs in the

Date: December 19, 2024.

functoriality and monoidality of Σ in [\(2\)](#page-0-1) induce the following sign in the Frobenius relation:

$$
(\Sigma \mu \otimes id) \circ (id \otimes \Sigma \nu) = (-1) \nu \circ \mu = (id \otimes \Sigma \mu) \circ (\Sigma \nu \otimes id).
$$

Therefore, the natural concept of graded Frobenius algebras will have signs appearing in the relations.

1.1. The Graded Frobenius PROP. In this paper, we give a definition of graded Frobenius algebras with a multiplication μ in degree c and a comultiplication ν in degree d for any choice $c, d \in \mathbb{Z}$ (see Definition [4.4\)](#page-15-0). This definition solves the two problems described above: we can describe non-trivial examples for any choice of $c, d \in \mathbb{Z}$ and the signs that appear in this definition are stable under suspending algebras. We make this second claim precise when we discuss the relations in subsection [1.3.](#page-2-0)

We describe a category of graph cobordisms in Definition [2.7.](#page-8-0) This lets us model every cobordism by an equivalence class of graphs.

For a graph G representing a cobordism, we consider a twisting by the determinant $\det(G, \partial_{in})$ of the homology relative to the in-coming boundary. The determinant $\det(G, \partial_{in})$ is a graded rank one free abelian group concentrated in degree $-\chi(G, \partial_{in})$, the negative Euler characteristic. Analogously, we consider $\det(G, \partial_{out})$.

This lets us define the category of graded graph cobordism $GCob_{c,d}$ and fat graph cobordisms $fGCob_{c,d}$ in Definition [4.3.](#page-15-1) This category has morphisms given by elements in the following graded abelian groups

$$
\mathbf{det}_{c,d}(G):=\mathbf{det}(G,\partial_{in})^{\otimes c}\otimes \mathbf{det}(G,\partial_{out})^{\otimes d}.
$$

In this category, there is a morphism $\{*\} \sqcup \{*\} \rightarrow \{*\}$. It models a multiplication of the graded Frobenius algebra $\{*\}$. It is given by a generator $\omega(\rightarrow) \in \det_{c,d}(\rightarrow)$. Here \succ is the graph with two inputs to the left, one output to the right and one three-valent vertex in the middle. It has relative Euler characteristic $-\chi(\rightarrow,\partial_{in}) = 1$ and $-\chi(\rightarrow,\partial_{out}) = 0$. Therefore $\det_{c,d}(\rightarrow)$ is concentrated in degree c and the multiplication $\omega(\rightarrow)$ is in degree c. Analogously, the comultiplication is given by a generator $\omega(\cdot) \in \det_{c,d}(\cdot)$ in degree d.

The categories $\text{GCob}_{c,d}$ and $\text{fGCob}_{c,d}$ lets us thus attain any combination in degrees c, d. Moreover, $GCob_{c,d}$ is modelled by graphs and describes (c, d) -graded *commutative* Frobenius algebras while $fGCob_{c,d}$ has morphisms given by fat graphs and describes (c, d) -graded symmetric Frobenius algebras.

This twisting was previously described by Godin in [\[God07\]](#page-47-0) and by Wahl and Westerland in [\[WW16\]](#page-47-1). However, in contrast to Godin who models commutative Frobenius algebras by closed surfaces, we work with graphs. In particular, twisting a surface S by $\det(S, \partial_{in})$ corresponds to a twisting by $\det(G, \partial_{in}) \otimes \det(G, \partial_{out})$ where G is a graph modelling the cobordism S. We prove this relation between the twistings in Proposition [6.4.](#page-28-0) In particular, working with graphs rather than closed surfaces for commutative graded Frobenius algebras allows us to describe examples of algebras where $c + d$ is odd.

1.2. Suspending PROPs. Above we observed that suspending a algebra with multiplication in degree 0 and comultiplication in degree 0 gives an algebra with multiplication in degree −1 and a comultiplication in degree 1. These shifts in degree hold for any choice in degrees c, d .

This can be seen on the level of PROPs: we define the usual suspension PROP Σ in Definition [4.13.](#page-17-0) This PROP has the following universal property: a graded abelian group A has the structure of an algebra over a PROP P if and only if ΣA has the structure of an algebra over $\Sigma \otimes P$ (see Proposition [4.14\)](#page-17-1). On the other hand, we give natural isomorphisms of PROPs

$$
\Sigma\otimes \operatorname{GCob}_{c,d}\cong \operatorname{GCob}_{c-1,d+1},
$$

$$
\Sigma\otimes \operatorname{fGCob}_{c,d}\cong \operatorname{fGCob}_{c-1,d+1}
$$

in Proposition [4.16.](#page-18-0) This, in particular, shows that suspending a (c, d) -graded Frobenius algebra gives a $(c-1, d+1)$ -graded Frobenius algebra.

This is in contrast to the relation between (c, d) -graded Frobenius algebras and $(c + 1, d)$ -graded Frobenius algebras. In Remark [4.17,](#page-19-0) we give a construction that produces a $(c + 1, d)$ -graded Frobenius algebra from a (c, d) -graded Frobenius algebra. This construction could be called a onesided suspension of graded Frobenius algebras as it only affects the degree of the multiplication and not the coproduct. However in contrast to the usual suspension, this one-sided suspension is not an equivalence of categories.

1.3. Maps, Relations and Signs. In the ungraded case, there are several equivalent descriptions of a Frobenius algebra. We contrast two different types of definitions:

- (1) Descriptions in terms of maps with relations: an object A together with a selection of the following maps: a multiplication μ , a unit η , a comultiplication ν and a counit ε that satisfy certain relations.
- (2) Descriptions in terms of PROPs: a 2-dimensional topological quantum field theory (TQFT). This is a symmetric monoidal functor out of a 2-dimensional cobordism category.

There is a myriad of different variants of the two flavours of definitions with optional commutativity, unitality, counitality. The equivalence between these two types of definitions is described in [\[Abr96,](#page-47-2) [LP08,](#page-47-3) [Koc04\]](#page-47-4).

We explained above that we define a (c, d) -graded Frobenius algebra in terms of a PROP. In our main theorem, we relate our definition to a description in terms of maps and a choice of signs appearing in the relations:

Main Theorem (Theorem [5.1\)](#page-20-0). Let $(C, \otimes, \mathbb{1}, \tau)$ be a symmetric monoidal category enriched over graded abelian groups and let $c, d \in \mathbb{Z}$. The data $(A, \mu, \eta, \nu, \varepsilon)$ where

 $A \in \mathcal{C}$,

uniquely defines a symmetric monoidal functor out of $\text{GCob}_{c,d}$ into C if and only if the following relations are satisfied:

(i) graded associativity $\mu \circ (\mu \otimes id) = (-1)^c \mu \circ (id \otimes \mu);$

(ii) graded unitality $(-1)^c \mu \circ (\eta \otimes id) = (-1)^{\frac{c(c-1)}{2}} id = \mu \circ (id \otimes \eta)$;

(iii) graded coassociativity (id $\otimes \nu$) $\circ \nu = (-1)^d (\nu \otimes id) \circ \nu$;

(iv) graded counitality $(-1)^d$ (id $\otimes \varepsilon$) $\circ \nu = (-1)^{\frac{d(d-1)}{2}}$ id $= (\varepsilon \otimes id) \circ \nu$;

(v) graded Frobenius relation $(\mu \otimes id) \circ (id \otimes \nu) = (-1)^{cd} \nu \circ \mu = (id \otimes \mu) \circ (\nu \otimes id);$

(vi) graded commutativity $\mu \circ \tau = (-1)^c \mu$;

and it uniquely defines a (c, d) -graded symmetric Frobenius algebra if and only if the maps satisfy the same relations replacing (vi) by

(vi') graded symmetry $\varepsilon \circ \mu \circ \tau = (-1)^c \varepsilon \circ \mu$.

These signs are not canonical. They depend on choices of generators $\omega(\rightarrow) \in \det_{c,d}(\rightarrow)$ and $\omega(\mathcal{A}) \in \det_{c,d}(\mathcal{A})$. We explain in Remark [5.3](#page-20-1) our choices. However, no choice can make the signs

disappear.

We remarked above that our signs are stable under suspending algebras. This means that our choices of signs are such that if $(A, \mu, \eta, \nu, \varepsilon)$ satisfies the relations of (c, d) -graded Frobenius algebra, then $(\Sigma A, \Sigma u, \Sigma \eta, \Sigma \nu, \Sigma \varepsilon)$ satisfies the relations of a $(c-1, d+1)$ -graded Frobenius algebra. This is proved in Proposition [5.11.](#page-24-0)

Our signs almost coincide with the signs found in the definition of a biunital coFrobenius as defined in [\[CO22b\]](#page-47-5). We compare this in Remark [5.4.](#page-21-0)

1.4. Examples. In the literature, graded Frobenius algebras appear with different characterisations. Some have a more PROP-theoretic description and some have a description in terms of maps and relations. Our main theorem lets us relate the two types of description. In section [6,](#page-24-1) we discuss a selection of the examples found in the literature.

The first example we discuss is the cohomology ring of an oriented manifold M of dimension d (see Example [6.1\)](#page-25-0). The intersection coproduct defines a degree d comultiplication on cohomology. In particular, we discuss different sign conventions for the intersection coproduct found in the literature.

A second example, we discuss, is the Hochschild homology of a graded Frobenius algebra A (see Example [6.3\)](#page-28-1). Wahl and Westerland give a description of a Frobenius algebra on $\overline{HH}_{*}(A)$ twisted by $\det(S, \partial_{in})^{\otimes d}$. As discussed above, the twisting by the relative homology of the surface gives a (d, d) -graded Frobenius algebra.

In a similar vein, we discuss the homology of the loop space with the Chas-Sullivan product and the trivial coproduct. Godin gives operations which are twisted by $\det(S, \partial_{in})^{\otimes d}$ in [\[God07\]](#page-47-0). By our Proposition [6.4,](#page-28-0) this corresponds to a twisting by $\det(G, \partial_{in})^{\otimes d} \otimes \det(G, \partial_{out})^{\otimes d}$ (see Example [6.5\)](#page-31-0).

Finally, we discuss the relation between the Chas-Sullivan product and the Goresky-Hingston coproduct in Rabinowitz loop homology as in the work by Cieliebak, Hingston and Oancea in [\[CHO20,](#page-47-6) [CO22b,](#page-47-5) [CO22a,](#page-47-7) [CHO23,](#page-47-8) [CO24\]](#page-47-9).

We discuss the language of Tate vector spaces as presented in [\[CO24\]](#page-47-9). Concretely for A a graded Tate vector space, we show that the endomorphisms $\text{End}_A(X, Y) = \text{Hom}(A^{\widehat{\otimes}^* X}, A^{\widehat{\otimes}^! Y})$ have a canonical dioperad structure in Proposition [6.6.](#page-33-0)

On the other hand, we define the canonical subdioperad of $GCob_{c,d}$ in Definition [4.19.](#page-19-1) This dioperad has the same operations as the full PROP in the case of a field of characteristic $\neq 2$. This is a consequence of our description of the subdioperad in Proposition [4.20.](#page-19-2) We can thus describe the Rabinowitz loop homology as an algebra over the subdioperad in Example [6.8.](#page-33-1)

Organisation of the Paper. In Section [2,](#page-5-0) we describe a category of graph cobordisms modelling the category of 2-dimensional cobordisms. In Section [3,](#page-9-0) we recall the classical relations of Frobenius algebras and show that they force algebras to be trivial in the odd dimensional graded case. In Section [4,](#page-12-0) we construct the categories $\text{GCob}_{c,d}$ and $\text{fGCob}_{c,d}$ and study graded Frobenius algebras in terms of PROPs. In section [5,](#page-20-2) we state the main theorem and use it to study graded Frobenius algebras in terms of maps and relations. In Section [6,](#page-24-1) we review and relate examples from the literature of graded Frobenius algebras. In Section [7,](#page-33-2) we set up machinery that helps us then prove the main theorem.

Conventions. Throughout this paper, we use the convention that maps act from the left. This in particular implies that in a graded context, we have the following signs

$$
(f \otimes g)(a \otimes b) = (-1)^{|g||a|} f(a) \otimes g(b)
$$

and that the canonical composition map on morphism spaces is order-reversing

$$
Hom(Y, Z) \otimes Hom(X, Y) \to Hom(X, Z),
$$

$$
f \otimes g \mapsto f \circ g.
$$

We define the suspension of a graded abelian A by ΣA . It is defined by $(\Sigma A)_{i+1} = A_i$. We denote by 1 the unit of the category of graded abelian groups: $\mathbb Z$ concentrated in degree 0. We denote the dual of a graded abelian group by A^{\vee} defined as $A_k^{\vee} := \text{Hom}_{\bullet}(A_{-k}, \mathbb{1}).$ The natural isomorphism from the dual of a tensor product to the product of the duals is orderreversing, we have

$$
(A \otimes B)^{\vee} \cong B^{\vee} \otimes A^{\vee},
$$

$$
(a \otimes b)^{\vee} \mapsto b^{\vee} \otimes a^{\vee}.
$$

This can be seen as $b^{\vee} \otimes a^{\vee}$ evaluated on $a \otimes b$ gives 1. But $a^{\vee} \otimes b^{\vee}$ evaluated on $a \otimes b$ gives $(-1)^{|a||b|}$.

Acknowledgments. I thank my PhD advisor Nathalie Wahl for suggesting the topic of signs in Frobenius algebras, many interesting discussions and reading my various drafts. I also like to thank Jan Steinebrunner for helpful discussions about operads, PROPs and Frobenius algebras. I further like to thank Kai Cieliebak and Alexandru Oancea for sharing their draft of [\[CO22b\]](#page-47-5) and their insights on graded Frobenius algebras and its relation to string topology. Furthermore, I thank Alex Takeda for explaining the relation of our conventions to pre-Calabi-Yau structures. Finally, I thank Isaac Moselle and Janine Roshardt for proof-reading and encouraging discussions. I was supported by the Danish National Research Foundation through the Copenhagen Centre for Geometry and Topology (DNRF151).

CONTENTS

2. Cobordisms and Graphs

In this section, we recall the definition of closed and open 2D-cobordism categories. Moreover, we describe an equivalent category coming from a 2-category of graph cobordism.

2.1. Cobordisms.

Definition 2.1. The category of *closed 2D-cobordisms* $\text{Cob}_2^{\text{closed}}$ is defined as

$$
\text{Ob}(\text{Cob}_2^{\text{closed}}) := \left\{ \bigsqcup_X S^1 \middle| X \text{ is a finite set} \right\},
$$

$$
\text{Map}_{\text{Cob}_2^{\text{closed}}} \left(\bigsqcup_X S^1, \bigsqcup_Y S^1 \right) := \left\{ S \text{ surface} \middle| \partial S = \bigsqcup_X S^1 \sqcup \bigsqcup_Y S^1 \right\} / \cong.
$$

See Figure [1.](#page-6-1)

The category of *open 2D-cobordisms* $\mathrm{Cob}_2^{\mathrm{open}}$ is defined as

$$
Ob(Cob_2^{\text{open}}) = \left\{ \bigsqcup_X [0,1] \, \middle| \, X \text{ is a finite set} \right\},
$$

$$
Map_{Cob_2^{\text{open}}} \left(\bigsqcup_X [0,1], \bigsqcup_Y [0,1] \right) = \left\{ S \text{ surface} \, \middle| \, \partial S = \bigsqcup_X S^1 \sqcup \bigsqcup_Y S^1 \sqcup \partial_{free} S \right\} / \cong.
$$

See Figure [2.](#page-6-2)

FIGURE 1. An example of a closed cobordism from $S^1 \sqcup S^1 \sqcup S^1$ to $S^1 \sqcup S^1$. We use the convention of drawing the in-coming boundary to the left and the out-going boundary to the right.

FIGURE 2. An example of an open cobordism from $[0, 1] \sqcup [0, 1]$ to $[0, 1]$.

2.2. Graphs. We later give a description of those two categories using graphs. For this, we introduce what we mean by a graph.

Definition 2.2. A graph $G = (V, H, \sigma, s, L_{in}, L_{out})$ consists of the following data:

- a finite set $V(G) := V$ called the vertices;
- a finite set $H(G) := H$ called the *half-edges*;
- an involution $\sigma: H \to H$ with no fixed points, the orbits $\{h, \sigma h\}$ are called the *edges* and we denote the set $E(G) := E := H_{\sigma};$
- a map $s: H \to V$, where for a vertex $v \in V$ the cardinality $|s^{-1}(v)|$ is called its *arity*;
- disjoint subsets $L_{in}(G) := L_{in}$, $L_{out}(G) := L_{out} \subseteq V(G)$ of the set of vertices of arity ≤ 1 .

A fat graph is a graph together with a cyclic ordering of the half-edges in $s^{-1}(v)$ for all v.

We call the vertices in L_{in} and L_{out} the external vertices. The other vertices are called the internal *vertices*. In our conventions, we allow for internal vertices of arity ≤ 1 .

A graph defines a 1-dimensional CW-complex, called the geometric realization, denoted $|G|$: the zero-cells are the vertices and the one-cells are the edges. Choosing an orientation on each one-cell corresponds to an ordering of the two half-edges making up the edge. Then the map s describes how to glue the edges to the vertices.

We give some additional terminology which is useful, when talking about graphs.

Definition 2.3. Let G be a graph.

- The graph G is *connected* if $|G|$ is connected.
- The graph G is a *forest* if $\pi_1(|G|) = 1$ at every basepoint.
- An edge $e = \{h, \sigma h\}$ is called a *tadpole* if $s(h) = s(\sigma h)$.

Definition 2.4. A morphism of graphs $f: G \to G'$ is a cellular map of the underlying CW-complexes $|G| \to |G'|$ which is on edges either the collapse or the identity and is a bijection on L_{in} and L_{out} .

Remark 2.5. Our definition of morphism of graphs is such that every morphism of graph is given by a sequence of edge collapses and a graph isomorphism. Importantly, the edges that are collapsed are not tadpoles and are not adjacent to an external vertex.

FIGURE 3. A morphism of graphs sends L_{in} to L_{in} and L_{out} to L_{out} and possibly collapses edges. Here the graph morphism collapses two edges.

2.3. Graph Cobordisms. We can now construct the 2-categories GCob and fGCob. This gives models for the cobordism categories.

Definition 2.6. The *graph cobordism category* GCob is a 2-category i.e. a category enriched over 1-categories with

$$
Ob(\underline{GCob}) = \{\text{finite sets}\},\
$$

and the morphism categories $G\text{Cob}(X, Y)$ are given by the category of graphs G with labellings $L_{in}(G) \cong X$ and $L_{out}(G) \cong Y$ and 2-morphisms graph morphisms which respect the labellings. The fat graph cobordism category is the 2-category $fG\dot{C}ob$ with morphism categories $fG\dot{C}ob(X, Y)$ are given by the analogous category of fat graphs with labellings and morphisms are fat graph morphisms which respect the labellings.

This is well-defined as the composition along objects is given as follows: let X, Y, Z be finite. We define the composition as

$$
\underline{\text{GCob}}(Y, Z) \times \underline{\text{GCob}}(X, Y) \to \underline{\text{GCob}}(X, Z)
$$

which sends (G', G) to the graph $G' \circ G := G' \cup_Y G$. This is defined as the graph obtained from the disjoint union of G and G' where we associate the elements in $L_{out}(G)$ with $L_{in}(G')$ via the labelling by Y. In more visual terms, we glue the outgoing legs of G to the incoming legs of G' (as in Figure [4\)](#page-7-1).

FIGURE 4. Composition of a graph cobordism between $\{a, b, c\}$ and $\{x\}$ and a graph cobordism between $\{x\}$ and $\{y\}$.

We note that this defines a functor as morphisms in $GCob(X, Y)$ and $GCob(Y, Z)$ can be glued together in a similar manner.

Analogously, we can define a composition map

$$
\underline{\mathrm{fGCob}}(Y, Z) \times \underline{\mathrm{fGCob}}(X, Y) \to \underline{\mathrm{fGCob}}(X, Z)
$$

for fat graphs.

This works as the newly combined vertices coming from $L_{out}(G) \cong L_{in}(G')$ have arity ≤ 2 and thus there is only one cyclic ordering of the half-edges at these vertices. The fat graph structure of $G' \circ G$ is thus well-defined.

In [\[Chi14\]](#page-47-12)[Section 1.3], it is described that the inclusion of 1-categories into 2-categories has a left adjoint τ_{ι} given on a 2-category $\underline{\mathcal{C}}$ by

Ob
$$
(\tau_{\iota}(\underline{\mathcal{C}})) = Ob(\underline{\mathcal{C}}),
$$

 $\tau_{\iota}(\underline{\mathcal{C}})(X, Y) = \pi_0(\underline{\mathcal{C}}(X, Y)).$

Definition 2.7. We define GCob := τ_L (GCob) and fGCob := τ_L (fGCob) for the associated 1categories.

We can now construct equivalences of categories $\text{GCob} \to \text{Cob}_2^{\text{closed}}$ and $\text{fGCob} \to \text{Cob}_2^{\text{open}}$. A morphism in GCob (or $fGCob$) is given by an equivalence class $[G]$ of (fat) graphs up to zig-zags of (fat) graph isomorphisms.

Let $G \in \underline{GCob}(X, Y)$ be a graph cobordism. We construct the corresponding closed cobordism $Cob(G)$ as follows. Each edge gives a copy of the cylinder $S^1 \times [0, 1]$. Each external vertex is labelled by an element in X or Y. This gives a labelled copy of $S¹$ which we glue to the corresponding cylinder. Finally, we can glue the edges at internal vertices to copies of $S^2 \setminus (\sqcup_{s^{-1}(v)} D^2)$ (see Figure [5\)](#page-8-1). The homeomorphism type of this construction is invariant under edge collapses and graph isomorphisms. Therefore Cob(G) only depends on the class that G represents in $\pi_0(\text{GCob}(X, Y))$. We thus get a well defined map

$$
G\mathrm{Cob}(X,Y) = \pi_0(\underline{G\mathrm{Cob}}(X,Y)) \to \mathrm{Cob}_2^{\mathrm{closed}}(\bigsqcup_X S^1, \bigsqcup_Y S^1),
$$

$$
[G] \mapsto \mathrm{Cob}(G).
$$

Moreover, we note that this is a bijection of sets.

FIGURE 5. The closed cobordism $Cob(G)$ corresponding to a graph G.

Similarly we can construct for a fat graph cobordism between X and Y an open cobordism $f\text{Cob}(G)$ between $\bigsqcup_{X}[0,1]$ and $\bigsqcup_{Y}[0,1]$. It is given by the following construction: each edge can be thickened to $[0, 1] \times [0, 1]$. Then each external vertex corresponds to a closed interval. This gives us copies of the interval $[0, 1]$ labelled by X and Y. Finally, the cyclic ordering at each internal vertex gives a way to glue the thick edges to a closed disc D^2 (see Figure [6\)](#page-9-1). The homeomorphism type fCob(G) only depends on the equivalence class $[G] \in \pi_0(\underline{fGCob}(X, Y))$, because the construction is invariant under fat edge collapses and graph isomorphisms.

We thus find a bijection of sets:

$$
fGCob(X,Y) = \pi_0(\underline{fGCob}(X,Y)) \to Cob_2^{open}(\bigsqcup_X S^1, \bigsqcup_Y S^1),
$$

$$
[G] \mapsto fCob(G).
$$

FIGURE 6. The open cobordism $fCob(G)$ corresponding to a graph G.

We thus proved the following result.

Proposition 2.8. There are equivalences of categories

$$
GCob \to Cob_2^{closed},
$$

\n
$$
X \mapsto \sqcup_X S^1
$$

\n
$$
[G] \mapsto Cob(G)
$$

and

$$
\begin{aligned} \text{fGCob} &\rightarrow \text{Cob}_2^{\text{open}}, \\ X &\rightarrow \sqcup_X[0,1] \\ [G] &\rightarrow \text{fCob}(G). \end{aligned}
$$

3. Frobenius Algebras

In this section, we recall the definition of commutative and symmetric Frobenius algebras. We give definitions both in terms of maps and generators and, on the other hand, as functors out of cobordism categories. Then in [3.1,](#page-11-1) we apply those definitions to the graded context and show in Proposition [3.6](#page-11-0) that in the odd dimensional case, there exist only trivial Frobenius algebras.

Definition 3.1. Let $(C, \otimes, \mathbb{1}, \tau)$ be a symmetric monoidal category.

A commutative Frobenius algebra $(A, \mu, \eta, \nu, \varepsilon)$ in C is an object A in C together with maps

satisfying the following relations:

(i) associativity $\mu \circ (\mu \otimes id) = \mu \circ (id \otimes \mu);$

(ii) unitality $\mu \circ (\eta \otimes id) = id_A = \mu \circ (id \otimes \eta);$

(iii) coassociativity (id $\otimes \nu$) $\circ \nu = (\nu \otimes id) \circ \nu;$

(iv) counitality (id $\otimes \varepsilon$) $\circ \nu = id = (\varepsilon \otimes id) \circ \nu;$

(v) Frobenius relation $(\mu \otimes id) \circ (id \otimes \nu) = \nu \circ \mu = (id \otimes \mu) \circ (\nu \otimes id);$

(vi) commutativity $\mu \circ \tau = \mu$.

Denote by $ComFrob(\mathcal{C})$ the category of commutative Frobenius algebras in \mathcal{C} .

A symmetric Frobenius algebra $(A, \mu, \eta, \nu, \varepsilon)$ in C is an object A in C together with maps satisfying the same relations but replacing (vi) by

(vi') symmetry $\varepsilon \circ \mu \circ \tau = \varepsilon \circ \mu$.

Denote by $SymmFrob(\mathcal{C})$ the category of symmetric Frobenius algebras in \mathcal{C} .

It is a well-known fact that an equivalent description of commutative Frobenius algebras is as symmetric monoidal functor out of a closed 2D-cobordism category (see [\[Abr96,](#page-47-2) [Koc04\]](#page-47-4)). Similarly, a symmetric Frobenius algebra can be characterised as a symmetric monoidal functor out of a open 2D-cobordism category (see [\[LP08\]](#page-47-3)).

Proposition 3.2 ([\[LP08,](#page-47-3) Corollary 4.5, Corollary 4.6]). There are natural equivalences of categories $\text{Fun}^{\otimes}(\text{Cob}_2^{\text{closed}}, \mathcal{C}) \to \text{ComFrob}(\mathcal{C}),$

$$
F\mapsto (F\left(\bigcirc\right),F\left(\bigcirc\right))\bigcirc F\left(\bigcirc\right),F\left(\bigcirc\right),F\left(\bigcirc\right)\bigcirc F\left(\bigcirc\right))
$$

and

 $\text{Fun}^{\otimes}(\text{Cob}^{\text{open}}_2, \mathcal{C}) \to \text{SymFrob}(\mathcal{C}),$

$$
F\mapsto (F\left(\!\!\left|\!\!\left|\!\!\left|\!\!\right\rangle\right.\!,F\left(\!\!\left|\!\!\right\rangle\!\!\right.\right),F\left(\bigcup\!\!\right),F\left(\bigcup\!\!\right),F\left(\bigcup\!\!\right),F\left(\!\!\left|\!\!\right\rangle\!\!\right),F\left(\!\!\left|\!\!\right\rangle\!\!\right)).
$$

We can also apply Proposition [2.8](#page-9-2) to get a characterisation in terms of graph cobordisms.

Corollary 3.3. Let C be a symmetric monoidal category. There are natural equivalences of categories

Fun[®](GCob,
$$
C
$$
) \rightarrow ComFrob(C),
\n $F \mapsto (F({*}), F(\rightarrow), F(\rightarrow), F(\rightarrow), F(\rightarrow))$

and

Fun[®](fGCob, C)
$$
\rightarrow
$$
 SymFrob(C),
\n $F \mapsto (F(\{\ast\}), F(\succ), F(\blacklozenge), F(\prec), F(\rightarrow)).$

 $Here \rightarrow$ denotes the following graph

where we do not denote the external vertices. Similarly, \bullet denotes the following graph

without the external vertex. The graphs \prec and \rightarrow are defined as their respective duals. For a functor F out of $GCob$ (or $fGCob$), we define the following notation:

One advantage of the graph models of cobordism categories GCob and fGCob compared to the geometric cobordism categories $\mathrm{Cob}^{\mathrm{closed}}_2$ and $\mathrm{Cob}^{\mathrm{open}}_2$ is that it becomes convenient to state in the characterisation via PROPs that all commutative Frobenius algebras are symmetric.

Proposition 3.4. There is a canonical symmetric monoidal functor $fGCob \rightarrow GCob$ given by forgetting the fat graph structure. Precomposition with this functor encodes the forgetful functor $ComFrob(\mathcal{C}) \rightarrow SymFrob(\mathcal{C})$ for all symmetric monoidal categories \mathcal{C} .

3.1. Graded Context. A first attempt to define a Frobenius algebra in a graded context may be to give a functor out of the cobordism categories we described until now. This fails to describe certain interesting examples in the graded context. We first specify what we mean by the graded context.

Let R be a commutative ring. We consider two notions of categories of graded R -module. Let RMod• be the category of Z-graded R-modules $A = \{A_i\}_{i \in \mathbb{Z}}$ with morphisms

$$
R\text{Mod}_{\bullet}(A, B) = \{R\text{Mod}_{d}(A, B)\}_{d \in \mathbb{Z}},
$$

$$
RModd(A, B) = \{ \phi = (\phi_i)_{i \in \mathbb{Z}} \mid \phi_i \colon A_i \to B_{i+d} \text{ } R \text{-linear} \}.
$$

On the other hand, we denote by $R\text{Mod}_0$ the category of Z-graded R-modules with morphisms $RMod_0(A, B)$.

Both are symmetric monoidal categories with unit

$$
\mathbb{1}_d := \begin{cases} \mathbb{Z} & \text{if } d = 0, \\ 0 & \text{else,} \end{cases}
$$

and the tensor product:

$$
(A \otimes B)_k = \bigoplus_{i+j=k} A_i \otimes_R B_j
$$

with the Koszul twist induced by

$$
\tau: A_i \otimes B_j \to B_j \otimes A_i,
$$

$$
a \otimes b \mapsto (-1)^{ij}b \otimes a.
$$

Recall that we denote the shift ΣA as the object with $(\Sigma A)_{i+1} = A_i$. We note that both categories $R\text{Mod}_9$ and $R\text{Mod}_9$ are enriched over $R\text{Mod}_9$. Furthermore, if R is an S-algebra, both are enriched over $SMod_0$.

We fix a monoidal functor $F: GCob \to RMod_{\bullet}$ (or $F: fGCob \to RMod_{\bullet}$) enriched over sets. In particular, we allow all degrees for $\mu = F(\rightarrow)$ and $\nu = F(\rightarrow)$. We note that unitality implies that $|\mu| + |\eta| = |id| = 0$ and counitality implies $|\nu| + |\varepsilon| = |id| = 0$.

Proposition 3.5. Let R be a principal ideal domain. Let F: fGCob \rightarrow RMod_• be a symmetric monoidal functor. Then A is free of finite rank.

Proof. A consequence of unitality, counitality and the Frobenius relation is that

(3) $(p \otimes id) \circ (id \otimes c) = id = (id \otimes p) \circ (c \otimes id).$

This means that A is dualizable which is equivalent to A being projective and finitely generated. As R is a principal ideal domain, this is equivalent to A being free of finite rank. \Box

Proposition 3.6. Let R be a principal ideal domain of characteristic not 2.

Let F: fGCob \rightarrow RMod_• or F: fGCob \rightarrow RMod_• be a symmetric monoidal functor. If $|\mu| - |\nu|$ is odd, then A is the zero module.

Proof. By Proposition [3.4,](#page-10-0) it suffices to show the statement for fGCob and, by Proposition [3.5,](#page-11-2) we can fix a basis $\alpha_1, \ldots, \alpha_k$ of A.

We note that $n := |p| = |\mu| + |\varepsilon| = |\mu| - |\nu|$ and $|c| = |\nu| + |\eta| = |\nu| - |\mu| = -n$ are both of odd degree. We write

$$
c(1) = \sum_j \alpha_j \otimes \beta_j
$$

for some $\beta_j \in A$. By duality, β_1, \ldots, β_k also defines a basis. The fact that $|c| = -n$ shows that $|\beta_i| = -|\alpha_i| - n.$

If $A \neq 0$, we may assume that $k > 0$ and thus $\alpha_1 \neq 0$. We want to exploit [\(3\)](#page-11-3) for α_1 and β_1 . We first apply the right hand side to α_1 :

$$
\alpha_1 = (\mathrm{id} \otimes p) \circ (c \otimes \mathrm{id})(\alpha_1) = (\mathrm{id} \otimes p) \left(\sum_j \alpha_j \otimes \beta_j \otimes \alpha_1 \right) = \sum_j (-1)^{n |\alpha_j|} \alpha_j p(\beta_j \otimes \alpha_1).
$$

As $\alpha_1, \ldots, \alpha_k$ is a basis, we must have

(4)
$$
p(\beta_1 \otimes \alpha_1) = (-1)^{n|\alpha_1|}.
$$

On the other hand, when we apply the left hand side of [\(3\)](#page-11-3) to β_1 , we find

$$
\beta_1 = (p \otimes id) \circ (id \otimes c)(\beta_1) = (p \otimes id) \left((-1)^{n|\beta_1|} \beta_i \otimes \sum_j \alpha_j \otimes \beta_j \right) = (-1)^{n|\beta_i|} \sum_j p(\beta_1 \otimes \alpha_j) \beta_j.
$$

As β_1, \ldots, β_k is a basis, we must have

(5)
$$
p(\beta_1 \otimes \alpha_1) = (-1)^{n(-n-|\alpha_1|)} = (-1)^{n+n|\alpha_1|}.
$$

As n is odd and $(-1)^n = -1 \neq 1$ in R, we have the desired contradiction between [\(4\)](#page-12-2) and [\(5\)](#page-12-3). Therefore there is no element $\alpha_1 \neq 0$ and thus $A = 0$.

In section [6,](#page-24-1) we give examples of structures that look like Frobenius algebras that have $|\mu| - |\nu|$ odd and non-trivial. In other words, this characterisation of a Frobenius algebra misses key examples.

4. Graded Frobenius Algebras: PROPs

The goal of this section is to define cobordism categories which already encode the signs and degrees. Concretely, we define categories $\text{GCob}_{c,d}$ and $\text{fGCob}_{c,d}$ enriched over graded abelian groups where the product \rightarrow has degree c and the coproduct \rightarrow degree d. This gives us a new definition of graded commutative or symmetric Frobenius algebras as functors out of $\text{GCob}_{c,d}$ or $\text{fGCob}_{c,d}$, respectively, to a category enriched over graded abelian groups \mathbb{ZMod}_{0} .

In subsection [4.2,](#page-15-2) we prove some properties immediately from the PROP definition: suspending an algebra over $\text{GCob}_{c,d}$ (or $\text{fGCob}_{c,d}$) gives an algebra in $\text{GCob}_{c-1,d+1}$ (or $\text{fGCob}_{c-1,d+1}$). Tensoring algebras over $\text{GCob}_{c,d}$ and $\text{GCob}_{c',d'}$ (or $\text{fGCob}_{c,d}$ and $\text{fGCob}_{c',d'}$) gives an algebra over $\text{GCob}_{c+c',d+d'}$ (or $\text{fGCob}_{c+c',d+d'}$).

In subsection [4.3,](#page-19-3) we study the canonical subdioperad of $GCob_{c,d}$ which is generated by forests. If $c + d$ is odd, then this subdioperad has 2-torsion cokernel in the full PROP. This dioperad naturally shows up to describe an infinite-dimensional structure carrying a Frobenius relation as in [6.4.](#page-32-0)

4.1. Definition. In the following computations and definitions, we heavily use the concept of a top exterior power of an abelian group. Unfortunately, we require multiple slightly different versions of a top exterior power. We hope to ameliorate this situation by giving a clear overview of the versions we use.

- Let M be a free abelian group of rank n , we denote
	- $\text{Or}(M) := \Lambda^n M$ the top exterior power as a graded abelian group concentrated in degree 0;
	- det(M) := Σ^{-n} Or(M) the top exterior power as a graded abelian group concentrated in degree $-n$.
- Let A be a free graded abelian group of rank 1, it is invertible with its inverse A^{-1} := $\mathbb{ZMod}_{\bullet}(A, \mathbb{1}).$

• Let $A = \{A_i\}_{i \in \mathbb{Z}}$ be a free graded abelian group of finite rank, we denote

$$
\det(A) := \bigotimes_{i \in \mathbb{Z}} \det(A_i)^{(-1)^i}.
$$

- Let G be a graph and $c, d \in \mathbb{Z}$, we denote
	- $-$ det $(G, \partial_{in}) :=$ det $(H_*(|G|, |L_{in}(G)|));$
	- $\det(G, \partial_{out}) := \det(H_*(|G|, |L_{out}(G)|));$
	- $\,\, \det_{c,d} (G) := \det (G, \partial_{in})^{\otimes c} \otimes \det (G, \partial_{out})^{\otimes d}.$

Lemma 4.1. Let $A_{\bullet}, B_{\bullet}, C_{\bullet} \in \mathbb{Z}$ Mod_{\bullet} free of finite total rank. Then the following holds:

- (a) the determinant $\det(A_{\bullet})$ is concentrated in degree $-\chi(A)$ where $\chi(A)$ is the Euler characteristic of A_{\bullet} :
- (b) there exists a natural isomorphism

$$
\det(A_{\bullet} \oplus B_{\bullet}) \cong \det(A_{\bullet}) \otimes \det(B_{\bullet});
$$

(c) there exists a natural isomorphism

$$
\det(A_{\bullet} \oplus \Sigma A_{\bullet}) \cong \mathbb{1}.
$$

This implies a natural isomorphism $\det(A)^{-1} \cong \det(\Sigma A)$;

- (d) if $0 \to A_{\bullet} \to B_{\bullet} \to C_{\bullet} \to 0$ is a short exact sequence, there exists a canonical isomorphism $\det(A_{\bullet}) \otimes \det(C_{\bullet}) \cong \det(B_{\bullet});$
- (e) if ∂ a differential on A, such that $H_{\bullet}(A) = H_{\bullet}(A, \partial)$ and A_{\bullet} are finitely generated free Zmodules, then there exists a natural isomorphism:

$$
\det(A_{\bullet}) \cong \det(H_{\bullet}(A)).
$$

Proof. Statements [\(a\)](#page-13-0), [\(b\)](#page-13-1) and [\(c\)](#page-13-2) follow directly from the definition and the following facts for finitely generated free Z-modules M_1, \ldots, M_k :

$$
\det\left(\bigoplus_{i=1}^k M_i\right) \cong \bigotimes_{i=1}^k \det(M_i) \text{ and } \det(\Sigma M_i) \cong \det(M_i)^{-1}.
$$

For statement [\(d\)](#page-13-3), we note that we can choose a splitting $f: C_{\bullet} \to B_{\bullet}$ because C_{\bullet} is free. This gives an isomorphism

$$
\det(A_{\bullet}) \otimes \det(C_{\bullet}) \cong \det(A_{\bullet} \oplus C_{\bullet}) \cong \det(B_{\bullet}).
$$

This is independent of the choice of splitting because two splittings of $B_{\bullet} \to C_{\bullet}$ differ by elements in A_{\bullet} and thus the difference between the two induced isomorphisms is zero after tensoring with $\det(A_{\bullet}).$

For statement [\(e\)](#page-13-4), the short exact sequence

$$
0 \to \text{im}(\partial: A_{\bullet+1} \to A_{\bullet}) \to \text{ker}(\partial: A_{\bullet} \to A_{\bullet-1}) \to H_{\bullet}(A) \to 0
$$

gives

$$
\begin{aligned}\n\det(\ker(\partial \colon A_{\bullet} \to A_{\bullet-1})) &\stackrel{\text{(d)}}{\cong} \det(\text{im}(\partial \colon A_{\bullet+1} \to A_{\bullet})) \otimes \det(H_{\bullet}(A)) \\
&\implies \det(H_{\bullet}(A)) \cong \det(\ker(\partial \colon A_{\bullet} \to A_{\bullet-1})) \otimes \det(\text{im}(\partial \colon A_{\bullet+1} \to A_{\bullet}))^{-1} \\
&\stackrel{\text{(c)}}{\cong} \det(\ker(\partial \colon A_{\bullet} \to A_{\bullet-1})) \otimes \det(\Sigma \text{ im}(\partial \colon A_{\bullet+1} \to A_{\bullet})) \\
&\cong \det(\ker(\partial \colon A_{\bullet} \to A_{\bullet-1})) \otimes \det(\text{im}(\partial \colon A_{\bullet} \to A_{\bullet-1})).\n\end{aligned}
$$

On the other hand, the short exact sequence

$$
0 \to \ker(\partial \colon A_{\bullet} \to A_{\bullet-1}) \to A_{\bullet} \stackrel{\partial}{\to} \text{im}(\partial \colon A_{\bullet} \to A_{\bullet-1}) \to 0
$$

gives

$$
\det(A) \stackrel{\text{(c)}}{\cong} \det(\ker(\partial \colon A_\bullet \to A_{\bullet-1})) \otimes \det(\text{im}(\partial \colon A_\bullet \to A_{\bullet-1}))
$$

which concludes the proof. \Box

We note that a morphism of graph cobordisms $G \to G'$ induces homotopy equivalences

$$
(|G|, |L_{in}(G)|) \to (|G'|, |L_{in}(G')|),
$$

$$
(|G|, |L_{out}(G)|) \to (|G'|, |L_{out}(G')|)
$$

and thus isomorphisms

$$
\det(G, \partial_{in}) \to \det(G', \partial_{in}),
$$

\n
$$
\det(G, \partial_{out}) \to \det(G', \partial_{out}),
$$

\n
$$
\det_{c,d}(G) \to \det_{c,d}(G').
$$

Therefore $\det_{c,d}(G)$ only depends on the class $[G] \in \pi_0(\underline{GCob}(X,Y)) = GCob(X,Y)$. Moreover, we can glue orientations together:

Lemma 4.2. Let G, G' be composable graphs in \underline{GCob} , there is a natural isomorphism

 $\mathrm{comp}\colon \mathbf{det}_{c,d}(G')\otimes \mathbf{det}_{c,d}(G)\cong \mathbf{det}_{c,d}(G'\circ G).$

Proof. We recall:

$$
\begin{aligned} \text{det}_{c,d}(G')\otimes\text{det}_{c,d}(G)&\cong\text{det}(G',\partial_{in})^{\otimes c}\otimes\text{det}(G',\partial_{out})^{\otimes d}\otimes\text{det}(G,\partial_{in})^{\otimes c}\otimes\text{det}(G,\partial_{out})^{\otimes d}\\ &\cong\text{det}(G',\partial_{in})^{\otimes c}\otimes\text{det}(G,\partial_{in})^{\otimes c}\otimes\text{det}(G',\partial_{out})^{\otimes d}\otimes\text{det}(G,\partial_{out})^{\otimes d}\end{aligned}
$$

and

$$
\mathbf{det}_{c,d}(G' \circ G) \cong \mathbf{det}(G' \circ G, \partial_{in})^{\otimes c} \otimes \mathbf{det}(G' \circ G, \partial_{out})^{\otimes d}.
$$

It thus suffices to give isomorphisms

$$
\begin{aligned}\n\det(G',\partial_{in}) \otimes \det(G,\partial_{in}) &\cong \det(G' \circ G,\partial_{in}) \\
\det(G',\partial_{out}) \otimes \det(G,\partial_{out}) &\cong \det(G' \circ G,\partial_{out})\n\end{aligned}
$$

We give the first isomorphism, the second one is constructed analogously. We can canonically embed $|G| \subseteq |G' \circ G|$. Let $C_{\bullet}(|G|, |\partial_{in}|)$, $C_{\bullet}(|G'|, |\partial_{in}|)$, $C_{\bullet}(|G' \circ G|, |\partial_{in}|)$ and $C_{\bullet}(|G' \circ G|, |G|)$ be the cellular chain complexes of the pairs. We note that the inclusion

$$
C_{\bullet}(|G'\circ G|,|G|)\to C_{\bullet}(|G'|,|\partial_{in}|)
$$

is an isomorphism.

The triple of spaces $(|G' \circ G|, |G|, |\partial_{in}|)$ gives a short exact sequence of chain complexes:

$$
0 \longrightarrow C_{\bullet}(|G|,|\partial_{in}|) \longrightarrow C_{\bullet}(|G' \circ G|,|\partial_{in}|) \longrightarrow \overbrace{C_{\bullet}(|G' \circ G|,|G|)}^{C_{\bullet}(|G'|,|\partial_{in}|)} \longrightarrow 0.
$$

Lemma [4.1](#page-13-5) [\(d\)](#page-13-3) and [\(e\)](#page-13-4) now give the desired isomorphism. \Box

Definition 4.3. Let $c, d \in \mathbb{Z}$.

The category $\text{GCob}_{c,d}$ is defined with objects finite sets and morphism spaces

$$
\mathrm{GCob}_{c,d}(X,Y) = \underset{\underline{\mathrm{GCob}}(X,Y)}{\mathrm{colim}} \mathrm{\bf det}_{c,d}(G).
$$

The category $fGCob_{c,d}$ is defined with objects finite sets and morphism spaces

$$
\mathrm{fGCob}_{c,d}(X,Y)=\operatornamewithlimits{colim}_{\underline{\mathrm{fGCob}}(X,Y)}\mathbf{det}_{c,d}(G).
$$

Lemma [4.2](#page-14-0) lets us define the composition in these categories.

We can now define a graded Frobenius algebra:

Definition 4.4. Let C be a symmetric monoidal category enriched over $\mathbb{Z} \text{Mod}_0$ and let $c, d \in \mathbb{Z}$. A (c, d) -graded commutative Frobenius algebra is a symmetric monoidal functor $F: {\rm GCob}_{c,d} \to \mathcal{C}$ enriched over \mathbb{ZMod}_0 i.e. degree-preserving on morphism spaces. The category is denoted by

$$
\mathrm{grComFrob}_{c,d}(\mathcal{C}) = \mathrm{Fun}_{\mathbb{Z}\mathrm{Mod}_0}^{\otimes}(\mathrm{GCob}_{c,d}, \mathcal{C}).
$$

Analogously, a (c, d) -graded symmetric Frobenius algebra is given by such a functor $F: GCob_{c,d} \to \mathcal{C}$ and the category is denoted by

$$
\mathrm{gr} \mathrm{SymFrob}_{c,d}(\mathcal{C}) = \mathrm{Fun}^\otimes_{\mathbb{Z}\mathrm{Mod}_0}(\mathrm{fGCob}_{c,d}, \mathcal{C}).
$$

4.2. Properties. In this subsection, we prove some first properties of these category. The definitions of the categories $\text{GCob}_{c,d}$ and $\text{fGCob}_{c,d}$ we gave are admittedly hard to work with. However, the morphism spaces can be described in a more convenient way.

Proposition 4.5. There exist isomorphisms

$$
\mathrm{GCob}_{c,d}(X,Y)\cong \bigoplus_{[G]\in \mathrm{GCob}(X,Y))}\mathbf{det}_{c,d}(G)_{\pi_1(|\underline{\mathrm{GCob}}(X,Y)|,G)}
$$

and

$$
\mathrm{fGCob}_{c,d}(X,Y)\cong \bigoplus_{[G]\in \mathrm{fGCob}(X,Y))}\mathbf{det}_{c,d}(G)_{\pi_1(|\underline{\mathrm{fGCob}}(X,Y)|,G)}.
$$

Proof. We note that the functor $\det_{c,d}$ sends all maps in $\text{GCob}(X, Y)$ and $\text{fGCob}(X, Y)$ to isomorphisms. Therefore, $\det_{c,d}$ factors through the localisations $GCob(X, Y) \to |GCob(X, Y)|$ and $fG\text{Cob}(X, Y) \to [fG\text{Cob}(X, Y)]$. Here |C| is interpreted as the ∞ -groupoid given by the category C with all morphisms formally inverted.

We can thus compute the colimit giving the morphism space between finite sets X and Y as follows:

$$
\operatorname{colim}_{G \in \underline{\mathrm{GCob}}(X,Y)} \mathbf{det}_{c,d}(G) = \operatorname{colim}_{G \in |\underline{\mathrm{GCob}}(X,Y)|} \mathbf{det}_{c,d}(G)
$$

The functor $\det_{c,d}$ lands in a 1-category. Moreover, two graphs $G, G' \in \underline{GCob}(X, Y)$ are equivalent in $|GCob(X, Y)|$ if and only if they represent the same element in $GCob(X, Y) = \pi_0(GCob(X, Y)).$ This shows

$$
\mathrm{GCob}_{c,d}(X,Y)\cong \bigoplus_{[G]\in \mathrm{GCob}(X,Y))}\mathbf{det}_{c,d}(G)_{\pi_1(|\underline{\mathrm{GCob}}(X,Y)|,G)}.
$$

Similarly for fat graphs, we have

$$
\mathrm{fGCob}_{c,d}(X,Y)\cong \bigoplus_{[G]\in \mathrm{fGCob}(X,Y)}\mathbf{det}_{c,d}(G)_{\pi_1(|\underline{\mathrm{fGCob}}(X,Y)|,G)}.
$$

Remark 4.6. The group $\pi_1(|\text{GCob}(X, Y)|, G)$ is isomorphic to $\pi_0(h\text{Aut}(|G|))$: a zig-zag of graphs morphisms is the same as a simple homotopy equivalence from the CW-complex $|G|$ to itself. For the 1-dimensional CW complex $|G|$, any homotopy equivalence is homotopic to a simple homotopy equivalence.

On the other hand, the group $\pi_1(|\text{GCob}(X, Y)|, G)$ is isomorphic to the mapping class group $\pi_0(\text{Homeo}^+(\text{fCob}(G), \partial_{in}(\text{fCob}(G))) \cup \partial_{out}(\text{fCob}(G)))$ that fixes the boundary point-wise. This can be seen for example by the open part of Theorem A in [\[San15\]](#page-47-13).

Example 4.7. In the case $c = d = 0$, we have $\det_{0,0}(G) = 1$ and the automorphisms of G act trivially. We therefore compute:

$$
\mathrm{GCob}_{0,0}(X,Y)\cong\bigoplus_{[G]\in\mathrm{GCob}(X,Y)}\mathbb{1}
$$

and

$$
\mathrm{fGCob}_{0,0}(X,Y)\cong \bigoplus_{[G]\in \mathrm{fGCob}(X,Y]}{\mathbb{1}}.
$$

We find

 $\mathrm{grComFrob}_{0,0}(R\mathrm{Mod}_\bullet)=\mathrm{Fun}^\otimes_{{\mathbb Z}\mathrm{Mod}_0}(\mathrm{GCob}_{0,0},R\mathrm{Mod}_\bullet)$ $=\text{Fun}^{\otimes}(\text{GCob}, R\text{Mod}_0)$ $=\text{ComFrob}(R\text{Mod}_0).$

Therefore a $(0, 0)$ -graded commutative Frobenius algebra coincides with the classical definition of a commutative Frobenius algebras with maps in degree 0. An analogous statement holds for symmetric Frobenius algebras.

In the following, we study the relation between $\mathrm{GCob}_{c,d}$ for different choices of $c, d \in \mathbb{Z}$. For this, we use the symmetric monoidal structure on PROPs.

Notation 4.8. We denote the *tensor product* of two PROPs in \mathbb{ZMod}_0 for the coordinate-wise tensor product: for PROPs P, Q

$$
(P \otimes Q)(X, Y) := P(X, Y) \otimes Q(X, Y)
$$

for finite sets X, Y with the component-wise composition maps.

Example 4.9. Let $(C, \otimes, 1, \tau)$ be a symmetric monoidal category enriched over ZMod₀ and let $A, B \in \mathcal{C}$ be objects. There is an isomorphism of PROPs

$$
\mathrm{End}_A\otimes\mathrm{End}_B\cong\mathrm{End}_{A\otimes B}
$$

given on maps $f: A^{\otimes k} \to A^{\otimes l}$ and $g: B^{\otimes k} \to B^{\otimes l}$ by

$$
(A \otimes B)^{\otimes k} \to A^{\otimes k} \otimes B^{\otimes k} \overset{f \otimes g}{\to} A^{\otimes l} \otimes B^{\otimes l} \to (A \otimes B)^{\otimes l}.
$$

Indeed, this map is equivariant and commutes with composition and the product ⊗.

Proposition 4.10. There exist canonical maps of PROPs

 $\mathrm{GCob}_{c+c',d+d'} \to \mathrm{GCob}_{c,d} \otimes \mathrm{GCob}_{c',d'}$

and

$$
\text{fGCob}_{c+c',d+d'} \to \text{fGCob}_{c,d} \otimes \text{fGCob}_{c',d'}.
$$

Proof. There are canonical maps

$$
\det(G, \partial_{in})^{\otimes c+c'} \otimes \det(G, \partial_{out})^{\otimes d+d'}
$$

\n
$$
\cong \det(G, \partial_{in})^{\otimes c} \otimes \det(G, \partial_{out})^{\otimes d} \otimes \det(G, \partial_{in})^{\otimes c'} \otimes \det(G, \partial_{out})^{\otimes d'}.
$$

This then gives maps

$$
\textbf{det}_{c+c',d+d'}(G)\to\operatornamewithlimits{colim}_{G'\in \underline{\textbf{GCob}}(X,Y)}\textbf{det}_{c,d}(G')\otimes\operatornamewithlimits{colim}_{G'\in \underline{\textbf{GCob}}(X,Y)}\textbf{det}_{c',d'}(G')
$$

for all $G \in \text{GCob}(X, Y)$. Taking the colimit over $\text{GCob}(X, Y)$ thus gives the desired map for the GCob. The proof for fGCob is analogous. □

Remark 4.11. Even though the map [\(6\)](#page-17-2) used in the construction is an isomorphism, the resulting map of PROPs is not a isomorphism in general. For example if $c+d$ and $c'+d'$ are odd, Proposition [4.20](#page-19-2) shows that $\text{GCob}_{c,d}$ and $\text{GCob}_{c',d'}$ is 2-torsion on summands coming from graphs which are not forests. This is not true for $\text{GCob}_{c+c',d+d'}$.

Corollary 4.12. Let A be an algebra over $\mathrm{GCob}_{c,d}$ (or $\mathrm{fGCob}_{c,d}$) and B be an algebra over $\mathrm{GCob}_{c',d'}$ (or $\mathrm{fGCob}_{c',d'}$). Then $A \otimes B$ inherits the structure of an algebra over $\mathrm{GCob}_{c+c',d+d'}$ (or $fGCob_{c+c',d+d'}$, respectively).

Proof. We show the statement for $GCob_{c,d}$. The proof for fGCob is analogous.

Giving a GCob_{c,d}-algebra structure on A is the same thing as giving a map of PROPs GCob_{c,d} \rightarrow End_A (see [\[Mar08\]](#page-47-14)). Similarly, we get a map of PROPs $\text{GCob}_{c',d'} \to \text{End}_B$. Precomposing with the map from Proposition [4.10](#page-16-0) and postcomposing with the map from Example [4.9,](#page-16-1) we thus get a map of PROPs

$$
\mathrm{GCob}_{c+c',d+d'} \to \mathrm{GCob}_{c,d} \otimes \mathrm{GCob}_{c',d'} \to \mathrm{End}_A \otimes \mathrm{End}_B \to \mathrm{End}_{A \otimes B}
$$

and thus a $\text{GCob}_{c+c',d+d'}$ -algebra structure on $A \otimes B$.

Another relation between $\text{GCob}_{c,d}$ -algebras for different c, d comes from suspending algebras. This operation has a corresponding operation on the level of PROPs:

Definition 4.13. The *suspension PROP* in \mathbb{ZMod}_{0} is defined as the endomorphism PROP End_{Σ 1} of Σ 1 in \mathbb{Z} Mod. Explicitly, it is given by

$$
\Sigma(X, Y) := \mathbb{Z} \text{Mod}_{\bullet}((\Sigma \mathbb{1})^{\otimes X}, (\Sigma \mathbb{1})^{\otimes Y})
$$

for finite sets X, Y .

For a PROP P in \mathbb{Z} Mod₀, the *suspended PROP* is defined as $\Sigma P := \Sigma \otimes P$.

Proposition 4.14. Let P be a PROP in $\mathbb{Z} \text{Mod}_0$ and $A \in \mathbb{Z} \text{Mod}_\bullet$. Then giving a P-algebra structure on A is equivalent to giving a ΣP -algebra structure on ΣA .

Proof. We again use the characterisation of A being a P-algebra as a PROP maps into End_A . By definition, this is equivalent to giving for all finite sets X, Y equivariant maps

$$
P(X, Y) \to \mathcal{C}(A^{\otimes X}, A^{\otimes Y})
$$

which commute with composition ∘ and the symmetric monoidal product \otimes . This is equivalent to giving equivariant maps

$$
\mathbb{Z}\mathrm{Mod}_{\bullet}((\Sigma\mathbb{1})^{\otimes X},(\Sigma\mathbb{1})^{\otimes Y})\otimes P(X,Y)\rightarrow \mathbb{Z}\mathrm{Mod}_{\bullet}((\Sigma\mathbb{1})^{\otimes X},(\Sigma\mathbb{1})^{\otimes Y})\otimes \mathbb{Z}\mathrm{Mod}_{\bullet}(A^{\otimes X},A^{\otimes Y})
$$

which commute with \circ and \otimes . After postcomposing with the isomorphism from Example [4.9,](#page-16-1) we get equivariant maps

$$
\Sigma P(X, Y) \to \mathbb{Z}Mod_{\bullet}((\Sigma A)^{\otimes X}, (\Sigma A)^{\otimes Y})
$$

which commute with ∘ and ⊗. This is by definition equivalent to giving a PROP morphism $\Sigma P \rightarrow$ End_{ΣA}, which is equivalent to giving a ΣP structure on ΣA . \square

Remark 4.15. This construction is completely analogous as for operads, cyclic operads, dioperads and properads. However in the category of modular operads, there exists no suspension modular operad in ZMod₀. Indeed there is a canonical pairing $\Sigma \mathbb{1} \otimes \Sigma \mathbb{1} \to \Sigma \mathbb{1}$ in degree -1. This lets us identify

$$
\mathrm{End}_{\Sigma \mathbb{1}}(X) = \mathbb{Z} \mathrm{Mod}_{\bullet}((\Sigma \mathbb{1})^{\otimes X}, \Sigma \mathbb{1}) \cong \Sigma^{1-|X|} \mathbb{1}.
$$

For any elements $x \in X$ and $y \in Y$, this gives a map

$$
\mathrm{End}_{\Sigma_1}(X)\otimes \mathrm{End}_{\Sigma_1}(Y)\to \mathrm{End}_{\Sigma_1}((X\setminus x)\sqcup (Y\setminus y))
$$

in $\mathbb{Z} \text{Mod}_0$. However, the map for elements $x \neq x' \in X$

$$
End_{\Sigma_1}(X) \to End_{\Sigma_1}(X \setminus \{x, x'\})
$$

is of degree -1 and thus not in $\mathbb{Z}Mod_0$. One way to amend this is to work with the language of twisted modular operads as introduced in [\[GK98\]](#page-47-15).

Proposition 4.16. For integers $c, d \in \mathbb{Z}$, there exist canonical isomorphisms

$$
\Sigma \mathrm{GCob}_{c,d} \cong \mathrm{GCob}_{c-1,d+1}
$$

and

$$
\Sigma f \text{GCob}_{c,d} \cong f \text{GCob}_{c-1,d+1}.
$$

In particular, suspension defines equivalences of categories

$$
\Sigma \colon \mathrm{grComFrob}_{c,d}(\mathbb{ZMod}_{\bullet}) \cong \mathrm{grComFrob}_{c-1,d+1}(\mathbb{ZMod}_{\bullet}),
$$

$$
\Sigma \colon \mathrm{grSymFrob}_{c,d}(\mathbb{ZMod}_{\bullet}) \cong \mathrm{grSymFrob}_{c-1,d+1}(\mathbb{ZMod}_{\bullet}).
$$

Proof. We fix integers $c, d \in \mathbb{Z}$ and finite sets X, Y. The key observation is that for every graph cobordism G between X and Y there is an isomorphism

(7)
$$
\mathbf{det}_{c-1,d+1}(G) \cong \mathbb{Z} \mathbf{Mod}_{\bullet}((\Sigma \mathbb{1})^{\otimes X}, (\Sigma \mathbb{1})^{\otimes Y}) \otimes \mathbf{det}_{c,d}(G)
$$

which is natural in X and Y .

To see the isomorphism, we observe

$$
\begin{aligned} \det_{c-1,d+1}(G)&=\det(G,\partial_{in})^{\otimes c-1}\otimes \det(G,\partial_{out})^{\otimes d+1}\\ &\cong \det(G,\partial_{in})^{-1}\otimes \det(G,\partial_{out})\otimes \det_{c,d}(G). \end{aligned}
$$

We consider the short exact sequences

$$
0 \longrightarrow C_{\bullet}(|\partial_{in}|) \longrightarrow C_{\bullet}(|G|) \longrightarrow C_{\bullet}(|G|, |\partial_{in}|) \longrightarrow 0,
$$

$$
0 \longrightarrow C_{\bullet}(|\partial_{out}|) \longrightarrow C_{\bullet}(|G|) \longrightarrow C_{\bullet}(|G|, |\partial_{out}|) \longrightarrow 0.
$$

Then Lemma [4.1](#page-13-5) [\(d\)](#page-13-3) gives the natural isomorphisms

$$
\begin{aligned}\n\det(G,\partial_{in})^{-1}\otimes \det(G,\partial_{out}) &\cong \det(|\partial_{in}|)\otimes \det(|G|)^{-1}\otimes \det(|G|)\otimes \det(|\partial_{out}|)^{-1} \\
&\cong \det(|\partial_{in}|)\otimes \det(|\partial_{out}|)^{-1} \\
&\cong \det(X)\otimes \det(Y)^{-1} \\
&\cong ((\Sigma\mathbb{1})^{\otimes X})^{-1}\otimes (\Sigma\mathbb{1})^{\otimes Y} \\
&\cong \mathbb{ZMod}_{\bullet}((\Sigma\mathbb{1})^{\otimes X}, (\Sigma\mathbb{1})^{\otimes Y}) = \Sigma(X,Y).\n\end{aligned}
$$

We recall that, in our convention, the determinant lives in the degree equal to the negative rank. Thus, we have for a finite set X a natural isomorphism $\det(X) \cong ((\Sigma \mathbb{1})^X)^{-1}$. This gives [\(7\)](#page-18-1) and thus the proof. \Box

Remark 4.17. We note that $|\mu| + |\nu| = |\Sigma(\mu)| + |\Sigma(\nu)|$. Therefore this defines an invariant under suspension. In particular if $|\mu| + |\nu| \neq 0$, the algebra is not a suspension of an algebra with operations in degree 0. The machinery of twisting by $\det(G, \partial_{in})$ (or $\det(G, \partial_{out})$) helps us describe other algebras because this only changes the degree of μ and not ν (or vice-versa).

4.3. Canonical Dioperad. The PROP $GCob_{c,d}$ has a canonical subdioperad generated by forests. In subsection [6.4,](#page-32-0) we study an example where this dioperad shows up naturally rather than the whole PROP. We study the structure of this canonical subdioperad.

For this, we recall the definition of a dioperad:

Definition 4.18. A *dioperad* consists of the following data a collection $\mathcal{D}(X, Y)$ for all finite sets X, Y with

- a (S_X, S_Y) -bimodule structure on $\mathcal{D}(X, Y)$;
- a unit

$$
1\colon 1\to \mathcal{D}(\{*\}, \{*\});
$$

• composition maps

$$
{}_{y}\circ_{x}:\mathcal{D}(Y,Z)\otimes\mathcal{D}(W,X)\to\mathcal{D}((Y\setminus\{y\})\sqcup W,Z\sqcup(X\setminus\{x\}))
$$

for all finite sets W, X, Y, Z and elements $x \in X, y \in Y$.

This data has to satisfy some unitality, associativity and equivariance relations. See [\[Gan02\]](#page-47-16) for the precise formulation of those relations.

Definition 4.19. The forest dioperad $\text{GCob}_{c,d}^{forest}$ of $\text{GCob}_{c,d}$ is given by the collections

(8)
$$
\mathrm{GCob}_{c,d}^{forest}(X,Y) \subseteq \mathrm{GCob}_{c,d}(X,Y)
$$

generated by the graph cobordisms between X and Y which are forests with the restricted composition.

In general this dioperad carries less information than the PROP. But in some cases, we have the following proposition:

Proposition 4.20. If $c, d \in \mathbb{Z}$ are of different parity, then the PROP GCob_{c,d} is an extension by $\mathbb{Z}/2$ factors of dioperad $\mathrm{GCob}^{forest}_{c,d}$. In explicit terms, the inclusion [\(8\)](#page-19-4) finite sets X, Y

$$
\text{GCob}^{forest}_{c,d}(X,Y) \to \text{GCob}_{c,d}(X,Y)
$$

has cokernel which is 2-torsion.

Proof. We fix $c, d \in \mathbb{Z}$ of different parity. We use the description

$$
\mathrm{GCob}_{c,d}(X,Y) \cong \bigoplus_{[G]\in \mathrm{GCob}(X,Y)} \mathbf{det}_{c,d}(G)_{\pi_1(|\underline{\mathrm{GCob}}(X,Y)|,G)}
$$

as in Proposition [4.5.](#page-15-3) We have an analogous description:

$$
\mathrm{GCob}^{forest}_{c,d}(X,Y)\cong \bigoplus_{\substack{[G]\in \mathrm{GCob}(X,Y)\\G\text{ a forest}}}\det{}_{c,d}(G)_{\pi_1(|\underline{\mathrm{GCob}}(X,Y)|,G)}.
$$

Let $[G] \in \text{GCob}(X, Y)$ be an equivalence class of graphs cobordisms between X and Y with G not a forest. We want to show that the summand

$$
\mathbf{det}_{c,d}(G)_{\pi_1(|\mathbf{G}\mathbf{Cob}(X,Y)|,G)}
$$

is $\mathbb{Z}/2$.

After collapsing some edges, we can assume that G has a tadpole. Then there is a graph automorphism of G flipping the direction of the tadpole. This represents an element in $\pi_1(|\text{GCob}(X, Y)|, G)$. This automorphism acts by $(-1)^c$ on $\det(G, \partial_{in})^{\otimes c}$ and by $(-1)^d$ on $\det(G, \partial_{out})^{\otimes d}$. It thus acts by $(-1)^{c+d} = -1$ on the one-dimensional space $\det_{c,d}(G)$. This shows $\det_{c,d}(G)_{\pi_1(|\text{GCob}(X,Y)|,G)} = \mathbb{Z}/2$ and thus concludes the proof. \Box

5. Graded Frobenius Algebra: Maps and Relations

In this section, we state the main theorem which gives a description of graded Frobenius algebras in terms of maps and relations. We will use this theorem to study some of the simplest examples of graded Frobenius algebras. Moreover, we discuss the suspension of algebras from the perspective of maps and relations.

Theorem 5.1. Let $(C, \otimes, 1, \tau)$ be a symmetric monoidal category enriched over ZMod₀ and let $c, d \in \mathbb{Z}$.

The data $(A, \mu, \eta, \nu, \varepsilon)$ where

$$
A \in \mathcal{C},
$$

\n
$$
\mu \in \mathcal{C}_c(A \otimes A, A),
$$

\n
$$
\eta \in \mathcal{C}_{-c}(\mathbb{1}, A),
$$

\n
$$
\eta \in \mathcal{C}_{-c}(\mathbb{1}, A),
$$

\n
$$
\varepsilon \in \mathcal{C}_{-d}(A, \mathbb{1})
$$

uniquely defines a (c, d) -graded commutative Frobenius algebra if and only if the relations [\(i\)](#page-2-1)-[\(vi\)](#page-2-2) as stated in the introduction are satisfied and defines a (c, d)-graded symmetric Frobenius algebra if and only if the maps satisfy the same relations replacing [\(vi\)](#page-2-2) by (vi') as stated in the introduction.

We postpone the proof of this theorem to section [7](#page-33-2) as it involves a lot of technical details. Instead we focus on what are the choices involved in those signs and what are the consequences. The strategy for the theorem is to choose generators of the categories $\text{GCob}_{c,d}$ and $\text{fGCob}_{c,d}$ and study the generating relations between them. For this, we choose generators in $\det_{c,d}(G)$ for a list

of graphs G such that all other graphs can be decomposed into them.

Definition 5.2. A generator $\omega(G) \in \det_{c,d}(G)$ is called an *orientation* of the graph G.

Remark 5.3 (Choices of Orientations). The relations are not canonical. Namely, they depend on a choice of orientations

 $\omega(\rightarrow) \in \det_{c,d}(\rightarrow), \ \omega(\bullet) \in \det_{c,d}(\bullet), \ \omega(\rightarrow) \in \det_{c,d}(\rightarrow), \ \omega(\rightarrow) \in \det_{c,d}(\bullet).$

By definition, this depends on generators of

We thus choose generators $\omega_{in}(G) \in \det(G, \partial_{in})$ and $\omega_{out}(G) \in \det(G, \partial_{out})$ for those four graphs and then define $\omega(G) := \omega_{in}(G)^{\otimes c} \otimes \omega_{out}(G)^{\otimes d}$.

We note that $\det(\rightarrow, \partial_{out}) \cong \det(\leftarrow, \partial_{out}) \cong \det(\leftarrow, \partial_{in}) \cong \det(\leftarrow, \partial_{in}) \cong \mathbb{1}$. We thus choose the canonical generator 1. Moreover for $\det(\div, \partial_{in})$ and $\det(\div, \partial_{out})$, we choose the canonical generator

coming from the canonical generator of $H_0(\bullet, \partial_{in})$ and $H_0(\bullet, \partial_{out})$. However, there is no canonical choice for generators

$$
\omega_{in}(\mathbf{L}) \in \mathbf{det}(\mathbf{L}, \partial_{in}), \qquad \omega_{out}(\mathbf{L}) \in \mathbf{det}(\mathbf{L}, \partial_{out}).
$$

Therefore, there exists no canonical choice of orientation

$$
\omega(\succ):=\omega_{in}(\succ)^{\otimes c}\otimes 1^{\otimes d}
$$

for c odd and no canonical choice of orientation

$$
\omega(\mathbf{a}) := 1^{\otimes c} \otimes \omega_{out}(\mathbf{a})^{\otimes d}
$$

for d odd. We make a choice in the proof in [7.2.](#page-37-0) A different choice would lead to different signs in the unitality and counitality relation.

Remark 5.4. In [\[CO22b\]](#page-47-5), Cieliebak and Oancea introduce a type of algebra called biunital coFrobenius. In the finite dimensional case, this corresponds to our notion of graded Frobenius algebra (see [\[CO22b,](#page-47-5) Proposition 5.14]). The infinite dimensional case is discussed in [6.4.](#page-32-0) They define the graded unitality relation

$$
\mu \circ (\eta \otimes id) = id = (-1)^c \mu \circ (id \otimes \eta)
$$

compared to our graded unitality relation

$$
(-1)^{c}\mu \circ (\eta \otimes id) = (-1)^{\frac{c(c-1)}{2}}id = \mu \circ (id \otimes \eta).
$$

The two relations agree for $c \equiv 3, 4 \pmod{4}$ and are different for $c \equiv 1, 2 \pmod{4}$. Similarly, the signs in their graded counitality relation agree with our signs for $d \equiv 3.4 \pmod{4}$ and are different for $d \equiv 1, 2 \pmod{4}$.

For $(A, \mu, \eta, \nu, \varepsilon)$ satisfying the signs in our relations, the product

$$
\mu' := (-1)^{\frac{c(c+1)}{2}} \mu
$$

and the coproduct

$$
\nu' := (-1)^{\frac{d(d+1)}{2}} \nu
$$

together with the same unit and counit satisfy the signs in their relations.

Corollary 5.5. Let $(C, \otimes, \mathbb{1}, \tau)$ be a symmetric monoidal category enriched over $\mathbb{Z}Mod_0$ and let $c, d \in \mathbb{Z}$.

Let the data $(A, \mu, \eta, \varepsilon)$ be as in Theorem [5.1.](#page-20-0) This data uniquely defines a (c, d) -graded commutative Frobenius algebra if and only if it satisfies graded associativity as in [\(i\)](#page-2-1), graded unitality as in [\(ii\)](#page-2-3), graded commutativity as in [\(vi\)](#page-2-2) and the map

$$
A\otimes A\stackrel{\mu}{\to}A\stackrel{\varepsilon}{\to}\mathbbm{1}
$$

is a non-degenerate pairing.

Moreover, it uniquely defines a (c, d) -graded symmetric Frobenius algebra if and only if it satisfies graded associativity, graded unitality, graded symmetry and the map

$$
A\otimes A\stackrel{\mu}{\to} A\stackrel{\varepsilon}{\to}\mathbbm{1}
$$

is a non-degenerate pairing.

Proof. The proof strategy is analogous to the ungraded situation as in [\[Koc04,](#page-47-4) Section 2.4]: We first use Theorem [5.1.](#page-20-0) To show that a Frobenius algebra $(A, \mu, \eta, \nu, \varepsilon)$ as in the theorem induces a non-degenerate pairing $p := \varepsilon \circ \mu$, one can show that $q := (-1)^{cd + \frac{c(c+1)}{2} + \frac{d(d+1)}{2}} \nu \circ \eta$ is the corresponding copairing that satisfies

$$
(\mathrm{id} \otimes p) \circ (q \otimes \mathrm{id}) = \mathrm{id}.
$$

On the other hand, we assume that the data $(A, \mu, \eta, \varepsilon)$ is as in the statement. Let $q: 1 \to A \otimes A$ be the associated copairing. Then one can show that the comultiplication $\nu := (-1)^{\frac{d(d+1)}{2}} (id \otimes \mu) \circ (q \otimes id)$ satisfies coassociativity as in [\(iii\)](#page-2-4) and counitality as in [\(iv\)](#page-2-5).

Remark 5.6. Another perspective on graded Frobenius algebras is the following. Assume, we are given a degree 0 multiplication μ and a non-degenerate pairing p with associated copairing q in degree $-d$ and d. Then one can use the language of oriented marked ribbon quivers as in [\[KTV21,](#page-47-17) Section 6. In this language, one can define a coproduct as the following quiver with a 0-orientation:

The signs in coassociativity and the Frobenius relation then come from permutations in the orientations.

We can now study the simplest non-trivial $\text{GCob}_{c,d}$ -algebras. The simplest example is given for $c = d = 0$:

Example 5.7. In C, the unit 1 has a canonical structure of a $GCob_{0.0}$ -algebra given by the canonical maps

$$
\mu: \mathbb{1} \otimes \mathbb{1} \to \mathbb{1},
$$

\n
$$
\nu: \mathbb{1} \to \mathbb{1} \otimes \mathbb{1},
$$

\n
$$
\varepsilon: \mathbb{1} \to \mathbb{1}.
$$

\n
$$
\varepsilon: \mathbb{1} \to \mathbb{1}.
$$

Example 5.8. Let A be a $GCob_{c,d}$ in RMod. for some principal ideal domain R. We have unit maps $R \to A_{-c}$ and counit map $A_d \to R$. If A is not trivial, non-degeneracy implies thus that we have a factor R in degree $-c$ generated by $x := \eta(1)$ and a factor R in degree d generated by y with $\varepsilon(y) = 1_R.$

We distinguish between two cases: $c + d = 0$ and $c + d \neq 0$.

For $c + d = 0$, we apply Proposition [4.16](#page-18-0) to Example [5.7](#page-22-0) to find that $\Sigma^d \mathbb{1}$ inherits the structure of a $GCob_{c,d}$ -algebra. This is thus the simplest non-trivial $GCob_{c,d}$ -algebra in $RMod_{\bullet}$. Moreover, it is the simplest non-trivial $fGCob_{c,d}$ -algebra.

For $c+d \neq 0$, the algebra A is not the suspension of a $\text{GCob}_{0,0}$ -algebra. Moreover, every non-trivial $\text{GCob}_{c,d}$ -algebra in $R\text{Mod}_{\bullet}$ thus contains at least $\Sigma^{-c}R \oplus \Sigma^{d}R$. It turns out that we can define a $\text{GCob}_{c,d}$ -algebra structure on $R_{c,d} := \Sigma^{-c} R \oplus \Sigma^{d} R$.

We denote the generator of $\Sigma^{-c}R$ by x and the generator of $\Sigma^{d}R$ by y. By assumption, we have $\eta(1) = x$ and $\varepsilon(y) = 1$. The multiplication μ is then uniquely defined by unitality

$$
\mu(x \otimes y) = (-1)^{\frac{c(c+1)}{2}}y, \qquad \mu(y \otimes x) = (-1)^{cd + \frac{c(c-1)}{2}}y, \qquad \mu(x \otimes x) = (-1)^{\frac{c(c+1)}{2}}x
$$

and degree reasons

$$
\mu(y\otimes y)=0.
$$

On the other hand, the comultiplication ν is defined by counitality as

$$
\nu(x) = (-1)^{cd + \frac{d(d+1)}{2}} x \otimes y + (-1)^{\frac{d(d-1)}{2}} y \otimes x, \qquad \nu(y) = (-1)^{\frac{d(d-1)}{2}} y.
$$

One can check that this satisfies graded associativity, graded coassociativity, the graded Frobenius relation and graded commutativity.

This is also the simplest non-trivial $fGCob_{c,d}$ -algebra.

Remark 5.9. As discussed in Remark [4.17,](#page-19-0) twisting by $\det(G, \partial_{in})$ and $\det(G, \partial_{out})$ changes only the degree of one operation, either multiplication or comultiplication while suspension changes degrees of both. On the level of algebras, suspension is simpler than twisting by $\det(G, \partial_{in})$. We have an equivalence

$$
\mathrm{grComFrob}_{c,d}(\mathbb{ZMod}_{\bullet}) \cong \mathrm{grComFrob}_{c-1,d+1}(\mathbb{ZMod}_{\bullet})
$$

given by suspending the algebra. One can think of this in the following way. We tensor an algebra A in grComFrob_c_d(ZMod_•) with the simplest non-trivial GCob_{−1,1}-algebra Σ 1. We then apply Corollary [4.12](#page-17-3) to get a $\text{GCob}_{c-1,d+1}$ -algebra structure on $\Sigma \otimes A \cong \Sigma A$. We can do an analogous construction to get a functor

$$
\mathrm{grComFrob}_{c,d}(\mathbb{ZMod}_{\bullet}) \to \mathrm{grComFrob}_{c,d+1}(\mathbb{ZMod}_{\bullet}).
$$

For this we tensor a $\text{GCob}_{c,d}$ -algebra A with the simplest non-trivial $\text{GCob}_{0,1}$ -algebra $\mathbb{Z}_{0,1} \cong \mathbb{1} \oplus \Sigma \mathbb{1}$ as defined in Example [5.8.](#page-22-1) By Corollary [4.12,](#page-17-3) this gives a $\text{GCob}_{c,d+1}$ -algebra structure on $\mathbb{Z}_{0,1} \otimes A$ which has underlying module $A \oplus \Sigma A$.

In contrast to suspension, this functor is not an equivalence. This can be seen as the simplest nontrivial GCob_{0,1}-algebra $\mathbb{1} \oplus \Sigma \mathbb{1}$ is not ⊗-invertible in contrast to the simplest non-trivial GCob_{1,-1}algebra Σ1.

The suspension of Frobenius algebras can also be studied on the level of maps and relations. Proposition [4.16](#page-18-0) shows that a shifted $GCob_{c,d}$ -algebra is a $GCob_{c-1,d+1}$ -algebra. This can also be stated in terms of maps and relations. However, it is a priori not clear that the suspended maps $\Sigma\mu$, $\Sigma\eta$, $\Sigma\nu$ and $\Sigma\eta$ satisfy the signs as in our main theorem because those signs depend on choices of orientations. It turns out that our choices are stable under suspending algebras in this sense.

To make this precise, we study how a map from $A^{\otimes k} \to A^{\otimes l}$ induces a map $(\Sigma A)^{\otimes k} \to (\Sigma A)^{\otimes l}$. For this, we note that there is a canonical generator $\sigma_{k,l}$ in $\mathbb{ZMod}_{\bullet}((\Sigma \mathbb{1})^{\otimes k}, (\Sigma \mathbb{1})^{\otimes l})$ which sends the canonical generator $s(1) \otimes \cdots \otimes s(1) \in (\Sigma \mathbb{1})^{\otimes k}$ to the canonical generator $s(1) \otimes \cdots \otimes s(1) \in (\Sigma \mathbb{1})^{\otimes l}$. We thus get a map

$$
\Sigma \colon \mathbb{Z} \text{Mod}_{\bullet}(A^{\otimes k}, A^{\otimes l}) \to \mathbb{Z} \text{Mod}_{\bullet}((\Sigma \mathbb{1})^{\otimes k}, (\Sigma \mathbb{1})^{\otimes l}) \otimes \mathbb{Z} \text{Mod}_{\bullet}(A^{\otimes k}, A^{\otimes l}) \cong \mathbb{Z} \text{Mod}_{\bullet}((\Sigma A)^{\otimes k}, (\Sigma A)^{\otimes l}),
$$

 $f \mapsto \sigma_{k,l} \otimes f.$

We note that this is coherent with our choice that maps (here tensoring with $\sigma_{k,l}$) act from the left. However, this map is functorial and monoidal only up to sign:

Lemma 5.10. Let $f: A^{\otimes k} \to A^{\otimes l}$ and $g: A^{\otimes m} \to A^{\otimes n}$ be maps. Then we have

- (a) $\Sigma f \otimes \Sigma g = (-1)^{(n-m)(|f|+k)} \Sigma (f \otimes g);$
- (b) $\Sigma f \circ \Sigma g = (-1)^{(n-m)|f|} \Sigma (f \circ g)$ if $n = k$;
- (c) $\sum id_{A^{\otimes k}} = id_{(\sum A)^{\otimes k}};$
- (d) $\Sigma \tau_A = -\tau_{\Sigma A}$ for $\tau_A : A \otimes A \to A \otimes A$ the braiding morphism.

Proof. We first note that the maps $\sigma_{k,l}$ are in degree $l - k$ and satisfy

$$
\sigma_{k,l}\otimes \sigma_{m,n}(s(1)^{\otimes k}\otimes s(1)^{\otimes m})=(-1)^{(n-m)k}s(1)^{\otimes l}\otimes s(1)^{\otimes n}=(-1)^{(n-m)k}\sigma_{k+m,l+n}(s(1)^{\otimes k+m})
$$

and

$$
\sigma_{k,l} \circ \sigma_{m,k} = \sigma_{m,l}.
$$

We can thus compute

$$
\Sigma f \otimes \Sigma g = (\sigma_{k,l} \otimes f) \otimes (\sigma_{m,n} \otimes g) = (-1)^{(n-m)|f|} (\sigma_{k,l} \otimes \sigma_{m,n}) \otimes (f \otimes g)
$$

= $(-1)^{(n-m)(|f|+k)} (\sigma_{k+m,l+n}) \otimes (f \otimes g) = (-1)^{(n-m)(|f|+k)} \Sigma (f \otimes g)$

and if $n = k$

$$
\Sigma f \circ \Sigma g = (\sigma_{k,l} \otimes f) \circ (\sigma_{m,n} \otimes g) = (-1)^{|\sigma_{m,n}||f|} (\sigma_{k,l} \circ \sigma_{m,n}) \otimes (f \circ g)
$$

= $(-1)^{(n-m)|f|} \sigma_{m,l} \otimes (f \circ g) = (-1)^{(n-m)|f|} \Sigma (f \circ g).$

The equation [\(c\)](#page-24-2) follows directly from [\(b\)](#page-24-3). For [\(d\)](#page-24-4), we denote $s(a) \in (\Sigma A)_{i+1}$ for the element coming from $a \in A_i$ and similarly $s(b) \in A_{j+1}$. The map $\Sigma \tau_A$ is defined as the composition

$$
s(a) \otimes s(b) \mapsto (-1)^{|a|} s(1) \otimes s(1) \otimes a \otimes b
$$

\n
$$
\xrightarrow{\sigma_{2,2} \otimes \tau_A} (-1)^{|a||b|+|a|} (s(1) \otimes s(1)) \otimes (b \otimes a)
$$

\n
$$
\mapsto (-1)^{|a||b|+|a|+|b|} s(b) \otimes s(a)
$$

\n
$$
= (-1)^{|s(a)||s(b)|-1} s(b) \otimes s(a) = -\tau_{\Sigma A}(s(a) \otimes s(b)).
$$

Proposition [5.1](#page-20-0)1. Let $(A, \mu, \eta, \nu, \varepsilon)$ in $\mathbb{Z} \mathbf{Mod}_{\bullet}$ be as in Theorem 5.1 and c, $d \in \mathbb{Z}$.

This data satisfies the relations of a (c, d) -graded commutative (or symmetric) Frobenius algebra if and only if $(\Sigma A, \Sigma \mu, \Sigma \eta, \Sigma \nu, \Sigma \varepsilon)$ satisfies the relations of a $(c-1, d+1)$ -graded commutative (or symmetric) Frobenius algebra.

Proof. This is a direct application of Lemma [5.10.](#page-23-0) For example, if $(A, \mu, \eta, \nu, \varepsilon)$ satisfies (c, d) -graded left unitality, then we check compute $(c-1, d+1)$ -graded left unitality for $(\Sigma A, \Sigma \mu, \Sigma \eta, \Sigma \nu, \Sigma \varepsilon)$:

$$
(-1)^{c-1} \Sigma \mu \circ (\Sigma \eta \otimes \mathrm{id}_{\Sigma A}) \stackrel{(c)}{=} (-1)^{c-1} \Sigma \mu \circ (\Sigma \eta \otimes \Sigma \mathrm{id}_A) \stackrel{(a)}{=} (-1)^{c-1} \Sigma \mu \circ \Sigma (\eta \otimes \mathrm{id}_A)
$$

$$
\stackrel{(b)}{=} (-1)^{c-1+c} \Sigma (\mu \circ (\eta \otimes \mathrm{id}_A)) = (-1)^{c-1+\frac{c(c-1)}{2}} \Sigma \mathrm{id}_A
$$

$$
\stackrel{(c)}{=} (-1)^{\frac{(c-1)(c-1-1)}{2}} \mathrm{id}_{\Sigma A}.
$$

6. Examples

In this section, we discuss different examples of graded Frobenius algebras found in the literature. Some of the examples have a description more akin to our Definition [4.4](#page-15-0) in terms of PROPs, while some of the examples have a description in terms of maps and relations similar to Theorem [5.1.](#page-20-0)

Throughout this section, we use Sweedler notation of a coproduct ν on an element a:

$$
\nu(a) = \sum_{(a)} a' \otimes a''.
$$

6.1. Cohomology of an Oriented Manifold. Let M be a connected, closed manifold of dimension d and let K be a field. Denote by $H^*(M)$ the cohomology of M with coefficients in K. Assume that M is K-orientable and fix an orientation $[M] \in H_d(M)$. We denote

$$
\mu\colon H^*(M)\otimes H^*(M)\to H^*(M),
$$

$$
\alpha\otimes\beta\mapsto\alpha\cup\beta
$$

and

$$
\varepsilon: H^*(M) \to K,
$$

\n
$$
\alpha \mapsto \begin{cases} \alpha([M]) & \text{if } * = d; \\ 0 & \text{else.} \end{cases}
$$

The multiplication μ has degree 0 and ε has degree $-d$. Moreover, μ is associative and has unit map

$$
\eta: K \to H^*(M),
$$

$$
1 \mapsto 1 \in H^0(M).
$$

Moreover, it is graded commutative as it satisfies $\mu \circ \tau = \mu$. Poincaré duality implies that the pairing $\varepsilon \circ \mu$ is non-degenerate. We can thus apply Corollary [5.5](#page-21-1) and get the following example

Example 6.1. The data $(H^*(M), \mu, \eta, \varepsilon)$ defines a $(0, d)$ -graded commutative Frobenius algebra

The associated comultiplication ν is the intersection coproduct. There are two common constructions of the intersection coproduct leading to different signs: we refer to them as the Poincaré coproduct ν_P and the Thom coproduct ν_{Th} (see e.g. [\[HW17,](#page-47-18) Appendix B]).

We first define the Poincaré coproduct. As we work with field coefficients, the Künneth maps

$$
\times: H_*(M) \otimes H_*(M) \to H_*(M^2),
$$

$$
\times: H^*(M) \otimes H^*(M) \to H^*(M^2)
$$

are isomorphisms. Denote by θ their inverses. We thus have a coproduct on *homology* given by

$$
\nu_{Hom} := \theta \circ \Delta_* \colon H_*(M) \to H_*(M^2) \to H_*(M) \otimes H_*(M).
$$

We denote the Poincaré duality by

$$
PD: H^*(M) \stackrel{\cong}{\to} H_{d-*}(M),
$$

$$
\alpha \mapsto \alpha \cap [M].
$$

We then define the Poincaré coproduct as Poincaré dual to the homology coproduct ν_{Hom}

$$
\nu_P(\alpha) = \sum_{(PD(\alpha))} PD^{-1}(PD(\alpha)') \otimes PD^{-1}(PD(\alpha)'')
$$

where sum comes from ν_{Hom} . This formula in terms of elements seems rather straight forward. But if we describe it as a composition of maps, we need the tensor product without Koszul sign: for graded maps $f: V \to V'$ and $g: W \to W'$ denote $f \widetilde{\otimes} g$ as the as the map

$$
(f \otimes g)(a \otimes b) = f(a) \otimes g(b).
$$

Then we write the Poincaré coproduct as

$$
\nu_P := (PD^{-1}\widetilde{\otimes} PD^{-1}) \circ \theta \circ \Delta_* \circ PD.
$$

The Thom coproduct is defined via an orientation $[M^2]$ of the manifold M^2 . To get the signs as in our main theorem, we define $[M^2] := (-1)^{\frac{n(n-1)}{2}} [M] \times [M]$.

This sign convention might seem surprising. It reflects the fact that we are shuffling the *i*-th dimension of the first copy of M to be the $2i-1$ -th dimension of M^2 and the i-th dimension of the second copy of M to be the 2*i*-th dimension of M^2 . We call such a shuffle of two ordered sets of the same size a riffle shuffle.

In terms of coordinates, take a ordered basis (v_1, \ldots, v_n) of \mathbb{R}^n . We then order the basis

$$
\{(v_1,0),\ldots,(v_n,0),(0,v_1),\ldots,(0,v_n)\}
$$

of $\mathbb{R}^n \oplus \mathbb{R}^n$ as

$$
((v_1,0),(0,v_1),(v_2,0),\ldots,(0,v_n)).
$$

This choice has the advantage that for two oriented manifolds M, N of dimension d, d' , we have $[(M \times N)^2] = [M^2] \times [N^2]$. Corollary [4.12](#page-17-3) then gives that $H^*(M) \otimes H^*(N)$ inherits the structure of a $\mathrm{GCob}_{0,d+d'}$ -algebra. Our choice of signs for ν_{Th} then ensures that

$$
H^*(M)\otimes H^*(N)\to H^*(M\times N)
$$

is an map of coalgebras.

We consider the diagonal $\Delta: M \to M \times M$ and fix a tubular neighbourhood $U \subset M \times M$ of $\Delta(M)$. We denote $j: M \to U$ and $i: U \to M \times M$ for the canonical inclusions. We thus have $\Delta = i \circ j$. We define $[U] \in H_{2d}(U, \partial U)$ as the image of $[M^2]$ under the map

$$
H_{2d}(M^2) \to H_{2d}(M^2, \Delta(M)^c) \stackrel{(i_*)^{-1}}{\cong} H_{2d}(U, \partial U).
$$

Let $\tau \in H^d(U, \partial U)$ be the Thom class that satisfies

$$
\tau \cap [U] = j_*([M]).
$$

Now, we can define the Thom coproduct as the following composition

$$
\nu_{Th}\colon H^*(M)\cong H^*(U)\stackrel{\tau\cup}{\to}H^{*+d}(U,\partial U)\cong H^{*+d}(M^2,\Delta(M)^c)\to H^{*+d}(M^2)\cong (H^{\bullet}(M)^{\otimes 2})^{*+d}.
$$

We point the readers attention to the fact that this coherent with our convention that maps act from the left as we are cupping with τ from the left in the second map.

To compare the signs that arise in the two definitions, we consider this diagram

$$
H^{k}(M) \xleftarrow{j^{*}} H^{k}(U) \xleftarrow{(-1)^{dk} H^{k+d}(U, \partial U) \xleftarrow{i^{*}} H^{k+d}(M^{2}, \Delta(M)^{c})
$$
\n
$$
\bigcap [M] \bigcup_{\substack{\bigcup f \to \bigcup f \
$$

- The composition along the bottom left gives the Poincaré coproduct ν_P .
- The composition along the top right gives the Thom coproduct ν_{Th} .
- The rectangle on the left hand side and the hexagon on the top right commute due to the naturality of the cap product.
- The triangle in the centre commutes by the definition of the Thom class τ .
- The sign in the bottom right rectangle means that, on elements in $H^{i}(M) \otimes H^{d+k-i}(M)$, we have the sign $(-1)^{d+dk+di}$ because the cross product and the cap product commute:

$$
(\alpha \cap [M]) \times (\beta \cap [M]) = (-1)^{d|\beta|} (\alpha \times \beta) \cap ([M] \times [M]).
$$

Combining this, we get on the $(i, d + k - i)$ -coordinate in the output the sign

(9)
$$
\nu_P = (-1)^{\frac{d(d-1)}{2} + d + di} \nu_{Th}.
$$

One can check that $(A, \mu, \eta, \nu_{Th}, \varepsilon)$ satisfies the relations of a $(0, d)$ -graded commutative Frobenius algebra as in Theorem [5.1.](#page-20-0) Cieliebak and Oancea prove this with their slightly different sign conventions in [\[CO22b,](#page-47-5) Proposition 7.1].

The signs of the Poincaré coproduct can be explained in the following way: as discussed in Proposi-tion [4.16](#page-18-0) if A has the structure of a (c, d) -graded commutative Frobenius algebra then ΣA inherits the structure of a $(c-1, d+1)$ -graded commutative Frobenius algebra. In this example, this means that

$$
\Sigma^{-d} H^*(M) = H^{*+d}(M)
$$

inherits the structure of a $(d, 0)$ -graded Frobenius algebra. Then the coproduct one obtains is given on an element $s^{-d}(\alpha) \in H^{*+d}(M)$ by:

$$
\Sigma^{d} \nu_{Th}(s^{-d}(\alpha)) = (\sigma_{1,2}^{\otimes -d} \otimes \nu_{Th})(s(1)^{\otimes -d} \otimes \alpha)
$$

\n
$$
= (-1)^{d} \sigma_{1,2}^{\otimes -d}(s(1)^{\otimes -d}) \otimes \nu_{Th}(\alpha)
$$

\n
$$
= (-1)^{d} \sigma_{1,2}(s(1))^{\otimes -d} \otimes \sum_{(\alpha)} \alpha' \otimes \alpha''
$$

\n
$$
= (-1)^{d} (s(1)^{\otimes 2})^{\otimes -d} \otimes \sum_{(\alpha)} \alpha' \otimes \alpha''
$$

\n
$$
= \sum_{(\alpha)} (-1)^{d+|\alpha'|+|s(\alpha')|+\cdots+|s^{d-1}(\alpha')|} s^{-d}(\alpha') \otimes s^{-d}(\alpha'')
$$

\n
$$
= \sum_{(\alpha)} (-1)^{d+d|\alpha'|+ \frac{d(d-1)}{2}} s^{-d}(\alpha') \otimes s^{-d}(\alpha'').
$$

Here $\sigma_{1,2}^{\otimes -d}$ denotes the d-fold tensor product of the canonical map $\Sigma^{-1} \mathbb{1} \to \Sigma^{-1} \mathbb{1} \otimes \Sigma^{-1} \mathbb{1}$. These are precisely the signs, we observed in [\(9\)](#page-27-0). We thus conclude $\Sigma^{-d} \nu_{Th} = \nu_P$. Therefore, the Poincaré coproduct is the suspended Thom coproduct on $\Sigma^{-d}H^*(M) \cong H^{*+d}(M)$. In $H^{*+d}(M)$, the coproduct is a degree 0 map, which makes the signs appearing in the graded coassociativity and counitality considerably more comfortable. One pays for this comfort by getting less appealing signs for the associativity and unitality of the corresponding shifted multiplication.

6.2. Hochschild Homology of a Frobenius Algebra.

Definition 6.2. Let $\mathrm{GCob}_{c,d}^{\partial_{out}\neq\varnothing}$ be the subcategory of $\mathrm{GCob}_{c,d}$ which only has morphisms coming from graphs G satisfying the plumber's condition i.e. all connected components of G have at least one out-going external vertex.

Let $\overline{GCo}_{c,d}^{\partial_{in} \neq \varnothing}$ be the subcategory of $\overline{GCo}_{c,d}$ which only has morphisms coming from graphs G satisfying the reverse plumber's condition i.e. all connected components of G have at least one in-going external vertex.

In [\[WW16\]](#page-47-1), Wahl and Westerland show that if A is a $(0, d)$ -graded symmetric Frobenius algebra, then the normalized Hochschild homology $\overline{HH}_*(A)$ and A has the structure of an open-closed TQFT. Isolating the closed part, we get a TQFT twisted by $\det(S, \partial_{out}S)^{\otimes d}$. We can restate this in our language.

Example 6.3. Let A be a $(0, d)$ -graded symmetric Frobenius algebra. Then $\overline{HH}_{*}(A)$ has the structure of a $\mathrm{GCob}^{\partial_{in} \neq \varnothing}_{d,d}$ -algebra.

The key input for this is the following observation: the twisting $\det(\text{Cob}(G), \partial_{out}(\text{Cob}(G)))^{\otimes d}$ corresponds to our (d, d) -twisting: $\det(G, \partial_{in})^{\otimes d} \otimes \det(G, \partial_{out})^{\otimes d}$. (Recall that $Cob(G)$ is the corresponding closed surface cobordism represented by the graph G .) This is given by the following proposition:

Proposition 6.4. The three functors of 2-categories

$$
\mathbf{det}_{1,1}, \mathbf{det}(\mathrm{Cob}(-), \partial_{in}), \mathbf{det}(\mathrm{Cob}(-), \partial_{out}) \colon \underline{\mathrm{GCob}} \to B\mathrm{Pic}(\mathbb{Z})
$$

are isomorphic.

Explicitly, there exist natural isomorphisms

(10)
$$
\det(G, \partial_{in}) \otimes \det(G, \partial_{out}) \cong \det(\text{Cob}(G), \partial_{in}(\text{Cob}(G)))
$$

$$
\det(G, \partial_{in}) \otimes \det(G, \partial_{out}) \cong \det(\text{Cob}(G), \partial_{out}(\text{Cob}(G)))
$$

for all $G \in \text{GCob}(X, Y)$ which commute with gluing of graphs.

Proof. We only prove the first isomorphism. The second one is proved analogously. Both the left and right hand side of [\(10\)](#page-29-0) represent functors out of $\text{GCob}(X, Y)$:

$$
\det(-,\partial_{in}) \otimes \det(-,\partial_{out}), \qquad \qquad \det(\mathrm{Cob}(-),\partial_{in}(\mathrm{Cob}(-))).
$$

After precomposition with the forgetful functor $\underline{fGCob}(X, Y) \rightarrow \underline{GCob}(X, Y)$, they define functors out of $fGCob(X, Y)$. What we construct in the following is a natural isomorphism of the two functors out of $fGCob(X, Y)$.

This lifts to a natural transformation of functors out of $G\text{Cob}(X, Y)$. This can be seen as all edge collapses can be lifted to an edge collapse in a fat graph. It remains to check that graph isomorphisms can be lifted. For this we note that all graph isomorphisms can be decomposed into a graph isomorphism that can be lifted to a fat graph isomorphism and a graph automorphism. Finally all graph automorphisms act trivially on the left and right hand side of [\(10\)](#page-29-0). Therefore the following isomorphism can be lifted to a natural transformation of functors out of $GCob(X, Y)$.

Let $G \in \mathrm{fGCob}(X, Y)$ be a fat graph. We construct isomorphisms which commute with gluing of graphs:

$$
f_G \colon \mathbf{det}(\mathrm{fCob}(G), \partial_{out}) \stackrel{\cong}{\to} \mathbf{det}(\mathrm{fCob}(G), \partial_{in} \cup \partial_{free});
$$

$$
g_G \colon \mathbf{det}(\mathrm{fCob}(G), \partial_{in}) \otimes \mathbf{det}(\mathrm{fCob}(G), \partial_{in} \cup \partial_{free}) \stackrel{\cong}{\to} \mathbf{det}(\mathrm{Cob}(G), \partial_{in}).
$$

We recall that ∂_{free} is the part of the boundary which is neither in the in-coming boundary nor in the out-going boundary. Then the proposition follows by taking $g_G \circ (\mathrm{id} \otimes f_G)$.

For f_G , we first note that $H_*(|G|, |L_{out}|)$ is a free graded Z-module. Therefore any isomorphism

 $H_*(|G|, |L_{out}|) \cong (H_*(|G|, |L_{out}|))^{\vee} \cong H^*(|G|, |L_{out}|)$

gives an isomorphism

$$
\det(G, \partial_{out}) = \det(H_*(|G|, |L_{out}|)) \cong \det(H^*(|G|, |L_{out}|)).
$$

Next, we note that the homotopy equivalence of pairs

$$
(|G|, |L_{out}|) \simeq (fCob(G), \partial_{out})
$$

gives an isomorphism

$$
\det(H^*(|G|, |L_{out}|)) \cong \det(H^*(\mathrm{fCob}(G), \partial_{out})).
$$

All those isomorphisms so far give an isomorphism

$$
\det(G, \partial_{out}) \cong \det(H^*(\operatorname{fCob}(G), \partial_{out}))
$$

which commutes with gluing.

For the next step, we use generalized Lefschetz duality (see for example [\[Hat02,](#page-47-19) Theorem 3.43]). This gives an isomorphism

$$
H^*(\mathrm{fCob}(G), \partial_{out}) \cong H_{2-*}(\mathrm{fCob}(G), \partial_{in} \cup \partial_{free})
$$

induced by capping with $\lbrack \text{fCob}(G) \rbrack$, the orientation in $H_2(\text{fCob}(G), \partial \text{fCob}(G))$. As the fat graphs induce oriented open cobordisms, this isomorphism also commutes with gluing. We thus get an isomorphism

 $\det(H^*(\operatorname{fCob}(G),\partial_{out})) \cong \det(H_{2-*}(\operatorname{fCob}(G),\partial_{in} \cup \partial_{free}) \cong \det(H_*(\operatorname{fCob}(G),\partial_{in} \cup \partial_{free}))$

which commutes with gluing. This gives f_G .

For g_G , we consider $fCob(G)$ as a subspace of $Cob(G)$. Figuratively speaking, we consider it as the lower half. See Figure [7.](#page-30-0)

FIGURE 7. Embedding $fCob(G)$ in grey as the bottom half of $Cob(G)$.

Consider the triple of spaces $(\text{Cob}(G), \text{fCob}(G) \cup \partial_{in}(\text{Cob}(G)), \partial_{in}(\text{Cob}(G)))$. We note that by excision

$$
H_*(\operatorname{fCob}(G), \partial_{in}(\operatorname{fCob}(G))) \cong H_*(\operatorname{fCob}(G) \cup \partial_{in}(\operatorname{Cob}(G)), \partial_{in}(\operatorname{Cob}(G)))
$$

$$
H_*(\mathrm{fCob}(G), \partial_{in}(\mathrm{fCob}(G)) \cup \partial_{free}(\mathrm{fCob}(G))) \cong H_*(\mathrm{Cob}(G), \mathrm{fCob}(G) \cup \partial_{in}(\mathrm{Cob}(G))).
$$

We thus have

$$
\textbf{let}(\text{fCob}(G), \partial_{in}(\text{fCob}(G))) \otimes \textbf{det}(\text{fCob}(G), \partial_{in}(\text{fCob}(G))) \cup \partial_{free}(\text{fCob}(G)))
$$

$$
\stackrel{\cong}{\to} \det(\mathrm{fCob}(G) \cup \partial_{in}(\mathrm{Cob}(G)), \partial_{in}(\mathrm{Cob}(G))) \otimes \det(\mathrm{Cob}(G), \mathrm{fCob}(G) \cup \partial_{in}(\mathrm{Cob}(G))).
$$

We note that collapsing the in-coming boundary and the lower half in $Cob(G)$ is the same space as collapsing the free and in-coming boundary of the upper half. See Figure [7.](#page-30-0) The short exact sequence of the triple of spaces and Lemma [4.1](#page-13-5) [\(d\)](#page-13-3) then gives

$$
\det(\mathrm{fCob}(G)\cup \partial_{in}(\mathrm{Cob}(G)),\partial_{in}(\mathrm{Cob}(G)))\otimes \det(\mathrm{Cob}(G),\mathrm{fCob}(G)\cup \partial_{in}(\mathrm{Cob}(G)))
$$

$$
\stackrel{\cong}{\to} \det(\mathrm{Cob}(G), \partial_{in}(\mathrm{Cob}(G))).
$$

This gives g_G and thus concludes the proof. □

We note that the category $\text{GCob}_{d,d}^{\partial_{in} \neq \emptyset}$ is generated by $\omega(\rightarrow)$, $\omega(\rightarrow)$ and $\omega(\rightarrow)$. A symmetric monoidal functor $F: \text{GCob}_{d,d} \to K\text{Mod}_{\bullet}$ is uniquely determined by

$$
\mu_{HH} := F(\omega(\rightarrow)) \in \mathbb{Z} \text{Mod}_d(\overline{HH}_*(A) \otimes \overline{HH}_*(A), \overline{HH}_*(A)),
$$

\n
$$
\nu_{HH} := F(\omega(\rightarrow)) \in \mathbb{Z} \text{Mod}_d(\overline{HH}_*(A), \overline{HH}_*(A) \otimes \overline{HH}_*(A)),
$$

\n
$$
\varepsilon_{HH} := F(\omega(\rightarrow)) \in \mathbb{Z} \text{Mod}_{-d}(\overline{HH}_*(A), \mathbb{1}).
$$

Those maps satisfy graded associativity, coassociativity, counitality and commutativity and the graded Frobenius relation as in Theorem [5.1.](#page-20-0)

Following [\[WW16,](#page-47-1) Section 6.5] using our sign conventions that maps act from the left, we have the description

$$
\mu_{HH}((a_0 \otimes \cdots \otimes a_k) \otimes (b_0 \otimes \cdots \otimes b_l)) = \begin{cases} 0 & \text{if } k > 0, \\ \sum_{(a_0)} (-1)^{(|a'_0| + d)|a''_0} a''_0 a'_0 b_0 \otimes b_1 \otimes \cdots \otimes b_l & \text{if } k = 0, \end{cases}
$$

$$
\nu_{HH}(a_0 \otimes \cdots \otimes a_k) = \sum_{\substack{i=0,\ldots,k \\ (a_0)}} (-1)^{(|a'_0| + k - i)(|a''_0| + |a_1| + \cdots + |a_i|)}
$$

$$
(a''_0 \otimes a_1 \otimes \cdots \otimes a_i) \otimes (a'_0 \otimes a_{i+1} \otimes \cdots \otimes a_k),
$$

$$
\varepsilon_{HH}(a_0 \otimes \cdots \otimes a_k) = \begin{cases} 0 & \text{if } k > 0, \\ \varepsilon(a_0) & \text{if } k = 0. \end{cases}
$$

We note that these are different signs compared to [\[WW16\]](#page-47-1). One reason is that they have a different sign convention that maps act from the right. On the other hand, the signs in their coproducts are missing the Koszul sign coming from moving a'_0 past $a''_0 \otimes a_1 \otimes \cdots \otimes a_i$.

6.3. **Homology of the Loop Space.** Let M be a connected, closed, oriented manifold of dimension d and let K be a field. Let $LM = \text{Map}(S^1, M)$ be its loop space.

In [\[CG04\]](#page-47-20), Cohen and Godin define operations μ_S on $H_*(LM) := H_*(LM;K)$ for every closed cobordism S such that each connected component has at least one out-going boundary. In [\[God07\]](#page-47-0), Godin moreover explains that this fits into a positive boundary open-closed TQFT which is twisted by $\det(S, \partial_{in}S)^{\otimes -d}$. This minus sign comes from our convention that the determinant is in negative degree while Godin has the convention that the determinant is in positive degree.

We only focus on the closed underived part of this TQFT. Using the first isomorphism in Proposition [6.4,](#page-28-0) we can rephrase this in our language as the following example:

Example 6.5. The homology of the loop space $H_*(LM)$ has the structure of a $\text{GCob}^{\partial_{out} \neq \emptyset}_{-d,-d}$ -algebra with operations given by the Chas-Sullivan product and the trivial coproduct.

We thus get a multiplication on $H_*(LM)$, that we denote by μ_{Go} , and a comultiplication on $H_*(LM)$, that we denote by ν_{Go} .

We can compare this to the results of Tamanoi in [\[Tam10\]](#page-47-21). Tamanoi defines a coproduct

$$
\Psi: H_{*+d}(LM) \to H_*(LM \times LM).
$$

As we work over a field, we can assume that the Künneth map $\times : H_*(LM) \otimes H_*(LM) \to H_*(LM)$ LM) is invertible. We therefore can denote $\nu_{Ta} := \times^{-1} \circ \Psi$ a comultiplication in our sense. Tamanoi moreover defines multiplications

$$
\mu_{Ta}: H_*(LM) \otimes H_*(LM) \to H_*(LM),
$$

$$
\cdot: H_*(LM) \otimes H_*(LM \times LM) \to H_*(LM \times LM),
$$

$$
\cdot: H_*(LM \times LM) \otimes H_*(LM) \to H_*(LM \times LM)
$$

which are defined such that for $a, b, c \in H_*(LM)$, we have

$$
a \cdot (b \times c) = \mu_{Ta}(a \otimes b) \times c
$$

$$
(a \times b) \cdot c = a \times \mu_{Ta}(b \otimes c).
$$

In [\[Tam10,](#page-47-21) Theorem 2.2], Tamanoi observes the following Frobenius relation

$$
\Psi(\mu_{Ta}(a\otimes b)) = (-1)^{d(|a|-d)}a \cdot \Psi(b) = \Psi(a) \cdot b \in H_*(LM \times LM).
$$

We interpret everything as living in $H_{*+d}(LM)$ and $H_{*+d}(LM \times LM)$. Then the Künneth map is a degree d map and its inverse a degree −d map. We can thus rewrite the Frobenius relation as

$$
\nu_{Ta} \circ \mu_{Ta} = (\mu_{Ta} \otimes id) \circ (id \otimes \nu_{Ta}) = (id \otimes \mu_{Ta}) \circ (\nu_{Ta} \otimes id).
$$

This is exactly the relation, we expect for a the $\text{GCob}^{\partial_{out} \neq \emptyset}_{0,-2d}$ -algebra structure on $H_{*+d}(LM)$ $\Sigma^{-d}H_*(LM)$. This structure is predicted by shifting the $\text{GCob}_{-d,-d}^{\partial_{out}\neq\varnothing}$ -structure in Example [6.5](#page-31-0) and applying Proposition [4.16.](#page-18-0)

It is worth noting that the coproducts, ν_{Go} and its shifted version ν_{Ta} , are the so called trivial coproduct on $H_*(LM)$. This is in contrast to the Goresky-Hingston coproduct constructed in [\[GH09,](#page-47-22) [HW17\]](#page-47-18).

Explicitly, Hingston and Wahl give a definition for the Chas-Sullivan product and the Goresky-Hingston product in [\[HW17\]](#page-47-18).

Their definition of the product is graded associative, unital and commutative. This lifts from the respective properties of the intersection product ν_{Th} discussed in [6.1](#page-25-1) (see the proof of [\[HW17,](#page-47-18) Theorem 2.5]). This is in line with Example [6.5.](#page-31-0)

For the coproduct, we slightly amend their definition. Hingston and Wahl define in [\[HW17,](#page-47-18) Definition 1.4] the loop coproduct as the degree $1 - n$ map given for a chain $A \in C_*(LM, M)$ by

$$
\mathrm{AW}(\mathrm{cut}(R_\mathrm{GH}([\tau_{\mathrm{GH}}\cap](A\times I))))
$$

For a detailed explanation of this notation, see [\[HW17\]](#page-47-18). We note that the first map in this composition $\times I$ acts from the right while the second map $[\tau_{GH} \cap]$ acts from the left.

As we chose the convention that maps act from the left, we define the map as the composition

$$
\vee (A) := \text{AW}(\text{cut}(R_{\text{GH}}([\tau_{\text{GH}} \cap](I \times A))))
$$

An analogous proof strategy as for [\[HW17,](#page-47-18) Theorem 2.14] shows that \vee is a degree $1 - n$ coproduct that satisfies the graded coassociativity and cocommutativity relation as in Theorem [5.1.](#page-20-0)

However, the Chas-Sullivan product and the Goresky-Hingston coproduct do not satisfy a Frobenius relation in this setting. In particular, the product is an operation on $H_*(LM)$ while the coproduct is an operation on $H_*(LM, M)$ the homology relative the constant loops.

6.4. Rabinowitz Loop Homology. In the series of articles [\[CHO20,](#page-47-6) [CO22b,](#page-47-5) [CO22a,](#page-47-7) [CHO23,](#page-47-8) [CO24\]](#page-47-9), Cieliebak, Hingston and Oancea give an interpretation where the Chas-Sullivan product and the Goresky-Hingston coproduct can be interpreted in a Frobenius type structure, what Cieliebak and Oancea call a biunital coFrobenius algebra. As discussed in Remark [5.4,](#page-21-0) the signs in Theorem [5.1](#page-20-0) correspond to the signs appearing in the definition of biunital coFrobenius algebras. This gives a correspondence of their notion of a biunital coFrobenius algebras and our graded Frobenius algebras in the finite dimensional case. But for the key example of Rabinowitz loop homology, Cieliebak and Oancea need infinite dimensional algebras. For this they use the language of graded Tate vector spaces to get a Frobenius relation.

They give two tensor products on graded Tate vector spaces: $\hat{\otimes}^*, \hat{\otimes}^!$. All maps that appear in their relations have source $A^{\widehat{\otimes}^*k}$ and target $A^{\widehat{\otimes}^!l}$ for some $k, l \geq 0$. The framework of PROPs set up so far fails to describe those algebras. Indeed for a graded Tate vector space A, the collection

$$
\text{End}_A(X, Y) := \text{Hom}(A^{\widehat{\otimes}^* X}, A^{\widehat{\otimes}^! Y})
$$

for X, Y finite sets does not have a canonical structure of a PROP. This is because in general there is no canonical map $A^{\widehat{\otimes}^{\dagger}X} \to A^{\widehat{\otimes}^*X}$ and thus no composition.

Our remedy for this problem is to work with dioperads and the following proposition:

Proposition 6.6. Let A be a graded Tate vector. The collection $\text{End}_A(X, Y)$ has a canonical structure of a dioperad in $\mathbb{Z}Mod_0$.

Proof. We recall from subsection [4.3](#page-19-3) that a dioperad structure consists of a (S_X, S_Y) -bimodule structure, a unit and composition maps.

The objects $\text{End}_A(X, Y)$ have a canonical (S_X, S_Y) -bimodule structure given by permuting the coordinates.

The unit is given by $id_A \in \text{End}_A(\{*\}, \{*\}) = \text{Hom}(A, A).$

The composition map uses the map

$$
\alpha\colon A\widehat{\otimes}^*(B\widehat{\otimes}^!C)\to (A\widehat{\otimes}^*B)\widehat{\otimes}^!C
$$

for graded Tate vector spaces A, B, C as described in [\[CO24,](#page-47-9) Proposition 5.5]. We fix some sets W, X, Y, Z and elements $x \in X, y \in Y$ and an element $f \in \text{End}_A(W, X)$ and $g \in \text{End}_A(Y, Z)$. We define $g_y \circ_x f \in \text{End}_A((Y \setminus \{y\}) \sqcup W, Z \sqcup (X \setminus \{x\}))$ as the composition

$$
A^{\widehat{\otimes}^*(Y\backslash\{y\})\sqcup W} \cong (A^{\widehat{\otimes}^*Y\backslash\{y\}})\widehat{\otimes}^*(A^{\widehat{\otimes}^*W}) \xrightarrow{f} (A^{\widehat{\otimes}^*Y\backslash\{y\}})\widehat{\otimes}^*(A^{\widehat{\otimes}^*X})
$$

$$
\xrightarrow{\alpha} (A^{\widehat{\otimes}^*Y})\widehat{\otimes}^!(A^{\widehat{\otimes}^*X\backslash\{x\}}) \xrightarrow{g} (A^{\widehat{\otimes}^*Z})\widehat{\otimes}^!(A^{\widehat{\otimes}^*X\backslash\{x\}}) \cong A^{\widehat{\otimes}^*Z\sqcup(X\backslash\{x\})}.
$$

Checking that this data satisfies the unitality, associativity and equivariance relations is analogous to checking that the endomorphism dioperad of an object in any symmetric monoidal category is a dioperad. □

This allows us to define a Tate vector space having the structure of an algebra over a dioperad:

Definition 6.7. Let D a dioperad in \mathbb{ZMod}_0 and let A be a Tate vector space. A D -algebra structure on A is a morphism of dioperads

 $\mathcal{D} \to \text{End}_A$

enriched over \mathbb{ZMod}_0 i.e. a morphism of dioperads which is linear and preserves degree.

We recall that Proposition [4.20](#page-19-2) shows that if c, d are of different parity working with the forest dioperad does not lose operations compared to the PROP in characteristic $\neq 2$. The main example of Rabinowitz loop homology is such a case. We can thus state the closed part of [\[CO22b,](#page-47-5) Corollary 9.3 (c)] in the following example.

Example 6.8. Let M be an oriented d -dimensional manifold. The Rabinowitz loop homology $\widehat{H}_*(LM)$ has the structure of a $\text{GCob}^{forest}_{d,d+1}$ -algebra.

7. Proof of Main Theorem

In this section, we introduce some computational tools which lets us more easily compute orientations in $\det_{c,d}(G)$. With this machinery we can prove the main theorem in subsection [7.2.](#page-37-0)

7.1. Computational Tools. We recall that a generator $\omega(G) \in \det_{c,d}(G)$ is called an orientation. For the proof, we need to be able to do three things:

- (I) compute how (fat) graph isomorphisms act on orientations;
- (II) compute how edge collapses act on orientations;
- (III) compute the composition of two orientations.

We recall that the composition in (III) was defined in Lemma [4.2](#page-14-0) using a splitting of the short exact sequence:

$$
0 \longrightarrow C_{\bullet}(|G|, |\partial_{in}|) \longrightarrow C_{\bullet}(|G' \circ G|, |\partial_{in}|) \longrightarrow C_{\bullet}(|G'|, |\partial_{in}|) \longrightarrow 0.
$$

To explicitly compute [\(I\)](#page-33-5), [\(II\)](#page-33-6) and [\(III\)](#page-33-4), we use Proposition [7.1](#page-34-0) inspired by [\[GK98,](#page-47-15) Proposition 4.14] and [\[God07,](#page-47-0) Lemma 14]. This gives us an expression of orientations in terms of top exterior powers of edges, half-edges and vertices:

For X a finite set, we use the following notation

- $Or(X) := Or(\mathbb{Z}^X)$ the top exterior concentrated in degree 0;
- det(X) := det(\mathbb{Z}^X) the top exterior concentrated in degree $-|X|$.

Proposition 7.1. Let G be a graph. There is a natural isomorphism

$$
\det(G, \partial_{in}) \cong \det(E)^{-1} \otimes \text{Or}(H) \otimes \det(V \setminus L_{in}).
$$

Moreover, an analogous statement for $\det(G, \partial_{out})$ holds after replacing L_{in} by L_{out} .

Proof. We set $L := L_{in}$ or $L := L_{out}$ to obtain the two statements. Consider the cellular chain complex $C_{\bullet}(|G|, |L|)$ for $|G|$ relative $|L|$:

$$
0 \longrightarrow \bigoplus_{e \in E} \text{Or}(\{h_e, \sigma h_e\}) \longrightarrow \mathbb{Z}^V/\mathbb{Z}^L \longrightarrow 0,
$$

$$
h_e \wedge \sigma h_e \longmapsto s(\sigma h) - s(h).
$$

We have that the homology of this complex is $H_*(G, L)$. By Lemma [4.1,](#page-13-5) we have a sequence of natural isomorphisms

$$
\det(G, L) \stackrel{\text{(e)}}{\cong} \det(C_{\bullet}(G, L))
$$
\n
$$
\stackrel{\text{(b)}}{\cong} \det\left(\Sigma \bigoplus_{e \in E} \text{Or}(\{h_e, \sigma h_e\})\right) \otimes \det(\mathbb{Z}^V / \mathbb{Z}^L)
$$
\n
$$
\cong \det\left(\Sigma \bigoplus_{e \in E} \text{Or}(\{h_e, \sigma h_e\})\right) \otimes \det(V \setminus L)
$$
\n
$$
\stackrel{\text{(c)}}{\cong} \det\left(\bigoplus_{e \in E} \text{Or}(\{h_e, \sigma h_e\})\right)^{-1} \otimes \det(V \setminus L)
$$
\n
$$
\stackrel{\text{(b)}}{\cong} \bigotimes_{e \in E} \det(\text{Or}(\{h_e, \sigma h_e\}))^{-1} \otimes \det(V \setminus L)
$$

We note that $\det(\text{Or}(\{h_e, \sigma h_e\}))^{-1}$ is a free group of rank 1 concentrated in degree 1. Choosing a generator of this group is equivalent to choosing a generator in the degree 0 group $\text{Or}(\{h_e, \sigma h_e\})$. We thus find

$$
\det(\text{Or}(\{h_e, \sigma h_e\})^{-1} \cong \det(\{e\})^{-1} \otimes \text{Or}(\{h_e, \sigma h_e\})
$$

where tensoring with $\det({e})^{-1}$ ensures that the right hand side is in degree 1. We thus find

$$
\det(G, L) \cong \bigotimes_{e \in E} (\det(\{e\})^{-1} \otimes \text{Or}(\{h_e, \sigma h_e\})) \otimes \det(V \setminus L)
$$

$$
\stackrel{\text{(b)}}{\cong} \det(E)^{-1} \otimes \text{Or}(H) \otimes \det(V \setminus L)
$$

where the last isomorphism comes from the fact that $H = \bigcup_{e \in E} \{h_e, \sigma h_e\}.$

Remark 7.2. Explicitly, for $E(G) = \{e_1, \ldots, e_\alpha\}$, $e_i = \{h_i, \sigma h_i\}$ and $V(G) \setminus L_{in}(G) = \{v_1, \ldots, v_\beta\}$ a generator of $\det(G, \partial_{in})$ can be written as

(11)
$$
(e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes h_1 \wedge \sigma h_1 \wedge \cdots \wedge \sigma h_\alpha \otimes v_1 \wedge \cdots \wedge v_\beta.
$$

A graph isomorphism induces a map of edges, half-edges and vertices. This lets us compute isomorphism on orientations which gives [\(I\)](#page-33-5) on our wish list. For computing edge collapses as in [\(II\)](#page-33-6), we use Lemma [7.3](#page-35-0) which lets us compute an edge collapse on an orientation expressed in terms of edges, half-edges and vertices. For computing compositions as in [\(III\)](#page-33-4), we use Lemma [7.4.](#page-36-0) The proofs of both lemmas are quite technical. But once they are proven doing [\(I\)](#page-33-5), [\(II\)](#page-33-6) and [\(III\)](#page-33-4) are straight-forward computations.

Lemma 7.3. Let G be a graph and $E = \{e_0, e_1, \ldots, e_\alpha\}$, $e_i = \{h_i, \sigma h_i\}$ and $V = \{v_0, v_1, \ldots, v_\beta\}$. Assume that e_0 is not a tadpole and $s(\sigma h_0) = v_0 \notin L_{in}(G)$. The canonical map $c: |G| \to |G|/e_0$ does the following to generators as in [\(11\)](#page-35-1):

$$
\det(c): \det(E(G))^{-1} \otimes \text{Or}(H(G)) \otimes \det(V(G) \setminus L_{in}(G))
$$

\n
$$
\to \det(E(G/e_0))^{-1} \otimes \text{Or}(H(G/e_0)) \otimes \det(V(G/e_0) \setminus L_{in}(G/e_0))
$$

\n
$$
(e_0 \wedge e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \cdots \wedge \sigma h_\alpha \otimes v_0 \wedge v_1 \wedge \cdots \wedge v_\beta
$$

\n
$$
\mapsto (e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes h_1 \wedge \cdots \wedge \sigma h_\alpha \otimes v_1 \wedge \cdots \wedge v_\beta.
$$

Here we identify the newly created vertex in G/e_0 with $s(h_0)$ which gives an identification

$$
V(G) \setminus \{v_0\} \cong V(G/e_0)
$$

together with the canonical identification

$$
E(G) \setminus \{e_0\} \cong E(G/e_0), \qquad H(G) \setminus \{h_0, \sigma h_0\} \cong H(G/e_0).
$$

Moreover, an analogous statement for $\det(G, \partial_{out})$ holds after replacing L_{in} by L_{out} .

Proof. We set $L := L_{in}$ or $L := L_{out}$ and $\partial := \partial_{in}$ or $\partial := \partial_{out}$. The map c induces a map of CW-complexes. This in turn induces a map of cellular chain complexes

$$
0 \longrightarrow \bigoplus_{e \in E(G)} \text{Or}(e) \longrightarrow \mathbb{Z}^{V(G)}/\mathbb{Z}^L \longrightarrow 0
$$

$$
\downarrow_{c|E} \qquad \qquad \downarrow_{c|V}
$$

$$
0 \longrightarrow \bigoplus_{e \in E(G/e_0)} \text{Or}(e) \longrightarrow \mathbb{Z}^{V(G/e_0)}/\mathbb{Z}^L \longrightarrow 0
$$

where $c|_E$ sends $Or(e_0)$ to zero and all other edges to themselves and $c|_V$ sends v_0 to $v := s(h_0)$ and all other vertices to themselves. This is a surjective map of chain complexes with kernel:

$$
0 \longrightarrow \text{Or}(\{h_0, \sigma h_0\}) \longrightarrow \langle v_0 - v \rangle \longrightarrow 0
$$

$$
h_0 \wedge \sigma h_0 \longmapsto v_0 - v
$$

we denote this complex by C_{\bullet} . Because e_0 is not a tadpole $v_0 - v \neq 0$ and thus C_{\bullet} has trivial homology. Lemma [4.1](#page-13-5) [\(e\)](#page-13-4) gives the isomorphism

$$
\mathbb{1} \cong \det(C_{\bullet}) \cong \det(\{e_0\})^{-1} \otimes \text{Or}(e_0) \otimes \det(\langle v_0 - v \rangle)
$$

given by

(12)
$$
1 \mapsto e_0^{-1} \otimes h_0 \wedge \sigma h_0 \otimes (v_0 - v).
$$

The geometric map $\det(c)$: $\det(G, \partial) \to \det(G/e_0, \partial)$ is given by the composition

$$
\det(H_{\bullet}(G,\partial))\cong \det(C_{\bullet})\otimes \det(H_{\bullet}(G/e_{0},\partial))\cong \mathbb{1}\otimes \det(H_{\bullet}(G/e_{0},\partial))\cong \det(H_{\bullet}(G/e_{0},\partial)).
$$

The isomorphism $\det(H_{\bullet}(G,\partial)) \cong \det(C_{\bullet}) \otimes \det(H_{\bullet}(G/e_0,\partial))$ induces

$$
det(E(G))^{-1} \otimes \text{Or}(H(G)) \otimes det(V(G) \setminus L)
$$

\n
$$
\rightarrow det(E(G/e_0))^{-1} \otimes det(\lbrace e_0 \rbrace)^{-1} \otimes \text{Or}(e_0) \otimes \text{Or}(H(G/e_0)) \otimes det(\lbrace v_0 - v \rbrace) \otimes det(V(G/e_0) \setminus L)
$$

\n
$$
\rightarrow det(\lbrace e_0 \rbrace)^{-1} \otimes \text{Or}(e_0) \otimes det(\lbrace v_0 - v \rbrace) \otimes det(E(G/e_0))^{-1} \otimes \text{Or}(H(G/e_0)) \otimes det(V(G/e_0) \setminus L).
$$

This isomorphism is given by the identifications

$$
E(G) = \{e_0\} \sqcup E(G/e_0), \ H(G) = e_0 \sqcup H(G/e_0), \ \mathbb{Z}^{V(G) \setminus L} \cong \mathbb{Z}^{v_0 - v} \oplus \mathbb{Z}^{V(G/e_0) \setminus L}
$$

and then permuting the factors. We recall that in our conventions for graded abelian groups, the canonical isomorphism between the dual of the tensor product and the tensor product of the dual is order reversing. Thus the map on edges is order reversing:

$$
\det(E(G))^{-1} \to \det(E(G/e_0)) \otimes \det(\{e_0\})^{-1},
$$

$$
(e_0 \wedge e_1 \wedge \cdots \wedge e_\alpha)^{-1} \mapsto (e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes e_0^{-1}.
$$

Concretely on generators, we thus get the map

$$
(e_0 \wedge e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \cdots \wedge \sigma h_\alpha \otimes v_0 \wedge v_1 \wedge \cdots \wedge v_\beta
$$

= $(e_0 \wedge e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \cdots \wedge \sigma h_\alpha \otimes (v_0 - v) \wedge v_1 \wedge \cdots \wedge v_\beta$
 $\rightarrow (e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes e_0^{-1} \otimes h_0 \wedge \sigma h_0 \otimes h_1 \wedge \cdots \wedge \sigma h_\alpha \otimes (v_0 - v) \otimes v_1 \wedge \cdots \wedge v_\beta$
 $\rightarrow e_0^{-1} \otimes h_0 \wedge \sigma h_0 \otimes (v_0 - v) \otimes (e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes h_1 \wedge \cdots \wedge \sigma h_\alpha \otimes v_1 \wedge \cdots \wedge v_\beta.$

The isomorphism $\det(C_{\bullet}) \otimes \det(H_{\bullet}(G/e_0, \partial)) \cong \det(H_{\bullet}(G/e_0, \partial))$ is now given by applying [\(12\)](#page-36-1). Concretely on generators, we compute

$$
e_0^{-1} \otimes h_0 \wedge \sigma h_0 \otimes (v_0 - v) \otimes (e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes h_1 \wedge \cdots \wedge \sigma h_\alpha \otimes v_1 \wedge \cdots \wedge v_\beta
$$

$$
\mapsto (e_1 \wedge \cdots \wedge e_\alpha)^{-1} \otimes h_1 \wedge \cdots \wedge \sigma h_\alpha \otimes v_1 \wedge \cdots \wedge v_\beta.
$$

Composing these two maps thus gives the equation we wanted to compute. $\hfill \Box$

Lemma 7.4. Let G, G' be composable graphs in \underline{GCob} . The composition map

$$
\mathrm{comp} \colon \mathbf{det}(G',\partial_{in}) \otimes \mathbf{det}(G,\partial_{in}) \cong \mathbf{det}(G' \circ G,\partial_{in})
$$

given in Lemma [4.2](#page-14-0) is computed via

$$
\det(E(G'))^{-1} \otimes \operatorname{Or}(H(G')) \otimes \det(V(G') \setminus L_{in}(G'))
$$

$$
\otimes \det(E(G))^{-1} \otimes \operatorname{Or}(H(G)) \otimes \det(V(G) \setminus L_{in}(G))
$$

$$
\cong \det(E(G' \circ G))^{-1} \otimes \operatorname{Or}(H(G' \circ G)) \otimes \det(V(G' \circ G) \setminus L_{in}(G' \circ G))
$$

where the map is given by the canonical identifications

(13)
$$
E(G') \sqcup E(G) \cong E(G' \circ G),
$$

$$
H(G') \sqcup H(G) \cong H(G' \circ G),
$$

$$
(V(G')\setminus L_{in}(G'))\sqcup (V(G)\setminus L_{in}(G))\cong V(G'\circ G)\setminus L_{in}(G'\circ G).
$$

Moreover, an analogous statement holds for $\det(G, \partial_{out})$ after replacing L_{in} by L_{out} .

Proof. We set $L := L_{in}$ or $L := L_{out}$ and $\partial := \partial_{in}$ or $\partial := \partial_{out}$. We consider the cellular chain complexes $C_{\bullet}(G, L), C_{\bullet}(G', L)$ and $C_{\bullet}(G' \circ G, L)$. The map comp is given by Lemma [4.1](#page-13-5) [\(d\)](#page-13-3) applied to the short exact sequence of complexes

$$
0 \to C_{\bullet}(G, L) \to C_{\bullet}(G' \circ G, L) \to C_{\bullet}(G', L) \to 0.
$$

One splitting of this short exact sequence is given by the maps on chains coming from the identifi-cation of the generating sets as in [\(13\)](#page-37-1). \Box

Remark 7.5. We write $E(G) = \{e_1, \ldots, e_k\}, e_i = \{h_i, \sigma h_i\}, V(G) \setminus L_{in}(G) = \{v_1, \ldots, v_l\}, E(G') =$ $\{e'_1,\ldots,e'_m\}, e'_i = \{h'_i,\sigma h'_i\}$ and $V(G') \setminus L_{in}(G') = \{v'_1,\ldots,v'_n\}.$ Then we compute the following Koszul signs:

comp
$$
(\langle e'_1 \wedge \cdots \wedge e'_m \rangle^{-1} \otimes h'_1 \wedge \sigma h'_1 \wedge \cdots \wedge \sigma h'_m \otimes v'_1 \wedge \cdots \wedge v'_n
$$

\n
$$
\otimes (\langle e_1 \wedge \cdots \wedge e_k \rangle^{-1} \otimes h_1 \wedge \sigma h_1 \wedge \cdots \wedge \sigma h_k \otimes v_1 \wedge \cdots \wedge v_l)
$$

\n
$$
= (-1)^{(m+n)k} (\langle e'_1 \wedge \cdots \wedge e'_m \wedge e_1 \wedge \cdots \wedge e_k)^{-1} \otimes h'_1 \wedge \cdots \wedge \sigma h'_m \wedge h_1 \wedge \cdots \wedge \sigma h_k
$$

\n
$$
\otimes v'_1 \wedge \cdots \wedge v'_n \wedge v_1 \wedge \cdots \wedge v_l
$$

in det $(E(G' \circ G))^{-1} \otimes \text{Or}(H(G' \circ G)) \otimes \text{det}(V(G' \circ G) \setminus L_{in}(G' \circ G)).$

7.2. Proof. With the calculation tools in Proposition [7.1,](#page-34-0) Lemma [7.3](#page-35-0) and Lemma [7.4,](#page-36-0) we are in a position to prove Theorem [5.1.](#page-20-0)

Proof of Theorem [5.1.](#page-20-0) We recall that Theorem [5.1](#page-20-0) gives a description of (c, d) -graded commutative (or symmetric) Frobenius algebras in terms of maps and relations that those maps satisfy. We defined a (c, d) -graded commutative (or symmetric) Frobenius algebra as a symmetric monoidal functor preserving degrees out of $\mathrm{GCob}_{c,d}$ (or $\mathrm{fGCob}_{c,d}$).

For the parts of the proof that apply to both categories, we denote $\mathcal{G}_{c,d}$ for the category $\mathrm{GCob}_{c,d}$ or $fGCob_{c,d}$. This is the category enriched in graded abelian groups. We denote by G the corresponding $(2, 2)$ -category of graph cobordisms GCob or fGCob. Finally, we denote by G the associated 1-categories GCob and fGCob where morphisms are equivalence classes of graph cobordisms as in Definition [2.7.](#page-8-0)

We recall that Proposition [4.5](#page-15-3) gives isomorphisms

(14)
$$
\mathcal{G}_{c,d}(X,Y) \cong \bigoplus_{[G]\in\mathcal{G}(X,Y)} \det_{c,d}(G)_{\pi_1(|\underline{\mathcal{G}}(X,Y)|,G)}.
$$

An element $[G] \in \mathcal{G}(X, Y)$ is an equivalence class of a graph cobordism up to zig-zag of graph morphisms. Such an equivalence class can be represented by a graph that can be written as a

composition of the four elementary graphs \rightarrow , \leftarrow , \rightarrow . Indeed, expand the graph such that it has only graphs of valence ≤ 3 and no two valence 3-vertex are next to each other. Direct the edges such that no vertex of valence 2 or 3 has only in-coming or only out-going edges. To make this possible, one has to possibly introduce some units or counits. See Figure [8.](#page-38-0)

Figure 8. The graph on the bottom can be lifted to a graph which can be decomposed into the four elementary graphs. This graph can for example be written as $(\text{-}\otimes)$ o $(\text{-}\otimes \text{id} \otimes \text{id})$ o $(\text{id} \otimes \text{id})$ o $(\text{-}\otimes \text{id})$ o $(\text{-}\otimes)$ o $(\text{-}\otimes)$ o $(\text{-}\otimes \text{id})$. (Note the order-reversing convention for the composition \circ .)

This shows that $\mathcal{G}_{c,d}$ is generated (under composition and symmetric monoidal product) by chosen generators

$$
\begin{aligned} &\omega(\succ) \in \textbf{det}_{c,d}(\succ)_{\pi_1(|\underline{\mathcal{G}}(\{\ast\};\{\ast\});\{\ast\})|,\succ)}, &\hspace{1cm} \omega(\blacktriangleleft) \in \textbf{det}_{c,d}(\blacktriangleleft_{\pi_1(|\underline{\mathcal{G}}(\mathcal{S},\{\ast\})|,\blacktriangle)},\\ &\omega(\blacktriangleleft) \in \textbf{det}_{c,d}(\blacktriangleleft_{\pi_1(|\underline{\mathcal{G}}(\{\ast\};\{\ast\})|,\blacktriangleleft),\succ}), &\hspace{1cm} \omega(\blacktriangleleft) \in \textbf{det}_{c,d}(\blacktriangleleft_{\pi_1(|\underline{\mathcal{G}}(\{\ast\};\mathcal{S})|,\blacktriangleleft),\succ}). \end{aligned}
$$

We note that those four elements are non-trivial for all $c, d \in \mathbb{Z}$. This can be seen as there exists examples of (c, d) -graded commutative Frobenius algebras with non-trivial μ, η, ν and ε (see Example [5.8\)](#page-22-1). However, our proof does not use this fact.

Any symmetric monoidal functor $F: \mathcal{G}_{c,d} \to \mathcal{C}$ is thus uniquely determined by the images $\mu =$ $F(\omega(\)$, $\eta = F(\omega(\bullet)), \nu = F(\omega(\bullet))$ and $\varepsilon = F(\omega(\bullet)).$

We now fix explicit orientations $\omega(\rightarrow)$, $\omega(\rightarrow)$, $\omega(\rightarrow)$ and $\omega(\rightarrow)$: we label the graphs \succ , \leftarrow , \prec and \rightarrow as in Figures [9,](#page-39-0) [10,](#page-39-1) [11](#page-39-2) and [12.](#page-39-3) We choose the convention that all half-edges named σh_i are glued to the internal vertex. This is convenient when we later apply Lemma [7.3](#page-35-0) to compute edge collapses.

FIGURE 9. The graph \rightarrow with v the lone internal vertex, v_0 the lone outgoing vertex and v_1 and v_2 the incoming vertices. The vertex v_1 corresponding to the first coordinate of the input and v_2 to the second coordinate.

FIGURE 11. The graph \bullet with v the lone internal vertex and v_0 the lone outgoing vertex.

FIGURE 10. The graph \prec with v the lone internal vertex, v_0 the lone incoming vertex and v_1 and v_2 the outgoing vertices. The vertex v_1 corresponding to the first coordinate of the output and v_2 to the second coordinate.

FIGURE 12. The graph \rightarrow with v the lone internal vertex and v_0 the lone incoming vertex.

Using the identification as in Proposition [7.1,](#page-34-0) we define generators

$$
\omega_{in}(\mathbf{H}) := (e_2 \wedge e_1 \wedge e_0)^{-1} \otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \otimes v \wedge v_0 \in \det(\mathbf{H}, \partial_{in}),
$$

\n
$$
\omega_{out}(\mathbf{H}) := 1 \in \det(\mathbf{H}, \partial_{out}) \cong \mathbb{I},
$$

\n
$$
\omega_{in}(\mathbf{H}) := e_0^{-1} \otimes h_0 \wedge \sigma h_0 \otimes v \wedge v_0 \in \det(\mathbf{H}, \partial_{in}),
$$

\n
$$
\omega_{out}(\mathbf{H}) := 1 \in \det(\mathbf{H}, \partial_{out}) \cong \mathbb{I},
$$

\n
$$
\omega_{in}(\mathbf{H}) := (e_2 \wedge e_1 \wedge e_0)^{-1} \otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \otimes v \wedge v_0 \in \det(\mathbf{H}, \partial_{out}),
$$

\n
$$
\omega_{in}(\mathbf{H}) := 1 \in \det(\mathbf{H}, \partial_{in}) \cong \mathbb{I},
$$

\n
$$
\omega_{out}(\mathbf{H}) := e_0^{-1} \otimes h_0 \wedge \sigma h_0 \otimes v \wedge v_0 \in \det(\mathbf{H}, \partial_{out}).
$$

The generator $\omega_{in}(\bullet)$ is the canonical generators coming from the isomorphism

$$
\det(H_{\bullet}(\bullet,\partial_{in})) \stackrel{4.1(e)}{\rightarrow} \det(C_{\bullet}(\bullet,\partial_{in})) \stackrel{7.1}{\rightarrow} \det(E(\bullet))^{-1} \otimes \text{Or}(H(\bullet)) \otimes \det(V(\bullet) \setminus L_{in}(\bullet)),
$$

$$
v_0 \mapsto (h_0 \otimes \sigma h_0)^{-1} \otimes v \wedge v_0 \mapsto e_0^{-1} \otimes h_0 \wedge \sigma h_0 \otimes v \wedge v_0.
$$

An analogous statement holds for the chosen generator $\omega_{out}(\rightarrow)$.

We note that $\omega_{out}(\rightarrow), \omega_{out}(\rightarrow), \omega_{in}(\rightarrow)$ and $\omega_{in}(\rightarrow)$ all have degree zero. The generators $\omega_{in}(\rightarrow)$ and $\omega_{out}(\mathcal{A})$ are in degree 1 while $\omega_{in}(\mathcal{A})$ and $\omega_{out}(\mathcal{A})$ are in degree -1. Then we set for fixed $c, d \in \mathbb{Z}$

the orientations

$$
\omega(\boldsymbol{\rangle}) := \omega_{in}(\boldsymbol{\rangle})^{\otimes c} \otimes \omega_{out}(\boldsymbol{\rangle})^{\otimes d} \in \det_{c,d}(\boldsymbol{\rangle}), \qquad \omega(\bullet) := \omega_{in}(\bullet)^{\otimes c} \otimes \omega_{out}(\bullet)^{\otimes d} \in \det_{c,d}(\bullet),
$$

$$
\omega(\mathbf{A}) := \omega_{in}(\mathbf{A})^{\otimes c} \otimes \omega_{out}(\mathbf{A})^{\otimes d} \in \det_{c,d}(\mathbf{A}), \qquad \omega(\bullet) := \omega_{in}(\bullet)^{\otimes c} \otimes \omega_{out}(\bullet)^{\otimes d} \in \det_{c,d}(\bullet).
$$

We note that $\omega(\rightarrow)$ is in degree c and $\omega(\rightarrow)$ is in degree $-c$ while $\omega(\rightarrow)$ is in degree d and $\omega(\rightarrow)$ is in degree $-d$. As F is enriched over \mathbb{ZMod}_0 , it preserves degrees of maps. We thus have $|\mu| = c$, $|\eta| = -c$, $|\nu| = d$ and $|\varepsilon| = -d$.

We have proven so far that a symmetric monoidal functor $F: \mathcal{G}_{c,d} \to \mathcal{C}$ is uniquely determined by the data

$$
A = F(*) \in C,
$$

\n
$$
\mu = F(\omega(\rightarrow)) \in C_c(A \otimes A, A),
$$

\n
$$
\eta = F(\omega(\rightarrow)) \in C_{-c}(1, A),
$$

\n
$$
\tau = F(\omega(\rightarrow)) \in C_{-d}(A, A \otimes A),
$$

\n
$$
\epsilon = F(\omega(\rightarrow)) \in C_{-d}(A, 1).
$$

The theorem we are proving states that this data defines a functor if and only if certain relations are satisfied. We first argue the only if part of the statement assuming the following lemma:

Lemma 7.6. In the category $\mathcal{G}_{c,d}$, the following identities hold:

$$
\omega(\mathbf{H}) \circ (\omega(\mathbf{H})) \otimes 1) = (-1)^{c} \omega(\mathbf{H}) \circ (1 \otimes \omega(\mathbf{H})) \in \det_{c,d}(\mathbf{H}),
$$

$$
(-1)^{c} \omega(\mathbf{H}) \circ (\omega(\mathbf{H})) \otimes 1) = (-1)^{\frac{c(c-1)}{2}} = \omega(\mathbf{H}) \circ (1 \otimes \omega(\mathbf{H})) \in \det_{c,d}(\mathbf{H}) \cong 1,
$$

$$
(\omega(\mathbf{H})) \otimes 1) \circ \omega(\mathbf{H}) = (-1)^{d} (1 \otimes \omega(\mathbf{H})) \circ \omega(\mathbf{H}) \in \det_{c,d}(\mathbf{H}),
$$

$$
(-1)^{d} (1 \otimes \omega(\mathbf{H})) \circ \omega(\mathbf{H}) = (-1)^{\frac{d(d-1)}{2}} = (\omega(\mathbf{H}) \otimes 1) \circ \omega(\mathbf{H}) \in \det_{c,d}(\mathbf{H}) \cong 1,
$$

$$
(\omega(\mathbf{H})) \otimes 1) \circ (1 \otimes \omega(\mathbf{H})) = (-1)^{c d} \omega(\mathbf{H}) \circ \omega(\mathbf{H}) = (1 \otimes \omega(\mathbf{H})) \circ (\omega(\mathbf{H}) \otimes 1) \in \det_{c,d}(\mathbf{H})
$$

Moreover in the category $\text{GCob}_{c,d}$, the automorphism of the graph \rightarrow which switches the two incoming edger acts by $(-1)^c$ on $\det_{c,d}(\succ)$ and, in the category fGCob_{c,d}, the fat graph automorphism of the fat graph > which switches the two incoming edges acts by $(-1)^c$ on $\det_{c,d}(\mathcal{E})$.

These identities are preserved under F because F is linear and monoidal. The lemma thus shows that the relations are satisfied by μ , η , ν and ε if they come from a functor F. This thus concludes the only if part.

Before proving the lemma, we show the if part of the statement that the data $(A, \mu, \eta, \nu, \varepsilon)$ satisfying the relations defines a functor:

Consider a morphism $f \in \mathcal{G}(X, Y)$. As we have

$$
\mathcal{G}_{c,d}(X,Y) \cong \bigoplus_{[G]\in\mathcal{G}(X,Y)} \det_{c,d}(G)_{\pi_1(|\underline{\mathcal{G}}(X,Y)|,G)},
$$

we may assume that f is a homogeneous generator i.e. it is given by the $\pi_1(|\mathcal{G}(X, Y)|, G)$ -orbit of $\omega(G)$ for some generator $\omega(G) \in \det_{c,d}(G)$. We write $[G]$ as a composition of the graphs \rightarrow , \leftarrow , and \bullet as in Figure [8.](#page-38-0) We then define $F(\omega(G))$ as the corresponding composition of μ , η , ν and ε . The map F is by definition linear and degree-preserving i.e. it is enriched over \mathbb{ZMod}_0 . Moreover up to choosing the right decomposition of G, this is functorial and symmetric monoidal. However, our definition of $F(f)$ depends on two choices: the decomposition of G and the choice of representative of the $\pi_1(|\mathcal{G}(X, Y)|, G)$ -orbit.

We first argue why the definition is independent of the choice of decomposition: in [\[Koc04,](#page-47-4) [LP08\]](#page-47-3), Kock and Lauda and Pfeiffer describe a normal form for a decomposition of a closed or an open surface. Moreover, they show that any decomposition can be brought into this normal form using the six generating relations: associativity, unitality, coassociativity, counitality, the Frobenius relation

and either commutativity or symmetry. Thus any two decompositions are related to each other by a sequence of those six relations.

An analogous argument for (fat) graphs shows that any two decompositions into the four graphs \rightarrow \bullet , \prec and \bullet are related by a sequence of those six relations.

Therefore, any two decompositions are related by applying the relations in Lemma [7.6.](#page-40-0) If the morphisms μ , η , ν and ε satisfy the corresponding relation in C, the definition of $F(\omega(G))$ as above is independent of the choice of decomposition.

It remains to show that our definition of F is independent of choice of representative in the $\pi_1(|\mathcal{G}(X, Y)|, G)$ -orbit. This is equivalent to showing that our definition of $F(\omega(G))$ is invariant under the $\pi_1(|\mathcal{G}(X, Y)|, G)$ -action:

Any element in $\pi_1(|\mathcal{G}(X, Y)|, G)$ is represented by a zig-zag of edge collapses and (fat) graph isomorphisms:

 $G = G_0 \longrightarrow G_1 \longleftarrow G_2 \longrightarrow G_3 \longleftarrow \dots \longrightarrow G_{n-1} \longleftarrow G_n = G.$

Applying one of the relations in Lemma [7.6](#page-40-0) to a decomposition G produces a new decomposition G′ . The relation then specifies a zig-zag

 $G \longrightarrow G'' \longleftarrow G'$

where G and G' are given by a different decomposition. We show that all zig-zags can be lifted to a sequence of zig-zags coming from the relations in Lemma [7.6.](#page-40-0)

We can lift each graph G_{2i} to a decomposition into the four elementary graphs: $\widetilde{G}_{2i} \rightarrow G_{2i}$ where G_{2i} has a decomposition into the four elementary graphs. We fix such a decomposition for all G_{2i} . We thus have the following diagram:

$$
\widetilde{G}_0 \qquad \qquad \widetilde{G}_2 \qquad \qquad \cdots \qquad \qquad \widetilde{G}_n
$$
\n
$$
\parallel \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \parallel
$$
\n
$$
G \longrightarrow G_1 \longleftarrow G_2 \longrightarrow G_3 \longleftarrow \cdots \longrightarrow G_{n-1} \longleftarrow G.
$$

Each $\tilde{G}_{2i} \rightarrow G_{2i} \rightarrow G_{2i\pm 1}$ gives a decomposition of $G_{2i\pm 1}$. We thus can rewrite the above diagram suggestively as

$$
\widetilde{G}_0 \longrightarrow \widetilde{G}_0 \qquad \widetilde{G}_2 \longrightarrow \widetilde{G}_2 \longrightarrow \widetilde{G}_2 \qquad \dots \qquad \widetilde{G}_n \longrightarrow \widetilde{G}_n
$$
\n
$$
\downarrow \qquad \downarrow \
$$

The decompositions $\widetilde{G}_{2i} \to G_{2i+1}$ and $\widetilde{G}_{2i+2} \to G_{2i+1}$ are two decompositions of G_{2i+1} . Therefore \tilde{G}_{2i} and \tilde{G}_{2i+2} are related by a sequence of the relations in Lemma [7.6.](#page-40-0) Thus there exists a zig-zag coming from those relations fitting into the lifting problem

This lifts the whole zig-zag to a zig-zag coming from the relations in Lemma [7.6.](#page-40-0) The action of the element of $\pi_1(|\underline{\mathcal{C}}(X, Y)|, G)$ can thus be computed by applying the relations in Lemma [7.6.](#page-40-0)

Consequently, if μ , η , ν and ε satisfy the corresponding relations in C, our definition of $F(\omega(G))$ does not depend on the representative of the $\pi_1(\mathcal{G}(X, Y), G)$ -orbit as it is invariant under the $\pi_1(\mathcal{G}(X, Y), G)$ -action.

This shows that the data μ , η , ν and ε satisfying the relations lets us define a functor F and thus concludes the only if part of the statement. The only thing left to prove is Lemma [7.6.](#page-40-0) The proof makes up the rest of the paper.

Proof of Lemma [7.6.](#page-40-0) We prove the seven identities using Lemmas [7.3](#page-35-0) and [7.4.](#page-36-0)

Graded associativity. We show that

(15)
$$
\omega(\rightarrow) \circ (\omega(\rightarrow) \otimes 1) = (-1)^{c} \omega(\rightarrow) \circ (1 \otimes \omega(\rightarrow)) \in \mathbf{det}_{c,d}(\rightarrow)
$$

holds in $\mathcal{G}_{c,d}$. We start computing

 $\omega(\succ) \circ (\omega(\succ) \otimes 1) = (\omega_{in}(\succ)^{\otimes c} \otimes \omega_{out}(\succ)^{\otimes d}) \circ (\omega_{in}(\succ)^{\otimes c} \otimes \omega_{out}(\succ)^{\otimes d} \otimes 1)$ $=\text{comp}_1(\omega_{in}(\text{)-})^{\otimes c}\otimes \omega_{in}(\text{)-})^{\otimes c})\otimes \text{comp}_1(\omega_{out}(\text{)-})^{\otimes d}\otimes \omega_{out}(\text{)-})^{\otimes d})$

$$
=(-1)^{\frac{c(c-1)}{2}}\text{comp}_1(\omega_{in}(\mathcal{F})\otimes\omega_{in}(\mathcal{F}))^{\otimes c}\otimes\text{comp}_1(\omega_{out}(\mathcal{F})\otimes\omega_{out}(\mathcal{F}))^{\otimes d}.
$$

Here comp₁ is the composition we get by gluing one copy of \rightarrow at v_0 to v_1 of the other copy. The sign $(-1)^{\frac{c(c-1)}{2}}$ comes from the fact that we have to reshuffle $\omega_{in}(\succ)^{\otimes c}\otimes \omega_{in}(\succ)^{\otimes c}$ into c pairs of $\omega_{in}(\)$ $\otimes \omega_{in}(\)$ (via a riffle shuffle). For $\omega_{out}(\)$ no such sign appears as they are in degree 0. Analogously, we compute

$$
\omega(\boldsymbol{\succ}) \circ (1 \otimes \omega(\boldsymbol{\succ})) = (-1)^{\frac{c(c-1)}{2}} \text{comp}_2(\omega_{in}(\boldsymbol{\succ}) \otimes \omega_{in}(\boldsymbol{\succ}))^{\otimes c} \otimes \text{comp}_2(\omega_{out}(\boldsymbol{\succ}) \otimes \omega_{out}(\boldsymbol{\succ}))^{\otimes d}.
$$

Here comp₂ is the composition we get by gluing one copy of \rightarrow at v_0 to v_2 of the other copy because v_2 is the vertex corresponding to the second input.

It remains to compare $\text{comp}_1(\omega_{in}(\text{)-}) \otimes \omega_{in}(\text{)-})$ to $\text{comp}_2(\omega_{in}(\text{)-}) \otimes \omega_{in}(\text{)-})$ and $\text{comp}_1(\omega_{out}(\text{)-}) \otimes \omega_{in}(\text{)-})$ $\omega_{out}(\rangle)$ to $comp_2(\omega_{out}(\rangle) \otimes \omega_{out}(\rangle)).$

We note that because of triviality of the $\omega_{out}(\rightarrow)$ orientations, we have

comp₁
$$
(\omega_{out}(\rangle \rightarrow) \otimes \omega_{out}(\rangle \rightarrow)) = comp_{2}(\omega_{out}(\rangle \rightarrow) \otimes \omega_{out}(\rangle \rightarrow)).
$$

Equation [\(15\)](#page-42-1) then follows from the following identity

(16)
$$
\operatorname{comp}_1(\omega_{in}(\succ) \otimes \omega_{in}(\succ)) = -\operatorname{comp}_2(\omega_{in}(\succ) \otimes \omega_{in}(\succ)).
$$

To show this identity, we start with $\text{comp}_1(\omega_{in}(\text{)-}) \otimes \omega_{in}(\text{)-})$. For this, we denote the edges, halfedges and vertices of the left copy of \rightarrow by e, e_0, e_1, e_2 and so on, as before. This represents the multiplication that is applied second. We denote the edges, half-edges and vertices of the right copy of \rightarrow by e' , e'_{0} , e'_{1} , e'_{2} and so on. This represents the multiplication that is applied first. We compute using Lemma [7.4:](#page-36-0)

comp₁(
$$
\omega_{in}
$$
(\rightarrow) $\otimes \omega_{in}$ (\rightarrow)) = comp₁(($e_2 \wedge e_1 \wedge e_0$)⁻¹ $\otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \otimes v \wedge v_0$
\n $\otimes (e'_2 \wedge e'_1 \wedge e'_0)^{-1} \otimes h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v' \wedge v'_0)$
\n= $(e_2 \wedge e_1 \wedge e_0 \wedge e'_2 \wedge e'_1 \wedge e'_0)^{-1}$
\n $\otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \wedge h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2$
\n $\otimes v \wedge v_0 \wedge v' \wedge v'_0$

Figure 13. The two composition of two multiplications give the same graph after collapsing two edges.

As we glued v'_0 to v_1 , we can collapse e'_0 and e_1 to obtain the graph \rightarrow (see top of Figure [13\)](#page-43-0). We recall the naming convention that $s(\sigma h_i)$ is an internal vertex. Therefore when collapsing e'_0 , we can apply Lemma [7.3](#page-35-0) using that $s(\sigma h'_0) = v'$ is the vertex we want to delete from the orientation. Thereafter deleting e_1 , we note that $s(\sigma h_1) = v$ is the vertex we delete in the process. Importantly to apply Lemma [7.3,](#page-35-0) we must reshuffle the edge and vertex, we want to delete to the first position. This gives some signs. We do the reshuffle explicitly in the first computation and thereafter only denote the corresponding sign after the edge collapse. We find:

$$
-(e_2 \wedge e_1 \wedge e_0 \wedge e'_2 \wedge e'_1 \wedge e'_0)^{-1}
$$

\n
$$
\otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \wedge h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v \wedge v_0 \wedge v' \wedge v'_0
$$

\n
$$
=(e'_0 \wedge e_2 \wedge e_1 \wedge e_0 \wedge e'_2 \wedge e'_1)^{-1}
$$

\n
$$
\otimes h'_0 \wedge \sigma h'_0 \wedge h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v' \wedge v \wedge v_0 \wedge v'_0
$$

\n
$$
\mapsto (e_2 \wedge e_1 \wedge e_0 \wedge e'_2 \wedge e'_1)^{-1}
$$

\n
$$
\otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v \wedge v_0 \wedge v'_0
$$

\n
$$
\mapsto -(e_2 \wedge e_0 \wedge e'_2 \wedge e'_1)^{-1} \otimes h_0 \wedge \sigma h_0 \wedge h_2 \wedge \sigma h_2 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v_0 \wedge v'_0.
$$

The edges of \rightarrow are here labelled (from top to bottom) by e'_1, e'_2 and e_2 as in the top right of Figure [13.](#page-43-0)

Analogously, we compute

comp₂(
$$
\omega_{in}(\rightarrow)
$$
) $\otimes \omega_{in}(\rightarrow)$) = - ($e_2 \wedge e_1 \wedge e_0 \wedge e'_2 \wedge e'_1 \wedge e'_0$)⁻¹
 $\otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \wedge h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2$
 $\otimes v \wedge v_0 \wedge v' \wedge v'_0$.

But as we glued v'_0 to v_2 , we can now collapse e'_0 and e_2 (see bottom of Figure [13\)](#page-43-0). In this process, we first delete the vertex $s(\sigma h'_0) = v'$ and then the vertex $s(\sigma h_2) = v$. We thus find

$$
-(e_2 \wedge e_1 \wedge e_0 \wedge e'_2 \wedge e'_1 \wedge e'_0)^{-1}
$$

\n
$$
\otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \wedge h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v \wedge v_0 \wedge v' \wedge v'_0
$$

\n
$$
\rightarrow (e_2 \wedge e_1 \wedge e_0 \wedge e'_2 \wedge e'_1)^{-1}
$$

\n
$$
\otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v \wedge v_0 \wedge v'_0
$$

\n
$$
\mapsto (e_1 \wedge e_0 \wedge e'_2 \wedge e'_1)^{-1} \otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v_0 \wedge v'_0.
$$

The edges of \rightarrow are here labelled (from top to bottom) by e_1 , e'_1 and e'_2 as in the bottom right of Figure [13.](#page-43-0) This differs from the naming scheme of the other composition by an even permutation. This shows that the two orientations differ by the sign -1 . We have proved [\(16\)](#page-42-2) and thus [\(15\)](#page-42-1).

Graded unitality. We want to compute

(17)
$$
(-1)^{c} \omega(\mathbf{r}) \circ (\omega(\mathbf{r}) \otimes 1) = (-1)^{\frac{c(c-1)}{2}} = \omega(\mathbf{r}) \circ (1 \otimes \omega(\mathbf{r})) \in \mathbf{det}_{c,d}(\mathbf{r}) \cong \mathbb{1}.
$$

As for graded associativity, we compute the signs from the riffle shuffle

$$
\omega(\boldsymbol{\succ}) \circ (\omega(\mathbf{e}) \otimes 1) = (-1)^{\frac{c(c-1)}{2}} \text{comp}_1(\omega_{in}(\boldsymbol{\succ}) \otimes \omega_{in}(\mathbf{e}))^{\otimes c} \otimes \text{comp}_1(\omega_{out}(\boldsymbol{\succ}) \otimes \omega_{out}(\mathbf{e}))^{\otimes d}
$$

and

$$
\omega(\boldsymbol{\mathcal{F}})\circ(1\otimes\omega(\bullet))=(-1)^{\frac{c(c-1)}{2}}\text{comp}_2(\omega_{in}(\boldsymbol{\mathcal{F}})\otimes\omega_{in}(\bullet))^{\otimes c}\otimes\text{comp}_2(\omega_{out}(\boldsymbol{\mathcal{F}})\otimes\omega_{out}(\bullet))^{\otimes d}.
$$

We note that, by triviality of the ω_{out} -orientations, we have

$$
\mathrm{comp}_1(\omega_{out}(\boldsymbol{\cdot}) \otimes \omega_{out}(\bullet)) = 1 = \mathrm{comp}_2(\omega_{out}(\boldsymbol{\cdot}) \otimes \omega_{out}(\bullet)) \in \mathbf{det}(\bullet, \partial_{out}).
$$

The equation [\(17\)](#page-44-1) then follows from the following identity

(18)
$$
-\text{comp}_1(\omega_{in}(\boldsymbol{\cdot}) \otimes \omega_{in}(\bullet)) = 1 = \text{comp}_2(\omega_{in}(\boldsymbol{\cdot}) \otimes \omega_{in}(\bullet)) \in \text{det}(\bullet, \partial_{in}) \cong \mathbb{1}.
$$

To show this, we denote the edges, half-edges and vertices of \leftarrow by e_0 and so on. On the other hand, we denote the edges, half-edges and vertices of \succ by e'_{0} , e'_{1} , e'_{2} and so on. We thus compute by Lemma [7.4:](#page-36-0)

$$
\text{comp}_1(\omega_{in}(\mathbf{r}) \otimes \omega_{in}(\mathbf{r})) = \text{comp}_1((e'_2 \wedge e'_1 \wedge e'_0)^{-1} \otimes h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v' \wedge v'_0
$$

$$
\otimes (e_0)^{-1} \otimes h_0 \wedge \sigma h_0 \otimes v \wedge v_0)
$$

$$
= -(e'_2 \wedge e'_1 \wedge e'_0 \wedge e_0)^{-1}
$$

$$
\otimes h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \wedge h_0 \wedge \sigma h_0 \otimes v' \wedge v'_0 \wedge v \wedge v_0.
$$

Figure 14. The two composition of a unit and a multiplications give the graph with one vertex representing id after collapsing four edges.

Now we can collapse e_0 and e'_1 as in the top of Figure [14.](#page-45-1) We use Lemma [7.3](#page-35-0) and note that when we collapse e_0 , we have to delete $s(\sigma h_0) = v$. If we then collapse e'_1 , we note that $s(\sigma h'_1) = v'$. We compute:

$$
-(e'_2 \wedge e'_1 \wedge e'_0 \wedge e_0)^{-1} \otimes h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \wedge h_0 \wedge \sigma h_0 \otimes v' \wedge v'_0 \wedge v \wedge v_0
$$

\n
$$
\rightarrow (e'_2 \wedge e'_1 \wedge e'_0)^{-1} \otimes h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \otimes v' \wedge v'_0 \wedge v_0
$$

\n
$$
\rightarrow -(e'_2 \wedge e'_0)^{-1} \otimes h'_0 \wedge \sigma h'_0 \wedge h'_2 \wedge \sigma h'_2 \otimes v'_0 \wedge v_0.
$$

Next we collapse e'_0 which deletes $s(\sigma h'_0) = v_0$:

$$
-(e'_2 \wedge e'_0)^{-1} \otimes h'_0 \wedge \sigma h'_0 \wedge h'_2 \wedge \sigma h'_2 \otimes v'_0 \wedge v_0
$$

$$
\mapsto -(e'_2)^{-1} \otimes h'_2 \wedge \sigma h'_2 \otimes v'_0.
$$

Finally collapsing e'_2 , we delete $s(\sigma h'_2) = v'_0$ and find:

$$
-(e_2')^{-1} \otimes h_2' \wedge \sigma h_2' \otimes v_0' \mapsto -1.
$$

This shows the left equality in (18) . For the other equation, we compute:

comp₂(
$$
\omega_{in}(\rightarrow)
$$
) $\otimes \omega_{in}(\rightarrow)$) = - ($e'_2 \wedge e'_1 \wedge e'_0 \wedge e_0$)⁻¹
 $\otimes h'_0 \wedge \sigma h'_0 \wedge h'_1 \wedge \sigma h'_1 \wedge h'_2 \wedge \sigma h'_2 \wedge h_0 \wedge \sigma h_0 \otimes v' \wedge v'_0 \wedge v \wedge v_0$.

But now we collapse first e_0 and at the same time deleting $s(\sigma h_0) = v$. Then we collapse e'_2 and delete $s(\sigma h'_2) = v'$. Then we delete e'_0 and delete $s(\sigma h'_0) = v_0$. Finally, we collapse e'_1 and delete $s(\sigma h'_1) = v'_0$ as in the bottom of Figure [14.](#page-45-1) We thus find

$$
-(e_0\wedge e'_2\wedge e'_1\wedge e'_0)^{-1}\otimes h_0\wedge \sigma h_0\wedge h'_0\wedge \sigma h'_0\wedge h'_1\wedge \sigma h'_1\wedge h'_2\wedge \sigma h'_2\otimes v\wedge v_0\wedge v'\wedge v'_0\mapsto 1.
$$

One can see that by either doing the calculation explicitly. Or one can notice that we are deleting the vertices in the same order as before but switched the order of the two edges e'_1 and e'_2 in which we delete them (see bottom of Figure [14\)](#page-45-1). This shows [\(18\)](#page-44-2) and thus [\(17\)](#page-44-1).

Graded coassociativity. This computation is dual to the computation of graded associativity, we thus obtain:

$$
(\omega(\mathcal{A}) \otimes 1) \circ \omega(\mathcal{A}) = (-1)^d (1 \otimes \omega(\mathcal{A})) \circ \omega(\mathcal{A}) \in \mathbf{det}_{c,d}(\mathcal{A}).
$$

Graded counitality. This computation is dual to the computation of graded unitality, we thus obtain:

$$
(-1)^d (1 \otimes \omega(\mathbin{\blacklozenge})) \circ \omega(\mathbin{\blacklozenge}) = (-1)^{\frac{d(d-1)}{2}} = (\omega(\mathbin{\blacklozenge}) \otimes 1) \circ \omega(\mathbin{\blacklozenge}) \in \textnormal{det}_{c,d}(\mathbin{\blacklozenge}) \cong \mathbb{1}.
$$

We note that the dual term to $\omega(\rightarrow) \circ (\omega(\rightarrow) \otimes 1)$ is $(1 \otimes \omega(\rightarrow)) \circ \omega(\rightarrow)$.

The graded Frobenius relation. We want to show that

(19)
$$
(\omega(\mu) \otimes 1) \circ (1 \otimes \omega(\nu)) = (-1)^{cd} \omega(\nu) \circ \omega(\mu) = (1 \otimes \omega(\mu)) \circ (\omega(\nu) \otimes 1) \in \mathbf{det}_{c,d}(\mathbf{X}).
$$

We start computing

$$
(\omega(\mu) \otimes 1) \circ (1 \otimes \omega(\nu)) = (\omega_{in}(\mu)^{\otimes c} \otimes \omega_{out}(\mu)^{\otimes d} \otimes 1) \circ (1 \otimes \omega_{in}(\nu)^{\otimes c} \otimes \omega_{out}(\nu)^{\otimes d})
$$

=
$$
(\omega_{in}(\mu)^{\otimes c} \otimes 1) \circ (1 \otimes \omega_{in}(\nu)^{\otimes c}) \otimes (\omega_{out}(\mu)^{\otimes d} \otimes 1) \circ (1 \otimes \omega_{out}(\nu)^{\otimes d})
$$

=
$$
\text{comp}_{1,2}(\omega_{in}(\mu) \otimes \omega_{in}(\nu))^{\otimes c} \otimes \text{comp}_{1,2}(\omega_{out}(\mu) \otimes \omega_{out}(\nu))^{\otimes d}.
$$

Here $\text{comp}_{1,2}$ is the composition given by gluing together the first outgoing vertex of \prec to the second incoming vertex of \rangle .

Analogously, we compute

$$
(1\otimes \omega(\mu))\circ (\omega(\nu)\otimes 1) = \mathrm{comp}_{2,1}(\omega_{in}(\mu)\otimes \omega_{in}(\nu))^{\otimes c}\otimes \mathrm{comp}_{2,1}(\omega_{out}(\mu)\otimes \omega_{out}(\nu))^{\otimes d}.
$$

Here $\text{comp}_{2,1}$ is the composition given by gluing together the second outgoing vertex of \prec to the first incoming vertex of \rightarrow .

In contrast, we compute

$$
\omega(\nu) \circ \omega(\mu) = (\omega_{in}(\nu)^{\otimes c} \otimes \omega_{out}(\nu)^{\otimes d}) \circ (\omega_{in}(\mu)^{\otimes c} \otimes \omega_{out}(\mu)^{\otimes d})
$$

= $(-1)^{cd}(\omega_{in}(\nu)^{\otimes c} \circ \omega_{in}(\mu)^{\otimes c}) \otimes (\omega_{out}(\nu)^{\otimes d} \circ \omega_{out}(\mu)^{\otimes d}).$

The sign $(-1)^{cd}$ comes from the fact that we have to move $\omega_{out}(\nu)^{\otimes d}$ which is in degree d past $\omega_{in}(\mu)^{\otimes c}$ which is in degree c. By definition, we have:

$$
(-1)^{cd}(\omega_{in}(\nu)^{\otimes c} \circ \omega_{in}(\mu)^{\otimes c}) \otimes (\omega_{out}(\nu)^{\otimes d} \circ \omega_{out}(\mu)^{\otimes d})
$$

=
$$
(-1)^{cd} \text{comp}(\omega_{in}(\nu) \otimes \omega_{in}(\mu))^{\otimes c} \otimes \text{comp}(\omega_{out}(\nu) \otimes \omega_{out}(\mu))^{\otimes d}.
$$

The equation [\(19\)](#page-46-3) now follows from the following identities It holds that

$$
\text{comp}_{1,2}(\omega_{in}(\mu) \circ \omega_{in}(\nu)) = \text{comp}(\omega_{in}(\nu) \otimes \omega_{in}(\mu)) = \text{comp}_{2,1}(\omega_{in}(\mu) \circ \omega_{in}(\nu))
$$

and

$$
\text{comp}_{1,2}(\omega_{out}(\mu) \circ \omega_{out}(\nu)) = \text{comp}(\omega_{out}(\nu) \otimes \omega_{out}(\mu)) = \text{comp}_{2,1}(\omega_{out}(\mu) \circ \omega_{out}(\nu)).
$$

This can be seen either by doing the hands-on computation using Lemmas [7.3](#page-35-0) and [7.4](#page-36-0) or by noting that $\omega_{in}(\nu)$ and $\omega_{out}(\mu)$ are trivial. Composition with them thus gives the trivial map.

Graded Commutativity. We want to show that the automorphism of the graph \rightarrow which switches the two incoming edges acts by $(-1)^c$ on $\omega(\)$. For this, we note that this automorphism acts trivially on $\det(\rightarrow, \partial_{out}) \cong \mathbb{1}$. On the other hand, we compute the action $\det(\rightarrow, \partial_{in})$ as

$$
\textbf{det}(\bm{\succ},\partial_{in})\rightarrow \textbf{det}(\bm{\succ},\partial_{in}),
$$

$$
(e_2 \wedge e_1 \wedge e_0)^{-1} \otimes h_0 \wedge \sigma h_0 \wedge h_1 \wedge \sigma h_1 \wedge h_2 \wedge \sigma h_2 \otimes v \wedge v_0
$$

$$
\rightarrow (e_1 \wedge e_2 \wedge e_0)^{-1} \otimes h_0 \wedge \sigma h_0 \wedge h_2 \wedge \sigma h_2 \wedge h_1 \wedge \sigma h_1 \otimes v \wedge v_0.
$$

It only changes the order of e_1 and e_2 and their half-edges. In other words, it acts by -1 on $\det(\mathcal{F}, \partial_{in})$. This combines to a $(-1)^c$ -action on $\det_{c,d}(\mathcal{F})$.

Graded Symmetry. We note that flipping the two incoming vertices of \vdash is not a fat graph automorphism. As before, we call the edges, half-edges and vertices of $\rightarrow e_0, e_1, e_2$ and so on. On the other hand, we call the edges, half-edges and vertices of $\rightarrow e'_0$, h_0 , σh_0 and v'_0 , v' . We glue v_0 to v'_0 , then we can collapse the edge e_0 and e'_0 . On the resulting graph), flipping v_1 with v_2 is a fat graph automorphism. This acts on $\det(\lambda, \partial_{in})$ by -1 and on $\det(\lambda, \partial_{out})$ trivially. This combines to a $(-1)^c$ -action on $\det_{c,d}(\lambda)$.

Therefore, we have computed all signs appearing in the identities in Lemma [7.6.](#page-40-0) This concludes the proof of Lemma [7.6,](#page-40-0) the proof of Theorem [5.1](#page-20-0) and thus the paper. $□$

□

REFERENCES

- [Abr96] Lowell Abrams. Two-dimensional topological quantum field theories and Frobenius algebras. Journal of Knot theory and its ramifications, 5(05):569–587, 1996.
- [CG04] Ralph L Cohen and Véronique Godin. A polarized view of string topology. London Mathematical Society Lecture Note Series, 308:127, 2004.
- [Chi14] Jonathan Chiche. La théorie de l'homotopie des 2-catégories. arXiv preprint arXiv:1411.6936, 2014.
- [CHO20] Kai Cieliebak, Nancy Hingston, and Alexandru Oancea. Poincaré duality for loop spaces. arXiv preprint arXiv:2008.13161, 2020.
- [CHO23] Kai Cieliebak, Nancy Hingston, and Alexandru Oancea. Loop coproduct in Morse and Floer homology. Journal of Fixed Point Theory and Applications, 25(2):59, 2023.
- [CO22a] Kai Cieliebak and Alexandru Oancea. Reduced symplectic homology and the secondary continuation map. arXiv preprint arXiv:2209.02972, 2022.
- [CO22b] Kai Cieliebak and Alexandru Oancea. Topological Frobenius Algebras. arXiv preprint arXiv:2209.02979, 2022.
- [CO24] Kai Cieliebak and Alexandru Oancea. Rabinowitz Floer homology as a Tate vector space. arXiv preprint arXiv:2407.21741, 2024.
- [Gan02] Wee Liang Gan. Koszul duality for dioperads. arXiv preprint math/0201074, 2002.
- [GH09] Mark Goresky and Nancy Hingston. Loop products and closed geodesics. Duke Mathematical Journal, 150(1):117, 2009.
- [GK98] Ezra Getzler and Mikhail M Kapranov. Modular operads. Compositio Mathematica, 110(1):65–125, 1998.
- [God07] Véronique Godin. Higher string topology operations. $a rXiv$ preprint $a rXiv$:0711.4859, 2007.
- [Hat02] Allen Hatcher. Algebraic Topology. Cambridge University Press, 2002.
- [HW17] Nancy Hingston and Nathalie Wahl. Product and coproduct in string topology. arXiv preprint arXiv:1709.06839, 2017.
- [Koc04] Joachim Kock. Frobenius algebras and 2-d topological quantum field theories. Number 59 in London Mathematical Society Student Texts. Cambridge University Press, 2004.
- [KTV21] Maxim Kontsevich, Alex Takeda, and Yiannis Vlassopoulos. Pre-Calabi-Yau algebras and topological quantum field theories. arXiv preprint arXiv:2112.14667, 2021.
- [LP08] Aaron D Lauda and Hendryk Pfeiffer. Open–closed strings: Two-dimensional extended TQFTs and Frobenius algebras. Topology and its Applications, 155(7):623–666, 2008.
- [Mar08] Martin Markl. Operads and PROPs. Handbook of algebra, 5:87–140, 2008.
- [San15] Daniela Egas Santander. Comparing fat graph models of moduli space. arXiv preprint arXiv:1508.03433, 2015.
- [Tam10] Hirotaka Tamanoi. Loop coproducts in string topology and triviality of higher genus TQFT operations. Journal of Pure and Applied Algebra, 214(5):605–615, 2010.
- [WW16] Nathalie Wahl and Craig Westerland. Hochschild homology of structured algebras. Advances in Mathematics, 288:240–307, 2016.