Preconditioned Subspace Langevin Monte Carlo

Tyler Maunu, Jiayi Yao Brandeis University maunu@brandeis.edu, jiayiyao@brandeis.edu

We develop a new efficient method for high-dimensional sampling called Subspace Langevin Monte Carlo. The primary application of these methods is to efficiently implement Preconditioned Langevin Monte Carlo. To demonstrate the usefulness of this new method, we extend ideas from subspace descent methods in Euclidean space to solving a specific optimization problem over Wasserstein space. Our theoretical analysis demonstrates the advantageous convergence regimes of the proposed method, which depend on relative conditioning assumptions common to mirror descent methods. We back up our theory with experimental evidence on sampling from an ill-conditioned Gaussian distribution.

1. Introduction

The Langevin diffusion and its variants have become a fundamental object of study in modern machine learning. On the mathematical side, these diffusions have a deep connection to Wasserstein gradient flows. This connection has been used to study their convergence and to consequently develop new and more efficient diffusions. Practically, discretizations of the Langevin diffusion are highly scalable for generating samples from complex, high-dimensional target distribution. Many examples of the successful application of these methods exist, including denoising diffusion models [1, 2], characterization of complex Bayesian posteriors [3], and differential privacy mechanisms [4].

As mentioned in the previous paragraph, the Langevin diffusion solves an optimization problem in Wasserstein space. Taking this point of view, a theme in recent research has been translating ideas from gradient-based optimization over Euclidean spaces to optimization over Wasserstein spaces. In particular, this has resulted in a growing zoo of optimization methods over Wasserstein space, which include notions of gradient descent [5], mirror descent and Newton methods [6–9], proximal algorithms [10], preconditioned methods [11, 12], and more.

One class of methods that are recently gaining steam in the optimization literature are blockcoordinate and subspace descent methods [13–20]. These methods incorporate low-dimensional updates, and so can be highly efficient and scalable in high dimensions. The projected version of the gradient has advantages in terms of directional derivative oracle calls [21] as well as memory [20]. Furthermore, one can efficiently incorporate higher-order information, which is present in adaptive methods [20] and Newton-type methods [15]. Importantly, analogs of most of these methods have not yet been extended to Wasserstein space outside of coordinate descent [22, 23].

Since in many modern applications, the Langevin algorithm is run in high-dimensional spaces, it stands to reason that it could benefit from further studies into more efficient updating schemes. This work represents a significant step in achieving this goal. Our key contributions are:

- 1. We discuss the convergence of Preconditioned Langevin Monte Carlo (PLMC) from the perspective of Mirror Langevin Monte Carlo. This specific connection has not been discussed before, and in some simple examples, we can show how it outperforms Langevin Monte Carlo (LMC) due to its dependence on notions of relative conditioning of the target distribution.
- 2. The implementation of Preconditioned Langevin Monte Carlo involves the multiplication of a $d \times d$ with the gradient at each iteration, which scales poorly in high dimensions both in terms of complexity and memory. While this can be alleviated with an initial eigenvalue

decomposition, it can also be alleviated by subsampling. We develop a random subspace approximation of the Preconditioned Langevin Monte Carlo method, which we call Subspace Langevin Monte Carlo (SLMC). SLMC is a significant generalization of past work on Random Coordinate Langevin Monte Carlo (RCLMC) [22], since it has a low-rank preconditioner rather than just updating individual coordinates.

- 3. We show the achieved theoretical convergence bounds for SLMC outperform past methods of PLMC, LMC, and RCLMC in certain regimes. Furthermore, it can operate with flexible memory cost depending on the rank of the updates.
- 4. We finish with some simple experiments on an ill-conditioned Gaussian distribution with a block-covariance. These experiments are meant to demonstrate the flexibility of the method developed here for fast estimation in such settings.

1.1. Related Work

The most directly related works to ours are the sequence of works studying Random Coordinate Langevin Monte Carlo (RCLMC) and its reduced variance versions [22, 23]. In these, the authors randomly sample coordinates and take a Langevin step along that coordinate only at each iteration. They also incorporate variance reduction techniques. Another work in this vein is Roy et al. [24], which uses random directions to compute zeroth order approximations to the gradient that are then used for a Langevin method.

Another relevant work considers Subspace Diffusion Models [25]. The authors train diffusion models that learn over a sequence of nested subspaces. Since the SDE in each time interval has a solution constrained to each subspace, they can learn the score function restricted to these subspaces, which results in efficiency and accuracy gains. Other recent work has explored low-dimensionality within diffusion models themselves. For example, Wang et al. [26] shows that diffusion models trained on mixtures of low-rank Gaussians do not incur the curse of dimensionality. Chen et al. [27] use low-rank structures of the Jacobian of the posterior mean prediction to edit diffusion models.

Random projections have a long history in data science, beginning with the seminal work of Johnson et al. [28]. Whereas the initial work used Gaussian projections, more recent sketching algorithms have utilized structured random projection matrices to make algorithms more efficient. Variants of sketching algorithms include CountSketch [29], sparse JL transform [30], Subsampled Randomized Hadamard Transform [31, 32], and more. See Halko et al. [33] for an overview.

Our work draws inspiration from the many existing works studying subspace methods for optimization. The oldest work on random subspace methods are those on random coordinate and block coordinate descent; see the discussion in Nesterov [13], Wright [34]. Early works [35, 36] use randomized methods for solving linear systems. Frongillo and Reid [37] analyzed subspace descent as a unification of coordinate and block coordinate descent methods. Nesterov and Spokoiny [38] uses random Gaussian search directions for gradient-free optimization. Later, Kozak et al. [16] generalize this to random subspaces and study the effect of uniformly random projections on gradient descent.Gower et al. [14] study a Newton method on random subspaces, and Hanzely et al. [15] develop a random subspace-based Cubic Newton Method. Other recent work has used sketching for more efficient SDP solvers [39]. Ivkin et al. [40] uses gradient sketches for efficient distributed optimization. Variance reduction with gradient sketching has also been studied [41, 42].

Other works consider the computation of a good subspace rather than using random subspaces for optimization. Cosson et al. [21] compute a PCA on sampled gradients of a function and then run gradient descent constrained to the found subspace. Other works seek to find memory-efficient methods for training large language models by using adaptive projections [18–20]. Our work can be seen as a hybrid of random subspace updates with some adaptation since the subspaces are taken from the eigendecomposition of a preconditioning matrix A.

Related works also include studies into the low-dimensionality of stochastic gradient descent iterates in high-dimensional nonconvex optimization. Earlier works of Sagun et al. [43, 44] showed this was empirically the case. Later, Jastrzębski et al. [45] showed that SGD dynamics tend to follow sharp directions in the loss landscape. Arous et al. [46] show that training aligns with principal subspaces of the Hessian or Gram matrices in multi-class logistic regression or XOR classification. On the other hand, Song et al. [47] argues that for general deep learning tasks, learning by SGD does not occur in the top subspace of the Hessian, but rather in the bulk subspace. Li et al. [48] and Gressmann et al. [49] use projections to random subspaces to get around this, and Li et al. [50] uses a carefully chosen subspace to improve DNN training.

Another strategy to make sampling more efficient involves running a diffusion in a latent space Vahdat et al. [51], Rombach et al. [52]. However, for such strategies, one must train a variational autoencoder to perform the embedding. This additional step has made it difficult to give useful theoretical bounds on the mixing time of the resulting diffusion method, although some limited results exist Tzen and Raginsky [53]. Also, the dimension of latent spaces is typically still high.

Finally, random projections have been used in the study of Wasserstein space, particularly in the study of Sliced Wasserstein Distances [54]. These offer a scalable way of estimating distances between high-dimensional distributions and have also been used in the efficient implementation of gradient flows [55]. We note that this path is distinct from the one we take here: *they develop gradient flows of the space of probability measures equipped with a different metric (the sliced Wasserstein distance).* We seek to approximate the Wasserstein gradient flow itself with random projections.

1.2. Notation

For a positive definite matrix A, we define an inner product $\langle x, y \rangle_A = x^T A y$. The norm $\|\cdot\|_A^2$ is defined similarly. Random vectors are defined using upper case letters, and matrices are defined with bold upper case letters. Fixed vectors and scalars are lowercase letters, and it should be clear from the context which is which. The Euclidean norm for vectors is $\|\cdot\|$, and the spectral (operator) norm for a matrix A is $\|A\|_2$.

2. The Subspace Langevin Algorithm

As was discussed in the introduction, the many variants of subspace descent have increasingly been used to scale optimization methods in high-dimensional settings. Our focus in this paper is to begin to bring these ideas to optimization over Wasserstein space by focusing first on the particular problem of minimizing the Kullback-Leibler (KL) divergence between a proposal and a target measure.

We begin by outlining subspace gradient methods in Euclidean space in Section 2.1. After this, we discuss optimization over Wasserstein space and how LMC can be thought of as a discretization of a Wasserstein gradient flow in Section 2.2. Then, Section 2.3 discusses how the addition of preconditioning can improve convergence bounds for the Wasserstein gradient flow and its discretization. We finish in Section 2.4 with our proposed method SLMC.

2.1. Euclidean Subspace Descent

Suppose that we wish to minimize a function f over \mathbb{R}^d . Subspace gradient methods in Euclidean spaces have both a continuous and discrete formulation. In continuous time, for a sequence of projection matrices P_t , $t \in [0, \infty)$, the subspace gradient flow of f initialized at x_0 solves the ODE

$$\dot{x}_t = -\boldsymbol{P}_t \nabla f(x_t). \tag{1}$$

The forward Euler discretization of this flow is

$$x_{k+1} = x_k - h\boldsymbol{P}_k \nabla f(x_k). \tag{2}$$

for a sequence of low-rank matrices $(\mathbf{P}_k)_{k \in \mathbb{N}}$ and step sizes h. Convergence of the discrete method is considered in [16], where under the assumption the matrices \mathbf{P}_k are rank r, $\mathbb{E}\mathbf{P}_k = \mathbf{I}$ (which means that \mathbf{P}_k are actually scaled projection matrices), f is β -smooth (it has a quadratic upper bound at

all points in the domain), and satisfies a PL inequality, $f(x) - f(x_*) \leq \frac{1}{2\alpha} \|\nabla f(x)\|^2$, one can show

$$\mathbb{E}f(x_k) - f(x_\star) \le \omega^k (f(x_0) - f(x_\star)) \tag{3}$$

for $\omega = 1 - \frac{r\beta}{d\alpha}$. The iteration complexity in this setting is $O(\frac{d\alpha}{r\beta}\log(1/\epsilon))$ to achieve ϵ expected error. We include a guarantee of convergence in continuous time in Appendix B

We notice that there is no free lunch regarding computational cost here: per-iteration complexity of subspace gradient descent requires r directional derivative computations. In contrast, the periteration complexity of gradient descent requires d directional derivatives. Therefore, both methods have an overall complexity in terms of directional derivatives $O(d\kappa \log(1/\epsilon))$. Even though there is no improvement in terms of the overall complexity, there are three key advantages of subspace descent. First, the smoothness parameter β can be traded for a generally smaller directional smoothness that we outline later in Assumption 3. This is analogous to what is done in coordinate descent methods [34]. The second advantage is in terms of memory. In particular, the method does not need to store the full d-dimensional gradient at each iteration; it can just store the r-dimensional directional gradient. Third, in settings such as PDE-constrained optimization [16], computation of a high-dimensional gradient is expensive. If our updates are constrained to low-dimensional subspaces, we can efficiently solve the PDE in said space and thus avoid the curse of dimensionality.

2.2. Wasserstein Gradient Flows and Langevin Monte Carlo

We now briefly outline Wasserstein gradient flows and their connection to Langevin Monte Carlo. For a more detailed discussion of these topics, we refer the reader to [56, 57]. For a friendly introduction to Wasserstein gradient flows, we recommend [5, 58]. Throughout, we restrict to measures with finite second moment and densities with respect to the Lebesgue measure, $\mathcal{P}_{2,ac}(\mathbb{R}^d)$. The 2-Wasserstein distance between measures μ and ν is

$$W_{2}^{2}(\mu,\nu) = \inf_{\gamma \in \Gamma(\mu,\nu)} \int \|x - y\|^{2} \,\mathrm{d}\gamma(x,y),$$
(4)

where $\Gamma(\mu, \nu)$ is the set of all couplings between μ and ν . $\mathcal{P}_{2,ac}$ endowed with this metric defines a geodesic metric space. The function that we consider minimizing over Wasserstein is the KL divergence between a variable measure μ and a target measure $\pi \propto \exp(-V)$, which is defined by

$$\mathcal{F}(\mu) = \mathsf{KL}(\mu|\pi) = \int \log\left(\frac{\pi}{\mu}\right) \pi$$
 (5)

This setup is typical in sampling problems where we wish to generate samples from π . The gradient flow associated with the minimization of this functional is [5, Section 4.3]

$$\partial_t \mu_t = \operatorname{div}\left(\mu_t \nabla \ln \frac{\mu_t}{\pi}\right).$$
 (6)

Here, $\nabla \ln \frac{\mu_t}{\pi} = \nabla_{W_2} \mathcal{F}(\mu_t)$ is the Wasserstein gradient of \mathcal{F} .

A typical way of implementing the gradient flow associated with this cost functional is through the Langevin diffusion [59]

$$dX_t = -\nabla V(X_t)dt + \sqrt{2}dB_t.$$
(7)

In particular, one can show that $Law(X_t) = \mu_t$, where μ_t solves (6). Therefore, one can generate samples from π by solving this stochastic differential equation. The Euler-Maruyama discretization of this SDE is known as Langevin Monte Carlo (LMC)

$$X_{k+1} = X_k - h\nabla V(X_t) + \sqrt{2h}\xi_k,$$
(LMC)

where $\xi_k \sim N(0, I)$. One can view this as a certain discretization of (6) using a forward-flow [60].

2.3. Preconditioned Langevin Methods

We can define a preconditioned version of the Wasserstein gradient flow (6) as

$$\partial_t \mu_t = \operatorname{div}\left(\mu_t \boldsymbol{A} \nabla \ln \frac{\mu_t}{\pi}\right) \tag{8}$$

for a full rank matrix $A \succ 0$. A more general version of this was studied by [61], and more recently, preconditioned methods have been shown to be more efficient than standard LMC [11, 62].

It turns out that this is a specific example of an important more general class of Langevin diffusions that are called mirror Langevin Diffusions [7, 8, 63]. One can view preconditioned Langevin diffusion as a mirror Langevin diffusion with a linear mirror map. In particular, if one defines $\phi(x) = \frac{1}{2}x^T A^{-1}x$, one defines the corresponding mirror Langevin diffusion as

$$dX_t = -\mathbf{A}\nabla V(X_t)dt + \sqrt{2}\mathbf{A}^{1/2}dB_t.$$
 (PLD)

We note that this is also a Wasserstein gradient flow with respect to the modified metric $W_{2,A^{-1}}$, which is W_2 as defined in (4) with the norm replaced by $\|\cdot\|_{A^{-1}}$. The discretization of this is Preconditioned Langevin Monte Carlo (PLMC),

$$X_{k+1} = X_k - h\mathbf{A}\nabla V(X_t) + \sqrt{2h}\mathbf{A}^{1/2}\xi_k.$$
 (PLMC)

The diffusion (PLMC) has π as a stationary distribution because it is a special case of Theorem 1 of Ma et al. [61].

Theorem 1 (Theorem 1 of Ma et al. [61]). *If* $A \succeq 0$ *then,* (PLMC) *has* $\pi(-V)$ *as a stationary distribution.*

2.4. Subspace Langevin Monte Carlo

In the previous section, we discussed PLMC, which generalizes LMC and is a special case of mirror LMC. A disadvantage of PLMC is that at each iteration, we must multiply a $d \times d$ positive definite matrix A by the gradient, which makes the per-iteration complexity and memory significant in high dimensions.

To make this method more efficient, in analogy to the subspace gradient flow and subspace gradient descent in Euclidean space, we consider a subspace version in Wasserstein space. For a sequence of low-rank matrices $(\mathbf{P}_t)_{t \in [0,\infty)}$, we define the subspace Wasserstein gradient flow of the KL divergence as

$$\partial_t \mu_t = \operatorname{div}\left(\mu_t \boldsymbol{P}_t \nabla \ln \frac{\mu_t}{\pi}\right). \tag{9}$$

For certain choices of P_t , we can view this as a low-rank approximation to the full gradient flow (PLD). We do not study this object further here and leave its properties for future work.

For a sequence of low-rank matrices P_k , the discrete-time process is

$$X_{k+1} = X_k - h_k \boldsymbol{P}_k \nabla V(X_k) + \sqrt{2h_k \boldsymbol{P}_k^{1/2}} \boldsymbol{\xi}_k.$$
(SLMC)

which we call the *Subspace Langevin Monte Carlo* algorithm (SLMC). A specific case of SLMC is RCLMC [22], which happens when P_k is the projection onto a random coordinate. Another specific case of SLMC is Block-Coordinate Langevin Monte Carlo (BCLMC), where P_k is a projection onto a block of coordinates. This method has not yet been studied and is a special case of our later analysis.

3. Theoretical Analysis of Subspace Langevin Monte Carlo

In this section, we conduct a discrete-time analysis of (SLMC) with the goal of developing a comparison between the complexity of LMC, PLMC, and SLMC. Our analysis of SLMC is inspired by Ding et al. [22], with an added twist that we use relative notions of strong convexity and smoothness that are a special case of those in [63]. We make the following assumption throughout.

Assumption 1. The potential V is m relatively strongly convex and M relatively smooth with respect to $\|\cdot\|_{B'}$

$$V(x) + \langle \nabla V(x), y - x \rangle + \frac{m}{2} \|y - x\|_{B}^{2} \le V(y) \le V(x) + \langle \nabla V(x), y - x \rangle + \frac{M}{2} \|y - x\|_{B}^{2}.$$

When *V* is twice differentiable, this is equivalent to the condition $m\mathbf{B} \preceq \nabla^2 V \preceq M\mathbf{B}$. We note that these are special cases of the conditions discussed in [64], where the reference function is $\|\cdot\|_{\mathbf{B}}^2$.

We note that the primary goal of this paper is to have a clear picture of the complexity of the discussed methods. We choose to measure the complexity in terms of calls to a directional derivative oracle. Therefore, for example, computing the gradient of a function $f : \mathbb{R}^d \to \mathbb{R}$ requires d calls to this oracle while computing the derivative along r directions requires r calls to this oracle.

In Section 3.1, we recall the state-of-the-art bounds for LMC under the assumptions of strong logconcavity and smoothness. Then, in Section 3.2, we specialize the result of [63] to give a complexity bound for PLMC. In Section 3.3, we present our convergence result for SLMC, and we finish with a discussion of all of these results in Section 3.4.

3.1. Convergence of LMC

We say that the potential *V* is α -strongly convex and β -smooth, which means Assumption 1 holds with $\alpha = m$, $\beta = M$, and B = I. State-of-the-art bounds for the mixing of LMC in this setting can be found in Durmus et al. [65], where the complexity required to achieve ϵ error in 2-Wasserstein distance is

$$N_{\rm LMC} = O\left(\frac{d^2\beta}{\epsilon^2 \alpha^2} \log \frac{W_2(\mu_0, \pi)}{\epsilon}\right).$$
(10)

Here, β/α is the condition number, and the extra factor of *d* over the result in Durmus et al. [65] comes from the fact that one must compute *d*-directional derivatives at each iteration of the method.

3.2. Convergence of PLMC

To prove convergence results for PLMC, we can follow the work of Ahn and Chewi [63], which proves convergence of mirror LMC under the assumptions of relative strong convexity and smoothness. Here, we now assume that Assumption 1 holds with $B = A^{-1}$. The relative strong convexity allows us to prove exponential convergence of (PLD), which is given in Lemma 1 in the Appendix.

We have the following theorem on the convergence of the PLMC algorithm. This is a special case of Theorem 2 of [63]. We include a simplified proof of this theorem in Appendix A.2.

Theorem 2 (Ahn and Chewi [63]). Suppose that Assumption 1 holds with $B = A^{-1}$. Then PLMC achieves the convergence bound

$$W_{2,\mathbf{A}^{-1}}(\mu_{Nh},\pi) \le (1-mh)^{N/2} W_{2,\mathbf{A}^{-1}}(\mu_0,\pi) + O(\sqrt{\frac{M}{m}}dh).$$
(11)

It is possible to go from $W_{2,A^{-1}}$ bounds to W_2 bounds at the cost of a factor $||A||_2$. If one precomputes the eigenvalue decomposition of A to compute directional derivatives along the eigenspace at each iteration, then we find the iteration complexity to achieve ϵ error to be

$$N_{\text{PLMC}} = O\left(\frac{d^2 M \|\boldsymbol{A}\|_2}{\epsilon^2 m^2} \log \frac{W_{2,\boldsymbol{A}^{-1}}(\mu_0,\pi) \|\boldsymbol{A}\|_2}{\epsilon}\right).$$
(12)

If we did not use the eigenvalue decomposition of *A*, then there would be an additional factor of *d*.

3.3. Convergence of SLMC

We now present our main theoretical result on the convergence of SLMC. It relies on the following assumptions. First, we assume that the projections P_k are generated in the following way.

Assumption 2. At iteration k, we sample P_k in the following manner. We assume that the preconditioner A has eigenvalue decomposition WDW^T that is divided into d/r blocks W_i , for i = 1, ..., d/r, with corresponding $r \times r$ eigenvalue blocks D_i . We then set $P_k = W_i D_i W_i^T$ and $h_k = h/\phi_i$ with probability ϕ_i . We thus see that $hA = \mathbb{E}h_k P_k$.

We note that this procedure is essentially preconditioned block coordinate descent after applying a rotation W^T . In the case of W = D = I and r = 1, this is just the procedure in RCLMC [22].

Assumption 3. The function V has directional smoothness properties. Along any linear subspace spanned by $U \in O(d, r)$, V is smooth with parameter M(U) if for all $y = x + UU^T \delta$,

$$\|\boldsymbol{U}\boldsymbol{U}^{T}(\nabla V(x) - \nabla V(y))\|_{\boldsymbol{A}} \le M(\boldsymbol{U})\|\boldsymbol{U}\boldsymbol{U}^{T}(x-y)\|_{\boldsymbol{A}^{-1}}.$$

For the blocks $W_1, \ldots, W_{d/r}$, we let $M_i = M(W_i)$, and we note that when Assumption 1 holds $M_i \leq M$. **Remark 1.** A more general version of our theory holds when one also changes A at each iteration. For simplicity, we study the case of fixed A here and leave the extended discussion of time-varying A to future work.

Theorem 3. Under Assumptions 1-3, where we use $B = A^{-1}$, and $h \leq \min \phi_i / M$, we have

$$W_{2,\mathbf{A}^{-1}}(\mu_N,\pi) \lesssim \exp(-\frac{hmN}{4})W_{2,\mathbf{A}^{-1}}(\mu_0,\pi) + \frac{\sqrt{rh}}{m}\sqrt{\sum_i \frac{M_i^2}{\phi_i}}.$$

The proof of this theorem is given in Appendix A.3 and is similar to that of the main theorem of Ding et al. [22] with a few twists. The proof follows a Wasserstein coupling style argument developed by Dalalyan and Karagulyan [66].

Again, we can go from $W_{2,A^{-1}}$ bounds to W_2 bounds at the cost of a factor $||A||_2$. if we take $h \simeq \frac{\epsilon^2}{r\sum_{i} \kappa_i^2/\phi_i}$ in Theorem 3, we see that the complexity of (SLMC) is

$$N_{\text{SLMC}} = O\left(\frac{r^2 \|\boldsymbol{A}\|_2 \sum_{i=1}^{d/r} \kappa_i^2 / \phi_i}{\epsilon^2 m} \log \frac{W_2(\mu_0, \pi) \|\boldsymbol{A}\|_2}{\epsilon}\right)$$
(13)

Note that there is no extra factor of d since we assume that the eigenvalue decomposition of A is precomputed. We discuss and further compare all of these results in the next section. In this case, the method reduces to block-coordinate descent in the eigenspace of A.

3.4. Discussion

Here, we present a theoretical comparison of the various methods. At a high level, due to the preconditioning, PLMC and SLMC depend on a relaxation of the strong convexity condition to a relatively strong monotonicity, which allows more adaptation than existing results for LMC and RCLMC.

We now illustrate some simple cases when Assumptions 1 is satisfied. Consider a first example of $D_i = hI$, $\phi_i = r/d$. In this case, $D_k = rh/dI$, h = rh/d, and $h_i = h$. We have that the Assumption 1 holds when V is m-strongly convex and M-smooth. Now suppose $V(x) = \frac{1}{2}x^T \Sigma^{-1}x$ – the case of a centered Gaussian distribution. Suppose that we take W_k and D_k such that $A = \Sigma$. Then, since V is twice, differentiable, we see that Assumption 1 trivially holds with m = M = 1, and the bound (13) no longer depends on the conditioning of Σ^{-1} !

While we can think of our method as forming a subspace approximation to Preconditioned Langevin [11, 62], our analysis is more comparable to existing bounds for LMC. This allows us to directly see how preconditioning can adapt to the problem at hand.

In order to compare complexities of these four methodologies, we also must define the result for RCLMC [22]. Let β_i denote the smoothness in the *i*th coordinate direction, as in [22], and suppose that *V* is α -strongly convex. Denote $\tilde{\kappa}_i = \beta_i / \alpha$ as the condition number along the *i*th direction with respect to the strong convexity parameter α , and $\tilde{\kappa} = \beta / \alpha$. The complexity of RCLMC, in this case, is

$$N_{\mathsf{RCLMC}} = O\left(\frac{\sum_{i=1}^{d} \tilde{\kappa}_i^2 / \phi_i}{\epsilon^2 \alpha} \log \frac{W_2(\mu_0, \pi)}{\epsilon}\right)$$

For SLMC, let $\kappa_{rel,i} = M_i/m$ as the condition number with respect to the *i*th block of **A**, and for PLMC, let $\kappa_{rel} = M/m$ be the relative condition number.

The simplest comparison between SLMC and LMC occurs when there is no adaptation, where A = I and we use uniform random sampling, $\phi_i = r/d$. In this case, relative strong convexity and smoothness reduce to normal strong convexity and smoothness, and so $m = \alpha$ and $M = \beta$. If we then further use the worst case bound to set $\kappa_{\text{rel},i} = \kappa$, then we see that SLMC is worse than LMC by a factor of κ . However, even in this worst-case situation, we note that SLMC still has the benefit of *lower memory usage* than LMC. On the other hand, if we keep our relative constants, which we expect to potentially be better than those of LMC in general, and follow the strategy in [22] and take $\phi_i = \kappa_{\text{rel},i} / \sum_j \kappa_{\text{rel},j}$, then we find that SLMC outperforms LMC once $r^2 \frac{(\sum_{i=1}^{d/r} \kappa_{\text{rel},i})^2}{m} \leq \frac{d^2 \kappa}{\alpha}$. When the function is highly skewed and A is well-adapted to it ($\kappa_{\text{rel},i} \ll \kappa$ and $m \gg \alpha$), we see that there are definitive gains in performance for (SLMC) on top of the lower memory usage.

We finish this section by summarizing the complexity results in Table 1.

	LMC	PLMC	RCLMC	SLMC
Relative or standard conditioning	standard	relative	standard	relative
Non-random	$\tilde{O}(\frac{d^2\tilde{\kappa}}{\epsilon^2\alpha})$	$\tilde{O}(\frac{d^2\kappa_{rel}}{\epsilon^2m})$	-	-
$\phi_i = r/d$	-	-	$\tilde{O}(\frac{d\sum_{i=1}^d \kappa_i^2}{\epsilon^2 \alpha})$	$\tilde{O}(\frac{dr\sum_{i=1}^{d/r}\kappa_{rel,i}^2}{\epsilon^2m})$
$\phi_i = M_i / \sum_i M_i$	-	-	$\tilde{O}(\frac{d(\sum_{i=1}^d \tilde{\kappa}_i)^2}{\epsilon^2 \alpha})$	$\tilde{O}\left(\frac{dr(\sum_{i=1}^{d/r}\kappa_{rel,i})^2}{\epsilon^2 m}\right)$

Table 1: Table comparing complexity bounds for LMC, PLMC, RCLMC, and SLMC measured in terms of total number of directional derivative computations. Our analysis of SLMC is strictly more general than that of RCLMC and allows for a method that can be much more efficient than vanilla LMC. The \tilde{O} removes constants and dependence log factors, such as $\log W_2(\mu_0, \pi)/\epsilon$.

4. Experiments

The experiments in this section are meant to demonstrate the range of potential SLMC methods that can be used for sampling on a specific example of sampling from an ill-conditioned Gaussian distribution. The experiment is inspired by Ding and Li [23]. In these experiments, we will compare different variants of SLMC with LMC, PLMC, and RCLMC that demonstrate its convergence. We set up SLMC to be a block coordinate method, where for a given basis \boldsymbol{W} , the blocks are defined by a sequence of columns $\boldsymbol{W}_i = (w_{r(i-1)+1}, \dots, w_{r(i+1)})$. We try to sample from a Gaussian distribution in dimension d = 20. We let the target distribution have potential $V(x) = \frac{1}{2}x^T \Sigma^{-1}x$, where Σ^{-1} is formed as follows. For a random 5×5 Gaussian matrix \boldsymbol{G} and a 10×10 random orthogonal matrix \boldsymbol{U} . The the lower right 10×10 block of Σ^{-1} is \boldsymbol{I} and the upper left 10×10 block is $\boldsymbol{U} \begin{pmatrix} (\boldsymbol{G} + 10\boldsymbol{I}_5)(\boldsymbol{G} + 10\boldsymbol{I}_5)^T & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{I}_5 \end{pmatrix} \boldsymbol{U}^T$. In the following, we define a test function $\phi(x) = |\mathbf{1}^T x|$, and compute the error as $\operatorname{Err} = |\frac{1}{N} \sum_i \phi(X_i) - \mathbb{E}_{\pi} \phi|$. The initial to be $N(1, \boldsymbol{I})$ and for each method we run N = 20000 particles.

In Figure 1 we give two experiments comparing the use of no preconditioning and preconditioning in the application of SLMC. As we can see, in the case of no preconditioning, and W = D = I SLMC is still able to match the performance of LMC when the block size is adapted to the block structure of the covariance Σ since it is a 10×10 block in the upper left. In the second experiment, we see that preconditioning can enable fast initial convergence, especially when the block size is also adapted to the covariance structure.

In Figure 2, we run an experiment to demonstrate how the choice of basis can assist SLMC in converging faster. These correspond to cases where the relative constants end up being better for SLMC, and so it converges faster than LMC and RCLMC. Depending on the choice of basis W, we see that the best for SLMC is when the selected blocks adapt to a block structure in $W^T \Sigma^{-1} W$. That is why the rotated version of SLMC performs better in the right plot of Figure 2 since the rotation undoes the rotation by U to create a matrix with a 5×5 upper left block.



Figure 1: Experiments demonstrating the convergence of SLMC and PLMC compared to LMC and RCLMC for a diagonal preconditioner. The step size in both experiments is h = 0.01. Left: No preconditioning. As we can see, setting the dimension equal to the size of the upper left block allows SLMC to converge as fast as LMC. Right: SLMC and PLMC have a diagonal preconditioner that is I on the upper left 10×10 block and 10I on the lower 10×10 block. As we can see, using a nonuniform step size allows for faster initial convergence since it is adapted to the covariance structure. This step size incurs a larger final bias.



Figure 2: Figure experiments where SLMC uses a different basis. Left: SLMC blocks are taken from the eigenvalue decomposition of the covariance. The step size for SLMC is considered to be larger at h = 0.5, while for LMC and RCLMC, it is 0.01. We see that the adaptation allows SLMC to rapidly converge at the onset while having a larger bias due to a larger effective step size. Right: SLMC now uses blocks from the rotation such that the top left block is 5×5 . As we can see, now the SLMC method with r = 5 can adapt to the blocks and converges faster than r = 10, as we saw in Figure 1.

5. Conclusion

We have presented Subspace Langevin Monte Carlo, a version of Langevin Monte Carlo that updates along low-dimensional subspaces. This method generalizes Random Coordinate Langevin Monte Carlo as well as its block coordinate analog, as well as preconditioned Langevin [62]. Our theoretical results show that through proper adaptation of the selected subspaces and preconditioner, the SLMC method can significantly outperform LMC in terms of overall complexity at a reduced memory cost. Our experiments demonstrate this distinction in practice.

We believe that our work opens many interesting future avenues for research. First, there are many interesting theoretical extensions of SLMC that one can consider. For example, it would be interesting to extend the theory to more general settings, such as convergence under log-Sobolev and Poincaré inequalities. It would also be interesting to develop subspace versions of other variances of LMC, such as Underdamped Langevin Monte Carlo. Finally, one can feasibly create preconditioners that adapt to the problem at hand as the method iterates and also study time-varying preconditioners. Another interesting avenue of future research involves the prospect of SLMC as subspace descent in Wasserstein space. It would be interesting to consider subspace descent methods over Wasserstein gradient flows [55]. Finally, it would be interesting to incorporate these ideas into the training of score-based diffusion models to further the ideas in put forth in [25] with more principled ways of choosing subspaces. This would open up broader applications for the methods and theory developed in this paper.

References

- [1] Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. *Advances in neural information processing systems*, 33:6840–6851, 2020.
- [2] Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations*, 2021.
- [3] Alain Durmus and Eric Moulines. High-dimensional bayesian inference via the unadjusted langevin algorithm. *Bernoulli*, 25(4A):2854–2882, 2019.
- [4] Rishav Chourasia, Jiayuan Ye, and Reza Shokri. Differential privacy dynamics of langevin diffusion and noisy gradient descent. *Advances in Neural Information Processing Systems*, 34: 14771–14781, 2021.
- [5] Filippo Santambrogio. {Euclidean, metric, and Wasserstein} gradient flows: an overview. *Bulletin of Mathematical Sciences*, 7:87–154, 2017.
- [6] Ya-Ping Hsieh, Ali Kavis, Paul Rolland, and Volkan Cevher. Mirrored langevin dynamics. *Advances in Neural Information Processing Systems*, 31:2878–2887, 2018.
- [7] Kelvin Shuangjian Zhang, Gabriel Peyré, Jalal Fadili, and Marcelo Pereyra. Wasserstein control of mirror langevin monte carlo. In *Conference on learning theory*, pages 3814–3841. PMLR, 2020.
- [8] Sinho Chewi, Thibaut Le Gouic, Chen Lu, Tyler Maunu, Philippe Rigollet, and Austin Stromme. Exponential ergodicity of mirror-Langevin diffusions. *Advances in Neural Information Processing Systems*, 33, 2020.
- [9] Yifei Wang and Wuchen Li. Information newton's flow: second-order optimization method in probability space. *arXiv preprint arXiv:2001.04341*, 2020.
- [10] Yongxin Chen, Sinho Chewi, Adil Salim, and Andre Wibisono. Improved analysis for a proximal algorithm for sampling. In *Conference on Learning Theory*, pages 2984–3014. PMLR, 2022.
- [11] Michalis Titsias. Optimal preconditioning and fisher adaptive langevin sampling. *Advances in Neural Information Processing Systems*, 36, 2024.
- [12] Clément Bonet, Théo Uscidda, Adam David, Pierre-Cyril Aubin-Frankowski, and Anna Korba. Mirror and preconditioned gradient descent in wasserstein space. *arXiv preprint arXiv:*2406.08938, 2024.
- [13] Yu Nesterov. Efficiency of coordinate descent methods on huge-scale optimization problems. *SIAM Journal on Optimization*, 22(2):341–362, 2012.
- [14] Robert Gower, Dmitry Kovalev, Felix Lieder, and Peter Richtárik. Rsn: randomized subspace newton. Advances in Neural Information Processing Systems, 32, 2019.
- [15] Filip Hanzely, Nikita Doikov, Yurii Nesterov, and Peter Richtarik. Stochastic subspace cubic newton method. In *International Conference on Machine Learning*, pages 4027–4038. PMLR, 2020.
- [16] David Kozak, Stephen Becker, Alireza Doostan, and Luis Tenorio. A stochastic subspace approach to gradient-free optimization in high dimensions. *Computational Optimization and Applications*, 79(2):339–368, 2021.
- [17] Romain Cosson, Ali Jadbabaie, Anuran Makur, Amirhossein Reisizadeh, and Devavrat Shah. Low-rank gradient descent. *IEEE Open Journal of Control Systems*, 2023.
- [18] Vladimir Feinberg, Xinyi Chen, Y Jennifer Sun, Rohan Anil, and Elad Hazan. Sketchy: Memory-efficient adaptive regularization with frequent directions. *Advances in Neural Information Processing Systems*, 36, 2024.

- [19] Jiawei Zhao, Zhenyu Zhang, Beidi Chen, Zhangyang Wang, Anima Anandkumar, and Yuandong Tian. Galore: Memory-efficient llm training by gradient low-rank projection. arXiv preprint arXiv:2403.03507, 2024.
- [20] Kaizhao Liang, Bo Liu, Lizhang Chen, and Qiang Liu. Memory-efficient llm training with online subspace descent. *arXiv preprint arXiv:2408.12857*, 2024.
- [21] Romain Cosson, Ali Jadbabaie, Anuran Makur, Amirhossein Reisizadeh, and Devavrat Shah. Gradient descent with low-rank objective functions. In 2023 62nd IEEE Conference on Decision and Control (CDC), pages 3309–3314. IEEE, 2023.
- [22] Zhiyan Ding, Qin Li, Jianfeng Lu, and Stephen J Wright. Random coordinate langevin monte carlo. In *Conference on learning theory*, pages 1683–1710. PMLR, 2021.
- [23] Zhiyan Ding and Qin Li. Langevin monte carlo: random coordinate descent and variance reduction. *Journal of machine learning research*, 22(205):1–51, 2021.
- [24] Abhishek Roy, Lingqing Shen, Krishnakumar Balasubramanian, and Saeed Ghadimi. Stochastic zeroth-order discretizations of langevin diffusions for bayesian inference. *Bernoulli*, 28(3): 1810–1834, 2022.
- [25] Bowen Jing, Gabriele Corso, Renato Berlinghieri, and Tommi Jaakkola. Subspace diffusion generative models. In European Conference on Computer Vision, pages 274–289. Springer, 2022.
- [26] Peng Wang, Huijie Zhang, Zekai Zhang, Siyi Chen, Yi Ma, and Qing Qu. Diffusion models learn low-dimensional distributions via subspace clustering. arXiv preprint arXiv:2409.02426, 2024.
- [27] Siyi Chen, Huijie Zhang, Minzhe Guo, Yifu Lu, Peng Wang, and Qing Qu. Exploring lowdimensional subspaces in diffusion models for controllable image editing. arXiv preprint arXiv:2409.02374, 2024.
- [28] William B Johnson, Joram Lindenstrauss, and Gideon Schechtman. Extensions of lipschitz maps into banach spaces. *Israel Journal of Mathematics*, 54(2):129–138, 1986.
- [29] Moses Charikar, Kevin Chen, and Martin Farach-Colton. Finding frequent items in data streams. *Theoretical Computer Science*, 312(1):3–15, 2004.
- [30] Anirban Dasgupta, Ravi Kumar, and Tamás Sarlós. A sparse johnson: Lindenstrauss transform. In *Proceedings of the forty-second ACM symposium on Theory of computing*, pages 341–350, 2010.
- [31] Nir Ailon and Bernard Chazelle. The fast johnson–lindenstrauss transform and approximate nearest neighbors. *SIAM Journal on computing*, 39(1):302–322, 2009.
- [32] Joel A Tropp. Improved analysis of the subsampled randomized hadamard transform. *Advances in Adaptive Data Analysis*, 3(01n02):115–126, 2011.
- [33] Nathan Halko, Per-Gunnar Martinsson, and Joel A Tropp. Finding structure with randomness: Probabilistic algorithms for constructing approximate matrix decompositions. *SIAM review*, 53(2):217–288, 2011.
- [34] Stephen J Wright. Coordinate descent algorithms. *Mathematical programming*, 151(1):3–34, 2015.
- [35] Robert M Gower and Peter Richtárik. Randomized iterative methods for linear systems. SIAM Journal on Matrix Analysis and Applications, 36(4):1660–1690, 2015.
- [36] Robert Mansel Gower and Peter Richtárik. Stochastic dual ascent for solving linear systems. *arXiv preprint arXiv:1512.06890*, 2015.

- [37] Rafael Frongillo and Mark D Reid. Convergence analysis of prediction markets via randomized subspace descent. *Advances in Neural Information Processing Systems*, 28, 2015.
- [38] Yurii Nesterov and Vladimir Spokoiny. Random gradient-free minimization of convex functions. *Foundations of Computational Mathematics*, 17(2):527–566, 2017.
- [39] Alp Yurtsever, Joel A Tropp, Olivier Fercoq, Madeleine Udell, and Volkan Cevher. Scalable semidefinite programming. SIAM Journal on Mathematics of Data Science, 3(1):171–200, 2021.
- [40] Nikita Ivkin, Daniel Rothchild, Enayat Ullah, Ion Stoica, Raman Arora, et al. Communicationefficient distributed sgd with sketching. *Advances in Neural Information Processing Systems*, 32, 2019.
- [41] Filip Hanzely, Konstantin Mishchenko, and Peter Richtárik. Sega: Variance reduction via gradient sketching. Advances in Neural Information Processing Systems, 31, 2018.
- [42] Robert M Gower, Peter Richtárik, and Francis Bach. Stochastic quasi-gradient methods: Variance reduction via jacobian sketching. *Mathematical Programming*, 188(1):135–192, 2021.
- [43] Levent Sagun, Leon Bottou, and Yann LeCun. Eigenvalues of the hessian in deep learning: Singularity and beyond. *arXiv preprint arXiv:1611.07476*, 2016.
- [44] Levent Sagun, Utku Evci, V Ugur Guney, Yann Dauphin, and Leon Bottou. Empirical analysis of the hessian of over-parametrized neural networks. *arXiv preprint arXiv:1706.04454*, 2017.
- [45] Stanisław Jastrzębski, Zachary Kenton, Nicolas Ballas, Asja Fischer, Yoshua Bengio, and Amos Storkey. On the relation between the sharpest directions of dnn loss and the sgd step length. In *International Conference on Learning Representations*, 2019.
- [46] Gerard Ben Arous, Reza Gheissari, Jiaoyang Huang, and Aukosh Jagannath. Highdimensional sgd aligns with emerging outlier eigenspaces. In *The Twelfth International Conference on Learning Representations*, 2024.
- [47] Minhak Song, Kwangjun Ahn, and Chulhee Yun. Does sgd really happen in tiny subspaces? *arXiv preprint arXiv*:2405.16002, 2024.
- [48] Chunyuan Li, Heerad Farkhoor, Rosanne Liu, and Jason Yosinski. Measuring the intrinsic dimension of objective landscapes. In *International Conference on Learning Representations*, 2018.
- [49] Frithjof Gressmann, Zach Eaton-Rosen, and Carlo Luschi. Improving neural network training in low dimensional random bases. *Advances in Neural Information Processing Systems*, 33:12140– 12150, 2020.
- [50] Tao Li, Lei Tan, Zhehao Huang, Qinghua Tao, Yipeng Liu, and Xiaolin Huang. Low dimensional trajectory hypothesis is true: Dnns can be trained in tiny subspaces. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 45(3):3411–3420, 2022.
- [51] Arash Vahdat, Karsten Kreis, and Jan Kautz. Score-based generative modeling in latent space. *Advances in neural information processing systems*, 34:11287–11302, 2021.
- [52] Robin Rombach, Andreas Blattmann, Dominik Lorenz, Patrick Esser, and Björn Ommer. Highresolution image synthesis with latent diffusion models. In *Proceedings of the IEEE/CVF conference on computer vision and pattern recognition*, pages 10684–10695, 2022.
- [53] Belinda Tzen and Maxim Raginsky. Theoretical guarantees for sampling and inference in generative models with latent diffusions. In *Conference on Learning Theory*, pages 3084–3114. PMLR, 2019.
- [54] Nicolas Bonneel, Julien Rabin, Gabriel Peyré, and Hanspeter Pfister. Sliced and radon wasserstein barycenters of measures. *Journal of Mathematical Imaging and Vision*, 51:22–45, 2015.

- [55] Clément Bonet, Nicolas Courty, François Septier, and Lucas Drumetz. Efficient gradient flows in sliced-wasserstein space. *Transactions on Machine Learning Research*, 2022. ISSN 2835-8856. URL https://openreview.net/forum?id=Au1LNKmRvh.
- [56] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. *Gradient flows: in metric spaces and in the space of probability measures.* Springer Science & Business Media, 2008.
- [57] Cédric Villani. Optimal transport: old and new, volume 338. Springer, 2009.
- [58] Sinho Chewi. Log-concave sampling. Book draft available at https://chewisinho. github. io, 2024.
- [59] Richard Jordan, David Kinderlehrer, and Felix Otto. The variational formulation of the fokker– planck equation. *SIAM journal on mathematical analysis*, 29(1):1–17, 1998.
- [60] Andre Wibisono. Sampling as optimization in the space of measures: The langevin dynamics as a composite optimization problem. In *Conference on Learning Theory*, pages 2093–3027. PMLR, 2018.
- [61] Yi-An Ma, Tianqi Chen, and Emily Fox. A complete recipe for stochastic gradient mcmc. *Advances in neural information processing systems*, 28, 2015.
- [62] Riddhiman Bhattacharya and Tiefeng Jiang. Fast sampling and inference via preconditioned langevin dynamics. *arXiv preprint arXiv:2310.07542*, 2023.
- [63] Kwangjun Ahn and Sinho Chewi. Efficient constrained sampling via the mirror-langevin algorithm. *arXiv preprint arXiv:2010.16212*, 2020.
- [64] Haihao Lu, Robert M Freund, and Yurii Nesterov. Relatively smooth convex optimization by first-order methods, and applications. *SIAM Journal on Optimization*, 28(1):333–354, 2018.
- [65] Alain Durmus, Szymon Majewski, and Błażej Miasojedow. Analysis of langevin monte carlo via convex optimization. *Journal of Machine Learning Research*, 20(73):1–46, 2019.
- [66] Arnak S Dalalyan and Avetik Karagulyan. User-friendly guarantees for the langevin monte carlo with inaccurate gradient. *Stochastic Processes and their Applications*, 129(12):5278–5311, 2019.

A. Supplementary Proofs

A.1. Contraction Lemma for (PLD)

Lemma 1. Let Z_t and Z'_t be two copies of the diffusion (PLD) with the same Brownian motion. Assume that *V* and *A* satisfy Assumption 1. Then, we have the following contraction.

$$||Z_t - Z'_t||^2_{\mathbf{A}^{-1}} \le \exp(-mt)\mathbb{E}||Z_0 - Z'_0||^2_{\mathbf{A}^{-1}}.$$

Remark 2. Note that if we optimally couple Z_0 and Z'_0 , this becomes an exponential contraction in the $W_{2,\mathbf{A}^{-1}}$ distance. Since the Wasserstein distance is an expectation, we could get away with a weaker condition than that in Assumption 1 since we only require a notion of relatively strong monotonicity in expectation along the flows Law(Z_t) and Law(Z'_t). An interesting line of future work could study this assumption in more detail.

Proof. Applying Itô's lemma to $f(z, z') = ||z - z'||^2_{A^{-1}}$

$$f(Z_t, Z'_t) - f(Z_0, Z'_0) = 2 \int_0^t \langle Z_s - Z'_s, \nabla V(Z_s) - \nabla V(Z'_s) \rangle ds$$
$$\leq 2 \int_0^t m f(Z_s, Z'_s) ds.$$

Thus we can apply Grönwall's lemma to $f(Z_s, Z'_s)$ to find

$$f(Z_t, Z'_t) \le \exp(-2\int_0^t m ds) f(Z_0, Z'_0).$$

A.2. Proof of Theorem 2

Proof. In the case of a fixed A, this theorem is just a particular case of Theorem 2 in [63]. We replicate the proof here with our simplified setting, which avoids needing to work with abstract Bregman divergences.

The proof technique works by separating the Langevin update into a gradient update followed by the addition of a Gaussian, which corresponds to a step with respect to the potential energy $\mathbb{E}_{\mu}V$ followed by a step with respect to the entropy $\mathbb{E}_{\mu} \ln \mu$. It then proceeds in three steps: bounding the decrease in the potential energy, showing that the entropy step does not increase the potential energy too much, and then showing that the entropy step decreases the entropy.

We first preconditioned gradient descent,

$$x_{k+1} = x_k - h\mathbf{A}\nabla V(x_k).$$

Assuming that *V* is β smooth relative to $\|\cdot\|^2_{A^{-1}}$,

$$V(x_{k+1}) \le V(x_k) + \langle \nabla V(x_k), x_{k+1} - x_k \rangle + \frac{M}{2} \|x_{k+1} - x_k\|_{A^{-1}}^2$$

= $V(x_k) + \left(\frac{Mh^2}{2} - h\right) \|\nabla V(x_k)\|_A^2.$

This gives us a relative descent lemma. For example, if one sets $h \leq 1/M$, we get a descent lemma

$$\|\nabla V(x_k)\|_{\boldsymbol{A}}^2 \le 2hM(V(x_k) \le V(x_{k+1}))$$

Similar to [56, 58], when $h \le 1/M$, we can prove an evolution variational inequality (EVI)

$$\begin{aligned} \|x_{k+1} - y\|_{\mathbf{A}^{-1}}^2 &= \|x_k - y\|_{\mathbf{A}^{-1}}^2 - 2h\langle \nabla V(x_k), x_k - y \rangle + h^2 \|\nabla V(x_k)\|_{\mathbf{A}^{-1}}^2 \\ &\leq (1 - mh) \|x_k - y\|_{\mathbf{A}^{-1}}^2 - 2h(V(x_k) - V(y)) + 2h(V(x_k) - V(x_{k+1})) \\ &\leq (1 - mh) \|x_k - y\|_{\mathbf{A}^{-1}}^2 - 2h(V(x_{k+1}) - V(y)). \end{aligned}$$

Applying this EVI to X_{kh}^+ , X_{kh} and Z, and taking an expectation, we get contraction of the energy with respect to the $W_{2,A^{-1}}$ distance

$$\mathcal{E}(\mu_{kh}^{+}) - \mathcal{E}(\pi) \le \frac{1}{2h} \left[(1 - mh) W_{2,\mathbf{A}^{-1}}^2(\mu_{kh},\pi) - W_{2,\mathbf{A}^{-1}}^2(\mu_{kh}^{+},\pi) \right].$$
(14)

We further have that the energy does not increase too much over the entropy step following the same computation in [65], which uses the *M*-relative smoothness,

$$\mathcal{E}(\mu_{(k+1)h}) - \mathcal{E}(\mu_{kh}^+) = Mdh.$$
(15)

Finally, we have that the entropy \mathcal{H} is relatively convex [63]. In particular, for $X \sim \mu$ and $Y \sim \nu$ optimally coupled in terms of $W_{2,A^{-1}}$, we have

$$\mathcal{H}(\nu) \ge \mathcal{H}(\mu) + \langle \nabla_{W_2} \mathcal{H}(\mu)(X), Y - X \rangle.$$

If we let *X* be the output of a full step and *Y* ~ π , this implies that

$$\mathcal{H}(\pi) \ge \mathcal{H}(\mu_{(k+1)h}) + \langle \nabla_{W_2} \mathcal{H}(\mu_{(k+1)h})(X), Y - X \rangle.$$

Let Q_t denote the semigroup

$$Q_t f(x) = \mathbb{E} f(x + \sqrt{2} \mathbf{A}^{1/2} B_t),$$

so that $\mu_{(k+1)h} = \mu_{kh}^+ Q_h$. We claim that $\mathcal{H}(\mu_{kh}^+ Q_t)$ is the gradient flow of \mathcal{H} with respect to the $W_{2,\mathbf{A}^{-1}}$ geometry, and is therefore nonincreasing. Indeed, the $W_{2,\mathbf{A}^{-1}}$ gradient of \mathcal{H} is

$$\nabla_{W_2} \mathbf{A}^{-1} \mathcal{H}(\mu) = \mathbf{A} \nabla_{W_2} \ln(\mu)$$

And so the $W_{2, \mathbf{A}^{-1}}$ gradient flow of \mathcal{H} is

$$\partial_t \mu_t = \operatorname{div}(\mu_t A \nabla \ln(\mu_t)).$$

This is precisely the same as what is found for the Fokker-Planck equation corresponding to the diffusion $\sqrt{2}A^{1/2}dB_t$. Therefore, for X^+_{kh+t} and Z optimally coupled for the $W_{2,A^{-1}}$ distance, and since \mathcal{H} is decreasing along $\mu^+_{kh}Q_t$,

$$\partial_t W_{2,\mathbf{A}^{-1}}^2(\mu_{kh}^+ Q_t, \pi) \le 2 \langle \nabla_{W_2} \mathcal{H}(\mu_{(k+1)h})(X_{kh+t}^+), Z - X_{kh+t}^+ \rangle \le \mathcal{H}(\pi) - \mathcal{H}(\mu_{(k+1)h}).$$

Integrating from 0 to h yields

$$W_{2,\mathbf{A}^{-1}}^{2}(\mu_{(k+1)h},\pi) - W_{2,\mathbf{A}^{-1}}^{2}(\mu_{kh}^{+},\pi) \le h[\mathcal{H}(\pi) - \mathcal{H}(\mu_{(k+1)h})]$$
(16)

Putting together (14), (15), and (16),

$$W_{2,\mathbf{A}^{-1}}^2(\mu_{(k+1)h},\pi) \le (1-mh)W_{2,\mathbf{A}^{-1}}^2(\mu_{kh},\pi) + 2Mdh^2.$$

Iterating the inequality,

$$W_{2,\mathbf{A}^{-1}}^2(\mu_{Nh},\pi) \le (1-mh)^N W_{2,\mathbf{A}^{-1}}^2(\mu_0,\pi) + \frac{2Mdh}{m}.$$

A.3. Proof of Theorem 3

We note that assumption 1 implies that

$$\|\nabla V(x) - \nabla V(y)\|_{\mathbf{A}}^2 \le m \|y - x\|_{\mathbf{A}^{-1}}^2.$$

We begin with a lemma.

Lemma 2. Suppose that Assumption 1 holds with $B = A^{-1}$ and Assumption 3 holds. If W_i is a block of eigenvectors from A with corresponding relative smoothness parameter M_i , then

$$\mathbb{E}_{\pi} \| \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \nabla V(Z) \|_{\boldsymbol{A}}^{2} \leq M_{i} r$$

Proof. Suppose that we are running (PLD) with the preconditioner $W_i W_i^T A W_i W_i^T = W_i D_i W_i^T$. Let \mathcal{L} be the generator of this process, which takes the form

$$\mathcal{L}f = \operatorname{Tr}(\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\boldsymbol{A}\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\nabla^{2}V) - \langle \boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\boldsymbol{A}\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\nabla V, \nabla f \rangle.$$

Since $\pi \propto \exp(-V)$ is stationary, we have

$$0 = \mathbb{E}_{\pi} \mathcal{L} V = \mathbb{E} \operatorname{Tr}(\boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \boldsymbol{A} \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \nabla^{2} V) - \langle \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \boldsymbol{A} \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \nabla V, \nabla V \rangle.$$

We have

$$\begin{aligned} \operatorname{Tr}(\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\boldsymbol{A}\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\nabla^{2}V) &= \operatorname{Tr}((\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\boldsymbol{A}\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T})^{1/2}\nabla^{2}V(\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\boldsymbol{A}\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T})^{1/2}) \\ &\leq M_{i}\operatorname{Tr}((\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\boldsymbol{A}\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T})^{1/2}\boldsymbol{A}^{-1}(\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\boldsymbol{A}\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T})^{1/2}) \\ &= M_{i}r\end{aligned}$$

We begin now restate our main theorem.

Theorem 4. Under Assumptions 1-3, and $h \leq \min(\min \phi_i/M,)$

$$W_{2,\boldsymbol{A}^{-1}}(\mu_N,\pi) \lesssim \exp(-\frac{hmN}{4})W_{2,\boldsymbol{A}^{-1}}(\mu_0,\pi) + \frac{\sqrt{rh}}{m}\sqrt{\sum_i \frac{M_i^2}{\phi_i}}.$$

Proof. Let $P_k = W_{i_k} D_{i_k} W_{i_k}^T$ be the projection chosen at the *k*th iteration, where $W_{i_k} \in O(d, r)$ and D_{i_k} is diagonal.

We define the auxiliary process

$$Z_{k+1} = Z_k(h) = Z_k(0) - \int_0^{h_i} \mathbf{P}_k \nabla V(Z_k(s)) ds + \sqrt{2h} \mathbf{P}_k^{1/2} \xi_k.$$
 (17)

Here, ξ_k is the same noise as (SLMC). Let $\Delta_k = Z_k - X_k$. We have

$$\Delta_{k+1} = \Delta_k - \mathbf{P}_k \Big[\int_0^{h_i} \nabla V(Z_k(s)) ds - \nabla V(X_k) \Big].$$

After multiplying by U_k^T , we see that really we are looking at a difference in block-coordinate updates between Z_k and X_k in the new basis:

$$\boldsymbol{U}_{k}^{T} \boldsymbol{\Delta}_{k+1} = \boldsymbol{U}_{k}^{T} \boldsymbol{\Delta}_{k} - \boldsymbol{D}_{i} \boldsymbol{U}_{k}^{T} \Big[\int_{0}^{h_{i}} \nabla V(\boldsymbol{Z}_{k}(s)) ds - \nabla V(\boldsymbol{X}_{k}) \Big].$$

By assumption, W is the full basis from which we select W_{i_k} , which is divided into d/r blocks, each of which we select with probability ϕ_i . These blocks are denoted by W_i . We have

$$\mathbb{E} \| \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \Delta_{k+1} \|_{\boldsymbol{A}^{-1}}^{2} = \phi_{i} \mathbb{E} [\| \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \Delta_{k+1} \|_{\boldsymbol{A}^{-1}}^{2} | i_{k} = i] + (1 - \phi_{i}) \mathbb{E} \| \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \Delta_{k} \|_{\boldsymbol{A}^{-1}}^{2}.$$

Going term by term, given $i_k = i$ is chosen,

$$\mathbb{E}[\|\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\boldsymbol{\Delta}_{k+1}\|^{2}] = (1+a)\mathbb{E}\|\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\boldsymbol{\Delta}_{k+1} + \boldsymbol{W}_{i}\boldsymbol{D}_{i}\boldsymbol{W}_{i}^{T}\int_{0}^{h_{i}}\nabla V(Z_{k}(s) - \nabla V(Z_{k})ds\|_{\boldsymbol{A}^{-1}}^{2} + (18))$$

$$\left(1 + \frac{1}{a}\right)\mathbb{E}\|\boldsymbol{W}_{i}\boldsymbol{D}_{i}\boldsymbol{W}_{i}^{T}\int_{0}^{h_{i}}\nabla V(Z_{k}(s) - \nabla V(Z_{k})ds\|_{\boldsymbol{A}^{-1}}^{2}.$$

The first term of this is then bounded by

$$\mathbb{E} \| \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \boldsymbol{\Delta}_{k+1} + \boldsymbol{W}_{i} \boldsymbol{D}_{i} \boldsymbol{W}_{i}^{T} \int_{0}^{h_{i}} \nabla V(Z_{k}(s) - \nabla V(Z_{k}) ds \|^{2} \leq \mathbb{E} \| \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \boldsymbol{\Delta}_{k} \|_{\boldsymbol{A}^{-1}}^{2} - 2h_{i} \mathbb{E} \langle \boldsymbol{W}_{i}^{T} \boldsymbol{\Delta}_{k}, \boldsymbol{D}_{i} \boldsymbol{W}_{i}^{T} (\nabla V(Z_{k}) - \nabla V(X_{k})) \rangle_{\boldsymbol{A}^{-1}} + h_{i}^{2} \mathbb{E} \| \boldsymbol{W}_{i} \boldsymbol{D}_{i} \boldsymbol{W}_{i}^{T} (\nabla V(Z_{k}) - \nabla V(X_{k})) \|_{\boldsymbol{A}^{-1}}^{2}.$$

Again, given that $i_k = i$, following the argument in the proof of Lemma 12 of [22], and using Lemma 2, while being careful with our notion of relative smoothness, the second term of (18) is bounded

$$\begin{split} \mathbb{E} \| \boldsymbol{W}_{i} \boldsymbol{D}_{i} \boldsymbol{W}_{i}^{T} \int_{0}^{h_{i}} \nabla V(Z_{k}(s)) - \nabla V(Z_{k}) \, \mathrm{d}s \|_{\boldsymbol{A}^{-1}}^{2} \\ &\leq h_{i} \int_{0}^{h_{i}} \mathbb{E} \| \boldsymbol{W}_{i} \boldsymbol{D}_{i} \boldsymbol{W}_{i}^{T} (\nabla V(Z_{k}(s)) - \nabla V(Z_{k})) \|_{\boldsymbol{A}^{-1}}^{2} \, \mathrm{d}s \\ &= h_{i} \int_{0}^{h_{i}} \mathbb{E} \| \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} (\nabla V(Z_{k}(s)) - \nabla V(Z_{k})) \|_{\boldsymbol{A}}^{2} \, \mathrm{d}s \\ &\leq h_{i} M_{i}^{2} \int_{0}^{h_{i}} \mathbb{E} \| Z_{k}(s) - Z_{k} \|_{\boldsymbol{A}^{-1}}^{2} \, \mathrm{d}s \\ &= h_{i} M_{i}^{2} \int_{0}^{h_{i}} \mathbb{E} \| \int_{0}^{s} \boldsymbol{P}_{k} \nabla V(Z_{k}(\tau)) \, \mathrm{d}\tau + \sqrt{2} \boldsymbol{P}_{k}^{1/2} \xi_{k} \|_{\boldsymbol{A}^{-1}}^{2} \, \mathrm{d}s \\ &\leq 2h_{i} M_{i}^{2} \int_{0}^{h_{i}} \mathbb{E} \| \int_{0}^{s} \boldsymbol{P}_{k} \nabla V(Z_{k}(\tau)) \, \mathrm{d}\tau \|_{\boldsymbol{A}^{-1}}^{2} \, \mathrm{d}s + 2h_{i} M_{i}^{2} \int_{0}^{h_{i}} \mathbb{E} \| \sqrt{2h_{i}} \boldsymbol{P}_{k}^{1/2} \xi_{k} \|_{\boldsymbol{A}^{-1}}^{2} \, \mathrm{d}s \\ &\leq 2h_{i}^{2} M_{i}^{2} \int_{0}^{h_{i}} \int_{0}^{s} \mathbb{E} \| \boldsymbol{P}_{k} \nabla V(Z_{k}(\tau)) \|_{\boldsymbol{A}^{-1}}^{2} \, \mathrm{d}\tau \, \mathrm{d}s + 4r h_{i}^{3} M_{i}^{2} \\ &\leq h_{i}^{4} M_{i}^{2} \mathbb{E}_{\pi} \| \boldsymbol{W}_{i} \boldsymbol{W}_{i}^{T} \nabla V(Z) \|_{\boldsymbol{A}}^{2} + 4r h_{i}^{3} M_{i}^{2} \\ &\leq h_{i}^{4} M_{i}^{3} r + 4r h_{i}^{3} M_{i}^{2}. \end{split}$$

Here, we use the fact that $P_k^{1/2} A^{-1} P_k^{1/2} = W_i W_i^T$ and $\mathbb{E}_{\pi} || W_i W_i^T \nabla V(Z) ||_A^2 \le M_i r$ by Lemma 2. Using $h_i = h/\phi_i$, we can put these together to yield

$$\begin{split} \mathbb{E} \|\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\Delta_{k+1}\|_{\boldsymbol{A}^{-1}}^{2} &\leq (1+a\phi_{i})\mathbb{E} \|\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\Delta_{k}\|_{\boldsymbol{A}^{-1}}^{2} \\ &\quad -2(1+a)h\mathbb{E}\langle\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\Delta_{k},\boldsymbol{W}_{i}\boldsymbol{D}_{i}\boldsymbol{W}_{i}^{T}(\nabla V(Z_{k})-\nabla V(X_{k}))\rangle_{\boldsymbol{A}^{-1}} \\ &\quad +(1+a)\frac{h^{2}}{\phi_{i}}\mathbb{E} \|\boldsymbol{W}_{i}\boldsymbol{D}_{i}\boldsymbol{W}_{i}^{T}(\nabla V(Z_{k})-\nabla V(X_{k}))\|_{\boldsymbol{A}^{-1}}^{2} \\ &\quad +\left(1+\frac{1}{a}\right)\phi_{i}[\frac{rh^{4}M_{i}^{3}r}{\phi_{i}^{4}}+\frac{4rh^{3}M_{i}^{2}}{\phi_{i}^{3}}] \end{split}$$

Letting $a = hm/\phi_i$, $1 + 1/a \lesssim \phi_i/hm$, $h \le \min \phi_i/M$,

$$\begin{split} \mathbb{E} \|\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\Delta_{k+1}\|_{\boldsymbol{A}^{-1}}^{2} &\leq (1+hm)\mathbb{E} \|\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\Delta_{k}\|_{\boldsymbol{A}^{-1}}^{2} \\ &\quad -2(1+hm/\phi_{i})h\mathbb{E}\langle\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\Delta_{k},\boldsymbol{W}_{i}\boldsymbol{D}_{i}\boldsymbol{W}_{i}^{T}(\nabla V(Z_{k})-\nabla V(X_{k}))\rangle_{\boldsymbol{A}^{-1}} \\ &\quad +(1+hm/\phi_{i})\frac{h^{2}}{\phi_{i}}\mathbb{E} \|\boldsymbol{W}_{i}\boldsymbol{D}_{i}\boldsymbol{W}_{i}^{T}(\nabla V(Z_{k})-\nabla V(X_{k}))\|_{\boldsymbol{A}^{-1}}^{2} \\ &\quad +\frac{1}{m}[\frac{rh^{3}M_{i}^{3}}{\phi_{i}^{2}}+\frac{4rh^{2}M_{i}^{2}}{\phi_{i}}] \\ &\leq (1+hm)\mathbb{E} \|\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\Delta_{k}\|_{\boldsymbol{A}^{-1}}^{2} \\ &\quad -2\mathbb{E}\langle\boldsymbol{W}_{i}\boldsymbol{W}_{i}^{T}\Delta_{k},\boldsymbol{W}_{i}\boldsymbol{D}_{i}\boldsymbol{W}_{i}^{T}(\nabla V(Z_{k})-\nabla V(X_{k}))\rangle_{\boldsymbol{A}^{-1}} \\ &\quad +\frac{2}{\phi_{i}}\mathbb{E} \|\boldsymbol{W}_{i}\boldsymbol{D}_{i}\boldsymbol{W}_{i}^{T}(\nabla V(Z_{k})-\nabla V(X_{k}))\|_{\boldsymbol{A}^{-1}}^{2} \\ &\quad +\frac{1}{m}[\frac{rh^{3}M_{i}^{3}}{\phi_{i}^{2}}+\frac{4rh^{2}M_{i}^{2}}{\phi_{i}}] \end{split}$$

Summing, using $\boldsymbol{A} = \sum_{i} \boldsymbol{W}_{i} \boldsymbol{D}_{i} \boldsymbol{W}_{i}^{T}$,

$$\mathbb{E} \|\Delta_{k+1}\|_{\boldsymbol{A}^{-1}}^2 \leq (1+hm)\mathbb{E} \|\Delta_k\|_{\boldsymbol{A}^{-1}}^2$$
$$-2h\mathbb{E} \langle \Delta_k, \nabla V(Z_k) - \nabla V(X_k) \rangle$$
$$+ \frac{2h^2}{\min \phi_i} \mathbb{E} \| (\nabla V(Z_k) - \nabla V(X_k)) \|_{\boldsymbol{A}}^2$$
$$+ \frac{1}{m} \sum_i [\frac{rh^3 M_i^3}{\phi_i^2} + \frac{4rh^2 M_i^2}{\phi_i}]$$

The smoothness in Assumption 1 implies that

$$\|\nabla V(Z_k) - \nabla V(X_k)\|_{\boldsymbol{A}}^2 \le M^2 \|\nabla V(Z_k) - \nabla V(X_k)\|_{\boldsymbol{A}^{-1}}^2$$

Therefore, using the previous display and using strong relative monotonicity implied by Assumption 1,

$$\mathbb{E} \|\Delta_{k+1}\|_{\boldsymbol{A}^{-1}}^2 \le (1 + hm - 2hm + \frac{2}{\min \phi_i} h^2 M^2) \mathbb{E} \|\Delta_k\|_{\boldsymbol{A}^{-1}}^2 \\ + \frac{1}{m} \sum_i \left[\frac{rh^3 M_i^3}{\phi_i^2} + \frac{4rh^2 M_i^2}{\phi_i}\right]$$

Since $h \lesssim \frac{\min \phi_i}{M}$,

$$\mathbb{E}\|\Delta_{k+1}\|_{\boldsymbol{A}^{-1}}^2 \le (1 - \frac{hm}{2})\mathbb{E}\|\Delta_k\|_{\boldsymbol{A}^{-1}}^2 + \frac{1}{m}\sum_i [\frac{4rh^2 M_i^2}{\phi_i}].$$

Unrolling,

$$\mathbb{E} \|\Delta_{k+1}\|_{\boldsymbol{A}^{-1}}^2 \lesssim (1 - \frac{hm}{2})^k \mathbb{E} \|\Delta_0\|_{\boldsymbol{A}^{-1}}^2 + \frac{1}{m} \sum_j (1 - \frac{hm}{2})^{k-j} \sum_i [\frac{4rh^2 M_{ji}^2}{\phi_{ji}}]$$

$$\leq (1 - \frac{hm}{2})^k \mathbb{E} \|\Delta_0\|_{\boldsymbol{A}^{-1}}^2 + \frac{1}{m^2 h} \sum_i [\frac{4rh^2 M_i^2}{\phi_i}]$$

$$= (1 - \frac{hm}{2})^k \mathbb{E} \|\Delta_0\|_{\boldsymbol{A}^{-1}}^2 + \frac{1}{m^2} \sum_i [\frac{4rh M_i^2}{\phi_i}].$$

where $\sum_{i} \left[\frac{4rh^2 M_i^2}{\phi_i}\right]$ is an upper bound on $\sum_{i} \left[\frac{4rh^2 M_i^2}{\phi_i}\right]$. Optimally coupling Δ_0 and taking a square root yields

$$W_{2,\mathbf{A}^{-1}}(\mu_N,\pi) \lesssim \exp(-\frac{hmN}{4})W_{2,\mathbf{A}^{-1}}(\mu_0,\pi) + \frac{\sqrt{rh}}{m}\sqrt{\sum_i \frac{M_i^2}{\phi_i}}.$$

B. Continuous Time Subspace Descent

We can prove a similar result to the discrete-time convergence of subspace descent to its continuous-time counterpart (1).

Proposition 1. Suppose that f satisfies a PL inequality with parameter α . Then, the continuous time flow (1) satisfies

$$f(x_t) - f(x_\star) \le \exp\left(-\int_0^t m_t \alpha dt\right) (f(x_0) - f(x_\star)),$$

where

$$m_t = \frac{\langle \nabla f(x_t), \boldsymbol{P}_t \nabla f(x_t) \rangle}{\|\nabla f(x_t)\|^2} \ge 0.$$

Furthermore, if $\mathbb{E} \mathbf{P}_t = c \mathbf{I}$, then

$$\mathbb{E}f(x_t) - f(x_\star) \le \exp\left(-t\alpha c\right) \left(f(x_0) - f(x_\star)\right),$$

Proof.

$$\partial_t (f(x_t) - f(x_\star)) = \langle \nabla f(x_t), \mathbf{P}_t \nabla f(x_t) \rangle$$

$$\leq -m_t \| \nabla f(x_t) \|^2$$

$$\leq -m_t \alpha (f(x_t) - f(x_\star)).$$

Therefore by Grönwall's inequality,

$$f(x_t) - f(x_\star) \le \exp\left(-\int_0^t m_t \alpha dt\right) (f(x_0) - f(x_\star)).$$

If we instead take an expectation,

$$\partial_t (\mathbb{E}f(x_t) - f(x_\star)) = \mathbb{E}\langle \nabla f(x_t), \boldsymbol{P}_t \nabla f(x_t) \rangle$$

$$\leq -c \mathbb{E} \| \nabla f(x_t) \|^2$$

$$\leq -c \alpha (\mathbb{E}f(x_t) - f(x_\star)).$$

Notice that in expectation there is no loss in convergence rate when compared with the standard result for gradient flow under a PL inequality, provided that c = 1. Furthermore, if P_t is adapted to the flow x_t in the sense that m_t is close to 1, then we see that there rate of convergence for the subspace flow is close to that of the deterministic full-dimensional gradient flow.