ON THE PRINCIPLE OF LINEARIZED STABILITY FOR QUASILINEAR EVOLUTION EQUATIONS IN TIME-WEIGHTED SPACES

BOGDAN–VASILE MATIOC, LINA SOPHIE SCHMITZ, AND CHRISTOPH WALKER

ABSTRACT. Quasilinear (and semilinear) parabolic problems of the form $v' = A(v)v + f(v)$ with strict inclusion dom(f) \subsetneq dom(A) of the domains of the function $v \mapsto f(v)$ and the quasilinear part $v \mapsto A(v)$ are considered in the framework of time-weighted function spaces. This allows one to establish the principle of linearized stability in intermediate spaces lying between $dom(f)$ and $dom(A)$ and yields a greater flexibility with respect to the phase space for the evolution. In applications to differential equations such intermediate spaces may correspond to critical spaces exhibiting a scaling invariance. Several examples are provided to demonstrate the applicability of the results.

1. INTRODUCTION

The principle of linearized stability is a widely recognized method for various nonlinear parabolic evolution equations to derive stability or instability properties of an equilibrium from the real parts of the spectral points of the linearization at the equilibrium. Extensive research has been conducted on this subject under different assumptions and using various techniques, see e.g. $[6-8, 10, 11, 13, 15-19]$ $[6-8, 10, 11, 13, 15-19]$ $[6-8, 10, 11, 13, 15-19]$ $[6-8, 10, 11, 13, 15-19]$ $[6-8, 10, 11, 13, 15-19]$ $[6-8, 10, 11, 13, 15-19]$ $[6-8, 10, 11, 13, 15-19]$ $[6-8, 10, 11, 13, 15-19]$, though this list is by no means exhaustive.

In this research we focus on quasilinear parabolic problems

$$
v' = A(v)v + f(v), \quad t > 0, \qquad v(0) = v^0,
$$
\n(1.1)

assuming a strict inclusion dom(f) \subsetneq dom(A) for the domains of the function $v \mapsto f(v)$ and the quasilinear part $v \mapsto A(v)$. We shall establish herein the principle of linearized stability in phase spaces for the initial values that lie between $dom(f)$ and $dom(A)$ enabling greater flexibility in applications. For this purpose we rely on previous results [\[12,](#page-18-7) [14\]](#page-18-8) on the well-posedness of [\(1.1\)](#page-0-0) in time-weighted function spaces.

To be more precise, let E_0 and E_1 be Banach spaces over $\mathbb{K} \in \{ \mathbb{R}, \mathbb{C} \}$ with continuous and dense embedding

$$
E_1 \stackrel{d}{\hookrightarrow} E_0.
$$

Given $\theta \in (0, 1)$, we fix an admissible interpolation functor $(\cdot, \cdot)_{\theta}$ of exponent θ (see [\[4,](#page-18-9) I.Sections 2.1, 2.11]) and set $E_{\theta} := (E_0, E_1)_{\theta}$ and $\|\cdot\|_{\theta} := \|\cdot\|_{E_{\theta}}$. Then

$$
E_1 \stackrel{d}{\hookrightarrow} E_{\theta} \stackrel{d}{\hookrightarrow} E_0.
$$

Moreover, we fix numbers

$$
0 < \gamma < \beta < \xi < 1, \qquad q \ge 1,\tag{1.2a}
$$

and assume for the quasilinear part in [\(1.1\)](#page-0-0) that

$$
A \in C^{1-}(O_{\beta}, \mathcal{H}(E_1, E_0)), \qquad (1.2b)
$$

where

$$
\emptyset \neq O_{\beta} \text{ is an open subset of } E_{\beta}. \tag{1.2c}
$$

By $\mathcal{H}(E_1, E_0)$ we denote the open subset of the bounded linear operators $\mathcal{L}(E_1, E_0)$ consisting of generators of strongly continuous analytic semigroups on E_0 . The semilinear part $f: O_{\xi} \to E_{\gamma}$ is assumed to belong to

²⁰²⁰ Mathematics Subject Classification. 35B35; 35B40; 35K59.

Key words and phrases. Quasilinear parabolic problem; Principle of linearized stability; Interpolation spaces.

the class of locally Lipschitz continuous mappings from the open subset $O_{\xi} := O_{\beta} \cap E_{\xi}$ of E_{ξ} into E_{γ} . More precisely, we assume that for each $R > 0$ there is $c(R) > 0$ such that, for all $w, v \in O_{\xi} \cap \mathbb{B}_{E_{\beta}}(0, R)$,

$$
||f(w) - f(v)||_{\gamma} \le c(R) \left[1 + ||w||_{\xi}^{q-1} + ||v||_{\xi}^{q-1} \right] \left[\left(1 + ||w||_{\xi} + ||v||_{\xi} \right) ||w - v||_{\beta} + ||w - v||_{\xi} \right].
$$
 (1.2d)

Moreover, we define a critical value

$$
\alpha_{\rm crit} := \frac{q\xi - 1 - \gamma}{q - 1} \quad \text{if } q > 1, \qquad \alpha_{\rm crit} := -\infty \quad \text{if } q = 1,
$$
\n(1.2e)

with $\alpha_{\rm crit} < \xi$, and consider initial values $v^0 \in O_\alpha := O_\beta \cap E_\alpha$ with

$$
\alpha_{\rm crit} \le \alpha \in (\beta, \xi). \tag{1.2f}
$$

The well-posedness theory for [\(1.1\)](#page-0-0) developed in [\[12\]](#page-18-7) in the critical case $\alpha = \alpha_{\rm crit} \in (\beta, \xi)$ relies on the assumption that there are interpolation functors $\{\cdot,\cdot\}_{\alpha/\xi}$ and $\{\cdot,\cdot\}_{\gamma/\eta}$ of exponents α/ξ and γ/η , respectively, for $\eta \in \{\alpha, \beta, \xi\}$, such that

$$
E_{\alpha} \doteq \{E_0, E_{\xi}\}_{{\alpha}/{\xi}}, \qquad E_{\gamma} \doteq \{E_0, E_{\eta}\}_{{\gamma}/{\eta}}, \quad {\eta \in {\alpha}, \beta, \xi}.
$$
 (1.3)

Assumption [\(1.3\)](#page-1-0) is not really restrictive in applications:

Remark 1.1. Assumption [\(1.3\)](#page-1-0) is automatically satisfied if, for each $\theta \in \{\gamma, \beta, \alpha, \xi\}$, the interpolation functor $(\cdot, \cdot)_{\theta}$ is chosen to be always either the complex interpolation functor $[\cdot, \cdot]_{\theta}$, or the continuous interpolation functor $(\cdot, \cdot)_{\theta,\infty}^0$, or the real interpolation functor $(\cdot, \cdot)_{\theta} = (\cdot, \cdot)_{\theta,p}$ with parameter $p \in [1,\infty]$. This is a consequence of the reiteration theorems for these functors, see, e.g., [\[4,](#page-18-9) I.Remarks 2.11.2 (b)].

It is also worth emphasizing that [\(1.2d\)](#page-1-1) is satisfied in the particular case when

$$
||f(w) - f(v)||_{\gamma} \le c [||w||_{\xi}^{q-1} + ||v||_{\xi}^{q-1}] ||w - v||_{\xi}, \qquad w, v \in O_{\xi}.
$$
 (1.4)

The special case [\(1.4\)](#page-1-2) of [\(1.2d\)](#page-1-1) frequently arises in applications; see, for instance, the examples provided in Section [4.](#page-12-0)

Let us point out with respect to the parameters in $(1.2a)$ that the parameter $q \ge 1$ measures the growth of the nonlinearity f with respect to the E_{ξ} -terms, and its value is generally fixed and usually cannot be adjusted freely. In certain applications, such as considered in Examples [4.2](#page-13-0) and [4.3,](#page-16-0) there may be, however, some flexibility with respect to the choice of the interpolation exponents β , γ , and ξ . These parameters then define the range of $\alpha \geq \alpha_{\rm crit}$ for the phase space E_{α} with critical value $\alpha = \alpha_{\rm crit}$. In certain applications, the critical space $E_{\alpha_{\rm crit}}$ can also be identified due to scaling invariance properties (see Example [4.3\)](#page-16-0).

For simplicity, we only consider herein the situation that $\alpha \in (\beta, \xi)$ since the principle of linearized stability for the case $\alpha > \xi$ is treated in [\[13\]](#page-18-4). We emphasize that $E_{\xi} \hookrightarrow E_{\alpha} \hookrightarrow E_{\beta}$ in the setting herein and hence, the semilinear part f, being defined on O_{ξ} , does not necessarily need to be defined on the phase space E_{α} and requires possibly more regularity than the quasilinear part A.

We first recall the well-posedness of the quasilinear problem (1.1) .

Well-Posedness. Existence and uniqueness of solutions to problem [\(1.1\)](#page-0-0) have been derived in [\[14,](#page-18-8) Theorem 1.1] for $\alpha > \alpha_{\rm crit}$ assuming [\(1.2\)](#page-0-2), respectively in [\[12,](#page-18-7) Theorem 1.2] for the critical case $\alpha = \alpha_{\rm crit}$ assum-ing [\(1.2\)](#page-0-2)-[\(1.3\)](#page-1-0). The key ingredient to establish well-posedness for initial values v^0 in the phase space E_α is the use of time-weighted spaces of continuous functions $v : (0, T] \to E_{\xi}$ satisfying

$$
\sup_{t \in (0,T]} t^{\mu} ||v(t)||_{\xi} < \infty \quad \text{and} \quad \lim_{t \to 0} t^{\mu} ||v(t)||_{\xi} = 0
$$

for some $\mu \geq \xi - \alpha$, which are adapted to the regularizing effects of the parabolic operator A.

The relevant aspects of the results from [\[12,](#page-18-7)[14\]](#page-18-8) that are essential for our present purposes are summarized in Theorem [1.2.](#page-2-0) To unify these aspects into a single statement applicable to both critical and noncritical regimes, we assume throughout (1.2) – (1.3) . This approach provides a slight refinement of the well-posedness result established in [\[14,](#page-18-8) Theorem 1.1], as detailed below.

Theorem 1.2 (Well-Posedness). Assume [\(1.2\)](#page-0-2)-[\(1.3\)](#page-1-0). Then, given any $v^0 \in O_\alpha$, the Cauchy problem [\(1.1\)](#page-0-0) possesses a unique maximal strong solution

$$
v(\cdot; v^0) \in C^1((0, t^+(v^0)), E_0) \cap C((0, t^+(v^0)), E_1) \cap C([0, t^+(v^0)), O_\alpha) \cap C^{\alpha-\beta}([0, t^+(v^0)), E_\beta)
$$
(1.5a)

with $t^+(v^0) \in (0,\infty]$, such that

$$
\lim_{t \to 0} t^{\xi - \alpha} \|v(t; v^0)\|_{E_{\xi}} = 0.
$$
\n(1.5b)

Moreover, if $v^0 \in O_\alpha$ is such that $t^+(v^0) < \infty$ and $v(\cdot; v^0) : [0, t^+(v^0)) \to E_\beta$ is uniformly Hölder continuous, then

$$
\limsup_{t \nearrow t^+(v^0)} ||f(v(t;v^0))||_0 = \infty \quad or \quad \lim_{t \nearrow t^+(v^0)} \text{dist}_{E_\beta}(v(t;v^0), \partial O_\beta) = 0. \tag{1.6}
$$

Proof. The existence and uniqueness of the maximal strong solution is established in [\[12,](#page-18-7) Theorem 1.2] for the critical value $\alpha_{\rm crit} = \alpha \in (\beta, \xi)$. Moreover, in the noncritical case $\alpha_{\rm crit} < \alpha \in (\beta, \xi)$, it is proven in [\[14,](#page-18-8) Theorem 1.1 (without assuming (1.3)) that (1.1) has a unique maximal strong solution which satisfies $(1.5a)$ together with

$$
\lim_{t \to 0} t^{\mu} \| v(t; v^{0}) \|_{E_{\xi}} = 0 \tag{1.7}
$$

for all $\mu > \xi - \alpha$ (instead of the stronger property [\(1.5b\)](#page-2-2)). This solution fulfills the variation-of-constants formula

$$
v(t) = U_{A(v)}(t,0)v^{0} + \int_{0}^{t} U_{A(v)}(t,\tau) f(v(\tau)) d\tau, \qquad t \in (0,t^{+}(v^{0})),
$$

where $U_{A(v)}(\cdot, \cdot)$ is the evolution operator associated with $A(v(\cdot))$. Assuming [\(1.3\)](#page-1-0), it readily follows that $t^{\xi-\alpha}U_{A(v)}(t,0)v^0\to 0$ in E_{ξ} as $t\to 0$, see e.g. the proof of [\[12,](#page-18-7) Proposition 2.1]. Moreover, arguing as in the proof of $[14,$ Proposition 2.1, Eq. (2.18) , we infer from (1.7) that

$$
t^{\xi-\alpha} \int_0^t U_{A(v)}(t,\tau) f(v(\tau)) d\tau \xrightarrow[t\to 0]{} 0 \quad \text{in } E_\xi,
$$

which proves $(1.5b)$.

Finally, the blow-up criterion [\(1.6\)](#page-2-4) is provided for $\alpha_{\rm crit} = \alpha \in (\beta, \xi)$ in [\[12,](#page-18-7) Theorem 1.2 (v) (b)]. The proof of [\(1.6\)](#page-2-4) in the case when $\alpha_{\rm crit} < \alpha \in (\beta, \xi)$ is identical to that of [\[12,](#page-18-7) Theorem 1.2 (v) (b)] and therefore we omit herein the details. \Box

Exponential Stability. We shall now present the principle of linearized stability in the phase space E_{α} for a certain range of exponents α . With regard to applications it proves crucial to state sharp results including certain special cases. To this end, let

$$
v_* \in O_1 := O_\beta \cap E_1 \quad \text{with} \quad A(v_*)v_* + f(v_*) = 0 \tag{1.8a}
$$

be an equilibrium solution to [\(1.1\)](#page-0-0). In order to derive stability properties for $v_*,$ we further assume that

$$
f: O_{\xi} \to E_0
$$
 and $A(\cdot)v_*: O_{\xi} \to E_0$ are Fréchet differentiable at v_* (1.8b)

with Fréchet derivatives $\partial f(v_*) \in \mathcal{L}(E_{\xi}, E_0)$ and $(\partial A(v_*)[\cdot])v_* \in \mathcal{L}(E_{\xi}, E_0)$, respectively, such that there are

$$
\gamma_* \in [0, \gamma], \qquad q_* > 1,\tag{1.8c}
$$

and $r_*, c_* > 0$ with

$$
|| f(w + v_*) - f(v_*) - \partial f(v_*)w ||_{\gamma_*} \le c_* ||w||_{\xi}^{q_*}, \qquad w \in \widehat{O}_{\xi} \cap \overline{\mathbb{B}}_{E_{\alpha}}(0, r_*), \qquad (1.8d)
$$

and

$$
||A(w + v_*)v_* - A(v_*)v_* - (\partial A(v_*)[w])v_*||_{\gamma_*} \leq c_* ||w||_{\xi}^{q_*}, \qquad w \in \widehat{O}_{\xi} \cap \overline{\mathbb{B}}_{E_{\alpha}}(0, r_*) ,
$$
 (1.8e)

where $\widehat{O}_{\xi} := O_{\xi} - v_*$. Moreover, we assume that the linearized operator

$$
\mathbb{A} := A(v_*) + (\partial A(v_*)[\cdot])v_* + \partial f(v_*) \in \mathcal{L}(E_1, E_0)
$$

has a negative spectral bound, that is,

$$
-\omega_0 := s(\mathbb{A}) := \sup \{ \text{Re } \lambda : \lambda \in \sigma(\mathbb{A}) \} < 0. \tag{1.8f}
$$

We shall prove the following result regarding the asymptotic exponential stability of the equilibrium v_* . **Theorem 1.3** (Stability). Assume (1.2) , (1.3) , and (1.8) and set

$$
\alpha^*_{\textnormal{crit}}:=\frac{q_*\xi-1-\gamma_*}{q_*-1}<\xi\,.
$$

Moreover, assume $\alpha \geq \alpha_{\text{crit}}^*$ with strict inequality $\alpha > \alpha_{\text{crit}}^*$ in case that $\gamma_* \in (0, \gamma)$. Then, the equilibrium v_* is asymptotically exponentially stable in E_{α} . More precisely, given any $\omega \in (0, \omega_0)$, there exist $\varepsilon_0 > 0$ and $M \geq 1$ such that, for each $v^0 \in \overline{\mathbb{B}}_{E_\alpha}(v_*, \varepsilon_0)$, the solution to [\(1.1\)](#page-0-0) exists globally in time and

$$
||v(t; v^{0}) - v_{*}||_{\alpha} + t^{\xi - \alpha} ||v(t; v^{0}) - v_{*}||_{\xi} \leq Me^{-\omega t} ||v^{0} - v_{*}||_{\alpha}, \qquad t \geq 0.
$$
 (1.9)

Remark 1.4. In the noncritical case $\max\{\alpha_{\rm crit}, \alpha_{\rm crit}^*\} < \alpha \in (\beta, \xi)$ one may actually drop the assumption (1.3) in Theorem [1.3.](#page-3-0) The claim remains valid (with a similar proof) provided [\(1.9\)](#page-3-1) is replaced by

$$
||v(t; v0) - v_*||_{\alpha} + t^{\mu}||v(t; v0) - v_*||_{\xi} \leq Me^{-\omega t}||v0 - v_*||_{\alpha}, \qquad t \geq 0,
$$

for some fixed (but arbitrary) $\mu > \xi - \alpha$. Since in many applications [\(1.3\)](#page-1-0) is automatically satisfied (see Remark [1.1\)](#page-1-3), we chose to present the (slightly) less general but more concise result in Theorem [1.3.](#page-3-0)

With respect to the assumption [\(1.8e\)](#page-2-6), note that, if $v_* = 0$ and $\partial f(v_*) = 0$, then one may take, in the particular context of [\(1.4\)](#page-1-2), $\gamma_* = \gamma$ and $q_* = q$ in [\(1.8c\)](#page-2-7)-[\(1.8e\)](#page-2-6) (and hence $\alpha_{\rm crit} = \alpha_{\rm crit}^*$). This observation proves very useful in the critical case, as illustrated in Example [4.3.](#page-16-0) Moreover, even if v_* is non-zero, since $A(v_*)v_* = -f(v_*) \in E_\gamma$ by [\(1.8a\)](#page-2-8), the equilibrium v_* (and consequently the linearizations $(\partial A(v_*)[\cdot])v_*$ and $\partial f(v_*)$) may exhibit higher regularity properties in applications. As a consequence, one may more broadly expect to have $\gamma_* > 0$ in [\(1.8c\)](#page-2-7)-[\(1.8e\)](#page-2-6).

We also point out that though (γ_*, q_*) and (γ, q) need not be related, in applications it often turns out that $(\gamma_*, q_*) = (\gamma, q)$ and therefore $\alpha_{\rm crit} = \alpha_{\rm crit}^*$, see the examples in Section [4.](#page-12-0)

Semilinear Evolution Equations. The previous results hold, of course, also for the semilinear evolution problem

$$
v' = Av + f(v), \quad t > 0, \qquad v(0) = v^0,
$$
\n(1.10)

with

$$
A \in \mathcal{H}(E_1, E_0). \tag{1.11a}
$$

However, for this particular case, we present a sharper version of the exponential stability result in Theorem [1.3.](#page-3-0) Indeed, we now only assume that

$$
0 \le \gamma < \xi \le 1, \qquad (\gamma, \xi) \ne (0, 1), \qquad q \ge 1, \tag{1.11b}
$$

and define again

$$
\alpha_{\rm crit} := \frac{q\xi - 1 - \gamma}{q - 1} \quad \text{if } q > 1, \qquad \alpha_{\rm crit} := -\infty \quad \text{if } q = 1,
$$
\n(1.11c)

noticing that $\alpha_{\rm crit} < \xi$. We assume that

either
$$
\alpha_{\rm crit} < \alpha \in [\gamma, \xi)
$$
 or $\alpha_{\rm crit} = \alpha \in (\gamma, \xi)$, $(1.11d)$

and let O_{α} be an arbitrary open subset of E_{α} . The semilinearity $f: O_{\xi} := O_{\alpha} \cap E_{\xi} \to E_{\gamma}$ is again assumed to be locally Lipschitz continuous in the sense that for each $R > 0$ there is a constant $c(R) > 0$ such that

$$
||f(w) - f(v)||_{\gamma} \le c(R) \left[1 + ||w||_{\xi}^{q-1} + ||v||_{\xi}^{q-1}\right] \left[(1 + ||w||_{\xi} + ||v||_{\xi})||w - v||_{\alpha} + ||w - v||_{\xi}\right] \tag{1.11e}
$$

for all $w, v \in O_{\xi} \cap \mathbb{B}_{E_{\alpha}}(0, R)$.

The well-posedness of [\(1.1\)](#page-0-0) is established in [\[12\]](#page-18-7) for the critical case $\alpha = \alpha_{\rm crit} \in (\gamma, \xi)$ under the assumption that, if $\xi < 1$, there exists an interpolation functor $\{\cdot,\cdot\}_{\alpha/\xi}$ of exponent α/ξ , and, if $\gamma > 0$, that for $\eta \in \{\alpha, \xi\} \setminus \{1\}$ there are interpolation functors $\{\cdot, \cdot\}_{\gamma/\eta}$ of exponent γ/η such that

$$
E_{\alpha} \doteq \{E_0, E_{\xi}\}_{{\alpha}/{\xi}} \text{ if } {\xi} < 1, \qquad E_{\gamma} \doteq \{E_0, E_{\eta}\}_{{\gamma}/{\eta}} \text{ if } {\gamma} > 0, \ \eta \in {\{\alpha, \xi\}} \setminus {\{1\}}. \tag{1.12}
$$

Under these assumptions the Cauchy problem [\(1.10\)](#page-3-2) is locally well-posed in O_{α} . Similarly to the quasilinear problem [\(1.1\)](#page-0-0), we also assume in the noncritical case the interpolation property [\(1.12\)](#page-4-0). This leads to a unified and slightly refined well-posedness result for [\(1.10\)](#page-3-2), compared to [\[12,](#page-18-7) Theorem 1.3].

Theorem 1.5 (Well-Posedness). Assume [\(1.11\)](#page-3-3)-[\(1.12\)](#page-4-0). Then, given any $v^0 \in O_\alpha$, the semilinear Cauchy problem [\(1.10\)](#page-3-2) possesses a unique maximal strong solution

$$
v(\cdot; v^0) \in C^1((0, t^+(v^0)), E_0) \cap C((0, t^+(v^0)), E_1) \cap C([0, t^+(v^0)), O_\alpha)
$$
\n(1.13a)

with $t^+(v^0) \in (0,\infty]$, such that

$$
\lim_{t \to 0} t^{\xi - \alpha} \|v(t; v^0)\|_{E_{\xi}} = 0.
$$
\n(1.13b)

Moreover, if $v^0 \in O_\alpha$ is such that $t^+(v^0) < \infty$, then

$$
\limsup_{t \nearrow t^+(v^0)} ||f(v(t;v^0))||_0 = \infty \qquad or \qquad \lim_{t \nearrow t^+(v^0)} \operatorname{dist}_{E_\alpha}\left(v(t;v^0), \partial O_\alpha\right) = 0 \tag{1.14}
$$

Proof. This result is in [\[12,](#page-18-7) Theorem 1.3] in the critical case $\alpha = \alpha_{\rm crit} \in (\gamma, \xi)$. For $\alpha_{\rm crit} < \alpha \in [\gamma, \xi)$ the claim follows by arguing along the lines of the proof of [\[12,](#page-18-7) Theorem 1.3] (we omit therefore the details). For an alternative proof in the case $\alpha_{\rm crit} < \alpha \in [\gamma, \xi]$ under the restriction $O_\alpha = E_\alpha$ (but with a stronger blow-up criterion than in (1.14)) we refer to [\[14,](#page-18-8) Theorem 1.2].

Let again

$$
v_* \in O_1 := O_\alpha \cap E_1 \quad \text{with} \quad Av_* + f(v_*) = 0 \tag{1.15a}
$$

be an equilibrium solution to [\(1.10\)](#page-3-2) such that

$$
f: O_{\xi} \to E_0 \qquad \text{is Fréchet differentiable at } v_* \,. \tag{1.15b}
$$

For the Fréchet derivative $\partial f(v_*) \in \mathcal{L}(E_{\xi}, E_0)$ we assume that there exist

$$
\gamma_* \in [0, \gamma], \qquad q_* > 1 \,, \tag{1.15c}
$$

and constants $r_*, c_* > 0$ such that

$$
|| f(w + v_*) - f(v_*) - \partial f(v_*)w ||_{\gamma_*} \le c_* ||w||_{\xi}^{q_*}, \qquad w \in \widehat{O}_{\xi} \cap \overline{\mathbb{B}}_{E_{\alpha}}(0, r_*) ,
$$
 (1.15d)

where again $\widehat{O}_{\xi} := O_{\xi} - v_*$. Finally, we assume that the linearized operator

$$
\mathbb{A} := A + \partial f(v_*) \in \mathcal{L}(E_1, E_0)
$$

has a negative spectral bound, that is

$$
-\omega_0 := s(\mathbb{A}) := \sup \{ \text{Re } \lambda : \lambda \in \sigma(\mathbb{A}) \} < 0. \tag{1.15e}
$$

We can now state the exponential stability of the equilibrium solution v_* to the semilinear problem [\(1.10\)](#page-3-2): **Theorem 1.6** (Stability). Assume (1.11) , (1.12) , and (1.15) and set

$$
\alpha^*_{\textnormal{crit}}:=\frac{q_*\xi-1-\gamma_*}{q_*-1}<\xi\,.
$$

Moreover, assume $\alpha \geq \alpha_{\text{crit}}^*$ with strict inequality $\alpha > \alpha_{\text{crit}}^*$ in case that $\gamma_* \in (0, \gamma)$ or $\alpha = \gamma$. Then, the equilibrium v_* is asymptotically exponentially stable in E_α . More precisely, given any $\omega \in (0, \omega_0)$, there are $\varepsilon_0 > 0$ and $M \ge 1$ such that, for each $v^0 \in \overline{\mathbb{B}}_{E_\alpha}(v_*, \varepsilon_0)$, the solution to [\(1.10\)](#page-3-2) exists globally in time and

$$
||v(t; v^{0}) - v_{*}||_{\alpha} + t^{\xi - \alpha} ||v(t; v^{0}) - v_{*}||_{\xi} \leq Me^{-\omega t} ||v^{0} - v_{*}||_{\alpha}, \qquad t \geq 0.
$$
 (1.16)

Remark 1.7. As for the quasilinear problem [\(1.1\)](#page-0-0), in the noncritical case $\max\{\alpha_{\rm crit}, \alpha_{\rm crit}^*\} < \alpha \in [\gamma, \xi)$, the assumption [\(1.12\)](#page-4-0) in Theorem [1.6](#page-4-3) can be omitted, and the claim remains valid (with a similar proof) provided that [\(1.16\)](#page-4-4) is replaced by

$$
||v(t; v0) - v_*||_{\alpha} + t^{\mu}||v(t; v0) - v_*||_{\xi} \leq Me^{-\omega t}||v0 - v_*||_{\alpha}, \qquad t \geq 0,
$$

for some fixed (but arbitrary) $\mu > \xi - \alpha$. For similar reasons as in Remark [1.4,](#page-3-4) we present herein the (slightly) less general but more concise result in Theorem [1.6.](#page-4-3)

As in the quasilinear case, in applications it often has $(\gamma_*, q_*) = (\gamma, q)$ and $\alpha_{\rm crit} = \alpha_{\rm crit}^*$, see the examples in Section [4.](#page-12-0)

Instability. For the sake of completeness we also state conditions sufficient for instability of an equilibrium solution $v_* \in O_1 \subset E_1$ to the quasilinear problem [\(1.1\)](#page-0-0) or the semilinear problem [\(1.10\)](#page-3-2). Specifically, we assume that

$$
(A, f) \in C^{2-}(O_1, \mathcal{L}(E_1, E_0) \times E_0)
$$
\n(1.17a)

with Fréchet derivatives

$$
\partial f(v_*) , (\partial A(v_*)[\cdot])v_* \in \mathcal{L}(E_\eta, E_0) \quad \text{for some } \eta \in [\beta, 1). \tag{1.17b}
$$

Moreover, we require that the linearized operator

$$
\mathbb{A} = A(v_*) + (\partial A(v_*)[\cdot])v_* + \partial f(v_*) \in \mathcal{H}(E_1, E_0)
$$

satisfies

$$
\sigma_{+}(\mathbb{A}) := \{ \lambda \in \sigma(\mathbb{A}) : \text{Re}\,\lambda > 0 \} \neq \emptyset, \qquad \inf\{\text{Re}\,\lambda : \lambda \in \sigma_{+}(\mathbb{A})\} > 0. \tag{1.17c}
$$

Conditions [\(1.17\)](#page-5-0) guarantee the instability of v_* in the phase space E_α :

Theorem 1.8 (Instability). Assume [\(1.17\)](#page-5-0) and that the assumptions of Theorem [1.2](#page-2-0) are satisfied for the quasilinear problem [\(1.1\)](#page-0-0) or, alternatively, the assumptions of Theorem [1.5](#page-4-5) are satisfied for the semilinear problem [\(1.10\)](#page-3-2).

Then, the equilibrium v_* is unstable in E_α . More precisely, there exists a neighborhood U of v_* in O_α such that for each $n \in \mathbb{N}^*$ there exists $v_n^0 \in \mathbb{B}_{E_\alpha}(v_*, 1/n) \cap O_\alpha$ such that the corresponding solution $v(\cdot; v_n^0)$ to (1.1) or (1.10) satisfies

$$
v(t; v_n^0) \notin U \quad \text{for some } t \in (0, t^+(v_n^0)).
$$

The proof of this result is similar to the one in the classical case $\xi = \beta$, see [\[13,](#page-18-4) Theorem 1.4], and is based on an instability result for fully-nonlinear parabolic problems established in [\[11,](#page-18-3) Theorem 9.1.3], where instability in E_1 is proven.

Outline. The outline of the paper is as follows. Section [2](#page-5-1) is dedicated to the quasilinear problem [\(1.1\)](#page-0-0). We provide a detailed proof of Theorem [1.3.](#page-3-0) In Section [3](#page-11-0) we sketch the proof of Theorem [1.6](#page-4-3) for the semilinear problem [\(1.10\)](#page-3-2). In Section [4](#page-12-0) we present examples that illustrate our previous findings both for the critical case $\alpha = \alpha_{\rm crit}$ and the non-critical case $\alpha > \alpha_{\rm crit}$, respectively, including quasilinear problems with quadratic semilinearities, a parabolic-parabolic chemotaxis system, and a quasilinear evolution equation in critical spaces with scaling invariance.

2. Quasilinear Problem: Proof of Theorem [1.3](#page-3-0)

Before establishing the proofs of Theorem [1.3](#page-3-0) and Theorem [1.8](#page-5-2) we recall the following consequence of (the proof of) Theorem [1.2](#page-2-0) that relies on the semiflow property of the solution map associated with problem [\(1.1\)](#page-0-0). Corollary 2.1. Assume [\(1.2\)](#page-0-2), [\(1.3\)](#page-1-0), and [\(1.8a\)](#page-2-8). Then, given $t_* > 1$, there are a neighborhood V_α of v_* in O_{α} and $k_0 \geq 1$ such that $t_* < t^+(v^0)$ and

$$
||v(t; v0) - v_*||_{\alpha} \le k_0 ||v0 - v_*||_{\alpha}, \quad 0 \le t \le t_*,
$$
\n(2.1)

for each $v^0 \in V_\alpha$. Moreover, there exist $t_0 \in (0,1)$ and $k_1 > 0$ such that

$$
||v(t; v0) - v(s; v0)||_{\beta} \le k_1(t - s)^{\alpha - \beta}, \quad 0 \le s \le t \le t_0, \quad v0 \in V_{\alpha}.
$$
 (2.2)

Proof. Property [\(2.1\)](#page-6-0) is a consequence of the continuous dependence in E_α stated in [\[14,](#page-18-8) Theorem 1.1 (iii)] for $\alpha_{\rm crit} < \alpha \in (\beta, \xi)$, respectively in [\[12,](#page-18-7) Theorem 1.2 (iii)] in the critical case $\alpha_{\rm crit} = \alpha \in (\beta, \xi)$ (see the proof thereof). The second statement is shown in the proof of [\[14,](#page-18-8) Proposition 2.1] (see (2.13) therein) for $\alpha_{\rm crit} < \alpha \in (\beta, \xi)$, respectively in the proof of [\[12,](#page-18-7) Proposition 2.1] (see (2.19)-(2.22) therein) in the case $\alpha_{\rm crit} = \alpha \in (\beta, \xi)$.

The proof of Theorem [1.3](#page-3-0) is performed in the framework of time-weighted spaces. Recall that, given a Banach space E, $\mu \in \mathbb{R}$, and $T > 0$, the space

$$
C_{\mu}\big((0,T],E\big):=\big\{u\in C((0,T],E)\,:\,t^{\mu}\|u(t)\|_{E}\to 0\text{ for }t\to 0\big\}
$$

is a Banach space with norm

$$
||u||_{C_{\mu}((0,T],E)} := \sup_{t \in (0,T]} t^{\mu} ||u(t)||_{E}.
$$

We will use this notation in the subsequent analysis.

Proof of Theorem [1.3.](#page-3-0) Let the assumptions of Theorem [1.3](#page-3-0) hold.

(a) Preliminaries and Notation. Let $e_{\alpha,\beta} > 0$ be the norm of the continuous embedding $E_{\alpha} \hookrightarrow E_{\beta}$. Since O_β is open in E_β , there exists $\varepsilon > 0$ such that

$$
\overline{\mathbb{B}}_{E_{\alpha}}(v_*, 2\varepsilon/\mathsf{e}_{\alpha,\beta}) \subset \overline{\mathbb{B}}_{E_{\beta}}(v_*, 2\varepsilon) \subset O_{\beta}.
$$
\n(2.3)

We define

$$
\widehat{A}(w) := A(w + v_*) + (\partial A(v_*)[\cdot])v_* + \partial f(v_*), \qquad w \in \widehat{O}_{\beta},
$$

and

$$
\widehat{f}(w) := f(w + v_*) + A(w + v_*)v_* - \partial f(v_*)w - (\partial A(v_*)[w])v_*, \qquad w \in \widehat{O}_{\xi},
$$

where we set $\widehat{O}_{\theta} := O_{\theta} - v_*$ for $\theta \in \{\beta, \xi\}$. The functions \widehat{f} and \widehat{A} will appear in the reformulation [\(2.16\)](#page-8-0) below of [\(1.1\)](#page-0-0). Since $(\partial A(v_*)[\cdot])v_* - \partial f(v_*) \in \mathcal{L}(E_{\xi}, E_0)$ with $\xi < 1$, we may use a perturbation argument [\[4,](#page-18-9) I.Theorem 1.3.1] to infer from [\(1.2b\)](#page-0-3) that

$$
\widehat{A} \in C^{1-}(\widehat{O}_{\beta}, \mathcal{H}(E_1, E_0)).
$$
\n(2.4)

Let $\rho \in (0, \alpha - \beta)$ be fixed and note that $\alpha \geq \alpha^*_{\text{crit}}$ implies

$$
\mu := \xi - \alpha \le \frac{1 + \gamma_* - \alpha}{q_*}.
$$
\n
$$
(2.5)
$$

In fact, the inequality in [\(2.5\)](#page-6-1) turns into a strict inequality if $\alpha > \alpha^*_{\rm crit}$, that is, if $\gamma_* \in (0, \gamma)$ by assumption. If $\gamma_* \in (0, \gamma)$, we may thus choose $\gamma_0 \in (0, \gamma_*)$ such that $\mu q_* < 1 + \gamma_0 - \alpha$, while we set $\gamma_0 := \gamma_*$ if $\gamma_* \in \{0, \gamma\}$. We then have in any case that

$$
\mu q_* \le 1 + \gamma_0 - \alpha. \tag{2.6}
$$

Further, fix an arbitrary $\omega \in (0, \omega_0)$ with $\omega_0 - \omega = 4\delta > 0$. Noticing that $\hat{A}(0) = \mathbb{A}$, we infer from [\(2.4\)](#page-6-2) and [\[4,](#page-18-9) I.Proposition [1](#page-7-0).4.2] that there are $\kappa \ge 1$ and $\ell > 0$ such that (making $\varepsilon > 0$ smaller, if necessary)¹

$$
\omega_0 - \delta + \tilde{A}(w) \in \mathcal{H}(E_1, E_0; \kappa, \delta), \qquad w \in \overline{\mathbb{B}}_{E_\beta}(0, 2\varepsilon), \tag{2.7}
$$

and

$$
\|\widehat{A}(w_1) - \widehat{A}(w_2)\|_{\mathcal{L}(E_1, E_0)} \le \ell \|w_1 - w_2\|_{\beta}, \qquad w_1, w_2 \in \overline{\mathbb{B}}_{E_{\beta}}(0, 2\varepsilon).
$$
\n(2.8)

For $T \in (0, \infty)$, we define

$$
\mathcal{M}(T) := \left\{ w \in C([0,T], \overline{\mathbb{B}}_{E_{\alpha}}(0, 2\varepsilon/\mathsf{e}_{\alpha,\beta})) : ||w(t) - w(s)||_{\beta} \leq \frac{N}{\ell} |t - s|^{\rho}, 0 \leq s, t \leq T \right\},\
$$

where $N > 0$ will be chosen below. More precisely, given $w \in \mathcal{M}(T)$, we derive from [\(2.7\)](#page-7-1)-[\(2.8\)](#page-7-2) that

$$
\omega_0 - \delta + A(w(t)) \in \mathcal{H}(E_1, E_0; \kappa, \delta), \qquad t \in [0, T], \tag{2.9a}
$$

and

$$
\widehat{A}(w) \in C^{\rho}([0, T], \mathcal{L}(E_1, E_0)) \quad \text{with} \quad \sup_{0 \le s < t \le T} \frac{\|A(w(t)) - A(w(s))\|_{\mathcal{L}(E_1, E_0)}}{(t - s)^{\rho}} \le N. \tag{2.9b}
$$

In view of [\(2.9\)](#page-7-3) we may apply the results of [\[4,](#page-18-9) II.Section 5] to the family $\mathcal{A} := {\{\hat{A}(w) : w \in \mathcal{M}(T)\}}.$ Letting $c_0(\rho) > 0$ be the constant from [\[4,](#page-18-9) II.Theorem 5.1.1] (which is independent of N), we choose $N > 0$ such that $c_0(\rho)N^{1/\rho} = \delta$. Then, by [\[4,](#page-18-9) II.Theorem 5.1.1, II.Lemma 5.1.3] there exists for each $w \in \mathcal{M}(T)$ a unique evolution operator $U_{\widehat{A}(w)}$ for $A(w)$ satisfying

$$
||U_{\widehat{A}(w)}(t,s)||_{\mathcal{L}(E_{\theta})} + (t-s)^{\theta-\vartheta_0} ||U_{\widehat{A}(w)}(t,s)||_{\mathcal{L}(E_{\vartheta},E_{\theta})} \le \frac{M_1}{2} e^{-\nu(t-s)}, \qquad 0 \le s \le t \le T,
$$
 (2.10a)

with a constant $M_1 > 0$ (independent of $T > 0$) and

$$
-\nu := c_0(\rho)N^{1/\rho} - \omega_0 + \delta + \delta = -\omega - \delta < -\omega < 0.
$$

The estimate [\(2.10a\)](#page-7-4) holds for $0 \le \vartheta_0 \le \vartheta \le 1$ with $\vartheta_0 < \vartheta$ if $0 < \vartheta < \vartheta < 1$, or, thanks to [\(1.3\)](#page-1-0), for $(\vartheta, \theta) \in \{(\alpha, \xi), (\gamma, \alpha), (\gamma, \beta), (\gamma, \xi)\}\$ with $\vartheta_0 = \vartheta$. Since we use [\(2.10a\)](#page-7-4) only for a finite number of pairs (ϑ, θ) , we may assume that the constant M_1 does not depend on (ϑ, θ) . Moreover, [\[4,](#page-18-9) II. Theorem 5.3.1] (with $f = 0$ therein) implies that there exists $M_2 > 0$ (independent of $T > 0$) with

$$
||U_{\widehat{A}(w)}(t,0) - U_{\widehat{A}(w)}(s,0)||_{\mathcal{L}(E_{\alpha},E_{\beta})} \le M_2(t-s)^{\alpha-\beta}, \quad 0 \le s \le t \le T,
$$
\n(2.10b)

while we may assume due to $[4, II.$ Equation $(5.3.8)$ that

$$
||U_{\widehat{A}(w)}(t,s) - 1||_{\mathcal{L}(E_{\alpha},E_{\beta})} \le M_2(t-s)^{\alpha-\beta}, \quad 0 \le s \le t \le T.
$$
\n(2.10c)

Recalling [\(2.6\)](#page-6-3), we may set

$$
c_0 := \sup_{r>0} \left(r^{1+\gamma_0 - \alpha - \mu q_*} e^{-\nu r} \right) < \infty,
$$

and, with B denoting the Beta function and noticing that $1 - \mu q_* \ge \alpha - \gamma_0 \ge \alpha - \gamma_* \ge \alpha - \gamma > 0$,

$$
\mathsf{B}_{\theta} := \mathsf{B}(\theta, 1 - \mu q_*) , \qquad \theta > 0 .
$$

$$
\kappa^{-1} (|\lambda| \|x\|_0 + \|x\|_1) \le \|(\lambda - A)x\|_0 \le \kappa (|\lambda| \|x\|_0 + \|x\|_1), \qquad x \in E_1, \quad \text{Re}\,\lambda \ge \infty.
$$

¹Given $\kappa \ge 1$ and $\varpi > 0$, the set $\mathcal{H}(E_1, E_0; \kappa, \varpi) \subset \mathcal{H}(E_1, E_0)$ consists of the bounded operators $A \in \mathcal{L}(E_1, E_0)$ with the property $\varpi - A \in \mathcal{L}(E_1, E_0)$ is an isomorphism and

Since $q_* > 1$ and $\mu q_* \leq 1 + \gamma_0 - \alpha$, we can choose $L > 0$ and $\varepsilon_0 \in (0, \varepsilon)$ such that

$$
\frac{\varepsilon_0 M_1}{2} + c_* c_0 M_1 L^{q_*} \mathsf{B}_{\mu q_*} \le \frac{\varepsilon}{\mathsf{e}_{\alpha,\beta}},\tag{2.11a}
$$

$$
\frac{\varepsilon_0 M_1}{2} + c_* c_0 M_1 L^{q_*} \mathsf{B}_{\mu(q_*-1)} \le \frac{L}{2},\tag{2.11b}
$$

$$
\left[\varepsilon_0 M_2 + c_* c_0 M_1 L^{q_*} \left(M_2 \mathsf{B}_{\mu q_*} + \mathsf{B}_{\alpha-\beta+\mu q_*}\right)\right] \left(\frac{4\varepsilon \ell}{N}\right)^{\frac{\alpha-\beta-\rho}{\rho}} \le \frac{N}{2\ell},\tag{2.11c}
$$

$$
c_* M_1 L^{q_*-1} \left(\mathsf{B}_{\mu q_*} + \mathsf{B}_{\mu (q_*-1)} \right) \left(\sup_{r>0} r^{1+\gamma_0 - \alpha - \mu q_*} e^{(\omega - \nu)r} \right) \le \frac{1}{2} \,. \tag{2.11d}
$$

Also note from [\(1.8d\)](#page-2-9)-[\(1.8e\)](#page-2-6) that (assuming without loss of generality that $2\varepsilon/\mathsf{e}_{\alpha,\beta} \leq r_*)$

$$
\|\widehat{f}(w)\|_{\gamma_*} \le 2c_* \|w\|_{\xi}^{q_*}, \qquad w \in \widehat{O}_{\xi} \cap \overline{\mathbb{B}}_{E_{\alpha}}(0, 2\varepsilon/\mathsf{e}_{\alpha,\beta}). \tag{2.12}
$$

We will also use frequently the observation that

$$
\sup_{t>0} \left(t^a \int_0^1 (1-s)^{-b} e^{-\varpi t (1-s)} s^{-\mu q_*} ds \right) \le \left(\sup_{r>0} r^a e^{-\varpi r} \right) \mathsf{B}_{1-a-b}, \qquad 0 \le a < 1-b, \quad \varpi > 0. \tag{2.13}
$$

(b) Global Existence and Uniform Estimates. Let now $v^0 \in \overline{\mathbb{B}}_{E_\alpha}(v_*, \varepsilon_0)$ be fixed. Then, the evolu-tion problem [\(1.1\)](#page-0-0) has a unique maximal strong solution $v(\cdot; v^0)$ on $[0, t^+(v^0))$ according to Theorem [1.2](#page-2-0) with regularity properties stated in [\(1.5\)](#page-2-10). Moreover, invoking Corollary [2.1](#page-6-4) and making $\varepsilon_0 > 0$ smaller, if necessary, we may assume that $\overline{\mathbb{B}}_{E_\alpha}(v_*, \varepsilon_0) \subset V_\alpha$ and $t^+(v^0) > 1$ with

$$
||v(t; v^{0}) - v_{*}||_{\alpha} \le k_{0} \varepsilon_{0} \le 2\varepsilon/\mathsf{e}_{\alpha,\beta}, \qquad t \in [0, 1], \quad v^{0} \in \overline{\mathbb{B}}_{E_{\alpha}}(v_{*}, \varepsilon_{0}). \tag{2.14}
$$

Moreover, Corollary [2.1](#page-6-4) ensures that there exist $t_0 \in (0,1)$ and a constant $k_1 > 0$ such that

$$
||v(t; v^{0}) - v(s; v^{0})||_{\beta} \le k_{1}(t - s)^{\alpha - \beta}, \qquad 0 \le s \le t \le t_{0}, \quad v^{0} \in \overline{\mathbb{B}}_{E_{\alpha}}(v_{*}, \varepsilon_{0}). \tag{2.15}
$$

We set $u^0 := v^0 - v_* \in \overline{\mathbb{B}}_{E_\alpha}(0, \varepsilon_0)$ and $u := v(\cdot; v^0) - v_*$. Then u is a strong solution to the Cauchy problem

$$
u' = \hat{A}(u)u + \hat{f}(u), \quad t > 0, \qquad u(0) = u^{0}.
$$
 (2.16)

Moreover, since $\rho < \alpha - \beta$, we can make $t_0 \in (0, 1)$ smaller, if necessary, to infer from [\(2.15\)](#page-8-1) that

$$
||u(t) - u(s)||_{\beta} = ||v(t; v^{0}) - v(s; v^{0})||_{\beta} \le k_{1}(t - s)^{\alpha - \beta} \le \frac{N}{\ell}(t - s)^{\rho}, \qquad 0 \le s \le t \le t_{0}, \quad v^{0} \in \overline{\mathbb{B}}_{E_{\alpha}}(v_{*}, \varepsilon_{0}).
$$

Recalling [\(2.14\)](#page-8-2), we deduce $u|_{[0,t_0]} \in \mathcal{M}(t_0)$. Moreover, since $u \in C_{\mu}((0,t_0], E_{\xi})$ by Theorem [1.2,](#page-2-0) we may assume (making again $t_0 > 0$ smaller, if necessary) that $||u||_{C_\mu((0,t_0],E_\xi)} \leq L$. Consequently, it holds that

$$
t_1:=\sup\left\{t
$$

Let $t \in (0, t_1)$. Then, $||u(s)||_{\xi} \leq s^{-\mu} L$ for $s \in (0, t]$, and (2.10) holds for $\widehat{A}(u)$ and $T = t$ since $u|_{[0,t]} \in \mathcal{M}(t)$. Moreover, u satisfies the variation-of-constants formula

$$
u(\tau) = U_{\widehat{A}(u)}(\tau,0)u^0 + \int_0^{\tau} U_{\widehat{A}(u)}(\tau,s)\widehat{f}(u(s)) \,ds \,, \quad 0 \le \tau \le t \,,
$$

according to [\(1.5\)](#page-2-10) and [\(2.16\)](#page-8-0). Therefore, we deduce from [\(2.10a\)](#page-7-4) and [\(2.12\)](#page-8-3) for $\theta \in {\alpha, \xi}$ that

$$
||u(t) - U_{\widehat{A}(u)}(t,0)u^0||_{\theta} \le \int_0^t ||U_{\widehat{A}(u)}(t,s)||_{\mathcal{L}(E_{\gamma_*},E_{\theta})} ||\widehat{f}(u(s))||_{\gamma_*} ds
$$

$$
\le c_* M_1 L^{q_*} \int_0^t (t-s)^{\gamma_0-\theta} e^{-\nu(t-s)} s^{-\mu q_*} ds
$$

$$
= c_* M_1 L^{q_*} t^{1+\gamma_0-\theta-\mu q_*} \int_0^1 (1-s)^{\gamma_0-\theta} e^{-\nu t(1-s)} s^{-\mu q_*} ds.
$$
 (2.17)

Choosing $\theta = \alpha$ in [\(2.17\)](#page-9-0), we infer from [\(2.10a\)](#page-7-4) and [\(2.13\)](#page-8-4) that

$$
||u(t)||_{\alpha} \le \frac{M_1}{2} e^{-\nu t} ||u^0||_{\alpha} + c_* M_1 L^{q_*} t^{1+\gamma_0-\alpha-\mu q_*} \int_0^1 (1-s)^{\gamma_0-\alpha} e^{-\nu t(1-s)} s^{-\mu q_*} ds
$$

$$
\le \frac{\varepsilon_0 M_1}{2} + c_* M_1 L^{q_*} \left(\sup_{r>0} r^{1+\gamma_0-\alpha-\mu q_*} e^{-\nu r} \right) B_{\mu q_*} ,
$$

and, recalling [\(2.11a\)](#page-8-5), we thus have

$$
||u(t)||_{\alpha} \leq \frac{\varepsilon}{\mathsf{e}_{\alpha,\beta}}, \qquad t \in (0, t_1).
$$
 (2.18)

Now, since

$$
||U_{\widehat{A}(u)}(t,0)u^0||_{\xi} \le \frac{M_1}{2}t^{-\mu}e^{-\nu t}||u^0||_{\alpha},
$$

due to [\(2.10a\)](#page-7-4), we obtain from [\(2.17\)](#page-9-0) (with $\theta = \xi$), using [\(2.13\)](#page-8-4), that

$$
t^{\mu} \|u(t)\|_{\xi} \leq \frac{M_1}{2} e^{-\nu t} \|u^0\|_{\alpha} + c_* M_1 L^{q_*} t^{1+\gamma_0+\mu-\xi-\mu q_*} \int_0^1 (1-s)^{\gamma_0-\xi} e^{-\nu t(1-s)} s^{-\mu q_*} ds
$$

$$
\leq \frac{\varepsilon_0 M_1}{2} + c_* M_1 L^{q_*} \Big(\sup_{r>0} r^{1+\gamma_0-\alpha-\mu q_*} e^{-\nu r} \Big) \mathsf{B}_{\mu(q_*-1)}.
$$

Therefore, [\(2.11b\)](#page-8-6) implies that

$$
||u||_{C_{\mu}((0,t],E_{\xi})} \le \frac{L}{2}, \qquad t \in (0,t_1).
$$
\n(2.19)

Next, let $0 \le s < t < t_1$. If $|t - s|^{\rho} \ge 4\varepsilon \ell/N$, then (2.18) yields

$$
||u(t) - u(s)||_{\beta} \le 2\varepsilon = \frac{2\varepsilon}{|t - s|^\rho} |t - s|^\rho \le \frac{N}{2\ell} |t - s|^\rho, \qquad |t - s|^\rho \ge \frac{4\varepsilon\ell}{N}.
$$

Conversely, consider $|t - s|$ ^{$\rho \leq 4\varepsilon \ell/N$ and note that}

$$
||u(t) - u(s)||_{\beta} \le ||U_{\widehat{A}(u)}(t,0)u^{0} - U_{\widehat{A}(u)}(s,0)u^{0}||_{\beta} + \int_{0}^{s} ||U_{\widehat{A}(u)}(t,\tau) - U_{\widehat{A}(u)}(s,\tau)||_{\mathcal{L}(E_{\gamma_{*}},E_{\beta})} ||\widehat{f}(u(\tau))||_{\gamma_{*}} d\tau + \int_{s}^{t} ||U_{\widehat{A}(u)}(t,\tau)||_{\mathcal{L}(E_{\gamma_{*}},E_{\beta})} ||\widehat{f}(u(\tau))||_{\gamma_{*}} d\tau =: I_{1} + I_{2} + I_{3}.
$$
 (2.21a)

From [\(2.10b\)](#page-7-6) it follows that

$$
I_1 \le M_2 \|u^0\|_{\alpha} (t-s)^{\alpha-\beta} \le \varepsilon_0 M_2 \left(\frac{4\varepsilon\ell}{N}\right)^{\frac{\alpha-\beta-\rho}{\rho}} (t-s)^{\rho},\tag{2.21b}
$$

while [\(2.10a\)](#page-7-4), [\(2.10c\)](#page-7-7), and [\(2.12\)](#page-8-3) entail

$$
I_2 \leq \int_0^s \|U_{\widehat{A}(u)}(t,s) - 1\|_{\mathcal{L}(E_{\alpha},E_{\beta})} \|U_{\widehat{A}(u)}(s,\tau)\|_{\mathcal{L}(E_{\gamma_*},E_{\alpha})} \|f(u(\tau))\|_{\gamma_*} d\tau
$$

$$
\leq c_* M_1 M_2 L^{q_*}(t-s)^{\alpha-\beta} \int_0^s (s-\tau)^{\gamma_0-\alpha} e^{-\nu(s-\tau)} \tau^{-\mu q_*} d\tau
$$

$$
= c_* M_1 M_2 L^{q_*}(t-s)^{\alpha-\beta} s^{1+\gamma_0-\alpha-\mu q_*} \int_0^1 (1-\tau)^{\gamma_0-\alpha} e^{-\nu s(1-\tau)} \tau^{-\mu q_*} d\tau
$$

and hence, using [\(2.13\)](#page-8-4),

$$
I_2 \le c_* c_0 M_1 M_2 L^{q_*} \mathsf{B}_{\mu q_*} \left(\frac{4\varepsilon \ell}{N}\right)^{\frac{\alpha-\beta-\rho}{\rho}} (t-s)^\rho \,. \tag{2.21c}
$$

Finally, from $(2.10a)$ and (2.12) we get

$$
I_3 \le c_* M_1 L^{q_*} \int_s^t (t-\tau)^{\gamma_0-\beta} e^{-\nu(t-\tau)} \tau^{-\mu q_*} d\tau
$$

$$
\le c_* M_1 L^{q_*} (t-s)^{\alpha-\beta} (t-s)^{1+\gamma_0-\alpha-\mu q_*} \int_0^1 (1-\tau)^{\gamma_0-\beta} e^{-\nu(t-s)(1-\tau)} \tau^{-\mu q_*} d\tau
$$

and therefore, invoking again [\(2.13\)](#page-8-4),

$$
I_3 \le c_* c_0 M_1 L^{q_*} \mathsf{B}_{\alpha-\beta+\mu q_*} \left(\frac{4\varepsilon \ell}{N}\right)^{\frac{\alpha-\beta-\rho}{\rho}} (t-s)^\rho \,. \tag{2.21d}
$$

Consequently, gathering $(2.20)-(2.21)$ $(2.20)-(2.21)$ and then using $(2.11c)$, we deduce that

$$
||u(t) - u(s)||_{\beta} \le \frac{N}{2\ell}(t - s)^{\rho}, \qquad 0 \le s \le t < t_1.
$$
 (2.22)

Summarizing, we obtain from the estimates [\(2.18\)](#page-9-1), [\(2.19\)](#page-9-4), and [\(2.22\)](#page-10-0) that $t_1 = t^+(v^0)$. In particular, the solution $v(\cdot; v^0) : [0, t^+(v^0)) \to E_\beta$ is uniformly continuous. Moreover, recalling [\(1.2d\)](#page-1-1) and [\(2.18\)](#page-9-1), and noticing from [\(2.19\)](#page-9-4) that

$$
||v(t; v0)||\xi \le ||v_*||\xi + Lt-\mu, \t t \in (0, t+(v0)),
$$

we also have

$$
\limsup_{t \nearrow t^+(v^0)} \|f(v(t; v^0))\|_0 < \infty
$$

while [\(2.3\)](#page-6-5) ensures that

$$
\liminf_{t \nearrow t^+(v^0)} \operatorname{dist}_{E_\beta}\left(v(t;v^0), \partial O_\beta\right) > 0.
$$

Consequently, Theorem [1.2](#page-2-0) implies that $t^+(v^0) = \infty$. In particular, the evolution system $U_{\hat{A}(u)}(t,s)$ is defined for all $0 \le s \le t < \infty$ and satisfies [\(2.10a\)](#page-7-4) for all $0 \le s \le t < \infty$.

(c) Stability. Set

$$
z(t) := \sup_{\tau \in (0,t]} (||u(\tau)||_{\alpha} + \tau^{\mu}||u(\tau)||_{\xi})e^{\omega \tau}, \qquad t > 0,
$$

and note from [\(2.19\)](#page-9-4) that

$$
\|u(\tau)\|^{q_*}_{\xi} \leq \tau^{-\mu q_*} e^{-\omega \tau} L^{q_*-1} z(t)\,, \qquad 0<\tau\leq t\,.
$$

Using the latter along with $(2.10a)$ $(2.10a)$ $(2.10a)$ and (2.12) , it follows analogously as before² when deriving (2.18) and [\(2.19\)](#page-9-4) that for $0 < \tau < t$ we have

$$
\begin{aligned} \left(\|u(\tau)\|_{\alpha} + \tau^{\mu} \|u(\tau)\|_{\xi} \right) e^{\omega \tau} \\ &\leq \frac{c_1}{2} \|u^0\|_{\alpha} + c_* M_1 L^{q_*-1} \left(\mathsf{B}_{\mu q_*} + \mathsf{B}_{\mu(q_*-1)} \right) \left(\sup_{r>0} r^{1+\gamma_0-\alpha-\mu q_*} e^{(\omega-\nu)r} \right) z(t) \,, \end{aligned}
$$

where $c_1 := 2M_1$. Invoking [\(2.11d\)](#page-8-8), we thus have

$$
(\|u(t)\|_{\alpha} + t^{\mu} \|u(t)\|_{\xi})e^{\omega t} \leq z(t) \leq c_1 \|u^0\|_{\alpha}, \qquad t > 0,
$$

and, since $v(t; v^0) = u(t) + v_*$ and $v^0 = u^0 + v_*$, the proof of Theorem [1.3](#page-3-0) is complete.

Concerning the instability result Theorem [1.8,](#page-5-2) we note the following:

Proof of Theorem [1.8.](#page-5-2) The proof is identically to that of [\[13,](#page-18-4) Theorem 1.4].

3. Semilinear Problem: Proof of Theorem [1.6](#page-4-3)

The proof of Theorem [1.6](#page-4-3) follows along the lines of the proof of Theorem [1.3.](#page-3-0) We thus only sketch it.

Proof of Theorem [1.6.](#page-4-3) Let the assumptions of Theorem [1.6](#page-4-3) hold. To start with, we note that since by assumption $\alpha \geq \alpha^*_{\text{crit}}$, we have

$$
\mu := \xi - \alpha \le \frac{1 + \gamma_* - \alpha}{q_*},
$$

the inequality turning into a strict inequality if $\alpha > \alpha^*_{\text{crit}}$, that is, if $\gamma_* \in (0, \gamma)$ or $\alpha = \gamma$ by assumption. If $\gamma_* \in (0, \gamma)$, we may thus choose $\gamma_0 \in (0, \gamma_*)$ with $\mu q_* < 1 + \gamma_0 - \alpha$, while we set $\gamma_0 = \gamma_*$ for $\gamma_* \in \{0, \gamma\}$. We then have in any case that

 $\mu q_* \leq 1 + \gamma_0 - \alpha$. Let $R \in (0, r_*)$, see [\(1.15d\)](#page-4-6), be chosen such that $\mathbb{B}_{E_\alpha}(v_*, R) \subset O_\alpha$ and fix $\omega_0 > \zeta > \omega > 0$.

Since $\mu q_* < 1$ and $\mu q_* \leq 1 + \gamma_0 - \alpha = 1 + \gamma_0 + \mu - \xi$, the constant

$$
c_0:=1+\big(\mathsf{B}_{\mu q_*}+\mathsf{B}_{\mu(q_*-1)}\big)\Big(\sup_{r>0}r^{1+\gamma_0-\alpha-\mu q_*}e^{(\omega-\zeta)r}\Big)\,,
$$

with $B_{\theta} := B(\theta, 1 - \mu q_*)$ for $\theta > 0$, is well-defined. Assumption [\(1.15e\)](#page-4-7) together with [\[4,](#page-18-9) II.Lemma 5.1.3] ensures the existence of a constant $M \geq 1$ such that the strongly continuous analytic semigroup $(e^{tA})_{t\geq 0}$ generated by A satisfies

$$
||e^{t\mathbb{A}}||_{\mathcal{L}(E_{\theta})} + t^{\theta - \vartheta_0}||e^{t\mathbb{A}}||_{\mathcal{L}(E_{\vartheta}, E_{\theta})} \le \frac{M}{4c_0}e^{-\zeta t}, \qquad t > 0,
$$
\n(3.1)

for $0 \le \vartheta_0 \le \vartheta \le 1$ with $\vartheta_0 < \vartheta$ if $0 < \vartheta < \theta < 1$ or, thanks to (1.12) , for $(\vartheta, \theta) \in \{(\alpha, \xi), (\gamma, \alpha), (\gamma, \xi)\}\$ with $\vartheta_0 = \vartheta$. We may then fix $L \in (0, R)$ such that

$$
c_* M L^{q_*-1} \le 1, \tag{3.2}
$$

where $c_* > 0$ stems from [\(1.15d\)](#page-4-6). We now set

$$
\widehat{f}(w) := f(w + v_*) - f(v_*) - \partial f(v_*)w, \qquad w \in \widehat{O}_{\xi} := O_{\xi} - v_*\,,
$$

and note, for $v^0 \in \overline{\mathbb{B}}_{E_\alpha}(v_*, L/M)$, that $u := v(\cdot; v^0) - v_*$ is a strong solution to the evolution problem

$$
u' = \mathbb{A}u + \hat{f}(u), \quad t > 0, \qquad u(0) = u^0 := v^0 - v_* \in \overline{\mathbb{B}}_{E_\alpha}(0, L/M). \tag{3.3}
$$

Set

$$
t_1 := \sup \{ t \in (0, t^+(v^0)) : ||u||_{C_{\mu}((0,t], E_{\xi})} < L \text{ and } ||u||_{C([0,t], E_{\alpha})} < R \} > 0
$$

²The exponential $e^{-\nu r}$ has now to be replaced everywhere by $e^{(\omega-\nu)r}$.

noticing $||u||_{C_{\mu}((0,t],E_{\xi})} \to 0$ as $t \to 0$ and $||u(0)||_{\alpha} \leq L < R$. Let $t \in (0,t_1)$. Then, $||u(s)||_{\xi} \leq s^{-\mu}L$ for all $s \in (0, t]$ and u satisfies in view of (1.13) and (3.3) the variation-of-constants formula

$$
u(\tau) = e^{\tau \mathbb{A}} u^0 + \int_0^{\tau} e^{(\tau - s)\mathbb{A}} \hat{f}(u(s)) ds, \qquad 0 \le \tau \le t,
$$

with

$$
\|\widehat{f}(u(s))\|_{\gamma_*} \le c_* \|u(s)\|_{\xi}^{q_*} \le c_* L^{q_*} s^{-\mu q_*}, \qquad s \in (0, t],
$$
\n(3.4)

see $(1.15d)$. The estimates (2.13) , (3.1) , and (3.4) lead to

$$
||u(t)||_{\alpha} \leq ||e^{t\mathbb{A}}||_{\mathcal{L}(E_{\alpha})} ||u^{0}||_{\alpha} + \int_{0}^{t} ||e^{(t-s)\mathbb{A}}||_{\mathcal{L}(E_{\gamma_{*}}, E_{\alpha})} ||\hat{f}(u(s))||_{E_{\gamma_{*}}} ds
$$

$$
\leq \frac{M}{4} ||u^{0}||_{\alpha} + \frac{c_{*}ML^{q_{*}}}{4},
$$
 (3.5)

and similarly

$$
t^{\mu} \|u(t)\|_{\xi} \leq t^{\mu} \|e^{t\mathbb{A}}\|_{\mathcal{L}(E_{\alpha}, E_{\xi})} \|u^{0}\|_{\alpha} + t^{\mu} \int_{0}^{t} \|e^{(t-s)\mathbb{A}}\|_{\mathcal{L}(E_{\gamma_{*}}, E_{\xi})} \|\widehat{f}(u(s))\|_{E_{\gamma_{*}}} ds
$$

$$
\leq \frac{M}{4} \|u^{0}\|_{\alpha} + \frac{c_{*}ML^{q_{*}}}{4}.
$$
 (3.6)

It now follows from (3.2) , (3.5) , and (3.6) that

$$
||u||_{C_{\mu}((0,t],E_{\xi})} \leq \frac{L}{2}
$$
 and $||u||_{C([0,t],E_{\alpha})} \leq \frac{R}{2}$

for each $t \in (0, t_1)$, hence $t_1 = t^+(v^0)$ and, together with [\(1.11e\)](#page-3-5) and [\(1.14\)](#page-4-1), we conclude, similarly as in the proof of Theorem [1.3,](#page-3-0) that $t^+(v^0) = \infty$.

Define now

$$
z(t) := \sup_{\tau \in (0,t]} (||u(\tau)||_{\alpha} + \tau^{\mu}||u(\tau)||_{\xi})e^{\omega \tau}, \qquad t > 0.
$$

Given $0 < \tau < t$, we then have

$$
||u(\tau)||_{\xi}^{q_*} \le L^{q_*-1} z(t) \,\tau^{-\mu q_*} \, e^{-\omega \tau} \,,
$$

and together with $(1.15d)$, (3.1) , and the latter estimate we deduce, analogously to $(3.5)-(3.6)$ $(3.5)-(3.6)$, that

$$
z(t) \le \frac{M}{2} \|u^0\|_{\alpha} + \frac{c_* M L^{q_*-1}}{2} z(t)
$$

and therefore, by the choice of L from (3.2) ,

$$
z(t) \le M \|u^0\|_{\alpha}, \qquad t > 0,
$$

that is,

$$
||u(t)||_{\alpha} + t^{\mu}||u(t)||_{\xi} \leq M e^{-\omega t} ||u^{0}||_{\alpha}, \qquad t > 0.
$$

This completes the proof.

4. Applications

In this section we apply our theory to various semilinear and quasilinear evolution problems of parabolic type, examining both critical and non-critical regimes, to explore the stability properties of their equilibrium solutions. The applications include quasilinear problems with quadratic semilinearities (see Example [4.1\)](#page-13-1), a parabolic-parabolic chemotaxis system (Example [4.2\)](#page-13-0), and a quasilinear evolution equation in critical spaces exhibiting scaling invariance (Example [4.3\)](#page-16-0).

4.1. Quaslilinear Problems with Quadratic Semilinearity. To give a first flavor of our results we consider for densely embedded Banach spaces $E_1 \hookrightarrow E_0$ and complex interpolation spaces $E_\theta := [E_0, E_1]_\theta$ with $\theta \in (0, 1)$, the quasilinear problem

$$
u' = A(u)u + Q(u, u), \quad t > 0, \qquad u(0) = u^0,
$$
\n(4.1)

with

$$
A \in C^{1-}(E_{\beta}, \mathcal{H}(E_1, E_0))
$$
\n
$$
(4.2a)
$$

and a quadratic bilinear term

$$
Q \in \mathcal{L}^2(E_{\xi}, E_{\gamma}). \tag{4.2b}
$$

We are interested in the stability properties of the equilibrium solution $u_* = 0 \in E_1$ to [\(4.1\)](#page-13-2). In the context of [\(4.1\)](#page-13-2), it is appropriate to choose

$$
0 < \gamma = \gamma_* < \beta < \xi < 1 \qquad \text{and} \qquad 2\xi - 1 - \gamma \le \alpha \in (\beta, \xi), \tag{4.2c}
$$

noticing that $\alpha_{\rm crit} = \alpha_{\rm crit}^* = 2\xi - 1 - \gamma$. Then, $f : E_{\xi} \to E_{\gamma}$ with $f(u) := Q(u, u)$ satisfies [\(1.4\)](#page-1-2) and [\(1.8d\)](#page-2-9), both with $q = q_* = 2$. Consequently, we obtain from Remark [1.1,](#page-1-3) Theorem [1.3,](#page-3-0) Theorem [1.6,](#page-4-3) and Theorem [1.8:](#page-5-2)

Theorem 4.1. Assume [\(4.2\)](#page-13-3). If $s(A(0)) < 0$, then $u_* = 0$ is asymptotically exponentially stable in E_α for the Cauchy problem (4.1) . If A is Fréchet differentiable with Lipschitz continuous derivative and

$$
\sigma_+(A(0)) := \{\lambda \in \sigma(A(0)) : \text{Re}\,\lambda > 0\} \neq \emptyset, \qquad \inf\{\text{Re}\,\lambda : \lambda \in \sigma_+(A(0))\} > 0,
$$

then $u_* = 0$ is unstable in E_α for the Cauchy problem [\(4.1\)](#page-13-2). In the semilinear case when A is independent of u, the same results hold if $(4.2c)$ is replaced by

$$
0\leq \gamma < \xi \leq 1\,,\quad (\gamma,\xi)\neq (0,1)\,,
$$

and either

$$
2\xi - 1 - \gamma < \alpha \in [\gamma, \xi) \qquad \text{or} \qquad 2\xi - 1 - \gamma = \alpha \in (\gamma, \xi).
$$

Of course, as pointed out in Remark [1.1,](#page-1-3) also other interpolation functors than the complex functor $[\cdot, \cdot]_\theta$ (such as real or continuous interpolation functors) can be considered.

In the critical case $2\xi - 1 - \gamma = \alpha \in (\gamma, \xi)$, Theorem [4.1](#page-13-5) has been established previously in the particular context of a semilinear asymptotic model for atmospheric flows describing morning glory clouds, see [\[12,](#page-18-7) Theorem 4.1]. For similar results for semilinear parabolic problems with general superlinear nonlinearites we refer to [\[12,](#page-18-7) Corollary 1.4] and [\[12,](#page-18-7) Example 4.2]. We also refer to [\[17,](#page-18-10) Corollary 2.2] for a related exponential stability result in the semilinear case within the framework of maximal L_p -regularity.

4.2. A Parabolic-Parabolic Chemotaxis System. We present an application of the exponential stability result in the semilinear case, cf. Theorem [1.6,](#page-4-3) and of the instability result in Theorem [1.8](#page-5-2) in the context of a parabolic-parabolic chemotaxis system with logistic source, see e.g. [\[5,](#page-18-11) [22\]](#page-18-12),

$$
\partial_t u = \text{div}(\nabla u - \chi u \nabla v) + \kappa u (1 - u), \qquad t > 0, \quad x \in \Omega,
$$
\n(4.3a)

$$
\partial_t v = \Delta v + u - v, \qquad t > 0, \quad x \in \Omega, \qquad (4.3b)
$$

where κ and χ are positive constants, subject to the initial conditions

$$
u(0, x) = u^{0}(x), \quad v(0, x) = v^{0}(x), \qquad x \in \Omega,
$$
\n(4.3c)

and homogeneous Neumann boundary conditions

$$
\partial_{\nu}u = \partial_{\nu}v = 0 \qquad \text{on } \partial\Omega. \tag{4.3d}
$$

The functions $u^0, v^0: \Omega \to \mathbb{R}$ are given and $\Omega \subset \mathbb{R}^n$, $n \geq 1$, is a smooth bounded domain with outward unit normal ν . Some constants from [\[5,](#page-18-11)[22\]](#page-18-12), which are qualitatively irrelevant to the analysis, have been replaced by 1.

We observe that problem (4.3) has exactly two constant equilibrium solutions, namely

$$
(u_1, v_1) = (0, 0)
$$
 and $(u_2, v_2) = (1, 1)$.

It is well-known that for sufficiently smooth non-negative initial data [\(4.3\)](#page-13-6) possesses a unique bounded global classical solution provided that Ω is convex and κ is sufficiently large, see e.g. [\[22\]](#page-18-12). We prove herein that [\(4.3\)](#page-13-6) is locally well-posed for more general initial data and that the equilibrium solution $(u_1, v_1) = (0, 0)$ is always unstable. Moreover, under certain restrictions of the coefficients χ and κ , we establish the exponential stability of the equilibrium $(u_2, v_2) = (1, 1)$.

Before presenting our result in Theorem [4.2,](#page-14-0) we define for $p \in (1,\infty)$ the Banach spaces

$$
F_0 := L_p(\Omega)
$$
, $F_1 := W_{p,N}^2(\Omega) = H_{p,N}^2(\Omega) = \{ v \in H_p^2(\Omega) : \partial_\nu v = 0 \text{ on } \partial\Omega \},$

and set, see [\[3,](#page-18-13) §4],

$$
B_0 := \Delta_N := \Delta\big|_{W^2_{p,N}(\Omega)} \in \mathcal{H}\big(W^2_{p,N}(\Omega), L_p(\Omega)\big) .
$$

Let

$$
\big\{ (F_\theta, B_\theta) \,:\, -1 \leq \theta < \infty \big\}
$$

be the interpolation-extrapolation scale generated by (F_0, B_0) and the complex interpolation functor $[\cdot, \cdot]_\theta$ (see [\[3,](#page-18-13) §6] and [\[4,](#page-18-9) §V.1]). Then,

$$
B_{\theta} \in \mathcal{H}(F_{1+\theta}, F_{\theta}), \qquad -1 \le \theta < \infty, \tag{4.4}
$$

where

$$
F_{\theta} \doteq H_{p,N}^{2\theta}(\Omega) := \begin{cases} \{v \in H_p^{2\theta}(\Omega) : \partial_{\nu}v = 0 \text{ on } \partial\Omega\}, & 1 + \frac{1}{p} < 2\theta < 3 + \frac{1}{p}, \\ H_p^{2\theta}(\Omega), & -1 + \frac{1}{p} < 2\theta < 1 + \frac{1}{p}, \end{cases}
$$
(4.5)

see [\[3,](#page-18-13) Theorem 7.1; Equation (7.5)]^{[3](#page-14-1)} Moreover, since $\Delta_N - 1$ has bounded imaginary powers, see e.g. from $[4, III.$ Examples $4.7.3$ (d), we infer from $[3,$ Remarks 6.1 (d) that

$$
[F_{\beta}, F_{\alpha}]_{\theta} \doteq F_{(1-\theta)\beta+\theta\alpha}, \qquad -1 \leq \beta < \alpha, \quad \theta \in (0,1). \tag{4.6}
$$

Theorem [4.2](#page-14-0) below provides the aforementioned stability result for problem [\(4.3\)](#page-13-6), which, despite the quasilinear term div $(u\nabla v)$ in [\(4.3a\)](#page-13-7), is treated as a semilinear evolution problem in the subcritical regime when $\alpha_{\rm crit} < \alpha$, see [\(1.11d\)](#page-3-6).

Theorem 4.2. Fix $\kappa, \chi \in (0, \infty)$, $n \geq 1$, $p \in (1, \infty)$ with $p > n/2$. Then, [\(4.3\)](#page-13-6) generates a semiflow on $L_p(\Omega) \times H^1_{2p}(\Omega)$. Moreover, the following hold:

(i) If the maximal existence time $t^+ \in (0,\infty]$ of the solution (u, v) to (4.3) is finite, then

$$
\limsup_{t \nearrow t^+} \|u(t)\|_{L_p} = \infty.
$$

- (ii) The equilibrium solution $(u_1, v_1) = (0, 0)$ is unstable in $L_p(\Omega) \times H^1_{2p}(\Omega)$.
- (iii) If $\chi \leq 2$ and $\kappa > 1/4$, then $(u_2, v_2) = (1, 1)$ is exponentially stable in $L_p(\Omega) \times H^1_{2p}(\Omega)$. More precisely, given

$$
0 < 2\varepsilon < \min\left\{1 - \frac{1}{p}, \, 1 - \frac{n}{2p}\right\} \qquad \text{and} \qquad \mu > \frac{1 - \varepsilon}{2},
$$

there exist constants $r, \omega \in (0,1)$ and $M \geq 1$ such that for all initial data $(u_0, v_0) \in L_p(\Omega) \times H^1_{2p}(\Omega)$ with $\|(u_0, v_0)\|_{L_p \times H_{2p}^1} \leq r$, the solution (u, v) to (4.3) is globally defined and

$$
||(u(t)-1,v(t)-1)||_{L_p\times H^1_{2p}}+t^{\mu}||(u(t)-1,v(t)-1)||_{H^{1-\varepsilon}_p\times H^{2-\varepsilon}_{2p}}\leq Me^{-\omega t}||(u_0,v_0)||_{L_p\times H^1_{2p}}, \qquad t>0. \tag{4.7}
$$

³This property is stated in [\[3\]](#page-18-13) for $-1+\frac{1}{p} < 2\theta \le 2$. However, since $(1-\Delta_N)^{-1} \in \mathcal{L}(H_p^{2\theta-2}(\Omega), H_p^{2\theta}(\Omega))$ for $2 < 2\theta < 3+1/p$, see [\[21,](#page-18-14) Theorem 5.5.1], we obtain the full range in [\(4.5\)](#page-14-2).

Proof. The local well-posedness of (4.3) (together with (i)) has been established in [\[14,](#page-18-8) Theorem 5.1] in a slightly more general context and we therefore only sketch the proof of this result. Set

$$
E_0 := H_{p,N}^{-2\varepsilon}(\Omega) \times H_{2p,N}^{1-2\varepsilon}(\Omega) , \qquad E_1 := H_{p,N}^{2-2\varepsilon}(\Omega) \times H_{2p,N}^{3-2\varepsilon}(\Omega) ,
$$

so that, by $(4.5)-(4.6)$ $(4.5)-(4.6)$,

$$
E_{\theta} = H_{p,N}^{2\theta - 2\varepsilon}(\Omega) \times H_{q,N}^{1+2\theta - 2\varepsilon}(\Omega), \qquad 2\theta \in [0,2] \setminus \{2\varepsilon + 1 + 1/p, 2\varepsilon + 1/2p\}.
$$

Choosing

$$
q = q_* = 2
$$
 and $0 < \gamma = \gamma_* := \frac{\varepsilon}{3} < \alpha := \varepsilon < \xi = \frac{1+\varepsilon}{2} < 1$,

we have

$$
\alpha_{\rm crit} = \alpha_{\rm crit}^* = 2\xi - 1 - \gamma = \frac{2\varepsilon}{3} < \alpha \in (\gamma, \xi),
$$

see [\(1.11c\)](#page-3-7), and

$$
E_{\xi} = H_{p,N}^{1-\varepsilon}(\Omega) \times H_{2p,N}^{2-\varepsilon}(\Omega) \hookrightarrow E_{\alpha} = L_p(\Omega) \times H_{2p}^1(\Omega) \hookrightarrow E_{\gamma} = H_{p,N}^{-4\varepsilon/3}(\Omega) \times H_{2p,N}^{1-4\varepsilon/3}(\Omega).
$$

Since $H_{p,N}^{2-2\varepsilon}(\Omega) \hookrightarrow H_{2p,N}^{1-2\varepsilon}(\Omega)$, we obtain from [\[4,](#page-18-9) I. Theorem 1.6.1] and [\(4.4\)](#page-14-4)-[\(4.5\)](#page-14-2) that

$$
A:=\begin{pmatrix} \Delta+\kappa & 0 \\ 1 & \Delta-1 \end{pmatrix} \in \mathcal{H}(E_1,E_0)\,.
$$

Let $f: E_{\xi} \to E_{\gamma}$ be given by

$$
f(w) := -(\chi \operatorname{div} (u \nabla v) + \kappa u^2, 0), \qquad w = (u, v) \in E_{\xi}.
$$

Using the continuity of the multiplications

$$
H_{p,N}^{1-\varepsilon}(\Omega) \bullet H_{2p,N}^{1-\varepsilon}(\Omega) \longrightarrow H_{p,N}^{1-4\varepsilon/3}(\Omega) \quad \text{and} \quad H_{p,N}^{1-\varepsilon}(\Omega) \bullet H_{p,N}^{1-\varepsilon}(\Omega) \longrightarrow L_p(\Omega),
$$

see [\[2,](#page-18-15) Theorem 4.1], it is not difficult to conclude that there is a constant $C > 0$ such that

$$
||f(w) - f(\bar{w})||_{E_{\gamma}} \le C[||w||_{E_{\xi}} + ||\bar{w}||_{E_{\xi}}]||w - \bar{w}||_{E_{\xi}}, \qquad w, \, \bar{w} \in E_{\xi}.
$$
\n(4.8)

The local well-posedness of [\(4.3\)](#page-13-6) and the blow-up criterion (i) follow by using [\[14,](#page-18-8) Theorem 1.2] as in the proof of [\[14,](#page-18-8) Theorem 5.1].

In order to address the stability properties of the equilibria $(0,0)$ and $(1,1)$ we note that $f \in C^{2-}(E_{\xi}, E_{\gamma})$ with

$$
\partial f(\bar{w})w = -(\chi \text{div}\big(u\nabla \bar{v} + \bar{u}\nabla v) + 2\kappa u\bar{u}, 0\big), \qquad w = (u, v), \ \bar{w} = (\bar{u}, \bar{v}) \in E_{\xi}.
$$

Moreover, since $A + \partial f(w) \in \mathcal{H}(E_1, E_0)$ for all $w \in E_{\xi}$ with compact embedding $E_1 \hookrightarrow E_0$, the spectrum of $A + \partial f(w)$ consists entirely of isolated eigenvalues with finite algebraic multiplicities [\[9,](#page-18-16) Theorem III.6.29].

For $w_1 = (0,0)$ we have $\partial f(w_1) = 0$ and the linearization $A + \partial f(w_1) = A$ has the positive eigenvalue $\lambda = \kappa$ (with a constant eigenvector). Recalling Remark [1.1](#page-1-3) and [\(4.6\)](#page-14-3), we are in a position to apply Theorem [1.8](#page-5-2) and deduce that $(0, 0)$ is an unstable equilibrium for (4.3) .

For $w_2 = (1, 1)$ we have

$$
\partial f(w_2)w = \big(-\chi \Delta v - 2\kappa u, 0\big), \qquad w = (u, v) \in E_{\xi},
$$

hence [\(4.8\)](#page-15-0) (with $\bar{w}=0$) implies that

$$
|| f(w + w_2) - f(w_2) - \partial f(w_2)w||_{\gamma} = || f(w)||_{\gamma} \le C||w||_{\xi}^2, \qquad w \in E_{\xi}.
$$

Thus, the assumptions [\(1.15a\)](#page-4-9)-[\(1.15d\)](#page-4-6) are all satisfied within this framework.

It remains to verify that [\(1.15e\)](#page-4-7) is also satisfied in the context of [\(4.3\)](#page-13-6). Let thus $\lambda = \text{Re }\lambda + i \text{Im }\lambda \in \mathbb{C}$ be an eigenvalue of $A + \partial f(w_2)$ with eigenvector $0 \neq w = (u, v) \in H_N^2(\overline{\Omega})^2$ by elliptic regularity, see e.g. [\[1,](#page-18-17) Theorem 15.2]. It then holds

$$
\Delta u - \chi \Delta v - \kappa u = \lambda u \quad \text{in } \Omega \,, \tag{4.9}
$$

$$
\Delta v + u - v = \lambda v \quad \text{in } \Omega \,. \tag{4.10}
$$

Let $u = u_1 + iu_2$ and $v = v_1 + iv_2$. Testing the real part of [\(4.9\)](#page-16-1) with u_1 and the imaginary part with u_2 and proceeding in the same way with (4.10) (where we test with v_1 and v_2 , respectively), we arrive at

$$
-\sum_{i=1}^{2} \int_{\Omega} \left[\left(|\nabla u_i|^2 - \chi \nabla u_i \cdot \nabla v_i + |\nabla v_i|^2 \right) + \left(\kappa |u_i|^2 - u_i v_i + v_i^2 \right) \right] dx = \text{Re } \lambda \int_{\Omega} (|u|^2 + |v|^2) dx.
$$

Since $\chi \le 2$ an $\kappa > 1/4$, Young's inequality and the observation that $v \ne 0$ (otherwise $w = 0$ by [\(4.10\)](#page-16-2)) leads us now to

$$
\operatorname{Re}\lambda\int_{\Omega}(|u|^2+|v|^2)\,\mathrm{d}x\le\left(\frac{1}{4\kappa}-1\right)\int_{\Omega}|v|^2\,\mathrm{d}x<0,
$$

hence $\text{Re } \lambda < 0$, which proves [\(1.15e\)](#page-4-7). Assertion (iii) is now a direct consequence of Theorem [1.6.](#page-4-3)

4.3. A Quasilinear Problem with Scaling Invariance. We shall apply Theorem [1.3](#page-3-0) in the context of a quasilinear evolution equation from [\[17,](#page-18-10) [20\]](#page-18-18):

$$
\partial_t u = \text{div}(a(u)\nabla u) + |\nabla u|^\kappa \qquad \text{in } \Omega \,,\ t > 0 \,, \tag{4.11a}
$$

subject to homogeneous Dirichlet boundary conditions

$$
u = 0 \qquad \text{on } \partial\Omega \,, \quad t > 0 \,, \tag{4.11b}
$$

and the initial condition

$$
u(0) = u_0, \t\t(4.11c)
$$

where $\kappa > 3$, $u_0 : \Omega \to \mathbb{R}$ is a given function, and $\Omega \subset \mathbb{R}^n$ with $n \geq 1$ is a smooth bounded domain. In order to recast [\(4.11\)](#page-16-3) in an appropriate framework we set for $p \in (1,\infty)$

$$
H_{p,D}^{2\theta}(\Omega):=\left\{\begin{array}{ll} \{v\in H_{p}^{2\theta}(\Omega)\,:\,v=0\,\,{\rm on}\,\,\partial\Omega\}\,,&\frac{1}{p}<2\theta\leq 2\,,\\ H_{p}^{2\theta}(\Omega)\,,&-2+\frac{1}{p}<2\theta<\frac{1}{p}\,. \end{array}\right.
$$

As observed in [\[12,](#page-18-7)[17\]](#page-18-10), the space $H_{p,D}^{s_c}(\Omega)$ with

$$
s_c:=\frac{n}{p}+\frac{\kappa-2}{\kappa-1}
$$

can be identified (via a scaling invariance argument) as a critical space for [\(4.11\)](#page-16-3). The next results establishes the exponential stability of the zero solution to [\(4.11\)](#page-16-3) in this critical space.

Theorem 4.3. Let $\kappa > 3$ and let $a \in C^1(\mathbb{R})$ be a strictly positive function with uniformly Lipschitz continuous derivative. We choose $p \in (2n, (\kappa - 1)n)$ with $p \neq (n - 1)(\kappa - 1)$ and $\tau \in (0, 1)$ such that

$$
\frac{1}{2} < 2\tau < 1 - \frac{n}{p}
$$

and set

$$
0 < \bar{s} := 2\tau + \frac{n}{p} < s_c < s := 1 + \frac{n(\kappa - 1)}{p\kappa} < 2 - 2\tau \,,
$$

as well as

$$
\mu := \frac{1}{2(\kappa - 1)} - \frac{n}{2p\kappa} \in (0, 1).
$$

Then, [\(4.11\)](#page-16-3) is locally well-posed in $H^{s_c}_{p,D}(\Omega)$. Moreover, there exist constants $r, \omega \in (0,1)$ and $M \ge 1$ such that for all $||u_0||_{H^{s_c}_{p,D}} \leq r$ the solution $u = u(\cdot; u_0)$ to [\(4.11\)](#page-16-3) is globally defined and

$$
||u(t)||_{H_p^{s_c}} + t^{\mu}||u(t)||_{H_p^s} \le M e^{-\omega t} ||u_0||_{H_p^{s_c}}, \qquad t > 0.
$$
\n(4.12)

Proof. The local well-posedness result follows by arguing as in [\[12,](#page-18-7) Example 3], where homogeneous Neumann conditions were considered instead. Therefore we only sketch the proof of the local well-posedness result. Set

$$
F_0 := L_p(\Omega), \qquad F_1 := W_{p,D}^2(\Omega) = H_{p,D}^2(\Omega),
$$

and note from [\[3,](#page-18-13) §4] that

$$
B_0 := \Delta_D := \Delta|_{W^2_{p,D}(\Omega)} \in \mathcal{H}(W^2_{p,D}(\Omega), L_p(\Omega))
$$

Let further

$$
\{(F_{\theta}, B_{\theta}) : -1 \leq \theta < \infty\}
$$

be the interpolation-extrapolation scale generated by (F_0, B_0) and the complex interpolation functor $[\cdot, \cdot]_\theta$ (see [\[3,](#page-18-13) §6] and [\[4,](#page-18-9) §V.1]), that is,

$$
B_{\theta} \in \mathcal{H}(F_{1+\theta}, F_{\theta}), \quad -1 \le \theta < \infty, \tag{4.13}
$$

.

with (see $[3,$ Theorem 7.1; Equation (7.5)))

$$
F_{\theta} \doteq H_{p,D}^{2\theta}(\Omega), \qquad 2\theta \in \left(-2 + \frac{1}{p}, 2\right] \setminus \left\{\frac{1}{p}\right\}.
$$
\n
$$
(4.14)
$$

Moreover, since Δ_D has bounded imaginary powers (see [\[4,](#page-18-9) III. Examples 4.7.3 (d)]), we infer from [\[3,](#page-18-13) Remarks 6.1 (d)] that

$$
[F_{\beta}, F_{\alpha}]_{\theta} \doteq F_{(1-\theta)\beta + \theta\alpha}, \qquad -1 \leq \beta < \alpha, \quad \theta \in (0,1). \tag{4.15}
$$

With p and τ fixed in the statement, we set

$$
E_{\theta} := H_{p,D}^{2\theta - 2\tau}(\Omega) , \qquad 2\tau + \frac{1}{p} \neq 2\theta \in [0,2] ,
$$

and point out that $E_{\theta} = [E_0, E_1]_{\theta}$ and that none of the constants \bar{s} , s, and s_c is equal to $1/p$. Let

$$
q=q_*:=\kappa>3
$$

and

$$
0 < \gamma = \gamma_* := \tau < \beta := \tau + \frac{\bar{s}}{2} < \alpha := \alpha_{\text{crit}} < \xi := \tau + \frac{s}{2} < 1,
$$
\n(4.16)

where

$$
\alpha_{\rm crit} = \alpha_{\rm crit}^* := \frac{\kappa \xi - 1 - \gamma}{\kappa - 1} = \tau + \frac{s_c}{2} \in (\beta, \xi).
$$

Note that

$$
E_{\xi} = H_{p,D}^{s}(\Omega) \hookrightarrow E_{\alpha} = H_{p,D}^{s_c}(\Omega) \hookrightarrow E_{\beta} = H_{p,D}^{\overline{s}}(\Omega) \hookrightarrow E_{\gamma} = L_p(\Omega).
$$

We may now formulate (4.11) as the quasilinear evolution problem

$$
u' = A(u)u + f(u), \quad t > 0, \qquad u(0) = u_0,
$$
\n(4.17)

where $A: E_\beta \to \mathcal{L}(E_1, E_0)$ is defined by

$$
A(u)v := \operatorname{div}(a(u)\nabla v), \qquad v \in E_1, \quad u \in E_\beta,
$$

and $f: E_{\xi} \to E_{\gamma}$ is defined by

$$
f(u) := |\nabla u|^\kappa, \qquad u \in E_\xi.
$$

As shown in [\[12,](#page-18-7) Example 3],

$$
||f(u) - f(v)||_{E_{\gamma}} \le \kappa (||u||_{E_{\xi}}^{\kappa - 1} + ||v||_{E_{\xi}}^{\kappa - 1}) ||u - v||_{E_{\xi}}, \qquad u, v \in E_{\xi}, \qquad (4.18)
$$

and moreover $A \in C^{1-}(E_\beta, \mathcal{H}(E_1, E_0))$. The latter property together with [\(4.14\)](#page-17-0)-[\(4.16\)](#page-17-1) and [\(4.18\)](#page-17-2) enables us to apply Theorem [1.2](#page-2-0) to [\(4.17\)](#page-17-3) and deduce the local well-posedness of this problem in $H_{p,D}^{s_c}(\Omega)$.

We next verify [\(1.8\)](#page-2-5) for the stationary solution $v_* := 0 \in E_1$. To this end we first infer from [\(4.18\)](#page-17-2) that the map $f : E_{\xi} \to E_{\gamma}$ is Fréchet differentiable in 0 with $\partial f(0) = 0$ and

$$
||f(w)||_{\gamma} \leq \kappa ||w||_{\xi}^{\kappa}, \qquad w \in E_{\xi}.
$$

Since the spectral bound $s(\Delta_D) = \sup\{\text{Re }\lambda : \lambda \in \sigma(\Delta_D)\}\$ is negative and $v_* = 0$ (in which case the left hand side of $(1.8e)$ vanishes), assumptions (1.8) are satisfied. We are thus in a position to apply Theorem [1.3](#page-3-0) and establish in this way the exponential stability of the zero solution to [\(4.17\)](#page-17-3) in $H^{s_c}_{p,D}(\Omega)$, see [\(4.12\)](#page-17-4). \Box

REFERENCES

- [1] S. AGMON, A. DOUGLIS, AND L. NIRENBERG, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I., Comm. Pure Appl. Math., 12 (1959), pp. 623–727.
- [2] H. Amann, Multiplication in Sobolev and Besov spaces, in Nonlinear analysis, Sc. Norm. Super. di Pisa Quaderni, Scuola Norm. Sup., Pisa, 1991, pp. 27–50.
- [3] \longrightarrow , Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems, in Function spaces, differential operators and nonlinear analysis (Friedrichroda, 1992), vol. 133 of Teubner-Texte Math., Teubner, Stuttgart, 1993, p. 9–126.
- [4] , Linear and Quasilinear Parabolic Problems. Vol. I, vol. 89 of Monographs in Mathematics, Birkhäuser Boston, Inc., Boston, MA, 1995. Abstract linear theory.
- [5] X. Bai and M. Winkler, Equilibration in a fully parabolic two-species chemotaxis system with competitive kinetics, Indiana Univ. Math. J., 65 (2016), pp. 553–583.
- [6] G. DA PRATO AND A. LUNARDI, Stability, instability and center manifold theorem for fully nonlinear autonomous parabolic equations in Banach space, Arch. Rational Mech. Anal., 101 (1988), p. 115–141.
- [7] A.-K. DRANGEID, The principle of linearized stability for quasilinear parabolic evolution equations, Nonlinear Anal., 13 (1989), p. 1091–1113.
- [8] D. Guidetti, Convergence to a stationary state and stability for solutions of quasilinear parabolic equations, Ann. Mat. Pura Appl. (4), 151 (1988), p. 331–358.
- [9] T. Kato, Perturbation Theory for Linear Operators, Springer Verlag, Berlin, 1995.
- [10] A. Lunardi, Asymptotic exponential stability in quasilinear parabolic equations, Nonlinear Anal., 9 (1985), p. 563–586.
[11] _____, Analytic Semigroups and Optimal Reqularity in Parabolic Problems, Progress in Nonline $-$, Analytic Semigroups and Optimal Regularity in Parabolic Problems, Progress in Nonlinear Differential Equations and their Applications, 16, Birkhäuser Verlag, Basel, 1995.
- [12] B.-V. MATIOC, L. ROBERTI, AND CH. WALKER, Quasilinear parabolic equations with superlinear nonlinearities in critical spaces, Submitted, (Preprint 2024).
- [13] B.-V. MATIOC AND CH. WALKER, On the principle of linearized stability in interpolation spaces for quasilinear evolution equations, Monatsh. Math., 191 (2020), pp. 615–634.
- [14] $______________________\.______\._$ Mell-posedness of quasilinear parabolic equations in time-weighted spacess, Proc. Roy. Soc. Edinburgh Sect. A, (2023). Preprint.
- [15] M. POTIER-FERRY, The linearization principle for the stability of solutions of quasilinear parabolic equations. I, Arch. Rational Mech. Anal., 77 (1981), p. 301–320.
- [16] J. Prüss, Maximal regularity for evolution equations in Lp-spaces, Conf. Semin. Mat. Univ. Bari, (2002), p. 1–39 (2003).
- [17] J. PRÜSS, G. SIMONETT, AND M. WILKE, Critical spaces for quasilinear parabolic evolution equations and applications, J. Differential Equations, 264 (2018), p. 2028–2074.
- [18] J. PRÜSS, G. SIMONETT, AND R. ZACHER, On convergence of solutions to equilibria for quasilinear parabolic problems, J. Differential Equations, 246 (2009), p. 3902–3931.
- [19] , On normal stability for nonlinear parabolic equations, Discrete Contin. Dyn. Syst., (2009), p. 612–621.
- [20] P. Quittner and P. Souplet, Superlinear parabolic problems, Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks], Birkhäuser/Springer, Cham, second ed., 2019. Blow-up, global existence and steady states.
- [21] H. Triebel, Interpolation Theory, Function Spaces, Differential Operators, North-Holland, Amsterdam, 1978.
- [22] M. Winkler, Boundedness in the higher-dimensional parabolic-parabolic chemotaxis system with logistic source, Comm. Partial Differential Equations, 35 (2010), pp. 1516–1537.

Fakultät für Mathematik, Universität Regensburg, 93053 Regensburg, Deutschland. Email address: bogdan.matioc@ur.de

LEIBNIZ UNIVERSITÄT HANNOVER, INSTITUT FÜR ANGEWANDTE MATHEMATIK, WELFENGARTEN 1, 30167 HANNOVER, Germany

Email address: schmitz@ifam.uni-hannover.de

LEIBNIZ UNIVERSITÄT HANNOVER, INSTITUT FÜR ANGEWANDTE MATHEMATIK, WELFENGARTEN 1, 30167 HANNOVER, $\mathcal G_{\rm ERMANY}$

 $Email \;address\colon$ $\texttt{walker@ifam.uni-hannover.de}$