A FEW SHARP ESTIMATES OF HARMONIC FUNCTIONS WITH APPLICATIONS TO STEKLOV EIGENFUNCTIONS

XING WANG AND CHENG ZHANG

ABSTRACT. On smooth compact manifolds with smooth boundary, we first establish the sharp lower bounds for the restrictions of harmonic functions in terms of their frequency functions, by using a combination of microlocal analysis and frequency function techniques by Almgren [1] and Garofalo-Lin [21]. The lower bounds can be saturated by Steklov eigenfunctions on Euclidean balls and a family of symmetric warped product manifolds. Moreover, as in Sogge [35] and Taylor [38], we analyze the interior behavior of harmonic functions by constructing a parametrix for the Poisson integral operator and calculate its composition with the spectral cluster. By using microlocal analysis, we obtain several sharp estimates for the harmonic functions whose traces are quasimodes on the boundary. As applications, we establish the almost-orthogonality, bilinear estimates and transversal restriction estimates for Steklov eigenfunctions, and discuss the numerical approximation of harmonic functions.

1. Introduction

Let (Ω, h) be a connected smooth compact manifold with smooth boundary (M, g), where $\dim \Omega = n + 1 \geq 2$ and $h|_M = g$. The Laplace operators on Ω and M are denoted by Δ and Δ_g respectively. Let u be a harmonic function on Ω with boundary datum f, namely

(1.1)
$$\begin{cases} \Delta u = 0 \text{ on } \Omega, \\ u = f \text{ on } \partial \Omega = M. \end{cases}$$

Let \mathcal{H} be the harmonic extension operator (or Poisson integral operator) on Ω . Then $u = \mathcal{H}f$. Let $P \in OPS^1(M)$ be classical and self-adjoint with principal symbol $p(x,\xi) = |\xi|_{g(x)}$ on the boundary. Two important examples of P are $\sqrt{-\Delta_g}$ and the Dirichlet-to-Neumann (DtN) operator \mathcal{N} . If e_{λ} is an eigenfunction of \mathcal{N} associated with the eigenvalue λ , then its harmonic extension $u_{\lambda} = \mathcal{H}e_{\lambda}$ is called a Steklov eigenfunction. The Steklov eigenvalue problem has seen a surge of interest in the past few decades, and there are many open problems in this rapidly expanding area. For the historical background and recent developments, we recommend the survey papers by Girouard-Polterovich [22] and Colbois-Girouard-Gordon-Sher [16].

In this paper, we mainly use microlocal analysis to investigate the interior behavior of u with respect to the frequency of f. Specifically, we shall use two types of frequencies. The first type is the frequency function as in the seminal works of Almgren [1] and Garofalo-Lin [21]. The second type is the spectral frequency described by the associated eigenvalues in its spectral decomposition with respect to P. These two types of frequencies coincide when the boundary datum f is an eigenfunction of the Dirichlet-to-Neumann operator on M. We shall analyze them by using the microlocal analysis techniques in Sogge [35] and Taylor [38].

1

1.1. **Geometric quantities.** First, we introduce the geometric quantities that will be used to state our results. Let $d_h(\cdot,\cdot)$, $d_q(\cdot,\cdot)$ be the Riemannian distance functions on Ω and M respectively. Let

(1.2)
$$\Omega_t = \{ \omega \in \Omega : d_h(\omega, \partial \Omega) \ge t \}, \quad \Omega_t^{\mathsf{c}} = \Omega \setminus \Omega_t, \quad \Sigma_t = \partial \Omega_t.$$

Then t is the distance from Σ_t to the boundary.

For each $x \in M$, we have a unique inward geodesic $\gamma_x(t) \subset \Omega$ normal to M such that $\gamma_x(0) = x$, $|\nabla \gamma_x(t)| = 1$. Let $\phi(t, x) = \gamma_x(t)$. Let

(1.3)
$$\delta = \sup\{\tau > 0: \ \phi: [0,\tau] \times M \to \Omega \text{ is injective}\}.$$

Then for any $t < \delta$, Σ_t is a smooth hypersurface, and $\gamma_x(t)$ is normal to Σ_t for any $x \in M$.

By (1.3), we get a bijection $\phi:[0,\delta)\times M\to\Omega^{\mathsf{c}}_{\delta}$, so we can find a local coordinates system near the boundary adapted to the geodesics normal to boundary. Let $\phi^{-1}=(g_1,g_2)$ with $g_1:\Omega^{\mathsf{c}}_{\delta}\to M$ and $g_2:\Omega^{\mathsf{c}}_{\delta}\to[0,\delta)$. We choose a finite covering $M=\cup U_{\nu}$ and the corresponding coordinate maps $\kappa_{\nu}:U_{\nu}\to \tilde{U}_{\nu}\subset\mathbb{R}^n$. Let

$$\tilde{\kappa}_{\nu}: \Omega_{\nu} = \phi([0,\delta) \times U_{\nu}) \to \mathbb{R}^{n+1}, \quad p \mapsto (\kappa_{\nu}(g_2(p)), g_1(p)).$$

Under this setting, let $x = (x_1, x_2, ..., x_n)$ be local coordinates on some open set of M. Then for $t \in [0, \delta)$ and fixed $x, (t, x_1, x_2, ..., x_n)$ will be a geodesic curve starting from x and normal to both M and Σ_t , so ∂_t is the inward normal derivative of Σ_t . Specifically, the metric tensor of Ω can be written as

(1.4)
$$h = dt^2 + g_{ij}(t, x)dx_i dx_j = dt^2 + g_t,$$

where g_t is the metric on Σ_t , $g_0 = g$. Then the Laplace operator on Ω can be written as

(1.5)
$$\Delta = \partial_t^2 + a(t, x)\partial_t + \Delta_{g_t}$$

where

(1.6)
$$a(t,x) = \frac{1}{2} \partial_t \det(g_{ij}(t,x)) / \det(g_{ij}(t,x))$$

and Δ_{g_t} is the Laplace operator on Σ_t .

For each $t \in [0, \delta)$ and $x \in \Sigma_t$, we denote the Weingarten map with respect to the inward normal unit vectors of Σ_t as

$$(1.7) W_{t,x}: T_x \Sigma_t \to T_x \Sigma_t,$$

and $W_{t,x}^*: T_x^*\Sigma_t \to T_x^*\Sigma_t$ is its dual. Let $k_1(x) \leq ... \leq k_n(x)$ be the principal curvatures at x, and

(1.8)
$$\tilde{k}_t(x) = -\text{Tr}\mathcal{W}_{t,x} + \sup_{\xi \neq 0} \frac{\langle \mathcal{W}_{t,x}^* \xi, \xi \rangle_t}{\langle \xi, \xi \rangle_t} = -\sum_{i=1}^{n-1} k_j(x),$$

where the inner product $\langle \cdot, \cdot \rangle_t$ on $T_x^* \Sigma_t$ is defined as

$$\langle \xi, \eta \rangle_t = \sum_{i,j} g^{ij}(t, x) \xi_i \eta_j.$$

Let $|\xi|_{g_t(x)} = \sqrt{\langle \xi, \xi \rangle_t}$. Then we define $\Theta: [0, \delta) \to \mathbb{R}$ as

(1.9)
$$\Theta(t) = \sup_{x \in \Sigma_t} \tilde{k}_t(x) + \sup_{x \in \Sigma_t} \text{Tr} \mathcal{W}_{t,x},$$

and let

(1.10)
$$K(t) = \int_0^t e^{\int_0^s \Theta(\tau)d\tau} ds.$$

By Taylor's expansion, we have

$$K(t) = t + \frac{\Theta(0)}{2}t^2 + O(t^3).$$

Let

(1.11)
$$r(t) = \inf_{x \in M, \ \xi \neq 0} \frac{|\xi|_{g_t(x)}}{|\xi|_{g(x)}}, \quad G(t) = \int_0^t r(s)ds.$$

We have r(t) = 1 + O(t) and $G(t) = t + O(t^2)$. We will see that G(t) = K(t) when Ω is a Euclidean ball or the warped product manifold in Section 3.

1.2. Lower bounds. We first prove the lower bounds for the L^2 restrictions of harmonic functions.

Theorem 1. Fix any $0 < \delta_0 < \delta$. There is a constant C > 0 dependent on (Ω, h) and δ_0 , such that for any $0 \le t \le \delta_0$, and any $f \in L^2(M)$, we have

(1.12)
$$\|\mathcal{H}f\|_{L^2(\Sigma_t)} \ge Ce^{-\Lambda K(t)} \|f\|_{L^2(M)},$$

where the frequency $\Lambda = \int_{\Omega} |\nabla u|^2 / \int_{M} u^2$ with $u = \mathcal{H}f$, and $K(t) = t + O(t^2)$ is defined in (1.10).

The lower bounds are sharp in the sense that they can be saturated by Steklov eigenfunctions on Euclidean balls and a family of symmetric warped product manifolds, see Section 3. When $f \in L^2(M) \setminus H^{\frac{1}{2}}(M)$, we have $\Lambda = \infty$ and (1.12) is trivial in this case. Moreover, a simple consequence is the lower bound for the L^p norm with p > 2

(1.13)
$$\|\mathcal{H}f\|_{L^p(\Sigma_t)} \ge Ce^{-\Lambda K(t)} \|f\|_{L^2(M)}.$$

It is also sharp, since it can be saturated by Example 2 in Section 3.

As an application, using $\int_{\Omega} |\nabla u|^2 = \int_{M} u \partial_{\nu} u$, we get the following exponential lower bound for the Steklov eigenfunctions on smooth manifolds. As before, let e_{λ} be a DtN eigenfunction and $u_{\lambda} = \mathcal{H}e_{\lambda}$ be the Steklov eigenfunction.

Corollary 1. Fix any $0 < \delta_0 < \delta$. There is a constant C > 0 dependent on (Ω, h) and δ_0 , such that for any $0 \le t \le \delta_0$, we have

(1.14)
$$||u_{\lambda}||_{L^{2}(\Sigma_{t})} \geq Ce^{-\lambda K(t)} ||e_{\lambda}||_{L^{2}(M)}.$$

For real analytic Ω and M, Galkowski-Toth [20] have proved some lower bounds of this type. Indeed, it follows from [20, Theorem 1] that for any $\varepsilon > 0$, $\lambda > \lambda_0(\varepsilon)$ and $t \leq \varepsilon_0$, there is c > 0 such that

(1.15)
$$||u_{\lambda}||_{L^{2}(\Sigma_{t})} \gtrsim e^{-\varepsilon \lambda} e^{-(t+ct^{2})\lambda} ||e_{\lambda}||_{L^{2}(M)}.$$

In comparison, our estimate (1.14) is sharp and has no ε -loss. Moreover, we do not need the analyticity assumptions.

1.3. **Upper bounds.** From the examples of Steklov eigenfunctions (see Section 3), we naturally expect that any harmonic function should rapidly decay with respect to λt , where t is the distance to the boundary and λ is its "lowest spectral frequency". We shall give a rigorous justification for this observation.

Let $P \in OPS^1(M)$ be classical and self-adjoint with principal symbol $p(x,\xi) = |\xi|_{g(x)}$. Let $\mathbb{1}_{\geq \lambda}$ be the indicator function of $[\lambda, \infty)$. We shall prove the following interior decay property when the spectral frequency of the boundary datum f is bounded below by λ .

Theorem 2. Let $0 \le \delta_0 < \delta$, 0 < c < 1, N > 0 and $\lambda > 0$. Suppose that $f = \mathbb{1}_{\ge \lambda}(P)f$. Then for $1 and <math>0 \le t \le \delta_0$, we have

(1.16)
$$\|\mathcal{H}f\|_{L^p(\Sigma_t)} \leq C_1 e^{-c\lambda G(t)} \|f\|_{L^p(M)} + C_2 \lambda^{-N} \|f\|_{L^1(M)},$$
 where $C_1 = C_1(\delta_0, c, p), C_2 = C_2(\delta_0, c, N), \text{ and } G(t) = t + O(t^2) \text{ is defined in (1.11)}.$

The exponential term is essentially sharp, in the sense that G(t) = K(t) when Ω is a Euclidean ball or the warped product manifold in Section 3. The remainder term $O(\lambda^{-N})$ comes from the smoothing operators that are natural in the theory of pseudo-differential operators on smooth manifolds. The crux of the proof is the parametrix for the Poisson integral operator in Lemma 2. From the proof, we shall notice that (1.16) is valid for all $1 \le p \le \infty$ if the boundary datum f has spectral frequency $\approx \lambda$. Moreover, the proof also implies the derivative estimates. Indeed, if $f = 1 \ge \lambda(P) f$ then we have for 1

Remark 1. In comparison to Theorem 1, we remark that the upper bound of the form with C, c > 0 (1.18) $\|\mathcal{H}f\|_{L^2(\Sigma_t)} \leq Ce^{-c\Lambda t} \|f\|_{L^2(M)}$

cannot hold for general boundary datum f, where the frequency $\Lambda = \int_{\Omega} |\nabla u|^2 / \int_{M} u^2$ with $u = \mathcal{H}f$. For example, let $f = 1 + e_{\lambda}$ and $u = \mathcal{H}f = 1 + u_{\lambda}$, where e_{λ} is an L^2 normalized DtN eigenfunction. Then both u and u_{λ} have frequencies $\Lambda \approx \lambda$. However, when $\lambda t \gg 1$, we notice that $||u||_{L^2(\Sigma_t)} \approx 1$ shows no decay while $||u_{\lambda}||_{L^2(\Sigma_t)} \lesssim (1+\lambda t)^{-N}$ decays rapidly. This key observation suggests that the interior decay property of harmonic functions should only rely on the "lowest spectral frequency", as described in Theorem 2.

Recall Sogge's seminal work [34] on L^p estimates. If $\mathbb{1}_{(\lambda-1,\lambda]}$ is the indicator function of $(\lambda-1,\lambda]$, then

(1.19)
$$\|\mathbb{1}_{(\lambda-1,\lambda]}(P)f\|_{L^p(M)} \lesssim \lambda^{\sigma(p)} \|f\|_{L^2(M)}, \quad 2 \le p \le \infty,$$

where

(1.20)
$$\sigma(p) = \begin{cases} \frac{n-1}{2} \left(\frac{1}{2} - \frac{1}{p}\right), & 2 \le p < \frac{2(n+1)}{n-1} \\ \frac{n-1}{2} - \frac{n}{p}, & \frac{2(n+1)}{n-1} \le p \le \infty. \end{cases}$$

See [35, Theorem 5.1.1]. The L^{∞} bound is due to Hörmander [25]. The estimates are sharp and they can be saturated by some spherical harmonics on the standard sphere. So we have the following interior decay estimates for Steklov eigenfunctions on smooth manifolds with smooth boundary.

Corollary 2. Let $0 \le \delta_0 < \delta$, 0 < c < 1, N > 0 and $\lambda > 0$. Then for $2 \le p \le \infty$ and $t \in [0, \delta_0]$ we have

$$(1.21) ||u_{\lambda}||_{L^{p}(\Sigma_{t})} \lesssim (\lambda^{\sigma(p)} e^{-c\lambda G(t)} + \lambda^{-N}) ||e_{\lambda}||_{L^{2}(M)}, \ \forall N.$$

This implies

(1.22)
$$||u_{\lambda}||_{L^{p}(\Sigma_{t})} \lesssim \lambda^{\sigma(p)} (1 + \lambda t)^{-N} ||e_{\lambda}||_{L^{2}(M)}, \ \forall N.$$

Here $\sigma(p)$ is Sogge's exponent in (1.19).

Our estimates (1.21) and (1.22) are at least sharp for small t, by the sharpness of Sogge's estimates (1.19). We also have the derivative estimates for $2 \le p \le \infty$

(1.23)
$$\|\partial_{t,x}^{\alpha} u_{\lambda}\|_{L^{p}(\Sigma_{t})} \lesssim \lambda^{\sigma(p)+|\alpha|} (1+\lambda t)^{-N} \|e_{\lambda}\|_{L^{2}(M)}, \ \forall \alpha, \forall N.$$

The interior decay properties of Steklov eigenfunctions have been studied by Hislop-Lutzer [24], Polterovich-Sher-Toth [33], Galkowski-Toth [19], Di Cristo-Rondi [18], Daudé-Helffer-Nicoleau [17], Helffer-Kachmar [23], Levitin-Parnovski-Polterovich-Sher [31]. Now we go over the previous results and compare them with ours.

In 2001, Hislop-Lutzer [24] proved when the boundary is smooth, Steklov eigenfunctions decay super-polynomially into the interior, namely

(1.24)
$$||u_{\lambda}||_{H^{1}(K)} = O(\lambda^{-N}), \ \forall N,$$

on any compact subset K of the interior of Ω , and they also conjectured that the decay is actually of order $e^{-\lambda t}$ if Ω and M are real analytic. Under the analyticity assumptions, recently Polterovich-Sher-Toth [33] (dim $\Omega = 2$) and Galkowski-Toth [19] (dim $\Omega \geq 2$) established pointwise bounds with exponential decay. Specifically, Galkowski-Toth [19, Theorem 1] obtained

(1.25)
$$||u_{\lambda}||_{L^{\infty}(\Sigma_{t})} \lesssim \lambda^{\sigma(\infty) + \frac{1}{4}} e^{-\lambda(t - ct^{2})} ||u_{\lambda}||_{L^{2}(M)}.$$

The proof are based on analytic microlocal analysis and thus use the analyticity hypotheses in extremely strong ways. So one would perhaps believe that the exponential decay property cannot hold in the smooth case. For smooth manifolds, Helffer-Kachmar [23, Theorem 1] proved

(1.26)
$$||u_{\lambda}||_{L^{\infty}(\Sigma_{t})} \lesssim \lambda^{n-1} (1+\lambda t)^{-N} ||e_{\lambda}||_{L^{2}(M)}, \ \forall N.$$

Di Cristo-Rondi [18] considered the problems of interior decay for very general elliptic equations. The following estimates can be derived from [18, Theorem 3.3]

(1.27)
$$||u_{\lambda}||_{L^{2}(\Sigma_{t})} \lesssim (1+\lambda t)^{-\frac{1}{2}} ||e_{\lambda}||_{L^{2}(M)}.$$

In comparison, our results (1.21) and (1.22) are better than (1.25) when $t \lesssim \lambda^{-1} \log \lambda$, and we do not need the analyticity hypotheses. Moreover, our results are stronger than (1.26) and (1.27). By the maximum principle, our results imply (1.24).

1.4. Comparable interior and boundary norms. By Theorem 2, if the spectral frequency is bounded below by λ , i.e. $f = \mathbb{1}_{\geq \lambda}(P)f$, then we can integrate in t to get the following interior norm estimates for 1

(1.28)
$$\|\mathcal{H}f\|_{L^{p}(\Omega)} \lesssim \lambda^{-\frac{1}{p}} \|f\|_{L^{p}(M)}.$$

Next, we further prove that if the boundary datum f has spectral frequency $\approx \lambda$, then the interior norm of its harmonic extension $\mathcal{H}f$ is comparable to the norm of f on the boundary up to a factor of the frequency. As before, let $P \in OPS^1(M)$ be classical and self-adjoint with principal symbol $p(x,\xi) = |\xi|_{q(x)}$.

Theorem 3. Let $\lambda > 0$ and $\beta \in C_0^{\infty}(\mathbb{R}^+)$. Let $f_{\lambda} = \beta(P/\lambda)f$. Then for any $1 \leq p \leq \infty$, we have

(1.29)
$$\|\mathcal{H}f_{\lambda}\|_{L^{p}(\Omega)} \approx \lambda^{-\frac{1}{p}} \|f_{\lambda}\|_{L^{p}(M)},$$

i.e. there is a constant $C = C(\Omega) > 0$ such that

$$(1.30) C^{-1}\lambda^{-\frac{1}{p}} \|f_{\lambda}\|_{L^{p}(M)} \leq \|\mathcal{H}f_{\lambda}\|_{L^{p}(\Omega)} \leq C\lambda^{-\frac{1}{p}} \|f_{\lambda}\|_{L^{p}(M)}.$$

The power gain $\lambda^{-\frac{1}{p}}$ is due to the rapid decay property in λt , like $(1+\lambda t)^{-N}$. This result is sharp in the sense that it can be saturated by Steklov eigenfunctions on Euclidean balls. As a corollary, we have the following two-sided interior L^p estimates for Steklov eigenfunctions.

Corollary 3. For $1 \le p \le \infty$, we have

(1.31)
$$||u_{\lambda}||_{L^{p}(\Omega)} \approx \lambda^{-\frac{1}{p}} ||e_{\lambda}||_{L^{p}(M)}.$$

Motivated by the study of nodal sets, the upper bounds in (1.31) were first obtained in our joint work with Huang and Sire [28] for generalized Steklov eigenfunctions with rough potentials.

1.5. Almost-orthogonality, transversal restrictions and bilinear estimates. We denote the interior and boundary inner products as

$$\langle u, v \rangle_{\Omega} = \int_{\Omega} u \bar{v}$$
 and $\langle f, g \rangle_{M} = \int_{M} f \bar{g}$.

It is easy to see the gradients of Steklov eigenfunctions are orthogonal on Ω . Indeed,

$$\langle \nabla u_{\lambda}, \nabla u_{\mu} \rangle_{\Omega} = \langle \mathcal{N} e_{\lambda}, e_{\mu} \rangle_{M} = 0, \ \forall \lambda \neq \mu.$$

But for general Ω other than Euclidean balls, Steklov eigenfunctions are not orthogonal in $L^2(\Omega)$. However, we can show they are almost-orthogonal. Let $\chi \in \mathcal{S}(\mathbb{R})$ satisfy

supp
$$\hat{\chi} \subset (\frac{1}{2}, 1), \quad \chi(0) = 1.$$

As before, let $P \in OPS^1(M)$ be classical and self-adjoint with principal symbol $p(x,\xi) = |\xi|_{g(x)}$.

Theorem 4 (Almost-orthogonality). Let $v_{\lambda} = \mathcal{H} \circ \chi(P - \lambda)f$, $w_{\mu} = \mathcal{H} \circ \chi(P - \mu)g$. We have

$$(1.32) |\langle v_{\lambda}, w_{\mu} \rangle_{\Omega}| \lesssim (1 + |\lambda + \mu|)^{-1} (1 + |\lambda - \mu|)^{-N} ||f||_{L^{2}(M)} ||g||_{L^{2}(M)}, \forall N.$$

The crux of the proof is the kernel formula in Lemma 3. The question on the L^2 -orthogonality of Steklov eigenfunctions has been considered by Auchmuty [2], Auchmuty-Rivas [9], Cho-Rivas [15]. Specifically, Cho-Rivas [15] analyzed the problem on rectangles by the explicit formula for the inner products in $L^2(\Omega)$, and provide accompanying numerics. However, a complete description of the orthogonality in $L^2(\Omega)$ has not been made so far. We address this issue in the following corollary.

Corollary 4. For Steklov eigenfunctions u_{λ} and u_{μ} , we have

$$(1.33) |\langle u_{\lambda}, u_{\mu} \rangle_{\Omega}| \lesssim (1 + \lambda + \mu)^{-1} (1 + |\lambda - \mu|)^{-N} ||e_{\lambda}||_{L^{2}(M)} ||e_{\mu}||_{L^{2}(M)}, \quad \forall N.$$

Thus, by Theorem 3 we have for $\mu > \lambda > 1$

$$(1.34) |\langle u_{\lambda}, u_{\mu} \rangle_{\Omega}| \lesssim (\lambda/\mu)^{\frac{1}{2}} (1 + |\lambda - \mu|)^{-N} ||u_{\lambda}||_{L^{2}(\Omega)} ||u_{\mu}||_{L^{2}(\Omega)}, \quad \forall N.$$

This significantly improves the trivial bounds by Cauchy-Schwarz, and provide a rigorous justification for the almost-orthogonality of Steklov eigenfunctions in $L^2(\Omega)$.

Similarly, we can establish the bilinear estimates and transversal restriction estimates.

Theorem 5 (Bilinear estimates). Let $v_{\lambda} = \mathcal{H} \circ \chi(P - \lambda)f$, $w_{\mu} = \mathcal{H} \circ \chi(P - \mu)g$ with $\mu \geq \lambda \geq 1$. Then we have

$$||v_{\lambda}w_{\mu}||_{L^{2}(\Omega)} \lesssim \mu^{-\frac{1}{2}}B||f||_{L^{2}(M)}||g||_{L^{2}(M)},$$

where

(1.36)
$$B = \begin{cases} \lambda^{\frac{1}{4}}, & n = 2\\ \lambda^{\frac{1}{2}}\sqrt{\log \lambda}, & n = 3\\ \lambda^{\frac{n-2}{2}}, & n \ge 4. \end{cases}$$

Bilinear and multilinear eigenfunction estimates on closed manifolds have been proved by Burq-Gérard-Tzvetkov [11, 12]. Our result (1.35) shows that the same type of estimates still hold for harmonic functions. In comparison to the results in [11, 12], there is a power gain $\mu^{-\frac{1}{2}}$ that only depends on the relatively high frequency μ . This is due to the rapid decay property of harmonic functions. Moreover, the bound (1.35) is sharp in the sense that it can be saturated by Steklov eigenfunctions on Euclidean balls.

Corollary 5. If B is defined as in Theorem 6, then we have the bilinear estimates for Steklov eigenfunctions

$$||u_{\lambda}u_{\mu}||_{L^{2}(\Omega)} \lesssim \mu^{-\frac{1}{2}}B||e_{\lambda}||_{L^{2}(M)}||e_{\mu}||_{L^{2}(M)}.$$

Equivalently, by Theorem 3 we have for $\mu \geq \lambda \geq 1$

The bound $\lambda^{\frac{1}{2}}B$ is sharp on Euclidean balls and only depends on the relatively low frequency λ .

Theorem 6 (Transversal restriction estimates). Let $0 \le k \le n-1$. Let Σ be a (k+1)-dimensional submanifold in Ω , intersecting the boundary $\partial\Omega=M$ transversally, with $\partial\Sigma\subset\partial\Omega$. For $\lambda\geq 1$, let $v_{\lambda} = \mathcal{H} \circ \chi(P - \lambda)f$. Then we have

(1.39)
$$||v_{\lambda}||_{L^{p}(\Sigma)} \lesssim \lambda^{-\frac{1}{p}} A ||f||_{L^{2}(M)}, \quad 2 \leq p \leq \infty.$$

When n > 2,

- If $k \leq n-3$, then $A = \lambda^{\frac{n-1}{2} \frac{k}{p}}$ for $2 \leq p \leq \infty$ If k = n-2, then $A = \lambda^{\frac{n-1}{2} \frac{k}{p}}$ for $2 , and <math>A = \lambda^{\frac{1}{2}} (\log \lambda)^{\frac{1}{2}}$ for p = 2• If k = n-1, then $A = \lambda^{\frac{n-1}{2} \frac{k}{p}}$ for $\frac{2n}{n-1} \leq p \leq \infty$, and $A = \lambda^{\frac{n-1}{4} \frac{n-2}{2p}}$ for $2 \leq p < \frac{2n}{n-1}$

and when n = 1 and k = 0, we have A = 1 for 2 .

Eigenfunction restriction estimates on closed manifolds were obtained by Burq-Gérard-Tzvetkov [13] and Hu [27]. See also Tataru [37] for the first appearance of this type of estimates. The transversal restriction estimates of harmonic functions on hypersurfaces can be found in Taylor [38, Exercise 4 in Section 11 of Chapter 7]. Our result (1.39) precisely determines the dependence on the spectral frequency, and it is sharp in the sense that it can be saturated by Steklov eigenfunctions on Euclidean balls.

Corollary 6. If Σ and A are defined as in Theorem 5, then we have the transversal restriction estimates for Steklov eigenfunctions

(1.40)
$$||u_{\lambda}||_{L^{p}(\Sigma)} \lesssim \lambda^{-\frac{1}{p}} A ||e_{\lambda}||_{L^{2}(M)}.$$

Equivalently, by Theorem 3 we have

(1.41)
$$||u_{\lambda}||_{L^{p}(\Sigma)} \lesssim \lambda^{\frac{1}{2} - \frac{1}{p}} A ||u_{\lambda}||_{L^{2}(\Omega)}.$$

The bound $\lambda^{\frac{1}{2}-\frac{1}{p}}A$ is sharp on Euclidean balls.

- 1.6. Organization of the paper. In Section 2, we introduce the symbol classes and the Dirichlet-to-Neumann operator. In Section 3, we prove the Theorem 1 and its sharpness. In Section 4, we construct a parametrix for the Poisson integral operator and calculate the kernel of the its composition with the spectral cluster. In Section 5, we prove Theorem 2 and Theorem 3 by the parametrix in Lemma 2. In Section 6, we prove Theorem 4 by Lemma 3 and give a proof sketch for Theorems 5 and 6, and discuss related results on the numerical approximation of harmonic functions with various boundary conditions. In the Appendix, we prove Lemma 5.
- 1.7. **Notations.** Throughout this paper, $X \lesssim Y$ means $X \leq CY$ for some positive constants C that are independent of key parameters: the distance t to the boundary, and the frequencies λ , μ . If $X \lesssim Y$ and $Y \lesssim X$, we denote $X \approx Y$.

2. Preliminaries

In this section, we introduce some background on pseudo-differential operators. Let $D_x = \frac{1}{i}\partial_x$. Symbol class S^m is the set of smooth functions $p(x,\xi)$ satisfying

$$|D_x^{\beta} D_{\xi}^{\alpha} p(x,\xi)| \le C_{\alpha\beta} (1+|\xi|)^{m-|\alpha|}, \ \forall \alpha, \beta.$$

Let OPS^m be the set of pseudo-differential operators with symbols in S^m . Let $\delta_0 > 0$. We call $p(t, x, \xi) \in C^{\infty}([0, \delta_0] \times \mathbb{R}^n \times \mathbb{R}^n)$ a smooth family of symbols in S^m if

$$(2.2) |D_t^j D_x^{\beta} D_{\xi}^{\alpha} p(t, x, \xi)| \le C_{j\alpha\beta} (1 + |\xi|)^{m - |\alpha|}, \quad t \in [0, \delta_0], \ \forall j, \alpha, \beta.$$

The associated family of operators are called a smooth family in OPS^m .

Symbol class \mathcal{P}^m is the set of functions $p(t, x, \xi)$ in $C^{\infty}([0, \delta_0] \times \mathbb{R}^n \times \mathbb{R}^n)$ such that

$$|D_x^{\beta} D_{\varepsilon}^{\alpha}(t^k D_t^{\ell} p(t, x, \xi))| \le C_{\alpha\beta} (1 + |\xi|)^{m-k+\ell-|\alpha|}, \ t \in [0, \delta_0], \ \forall k, \ell, \alpha, \beta.$$

Symbol class \mathcal{P}_e^m is the set of functions $p(t, x, \xi) \in C^{\infty}([0, \delta_0] \times \mathbb{R}^n \times \mathbb{R}^n)$ such that

(2.3)
$$p(t, x, \xi) = q(t, x, \xi)e^{-ct\langle \xi \rangle}$$

with
$$\langle \xi \rangle = \sqrt{1 + |\xi|^2}$$
 for some $c > 0$ and $q(t, x, \xi) \in \mathcal{P}^m$. See e.g. Taylor [38, Chapter 7].

For any $f \in L^2(M)$, we denote its harmonic extension on Ω by $u = \mathcal{H}f$ that solves the boundary value problem (1.1). Then the DtN operator on M is defined as $\mathcal{N}f = \partial_{\nu}u|_{M}$, where ν is the outward unit norm vector. In addition, \mathcal{N} is self-adjoint.

Using the Fermi coordinates, for each $0 \le t \le \delta_0 < \delta$, we can define a DtN operator \mathcal{N}_t on Σ_t . By [38, Section C of Chapter 12], the Laplacian Δ on Ω in local coordinates with respect to the boundary has the form

(2.4)
$$\Delta = (\partial_t - A_1(t))(\partial_t + A(t)) + B(t)$$

where the smooth families A(t), $A_1(t) \in OPS^1(M)$ and $B(t) \in OPS^{-\infty}(M)$. Moreover, $A(t) = \mathcal{N}_t$ modulo smoothing operators, then \mathcal{N}_t is a smooth family in $OPS^1(M)$. By [38, Proposition C.1 in Chapter 12], we have explicit formulas for the principal and sub-principal symbols of \mathcal{N}_t .

Lemma 1. The Dirichlet-to-Neumann operator satisfies

$$\mathcal{N}_t = \sqrt{-\Delta_{g_t}} - \mathcal{B}_t$$

where $\mathcal{B}_t \in OPS^0(\Sigma_t)$ has principal symbol

(2.5)
$$\sigma_t(x,\xi) = \frac{1}{2} \left(Tr \mathcal{W}_{t,x} - \frac{\langle \mathcal{W}_{t,x}^* \xi, \xi \rangle_t}{\langle \xi, \xi \rangle_t} \right).$$

Here we define DtN operator by using the outward unit normal vector, so there is a difference in the sign compared to the one in [38, Section C of Chapter 12].

3. Exponential lower bounds

We first prove Theorem 1 based on the frequency function in Almgren [1] and Garofalo-Lin [21], and microlocal analysis techniques involving Lemma 1. See also Lin [32] and Di Cristo-Rondi [18]. And then we show the sharpness of Theorem 1 by constructing examples.

3.1. Proof of Theorem 1. Let

(3.1)
$$H(t) = \int_{\Sigma_t} u^2, \quad D(t) = -\int_{\Sigma_t} u \partial_t u, \quad N(t) = D(t)/H(t).$$

Let (t, x) be the Fermi coordinates described before and suppose $0 \le t \le \delta_0 < \delta$. We can identify each Σ_t as M with metric tensor g_t , so we can rewrite H(t), D(t) as

$$H(t) = \int_{\Sigma_t} u^2 = \int_M u(t, x)^2 d\omega_t(x)$$
$$D(t) = -\int_{\Sigma_t} u \partial_t u = -\int_M u(t, x) \partial_t u(t, x) d\omega_t(x)$$

where $d\omega_t(x) = \sqrt{\det(g_{ij}(t,x))}dx$ in local coordinates. Then

(3.2)
$$H'(t) = 2 \int_M u(t,x) \partial_t u(t,x) d\omega_t(x) + \int_M u^2(t,x) \partial_t (d\omega_t(x)).$$

For the volume derivative term $\partial_t(d\omega_t(x))$, As in [38, (C.26) in Section C of Chapter 12], under Fermi coordinate, we have

(3.3)
$$\frac{\partial_t \det(g_{ij}(t,x))}{\det(g_{ij}(t,x))} = -2\operatorname{Tr} \mathcal{W}_{t,x}.$$

Then we have

(3.4)
$$\partial_{t}(d\omega_{t}(x)) = \partial_{t}(\sqrt{\det(g_{ij}(t,x))}dx)$$

$$= \frac{\partial_{t}(\det(g_{ij}(t,x)))}{2\sqrt{\det(g_{ij}(t,x))}}dx$$

$$= \frac{\partial_{t}(\det(g_{ij}(t,x)))}{2\det(g_{ij}(t,x))}d\omega_{t}(x)$$

$$= -(\operatorname{Tr}\mathcal{W}_{t,x})d\omega_{t}(x).$$

Thus.

(3.5)
$$H'(t) = 2 \int_{\Sigma_t} u \partial_t u - \int_{\Sigma_t} (\operatorname{Tr} \mathcal{W}_{t,x}) u^2 = -2D(t) - \int_{\Sigma_t} (\operatorname{Tr} \mathcal{W}_{t,x}) u^2.$$

Next, we have

(3.6)
$$D'(t) = -\int_{M} |\partial_{t}u|^{2} d\omega_{t}(x) - \int_{M} u \partial_{t}^{2} u d\omega_{t}(x) - \int_{M} u \partial_{t} u \partial_{t} (d\omega_{t}(x)).$$

By (1.5),

$$0 = \Delta u = \partial_t^2 u(t, x) + a(t, x) \partial_t u(t, x) + \Delta_{q_t} u(t, x)$$

and we use (1.6), (3.4) to obtain

$$\int_{M} u \partial_{t}^{2} u d\omega_{t}(x) = -\int_{M} u \partial_{t} u a(t, x) d\omega_{t}(x) - \int_{M} u \Delta_{g_{t}} u d\omega_{t}(x)
= -\int_{M} u \partial_{t} u \frac{\partial_{t} (\det(g_{ij}(t, x)))}{2 \det(g_{ij}(t, x))} d\omega_{t}(x) - \int_{M} u \Delta_{g_{t}} u d\omega_{t}(x)
= -\int_{M} u \partial_{t} u \partial_{t} (d\omega_{t}(x)) - \int_{M} u \Delta_{g_{t}} u d\omega_{t}(x).$$
(3.7)

Plugging this into (3.6), we get

(3.8)
$$D'(t) = -\int_{\Sigma_t} |\partial_t u|^2 + \int_{\Sigma_t} u \Delta_{g_t} u.$$

To handle the second term on the right, we use the fact that $\sqrt{-\Delta_{g_t}}$ is self-adjoint and Lemma 1 to obtain

(3.9)
$$\int_{\Sigma_t} u \Delta_{g_t} u = -\int_{\Sigma_t} |\sqrt{-\Delta_{g_t}} u|^2$$

$$= -\int_{\Sigma_t} |(\mathcal{N}_t + \mathcal{B}_t) u|^2$$

$$= -\int_{\Sigma_t} |\mathcal{N}_t u|^2 - 2\int_{\Sigma_t} \mathcal{N}_t u \mathcal{B}_t u - \int_{\Sigma_t} |\mathcal{B}_t u|^2.$$

Since u is harmonic on Σ_t and $\mathcal{N}_t u = \partial_t u$ on Σ_t , and \mathcal{N}_t is self-adjoint, we have

(3.10)
$$D'(t) = -2 \int_{\Sigma_t} |\partial_t u|^2 - 2 \int_{\Sigma_t} \mathcal{N}_t u \mathcal{B}_t u - \int_{\Sigma_t} |\mathcal{B}_t u|^2$$
$$= -2 \int_{\Sigma_t} |\partial_t u|^2 - 2 \int_{\Sigma_t} u \mathcal{N}_t \mathcal{B}_t u - \int_{\Sigma_t} |\mathcal{B}_t u|^2.$$

Now we are ready to compute the derivative of N(t). Since $\mathcal{B}_t \in OPS^0$, it is bounded on L^2 and then $\int_{\Sigma_t} |\mathcal{B}_t u|^2 / \int_{\Sigma_t} |u|^2 = O(1)$. So by (3.1), (3.5) and (3.10) we have

$$N'(t) = \frac{D'(t)}{H(t)} - \frac{D(t)H'(t)}{H(t)^{2}}$$

$$= \frac{-2\int_{\Sigma_{t}} |\partial_{t}u|^{2} - 2\int_{\Sigma_{t}} u\mathcal{N}_{t}\mathcal{B}_{t}u}{\int_{\Sigma_{t}} u^{2}} + \frac{2(\int_{\Sigma_{t}} u\partial_{t}u)^{2}}{(\int_{\Sigma_{t}} u^{2})^{2}} + \frac{\int_{\Sigma_{t}} (\text{Tr}\mathcal{W}_{t,x})u^{2}}{\int_{\Sigma_{t}} u^{2}} N(t) + O(1)$$

$$= \frac{-2\int_{\Sigma_{t}} |\partial_{t}u|^{2} \int_{\Sigma_{t}} u^{2} + 2(\int_{\Sigma_{t}} u\partial_{t}u)^{2}}{(\int_{\Sigma_{t}} u^{2})^{2}} - \frac{2\int_{\Sigma_{t}} u\mathcal{N}_{t}\mathcal{B}_{t}u}{\int_{\Sigma_{t}} u^{2}} + \frac{\int_{\Sigma_{t}} (\text{Tr}\mathcal{W}_{t,x})u^{2}}{\int_{\Sigma_{t}} u^{2}} N(t) + O(1)$$

$$\leq -\frac{2\int_{\Sigma_{t}} u\mathcal{N}_{t}\mathcal{B}_{t}u}{\int_{\Sigma_{t}} u^{2}} + \frac{\int_{\Sigma_{t}} (\text{Tr}\mathcal{W}_{t,x})u^{2}}{\int_{\Sigma_{t}} u^{2}} N(t) + O(1).$$

The last step follows from Cauchy–Schwartz inequality.

For the rest terms, by Lemma 1, the principal symbol of $-2\mathcal{N}_t\mathcal{B}_t \in OPS^1$ is

$$-2|\xi|_{g_t}\sigma_t(x,\xi) \le |\xi|_{g_t}\tilde{k}_t(x) \le |\xi|_{g_t} \sup_{x \in \Sigma_t} \tilde{k}_t(x).$$

So the symbol of $(\sup_{x \in \Sigma_t} \tilde{k}_t(x)) \mathcal{N}_t + 2 \mathcal{N}_t \mathcal{B}_t$ is the sum of a non-negative principal symbol and a symbol of order 0. Combining with the L^2 boundedness of operators of order 0, we can apply the sharp Gårding inequality [26, Theorem 18.1.14] to the principal symbol and obtain

$$-\frac{2\int_{\Sigma_t} u \mathcal{N}_t \mathcal{B}_t u}{\int_{\Sigma_t} u^2} \le N(t) \sup_{x \in \Sigma_t} \tilde{k}_t(x) + O(1).$$

It is easy to see

(3.12)
$$\frac{\int_{\Sigma_t} (\operatorname{Tr} \mathcal{W}_{t,x}) u^2}{\int_{\Sigma_t} u^2} N(t) \le N(t) \sup_{x \in \Sigma_t} \operatorname{Tr} \mathcal{W}_{t,x}.$$

We finally have

$$(3.13) N'(t) \le (\sup_{x \in \Sigma_t} \tilde{k}_t(x) + \sup_{x \in \Sigma_t} \operatorname{Tr} \mathcal{W}_{t,x}) N(t) + O(1) = \Theta(t) N(t) + O(1).$$

So we have,

(3.14)
$$N(t) \le e^{\int_0^t \Theta(y)dy} N(0) + O(1).$$

By (3.5), we have

$$(\log H(t))' = \frac{H'(t)}{H(t)} \ge -2N(t) - \sup_{x \in \Sigma_t} \text{Tr} \mathcal{W}_{t,x}$$

again integrating both sides

(3.15)
$$\log H(t) - \log H(0) \ge -2 \int_0^t N(s) ds - \int_0^t \sup_{x \in \Sigma_s} \text{Tr} \mathcal{W}_{s,x} ds$$
$$\ge -2N(0) \int_0^t e^{\int_0^t \Theta(y) dy} dt + O(t)$$

and this gives us the desired bound

$$||u||_{L^2(\Sigma_t)} \gtrsim e^{-N(0)K(t)}||u||_{L^2(M)}.$$

3.2. Sharpness of Theorem 1. Now we are going to show the exponential lower bound can be saturated by Euclidean balls and a family of symmetric warped product manifolds.

Let $\Omega = [-R, R] \times M_0$, where (M_0, g) is a connected closed manifold, and Ω is equipped with the metric $h = \rho(s)^2 g + ds^2$ for some positive $\rho \in C^{\infty}([-R, R])$. First, we shall compute some geometric quantities and the Weingarten map for some special cases.

Next, we compute Weingarten maps of manifolds (Ω, h) in the setting as in Proposition 1 without the symmetry condition $\rho(s) = \rho(-s)$. For $0 \le t < R$,

$$\Sigma_t = \{-R + t\} \times M_0 \cup \{R - t\} \times M_0 = \Sigma_t^- \cup \Sigma_t^+.$$

We shall only compute it at Σ_t^+ , under this setting $g_t = \rho (R-t)^2 g$.

As in [38, (4.68)–(4.70) of Appendix C], we consider the one-parameter family of submanifolds given by

$$\phi_{\tau}: \Sigma_t^+ \to \Omega, \ \phi_{\tau}(R-t,x) = (R-t-\tau,x), \ \forall x \in M_0.$$

Let \tilde{g}_{τ} be the induced metric on the submanifold $\phi_{\tau}(\Sigma_t^+)$ and $x_+ = (R - t, x)$. Then for any tangent vectors v_1 , v_2 on Σ_t^+ at x_+ , we have

$$\partial_{\tau} \tilde{g}_{\tau}(v_1, v_2)|_{\tau=0} = -2\langle \partial_t, \mathbb{I}_{x_+}(v_1, v_2) \rangle = -2\langle \mathcal{W}_{t, x_+} v_1, v_2 \rangle_t$$

where the second fundamental form $\mathbb{I}_{x_+}(v_1, v_2) = \langle \mathcal{W}_{t, x_+} v_1, v_2 \rangle_t \partial_t$, and ∂_t is the inward unit normal vector on Σ_t^+ .

Since $\tilde{g}_{\tau} = \rho (R - t - \tau)^2 g$, we have

$$\begin{split} \partial_{\tau} \tilde{g}_{\tau}(v_1, v_2)|_{\tau=0} &= -2\rho'(R-t)\rho(R-t)g(v_1, v_2) \\ &= -\frac{2\rho'(R-t)}{\rho(R-t)}g_t(v_1, v_2) \\ &= -\frac{2\rho'(R-t)}{\rho(R-t)}\langle v_1, v_2 \rangle_t. \end{split}$$

Thus, for $x \in \Sigma_t^+$

(3.16)
$$\mathcal{W}_{t,x} = \frac{\rho'(R-t)}{\rho(R-t)}I$$

where I is the identity matrix. Similarly for $x \in \Sigma_t^-$,

$$\mathcal{W}_{t,x} = -\frac{\rho'(-R+t)}{\rho(-R+t)}I.$$

So the principal curvatures $k_1(x)$, \cdots , $k_n(x)$ are all equal to $\frac{\rho'(R-t)}{\rho(R-t)}$ on Σ_t^+ , and $-\frac{\rho'(-R+t)}{\rho(-R+t)}$ on Σ_t^- , and

$$\Theta(t) = \max \left\{ \frac{\rho'(R-t)}{\rho(R-t)}, -\frac{\rho'(-R+t)}{\rho(-R+t)} \right\}.$$

Specially if ρ is symmetric about 0, then

(3.17)
$$k_1(x) = k_2(x) = \dots = k_n(x) = \frac{\rho'(R-t)}{\rho(R-t)}, \quad \tilde{k}_t(x) = -\frac{(n-1)\rho'(R-t)}{\rho(R-t)},$$

and

(3.18)
$$\sigma_t(x,\xi) = \frac{(n-1)\rho'(R-t)}{2\rho(R-t)}, \quad \text{Tr}\mathcal{W}_{t,x} = \frac{n\rho'(R-t)}{\rho(R-t)}, \quad \Theta(t) = \frac{\rho'(R-t)}{\rho(R-t)}$$

for all $(x,\xi) \in T^*\Sigma_t$. Thus,

(3.19)
$$K(t) = \int_0^t \frac{\rho(R)}{\rho(R-s)} ds.$$

Example 1. Euclidean balls.

Let $\Omega = B(0,R) \in \mathbb{R}^{n+1}$ be the ball of radius of R centered at the origin. Let g, Δ, \mathcal{N} be the induced metric, Laplace operator and DtN operator on $\partial\Omega$ (a sphere of radius R). We have

$$\mathcal{N} = \sqrt{-\Delta + \frac{(n-1)^2}{4R^2}} - \frac{n-1}{2R},$$

see [38, Section 4 of Chapter 8]. Under Fermi coordinates, for $0 \le t < \delta = R$ we have

$$g_t = \frac{(R-t)^2}{R^2}g, \ \Delta_t = \frac{R^2}{(R-t)^2}\Delta, \ \mathcal{N}_t = \frac{R}{R-t}\mathcal{N}.$$

Suppose $\{\mu_k\}_{k=0}^{\infty}$ are the eigenvalues (counting multiplicity) of $\sqrt{-\Delta}$ and $\{e_k\}_{k=0}^{\infty}$ are the corresponding eigenfunctions. Then

$$\lambda_k = \sqrt{\mu_k^2 + \frac{(n-1)^2}{4R^2}} - \frac{n-1}{2R}, \quad \ u_k = (1 - \frac{t}{R})^{R\lambda_k} e_k, \ k = 0, 1, 2, \dots$$

are Steklov eigenvalues and eigenfunctions. Thus,

$$(3.20) ||u_k||_{L^p(\Sigma_t)} = \frac{(R-t)^{n/p}}{R^{n/p}} (1 - \frac{t}{R})^{R\lambda_k} ||e_k||_{L^p(M)} \approx e^{-\lambda_k R \log(R/(R-t))} ||e_k||_{L^p(M)}.$$

In this case, the metric has the form of $ds^2 + \rho(s)^2 g$ under polar coordinate with $\rho(s) = \frac{s}{R}$ when $0 < s \le R$, and the boundary has only one component. Thus

$$\Theta(t) = \frac{\rho'(R-t)}{\rho(R-t)} = \frac{1}{R-t}$$

and

$$K(t) = \int_0^t e^{\int_0^s (R - \tau)^{-1} d\tau} ds = R \log(\frac{R}{R - t}),$$

which is exactly the same as the coefficient of $-\lambda_k$ in (3.20).

Proposition 1. Let $\Omega = [-R, R] \times M_0$, where (M_0, g) is a connected closed manifold, and Ω is equipped with the metric $h = \rho(s)^2 g + ds^2$ for some positive $\rho \in C^{\infty}([-R, R])$ with $\rho(-s) = \rho(s)$. Then for any $\delta_0 < \delta = R$, there is a constant C > 0 depends on M_0 , ρ and δ_0 , such that, for each Steklov eigenvalue λ , there exists a Steklov eigenfunction u_{λ} satisfying

(3.21)
$$C^{-1}e^{-\lambda K(t)}\|u_{\lambda}\|_{L^{p}(M)} \le \|u_{\lambda}\|_{L^{p}(\Sigma_{t})} \le Ce^{-\lambda K(t)}\|u_{\lambda}\|_{L^{p}(M)},$$

for any $1 \leq p \leq \infty$.

The upper bound in (3.21) was also obtained in [17, Theorem 1.5] in a slightly different form. Here we give a direct proof.

Proof of Proposition 1. By the symmetry of ρ , there is a Steklov eigenfunction of the form

$$u_{\lambda} = b(s)e_{\mu}(x)$$

associated with the Steklov eigenvalue λ , where b(s) is symmetric or anti-symmetric about 0 and $e_{\mu}(x)$ is an L^2 normalized eigenfunction on M_0 .

We just need to handle p=2, since it follows from $u_{\lambda}=b(s)e_{\mu}(x)$ that

$$\frac{\|u_{\lambda}\|_{L^{p}(\Sigma_{t})}}{\|u_{\lambda}\|_{L^{p}(M)}} = \frac{\|u_{\lambda}\|_{L^{2}(\Sigma_{t})}}{\|u_{\lambda}\|_{L^{2}(M)}}.$$

The argument is similar to the proof of Theorem 1. By (3.11), we have

$$(3.22) N'(t) = R_1(t) + R_2(t) + R_3(t) + O(1),$$

where

$$R_{1} = \frac{-2\int_{\Sigma_{t}} |\partial_{t}u_{\lambda}|^{2} \int_{\Sigma_{t}} u_{\lambda}^{2} + 2(\int_{\Sigma_{t}} u_{\lambda} \partial_{t}u_{\lambda})^{2}}{(\int_{\Sigma_{t}} u_{\lambda}^{2})^{2}},$$

$$R_{2} = -\frac{2\int_{\Sigma_{t}} u_{\lambda} \mathcal{N}_{t} \mathcal{B}_{t} u_{\lambda}}{\int_{\Sigma_{t}} u_{\lambda}^{2}}, \qquad R_{3} = \frac{\int_{\Sigma_{t}} (\text{Tr} \mathcal{W}_{t,x}) u_{\lambda}^{2}}{\int_{\Sigma_{t}} u_{\lambda}^{2}} N(t).$$

We notice $R_1 = 0$, by the symmetry of b(s), and the fact that $u_{\lambda} = b(s)e_{\mu}(x)$.

By (3.18), the principal symbol of \mathcal{B}_t is

$$\sigma_t(x,\xi) = \frac{(n-1)\rho'(R-t)}{2\rho(R-t)} = -\frac{1}{2}\tilde{k}_t(x),$$

which is constant for all x, ξ on $T^*\Sigma_t$. So $2\mathcal{N}_t\mathcal{B}_t + \tilde{k}_t\mathcal{N}_t \in OPS^0$, and we can write

$$R_2 = -\frac{(n-1)\rho'(R-t)}{\rho(R-t)}N(t) + O(1).$$

Again by (3.18), we have

$$R_3 = \frac{n\rho'(R-t)}{\rho(R-t)}N(t).$$

Thus, by (3.22) and (3.18) we get

$$N'(t) = \frac{\rho'(R-t)}{\rho(R-t)}N(t) + O(1)$$
$$= \Theta(t)N(t) + O(1).$$

Then

(3.23)
$$N(t) = e^{\int_0^t \Theta(y)dy} N(0) + O(1).$$

By (3.5), we have

$$(\log H(t))' = \frac{H'(t)}{H(t)} = -2N(t) - O(1),$$

which gives

(3.24)
$$\log H(t) - \log H(0) = -2N(0) \int_0^t e^{\int_0^s \Theta(y)dy} ds + O(1).$$

This implies the desired bound

$$\|u_{\lambda}\|_{L^{2}(\Sigma_{t})}^{2} \approx e^{-2N(0)K(t)}\|u_{\lambda}\|_{L^{2}(M)}^{2} = e^{-2\lambda K(t)}\|u_{\lambda}\|_{L^{2}(M)}^{2}.$$

Next, we apply Proposition 1 to construct some examples with various decaying speeds, compared to original conjetured decaying speed $e^{-\lambda t}$ by Hislop-Lutzer [24]. These phenomena have been observed by Galkowski-Toth [19, 20].

Example 2. Quasi-cylinder with convex boundary.

Let $M_0 = \mathbb{R}^n/(2\pi\mathbb{Z}^n) = \mathbb{T}^n$ be the standard flat torus, and $\Omega = [-1, 1] \times M_0$ with the metric $h = ds^2 + (1 + s^2)dx^2$. Then

$$\Theta(t) = \frac{\rho'(R-t)}{\rho(R-t)} = \frac{2(1-t)}{1+(1-t)^2}.$$

By Proposition 1, for each Steklov eigenvalue λ , we can find some $\xi \in \mathbb{Z}^n$ such that $u_{\lambda} = b(s)e^{i\xi \cdot x}$. Then by (3.21), we get

$$||u_{\lambda}||_{L^{p}(\Sigma_{t})} \approx e^{-\lambda(t+t^{2}+O(t^{3}))} ||u_{\lambda}||_{L^{p}(M)},$$

which decays faster than $e^{-\lambda t} \|u_{\lambda}\|_{L^{p}(M)}$.

Example 3. Quasi-cylinder with concave boundary.

Let $M_0 = \mathbb{S}^n$ be the standard sphere and $\Omega = \left[-\frac{\pi}{3}, \frac{\pi}{3} \right] \times M_0$ with the metric $h = ds^2 + (\cos s) dx^2$. Then

$$\Theta(t) = -\tan(\frac{\pi}{3} - s).$$

By Proposition 1 for each Steklov eigenvalue λ , we can find a Steklov eigenfunction $u_{\lambda} = b(s)e_{\mu}(x)$ such that

$$||u_{\lambda}||_{L^{p}(\Sigma_{t})} \approx e^{-\lambda(t-\frac{\sqrt{3}}{2}t^{2}+O(t^{3}))}||u_{\lambda}||_{L^{p}(M)},$$

which decays slower than $e^{-\lambda t} \|u_{\lambda}\|_{L^{p}(M)}$.

4. Parametrix for the Poisson integral operator

As in Taylor [38, Section 12 of Chapter 7], we can construct a parametrix for the Poisson integral operator $u = \mathcal{H}f$ near the boundary for the boundary value problem

(4.1)
$$\begin{cases} \Delta u(t,x) = 0, & (t,x) \in \Omega, \\ u(0,x) = f(x), & x \in \partial \Omega. \end{cases}$$

See also Boutet de Monvel [10], Lee-Uhlmann [30].

Let \mathcal{N} be the DtN operator on $M = \partial \Omega$. As in Section 2, the Laplacian Δ on Ω in local coordinates with respect to the boundary has the form

(4.2)
$$\Delta = (\partial_t - A_1(t))(\partial_t + A(t)) + B(t)$$

where the smooth families $A(t), A_1(t) \in OPS^1(M)$ and $B(t) \in OPS^{-\infty}(M)$. Moreover, $A(0) = \mathcal{N}$ modulo a smoothing operator, and A(t) has the symbol $a(t, x, \xi) \in S^1$ with principal symbol $|\xi|_{g_t(x)}$. We have

$$|\partial_t^k \partial_x^\beta \partial_\xi^\alpha a(t, x, \xi)| \le C_{\alpha, \beta, k} (1 + |\xi|)^{1 - |\alpha|}, \quad \forall \alpha, \beta, k.$$

Lemma 2 (Parametrix for the Poisson integral operator). For any $0 < \delta_0 < \delta$ and $t \in [0, \delta_0]$, under Fermi coordinates, there exist a parametrix P_t locally given by

$$(4.3) P_t f(x) = \frac{1}{(2\pi)^n} \int e^{ix\cdot\xi} b(t, x, \xi) \hat{f}(\xi) d\xi, \ b \in \mathcal{P}_e^0,$$

and a smooth family of smoothing operators R_t such that the Poisson integral operator

(4.4)
$$\mathcal{H}f(t,x) = P_t f(x) + R_t f(x).$$

Moreover, the symbol

$$b(t, x, \xi) \sim b_0(t, x, \xi) + \sum_{j \ge 1} \sum_{2 \le k \le 2j} b_0(t, x, \xi) t^k q_{j,k}(t, x, \xi)$$

with $q_{j,k}(t)$ are smooth families in S^{k-j} and $b_0(t,x,\xi) = e^{-\int_0^t a(s,x,\xi)ds} \in \mathcal{P}_e^0$.

The symbol class \mathcal{P}_{e}^{0} is defined in Section 2.

Proof. Let $\mathbb{B}_r = \{x \in \mathbb{R}^n : |x| < r\}$. Let $\{(U_j, B_j, \kappa_j)\}_{j=1}^K$ be a collection of open sets on M. Each (U_j, κ_j) is a local coordinate chart, $\kappa_j(B_j) = \mathbb{B}_1$ and $\mathbb{B}_3 \subset \kappa_j(U_j)$. We may assume that there is a partition of unity $\{\phi_j^0\}_{j=1}^K$ adapted to $\{B_j\}_{j=1}^K$. Let $\phi \in C_0^\infty(\mathbb{R}^n)$ so that $\phi(x) = 1$ when |x| < 2 and $\phi(x) = 0$ when |x| > 3, denote $\phi_j^1 = \kappa_j^{-1}(\mathbb{B}_3)$. For each j, we construct a parametrix $P_j(t)$ in the form of (4.3) in U_j , then

$$P_t = \sum_{i=1}^K \phi_j^1 P_j(t) \phi_j^0$$

is a parametrix for the Poisson integral operator \mathcal{H} near the boundary. So we will only work in local coordinates on M. As in [38, Section 12 of Chapter 7], we only need to construct a parametrix P_t for the following evolution equation

$$\partial_t u + A(t)u = 0, \ u(0) = f,$$

since P_t also gives a parametrix for the Poisson integral operator \mathcal{H} by (4.2).

We build the symbol of P_t by the series $b(t, x, \xi) \sim \sum_{j\geq 0} b_j(t, x, \xi)$, which should satisfy the equation

$$\partial_t b + \sum_{\alpha} \frac{1}{\alpha!} D_{\xi}^{\alpha} a \partial_x^{\alpha} b = 0, \quad b(0, x, \xi) = 1.$$

To determine each term of the series, we require that

$$\partial_t b_0 + ab_0 = 0, \quad b(0, x, \xi) = 1,$$

and for $j \geq 1$,

(4.6)
$$\partial_t b_j + \sum_{\ell=0}^{j} \sum_{|\alpha|=j-\ell} \frac{1}{\alpha!} D_{\xi}^{\alpha} a \partial_x^{\alpha} b_{\ell} = 0, \quad b_j(0, x, \xi) = 0.$$

Solving (4.5) gives

(4.7)
$$b_0(t, x, \xi) = e^{-\int_0^t a(s, x, \xi) ds} \in \mathcal{P}_e^0,$$

since a(s) is a smooth family in S^1 . To solve (4.6), we let

$$r_j(t, x, \xi) = -\sum_{\ell=0}^{j-1} \sum_{|\alpha|=j-\ell} \frac{1}{\alpha!} D_{\xi}^{\alpha} a \partial_x^{\alpha} b_{\ell}.$$

Then (4.6) is equivalent to

$$\partial_t b_i + ab_i = r_i, \quad b_i(0, x, \xi) = 0$$

which gives the recursive formula

$$b_{j}(t, x, \xi) = b_{0}(t, x, \xi) \int_{0}^{t} r_{j}(s, x, \xi) b_{0}(s, x, \xi)^{-1} ds$$

$$= -b_{0}(t, x, \xi) \sum_{\ell=0}^{j-1} \sum_{|\alpha|=j-\ell} \frac{1}{\alpha!} \int_{0}^{t} D_{\xi}^{\alpha} a(s, x, \xi) \partial_{x}^{\alpha} b_{\ell}(s, x, \xi) \cdot b_{0}(s, x, \xi)^{-1} ds.$$

We claim that for $j \geq 1$

(4.8)
$$b_j(t, x, \xi) = b_0(t, x, \xi) \sum_{2 \le k \le 2j} t^k q_{j,k}(t, x, \xi),$$

where $q_{j,k}(t)$ are smooth families in S^{k-j} . Then we have $b_j \in \mathcal{P}_e^{-j}$ by (4.7). So we can build a symbol $b \sim \sum b_j \in \mathcal{P}_e^0$ (see e.g. [35, Lemma 3.1.2]). Now we prove the claim (4.8) by induction. For j = 1,

$$b_{1}(t,x,\xi) = -b_{0}(t,x,\xi) \sum_{|\alpha|=1} \int_{0}^{t} D_{\xi}^{\alpha} a(s,x,\xi) \partial_{x}^{\alpha} b_{0}(s,x,\xi) \cdot b_{0}(s,x,\xi)^{-1} ds$$

$$= -b_{0}(t,x,\xi) \sum_{|\alpha|=1} \int_{0}^{t} D_{\xi}^{\alpha} a(s,x,\xi) \int_{0}^{s} \partial_{x}^{\alpha} a(\tau,x,\xi) d\tau ds$$

$$= b_{0}(t,x,\xi) t^{2} q_{1,2}(t,x,\xi)$$

where $q_{1,2}(t)$ is a smooth family in S^1 . Suppose (4.8) holds for b_ℓ with $1 \le \ell < j$. Then for $|\alpha| \ge 1$

$$\partial_x^{\alpha} b_0(t, x, \xi) = b_0(t, x, \xi) \sum_{m=1}^{|\alpha|} t^m q_m(t, x, \xi)$$

and for $1 \le \ell < j$ and $|\alpha| \ge 0$

$$\begin{split} \partial_x^\alpha b_\ell(t,x,\xi) &= \sum_{2 \leq k \leq 2\ell} \sum_{|\alpha_1| + |\alpha_2| = |\alpha|} t^k \partial_x^{\alpha_1} b_0(t,x,\xi) \partial_x^{\alpha_2} q_{\ell,k}(t,x,\xi) \\ &= b_0(t,x,\xi) \sum_{2 \leq k \leq 2\ell + |\alpha|} t^k \tilde{q}_{k-\ell}(t,x,\xi) \end{split}$$

where $q_m(t)$, $\tilde{q}_m(t)$ are smooth families in S^m . Thus, we have

$$b_{j}(t, x, \xi) = -b_{0}(t, x, \xi) \sum_{\ell=0}^{j-1} \sum_{|\alpha|=j-\ell} \frac{1}{\alpha!} \int_{0}^{t} D_{\xi}^{\alpha} a(s, x, \xi) \partial_{x}^{\alpha} b_{\ell}(s, x, \xi) \cdot b_{0}(s, x, \xi)^{-1} ds$$

$$= -b_{0}(t, x, \xi) \sum_{|\alpha|=j} \sum_{m=1}^{j} \frac{1}{\alpha!} \int_{0}^{t} D_{\xi}^{\alpha} a(s, x, \xi) s^{m} q_{m}(s, x, \xi) ds$$

$$-b_{0}(t, x, \xi) \sum_{\ell=1}^{j-1} \sum_{|\alpha|=j-\ell} \sum_{2 \le k \le j+\ell} \frac{1}{\alpha!} \int_{0}^{t} D_{\xi}^{\alpha} a(s, x, \xi) s^{k} \tilde{q}_{k-\ell}(s, x, \xi) ds$$

$$= b_{0}(t, x, \xi) \sum_{2 \le k \le j} t^{k} q_{j,k}(t, x, \xi).$$

where $q_{j,k}(t)$ are smooth families in S^{k-j} . So the claim (4.8) is proved.

4.1. **Kernel of the composition operator.** In the following, let $P \in OPS^1(M)$ be classical and self-adjoint with principal symbol $p(x,\xi) = |\xi|_{g(x)}$. Let $\chi \in \mathcal{S}(\mathbb{R})$ satisfy supp $\hat{\chi} \subset (\frac{1}{2},1)$ and $\chi(0) = 1$. By [35, Lemma 5.1.3], we can write the kernel of the spectral cluster as

$$\chi(P-\lambda)(x,y) = \lambda^{\frac{n-1}{2}} e^{i\lambda d_g(x,y)} a_\lambda(x,y) + r_\lambda(x,y)$$

where $a_{\lambda}(x,y)$ is supported in $\{d_g(x,y)\approx 1\}$ and $|\partial_{x,y}^{\alpha}a_{\lambda}(x,y)|\leq C_{\alpha}, \ \forall \alpha.$ And the remainder term is smooth and satisfies

$$|\partial_{x,y}^{\alpha} r_{\lambda}(x,y)| \le C_{\alpha} \lambda^{-N}, \quad \forall N, \alpha.$$

Recall g_t is the metric on Σ_t under Fermi coordinates and $g = g_0$. Let

(4.9)
$$r(t) = \inf_{x \in M, \ \xi \neq 0} \frac{|\xi|_{g_t(x)}}{|\xi|_{g(x)}}, \quad G(t) = \int_0^t r(s)ds.$$

We have r(t) = 1 + O(t) and $G(t) = t + O(t^2)$. One can check that G(t) = K(t) if Ω is a Euclidean ball or the warped product manifold in Proposition 1. Now we can calculate the kernel of $\mathcal{H} \circ \chi(P - \lambda)$, i.e. the composition of the Poisson integral operator with the spectral cluster, by applying the parametrix in Lemma 2.

Lemma 3. Fix $0 < \delta_0 < \delta$. For any $0 \le t \le \delta_0$, we have

$$(4.10) \qquad \mathcal{H}(\chi(P-\lambda)f)(t,x) = \lambda^{\frac{n-1}{2}}e^{-c\lambda G(t)} \int e^{i\lambda d_g(x,y)}a_\lambda(t,x,y)f(y)dz + \int r_\lambda(t,x,y)f(y)dy,$$

where $a_{\lambda}(t,x,y)$ is supported in $\{d_g(x,y)\approx 1\}$ and $|\partial_t^k\partial_{x,y}^{\alpha}a_{\lambda}(t,x,y)|\leq C_{\alpha,k}\lambda^k$, $\forall \alpha,k$. And the remainder term has a smooth kernel satisfying

$$|\partial_t^k \partial_{x,y}^{\alpha} r_{\lambda}(t,x,y)| \le C_{\alpha,k} \lambda^{-N}, \quad \forall N, \alpha, k.$$

The positive constants $c, C_{\alpha,k}$ are independent of λ .

Proof. For convenience, we choose x to be geodesic normal coordinates at x_0 and work in a small neighborhood of x_0 . Let $\chi_{\lambda} = \chi(P - \lambda) - r_{\lambda}$. By Lemma 2 we can write

$$\mathcal{H}(\chi(P-\lambda)f)(t,x) = P_t \chi(P-\lambda)f(x) + R_t \chi(P-\lambda)f(x)$$

= $P_t \chi_\lambda f(x) + P_t r_\lambda f(x) + R_t \chi(P-\lambda)f(x)$.

Here

$$P_t \chi_{\lambda} f(x) = (2\pi)^{-n} \iint e^{i(x-y)\cdot\xi} b(t,x,\xi) \chi_{\lambda} f(y) dy d\xi$$
$$= (2\pi)^{-n} \lambda^{\frac{n-1}{2}} \iiint e^{i(x-y)\cdot\xi + i\lambda d_g(y,z)} b(t,x,\xi) a_{\lambda}(y,z) f(z) dz dy d\xi.$$

Since P_t is a smooth family of operators in OPS^0 , it is smoothing off diagonal, and the contribution of y away from x is $O(\lambda^{-N})$. Thus, we can assume the integral with respect to y in the expression of P_t is taking near diagonal and $\max\{d_g(x,x_0),d_g(y,x)\}\ll d_g(x_0,z)$. Since the kernel of R_t is smooth, we can integrate by parts to see that the kernel of $R_t\chi(P-\lambda)$ is smooth and $O(\lambda^{-N})$. Similarly, the kernel of P_tr_λ is also smooth and $O(\lambda^{-N})$. Let $\beta\in C_0^\infty(\mathbb{R})$ satisfy

$$1\!\!1_{[\frac{1}{2},2]} \leq \beta \leq 1\!\!1_{[\frac{1}{4},4]}.$$

Let

$$R_t^{\lambda} f(x) = (2\pi)^{-n} \lambda^{\frac{n-1}{2}} \iiint e^{i(x-y)\cdot\xi + i\lambda d_g(y,z)} b(t,x,\xi) (1 - \beta(|\xi|_{g(x)}/\lambda)) a_{\lambda}(y,z) f(z) dz dy d\xi.$$

On the support of $1 - \beta(|\xi|_{g(x)}/\lambda)$, and $g_{ij}(x) \approx g_{ij}(x_0) = \delta_{ij}$, $d_g(y,z) \approx |y-z|$, we have $|\partial_y((x-y)\cdot\xi + \lambda d_g(y,z))| \gtrsim \lambda + |\xi|$.

So we integrate by parts in y to see the kernel of R_t^{λ} is smooth and $O(\lambda^{-N})$. Let

$$P_t^{\lambda}f(x) = (2\pi)^{-n}\lambda^{\frac{n-1}{2}+n} \iiint e^{i\lambda((x-y)\cdot\xi + d_g(y,z))}b(t,x,\lambda\xi)\beta(|\xi|_{g(x)})a_{\lambda}(y,z)f(z)dyd\xi dz.$$

By the support property of $a_{\lambda}(y,z)$, we can integrate by parts in ξ to see that the kernel

$$\left| \iint e^{i\lambda((x-y)\cdot\xi + d_g(y,z))} (1 - \eta(x,z)) b(t,x,\lambda\xi) \beta(|\xi|_{g(x)}) a_\lambda(y,z) dy d\xi \right| \lesssim \lambda^{-N}$$

if $\eta \in C^{\infty}$ equals 1 when $|x-z| \approx 1$ and 0 otherwise. Since $a(t,x,\xi) = |\xi|_{g_t(x)} \mod S^0$, by (4.9) on the support of $\beta(|\xi|_{g(x)})$ we have

$$\beta(|\xi|_{q(x)}) \ge \lambda r(t)|\xi|_{q(x)} + O(1) \gtrsim \lambda r(t).$$

Then for some c > 0 we can write

(4.11)
$$b(t, x, \lambda \xi) = e^{-c\lambda G(t)} \tilde{b}(t, x, \lambda \xi),$$

where on the support of $\beta(|\xi|_{g(x)})$ we have

$$|\partial_t^k \partial_x^\beta \partial_\xi^\alpha \tilde{b}(t, x, \lambda \xi)| \le C_{\alpha, \beta, k} \lambda^k, \ \forall \alpha, \beta, k.$$

So it suffices to handle the operator

$$\begin{split} \tilde{P}_t^{\lambda}f(x) &= (2\pi)^{-n}\lambda^{\frac{n-1}{2}+n} \iiint e^{i\lambda((x-y)\cdot\xi+d_g(y,z))}\eta(x,z)b(t,x,\lambda\xi)\beta(|\xi|_{g(x)})a_{\lambda}(y,z)f(z)dyd\xi dz \\ &= (2\pi)^{-n}\lambda^{\frac{n-1}{2}+n}e^{-c\lambda G(t)} \iiint e^{i\lambda((x-y)\cdot\xi+d_g(y,z))}\eta(x,z)\tilde{b}(t,x,\lambda\xi)\beta(|\xi|_{g(x)})a_{\lambda}(y,z)f(z)dyd\xi dz. \end{split}$$

We shall use stationary phase in (y, ξ) . Indeed,

$$\partial_{y,\xi}((x-y)\cdot\xi+d_q(y,z))=0 \Rightarrow y=x, \ \xi=\partial_y d_q(y,z),$$

and

$$\partial^2_{y,\xi}((x-y)\cdot\xi+d_g(y,z)) = \begin{bmatrix} \partial^2_y d_g(y,z) & -I \\ -I & 0 \end{bmatrix}.$$

Thus, $(y,\xi) = (x,\partial_y d_g(y,z))$ is a non-degenerate critical point of the phase function. By stationary phase [35, Corollary 1.1.8], we can write

$$\tilde{P}_t^{\lambda} f(x) = \lambda^{\frac{n-1}{2}} e^{-c\lambda G(t)} \int e^{i\lambda d_g(x,z)} a_{\lambda}(t,x,z) f(z) dz,$$

where $a_{\lambda}(t,x,z)$ is supported in $\{d_g(x,z)\approx 1\}$ and $|\partial_t^k\partial_{x,y}^{\alpha}a_{\lambda}(t,x,y)|\leq C_{\alpha,k}\lambda^k, \ \forall \alpha,k.$

By using an essentially the same argument, we can obtain the kernel of the composition operator $Q \circ \chi(P - \lambda)$ for any $Q \in OPS^m(M)$.

Lemma 4. Let $m \in \mathbb{R}$ and $Q \in OPS^m(M)$. We have

$$(4.12) (Q \circ \chi(P - \lambda))(x, y) = \lambda^{\frac{n-1}{2} + m} e^{i\lambda d_g(x, y)} \tilde{a}_{\lambda}(x, y) + \tilde{r}_{\lambda}(x, y)$$

where $\tilde{a}_{\lambda}(x,y)$ is supported in $\{d_g(x,y)\approx 1\}$ and $|\partial_{x,y}^{\alpha}\tilde{a}_{\lambda}(x,y)|\leq C_{\alpha}, \ \forall \alpha.$ And the remainder term is smooth and satisfies

$$|\partial_{x,y}^{\alpha} \tilde{r}_{\lambda}(x,y)| \leq C_{\alpha}, \quad \forall \alpha.$$

The positive constants C_{α} are independent of λ .

This lemma implies that the kernel of $Q \circ \chi(P - \lambda)$ has essentially the same form as the kernel of $\chi(P-\lambda)$. So one may use it to obtain natural eigenfunction estimates under the operations in OPS^{m} , which extend the seminal works by Sogge [35] and Burq-Gérard-Tzvetkov [13, 11, 12].

Proposition 2. Let $m \in \mathbb{R}$ and $Q \in OPS^m(M)$. For $\lambda \geq 1$, we have

where

$$\sigma(p) = \begin{cases} \frac{n-1}{2} (\frac{1}{2} - \frac{1}{p}), & 2 \le p < \frac{2(n+1)}{n-1} \\ \frac{n-1}{2} - \frac{n}{p}, & \frac{2(n+1)}{n-1} \le p \le \infty. \end{cases}$$

Proposition 3. (Bilinear estimates) Let $m_j \in \mathbb{R}$ and $Q_j \in OPS^{m_j}(M)$ for j = 1, 2. If $\mu \geq \lambda \geq 1$, then we have

$$(4.14) ||Q_1 \circ \chi(P - \lambda)f \cdot Q_2 \circ \chi(P - \mu)g||_{L^2(M)} \lesssim \lambda^{m_1} \mu^{m_2} B||f||_{L^2(M)} ||g||_{L^2(M)},$$

where

$$B = \begin{cases} \lambda^{\frac{1}{4}}, & n = 2\\ \lambda^{\frac{1}{2}}\sqrt{\log \lambda}, & n = 3\\ \lambda^{\frac{n-2}{2}}, & n \ge 4. \end{cases}$$

Proposition 4. (Restriction estimates) Let $m \in \mathbb{R}$ and $Q \in OPS^m(M)$. Let Σ be a k-dimensional submanifold in M. For $\lambda \geq 1$, we have

$$(4.15) ||Q \circ \chi(P - \lambda)f||_{L^p(\Sigma)} \lesssim \lambda^m A ||f||_{L^2(M)}, \quad 2 \le p \le \infty,$$

where

- If $k \le n-3$, then $A = \lambda^{\frac{n-1}{2} \frac{k}{p}}$ for $2 \le p \le \infty$. If k = n-2, then $A = \lambda^{\frac{n-1}{2} \frac{k}{p}}$ for $2 , and <math>A = \lambda^{\frac{1}{2}} (\log \lambda)^{\frac{1}{2}}$ for p = 2. If k = n-1, then $A = \lambda^{\frac{n-1}{2} \frac{k}{p}}$ for $\frac{2n}{n-1} \le p \le \infty$, and $A = \lambda^{\frac{n-1}{4} \frac{n-2}{2p}}$ for $2 \le p < \frac{2n}{n-1}$.

Let $\mathbb{I}_{(\lambda-1,\lambda]}$ be the indicator function of $(\lambda-1,\lambda]$. Since we may assume χ is non-vanishing on [-1,1], we have $\chi(P-\lambda)\tilde{f}=\mathbb{1}_{(\lambda-1,\lambda]}(P)f$ if $\tilde{f}=\chi(P-\lambda)^{-1}\mathbb{1}_{(\lambda-1,\lambda]}(P)f$. Therefore, the propositions above still hold if we replace $\chi(P-\lambda)$ by $\mathbb{1}_{(\lambda-1,\lambda]}(P)$.

5. Interior decay and comparable norms

In this section, we prove Theorem 2 and Theorem 3, by using microlocal analysis techniques involving the parametrix of the Poisson integral operator in Lemma 2. Note that the symbol of the parametrix P_t has an exponential decay factor $\sim e^{-ct\langle\xi\rangle}$, which formally gives the desired exponential decaying property if the frequency $|\xi| \gtrsim \lambda$. This is basic idea in the proof.

Let $P \in OPS^1$ be classical and self-adjoint with principal symbol $p(x,\xi)$ positive on $T^*M \setminus 0$. Let $\mu \in \mathbb{R}$ and let $m \in C^{\infty}(\mathbb{R})$ satisfy

$$|m^{(k)}(\tau)| \le C_k (1+|\tau|)^{\mu-k}, \ k=0,1,2,\dots$$

Let $\lambda \geq 1$ and $m_{\lambda}(\tau) = m(\tau/\lambda)$. We have $m_{\lambda} \in S^{\mu_1}$ if $\mu_1 = \max\{\mu, 0\}$, since

(5.2)
$$|m_{\lambda}^{(k)}(\tau)| \le C_k (1+|\tau|)^{\mu_1-k}, \ k=0,1,2,\dots$$

The constant C_k is independent of λ . We need the following lemma on the kernel of $m_{\lambda}(P)$, which can be viewed as a more precise version of [35, Theorem 4.3.1]. We shall prove it in the appendix.

Lemma 5. Let μ , μ_1 , m, m_{λ} defined as above. We have

$$m_{\lambda}(P)(x,y) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} M_{\lambda}(x,\xi) d\xi + R_{\lambda}(x,y)$$

where the symbol $M_{\lambda} \in S^{\mu_1}$ satisfies

$$(5.3) M_{\lambda}(x,\xi) \sim \sum_{\ell=0}^{\infty} m_{\lambda}^{(\ell)}(p(x,\xi)) q_{\ell}(x,\xi), \quad q_{\ell} \in S^{\lfloor \ell/2 \rfloor},$$

and the remainder term R_{λ} is smooth with

$$|\partial_{x,y}^{\alpha} R_{\lambda}(x,y)| \le C_{\alpha}, \quad \forall \alpha.$$

The symbols q_{ℓ} and the constant C_{α} are independent of λ .

In the following, we assume $P \in OPS^1(M)$ is classical and self-adjoint with principal symbol $p(x,\xi) = |\xi|_{g(x)}$. Let $\mathbb{1}_{\geq \lambda}$ be the indicator function of $[\lambda, \infty)$, and let $f_{\geq \lambda} = \mathbb{1}_{\geq \lambda}(P)f$.

Proof of Theorem 2. Since $(\mathcal{H}f)(t,x) = P_t f(x) + R_t f(x)$ with $R_t \in OPS^{-\infty}$ by Lemma 2, we shall estimate the norms $\|P_t f_{\geq \lambda}\|_{L^p(M)}$ and $\|R_t f_{\geq \lambda}\|_{L^p(M)}$.

If $\{e_j\}$ is an orthonormal eigenbasis associated with the eigenvalues $\{\lambda_j\}$ of the operator P, then by the Hörmander's $L^2 - L^{\infty}$ bounds (1.19) we have

$$\sum_{\lambda_j \ge \lambda} (1 + \lambda_j)^{-N-n} |e_j(x)|^2 \lesssim \sum_{k \ge \lambda} k^{-N-1} \lesssim \lambda^{-N}, \ \forall N > 0.$$

Since R_t is smoothing.

$$||R_t f_{\geq \lambda}||_{L^{\infty}(M)} = ||R_t (1+P)^{N+n} (1+P)^{-N-n} f_{\geq \lambda}||_{L^{\infty}(M)}$$

$$\lesssim ||(1+P)^{-N-n} f_{\geq \lambda}||_{L^{\infty}(M)}$$

$$\lesssim \lambda^{-N} ||f_{\geq \lambda}||_{L^1(M)}, \quad \forall N > 0.$$

Now we begin to handle P_t . By Lemma 2, we have

$$P_t f(x) = (2\pi)^{-n} \int e^{ix\cdot\xi} b(t, x, \xi) \hat{f}(\xi) d\xi$$

where the symbol

$$b(t, x, \xi) \sim b_0(t, x, \xi) + \sum_{j \ge 1} \sum_{2 \le k \le 2j} b_0(t, x, \xi) t^k q_{j,k}(t, x, \xi)$$

with $q_{j,k}(t)$ are smooth families in S^{k-j} and $b_0(t,x,\xi) = e^{-\int_0^t a(s,x,\xi)ds} \in \mathcal{P}_e^0$. Let $0 < \varepsilon < \frac{1}{100}$ and $\eta \in C^{\infty}(\mathbb{R})$ satisfy

$$\mathbb{1}_{[1,\infty)} \leq \eta \leq \mathbb{1}_{[1-\varepsilon,\infty)}$$
.

Let $\eta_1 \in C^{\infty}(\mathbb{R})$ satisfy

$$\mathbf{1}_{[1-\varepsilon,\infty)} \le \eta_1 \le \mathbf{1}_{[1-2\varepsilon,\infty)}.$$

Clearly, we have $\eta(P/\lambda)f_{\geq \lambda} = f_{\geq \lambda}$. We split

$$P_t f_{\geq \lambda}(x) = \int e^{ix \cdot \xi} b(t, x, \xi) \eta_1(p(x, \xi)/\lambda) (f_{\geq \lambda})^{\wedge}(\xi) d\xi + \int e^{ix \cdot \xi} b(t, x, \xi) (1 - \eta_1(p(x, \xi)/\lambda)) (\eta(P/\lambda) f_{\geq \lambda})^{\wedge}(\xi) d\xi$$
$$:= \tilde{P}_t f_{>\lambda}(x) + \tilde{R}_t f_{>\lambda}(x).$$

The operator \tilde{R}_t is smoothing by Lemma 5, since the supports of $1 - \eta_1$ and η are disjoint. Then we also have

(5.4)
$$\|\tilde{R}_t f_{\geq \lambda}\|_{L^{\infty}(M)} \lesssim \lambda^{-N} \|f_{\geq \lambda}\|_{L^1(M)}, \ \forall N.$$

Next, we handle \tilde{P}_t . The kernel of \tilde{P}_t is

$$K_t(x,y) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} b(t,x,\xi) \eta_1(p(x,\xi)/\lambda) d\xi.$$

On the support of $\eta_1(p(x,\xi)/\lambda)$ we have $p(x,\xi) = |\xi|_{g(x)} \ge (1-2\varepsilon)\lambda$. Since the principal symbol of $a(t,x,\xi)$ is $|\xi|_{q_t(x)}$, by (4.9) we have

$$|\xi|_{q_t(x)} \ge r(t)|\xi|_{q(x)} \ge (1 - 2\varepsilon)\lambda r(t).$$

Recall (4.9). Then we can write

$$(5.5) b(t, x, \xi) = e^{-(1-3\varepsilon)\lambda G(t)}\tilde{b}(t, x, \xi),$$

where

$$|\partial_t^k \partial_x^\beta \partial_{\xi}^\alpha \tilde{b}(t, x, \xi)| \le C_{\alpha, \beta, k} (1 + |\xi|)^{k - |\alpha|}, \ \forall \alpha, \beta, k.$$

Since $\tilde{b}(t, x, \xi)\eta_1(p(x, \xi)/\lambda) \in S^0$, we have for 1

(5.6)
$$\|\tilde{P}_t\|_{L^p(M)\to L^p(M)} \lesssim e^{-(1-3\varepsilon)\lambda G(t)}.$$

For $p = \infty$ or 1, we need to calculate the kernel. We have

$$(5.7) |K_t(x,y)| \lesssim e^{-(1-3\varepsilon)\lambda G(t)} \int_{|\xi| > \lambda} e^{-\varepsilon t|\xi|/2} d\xi \lesssim e^{-(1-3\varepsilon)\lambda G(t)} t^{-n}.$$

But integration by parts gives

$$(5.8) |K_t(x,y)| \lesssim e^{-(1-3\varepsilon)\lambda G(t)} |x-y|^{-N} \int_{|\xi| > \lambda} |\xi|^{-N} \lesssim e^{-(1-3\varepsilon)\lambda G(t)} \lambda^n (\lambda |x-y|)^{-N}.$$

So by splitting the integral into two parts: $|x-y| \le \lambda^{-1}$ and $|x-y| > \lambda^{-1}$ we have

$$\int |K_t(x,y)| dx \lesssim e^{-(1-3\varepsilon)\lambda G(t)} (1+(\lambda t)^{-n}).$$

By Young's inequality, for $p = \infty$ or 1

(5.9)
$$\|\tilde{P}_t\|_{L^p(M)\to L^p(M)} \lesssim e^{-(1-3\varepsilon)\lambda G(t)} (1+(\lambda t)^{-n}).$$

Then for $p = \infty$ or 1

But by the maximum principle, when $t \leq \lambda^{-1}$,

$$\|\mathcal{H}f_{\geq \lambda}\|_{L^{\infty}(\Sigma_t)} \leq \|f_{\geq \lambda}\|_{L^{\infty}(M)}$$
.

So we always have for 1

(5.11)
$$\|\mathcal{H}f_{\geq \lambda}\|_{L^{p}(\Sigma_{t})} \lesssim e^{-(1-3\varepsilon)\lambda G(t)} \|f_{\geq \lambda}\|_{L^{p}(M)} + \lambda^{-N} \|f_{\geq \lambda}\|_{L^{1}(M)}.$$

Furthermore, if f has spectral frequency $\approx \lambda$, then (5.11) is valid for all $1 \le p \le \infty$ by repeating the argument above, since (5.7) can be improved in this case. So we finish the proof of Theorem 2.

Next, we show that if f has spectral frequency $\approx \lambda$, than $\mathcal{H}f$ also has a lower bound near the boundary.

Lemma 6. Let $\beta \in C_0^{\infty}(\mathbb{R}^+)$ and $f_{\lambda} = \beta(P/\lambda)f$. For $t \leq \lambda^{-1}$, we have for any $1 \leq p \leq \infty$, (5.12) $\|\mathcal{H}f_{\lambda}\|_{L^p(\Sigma_t)} \gtrsim \|f_{\lambda}\|_{L^p(M)}.$

Proof. We may assume supp $\beta \subset (\frac{1}{2},1)$. Since $(\mathcal{H}f)(t,x) = P_t f(x) + R_t f(x)$ with $R_t \in OPS^{-\infty}$ by Lemma 2, we shall estimate the norms $\|P_t f_{\lambda}\|_{L^p(M)}$ and $\|R_t f_{\lambda}\|_{L^p(M)}$. As before, we have

$$||R_t f_{\lambda}||_{L^{\infty}(M)} \lesssim \lambda^{-N} ||f_{\lambda}||_{L^1(M)}, \ \forall N.$$

Here we use the fact that f_{λ} has spectral frequency bounded below.

To handle P_t , we split $b(t, x, \xi) = b_0(t, x, \xi) + r(t, x, \xi)$ with $r(t, x, \xi) \in S^{-1}$. Let

$$r_t f(x) = (2\pi)^{-n} \int e^{ix \cdot \xi} r(t, x, \xi) \hat{f}(\xi) d\xi.$$

Let $\beta_1 \in C_0^{\infty}$ satisfy

$$1_{\left[\frac{1}{2},1\right]} \le \beta_1 \le 1_{\left[\frac{1}{4},2\right]}.$$

Clearly, $\beta_1 \beta = \beta$. Since $t \leq \lambda^{-1}$, we can obtain the kernel estimate

$$|r_t \beta_1(P/\lambda)(x,y)| \lesssim \lambda^{n-1} (1+\lambda|x-y|)^{-N}, \ \forall N.$$

Then for $1 \le p \le \infty$ by Young's inequality we have

$$||r_t f_{\lambda}||_{L^p(M)} = ||r_t \beta_1(P/\lambda) f_{\lambda}||_{L^p(M)} \lesssim \lambda^{-1} ||f_{\lambda}||_{L^p(M)}.$$

Next, let $\beta_2 \in C_0^{\infty}$ satisfy

$$1_{\left[\frac{1}{8},4\right]} \le \beta_2 \le 1_{\left[\frac{1}{16},8\right]}$$
.

We split $b_0 = \beta_2(|\xi|_{q(x)}/\lambda))b_0 + (1 - \beta_2(|\xi|_{q(x)}/\lambda)))b_0$. Let

$$\tilde{R}_t f(x) = (2\pi)^{-n} \int e^{ix\cdot\xi} (1 - \beta_2(|\xi|_{g(x)}/\lambda)) b_0(t, x, \xi) \hat{f}(\xi) d\xi.$$

Since supp $(1 - \beta_2)$ is disjoint with supp β_1 , the operator $\tilde{R}_t \beta_1(P/\lambda)$ is smoothing by Lemma 5, and then

$$\|\tilde{R}_t f_{\lambda}\|_{L^{\infty}(M)} = \|\tilde{R}_t \beta_1(P/\lambda) f_{\lambda}\|_{L^{\infty}(M)} \lesssim \lambda^{-N} \|f_{\lambda}\|_{L^1(M)}.$$

Let

(5.13)
$$P_t^{\lambda} f(x) = (2\pi)^{-n} \int e^{ix \cdot \xi} b_0(t, x, \xi) \beta_2(|\xi|_{g(x)}/\lambda) \hat{f}(\xi) d\xi.$$

Notice that $b_0(t, x, \xi)$ behaves like an elliptic symbol in S^0 near $|\xi|_{g(x)} \sim \lambda$ when $t \leq \lambda^{-1}$, one can construct a microlocal inverse of P_t^{λ} as

(5.14)
$$E_t^{\lambda} f(x) = (2\pi)^{-n} \int e^{ix\cdot\xi} b_0(t, x, \xi)^{-1} \beta_2(|\xi|_{g(x)}/\lambda) \hat{f}(\xi) d\xi.$$

Since $t \leq \lambda^{-1}$, we have the kernel estimate

$$|E_t^{\lambda}(x,y)| \lesssim \lambda^n (1+\lambda|x-y|)^{-N}.$$

Then by Young's inequality for $1 \le p \le \infty$

$$||E_t^{\lambda}||_{L^p(M)\to L^p(M)}\lesssim 1.$$

Then we can write

$$E_t^{\lambda} P_t^{\lambda} f(x) = f(x) + (2\pi)^{-n} \int e^{ix \cdot \xi} (1 - \beta_2 (|\xi|_{g(x)}/\lambda)^2) \hat{f}(\xi) d\xi + (2\pi)^{-n} \int e^{ix \cdot \xi} q_{\lambda}(t, x, \xi) \hat{f}(\xi) d\xi$$
$$:= f(x) + R_t^{\lambda} f(x) + Q_t^{\lambda} f(x)$$

where the symbol $q_{\lambda} \in S^{-1}$ with $|\xi| \approx \lambda$. Then integration by parts gives the kernel estimate

$$|Q_t^{\lambda}(x,y)| \lesssim \lambda^{n-1} (1+\lambda|x-y|)^{-N}, \ \forall N.$$

By Young's inequality, this implies

(5.15)
$$||Q_t^{\lambda}||_{L^p(M) \to L^p(M)} \lesssim \lambda^{-1}.$$

Since $R_t^{\lambda}\beta_1(P/\lambda)$ is smoothing, we have

$$||R_t^{\lambda} f_{\lambda}||_{L^{\infty}(M)} = ||R_t^{\lambda} \beta_1(P/\lambda) f_{\lambda}||_{L^{\infty}(M)} \lesssim \lambda^{-N} ||f_{\lambda}||_{L^1(M)}.$$

Thus,

$$||P_t^{\lambda} f_{\lambda}||_{L^p(M)} \gtrsim ||E_t^{\lambda} P_t^{\lambda} f_{\lambda}||_{L^p(M)}$$

$$\geq ||f_{\lambda}||_{L^p(M)} + O(\lambda^{-1})||f_{\lambda}||_{L^p(M)}$$

Combing the estimates above, we obtain (5.12) for λ larger than a constant λ_0 . For $\lambda < \lambda_0$, we can apply the exponential lower bound in Theorem 1 and the maximum principle, since

$$||f_{\lambda}||_{L^{1}(M)} \approx ||f_{\lambda}||_{L^{2}(M)} \approx ||f_{\lambda}||_{L^{\infty}(M)}$$

in this case. \Box

Proof of Theorem 3. Take $\delta_0 = \delta/2$, we split the L^p norm into two parts:

$$\|\mathcal{H}f_{\lambda}\|_{L^{p}(\Omega)}^{p} = \|\mathcal{H}f_{\lambda}\|_{L^{p}(\Omega_{\delta_{0}})}^{p} + \|\mathcal{H}f_{\lambda}\|_{L^{p}(\Omega_{\delta_{0}}^{c})}^{p}.$$

For the lower bound, we only need to consider the contribution of the boundary part. If $\lambda \geq \delta_0^{-1}$, then by Lemma 6 and the co-area formula we have

(5.16)
$$\int_{\Omega_{\delta_0}^c} |\mathcal{H}f_{\lambda}|^p = \int_0^{\delta_0} \int_{\Sigma_t} |\mathcal{H}f_{\lambda}|^p d\omega_t dt$$
$$\gtrsim \int_0^{\lambda^{-1}} dt \|f_{\lambda}\|_{L^p(M)}^p$$
$$\gtrsim \lambda^{-1} \|f_{\lambda}\|_{L^p(M)}^p.$$

The case that $\lambda < \delta_0^{-1}$ can be handled similarly since δ_0 is a constant. For the upper bound, the interior part follows from Theorem 2 and the maximum principle, since

$$\|\mathcal{H}f_{\lambda}\|_{L^{\infty}(\Omega_{\delta_0})} \lesssim \lambda^{-N} \|f_{\lambda}\|_{L^1(M)}, \ \forall N.$$

The boundary part follows from Theorem 2 and the co-area formula, since for $1 \le p \le \infty$

$$\|\mathcal{H}f_{\lambda}\|_{L^{p}(\Sigma_{t})} \lesssim (e^{-c\lambda t} + \lambda^{-N})\|f_{\lambda}\|_{L^{p}(M)}, \ \forall N.$$

6. Applications to Steklov eigenfunctions and numerical approximation

Over the past two decades, efficient spectral methods have been developed in the field of computational mathematics, for the approximation of solutions to Laplace's equations using Steklov eigenfunctions. See e.g., Auchmuty [2, 3, 4, 5, 6], Auchmuty-Cho [7, 8], Auchmuty-Rivas [9], Cho [14], Cho-Rivas [15], Klouček-Sorensen-Wightman [29]. Among the key problems in this area are the error estimates for approximations and the orthogonality properties of the Steklov eigenfunctions. In this section, we shall use the results established in the preceding sections to address these issues.

We first prove the almost-orthogonality in Theorem 4.

Proof of Theorem 4. Suppose $\mu \geq \lambda$. If $\mu \leq 10$ or $|\lambda - \mu| \leq 10$, (1.32) just follows from Cauchy-Schwarz, Lemma 3 and Theorem 2. In the case of $\mu \geq 10$ and $\lambda \leq 10$, since the derivatives of the kernel of $\chi(P - \lambda)$ are bounded by constants independent with λ , we have

$$\langle v_{\lambda}, w_{\mu} \rangle_{\Omega} = O(\mu^{-N}) ||f||_{L^{2}(M)} ||g||_{L^{2}(M)}, \ \forall N.$$

Now suppose $\lambda \ge 10$ and $|\lambda - \mu| > 10$. Let $\gamma = \lambda/\mu \in (0,1)$. As before, we only need to handle the contribution near boundary. It suffices to show that on each slice Σ_t

$$(6.1) |\langle v_{\lambda}(t), w_{\mu}(t) \rangle_{\Sigma_{t}}| \lesssim e^{-ct(\lambda+\mu)} (1+|\lambda-\mu|)^{-N} ||f||_{L^{2}(M)} ||g||_{L^{2}(M)}, \quad \forall N,$$

and then integration with respect to t gives the desired bound. By Lemma 3 and Theorem 2, it suffices to handle the kernel

$$B_{\lambda,\mu}(y,z) = (\lambda\mu)^{\frac{n-1}{2}} \int e^{i\mu(\gamma d_g(x,y) - d_g(x,z))} a_{\lambda}(x,y) \overline{a_{\mu}(x,z)} dx$$

since the contributions from other terms are $O(\mu^{-N})$. By Cauchy-Schwarz, we only need to prove

(6.2)
$$\iint |B_{\lambda,\mu}(y,z)|^2 dy dz \lesssim (1+|\lambda-\mu|)^{-N}, \ \forall N,$$

Fix any $\varepsilon \in (0, \frac{1}{100})$. Let ∇ be the covariant derivative with respect to the metric g. Then we have $|\nabla_x d(x,y)|_{g(x)} = 1$. For convenience we take geodesic normal coordinates at x. If $\gamma < 1 - \varepsilon$, then

$$|\nabla_x(\gamma d_g(x,y) - d_g(x,z))| \ge \varepsilon.$$

Then integration by parts gives $B_{\lambda,\mu}(y,z) = O(\mu^{-N})$.

Suppose
$$1 - \varepsilon < \gamma < 1$$
. If $|\nabla_x d_a(x, y) - \nabla_x d_a(x, z)| > 2\varepsilon$, then

$$|\nabla_x(\gamma d_g(x,y) - d_g(x,z))| \ge |\nabla_x(d_g(x,y) - d_g(x,z))| - (1-\gamma) \ge \varepsilon.$$

Then integration by parts gives $B_{\lambda,\mu}(y,z) = O(\mu^{-N})$. Suppose $|\nabla_x d_g(x,y) - \nabla_x d_g(x,z)| < 2\varepsilon$. We may assume that these two unit vectors are close to $e_1 = (1,0,...,0)$. Then

$$|\nabla_x d_q(x,y) - \nabla_x d_q(x,z)| \gtrsim |y'-z'|$$

if $y = (y_1, y')$ and $z = (z_1, z')$. So we have

$$|\nabla_x(\gamma d_g(x,y) - d_g(x,z))| \gtrsim (1-\gamma) + |y'-z'|.$$

Then integration by parts gives

$$|B_{\lambda,\mu}(y,z)| \lesssim \mu^{n-1} (1+|\lambda-\mu|)^{-N} (1+\mu|y'-z'|)^{-N}, \ \forall N.$$

Then a direct calculation gives (6.2).

Proof Sketch of Theorems 5 and 6. The proofs of Theorems 5 and 6 are similar, so we only give a proof sketch. We can reduce the estimates in Ω to the bounds on each slice Σ_t , which follow from Lemma 3 and the arguments in [13] and [11, 12]. The power gains in Theorems 5 and 6 come from the integration of the exponential decay factor in the kernel formula in Lemma 3.

Now we shall discuss the numerical approximation problems of harmonic functions with various boundary conditions. Klouček-Sorensen-Wightman [29, Section 4.3] and Auchmuty-Cho [8, Sections 4 and 6] have obtained some basic error estimates and provided accompanying numerics. As an application of our results, we provide both L^2 and pointwise error estimates. For $x \in \Omega$, let

$$d(x) = \operatorname{dist}(x, \partial\Omega) = \operatorname{dist}(x, M).$$

Let $\{e_j\}_{j=0}^{\infty}$ be the orthonormal DtN eigenbasis in $L^2(M)$ associated with the Steklov eigenvalues

$$0 = \lambda_0 < \lambda_1 \le \lambda_2 \le ...,$$

which are arranged in increasing order and we account for multiplicity. The first eigenvalue $\lambda_0 = 0$ is simple since Ω is connected. Let $u_j = \mathcal{H}e_j$ be the associated Steklov eigenfunctions.

Theorem 7. Let u be the solution to Laplace equation (1.1) with Dirichlet boundary condition. Let $k \in \mathbb{N}_+$, $\lambda = \lambda_{k+1}$, and $c_j = \langle f, e_j \rangle_M$. Let $\tilde{f}_k = \sum_{j \leq k} c_j e_j$ and $\tilde{u}_k = \sum_{j \leq k} c_j u_j$. Then we have the L^2 error estimates

(6.3)
$$||u - \tilde{u}_k||_{L^2(\Omega)}^2 \lesssim \lambda^{-1} \left(||f||_{L^2(M)}^2 - ||\tilde{f}_k||_{L^2(M)}^2 \right)$$

and the pointwise error estimates

$$(6.4) |u(x) - \tilde{u}_k(x)|^2 \lesssim \left((\lambda + d(x)^{-1})^n e^{-\lambda d(x)} + \lambda^{-N} \right) \left(||f||_{L^2(M)}^2 - ||\tilde{f}_k||_{L^2(M)}^2 \right), \quad \forall N.$$

The bound (6.3) immediately follows from (1.28), since

$$||f||_{L^{2}(M)}^{2} - ||\tilde{f}_{k}||_{L^{2}(M)}^{2} = \sum_{\lambda_{i} > \lambda} |c_{j}|^{2} = ||f - \tilde{f}_{k}||_{L^{2}(M)}^{2}.$$

To get (6.4), we use Theorem 2 and Hörmander's L^{∞} bound (1.19). Indeed, for $0 < c < \frac{3}{4}$ we have

$$|u(x) - \tilde{u}_k(x)| \leq \sum_{2^{\ell} \geq \lambda} |\sum_{\lambda_j \approx 2^{\ell}} c_j u_j|$$

$$\lesssim \sum_{2^{\ell} \geq \lambda} (e^{-cd(x)2^{\ell}} 2^{\frac{n}{2}\ell} + 2^{-N\ell}) (\sum_{\lambda_j \approx 2^{\ell}} |c_j|^2)^{\frac{1}{2}}$$

$$\lesssim \left(\sum_{2^{\ell} \geq \lambda} (e^{-2cd(x)2^{\ell}} 2^{n\ell} + 2^{-2N\ell})\right)^{\frac{1}{2}} (\sum_{\lambda_j \geq \lambda} |c_j|^2)^{\frac{1}{2}}.$$

Then a direct calculation gives (6.4).

Next, let $b \ge 0$ be a fixed number and we consider the Laplace equation with Neumann (b = 0) or Robin (b > 0) boundary condition

(6.5)
$$\begin{cases} \Delta u(x) = 0, \ x \in \Omega, \\ \partial_{\nu} u(x) + bu(x) = f(x), \ x \in \partial \Omega = M. \end{cases}$$

Theorem 8. Let u be the solution to Laplace equation (6.5) with Neumann or Robin boundary condition. Let $k \in \mathbb{N}_+$, $\lambda = \lambda_{k+1}$, and $c_j = \langle f, e_j \rangle_M$. Let $\tilde{f}_k = \sum_{j \leq k} c_j e_j$. When b = 0, let

$$\tilde{u}_k = \sum_{1 \le j \le k} \frac{c_j}{\lambda_j} u_j.$$

When b > 0, let

$$\tilde{u}_k = \sum_{j \le k} \frac{c_j}{\lambda_j + b} u_j.$$

Then we have the L^2 error estimates

(6.6)
$$||u - \tilde{u}_k||_{L^2(\Omega)}^2 \lesssim \lambda^{-3} \Big(||f||_{L^2(M)}^2 - ||\tilde{f}_k||_{L^2(M)}^2 \Big).$$

and the pointwise error estimates

$$(6.7) |u(x) - \tilde{u}_k(x)|^2 \lesssim \left((\lambda + d(x)^{-1})^{n-2} e^{-\lambda d(x)} + \lambda^{-N} \right) \left(||f||_{L^2(M)}^2 - ||\tilde{f}_k||_{L^2(M)}^2 \right), \quad \forall N.$$

These estimates immediately follow from the same proof as Theorem 7 and the observation that

(6.8)
$$c_j = \langle f, e_j \rangle_M = \langle \partial_\nu u + bu, e_j \rangle_M = (\lambda_j + b) \langle u, e_j \rangle_M.$$

7. Appendix: Proof of Lemma 5

We prove Lemma 5 in the appendix. We essentially follow the strategy in the proof of [35, Theorem 4.3.1]. But Lemma 5 is more precise than [35, Theorem 4.3.1], and we need to carefully handle the dependence on λ . So we shall give a detailed proof for completeness.

Let $\mu_1 = \max\{\mu, 0\}$. We fix $\rho \in \mathcal{S}(\mathbb{R})$ with $\hat{\rho}$ supported in $\{|t| < 1\}$ and equal to 1 in $\{|t| < \frac{1}{2}\}$. We decompose

$$m_{\lambda}(P) = (2\pi)^{-1} \int \hat{\rho}(t) \hat{m}_{\lambda}(t) e^{itP} dt + (2\pi)^{-1} \int (1 - \hat{\rho}(t)) \hat{m}_{\lambda}(t) e^{itP} dt$$

:= $\tilde{m}_{\lambda}(P) + r_{\lambda}(P)$.

Here $r_{\lambda}(P)$ is a smoothing operator, since

$$\begin{split} |r_{\lambda}(\tau)| &= |(2\pi)^{-1} \int (1-\hat{\rho}(t)) \hat{m}_{\lambda}(t) e^{it\tau} dt| \\ &= |(2\pi)^{-1} \int \int (1-\hat{\rho}(t)) m_{\lambda}(s) e^{-its} e^{it\tau} ds dt| \\ &= C_N |\int \int (1-\hat{\rho}(t)) t^{-N} m_{\lambda}^{(N)}(s) e^{-its} e^{it\tau} ds dt| \\ &\lesssim \int (1+|s-\tau|)^{-N_1} |m_{\lambda}^{(N)}(s)| ds \\ &\lesssim \int (1+|s-\tau|)^{-N_1} (1+|s|)^{\mu_1-N} ds \\ &\lesssim (1+\tau)^{\mu_1-N} \end{split}$$

and its derivatives also satisfy similar bounds.

Without loss of generality, we assume the parametrix for half-wave kernel ([35, Theorem 4.1.2]) holds for $|t| \le 1$, that is

$$e^{itP}(x,y) = Q(t,x,y) + R(t,x,y)$$

where

$$Q(t,x,y) = (2\pi)^{-n} \int e^{i\varphi(x,y,\xi) + itp(y,\xi)} q(t,x,y,\xi) d\xi$$

and $R(t,x,y) \in C^{\infty}([-1,1] \times M \times M)$ and Q(t,x,y) is supported in a small neighborhood of the diagonal in $M \times M$. Moreover, $\varphi \in S^1$ and satisfies

$$\varphi(x, y, \xi) = (x - y) \cdot \xi + O(|x - y|^2 |\xi|)$$

and

$$q(0, x, x, \xi) - 1 \in S^{-1}$$
.

The integral of R(t) is a smoothing operator since

$$\left| \int \hat{\rho}(t)\hat{m}_{\lambda}(t)R(t,x,y)dt \right| = \left| \int \int m_{\lambda}(s)\hat{\rho}(t)R(t)e^{-its}dtds \right|$$

$$\lesssim \int |m(s/\lambda)|(1+|s|)^{-N}ds$$

$$\lesssim 1$$

and its derivatives also satisfy similar bounds.

Next, we consider the integral of Q(t, x, y). By Taylor expansion, we have

$$q(t) = q(0) + tq'(0) + \dots + \frac{t^N}{N!}q^{(N)}(0) + \frac{t^{N+1}}{N!} \int_0^1 q^{(N+1)}(\theta t)(1-\theta)^N d\theta.$$

For the remainder term, it contributes the kernel

(7.1)
$$\int_0^1 \iint \hat{\rho}(t) \hat{m}_{\lambda}(t) e^{i\varphi(x,y,\xi) + itp(y,\xi)} t^{N+1} q^{(N+1)}(\theta t, x, y, \xi) d\xi dt (1-\theta)^N d\theta.$$

Let

$$E_{\lambda,\theta}(x,y,\xi) = \int \hat{\rho}(t)\hat{m}_{\lambda}(t)e^{itp(y,\xi)}t^{N+1}q^{(N+1)}(\theta t,x,y,\xi)dt$$

Since $p(y,\xi) \approx |\xi|$ and $(\hat{\rho}q^{(N+1)}(\theta \cdot))^{\vee} \in \mathcal{S}$, we have

$$|E_{\lambda,\theta}(x,y,\xi)| \approx \left| \int \hat{\rho}(t) (m_{\lambda}^{(N+1)})^{\wedge}(t) e^{itp(y,\xi)} q^{(N+1)}(\theta t, x, y, \xi) dt \right|$$

$$\approx |m_{\lambda}^{(N+1)} * (\hat{\rho}q^{(N+1)}(\theta \cdot))^{\vee}(p(y,\xi))|$$

$$\lesssim \int \lambda^{-\mu} (\lambda + |\tau|)^{\mu - N - 1} (1 + |p(y,\xi) - \tau|)^{-N_1} d\tau$$

$$\lesssim (1 + |\xi|)^{\mu_1 - N - 1}$$

Similarly,

$$|\partial_{x,y}^{\beta}\partial_{\xi}^{\alpha}E_{\lambda,\theta}(x,y,\xi)| \lesssim (1+|\xi|)^{\mu_1-N-1-|\alpha|}, \quad \forall \alpha.$$

Thus, (7.1) is the kernel of an operator in OPS^{μ_1-N-1} by [35, Theorem 3.2.1].

For $\ell = 0, 1, 2, ...N$, we have

$$\iint \hat{\rho}(t)\hat{m}_{\lambda}(t)e^{i\varphi(x,y,\xi)+itp(y,\xi)}t^{\ell}q^{(\ell)}(0,x,y,\xi)d\xi dt$$

$$= \int \tilde{m}_{\lambda}^{(\ell)}(p(y,\xi))q^{(\ell)}(0,x,y,\xi)e^{i\varphi(x,y,\xi)}d\xi$$

$$= \int m_{\lambda}^{(\ell)}(p(y,\xi))q^{(\ell)}(0,x,y,\xi)e^{i\varphi(x,y,\xi)}d\xi + R_{\ell,\lambda}(x,y)$$

where

$$|R_{\ell,\lambda}(x,y)| = \left| \int r_{\lambda}^{(\ell)}(p(y,\xi))q^{(\ell)}(0,x,y,\xi)e^{i\varphi(x,y,\xi)}d\xi \right|$$

$$\lesssim \int (1+|\xi|)^{-N}d\xi$$

$$\lesssim 1$$

since $m_{\lambda} - \tilde{m}_{\lambda} = r_{\lambda} \in \mathcal{S}$. Its derivatives also satisfy similar bounds.

Let $\varphi_0 = \varphi$ and $\varphi_1 = (x - y) \cdot \xi$ and $\varphi_t = (1 - t)\varphi_0 + t\varphi_1$ with $t \in [0, 1]$. Let $P(x, y, \xi) = m_{\lambda}^{(\ell)}(p(y, \xi))q^{(\ell)}(0, x, y, \xi)$. Let

$$(P_t u)(x) = (2\pi)^{-n} \iint e^{i\varphi_t(x,y,\xi)} P(x,y,\xi) u(y) d\xi dy.$$

Since

(7.2)
$$|\partial_{\xi}^{\alpha}(\varphi_0 - \varphi_1)| \lesssim |x - y|^2 |\xi|^{1 - |\alpha|}, \ \forall \alpha,$$

we have

$$|\nabla_{\xi}\varphi_t| \gtrsim |x-y|$$

near the diagonal.

By (7.2), we can define
$$\phi(x, y, \xi) = (\varphi_1 - \varphi_0)/|x - y|^2 \in S^1$$
. So
$$\partial_t^j P_t u = (2\pi)^{-n} \iint e^{i\varphi_t} (i(\varphi_1 - \varphi_0))^j P(x, y, \xi) u(y) d\xi dy$$

and then

$$\begin{split} \partial_t^j|_{t=1} P_t u &= (2\pi)^{-n} \iint e^{i(x-y)\cdot\xi} (i(\varphi_1 - \varphi_0))^j P(x, y, \xi) u(y) d\xi dy \\ &= (2\pi)^{-n} (-1)^j \iint e^{i(x-y)\cdot\xi} \Delta_{\xi}^j \Big((i\phi(x, y, \xi))^j P(x, y, \xi) \Big) u(y) d\xi dy \\ &:= (2\pi)^{-n} (-1)^j \iint e^{i(x-y)\cdot\xi} Q_j(x, y, \xi) u(y) d\xi dy \end{split}$$

By Taylor formula we have

$$Q_j(x, y, \xi) = \sum_{|\alpha| \le N} \frac{1}{\alpha!} \partial_y^{\alpha}|_{y=x} Q(x, y, \xi) \cdot (y - x)^{\alpha} + R_N(x, y, \xi)$$

where R_N vanishes of order N+1 when x=y. For the remainder term, we use the reproducing operator (see [36, Lemma 4.1.1])

$$Le^{i\phi} = e^{i\phi}, \text{ if } L = \frac{1}{i} \frac{\nabla \phi}{|\nabla \phi|^2} \cdot \nabla$$

to rewrite the kernel

$$\int e^{i(x-y)\cdot\xi} R_N(x,y,\xi) d\xi = \int e^{i(x-y)\cdot\xi} (L^*)^{N+1} R_N(x,y,\xi) d\xi$$

Since that the adjoint of L is

$$L^*f = i\frac{(x-y)\cdot\nabla_{\xi}f}{|x-y|^2}$$

we have

$$|(L^*)^{N+1}R_N(x,y,\xi)| \lesssim (1+|\xi|)^{\mu_1-\ell-N-1}$$

Similarly,

$$|\partial_{x,y}^{\beta}\partial_{\xi}^{\alpha}(L^*)^{N+1}R_N(x,y,\xi)| \lesssim (1+|\xi|)^{\mu_1-\ell-N-1-|\alpha|}, \ \forall \alpha,\beta.$$

So the remainder term is an operator in $OPS^{\mu_1-\ell-N-1}$. Thus, by [35, (3.1.11)] the operator $\partial_t^j|_{t=1}P_t$ is a pseudo-differential operator with symbol

$$Q_j(x,\xi) \sim (-1)^j \sum_{\alpha} \frac{1}{\alpha!} D_{\xi}^{\alpha} \partial_y^{\alpha}|_{y=x} Q_j(x,y,\xi)$$

modulo a smoothing operator. Then by Taylor formula

$$P_0 = \sum_{i=0}^{N} \frac{(-1)^j}{j!} \partial_t^j |_{t=1} P_t + \frac{(-1)^{N-1}}{N!} \int_0^1 t^N \partial_t^{N+1} P_t dt$$

For the remainder term, we use the reproducing operator $Le^{i\varphi_t} = e^{i\varphi_t}$ and rewrite the kernel

$$\int e^{i\varphi_t} (i(\varphi_1 - \varphi_0))^{N+1} P(x, y, \xi) d\xi = \int e^{i\varphi_t} (L^*)^{2N+2} ((i(\varphi_1 - \varphi_0))^{N+1} P(x, y, \xi)) d\xi$$
$$:= \int e^{i\varphi_t} E_N(x, y, \xi) d\xi$$

Since the adjoint of L is

$$L^*f = i\sum_{j} \partial_{\xi_j} \left(\frac{\partial_{\xi_j} \varphi_t}{|\nabla_{\xi} \varphi_t|^2} \cdot f \right)$$

we have the symbol

$$|E_N(x,y,\xi)| \lesssim (1+|\xi|)^{\mu_1-\ell-N-1}$$

Similarly,

$$|\partial_{x,y}^{\beta}\partial_{\xi}^{\alpha}E_{N}(x,y,\xi)| \lesssim (1+|\xi|)^{\mu_{1}-\ell-N-1-|\alpha|}, \quad \forall \alpha, \beta.$$

So the remainder term is an operator in $OPS^{\mu_1-\ell-N-1}$ by [35, Theorem 3.2.1]. Thus, P_0 is a pseudo-differential operator with symbol given by the formal series

$$Q(x,\xi) \sim \sum_{j=0}^{\infty} \frac{1}{j!} Q_j(x,\xi)$$

modulo a smoothing operator.

Thus, $m_{\lambda}(P)$ is a pseudo-differential operator with symbol given by the formal series

$$M_{\lambda}(x,\xi) \sim \sum_{\ell=0}^{\infty} \sum_{j=0}^{\infty} \sum_{\alpha} \frac{1}{\ell! j! \alpha!} D_{\xi}^{\alpha} \partial_{y}^{\alpha}|_{y=x} \Delta_{\xi}^{j} \Big((i(\phi(x,y,\xi))^{j} m_{\lambda}^{(\ell)}(p(y,\xi)) q^{(\ell)}(0,x,y,\xi) \Big)$$

modulo a smoothing operator. Here $q^{(\ell)}(0) \in S^0$ and

$$\phi(x, y, \xi) = (\varphi_1 - \varphi_0)/|x - y|^2 \in S^1.$$

We can further simplify the formal series. For convenience, we use q_k to denote some symbol in S^k and it may vary from line to line. Then

$$\begin{split} M_{\lambda}(x,\xi) &\sim \sum_{\ell=0}^{\infty} \sum_{j=0}^{\infty} \sum_{\alpha} \sum_{0 \leq \alpha_{1} \leq \alpha} \sum_{0 \leq \alpha_{2} \leq \alpha+2j} m_{\lambda}^{(\ell+\alpha_{1}+\alpha_{2})}(p(x,\xi)) q_{\alpha_{1}+j-(\alpha+2j-\alpha_{2})}(x,\xi) \\ &= \sum_{\ell=0}^{\infty} \sum_{j=0}^{\infty} \sum_{\alpha} \sum_{0 \leq k \leq 2\alpha+2j} m_{\lambda}^{(\ell+k)}(p(x,\xi)) q_{k-\alpha-j}(x,\xi) \\ &= \sum_{\ell=0}^{\infty} \sum_{j=0}^{\infty} \sum_{\alpha} \sum_{0 \leq k \leq \min(\ell,2\alpha+2j)} m_{\lambda}^{(\ell)}(p(x,\xi)) q_{k-\alpha-j}(x,\xi) \\ &= \sum_{\ell=0}^{\infty} m_{\lambda}^{(\ell)}(p(x,\xi)) \tilde{q}_{\ell}(x,\xi) \end{split}$$

where $\tilde{q}_{\ell} \in S^{\lfloor \ell/2 \rfloor}$. The leading term only comes from the term with $\ell = j = \alpha = 0$, so it is exactly

$$m_{\lambda}(p(x,\xi))q(0,x,x,\xi)$$

which is $m_{\lambda}(p(x,\xi)) \mod S^{\mu_1-1}$.

References

- Almgren, Frederick J., Jr. Dirichlet's problem for multiple valued functions and the regularity of mass minimizing integral currents. Minimal submanifolds and geodesics (Proc. Japan-United States Sem., Tokyo, 1977), pp. 1–6 North-Holland Publishing Co., Amsterdam-New York, 1979
- [2] Auchmuty, Giles. Steklov eigenproblems and the representation of solutions of elliptic boundary value problems, Numer. Funct. Anal. Optim., 25 (2004), 321-348.
- [3] Auchmuty, Giles. Spectral characterization of the trace spaces $H^s(\partial\Omega)$. SIAM J. Math. Anal. 38 (2006), no. 3, 894–905.
- [4] Auchmuty, Giles. The S.V.D. of the Poisson kernel. J. Fourier Anal. Appl. 23 (2017), no. 6, 1517–1536.
- [5] Auchmuty, Giles. Steklov representations of Green's functions for Laplacian boundary value problems. Appl. Math. Optim. 77 (2018), no. 1, 173–195.
- [6] Auchmuty, Giles. Robin approximation of Dirichlet boundary value problems. Numer. Funct. Anal. Optim. 39 (2018), no. 10, 999–1010.
- [7] Auchmuty, Giles; Cho, Manki. Boundary integrals and approximations of harmonic functions. Numer. Funct. Anal. Optim. 36 (2015), no. 6, 687–703.
- [8] Auchmuty, Giles; Cho, Manki. Steklov approximations of harmonic boundary value problems on planar regions.
 J. Comput. Appl. Math. 321 (2017), 302–313.
- [9] Auchmuty, Giles; Rivas, M. A. Laplacian eigenproblems on product regions and tensor products of Sobolev spaces.
 J. Math. Anal. Appl. 435 (2016), no. 1, 842–859.
- [10] Boutet de Monvel, Louis. Boundary problems for pseudo-differential operators. Acta Math. 126 (1971), no. 1-2, 11-51.
- [11] Burq N, Gérard P, Tzvetkov N. Bilinear eigenfunction estimates and the nonlinear Schrödinger equation on surfaces. Inventiones mathematicae. 2005 Jan;159(1):187-223.
- [12] Burq N, Gérard P, Tzvetkov N. Multilinear estimates for the Laplace spectral projectors on compact manifolds. Comptes Rendus Mathematique. 2004 Mar 1;338(5):359-64.
- [13] Burq, N., Gérard, P. and Tzvetkov, N., Restrictions of the Laplace-Beltrami eigenfunctions to submanifolds, Duke Math. J. 138 (2007), 445–486.
- [14] Cho, Manki. Steklov expansion method for regularized harmonic boundary value problems. Numer. Funct. Anal. Optim. 41 (2020), no. 15, 1871–1886.

- [15] Cho, Manki and Rivas, Mauricio A. On the L²-orthogonality of Steklov eigenfunctions. Proceedings of the 2021 UNC Greensboro PDE Conference, 45–58. Electron. J. Differ. Equ. Conf., 26
- [16] Colbois, Bruno; Girouard, Alexandre; Gordon, Carolyn; Sher, David. Some recent developments on the Steklov eigenvalue problem. Rev. Mat. Complut. 37 (2024), no. 1, 1–161.
- [17] Daudé, T., Helffer, B., Nicoleau, F.: Exponential localization of Steklov eigenfunctions on warped product manifolds: the flea on the elephant phenomenon. Ann. Math, Québec (2021)
- [18] Di Cristo, M., and L. Rondi. Interior decay of solutions to elliptic equations with respect to frequencies at the boundary. Indiana University Mathematics Journal 70.4 (2021): 1303-1334.
- [19] Galkowski, J., Toth, J.A.. Pointwise bounds for Steklov eigenfunctions. J. Geom. Anal. 29(1), 142–193 (2019)
- [20] Galkowski, J., Toth, J.A.. Lower bounds for Steklov eigenfunctions. Pure Appl. Math. Q. 19 (2023), no. 4, 1873–1898.
- [21] Garofalo, Nicola, and Lin, Fang-Hua. Monotonicity properties of variational integrals, A_p weights and unique continuation. Indiana University Mathematics Journal 35, no. 2 (1986): 245-268.
- [22] Girouard, A., Polterovich, I.: Spectral geometry of the Steklov problem (survey article). J. Spectr. Theory 7(2), 321–359 (2017)
- [23] Helffer, B., Kachmar, A.: Semi-classical edge states for the robin laplacian. Mathematika 68(2), 454-485 (2022)
- [24] Hislop, Peter D., and Carl V. Lutzer. Spectral asymptotics of the Dirichlet-to-Neumann map on multiply connected domains in R^d. Inverse Problems 17.6 (2001): 1717.
- [25] Hörmander, Lars. The spectral function of an elliptic operator. Acta Math. 121 (1968), 193–218.
- [26] Hörmander, Lars. The Analysis of Linear Partial Differential Operators III Pseudo-Differential Operators, Springer-Verlag, Berlin, 1994.
- [27] R. Hu. L^p norm estimates of eigenfunctions restricted to submanifolds. Forum Math., 6:1021–1052, 2009.
- [28] X. Huang, Y. Sire, X. Wang and C. Zhang. Sharp L^p estimates and size of nodal sets for generalized Steklov eigenfunctions, arXiv preprint arXiv:2301.00095
- [29] Klouček, Petr; Sorensen, Danny C.; Wightman, Jennifer L. The approximation and computation of a basis of the trace space H^{1/2}. J Sci Comput. 2007; 32(1):73–108.
- [30] Lee, John M.; Uhlmann, Gunther. Determining anisotropic real-analytic conductivities by boundary measurements. Comm. Pure Appl. Math. 42 (1989), no. 8, 1097–1112.
- [31] Levitin, M., Parnovski, L., Polterovich, I., Sher, D.A.: Sloshing, Steklov and corners: asymptotics of Steklov eigenvalues for curvilinear polygons. Proc. Lond. Math. Soc. (3) 125(3), 359–487 (2022)
- [32] Lin, Fang-Hua. Nodal sets of solutions of elliptic and parabolic equations. Communications on Pure and Applied Mathematics 44, no. 3 (1991): 287-308.
- [33] Polterovich, I., Sher, D.A., Toth, J.A.: Nodal length of Steklov eigenfunctions on real-analytic Riemannian surfaces. J. Reine Angew. Math. 754, 17–47 (2019)
- [34] Sogge, Christopher D. Concerning the L^p norm of spectral clusters for second-order elliptic operators on compact manifolds. J. Funct. Anal. 77 (1988), no. 1, 123–138.
- [35] Sogge, Christopher D. Fourier integrals in classical analysis. Vol. 210. Cambridge University Press, 2017.
- [36] Sogge, Christopher D. Hangzhou lectures on eigenfunctions of the Laplacian, volume 188 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2014.
- [37] Tataru, Daniel. On the regularity of boundary traces for the wave equation. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 26 (1998), no. 1, 185–206.
- [38] Taylor, Michael. Partial differential equations II: Qualitative studies of linear equations. Vol. 116. Springer Science and Business Media, 2013.

DEPARTMENT OF MATHEMATICS, HUNAN UNIVERSITY, CHANGSHA, HN 410012, CHINA

Email address: xingwang@hnu.edu.cn

MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING, BJ 100084, CHINA

 $Email\ address: {\tt czhang98@tsinghua.edu.cn}$