On the image of a curve in a normal surface by a plane projection [∗]

F. Delgado † H. Maugendre ‡

Abstract

We consider a finite analytic morphism $\varphi = (f, g)$ defined from a complex analytic normal surface (Z, z) to \mathbb{C}^2 . We describe the topology of the image by φ of a reduced curve on (Z, z) by means of iterated pencils defined recursively for each branch of the curve from the initial one $\langle f, g \rangle$. This result generalizes the one obtained in a previous paper for the case in which (Z, z) is smooth and the curve irreducible. As a consequence of the methods we can describe also the topological type of the discriminant curve of φ , in particular the topological type of each branch of the discriminant can be obtained from the map without the previous knowledge of the critical locus.

Keywords Topological type - Pencils - Analytic morphisms - Discriminant curve - Critical locus.

Introduction

Let (Z, z) be the germ of a complex analytic normal surface and let $\varphi =$ $(f,g) : (Z, z) \to (\mathbb{C}^2, 0)$ be the germ of a finite complex analytic morphism. Let $\gamma \subset (Z, z)$ be the germ of a reduced curve, $\gamma = \bigcup_{i=1}^r \gamma_i$, with γ_i a branch in (Z, z) , and $\delta = \bigcup_{i=1}^r \delta_i$ with $\varphi(\gamma_i) = \delta_i, 1 \leq i \leq r$. If (Z, z) is equal to $(\mathbb{C}^2, 0)$ and γ is irreducible, in [4] we have described the topology of the plane

[∗]Math. Subject Class. 14H20, 32S05, 32S15, 32S45, 32S55

[†]Partially supported by grant PID2022-138906NB-C21 funded by MCIN/AEI/ 10.13039/501100011033 and by ERDF "A way of making Europe". The author is thankful to the Institut Fourier, Université de Grenoble I for hospitality.

Address: IMUVA (Instituto de Investigación en Matemáticas). University of Valladolid. Valladolid, Spain. E-mail: fdelgado@uva.es.

[‡]Partially supported by the ANR LISA Project ANR-17-CE40-0023.

Address: Institut Fourier, Université de Grenoble I, BP 74, F-38402 Saint-Martin d'Hères, France. E-mail: helene.maugendre@univ-grenoble-alpes.fr

branch δ by using iterated pencils of functions defined recursively from the initial one $\Phi = \langle f, g \rangle$. The purpose of this article is to show that this result can be generalized to the case of normal surfaces and when the curve γ is not necessarily irreducible; it means that we can describe the topological type of each branch $\delta_i, 1 \leq i \leq r$ (section 1), and moreover give the intersection multiplicities between each pair of branches δ_i , δ_j (section 2), which completely describe the topology of δ .

In section 1, we show that, as in the plane case, the topology of the image by φ of an irreducible curve γ can be determined by the degree of the restriction map $\varphi|_{\gamma} : \gamma \to \delta = \varphi(\gamma)$ and intersection multiplicities between γ and some particular fibres of a sequence of iterated pencils constructed recursively from $\langle f, g \rangle$ (see 1.5). This result is stated in Theorem 1.8. In section 2 we treat the case when γ is reduced but no more irreducible. It is well-known that the topological type of its plane curve image by φ , $\delta = \bigcup_{i=1}^r \delta_i$, is enterely determined by the topological type of each irreducible component $\delta_i = \varphi(\gamma_i)$ and the intersection multiplicity between each pair $(\delta_i, \delta_j), j \neq i$. As the topological type of each irreducible component image δ_i has been treated in section 1, it leaves to compute the intersection multiplicity between each pair of branches of δ . This result is stated in Theorem 2.12. The proof consists in first computing such intersection multiplicities when φ is the identity map from the plane to the plane (Theorem 2.8), then using the projection formula for the intersection multiplicity (see Proposition 1.1) and Theorem 2.8 we can compute the intersection multiplicity between δ_i and δ_j for the general case of a finite map $\varphi: (Z, z) \to (\mathbb{C}^2, 0).$

A remarkable fact is that, for the computation of the intersection multiplicity, one needs more precise informations than for the topological type of the branches (see the Example 2.11). Namely informations coming from the branches of the fibres and not only the contacts of the branches of γ with the whole fibres.

In [4] the process was in particular applied to the description of the branches of the discriminant locus of $\varphi : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$. To do that the key point was the study of the special fibres of the pencil $\langle f, g \rangle$ and their relations with the critical locus of φ developed in [2] as well as the relation with the behaviour of the Hironaka quotients established in [9] and [10]. For a normal singularity (Z, z) and a finite map $\varphi = (f, g) : (Z, z) \to (\mathbb{C}^2, 0)$ the relation of the special fibres with the critical locus was generalized in [5]. (For a reduced surface singularity the behaviour of the generic and special fibres has been recently treated in [1]). These relations allow us to extend the results of [4] and so apply the results of Sections 1 and 2 to describe the topological type of the discriminant curve of φ . This is the purpose of Section 3. In relation with the study of the discriminant curve one has to mention the recent paper [7] from García-Barroso and Popescu-Pampu in which they show that the Newton polynomial of the discriminant curve essentially depends only on the pencil $\langle f, q \rangle$ (see also [8] for the plane case). Althougth the results are different, some of the technics used there are very close to the ones we use here.

1 Case of an irreducible curve

Let $\gamma \subset (Z, z)$ be an irreducible curve (in short, a branch) and take $\eta : (\mathbb{C}, 0) \to$ (γ, z) a parametrization (uniformization) of (γ, z) . Let $\{\ell = 0\} \subset Z$ be the curve defined by the analytic function $\ell : (Z, z) \to (\mathbb{C}, 0)$. We define the intersection multiplicity of $\{\ell = 0\}$ and γ at z, denoted $I_z(\ell, \gamma)$, by:

$$
I_z(\ell, \gamma) = \text{ ord }_t(\ell \circ \eta(t)).
$$

Notice that the normalization of the local ring $\mathcal{O}_{\gamma,z}$ is isomorphic to $\mathbb{C}{t}$ and $\nu_{\gamma}(-) := I_z(-, \gamma)$ is the valuation ν_{γ} defined by γ . Moreover, we write $I_z(\ell, \gamma) = \infty$ if γ is a branch of $\{\ell = 0\}$, i.e. if the function ℓ vanishes on γ . If (Z, z) is smooth then $I_z(\ell, \gamma)$ coincides with the usual intersection multiplicity between the curve germs $L := \{ \ell = 0 \}$ and γ . In this case we use also the notation $[L, \gamma] = [\{\ell = 0\}, \gamma]$ (and $[L, \gamma]_z$ if we want to specify the point) to indicate the intersection multiplicity at $z \in Z$.

Let $\varphi^* : \mathcal{O}_{\mathbb{C}^2,0} \to \mathcal{O}_{Z,z}$, be the ring homomorphism associated to φ , i.e $h \mapsto \varphi^* h = h \circ \varphi$ for $h \in \mathcal{O}_{\mathbb{C}^2,0}$. If $\xi = \{h = 0\} \subset (\mathbb{C}^2,0)$ we use also $\varphi^* \xi$ to denote the local curve in (Z, z) defined by φ^*h . Notice that $\varphi^*\xi$ is a local Cartier divisor in (Z, z) .

Let $\gamma \subset (Z, z)$ be a branch and $\varphi(\gamma) = \delta$. The restriction map $\varphi|_{\gamma} : \gamma \to \delta$ has degree $k \geq 1$ and we define the *direct image* of γ as $\varphi_* \gamma := k \cdot \delta$.

Proposition 1.1 (Projection formula). *One has:*

$$
I_z(\varphi^*h, \gamma) = I_0(h, \varphi_*\gamma) (= kI_0(h, \delta))
$$
\n(1)

Proof. Let **p** be the prime ideal on $\mathcal{O}_{Z,z}$ defining the curve germ γ on (Z, z) , then the prime ideal $\mathfrak{q} = (\varphi^*)^{-1}(\mathfrak{p})$ is the ideal defining $\delta = \varphi(\gamma) \subset (\mathbb{C}^2, 0)$ and one has the finite extension of one-dimensional local rings $\mathcal{O}_{\delta,0} = \mathcal{O}_{\mathbb{C}^2,0}/\mathfrak{q} \hookrightarrow$ $\mathcal{O}_{\gamma,0} = \mathcal{O}_{Z,z}/\mathfrak{p}$. The degree of the above extension is just k: the one of the restriction map $\varphi|_{\gamma} : \gamma \to \delta$. Let $\mathbb{C}{t}$ (respectively $\mathbb{C}{\{\tau\}}$) be isomorphic to the normalization of $\mathcal{O}_{\gamma,z}$ (respectively to the one of $\mathcal{O}_{\delta,0}$), one has $\mathbb{C}\{\tau\} \subset \mathbb{C}\{t\}$ and $\tau(t) \in \mathbb{C}{t}$ has order k. In fact one could take $\tau = t^k$ with a convenient choice of the uniformizing parameters τ and t. Now, let $h \in \mathcal{O}_{\mathbb{C}^2,0}$ be an holomorphic function on \mathbb{C}^2 . One has that $\text{ord}_t(\varphi^*h(t)) = \text{ord}_t(h(t^k)) = k$. ord_{τ} $(h(\tau))$. \Box Remark 1.2. The above formula (1) extends the classical projection formula for a map φ between smooth surfaces (see, e.g. [6]). Notice also that one could extend the above formula for any local Weil divisor $\gamma = \sum_{i=1}^{\ell} n_i \gamma_i$ on (Z, z) . Finally, the formula (1) is obviously also true for a finite morphism $\varphi: (Z, z) \to (X, x)$ between normal surface singularities.

1.3. Let $\pi : (X, E) \to (Z, z)$ be a good resolution of (Z, z) , i.e. a resolution of the singularity (Z, z) such that the exceptional divisor $E = \bigcup_{\sigma \in \Gamma} E_{\sigma}$ is a union of smooth projective curves with normal crossings. For $\beta \in \Gamma$ and for each holomorphic function $h: (\mathbb{C}^2, 0) \to (\mathbb{C}, 0)$ let $\nu_\beta(h)$ denote the vanishing order of $\overline{h} = h \circ \pi : X \to \mathbb{C}$ along the irreducible exceptional curve E_{β} (ν_{β} is just the divisorial valuation defined by E_β). If $\gamma = \{h = 0\}$ is the curve defined by h we use also the notation $\nu(\gamma)$ instead $\nu(h)$, in particular this notation could be used for any curve germ if $(Z, z) = (\mathbb{C}^2, 0)$ and π is a modification of it.

The lifting $\overline{\psi} = \psi \circ \pi$ of the meromorphic function $\psi = g/f$ is a meromorphic function defined in a suitable neighbourhood of the exceptional divisor E but in a finite number of points. The irreducible component E_β of E is called a *dicritical* component of the pencil $\Phi = \langle f, g \rangle$ (or of the meromorphic function ψ) if the restriction of $\overline{\psi}: E_{\beta} \to \mathbb{C} \mathbb{P}^1$ is everywhere defined and not constant.

A good resolution of the pencil Φ is a good resolution $\pi : (X, E) \to (Z, z)$ of (Z, z) such that ψ is everywhere defined. In such a case, the zero locus ${h = 0}$ of the elements $h \in \Phi$ are equisingular and called the generic fibres of Φ, excepted a finite number of them called the special fibres. Moreover from [5], we know that the strict transforms of the generic fibres by π intersect the exceptional divisor smoothly and transversaly at smooth points of the dicritical components. In fact, a good resolution of (Z, z) is a good resolution of Φ if and only if is a resolution of all the fibers of Φ but a finite number.

Lemma 1.4. With the above notations. Let $h : (Z, z) \to (\mathbb{C}, 0)$ be an analytic *function,* $\eta = \{h = 0\}$ *and let* $(\gamma, z) \subset (Z, z)$ *be a branch such that its strict transform by* π , $\widetilde{\gamma}$, *intersects* $E_{\alpha} \subset E$ *at a smooth point* P. Then

$$
I_z(h,\gamma) = [\widetilde{\eta}, \widetilde{\gamma}]_P + [E_\alpha, \widetilde{\gamma}]_P \nu_\alpha(h) .
$$

In particular, if $\tilde{\eta} \cap \tilde{\gamma} = \emptyset$ *, then* $I_z(h, \gamma) = [E_{\alpha}, \tilde{\gamma}]_P \nu_{\alpha}(h)$.

Proof. The divisor (\overline{h}) defined by $\overline{h} = h \circ \sigma$ on X could be written as

$$
(\overline{h}) = (\widetilde{h}) + \sum_{\beta \in \Gamma} \nu_{\beta}(h) E_{\beta} ,
$$

where the local part $(\tilde{h}) = \tilde{\eta}$ is the strict transform of the germ $\{h = 0\}$ by π . For each $\sigma \in \Gamma$ one has the known Mumford formula (see [11]):

$$
(\overline{h}) \cdot E_{\sigma} = (\widetilde{h}) \cdot E_{\sigma} + \sum_{\beta} \nu_{\beta}(h) (E_{\beta} \cdot E_{\sigma}) = 0.
$$
 (2)

(Here " \cdot " stands for the intersection form on the smooth surface X).

Applying the equation (2) to the total transforms of γ and η we have:

$$
I_z(h,\gamma) = (\overline{\eta}) \cdot (\overline{\gamma}) = (\widetilde{h}) \cdot \widetilde{\gamma} + \sum_{\beta \in \Gamma} \nu_\beta(h) (E_\beta \cdot \widetilde{\gamma}) = [\widetilde{\eta}, \widetilde{\gamma}]_P + \nu_\alpha(h) [E_\alpha, \widetilde{\gamma}]_P.
$$

The above result will be used in the paper also in the case of $(Z, z) =$ $(\mathbb{C}^2, 0)$. Note that in this case every curve is defined by a function (i.e. is a Cartier divisor).

1.5 (Iterated pencils). We denote $\Phi = \langle g, f \rangle = \{ag - bf : a, b \in \mathbb{C}\}\$ the pencil defined by f and g. Let us assume that $I_z(g, \gamma) \geq I_z(f, \gamma)$. If $I_z(g, \gamma) =$ ∞ then $\varphi(\gamma)$ is the x axis $\{y=0\}$ and the procedure stops here. Otherwise, let $q/p = I_z(g, \gamma)/I_z(f, \gamma) \geq 1$ with $gcd(p, q) = 1$. Let Φ_1 be the pencil generated by g^p and f^q . From Proposition 1 of [5] the intersection multiplicity $I_z(ag^p - bf^q, \gamma)$ is constant (and equal to 1) for all $w = (a : b) \in \mathbb{CP}^1$, except for one value w_0 in \mathbb{CP}^1 . With the hypothesis that $q/p \geq 1$, it is obvious that we can write $w_0 = (1 : a_1)$. We denote $g_1 = g^p - a_1 f^q$.

Let $f_1 := f^q$, and $\varphi_1 = (f_1, g_1) : (Z, z) \longrightarrow (\mathbb{C}^2, 0)$. If $I_z(g_1, \gamma) = \infty$ then γ is a branch of ${g_1 = 0}$ and in this case one has $\varphi_1(\gamma) = L, L = {y_1 = 0}$ for $y_1 = y^p - a_1 x^q$, so $\varphi(\gamma) = \{y^p - a_1 x^q = 0\}$ and we stop here.

Else, we consider $\Phi_1 = \langle g_1, f_1 \rangle$ and we can define a new rational number $q_1/p_1 = I_z(g_1, \gamma)/I_z(f_1, \gamma) > 1$. In this case we can proceed to construct $\Phi_2 = \langle g_2, f_2 \rangle$, $g_2 \in \Phi_2$, and q_2/p_2 as we did for $\Phi_1, g_1, q_1/p_1$.

Recursively, we define a sequence $\mathcal{P}(f,g,\gamma) = \{(\Phi_i,g_i,q_i/p_i) : i \geq 0\}$ where, $(\Phi_0, g_0, q_0/p_0) = (\Phi, g, q/p)$ and, for $i \geq 1$, $\Phi_i = \langle g_i, f_i \rangle = \langle g_{i-1}^{p_{i-1}} \rangle$ $\vert_{i-1}^{p_{i-1}}, f_{i-1}^{q_{i-1}}\rangle$ is a pencil of functions on (Z, z) , g_i is the unique function of Φ_i such that $I_z(g_i, \gamma) > I_z(h, \gamma)$ ($h \in \Phi_i$ a generic function) and $q_i/p_i = I_z(g_i, \gamma)/I_z(f_i, \gamma)$ 1, with $gcd(q_i, p_i) = 1$. We denote also $g_{-1} = f$ and by ξ_i the curve on (Z, z) defined by $g_i = 0$. Note that, for $i \geq 1$, $f_i = f^{q_0 \cdots q_{i-1}}$ and also that the sequence is infinite unless there exists $l \in \mathbb{N}$ such that $q_l/p_l = \infty$ in which case the sequence is finite and ends at l.

Definition 1.6. The *iterated sequence of pencils* of the branch γ (with respect to $\varphi = (f, g)$ is the sequence $\mathcal{P}(f, g, \gamma)$ defined above.

Remark 1.7. Along the paper we use frequently the case $(Z, z) = (\mathbb{C}^2, 0)$ and $\varphi = (x, y)$ the identity map from $(\mathbb{C}^2, 0)$ to $(\mathbb{C}^2, 0)$. In this case we use $\{(\Lambda_i, y_i, q_i/p_i) : i \geq 0\}$ to denote the iterated pencil of a branch $\delta \subset (\mathbb{C}^2, 0)$. We will also denote λ_i the curve defined by $y_i = 0$.

Theorem 1.8. Let k be the degree of the restriction map $\varphi_{|\gamma} : \gamma \longrightarrow \delta$ and let $\varphi_*\gamma = k \cdot \delta$ be the direct image of γ . The sequence $\{I_z(g_i, \gamma) : i = -1, 0, \ldots\}$ *and the integer* k *determine the topological type of* δ*.*

And as an easy consequence one has:

Corollary 1.9. *The sequence of rational numbers* $\{q_i/p_i, i \geq 0\}$ *determines the topological type of* δ*.*

Proof. The proof of Theorem 1.8 and Corollary 1.9 repeats the scheme used in the Theorem 1 (see also Corollary 1) of $[4]$. The first step consists in the proof that the topological type (say e.g. the semigroup of values) of δ could be determined with the described procedure in the particular case in which we take the identity map $(x, y) : (\mathbb{C}^2, 0) \to (\mathbb{C}^2, 0)$. For the sake of completeness we will describe the procedure, the proof could be read in [4].

Assume that δ is not a coordinate axis and $I_0(x, \delta) \leq I_0(y, \delta)$. Let us consider the iterated sequence of pencils $\mathcal{P}(x, y, \delta) = \{(\Lambda_i, y_i, q_i/p_i) : i \geq 0\}$ of δ with respect to the identity.

Let m_{-1}, m_0, \ldots be the sequence of positive integers defined recursively as follows: $m_{-1} = I_0(x, \delta)$, $m_0 = I_0(y, \delta)$ and, if we assume that m_1, \ldots, m_i have been defined, then

$$
m_{i+1} := d_i m_i + I_0(y_{i+1}, \delta) - p_i I_0(y_i, \delta) , \qquad (3)
$$

where $d_i = \gcd(m_{-1}, \ldots, m_{i-1})/\gcd(m_{-1}, \ldots, m_i)$.

Theorem 2 in [4] says that the sequence m_{-1}, m_0, \ldots generates the semigroup of values of δ (see also the Remark below).

Now, using the ring homomorphism $\varphi^* : \mathcal{O}_{\mathbb{C}^2,0} \to \mathcal{O}_{Z,z}$ associated to φ , one can recover by pullback the iterated sequence of pencils $\mathcal{P}(f, g, \gamma)$ on (Z, z) : that means $\varphi^*(\Lambda_i) = \Phi_i$, $\varphi^*(y_i) = g_i$ for $i \geq 0$. The projection formula 1.1 applied to the equations 3 gives:

$$
m_{-1} = I_0(x, \delta) = I_z(f, \gamma)/k
$$
, $m_0 = I_0(y, \delta) = I_z(g, \gamma)/k$

and for $i \geq 1$:

$$
m_{i+1} = d_i m_i + I_0(y_{i+1}, \delta) - p_i I_0(y_i, \delta) = d_i m_i + \frac{I_z(g_{i+1}, \gamma)}{k} - p_i \frac{I_z(g_i, \gamma)}{k}.
$$

Thus, we recover the semigroup of values of the branch δ .

The proof of the Corollary is the same as the one of Corollary 1 in $[4]$, however is also included in the proof of Theorem 3.4. \Box

Remark 1.10. In the proof of Theorem 1.8 we recover the semigroup of values, in the same way one can recover the characteristic exponents of δ . Let $\mu_i =$ $I_z(g_i, \gamma)$ for $i \geq -1$. Let $m_{-1} = \widetilde{m}_{-1} = I_z(f, \gamma)/k$, $m_0 = \widetilde{m}_0 = I_z(g_0, \gamma)/k$ and for $i \geq 1$,

$$
m_{i+1} = d_i m_i + \frac{\mu_{i+1}}{k} - p_i \frac{\mu_i}{k} = d_i m_i + \frac{\mu_{i+1}}{k} - q_i \frac{I_z(f_i, \gamma)}{k}
$$

$$
\widetilde{m}_{i+1} = \widetilde{m}_i + \frac{\mu_{i+1}}{k} - p_i \frac{\mu_i}{k} = \widetilde{m}_i + \frac{\mu_{i+1}}{k} - q_i \frac{I_z(f_i, \gamma)}{k}
$$

where, $d_i = \gcd(m_{-1}, \ldots, m_{i-1})/\gcd(m_{-1}, \ldots, m_i)$. Note that

 $e_i := \gcd(m_{-1}, \ldots, m_i) = \gcd(\widetilde{m}_{-1}, \ldots, \widetilde{m}_i) = \gcd(\mu_{-1}, \ldots, \mu_i)/k$.

As a consequence, denoting $\epsilon_i := \gcd(\mu_{-1}, \ldots, \mu_i)$, one has also $d_i = \epsilon_{i-1}/\epsilon_i$.

From [4] one knows that the minimal set of generators $\{\beta_0, \ldots, \beta_g\}$ of the semigroup of the plane germ δ is a subset of $\{m_i \mid i \geq -1\}$. More specifically $\{\beta_0, \ldots, \beta_g\} = \{m_{-1}\} \cup \{m_i : d_i > 1\}.$ In the same way, the set of characteristic exponents $\{\beta_0, \ldots, \beta_g\}$ of δ is a subset of $\{\widetilde{m}_i \mid i \geq -1\}$, namelly the set $\{\beta_0, \ldots, \beta_g\} = \{\widetilde{m}_{-1}\} \cup \{\widetilde{m}_i : d_i > 1\}.$

In particular the computation of the sequences stops in a finite number of steps, just when $e_i = 1$ or equivalently $gcd(\mu_{-1}, \ldots, \mu_i) = k$. Notice that $k = \text{gcd}\{\mu_i : i \geq -1\},\$ i.e. the set $\mu = \{\mu_i : i \geq -1\}$ generates a finitely generated subsemigroup of $k\mathbb{Z}$. So theoretically one can recover the integer k and so the topological type of δ from μ , however it can not be computed in practice.

2 Case of several branches

In this section we treat the case in which $\gamma \subset (Z, z)$ is a reduced but not irreducible curve in (Z, z) . We denote $\gamma = \bigcup_{i=1}^r \gamma_i$, with γ_i a branch in (Z, z) , and its image by φ is $\delta = \bigcup_{i=1}^r \delta_i$ in such a way that $\varphi(\gamma_i) = \delta_i, 1 \le i \le r$. Pay attention that we could have $\delta_i = \delta_j$ for some $i \neq j$.

To describe the topological type of δ it suffices to describe the topological type of each branch $\delta_i, 1 \leq i \leq r$, and give the intersection multiplicities $[\delta_i, \delta_j]$ between pairs of different branches δ_i, δ_j . Thus it is enough to resolve the case of two branches.

Along the section $\gamma, \gamma' \subset (Z, z)$ are two irreducible curves in (Z, z) and we will assume that $\varphi(\gamma) = \delta$ is not equal to $\varphi(\gamma') = \delta'$.

2.1 Case of $(\mathbb{C}^2, 0)$

We will start, as in the irreducible case, describing the intersection multiplicity $[\delta, \delta']$ by the use of the iterated pencils starting with the pencil $\Lambda_0 = \langle x, y \rangle$ on \mathbb{C}^2 corresponding to the identity map from $(\mathbb{C}^2,0)$ to $(\mathbb{C}^2,0)$. So, we proceed by induction.

2.1.1 First step

Let assume that $I_0(\delta, y) \geq I_0(\delta, x)$ and let $I_0(\delta, y)/I_0(\delta, x) = q/p \geq 1$, with $gcd(p, q) = 1$. We have to consider several situations according to the value of $I_0(\delta', y)/I_0(\delta', x)$:

a) If $I_0(\delta', y) < I_0(\delta', x)$ then $\{x = 0\}$ is tangent to δ' and transversal to δ . So δ and δ' are transversal and we have $[\delta, \delta'] = I_0(\delta, x)I_0(\delta', y)$.

Otherwise we have $I_0(\delta', y)/I_0(\delta', x) = q'/p' \ge 1$, with $gcd(p', q') = 1$.

b) Let us suppose that $q'/p' \neq q/p$, and $q'/p' \geq 1$ and $q/p \geq 1$. Assume that $q'/p' > q/p \geq 1$.

Note that if $q/p = 1$ then δ and δ' are transversal and then

$$
[\delta, \delta'] = I_0(\delta, x)I_0(\delta', x) = I_0(\delta, y)I_0(\delta', x) < I_0(\delta', y)I_0(\delta, x) \; .
$$

Else we have $q'/p' > q/p > 1$ and in this case $\{y = 0\}$ is tangent to δ and δ' . Then, an easy computation (see also the proof of Proposition 2.3) allows to show that

$$
[\delta, \delta'] = \min \{ I_0(\delta, x) I_0(\delta', y), I_0(\delta, y) I_0(\delta', x) \} = I_0(\delta, y) I_0(\delta', x) .
$$

c) Endly, let us assume that $q/p = q'/p' \ge 1$. Let $\Lambda_1 = \langle y^p, x^q \rangle$ be the first iterated pencil. Notice that $\Lambda_1 = \Lambda'_1$. Taking into account the construction of (1.5), let $y_1 = y^p - ax^q \in \Lambda_1$, $a \neq 0$, be the only fibre of Λ_1 such that $I_0(\delta, y_1) > I_0(\delta, y^p) (= I_0(\delta, x^q))$ (resp. $y'_1 = y^p - a' x^q \in \Lambda_1, a' \neq 0$, the corresponding one for δ'). We have to distinguish two cases according to $a = a'$ or $a \neq a'$.

 $c-1) Case a \neq a'.$

Lemma 2.1. *If* $a \neq a'$ *then* $[\delta, \delta'] = I_0(\delta, x)I_0(\delta', y) = I_0(\delta, y)I_0(\delta', x)$ *.*

Proof. Let $\sigma: (X, E) \to (\mathbb{C}^2, 0)$ be the minimal resolution of Λ_1 . Notice that the fibers y_1, y'_1 are generic for $\Lambda_1 = \langle y^p, x^q \rangle$, in particular they are equisingular. The fact that $I_0(\delta, y_1) > I_0(\delta, y^p)$ implies that the strict transform of δ and y_1 by σ intersect in a point $P \in E_\alpha \subset E$. As y_1 is generic for Λ_1 it implies that E_{α} is a dicritical divisor for Λ_1 (it is the unique one in this particular case, see [2]) and P is a smooth point of E_{α} which is not a critical point. (In this case none smooth point of E_{α} is critical). In the same way one has that the strict transform of δ' and y'_1 by σ intersect in a point $P' \in E_\alpha \subset E$. As $a \neq a'$ we have $P \neq P'$ and it is known that $[\delta, \delta'] = I_0(\delta, x)I_0(\delta', y) = I_0(\delta, y)I_0(\delta', x)$. \Box

 $c-2) \text{Case } a = a'.$

Following the notations of the previous Lemma, as $gcd(p, q) = 1$, the fibres y_1 and y'_1 are irreducible, so the case $a = a'$ is equivalent to say $P = P'$ and $[\delta, \delta'] > I_0(\delta, x)I_0(\delta', y)$. In this case one has to iterate the process.

Remark 2.2. In all the described cases but in case c-2) we have proved that

$$
[\delta, \delta'] = \min \{ I_0(\delta, x) I_0(\delta', y), I_0(\delta, y) I_0(\delta', x) \} .
$$

On the other hand, in the case c-2) for the first iterated pencils $(\Lambda_1, y_1, q_1/p_1)$ and $(\Lambda'_1, y'_1, q'_1/p'_1)$ of δ and δ' respectively we have $\Lambda_1 = \Lambda'_1$, $y_1 = y'_1$ and $P = P'$. Moreover, for the same reasons as in **c-1**, the divisor E_{α} is a dicritical divisor of Λ_1 and $P \in E_\alpha$ is not a critical point.

2.1.2 General recursive step

For an index $i \geq 1$, let $(\Lambda_j, y_j, q_j/p_j)$ and $(\Lambda'_j, y'_j, q'_j/p'_j)$, $j \leq i$, be the sequence of iterated pencils for δ and δ' . Let us assume that, for $j \leq i$ one has $\Lambda_j = \Lambda'_j$ and $y_j = y'_j$ and moreover $q_j/p_j = q'_j/p'_j$ for $j < i$.

Let π : $(X, E) \to (\mathbb{C}^2, 0)$ be the minimal resolution of the pencil $\Lambda_i =$ $\langle x_i, y_i \rangle$. Let δ (resp. δ') be the strict transform of δ (resp. δ') by π . We suppose that $\delta \cap E = \delta' \cap E$ and we denote Q this point.

Following the same idea of the proof of Theorem 2 in [4], we assume the following properties:

- 1. $Q = \tilde{\delta} \cap E = \tilde{\delta}' \cap E$ is a smooth point of the exceptional divisor E. So, there exists a (unique) irreducible component $E_{\alpha} \subset E$ such that $Q = \widetilde{\delta} \cap E_{\alpha} = \widetilde{\delta}' \cap E_{\alpha}.$
- 2. The irreducible component E_{α} is dicritical for the pencil Λ_i .
- 3. There exists a (unique) branch ζ_i of $\lambda_i = \{y_i = 0\}$ such that it is a curvette at the point Q (i.e. its strict transform by π is smooth and transversal to E_{α} at the point Q.

Let $\sigma: (X', E') \to (\mathbb{C}^2, 0)$ be the composition of π with the minimal modification of (X, Q) until the strict transform of δ and δ' by σ meets the exceptional locus E' at smooth points, P, P', and also such that the strict transforms of ζ_i by σ do not meet P nor P'.

Proposition 2.3. *Following the above notations we obtain :*

1. If $P \neq P'$ one has :

$$
[\delta, \delta'] = \min\{ [E_{\alpha}, \delta]_Q [\delta', \zeta_i], [E_{\alpha}, \delta']_Q [\delta, \zeta_i] \} .
$$

2. If $P = P'$ then $q_i/p_i = q'_i/p'_i$, $\Lambda_{i+1} = \Lambda'_{i+1}$, $y_{i+1} = y'_{i+1}$ and we proceed with a new iteration. Moreover one has $P = P'$ if and only if $[\delta, \delta'] >$ $[E_{\alpha}, \delta]_Q [\delta', \zeta_i] = [E_{\alpha}, \delta']_Q [\delta, \zeta_i].$

Remark 2.4. In [4] the next property is also included in the above list, as fourth property, for the recursive step:

 \ll Let $m_i = [\delta, \zeta_i]$, the set $\{m_{-1}, \ldots, m_{i-1}\}$ contains all the maximal contact values of δ smaller than m_i . \gg

In our case one can add the corresponding one for δ' , $m'_i = [\delta', \zeta_i]$. (See also the proof of Theorem 1.8 for the definition of the integers m_i .) Here we do not need it because the semigroup of each branch is already computed in Theorem 1.8. However it is interesting to notice that (see $[4]$): $[E_{\alpha}, \delta]_Q = \gcd(m_{-1}, m_0, \dots, m_{i-1}) := e_{i-1}$ and so, for $P \neq P'$:

$$
[\delta,\delta']=\min\{e_{i-1}m'_i,e'_{i-1}m_i\}
$$

The next lemma will be useful for the proof of Proposition 2.3.

Lemma 2.5. *With the above notations, we have the following equivalence:*

$$
\frac{q_i'}{p_i'} > \frac{q_i}{p_i} \Longleftrightarrow \frac{[\tilde{\delta}', \tilde{\zeta}_i]_Q}{[E_\alpha, \tilde{\delta}']_Q} > \frac{[\tilde{\delta}, \tilde{\zeta}_i]_Q}{[E_\alpha, \tilde{\delta}]_Q}.
$$

Proof. For each irreducible component ζ of λ_i different from ζ_i one has, from lemma 1.4, $[\delta, \zeta] = [E_{\alpha}, \delta]_Q \nu_{\alpha}(\zeta)$ and $[\delta', \zeta] = [E_{\alpha}, \delta']_Q \nu_{\alpha}(\zeta)$ and so

$$
[\delta, \zeta]/[E_{\alpha}, \delta]_Q = [\delta', \zeta]/[E_{\alpha}, \delta']_Q.
$$

As

$$
\frac{q_i}{p_i} = \frac{[\delta, \lambda_i]}{I_0(\delta, x_i)} = \frac{[\delta, \zeta_i]}{I_0(\delta, x_i)} + \sum_{\zeta \in \lambda_i, \ \zeta \neq \zeta_i} \frac{[\delta, \zeta]}{[E_\alpha, \widetilde{\delta}]_Q \ \nu_\alpha(x_i)}
$$

and

$$
\frac{q_i'}{p_i'} = \frac{[\delta', \lambda_i]}{I_0(\delta', x_i)} = \frac{[\delta', \zeta_i]}{I_0(\delta', x_i)} + \sum_{\zeta \in \lambda_i, \ \zeta \neq \zeta_i} \frac{[\delta', \zeta]}{[E_\alpha, \widetilde{\delta'}]_Q \ \nu_\alpha(x_i)},
$$

one has

$$
\frac{q_i'}{p_i'} > \frac{q_i}{p_i} \Longleftrightarrow \frac{[\delta', \zeta_i]}{I_0(\delta', x_i)} > \frac{[\delta, \zeta_i]}{I_0(\delta, x_i)} \Longleftrightarrow \frac{[\delta', \zeta_i]}{q_0 \dots q_{i-1} I_0(\delta', x)} > \frac{[\delta, \zeta_i]}{q_0 \dots q_{i-1} I_0(\delta, x)}
$$
\n
$$
\Longleftrightarrow \frac{[\widetilde{\delta}', \widetilde{\zeta}_i]_Q + [E_\alpha, \widetilde{\delta}']_Q \nu_\alpha(\zeta_i)}{[E_\alpha, \widetilde{\delta}']_Q} > \frac{[\widetilde{\delta}, \widetilde{\zeta}_i]_Q + [E_\alpha, \widetilde{\delta}]_Q \nu_\alpha(\zeta_i)}{[E_\alpha, \widetilde{\delta}]_Q}
$$
\n
$$
\Longleftrightarrow \frac{[\widetilde{\delta}', \widetilde{\zeta}_i]_Q}{[E_\alpha, \widetilde{\delta}']_Q} > \frac{[\widetilde{\delta}, \widetilde{\zeta}_i]_Q}{[E_\alpha, \widetilde{\delta}]_Q}.
$$

Proof of Proposition 2.3.

1) Let us assume $P \neq P'$. We start by computing $[\delta, \delta']_Q$. Taking into account that ζ_i and E_α are both smooth and intersect transversaly at Q , we can choose a pair of local coordinates (u, v) at (X, Q) in such a way that $u = 0$ and $v = 0$ are the (local) equations of E_{α} and ζ_i respectively. Thus one has locally, at $Q \in X$, exactly the same situation as in the First step 2.1.1 for the pencil $\langle u, v \rangle$ instead $\Lambda = \langle x, y \rangle$. Thus, we have:

$$
P \neq P' \Longleftrightarrow [\widetilde{\delta}, \widetilde{\delta}']_Q = \min\{[E_\alpha, \widetilde{\delta}]_Q[\widetilde{\delta}', \widetilde{\zeta}_i]_Q, [E_\alpha, \widetilde{\delta}']_Q[\widetilde{\delta}, \widetilde{\zeta}_i]_Q\}.
$$

Now one has to compute $[\delta, \delta']$. By lemma 1.4 we have :

$$
[\delta, \delta'] = [\tilde{\delta}, \tilde{\delta}']_Q + [E_\alpha, \tilde{\delta}']_Q \nu_\alpha(\delta) = [\tilde{\delta}, \tilde{\delta}']_Q + [E_\alpha, \tilde{\delta}]_Q \nu_\alpha(\delta').
$$

Let assume that $[E_\alpha, \delta']_Q[\delta, \zeta_i]_Q < [E_\alpha, \delta]_Q[\delta', \zeta_i]_Q$, (from lemma 2.5 this is equivalent to $q'_i/p'_i > q_i/p_i$ then

$$
[\delta, \delta'] = [E_{\alpha}, \tilde{\delta}']_Q[\tilde{\delta}, \tilde{\zeta}_i]_Q + [E_{\alpha}, \tilde{\delta}']_Q \nu_{\alpha}(\delta)
$$

\n
$$
= [E_{\alpha}, \tilde{\delta}']_Q \left([\tilde{\delta}, \tilde{\zeta}_i]_Q + \nu_{\alpha}(\delta) \right)
$$

\n
$$
= [E_{\alpha}, \tilde{\delta}']_Q[\delta, \zeta_i] < [E_{\alpha}, \tilde{\delta}]_Q[\delta', \zeta_i].
$$

Notice that in case of $q_i/p_i = q'_i/p'_i$ one has $[\delta, \delta'] = [E_\alpha, \delta']_Q[\delta, \zeta_i] =$ $[E_{\alpha}, \delta]_Q[\delta', \zeta_i].$

2) Let $P = P' \in E_\beta \subset E'$. The strict transform of ζ_i by σ does not intersects E_β at P and obviously the same is true for all the branches of $\lambda_i = \{y_i = 0\}$. So by Lemma 1.4 we have $[\lambda_i, \delta] = [E_\beta, \delta]_P \nu_\beta(y_i)$ and $[\lambda_i, \delta'] =$ $[E_{\beta}, \delta']_P \nu_{\beta}(y_i)$. The same equalities are also true for x_i instead y_i . Then

$$
\frac{q_i}{p_i} = \frac{[\lambda_i, \delta]}{I_0(x_i, \delta)} = \frac{\nu_\beta(y_i)}{\nu_\beta(x_i)} = \frac{[\lambda_i, \delta']}{I_0(x_i, \delta')} = \frac{q_i'}{p_i'}
$$

and as a consequence $\Lambda_{i+1} = \Lambda'_{i+1}$. As $\nu_\beta(x_i^{q_i})$ $\binom{q_i}{i} = \nu_\beta(y_i^{p_i})$ i^{p_i} , the lifting of the meromorphic function $\psi = y_i^{p_i}$ $i^{p_i}/x_i^{q_i}$ is not equal to zero along E_β and has a zero at the point P, thus E_β is a dicritical divisor of Λ_{i+1} . As in the proof of Lemma 1 of [4] one can see that Q is not a critical point of Λ_{i+1} and so one has also that there exists an unique fibre $y_{i+1} = y_i^{p_i} - ax_i^{q_i} \in \Lambda_{i+1}$ $(a = \widetilde{\psi}(P) \in \mathbb{C}^*)$ such that $I_0(y_{i+1}, \delta) > I_0(y_i^{p_i})$ $\sum_{i=1}^{p_i}$, δ) and $I_0(y_{i+1}, \delta') > I_0(y_i^{p_i})$ i^{p_i}, δ'). The fact that E_β is dicritical and $P \in E_\beta$ is not a critical point implies that there exists an unique branch ζ_{i+1} of $\lambda_{i+1} = \{y_{i+1} = 0\}$ such that its strict transform is a curvette at the point P.

Notice that, if σ is not the minimal resolution of Λ_{i+1} then the new blowingups up to reach it do not affect the described situation on $P \in E_\beta$.

Notice also that in this case we have $[\delta, \delta'] > [E_\alpha, \delta']_Q[\delta, \zeta_i] = [E_\alpha, \delta]_Q[\delta', \zeta_i].$ \Box

Remark 2.6. Let us summarize the case when $P \neq P'$ and $q_i/p_i = q'_i/p'_i$. Note that at this step, $q_i/p_i = q'_i/p'_i$ is equivalent to say $\Lambda_{i+1} = \Lambda'_{i+1}$. In this case P and P' belong to the same dicritical component $E_\beta \subset E'$ of the minimal resolution of Λ_{i+1} . Let $y_{i+1} = y_i^{p_i} - ax_i^{q_i}$ $e_i^{q_i}, \lambda_a = \{y_{i+1} = 0\}$ (resp. $y'_{i+1} = y_i^{p_i} - a'x_i^{q_i}$ $\mathcal{L}_{i}^{q_i}, \lambda_{a'} = \{y'_{i+1} = 0\}$ be the unique fibre such that $I_0(\delta, \lambda_a) >$ $I_0(\delta,\lambda)$ (resp. $[\delta',\lambda_{a'}] > [\delta',\lambda]$), for any other fibre λ of Λ_{i+1} . Thus we have two possibilities:

Either $a \neq a'$, and in this case $y_{i+1} \neq y'_{i+1}$ and $\Lambda_{i+2} \neq \Lambda'_{i+2}$.

Otherwise $a = a'$. In this case there exists ζ, ζ' irreducible components of the same fibre λ_a such that the strict transform ζ of ζ (resp. ζ' of ζ') meets E_β at P (resp. at P'). By using lemma 1.4 it is easy to check that it is equivalent to

$$
\frac{[\delta,\zeta]}{[E_{\beta},\widetilde{\delta}]} > \frac{[\delta',\zeta]}{[E_{\beta},\widetilde{\delta}']} \left(\text{resp. } \frac{[\delta',\zeta']}{[E_{\beta},\widetilde{\delta}']} > \frac{[\delta,\zeta']}{[E_{\beta},\widetilde{\delta}]} \right) . \quad (*)
$$

As a consequence, always using lemma 1.4 we have $[\delta, \delta'] = \nu_\beta(\delta)[E_\beta, \tilde{\delta}']$ and as $\tilde{\delta} \cap \tilde{\zeta}' = \emptyset$ and ζ' is transversal to E_{β} , $\nu_{\beta}(\delta) = [\delta, \zeta']$ and we obtain

$$
[\delta,\delta']=[E_{\beta},\delta'][\delta,\zeta']\ .
$$

In the same way we have $[\delta, \delta'] = \nu_\beta(\delta')[E_\beta, \tilde{\delta}] = [E_\beta, \tilde{\delta}][\delta', \zeta].$

From the Proposition 2.3 and its proof it is obvious that the maximal integer κ such that the requirements of the General recursive step are satisfied can be determined in a more easy way: let $\mathcal{P}(x, y, \delta) = \{(\Lambda_i, y_i, q_i/p_i) : i \geq 0\}$ (resp. $\mathcal{P}(x, y, \delta') = \{(\Lambda'_i, y'_i, q'_i/p'_i) : i \ge 0\}$) be the sequence of iterated pencils for δ (resp. for δ'). Then one has:

Statement 2.7. *Let* κ *be the largest integer such that, if* π : $(X, E) \rightarrow (\mathbb{C}^2, 0)$ *is the minimal resolution of the pencil* Λ_{κ} , then the strict transforms of δ and δ' *intersects* E *at the same point* $Q \in E_\alpha \subset E$. Then E_α *is a dicritical component for* Λ_{κ} *and* Q *is not a critical point of* Λ_{κ} *.*

Moreover, if ζ *is the unique branch of* $\lambda_{\kappa} = \{y_{\kappa} = 0\}$ *such that its strict transform by* π *intersects* E *at the point* Q *one has:*

$$
[\delta, \delta'] = \min\{[E_{\alpha}, \widetilde{\delta}]_Q [\delta', \zeta], [E_{\alpha}, \widetilde{\delta}']_Q [\delta, \zeta]\} .
$$

Proof. The proof goes by induction on κ , the case $\kappa = 0$ is just the first step 2.1.1 and the inductive step is the General recursive step 2.1.2. □

Although the above statement completely solves the problem of determining the intersection multiplicity of δ and δ' , the computation of κ requires the resolution of the iterated pencils. Let us see that this computation can be set up in another way.

Let $\{m_i, i \geq -1\}$ and $\{m'_i, i \geq -1\}$ be the sequences defined in the proof of Theorem 1.8 and Remark 1.10 for δ and δ' . For $i \ge -1$, let $e_i = \gcd(m_{-1}, \ldots, m_i)$ and $e'_i = \gcd(m'_{-1}, \ldots, m'_i)$.

Theorem 2.8. *Let* ℓ *be the smallest integer such that there exists a branch* ζ *of the fibre* $\lambda_{\ell} = \{y_{\ell} = 0\}$ *of* Λ_{ℓ} *such that* $[\delta, \zeta]/I_0(\delta, x) \neq [\delta', \zeta]/I_0(\delta', x)$. *Then:*

$$
[\delta, \delta'] = \min\{e_{\ell-1}[\delta', \zeta], e'_{\ell-1}[\delta, \zeta]\}.
$$

Proof. Let κ be the integer defined in Statement 2.7, then one has that $[\delta, \zeta]/e_{i-1} = [\delta', \zeta]/e_{i-1}'$ for all branches ζ of the fibres $\{y_i = 0\}$ for $i < \kappa$. Let ℓ be equal to $\kappa + 1$ if we are in the case $a = a'$ described in Remark 2.6 and let $\ell = \kappa$ otherwise. Proposition 2.3 and Remark 2.6 implies that there exists a branch ζ of $\lambda_{\ell} = \{y_{\ell} = 0\}$ such that $[\delta, \zeta]/e_{i-1} \neq [\delta', \zeta]/e_{i-1}'$. Then, the integer ℓ could be defined as the smallest such that $[\delta, \zeta]/e_{i-1} \neq [\delta', \zeta]/e_{i-1}'$ for some branch ζ of λ_{ℓ} . The same results implies the stated equality for $[\delta, \delta']$.

Now, it is well known that, in our conditions, $d_i = e_{i-1}/e_i = e'_{i-1}/e'_i = d'_i$ for $i \leq \ell - 1$ (see e.g. [3]). Moreover, $e_{-1} = m_{-1} = I_0(\delta, x)$ and $e'_{-1} = m'_{-1} =$ $I_0(\delta', x)$. Then for the integer ℓ we have just defined above one has that also ℓ is the smallest integer such that $[\delta, \zeta]/I_0(\delta, x) \neq [\delta', \zeta]/I_0(\delta', x)$. \Box

Remark 2.9. Notice that the formula for $[\delta, \delta']$ in the above Theorem is not the same as in Remark 2.4. Let us assume that we are in the case $\ell = \kappa + 1$, then one has that $[\delta, \delta'] = e_{\kappa-1} m'_{\kappa} = e'_{\kappa-1} m_{\kappa}$ and also $[\delta, \delta'] = \min\{e_{\ell-1}[\delta', \zeta], e'_{\ell-1}[\delta, \zeta]\}.$ However $m'_{\ell} \neq [\delta', \zeta]$ because ζ does not go by P'. In fact it could happen even that $m'_{\ell}/e'_{\ell-1} = m_{\ell}/e_{\ell-1}$ (see the Example below). Thus in order to detect the integer ℓ (and so κ) it does not suffice with the sequences $\{m_i\}$ and $\{m'_i\}$.

Remark 2.10. If there exists an integer r such that $\Lambda_r \neq \Lambda'_r$ then there exists ζ , and ζ' branches of $\lambda_r = \{y_r = 0\}$ and $\lambda'_r = \{y'_r = 0\}$ such that $\delta \cap \zeta \neq \emptyset$ and $\delta' \cap \zeta' \neq \emptyset$ respectively. Then, one has:

$$
[\delta, \delta'] = e'_{r-1}[\delta, \zeta'] = e_{r-1}[\delta', \zeta].
$$

Note that the only condition is that $\Lambda_r \neq \Lambda'_r$. In particular the branches δ, δ' could be separated before the step r.

Example 2.11. The following example from [4] shows that the use we made of the branches of the fibres and not only of the fibres themselves is required. Let $\delta = \{4y^2 - 4x^3 - 4yx^3 + x^6 = 0\}, \delta' = \{4y^2 - 4x^3 + 4yx^3 + x^6 = 0\}.$ The Puiseux expansion of them are $x = t^2, y = t^3 + (1/2)t^6$ and $x = t^2, y = t^3 - (1/2)t^6$ respectively. Their intersection multiplicity $[\delta, \delta']$ is equal to 9. As one can prove in an easy way the sequence of pencils and fibres for both are the same (as well as the sequences $\{m_i\}$ and $\{m'_i\}$). For the first steps one founds:

$$
y_1 = y^2 - x^3
$$

\n
$$
y_2 = (y^2 - x^3)^2 - x^9
$$

\n
$$
y_3 = ((y^2 - x^3)^2 - x^9)^6 - (1/64)x^{63}
$$

\n
$$
y_4 = (((y^2 - x^3)^2 - x^9)^6 - (1/64)x^{63})^{42} - (3/256)^{42}(x^{63})^{43}
$$

and the sequences of $m_i = m'_i$ is $(m_{-1} = 2, m_0 = 3, m_1 = 9, m_2 = 12, m_3 = 15)$ and $(q_0/p_0 = q'_0/p'_0 = 3/2, q_1/p_1 = q'_1/p'_1 = 3/2, q_2/p_2 = q_2/p'_2 = 7/6, \ldots$. Note that $m_{-1} = 2$ and $m_0 = 3$ are enough to compute the semigroup of δ (see Corollary 1.9): the one generated by 2, 3. The same is true for δ' .

The fibre y_1 is irreducible. But at the second step, the fibre y_2 has two branches ζ_1, ζ_2 such that $[\delta, \zeta_1] = 12$ and $[\delta, \zeta_2] = 9$ and inversely $[\delta', \zeta_1] = 9$ and $[\delta', \zeta_2] = 12$. Thus following Theorem 2.8 we have that $\ell = 2$ and

Figure 1: Graph of the minimal resolution of $\delta \cup \delta' \cup \{y_2 = 0\}$

In such a way, as moreover $[\delta, \delta'] = 9$, at the sixth blow-up (the corresponding divisor E_6 is the dicritical one of the pencil Λ_2 , see Figure 1) the strict transforms of δ and δ' separate, as do the ones of ζ_1 and ζ_2 , while the stricts transforms of δ and ζ_1 , resp. of δ' and ζ_2 , separate three blow-ups later (the corresponding divisors E_9 and E'_9 are dicritical for the pencil Λ_3).

With the above notations it means that we have $P \neq P'$ at E_6 although the pencils of δ and δ' are the same.

The resolution graph of the minimal resolution of $\delta \cup \delta' \cup \{y_2 = 0\}$ is depicted in Figure 1.

2.2 Case of a normal singularity (Z, z)

Let γ, γ' be two irreducible curves in (Z, z) and $\varphi(\gamma) = \delta$ and $\varphi(\gamma') = \delta'$ their image in the plane \mathbb{C}^2 . We will assume that $\delta \neq \delta'$. Recall that $\varphi_* \gamma = k \delta$ and $\varphi_*(\gamma') = k'\delta'$ for some positive integers k, k' .

Let $\mathcal{P}(f,g,\gamma) = \{(\Phi_i, g_i, q_i/p_i) : i \geq 0\}$ and $\mathcal{P}(f,g,\gamma') = \{(\Phi'_i, g'_i, q'_i/p'_i) :$ $i \geq 0$ } be the iterated sequence of pencils of the branches γ and γ' w.r.t. $\varphi = (f, q).$

Theorem 2.12. *Let* ℓ *be the smallest integer such that there exists a branch* ρ *of the fibre* $\xi = \{g_\ell = 0\}$ *of* Φ_ℓ *such that*

$$
\frac{I_z(\gamma, \varphi^*(\varphi(\rho)))}{I_z(\gamma, f)} \neq \frac{I_z(\gamma', \varphi^*(\varphi(\rho)))}{I_z(\gamma', f)}.
$$

Then:

$$
[\delta, \delta'] = \min \left\{ \frac{e'_{\ell-1} I_z(\gamma, \varphi^*(\varphi(\rho)))}{k}, \frac{e_{\ell-1} I_z(\gamma', \varphi^*(\varphi(\rho)))}{k'} \right\}.
$$

Proof. Let $\mathcal{P}(x, y, \delta) = \{(\Lambda_i, y_i, q_i/p_i) : i \geq 0\}$ (resp. $\mathcal{P}(x, y, \delta') =$ $\{(\Lambda'_i, y'_i, q'_i/p'_i) : i \geq 0\}$) be the sequence of iterated pencils for δ (resp. for δ'). As a consequence of Proposition 1.1, for $j \geq 0$, $(\varphi^*(\Lambda_j), \varphi^* y_j, q_j/p_j) =$ $(\Phi_j, g_j, q_j/p_j).$

Let $\zeta \subset (\mathbb{C}^2, 0)$ be a plane branch. By the projection formula 1.1 one has $[\delta, \zeta] = I_z(\gamma, \varphi^* \zeta)/k$ and so

$$
\frac{[\delta,\zeta]}{I_0(\delta,x)} = \frac{I_z(\gamma,\varphi^*\zeta)}{I_z(\gamma,f)}.
$$

This equalities applied to the branches ζ of $\lambda_i = \{y_i = 0\}$ and both pairs (γ, δ) , (γ', δ') implies that the integer ℓ defined in Theorem 2.8 is charaterized as the one described in the statement.

Now, the equality for the intersection multiplicity of δ and δ' given in Theorem 2.8 is (by using the projection formula) the one described in the statement of the Theorem. \Box

Remark 2.13. Using remark 1.10, the expression of the formula of Theorem 2.12 in terms of intersections multiplicities is the following one :

$$
[\delta, \delta'] = \frac{1}{kk'} \min \left\{ \epsilon'_{\ell-1} I_z(\gamma, \varphi^*(\varphi(\rho))), \epsilon_{\ell-1} I_z(\gamma', \varphi^*(\varphi(\rho))) \right\}.
$$

3 The discriminant case

The results proved in [5] about the behaviour of the critical locus of the map φ and its relation with the special fibres of the pencil $\Phi = \langle f, g \rangle$ (see also [2] for the plane case) allows to determine the topology of the irreducible components of the discriminant curve, $D(\varphi)$, of the morphism φ . We recall that the discriminant curve is the image by φ of the critical locus, $C(\varphi)$, of the map φ (we consider only the one-dimensional components of the critical locus, so $C(\varphi)$ is the topological closure of the vanishing of the restriction of the jacobian determinant of (f, g) to $(Z, z) \setminus \{z\}$. Moreover, the sequence of rational numbers q_i/p_i , $i \geq 0$, for each branch of $C(\varphi)$ can be computed directly from the pencils without the knowledge of the concrete branches of $C(\varphi)$ (see [5]). We will show here how we obtain the whole topology type of the discriminant curve using only the construction of the pencils.

To make this paragraph self-contained, we recall here the construction of S (the set of sequences $(B_i)_{i\geq 0}$ associated to each branch of the discriminant curve) in a similar way to the one given in section 4 of [4].

3.1. Let π : $(X, E) \rightarrow (Z, z)$ be the minimal good resolution of (Z, z) which is also a resolution of the pencil $\langle g, f \rangle$ and of the curve $\{fg = 0\}$ (i.e. the minimal good resolution of the singularity (Z, z) such that $(g/f) \circ \pi$ is a morphism and the strict transform of ${fg = 0}$ is smooth and transversal to the exceptional locus, see $[5]$).

An irreducible component E_{α} of the exceptional divisor E is called a *nodal* component (called rupture component in [4]) if either E_{α} is non-rational or the number of connected components of $\overline{C}_{red} \backslash E_{\alpha}$ is at least three, where \overline{C}_{red} stands for the total transform of $C = \{fg = 0\}$ by π with reduced structure. A *nodal zone* R ⊂ E is a maximal connected union of irreducible exceptional components containing at least one nodal component and such that the ratio $\nu_{\alpha}(g)$ $\frac{\nu_{\alpha}(g)}{\nu_{\alpha}(f)}$ is constant for $E_{\alpha} \subset R$. The term r-nodal zone is used to indicate that $\frac{\nu_{\alpha}(g)}{\nu_{\alpha}(f)}=r.$

Let $R\mathcal{Z}$ be the set of all nodal zones R such that $Q(R) := \frac{\nu_{\alpha}(g)}{\nu_{\alpha}(f)} \neq 1$, $E_{\alpha} \subset$ R. Let $E^{(1)} = \bigcup_{\alpha} E_{\alpha} \subset E$ be the union of the irreducible components E_{α} of E such that $\frac{\nu_{\alpha}(g)}{\nu_{\alpha}(f)} = 1$. Notice that the union D of dicritical components of E is included in $E^{(1)}$. Let A be the set of connected components of $\overline{E^{(1)}\setminus \mathcal{D}}$ which contain a nodal component. For $A \in \mathcal{A}$ we define $Q(A) := \frac{\nu_{\alpha}(g)}{\nu_{\alpha}(f)} = 1$, $E_{\alpha} \subset A$. Moreover let $\mathcal{P}_{\mathcal{C}}$ be the set of points $P \in \mathcal{D}$ such that either P is a singular point of D or P is a smooth point of \overline{C}_{red} which is a critical point of the map $(g/f) \circ \pi_{|_{\mathcal{D}}} : \mathcal{D} \longrightarrow \mathbb{C} \mathbb{P}^1$. For $P \in \mathcal{P}_{\mathcal{C}}$ we put $Q(P) := 1$.

We denote $\mathcal{B}^0 = \mathcal{R} \mathcal{Z} \cup \mathcal{A} \cup \mathcal{P}_\mathcal{C}$ and fix $B \in \mathcal{B}^0$. We write $Q(B) = \frac{q}{p} \in \mathbb{Q}$ with $gcd(p, q) = 1$ and consider the pencil $\Phi_B = \langle g^p, f^q \rangle$. Notice that π is also

an embedded resolution of $\{f^q g^p = 0\}$. Thus by theorem 3 of [5] there exists a unique special fibre $\xi_B = \{g^p - af^q = 0\}$ of Φ_B whose strict transform $\tilde{\xi}_B$ by π intersects B .

Let us denote $\varphi_1 = (f_1, g_1) = (f^q, g^p - af^q)$ and σ_B the minimal sequence of blowing-ups of points such that $\pi_B := \sigma_B \circ \pi$ is an embedded resolution of ${f_1g_1 = 0}.$

One can construct the set $\mathcal{B}^0(\varphi_1)$ for φ_1 in the same way we have defined \mathcal{B}^0 for (f, g) and for each $B^1 \in \mathcal{B}^0(\varphi_1)$ we define $Q_1(B_1)$ as $Q(B_1)$ with respect to g_1, f_1 . Moreover we add the symbol B_{∞} when there exists a non reduced branch $r\zeta$ of $\{g_1 = 0\}$, $r > 1$, whose strict transform $r\zeta$ by π_B intersects σ_B^{-1} $B_B^{-1}(B)$. When B_{∞} exists we put $Q_1(B_{\infty}) := \infty$.

Let us denote :

$$
\mathcal{B}_{B}^{1} = \{ B_{1} \in \mathcal{B}^{0}(\varphi_{1}) / \sigma_{B}(B^{1}) \subset B \} \cup \{ B_{\infty} \}
$$

and we set $\mathcal{B}^1 = \cup_{B \in \mathcal{B}^0} \mathcal{B}_B^1$. Moreover one has the map $\psi_1 : \mathcal{B}^1 \longrightarrow \mathcal{B}^0$ defined by $\psi_1(B_1) = B$ if and only if $B_1 \in \mathcal{B}_B^1$. Thus we can follow the same process for each element $B_1 \in \mathcal{B}^1$ with $B_1 \neq B_\infty$ for any $B \in \mathcal{B}^0$ and so we inductively construct the collection of sets $\{\mathcal{B}^i\}_{i\geq 0}$ together with maps $\psi_i : \mathcal{B}^i \longrightarrow \mathcal{B}^{i-1}$ and $Q_i: \mathcal{B}^i \longrightarrow \mathbb{Q} \cup {\infty}$ in such a way that the definition of $\psi_i^{-1}(B)$ for $B \in \mathcal{B}^{i-1}$ (and the map $Q_{i|\psi_i^{-1}(B)}$) follows the same process described above for $B \in \mathcal{B}^0$.

In a such a way we obtain S the set of sequences $(B_i)_{i>0}$ such that $B_i \in \mathcal{B}^i$ and $\psi_i(B_i) = B_{i-1}$ for $i \geq 1$. We assume that if $B_i = (B_{i-1})_{\infty}$ then the sequence is finite and ends at B_i . For our purposes it is better to encode this information in the form of a weighted oriented graph $\mathcal T$ in the obvious form. That is, one puts a vertex for each $B \in \mathcal{B}^i$ $(i \geq 0)$ with weight equal to $Q_i(B)$. The set of oriented edges is the set of pairs $(B_{i-1}, B_i) \in \mathcal{B}^{i-1} \times \mathcal{B}^i$ such that $\psi_i(B_i) = B_{i-1}$. Thus, the graph $\mathcal T$ is the union of $\#\mathcal B^0$ trees, each one with root the corresponding element (vertex) of \mathcal{B}^0 . We will call the above defined graph the graph of φ . In this way, the set S corresponds to the maximal completely ordered (and weighted) subtrees of \mathcal{T} .

3.2 (The branches of the discriminant). Let $\mathcal{C}(\varphi)$ be the set of branches of $C(\varphi)$ which are not branches of $fg = 0$. Let also denote now by $\Delta(\varphi)$ the set of branches of the discriminant locus wich are the image of a branch of $\mathcal{C}(\varphi)$, so the map φ gives a surjective map $\varphi : \mathcal{C}(\varphi) \to \Delta(\varphi)$.

As a consequence of the results of [9], [10] and [5] one has that the set $\mathcal{C}(\varphi)$ can be decomposed as $\bigcup_{B\in\mathcal{B}^0}\mathcal{J}_B$, $\mathcal{J}_B\neq\emptyset$ for all $B\in\mathcal{B}^0$, and such that $\gamma\in\mathcal{J}_B$ if and only if its strict transform by π intersects B. Moreover, for $\gamma \in \mathcal{J}_B$ one has $I_z(g, \gamma)/I_z(f, \gamma) = Q(B)$. Note that $Q(B) = q/p$ (the rational number defined in 1.5).

Remark 3.3. The decomposition described in [5] relating the special values and the branches of the critical locus $\mathcal{C}(\varphi)$ is not the same as the one described here: in fact it could happen that a special component (i.e. a connected component of $E \setminus \mathcal{D}$ can be decomposed in several different r-nodal zones. On the other hand, the points of a divisorial component in $\mathcal{P}_{\mathcal{C}}$ offers a more precise decomposition than the one given by the corresponding 1-nodal zone described in [10]. In this sense the decomposition from the elements of β is a mixture of both decompositions.

Now, by Theorem 4 and Corollary 3 in [5], the special fiber $\xi_B = \{g^p - af^q = 0\}$ 0} of $\Phi = \langle g, f \rangle$ defined above just coincides with the fiber g_1 defined in 1.5. Thus the pencil $\Phi_1 = \langle f_1, g_1 \rangle$ is exactly the one corresponding to the map $\varphi_1 = (g_1, f_1)$. Moreover, the critical locus $\mathcal{C}(\varphi_1)$ is the same as $\mathcal{C}(\varphi)$, the one of the map φ (this is a straighforward computation, see also the proof of Theorem 4 in [4] or lemma 6.4 in [7]). That means that we can repeat the same process for the new map and so consider $B_1 \in \mathcal{B}^0(\varphi_1)$ such that the strict transform of γ by the corresponding resolution map intersects B_1 .

The above construction makes possible to define the map $F : \mathcal{C}(\varphi) \to \mathcal{S}$ such that for $\gamma \in \mathcal{C}(\varphi)$, $F(\gamma) \in \mathcal{S}$ is the unique sequence $(B_i)_{i \geq 0}$ such that the corresponding strict transform of γ intersects B_i for all i. Note that the fact that $\mathcal{J}_{B_i} \neq \emptyset$ for all i implies that the map F is surjective.

Theorem 3.4. Let $\gamma \in \mathcal{C}(\varphi)$, $\delta = \varphi(\gamma) \in \Delta(\varphi)$ and let $F(\gamma) = (B_i)_{i \geq 0}$. *Then* $(Q_i(B_i))_{i\geq 0}$ *together with the integer* k *such that* $\varphi_* \gamma = k \delta$ *determines the semigroup (and so the topological type) of the branch* δ*.*

Proof. One needs to compute the sequence $\{m_i : i \geq -1\}$ described in Theorem 1.8. For $i \geq 0$ let $r_i = q_i/p_i = Q_i(B_i)$, $gcd(q_i, p_i) = 1$. As $r_i = q_i/p_i = I_z(g_i, \gamma)/I_z(f_i, \gamma)$ and $f_i = f_{i-1}^{q_{i-1}} = f^{q_0...q_{i-1}}$ we deduce :

$$
I_z(g_i, \gamma) = r_i I_z(f_i, \gamma) = r_i q_{i-1} I_z(f_{i-1}, \gamma) = \left(\prod_{k=0}^{i-1} q_k\right) r_i I_z(f, \gamma) .
$$

Moreover, using notations of remark 1.10, $\mu_{-1} = I_z(f, \gamma)$ and for $i \geq 0$:

$$
\mu_i = I_z(g_i, \gamma) = I_z(\varphi^* y_i, \gamma) = I_0(y_i, \varphi_* \gamma) = k I_0(y_i, \delta) .
$$

Thus

$$
\mu_i/k = I_0(y_i, \delta) = I_z(g_i, \gamma)/k = (\prod_{k=0}^{i-1} q_k) r_i I_z(f, \gamma)/k = (\prod_{k=0}^{i-1} q_k) r_i \mu_{-1}/k.
$$

So for the elements $m_i, i \geq -1$ of the semigroup of δ we obtain:

$$
m_{i+1} = d_i m_i + \frac{\mu_{i+1}}{k} - p_i \frac{\mu_i}{k} = d_i m_i + (\prod_{k=0}^i q_k) r_{i+1} \frac{\mu_{-1}}{k} - p_i (\prod_{k=0}^{i-1} q_k) r_i \frac{\mu_{-1}}{k}
$$

= $d_i m_i + (r_{i+1} - 1) (\prod_{k=0}^i q_k) \frac{\mu_{-1}}{k}$.

(As in Remark 1.10 $\epsilon_i = \gcd(\mu_{-1}, \ldots, \mu_i)$ and $d_i = \epsilon_{i-1}/\epsilon_i$.)

Remark 3.5. Let S be the set of maximal completely ordered subtrees of T. Then $\mathcal{C}(\varphi) = \bigcup_{S \in \mathcal{S}} F^{-1}(S)$ is a partition of the set of branches $\mathcal{C}(\varphi)$. It is clear that, if $\gamma, \gamma \in F^{-1}(S)$ for some $S \in \mathcal{S}$, then $\varphi(\gamma)$ and $\varphi(\gamma')$ are equisingular: both have the same semigroup. Notice that $F^{-1}(S) \neq \emptyset$ for any $S \in \mathcal{S}$, however it could happen that $\#F^{-1}(S) > 1$ for some S.

 \Box

Moreover, if $F(\gamma) = F(\gamma')$ one has that the iterated sequences of pencils $\mathcal{P}(f,g,\gamma) = \{(\Phi_i,g_i,q_i/p_i) : i \geq 0\} \text{ and } \mathcal{P}(f,g,\gamma') = \{(\Phi'_i,g'_i,q'_i/p'_i) : i \geq 0\}$ are the same but not at the inverse, i.e. it could happen that $\mathcal{P}(f, g, \gamma) =$ $\mathcal{P}(f,g,\gamma')$ but $F(\gamma) \neq F(\gamma')$. A key point when $\gamma, \gamma' \in \mathcal{C}(\varphi)$ is that the iterated sequences of pencils can be determined without the previous knowledge of the branches γ and γ' provided they are from the critical locus.

For each $S = (B_i)_{i>0} \in S$ let $Q(S) = (Q_i(B_i))_{i>0}$. Theorem above implies that the set of sequences ${Q(S) : S \in S}$ theoretically permits to recover the set of topological types (the semigroups) of the branches of $\Delta(\varphi)$ (see the end of Remark 1.10). So in this sense both sets are equivalent.

3.6 (The intersection multiplicity). Let $\gamma, \gamma' \in C(\varphi)$ and let $\varphi_*(\gamma) = k\delta$, $\varphi_*(\gamma') = k'\delta'.$ Assume that $\delta \neq \delta'.$ The computation of the intersection multiplicity $[\delta, \delta']$ of δ and δ' can be made as in the Theorem 2.12 and so it depends of a more detailled knowledge of the branches of the fibers.

Theorem 3.7. *Let* ℓ *be the smallest integer such that there exists a branch* ρ *of the fibre* $\xi = \{g_\ell = 0\}$ *of* Φ_ℓ *such that*

$$
\frac{I_z(\gamma, \varphi^*(\varphi(\rho)))}{I_z(\gamma, f)} \neq \frac{I_z(\gamma', \varphi^*(\varphi(\rho)))}{I_z(\gamma', f)}.
$$

Then:

$$
[\delta, \delta'] = \min \left\{ \frac{e'_{\ell-1} I_z(\gamma, \varphi^*(\varphi(\rho)))}{k}, \frac{e_{\ell-1} I_z(\gamma', \varphi^*(\varphi(\rho)))}{k'} \right\}.
$$

Remark 3.8. Let $\gamma, \gamma' \in \mathcal{C}(\varphi)$ be such that $F(\gamma) = (B_i)_{i \geq 0} \neq F(\gamma') = (B_i')_{i \geq 0}$ and let ℓ be such that $B_i = B'_i$ for $i < \ell$ and $B_\ell \neq B'_\ell$. In this case there exists a branch ρ of $\xi = \{g_\ell = 0\}$ such that its strict transform intersects B_ℓ and does not B'_{ℓ} . For such a branch one has that

$$
\frac{I_z(\gamma, \varphi^*(\varphi(\rho)))}{I_z(\gamma, f)} > \frac{I_z(\gamma', \varphi^*(\varphi(\rho)))}{I_z(\gamma', f)}.
$$

So, in this case one has that

$$
[\delta,\delta'] = \frac{e_{\ell-1}I_z(\gamma',\varphi^*(\varphi(\rho)))}{k'}.
$$

References

- [1] Barranco Mendoza, G. & Snoussi, J. Equisingularity in pencils of curves on germs of reduced complex surfaces. Proc. Edinb. Math. Soc. (2), 67, 990-1012 (2024).
- [2] Delgado, F. & Maugendre, H. Special fibres and critical locus for a pencil of plane curve singularities. Compositio Math. 136, 69–87 (2003).
- [3] Delgado, F. An arithmetical factorization for the critical point set of some maps from \mathbb{C}^2 to \mathbb{C}^2 . In: Singularities–Lille 1991. Jean-Paul Brasselet (ed). London Mathematical Society Lecture Note Series, v.201. Cambridge University Press, 61–100 (1994).
- [4] Delgado, F. & Maugendre, H. On the topology of the image by a morphism of plane curve singularities. Rev. Mat. Complut. 27, 369–384 (2014).
- [5] Delgado, F. & Maugendre, H. Pencils and critical loci on normal surfaces. Rev. Mat. Complut. 34, 691-714 (2021).
- [6] Fulton, W. Intersection theory. Springer Verlag. New York, Heidelberg, Berlin. 1984.
- [7] García Barroso, E. & Popescu Pampu, P. A eudoxian study of discriminant curves associated to normal surface singularities. ArXiv [math-AG] 2410.21250 (2024).
- [8] Gryszka, B.; Gwozdziewicz, J. & Parusinski, A. Initial Newton polynomial of the discriminant. Bull. London Math. Soc., 54, 1584-1594 (2022).
- [9] Lˆe D˜ung Tr`ang, Maugendre, H. and Weber, C.. Geometry of critical loci. Journal of the L.M.S. 63 (2001), 533-552.
- [10] Michel, F. Jacobian curves for normal complex surfaces. Brasselet, J-P. (ed.) et al., Singularities II. Geometric and topological aspects. Proceedings of the international conference "School and workshop on the geometry and topology of singularities" in honor of the 60th birthday of Lê Dũng Tràng, Cuernavaca, Mexico, January 8-26, 2007. Providence, RI: American Mathematical Society (AMS). Contemporary Mathematics 475, 135-150 (2008).
- [11] Mumford, D. The topology of normal singularities of an algebraic surface and a criterion for simplicity. Publications Mathématiques de l'IHES, tome 9 (1961), 5-22.