A convexity-like structure for polar decomposition with an application to distributed computing

Foivos Alimisis Bart Vandereycken

University of Geneva, Switzerland

Abstract

We make a full landscape analysis of the (generally non-convex) orthogonal Procrustes problem. This problem is equivalent with computing the polar factor of a square matrix. We reveal a convexity-like structure, which explains the already established tractability of the problem and show that gradient descent in the orthogonal group computes the polar factor of a square matrix with linear convergence rate if the matrix is invertible and with an algebraic one if the matrix is singular. These results are similar to the ones of [5] for the symmetric eigenvalue problem. We present an instance of a distributed Procrustes problem, which is hard to deal by standard techniques from numerical linear algebra. Our theory though can provide a solution.

1 Introduction

The polar decomposition of a matrix is a standard factorization, where some matrix $C \in \mathbb{R}^{m \times n}$, $m \ge n$, must be written as the product of an orthonormal matrix $X \in \mathbb{R}^{n \times m}$ and a symmetric and positive semidefinite matrix $P \in \mathbb{R}^{n \times n}$, i.e.

$$C = XP.$$

Such a decomposition always exists and a good way to see that is through the singular value decomposition. If a singular value decomposition of C is

$$C = U\Sigma V^T$$
.

then the "polar factor" X of the polar decomposition is given as

 $X = UV^T$

and the symmetric positive semidefinite part P is given as

$$P = V\Sigma V^T$$

One can easily see that the polar decomposition of C is unique if and only if C is invertible, i.e. if and only if its singular values are all positive.

An interesting property of the polar factor is that it is the closest orthonormal matrix to the original matrix C (see [14], Theorem 8.4). This makes polar decomposition intimately related to the orthogonal Procrustes problem (see [14], Theorem 8.6). The Procrustes problem [27] is important in many areas of applied science [2, 11, 16]. It seeks for an orthogonal matrix $X \in \mathbb{O}(n)$, such that the quantity $||AX - B||_F^2$ for two matrices $A, B \in \mathbb{R}^{m \times n}$ is as small as possible. This problem admits the equivalent formulation

$$\min_{X \in \mathbb{O}(n)} - \operatorname{Tr}(CX),$$

with $C := B^T A$, and its solution is the polar factor of this matrix $C \in \mathbb{R}^{n \times n}$. This problem turns out to have a geodesic convexity-like structure in the orthogonal group, which we analyze in Section 4. This structure is similar to the one that has recently been proven for the symmetric eigenvalue problem [5]. Using this convexity-like structure, we analyze a Riemannian gradient descent algorithm in the orthogonal group for computing the polar factor of C. This algorithm is in general slow compared to the state-of-the-art and is presented mostly for theoretical purposes. It does provide though a principled way to deal with the case that C is singular (Theorem 11), without extra pivoted QR decompositions, as needed by the standard algorithms (see [14], bottom of page 196).

The main application of our theory is in a noisy version of the Procrustes problem. In general, polar factors behave quite badly with respect to perturbations of the original matrix. Let \tilde{C} be a perturbation of C, then the distance between the polar factors \tilde{X} and X can be upper bounded in general as (see Theorem 8.10 in [14])

$$\|X - \tilde{X}\|_F \le \frac{2}{\sigma_{\min}(C) + \sigma_{\min}(\tilde{C})} \|C - \tilde{C}\|_F.$$

This means that computing the polar factor of a perturbed version of C fast and in high accuracy does not mean much, especially in the case where C and its perturbed version are nearly singular. In other words, we cannot just take \tilde{C} and apply some of the classic algorithms (see Section 2) directly on it.

In Section 6, we present a structured noisy version of the polar factor computation problem. Namely, we count the bit complexity of a distributed orthogonal Procrustes problem. That is to say, the datasets we want to fit via an orthogonal transformation are distributed in computing nodes, which communicate messages in finite precision. As datasets become larger and larger and distribution vaster, it is in general agreed that the main computational bottleneck of such problem is the communication cost, and not the cost of local computations in each node. This is the reason that, in recent years, this so-called *bandwidth cost* has been analyzed for many different problems [15, 28]. We tackle this problem by developing a "quantized" version of gradient descent in the orthogonal group, which performs error analysis in the style of [21]. A similar result for the problem of computing the first leading eigenvector of a symmetric matrix can be found in [4]. It is revealed that one can trade-off bit complexity depending on $\sigma_{\min}(C)$ with the ability to approximate the polar factor of the exact matrix as accurately as one wishes (Theorem 15).

2 Related work

The most direct way to compute a polar decomposition is via the SVD. Clearly, this approach is too expensive. The numerical linear algebra community has developed plenty of faster algorithms to tackle the problem of computing a polar factor. The most basic one is the Newton method ([14], Section 8.3). The Newton method is in general fast in the late stage of convergence, but can be very slow in the beginning if the matrix C is ill-conditioned. Another prominent class of algorithms is the Padé family of iterations ([14], Section 8.5), which suffers more or less by the same issues.

Most of the effort in the last few years has been focused on scaling the basic Newton iteration, in order to obtain variants that do not suffer from slow convergence in the beginning of the iterations. The so-called "optimal" scaling [18] enjoys excellent theoretical behaviour, but the scaling factor depends on the (generally unknown) smallest and largest singular values in each iterate X_t . A more practical version, that however lacks convergence guarantees, can be found in [13]. A middle ground with a sub-optimal computable scaling that still enjoys some convergence guarantees can be found in [8].

The state-of-the art in this area comes probably from [22]. There, the Halley's method (which is a member of the Padè family of iterations) is scaled in a principled way. The Halley method has cubic asymptotic convergence, but the initial stage can be very slow for ill-conditioned matrices [12]. The scaling of [22] helps to improve its performance in the initial stage of convergence.

Other studies on meta-issues like the backward stability of the aforementioned algorithms have also been conducted [23], but this starts deviating from the purpose of our work.

Our work shows that the problem of computing the polar factor of a square matrix can be treated in the realm of convex optimization. That is because this problem enjoys a structure that we call weak-quasi-strong-convexity (WQSC) (Proposition 5). Versions of this property have appeared in [24] (Definition 1), [17](Appendix A) and [7]. WQSC has been derived also for the symmetric eigenvalue problem in [5]. It guarantees a linear convergence rate for gradient descent in terms of distances of the iterates to the optimum in the case that C is non-singular (Theorem 10). Actually, WQSC has been proven to be also a necessary property for having this kind of convergence [3]. This structure gives a principled way to work also in the case that the matrix C is singular (Theorem 11). Note that the classic algorithms we discuss above can indeed sometimes be applied to non-singular matrices, but, even then, we are not aware of any convergence results in this case. The way classic numerical linear algebra

has dealt with this issue is by successive pivoted QR decompositions until the problem is deduced to a non-singular one (see page 196 of [14]).

A problem that classic algorithms may completely fail to deal with, is the theoretical analysis of an orthogonal Procrustes setting where the two datasets to be fit are distributed among different computing nodes, each one having its own memory. In the most practical of such scenarios, one is interested in the total communication cost for solving the problem, counted in bits. While distributed SVD has received some attention [4, 9, 19, 26], we are not aware of any works that consider directly the computation of a polar factor in a distributed setting. A relevant paper is [20], but it deals with the distributed memory problem from an implementation point of view in a supercomputer. The reason we think that the classic algorithms for polar decomposition may fail to deal with such scenario is that, in general, they behave badly to perturbations of the matrix C (see introduction). In the aforementioned setting, C is perturbed since it is split among different computing nodes, which communicate messages in low precision. While gradient descent is amenable to a basic error analysis (see Section 6) Newton or Padé-type methods do not seem to behave the same way (at least, we are not aware of any principled way to do error analysis for these algorithms).

3 Geometry of the orthogonal group

 $\mathbb{O}(n)$ is a Lie group (thus also a manifold) consisted by orthogonal $n \times n$ matrices. It is consisted by two connected components, the orthogonal matrices with determinant +1 and the ones with determinant -1. We recall here basic aspects of the geometry of the orthogonal group. A more complete exposition can be found along the excellent text [25].

The tangent space at a point X is

$$T_X \mathbb{O}(n) = \{ X \Omega \mid \Omega \in \mathbb{R}^{n \times n} \text{ is skew-symmetric} \}.$$

The usual Riemannian metric one equips this space is just the Euclidean one:

$$\langle V, W \rangle_X = \operatorname{Tr}(W^T V).$$

The orthogonal projection of a matrix $Z \in \mathbb{R}^{n \times n}$ onto $T_X \mathbb{O}(n)$ is

$$P_X(Z) = X \operatorname{skew}(X^T Z). \tag{1}$$

The exponential map at a point X in the direction $X\Omega$ is defined as

$$\exp_X(X\Omega) = X \exp_m(\Omega),$$

where \exp_m is the matrix exponential.

The Riemannian logarithm is the inverse of the exponential map, when the

latter is invertible. We now examine when this is the case. The domain where the exponential map is a diffeomorphism is usually called injectivity domain. In order to identify it, we need to verify when the equation

$$\exp_X(X\Omega) = X \exp_m(\Omega) = Y$$

has a unique solution. This happens if and only if the equation

$$\exp_m(\Omega) = X^T Y$$

has a unique solution. Considering the eigenvalue decomposition $\Omega = U\Lambda U^{-1}$, where Λ is diagonal with entries of the form ir with $r \in (-\pi, \pi]$ (since Ω is skew-symmetric). This implies that the eigenvalue decomposition of $\exp_m(\Omega)$ is $U \exp_m(\Lambda) U^{-1}$ and $\exp_m(\Lambda)$ is diagonal featuring entries of the form e^{ir} . Thus, the previous equation is equivalent to the diagonal system $\exp_m(\Lambda) = U^{-1} X^T Y U$. Reading the diagonal entries, we obtain a series of equations of the form

$$e^{ir} = s$$

where s are the eigenvalues of $X^T Y$. These equations are well-defined and have a unique solution if and only if s is in the domain of a definition of the complex logarithm, $\mathbb{C} \setminus (-\infty, 0]$. In that case, r is allowed to be in $(-\pi, \pi)$. We can summarize the previous discussion as follows:

Lemma 1. • The domain of the orthogonal group where the exponential map is a diffeomorphishm is

$$\{X\Omega \mid \Omega^T = -\Omega, \|\Omega\|_2 < \pi\}.$$
 (2)

- Let $X, Y \in \mathbb{O}(n)$. If the phases r of the eigenvalues e^{ir} of $X^T Y$ satisfy $r \in (-\pi, \pi)$, then there is a unique geodesic connecting X and Y. In this case, it trivially holds that X and Y are in the same connected component of $\mathbb{O}(n)$.
- If some of the r's are equal to π , then it holds: if there is even number of r's equal to π , then X and Y are in the same connected component (and are connected by multiple geodesics). If there is odd number of r's equal to π , then X and Y are in different connected components (i.e. det(XY) = -1).

Let us now consider X and Y such that $X^T Y$ has eigenvalues with phases in $(-\pi, \pi)$. Then $\log_X(Y)$ is well-defined and

$$\exp_X(\log_X(Y)) = Y.$$

We can write $\log_X(Y) = X\Omega$ for some skew-symmetric Ω and we have

$$X \exp_m(\Omega) = Y,\tag{3}$$

which can be written as

$$\Omega = \log_m(X^T Y),$$

where \log_m is the matrix logarithm.

Thus,

$$\log_X(Y) = X \log_m(X^T Y). \tag{4}$$

Note that $\log_m(X^T Y)$ is indeed a skew-symmetric matrix since X and Y are orthogonal.

 $\log_X(Y)$ is a tangent direction at X that allows to move from X to Y along the geodesic connecting them. In Riemannian manifolds, there is a general rule to transport tangent vectors from one point to another one parallely to the geodesics of the manifold, called parallel transport. In the orthogonal group, the parallel transport from a point X to a point Y (denoted by Γ_X^Y), is given by

$$\Gamma_X^Y(X\Omega) = Y(X^T Y\Omega Y^T X).$$

Notice that $X^T Y \Omega Y^T X$ is a skew-symmetric matrix, since it is a conjugation of the slew-symmetric matrix Ω . This definition makes sense of course only if X and Y are in the same connected component of \mathbb{O}^n .

Since we have computed the Riemannian logarithm between two orthogonal matrices X and Y, we can also compute the Riemannian distance between such matrices based on it:

$$\operatorname{dist}^{2}(X,Y) = \|\log_{X}(Y)\|^{2} = \|X\log_{m}(X^{T}Y)\|^{2} = \|\log_{m}(X^{T}Y)\|^{2}.$$

In order to proceed, we decompose the orthogonal matrix $X^T Y$ into the socalled canonical form PDP^T , where P is orthogonal featuring the eigenvectors of $X^T Y$ in its columns and D is block diagonal. D is constructed as follows. When $X^T Y$ has an eigenvalue equal to 1, D has a diagonal entry equal to 1. When $X^T Y$ has an eigenvalue of the form e^{ir} for some $r \in (-\pi, 0) \cup (\pi, 0)$, then e^{-ir} is also an eigenvalue and D features the 2×2 block that is the 2-d rotation with angles r. That is $\begin{bmatrix} \cos r & -\sin r \\ \sin r & \cos r \end{bmatrix}$.

The matrix logarithm has the following convenient property. Given the above decomposition, we have

$$\log_m(PDP^T) = P\log_m(D)P^T.$$

Taking D as constructed previously, $\log_m(D)$ has 0 in the positions where D has 1 and $\begin{bmatrix} 0 & -r \\ r & 0 \end{bmatrix}$ where D has $\begin{bmatrix} \cos r & -\sin r \\ \sin r & \cos r \end{bmatrix}$. Since P is orthogonal, the distance between X and Y turns out to be equal to $\|\log_m(D)\|^2 = \operatorname{Tr}(\log_m(D)^T \log_m(D))$. $\log_m(D)^T \log_m(D)$ is again a 2×2 block diagonal matrix with 0's where $\log_m(D)$ has 0's and $\begin{bmatrix} r^2 & 0 \\ 0 & r^2 \end{bmatrix}$ where $\log_m(D)$ has $\begin{bmatrix} 0 & -r \\ r & 0 \end{bmatrix}$. Thus, the distance between X and Y is

$$dist(X,Y) = \left(\sum_{i=1}^{n} r_i^2\right)^{1/2},$$
 (5)

where e^{ir_i} are the eigenvalues of $X^T Y$. That is to say that

$$\operatorname{dist}(X,Y) = \|\phi\|_2$$

where $\phi = (r_1, \ldots, r_n)$. If $r_j = 0$, then it appears only once in ϕ , otherwise it appears as a couple with $-r_j$. Note that with a simple limit argument, we can conclude that the same formula still holds when some eigenvalues of $X^T Y$ have a phase equal to π .

We end this section on the geometry basics of the orthogonal group by discussing its sectional curvatures. The only fact about its curvatures that will be useful to us is that they are always nonnegative. This is folklore as the sectional curvatures of all Stiefel manifolds are nonegative, thus also the ones of the orthogonal group (which is a special case of a Stiefel manifold). This gives rise to the following useful geometric bound:

Proposition 2. Consider three points $X, Y, Z \in \mathbb{O}(n)$, such that they are connected by unique geodesics. Then, we have

- 1. dist²(X, Y) \leq dist²(Z, X) + dist²(Z, Y) 2(Log_Z(X), Log_Z(Y)).
- 2. dist $(X, Y) \le \|\log_Z(X) \log_Z(Y)\|.$

Proof. Both inequalities are a consequence of the famous Toponogov's theorem, taking into account that the sectional curvatures of $\mathbb{O}(n)$ are nonnegative. \Box

4 Convexity-like properties of orthogonal Procrustes

We investigate now thoroughly the orthogonal Procrustes problem. This problem concerns with finding orthogonal matrices X_1 and X_2 that best fit two other matrices $A, B \in \mathbb{R}^{k \times n}$:

$$\min_{X_1, X_2 \in \mathbb{O}(n)} \|AX_1 - BX_2\|^2.$$

Since this problem is invariant under simultaneous right multiplication of X_1 and X_2 with an orthogonal matrix $Q \in \mathbb{R}^{n \times n}$, we can fix X_2 to be identity and target only the matrix $X_1 \rightsquigarrow X$:

$$\min_{X\in\mathbb{O}(n)}\|AX-B\|^2.$$

This problem can be written equivalently as

$$\min_{X \in \mathbb{O}(n)} -\operatorname{Tr}(CX) =: f(X), \tag{6}$$

where

$$C := B^T A$$

It has a global solution and can be found in closed form [27]: if $C = U\Sigma V^T$ is an SVD of C, then a global solution is $X_* = VU^T$. The minimum $f_* := f(X_*)$ is the opposite of the sum of the singular values of C. We will use this structure to prove a quasi-convexity property for the function

$$f(X) = -\operatorname{Tr}(CX)$$

around X_* .

It is well known that the solution of the problem is **unique** if and only if all the singular values of C are strictly positive, i.e. if and only if C is invertible.

Riemannian gradient: To compute the Riemannian gradient of f, we need just to project the Euclidean gradient $\nabla f(X) = -C^T$ onto the tangent space $T_X \mathbb{O}(n)$. This results to

$$\operatorname{grad} f(X) = P_X(-C^T) = -X\operatorname{skew}(X^T C^T).$$
(7)

Riemannian Hessian: For a function f defined in the orthogonal group, we have (see [6])

$$D \operatorname{grad} f(X)[\dot{X}] = \dot{X} \operatorname{skew}(X^T \nabla f(X)) + X \operatorname{skew}(\dot{X}^T \nabla f(X) + \dot{X}^T \nabla^2 f(X)[\dot{X}]),$$

where $\dot{X} = X\Omega$ is an arbitrary tangent vector. In our case, $\nabla f(X) = -C^T$ and $\nabla^2 f(X) = 0$, thus

$$\operatorname{Hess} f(X)[\dot{X}] = -\dot{X}\operatorname{skew}(X^T C^T) - X\operatorname{skew}(\dot{X}^T C^T).$$
(8)

Proposition 3 (Geodesic weak-quasi-convexity). Let $X_* \in \mathbb{O}(n)$ a global optimum of the function $f : \mathbb{O}(n) \to \mathbb{R}$. Let also $X \in \mathbb{O}(n)$ such that the eigenvalues e^{ir} of $X^T X_*$ are such that $r \in (-\pi, \pi)$. If $|r|_{\text{max}}$ denotes the largest possible rotation induced by $X^T X_*$ in absolute value, then

$$\langle \operatorname{grad} f(X), -\log_X(X_*) \rangle \ge \frac{1}{2} (1 + \cos(|r|_{\max})) (f(X) - f_*).$$

Proof. The Riemannian gradient of f is given in equation 7. It remains to compute a convenient expression for the Riemannian logarithm.

According to equation (4), the Riemannian logarithm is given as

$$\log_X(X_*) = X \log_m(X^T X_*).$$

As in the introduction, we use the canonical form of the orthogonal matrix $X^T X^\ast$

$$X^T X_* = P D P^T.$$

Since the matrix logarithm is invariant under conjugate action, we have

$$\log_m(X^T X_*) = P \log_m(D) P^T$$

and $\log_m(D)$ is again a block diagonal matrix, with blocks being the logarithms of the blocks of D: when D has a diagonal entry equal to 1, $\log_m(D)$ has a diagonal entry equal to 0 and when D features a 2×2 block, which is a rotation of angle r, $\log_m(D)$ features the block $\begin{bmatrix} 0 & -r \\ r & 0 \end{bmatrix}$

Similarly, the skew-symmetric part of $X^T \vec{X}_*$ satisfies

$$\operatorname{skew}(X^T X_*) = P\operatorname{skew}(D)P^T$$

where skew(D) is again block diagonal and has a 0 diagonal entry when D has a 1 diagonal entry, while it has a block $\begin{bmatrix} 0 & -\sin r \\ \sin r & 0 \end{bmatrix}$ when D features a 2×2 rotation of angle r. Thus, it holds in general that

$$\log_m(D) = \operatorname{skew}(D) \frac{\phi}{\sin\phi},$$

where $\phi = (r_1, \ldots, r_n)$ is a vector capturing all the rotations induced by the orthogonal matrix $X^T X_*$. If r = 0, i.e. corresponds to a diagonal entry equal to 1, then it appears only once in ϕ , while if $r \in (-\pi, \pi) \setminus \{0\}$ it appears as a couple with -r.

 $\phi/\sin\phi$ is a diagonal matrix with diagonal elements $r_j/\sin r_j$. This convention is made for ease of notation.

Given that, we can write

$$\log_X(X_*) = X \log_m(X^T X_*) = X P \log_m(D) P^T = X P \operatorname{skew}(D) \frac{\phi}{\sin \phi} P^T$$
$$= X P \operatorname{skew}(D) P^T P \frac{\phi}{\sin \phi} P^T = P_X(X_*) P \frac{\phi}{\sin \phi} P^T.$$

Now we can finally deal with the desired inequality:

$$\langle \operatorname{grad} f(X), -\log_X(X_*) \rangle = \left\langle P_X(C^T), P_X(X_*) P \frac{\phi}{\sin \phi} P^T \right\rangle = \left\langle X \operatorname{skew}(X^T C^T), X_* P \frac{\phi}{\sin \phi} P^T \right\rangle = \operatorname{Tr} \left(P \frac{\phi}{\sin \phi} P^T X_*^T X \operatorname{skew}(X^T C^T) \right).$$

We pause to deal with the term $X_*^T X$ skew $(X^T C^T)$:

$$X_*^T X \text{skew}(X^T C^T) = X_*^T X \frac{X^T C^T - CX}{2} = \frac{X_*^T C^T - X_*^T X CX}{2}.$$

Remember that if $X_* = VU^T$, then $C = U\Sigma V^T$ is an SVD of C. Thus

$$X_*^T C^T = U \Sigma U^T$$

and

$$X_*^T X C X = P D^T P^T U \Sigma V^T V U^T P D^T P^T = P D^T P^T U \Sigma U^T P D^T P^T.$$

Plugging this expression in, we get

$$\begin{aligned} 2\langle \operatorname{grad} f(X), -\log_X(X_*) \rangle &= \operatorname{Tr} \left(P \frac{\phi}{\sin \phi} P^T X_*^T X \operatorname{skew}(X^T C^T) \right) \\ &= \operatorname{Tr} \left(P \frac{\phi}{\sin \phi} P^T (U \Sigma U^T - P D^T P^T U \Sigma U^T P D^T P^T) \right) \\ &= \operatorname{Tr} \left(\frac{\phi}{\sin \phi} (P^T U \Sigma U^T P - D^T P^T U \Sigma U^T P D^T) \right) \\ &= \operatorname{Tr} \left(\frac{\phi}{\sin \phi} (P^T U \Sigma U^T P) \right) - \operatorname{Tr} \left(\frac{\phi}{\sin \phi} (D^T P^T U \Sigma U^T P D^T) \right) \\ &= \operatorname{Tr} \left(\left(\frac{\phi}{\sin \phi} - D^T \frac{\phi}{\sin \phi} D^T \right) P^T U \Sigma U^T P \right). \end{aligned}$$

It suffices to show that

$$\operatorname{Tr}\left(\left(\frac{\phi}{\sin\phi} - D^T \frac{\phi}{\sin\phi} D^T\right) P^T U \Sigma U^T P\right) \ge (1 + \cos(|r|_{\max}))(f(X) - f_*) = \underbrace{(1 + \cos(|r|_{\max}))}_{:=c} \left(\underbrace{\operatorname{Tr}(P^T U \Sigma U^T P)}_{-f_*} - \underbrace{\operatorname{Tr}(D^T P^T U \Sigma U^T P)}_{-f(X)}\right).$$

This holds if

$$\operatorname{Tr}\left(\left(\frac{\phi}{\sin\phi} - D^T \frac{\phi}{\sin\phi} D^T + c(D^T - I)\right) \underbrace{P^T U \Sigma U^T P}_{:=A}\right) \ge 0.$$

Notice that the matrix A is symmetric and positive semi-definite.

 $D^{T} \text{ is a matrix with diagonal entries equal to 1 and <math>2 \times 2$ diagonal blocks of the form $\begin{bmatrix} \cos r & \sin r \\ -\sin r & \cos r \end{bmatrix}$, which essentially correspond to rotations with -r. Multiplying with the diagonal matrix $\phi/\sin\phi$ from the right, keeps the 1 diagonal entries of D^{T} unchanged, while it transforms the 2×2 diagonal blocks to $\begin{bmatrix} r/\tan r & r \\ -r & r/\tan r \end{bmatrix}$. The matrix $D^{T}\frac{\phi}{\sin\phi}D^{T}$ still keeps 1 in the entries that correspond to r = 0 and has 2×2 diagonal blocks associated with $r \in (-\pi, \pi) \setminus \{0\}$ that are $\begin{bmatrix} \frac{r}{\tan r}\cos r - r\sin r & \frac{r}{\tan r}\sin r & \frac{r}{\tan r}\cos r - r\sin r \end{bmatrix}$.

The matrix $\frac{\phi}{\sin \phi} - D^T \frac{\phi}{\sin \phi} D^T + c(D^T - I)$ has 1 in the diagonal entries that D^T has 1 (r = 0) and has 2×2 diagonal blocks that correspond to rotations with $r \in (-\pi, \pi) \setminus \{0\}$, which are

$$\begin{bmatrix} \frac{r}{\sin r} - \frac{r}{\tan r}\cos r + r\sin r + c(\cos r - 1) & -\frac{r}{\tan r}\sin r - r\cos r + c\sin r\\ \frac{r}{\tan r}\sin r + r\cos r - c\sin r & \frac{r}{\sin r} - \frac{r}{\tan r}\cos r + r\sin r + c(\cos r - 1) \end{bmatrix}$$

Notice that this last 2×2 matrix is of the form $\begin{bmatrix} \alpha & \beta \\ -\beta & \alpha \end{bmatrix}$. The expression $\operatorname{Tr}\left(\left(\frac{\phi}{\sin\phi} - D^T \frac{\phi}{\sin\phi} D^T + c(D^T - I)\right)A\right)$ that we want to prove nonnegative is the sum of the traces of the product of the diagonal entries of $\frac{\phi}{\sin\phi} - D^T \frac{\phi}{\sin\phi} D^T + c(D^T - I)$ that correspond to r = 0 (i.e. 1) with the corresponding diagonal entries of A and the 2 \times 2 diagonal blocks of $\frac{\phi}{\sin \phi}$ – $D^T \frac{\phi}{\sin \phi} D^T + c(D^T - I)$ with the corresponding 2×2 diagonal blocks of A. In the first case we get back the diagonal entries of A (which are nonnegative) and in the second case we have the product of a matrix of the form $\begin{bmatrix} \alpha & \beta \\ -\beta & \alpha \end{bmatrix}$ with one of the form $\begin{bmatrix} s & t \\ t & k \end{bmatrix}$, since A is symmetric. The diagonal entries of this product (which are the ones that contribute in the trace) are $\alpha s + \beta t$ and $-\beta t + \alpha k$. Their sum is $\alpha(s+t)$, thus it suffices to show that this expression is nonnegative, i.e. that α is nonnegative since s and t are nonnegative as diagonal entries of the positive semi-definite matrix A.

Remember that α has been taken as

$$\alpha := \frac{r}{\sin r} - \frac{r}{\tan r} \cos r + r \sin r + \underbrace{(1 + \cos(|r|_{\max}))}_{c} (\cos r - 1)$$
$$\geq \frac{r}{\sin r} - \frac{r}{\tan r} \cos r + r \sin r + (1 + \cos r)(\cos r - 1),$$

since $r \leq |r|_{\text{max}}$ and $\cos r - 1 \leq 0$. The last lower bound for α turns out to be positive for all $r \in (-\pi, \pi)$, thus our proof is complete.

We now examine a property for f known as quadratic growth. This property gives a non-trivial inequality only in the case that the Procrustes problem has a unique solution (i.e. if and only if C is non-singular).

Proposition 4 (Quadratic growth). Let $X_* \in \mathbb{O}(n)$ to be a global minimizer for f and $X \in \mathbb{O}(n)$ in the same connected component. Then f satisfies

$$f(X) - f_* \ge \frac{2\sigma_{min}(C)}{\pi^2} \operatorname{dist}^2(X, X_*),$$

where $\sigma_{\min}(C)$ is the smallest singular value of C.

Proof. Recall that if $C = U\Sigma V^T$ is an SVD of C, then $X_* = VU^T$ is a global minimizer. Consider again the canonical form of the orthogonal matrix $X^T X_*$:

$$X^T X_* = P D P^T.$$

Then, we have

$$f(X) - f_* = -\operatorname{Tr}(CX) + \operatorname{Tr}(CX_*) = \operatorname{Tr}(P^T U \Sigma U^T P) - \operatorname{Tr}(U \Sigma U^T P D^T P^T)$$
$$= \operatorname{Tr}((I - D^T) \underbrace{P^T U \Sigma U^T P}_{\text{pos. semi-definite}})$$

Let us denote again $A := P^T U \Sigma U^T P$, which is symmetric and positive semidefinite. The matrix $I - D^T$ has diagonal entries equal to 0 for rotations r = 0, diagonal entries equal to 2 for $r = \pi$ and 2×2 diagonal blocks of the form $\begin{bmatrix} 1 - \cos r & \sin r \\ -\sin r & 1 - \cos r \end{bmatrix}$ for rotations with angle $r \in (-\pi, \pi)$ not 0. Thus, the diagonal entries of the product $(I - D^T)P^T U \Sigma U^T P$ are either 0 for entries that correspond to no rotation, i.e. $(1 - \cos r)A_{ii}$, or the diagonal entries of a product of the form

$$\begin{bmatrix} 1 - \cos r & \sin r \\ -\sin r & 1 - \cos r \end{bmatrix} \begin{bmatrix} s & t \\ t & k \end{bmatrix}.$$

These are $(1 - \cos r)s + \sin rt$ and $-\sin rt + (1 - \cos r)k$. Since summing them makes the terms $\sin rt$ to cancel out, we get

$$\operatorname{Tr}((I - D^T)P^T U\Sigma U^T P) = \operatorname{Tr}((I - \cos\phi)P^T U\Sigma U^T P),$$

where $\phi = (r_1, \ldots, r_n)$ a vector capturing all the rotations between X and X_* . If $r_j = 0$ or π , then it appears only once, if $r_j \neq 0, \pi$ it appears coupled with its opposite -r. Notice that

$$\|\phi\| = \operatorname{dist}(X, X_*).$$

Since for all r it holds $r \in (-\pi, \pi]$, we have

$$1 - \cos r \ge \frac{2}{\pi^2} r^2.$$

By basic properties of the trace, we have

$$\operatorname{Tr}((I - \cos \phi)P^T U \Sigma U^T P) \ge \lambda_{\min}(P^T U \Sigma U^T P) \operatorname{Tr}(I - \cos \phi) \ge \frac{2\sigma_{\min}(C)}{\pi^2} \|\phi\|^2.$$

The last inequality completes the proof.

We can combine Propositions 3 and 4 to a more compact form, which we call weak-quasi-strong-convexity (WQSC) [3]. Similar versions of this property have also appeared in [24] (Definition 1) and [17] (Appendix A). WQSC looks a lot like strong convexity but it is much more general. Interestingly, the role of a strong convexity constant μ is played by a multiple of $\sigma_{\min}(C)$. That is to say, the further away from being singular C is, the stronger this property becomes. If Cis singular, the derived inequality reduces to weak-quasi-convexity (Proposition

3, but with slightly weaker parameters).

 \Box

Proposition 5 (Weak-quasi-strong-convexity). For any X satisfying the properties of Propositions 3, 4, f satisfies the following inequality:

$$f(X) - f^* \le \frac{1}{a(X)} \langle \operatorname{grad} f(X), -\log_X(X_*) \rangle - \frac{\mu}{2} \operatorname{dist}^2(X, X_*)$$

with $a(X) := \frac{1 + \cos(|r|_{max})}{4}$ and $\mu := \frac{4\sigma_{min}(C)}{\pi^2}$. $|r|_{max} < \pi$ is the largest rotation in absolute value induced by the orthogonal matrix $X^T X_*$.

Proof. For the specific choices of a(X) and μ , we have

$$\frac{\mu}{2}\operatorname{dist}^2(X, X_*) \le f(X) - f_* \le \frac{1}{2a(X)} \langle \operatorname{grad} f(X), -\log_X(X_*) \rangle.$$

The left inequality is derived by Proposition 4 and the right one by Proposition 3.

Now, again by Proposition 3, we have

$$f(X) - f_* \leq \frac{1}{2a(X)} \langle \operatorname{grad} f(X), -\log_X(X_*) \rangle + \frac{\mu}{2} \operatorname{dist}^2(X, X_*) - \frac{\mu}{2} \operatorname{dist}^2(X, X_*)$$
$$\leq \frac{1}{a(X)} \langle \operatorname{grad} f(X), -\log_X(X_*) \rangle - \frac{\mu}{2} \operatorname{dist}^2(X, X_*)$$

by substituting the previous inequality.

We now analyze a property for f usually referred as smoothness.

Definition 6. A function $f : \mathbb{O}(n) \to \mathbb{R}$ is called L-smooth, if for all X, Y in the same connected component of $\mathbb{O}(n)$, we have

$$\|\operatorname{grad} f(X) - \Gamma_Y^X \operatorname{grad} f(Y)\| \le L\operatorname{dist}(X, Y),\tag{9}$$

where Γ_Y^X is the parallel transport between X and Y.

Since f is twice differentiable, the previous property is equivalent with the eigenvalues of its Riemannian Hessian being upper bounded in absolute value uniformly by L. Using a standard Taylor expansion (see [1], Theorem 7.1.2), we easily see that this implies

$$f(Y) - f(X) \le \langle \operatorname{grad} f(X), \log_X(Y) \rangle + \frac{L}{2} \operatorname{dist}^2(X, Y), \tag{10}$$

for all $X, Y \in \mathbb{O}(n)$ that are connected by a unique geodesic.

A weaker, but useful, inequality comes by setting $Y = \exp_X \left(-\frac{1}{L}\operatorname{grad} f(X)\right)$ (see Proposition 9 for a proof that such Y is uniquely defined). Then, we get

$$f(Y) \le f(X) - \frac{1}{2L} \| \operatorname{grad} f(X) \|^2$$

and since $f^* \leq f(Y)$,

$$f(X) - f^* \ge \frac{1}{2L} \|\operatorname{grad} f(X)\|^2, \text{ for all } X \in \mathbb{O}(n).$$
(11)

Proposition 7 (Smoothness). f is geodesically $\sigma_{\max}(C)$ -smooth.

Proof. It suffices to show that the eigenvalues of the Riemannian Hessian Hess f(X) are in absolute value upper bounded by $\sigma_{\max}(C)$ for all $X \in \mathbb{O}(n)$. For our computations, we follow the exposition in [6]:

$$\langle \dot{X}, \operatorname{Hess} f(X)[\dot{X}] \rangle = \operatorname{Tr}(\dot{X}^T \dot{X} \operatorname{skew}(-X^T C^T)) + \operatorname{Tr}(\dot{X}^T X \operatorname{skew}(-\dot{X}^T C^T)).$$

The first term is 0 as the trace of the product of a symmetric and skewsymmetric matrix. The second term becomes

$$\operatorname{Tr}(\dot{X}^T X \operatorname{skew}(-\dot{X}^T C^T)) = \frac{1}{2} \operatorname{Tr}(\dot{X}^T X C \dot{X} - \dot{X}^T X \dot{X}^T C^T).$$

Substituting $\dot{X}^T X = \Omega^T$, we get

$$\frac{1}{2}\operatorname{Tr}(\dot{X}^T X C X - \dot{X}^T X \dot{X}^T C^T) = \frac{1}{2}\operatorname{Tr}(\Omega^T C \dot{X} - \Omega^T \dot{X}^T C^T) = \frac{1}{2}\operatorname{Tr}(\Omega^T C \dot{X} + \Omega \dot{X}^T C^T) = \operatorname{Tr}(\Omega^T C \dot{X}) = \operatorname{Tr}(\Omega^T C X \Omega) = \operatorname{Tr}(C X \Omega \Omega^T).$$

The last expression features the trace of the product of the matrix CX with the symmetric and positive semi-definite matrix $\Omega\Omega^T$. By basic facts in linear algebra (Von Neumann's trace inequality), we can upper bound the absolute value of this expression by $\sigma_{\max}(CX) \operatorname{Tr}(\Omega\Omega^T)$. Since X is orthogonal, we have that $\sigma_{\max}(CX) = \sigma_{\max}(C)$. Also $\operatorname{Tr}(\Omega\Omega^T) = \operatorname{Tr}(\dot{X}\dot{X}^T) = ||\dot{X}||^2$. Putting it all together, we get

$$|\langle X, \operatorname{Hess} f(X)[X] \rangle| \le \sigma_{\max}(C) ||X||^2$$

and the desired result follows.

As it is customary, we denote

$$L := \sigma_{\max}(C).$$

Remark. Since we have computed the quantity $\langle \dot{X}, \text{Hess}f(X)[\dot{X}] \rangle$, it is a good point to verify that f is indeed non-convex, even inside the same connected component. To see that, take $C = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$. It is easy to see that in this case, the minimizer of f is the identity matrix $X_* = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$. Consider a matrix, which is almost equal to $\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$, but not exactly, take for example $X := \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$, with any $\theta \in (\pi/2, \pi)$. The eigenvalues of $X^T X_*$ are $e^{i\theta}$ and $e^{-i\theta}$ and, since $|\theta| < \pi$, X and X^* are connected by a unique geodesic. The Riemannian Hessian at X satisfies

$$\langle \dot{X}, \operatorname{Hess} f(X)[\dot{X}] \rangle = \operatorname{Tr}(CX\Omega\Omega^T),$$

where $\dot{X} = X\Omega$. We have that $CX = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & 2\cos\theta \end{bmatrix}$ and set the symmetric and positive semidefinite matrix $\Omega\Omega^T$ to be $\begin{bmatrix} \alpha & \beta \\ \beta & \gamma \end{bmatrix}$, where α and γ are nonnegative and at least one is strictly positive.

The diagonal entries of the matrix $CX\Omega\Omega^T$ are $\alpha\cos\theta - \beta\sin\theta$ and $\beta\sin\theta + 2\gamma\cos\theta$. Thus,

$$\operatorname{Tr}(CX\Omega\Omega^T) = \underbrace{(\alpha + 2\gamma)}_{>0} \underbrace{\cos\theta}_{<0}$$

and we have found a point X where the Riemannian Hessian has negative eigenvalues (it is actually negative definite).

We conclude this section with a small technical lemma that allows us to show that gradient descent with a properly chosen step size is well-defined in the sense that the direction used for update belong in the injectivity domain (2).

Lemma 8. The Riemannian gradient of f evaluated at X is of the form $X\Omega$, for a skew-symmetric matrix Ω with

$$\|\Omega\|_2 \le \sigma_{\max}(C).$$

Proof. The Riemannian gradient of f at X is

$$\operatorname{grad} f(X) = X \operatorname{skew}(X^T C^T),$$

thus Ω is taken as skew $(X^T C^T)$. By the sub-additivity of the spectral norm and its invariance under multiplication with orthogonal matrices, we have

$$\|\Omega\|_{2} = \|\operatorname{skew}(X^{T}C^{T})\|_{2} = \leq \frac{\|X^{T}C^{T}\|_{2} + \|CX\|_{2}}{2} = \frac{\|C^{T}\|_{2} + \|C\|_{2}}{2}.$$

This gives the desired result.

5 Convergence of Riemannian gradient descent

Riemannian gradient descent applied to a function $f : \mathbb{O}(n) \to \mathbb{R}$ reads as

$$X_{t+1} = \exp_{X_t}(-\eta_t \operatorname{grad} f(X_t)), \tag{12}$$

with $\eta_t > 0$ being the step size.

The results of Section 4 guarantee a local (non-asymptotic) linear convergence rate for Riemannian gradient descent on f in the case that C is invertible, if ran with a properly chosen step size and the initial guess X_0 is sufficiently close to the optimum. **Proposition 9.** Let X_t and X_* be such that the largest rotation $|r|_{\max}$ induced by the orthogonal matrix $X_t^T X_*$ satisfies $|r|_{\max} < \pi$. Then, iteration (12) with $0 \le \eta_t \le a(X_t)/L$ satisfies

$$\operatorname{dist}^{2}(X_{t+1}, X_{*}) \leq \left(1 - \frac{4}{\pi^{2}} \sigma_{\min}(C) a(X_{t}) \eta_{t}\right) \operatorname{dist}^{2}(X_{t}, X_{*}),$$

with $a(X_t)$ defined as in Proposition 5.

Proof. We start by showing that iteration 12 is well-defined. By the assumption $|r|_{\max} < \pi$, we get that $0 < a(X_t) = \frac{1+\cos(|r|_{\max}))}{4} \leq \frac{1}{2}$. By Lemma 8, the tangent vector $\eta_t \operatorname{grad} f(X_t)$ that is used to update iteration 12 can be written as $X\Omega$, with $\|\Omega\|_2 \leq \eta_t \sigma_{\max}(C)$. By the definition of η_t , we have that

$$\|\Omega\|_2 \le \frac{a(X_t)}{L}\sigma_{\max}(C) = \frac{a(X_t)}{\sigma_{\max}(C)}\sigma_{\max}(C) \le \frac{1}{2}.$$

Thus, $\eta_t \operatorname{grad} f(X_t)$ is inside the injectivity domain (2) and, as a consequence, iteration (12) is well-defined.

We can now apply Proposition 2 to obtain

$$dist^{2}(X_{t+1}, X_{*}) \leq \| -\eta_{t} \operatorname{grad} f(X_{t}) - \log_{X_{t}}(X_{*}) \|^{2}$$
$$= \eta_{t}^{2} \| \operatorname{grad} f(X_{t}) \|^{2} + \operatorname{dist}^{2}(X_{t}, X_{*}) + 2\eta_{t} \sigma$$
(13)

with

$$\sigma := \langle \operatorname{grad} f(X_t), \log_{X_t}(X_*) \rangle.$$

Proposition 5 and equation (11) give

$$\frac{\sigma}{a(X_t)} \le f^* - f(X_t) - \frac{2\sigma_{\min}(C)}{\pi^2} \text{dist}^2(X_t, X_*) \\ \le -\frac{1}{2L} \|\text{grad}f(X_t)\|^2 - \frac{2\sigma_{\min}(C)}{\pi^2} \text{dist}^2(X_t, X_*).$$

Multiplying by $2a(X_t) \eta_t$ and using $\eta_t \leq a(X_t)/L$, we get

$$2\eta_t \,\sigma \leq -\frac{a(X_t)\,\eta_t}{L} \|\text{grad}f(X_t)\|^2 - \frac{4\sigma_{\min}(C)}{\pi^2} a(X_t)\,\eta_t\,\text{dist}^2(X_t, X_*) \\ \leq -\eta_t^2 \|\text{grad}f(X_t)\|^2 - \frac{4\sigma_{\min}(C)}{\pi^2} a(X_t)\,\eta_t\,\text{dist}^2(X_t, X_*).$$

Substituting into (13), we obtain the desired result.

Theorem 10 (Convergence of RGD for the Procrustes problem). Let C be invertible ($\sigma_{\min}(C) > 0$) and X_* the (unique) minimizer of f. Then, Riemannian gradient descent (12) in the orthogonal group, starting by a point $X_0 \in \mathbb{O}(n)$ such that

$$\operatorname{dist}(X_0, X_*) < \pi,$$

and ran with fixed step size

$$\eta_t \equiv \eta \le \frac{1 + \cos(\operatorname{dist}(X_0, X_*))}{4\sigma_{\max}(C)},$$

produces iterates X_t that satisfy

$$\operatorname{dist}^{2}(X_{t}, X_{*}) \leq \left(1 - \frac{1}{\pi^{2}} (1 + \cos(\operatorname{dist}(X_{0}, X_{*}))) \sigma_{\min}(C) \eta\right)^{t} \operatorname{dist}^{2}(X_{0}, X_{*}).$$

Proof. We do the proof by induction.

For t = 0, the inequality is trivially true.

We now assume that the inequality is true for t and we wish to show that it is true also for t + 1.

Since $\operatorname{dist}(X_t, X_*) \leq \operatorname{dist}(X_0, X_*)$, we also get that the largest possible rotation $|r(X_t, X_*)|_{\max}$ induced by $X_t^T X_*$ satisfies

$$|r(X_t, X_*)|_{\max} \le \sqrt{\sum_{i=1}^n r_i(X_t, X_*)^2} = \operatorname{dist}(X_t, X_*) \le \operatorname{dist}(X_0, X_*),$$

where $r_i(X_t, X_*)$ are the rotations induced by the matrix $X_t^T X_*$. The first equality comes from equation (5).

By the definition of $a(X_t)$ in Proposition (5), we have

$$a(X_t) = \frac{1 + \cos(|r(X_t, X_*)|_{\max})}{4} \ge \frac{1 + \cos(\operatorname{dist}(X_0, X_*))}{4}$$

thus $\eta \leq a(X_t)/L$.

Since η satisfies the previous bound, the outcome of Proposition 9 holds, and combining it with the induction hypothesis, we get

$$dist^{2}(X_{t+1}, X_{*}) \leq \left(1 - \frac{4}{\pi^{2}}\sigma_{\min}(C)a(X_{t})\eta\right) dist^{2}(X_{t}, X_{*}) \leq \left(1 - \frac{1}{\pi^{2}}(1 + \cos(dist(X_{0}, X_{*})))\sigma_{\min}(C)\eta\right) dist^{2}(X_{t}, X_{*}) \leq \left(1 - \frac{1}{\pi^{2}}(1 + \cos(dist(X_{0}, X_{*})))\sigma_{\min}(C)\eta\right)^{t+1} dist^{2}(X_{0}, X_{*}).$$

This concludes the induction.

Remark. If C is singular, then the previous theorem only states that the distances of the iterates of gradient descent to the set of optima do not increase. In that case we can still prove an algebraic convergence rate for the function values of Riemannian gradient descent based only on weak-quasi-convexity. **Remark.** The assumption $\operatorname{dist}(X_0, X_*) < \pi$ allows to bound globally $|r(X_t, X_*)|_{\max}$ from above by $\operatorname{dist}(X_0, X_*)$ and as a result keep the quantity $1 + \cos(|r(X_t, X_*)|_{\max})$ far away from 0 over the course of gradient descent. Intuitively, it does not allow the algorithm to go too close to non-optimal critical points. Gradient descent would not stick to non-optimal critical points, but it would probably slow down a lot.

We now show an algebraic convergence rate for gradient descent that covers also the case that C is singular.

Theorem 11. Gradient descent applied to f for any square non-zero matrix C, starting from $X_0 \in \mathbb{O}(n)$ such that

$$\operatorname{dist}(X_0, X_*) < \pi$$

and with fixed step size

$$\eta \le \frac{1 + \cos(\operatorname{dist}(X_0, X_*))}{4\sigma_{\max}(C)},$$

produces iterates X_t that satisfy

$$f(X_t) - f^* \le \frac{2L + \frac{1}{\eta}}{(1 + \cos(\operatorname{dist}(X_0, X_*)))t + 4} \operatorname{dist}^2(X_0, X_*) = \mathcal{O}\left(\frac{1}{t}\right)$$

Proof. Since we still satisfy all the hypotheses of Theorem 10, we know that for all $t \ge 0$ it holds $dist(X_t, X_*) \le dist(X_0, X_*) < \pi$. This implies that

$$a(X_t) \ge \frac{1 + \cos(\operatorname{dist}(X_0, X_*))}{4} > 0,$$

which implies that the function f is weakly-quasi-convex (Proposition 3) at every X_t such that:

$$\langle \operatorname{grad} f(X_t), -\log_X(X_*) \rangle \ge \frac{1}{2} (1 + \cos(\operatorname{dist}(X_0, X_*)))(f(X_t) - f_*).$$

Denoting $C_0 := \frac{1 + \cos(\operatorname{dist}(X_0, X_*))}{4}$ and $\Delta_t := f(X_t) - f^*$, we can write

$$2C_0\Delta_t \le \langle \operatorname{grad} f(X_t), -\log_{X_t}(X_*) \rangle.$$
(14)

Similar to the proof of Theorem 10, by the hypothesis on the step size η_t , Lemma 9 shows that $-\eta_t X_{t+1}$ is in the injectivity domain of exp at X_t . Hence, by the definition of Riemannian gradient descent, we have

$$\log_{X_t}(X_{t+1}) = -\eta \operatorname{grad} f(X_t). \tag{15}$$

In addition, the smoothness property of f (equation 10) gives

$$\Delta_{t+1} - \Delta_t \le \langle \operatorname{grad} f(X_t), \log_{X_t}(X_{t+1}) \rangle + \frac{L}{2} \operatorname{dist}^2(X_t, X_{t+1}).$$

Substituting (15), we obtain

$$\Delta_{t+1} - \Delta_t \le \left(-\eta + \frac{L}{2}\eta^2\right) \|\operatorname{grad} f(X_t)\|^2 \le 0.$$
(16)

By Proposition 2, we have

$$\begin{split} \operatorname{dist}^2(X_{t+1},X_*) &\leq \operatorname{dist}^2(X_t,X_{t+1}) + \operatorname{dist}^2(X_t,X_*) - 2\langle \log_{X_t}(X_{t+1}), \log_{X_t}(X_*) \rangle. \\ \text{Substituting (15) into the above and rearranging terms gives} \\ &2\eta \langle \operatorname{grad} f(X_t), -\log_{X_t}(X_*) \rangle \leq \operatorname{dist}^2(X_t,X_*) - \operatorname{dist}^2(X_{t+1},X_*) + \eta^2 \|\operatorname{grad} f(X_t)\|^2. \\ \text{Combining with (14), we get} \end{split}$$

$$\Delta_t \le \frac{1}{4C_0\eta} (\operatorname{dist}^2(X_t, X_*) - \operatorname{dist}^2(X_{t+1}, X_*)) + \frac{\eta}{4C_0} \|\operatorname{grad} f(X_t)\|^2.$$
(17)

Now multiplying (16) by $\frac{1}{C_0}$ and summing with (17) gives

$$\frac{1}{C_0} \Delta_{t+1} - \left(\frac{1}{C_0} - 1\right) \Delta_t \leq \frac{1}{4C_0 \eta} (\operatorname{dist}^2(X_t, X_*) - \operatorname{dist}^2(X_{t+1}, X_*)) \\ + \frac{1}{C_0} \left(-\eta + \frac{L}{2} \eta^2 + \frac{\eta}{4}\right) \|\operatorname{grad} f(X_t)\|^2.$$
(18)

By assumption, we have $\eta \leq C_0/L$, where $0 < C_0 = (1 + \cos(\operatorname{dist}(X_0, X_*)))/4 \leq 1$ and L > 0. Since

$$\frac{\eta}{C_0} \left(-1 + \frac{L}{2}\eta + \frac{1}{4} \right) \le \frac{\eta}{C_0} \left(\frac{C_0}{2} - \frac{3}{4} \right) \le -\frac{1}{4} \frac{\eta}{C_0} < 0.$$

Inequality (18) can be simplified to

$$\frac{1}{C_0}\Delta_{t+1} - \left(\frac{1}{C_0} - 1\right)\Delta_t \le \frac{1}{4C_0\eta}(\operatorname{dist}^2(X_t, X_*) - \operatorname{dist}^2(X_{t+1}, X_*)).$$

Summing from 0 to t - 1 gives

$$\frac{1}{C_0}\Delta_t + \sum_{s=1}^{t-1}\Delta_s - \left(\frac{1}{C_0} - 1\right)\Delta_0 \le \frac{1}{4C_0\eta} \left(\operatorname{dist}^2(X_0, X_*) - \operatorname{dist}^2(X_t, X_*)\right).$$

From the smoothness property (10) with $Y \rightsquigarrow X$ and $X \rightsquigarrow X_*$, we get

$$\Delta_0 \le \frac{L}{2} \operatorname{dist}^2(X_0, X_*).$$

Combining these two inequalities leads to

$$\frac{1}{C_0} \Delta_t + \sum_{s=0}^{t-1} \Delta_s \le \frac{1}{C_0} \Delta_0 + \frac{1}{4C_0 \eta} \text{dist}^2(X_0, X_*) \\ \le \frac{1}{2C_0} \left(L + \frac{1}{2\eta} \right) \text{dist}^2(X_0, X_*).$$

Since (16) holds for all $t \ge 0$, it also implies $\Delta_t \le \Delta_s$ for all $0 \le s \le t$. Substituting

$$t\Delta_t \le \sum_{s=0}^{t-1} \Delta_s$$

into the inequality from above, we obtain

$$\Delta_t \le \frac{1}{2C_0} \frac{L + \frac{1}{2\eta}}{\frac{1}{C_0} + t} \text{dist}^2(X_0, X_*) = \frac{L + \frac{1}{2\eta}}{2(C_0 t + 1)} \text{dist}^2(X_0, X_*),$$

we obtain the desired result.

6 Distributed orthogonal Procrustes with limited communication

As an instance of a problem which cannot be solved with the traditional numerical linear algebra techniques, we consider a setting where the datasets $A \in \mathbb{R}^{k \times n}$ and $B \in \mathbb{R}^{k \times n}$ are distributed among *p*-many different computing nodes in an arbitrary way. These nodes communicate messages with a *master node* in a small bit precision. As the communication cost is now widely considered as the main bottleneck of such task, we consider the problem of computing the total number of bits that the nodes need to communicate in order to compute an orthogonal transformation $X \in \mathbb{O}(n)$, such that $||AX - B||^2 < \epsilon$. Such an analysis is possible only when there is an available algorithm that can solve the non-distributed version of the problem with linear convergence. Thus, from now on we restrict ourselves in the case that C is invertible, i.e. $\sigma_{\min}(C) > 0$.

Every node holds a pair of datasets $A_i, B_i \in \mathbb{R}^{k_i \times n}$, which give rise to a local optimization problem of $-\operatorname{Tr}(C_iX) := f_i(X)$, where $C_i = B_i^T A_i$. Notice that it holds

$$\sum_{i=1}^{p} f_i(X) = -\operatorname{Tr}\left(\left(\sum_{i=1}^{p} C_i\right) X\right) = -\operatorname{Tr}\left(\left(\sum_{i=1}^{p} B_i^T A_i\right) X\right).$$

Regardless of the structure of the data partition, we have

$$\sum_{i=1}^{p} B_i^T A_i = B^T A.$$

This implies that

$$f(X) = \sum_{i=1}^{p} f_i(X)$$

and as a concequence

$$\operatorname{grad} f(X) = \sum_{i=1}^{p} \operatorname{grad} f_i(X).$$

We tackle this problem with a distributed version of gradient descent, where each node computes the local gradient based on its datasets A_i, B_i , then compresses it using some standard compression scheme and then transmits the compressed message to the master node. The master node sums all the compressed local gradients and send a compressed version of this sum to all local nodes. The nodes then use it as a descend direction to calculate the new iterate and the process goes on.

First, we give a small overview of the compression (quantization) strategy we use. Quantization schemes is an active area of research, which is orthogonal with our work. Note that our algorithm can be combined with many different quantization schemes. However, the one we choose to use comes from [10] (lattice-based quantization). Note that [10] develops primarily a random strategy of quantization, but we can easily make it deterministic by just choosing the closest point in the constructed lattice. The derived quantization scheme has the following properties:

Proposition 12. [10] Denoting by b the number of bits that each node uses to communicate, there exists a quantization function

$$Q: \mathbb{R}^{n(n-1)/2} \times \mathbb{R}^{n(n-1)/2} \times \mathbb{R}_+ \times \mathbb{R}_+ \to \mathbb{R}^{n(n-1)/2}.$$

which, for each w, y > 0, consists of an encoding function $\operatorname{enc}_{w,y} : \mathbb{R}^{n(n-1)/2} \to \{0,1\}^b$ and a decoding one $\operatorname{dec}_{w,y} : \{0,1\}^b \times \mathbb{R}^{n(n-1)/2} \to \mathbb{R}^{n(n-1)/2}$, such that, for all $x, x' \in \mathbb{R}^{n(n-1)/2}$, we have

- $\operatorname{dec}_{w,y}(\operatorname{enc}_{w,y}(x), x') = Q(x, x', y, w), \text{ if } ||x x'|| \le y.$
- $||Q(x, x', y, w) x|| \le w$, if $||x x'|| \le y$.
- If y/w > 1, the cost of the quantization procedure in number of bits satisfies

$$b = \mathcal{O}\left(\frac{n(n-1)}{2}\log_2\left(\frac{y}{w}\right)\right) = \mathcal{O}\left(\frac{n(n-1)}{2}\log\left(\frac{y}{w}\right)\right).$$

The reason that the number $\frac{n(n-1)}{2}$ appears is that we will be quantizing tangent vectors of $\mathbb{O}(n)$. As discussed previously, the dimension of $\mathbb{O}(n)$ (thus also the one of every tangent space) is $\frac{n(n-1)}{2}$.

For quantizing a tangent vector in $T_X \tilde{\mathbb{O}}(n)$, recall that it is of the form $X\Omega$ for some skew-symmetric matrix Ω . Since X is fixed and stored in the master node, we will be quantizing only the individual Ω matrices (which depend on the batch that each node holds). For quantizing a skew-symmetric matrix, we first need to vectorize it. The simplest way to do so is by keeping the upper triangle of the matrix and then flatten it. We call such a mapping from the space of skew-symmetric matrices skew(n) as ρ .

Lemma 13. Consider

$$\rho: \operatorname{skew}(n) \to \mathbb{R}^{\frac{n(n-1)}{2}}$$

 $defined \ by$

$$\rho(\Omega) = (\omega_{12}, \dots, \omega_{1n}, \omega_{23}, \dots, \omega_{2n}, \dots, \omega_{n-1n}),$$

where $\Omega = (\omega_{ij})_{i,j=1}^n$.

Then we have $\tilde{}$

$$\|\Omega_1 - \Omega_2\|_F = \sqrt{2} \|\rho(\Omega_1) - \rho(\Omega_2)\|_2$$

for any $\Omega_1, \Omega_2 \in \text{skew}(n)$.

Proof. The proof is direct from the definition of the Frobenius norm of a matrix and the ℓ_2 -norm of a vector. Both are the square roots of the sum of squares of the entries, thus

$$\|\Omega\|_{F}^{2} = 2\|\rho(\Omega)\|_{2}^{2}$$

since the diagonal of Ω must contain only zeros.

Taking into account also that ρ is a linear map, we get the desired result. \Box

We now present a version of gradient descent with quantized gradients. It is inspired by the original Euclidean algorithm presented in [21] and is similar to the Riemannian quantized gradient descent presented in [4]. In the previous parts of the text, we have denoted $L := \sigma_{max}(C)$ to be the smoothness constant of f. Now, the individual f_i 's have their own smoothness constants $L_i :=$ $\sigma_{max}(C_i)$. Since now $\sigma_{max}(C)$ is not known in the central node, we shall use the overapproximation $\sigma_{max}(C) \leq p \max_{i=1,\dots,p} \sigma_{max}(C_i)$ and denote

$$L := p \max_{i=1,\dots,p} \sigma_{\max}(C_i)$$

for the rest of this section.

Note also that the quantization of tangent vectors of the orthogonal group is done always through a quantization of their skew-symmetric parts.

- 1. Choose an arbitrary machine to be the master node, let it be i_0 .
- 2. Choose $X_0 \in \mathbb{O}(n)$.
- 3. Consider the following parameters

$$\sigma := 1 - \frac{1 + \cos(D)}{4} \mu \eta, \quad K := \frac{2}{\sqrt{\sigma}}, \quad \theta := \frac{\sqrt{\sigma}(1 - \sqrt{\sigma})}{4},$$
$$\sqrt{\xi} := \theta K + \sqrt{\sigma}, \quad R_t = LK \left(\sqrt{\xi}\right)^t D, \quad \eta > 0$$

where D is an over-approximation for $dist(X_0, X_*)$.

We assume that $\frac{1+\cos(D)}{4}\mu\eta \leq \frac{1}{2}$, otherwise we run the algorithm with $\sigma = \frac{1}{2}$.

In
$$T_{X_0}\mathbb{O}(n)$$
:

- 4. Compute the local Riemannian gradient $\operatorname{grad} f_i(X_0) := X_0 \Omega_{i,0}$ at X_0 in each node.
- 5. If $\operatorname{grad} f_{i_0}(X_0) =: X_0 \Omega_{i_0,0}$, encode $\rho(\Omega_{i,0})$ in each node and decode in the master node using its local information $\rho(\Omega_{i_0,0})$:

$$\omega_{i,0} = Q\left(\rho(\Omega_{i,0}), \rho(\Omega_{i_0,0}), \frac{2}{\sqrt{2}}L\pi, \frac{\theta R_0}{2\sqrt{2}p}\right).$$

Compute a quantized local gradient as

$$q_{i,0} := X_0 \rho^{-1}(\omega_{i,0}).$$

6. Sum the local quantized gradients in the master node:

$$R_0 = \sum_{i=1}^{p} q_{i,0} =: X_0 \tilde{\Omega}_0.$$

7. Encode $\rho(\tilde{\Omega}_0)$ in the master node and decode in each machine *i* using its local information $\rho(\Omega_{i,0})$:

$$\omega_0 = Q\left(\rho(\tilde{\Omega}_0), \rho(\Omega_{i,0}), \frac{1}{\sqrt{2}}\left(\frac{\theta R_0}{2} + 2L\pi\right), \frac{\theta R_0}{2\sqrt{2}}\right).$$

Compute the global quantized gradient as

$$q_0 := X_0 \rho^{-1}(\omega_0).$$

For $t \ge 0$:

8. Take a gradient step using the exponential map:

$$X_{t+1} = \exp_{X_t}(-\eta q_t)$$

with step size η .

In $T_{X_{t+1}}\mathbb{O}(n)$:

- 9. Compute the local Riemannian gradient $\operatorname{grad} f_i(X_{t+1}) =: X_{t+1}\Omega_{i,t+1}$ at X_{t+1} in each node.
- 10. Encode $\rho(\Omega_{i,t+1})$ in each node and decode in the master node using its (parallelly transported) local information from the previous step $\Gamma_{X_t}^{X_{t+1}}q_{i,t} =: X_{t+1}\tilde{\Omega}_{i,t}$:

$$\omega_{i,t+1} = Q_{X_{t+1}} \left(\rho(\Omega_{i,t+1}), \rho(\tilde{\Omega}_{i,t}), \frac{R_{t+1}}{\sqrt{2p}}, \frac{\theta R_{t+1}}{2\sqrt{2p}} \right).$$

Compute the local quantized gradients as

$$q_{i,t+1} := X_{t+1}\rho^{-1}(\omega_{i,t+1})$$

11. Sum the decoded vectors in the master node:

$$R_{t+1} = \sum_{i=1}^{p} q_{i,t+1} =: X_{t+1} \tilde{\Omega}_{t+1}.$$

12. Encode the sum in the master node and decode in each machine using its local information in the previous step $\Gamma_{X_t}^{X_{t+1}}q_t =: X_{t+1}\tilde{\Omega}_t$:

$$\omega_{t+1} = Q_{X_{t+1}}\left(\rho(\tilde{\Omega}_{t+1}), \rho(\tilde{\Omega}_t), \frac{1}{\sqrt{2}}\left(1 + \frac{\theta}{2}\right)R_{t+1}, \frac{\theta R_{t+1}}{2\sqrt{2}}\right).$$

Compute the global quantized gradient as

$$q_{t+1} = X_{t+1}\rho^{-1}(\omega_{t+1}).$$

The essential step in order to analyze the convergence behaviour of the previous algorithm is an induction that controls simultaneously the convergence of iterates to the optimum and the error in the quantization of the gradients. A prototype of this technique can be found in [21].

Proposition 14. Let the initial guess X_0 be such that

$$\operatorname{dist}(X_0, X_*) \le D < \pi$$

and the step size η such that

$$\eta \le \frac{1 + \cos(D)}{4L}.$$

Then, the previous quantized gradient descent algorithm produces iterates X_t and quantized gradients q_t that satisfy

$$(i)$$
dist² $(X_t, X_*) \le \xi^t D^2, \ (ii) ||q_{i,t} - \text{grad}f_i(X_t)|| \le \frac{\theta R_t}{2p}, \ (iii) ||q_t - \text{grad}f(X_t)|| \le \theta R_t.$

Proof. We do the proof by induction starting from the case t = 0.

For t = 0, (i) is direct by the definition of D. For showing (ii), we can write

$$\|\operatorname{grad} f_i(X_0) - q_{i,0}\| = \|X_0 \Omega_{i,0} - X_0 \rho^{-1}(\omega_{i,0})\| = \|\Omega_{i,0} - \rho^{-1}(\omega_{i,0})\| = \sqrt{2} \|\rho(\Omega_{i,0}) - \omega_{i,0}\|.$$

This is by X_0 being orthogonal and Lemma 13.

Using again Lemma (13) and L-smoothness (equation 9), we have

$$\begin{aligned} \|\rho(\Omega_{i,0}) - \rho(\Omega_{i_0,0})\| &= \frac{1}{\sqrt{2}} \|\Omega_{i,0} - \Omega_{i_0,0}\| = \frac{1}{\sqrt{2}} \|X_0 \Omega_{i,0} - X_0 \Omega_{i_0,0}\| \\ &= \frac{1}{\sqrt{2}} (\|\operatorname{grad} f_i(X_0)\| + \|\operatorname{grad} f_{i_0}(X_0)\|) \le \frac{2}{\sqrt{2}} L\pi \end{aligned}$$

and by the definition of quantization (step 5), we get

$$\|\rho(\Omega_{i,0}) - \omega_{i,0}\| \le \frac{\theta R_0}{2\sqrt{2p}}.$$

This implies (by Lemma 13 and X_0 being orthogonal) that

$$\|\operatorname{grad} f_i(X_0) - q_{i,0}\| = \sqrt{2} \|\rho(\Omega_{i,0}) - \omega_{i,0}\| \le \frac{\theta R_0}{2p}.$$

For (iii) we proceed similarly:

$$\|\operatorname{grad} f(X_0) - R_0\| \le \sum_{i=1}^p \|\operatorname{grad} f_i(X_0) - q_{i,0}\| \le \frac{\theta R_0}{2}.$$
 (19)

We need also to examine the quantity $\|\rho(\tilde{\Omega}_0) - \rho(\Omega_{i,0})\|$:

$$\|\rho(\tilde{\Omega}_{0}) - \rho(\Omega_{i,0})\| = \frac{1}{\sqrt{2}} \|R_{0} - \operatorname{grad} f_{i}(X_{0})\| = \frac{1}{\sqrt{2}} (\|R_{0} - \operatorname{grad} f(X_{0})\| + \|\operatorname{grad} f(X_{0}) - \operatorname{grad} f_{i}(X_{0}))\|) \le \frac{1}{\sqrt{2}} \left(\frac{\theta R_{0}}{2} + 2L\pi\right)$$

The first equality follows from Lemma 13 and the inequality by the previously derived inequality (19) and L-smoothness (equation 9).

By the definition of the quantization of step 7, we have

$$\|\omega_0 - \rho(\tilde{\Omega}_0)\| \le \frac{\theta R_0}{2\sqrt{2}}.$$

Using Lemma 13 again, we can write

$$||q_0 - R_0|| = \sqrt{2} ||\omega_0 - \rho(\tilde{\Omega}_0)|| \le \frac{\theta R_0}{2}.$$
 (20)

We can now write

$$||q_0 - \operatorname{grad} f(X_0)|| \le ||q_0 - R_0|| + ||R_0 - \operatorname{grad} f(X_0)|| \le \theta R_0,$$

by combining inequalities (19) and (20). This concludes the induction for t = 0.

We assume now that the inequalities hold for t and we wish to prove that they continue to hold for t + 1.

We start with (i) and denote by \tilde{X}_{t+1} the iteration of exact gradient descent (12) starting from X_t , with $\operatorname{dist}(X_t, X_*) \leq D$ where $D < \pi$.

We have

$$\operatorname{dist}(X_{t+1}, X_*) \leq \operatorname{dist}(X_{t+1}, \tilde{X}_{t+1}) + \operatorname{dist}(\tilde{X}_{t+1}, X_*)$$
$$\leq \|\eta \operatorname{grad} f(X_t) - \eta q_t\| + \sqrt{\sigma} \operatorname{dist}(X_t, X_*).$$

We have the last inequality, because dist(\tilde{X}_{t+1}, X_*) $\leq \sqrt{\sigma}$ dist(X_t, X_*) by Theorem 10 and ______

 $dist(X_{t+1}, \tilde{X}_{t+1}) \leq \|\log_{X_t}(X_{t+1}) - \log_{X_t}(\tilde{X}_{t+1})\| = \|\eta \operatorname{grad} f(X_t) - \eta q_t\|$ by Proposition 2.

By the definition of R_t and since $\eta \leq 1/L$, we have

$$\operatorname{dist}(X_{t+1}, X_*) \leq \eta \theta R_t + \sqrt{\sigma} \left(\sqrt{\xi}\right)^t D \leq \theta K \left(\sqrt{\xi}\right)^t D + \sqrt{\sigma} \left(\sqrt{\xi}\right)^t D$$
$$\leq \left(\theta K + \sqrt{\sigma}\right) \left(\sqrt{\xi}\right)^t D \leq \left(\sqrt{\xi}\right)^{t+1} D$$

which concludes the induction for the first inequality. Of course, we also get that $dist(X_{t+1}, X_*) \leq D < \pi$.

For (ii), we start by using Lemma 13, in order to write

$$\|\rho(\Omega_{i,t+1}) - \rho(\tilde{\Omega}_{i,t})\| = \frac{1}{\sqrt{2}} \left\| \operatorname{grad} f_i(X_{t+1}) - \Gamma_{X_t}^{X_{t+1}} q_{i,t} \right\|.$$
(21)

For the quantity in the right hand side, we have

$$\begin{aligned} \left\| \operatorname{grad} f_i(X_{t+1}) - \Gamma_{X_t}^{X_{t+1}} q_{i,t} \right\| &\leq \left\| \operatorname{grad} f_i(X_{t+1}) - \Gamma_{X_t}^{X_{t+1}} \operatorname{grad} f_i(X_t) \right\| \\ &+ \left\| \Gamma_{X_t}^{X_{t+1}} \operatorname{grad} f_i(X_t) - \Gamma_{X_t}^{X_{t+1}} q_{i,t} \right\| \\ &\leq L_i \operatorname{dist}(X_{t+1}, X_t) + \left\| \operatorname{grad} f_i(X_t) - q_{i,t} \right\| \\ &\leq 2 \frac{L}{p} \left(\sqrt{\xi} \right)^t D + \theta \frac{R_t}{p} \\ &= 2 \frac{L}{p} \left(\sqrt{\xi} \right)^t D + \frac{\theta L K \left(\sqrt{\xi} \right)^t}{p} \\ &= (2/K + \theta) K L \left(\sqrt{\xi} \right)^t D/p \\ &\leq (\sqrt{\sigma} + \theta K) K L \left(\sqrt{\xi} \right)^t D/p \\ &= \frac{R_{t+1}}{p}. \end{aligned}$$

By combining with equation 21, we get

$$\|\rho(\Omega_{i,t+1}) - \rho(\tilde{\Omega}_{i,t})\| \le \frac{R_{t+1}}{\sqrt{2p}}$$

and by the definition of the quantization (step 10), we have

$$\|\omega_{i,t+1} - \rho(\Omega_{i,t+1})\| \le \frac{\theta R_{t+1}}{2\sqrt{2p}}.$$

Again by Lemma 13 and using that X_{t+1} is orthogonal, we finally get

$$||q_{i,t+1} - \operatorname{grad} f_i(X_{t+1})|| = \sqrt{2} ||\omega_{i,t+1} - \rho(\Omega_{i,t+1})|| \le \frac{\theta R_{t+1}}{2p}.$$

For (iii), we have

$$\|R_{t+1} - \operatorname{grad} f(X_{t+1})\| \le \sum_{i=1}^{p} \|q_{i,t+1} - \operatorname{grad} f_i(X_{t+1})\| \le \frac{\theta R_{t+1}}{2}$$
(22)

and

$$\begin{aligned} \|R_{t+1} - \Gamma_{X_t}^{X_{t+1}} q_t\| &\leq \|R_{t+1} - \operatorname{grad} f(X_{t+1})\| + \|\operatorname{grad} f(X_{t+1}) - \Gamma_{X_t}^{X_{t+1}} \operatorname{grad} f(X_t)\| \\ &+ \|\Gamma_{X_t}^{X_{t+1}} \operatorname{grad} f(X_t) - \Gamma_{X_t}^{X_{t+1}} q_t\| \\ &\leq \frac{\theta R_{t+1}}{2} + L \operatorname{dist}(X_{t+1}, X_t) + \theta R_t \\ &\leq \frac{\theta R_{t+1}}{2} + R_{t+1} = \left(1 + \frac{\theta}{2}\right) R_{t+1}. \end{aligned}$$

by the same argument used to derive the inequality $\left\| \operatorname{grad} f_i(X_{t+1}) - \Gamma_{X_t}^{X_{t+1}} q_{i,t} \right\| \leq R_{t+1}/p$ previously. Using this bound and Lemma 13, we have

$$\|\rho(\tilde{\Omega}_{t+1}) - \rho(\tilde{\Omega}_t)\| = \frac{1}{\sqrt{2}} \|\tilde{\Omega}_{t+1} - \tilde{\Omega}_t\| = \frac{1}{\sqrt{2}} \|R_{t+1} - \Gamma_{X_t}^{X_{t+1}} q_t\| \le \frac{1}{\sqrt{2}} \left(1 + \frac{\theta}{2}\right) R_{t+1}$$

The last inequality implies that

$$\|\omega_{t+1} - \rho(\tilde{\Omega}_{t+1})\| \le \frac{\theta R_{t+1}}{2\sqrt{2}},$$

by the definition of quantization (step 12).

Applying again Lemma 13, we get

$$\|\operatorname{grad} f(X_{t+1}) - q_{t+1}\| = \|\tilde{\Omega}_{t+1} - \rho^{-1}(\omega_{t+1})\| = \sqrt{2} \|\rho(\tilde{\Omega}_{t+1}) - \omega_{t+1}\| \le \frac{\theta R_{t+1}}{2}.$$
(23)

Summing inequalities (22) and (23), we finally showed inequality (iii) for t + 1. This concludes the induction.

Based on Proposition 14, we can count the total number of bits the algorithm needs in order to achieve certain accuracy. This is the main result of this section.

Theorem 15. Let $\eta \leq \frac{1+\cos(D)}{4L}$. Then, the previous quantized gradient descent algorithm needs at most

$$b = \mathcal{O}\left(pn\frac{1}{(1+\cos(D))\eta\sigma_{\min}(C)}\log\left(\frac{p}{(1+\cos(D))\eta\sigma_{\min}(C)}\right)\log\left(\frac{D}{\epsilon}\right)\right)$$

bits in total to estimate the optimal orthogonal transformation X_* with accuracy ϵ measured in intrinsic distance.

Remark. As expected, this bit complexity depends linearly on the "condition number" of the problem $\sigma_{\max}(C)/\sigma_{\min}(C)$, linearly also to the dimension n of the problem, and logarithmically to the inverse of desired accuracy $1/\epsilon$. This is in line with the results of [21] and [4].

Proof. For computing the cost of quantization at each step, we use Proposition 12 (third bullet point).

The communication cost of encoding each $\operatorname{grad} f_i$ at t = 0 (step 5) is

$$\mathcal{O}\left(n\log\frac{\frac{2L\pi}{\sqrt{2}}}{\frac{\theta R_0}{2\sqrt{2}p}}\right) = \mathcal{O}\left(n\log\frac{4pL\pi}{\theta LK\pi}\right) \le \mathcal{O}\left(n\log\frac{2p}{\theta}\right).$$

Now we use that $\sigma \geq \frac{1}{2}$ and have

$$\frac{1}{\theta} = \frac{4}{\sqrt{\sigma}(1-\sqrt{\sigma})} \le \frac{12}{1-\sigma} = \frac{48}{(1+\cos(D))\eta\mu}$$

Thus, the previous cost becomes

$$\mathcal{O}\left(n\log\frac{p}{(1+\cos(D))\eta\mu}\right).$$

The communication cost of decoding each $q_{i,0}$ in the master node (step 7) is

$$\mathcal{O}\left(n\log\frac{\frac{1}{\sqrt{2}}\left(2L\pi + \frac{\theta R_0}{2}\right)}{\frac{\theta R_0}{2\sqrt{2}}}\right) \le \mathcal{O}\left(n\log\frac{2L\pi}{\frac{\theta R_0}{2}}\right) = \mathcal{O}\left(n\log\frac{1}{(1+\cos(D))\eta\mu}\right).$$

This is because $2L\pi \ge \frac{\theta R_0}{2}$. Thus, the total communication cost at t = 0 is

$$\mathcal{O}\left(pn\log\frac{p}{(1+\cos(D))\eta\mu}\right).$$

For t > 0, the cost of encoding grad f_i 's (step 10) is

$$\mathcal{O}\left(pn\log\frac{R_{t+1}/(\sqrt{2}p)}{\theta R_{t+1}/(2\sqrt{2}p)}\right) = \mathcal{O}\left(pn\log\frac{2}{\theta}\right) = \mathcal{O}\left(pn\log\frac{1}{(1+\cos(D))\eta\mu}\right).$$

as before.

The cost of decoding in the master node (step 12) is

$$\mathcal{O}\left(pn\log\frac{\frac{1}{\sqrt{2}}\left(1+\frac{\theta}{2}\right)R_{t+1}}{\frac{\theta R_{t+1}}{2\sqrt{2}}}\right) = \mathcal{O}\left(pn\log\frac{4}{\theta}\right) = \mathcal{O}\left(pn\log\frac{1}{(1+\cos(D))\eta\mu}\right).$$

because $\theta/2 \leq 1$.

Thus, the cost in each round of communication is in general bounded by

$$\mathcal{O}\left(pn\log\frac{p}{(1+\cos(D))\eta\mu}\right).$$

Our algorithm reaches accuracy ϵ if

 $\operatorname{dist}(X_t, X_*) \le \epsilon.$

We can now use Proposition 14 and write

$$\operatorname{dist}^{2}(X_{t}, X_{*}) \leq \xi^{t} D^{2} \leq e^{-(1-\xi)t} D^{2}$$

Thus, we need to run our algorithm for

$$\mathcal{O}\left(\frac{1}{1-\xi}\log\frac{D}{\epsilon}\right) \le \mathcal{O}\left(\frac{1}{(1+\cos(D))\mu\eta}\log\frac{D}{\epsilon}\right)$$

many iterates to reach accuracy ϵ .

The total communication cost is thus the total number of iterations (communication rounds) times the total number of bits communicated in each round:

$$\mathcal{O}\left(\frac{1}{(1+\cos(D))\mu\eta}\log\frac{D}{\epsilon}pn\log\frac{p}{(1+\cos(D))\eta\mu}\right) = \mathcal{O}\left(pn\frac{1}{(1+\cos(D))\mu\eta}\log\frac{p}{(1+\cos(D))\mu\eta}\log\frac{D}{\epsilon}\right)$$

Substituting

$$\mu = \frac{4\sigma_{min}(C)}{\pi^2}$$

by Proposition 5, we get

$$\mathcal{O}\left(pn\frac{1}{(1+\cos(D))\eta\sigma_{\min}(C)}\log\frac{p}{(1+\cos(D))\eta\sigma_{\min}(C)}\log\frac{D}{\epsilon}\right)$$

many bits in total.

7 Discussion

In this work, we showed that the (non-convex) problem of computing the polar factor of a square matrix satisfies a convexity-like structure in the orthogonal group. This structure allows the analysis of classic algorithms like gradient descent, which are not competitive theoretically and empirically compared to the state-of-the-art. In cases however that the matrix of which we wish to compute a polar factor is not known exactly but only up to some perturbation, our theory comes into play. An interesting direction for future work could be to explore other cases of "noisy" polar decomposition, for example robust polar decomposition in the sense of solving the problem

$$\min_{X \in \mathbb{O}(n)} \max_{C \in \mathbb{R}^{n \times n}} \left(-\operatorname{Tr}(CX) - \beta \sum_{i=1}^{s} \|C - C_i\|^2 \right),$$

where $\{C_i\}_{i=1}^s$ is a set of independent observations for C and $\beta > 0$ is a regularizer. To the best of our knowledge, traditional linear algebra techniques cannot be applied to such problem. A more viable approach would be min-max optimization (for instance gradient descent ascent), for which our theory could be of value.

In general, we think that the direction of discovering convexity structures for non-convex but still tractable problems in linear algebra is promising (as highlighted also by the work [5]) and can shed light to many aspects of algorithmic computation in this field.

References

- P-A Absil, Robert Mahony, and Rodolphe Sepulchre. Optimization algorithms on matrix manifolds. Princeton University Press, 2009.
- [2] Devrim Akca. Generalized procrustes analysis and its applications in photogrammetry. Technical report, ETH Zurich, 2003.
- [3] Foivos Alimisis. Characterization of optimization problems that are solvable iteratively with linear convergence. MTNS 2024, 2024.
- [4] Foivos Alimisis, Peter Davies, Bart Vandereycken, and Dan Alistarh. Distributed principal component analysis with limited communication. Advances in Neural Information Processing Systems, 34:2823–2834, 2021.
- [5] Foivos Alimisis and Bart Vandereycken. Geodesic convexity of the symmetric eigenvalue problem and convergence of steepest descent. *Journal of Optimization Theory and Applications*, pages 1–40, 2024.
- [6] Nicolas Boumal. Optimality conditions on the orthogonal group. https://www.racetothebottom.xyz/posts/optimality-orthogonal/, 2023.
- [7] Jingjing Bu and Mehran Mesbahi. A note on nesterov's accelerated method in nonconvex optimization: a weak estimate sequence approach. arXiv preprint arXiv:2006.08548, 2020.
- [8] Ralph Byers and Hongguo Xu. A new scaling for newton's iteration for the polar decomposition and its backward stability. SIAM Journal on Matrix Analysis and Applications, 30(2):822–843, 2008.
- [9] Vasileios Charisopoulos, Austin R Benson, and Anil Damle. Communication-efficient distributed eigenspace estimation. SIAM Journal on Mathematics of Data Science, 3(4):1067–1092, 2021.
- [10] Peter Davies, Vijaykrishna Gurunathan, Niusha Moshrefi, Saleh Ashkboos, and Dan Alistarh. New bounds for distributed mean estimation and variance reduction. arXiv preprint arXiv:2002.09268, 2020.
- [11] Shira Faigenbaum-Golovin and Ingrid Daubechies. Studying morphological variation: Exploring the shape space in evolutionary anthropology. arXiv preprint arXiv:2410.20040, 2024.

- [12] Walter Gander. On halley's iteration method. The American Mathematical Monthly, 92(2):131–134, 1985.
- [13] Nicholas J Higham. Computing the polar decomposition—with applications. SIAM Journal on Scientific and Statistical Computing, 7(4):1160– 1174, 1986.
- [14] Nicholas J Higham. Functions of matrices: Theory and computation, 2008.
- [15] Rustem Islamov, Xun Qian, and Peter Richtárik. Distributed second order methods with fast rates and compressed communication. arXiv preprint arXiv:2102.07158, 2021.
- [16] Wolfgang Kabsch. A solution for the best rotation to relate two sets of vectors. Acta Crystallographica Section A: Crystal Physics, Diffraction, Theoretical and General Crystallography, 32(5):922–923, 1976.
- [17] Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-lojasiewicz condition. In Machine Learning and Knowledge Discovery in Databases: European Conference, ECML PKDD 2016, Riva del Garda, Italy, September 19-23, 2016, Proceedings, Part I 16, pages 795–811. Springer, 2016.
- [18] Charles Kenney and Alan J. Laub. On scaling newton's method for polar decomposition and the matrix sign function. In 1990 American Control Conference, pages 2560–2564, 1990.
- [19] Xiang Li, Shusen Wang, Kun Chen, and Zhihua Zhang. Communicationefficient distributed svd via local power iterations. In *International Conference on Machine Learning*, pages 6504–6514. PMLR, 2021.
- [20] Hatem Ltaief, Dalal Sukkari, Aniello Esposito, Yuji Nakatsukasa, and David Keyes. Massively parallel polar decomposition on distributedmemory systems. 6(1), 2019.
- [21] Sindri Magnússon, Hossein Shokri-Ghadikolaei, and Na Li. On maintaining linear convergence of distributed learning and optimization under limited communication. *IEEE Transactions on Signal Processing*, 68:6101–6116, 2020.
- [22] Yuji Nakatsukasa, Zhaojun Bai, and François Gygi. Optimizing halley's iteration for computing the matrix polar decomposition. SIAM Journal on Matrix Analysis and Applications, 31(5):2700–2720, 2010.
- [23] Yuji Nakatsukasa and Nicholas J Higham. Backward stability of iterations for computing the polar decomposition. SIAM Journal on Matrix Analysis and Applications, 33(2):460–479, 2012.
- [24] Ion Necoara, Yurii Nesterov, and Francois Glineur. Linear convergence of first order methods for non-strongly convex optimization. *Mathematical Programming*, 175:69–107, 2019.

- [25] Joel Robbin and Dietmar Salamon. Introduction to Differential Geometry. ETH lecture notes, 2022.
- [26] Drew Schmidt. A survey of singular value decomposition methods for distributed tall/skinny data. In 2020 IEEE/ACM 11th Workshop on Latest Advances in Scalable Algorithms for Large-Scale Systems (ScalA), pages 27–34. IEEE, 2020.
- [27] Peter H Schönemann. A generalized solution of the orthogonal procrustes problem. *Psychometrika*, 31(1):1–10, 1966.
- [28] Ananda Theertha Suresh, X Yu Felix, Sanjiv Kumar, and H Brendan McMahan. Distributed mean estimation with limited communication. In *International Conference on Machine Learning*, pages 3329–3337. PMLR, 2017.