A FROBENIUS SPLITTING AND COHOMOLOGY VANISHING FOR THE COTANGENT BUNDLES OF THE FLAG VARIETIES OF GL_n

RUDOLF TANGE

ABSTRACT. Let k be an algebraically closed field of characteristic $p > 0$, let $G = GL_n$ be the general linear group over k, let P be a parabolic subgroup of G , and let \mathfrak{u}_P be the Lie algebra of its unipotent radical. We show that the Kumar-Lauritzen-Thomsen splitting of the cotangent bundle $G \times^P \mathfrak{u}_P$ of G/P has top degree $(p-1)\dim(G/P)$. The component of that degree is therefore given by the $(p-1)$ -th power of a function f. We give a formula for f and deduce that it vanishes on the exceptional locus of the resolution $G \times^P \mathfrak{u}_P \to \overline{\mathcal{O}}$ where $\overline{\mathcal{O}}$ is the closure of the Richardson orbit of P. As a consequence we obtain that the higher cohomology groups of a line bundle on $G \times^P u_P$ associated to a dominant weight are zero. The splitting of $G \times^P u_P$ given by f^{p-1} can be seen as a generalisation of the Mehta-Van der Kallen splitting of $G \times^B \mathfrak{u}$.

INTRODUCTION

Let G be a reductive group over an algebraically closed field k of positive characteristic p. For a parabolic P containing the positive Borel and P -module M, we denote by $H^i(G/P, M)$ the *i*-th chomology group of the sheaf $\mathcal{L}_{G/P}(M)$ on G/P associated to M. It is an open problem whether we have for all parabolic subgroups P and all dominant characters λ of P that

$$
H^{i}(G/P, S(\mathfrak{u}_P^*) \otimes k_{-\lambda}) = 0 \quad \text{for all } i > 0, \tag{*}
$$

where the most important case is $\lambda = 0$, see e.g. [\[2,](#page-6-0) Introduction to Ch 5]. In characteristic 0 this is an easy consequence of the Grauert-Riemenschneider Theorem, see [\[4,](#page-6-1) Thm 2.2]. In characteristic $p(*)$ is known for $P = B$, for arbitrary P and "P-regular" dominant λ , see [\[12\]](#page-7-0), and for P corresponding to sets of pairwise orthogonal short simple roots and $\lambda = 0$, see [\[15\]](#page-7-1).

It is easy to write a formula for the Euler character

$$
\sum_{i\geq 0} (-1)^i \mathrm{ch} \, H^i(G/P, S(\mathfrak{u}_P^*) \otimes k_{-\lambda}),
$$

see $[10, Sect 8.14-8.16]$ and $[3, Prop 2.1]$, so if (8) holds we get a formula for ch $H^0(G/P, S(\mathfrak{u}^*)\otimes k_{-\lambda}).$

When $\mathcal{L}_{G/P}(\lambda) = \mathcal{L}_{G/P}(k_{-\lambda})$ is ample, i.e. λ "P-regular" dominant, one gets (*) from the fact that $G \times^P u_P$ is Frobenius split. One can also use Frobenius

²⁰²⁰ Mathematics Subject Classification. 14F17,14M15,14L30.

Key words and phrases. cohomology, cotangent bundle, Frobenius splitting.

splittings to prove (*) for $\lambda = 0$ via a characteristic p-version of of the Grauert-Riemenschneider Theorem [\[13,](#page-7-3) Thm 1.2], since the canonical bundle of $G \times^P \mathfrak{u}_P$ is trivial. But then the map from $G \times^P u_P$ to the Richardson orbit closure has to be birational and the splitting has to be a $(p-1)$ -th power of a section σ of the anti-canonical bundle which vanishes on the excptional locus. This is the approach we will follow.

When I asked Thomsen about the case $G = GL_n$, he told me he expected that the pushforward to $G \times^P u_P$ of the splitting of $G \times^B u_P$ induced by the "MVdK-splitting" of $G \times^B \mathfrak{u}$ from [\[14\]](#page-7-4) is the homogeneous component of degree $(p-1)\dim(G/P)$ of the "KLT-splitting", see Section [1.2,](#page-2-0) from [\[12\]](#page-7-0). Although we can not prove this conjecture, we can show that the above component is in fact the top degree component and therefore a $(p-1)$ -th power. From this we can then deduce that this homogeneous splitting vanishes on the exceptional locus of the resolution $\varphi : [g, X] \mapsto gXg^{-1} : G \times^P \mathfrak{u}_P \to \overline{\mathcal{O}}$, where $\overline{\mathcal{O}}$ is the closure of the Richardson orbit corresponding to P , see Theorem [1](#page-5-0) in Section [2.](#page-3-0) Finally, we then deduce that $(*)$ holds in type A, see Theorem [2.](#page-6-3) In fact we can formulate this as a result for arbitrary reductive groups.

The main idea of the proof is as follows. The "KLT-splitting" from [\[12\]](#page-7-0) is the $(p-1)$ -th power of the pullback along φ of the function which maps an $n \times n$ matrix X to

$$
\prod_{i=1}^{n-1} \det \left((I_n + X)_{\leq i, \leq i} \right),\tag{1}
$$

where $Y_{\leq i,\leq i}$ denotes the submatrix of Y given by the first i rows and columns, see [\[2,](#page-6-0) Example 5.[1](#page-1-0).15].¹ Unlike in the case $P = B$, the degree of the *i*-th factor may be less than i. In Lemma's [1\(](#page-4-0)ii) and [3](#page-5-1) we determine the degree of the i-th factor and from that it follows that the product [\(1\)](#page-1-1) has degree $\dim(G/P)$.

1. Preliminaries

1.1. **Notation.** Let k be an algebraically closed field of characteristic $p > 0$ and let G be a reductive group over k. We fix a Borel subgroup $B \leq G$ and maximal torus $T \leq B$. We denote by R the set of roots of T in the Lie algebra $\mathfrak{g} = \text{Lie}(G)$ of G, and we denote the unipotent radical of B by U. We call the roots of T in $\mathfrak{u} = \text{Lie}(U)$ positive and we denote the corresponding set of simple roots by S . For a subset I of S we denote the root system spanned by I by R_I . Furthermore, we denote the corresponding parabolic subgroup containing B and its Levi subgroup containing T by P_I and L_I . Denote the character group of an algebraic group H by $X(H)$. For $I \subseteq S$ we identify $X(P_I)$ and $X(L_I)$ with $\{\lambda \in X(T) \,|\, \langle \lambda, \alpha^{\vee} \rangle = 0$ for all $\alpha \in I\}.$

For P a parabolic of G and M a P-module we write $\mathcal{L}(M)$ for the G-linearised sheaf on G/P associated to M. For $\lambda \in X(P) \leq X(T)$ we put $\mathcal{L}(\lambda) = \mathcal{L}(k_{-\lambda}),$ it is the sheaf of sections of the line bundle $G \times^P k_{-\lambda}$ on G/P . We use the same symbol $\mathcal{L}(\lambda)$ to denote the sheaf of sections of the pullback of this line bundle

 $1_{\rm Apart}$ from the degree computation, the arguments there work for any parabolic.

to $G \times^P V$ for any P-variety V. We also write $H^i(G/P, M)$ for

 $H^i(G/P, \mathcal{L}(M)) \simeq R^i {\rm ind}_P^G(M)$,

see [\[9,](#page-7-5) I.5.12]. We have that

$$
H^i(G\times^P \mathfrak{u}_P, \mathcal{L}(\lambda))=H^i(G/P, k[\mathfrak{u}_P]\otimes k_{-\lambda}),
$$

see [\[2,](#page-6-0) Lem 5.2.2].

If $p = \text{char } k$ is good for G, then we have $\mathfrak{g}/\mathfrak{p} \simeq \mathfrak{u}_P^*$ as P-modules and $G \times^P \mathfrak{u}_P$ is the cotangent bundle $T^{\vee}(G/P)$ of G/P , see [\[2,](#page-6-0) 5.1.8-11].

1.2. Frobenius splittings. By [\[2,](#page-6-0) Lem 5.1.1] the canonical bundle of $G \times^P u_P$ is trivial, so we can choose a nowhere zero global section: a volume form. It is easy to see that such a section is unique up to a scalar multiple, see [\[2,](#page-6-0) 5.1.2]. This means that we can think of Frobenius splittings (up to a scalar multiple) of $G \times^P \mathfrak{u}_P$ as certain regular functions on $G \times^P \mathfrak{u}_P$.

In [\[12,](#page-7-0) Thm 1] it was proved that, when p is good for G , the cotangent bundle $T^{\vee}(G/P)$ of G/P is Frobenius split, see also [\[2,](#page-6-0) Thm 5.1.3]. We will refer to the B-canonical splitting $\psi_P(f_-\otimes f_+)$ as the "KLT-splitting" of $T^{\vee}(G/P)$, where ψ_{P}, f_{-}, f_{+} are as defined in [\[2,](#page-6-0) Ch 5]. Actually this is only a splitting up to a scalar multiple, but in the case $G = GL_n$ we assume that the chosen volume form on $T^{\vee}(G/P)$ is such that the pullback along φ of the function given by $(1), \varphi$ $(1), \varphi$ defined as in the introduction, defines a splitting. That formula is all we need to know about the KLT-splitting in this paper.

The standard grading of $k[\mathfrak{u}_P] = S(\mathfrak{u}_P^*)$ gives a grading on $k[G \times^P \mathfrak{u}_P]$, and in [\[2,](#page-6-0) 5.1.14] it is explained that the homogeneous component of degree (p − 1) dim(G/P) of a splitting σ of $G \times^P u_P$ is again a splitting of $G \times^P u_P$. This component is B-canonical if σ is B-canonical.

1.3. A result on cohomology vanishing. The following result is probably well-known, see e.g. [\[1,](#page-6-4) Sect 7.2], but for lack of reference we give a proof.

Proposition 1. Asume p is good for G, let P be a parabolic of G, let $\lambda \in X(P)$ be dominant, let Q be the parabolic of G containing P such that $\lambda \in X(Q)$ and $\mathcal{L}_{G/O}(\lambda)$ is ample, and let L be the Levi subgroup of Q containing T. If $H^{i}(L/L \cap P, S(l/\mathfrak{l} \cap \mathfrak{p})) = 0$ for all $i > 0$, then $H^{i}(G/P, S(\mathfrak{g}/\mathfrak{p}) \otimes k_{-\lambda})$ for all $i > 0$.

Proof. By [\[12,](#page-7-0) Cor 3 to Thm 4] or [\[2,](#page-6-0) Thm 5.3] $G \times^P u_P$ is Frobenius split, so by [\[2,](#page-6-0) Lemma 1.2.7(i)] it is enough to show the vanishing for $m\lambda$, $m \gg 0$ (in fact we only need it for $p^m \lambda$ and some $m \ge 0$.

Some of the arguments below are adaptations of arguments from the proof of [\[2,](#page-6-0) Lem 5.2.7].

Each $S^j(\mathfrak{g}/\mathfrak{p})$ has a filtration with sections $S^r(\mathfrak{q}/\mathfrak{p}) \otimes S^s(\mathfrak{g}/\mathfrak{q})$, $r + s = j$, so it is enough to show that R^i ind ${}_P^G(S(\mathfrak{q}/\mathfrak{p}) \otimes S(\mathfrak{g}/\mathfrak{q}) \otimes k_{-\lambda}) = 0$ for all $i > 0$. We have $P = (L \cap P)U_Q$ and $\mathfrak{q}/\mathfrak{p} \simeq 1/\mathfrak{l} \cap \mathfrak{p}$. Note that U_Q acts trivially on $\mathfrak{q}/\mathfrak{p}$. Combining [\[9,](#page-7-5) I.6.11] and our assumption with a standard spectral sequence argument, we have

 $R^i{\rm ind}_P^G(S(\mathfrak{q}/\mathfrak{p})\otimes S(\mathfrak{g}/\mathfrak{q})\otimes k_{-\lambda})\simeq R^i{\rm ind}_Q^G({\rm ind}_P^Q S(\mathfrak{q}/\mathfrak{p})\otimes S(\mathfrak{g}/\mathfrak{q})\otimes k_{-\lambda})$. (2)

4 R. TANGE

We have $\mathfrak{p} = \mathfrak{l} \cap \mathfrak{p} \oplus \mathfrak{u}_Q$, $\mathfrak{u}_P = \mathfrak{u}_{L \cap P} \oplus \mathfrak{u}_Q$, $(\mathfrak{g}/\mathfrak{p})^* \simeq \mathfrak{u}_P$, $(\mathfrak{g}/\mathfrak{q})^* \simeq \mathfrak{u}_Q$, and $(\mathfrak{q}/\mathfrak{p})^* \simeq \mathfrak{u}_{L\cap P}$. By the arguments of [\[10,](#page-7-2) p94] there exists an affine Q-variety V₀ such that $k[V_0] \simeq k[Q \times^P \mathfrak{u}_{L \cap P}] = \text{ind}_{P}^Q S(\mathfrak{q}/\mathfrak{p})$, Q-equivariantly $(U_Q \text{ acting})$ trivially). Put $V = V_0 \times \mathfrak{u}_Q$. Then $k[V] = \text{ind}_P^Q S(\mathfrak{q}/\mathfrak{p}) \otimes S(\mathfrak{g}/\mathfrak{q})$. Now the morphism $G \times^Q V \to G/Q$ is affine, so by [\[7,](#page-7-6) Ex III.8.2] the RHS of [\(2\)](#page-2-1) is isomorphic to

$$
H^i(G \times^Q V, \mathcal{L}(\lambda)).
$$
\n(3)

By [\[6,](#page-7-7) 5.1.12] $\mathcal{L}(\lambda)$ is ample on $G \times^Q V$, since $G \times^Q V \to G/Q$ is affine. The morphism $V_0 \rightarrow \overline{Q \cdot \mathfrak{u}_{L \cap P}}$ is finite, see [\[10,](#page-7-2) p94], so the same is true for the morphisms $V \to \overline{Q \cdot \mathfrak{u}_{L \cap P}} \times \mathfrak{u}$ and $G \times^Q V \to G \times^Q (\overline{Q \cdot \mathfrak{u}_{L \cap P}} \times \mathfrak{u}_Q)$. So composing the latter with the projective morphism $G \times^Q (\overline{Q \cdot \mathfrak{u}_{L \cap P}} \times \mathfrak{u}_Q) \to \mathfrak{g}$, given by the embedding of $\overline{Q \cdot \mathfrak{u}_{L\cap P}} \times \mathfrak{u}_Q$ in g, we obtain a proper morphism $G \times^Q V \to \mathfrak{g}$. Now [\[7,](#page-7-6) III.5.3] tells us that [\(3\)](#page-3-1) is 0 if we replace λ by $m\lambda$, $m \gg 0.$

2. The main results

Throughout this section, except in Theorem [2](#page-6-3) and its proof, $G = GL_n$ $GL(k^n)$ and T is the subgroup of diagonal matrices in G. As simple roots we choose the usual characters $\varepsilon_i - \varepsilon_{i+1}$, $1 \leq i \leq n-1$, where we used additive notation for characters, and ε_i is the *i*-th coordinate function on T. Then B consists of the upper triangular matrices in G . As is well-known, the conjugacy classes of parabolic subgroups of G are labelled by the compositions of n , see e.g.[\[8,](#page-7-8) 3.2]. By ν we denote a composition of n and $P = P_{\nu} \supseteq B$ is the standard parabolic whose block sizes are given in order by ν . If A_{ν} is the set $\{\nu_1, \nu_1 + \nu_2, \dots, \sum_{j=1}^{s-1} \nu_j\}$, s the length of ν , then $P_{\nu} = P_{I_{\nu}}$, the parabolic associated to the set of simple roots $I_{\nu} = \{\varepsilon_i - \varepsilon_{i+1} | i \in \{1, ..., n-1\} \setminus A_{\nu}\}.$ We denote by λ the transposed partition of the weakly descending sorted version of ν . It is well-kown that the Richardson orbit of P_{ν} is \mathcal{O}_{λ} , the nilpotent orbit whose Jordan block sizes are given by λ , see e.g. [\[8,](#page-7-8) Thm 3.3(a)].

It is well-known that the map $\varphi : [g, X] \mapsto gXg^{-1} : G \times^P \mathfrak{u}_P \to \overline{\mathcal{O}_{\lambda}}$ is birational. Indeed the group centraliser G_X of any $X \in \mathfrak{g}$ is the set of invertible elements in the Lie algebra centraliser g_X , so is connected. Now see [\[10,](#page-7-2) 4.9] and 8.8 Remark]. It is also well-known that $\overline{\mathcal{O}_{\lambda}}$ is normal, see e.g. [\[5\]](#page-7-9) or [\[14,](#page-7-4) Sect 4.7].

For $i \in \{1, \ldots, n-1\}$ we denote by $d_{\lambda,i}$ the number of nonzero positions on the $(n-i)$ -th upper codiagonal of up. So for $\nu = (2,1,2)$ we have $d_{\lambda,1}, d_{\lambda,2}, d_{\lambda,3}, d_{\lambda,4} =$ $1, 2, 3, 2$, see the figure of \mathfrak{u}_p below.

Since each diagonal $j \times j$ block of P takes away $j - i$ nonzero positions from the *i*-th upper codiagonal, we have, if j occurs m_j times in ν ,

$$
d_{\lambda,n-i} = n - i - \sum_{j>i} (j-i)m_j = n - i - \sum_{j>i} \lambda_j = - \sum_{j=i+1}^n (\lambda_j - 1) = \sum_{j=1}^i (\lambda_j - 1).
$$

Therefore, $d_{\lambda,i} = i - \sum_{j>n-i} \lambda_j = \sum_{j=1}^{n-i} (\lambda_j - 1)$. So indeed the $d_{\lambda,i}$ only depend on λ , moreover, they determine λ .

For a square matrix X we denote by $X_{\leq i,\leq i}$ the submatrix of X given by the first i rows and columns. For an $i \times i$ matrix Y we denote by $s_i(Y)$ the trace of the j-th exterior power of Y, i.e. the sum of the diagonal $j \times j$ minors of Y. As is well-known, $\det(aI_i - Y) = a^i + \sum_{j=1}^i (-1)^j a^{i-j} s_j(Y)$, where I_i is the $i \times i$ identity matrix. So the largest j with $s_i(Y) \neq 0$ is the number of nonzero eigenvalues of Y , counted with (algebraic) multiplicity. This number also equals the rank of Y^l for l sufficiently big. We will call it the *stable rank* of Y .

Lemma 1. Let $X \in \mathcal{O}_{\lambda}$.

- (i) Any i-dimensional subspace W of $V = k^n$ contains an X-invariant subspace U of dimension $\geq \sum_{j>n-i} \lambda_j$.
- (ii) $X_{\leq i,\leq i}$ has stable rank $\leq d_{\lambda,i}$.

Proof. (i). We show this by induction on n. It is trivial when $i \leq n-r$, r the length of λ , in particular when $n = 0$. Assume $i > n - r$. Then W has nonzero intersection with $\text{Ker}(X)$ for dimension reasons. Pick v nonzero in that intersection. First note that the transformation \overline{X} induced on V/kv by X has partition μ which is obtained from λ by subtracting 1 from one part of λ and then sorting the result in weakly descending order. Indeed if we decompose V as a direct sum of X-Jordan blocks and we pick a X-Jordan block of minimal size with the property that v has nonzero component in it, then we can replace that X-Jordan block by an X-Jordan block of the same size which contains v. Now we apply the induction hypothesis to V/kv and W/kv , noting that $(n-1)-(i-1)=n-i$, to obtain an X-invariant subspace U/kv of W/kv of dimension $\geq \sum_{j>n-i} \mu_j \geq \sum_{j>n-i} \lambda_j - 1$. Now U is the X-invariant subspace we want.

(ii). The linear map $(X_{\leq i,\leq i})^i$ coincides with X^i on any X-invariant subspace U of $k^i \leq k^n$ and therefore kills it. Choosing U as in (i), it induces a linear map $k^{i}/U \to k^{i}$ and therefore has rank $\leq i - \sum_{j>n-i} \lambda_{j} = d_{\lambda,i}$.

Lemma [3](#page-5-1) below follows from Lemma [1\(](#page-4-0)ii) and the existence of the KLTsplitting, but we prefer to give a direct proof.

Lemma 2. For any $h \in \{1, \ldots, i-1\}$ there exists a regular nilpotent $i \times i$ matrix X such that $X_{\leq h,\leq h}$ is invertible.

Proof. Let (e_1, \ldots, e_i) be the standard basis of k^i . Then the regular nilpotent matrix X given by $X(e_i) = e_{i-1}$ for $2 < j \leq i$, $X(e_2) = e_1 + e_{h+1}$ and $X(e_1 + e_{h+1}) = 0$ has the desired property.

6 R. TANGE

Remark 2.1. Of course it follows from Lemma [2](#page-4-1) that there exsists a regular $i \times i$ matrix X such that $X_{\leq h,\leq h}$ is invertible for all $h \in \{1, \ldots, i-1\}$, but we won't need this.

Lemma 3. There exists $X \in \mathcal{O}_{\lambda}$ such that $X_{\leq i, \leq i}$ has stable rank $d_{\lambda,i}$.

Proof. First choose any $Y \in \mathfrak{g}$ nilpotent with partition λ and decompose k^n into Y-Jordan blocks with sizes $\lambda_1, \lambda_2, \ldots, \lambda_r$, where r is the length of λ . It suffices to find an ordered basis $\mathcal B$ of k^n such that the upper left $i \times i$ -block Z of the matrix of Y relative to this basis has stable rank $d_{\lambda,i}$.

Determine $s \leq r$ maximal with $\sum_{j=1}^{s} (\lambda_j - 1) \leq i$ and put $h = i - \sum_{j=1}^{s} (\lambda_j - 1)$. Using Lemma [2](#page-4-1) choose for each $j \leq s$ a basis of the j-th block such that the upper left $(\lambda_j - 1) \times (\lambda_j - 1)$ block of the matrix of Y relative to this basis is invertible, if $s < r$ and $h > 0$ choose a basis of the $(s + 1)$ -th block such that the upper left $h \times h$ block of the matrix of Y relative to this basis is invertible, and for the remaining blocks choose any basis.

We now form B as follows. First consider the case $i \leq n-r$. For each $j \leq s$ we pick the first $\lambda_j - 1$ basis vectors from the j-th block, if $s < r$ we append the first h basis vectors from the $(s + 1)$ -th block, and finally we append all remaining $n - i$ basis vectors. Now Z is in block diagonal form with invertible diagonal block of sizes $\lambda_1 - 1, \ldots, \lambda_s - 1, h$, where h has to be omitted if $h = 0$. Now consider the case $i > n - r$. For each $j \leq n - i$ we pick the first $\lambda_j - 1$ basis vectors from the j-th block, then we append the basis vectors from the next $r - (n - i)$ blocks, and finally we append all remaining $n - i$ basis vectors. Now Z is in block diagonal form with diagonal block sizes $\lambda_1 - 1, \ldots, \lambda_{n-i} - 1, \lambda_{n-i+1}, \ldots, \lambda_r$ where the first $n-i$ blocks are invertible, and the others nilpotent. In both cases we obtain that Z has stable rank $d_{\lambda,i}$ (when $i \leq n-r$ we have $d_{\lambda,i} = i$).

Below we will denote a function $X \mapsto E(X)$ on a closed subvariety of g just by the expression $E(X)$.

Theorem 1. The degree $(p-1)\dim(G/P)$ component of the KLT splitting of $G \times^P \mathfrak{u}_P$ is the top degree component and equals the $(p-1)$ -th power of the pullback of $\prod_{i=1}^{n-1} s_{d_{\lambda,i}} (X_{\leq i,\leq i}) \in k[\overline{\mathcal{O}_{\lambda}}]$ along the resolution $\varphi: G \times^P \mathfrak{u}_P \to \overline{\mathcal{O}_{\lambda}}$. This pullback vanishes on the exceptional locus of φ .

Proof. The KLT splitting is the pullback along φ of the function given by [\(1\)](#page-1-1). Furthermore, we have $\det(I_i + Y) = \sum_{j=0}^{i} s_j(Y)$ for any $i \times i$ matrix Y, and, of course, $s_j(X_{\leq i,\leq i}) \neq 0$ on $\mathcal{O}_{\lambda} \iff s_j(X_{\leq i,\leq i}) \neq 0$ on $\overline{\mathcal{O}_{\lambda}}$. So by Lemma's $1(ii)$ and [3](#page-5-1) the top degree component of the *i*-th factor in [\(1\)](#page-1-1) is $s_{d_{\lambda,i}}(X_{\leq i,\leq i})$. So the KLT-splitting has top degree $p-1$ times $\sum_{i=1}^{n-1} d_{\lambda,i} =$ $\dim \mathfrak{u}_P = \dim(G/P)$, and the top degree component is the $(p-1)$ -th power of the pullback along φ of the function given by the stated formula.

To prove he second assertion, put $f_{\lambda,i}(X) = s_{d_{\lambda,i}}(X_{\leq i,\leq i})$ and $f_{\lambda} = \prod_{i=1}^{n-1} f_{\lambda,i}$. The exceptional locus is $\varphi^{-1}(\overline{\mathcal{O}_{\lambda}} \setminus \mathcal{O}_{\lambda})$, so it suffices to show that f_{λ} vanishes on any $\mathcal{O}_{\mu} \subseteq \overline{\mathcal{O}_{\lambda}} \setminus \mathcal{O}_{\lambda}$. We have $\dim(\mathfrak{u}_Q) = \frac{1}{2} \dim(\mathcal{O}_{\mu}) < \frac{1}{2} \dim(\mathcal{O}_{\lambda}) = \dim(\mathfrak{u}_P)$, where Q is a standard parabolic whose Richardson orbit is \mathcal{O}_{μ} , see [\[10,](#page-7-2) 4.9]. So for some *i* we must have $d_{\mu,i} < d_{\lambda,i}$ which means that $f_{\lambda,i}$ and therefore f_{λ} vanishes on \mathcal{O}_u .

Theorem 2. Let G be any reductive group for which p is good, let $\lambda \in X(T)$ be dominant, put $I = {\alpha \in S \mid \langle \lambda, \alpha^{\vee} \rangle = 0}.$ Then $H^{i}(T^{\vee}(G/P_J), \mathcal{L}(\lambda)) = 0$ for all $J \subseteq I$ such that R_J contains all irreducible components of R_I not of type A.

Proof. By Proposition [1](#page-2-2) we may assume that $\lambda = 0$ and that all irreducible components of R have type A . Since we are dealing with cotangent bundles we may assume that G is semsimple and simply connected. By the Künneth for-mula [\[11,](#page-7-10) Prop 9.2.4] we may then assume $G = SL_n$ and finally we may assume $G = GL_n$. Now the result follows from Theorem [1](#page-5-0) and [\[13,](#page-7-3) Thm 1.2], bearing in mind that the canonical bundle of $T^{\vee}(G/P)$ is trivial, see [\[2,](#page-6-0) Lem 5.1.1], and that $R^i\varphi_*(\mathcal{O}_{T^{\vee}(G/P)})$ is the sheaf associated with the cohomology group $H^{i}(T^{\vee}(G/P), \mathcal{O}_{T^{\vee}(G/P)})$, since φ is affine.

We remind the reader that a proper birational morphism $\psi : X \to Y$ is called a rational resolution if $\psi_* \mathcal{O}_X = \mathcal{O}_Y$ and the higher direct images of \mathcal{O}_X and ω_X are 0, see [\[2,](#page-6-0) Def 3.4.1].

Corollary. The resolution φ : $G \times^P \mathfrak{u}_P \to \overline{\mathcal{O}_\lambda}$ is rational.

Proof. This follows from a standard argument, see e.g. [\[9,](#page-7-5) Lem 14.5], and Theorem [2.](#page-6-3) \Box

Remarks 2.2. 1. If $P = B$, then $d_{(n),i} = i$ for all i, so the splitting from Theorem [1](#page-5-0) equals the $(p-1)$ -th power of the pullback of $\prod_{i=1}^{n-1} \det(X_{\leq i, \leq i})$ along $\varphi: G \times^B \mathfrak{u} \to \mathcal{N}$. This is the MVdK splitting of $G \times^B \mathfrak{u}$, see [\[14\]](#page-7-4).

2. Thomsen mentioned to me another proof of Lemma [1\(](#page-4-0)ii): One can easily deduce it from the following result which can be proved by induction on n . For $X \in \mathfrak{u}_P$ let $y_{ij} = \delta_{ij} + x_{ij}$ be the (i, j) -th entry of $I_n + X$. Then any monomial $y_{i_1j_1}y_{i_2j_2}\cdots y_{i_sj_s}$ with the i_l all distinct and the j_l all distinct has degree $\leq d_{\lambda,s}$ in the x_{ij} .

Conjecture (Thomsen). The pushforward to $G \times^P u_P$ of the splitting of $G \times^B u_P$ induced by the MVdK splitting is the top degree component of the KLT splitting.

Acknowledgement

I would like to thank Jesper Funch Thomsen for helpful discussions.

REFERENCES

- [1] C. Bendel, D. Nakano, B. Parshall, C. Pillen, Cohomology for quantum groups via the geometry of the nullcone, Mem. Amer. Math. Soc. 229 (2014), no. 1077.
- [2] M. Brion, S. Kumar, Frobenius splitting methods in geometry and representation theory, Progress in Mathematics, 231, Birkhäuser Boston, Inc., Boston, MA, 2005.
- [3] A. Broer, Line bundles on the cotangent bundle of the flag variety, Invent. Math. 113 (1993), no. 1, 1-20.
- [4] A. Broer, Normality of some nilpotent varieties and cohomology of line bundles on the cotangent bundle of the flag variety, in "Lie Theory and Geometry", Prog. Math. 123 (1994), Birkhäuser, Boston, 1-19.

8 R. TANGE

- [5] S. Donkin, The normality of closures of conjugacy classes of matrices, Invent. Math. 101 (1990), no. 3, 717-736.
- [6] A. Grothendieck, Eléments de géométrie algébrique II, Étude globale élémentaire de quelques classes de morphismes, Inst. Hautes Etudes Sci. Publ. Math. No. 8, 1961. ´
- [7] R. Hartshorne, Algebraic Geometry, GTM 52, Springer-Verlag, New York-Heidelberg, 1977.
- [8] W. H. Hesselink, *Polarizations in the classical groups*, Math. Z. 160 (1978), no. 3, 217-234.
- [9] J. C. Jantzen, Representations of algebraic groups, Second edition, American Mathematical Society, Providence, RI, 2003.
- [10] J. C. Jantzen, Nilpotent orbits in representation theory, "Lie theory", Progr. Math. 228, Birkhäuser Boston, 2004, 1-211.
- [11] G. R. Kempf, Algebraic Varieties, London Math. Soc. Lecture Note Ser. 172, Cambridge University Press, Cambridge, 1993.
- [12] S. Kumar, N. Lauritzen,J. F. Thomsen, Frobenius splitting of cotangent bundles of flag varieties, Invent. Math. 136 (1999), no.3, 603-621.
- [13] V. B. Mehta, W. van der Kallen, On a Grauert-Riemenschneider vanishing theorem for Frobenius split varieties in characteristic p, Invent. Math. 108 (1992), no. 1, 11-13.
- [14] V. B. Mehta, W. van der Kallen, A simultaneous Frobenius splitting for closures of conjugacy classes of nilpotent matrices, Compositio Math. 84 (1992), no. 2, 211-221.
- [15] J. F. Thomsen, Normality of certain nilpotent varieties in positive characteristic, J. Algebra 227 (2000), no. 2, 595-613.

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LS2 9JT, LEEDS, UK Email address: R.H.Tange@leeds.ac.uk

