TURBULENT SOLUTIONS OF THE BINORMAL FLOW AND THE 1D CUBIC SCHRÖDINGER EQUATION

VALERIA BANICA AND LUIS VEGA

ABSTRACT. In the last three decades there is an intense activity on the exploration of turbulent phenomena of dispersive equations, as for instance the growth of Sobolev norms since the work of Bourgain in the 90s. In general the 1D cubic Schrödinger equation has been left aside because of its complete integrability. In a series of papers of the last six years that we survey here for the special issue of the ICMP 2024 ([12],[13],[14],[15],[16],[7],[8]), we considered, together with the 1D cubic Schrödinger equation, the binormal flow, which is a geometric flow explicitly related to it. We displayed rigorously a large range of complex behavior as creation of singularities and unique continuation, Fourier growth, Talbot effects, intermittency and multifractality, justifying in particular some previous numerical observations. To do so we constructed a class of wellposedness for the 1D cubic Schrödinger equation included in the critical Fourier-Lebesgue space $\mathcal{F}L^{\infty}$ and in supercritical Sobolev spaces with respect to scaling. Last but not least we recall that the binormal flow is a classical model for the dynamics of a vortex filament in a 3D fluid or superfluid, and that vortex motions are a key element of turbulence.

CONTENTS

1.	Introduction	- 2
1.1.	Vortex filaments dynamics and the binormal flow	2
1.2.	Links between the binormal flow and Schrödinger equations	2
1.3.	Self-similar type solutions generating one singularity	4
1.4.	Turbulent features of solutions generating several singularities	5
2.	Existence of solutions generating several singularities and unique continuation	7
3.	Fourier growth	12
4.	Talbot effects	16
5.	Intermittency and multifractality	20
6.	Well-posedness, phase blow-up and unique continuation at critical regularity	27
7.	Appendix: Passing from 1D cubic Schrödinger equation to binormal flow	32
Ref	References	

Date: December 19, 2024.

1. INTRODUCTION

1.1. Vortex filaments dynamics and the binormal flow. The evolution of vortex filaments is a key element in fluid and superfluid turbulence. We consider the classical model of the binormal flow (BF, or LIA from "local induction approximation" or VFE from "vortex filament equation"), which is the oldest, simpler, and richer, formally derived model for one vortex filament dynamics in a 3D fluid governed by Euler equations, and which is also used in superfluids. If the vorticity at time t is concentrated along an arclength-parametrized curve $\chi(t)$ in \mathbb{R}^3 , its evolution in time is expected to evolve according to the binormal flow:

(1)
$$\chi_t = \chi_x \wedge \chi_{xx}.$$

By using the Frenet system, that characterizes the tangent, normal and binormal vectors in terms of the curvature and torsion, the equation writes as:

$$\chi_t = c \, b,$$

where c stands for the curvature and b for the binormal vector, which explains the binormal flow name. The model was derived formally by using Biot-Savart integral by Da Rios in 1906 following the works of Levi-Civita ([46]). This was justified rigorously by the "if" theorem of Jerrard and Seis in 2017 ([84]). Understanding when the vorticity propagates its initial structure of being concentrated along a curve is still a very difficult open problem.

We shall recall here only a representative few advances in this direction in the last five years. Concerning the Navier-Stokes equation the Cauchy problem was proved by Bedrossian, Germain and Harrop-Griffiths ([19]) to be locally well-posed for an initial filament data with no symmetry assumptions, but for small times not allowing to observe the binormal flow and to pass to the vanishing viscosity limit. Vanishing viscosity limit was proved by Gallay and Sverak ([61]) for the particular case of axisymmetric vortex rings. The binormal flow dynamic was recovered by Fontelos and Vega ([58]) for Giga-Miyakawa solutions with initial filament data with no symmetry assumptions, but this regime does not allow for passing to the vanishing viscosity limit for positive times. In Euler equations Donati, Lacave and Miot ([53]), and previously Dávila, Del Pino, Musso and Wei by other methods ([47]), constructed solutions with vorticity concentrated on helices, which are particular solutions with helical symmetry of the binormal flow. Thus we are still facing a gap towards the case of no-symmetry vortex filaments in Euler equations.

1.2. Links between the binormal flow and Schrödinger equations. We shall present now the link between the binormal flow and Schrödinger equations that is at the basis of our results. A detailed presentation is done in §7. It is easy to see that χ is a binormal flow solution then its tangent vector $T = \chi_x$ satisfies the Schrödinger map equation with values in the sphere \mathbb{S}^2 :

(2)
$$T_t = T \wedge T_{xx},$$

....

that is the Heisenberg continuous model arrising in ferro-magnetic theory. Moreover, solutions of the Schrödinger map, and thus corresponding solutions of the binormal flow equation, are related, via the Hasimoto transformation, to the 1D focusing cubic Schrödinger equation (NLS) on the line:

(3)
$$iu_t + u_{xx} + |u|^2 u = 0$$

Indeed Hasimoto discovered in 1972 ([81]) the following transform based on the Frenet system of curves given by tangent, normal and binormal vector. If χ is a binormal flow solution with non-vanishing curvature, it is easy to show¹ that the following fonction, called filament function:

$$c(t,x)e^{i\int_0^x \tau(t,s)ds},$$

where c, τ are the curvature and torsion of the curve, satisfies to the 1D Schrödinger equation

(4)
$$iu_t + u_{xx} + (|u|^2 - f) u = 0,$$

where f is a space independent function determined by $(c, \tau)(t, 0)$. In particular Hasimoto's transform can be seen as an inverse Madelung transform². Conversely, for any function f depending only on the time variable, for instance $f \equiv 0$, from a smooth solution u of (4), Hasimoto gave a method to construct frames whose first vector is a solution of (2) and thus a solution of the binormal flow (1).

The non-vanishing curvature condition was removed by Koiso in 1997 ([92]) by considering instead of the filament function the complexified normal development of the curve given by the coefficients appearing in the derivatives of parallel transport frames. For these frames the other vectors than the tangent are relatively parallel in the sense that their variation is in the direction of the tangent vector, see Bishop's article "There are more than one way to frame a curve" ([26]). In §7 we shall present in a detailed way the Hasimoto construction in this parallel transport frames framework.

We note that using Hasimoto's approach one may try to find binormal flow solutions that generate singularities in finite time by considering smooth solutions of the 1D cubic Schrödinger equation that generate singularities in finite time. However, half of the job is to find a 1D cubic NLS solution with some precise description, smooth for example on t > 0 and generating a singularity at t = 0, and half of the job is to describe geometrically the associated binormal flow solution on t > 0 in order to understand its behavior at t = 0.

$$\begin{cases} c_t = -2c_x \,\tau - c \,\tau_x, \\ \tau_t = \left(\frac{c_{xx} - c \,\tau^2}{c} + \frac{c^2}{2}\right)_x. \end{cases}$$

¹This is due to the fact that if $\chi(t, x)$ is a binormal flow solution then, by computing and identifying the crossed second derivatives of the tangent and the normal vector $T_{tx} = T_{xt}$, $n_{tx} = n_{xt}$, it follows that its curvature and torsion satisfy the system (called intrinsic equations):

²From the intrinsic equations it follows that c^2 and 2τ satisfy a Euler-Korteweg type equation, and Hasimoto's transform sends it to Gross-Pitaesvskii; Madelung transform ensures that if u solves Gross-Pitaesvskii, then $\rho := |u|^2$ and $v := 2\nabla_x Arg u$ solve an Euler type equation with the extra quantum pressure.

1.3. Self-similar type solutions generating one singularity. An important class of solutions of the binormal flow are the self-similar solutions, that are smooth curves which develop in finite time a singularity in the shape of a corner. More precisely, arclength parametrized curves solutions of binormal flow are preserved by the rescaling $\lambda^{-1}\chi(\lambda^2 t, \lambda x)$. Therefore self-similar solutions of binormal flow are searched as $\chi(t,x) = \sqrt{t}G(\frac{x}{\sqrt{t}})$. The profile curve $G(s) = \chi(1,s)$ and its Frenet frame satisfy:

$$\frac{G(s)}{2} - \frac{sG_s(s)}{2} = G_s \wedge G_{ss} \Rightarrow -\frac{sT_s(s)}{2} = T \wedge T_{ss} \Rightarrow -\frac{scn}{2} = T \wedge (c_s n - c^2 T + c\tau b),$$

so this determines the curvature and torsion of the profile, $\tau(s) = \frac{s}{2}, c_s = 0$. Going back to the self-similar variables, the self-similar solutions form a 1-parameter family $\{\chi_a\}_{a \in \mathbb{R}^{*+}}$ with curvature $c_a(t, x) = \frac{a}{\sqrt{t}}$ and torsion $\tau_a(t, x) = \frac{x}{2t}$. Note that the filament function is $a \frac{e^{i\frac{x^2}{4t}}}{\sqrt{t}}$ with initial value $a\delta_0$, and it satisfies (4) with $f(t) = \frac{a^2}{t}$. These solutions were known and used by physicists from the 80s in the framework of reconnection of vortex filaments in ferromagnetics and superfluids (Schwarz [104], Lakshmanan and Daniel [96], Buttke [35]). They were rigorously studied by Gutiérrez, Rivas and Vega in 2003 ([77]), who proved that a corner is generated at t = 0 and the tangent vector of the profile curve G_a has a limit at infinity:

$$\exists A_a^{\pm} \in \mathbb{S}^2, \quad T_a(1,x) \xrightarrow{x \to \pm \infty} A_a^{\pm}.$$

Moreover, they proved that the value θ_a of the angle of the corner is related to the parameter a by the nonlinear formula:

(5)
$$\sin\frac{\theta_a}{2} = e^{-\frac{a^2}{2}}$$

This type of dynamics can be observed in fluids passing over a triangular obstacle and in fluid and superfluid vortex reconnection, see Figure 1. We also recall that the Constantin-Fefferman-Majda blow-up criterium ([45]) on the variation of the direction of vorticity writes $\int_0^t \|\nabla(\frac{\omega}{|\omega|})(\tau)\|_{L_x^\infty}^2 d\tau = \infty$. Here, having in mind that the tangent vector models the direction of vorticity, see also evidence in this sense in Theorem 1 in [58], we have



Figure 1. Left: vortices in a fluid flowing over a triangular obstacle, Werlé, 1963. Middle left: picture of trail vortices after reconnection 2024. Middle right: Numerical simulation of binormal flow selfsimilar solution (courtesy of Enrico Fonda 2014). Right: Direct observation of Kelvin

waves excited by quantized vortex reconnection, Fonda et al. 2014 ([57], see the two experiment movies on https://www.pnas.org/doi/suppl/10.1073/pnas.1312536110).

In a series of previous articles from 2009 to 2015 ([6],[9],[10],[11], see also [76]) we understood more generally the behaviour of curves generating one corner in finite time and smoothing immediately after. As a first step we constructed solutions for (4) with $f(t) = \frac{a^2}{t}$ that are small perturbations of $a \frac{e^{i\frac{x^2}{4t}}}{\sqrt{t}}$. Then these perturbations, that after pseudo-conformal transformation are long range scattering NLS solution, allowed us to construct via Hasimoto's method binormal flow solutions for t > 0, for which we determinate the behavior at time t = 0, and that we can continue for t < 0 in a unique way in the Hasimoto's framework.

1.4. **Turbulent features of solutions generating several singularities.** In the last three decades there is an intense activity on the analysis of turbulent behaviors of solutions of dispersive equations, as for instance the growth of Sobolev norms since the work of Bourgain in the 90s. In our works we have investigated turbulent dynamics through the 1D cubic Schrödinger completely integrable equation and its geometric version, the binormal flow, which as we have seen is an equation connected to fluids and superfluids. In the following we shall describe our framework.

As we have recalled in the previous subsection, an important class of solutions of the binormal flow are the self-similar solutions, that are smooth curves which develop in finite time a singularity in the shape of a corner. Making interact several corner singularities is a natural question that has been investigated first by physicists. Noncircular jets as square jets were studied since the 80s for the turbulent features they produce. For instance experiments were done by Todoya and Hussain ([109]), and numerics by Grinstein and De Vore ([71]), see Figure 2.



Figure 2. Axis switching in numerical simulation of square jets from [71].

At the level of the binormal flow this corresponds to consider as initial data a closed curve that is a regular polygon. Such a regular M-polygon $\chi_M(0, x)$ with corners located at $x \in \mathbb{Z}$ (we see here the closed curve as being parametrized, in a periodic way, by $x \in \mathbb{R}$) is expected to evolve by the binormal flow to skew Mq-polygons at times $\frac{p}{q} \in \mathbb{Q}$, as suggested by numerics by Jerrard and Smets ([85]), and by De la Hoz and Vega ([49]), see Figure 3. This can be seen as a Talbot effect. More precisely, in view of the generation of one corner of angle θ_a by the self-similar solution χ_a driven by the filament function $a \frac{e^{i\frac{x^2}{4t}}}{\sqrt{t}}$, and of the relation (5), one expects that the filament function of the M-polygon evolution

corresponds to considering NLS solutions with initial data of type Dirac comb distribution $\sum_{k \in \mathbb{Z}} \delta_{\frac{k}{M}}$, say instead $\sum_{k \in \mathbb{Z}} \delta_k$ to simplify the presentation, which is out of reach. However, by supposing Galilean invariance of solutions, De la Hoz and Vega proposed in [49] a filament candidate as $u(t) = \phi(t)e^{it\Delta} \sum_{k \in \mathbb{Z}} \delta_k$, which has an explicit expression at rational times $t = \frac{p}{q}$, since the fundamental solution of the periodic Schrödinger equation values at rational times:

$$e^{i\frac{p}{q}\Delta}\sum_{k\in\mathbb{Z}}\delta_k=\sum_{j\in\mathbb{Z}}\tau_j\delta(x-\frac{j}{q}),$$

with the coefficients τ_j given in terms of Gauss sums of exponentials³ (see the computation in §4). Then, they succeeded to integrate the parallel frame system (21) with filament function $u(\frac{p}{q})$ and proved that the corresponding curves are skew polygons with Mq sides, same angles, and torsion in terms of Gauss sums. This fits with the numerics performed on the square evolution, at the level of binormal flow and of the Schrödinger map (2).



Figure 3. Numerical binormal flow evolution of a square from [85], see https://www.ljll.fr/gallery/html/DSmets_LIA-fr.htm

Moreover, other turbulent features were observed numerically, as for instance the growth in phase space at the level of the tangent vectors which model the direction of vorticity, and the multifractality of the trajectories. The trajectories of corners $\chi_M(t,0)$ were numerically proved to behave as Riemann's complex function $\sum_{k \in \mathbb{Z}} \frac{e^{itk^2} - 1}{k^2}$ when M tends to infinity (De la Hoz and Vega [49], De la Hoz, Kumar and Vega [48], see Figure 4).



Figure 4. Left: Evolution by the binormal flow of a M-polygon with M = 15. Right: the trajectory in time of the solution at x = 0 (courtesy of Sandeep Kumar, see https://www.youtube.com/watch?v=bwbpKvqGk-o).

³Constants as for instance 2π that are irrelevant for the nature of the phenomena described will be neglected in general all along this article.

In a new series of papers of the last six years, that we survey here ([12], [13], [14], [15], [16], [7], [8]), by making interact several corners through the binormal flow, we displayed rigorously a large range of complex behavior as creation of singularities and unique continuation, Fourier growth, Talbot effects, intermittency and multifractality, justifying in particular the previous numerical observations. This has been done by constructing and analyzing binormal flow solutions with initial data of the shape of infinite polygonal lines with only two corners, or made by a regular M-polygon and two half-lines. To do so we constructed a class of well-posedness for the 1D cubic Schrödinger equation included in the critical Fourier-Lebesgue space $\mathcal{F}L^{\infty}$ and in supercritical Sobolev spaces with respect to scaling.

Acknowledgements: VB is partially supported by the ERC advanced grant GEOEDP and by the French ANR project BOURGEONS. LV is partially supported by MICINN (Spain) CEX2021-001142, PID2021-126813NB-I00 (ERDF A way of making Europe) and IT1247-19 (Gobierno Vasco).

2. Existence of solutions generating several singularities and unique continuation

As we have seen, a corner of angle θ_a is generated by the binormal flow self-similar solution χ_a , with a related to θ_a by (5). Moreover, the filament function of χ_a is $a \frac{e^{i\frac{x^2}{4t}}}{\sqrt{t}}$, solution of the 1D cubic Schrödinger equation (4) with $f = \frac{a^2}{t}$. Due to gauge invariance, as explained in the appendix, the solutions χ_a can be constructed from the filament function $ae^{ia^2 \log t} \frac{e^{i\frac{x^2}{4t}}}{\sqrt{t}}$, solution of the classical 1D cubic Schrödinger equation (3). Thus, if we want to construct a binormal flow evolution of a polygonal line with several corners, a first natural step is to look for a solution of (3), smooth for positive times, with data at t = 0 related to a superposition of Dirac masses. Then, the second step is to use Hasimoto's approach to construct from this solution a smooth binormal flow solution for positive times and see if it may have as a limit at t = 0 a polygonal line.

Looking for 1D cubic Schrödinger solutions on \mathbb{R} with data of Dirac mass type calls for a review on the state of the art of the Cauchy problem for (3). The rescaling that leaves invariant equation (3) is $\lambda u(\lambda^2 t, \lambda x)$, the invariant space with respect to this scaling on the Sobolev scale is $\dot{H}^{-\frac{1}{2}}$, and the one on the Fourier-Lebesgue scale is $\mathcal{F}L^{\infty}$, i.e. Fourier transform in L^{∞} . The equation is well-posed in H^s for $s \geq 0$ and the flow map is uniformly continuous on bounded sets of H^s (Ginibre and Velo [68], Tsutsumi [110], Cazenave and Weissler [39]). It turns out that this cannot hold in H^s for s < 0 (Kenig, Ponce and Vega [87], Christ, Coliander and Tao [42]). However, the Sobolev norms of Schwartz solutions have controlled growth for $-\frac{1}{2} < s < 0$ (Koch and Tataru [91] and Kilip, Visan and Zhang [88]). Finally, Harrop-Griffths, Killip and Visan proved in 2024 ([80]) the global wellposedness of (3) in H^s for s > -1/2 in the sense that the solution map on the Schwartz class admits a unique continuous extension to H^s , s > -1/2. Their result is sharp in the sense that for s < -1/2 a norm inflation with loss of regularity appear (Kishimoto [89], Carles and Kappeler [38], Oh [99]), in particular (3) is ill-posed in the Hadamard sense

in H^s , s < -1/2. In what concerns the Fourier-Lebesgue spaces, the problem is known to be locally well-posed on $\mathcal{F}L^p$ for $p < +\infty$ (Vargas and Vega [111], Grünrock [72]). In summary, we presently have a well-posedness theory that misses the critical spaces with respect to scaling.

Let us notice that the Dirac mass and its linear Schrödinger evolution, the fundamental Schrödinger solution, are of borderline regularity $H^{-\frac{1}{2}-\epsilon}$ for all $\epsilon > 0$ and $\mathcal{F}L^{\infty}$, and therefore do not fit in the scope of applicability of the previously described well-posedness results. Moreover, there are simple explicit solutions of the cubic Schrödinger equation (3) for t > 0, having the same kind of borderline regularity, given by

$$u_{\alpha}(t,x) = \alpha e^{i|\alpha|^2 \log t} e^{it\Delta} \delta_0 = \alpha e^{i|\alpha|^2 \log t} \frac{e^{i\frac{x^2}{4t}}}{\sqrt{t}}, \quad \alpha \in \mathbb{C}.$$

Actually these solutions were displayed in the proof of Kenig, Ponce and Vega in [87] to the non-uniqueness of the Cauchy problem of (3) with initial data a Dirac mass. Also, the self-similar solutions of the binormal flow are obtained by the Hasimoto approach also from them, as they differ from $|\alpha| \frac{e^{i\frac{x^2}{4t}}}{\sqrt{t}}$ just by a space-independent phase. This type of solutions allowed recently for a new insight on (3) at critical regularity. Also solutions which are smooth perturbations of u_{α} were constructed in our previous articles [6],[9],[10],[11], see also [76]. Then, following an ansatz of Kita proposed in [90] for subcritical nonlinearities (subcubic), we constructed for (3) solutions of type superpositions of u_{α} :

(6)
$$\sum_{j\in\mathbb{Z}}A_k(t)e^{it\Delta}\delta_k = \sum_{k\in\mathbb{Z}}A_k(t)\frac{e^{i\frac{(x-k)^2}{4t}}}{\sqrt{t}},$$

as follows.

Theorem 2.1. (Construction of critical NLS solutions, [14]) Let $s > \frac{1}{2}, 0 < \gamma < 1$, $\{\alpha_k\} \in l^{2,s}$. There exists T > 0 and a unique solution of (3) on (0,T) of the form

(7)
$$u_{\{\alpha_j\}}(t,x) = \sum_{k \in \mathbb{Z}} A_k(t) e^{it\Delta} \delta_k(x).$$

with

$$A_{k}(t) = e^{-i(|\alpha_{k}|^{2} - 2\sum_{j \in \mathbb{Z}} |\alpha_{j}|^{2}) \log t} (\alpha_{k} + R_{k}(t))$$

and

$$\sup_{0 < t < T} t^{-\gamma} \|\{R_k(t)\}\|_{l^{2,s}} + t \|\{\partial_t R_k(t)\}\|_{l^{2,s}} < C$$

Moreover, if $s \ge 1$ then the solution can be extended to $(0, \infty)$.

The $l^{2,s}$ hypothesis was relaxed to $l^{p,s}$ by Bravin and Vega ([33]). Also, solutions that are smooth perturbations of $u_{\{\alpha_i\}}$ were constructed by Guérin ([75]).

Let us note that by using the pseudo-conformal transformation

$$v(\tau, y) := \frac{e^{i\frac{y^2}{4\tau}}}{\sqrt{\tau}} \,\overline{u}(\frac{1}{\tau}, \frac{y}{\tau}),$$

equation (3) is transformed into

$$iv_t + v_{xx} + \frac{1}{t}|v|^2 v = 0.$$

Moreover, the ansatz (6) translates into simply being in the periodic setting of this equation. In particular, for this periodic 1D cubic NLS with time-variable coefficient we proved by Theorem 2.1 the existence of wave operators. We proved also asymptotic completeness in [8], that we review in $\S6$.

The proof goes as follows: plugging the ansatz $u(t,x) = \sum_{k \in \mathbb{Z}} A_k(t) e^{it\Delta} \delta_k(x)$ into equation (3) leads to a discrete nonautonomous Hamiltonian system on $\{A_k(t)\}$. We solve this discrete system by a fixed point argument, based on integrations by parts from the nonresonant phases. In $\S2.0.1$ we give more details on the proof.

The NLS solutions in Theorem 2.1 are the starting point, via the Hasimoto transform, that allowed us to construct the evolution of polygonal lines by the binormal flow as follows.

Theorem 2.2. (Evolution of polygonal lines by the binormal flow, [12]) We consider an arclength parametrized polygonal line χ_0 with corners located at $x = k \in \mathbb{Z}$, of angles⁴⁵ θ_k such that $a_k := \sqrt{-\frac{2}{\pi} \log\left(\sin\left(\frac{\theta_k}{2}\right)\right)} \in l^{2,\frac{3}{2}^+}$. Then, there exists $\chi(t)$ smooth solution of the binormal flow on \mathbb{R}^* , weak solution on \mathbb{R} , with

$$|\chi(t,x) - \chi_0(x)| \le C\sqrt{t}, \quad \forall x \in \mathbb{R}, |t| \le 1.$$

The solution is unique in the framework of curves having as filament functions the solutions in Theorem 2.1.

We note that the NLS solutions in Theorem 2.1 blow-up at time t = 0 with a loss of phase. Phase blow up phenomena were encountered for the Schrödinger equation since the works of Merle in the 90s ([97], see also [98]). Here, we see that despite this loss of phase the solutions of the binormal flow associated via Hasimoto's transform are uniquely continued after the singularity time t = 0. This can be seen as a way of continuation for NLS despite the phase loss. In $\S2.0.2$ we give details on the proof of Theorem 2.2.

2.0.1. Sketch of the proof of Theorem 2.1. By plugging the ansatz $u(t) = \sum_{k \in \mathbb{Z}} A_k(t) e^{it\Delta} \delta_k$ into equation (3) we obtain:

$$\sum_{k \in \mathbb{Z}} i\partial_t A_k(t) e^{it\Delta} \delta_k = |\sum_{j \in \mathbb{Z}} A_j(t) e^{it\Delta} \delta_j|^2 (\sum_{j \in \mathbb{Z}} A_j(t) e^{it\Delta} \delta_j).$$

⁴i.e. θ_k is the angle of the self-similar binormal flow solution χ_{a_k} , see (5). ⁵The fact that $\{a_k\} \in l^{2,3}$ implies that the angles θ_j tend to π as $|j| \to \infty$.

The family $e^{it\Delta}\delta_k(x) = \frac{e^{i\frac{(x-k)^2}{4t}}}{\sqrt{t}}$ is, modulo some constants, an orthonormal family of $L^2(0,4t)$, so by taking the scalar product with $e^{it\Delta}\delta_k$ we obtain the following nonautonomous Hamiltonian system of the coefficients $\{A_k(t)\}$:

$$\begin{split} i\partial_t A_k(t) &= \int_0^{4t} |\sum_{j\in\mathbb{Z}} A_j(t) \frac{e^{i\frac{(x-j)^2}{4t}}}{\sqrt{t}} |^2 (\sum_{j\in\mathbb{Z}} A_j(t) \frac{e^{i\frac{(x-j)^2}{4t}}}{\sqrt{t}}) \frac{-e^{i\frac{(x-k)^2}{4t}}}{\sqrt{t}} \, dx. \\ &= \frac{1}{4t} \sum_{k-j_1+j_2-j_3=0} e^{-i\frac{k^2-j_1^2+j_2^2-j_3^3}{4t}} A_{j_1}(t) \overline{A_{j_2}(t)} A_{j_3}(t). \end{split}$$

We split the summation indices into the following two sets:

$$NR_{k} = \{(j_{1}, j_{2}, j_{3}) \in \mathbb{Z}^{3}, k - j_{1} + j_{2} - j_{3} = 0, k^{2} - j_{1}^{2} + j_{2}^{2} - j_{3}^{2} \neq 0\},\$$

$$Res_{k} = \{(j_{1}, j_{2}, j_{3}) \in \mathbb{Z}^{3}, k - j_{1} + j_{2} - j_{3} = 0, k^{2} - j_{1}^{2} + j_{2}^{2} - j_{3}^{2} = 0\}.$$

As we are in one dimension, the resonant set is simply:

$$Res_k = \{(k, j, j), (j, j, k), j \in \mathbb{Z}\},\$$

so the system writes:

(8)

$$i\partial_t A_k(t) = \frac{1}{4t} \sum_{(j_1, j_2, j_3) \in NR_k} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^3}{4t}} A_{j_1}(t) \overline{A_{j_2}(t)} A_{j_3}(t) + \frac{A_k(t)}{4t} (2\sum_j |A_j(t)|^2 - |A_k(t)|^2).$$

The system preserves the "mass" $\sum_k |A_k(t)|^2$, since for any $a : \mathbb{Z} \to \mathbb{R}$ we can compute, by using a standard symmetrization argument and the fact that $a(k) - a(j_1) + a(j_2) - a(j_3)$ vanishes on the resonant set:

$$\partial_t \sum_k a(k) |A_k(t)|^2 = \frac{1}{4t} \sum_{k, NR_k} (a(k) - a(j_1) + a(j_2) - a(j_3)) e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^3}{4t}} A_{j_1}(t) \overline{A_{j_2}(t)} A_{j_3}(t)).$$

We shall obtain the existence of solutions of system (8) of the form

$$A_{k}(t) = e^{-i(|\alpha_{k}|^{2} - 2M)\log t}(\alpha_{k} + R_{k}(t)),$$

with R_k in

$$X^{\gamma} := \{\{f_k\} \in \mathcal{C}([0,T]; l^{2,s}), \sup_{0 \le t < T} t^{-\gamma} \|\{f_k(t)\}\|_{l^{2,s}} + t \|\{\partial_t f_k(t)\}\|_{l^{2,s}} < \infty\}.$$

By using the "mass" $\sum_j |A_j(t)|^2$ conservation, and the evolution law of $|A_k(t)|^2$, we reduce to proving that the operator $\Phi : \{R_k\} \to \{\Phi_k(\{R_j\})\}$ defined as

$$\Phi_k(\{R_j\})(t) = i \int_0^t g_k(\tau) d\tau - i \int_0^t \int_0^\tau \Im(g_k(s) \overline{(\alpha_k + R_k(s))}) ds \left(\alpha_k + R_k(\tau)\right) \frac{d\tau}{4\pi\tau},$$

where

$$g_k(t) = \frac{1}{4\pi t} \sum_{\substack{(j_1, j_2, j_3) \in NR_k}} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^2}{4t}} e^{-i\frac{|\alpha_k|^2 - |\alpha_{j_1}|^2 + |\alpha_{j_2}|^2 - |\alpha_{j_3}|^2}{4\pi}\log t} \times \frac{1}{2} \sum_{\substack{(j_1, j_2, j_3) \in NR_k}} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^2}{4t}} e^{-i\frac{|\alpha_k|^2 - |\alpha_{j_1}|^2 + |\alpha_{j_2}|^2 - |\alpha_{j_3}|^2}{4\pi}} \log t \times \frac{1}{2} \sum_{\substack{(j_1, j_2, j_3) \in NR_k}} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^2}{4t}} e^{-i\frac{|\alpha_k|^2 - |\alpha_{j_1}|^2 + |\alpha_{j_2}|^2 - |\alpha_{j_3}|^2}{4\pi}} \log t \times \frac{1}{2} \sum_{\substack{(j_1, j_2, j_3) \in NR_k}} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^2}{4t}} e^{-i\frac{|\alpha_k|^2 - |\alpha_{j_1}|^2 + |\alpha_{j_2}|^2}{4\pi}} \log t \times \frac{1}{2} \sum_{\substack{(j_1, j_2, j_3) \in NR_k}} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^2}{4t}} e^{-i\frac{|\alpha_k|^2 - |\alpha_{j_1}|^2 + |\alpha_{j_2}|^2}{4\pi}} \log t \times \frac{1}{2} \sum_{\substack{(j_1, j_2, j_3) \in NR_k}} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^2}{4t}} e^{-i\frac{|\alpha_k|^2 - |\alpha_{j_1}|^2}{4\pi}} \log t \times \frac{1}{2} \sum_{\substack{(j_1, j_2, j_3) \in NR_k}} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^2}{4t}} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^2}{4\pi}} \log t \times \frac{1}{2} \sum_{\substack{(j_1, j_2, j_3) \in NR_k}} e^{-i\frac{k^2 - j_1^2 + j_2^2 - j_3^2}{4\pi}} e^{-i\frac{k^2 - j_1^2 - j_3^2}{4\pi}} e^{-i\frac{k^2 - j_1^2 - j_1^2 - j_3^2}{4\pi}} \log t \times \frac{1}{2} \sum_{\substack{(j_1, j_2, j_3) \in NR_k}} e^{-i\frac{k^2 - j_1^2 - j_1^2 - j_1^2 - j_1^2}{4\pi}} e^{-i\frac{k^2 - j_1^2 -$$

$$\times (\alpha_{j_1} + R_{j_1}(t))(\alpha_{j_2} + R_{j_2}(t))(\alpha_{j_3} + R_{j_3}(t)),$$

is a contraction on a small ball in X^{γ} . The estimates in the fixed point arguments are based on integrations by parts from the nonresonant phases, which yield decay in time necessary for integration near zero.

2.0.2. Sketch of the proof of Theorem 2.2. The proof can be synthethized as follows:

Step 1: define $\alpha_k = a_k e^{i\gamma_k}$, with a_k, γ_k expressed in terms of the curvature angles and torsion angles of the polygonal line χ_0 in a way to be specified later (for instance a_k is determined from the curvature angle θ_k by formula (5)),

Step 2: use Theorem 2.1 to get a solution $u_{\{\alpha_k\}}$ of the NLS equation (3), smooth for t > 0,

Step 3: construct for t > 0 the binormal flow solution χ obtained from $u_{\{\alpha_k\}}$ by the Hasimoto construction detailed in §7.0.2. In particular, the frames (T, e_1, e_2) satisfy the ODEs in time and in space (23)-(24) with system matrices involving $u_{\{\alpha_k\}}$,

Step 4: get a trace $\chi(0)$ for $\chi(t)$ as t goes to zero with the rate of convergence \sqrt{t} , as from (26) we have

$$|\partial_t \chi(t)| \le 2|u_{\{\alpha_k\}}(t)| \le \frac{C(\|\alpha_j\|_{l^1})}{\sqrt{t}}$$

The goal is now to show that $\chi(0)$ and χ_0 coincide,

Step 5: for $x \notin \mathbb{Z}$ obtain a limit as t goes to zero for T(t, x), based on estimating oscilatory integrals involving $u_{\{\alpha_k\}}$,

Step 6: prove that the vectors T(0, x) are constant for k < x < k + 1, again based on estimating oscilatory integrals involving $u_{\{\alpha_k\}}$, so $\chi(0)$ is a polygonal line,

Step 7: recover a self-similar binormal flow profile χ_{a_k} on self-similar paths towards (0, k): first conclude by Arzela-Ascoli's theorem that there is a limit for the frames via self-similar path $(t_n^k, k + x\sqrt{t_n^k})$ for some sequence $t_n^k \stackrel{n \to \infty}{\to} 0$, and that the limit frame satisfies the EDO of the self-similar profile χ_{a_k} . Then conclude by uniqueness that there exists a rotation Θ_k such that

$$\lim_{n \to \infty} T(t_n^k, k + x\sqrt{t_n^k}) = \Theta_k(T_{a_k}(x)),$$

where T_{a_k} is the tangent vector of the profile curve $\chi_{a_k}(1)$, which has a limit $A_{a_k}^{\pm}$ at $\pm \infty$. This allows for recovering the curvature angle of χ_0 at x = k, since the values $T(0, k^{\pm})$ are limits of $T(0, k + x\sqrt{t_n^k})$, which in turn by Step 5 can be approximated by $T(t_n^k, k + x\sqrt{t_n^k})$ which eventually is close to $\Theta_k A_{a_k}^{\pm}$, by taking x large enough,

Step 8: recover the torsion angles of χ_0 by using also a similar analysis for the modulated normal vectors

$$\tilde{N}(t,x) = e^{i\sum_{j} |\alpha_{j}|^{2} \log \frac{x-j}{\sqrt{t}}} N(t,x).$$

At this stage the relation between the sequences $\{\alpha_k\}$ and $\{\gamma_k\}$ and the curvature and torsion angles of the initial curve χ_0 becomes explicit,

Step 9: unique continuation after singularity time: since now that the theorem is proved for positive time evolutions, we use the time invariance of binormal flow to construct χ for negative times by $\chi(t, x) = \tilde{\chi}(-t, -x)$, where $\tilde{\chi}$ is the evolution for positive times with initial data the polygonal line $\tilde{\chi}(0, x) = \chi(0, -x)$. Thus the continuation of χ for negative times is done by the evolution of $\chi(0)$ with the sense of parametrization inverted.

Let us recall that in the simple case of self-similar solutions χ_a , for which $\chi_a(0)$ is a curve with one corner of angle θ_a , for negative times the solution is a rotation and symmetry of the solution for positive times. This is due in particular to the fact that the filament function is the same for the evolution of $\chi_a(0, x)$ as well as for the evolution of $\chi_a(0, -x)$, namely $a \frac{e^{i\frac{x^2}{4t}}}{\sqrt{t}}$. In our case the inversion of the sense of parametrization has an evolution that is not geometrically trivially related to the evolution of $\chi(0, x)$ for positive times. This can be seen from the fact that the filament function used for constructing the evolution of $\chi(0, x)$ is

$$\sum_{k} e^{-i(|\alpha_{k}|^{2}-2\sum_{j\in\mathbb{Z}}|\alpha_{j}|^{2})\log t} (\alpha_{k} + R_{k}(t)) \frac{e^{i\frac{(x-k)^{2}}{4t}}}{\sqrt{t}},$$

while the filament function used for constructing the evolution of $\chi(0, -x)$ is

$$\sum_{k} e^{-i(|\alpha_{-k}|^2 - 2\sum_{j \in \mathbb{Z}} |\alpha_j|^2) \log t} (\alpha_{-k} + \tilde{R}_k(t)) \frac{e^{i\frac{(x-k)^2}{4t}}}{\sqrt{t}}.$$

As a byproduct this is a way to uniquely continuate the NLS solutions after a generation of singularities with phase loss.

3. Fourier growth

As said before, in the last decades there is an intense activity on the analysis of turbulent dynamics related to dispersive equations, as for instance the growth of Sobolev norms since the work of Bourgain in the 90s. To quote just a few, such studies were done for non-integrable equations as the linear Schrödinger equation with potential, the 2D cubic NLS, systems of 1D cubic NLS (Bourgain [30], [31], [32], Kuksin [94], [95], Colliander, Keel, Staffilani, Takaoka and Tao [44], Sohinger [106], Carles and Faou [36], Grébert, Paturel and Thomann [70], Delort [51], Hani [78], Hani, Pausader, Tzvetkov and Visciglia [79], Guardia and Kaloshin [73], Planchon, Tzvetkov and Visciglia [101], Bambusi, Grébert, Maspero and Robert [4], Carles and Gallagher [37], Deng and Germain [52], Thomann 2021 [108], Bambusi, Langella and Montalto 2022 [5], Giuliani and Guardia [69], Faou and Raphaël [56], Guardia, Hani, Haus, Maspero and Procesi [74], ...) In general the 1D cubic NLS and the Schrödinger map were left aside because of their complete integrability. We note however that turbulent behavior was observed for abstract integrable equations as of Szegö type (Gérard and Grellier [63], [64], Pocovnicu [102], Gérard, Lenzmann, Pocovnicu and Raphaël [67], Biasi and Evnin [25], Gérard, Grellier and He [65], Gérard and Lenzmann |66|,...).

In [13]-[14] we displayed a Fourier modes growth shifted for the Schrödinger map (2), that on one hand is directly connected via Hasimoto's transform with the 1D cubic NLS, and on the other hand is also directly connected to the binormal flow, which arises as a model for vortex filaments, as explained before.

More precisely, let us first recall that the Schrödinger map (2) has interaction energy:

$$\int |T_x(t,x)|^2 dx.$$

From now on we consider the binormal flow solutions of Theorem 2.2 that are constructed from the NLS solutions $u_{\{\alpha_j\}}$ in Theorem 2.1. Their tangent vector T satisfies the Schrödinger map (2) strongly on $(-1, 1) \setminus \{0\}$ and weakly on (-1, 1). These solutions have infinite interaction energy since, in view of (26), $|T_x(t, x)| = |u_{\{\alpha_j\}}(t, x)|^2$ is a periodic function. However, we shall prove they have a new type of finite energy.

We recall again that the Constantin-Fefferman-Majda blow-up criterium ([45]) on the variation of the direction of vorticity, writes $\int_0^t \|\nabla(\frac{\omega}{|\omega|})(\tau)\|_{L^\infty_x}^2 d\tau = \infty$. Here, having in mind that the tangent vector T models the direction of vorticity, see also evidence in this sense in Theorem 1 in [58], we have:

$$\int_0^t \|T_x(\tau)\|_{L^\infty}^2 d\tau = \int_0^t \|u_{\{\alpha_j\}}(\tau)\|_{L^\infty}^2 d\tau = \int_0^t \frac{\sum_k ((\alpha_k + R_k(\tau))e^{i\frac{(x-k)^2}{4\tau}})^2}{\tau} d\tau,$$

which is infinite for generic sequences $\{\alpha_j\}$ since $R_k(\tau)$ behaves as τ^{1^-} uniformly in k as τ goes to zero. In [13]-[14] we analyzed T_x in the phase variable. As a first result we identified a finite energy density framework which points out a growth at large frequencies as follows.

Theorem 3.1. (A finite energy density framework, [13]) Let T be the solution of (2) obtained as tangent vector of the binormal flow evolution of a polygonal line from Theorem 2.2. Then

$$\Xi(t) := \lim_{n \to \infty} \int_n^{n+1} |\widehat{T_x}(t,\xi)|^2 d\xi$$

is conserved for $t \in (0, 1)$ with a discontinuity at t = 0. Moreover, there is an instantaneous growth for positive times at large frequencies:

$$\forall n, \quad \Xi(0) = \int_{n}^{n+1} |\widehat{T_x}(0,\xi)|^2 d\xi = 4 \sum_{k} (1 - e^{-\pi \alpha_k^2}) < 4\pi \sum_{k} \alpha_k^2 = \Xi(t).$$

In particular we can see $|\widehat{T}_x(t,\xi)|^2$ as an asymptotic energy density in phase space. Moreover, we observed an energy cascade in the following precise sense, even for solutions of (2) obtained as tangent vector of the binormal flow evolution of a polygonal line with only two corners from Theorem 2.2.

Theorem 3.2. (Energy density growth, [14]) Let $T_0 : \mathbb{R} \to \mathbb{S}^2$, constant on $(-\infty, -1), (-1, 1)$ and $(1, +\infty)$, with jumps of same angle $\theta \in (0, \pi)$ at $x \in \{-1, 1\}$. Then there exists a solution of the Schrödinger map (2) on $t \in (0, 1)$, with T_0 as trace at time t = 0, satisfying: i) there exists $C_{\theta} > 0$ such that:

$$\sup_{\xi \in B(\pm \frac{1}{t}, \sqrt{t})} |\widehat{T_x}(t, \xi)| = C_{\theta} |\log t|.$$

ii) for $\xi \notin B(\frac{1}{t}, \frac{3}{4t}) \cup B(-\frac{1}{t}, \frac{3}{4t})$ we have an upper-bound of $|\widehat{T_x}(t, \xi)|$ depending only on θ .

The growth is valid also for polygonal lines with many corners. This results confirms the numeric results obtained by de la Hoz and Vega in [50] for periodic piecewise constant data, in the case when T_0 is the tangent vector of a regular polygon. Last but not least, let us underline that the growth of T_x is in terms of the Fourier-Lebesgue norm $\mathcal{F}L^{\infty}$, which is critical with respect to scaling for the 1D cubic NLS equation (3) to which (2) is linked via Hasimoto's method.

Let us also note that in view of equations (26) satisfied by T and of the particular oscillatory ansatz of u in Theorem 2.1 from which T is constructed, we have:

$$T_t = -\frac{x}{2t} \Re(\overline{u} N) + \text{ Remainder term}, \quad T_x = \Re(\overline{u} N).$$

Passing in Fourier gives:

$$\partial_t \widehat{T_x} = \frac{\xi}{2t} \partial_{\xi} \widehat{T_x} + \text{Remainder term.}$$

This phenomena is reminiscent of the works of Apolinário et al. [2]-[3] where they proposed abstract linear equations that mimic the phenomenology of energy cascades when the external force is a statistically homogeneous and stationary stochastic process. Indeed, these equations have a transport rate in wavenumber space, independent of time and frequency (see (2.1) in [2]). Existence of energy cascades for linear systems have been also proved by Colin de Verdière and Saint-Raymond in [43].

3.0.1. Sketch of the proof of Theorem 3.2. We consider χ_0 a polygonal line with two corners, having T_0 as tangent vector. We can thus apply Theorem 2.2 to get a binormal flow solution χ with data χ_0 , and a Schrödinger map solution $T = \partial_x \chi$ with initial data T_0 . From the construction in the proof of Theorem 2.2 we also get that T can be completed to a frame (T, e_1, e_2) such that T and $N = e_1 + ie_2$ satisfy (26) for some solution $u_{\{\alpha_j\}}$ of type (7). We aim to show a log t growth of $\widehat{T_x}(t,\xi)$ for frequencies $\xi \in B(\frac{1}{t}, \sqrt{t})$. We have, by using $T_x = \Re(\overline{u}N)$ from (26):

$$\widehat{T_x}(t,\xi) = \int e^{ix\xi} \Re(\sum_k \overline{A_k(t)} \frac{e^{-i\frac{(x-k)^2}{4t}}}{\sqrt{t}} N(t,x)) dx$$
$$= \frac{e^{it\xi^2}}{2\sqrt{t}} \sum_k e^{ik\xi} \overline{A_k(t)} \int e^{-i\frac{(x-k-2t\xi)^2}{4t}} N(t,x) dx + \text{ a similar term.}$$

We remove $B(k+2t\xi,\sqrt{t})$ from the domain of integration, as its contribution is of order 1. Then on the remaining domain we use integrations by parts and the expression $N_x = -uT$

14

from (26) to reduce to:

$$i\sum_{k,j} \overline{A_k(t)} A_j(t) e^{-i\frac{k^2 - j^2}{4t}} \int e^{i\frac{x(k-j+2t\xi)}{2t}} \frac{1_{|_{x-k-2t\xi>\sqrt{t}}}}{x-k-2t\xi} T(t,x) \, dx.$$

If $|k - j + 2t\xi| > \sqrt{t}$ the corresponding term is proved to be bounded also by using several integrations by parts involving (26). We thus get the general estimate, that indicate where to look for a potential growth:

$$\left| \widehat{T_x}(t,\xi) - i \sum_{\substack{|k-j+2t\xi| < \sqrt{t} \\ 2t}} \overline{\alpha_k + R_k(t)}(\alpha_j + R_j(t)) z \right| \\ \times \int e^{i\frac{x(k-j+2t\xi)}{2t}} \left(\frac{1_{|x-k-2t\xi| < \sqrt{t}}}{x-k-2t\xi} - \frac{1_{|x-j+2t\xi| < \sqrt{t}}}{x-j+2t\xi} \right) T(t,x) dx \right| \le C,$$

where

$$z = e^{-i(|\alpha_k|^2 - |\alpha_j|^2)\log t} e^{-i\frac{k^2 - j^2}{4t}}.$$

Considering $\xi \in B(\frac{1}{t}, \sqrt{t})$ implies $2t\xi \in B(2, 2t\sqrt{t})$, so the summation condition implies:

$$|k - j + 2t\xi| < \sqrt{t} \Longrightarrow j = k + 2.$$

Thus, restricting to polygonal lines with finite number of corners we end up with an estimate that writes for the two corners case (when $\alpha_j = 0$ for $|j| \ge 2$), for $\xi \in B(\frac{1}{t}, \sqrt{t})$:

$$\left|\widehat{T_x}(t,\xi) - i\overline{\alpha_{-1}}\alpha_1 z \int \left(\frac{1_{|_{x-1} > \sqrt{t}}}{x-1} - \frac{1_{|_{x+1} > \sqrt{t}}}{x+1}\right) T(t,x) dx\right| \le C.$$

In particular we have

$$|\widehat{T_x}(t,\xi) - i\overline{\alpha_{-1}}\alpha_1 z \int_{\sqrt{t} < |x-1| < \frac{1}{3}\} \cup \{\sqrt{t} < |x+1| < \frac{1}{3}\}} (\frac{1}{x-1} - \frac{1}{x+1}) T(t,x) \, dx| \le C.$$

Next we use the following result on the convergence of the tangent vector from [12]:

$$|T(t,x) - T(0,x)| \le C(\|\{\alpha_j\}\|_{l^{1,1}})(1+|x|)\sqrt{t}\left(\frac{1}{d(x,\frac{1}{2}\mathbb{Z})} + \frac{1}{d(x,\mathbb{Z})}\right),$$

to get

$$|\widehat{T_x}(t,\xi) - i\overline{\alpha_{-1}}\alpha_1 z \int_{\{\sqrt{t} < |x-1| < \frac{1}{3}\} \cup \{\sqrt{t} < |x+1| < \frac{1}{3}\}} (\frac{1}{x-1} - \frac{1}{x+1}) T(0,x) \, dx| \le C.$$

As T(0, x) is piecewise constant direct integration yields a log t growth.

4. Talbot effects

In 1836 Talbot, inventor of the photography independently of Daguerre, observed a diffraction effect of light. By illuminating a grating that has equally spaced transparent and opaque slits, the pattern of the grating can be observed away from the grating at a distance called nowadays Talbot distance. Also, at rational fractions $\frac{p}{q}$ of the Talbot distance the image observed consists of q overlapping copies of the pattern, see Figure 5. Then the Talbot effect has been forgotten, rediscovered by Lord Rayleight in 1881 and forgotten again. Nowadays it is well-known and also known to be related to a phenomenon in quantum physics called quantum revivals, that concerns reconstruction after a finite time of a wave packet, see Figure 5. For more details on this topic see the expository article [23] of Berry, Marzoli and Schleich.



Figure 5. Left: the optical effect (space and distance axis). Right: A quantum carpet i.e. plots of probability density for the propagation of a Gaussian wavepacket in a 1D box of length 1 on a revival time interval rescaled to 1 (space and time axis), from [23].

This effect can be explained by the following fact. Adopting the quantum framework, the Dirac comb evolution through the linear Schrödinger equation, i.e. the fundamental solution of the periodic Schrödinger equation, is, by using Poisson summation formula:

$$e^{it\Delta}(\sum_{k\in\mathbb{Z}}\delta_k)(x) = e^{it\Delta}(\sum_{k\in\mathbb{Z}}e^{i2\pi kx}) = \sum_{k\in\mathbb{Z}}e^{-it(2\pi k)^2 + i2\pi kx},$$

so for $t = \frac{1}{2\pi} \frac{p}{q}$ we have

$$e^{it\Delta}(\sum_{k\in\mathbb{Z}}\delta_k)(x) = \frac{1}{q}\sum_{l\in\mathbb{Z}}\sum_{m=0}^{q-1}G(-p,m,q)\delta_{l+\frac{m}{q}},$$

where G(-p, m, q) stands for the Gauss sum

$$G(-p,m,q) = \sum_{l=0}^{q-1} e^{2\pi i \frac{-pl^2 + ml}{q}}$$

The linear and nonlinear Schrödinger evolution on the torus of functions with bounded variation was proved to present Talbot effect features by Berry [22], Berry and Klein [21], Oskolkov [100], Kapitanski and Rodnianski [86], Rodnianski [103], Erdogan and Tzirakis [55]). We note that also other dispersive equations enjoy Talbot effect, as shown for instance by Boulton, Farmakis, Pelloni and Smith [29].

In the Dirac deltas more singular setting on the torus, De la Hoz and Vega, by supposing uniqueness of solutions, founded in [49] a solution of the 1D cubic Schrödinger equation (3) similar to the fundamental solution above, thus exhibiting a Talbot effect. Then they constructed curves, which have as filament function this solution at rational times $t = \frac{1}{2\pi} \frac{p}{q}$, that are skew polygons with corners located at $\frac{1}{q}$ distances, thus again a Talbot effect.

In [12] we placed ourselves in the Dirac deltas singular setting on \mathbb{R} . We proved a Talbot effect for the 1D cubic Schrödinger equation (3) at this rough regularity. To do so we first noted a Talbot effect at the level of linear solutions, as follows.

Theorem 4.1. (A Talbot effect for linear evolutions of Dirac type, [12]) Let $p \in \mathbb{N}, \eta \in$ (0,1) and u_0 be such that \hat{u}_0 is 2π -periodic with \hat{u}_0 supported modulo 2π in $B(0,\eta\frac{\pi}{p})$. For $x \in \mathbb{R}$ we define:

$$\xi_x := \frac{\pi q}{p} \operatorname{dist}\left(x, \frac{1}{q}\mathbb{Z}\right) \in [0, \frac{\pi}{p})$$

Then, for $t_{pq} = \frac{1}{2\pi} \frac{p}{q}$ with q odd there exists $\theta_{x,p,q} \in \mathbb{R}$ such that:

$$e^{it_{pq}\Delta}u_0(x) = \frac{1}{\sqrt{q}}\,\hat{u}_0(\xi_x)\,e^{-it_{pq}\,\xi_x^2 + ix\,\xi_x + i\theta_{x,p,q}}$$

In particular, $e^{it_{pq}\Delta}u_0(x)$ has $\frac{1}{q}$ -periodic modulus and vanishes if $d(x, \frac{1}{q}\mathbb{Z}) > \frac{\eta}{q}$. Moreover, linear evolutions can concentrate near $\frac{1}{q}\mathbb{Z}$. More precisely, there is a family of initial data $u_0^{\lambda} = \sum_{k \in \mathbb{Z}} \alpha_k^{\lambda} \delta_k$ such that there exists C > 0 and:

$$\left|\frac{e^{it_{p,q}\Delta}u_0^{\lambda}(0)}{e^{it_{p,q}\Delta}\alpha_0^{\lambda}\delta_0(0)}\right| = C\frac{\sqrt{p}}{q}\lambda \xrightarrow{\lambda \to \infty} \infty.$$

At the end of this section we shall give the sketch of the proof. The concentration phenomena is obtained by taking a sequence of initial data $\{u_0^{\lambda}\}$ that focus in Fourier variable near 2π -integers:

$$\widehat{u_0^{\lambda}}_{|(-\pi,\pi)}(\xi) = \lambda \psi(\lambda\xi) = \sum_k \alpha_k^{\lambda} e^{ik\xi},$$

with ψ a bump function. This can be seen as a sequence approaching a Dirac comb. We note that the Dirac comb satisfies the hypothesis of the periodicity and localization in Fourier, since it equals to its Fourier transform. Therefore we recover the known Talbot effect explained above. However, this kind of data does not enter the nonlinear framework. Nevertheless, we obtain a Talbot effect also at the nonlinear level, at the same rough regularity, by using the solutions in Theorem 2.2 that have as a leading term a linear evolution of a sum of Dirac deltas.

Theorem 4.2. (A Talbot effect for nonlinear evolutions of Dirac type, [12]) Let $p \in \mathbb{N}, \eta \in (0,1), s > \frac{1}{2}$ and $\epsilon \in (0,1)$. Let u_0 such that \hat{u}_0 is a 2π -periodic, supported modulo 2π in $B(0, \eta \frac{\pi}{p})$, and having a Fourier coefficient sequence $\{\alpha_k\}$ with $\|\{\alpha_k\}\|_{l^{2,s}} = \epsilon$. Let u(t, x) be the solution of the 1D cubic Schrödinger equation (3) on (0,T) obtained in Theorem 2.1 from the sequence $\{\alpha_k\}$. Then for $t = \frac{1}{2\pi} \frac{p}{q}$ with 1 < q odd such that $\epsilon^2 \sqrt{q} \log q < \frac{1}{2}$ the function u(t, x) almost vanishes, in the sense:

$$|u(t_{p,q},x)| \le \epsilon, \quad \text{if } d(x,\frac{1}{q}\mathbb{Z}) > \frac{\eta}{q}.$$

Moreover, nonlinear solutions can concentrate near $\frac{1}{q}\mathbb{Z}$ in the sense that there is a family of sequences $\{\alpha_k^{\lambda}\}$ with $\|\{\alpha_k^{\lambda}\}\|_{l^{2,s}} \xrightarrow{\lambda \to \infty} 0$, such that the solutions u^{λ} of (3) obtained in Theorem 2.1 from the sequence $\{\alpha_k\}$ satisfy:

$$\left|\frac{u^{\lambda}(t_{p,q},0)}{e^{it_{p,q}\Delta}\alpha_0^{\lambda}\delta_0(0)}\right| \stackrel{\lambda \to \infty}{\to} \infty.$$

Let us notice that despite the smallness condition on $\{\alpha_k\}$, the corresponding solution u(t) from Theorem 2.1 is large for small times in L^{∞} and L_{loc}^1 , due to its $\frac{1}{\sqrt{t}}$ factor. Going either backward or forward in time this, combined with the Talbot effect in Theorem 4.2, gives a phenomenon of constructive/destructive interference. More precisely, in Theorem 5.1 in [16] we proved the existence of rational times $\frac{1}{2\pi}\frac{p}{q}$ and $\frac{1}{2\pi}\frac{\tilde{p}}{\tilde{q}}$ such that on the interval $\left[-\frac{1}{2\tilde{q}}, \frac{1}{2\tilde{q}}\right]$ we observe at $\frac{1}{2\pi}\frac{p}{q}$ almost-periodic small waves and at $\frac{1}{2\pi}\frac{p}{\tilde{q}}$ a localized large-amplitude structure emerges. On one hand this can be seen as a creation of rogue waves in the sense of the dispersive blow-up, i.e. L^{∞} -pointwise blow-up. This phenomena of dispersive blow-up was proved by Bona and Saut in [27] for the linear and the cubic Schrödinger equation on \mathbb{R} by using smooth data of type $e^{ix^2}(1+x^2)^{-m}$ with $m \in (\frac{1}{4}, \frac{1}{2}]$. On the other hand, by using the pseudo-conformal transformation, our rogue waves result transfers to the periodic cubic Schrödinger equation (19), equation that has a coefficient $\frac{1}{t}$ in front of the nonlinearity. In the context of the classical periodic cubic Schrödinger equation with random data recent progress in the rogue waves phenomena was done by Garrido, Grande, Kurianski and Staffilani in [62].

Finally, as a consequence of Theorem 4.2, and by using Theorem 2.2, we constructed in [12] families of binormal flow evolutions of polygonal lines that have at rational-type times $2\pi \frac{p}{q}$ curvature concentrating more and more at \mathbb{Z}/q and getting smaller and smaller in between such locations.

4.0.1. Sketch of the proof of Theorem 4.1. Since $\hat{u}_0(\xi)$ is periodic we write:

$$e^{it\Delta}u_0(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ix\xi} e^{-it\xi^2} \hat{u}_0(\xi) \, d\xi = \frac{1}{2\pi} \sum_{k \in \mathbb{Z}} \int_{2\pi k}^{2\pi (k+1)} e^{ix\xi - it\xi^2} \hat{u}_0(\xi) \, d\xi$$
$$= \frac{1}{2\pi} \int_0^{2\pi} \hat{u}_0(\xi) \sum_{k \in \mathbb{Z}} e^{ix(2\pi k + \xi) - it(2\pi k + \xi)^2} \, d\xi$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \hat{u}_0(\xi) e^{-it\xi^2 + ix\xi} \sum_{k \in \mathbb{Z}} e^{-it(2\pi k)^2 + i2\pi k(x-2t\xi)} d\xi.$$

For $t_{p,q} = \frac{1}{2\pi} \frac{p}{q}$ we get using the linear periodic Talbot effect:

$$e^{it_{p,q}\Delta}u_0(x) = \frac{1}{q} \int_0^{2\pi} \hat{u}_0(\xi) e^{-it_{p,q}\xi^2 + ix\xi} \sum_{l \in \mathbb{Z}} \sum_{m=0}^{q-1} G(-p,m,q) \delta(x - 2t_{p,q}\xi - l - \frac{m}{q}) \, d\xi.$$

Since \hat{u}_0 is located modulo 2π in $B(0, \eta \frac{\pi}{p})$ then

$$e^{it_{p,q}\Delta}u_0(x) = \frac{1}{\sqrt{q}}\,\hat{u}_0(\xi_x)\,e^{-it_{p,q}\,\xi_x^2 + ix\,\xi_x + i\theta_{x,p,q}},$$

for some $\theta_{x,p,q} \in \mathbb{R}$ and $\xi_x := \frac{\pi q}{p} d(x, \frac{1}{q}\mathbb{Z}) \in [0, \frac{\pi}{p})$. For proving the concentration effect of Theorem 4.1 we shall construct a family of sequences $\{\alpha_k^{\lambda}\}$ such that $\sum_{k \in \mathbb{Z}} \alpha_k^{\lambda} \delta_k$ concentrates in the Fourier variable near the integers. To this purpose we consider ψ a real bounded function with support in $[-\frac{1}{2}, \frac{1}{2}]$ and $\psi(0) = 1$. We define

$$f^{\lambda}(\xi) = \lambda \psi(\lambda \xi), \forall \xi \in [-\pi, \pi]$$

Thus we can decompose

$$f^{\lambda}(\xi) = \sum_{k \in \mathbb{Z}} \alpha_k^{\lambda} e^{ik\xi},$$

and consider

$$u_0^{\lambda} = \sum_{k \in \mathbb{Z}} \alpha_k^{\lambda} \delta_k.$$

In particular, on $[-\pi,\pi]$, we have $\widehat{u_0^{\lambda}} = f^{\lambda}$. Given $t_{p,q} = \frac{1}{2\pi} \frac{p}{q}$, for $\lambda > p$, the restriction of $\widehat{u_0^{\lambda}}$ to $[-\pi,\pi]$ has support included in $B(0,\eta\frac{\pi}{p})$ for a $\eta \in]0,1[$. We then get by the first part of the statement

$$e^{it_{p,q}\Delta}u_0^{\lambda}(0) = \frac{1}{\sqrt{q}}\,\widehat{u_0^{\lambda}}(0)\,e^{-it_{p,q}\,\xi_x^2 + ix\,\xi_x + i\theta_{x,p,q}},$$

 \mathbf{SO}

$$|e^{it_{p,q}\Delta}u_0^{\lambda}(0)| = \frac{1}{\sqrt{q}}|f^{\lambda}(0)| = \frac{1}{\sqrt{q}}\lambda\psi(0) = \frac{1}{\sqrt{q}}\lambda.$$

On the other hand, at $t_{p,q} = \frac{1}{2\pi} \frac{p}{q}$ we have

$$|e^{it_{p,q}\Delta}\alpha_0^{\lambda}\delta_0(0)| = \sqrt{\frac{4q}{p}} |\alpha_0^{\lambda}| = \sqrt{\frac{4q}{p}} \frac{1}{2\pi} \left| \int_{-\pi}^{\pi} f^{\lambda}(\xi) d\xi \right| = C(\psi) \sqrt{\frac{q}{p}} \lambda^{-1}.$$

Therefore,

$$\left|\frac{e^{it_{p,q}\Delta}u_0^{\lambda}(0)}{e^{it_{p,q}\Delta}\alpha_0^{\lambda}\delta_0(0)}\right| = \frac{\sqrt{p}}{C(\psi)q}\lambda \stackrel{\lambda \to \infty}{\longrightarrow} \infty,$$

and the proof of Theorem 4.1 is complete.

5. INTERMITTENCY AND MULTIFRACTALITY

As recalled, in numerical simulations of the binormal flow evolution of a M-regular polygon, the trajectories in time of corners $\chi_M(t,0)$ were showed to behave as Riemann's complex function

(9)
$$R(t) = \sum_{k \in \mathbb{Z}} \frac{e^{itk^2} - 1}{k^2},$$

when M tends to infinity (De la Hoz and Vega [49], De la Hoz, Kumar and Vega [48], see Figure 4). In [14] we showed this kind of behavior by considering sequences of polygonal lines. More precisely, let $n \in \mathbb{N}^*$, $\nu \in]0,1]$, $\theta > 0$. We consider polygonal lines $\chi_n(0)$ with finite but possibly many corners located at $j \in \mathbb{Z}$ with $|j| \leq n^{\nu}$ and curvature angles θ_n such that

$$\theta_n = \pi - \frac{\theta}{n} + o\left(\frac{1}{n}\right),$$

 $\chi_n(0,0) = 0_{\mathbb{R}^3}$, and $\chi_n(0)$ lying in the *xy*-plane and symmetric with respect to the yz-plane, see Figure 6 for examples.



 $\chi_n(0)$ planar approximation of a line



 $\chi_n(0)$ planar approximation of a (multi-)loop

Figure 6. Examples of curves χ_n .

These polygonal lines with finite number of corners enter the framework of Theorem 2.2, but apriori their time of existence depends on n. We first refined in [14] the general analysis of Theorem 2.2 to the specific case of $\chi_n(0)$. We show that there exists T > 0 and $n_0 \in \mathbb{N}^*$, both depending only on θ , such that for all $n \ge n_0$ there exist smooth solutions $\chi_n(t)$ of the binormal flow on $(-T, T) \setminus \{0\}$, that are weak solutions on (-T, T), and at time t = 0 converge pointwise to $\chi_n(0)$. Then we proved the following result.

Theorem 5.1. (A multifractal trajectory, [14]) We have the following description of the trajectory of the corner $\chi_n(t,0)$, uniformly on (0,T):

(10)
$$n(\chi_n(t,0) - \chi_n(0,0)) - \theta(0, \Re(R(t)), \Im(R(t))) \stackrel{n \to \infty}{\longrightarrow} 0.$$

This theorem gives a non-obvious non-linear geometric interpretation of Riemann's function. It is valid also for $\chi_n(0)$ with same torsion angle ω_0 at all corners, and then the limit is given by the following Riemann-type function:

$$\Re_{\omega_0}(t) = \sum_{k \in \mathbb{Z}} \frac{e^{it(k-\omega_0)^2} - 1}{(k-\omega_0)^2}$$

We shall give the sketch of the proof of Theorem 5.1 in §5.0.1. For rational-type torsion ω_0 we also computed the spectrum of singularities of \Re_{ω_0} , showed that it is the same as the

one of Riemann's function, and proved that \Re_{ω_0} satisfies the Frisch-Parisi formalism⁶ and intermittency, notions that we recall in the following (see also the book of Frisch [60]).

We recall that Riemann's function $\sum_{k \in \mathbb{Z}^*} \frac{\sin(tk^2)}{k^2}$, and implicitly its complex versions $\sum_{k \in \mathbb{Z}^*} \frac{e^{itk^2}}{k^2}$ and R defined in (9), was studied by Jaffard in 1996 ([82]) from the point of view of multifractal analysis, first by computing its spectrum of singularities. The spectrum of singularities of a function $f : [0, 1] \to \mathbb{R}$ is given by the Hausdorff dimension of non-empty iso-Hölder sets,

$$d_f(\alpha) = \dim_{\mathcal{H}} \{ t \in [0, 1], \sup\{\beta, f \in \mathcal{C}^{\beta}(t)\} = \alpha \},\$$

with the convention $d_f(\alpha) = -\infty$ if the local α -Hölder regularity is not reached. Jaffard (see also Brouke and Vindas ([34]) for a recent different proof of computing pointwise the Hölder regularity) showed that

$$d_R(\alpha) = \begin{cases} 4\alpha - 2, \alpha \in [\frac{1}{2}, \frac{3}{4}], \\ 0, \alpha = \frac{3}{2}, \\ -\infty, \text{ otherwise.} \end{cases}$$

Moreover, he proved that it satisfies the multifractal formalism of Frisch-Parisi, a model for predicting the structure function exponents in turbulent flows, that was motivated by the experiments in Modane of Anselmet and all in 1984 ([1]) showing deviations from Kolmogorov 41 theory:

$$d_R(\alpha) = \inf_p(\alpha p - \eta_R(p) + 1), \quad \forall \alpha \in \left[\frac{1}{2}, \frac{3}{4}\right],$$

where η_R is determined in terms of Besov spaces:

$$\eta_R(p) := \sup\{s, R \in B_{p,\infty}^{\overline{p}}\}.$$

We note that recently Barral and Seuret ([17]-[18]) proved the validity of the multifractal formalism generically in Besov spaces. Again having in mind turbulent dynamics, Boritchev, Eceizabarrena and Da Rocha ([28]) proved that R is intermittent in small scales by showing that the flatness satisfies:

(11)
$$F_R(N) := \frac{\|P_{\geq N}R\|_4^4}{\|P_{\geq N}R\|_2^4} \xrightarrow{N \to \infty} \infty,$$

wher $P_{>N}$ denotes the high-pass filter of Fourier modes larger than N.

In collaboration with Daniel Eccizabarrena and Andrea Nahmod we extended in [7] the Theorem 5.1 to all locations $x_0 \in \mathbb{R}$, showing that the convergence (10) holds with

$$n(\chi_n(t,x_0) - \chi_n(0,x_0)) - \theta(0, \Re(R_{x_0}(t)), \Im(R_{x_0}(t))) \xrightarrow{n \to \infty} 0,$$

⁶The Frisch-Parisi multifractal formalism was originally proposed for the velocity in an Eulerian setting, but it can be equally proposed in the Lagrangian setting, to which Riemann's function is in our context more related to since it represents a time trajectory. See the work of Chevillard et al. [41] for a discussion on the differences between these two frameworks.

where

$$R_{x_0}(t) = \sum_{j \in \mathbb{Z}} \frac{e^{itj^2} - 1}{j^2} e^{ijx_0}.$$

In Figure 7 several such trajectories are represented, among which, for $x_0 = 0$, the one of the classical Riemann's complex function (9).



Figure 7. Trajectories of the limit curve evolution R_{x_0} for $t \in [0, \frac{1}{2\pi}]$, at $x_0 = 0$ (black), 0.1 (blue), 0.2 (green), 0.3 (cyan), 0.4 (red), 0.5 (magenta).

Many multifractal analysis studies have been done for various types of modifications of Riemann's function. For instance Chamizo and Ubis ([40]) and Seuret and Ubis ([105]) studied $\sum_n \frac{e^{iP(n)t}}{n^{\alpha}}$ for P polynomial. Also, Kapitanski and Rodnianski studied in [86] the behavior in space of $u_{t_0}(x) = \sum_n e^{in^2t_0 + inx}$ for t_0 fixed.

In [7] we have studied the function R_{x_0} , that is a natural extension of Riemann's function, linked with the fundamental solution of the periodic Schrödinger equation, and that came out in the study of trajectories of binormal flow evolutions, having in mind that the binormal flow is a model for non-smooth fluids presenting vortex filaments.

Theorem 5.2. (General multifractal study, [7]) For $x_0 \in \mathbb{R}$ the function R_{x_0} is multifractal, with infinitely many local Hölder exponents.

If $x_0 \in \mathbb{Q}$ then R_{x_0} has the same spectrum of singularities as Riemann's function:

$$d_{R_{x_0}}(\alpha) = \begin{cases} 4\alpha - 2, \alpha \in [\frac{1}{2}, \frac{3}{4}], \\ 0, \quad \alpha = \frac{3}{2}, \\ -\infty, \text{ otherwise,} \end{cases}$$

it satisfies the multifractal formalism of Frisch-Parisi, and it is intermittent in small scales.

The spectrum proof starts as in [82] and then follows the approach in [40] but ends up with new Diophantine sets that approximate the iso-Hölder sets. We measure these new sets using Duffin-Schaeffer theorem from 1941 finalized by Koukoulopoulos and Maynard in 2020 ([54]-[93]) and the Mass Transference Principle proved by Beresnevich and Velani in 2006 ([20]). We shall give a sketch of the computation of the spectrum in §5.0.2.

To show that R_{x_0} satisfies Frish-Parisi multifractal formalism we compute $\eta_{R_{x_0}}(p)$ by obtaining the L^p norms of dyadic blocs of R'_{x_0} , based on L^{∞} estimates of partial sums of the

22

fundamental solution of periodic 1D Schrödinger equation, i.e. Vinogradov trigonometrical series, near rational time and space locations, together with counting techniques.

5.0.1. Sketch of the proof of Theorem 5.1. The solutions of the binormal flow χ_n were constructed by using Hasimoto's method from the following solutions in Theorem 2.1:

$$u_{\{\alpha_{n,k}\}}(t,x) = \sum_{k \in \mathbb{Z}} e^{-i(|\alpha_{n,k}|^2 - 2\sum_{j \in \mathbb{Z}} |\alpha_{n,j}|^2) \log t} (\alpha_{n,k} + R_{n,k}(t)) e^{it\Delta} \delta_k(x),$$

with

$$\sup_{0 < t < T} t^{-\gamma} \| \{ R_{n,k}(t) \} \|_{l^{2,s}} + t \| \{ \partial_t R_{n,k}(t) \} \|_{l^{2,s}} < C,$$

and $s > \frac{1}{2}, 0 < \gamma < 1$. In view of the particular shape of $\chi_n(0)$ we have $\alpha_{n,j} = c_n \approx \frac{\theta}{n}$. Using the equations (26) we can compute the evolution of one arclength location:

$$\chi_n(t,0) - \chi_n(0,0) = \int_0^t \Im(\overline{u_{\{\alpha_{n,k}\}}} N_n(\tau,0)) \, d\tau$$
$$= \Im \int_0^t \sum_k (\overline{\alpha_{n,k} + R_{n,k}(\tau)}) \frac{e^{-i\frac{k^2}{4\tau}}}{\sqrt{\tau}} e^{-i(|\alpha_{n,k}|^2 - 2\sum_{j \in \mathbb{Z}} |\alpha_{n,j}|^2) \log t} N_n(\tau,0) \, d\tau.$$

The $R_{n,k}(\tau)$ constribution is $o(\frac{1}{n})$ due to the decay in time of its l^1 norm. In [12] we proved that a modulation of the normal vector $N_n(\tau, 0)$ has a limit $\tilde{N}_n(0, 0) \in \mathbb{S}^2 + i\mathbb{S}^2$ at $\tau = 0$, that allows us to estimate:

$$\chi_n(t,0) = \Im(e^{-i\sum_{1\le |j|\le n^{\nu}} |\alpha_{n,j}|^2 \log |k|} \tilde{N}_n(0,0) \int_0^t \sum_{|k|\le n^{\nu}} \overline{\alpha_{n,k}} \frac{e^{-i\frac{k^2}{4\tau}}}{\sqrt{\tau}} d\tau) + o(\frac{1}{n}).$$

Recalling that $\alpha_{n,k} = c_n \approx \frac{\theta}{n}$ and $\nu \in]0,1]$ we obtain:

$$\chi_n(t,0) = \frac{\theta}{n} \Im(\tilde{N}_n(0,0) \int_0^t \sum_{|k| \le n^{\nu}} \frac{e^{-i\frac{k^2}{4\tau}}}{\sqrt{\tau}} d\tau) + o(\frac{1}{n}).$$

On one hand, we prove convergence of the modulated normal vectors at (t, x) = (0, 0):

$$\lim_{n \to \infty} \tilde{N}_n(0,0) = (0, \frac{1-i}{\sqrt{2}}, \frac{-1-i}{\sqrt{2}}).$$

On the other hand, the summation in k can be taken over the whole set \mathbb{Z} and thanks to Poisson's summation formula $\sum_{k \in \mathbb{Z}} f(k) = \sum_{j \in \mathbb{Z}} \hat{f}(2\pi j)$ we have:

$$\sum_{k \in \mathbb{Z}} e^{i4\pi^2 tk^2} = \sum_{k \in \mathbb{Z}} \int e^{-i2\pi xk + i4\pi^2 tx^2} dx = \frac{1}{\sqrt{4\pi^2 t}} \sum_{j \in \mathbb{Z}} \int e^{-iy\frac{j}{\sqrt{t}} + iy^2} dy$$
$$= \frac{1}{2\pi\sqrt{t}} \sum_{j \in \mathbb{Z}} \widehat{e^{i\cdot^2}}(\frac{j}{\sqrt{t}}) = \frac{e^{i\frac{\pi}{4}}}{2\sqrt{\pi}\sqrt{t}} \sum_{j \in \mathbb{Z}} e^{-i\frac{j^2}{4t}},$$

Thus, neglecting again the 2π -factors for simplifying the presentation, we obtain uniformly on (0, T):

$$n \chi_n(t,0) - \theta(0, \Re(R(t)), \Im(R(t))) \xrightarrow{n \to \infty} 0$$

5.0.2. Sketch of the proof of Theorem 5.2 for $\alpha \in (\frac{1}{2}, \frac{3}{4}]$. We start by analyzing the variation of R_{x_0} at rationals. Splitting the sum in n modulo q and using Poisson summation formula we get:

$$R_{x_0}\left(\frac{p}{q}+h\right) - R_{x_0}\left(\frac{p}{q}\right) = -ih + \sum_{n \in \mathbb{Z}} e^{in^2 \frac{p}{q}} \frac{e^{in^2 h} - 1}{n^2} e^{inx_0}$$
$$= -ih + \frac{\sqrt{h}}{q} \sum_{m \in \mathbb{Z}} G(p, m, q) F\left(\frac{x_0 - \frac{m}{q}}{\sqrt{h}}\right),$$

with $G(p, s, q) = \sum_{r=0}^{q-1} e^{i\frac{p}{q}r^2 + i\frac{s}{q}r}$ i.e. Gauss sums, of size \sqrt{q} except if q even, q/2, s of different parity, and $F(x) = \mathcal{F}(\frac{e^{i\xi^2}-1}{\xi^2}) = O(\frac{1}{x^2}), F(0) \neq 0$. Thus the leading term is given by $m_{x_0,q}$ s.t. $x_0 - \frac{m_{x_0,q}}{q} = d(x_0, \frac{\mathbb{Z}}{q})$:

(12)
$$R_{x_0}\left(\frac{p}{q}+h\right) - R_{x_0}\left(\frac{p}{q}\right) = \frac{\sqrt{h}}{q}G(p, m_{x_0, q}, q)F\left(\frac{d(x_0, \frac{\mathbb{Z}}{q})}{\sqrt{h}}\right) - ih + O(\min\{\sqrt{q}h, q^{\frac{3}{2}}h^{\frac{3}{2}}\}).$$

Therefore, we obtain $R_{x_0} \in C^{\frac{1}{2}}(\frac{p}{q})$ if $G(p, m_{x_0,q}, q) \neq 0$ and $d(x_0, \frac{\mathbb{Z}}{q}) = 0$. These conditions are satisfied for rationals $x_0 = \frac{P}{Q}$ together with $q \in 4Q\mathbb{Z}$:

(13)
$$R_{\frac{P}{Q}} \in \mathcal{C}^{\frac{1}{2}}(\frac{p}{q}).$$

Now we shall look for upper and lower bounds for Hölder regularity at irrationals. To do so we have to recall the notion of exponent of irrationality of $t \notin \mathbb{Q}$:

$$\mu(t) = \sup\{\nu, t \in \mathbf{A}_{\nu}\},\$$

where

$$\mathbf{A}_{\nu} = \{ t \notin \mathbb{Q}, |t - \frac{p}{q}| \le \frac{1}{q^{\nu}} \text{ for infinitely many coprime pairs } (p,q) \in \mathbb{N} \times \mathbb{N} \}.$$

Also, we recall that the approximation by continuous fractions of t satisfies

$$|t - \frac{p_n}{q_n}| = \frac{1}{q_n^{\mu_n}} \le \frac{1}{q_{n+1}q_n} \text{ and } \mu(t) = \limsup_{n \to \infty} \mu_n.$$

Thus, for all h small there exists n such that

$$t - \frac{p_n}{q_n} | \le h \le |t - \frac{p_{n-1}}{q_{n-1}}|.$$

For estimating the variations at t we rely on variations at the rationals $\frac{p_n}{q_n}$, and use (12) to get:

$$|R_{x_0}(t+h) - R_{x_0}(t)| \le |R_{x_0}(\frac{p_n}{q_n} + (t - \frac{p_n}{q_n} + h)) - R_{x_0}(\frac{p_n}{q_n})| + |R_{x_0}(\frac{p_n}{q_n}) - R_{x_0}(\frac{p_n}{q_n} + (t - \frac{p_n}{q_n}))| \le |R_{x_0}(t+h) - R_{x_0}(t+h) - R_{x_0}(t+h)$$

$$\lesssim \frac{\sqrt{h}}{\sqrt{q_n}} + h + \min\{\sqrt{q_n}h, q_n^{\frac{3}{2}}h^{\frac{3}{2}}\} \lesssim h^{\frac{1}{2} + \frac{1}{2\mu_n}} + h^{\frac{1}{2} + \frac{1}{2\mu_{n-1}}} \lesssim h^{\frac{1}{2} + \frac{1}{2\mu} - \delta}, \quad \forall \delta > 0$$

Therefore we have a lower bound for the Hölder exponent of R_{x_0} at t:

(14)
$$\alpha_{R_{x_0}}(t) \ge \frac{1}{2} + \frac{1}{2\mu(t)}$$

To get an upper bound for Hölder regularity at irrational t we consider sets of irrationals well approximated by rationals $\frac{p_n}{q_n}, q_n \in 4Q\mathbb{Z}$, where the Hölder regularity is $\mathcal{C}^{\frac{1}{2}}$ by (13):

$$\mathbf{A}_{\mu,\mathbf{Q}} = \{t \notin \mathbb{Q}, |t - \frac{p}{q}| \le \frac{1}{q^{\mu}} \text{ for infinitely many coprime pairs } (p,q) \in \mathbb{N} \times 4Q\mathbb{N}\}.$$

We note that in particular if $t \in \mathbf{A}_{\mu,\mathbf{Q}}$ then $\mu \leq \mu(t)$. For $t \in \mathbf{A}_{\mu,\mathbf{Q}}$ we consider $(p_n, q_n) \in \mathbb{N} \times 4Q\mathbb{N}$ from the definition of $\mathbf{A}_{\mu,\mathbf{Q}}$. We define h_n, ν_n such that

$$h_n = t - \frac{p_n}{q_n}, \quad \frac{1}{q_n^{\nu_n}} = |h_n|.$$

In particular we have $\nu_n \ge \mu$. We use (12)-(13) to get the lower estimate:

$$|R_{x_0}(t+h_n) - R_{x_0}(t)| = |R_{x_0}(\frac{p_n}{q_n}) - R_{x_0}(\frac{p_n}{q_n} + h_n)| \gtrsim \frac{\sqrt{h_n}}{\sqrt{q_n}} = h_n^{\frac{1}{2} + \frac{1}{2\nu_n}} \ge h_n^{\frac{1}{2} + \frac{1}{2\mu}}.$$

Hence we have obtained an upper-bound for the Hölder regularity at $t \in \mathbf{A}_{\mu,\mathbf{Q}}$ that together with (14) gives the constraint:

(15)
$$\frac{1}{2} + \frac{1}{2\mu} \ge \alpha_{R_{x_0}}(t) \ge \frac{1}{2} + \frac{1}{2\mu(t)}, \quad \forall t \in \mathbf{A}_{\mu,\mathbf{Q}}$$

Now we shall start approximating the iso-Hölder sets. We remove from $\mathbf{A}_{\mu,\mathbf{Q}}$ the points that might have $\mu(t) > \mu$ by introducing the sets

$$\mathbf{B}_{\mu,\mathbf{Q}} = \mathbf{A}_{\mu,\mathbf{Q}} \setminus \Big(\cup_{\epsilon>0} \mathbf{A}_{\mu+\epsilon} \Big).$$

Then from $\mathbf{B}_{\mu,\mathbf{Q}} \subset \mathbf{A}_{\mu} \setminus \left(\cup_{\epsilon>0} \mathbf{A}_{\mu+\epsilon} \right)$ and the definition of $\mu(t)$ we get

(16)
$$\mu(t) = \mu, \quad \forall t \in \mathbf{B}_{\mu,\mathbf{Q}}$$

Therefore, we obtain the following approximation of the iso-Hölder sets:

$$\mathbf{B}_{\mu,\mathbf{Q}} \subset \{t, \alpha_{R_{x_0}}(t) = \frac{1}{2} + \frac{1}{2\mu}\} \subset \mathbf{A}_{\mu-\epsilon}, \forall \epsilon > 0.$$

Indeed, the first inclusion follows then by (15)-(16). From the general lower bound (15) it follows that a point t such that $\alpha_{R_{x_0}}(t) = \frac{1}{2} + \frac{1}{2\mu}$ satisfies $\mu \leq \mu(t)$. Thus $\mu - \epsilon < \mu(t) = \sup\{\nu, t \in \mathbf{A}_{\nu}\}$ and the second inclusion follows from the definition of \mathbf{A}_{ν} .

To prove the spectrum of singularities in Theorem 5.2 we need to show that for $\mu \ge 2$:

$$\dim_{\mathcal{H}} \{t, \alpha_{R_{x_0}}(t) = \frac{1}{2} + \frac{1}{2\mu} \} = \frac{2}{\mu}.$$

Since

$$\mathbf{B}_{\mu,\mathbf{Q}} = \mathbf{A}_{\mu,\mathbf{Q}} \setminus \left(\cup_n \mathbf{A}_{\mu+\frac{1}{n}} \right) \subset \{t, \alpha_{Rx_0}(t) = \frac{1}{2} + \frac{1}{2\mu}\} \subset \mathbf{A}_{\mu-\epsilon}$$

and since Jarník-Besicovitch theorem proved in 1931-1934 ([83]-[24]) states that $\dim_{\mathcal{H}} \mathbf{A}_{\mu} = \frac{2}{\mu}$, it is enough to show that:

$$\dim_{\mathcal{H}} \mathbf{B}_{\mu,\mathbf{Q}} \geq \frac{2}{\mu}$$

Moreover, as

$$\mathcal{H}^{\frac{2}{\mu}}(\mathbf{B}_{\mu,\mathbf{Q}}) = \mathcal{H}^{\frac{2}{\mu}}(\mathbf{A}_{\mu,\mathbf{Q}}) - \lim_{n \to \infty} \mathcal{H}^{\frac{2}{\mu}}(\mathbf{A}_{\mu+\frac{1}{\mathbf{n}}}) = \mathcal{H}^{\frac{2}{\mu}}(\mathbf{A}_{\mu,\mathbf{Q}}),$$

it is enough to prove $\mathcal{H}^{\frac{2}{\mu}}(\mathbf{A}_{\mu,\mathbf{Q}}) > 0$. Thus, to finish the proof it is enough to show

$$\mathcal{H}^{\frac{2}{\mu}}(\mathbf{A}_{\mu,\mathbf{Q}}) = +\infty$$

To do so we start with a Lebesgue measure information. We use Duffin-Schaeffer theorem ([54]-[93]) which states that if

$$\mathbf{A}_{\psi} = \{t, |t - \frac{p}{q}| \le \psi(q) \text{ for infinitely many coprime pairs } (p, q) \in \mathbb{N} \times \mathbb{N}\},\$$

and $\sum_{q\geq 1} \psi(q)\varphi(q) = \infty$, where φ is Euler's totient function $(\varphi(q) := \#\{1 \leq m \leq q, (m, q) = 1\})$, then $|\mathbf{A}_{\psi}| = 1$. We show that $\psi(q) := \frac{\mathbb{I}_{4Q\mathbb{N}}(q)}{q^2}$ satisfies the hypothesis, so we get

$$|\mathbf{A}_{2,\mathbf{Q}}| = |\mathbf{A}_{\psi}| = 1.$$

Then we use transform this Lebesgue information into a Hausdorff information by the Mass Transference Principle in [20], which states the following. Let $B_n(x_n, r_n)$ a sequence of balls in [0, 1] with $r_n \to 0$ and $\alpha < d$. If

$$|\limsup_{n} B_n(x_n, r_n^{\alpha})| = 1,$$

then

$$\dim_{\mathcal{H}} \limsup_{n} B_n(x_n, r_n) \ge \alpha, \quad \mathcal{H}^{\alpha}(\limsup_{n} B_n(x_n, r_n)) = +\infty$$

Since we have obtained in (17)

$$1 = |\mathbf{A}_{2,\mathbf{Q}}| = \Big|\limsup_{q} \bigcup_{p \le q, (p,q)=1} B\Big(\frac{p}{q}, \frac{\mathbb{I}_{4Q\mathbb{N}}(q)}{q^2}\Big)\Big|,$$

we apply the above result with $\alpha = \frac{2}{\mu}$ and radii $\frac{\mathbb{I}_{4Q\mathbb{N}}(q)}{q^2}$ to get:

$$+\infty = \mathcal{H}^{\frac{2}{\mu}}\left(\limsup_{q} \bigcup_{p \le q, (p,q)=1} B\left(\frac{p}{q}, \frac{\mathbb{I}_{4Q\mathbb{N}}(q)}{q^{\mu}}\right)\right) = \mathcal{H}^{\frac{2}{\mu}}(\mathbf{A}_{\mu,\mathbf{Q}}).$$

26

6. Well-posedness, phase blow-up and unique continuation at critical regularity

As we recalled in §2 the only known wellposedness results for 1D cubic Schrödinger equation are in the subscritical spaces with respect to the scaling, in H^s for $s > -\frac{1}{2}$ on the Sobolev scale, and $\mathcal{F}L^p$ for $p < +\infty$ ([80],[72]).

In collaboration with Renato Lucà and Nikolay Tzvetkov [8] we displayed a subset in the supercritical spaces H^s for $s < -\frac{1}{2}$ and in the critical space $\mathcal{F}L^{\infty}$ in which we obtained a well-posedness result in the following sense.

Theorem 6.1. (A well-posed critical subset for 1D cubic NLS and phase blow-up, [8]) Let r > 0 and $u_1(x)$ satisfying the following periodicity property at the level of the Fourier transform on \mathbb{R} :

(18)
$$\widehat{e^{-i\Delta u_1}}(\xi) = e^{i\xi^2}\widehat{u}_1(\xi) \in H^r(0, 2\pi).$$

Then there exists a unique solution of (3) on (0,1] such that $u(1) = u_1$ and

 $\widehat{e^{-it\Delta u}(t)}(\xi) = e^{it\xi^2}\widehat{u}(t,\xi) \in H^r(0,2\pi), \forall t \in (0,1].$

Moreover, by denoting $\{A_k(t)\}_{k\in\mathbb{Z}}$ the Fourier coefficients of $e^{it\xi^2}\hat{u}(t,\xi)$, we have the existence of a sequence $\{\alpha_k\}_{k\in\mathbb{Z}} \in l^{2,r}$ such that

$$|A_k(t) - e^{i(|\alpha_k|^2 - 2\|e^{i\xi^2}\widehat{u_1}(\xi)\|_{L^2(0,2\pi)}^2)\log t} \alpha_k| \le C(\|e^{i\xi^2}\widehat{u_1}(\xi)\|_{H^r(0,2\pi)})t, \quad \forall k \in \mathbb{Z}, t \in (0,1).$$

In particular, we have a blow-up in the sense that u(t) falls out from the periodicity functional framework at t = 0, as the Fourier coefficients $A_k(t)$ don't have a limit, due to the logarithmic phase loss.

The dependence with respect to the initial data is continuous in the following sense. Let $\{u_{1,n}\}_{n\in\mathbb{N}}$ be a sequence of initial data satisfying (18) such that $e^{i\xi^2}\widehat{u_{1,n}}(\xi)$ converges to $e^{i\xi^2}\widehat{u}(1,\xi)$ in $H^r(0,2\pi)$. Then for every $t \in (0,1]$ the sequence $e^{it\xi^2}\widehat{u_n}(t,\xi)$ converges to $e^{it\xi^2}\widehat{u}(t,\xi)$ in $H^r(0,2\pi)$, where $u_n(t)$ is the solution of (3) with data $u_{1,n}$ at t = 1. We have that $u \in C((0,1]; \mathcal{S}'(\mathbb{R}))$ but it blows-up at zero in the sense that $\lim_{t\to 0^+} u(t,\cdot)$ does not exist in $\mathcal{S}'(\mathbb{R})$.

Let us first note that for free Schrödinger solutions, if $e^{i\xi^2} \hat{u}(1,\xi)$ is 2π -periodic then $e^{it\xi^2} \hat{u}(t,\xi)$ is also 2π -periodic as

$$e^{it\xi^2}\widehat{u}(t,\xi) = e^{i\xi^2}\widehat{u}(1,\xi).$$

For the 1D cubic Schrödinger equation (3) the evolution law of $\omega(t,\xi) := e^{it\xi^2} \hat{u}(t,\xi)$ is:

$$\omega_t(\eta, t) = \frac{i}{8\pi^3} \int e^{it2(\xi_1 - \xi_2)(\xi_1 - \eta)} \omega(t, \xi_1) \bar{\omega}(t, \xi_2) \omega(t, \eta - \xi_1 + \xi_2) \, d\xi_1 d\xi_2,$$

thus compatible with periodicity of ω ; the solutions we construct in Theorem 6.1 are in this framework.

Phase blow up phenomena were encountered for the Schrödinger equation since the works of Merle in the 90s ([97], see also [98]). Here we shall prove that this loss of phase

does not affect the associated solutions of the binormal flow via Hasimoto's transform, and that they can be uniquely continued after the singularity time t = 0.

Let us now note that the periodicity condition in the theorem translates as follows:

$$\hat{u}(t,\xi) = \sum_{k \in \mathbb{Z}} A_k(t) e^{-it\xi^2} e^{ik\xi} = \sum_{k \in \mathbb{Z}} A_k(t) e^{it\Delta \delta_k},$$

so u(t, x) has the ansatz (6):

$$u(t,x) = \sum_{k} A_k(t) e^{it\Delta} \delta_k = \sum_{k} A_k(t) \frac{e^{i\frac{(x-k)^2}{4t}}}{\sqrt{t}}.$$

In particular u(t) belongs to the supercritical spaces $H^s(\mathbb{R})$ for $s < -\frac{1}{2}$, but not to the subcritical ones, and belongs also to the critical space $\mathcal{F}L^{\infty}(\mathbb{R})$ provided that $r > \frac{1}{2}$, but not to the subcritical ones. Also, the data in Theorem 6.1 is simply

$$u(1,x) = e^{ix^2} f(x),$$

with f a 4π -periodic function in $H^r(0, 4\pi)$ with r > 0.

Examples of solutions as in Theorem 6.1 were given in Theorem 2.1. We recall that by using the pseudo-conformal transformation

$$v(\tau, y) := \frac{e^{i\frac{y^2}{4\tau}}}{\sqrt{\tau}} \,\overline{u}(\frac{1}{\tau}, \frac{y}{\tau}),$$

equation (3) is transformed into

(19)
$$iv_t + v_{xx} + \frac{1}{t}|v|^2 v = 0$$

Moreover, the ansatz (6) translates into simply being in the periodic setting of (19) with data in $H^r(0, 2\pi)$ at time t = 1. Thus the local in time well-posedness in Theorem 6.1 is obtained by Bourgain's approach classical for the periodic 1D cubic Schrödinger equation, as the factor $\frac{1}{t}$ in (19) is harmless for this argument. The delicate point is to understand the asymptotics of the solution v(t) when t goes to infinity. In particular the result of Theorem 2.1 is the existence of wave operators for the above periodic 1-D cubic NLS equation with time-variable coefficient (19). In turn Theorem 6.1 corresponds to the much more delicate result of asymptotic completeness for (19), which translates to a blow-up for (3). We shall briefly sketch the proof at the end of this section.

With the help of Hasimoto's construction we obtained as a consequence of Theorem 6.1 and of the analysis in [12] the following result.

Theorem 6.2. (Criterium for generating binormal flow singularities, [8]) Let $\chi_1(x)$ be a curve with filament function $u_1(x)$ such that $e^{i\xi^2}\hat{u}_1(\xi) \in H^{\frac{3}{2}+}(0,2\pi)$. If its Frenet frame is well-defined, this condition means the curvature and torsion satisfy:

$$c_1(x) = g(x), \quad \tau_1(x) = \frac{x}{2} + h(x), \quad g, h \in H_{per}^{\frac{3}{2}^+}(0, 2\pi).$$

Then, there exists $\chi(t)$, with $\chi(1) = \chi_1$, a strong binormal flow solution on \mathbb{R}^* and weak solution on \mathbb{R} which generates several corner-singularities at t = 0. Uniqueness holds in the class of curves with filament functions in the functional frame of Theorem 6.1.

We note that the self-similar solutions χ_a of the binormal flow discussed in §1.3 enter the framework of this theorem with g = a and h = 0.

The proof of Theorem 6.2 relies in considering first the solutions in Theorem 6.1 with initial data u_1 at t = 1. We obtain a control in time of the $l^{2,\frac{3}{2}^+}$ -norms of the sequences $\{A_k(t)\}_{k\in\mathbb{Z}}$ since $e^{i\xi^2}\widehat{u_1}(\xi) \in H^{\frac{3}{2}^+}$, and also persistance of this regularity in the limit sequence $\{\alpha_k\}_{k\in\mathbb{Z}}$. Then we use Theorem 2.2 for this sequence $\{\alpha_k\}_{k\in\mathbb{Z}} \in l^{2,\frac{3}{2}^+}$ to obtain the solutions of the binormal flow $\chi(t)$ in Theorem 6.2.

6.0.1. Sketch of the proof of Theorem 6.1. Applying the pseudo-conformal transformation to (3) and to the ansatz (6) we look for periodic solutions on $[1, \infty)$ of (19) with data in $H^r(0, 2\pi)$ at t = 1. Thus the modulated Fourier coefficients:

$$B_k(t) := \mathcal{F}(e^{-it\Delta}v(t))(k),$$

which identifies to $\overline{A_k}(\frac{1}{t})$, must solve:

$$i\partial_t B_k(t) = \frac{1}{t} \sum_{k-j_1+j_2-j_3=0} e^{-it(k^2-j_1^2+j_2^2-j_3^2)} B_{j_1}(t) \overline{B_{j_2}(t)} B_{j_3}(t).$$

Let $b > \frac{1}{2}$. Following Bourgain's classical approach for

$$e^{-it\Delta}v(t)(x) = \sum_{k} B_k(t)e^{ikx},$$

we obtain H^r solutions v(t) of (19) on $[1, \infty)$ satisfying for any $\nu \in \mathbb{N}^*, t \in (\nu, \nu + 1)$:

$$\|\{B_k(t)\}\|_{l^{2,r}} = \|v(t)\|_{H^r} \le C \|v\|_{X_{\nu}^{r,b}} \le C \|v(\nu)\|_{H^r} = C \|\{B_k(\nu)\}\|_{l^{2,r}},$$

where the $X_{\nu}^{r,b}$ -norm is defined via a smooth localization at ν of B:

$$\|v\|_{X^{r,b}_{\nu}} := \left(\int \sum_{k} \langle k \rangle^{2r} \langle \lambda \rangle^{2b} |\widehat{B}_{k,\nu}(\lambda)|^2 d\lambda \right)^{\frac{1}{2}}.$$

Also, the mass is preserved:

$$M := \|\{B_k(t)\}\|_{l^2}^2 = \|v(t)\|_{L^2(0,2\pi)}^2 = \|v(1)\|_{L^2(0,2\pi)}^2 = \|e^{i\cdot^2}\hat{u}_1(\cdot)\|_{L^2(0,2\pi)}^2.$$

Now that the solution v is constructed on $[1, \infty)$ we look for further large time properties of $\{B_k(t)\}$. As a first result we obtain a uniform in k pointwise decay for the Fourier in time transform of $\partial_t B_k(t)$.

Lemma 6.3. (Control in time of $\partial_t B_k(t)$) For any $\nu \in \mathbb{N}^*$, and η_{ν} a smooth cutoff supported in $[\nu, \nu+1]$, we have for any $\epsilon > 0$ the following estimate on the Fourier transform in time:

$$\sup_{k \in \mathbb{Z}} \sup_{\lambda \in \mathbb{R}} |\mathcal{F}(\eta_{\nu}(\cdot)\partial_{t}B_{k}(\cdot))(\lambda)| \leq \frac{C(\epsilon, \|\{B_{j}(1)\}\|_{l^{2}}\|\xi^{1^{+}}\hat{\eta}_{\nu}(\xi)\|_{L^{\infty}})}{\nu} \|\{B_{j}(\nu)\}\|_{l^{2,\epsilon}}.$$

Proof. We have

$$\begin{aligned} \mathcal{F}(\eta_{\nu}(\cdot)\partial_{t}B_{k}(\cdot))(\lambda) &= \int e^{it\lambda}\eta_{\nu}(t)\frac{1}{t}\sum_{NR_{k}}e^{it(k^{2}-j_{1}^{2}+j_{2}^{2}-j_{3}^{2})}(B_{j_{1}}\overline{B}_{j_{2}}B_{j_{3}})(t)dt \\ &+ \int e^{it\lambda}\eta_{\nu}(t)\frac{1}{t}\left(2M - |B_{k}(t)|^{2}\right)B_{k}(t)dt. \end{aligned}$$

Recall that NR_k denotes the noresonant set:

$$NR_k := \{ (j_1, j_2, j_3), \, k - j_1 + j_2 - j_3 = 0, \, k^2 - j_1^2 + j_2^2 - j_3^2 \neq 0 \}.$$

The second term can be estimated straightforwardly to get the $\frac{1}{\nu}$ decay. In the first term we split the summation between the sets:

$$\Lambda_{k,m} = \{ (j_1, j_2) \in \mathbb{Z}^2, (k - j_1)(j_1 - j_2) = m \}, \quad m \in \mathbb{Z}^*,$$

and we pass in Fourier in time to get:

$$\int e^{it\lambda} \eta_{\nu}(t) \frac{1}{t} \sum_{NR_{k}} e^{it(k^{2}-j_{1}^{2}+j_{2}^{2}-j_{3}^{2})} (B_{j_{1}}\overline{B}_{j_{2}}B_{j_{3}})(t)dt$$

$$= \int e^{it\lambda} \eta_{\nu}(t) \frac{1}{t} \sum_{m} \sum_{j_{1},j_{2}\in\Lambda_{k,m}} e^{itm} (B_{j_{1}}\overline{B}_{j_{2}}B_{k-j_{1}+j_{2}})(t)dt$$

$$= \frac{1}{\nu} \int \sum_{m} \widehat{\eta}_{\nu}(\lambda - \lambda_{1} + \lambda_{2} - \lambda_{3} + m) \sum_{j_{1},j_{2}\in\Lambda_{k,m}} \hat{B}_{j_{1},\nu}(\lambda_{1})\widehat{\overline{B}}_{j_{2},\nu}(\lambda_{2})\hat{B}_{k-j_{1}+j_{2},\nu}(\lambda_{3})d\lambda_{1}d\lambda_{2}d\lambda_{3}.$$

To end the proof of the lemma via the $X_{\nu}^{s,b}$ -norms we perform Cauchy-Schwarz successively:

• in j_1, j_2 using that for all $j_1, j_2 \in \Lambda_{k,m}$:

$$#\Lambda_{k,m} \le C(\epsilon)m^{\epsilon} = C(\epsilon)(k-j_1)^{\epsilon}(j_1-j_2)^{\epsilon} \le C(\epsilon)\max\{\langle j_1\rangle, \langle j_2\rangle, \langle k-j_1+j_2\rangle\}^{2\epsilon},$$

- in *m* using the decay of $\hat{\tilde{\eta}}_{\nu}$,
- in $\lambda_1, \lambda_2, \lambda_3$ using the integrability of $\frac{1}{\lambda^{2b}}$.

In a more complicated way we obtain the control asymptotically in time of the low regularity H^r norms, as follows.

Lemma 6.4. (Control in time of the weighted-norms) For any $t \ge 1, r \in (0, \frac{1}{2})$ we have: $\|\{B_k(t)\}\|_{l^{2,r}} \le C(r, \|\{B_k(1)\}\|_{l^{2,r}}).$

30

Proof. We use the evolution of $l^{2,r}$ -norms dictated by the equation of the B_k s:

$$\sum_{k} \langle k \rangle^{2r} |B_k(t)|^2 - \sum_{k} \langle k \rangle^{2r} |B_k(1)|^2$$
$$= \int_1^t \sum_{k;NR_k} (\langle k \rangle^{2r} - \langle j_1 \rangle^{2r} + \langle j_2 \rangle^{2r} - \langle j_3 \rangle^{2r}) e^{i\tau(k^2 - j_1^2 + j_2^2 - j_3^2)} B_{j_1} \overline{B_{j_2}} B_{j_3} \overline{B_k}(\tau) \frac{d\tau}{\tau},$$

by performing integrations by parts:

$$\begin{split} \sum_{k} \langle k \rangle^{2r} |B_{k}(t)|^{2} &\leq \sum_{k} \langle k \rangle^{2r} |B_{k}(1)|^{2} + C \left[\sum_{k;NR_{k}} |\varphi_{k,j_{1},j_{2},j_{3}} B_{j_{1}} \overline{B_{j_{2}}} B_{j_{3}} \overline{B_{k}}(\tau) \frac{1}{\tau} | \right]_{1}^{t} \\ &+ C \left| \int_{1}^{t} \sum_{k;NR_{k}} \varphi_{k,j_{1},j_{2},j_{3}} e^{i\tau(k^{2}-j_{1}^{2}+j_{2}^{2}-j_{3}^{2})} B_{j_{1}} \overline{B_{j_{2}}} B_{j_{3}} \overline{\partial_{\tau} B_{k}}(\tau) \frac{d\tau}{\tau} \right|, \end{split}$$

where

$$\varphi_{k,j_1,j_2,j_3} := \frac{\langle k \rangle^{2r} - \langle j_1 \rangle^{2r} + \langle j_2 \rangle^{2r} - \langle j_3 \rangle^{2r}}{k^2 - j_1^2 + j_2^2 - j_3^2}.$$

We prove that

$$\sum_{k;(j_1,j_2,j_3)\in NR_k} |\varphi_{k,j_1,j_2,j_3}| N_{j_1}N_{j_2}N_{j_3}P_k \le C(r, \|N\|_{l^2}) \|N\|_{l^{2,r}}^{0^+} \|P\|_{l^2},$$

for $r < \frac{1}{2}$ and any two sequences $\{N_j\}$ and $\{P_j\}$ of positive numbers. The estimates in the cases $r \geq \frac{1}{2}$ were obtained in Lemma 2.5 in [59]. This settles the boundary term. For the integral term we develop $\partial_{\tau}B_k$ using its equation, which yields a six-product term, then use a ν -partition of unity in τ , pass to Fourier in τ , and split the *j*'s summation into different regions where we use different arguments to get the control of the Lemma.

Now we shall identify the asymptotics of $B_k(t)$. We first obtain for all $k \in \mathbb{Z}$ a first modulated limit:

$$\exists \beta_k := \lim_{t \to \infty} \tilde{B}_k(t), \quad |\tilde{B}_k(t) - \beta_k| \le \frac{C(\|\{B_j(1)\}\|_{l^{2,r}})}{t},$$

where

$$\tilde{B}_k(t) := e^{i2M\log t - i\int_1^t |B_k(\tau)|^2 \frac{d\tau}{\tau}} B_k(t).$$

To do so we integrate in time the evolution law of \tilde{B}_k , we use again integrations by parts, partition of unity in time, Fourier in time, splitting the nonresonant set NR_k into the sets $\Lambda_{k,m}$, and the two previous lemmas. Then we obtain

$$\exists \alpha_k := \lim_{t \to \infty} e^{i(2M - |\beta_k|^2) \log t} B_k(t),$$
$$|B_k(t) - \alpha_k e^{i(2M - |\alpha_k|^2) \log t}| \le \frac{C(\|\{B_j(1)\}\|_{l^{2,s}})}{t}.$$

7. Appendix: Passing from 1D cubic Schrödinger equation to binormal flow

Hasimoto's initial transformation in 1972 is based on the Frenet system of a curve:

$$\left(\begin{array}{c} T\\n\\b\end{array}\right)_{x}=\left(\begin{array}{ccc} 0&c&0\\-c&0&\tau\\0&-\tau&0\end{array}\right)\left(\begin{array}{c} T\\n\\b\end{array}\right),$$

where c represents the curvature and τ the torsion (see the book of Spivak [107]). His construction is valid for curves with non-vanishing curvature. This constraint was removed in 1997 by Koiso in [92] by using instead the notion of parallel transport of frames. Before explaining how the construction works we start with the (time-independent) notion of parallel transport of frames.

7.0.1. Parallel transport of frames and the filament functions of a curve. We call filament function of a curve χ a function u_{χ} obtained by the following parallel transport procedure. Denote the tangent vector $T = \partial_x \chi$. We consider the parallel frames $(T, e_1, e_2)(x)$ obtained by solving the ODEs:

$$\partial_x e_1(x) = -\langle \partial_x T, e_1 \rangle T, \quad \partial_x e_2(x) = -\langle \partial_x T, e_2 \rangle T,$$

with data at some $x_0 \in \mathbb{R}$ given by an orthonormal frame of \mathbb{R}^3 . This is always possible if the curve χ is regular enough, for instance if $\partial_x^2 \chi \in L^2_{loc}$. Indeed, this gives global existence in H^1_{loc} for the above ODEs. In addition, the orthonormal frame nature of $(T, e_1, e_2)(x)$ is preserved, since the matrix of the system of evolution in space of $(T, e_1, e_2)(x)$ is antisymmetric. Then we define:

$$u_{\chi} = \langle \partial_x T, e_1 \rangle + i \langle \partial_x T, e_2 \rangle.$$

The real and imaginary part of u_{χ} are a normal development of the curve χ (see [26]). We called them also filament function by referring to the notion introduced by Hasimoto [81] for curves χ with non-vanishing curvature:

$$c(x)e^{i\int_0^x \tau(s)ds}$$
.

For curves χ with non-vanishing curvature this function coincides with u_{χ} and

(20)
$$(e_1 + ie_2)(x) = (n + ib)(x)e^{i\int_{x_0}^{\infty} \tau(s)ds},$$

if in the construction of u_{χ} the initial orthonormal frame $(T, e_1, e_2)(x_0)$ is chosen to be the Frenet frame $(T, n, b)(x_0)$. Observe that even if the curvature vanishes, the expression in the right-hand side of (20) continue to make sense via the parallel frame construction.

We note that the only degree of freedom in the filament function u_{χ} construction is rotating the initial data $(T, e_1, e_2)(x_0)$, i.e. rotating (e_1, e_2) in the plane orthogonal to $T(x_0)$, which yields by this construction another filament function that is of type $u_{\chi}(x)e^{i\theta}$ (and changing x_0 boils down to the same argument).

We also note that u_{χ} is constructed exclusively from $T = \partial_x \chi$ so it does not depend on translations in space of χ , i.e. it does not depend on $\chi(x_0)$. Moreover, u_{χ} is uniquely determined by $T(x_0) = \partial_x \chi(x_0)$ modulo multiplication by $e^{i\theta}$. Therefore if two curves $\chi, \tilde{\chi}$ satisfy $\chi(x_0) = \tilde{\chi}(x_0), \partial_x \chi(x_0) = \partial_x \chi(x_0)$ and $u_{\chi} = u_{\tilde{\chi}}$, then $\chi = \tilde{\chi}$.

32

Eventually the ODE satisfied by the frame is:

(21)
$$\begin{pmatrix} T\\ e_1\\ e_2 \end{pmatrix}_x = \begin{pmatrix} 0 & -\operatorname{Im} u_{\chi} & \operatorname{Re} u_{\chi}\\ \operatorname{Im} u_{\chi} & 0 & 0\\ -\operatorname{Re} u_{\chi} & 0 & 0 \end{pmatrix} \begin{pmatrix} T\\ e_1\\ e_2 \end{pmatrix}.$$

7.0.2. The Hasimoto approach from NLS to the binormal flow. We recall here the classical Hasimoto approach to construct binormal flow solutions by using sufficiently regular solutions of the cubic Schrödinger equation (4):

$$iu_t + u_{xx} + (|u|^2 - f)u = 0,$$

for f a space-independent function, as for instance the null function. Let \mathcal{B} be an orthonormal basis of \mathbb{R}^3 , $x_0 \in \mathbb{R}$ and $P \in \mathbb{R}^3$. Let us assume that we have a smooth solution u of (4) on an open time interval I, and let $t_0 \in I$. Starting from u, the first step will be to construct orthonormal frames $(T, e_1, e_2)(t, x)$ such that the first vector T is a solution of the Schrödinger map. In order to construct these frames we solve the ODE in time:

(22)
$$\begin{pmatrix} T \\ e_1 \\ e_2 \end{pmatrix}_t (t, x_0) = \begin{pmatrix} 0 & -\operatorname{Im} u_x & \operatorname{Re} u_x \\ \operatorname{Im} u_x & 0 & -\frac{|u|^2 - f}{2} \\ -\operatorname{Re} u_x & \frac{|u|^2 - f}{2} & 0 \end{pmatrix} \begin{pmatrix} T \\ e_1 \\ e_2 \end{pmatrix} (t, x_0), \quad \forall t \in I,$$

with initial condition \mathcal{B} at time t_0 . The orthonormal frame character is preserved in the evolution due to the fact that the matrix of the system is antisymmetric. Then, for all $t \in I$, we solve the family of ODEs in space:

(23)
$$\begin{pmatrix} T\\ e_1\\ e_2 \end{pmatrix}_x (t,x) = \begin{pmatrix} 0 & -\operatorname{Im} u & \operatorname{Re} u\\ \operatorname{Im} u & 0 & 0\\ -\operatorname{Re} u & 0 & 0 \end{pmatrix} \begin{pmatrix} T\\ e_1\\ e_2 \end{pmatrix} (t,x), \quad \forall x \in \mathbb{R}.$$

with initial condition the frame $(T, e_1, e_2)(t, x_0)$ at the point x_0 . Using the fact that u is a solution of equation (4) one can prove⁷ that, at least for regular u, the ODE (22) is actually valid at any $x \in \mathbb{R}$:

$$(24) \quad \begin{pmatrix} T\\ e_1\\ e_2 \end{pmatrix}_t (t,x) = \begin{pmatrix} 0 & -\operatorname{Im} u_x & \operatorname{Re} u_x\\ \operatorname{Im} u_x & 0 & -\frac{|u|^2 - f}{2}\\ -\operatorname{Re} u_x & \frac{|u|^2 - f}{2} & 0 \end{pmatrix} \begin{pmatrix} T\\ e_1\\ e_2 \end{pmatrix} (t,x), \qquad \forall t \in I, x \in \mathbb{R}.$$

⁷Since $(T, e_1, e_2)(t, s)$ are orthonormal frames we can write for all t

$$\begin{pmatrix} T\\ e_1\\ e_2 \end{pmatrix}_t (t,x) = \begin{pmatrix} 0 & a & b\\ -a & 0 & c\\ -b & -c & 0 \end{pmatrix} \begin{pmatrix} T\\ e_1\\ e_2 \end{pmatrix} (t,x).$$

We also notice that $(a, b, c)(t, x_0) = (-\Im u_x, \Re u_x, -\frac{|u|^2 - f}{2})(t, x_0)$. By computing and identifying $T_{ts} = T_{st}, e_{1ts} = e_{1st}$ we get a system for (a, b, c) in terms of u, that together with the fact that u satisfies (4) allows for identifying $(a, b, c)(t, x) = (-\Im u_x, \Re u_x, -\frac{|u|^2 - f}{2})(t, x)$.

It is then easy to see that T solves the Schrödinger map equation. Indeed from (24) and (23) we obtain:

$$T \wedge T_{xx} = T \wedge (\operatorname{Re} u \, e_1 + \operatorname{Im} u \, e_2)_x = T \wedge (\operatorname{Re} u_x \, e_1 + \operatorname{Im} u_x \, e_2 + \operatorname{Re} u \, e_{1x} + \operatorname{Im} u \, e_{2x})$$
$$= \operatorname{Re} u_x \, e_2 - \operatorname{Im} u_x \, e_1 = T_t.$$

Moreover, one can easily check that χ defined as

(25)
$$\chi(t,x) := P + \int_{t_0}^t (T \wedge T_x)(s,x_0)ds + \int_{x_0}^x T(t,y)dy,$$

is a solution of the binormal flow equation, and that T is its tangent vector. Indeed, simply take the time derivative of (25) and use the divergence form $T_t = (T \wedge T_x)_x$. Finally let us note that u(t) is a filament function of $\chi(t)$.

By using the complex normal vector $N := e_1 + ie_2$ the equations write in a shorter way:

(26)
$$T_x = \Re \overline{u}N, \quad N_x = -uT, \quad T_t = \Im \overline{u_x}N, \quad N_t = -iu_xT + i\frac{|u|^2 - f}{2}N, \quad \chi_t = \Im \overline{u}N.$$

In particular, if u is solution of (4) that yields (T, N) by the above construction, then $u(t, x)e^{i\Phi(t)}$ is a solution of (4) with $f - \Phi$ instead of f that yields $(T, Ne^{i\Phi(t)})$ by the above construction, thus the same first vector, thus the same binormal flow solution.

To conclude, let us give some examples of known, rigorously or through experiments, vortex filament smooth dynamics that correspond to a binormal flow solution, and let us give also the solutions of 1D cubic Schrödinger equation (3) from which these binormal flow solutions are constructed via Hasimoto's approach:

• Straight still lines are parametrized by $\chi(t, x) = (0, 0, x)$ and correspond to u(t, x) = 0,

- Circles translating in the binormal direction (smoke rings) $\chi(t,s) = (\cos s, \sin s, t)$ are obtained from $u(t,x) = e^{-it}$,
- Helices with constant torsion translating on themselves are obtained from $u(t,x) = e^{-it-it\alpha^2 + i\alpha x}, \alpha \in \mathbb{R}$,

• Travelling waves solutions of the binormal flow were obtained by Hasimoto from $u(t, x) = e^{-it}e^{-it\alpha^2}e^{i\alpha x}\frac{1}{2\sqrt{2}\cosh(x-2\alpha t)}, \alpha \in \mathbb{R}.$

References

- Fabien Anselmet, Yves Gagne, Emil J. Hopfinger, and Robert A. Antonia. High-order velocity structure functions in turbulent shear flows. J. Fluid Mech., 140:63–89, 1984.
- [2] Gabriel B. Apolinário, Geoffrey Beck, Laurent Chevillard, Isabelle Gallagher, and Ricardo Grande. A linear stochastic model of turbulent cascades and fractional fields. To appear in Ann. Sc. Norm. Super. Pisa Cl. Sci., 2023.
- [3] Gabriel B. Apolinário, Laurent Chevillard, and Jean-Christophe Mourrat. Dynamical fractional and multifractal fields. J. Stat. Phys., 186:35, 2022.
- [4] Dario Bambusi, Benoît Grébert, Alberto Maspero, and Didier Robert. Reducibility of the quantum harmonic oscillator in *d*-dimensions with polynomial time-dependent perturbation. *Anal. PDE*, 11:775–799, 2018.
- [5] Dario Bambusi, Beatrice Langella, and Riccardo Montalto. Growth of Sobolev norms for unbounded perturbations of the Schrödinger equation on flat tori. J. Differ. Equations, 318:344–358, 2022.

34

- [6] V. Banica and L. Vega. On the stability of a singular vortex dynamics. Commun. Math. Phys., 286:593-627, 2009.
- [7] Valeria Banica, Daniel Eceizabarrena, Andrea Nahmod, and Luis Vega. Multifractality and intermittency in the limit evolution of polygonal vortex filaments. *Math. Ann.*, 2024.
- [8] Valeria Banica, Renato Lucà, Nikolay Tzvetkv, and Luis Vega. Blow-up for the 1D cubic NLS. Comm. Math. Phys., 405:21 pp., 2024.
- [9] Valeria Banica and Luis Vega. Scattering for 1d cubic NLS and singular vortex dynamics. J. Eur. Math. Soc. (JEMS), 14:209-253, 2012.
- [10] Valeria Banica and Luis Vega. Stability of the self-similar dynamics of a vortex filament. Arch. Ration. Mech. Anal., 210:673–712, 2013.
- [11] Valeria Banica and Luis Vega. The initial value problem for the binormal flow with rough data. Ann. Sci. Éc. Norm. Supér. (4), 48:1423–1455, 2015.
- [12] Valeria Banica and Luis Vega. Evolution of polygonal lines by the binormal flow. Ann. PDE, 6:53, 2020.
- [13] Valeria Banica and Luis Vega. On the energy of critical solutions of the binormal flow. Comm. PDE, 45:820–845, 2020.
- [14] Valeria Banica and Luis Vega. Riemann's non-differentiable function and the binormal curvature flow. Arch. Ration. Mech. Anal., 244:501–540, 2022.
- [15] Valeria Banica and Luis Vega. Unbounded growth of the energy density associated to the Schrödinger map and the binormal flow. Ann. Inst. Henri Poincaré, Anal. Non Linéaire, 39:927–946, 2022.
- [16] Valeria Banica and Luis Vega. New conservation laws and energy cascade for 1d cubic NLS and the Schrödinger map. Vietnam J. Math., 52:985–999, 2024.
- [17] Julien Barral and Stéphane Seuret. The Frisch-Parisi conjecture. I: Prescribed multifractal behavior, and a partial solution. J. Math. Pures Appl., 175:76–108, 2023.
- [18] Julien Barral and Stéphane Seuret. The Frisch-Parisi conjecture. II: Besov spaces in multifractal environment, and a full solution. J. Math. Pures Appl., 175:281–329, 2023.
- [19] Jacob Bedrossian, Pierre Germain, and Benjamin Harrop-Griffiths. Vortex filament solutions of the Navier-Stokes equations. Comm. Pure Appl. Math., 76:685–787, 2023.
- [20] Victor Beresnevich and Sanju Velani. A mass transference principle and the Duffin-Schaeffer conjecture for Hausdorff measures. Ann. of Math., 164:971–992, 2006.
- [21] M. V. Berry and S. Klein. Integer, fractional and fractal Talbot effects. J. Mod. Opt., 43:2139–2164, 1996.
- [22] Michael Berry. Quantum fractals in boxes. J. Phys. A, Math. Gen., 29:6617–6629, 1996.
- [23] Michael Berry and Wolfgang Schleich. Quantum carpets, carpets of light. Phys. World, 14:1–6, 2001.
- [24] A. S. Besicovitch. Sets of fractional dimensions. IV: On rational approximation to real numbers. J. Lond. Math. Soc., 9:126–131, 1934.
- [25] Anxo Biasi and Oleg Evnin. Turbulent cascades in a truncation of the cubic Szego equation and related systems. Analysis PDE, 15:217–243, 2022.
- [26] Richard L. Bishop. There are more than one way to frame a curve. Amer. Math. Mon., 82:246–251, 1975.
- [27] Jerry L. Bona and Jean-Claude Saut. Dispersive blow-up. II: Schrödinger-type equations, optical and oceanic rogue waves. Chin. Ann. Math., Ser. B, 31:793–818, 2010.
- [28] Alexandre Boritchev, Daniel Eceizabarrena, and Victor Vilaça Da Rocha. Intermittency of Riemann's non-differentiable function through the fourth-order flatness. J. Math. Phys., 62:Paper No. 093101, 14, 2021.
- [29] Lyonell Boulton, George Farmakis, Beatrice Pelloni, and David A. Smith. Jumps and cusps: a new revival effect in local dispersive PDEs, 2024.
- [30] Jean Bourgain. Aspects of long time behaviour of solutions of nonlinear Hamiltonian evolution equations. Geom. Funct. Anal., 5:105–140, 1995.

- [31] Jean Bourgain. On the growth in time of higher Sobolev norms of smooth solutions of Hamiltonian PDE. Int. Math. Res. Not., 1996:277–304, 1996.
- [32] Jean Bourgain. Growth of Sobolev norms in linear Schrödinger equations with quasi-periodic potential. Commun. Math. Phys., 204:207–247, 1999.
- [33] Marco Bravin and Luis Vega. On the one dimensional cubic NLS in a critical space. Discrete Contin. Dyn. Syst., 42:2563–2584, 2022.
- [34] Frederik Broucke and Jasson Vindas. The pointwise behavior of Riemann's function. J. Fractal Geom., 10:333–349, 2023.
- [35] Thomas F. Buttke. A numerical study of superfluid turbulence in the self-induction approximation. J. of Comp. Physics, 76:301–326, 1988.
- [36] Rémi Carles and Erwan Faou. Energy cascades for NLS on the torus. Discrete Contin. Dyn. Syst., 32:2063–2077, 2012.
- [37] Rémi Carles and Isabelle Gallagher. Universal dynamics for the defocusing logarithmic Schrödinger equation. Duke Math. J., 167:1761–1801, 2018.
- [38] Rémi Carles and Thomas Kappeler. Norm-inflation with infinite loss of regularity for periodic NLS equations in negative Sobolev spaces. Bull. Soc. Math. Fr., 145:623–642, 2017.
- [39] Thierry Cazenave and Fred B. Weissler. The Cauchy problem for the critical nonlinear Schrödinger equation in H^s. Nonlinear Anal., Theory Methods Appl., 14:807–836, 1990.
- [40] Fernando Chamizo and Adrián Ubis. Multifractal behavior of polynomial Fourier series. Adv. Math., 250:1–34, 2014.
- [41] Laurent Chevillard, Bernard Castaing, Alain Arneodo, Emmanuel Lévêque, Jean-François Pinton, and Stéphane G. Roux. A phenomenological theory of Eulerian and Lagrangian velocity fluctuations in turbulent flows. *Comptes Rendus. Physique*, 13:899–928, 2012.
- [42] Michael Christ, James Colliander, and Terrence Tao. Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations. Am. J. Math., 125:1235–1293, 2003.
- [43] Yves Colin de Verdière and Laure Saint-Raymond. Attractors for two-dimensional waves with homogeneous Hamiltonians of degree 0. Commun. Pure Appl. Math., 73:421–462, 2020.
- [44] James E. Colliander, Markus Keel, Gigliola Staffilani, Hideo Takaoka, and Terence Tao. Transfer of energy to high frequencies in the cubic defocusing nonlinear Schrödinger equation. *Invent. Math.*, 181:39–113, 2010.
- [45] Peter Constantin, Charles Fefferman, and Andrew J. Majda. Geometric constraints on potentially singular solutions for the 3 – D Euler equations. Commun. Partial Differ. Equations, 21:559–571, 1996.
- [46] Luigi Sante Da Rios. Sul moto d'un liquido indefinito con un filetto vorticoso di forma qualunque. Rend. Circ. Mat. Palermo, 22:117–135, 1906.
- [47] Juan Dávila, Manuel del Pino, Monica Musso, and Juncheng Wei. Travelling helices and the vortex filament conjecture in the incompressible Euler equations. *Calc. Var. Partial Differential Equations*, 61:Paper No. 119, 30, 2022.
- [48] Francisco de la Hoz, Sandeep Kumar, and Luis Vega. On the evolution of the vortex filament equation for regular *M*-polygons with nonzero torsion. *SIAM J. Appl. Math.*, 80:1034–1056, 2020.
- [49] Francisco de la Hoz and Luis Vega. Vortex filament equation for a regular polygon. Nonlinearity, 27:3031–3057, 2014.
- [50] Francisco de la Hoz and Luis Vega. On the relationship between the one-corner problem and the M-corner problem for the vortex filament equation. J. Nonlinear Sci., 28:2275–2327, 2018.
- [51] J.-M. Delort. Growth of Sobolev norms for solutions of time dependent Schrödinger operators with harmonic oscillator potential. Commun. Partial Differ. Equations, 39:1–33, 2014.
- [52] Yu Deng and Pierre Germain. Growth of solutions to NLS on irrational tori. Int. Math. Res. Not., 2019:2919–2950, 2019.
- [53] Martin Donati, Christophe Lacave, and Evelyne Miot. Dynamics of helical vortex filaments in non viscous incompressible flows. ArXiv:2403.00389, 2024.

36

- [54] Richard J. Duffin and Albert C. Schaeffer. Khintchine's problem in metric Diophantine approximation. Duke Math. J., 8:243–255, 1941.
- [55] M. B. Erdoğan and N. Tzirakis. Talbot effect for the cubic non-linear Schröedinger equation on the torus. Math. Res. Lett., 20:1081–1090, 2013.
- [56] Erwan Faou and Pierre Raphaël. On weakly turbulent solutions to the perturbed linear harmonic oscillator. Am. J. Math., 145:1465–1507, 2023.
- [57] Enrico Fonda, David P. Meichle, Nicolas T. Ouellette, and Daniel P. Lathrop. Direct observation of kelvin waves excited by quantized vortex reconnection. PNAS, 111:4707–4710, 2014.
- [58] Marco Antonio Fontelos and Luis Vega. Evolution of viscous vortex filaments and desingularization of the Biot-Savart integral. ArXiv:2311.12246, 2023.
- [59] Justin Forlano and Kihoon Seong. Transport of gaussian measures under the flow of one-dimensional fractional nonlinear schrödinger equations. Comm. PDE, 47:1296–1337, 2022.
- [60] Uriel Frisch. Turbulence: The legacy of A. N. Kolmogorov. Cambridge University Press, Cambridge, 1995.
- [61] Thierry Gallay and Vladimír Sverák. Vanishing viscosity limit for axisymmetric vortex rings. Invent. Math., 237:275–348, 2024.
- [62] Miguel Angel Garrido, Ricardo Grande, Kristin M. Kurianski, and Gigliola Staffilani. Large deviations principle for the cubic NLS equation. Commun. Pure Appl. Math., 76:4087–4136, 2023.
- [63] Patrick Gérard and Sandrine Grellier. Effective integrable dynamics for a certain nonlinear wave equation. Anal. PDE, 5:301–326, 2012.
- [64] Patrick Gérard and Sandrine Grellier. The cubic Szegő equation and Hankel operators, volume 389 of Astérisque. Paris: Société Mathématique de France (SMF), 2017.
- [65] Patrick Gérard, Sandrine Grellier, and Zihui He. Turbulent cascades for a family of damped Szegő equations. Nonlinearity, 35:4820–4849, 2022.
- [66] Patrick Gérard and Enno Lenzmann. The Calogero-Moser derivative nonlinear Schrödinger equation. Commun. Pure Appl. Math., 77:4008–4062, 2024.
- [67] Patrick Gérard, Enno Lenzmann, Oana Pocovnicu, and Pierre Raphaël. A two-soliton with transient turbulent regime for the cubic half-wave equation on the real line. Ann. PDE, 4:166, 2018. Id/No 7.
- [68] J. Ginibre and G. Velo. The global Cauchy problem for the nonlinear Schrödinger equation revisited. Ann. Inst. Henri Poincaré, Anal. Non Linéaire, 2:309–327, 1985.
- [69] Filippo Giuliani and Marcel Guardia. Sobolev norms explosion for the cubic NLS on irrational tori. Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods, 220:25, 2022. Id/No 112865.
- [70] Benoît Grébert, Éric Paturel, and Laurent Thomann. Beating effects in cubic Schrödinger systems and growth of Sobolev norms. *Nonlinearity*, 26:1361–1376, 2013.
- [71] Fernando F. Grinstein and Carl R. DeVore. Dynamics of coherent structures and transition to turbulence in free square jets. *Phys. Fluids*, 8:1237–1251, 1996.
- [72] Axel Grünrock. Bi- and trilinear Schrödinger estimates in one space dimension with applications to cubic NLS and DNLS. Int. Math. Res. Not., 2005:2525–2558, 2005.
- [73] M. Guardia and Vadim Kaloshin. Growth of Sobolev norms in the cubic defocusing nonlinear Schrödinger equation. J. Eur. Math. Soc., 17:71–149, 2015.
- [74] Marcel Guardia, Zaher Hani, Emanuele Haus, Alberto Maspero, and Michela Procesi. Strong nonlinear instability and growth of Sobolev norms near quasiperiodic finite gap tori for the 2d cubic NLS equation. J. Eur. Math. Soc., 25:1497–1551, 2023.
- [75] Anatole Guérin. A new class of critical solutions for 1d cubic NLS. ArXiv:2212.08564, 2022.
- [76] Anatole Guérin. On self-similar singularity formation for the binormal flow. J. Evol. Equ., 23:18, 2023.
- [77] Susana Gutiérrez, Judith Rivas, and Luis Vega. Formation of singularities and self-similar vortex motion under the localized induction approximation. *Comm. Partial Differential Equations*, 28:927– 968, 2003.
- [78] Zaher Hani. Long-time instability and unbounded Sobolev orbits for some periodic nonlinear Schrödinger equations. Arch. Ration. Mech. Anal., 211:929–964, 2014.

- [79] Zaher Hani, Benoit Pausader, Nikolay Tzvetkov, and Nicola Visciglia. Modified scattering for the cubic Schrödinger equation on product spaces and applications. Forum Math. Pi, 3:63, 2015.
- [80] Benjamin Harrop-Griffiths, Rowan Killip, and Monica Vişan. Sharp well-posedness for the cubic NLS and mKdV in $H^{s}(\mathbb{R})$. Forum Math. Pi, 12:86, 2024.
- [81] Hidenori Hasimoto. A soliton on a vortex filament. J. Fluid Mech., 51:477-485, 1972.
- [82] Stephane Jaffard. The spectrum of singularities of Riemann's function. Rev. Mat. Iberoamericana, 12:441–460, 1996.
- [83] Vojtěch Jarník. On simultaneous Diophantine approximations. Math. Z., 33:505–543, 1931.
- [84] Robert L. Jerrard and Christian Seis. On the vortex filament conjecture for Euler flows. Arch. Ration. Mech. Anal., 224:135–172, 2017.
- [85] Robert L. Jerrard and Didier Smets. On the motion of a curve by its binormal curvature. J. Eur. Math. Soc., 17:1487–1515, 2015.
- [86] Lev Kapitanski and Igor Rodnianski. Does a quantum particle know the time? In *Emerging applications of number theory (Minneapolis, MN, 1996)*, volume 109 of *IMA Vol. Math. Appl.*, pages 355–371. Springer, New York, 1999.
- [87] Carlos E. Kenig, Gustavo Ponce, and Luis Vega. On the ill-posedness of some canonical dispersive equations. Duke Math. J., 106:617–633, 2001.
- [88] Rowan Killip, Monica Vişan, and Xiaoyi Zhang. Low regularity conservation laws for integrable PDE. Geom. Funct. Anal., 28:1062–1090, 2018.
- [89] Nobu Kishimoto. Well-posedness of the Cauchy problem for the Korteweg-de Vries equation at the critical regularity. *Differ. Integral Equ.*, 22:447–464, 2009.
- [90] Naoyasu Kita. Mode generating property of solutions to the nonlinear Schrödinger equations in one space dimension. Nonlinear dispersive equations, Tokyo: Gakkotosho, pages 111–128, 2006.
- [91] Herbert Koch and Daniel Tataru. Conserved energies for the cubic nonlinear Schrödinger equation in one dimension. Duke Math. J., 167:3207–3313, 2018.
- [92] Norihito Koiso. The vortex filament equation and a semilinear Schrödinger equation in a Hermitian symmetric space. Osaka J. Math., 34:199–214, 1997.
- [93] Dimitris Koukoulopoulos and James Maynard. On the Duffin-Schaeffer conjecture. Ann. of Math., 192:251–307, 2020.
- [94] Sergei B. Kuksin. Growth and oscillations of solutions of nonlinear Schrödinger equation. Commun. Math. Phys., 178:265–280, 1996.
- [95] Sergei B. Kuksin. Oscillation in space-periodic nonlinear Schrödinger equations. Geom. Funct. Anal., 7:338–363, 1997.
- [96] Muthusamy Lakshmanan and Muthiah Daniel. On the evolution of higher dimensional Heisenberg continuum spin systems. *Physics A*, 107:533–552, 1981.
- [97] Frank Merle. On uniqueness and continuation properties after blow-up time of self-similar solutions of nonlinear Schrödinger equation with critical exponent and critical mass. Comm. Pure Appl. Math, 45:203–254, 1992.
- [98] Frank Merle, Pierre Raphaël, and Jérémie Szeftel. The instability of Bourgain-Wang solutions for the L² critical NLS. Amer. J. Math., 135:967–1017, 2013.
- [99] Tadahiro Oh. A remark on norm inflation with general initial data for the cubic nonlinear Schrödinger equations in negative Sobolev spaces. Funkc. Ekvacioj, Ser. Int., 60:259–277, 2017.
- [100] Konstantin I. Oskolkov. A class of I.M. Vinogradov's series and its applications in harmonic analysis. Springer Ser. Comput. Math., Springer, New York, pages 353–402, 1992.
- [101] Fabrice Planchon, Nikolay Tzvetkov, and Nicola Visciglia. On the growth of Sobolev norms for NLS on 2- and 3-dimensional manifolds. Anal. PDE, 10:1123–1147, 2017.
- [102] Oana Pocovnicu. Explicit formula for the solution of the Szegö equation on the real line and applications. Discrete Contin. Dyn. Syst., 31:607–649, 2011.
- [103] Igor Rodnianski. Fractal solutions of the Schrödinger equation. In Nonlinear PDE's, dynamics and continuum physics, pages 181–187. Providence, RI: American Mathematical Society (AMS), 2000.

- [104] Kevin W. Schwarz. Three-dimensional vortex dynamics in superfluid ⁴He: Line-line and line-boundary interactions. *Phys. Rev. B*, 31:5782–5804, 1981.
- [105] Stéphane Seuret and Adrián Ubis. Local L²-regularity of Riemann's Fourier series. Ann. Inst. Fourier, 67:2237–2264, 2017.
- [106] Vedran Sohinger. Bounds on the growth of high Sobolev norms of solutions to nonlinear Schrödinger equations on R. Indiana Univ. Math. J., 60:1487–1516, 2011.
- [107] Michael Spivak. A comprehensive introduction to differential geometry. Vol. 2. Waltham, Mass.: Brandeis University. ix, 425 p., 1970.
- [108] Laurent Thomann. Growth of Sobolev norms for linear Schrödinger operators. Ann. Henri Lebesgue, 4:1595–1618, 2021.
- [109] Kuniaki Todoya and Fazle Hussain. A study of the vortical structures of noncircular jets. Trans. Japan Soc. Mech. Eng. Ser. B, 55:1542–1545, 1989.
- [110] Yoshio Tsutsumi. L²-solutions for nonlinear Schrödinger equations and nonlinear groups. Funkc. Ekvacioj, Ser. Int., 30:115–125, 1987.
- [111] Ana Vargas and Luis Vega. Global wellposedness for 1D nonlinear Schrödinger equation for data with an infinite L^2 norm. J. Math. Pures Appl. (9), 80:1029–1044, 2001.

(V. Banica) SORBONNE UNIVERSITÉ, UNIVERSITÉ PARIS CITÉ, CNRS, INRIA, LABORATOIRE JACQUES-LOUIS LIONS, LJLL, F-75005 PARIS, FRANCE, VALERIA.BANICA@SORBONNE-UNIVERSITE.FR

(L. Vega) BCAM-UPV/EHU DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BASQUE COUNTRY, APDO 644, 48080 BILBAO, SPAIN, LUIS.VEGA@EHU.ES