

COMBINATORIAL INTERPRETATION OF THE COEFFICIENTS OF THE CAUSAL SET D’ALEMBERTIAN

KAREN YEATS

ABSTRACT. The causal set theory d’Alembertian has rational coefficients for which alternating expressions are known. Here, a combinatorial interpretation of these numbers is given.

1. INTRODUCTION

Causal set theory is an approach to quantum gravity where the causal relation between spacetime points and discreteness of spacetime are the two fundamental assumptions upon which one attempts to build everything else. Consequently, in this approach, one works with locally finite posets (also known as *causal sets*) as the spacetime where the partial order is the causal order in the spacetime, [4], for a review see [6].

To quickly set poset notation and language, a *poset* P is a set (by standard abuse of notation also called P), with a reflexive, antisymmetric, and transitive relation which we’ll write \leq . For $a, b \in P$ we write $a < b$ if $a \leq b$ and $a \neq b$. We say b *covers* a if $a < b$ but there is no $c \in P$ such that $a < c < b$. The *interval* from a to b is $[a, b] = \{c \in P : a \leq c \leq b\}$. Our intervals are by definition closed intervals, sometimes in causal set theory open intervals (not including a and b) are used instead; it is straightforward to convert between the different conventions. A poset is *locally finite* if all its intervals are finite sets.

Returning to the causal set context, while one ultimately would like to begin with causal sets alone, without reference to an underlying continuous spacetime, it is often useful to start with a Lorentzian manifold and choose points randomly on it according to a Poisson distribution, letting those points with the induced causal order be our causal set. Here the probability to find m poset elements in a volume V is $(\rho V)^m e^{-\rho V} / m!$, where ρ is a density parameter. Instead of working with ρ it is often convenient to work in terms of a length parameter, called the *discreteness scale*, ℓ defined by $\rho = \ell^{-d}$. Generating a causal set in this manner is called *sprinkling*, see Section 3 of [6].

One of the many operators which one has in the continuum case, which one would like an analogue of for causal sets, is the d’Alembertian or box operator, which in 4-dimensional Minkowski space is

$$\square = \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} - \frac{\partial^2}{\partial z^2}.$$

The author is supported by an NSERC Discovery grant and by the Canada Research Chairs program. The author is grateful to Max Planck Institute for Mathematics in Bonn for its hospitality and financial support. They also thank Rafael, Sorkin, Fay Dowker, Stav Zalel and Nick Olson-Harris for useful conversations.

Generally speaking, when moving to the discrete situation, one would expect differential operators, as in the classical d'Alembertian, to become difference operators. However, causal sets are only *locally* finite posets, so we can have infinitely many nearest neighbours of any given point, that is for a in the poset there can be infinitely many elements covering a and there can be infinitely many elements that a covers. Consequently, there's no reason to expect that a calculation involving nearest neighbours would converge in an infinite but locally finite poset, nor, if we take finite approximations is there a reason to expect a calculation involving nearest neighbours in the approximations to converge in any reasonable limit.

However, Sorkin in [5] observed that a particular alternating linear combination has the property that it approximates the d'Alembertian of 2-dimensional Minkowski space in the following sense. Let C be a causal set and let ϕ be a scalar field on C , which for us just means a function from C to \mathbb{C} . Define

$$B^{(2)}\phi(x) = \frac{1}{\ell^2} \left(-2\phi(x) + 4 \left(\sum_{y \in L_1(x)} \phi(y) - 2 \sum_{y \in L_2(x)} \phi(y) + \sum_{y \in L_3(x)} \phi(y) \right) \right)$$

where $L_i(x) = \{y \in C : |[y, x]| = i + 1\}$. Then letting $\langle B\phi(x) \rangle$ be the expectation with respect to Poisson sprinkling on 2-dimensional Minkowski space, Sorkin proves that

$$\lim_{\rho \rightarrow \infty} \frac{1}{\sqrt{\rho}} \langle B\phi(x) \rangle = \square\phi.$$

This result was generalized by Benincasa and Dowker to 4 dimensions [1], by Dowker and Glaser to general dimension [2] but without an explicit expression past $d = 7$, and then finally Glaser in [3] gave the following general form for the d -dimensional causal set d'Alembertian:

$$B^{(d)}\phi(x) = \frac{1}{\ell^2} \left(\alpha_d \phi(x) + \beta_d \sum_{i=1}^{\lfloor \frac{d}{2} \rfloor + 2} c_i^{(d)} \sum_{y \in L_i} \phi(y) \right)$$

where

$$\beta_d = \begin{cases} \frac{2\Gamma(\frac{d}{2} + 2)\Gamma(\frac{d}{2} + 1)}{\Gamma(\frac{d}{2})\Gamma(d)} c_d^{\frac{2}{d}} & \text{for even } d \\ \frac{d+1}{2^{d-1}\Gamma(\frac{d}{2} + 1)} c_d^{\frac{2}{d}} & \text{for odd } d \end{cases}$$

$$\alpha_d = \begin{cases} \frac{-2c_d^{\frac{2}{d}}}{\Gamma(\frac{d+2}{2})} & \text{for even } d \\ \frac{-c_d^{\frac{2}{d}}}{\Gamma(\frac{d+2}{d})} & \text{for odd } d \end{cases}$$

$$C_i^{(d)} = \begin{cases} \sum_{k=0}^{i-1} \binom{i-1}{k} (-1)^k \frac{\Gamma(\frac{d}{2}(k+1) + 2)}{\Gamma(\frac{d}{2} + 2)\Gamma(1 + \frac{dk}{2})} & \text{for even } d \\ \sum_{k=0}^{i-1} \binom{i-1}{k} (-1)^k \frac{\Gamma(\frac{d}{2}(k+1) + \frac{3}{2})}{\Gamma(\frac{d+3}{2})\Gamma(1 + \frac{dk}{2})} & \text{for odd } d \end{cases}$$

and $c_d = S_{d-2}/(d(d-1)2^{d/2-1})$ with S_{d-2} the volume of the $d-2$ dimensional unit sphere.

These $B^{(d)}$ when applied to sprinklings on d -dimensional Minkowski space likewise limit to the d -dimensional classical d'Alembertian [3].

These same coefficients also appear in the Benincasa-Dowker-Glaser (BDG) action in causal set theory [1]. This is because on a curved spacetime Benincasa, Dowker, and Glaser proved that $\lim_{\rho \rightarrow \infty} \langle B^{(d)} \phi(x) \rangle = \square \phi - \frac{1}{2} R(x) \phi(x)$ [1, 2, 3] where $R(x)$ is the Ricci scalar of the continuous spacetime. This gives a candidate for an analogue of the Ricci scalar for a causal set. Applying this to the constant field and summing over all elements of C then gives the analogue of the Einstein-Hilbert action for the causal set, which is the BDG action:

$$S^{(d)}(C) = -\alpha_d \ell^{d-2} \left(N + \frac{\beta_d}{\alpha_d} \sum_{i=1}^{\lfloor \frac{d}{2} \rfloor + 2} C_i^{(d)} N_i \right)$$

where $N = |C|$ and N_i is the number of intervals of size $i + 1$ in C .¹

Glaser comments at the end of Section 2 of [3], that the expressions for the $C_i^{(d)}$, while fast to compute, remain aesthetically unsatisfying since they could not simplify the sum. From an enumerative perspective, these sums cry out for a combinatorial explanation, one where the alternation can be realized as explicit cancellations. This is what the present paper accomplishes. Hopefully the reader will find this aesthetically satisfying, even without non-alternating expressions for the counts.

2. A CLASS OF NONCROSSING PARTIAL CHORD DIAGRAMS

We take a two phase approach to interpreting the $C_i^{(d)}$. The first phase is to define objects which give an interpretation for the ratio of gamma factors that appears in the sum. Furthermore, we want a unified interpretation that works for both the odd and even dimensions. The second phase is to modify these objects (in a way that's counted by the explicit binomial coefficients in the sum) so that the alternating sum can be realized by explicit cancellations, thus giving a class of objects which are counted by the $C_i^{(d)}$ directly.

The reader should note that the objects we define are certainly not the only objects which would have this property, but these ones seem reasonable and do the job.

2.1. Phase 1. Our main objects of interest are a particular class of coloured partial non-crossing rooted chord diagrams.

Definition 1. A *partial rooted chord diagram* with n chords on m points is a set partition of $\{1, 2, \dots, m\}$ where every part has size 1 or 2 and there are n parts of size 2. The parts of size 2 are called *chords*. A *rooted chord diagram* with n chords is a partial rooted chord diagram with n chords on $2n$ points, or equivalently where no parts have size 1.

When it is useful to emphasize a rooted chord diagram is not a partial diagram we'll call it a *complete* rooted chord diagram.

¹This differs from [3] by an overall factor independent of d , involving the rationalized Planck length, which is not important for our purposes.




FIGURE 1. A partial rooted chord diagram.

Definition 2. Two chords $\{a, b\}$ and $\{c, d\}$ with $a < b$ and $c < d$ *cross* if $a < c < b < d$ or $c < a < b < d$. A chord diagram is *noncrossing* if no two of its chords cross.

We will draw our partial rooted chord diagrams with m points on a circle, indicating point 1 with an \times and taking the rest in counterclockwise order, and indicating the chords by arcs inside the circle, see Figure 1. Often rooted chord diagrams are drawn with the points $\{1, 2, \dots, m\}$ from left to right on a line, but we will be making use of a notion of inside that is more visually appealing in the circle representation.

Call a point *bare* if it does not have any chord on it, or equivalently if it is a part of size 1 in the set partition.

Given a chord in a partial chord diagram, picking one end of the chord to be the *first end* lets us define the *inside* of the chord as the cyclic interval from immediately after the first end until immediately before the other end. For example, in Figure 1 if we take the chord from 4 to 7 as having 4 for its first end, then the inside of that chord includes the points 5 and 6, and so the chord from 5 to 6 is inside the chord from 4 to 7 for this choice of first end.

Definition 3. Let $\mathcal{B}_{n,m}$ be the set of partial rooted chord diagrams with n chords on m points with the following additional information:

- each chord is coloured, blue, red, or black,
- each blue and red chord has a designated first end,

and with the properties:

- the inside of each red or blue chord has every point in a black chord and no black chords exist which are not inside a red or a blue chord,
- the chord diagram is noncrossing.

Let

$$\mathcal{B} = \bigcup_{\substack{m \geq 2n \\ n \geq 0}} \mathcal{B}_{n,m}.$$

For an example of the definition see Figure 2.

There are a few things to observe about this somewhat unwieldy definition. First note that no red or blue chord can be in the inside of another red or blue chord.

Because of this, generically the information of which end of a red or blue chord is not extra information. If there exists at least one point which is not in any chord, this point must be on the outside of all red and blue chords and so unambiguously defines the first end of each red and blue chord. Similarly, if there exists a black chord, then this chord determines the inside of the red or blue chord it is contained




FIGURE 2. An example of an element of $\mathcal{B}_{8,12}$. The red and blue chords are also indicated with a heavier weight, and the first ends are circled.

by and is outside every other red or blue chord, hence defining the first end of each red and blue chord. Finally, if there is more than one red or blue chord, then since none can be inside another, the dual tree they define must be a star with one leaf for each red or blue chord, and a star with at least two leaves has a well defined centre vertex. This defines the outside of each red or blue chord and hence their first ends.

Consequently, the only case when the information of the first end cannot be deduced from the rest of the structure is the case when $n = 1$ and $m = 2$. None the less, it seems easier to keep this information in the definition since this way the $n = 1$ and $m = 2$ term does not need tweaking to agree with the physical problem we aim to solve. Additionally, the first end will be a useful notion later, and so given $b \in \mathcal{B}_{n,m}$, define $F(b) \subseteq \{1, 2, \dots, m\}$ be the set of first ends of b and note that the first ends inherit an order from the order $\{1, 2, \dots, m\}$.

Lemma 4. The generating function for \mathcal{B} is

$$B(x, y) = \sum_{\substack{m \geq 2n \\ n \geq 0}} \sum_{b \in \mathcal{B}_{n,m}} x^n y^m = \frac{\frac{y}{\sqrt{1-4xy^2}} + y^2(1+4x)}{1 - y^2(1+4x)}.$$

Proof. This is a fairly standard generating function argument.

Noncrossing completed rooted chord diagrams are well known to be counted by the Catalan numbers, $\text{Cat}_n = \frac{1}{n+1} \binom{2n}{n}$. Let $C(x)$ be the generating series for the Catalan numbers, equivalently for noncrossing rooted chord diagrams with x counting the number of chords. Then $C(xy^2)$ counts noncrossing rooted chord diagrams with x counting number of chords and y counting number of points. Adding a new outer chord around each noncrossing diagram multiplies the generating function by xy^2 . These are precisely the *indecomposable* noncrossing rooted chord diagrams, where here indecomposable means the diagram is nonempty and is not a concatenation of 2 or more nonempty diagrams.

Elements of \mathcal{B} are cyclic arrangements of indecomposable noncrossing rooted chord diagrams with the outer chord coloured red or blue (generating function $2xy^2C(xy^2)$) along with bare points (generating function y). If the root is a bare point, then such objects are formed by the root and a sequence of the building blocks described above, which has generating function

$$\frac{y}{1 - y - 2xy^2C(xy^2)}$$

If the root is a point in one of the indecomposable pieces then we choose a root in that piece and the rest is a sequence of the building blocks; this has generating function

$$\frac{y \frac{d}{dy} 2xy^2 C(xy^2)}{1 - y - 2xy^2 C(xy^2)}$$

Adding these two pieces together, using $C(x) = (1 - \sqrt{1 - 4x})/(2x)$, and rearranging, including rationalizing the denominator, we obtain the form of the generating function in the statement. \square

As is common in enumerative combinatorics, write $[x^n y^m]F(x, y)$ for the coefficient of $x^n y^m$ in $F(x, y)$.

Lemma 5. Let $B(x, y)$ be as in Lemma 4, then

$$[x^{\lfloor \frac{d}{2} \rfloor + 1} y^{2\lfloor \frac{d}{2} \rfloor + 2 + dk}]B(x, y) = \frac{(2dk + 4\lfloor \frac{d}{2} \rfloor + 4)(2dk + 4\lfloor \frac{d}{2} \rfloor) \cdots (2dk + 4)}{(\lfloor \frac{d}{2} \rfloor + 1)!}$$

where subsequent factors in the numerator differ by 4, and this is the ratio of gamma factors in the $C_i^{(d)}$ after scaling by $2^{2\lfloor \frac{d}{2} \rfloor + 2}$, namely

$$[x^{\lfloor \frac{d}{2} \rfloor + 1} y^{2\lfloor \frac{d}{2} \rfloor + 2 + dk}]B(x, y) = \begin{cases} 2^{d+2} \frac{\Gamma(\frac{d}{2}(k+1) + 2)}{\Gamma(\frac{d}{2} + 2) \Gamma(1 + \frac{dk}{2})} & \text{for even } d \\ 2^{d+1} \frac{\Gamma(\frac{d}{2}(k+1) + \frac{3}{2})}{\Gamma(\frac{d+3}{2}) \Gamma(1 + \frac{dk}{2})} & \text{for odd } d \end{cases}.$$

Proof. It is convenient to write out the odd and even d cases separately, so what we want to show is for even d

$$[x^{\frac{d}{2} + 1} y^{2+d+dk}]B(x, y) = \frac{(2dk + 2d + 4)(2dk + 2d) \cdots (2dk + 4)}{(\frac{d}{2} + 1)!} = 2^{d+2} \frac{\Gamma(\frac{d}{2}(k+1) + 2)}{\Gamma(\frac{d}{2} + 2) \Gamma(1 + \frac{dk}{2})}$$

and for odd d

$$[x^{\frac{d+1}{2}} y^{1+d+dk}]B(x, y) = \frac{(2dk + 2d + 2)(2dk + 2d - 2) \cdots (2dk + 4)}{(\frac{d+1}{2})!} = 2^{d+1} \frac{\Gamma(\frac{d}{2}(k+1) + \frac{3}{2})}{\Gamma(\frac{d+3}{2}) \Gamma(1 + \frac{dk}{2})}.$$

For even d , the arguments to the gamma functions are all integers so we can directly translate to factorials obtaining

$$\begin{aligned} 2^{d+2} \frac{\Gamma(\frac{d}{2}(k+1) + 2)}{\Gamma(\frac{d}{2} + 2) \Gamma(1 + \frac{dk}{2})} &= 2^{d+2} \binom{\frac{d}{2}k + \frac{d}{2} + 1}{\frac{d}{2} + 1} \\ &= 2^{d+2} \frac{(\frac{d}{2}k + \frac{d}{2} + 1)(\frac{d}{2}k + \frac{d}{2}) \cdots (\frac{d}{2}k + 1)}{\frac{d}{2} + 1} \\ &= \frac{(2dk + 2d + 4)(2dk + 2d) \cdots (2dk + 4)}{(\frac{d}{2} + 1)!} \end{aligned}$$

For odd d and even k , the arguments to the gamma functions are again integers and so similarly we obtain

$$2^{d+1} \frac{\Gamma(\frac{d}{2}(k+1) + \frac{3}{2})}{\Gamma(\frac{d+3}{2}) \Gamma(1 + \frac{dk}{2})} = 2^{d+1} \binom{\frac{d+1}{2} + dk}{\frac{d+1}{2}} = \frac{(2dk + 2d + 2)(2dk + 2d - 2) \cdots (2dk + 4)}{(\frac{d+1}{2})!}.$$

When d and k are both odd use the formula for the gamma function at half integers in terms of the double factorial: $\Gamma(\frac{1}{2} + n) = \frac{(2n-1)!!}{2^n} \sqrt{\pi}$. With this we obtain for d and k both odd

$$\begin{aligned} 2^{d+1} \frac{\Gamma(\frac{d}{2}(k+1) + \frac{3}{2})}{\Gamma(\frac{d+3}{2}) \Gamma(1 + \frac{dk}{2})} &= \frac{2^{\frac{d+1}{2}} (dk+d+1)!!}{(\frac{d+1}{2})! (dk)!!} \\ &= \frac{2^{\frac{d+1}{2}} (dk+d+1)(dk+d-1) \cdots (dk+2)}{(\frac{d+1}{2})!} \\ &= \frac{(2dk+2d+2)(2dk+2d-2) \cdots (2dk+4)}{(\frac{d+1}{2})!}. \end{aligned}$$

It remains now to do the coefficient extraction on $B(x, y)$. Note that the only place an odd power of y appears in $B(x, y)$ is in the term $y/\sqrt{1-4xy^2}$ in the numerator. Consider first the coefficient for even powers of y . For $i > 0$, $n \geq 0$ we have

$$[x^n y^{2i}] F(x, y) = [x^n y^{2i}] \frac{y^2(1+4x)}{1-y^2(1+4x)} = [x^n](1+4x)^i = 4^n \binom{i}{n}$$

For d even take $n = \frac{d}{2} + 1$, $i = \frac{d+2+dk}{2}$ and so

$$[x^{\frac{d}{2}+1} y^{d+2+dk}] F(x, y) = 4^{\frac{d}{2}+1} \binom{\frac{d+2+dk}{2}}{\frac{d}{2}+1} = \frac{(2dk+2d+4)(2dk+2d) \cdots (2dk+4)}{(\frac{d}{2}+1)!}$$

as desired, For odd d and even k take $n = \frac{d+1}{2}$, $i = \frac{d+1+dk}{2}$ and so

$$[x^{\frac{d+1}{2}} y^{d+1+dk}] F(x, y) = 4^{\frac{d+1}{2}} \binom{\frac{d+1+dk}{2}}{\frac{d+1}{2}} = \frac{(2dk+2d+2)(2dk+2d-2) \cdots (2dk+4)}{(\frac{d+1}{2})!}.$$

It remains to consider d and k both odd. In this case the relationship between the powers of x and y is important so let $t = xy^2$. Then for $n, i \geq 0$

$$\begin{aligned} [x^n y^{2i+1}] F(x, y) &= [x^n y^{2i}] \frac{1}{\sqrt{1-4xy^2(1-y^2(1+4x))}} \\ &= [t^n y^{2j}] \frac{1}{\sqrt{1-4t(1-(y^2+4t))}} \quad \text{where } j = i - n \\ &= [t^n](1-4t)^{-\frac{1}{2}} \sum_{k \geq j} \binom{k}{j} (4t)^{k-j} \\ &= [t^n](1-4t)^{-\frac{1}{2}} \sum_{\ell \geq 0} \binom{\ell+j}{\ell} (4t)^\ell \\ &= [t^n](1-4t)^{-\frac{1}{2}} (1-4t)^{-j-1} \\ &= (-4)^n \binom{-j-\frac{3}{2}}{n} \\ &= \frac{(4j+6)(4j+10) \cdots (4j+4n+2)}{n!} \end{aligned}$$

For d and k both odd we are interested in the case $n = \frac{d+1}{2}$ and $i = d + dk$ so $j = \frac{dk-1}{2}$, giving

$$[x^{\frac{d+1}{2}} y^{d+1+dk}]F(x, y) = \frac{(2dk + 2d + 2)(2dk + 2d - 2) \cdots (2dk + 4)}{(\frac{d+1}{2})!}$$

and completing the proof. \square

What we have accomplished in this phase is to define some objects, specifically the partial chord diagrams of \mathcal{B} , which are counted by the gamma function factor in the expressions for $C_i^{(d)}$ after scaling by $2^{2\lfloor \frac{d}{2} \rfloor + 2}$. Note that we only make use of elements of \mathcal{B} for certain choices of parameters, so it isn't accurate to say that $B(x, y)$ is the generating function for the gamma factors in the $C_i^{(d)}$ with appropriate scaling, since not all elements of \mathcal{B} are used in this interpretation.

2.2. Phase 2. For phase 2, we now want an interpretation for the $C_i^{(d)}$. Unfortunately, there are signs here. In general subtraction does not have a combinatorial interpretation since it depends on the sets of objects counted by the numbers on either side of the subtraction sign having an appropriate containment relation. Fortunately in this case, we can give a suitable interpretation where the containment makes sense.

Next note that in odd dimension the $C_i^{(d)}$ are not in general integers, so we will instead give an interpretation for $2^{2\lfloor \frac{d}{2} \rfloor + 2} C_i^{(d)}$ and will comment at the end of this section on how we can remove one power of 2 in all cases and in Section 3 on how we can remove all of the powers of two in even dimensions.

This phase is where we'll use the first ends of the red and blue chords, even though that information is redundant in almost all elements of \mathcal{B} . Recall that for $b \in \mathcal{B}$, $F(b)$ is the set of first ends and this set inherits a total order. Write $F(b) = \{e_1, e_2, \dots\}$ in this order. Given a red or blue chord c of some $b \in \mathcal{B}$, say c has *width* j if c has $j - 1$ black chords inside it. Write $w(e_j)$ for the width of the red or blue chord with first end e_j .

Definition 6. Let $\mathcal{B}_{n,m,c,d}$ be the set of elements of $\mathcal{B}_{n,m}$ with fewer than c consecutive bare points immediately preceding e_j for all j with $\sum_{\ell=1}^{j-1} w(e_\ell) < d$.

In the main proof, we'll be inserting bare points before the first ends of red and blue chords. If c has width j then c has j *insertion places* before its first end. We will view these j insertion places as distinct, so for a chord of width 2, inserting three bare points into its first insertion place and inserting three bare points into its second insertion place will be viewed as different acts of insertion even though the resulting partial chord diagrams are the same.

With this notion of insertion place in hand, another way to phrase the condition defining $\mathcal{B}_{n,m,c,d}$ is that there are fewer than c consecutive bare points immediately preceding each first end that corresponds to one of the the first d insertion places.

The $\mathcal{B}_{n,m,c,d}$ with appropriate parameters are the objects which the $2^{2\lfloor \frac{d}{2} \rfloor + 2} C_i^{(d)}$ count.

Theorem 7.

$$2^{2\lfloor \frac{d}{2} \rfloor + 2} C_i^{(d)} = (-1)^{i-1} |\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1), d, i-1}|.$$

Proof. Let's first pull out the sign in the expression for $C_i^{(d)}$, do a trivial rewriting of the binomial coefficients so that they're exactly in the form in which we'll interpret them, and apply Lemma 5.

$$\begin{aligned} 2^{2\lfloor \frac{d}{2} \rfloor + 2} C_i^{(d)} &= \begin{cases} 2^{2\lfloor \frac{d}{2} \rfloor + 2} (-1)^{i-1} \sum_{k=0}^{i-1} \binom{i-1}{k} (-1)^{k-i+1} \frac{\Gamma(\frac{d}{2}(k+1) + 2)}{\Gamma(\frac{d}{2} + 2) \Gamma(1 + \frac{dk}{2})} & \text{for even } d \\ 2^{2\lfloor \frac{d}{2} \rfloor + 2} (-1)^{i-1} \sum_{k=0}^{i-1} \binom{i-1}{k} (-1)^{k-i+1} \frac{\Gamma(\frac{d}{2}(k+1) + \frac{3}{2})}{\Gamma(\frac{d+3}{2}) \Gamma(1 + \frac{dk}{2})} & \text{for odd } d \end{cases} \\ &= (-1)^{i-1} \sum_{k=0}^{i-1} \binom{i-1}{i-1-k} (-1)^{k-i+1} |\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + dk}| \end{aligned}$$

So what we need to prove is

$$|\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1), d, i-1}| = \sum_{k=0}^{i-1} \binom{i-1}{i-1-k} (-1)^{k-i+1} |\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + dk}|$$

To that end, fix d and i , and consider an element $b \in \mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + dk}$ for some $0 \leq k \leq i-1$.

Observe that the sum of the widths of all red and blue chords is just the total number of chords of all three colours, so b has $\lfloor \frac{d}{2} \rfloor + 1$ insertion places total. Consider the multiset $M_{b,k}$ of partial chord diagrams resulting from choosing $i-1-k$ of the first $i-1$ insertion places of b and inserting d bare points into each of those insertion places. Each way of inserting adds a total of $d(i-1-k)$ points to b so each element of $M_{b,k}$ is an element of $\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + dk + d(i-1-k)} = \mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1)}$.

Thus we have

$$\sum_{k=0}^{i-1} \binom{i-1}{i-1-k} (-1)^{k-i+1} |\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + dk}| = \sum_{k=0}^{i-1} (-1)^{k-i+1} |M_k|,$$

where

$$M_k = \bigsqcup_{b \in \mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + dk}} M_{b,k}$$

is the multiset union of the $M_{b,k}$. Note that all the M_k are multisets whose elements are elements of the set $\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1)}$.

With these insertions all our objects live in the correct set. We only need to check that the signs and multiplicities conspire to cancel all elements of $\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1)}$ which are not in $\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1), d, i-1}$ and to give all elements that are in $\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1), d, i-1}$ exactly once.

The second part is simpler to check. All the insertions were of d bare points into $i-1-k$ of the first $i-1$ insertion places, so the only way to obtain an element of $\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1), d, i-1}$ is from the $k = i-1$ term (where there is no insertion) and that term is $(-1)^0 |\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1)}|$ which contributes each element of $\mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1), d, i-1}$ exactly once.

For the first part take $c \in \mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1)}$, $c \notin \mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1), d, i-1}$. By definition c has d or more bare points before at least one of its first $i-1$ insertion places, so c appears in at least one M_k for some $k < i-1$. Let ℓ be the minimum k for which c appears and take $b \in \mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d\ell}$ so that $c \in M_{b,\ell}$.

Now, we claim that b is the unique $b \in \mathcal{B}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d\ell}$ so that $c \in M_{b,\ell}$. If not then suppose b and b' were two such partial chord diagrams, then b and b' differ only in their bare points, so build b'' so as to agree with b and b' except that whenever b and b' have a different number of bare points before a given red or blue chord, take the minimum of these numbers as the number of bare points before this chord in b'' . In whichever of b or b' achieved the minimum at this point, these insertions were all valid, so doing so across b'' we get a valid insertion pattern giving c but b'' is smaller than b or b' contradicting the minimality of ℓ and proving the claim.

Write the first ends of b in their inherited order, $F(b) = \{e_1, e_2, \dots\}$ and let w_j be the number of the first $i - 1$ insertion places of b that are before e_j . Then for some t , w_1, w_2, \dots, w_{t-1} are the widths of their respective chords, $0 < w_t$ is bounded above by the width of its chord, and $w_{t+1} = w_{t+2} = \dots = 0$. Let i_1, \dots, i_t be the number of insertion places before e_1, \dots, e_t respectively that must be chosen in order to obtain c from b . Thus the multiplicity of c in $M_{b,\ell}$ is

$$\prod_{j=1}^t \binom{w_j}{i_j}.$$

Now for each $1 \leq j \leq t$ and any $0 \leq i'_j \leq i_j$ we can add $d(i_j - i'_j)$ base points to the partial chord diagram immediately before e_j giving a larger partial chord diagram for which i'_j insertion places before e_j still need to be inserted into in order to obtain c . Every such choice is counted separately in the multisets M_k since it comes from a different pair of a chord diagram and a pattern of insertions and every appearance of c in the M_k occurs in this way. The total signed contribution of these copies of c is

$$\sum_{\substack{0 \leq i'_1 \leq i_1 \\ \vdots \\ 0 \leq i'_t \leq i_t}} \prod_{j=1}^t \binom{w_j}{i'_j} (-1)^{i'_1 + i'_2 + \dots + i'_t} = \prod_{j=1}^t \sum_{0 \leq i'_j \leq i_j} \binom{w_j}{i'_j} (-1)^{i'_j} = 0$$

by the binomial theorem, which completes the proof of the result. \square

It would be nice not to have the initial power of 2 multiplying the $C_i^{(d)}$. In general, we do need some powers of 2 since the $C_i^{(d)}$ do sometimes have denominators. In the next section we'll look at removing the powers of 2 in the even dimensional case, but first observe that since every element of \mathcal{B} has at least one red or blue chord, at least one power of 2 is easy to remove in all cases.

Corollary 8. Let $\tilde{\mathcal{B}}_{n,m,a,b}$ be $\mathcal{B}_{n,m,a,b}$ where we identify partial chord diagrams that differ only by swapping all red chords for blue chords and vice versa. Then

$$2^{2\lfloor \frac{d}{2} \rfloor + 1} C_i^{(d)} = (-1)^{i-1} |\tilde{\mathcal{B}}_{\lfloor \frac{d}{2} \rfloor + 1, 2\lfloor \frac{d}{2} \rfloor + 2 + d(i-1), d, i-1}|.$$

3. SIMPLIFICATIONS IN EVEN DIMENSION

In even dimension the number of points in the partial chord diagram is even. In this situation things can be simplified and the power of 2 can be absorbed. We'll use the following map from elements of \mathcal{B} to binary strings.

Definition 9. Given $b \in \mathcal{B}_{n,2j}$, define $s(b)$ to be the binary string of length j whose ℓ th entry is 0 if point $2\ell + 1$ is bare in b and 1 otherwise.




FIGURE 3. An example of the sequence of indecomposable pieces making up a block of 1s in $s(b)$. The arrows indicate the points which directly contribute bits to $s(b)$.

Clearly the map s loses a lot of information about b . The next claim is that s is a 4^n -to-1 map, where n is the number of chords, and hence will absorb the power of 2 that we had in the previous section.

Lemma 10. For $b \in \mathcal{B}_{n,2j}$, $s(b)$ has exactly n 1s.

For every binary string w of length j with n 1s, $|s^{-1}(w)| = 4^n$.

Proof. For the first point, note that noncrossing complete rooted chord diagrams have every chord with one even indexed end and one odd indexed end. Since b is on an even number of points, the parity of the indices of the points is well defined cyclically. Further b is made of a cyclic sequence of complete noncrossing chord diagrams and bare points and hence the indexing of each complete noncrossing segment is changed by a constant shift. Thus b also has every chord with one even indexed end and one odd indexed end. Therefore every chord of b has exactly one end that becomes a 1 under s and no other 1s occur in $s(b)$ giving that $s(b)$ has exactly n 1s.

For the second part, consider how a block of 1s can appear in $s(b)$. Suppose first that there is at least one 0 in $s(b)$ and consider the blocks in $s(b)$ cyclically. Take a block of 1s in $s(b)$. There is some 0 before the block, corresponding to a bare point in b , then in b there is a sequence of indecomposable noncrossing chord diagrams with the outer chord red or blue and where the indexed points are in the even positions, then another bare point, then a sequence of indecomposable noncrossing chord diagrams with the outer chord red or blue and where the indexed points are in the odd positions, then another bare point, as illustrated in Figure 3. Either sequence of indecomposable diagrams may be empty, but both may not be empty. The first and last bare point may be the same point as we are working cyclically. Each block of 1s where $s(b)$ contains at least one 0 appears in this way, and each such configuration in b gives a block of 1s.

Again using $C(x)$ for the generating function for Catalan numbers, by standard generating function arguments we get that the generating function for a block of 1s is

$$\left(\frac{1}{1 - 2xC(x)} \right) \left(\frac{1}{1 - 2xC(x)} \right) - 1 = \frac{1}{\sqrt{1 - 4x}} \frac{1}{\sqrt{1 - 4x}} - 1 = \frac{4x}{1 - 4x}.$$

Extracting coefficients we get that every block of t 1s occurs 4^t times, so arguing similarly for each block of 1s, we get that each binary string w with at least one 0 occurs 4^n times, as desired.

It remains to consider the case that $s(b)$ has no 0s. In this case we simply have a cyclic sequence of indecomposable noncrossing chord diagrams with the outer chord coloured red or blue. As in Lemma 4 we can obtain the generating series for these, by choosing a point on one indecomposable diagram as the root and then taking a sequence of others. Note that since x counts the chords and each chord has two points, choosing a point is done by the operator $2x \frac{d}{dx}$ in this case. This gives the generating function

$$\frac{2x \frac{d}{dx} 2xC(x)}{1 - 2xC(x)} = \frac{4x}{\sqrt{1-4x}} \frac{1}{\sqrt{1-4x}} = \frac{4x}{1-4x}$$

giving this binary string also occurs 4^n times and completing the proof. \square

The previous lemma tells us that in the even dimensional case we can directly reinterpret the phase 1 numbers, that is the ratio of gamma functions appearing in the $C_i^{(d)}$ for even d , as the number of binary strings of length $d/2+1+dk/2$ and with $d/2+1$ 1s. We could also see that fact more directly by writing the ratio of gamma functions as a binomial coefficient in this case, but the lemma gives us the direct connection between the phase 1 construction and these strings, hence allowing us to carry over everything additional we know from the phase 1 construction.

From here we can redo the phase 2 argument adding 0s and checking the cancellation occurs. There is one important detail. Since we no longer know how each block of 1s came from a sequence of indecomposable noncrossing chord diagrams, we no longer know where the first ends of the red or blue chords were. However, we can now simply take an insertion place before every 1. The same argument for the cancellation holds, but now is even easier since we no longer need to worry about multiplicities from insertions into different places giving the same result. Taking this all together we obtain the following theorem.

Theorem 11. For d even $(-1)^{i-1}C_i^{(d)}$ is the number of binary strings of length $\frac{d}{2} + 1$ where there are fewer than $\frac{d}{2}$ 0s immediately before any of the first $i - 1$ 1s.

Proof. Use Lemma 10 and then argue as the the proof of Theorem 7 with insertion places before the first $i - 1$ 1s and with insertions of $\frac{d}{2}$ 0s. \square

These objects are arguably mildly more natural if we interpret the binary strings as lattice paths starting at the origin and where 0 becomes a step one unit up and 1 becomes a step one unit right. Then rephrasing we obtain the following corollary.

Corollary 12. For d even $(1)^{i-1}C_i^{(d)}$ is the number of lattice walks from $(0, 0)$ to $(\frac{d}{2} + 1, \frac{d(i-1)}{d})$ and using steps \rightarrow and \uparrow with the additional property that there are fewer than $\frac{d}{2}$ up steps with x -coordinate j for $0 \leq j \leq i - 2$.

Many other interpretations would be similarly possible.

4. DISCUSSION

We gave a combinatorial interpretation that captures the cancellations in the expression for the coefficients of the causal set d'Alembertian given in [3]. While this does not obviously yield a closed form expression, hopefully the reader finds this more satisfying hence in some sense an answer to Glaser's request for a more aesthetically satisfying expression. Combinatorial interpretations can be potentially useful even when they don't give nice formulas, as they can show structure that may

not be apparent otherwise and are well suited to proving identities or properties of the counts by combinatorial means.

From a combinatorial perspective it is tempting to rewrite the formulas for $B^{(d)}$ by pulling out a factor of β_d giving

$$B^{(d)}\phi(x) = \frac{\beta_d}{\ell^2} \left(\frac{\alpha_d}{\beta_d} \phi(x) + \sum_{i=1}^{\lfloor \frac{d}{2} \rfloor + 2} c_i^{(d)} \sum_{y \in L_i} \phi(y) \right).$$

We know [3] that

$$\begin{aligned} \frac{\alpha_d}{\beta_d} &= \begin{cases} -\frac{\Gamma(d)}{\Gamma(\frac{d}{2}+2)\Gamma(\frac{d}{2})} & \text{for even } d \\ -\frac{2^{d-1}}{d+1} & \text{for odd } d \end{cases} \\ &= \begin{cases} -\frac{1}{2} \text{Cat}_{\frac{d}{2}} & \text{for even } d \\ -\frac{2^{d-1}}{d+1} & \text{for odd } d \end{cases} \end{aligned}$$

where Cat_n is the n th Catalan number. This suggests some $i = 0$ version of our objects which are counted by Catalan numbers in the even dimensional case and by $\frac{1}{d+1}$ in the odd dimensional case, with scaling by powers of 2 as necessary. It is not clear how to fit this in to the picture described above. Likely an alternate interpretation would be needed.

Finally, one may ask what physical use these interpretations are. It is possible they are of no physical use, but it is worth noting that they do have some physically nice properties. For instance they do not have a dramatically different form in even and odd dimension. Ultimately one would like to have a way of understanding a causal set action without knowing the dimension in advance, so while the dimension is still an important part of the input here, at least how the behaviour changes with d is somewhat controlled. Additionally, since these are the coefficients of the number of intervals of size i , one could hope for some way to see them as the number of ways to do some specific thing in d dimensions with these intervals, or something similar.

REFERENCES

- [1] Dionigi M. T. Benincasa and Fay Dowker. Scalar curvature of a causal set. *Phys. Rev. Lett.*, 104:181301, May 2010.
- [2] Fay Dowker and L. Glaser. Causal set d'Alembertians for various dimensions. *Classical and Quantum Gravity*, 30(19):195016, sep 2013.
- [3] L. Glaser. A closed form expression for the causal set d'Alembertian. *Classical and Quantum Gravity*, 31(9):095007, apr 2014.
- [4] Bombelli L, Lee J, Meyer D, and Sorkin R. Space-time as a causal set. *Phys Rev Lett*, 59:521–524, 1987.
- [5] Rafael D. Sorkin. Does locality fail at intermediate length scales? In Daniele Oriti, editor, *Approaches to Quantum Gravity: Toward a New Understanding of Space, Time and Matter*, page 26–43. Cambridge University Press, 2009.
- [6] Sumati Surya. The causal set approach to quantum gravity. arXiv:1903.11544.