

The gluino condensate of large- N SUSY Yang–Mills

**Claudio Bonanno,^{a,*} Pietro Butti,^{a,b} Margarita García Pérez,^a
Antonio González-Arroyo,^{a,c} Ken-Ichi Ishikawa^{d,e} and Masanori Okawa^e**

^a*Instituto de Física Teórica UAM-CSIC, Calle Nicolás Cabrera 13-15,
Universidad Autónoma de Madrid, Cantoblanco, E-28049 Madrid, Spain*

^b*Departamento de Física Teórica, Facultad de Ciencias and
Centro de Astropartículas y Física de Altas Energías (CAPA),
Universidad de Zaragoza, Calle Pedro Cerbuna 12, E-50009, Zaragoza, Spain*

^c*Departamento de Física Teórica, Universidad Autónoma de Madrid,
Módulo 15, Cantoblanco, E-28049 Madrid, Spain*

^d*Core of Research for the Energetic Universe,
Graduate School of Advanced Science and Engineering,
Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan*

^e*Graduate School of Advanced Science and Engineering, Hiroshima University,
Higashi-Hiroshima, Hiroshima 739-8526, Japan*

*E-mail: claudio.bonanno@csic.es, pbutti@unizar.es,
margarita.garcia@csic.es, antonio.gonzalez-arroyo@uam.es,
ishikawa@theo.phys.sci.hiroshima-u.ac.jp, okawa@hiroshima-u.ac.jp*

We present the first lattice determination of the SUSY $SU(N)$ Yang–Mills gluino condensate at large N . We exploit large- N twisted volume reduction, and present two determinations based on the Banks–Casher relation and on a Gell–Mann–Oakes–Renner-like formula, both giving perfectly compatible results. By expressing the lattice results in the Novikov–Shifman–Vainshtein–Zakharov scheme, we are able for the first time to compare lattice and analytical computations, resolving a 40-year-long debate about the actual value and N -dependence of the gluino condensate.

*The 41st International Symposium on Lattice Field Theory (LATTICE2024)
28 July – 3 August 2024
Liverpool, UK*

*Speaker

1. Introduction

The $\mathcal{N} = 1$ supersymmetric (SUSY) $SU(N)$ Yang–Mills theory consists of a bosonic particle, the gluon, and its fermionic super-partner, the gluino. These particles are described in terms of a $SU(N)$ gauge field coupled to 1 massless adjoint Majorana field. The SUSY Yang–Mills (SYM) action enjoys a global $U(1)_V \otimes U(1)_A$ flavor chiral symmetry. At the quantum level, the vector symmetry is exactly realized à la Wigner–Weyl, while the axial sub-group is anomalously and spontaneously broken with the following pattern:

$$U(1)_A \xrightarrow[\text{Anomaly}]{} \mathbb{Z}_{2N} \xrightarrow[\text{Spontaneous}]{} \mathbb{Z}_2. \quad (1)$$

The theory is thus expected to be characterized by a non-vanishing gluino condensate $\langle \text{Tr} \lambda^2 \rangle$ due to the spontaneous breaking of the residual discrete \mathbb{Z}_{2N} symmetry.

The gluino condensate can be calculated exactly using analytic methods [1–4]. However, its actual value is the subject of a debate [5]. Two instanton-based calculations performed, respectively, in the Strong-Coupling (SC) [1–3] and in the Weak-Coupling (WC) [4] regimes yielded the following for the Renormalization-Group-Invariant (RGI) condensate:

$$\Sigma_{\text{RGI}} \equiv \frac{1}{(4\pi)^2 b_0 N} |\langle \text{Tr} \lambda^2 \rangle| = \begin{cases} 2e \Lambda_{\text{NSVZ}}^3 / N & \text{(SC)}, \\ \Lambda_{\text{NSVZ}}^3 & \text{(WC)}. \end{cases} \quad (2)$$

Here Λ_{NSVZ} is the dynamically-generated scale of SYM computed in the Novikov–Shifman–Vainshtein–Zakharov (NSVZ) scheme [1, 6]. Adopting the standard QCD convention (see [7] to trivially map it to the SUSY one), it reads:

$$\Lambda_{\text{NSVZ}}^3 = \frac{\mu^3}{b_0 \lambda_t^{(\text{NSVZ})}(\mu)} \exp\left(\frac{-8\pi^2}{\lambda_t^{(\text{NSVZ})}(\mu)}\right), \quad (3)$$

with $\lambda_t^{(\text{NSVZ})}(\mu)$ the renormalized 't Hooft coupling in the NSVZ scheme, and $b_0 = 3/(4\pi)^2$ the first universal coefficient of the SYM β -function. Recently, an alternative calculation of the gluino condensate, based on the use of fractional instantons [8–10] has been carried out in [11], based on ideas put forward in [12–15]. The authors found $\Sigma_{\text{RGI}} = 2\Lambda_{\text{NSVZ}}^3$ for gauge group $SU(2)$ and argue that $2 \rightarrow N$ for $SU(N)$, thus yielding yet another result.

This proceeding reports on the main results of our paper [16], where we performed the first non-perturbative first-principles calculation of the gluino condensate of large- N SYM theory via numerical Monte Carlo simulations of the lattice-discretized theory. Despite impressive recent progress in lattice simulations of SUSY theories [17–23], the lattice literature on this topic is quite limited [20, 21, 24, 25], and our paper presents the first comparison between numerical and analytical results.

We here anticipate that our value and N -dependence for the gluino condensate agree with the WC prediction. After the publication of our paper [16], the new study [26] from the same authors of [11] appeared, reporting a value and N -dependence for $\Sigma_{\text{RGI}}/\Lambda_{\text{NSVZ}}^3$ in agreement with ours.

2. From the lattice to the NSVZ scheme

To compare numerical and analytical results, we must compute two quantities: the RGI gluino condensate Σ_{RGI} and the SUSY scale Λ_{NSVZ} . We review their definition in the following.

2.1 The dynamically-generated scale

The Λ -parameter is a scheme-dependent quantity defined as the integration constant of the Callan–Symanzik equation for the renormalized 't Hooft coupling $\lambda_t^{(s)}$, expressed via the β -function:

$$\beta_s \left(\lambda_t^{(s)} \right) = \frac{d \lambda_t^{(s)}(\mu)}{d \log(\mu^2)}, \quad (4)$$

$$\Lambda_s = \mu \left[b_0 \lambda_t^{(s)}(\mu) \right]^{\frac{-b_1}{2b_0^2}} \exp \left(\frac{-1}{2b_0 \lambda_t^{(s)}(\mu)} \right) \exp \left[- \int_0^{\lambda_t^{(s)}(\mu)} dx \left(\frac{1}{2\beta_s(x)} + \frac{1}{2b_0 x^2} - \frac{b_1}{2b_0^2 x} \right) \right], \quad (5)$$

with $b_0 = 3/(4\pi)^2$, $b_1 = 6/(4\pi)^4$ the first two universal (i.e., scheme-independent) coefficients in the perturbative expansion of the β -function. In the NSVZ scheme, the exact β -function

$$\beta_{\text{NSVZ}} \left(\lambda_t^{(\text{NSVZ})} \right) = - \frac{b_0 \lambda_t^{(\text{NSVZ})^2}}{1 - \frac{b_1}{b_0} \lambda_t^{(\text{NSVZ})}} \quad (6)$$

yields the exact expression for Λ_{NSVZ} in Eq. (3).

2.2 The RGI gluino condensate

The Callan–Symanzik equation for the renormalized gluino mass $m_R^{(s)}$, expressed via the anomalous dimension τ -function, also features a *scheme-independent* integration constant:

$$\tau_s \left(\lambda_t^{(s)} \right) = \frac{d \log \left(m_R^{(s)}(\mu) \right)}{d \log(\mu)}. \quad (7)$$

$$m_{\text{RGI}} = \tilde{\mathcal{A}} m_R^{(s)}(\mu) \left[2b_0 \lambda_t^{(s)}(\mu) \right]^{-\frac{d_0}{2b_0}} \exp \left[- \int_0^{\lambda_t^{(s)}(\mu)} dx \left(\frac{\tau_s(x)}{2\beta_s(x)} - \frac{1}{x} \right) \right], \quad (8)$$

with $d_0 = 2b_0$ the first universal (i.e., scheme-independent) coefficient of the perturbative expansion of the τ -function, and $\tilde{\mathcal{A}}$ an arbitrary numerical constant. In the NSVZ scheme, also the τ -function is known exactly [27]:

$$\frac{\tau_{\text{NSVZ}}(x)}{2\beta_{\text{NSVZ}}(x)} = \frac{1}{x(1 - b_1 x/b_0)}. \quad (9)$$

Since the product $\Sigma_R^{(s)}(\mu) m_R^{(s)}(\mu)$ is RGI—with $\Sigma_R^{(s)}(\mu)$ the usual scheme-/scale-dependent fermion condensate—the RGI gluino condensate can be easily defined as [28, 29]:

$$\Sigma_{\text{RGI}} = \mathcal{A} \Sigma_R^{(s)}(\mu) \left[2b_0 \lambda_t^{(s)}(\mu) \right]^{\frac{d_0}{2b_0}} \exp \left[\int_0^{\lambda_t^{(s)}(\mu)} dx \left(\frac{\tau_s(x)}{2\beta_s(x)} - \frac{1}{x} \right) \right]. \quad (10)$$

Once the exact β and τ functions of the NSVZ scheme are inserted in Eq. (10), the arbitrary constant \mathcal{A} must be chosen as $\mathcal{A} = \frac{8\pi^2}{9N^2}$ in order for Eq. (10) to reproduce the known exact expression for the RGI condensate in terms of $\Sigma_R^{(\text{NSVZ})}(\mu)$ [6, 30]:

$$\Sigma_{\text{RGI}} = \frac{1}{(4\pi)^2 b_0 N} |\langle \text{Tr} \lambda^2 \rangle|, \quad |\langle \text{Tr} \lambda^2 \rangle| = \frac{\lambda_t^{(\text{NSVZ})}(\mu)}{N \left[1 - \lambda_t^{(\text{NSVZ})}(\mu)/(8\pi^2) \right]} \Sigma_R^{(\text{NSVZ})}(\mu). \quad (11)$$

3. Lattice setup

Our large- N calculation exploits large- N twisted volume reduction. By virtue of the well-known dynamical equivalence between space-time and color degrees of freedom unveiled in the pioneering paper of Eguchi and Kawai [31], it is possible to simulate $4d$ large- N gauge theories as a matrix model which can be interpreted as standard lattice gauge theory defined on a reduced one-point space-time lattice with twisted boundary conditions [32–34].

Reduced models have been extensively used in the last decade to study several large- N gauge theories [35–38], including also theories with dynamical adjoint fermions [39, 40]. This enabled in Ref. [41] the study of large- N SYM on the lattice using twisted volume reduction and the lattice techniques developed by the DESY–Jena–Regensburg–Münster collaboration, see Refs. [17–19]. More precisely, in Ref. [41] we generated several gauge configurations for a few values of inverse bare 't Hooft coupling $b = 1/\lambda_L$ and N according the following setup.

- Simulations are performed using a dynamical massive gluino, discretized using Wilson fermions. The gluino mass is controlled via the Wilson hopping parameter κ .
- The mild sign problem introduced by the non-positivity of the Pfaffian of the lattice Wilson–Dirac operator is bypassed via sign-quenched simulations. Since no occurrence of negative signs of $\text{Pf}(CD_W)$ was observed in [41], no reweighting was needed.
- The lattice regularization and the non-zero gluino mass explicitly break SUSY. According to the Kaplan–Curci–Veneziano prescription, the SUSY-restoration limit is achieved as the joint continuum and chiral (massless gluino) limit [42, 43].
- The massless gluino limit is obtained requiring a vanishing mass for the “adjoint pion”. This is an unphysical particle which is introduced by supplementing SYM with a quenched valence gluino. This approach can be rigorously justified within the theoretical framework of Partially Quenched Chiral Perturbation Theory [44].

The gluino condensate is computed adopting two different methods:

- Banks–Casher (BC) formula [45].

$$\frac{\Sigma_R^{(s)}(\mu)}{2\pi} = \lim_{\lambda \rightarrow 0} \lim_{m_R \rightarrow 0} \lim_{V \rightarrow \infty} \left[\rho_R^{(s)}(\mu) \right] (\lambda_R, m_R). \quad (12)$$

Here ρ_R stands for the spectral density of eigenmodes $i\lambda_R + m_R$ of the massive Dirac operator.

- Gell-Mann–Oakes–Renner (GMOR) relation [44].


$$m_\pi^2 = 2 \frac{\Sigma_R^{(s)}(\mu)}{F_\pi^2} m_R^{(s)}(\mu). \quad (13)$$

This relation involves the non-singlet adjoint pion mass m_π and its decay constant F_π .

4. Results

In this section we summarize the main results of Ref. [16], obtained for $b = 0.340, 0.345, 0.350$ and $N = 169, 289, 361$, using the gauge ensembles generated in Ref. [41]. Scale setting was performed in [41] using gradient flow through the standard reference scale $\sqrt{8t_0}$.

4.1 The gluino condensate from the BC relation



(a) Example of the linear fit to the mode number to extract the slope.

(b) Example of calculation of the ratio of non-singlet pseudoscalar and scalar renormalization constants.

Figure 1: Numerical results using spectral methods. Figures taken from [16].

The gluino condensate is obtained from the BC formula via the Giusti–Lüscher method [38, 46–50].

- We solved numerically $(\gamma_5 D_w[U]) u_\lambda = \lambda u_\lambda$ for the first $O(100)$ eigenvalues.
- Counting the number of modes below a certain threshold M we obtained the mode number $\langle \nu(M, m) \rangle = \langle \#\lambda \leq M \rangle$.
- The gluino condensate is obtained via (here $V = a^4$ with a the lattice spacing):

$$\Sigma_R = \frac{\pi}{4V} \sqrt{1 - \frac{m_R^2}{M_R^2}} \left[\frac{d \langle \nu_R(M_R) \rangle}{dM_R} \right] \quad \leftarrow \text{slope of } \langle \nu \rangle \text{ obtained from} \quad (14)$$

linear fit close to $M = \langle |\lambda_{\min}| \rangle \simeq m$.

- Renormalization: $\langle \nu_R \rangle = \langle \nu \rangle$, $M_R = M/Z_P$, $m_R = m/Z_S^{(0)} = r_m m/Z_S$, $r_m \equiv Z_S/Z_S^{(0)}$, where Z_S and Z_P are the non-singlet scalar/pseudo-scalar renormalization constants, while $Z_S^{(0)}$ is the singlet scalar renormalization constant.¹

On the lattice we can compute the following quantities:

- The slope of the mode number as a function of the bare threshold M , see Fig. 1a: $s \equiv \frac{d\langle \nu \rangle}{dM}$.
- The RGI ratio of non-singlet renormalization constants Z_P/Z_S using the following ratio of spectral sums [46–49], see Fig. 1b:

$$\left(\frac{Z_P}{Z_S} \right)^2 = \frac{\langle s_P(M) \rangle}{\langle \nu(M) \rangle} \quad s_P(M) \equiv \sum_{|\lambda|, |\lambda'| \leq M} \left| u_\lambda^\dagger \gamma_5 u_{\lambda'} \right|^2. \quad (15)$$

¹In our original paper [16] we overlooked that $r_m = 1 + O(\lambda^2)$ on the lattice due to explicit chiral symmetry breaking of Wilson fermions. We correct it here, with negligible impact on final results, and no change in the conclusions.

- Bare subtracted gluino mass: $am = 1/(2\kappa) - 1/(2\kappa_{\text{crit}}) = am_{\text{R}}Z_{\text{S}}^{(0)} = am_{\text{R}}Z_{\text{S}}/r_{\text{m}}$.

Combining these quantities we obtain:

$$\frac{\Sigma_{\text{R}}}{Z_{\text{S}}} = \frac{Z_{\text{P}}}{Z_{\text{S}}} \frac{\pi}{4V} \sqrt{1 - \left(\frac{Z_{\text{P}}}{Z_{\text{S}}} \frac{m}{M} r_{\text{m}}\right)^2} s. \quad (16)$$

Since we do not have non-perturbative estimates of the scheme-/scale-dependent constant Z_{S} and of the RGI ratio $r_{\text{m}} = Z_{\text{S}}/Z_{\text{S}}^{(0)}$, we relied on 2-loop perturbation theory to compute them in the $\overline{\text{MS}}$ scheme [51]:

$$Z_{\text{S}}^{(\overline{\text{MS}})} \left(\mu = \frac{1}{a}, \lambda_{\text{L}} \right) = 1 - \frac{12.9524103(1)}{(4\pi)^2} \lambda_{\text{L}} - \frac{60.68(10)}{(4\pi)^4} \lambda_{\text{L}}^2 + \mathcal{O}(\lambda_{\text{L}}^3) \quad (17)$$

$$Z_{\text{S}}^{(0)(\overline{\text{MS}})} \left(\mu = \frac{1}{a}, \lambda_{\text{L}} \right) = Z_{\text{S}}^{(\overline{\text{MS}})} \left(\mu = \frac{1}{a}, \lambda_{\text{L}} \right) - \frac{107.76(1)}{(4\pi)^4} \lambda_{\text{L}}^2 + \mathcal{O}(\lambda_{\text{L}}^3). \quad (18)$$

In these expressions, in order to accelerate the convergence of perturbation theory, we used the 1-loop perturbative expression of the lattice bare coupling in terms of improved couplings:

$$\lambda_{\text{L}} = \lambda_{\text{t}}^{(\text{s})} - 2b_0 \left(\lambda_{\text{t}}^{(\text{s})} \right)^2 \log(\Lambda_{\text{s}}/\Lambda_{\text{L}}), \quad (19)$$

with $\lambda_{\text{t}}^{(\text{i})} = 1/(bP)$, $\lambda_{\text{t}}^{(\text{E})} = 8(1-P)$, $\lambda_{\text{t}}^{(\text{E}')} = -8 \log(P)$, and P the expectation value of the plaquette. The ratio of Λ -parameters is given by [52, 53]:

$$\begin{aligned} \frac{\Lambda_{\text{L}}}{\Lambda_{\text{t}}} &= \frac{\Lambda_{\text{L}}}{\Lambda_{\overline{\text{MS}}}} \times 2.7373, \\ \frac{\Lambda_{\text{L}}}{\Lambda_{\text{E}}} &= \frac{\Lambda_{\text{L}}}{\Lambda_{\overline{\text{MS}}}} \times 29.005, \\ \frac{\Lambda_{\text{L}}}{\Lambda_{\text{E}'}} &= \frac{\Lambda_{\text{L}}}{\Lambda_{\overline{\text{MS}}}} \times 5.600, \\ \frac{\Lambda_{\overline{\text{MS}}}}{\Lambda_{\text{L}}} &= 73.467. \end{aligned} \quad (20)$$

4.2 The gluino condensate from the GMOR equation

Using standard techniques, in [41] we computed the pion mass m_{π} from the exponential time decay of the temporal pion-pion correlator, the pion decay constant,

$$\frac{F_{\pi}}{NZ_{\text{A}}} = \frac{1}{\sqrt{2}Nm_{\pi}} \langle 0 | A_4(x=0) | \pi(\vec{p}=0) \rangle, \quad (21)$$

and the PCAC (Partially Conserved Axial Current) gluino mass, related to the renormalized one by:

$$m_{\text{PCAC}} = \frac{Z_{\text{P}}}{Z_{\text{A}}} m_{\text{R}}. \quad (22)$$

Combining these quantities, and the ones computed before, one has:

$$\frac{F_{\pi}}{N} = \frac{F_{\pi}}{NZ_{\text{A}}} \frac{m}{m_{\text{PCAC}}} \frac{Z_{\text{P}}}{Z_{\text{S}}} r_{\text{m}}, \quad (23)$$

$$\frac{\Sigma_{\text{R}}}{Z_{\text{S}}} = \frac{1}{2} F_{\pi}^2 \frac{m_{\pi}^2}{m} \frac{1}{r_{\text{m}}}. \quad (24)$$

4.3 The Λ -parameter in the NSVZ scheme

We computed the Λ -parameter using 2-loop asymptotic scaling, and the 3 improved couplings introduced earlier (here a_χ denotes the lattice spacing extrapolated towards the massless gluino limit):

$$\sqrt{8t_0}\Lambda_{\text{NSVZ}} = \lim_{a_\chi \rightarrow 0} \frac{\Lambda_{\text{NSVZ}}}{\Lambda_{\overline{\text{MS}}}} \frac{\Lambda_{\overline{\text{MS}}}}{\Lambda_s} \frac{\sqrt{8t_0}}{a_\chi} \exp \left\{ -f \left(\lambda_i^{(s)} \right) \right\}, \quad (25)$$

$$f(x) = \frac{1}{2b_0} \left[\frac{1}{x} + \frac{b_1}{b_0} \log(b_0 x) \right].$$

where $\Lambda_{\text{NSVZ}}/\Lambda_{\overline{\text{MS}}} = e^{-1/18}$ [54].

The 3 improved couplings yield results in very good agreement, as shown in Fig. 2. We quote the following final result, obtained imposing a common continuum limit for the 3 different determinations:

$$\sqrt{8t_0}\Lambda_{\text{NSVZ}} = 0.376(25), \quad \sqrt{8t_0}\Lambda_{\overline{\text{MS}}} = 0.397(26). \quad (26)$$




Figure 2: Extrapolation of Λ_{NSVZ} towards the SUSY limit. Figure taken from [16].

4.4 Converting the renormalized condensate into the RGI one

In the previous subsections we described how we computed the scheme-/scale-dependent renormalized chiral condensate $\Sigma_{\text{R}}^{(s)}(\mu)$ in the $\overline{\text{MS}}$ scheme at the lattice scale $\mu = 1/a$, with a the lattice spacing. In order to compare our results with the analytic ones, we need to convert it into an RGI quantity.

To this end, we rely again on 2-loop perturbation theory as follows:

$$\Sigma_{\text{RGI}} = \mathcal{A} 2b_0 \lambda_i^{(\overline{\text{MS}})}(a\mu = 1) \left[1 + \frac{d_1^{(\overline{\text{MS}})} - 2b_1}{2b_0} \lambda_i^{(\overline{\text{MS}})}(a\mu = 1) \right] \Sigma_{\text{R}}^{(\overline{\text{MS}})}(a\mu = 1), \quad \mathcal{A} = 8\pi^2/(9N^2), \quad (27)$$

with $d_1^{(\overline{\text{MS}})} = 32/(4\pi)^4$ [55], and where the renormalized coupling in the $\overline{\text{MS}}$ scheme at the scale $\mu = 1/a$ was computed in 2-loop perturbation theory as follows:

$$2b_0 \lambda_i^{(\overline{\text{MS}})}(a\mu = 1) = -\frac{1}{\log(a\Lambda_{\overline{\text{MS}}})} - \frac{b_1}{2b_0^2} \frac{\log[-2\log(a\Lambda_{\overline{\text{MS}}})]}{\log^2(a\Lambda_{\overline{\text{MS}}})}. \quad (28)$$

The product $a\Lambda_{\overline{\text{MS}}}$ was practically computed as $a\Lambda_{\overline{\text{MS}}} = (a/\sqrt{8t_0}) \times \sqrt{8t_0}\Lambda_{\overline{\text{MS}}}$, with $\sqrt{8t_0}\Lambda_{\overline{\text{MS}}}$ the quantity in Eq. (26).

4.5 The N -dependence of the gluino condensate

We are now ready to compute the RGI gluino condensate from the BC and the GMOR relations in units of Λ_{NSVZ}^3 . Note that Eq. (23) was computed using the value of F_π extrapolated towards the SUSY limit (via a joint chiral/continuum extrapolation), cf. Fig. 3 (left panel):

$$\frac{F_\pi}{N\Lambda_{\text{NSVZ}}} = 0.101(15). \quad (29)$$

All our determinations of Σ_{RGI} are shown in Fig. 3 (central/right panels). Given the displayed N -dependence, our numerical results rule out all but the Weak Coupling (WC) calculation, cf. Eq. (2).




Figure 3: Left panel: calculation of F_π/N in the SUSY limit. Central and right panels: collection of our results for the third root of the RGI condensate in units of the SUSY scale in the NSVZ scheme Λ_{NSVZ} obtained with the GMOR and the BC formulas respectively. Figures adapted from [16].

5. Conclusions: the RGI gluino condensate in the SUSY limit

We extrapolate our results for $\Sigma_{\text{RGI}}^{1/3}/\Lambda_{\text{NSVZ}}$ towards the SUSY limit performing a joint chiral-continuum extrapolation as follows:

$$\left(\frac{\Sigma_{\text{RGI}}^{1/3}}{\Lambda_{\text{NSVZ}}}\right)(a, m_\pi) = \frac{\Sigma_{\text{RGI}}^{1/3}}{\Lambda_{\text{NSVZ}}} + c_1 \frac{a}{\sqrt{8t_0}} + c_2 (8t_0 m_\pi^2). \quad (30)$$

The SUSY-limit extrapolations, shown in Fig. 4, yield:

$$\begin{aligned} \frac{\Sigma_{\text{RGI}}}{\Lambda_{\text{NSVZ}}^3} &= [1.34(18)_{\text{stat}}(13)_{\text{syst}}]^3 \\ &= 2.39(97)_{\text{stat}}(72)_{\text{syst}}, \end{aligned} \quad (\text{BC}). \quad (31)$$

$$\begin{aligned} \frac{\Sigma_{\text{RGI}}}{\Lambda_{\text{NSVZ}}^3} &= [1.21(08)_{\text{stat}}(12)_{\text{syst}}]^3 \\ &= 1.77(35)_{\text{stat}}(53)_{\text{syst}}, \end{aligned} \quad (\text{GMOR}). \quad (32)$$

In the SUSY limit the two determinations from the GMOR and BC relations give agreeing results. Both are compatible with the WC instanton-calculation:

$$\frac{\Sigma_{\text{RGI}}}{\Lambda_{\text{NSVZ}}^3} = 1, \quad (\text{exact NSVZ analytic WC result}). \quad (33)$$

We quote the GMOR determination as our final lattice result, as this is the most precise one:

$$\frac{\Sigma_{\text{RGI}}}{\Lambda_{\text{NSVZ}}^3} = 1.77(35)_{\text{stat}}(53)_{\text{syst}} = 1.77(65), \quad (\text{final lattice result}). \quad (34)$$

We stress that we added a conservative 30% systematic error to our final extrapolations to take into account the perturbative renormalization we employed. This is the dominant source of uncertainty.




Figure 4: Extrapolation towards the SUSY limit of the RGI gluino condensate determined from the BC and the GMOR formulas for the largest value of N explored. Figure adapted from [16].

Acknowledgments

It is a pleasure to thank José L. F. Barbón, Georg Bergner, Gregorio Herdoíza, Nikolai Husung, Haralambos Panagopoulos, Carlos Pena and Erich Poppitz for useful discussions. This work is partially supported by the Spanish Research Agency (Agencia Estatal de Investigación) through the grant IFT Centro de Excelencia Severo Ochoa CEX2020-001007-S, funded by MCIN/AEI/10.13039/501100011033, and by grant PID2021-127526NB-I00, funded by MCIN/AEI/10.13039/501100011033 and by “ERDF A way of making Europe”. We also acknowledge support from the project H2020-MSCAITN-2018-813942 (EuroPLEx) and the EU Horizon 2020 research and innovation programme, STRONG-2020 project, under grant agreement No 824093. P. B. is supported by Grant PGC2022-126078NB-C21 funded by MCIN/AEI/10.13039/501100011033 and “ERDF A way of making Europe”. P. B. also acknowledges support by Grant DGA-FSE grant 2020-E21-17R Aragon Government and the European Union - NextGenerationEU Recovery and Resilience Program on ‘Astrofísica y Física de Altas Energías’ CEFCA-CAPAITAINNOVA. K.-I. I. is supported in part by MEXT as “Feasibility studies for the next-generation computing infrastructure”. M. O. is supported by JSPS KAKENHI Grant Number 21K03576. Numerical calculations have been performed on the Finisterrae III cluster at CESGA (Centro de Supercomputación de Galicia). We have also used computational resources of Oakbridge-CX at the University of Tokyo through the HPCI System Research Project (Project ID: hp230021 and hp220011).

References

- [1] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, *Instanton Effects in Supersymmetric Theories*, *Nucl. Phys. B* **229** (1983) 407.
- [2] G. C. Rossi and G. Veneziano, *Nonperturbative Breakdown of the Nonrenormalization Theorem in Supersymmetric QCD*, *Phys. Lett. B* **138** (1984) 195.
- [3] D. Amati, G. C. Rossi and G. Veneziano, *Instanton Effects in Supersymmetric Gauge Theories*, *Nucl. Phys. B* **249** (1985) 1.
- [4] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, *Supersymmetric Instanton Calculus (Gauge Theories with Matter)*, *Nucl. Phys. B* **260** (1985) 157.
- [5] T. J. Hollowood, V. V. Khoze, W.-J. Lee and M. P. Mattis, *Breakdown of cluster decomposition in instanton calculations of the gluino condensate*, *Nucl. Phys. B* **570** (2000) 241 [[hep-th/9904116](#)].
- [6] M. A. Shifman and A. I. Vainshtein, *Solution of the Anomaly Puzzle in SUSY Gauge Theories and the Wilson Operator Expansion*, *Nucl. Phys. B* **277** (1986) 456.
- [7] A. Armoni, M. Shifman and G. Veneziano, *QCD quark condensate from SUSY and the orientifold large N expansion*, *Phys. Lett. B* **579** (2004) 384 [[hep-th/0309013](#)].
- [8] G. ’t Hooft, *Some Twisted Selfdual Solutions for the Yang-Mills Equations on a Hypertorus*, *Commun. Math. Phys.* **81** (1981) 267.

- [9] M. García Pérez, A. González-Arroyo and C. Pena, *Perturbative construction of selfdual configurations on the torus*, *JHEP* **09** (2000) 033 [[hep-th/0007113](#)].
- [10] A. González-Arroyo, *Constructing $SU(N)$ fractional instantons*, *JHEP* **02** (2020) 137 [[1910.12565](#)].
- [11] M. M. Anber and E. Poppitz, *The gaugino condensate from asymmetric four-torus with twists*, *JHEP* **01** (2023) 118 [[2210.13568](#)].
- [12] E. Cohen and C. Gomez, *Chiral Symmetry Breaking in Supersymmetric Yang-Mills*, *Phys. Rev. Lett.* **52** (1984) 237.
- [13] A. R. Zhitnitsky, *Torons, Chiral Symmetry Breaking and the $U(1)$ Problem in the σ Model and in Gauge Theories*, *Nucl. Phys. B* **340** (1990) 56.
- [14] N. M. Davies, T. J. Hollowood, V. V. Khoze and M. P. Mattis, *Gluino condensate and magnetic monopoles in supersymmetric gluodynamics*, *Nucl. Phys. B* **559** (1999) 123 [[hep-th/9905015](#)].
- [15] N. M. Davies, T. J. Hollowood and V. V. Khoze, *Monopoles, affine algebras and the gluino condensate*, *J. Math. Phys.* **44** (2003) 3640 [[hep-th/0006011](#)].
- [16] C. Bonanno, P. Butti, M. García Pérez, A. González-Arroyo, K.-I. Ishikawa and M. Okawa, *Nonperturbative determination of the $\mathcal{N} = 1$ supersymmetric Yang-Mills gluino condensate at large N* , *Phys. Rev. D* **110** (2024) 074507 [[2406.08995](#)].
- [17] S. Ali, H. Gerber, I. Montvay, G. Münster, S. Piemonte, P. Scior et al., *Analysis of Ward identities in supersymmetric Yang–Mills theory*, *Eur. Phys. J. C* **78** (2018) 404 [[1802.07067](#)].
- [18] S. Ali, G. Bergner, H. Gerber, P. Giudice, I. Montvay, G. Münster et al., *The light bound states of $\mathcal{N} = 1$ supersymmetric $SU(3)$ Yang-Mills theory on the lattice*, *JHEP* **03** (2018) 113 [[1801.08062](#)].
- [19] S. Ali, G. Bergner, H. Gerber, I. Montvay, G. Münster, S. Piemonte et al., *Numerical results for the lightest bound states in $\mathcal{N} = 1$ supersymmetric $SU(3)$ Yang-Mills theory*, *Phys. Rev. Lett.* **122** (2019) 221601 [[1902.11127](#)].
- [20] G. Bergner, C. López and S. Piemonte, *Study of center and chiral symmetry realization in thermal $\mathcal{N} = 1$ super Yang-Mills theory using the gradient flow*, *Phys. Rev. D* **100** (2019) 074501 [[1902.08469](#)].
- [21] S. Piemonte, G. Bergner and C. López, *Monte Carlo simulations of overlap Majorana fermions*, *Phys. Rev. D* **102** (2020) 014503 [[2005.02236](#)].
- [22] G. Bergner, G. Münster and S. Piemonte, *Exploring Gauge Theories with Adjoint Matter on the Lattice*, *Universe* **8** (2022) 617 [[2212.10371](#)].

- [23] D. Schaich, *Lattice studies of supersymmetric gauge theories*, *Eur. Phys. J. ST* **232** (2023) 305 [2208.03580].
- [24] J. Giedt, R. Brower, S. Catterall, G. T. Fleming and P. Vranas, *Lattice super-Yang-Mills using domain wall fermions in the chiral limit*, *Phys. Rev. D* **79** (2009) 025015 [0810.5746].
- [25] JLQCD collaboration, S. W. Kim, H. Fukaya, S. Hashimoto, H. Matsufuru, J. Nishimura and T. Onogi, *Lattice study of $4d$ $N=1$ super Yang-Mills theory with dynamical overlap gluino*, *PoS LATTICE2011* (2011) 069 [1111.2180].
- [26] M. M. Anber and E. Poppitz, *Higher-order gaugino condensates on a twisted \mathbb{T}^4 : In the beginning, there was semi-classics* (2024) [2408.16058].
- [27] I. Jack and D. R. T. Jones, *The Gaugino Beta function*, *Phys. Lett. B* **415** (1997) 383 [hep-ph/9709364].
- [28] L. Giusti, F. Rapuano, M. Talevi and A. Vladikas, *The QCD chiral condensate from the lattice*, *Nucl. Phys. B* **538** (1999) 249 [hep-lat/9807014].
- [29] ALPHA collaboration, M. Della Morte, R. Hoffmann, F. Knechtli, J. Rolf, R. Sommer, I. Wetzorke et al., *Non-perturbative quark mass renormalization in two-flavor QCD*, *Nucl. Phys. B* **729** (2005) 117 [hep-lat/0507035].
- [30] J. Hisano and M. A. Shifman, *Exact results for soft supersymmetry breaking parameters in supersymmetric gauge theories*, *Phys. Rev. D* **56** (1997) 5475 [hep-ph/9705417].
- [31] T. Eguchi and H. Kawai, *Reduction of Dynamical Degrees of Freedom in the Large N Gauge Theory*, *Phys. Rev. Lett.* **48** (1982) 1063.
- [32] G. 't Hooft, *A Property of Electric and Magnetic Flux in Nonabelian Gauge Theories*, *Nucl. Phys. B* **153** (1979) 141.
- [33] A. González-Arroyo and M. Okawa, *The Twisted Eguchi-Kawai Model: A Reduced Model for Large N Lattice Gauge Theory*, *Phys. Rev. D* **27** (1983) 2397.
- [34] A. González-Arroyo and M. Okawa, *Large N reduction with the Twisted Eguchi-Kawai model*, *JHEP* **07** (2010) 043 [1005.1981].
- [35] A. González-Arroyo and M. Okawa, *The string tension from smeared Wilson loops at large N* , *Phys. Lett. B* **718** (2013) 1524 [1206.0049].
- [36] M. García Pérez, A. González-Arroyo, L. Keegan and M. Okawa, *The $SU(\infty)$ twisted gradient flow running coupling*, *JHEP* **01** (2015) 038 [1412.0941].
- [37] M. García Pérez, A. González-Arroyo and M. Okawa, *Meson spectrum in the large N limit*, *JHEP* **04** (2021) 230 [2011.13061].
- [38] C. Bonanno, P. Butti, M. García Pérez, A. González-Arroyo, K.-I. Ishikawa and M. Okawa, *The large- N limit of the chiral condensate from twisted reduced models*, *JHEP* **12** (2023) 034 [2309.15540].

- [39] A. González-Arroyo and M. Okawa, *Twisted space-time reduced model of large N QCD with two adjoint Wilson fermions*, *Phys. Rev. D* **88** (2013) 014514 [1305.6253].
- [40] M. García Pérez, A. González-Arroyo, L. Keegan and M. Okawa, *Mass anomalous dimension of Adjoint QCD at large N from twisted volume reduction*, *JHEP* **08** (2015) 034 [1506.06536].
- [41] P. Butti, M. García Pérez, A. González-Arroyo, K.-I. Ishikawa and M. Okawa, *Scale setting for large- N SUSY Yang–Mills on the lattice*, *JHEP* **07** (2022) 074 [2205.03166].
- [42] D. B. Kaplan, *Dynamical Generation of Supersymmetry*, *Phys. Lett. B* **136** (1984) 162.
- [43] G. Curci and G. Veneziano, *Supersymmetry and the Lattice: A Reconciliation?*, *Nucl. Phys. B* **292** (1987) 555.
- [44] G. Münster and H. Stüwe, *The mass of the adjoint pion in $\mathcal{N} = 1$ supersymmetric Yang-Mills theory*, *JHEP* **05** (2014) 034 [1402.6616].
- [45] H. Leutwyler and A. V. Smilga, *Spectrum of Dirac operator and role of winding number in QCD*, *Phys. Rev. D* **46** (1992) 5607.
- [46] L. Giusti and M. Lüscher, *Chiral symmetry breaking and the Banks-Casher relation in lattice QCD with Wilson quarks*, *JHEP* **03** (2009) 013 [0812.3638].
- [47] M. Lüscher and F. Palombi, *Universality of the topological susceptibility in the $SU(3)$ gauge theory*, *JHEP* **09** (2010) 110 [1008.0732].
- [48] C. Bonanno, G. Clemente, M. D’Elia and F. Sanfilippo, *Topology via spectral projectors with staggered fermions*, *JHEP* **10** (2019) 187 [1908.11832].
- [49] A. Athenodorou, C. Bonanno, C. Bonati, G. Clemente, F. D’Angelo, M. D’Elia et al., *Topological susceptibility of $N_f = 2 + 1$ QCD from staggered fermions spectral projectors at high temperatures*, *JHEP* **10** (2022) 197 [2208.08921].
- [50] C. Bonanno, F. D’Angelo and M. D’Elia, *The chiral condensate of $N_f = 2 + 1$ QCD from the spectrum of the staggered Dirac operator*, *JHEP* **11** (2023) 013 [2308.01303].
- [51] A. Skouroupathis and H. Panagopoulos, *Two-loop renormalization of scalar and pseudoscalar fermion bilinears on the lattice*, *Phys. Rev. D* **76** (2007) 094514 [Erratum *ibid* **78** (2008) 119901] [0707.2906].
- [52] P. Weisz, *On the Connection Between the Λ Parameters of Euclidean Lattice and Continuum QCD*, *Phys. Lett. B* **100** (1981) 331.
- [53] M. García Pérez, A. González-Arroyo and M. Okawa, *Perturbative contributions to Wilson loops in twisted lattice boxes and reduced models*, *JHEP* **10** (2017) 150 [1708.00841].
- [54] D. Finnell and P. Pouliot, *Instanton calculations versus exact results in four-dimensional SUSY gauge theories*, *Nucl. Phys. B* **453** (1995) 225 [hep-th/9503115].

- [55] J. A. M. Vermaseren, S. A. Larin and T. van Ritbergen, *The four loop quark mass anomalous dimension and the invariant quark mass*, *Phys. Lett. B* **405** (1997) 327 [[hep-ph/9703284](#)].