# Some estimates for generalized Wigner matrix linear spectral statistics

Benjamin Landon

University of Toronto Department of Mathematics blandon@math.toronto.edu December 19, 2024

Abstract: We consider the characteristic function of linear spectral statistics of generalized Wigner matrices. We provide an expansion of the characteristic function with error  $\mathcal{O}(N^{-1})$  around its limiting Gaussian form, and identify sub-leading non-Gaussian corrections of size  $\mathcal{O}(N^{-1/2})$ . Prior expansions with this error rate held only for Wigner matrices; only a weaker error rate was available for more general matrix ensembles. We provide some applications.

# Contents

1	Intr	oduction	3										
	1.1	Background	4										
	1.2	Definition of model	5										
	1.3	Main result	6										
2	Pre	liminaries	7										
	2.1	Notation	7										
		2.1.1 Stochastic domination and overwhelming probability	8										
		2.1.2 Quasi-analytic extension	8										
	2.2	Local laws	9										
	2.3	Regular matrices	9										
	2.4	Cumulant expansion	9										
	2.5	Variance matrix and properties of semicircle law 1	0										
	2.6	Helffer-Sjostrand formula 1	1										
	2.7	Characteristic function	2										
	2.8	Estimates for derivatives of characteristic function	2										
	2.9	Graphical notation	3										
	2.10	Real symmetric vs complex Hermitian matrices	5										
	2.11	Organization of remainder of paper	5										
3	Loops estimate												
	3.1	Self-consistent equation for loop graphs	6										
	3.2	Iterative expansion	3										
		3.2.1 Proof of Proposition 3.2	5										
	3.3	Better estimate in $L = 2$ case	5										

4	Line estimate 2	28
	4.1 Iterative expansion	31
	4.1.1 Proof of Proposition 4.2	32
	4.2 A better estimate for $G_{ij}$	32
5	Preliminary expansion for $\mathbb{E}[e_{\mathfrak{a}}G_{11}]$	34
	5.1 Fourth order terms	35
	5.2 Third order terms	37
	5.2.1 Diagonal terms	37
	5.2.2 Off-diagonal terms	37
	5.3 Intermediate expansion	39
	5.4 Better estimate for $G_{11}$	41
6	Estimates for $G_{12}G_{21}$	12
7	Estimates for $G_{11}G_{22}$	18
	7.1 Expansion for expectation	51
8	Characteristic function	54
	8.1 Computation of deterministic coefficients	57
	8.1.1 Chebyshev polynomials	57
	8.1.2 Easy coefficients	58
	8.1.3 Variance functional	58
	8.2 Stein's method	52
	8.3 Proof of main theorems in the real symmetric case	- 63
	8.3.1 Proof of Theorem 1.6	63
	8.3.2 Proof of Proposition 1.5	63
9	Application: max of log-characteristic polynomial	33
	9.1 Proof of Theorem 9.2	66
	9.2 Estimates for the CLT functionals for the logarithm	67
A	Changes for complex Hermitian Wigner matrices	38
	A.1 Loops estimate	69
	A.2 Line estimates $\ldots$	71
	A.3 Preliminary expansion for $\mathbb{E}[e_{\mathfrak{a}}G_{11}]$	71
	A.4 Estimates for $G_{12}G_{21}$	75
	A.5 Estimates for $G_{11}G_{22}$	75
	A.6 Characteristic function	76
В	Linear algebra results	78
	B.1 Derivatives of the resolvent wrt matrix entries	78
С	Various proofs	78
	C.1 Proof of Lemma 2.5	79
	C.2 Proof of $(2.22)$	79
	C.3. Proof of Lemma 2.10	70

C.4	Semic	ircle calculations		•										•	•	80
	C.4.1	Proof of Lemma 2.9														80

# **1** Introduction

This work is concerned with the behavior of eigenvalues of generalized Wigner matrices. Generalized Wigner matrices H are  $N \times N$  self-adjoint random matrices with independent centered entries, such that the matrix of variances  $S_{ij} := \mathbb{E}[H_{ij}^2]$  is doubly stochastic. Wigner's semicircle law states that as  $N \to \infty$ , the eigenvalues of H (in the scaling such that the entries  $S_{ij}$  are order  $\mathcal{O}(N^{-1})$ ) satisfy,

$$\lim_{N \to \infty} \frac{1}{N} \sum_{i=1}^{N} \delta_{\lambda_i}(E) = \frac{\sqrt{(4-E^2)_+}}{2\pi} dE =: \rho_{\rm sc}(E) dE$$
(1.1)

where the convergence is, e.g., weakly almost surely. It is a classical result that the fluctuations around the semicircle law are Gaussian in that for sufficiently nice  $f : \mathbb{R} \to \mathbb{R}$  we have that the random variable

$$LSS(f) := \sum_{i=1}^{N} f(\lambda_i) - N \int f(E)\rho_{\rm sc}(E) dE$$
(1.2)

converges weakly to a Gaussian as  $N \to \infty$  with a certain mean and variance E(f) and V(f) that will be defined below. This convergence can be quantified in several ways. In the present work, will be interested in deriving an expansion of the characteristic function of the form,

$$\varphi(\lambda) := \mathbb{E}[\mathrm{e}^{\mathrm{i}\lambda LSS(f)}] = \mathrm{e}^{-\lambda^2 V(f)/2 + \mathrm{i}\lambda \mathcal{E}(f) + N^{-1/2} P_f(\lambda)} + \mathcal{O}(N^{-1+\varepsilon}), \qquad (1.3) \quad \{\mathrm{eqn:intro-exp}\}$$

for any  $\varepsilon > 0$ . This is the content of our main result, Theorem 1.6 below. Above,  $N^{-1/2}P_f(\lambda)$ should be viewed as a sub-leading correction of size  $\mathcal{O}(N^{-1/2})$ ; without such a term, the above error rate would be  $\mathcal{O}(N^{-1/2})$ . In our result  $P_f(\lambda)$  explicit in that it depends only on the first few moments of the matrix entries of H and the coefficients of f in its expansion in Chebyshev polynomials. It contains both a non-Gaussian cubic correction  $i\lambda^3$  and a subleading correction to the variance V(f).

Our motivation in deriving the expansion (1.3) lies in our prior work with Sosoe [27]. In that work we derived almost-optimal regularity conditions on the function f so that the Gaussian convergence holds. A technical input into that work was the expansion (1.3) for Wigner matrices (which we established in [27]). By obtaining (1.3) for generalized Wigner matrices, in a forthcoming revision to [27] we will extend the result to this larger class of matrices.

In Section 9 we give a modest application of the CLT for generalized Wigner matrices. We extend one of the results of the recent work [11] of Bourgade, Lopatto and Zeitouni. In that work they computed the leading order of the maximum of Wigner matrix characteristic polynomials; we extend this, as well as their optimal rigidity result, to the generalized Wigner class. The bulk of the method of that work holds for generalized Wigner matrices; there is only a minor input they require for LSS, and we show how to provide that input for generalized Wigner matrices.

#### 1.1 Background

The CLT for LSS is a classical result of random matrix theory, and a vast literature has developed. We only review a few works and refer the reader to the references in these papers for a more complete bibliography. See, e.g., the works [5, 6, 21, 23, 24, 31, 32].

Direct analysis of the characteristic function was initiated by Shcherbina [33] who applied Stein's method to prove that it converged point-wise to that of a Gaussian. In [26] the author, together with Sosoe, adapted this approach together with the local semicircle law to prove the convergence of LSS even on mesoscopic scales (i.e., to functions f of the form  $f(x) = F(N^{\alpha}(x - E))$  for some  $0 < \alpha < 1$ ). The approach of the present paper is also based on Stein's method.

The two closest precedents to our work are [8] and [30]. Bao and He [8] studied the rate of convergence of LSS to their limiting Gaussian distributions in Kolmogorov-Smirnov distance; they find optimal rates of convergence in this metric by establishing upper and lower bounds. In terms of the characteristic function, they remove the contribution of the diagonal part of Hto LSS(f), which ensures an error of  $\mathcal{O}(N^{-1})$  but with no  $P_f(\lambda)$  term on the RHS. However, their analysis is restricted to the smaller class of Wigner matrices (i.e., when the matrix S is constant) and their result does not seem to imply an expansion for  $\varphi(\lambda)$  directly (i.e., for the full matrix) with error rate  $\mathcal{O}(N^{-1})$ . However, their analysis also allows them to control the characteristic function for very large  $\lambda \approx N$ .

Secondly, for generalized Wigner matrices, an estimate similar to (1.3) but with a polynomial error rate  $\mathcal{O}(N^{-c})$  was derived by Li and Xu in [30] and also for test functions on mesoscopic scales (with an error that degenerates as the scale shortens). Other prior work on CLTs for matrices that do not have a constant variance profile include [1,12,13] as well as the works [4,16,17,22,29,34] on band matrices.

The main difficulty in the approach based on Stein's method to computing the characteristic function in the generalized Wigner case is that one needs to estimate the leading order contributions of terms like,

$$\sum_{i,j} G_{ij}(z) G_{ji}(w) S_{ij}$$
(1.4) {eqn:intro-tij

where  $G(z) = (H - z)^{-1}$  is the resolvent of H. In the Wigner case  $S_{ij} = N^{-1}$ , this can be written as  $\frac{1}{N} \operatorname{tr} \frac{G(z) - G(w)}{z - w}$ , which allows one to use the local semicircle law to estimate this quantity. However for general S, no such simplification is available. In [30], Li and Xu derived a self consistent equation for this quantity, based on the methods of [18]. They achieved an estimate of  $\mathcal{O}(N^{-1/4})$  for this quantity (around its leading order deterministic contribution).<sup>1</sup> However, as can be seen from the Wigner setting, the optimal error should be  $\mathcal{O}(N^{-1})$ .

In order to improve on this estimate, we do not attempt to estimate (1.4) with high probability, and instead simply try to determine its expectation (or rather the expectation of  $e^{i\lambda LSS(f)}$  times this quantity), as this is what arises in Stein's method. We treat this quantity by means of the cumulant expansion. Our treatment has its origins in the work [28] of Lee-Schnelli, and adapted to proving CLTs via Stein's method in [26].

If one applies the cumulant expansion to an expression involving (1.4), one is led to expressions involving more resolvent entries. One can then iterate this process, resulting in quantities with even more resolvent entries. If one was interested in obtaining the CLT only

<sup>&</sup>lt;sup>1</sup>We ignore the dependence of the error rate on the imaginary parts of z and w for the present discussion.

for global LSS, then only a few repeated cumulant expansions would be necessary (say, three). However, we will also derive the CLT for mesoscopic LSS; in order to do so, one must expand to an arbitrarily high order<sup>2</sup> adapted to the mesoscopic scale of the function. In order to do this, we use a graphical notation to keep track of which kinds of terms arise. Such graphical expansions have appeared in many works, and we are partially inspired by [9, 18].

We first develop two general estimates for certain classes of graphs. The first are loops, which consist of (the expectation of) products of terms of the form  $G_{i_1i_2}(z_1)G_{i_2i_3}(z_2)\ldots G_{i_ki_1}(z_k)$ for distinct  $i_1,\ldots,i_k$ . The second are lines, of the form  $G_{i_1i_2}(z_1)G_{i_2i_3}(z_2)\ldots G_{i_ki_{k+1}}(z_k)$ , for distinct  $i_1,\ldots,i_{k+1}$ . By means of an iterated cumulant expansion, we obtain an estimate for these terms that is one order better than the naive estimate one would obtain from applying the entry-wise local semicircle law (see (2.10) below). These results are Propositions 3.2 and 4.2 below. At each step of the iteration, we find we are able to truncate the terms in the cumulant expansion arising from cumulants of order three and higher; due to this, the expansion "closes" in the sense that expanding loops only leads to expressions with loops (with more resolvent entries), and expanding lines just leads to larger lines.

After carrying out these expansions, we then proceed via Stein's method as in [26] and apply our estimates to treat the various terms that arise. We identify the sub-leading terms  $P_f(\lambda)$  and incorporate them into our expansion.

### 1.2 Definition of model

We now define our random matrix model.

**Definition 1.1.** A generalized Wigner matrix is an  $N \times N$  matrix H s.t. the entries  $\{H_{ij}\}_{i \leq j}$  are independent centered random variables. If we denote,

$$S_{ij} := \mathbb{E}[|H_{ij}|^2] \tag{1.5}$$

then we assume that,

$$\frac{1}{CN} \le S_{ij} \le \frac{C}{N} \tag{1.6}$$

for some C > 0 and that  $\sum_{i} S_{ij} = 1$  for all j. For every p we assume that there exists  $C_p > 0$  so that

$$\mathbb{E}[|\sqrt{N}H_{ij}|^p] \le C_p \tag{1.7}$$

for all i, j and N. We say that the generalized Wigner matrix is real symmetric if  $H_{ij} \in \mathbb{R}$ for all i, j. We say that the generalized Wigner matrix is complex Hermitian if the real and imaginary parts of the off-diagonal entries are independent and that  $\mathbb{E}[H_{ij}^2] = 0$  for all  $i \neq j$ .

We will denote by  $||f||_{p,w}$  the following weighted  $L^p$  norm of f,

$$||f||_{p,w} := \left( \int_{\mathbb{R}} |f(x)|^p \frac{\mathrm{d}x}{\sqrt{|4-x^2|}} \mathrm{d}x \right)^{1/p}$$
(1.8)

**Definition 1.2.** We will say that a sequence of test functions  $f = f_N \in C^2(\mathbb{R})$  is admissible if it is compactly supported in [-5,5] and the following holds. First, for all  $\varepsilon > 0$  we have that,

$$\|f'\|_{1,w} \le N^{\varepsilon} \tag{1.9}$$

<sup>&</sup>lt;sup>2</sup>The order of the expansion will be large but fixed as  $N \to \infty$ 

for N large enough. Secondly,

$$||f||_{1,w} \le C, \qquad ||f''||_{1,w} \le CN^{1-\mathfrak{b}}$$
 (1.10)

for some  $1 > \mathfrak{b} > 0$  and C > 0.

Define,

$$LSS(f) := \operatorname{tr} f(H) - N \int f(x) \rho_{\rm sc}(x) \mathrm{d}x.$$
(1.11)

#### 1.3 Main result

We now introduce the quantities that arise in our result on the characteristic function of the random variable LSS(f). We will denote by  $\kappa_k(X)$  denote the kth cumulant of a random variable X.

**Definition 1.3** (Chebyshev polynomials). We will define the nth Chebyshev polynomial  $T_n(x)$  of the first kind by,

$$T_n(2\cos(\theta)) = \cos(n\theta), \qquad (1.12) \quad \{\texttt{eqn:cheby-def}\}$$

for  $n \ge 0$ . The polynomials  $T_n(x)$  form an orthogonal basis of the space  $L^2([-2,2], (4-x^2)^{-1/2} dx)$ .<sup>3</sup> We denote the nth Chebyshev coefficient of a function f by,

$$t_n(f) := \frac{2}{\pi} \int_{-2}^{2} T_n(x) f(x) \frac{\mathrm{d}x}{\sqrt{4 - x^2}}.$$
(1.13)

**Definition 1.4** (Functionals in the CLT). For  $\beta = 1, 2$  define the variance functional,

$$V_{\beta}(f) := \frac{1}{2\beta} \sum_{j=1}^{\infty} j t_j(f)^2 \operatorname{tr}(S^j) - \frac{2-\beta}{4} t_1(f)^2 \operatorname{tr}(S) + \frac{\hat{s}_{4,\beta}}{2} t_2(f)^2 + \frac{\hat{s}_3}{2} t_2(f) t_1(f)$$
(1.14) {eqn:intro-V-d

where  $t_i(f)$  is the *j*th coefficient of f in the basis of Chebyshev polynomials and

$$\hat{s}_{4,1} := \sum_{i,j} \kappa_4(H_{ij}), \qquad \hat{s}_{4,2} := \sum_{i \neq j} \kappa_4(\operatorname{Re}[H_{ij}]) + \kappa_4(\operatorname{Im}[H_{ij}]) + \frac{1}{2} \sum_i \kappa_4(H_{ii})$$
(1.15)

and

$$\hat{s}_3 := \sum_i \kappa_3(H_{ii}).$$
 (1.16)

Denote also,

$$\mathcal{E}_{\beta}(f) := \hat{s}_{4,\beta} \frac{1}{2\pi} \int_{-2}^{2} f(x) \frac{x^{4} - 4x^{2} + 2}{\sqrt{4 - x^{2}}} dx + \hat{s}_{3} \frac{1}{8\pi} \int_{-2}^{2} f(x) \frac{x^{3} - x^{2} - 2x + 4}{\sqrt{4 - x^{2}}} dx + \mathbf{1}_{\{\beta=1\}} \left\{ \operatorname{tr}(S) \frac{1}{2\pi} \int_{-2}^{2} f(x) \frac{2 - x^{2}}{\sqrt{4 - x^{2}}} + \frac{f(2) + f(-2)}{4} + \frac{1}{\pi} \int_{-2}^{2} \frac{f(x)}{\sqrt{4 - x^{2}}} (\operatorname{Re}[m_{\mathrm{sc}}(x + \mathrm{i0})^{2} \operatorname{tr}(S(1 - m_{\mathrm{sc}}(x + \mathrm{i0})^{2}S)^{-1})]) dx \right\}$$
(1.17)

as well as,

$$B(f) := \frac{\hat{s}_3}{8} t_1(f)^3.$$
(1.18)

 $\{def:cheby\}$ 

<sup>&</sup>lt;sup>3</sup>We note that our convention for the  $T_n(x)$  differs from the usual convention, which defines the *n*th Chebyshev polynomial as  $T_n(2x)$  where  $T_n$  is as above (i.e., defined wrt a measure on (-1, 1) instead of (-2, 2)).

The following gives elementary properties of the functionals appearing above. In particular, it guarantees that the variance function is non-negative. It is proven in Section 8.3.2.

**Proposition 1.5.** If f is an admissible test function then,

$$0 \le V_{\beta}(f) \le C \|f'\|_{1,w}^2 \log(N)^2 \tag{1.19}$$

for some C > 0, and  $|\mathcal{E}_{\beta}(f)| \le C(||f||_{1,w} + |f(2)| + |f(-2)|)$ . Moreover,  $|B(f)| \le CN^{-1/2} ||f||_{1,w}^3$ .

The following is our main result. It is proven in Section 8.3.1.

**Theorem 1.6.** Let H be a real symmetric or complex Hermitian generalized Wigner matrix. If there is a c > 0 so that  $V_{\beta}(f) \ge c$  then for any  $\varepsilon > 0$  we have that for all  $|\lambda| \le N^{-\varepsilon} \sqrt{N \|f''\|_1^{-1}}$ that

$$\mathbb{E}\left[\exp(\mathrm{i}\lambda LSS(f))\right] = \mathrm{e}^{-\lambda^2 V_{\beta}(f)/2 + \mathrm{i}\lambda^3 B(f)/3 + \mathrm{i}\lambda \mathcal{E}_{\beta}(f)} + \mathcal{O}\left(\|f''\|_{1,w} N^{-1}(1+|\lambda|) + N^{-1}|\lambda|^2\right).$$
(1.20)

Otherwise,

$$\mathbb{E}\left[\exp(i\lambda LSS(f))\right] = e^{-\lambda^2 V_{\beta}(f)/2 + i\lambda^3 B(f)/3 + i\lambda \mathcal{E}_{\beta}(f)} + \mathcal{O}(N^{-1}(1+|\lambda|)^4 + (1+|\lambda|)^3 N^{-1}(1+\|f''\|_{1,w})).$$
(1.21)

We have also that,

$$\mathbb{E}[LSS(f)] = \mathcal{E}_{\beta}(f) + \mathcal{O}(N^{-1}(1 + ||f''||_{1,w}))$$
(1.22)

Acknowledgements. The author is supported by an NSERC Discovery grant and a Connaught New Researcher award.

# 2 Preliminaries

#### 2.1 Notation

For  $z \in \mathbb{C}$  we will use the notation  $z = E \pm i\eta$ , with  $E \in \mathbb{R}$  and  $\eta > 0$ . We introduce the control parameters,

$$\Psi(z) := \sqrt{\frac{\text{Im}[m_{\text{sc}}(z)]}{N\eta} + \frac{1}{N\eta}}, \qquad \Psi_1(z) := \frac{1}{\sqrt{N\eta}}$$
(2.1)

For an admissible test function f we will also define the control parameter,

$$\Phi = \Phi(f) := N^{-1/2} (1 + \|f''\|_1)^{1/2}, \qquad \Phi_w = \Phi_w(f) := N^{1/2} (1 + \|f''\|_{1,w})^{1/2}.$$
(2.2)

Note that  $\Psi(z), \Psi_1(z) \ge cN^{-1/2}$  for some c > 0.

For a generalized Wigner matrix H define the Green's function or resolvent by  $G(z) := (H - z)^{-1}$  and the empirical Stieltjes transform by,  $m_N(z) := \frac{1}{N} \sum_i G_{ii}(z)$ .

For  $A < B \in \mathbb{R}$  we set  $\llbracket A, B \rrbracket := \{n \in \mathbb{Z} : A \leq n \leq B\}$ . The notation  $A_N(i) \approx B_N(i)$ for two N-dependent functions of an index  $i \in \mathcal{I}$  (with  $\mathcal{I}$  an abstract index set) means that there is a constant C > 0 so that  $C^{-1}A_N(i) \leq B_N(i) \leq CA_N(i)$  for all N sufficiently large {prop:funct}

{thm:main}

and all *i*. We will denote the inner product on  $\mathbb{C}^N$  by  $\langle u, v \rangle = \sum_{i=1}^N \bar{u}_i v_i$  and the dot product by  $u \cdot v = \sum_{i=1}^n u_i v_i$  for  $u, v \in \mathbb{C}^N$ .

In order to lighten the notation, if we prove some estimate for an expression involving resolvent entries, we will often relabel the indices to just use  $\{1, 2, \ldots, k\}$ . E.g., instead of considering  $G_{i_1,i_2} \ldots G_{i_{k-1}i_k}$  we consider  $G_{12}G_{23} \ldots G_{k-1,k}$ , et cetera. Clearly this is no loss of generality, and we will not comment on this further in the paper.

#### 2.1.1 Stochastic domination and overwhelming probability

We now introduce the notion of high probability events that we will use in our paper.

**Definition 2.1** (Overwhelming probability). We say that an N-dependent family of events  $\{\Xi_u : u \in U\}$  (with  $U = U^{(N)}$  an abstract index set) holds with overwhelming probability if for any large D > 0 we have that

$$\sup_{u \in U} \mathbb{P}[\Xi_u^c] \le N^{-D} \tag{2.3}$$

for sufficiently large  $N \ge N_0(D)$ .

The following notion of stochastic domination will be useful. See [19, Section 6.3] for standard properties of stochastic domination.

**Definition 2.2** (Stochastic domination). Let  $X_u$  and  $Y_u$ , for  $u \in U$  be two families of nonnegative random variables and  $U = U^{(N)}$  is a possibly N-dependent index set. We say that X is stochastically dominated by Y if for all  $\varepsilon > 0$  and D > 0 it holds that

$$\sup_{u \in U} \mathbb{P}\left[X_u > N^{\varepsilon}Y\right] \le N^{-D} \tag{2.4}$$

for all  $N \ge N_0(\varepsilon, D)$  sufficiently large. If X is stochastically dominated by Y we write X < Y. If X is complex we write  $X = \mathcal{O}_{<}(Y)$  if |X| < Y. Similarly,  $X = Z + \mathcal{O}_{<}(Y)$  means |X - Z| < Y.

#### 2.1.2 Quasi-analytic extension

We will have use for the following notion of a quasi-analytic extension of a test function f to the complex plane.

**Definition 2.3** (Quasi-analytic extension). For  $f : \mathbb{R} \to \mathbb{R}$  that is  $C^2$ , we define its quasianalytic extension,  $\tilde{f}(z)$  by,

$$\hat{f}(x + iy) := \chi(y)(f(x) + iyf'(x))$$
(2.5)

where  $\chi(y)$  is a smooth, symmetric, non-negative function satisfying  $\chi(y) = 1$  for  $|y| \le 1$  and  $\chi(y) = 0$  for  $|y| \ge 2$ . The choice of  $\chi$  will be fixed throughout the paper.

We note that,

$$\partial_{\bar{z}}\tilde{f}(z) = \frac{\mathrm{i}y\chi(y)f''(x) + \mathrm{i}(f(x) + iyf'(x))\chi'(y)}{2}.$$
(2.6) {eqn:quasi-der

Here,  $\partial_{\bar{z}} = \frac{1}{2}(\partial_x + i\partial_y)$  and  $\partial_z = \frac{1}{2}(\partial_x - i\partial_y)$  are the usual Wirtinger derivatives.

#### 2.2 Local laws

We have the following local laws.

**Theorem 2.4.** Let  $\varepsilon > 0$  and fix  $t_i \in \mathbb{C}$  satisfying  $|t_i| \leq C$  for all *i*, for some fixed C > 0. Uniformly for all *z* satisfying  $\text{Im}[z] \geq N^{\varepsilon-1}$  and  $|z| \leq 10$  we have,

$$\left|\frac{1}{N}\sum_{i}t_{i}(G_{ii}(z)-m_{\rm sc}(z))\right| < \frac{1}{N\eta}.$$
(2.7) {eqn:fluct-av}

In particular,

$$|m_N(z) - m_{\rm sc}(z)| < \frac{1}{N\eta}.$$
 (2.8) {eqn:local-law

For any fixed unit vectors  $v, w \in \mathbb{C}^N$  we have uniformly in all z as above,

$$|v^*G(z)w - v^*wm_{\rm sc}(z)| < \Psi(z).$$
 (2.9) {eqn:iso}

In particular,

$$|G_{ij}(z) - \delta_{ij}m_{\rm sc}(z)| < \Psi(z)$$
(2.10) {eqn:entry-wis

**Proof.** The estimate (2.7) follows from, e.g., [3, Theorem 2.6] near the edge and [2, Corollary 1.8] near the bulk. The estimate (2.9) follows from [9, Theorem 2.12].

#### 2.3 Regular matrices

We say that a matrix M is  $\varepsilon$ -regular if the following hold for all z satisfying  $\text{Im}[z] \ge N^{\varepsilon-1}$ and  $|z| \le \varepsilon^{-1}$ :

$$\left|\frac{1}{N}\operatorname{tr}(M-z)^{-1} - m_{\rm sc}(z)\right| \le \frac{N^{\varepsilon}}{N\eta} \tag{2.11} \quad \{\operatorname{eqn:eps-regul}$$

and

$$|(M-z)_{ij} - \delta_{ij} m_{\rm sc}(z)| \le N^{\varepsilon} \Psi(z).$$
(2.12) {eqn:eps-regul

We have the following. The proof is straightforward and deferred to Appendix C.1.

**Lemma 2.5.** Let H be a generalized Wigner matrix for  $\theta \in \mathbb{C}$  and indices a, b define the matrix  $M^{(\theta)}$  by,

$$(M^{(\theta)})_{ij} = \begin{cases} \theta, & i = a, j = b\\ \bar{\theta}, & i = b, j = a\\ H_{ij}, & \text{otherwise} \end{cases}$$
(2.13)

If a = b further assume  $\theta \in \mathbb{R}$ . For any sufficiently small  $\delta > 0$  there is an event that holds with overwhelming probability on which every matrix  $M^{(\theta)}$  with  $|\theta| \leq N^{\delta - 1/2}$  is  $\varepsilon$ -regular.

#### 2.4 Cumulant expansion

We have the following, [28, Lemma 3.2].

 $\{\texttt{lem:eps-regul}$ 

{lem:cumu-exp}

**Lemma 2.6.** Fix  $\ell \in \mathbb{N}$  and  $F \in C^{\ell+1}(\mathbb{R}; \mathbb{C})$ . Let Y be a centered random variable with finite  $\ell + 2th$  moment. Then,

$$\mathbb{E}[YF(Y)] = \sum_{r=1}^{\ell} \frac{\kappa^{(r+1)}(Y)}{r!} \mathbb{E}[F^{(r)}(Y)] + \mathbb{E}[\Xi]$$
(2.14) {eqn:cumu-exp}

where the rth cumulant of Y is denoted by  $\kappa^{(r)}(Y)$ , and the error term  $\mathbb{E}[\Xi]$  satisfies,

$$|\mathbb{E}[\Xi]| \le C \left( \mathbb{E}[|Y|^{\ell+2} \sup_{|t|\le Q} |F^{(\ell+1)}(t)|] + \mathbb{E}[|Y|^{\ell+2} \mathbf{1}_{\{|Y|>Q\}} \sup_{t\in\mathbb{R}} |F^{(\ell+1)}(t)|] \right).$$
(2.15)

for any Q > 0.

We make the following convention for the terms in the expansion on the RHS of (2.14).

**Definition 2.7** (nth order terms). When we apply a cumulant expansion, we will refer to the term with the cumulant  $\kappa^{(r)}$  of order r as the "rth order term(s)."

# 2.5 Variance matrix and properties of semicircle law

**Lemma 2.8.** The matrix of variances  $S_{ij} := \frac{s_{ij}}{N}$  has the following properties. Let **e** be the constant vector with entries equal to 1. Then  $S = A + N^{-1} \mathbf{ee}^*$  with,

$$\|A\| \le 1 - c \tag{2.16}$$

for some c > 0.

**Proof.** The constant vector is an eigenvector of S with eigenvalue 1. The result then follows from [15, Lemma 7.4.2].

The following is proven in Appendix C.4.

**Lemma 2.9.** For any C > 0 there is a c > 0 so that, for all  $|z| \leq C$ 

$$c \le |m_{\rm sc}(z)| \le 1$$
 (2.17) {eqn:msc-upper

For any z and w satisfying  $|z| + |w| \le C$  we have that,

$$\left|\frac{m_{\rm sc}(z) - m_{\rm sc}(w)}{z - w}\right| \le \frac{C_1}{\left|\operatorname{Im}[z]\right| + \left|\operatorname{Im}[w]\right|} \tag{2.18} \quad {\rm {eqn:msc-diffe}}$$

for some  $C_1 > 0$ . Moreover,

$$\frac{m_{\rm sc}(z) - m_{\rm sc}(w)}{z - w} = \frac{m_{\rm sc}(z)m_{\rm sc}(w)}{1 - m_{\rm sc}(z)m_{\rm sc}(w)}$$
(2.19) {eqn:msc-dif-

 $\{\texttt{lem:spectral}-$ 

{lem:msc}

# 2.6 Helffer-Sjostrand formula

For a  $C^2$  function f with quasi-analytic extension  $\tilde{f}(x + \mathrm{i}y)$  the HS formula is,

$$\operatorname{tr} f(H) - \mathbb{E}[\operatorname{tr} f(H)] = \frac{1}{\pi} \int_{\mathbb{C}^2} (\partial_{\bar{z}} \tilde{f}(z)) \mathrm{d}z \mathrm{d}\bar{z}$$
$$= \frac{1}{2\pi} \int_{\mathbb{R}^2} \left( \mathrm{i}y \chi(y) f''(x) + \mathrm{i}f'(x) \chi'(y) \right) N(\mathbb{E}[m_N(x + \mathrm{i}y) - \mathbb{E}[m_N(x + \mathrm{i}y)]) \mathrm{d}x \mathrm{d}y.$$
(2.20)

Define,

$$\Omega_{\mathfrak{a}} := \{ z : |\mathrm{Im}[z]| \ge N^{\mathfrak{a}-1}, |\mathrm{Re}[z]| \le 6 \}.$$
(2.21)

A straightforward argument using (2.8) shows that (see Appendix C.2)

$$\begin{aligned} \operatorname{tr} f(H) &- \mathbb{E}[\operatorname{tr} f(H)] \\ &= \frac{1}{2\pi} \int_{\Omega_{\mathfrak{a}}} \left( \operatorname{i} y \chi(y) f''(x) + \operatorname{i} (f(x) + \operatorname{i} y f'(x)) \chi'(y) \right) N(m_N(z) - \mathbb{E}[m_N(z)]) \mathrm{d} x \mathrm{d} y \\ &+ \mathcal{O}_{\prec}(N^{\mathfrak{a}-1} \| f'' \|_1). \end{aligned}$$

$$(2.22) \quad \{ \operatorname{eqn}: \operatorname{HS-est} \}$$

Note that since we assume that  $\chi$  is symmetric, the integral on the RHS is real.

The following is straightforward, and the proof is deferred to Appendix C.3.

**Lemma 2.10.** We have the following. Assume that f' is supported in (-5,5). Suppose that H(z) is a holomorphic function on  $\mathbb{C}\setminus\mathbb{R}$ . Let  $S_H(z)$  denote the function,

$$S_H(x + iy) := \sup_{|w - (x + iy)| \le |y|/2} |H(w)|.$$
(2.23)

Then,

In particular, if,

$$|H(x + iy)| \le \sum_{l=1}^{K} \frac{A_l}{|y|^{s_l}}$$
(2.25) {eqn:H-upper}

{lem:H-est}

for some  $s_l \in [1, 2]$  and  $A_l > 0$  we have for admissible f that,

$$\left| \frac{1}{2\pi} \int_{\Omega_{a}} \left( iy \chi(y) f''(x) + if'(x) \chi'(y) \right) H(z) dx dy \right|$$

$$< \left( \sum_{l=1}^{K} A_{l} + \sum_{l=1}^{k} A_{l} \| f'' \|_{1}^{s_{l}-1} \right)$$

$$(2.26) \quad \{ eqn: H-est \}$$

Note also that if (2.25) holds then,

$$|\partial_z^k H(x+\mathrm{i}y)| \le C_k \sum_{l=1}^K \frac{A_l}{|y|^{s_l+k}} \tag{2.27} \quad \{\mathtt{eqn:del-z-H}\}$$

for some constant  $C_k > 0$ .

For future reference we record here Green's theorem in complex notation which states that,

$$\int_{\Omega} \partial_{\bar{z}} F(z) dz d\bar{z} = -\frac{i}{2} \int_{\partial \Omega} F(z) dz \qquad (2.28) \quad \{eqn:green-2\}$$

for any  $C^2$  function F and sufficiently regular domain  $\Omega$ .

#### 2.7 Characteristic function

We define,

$$e_{\mathfrak{a}}(\lambda) := \exp\left(\frac{\mathrm{i}\lambda}{2\pi} \int_{\Omega_{\mathfrak{a}}} \left(\mathrm{i}y\chi(y)f''(x) + \mathrm{i}f'(x)\chi'(y)\right) N(m_N(z) - \mathbb{E}[m_N(z)])\mathrm{d}x\mathrm{d}y\right)$$
(2.29)

Due to (2.22) we have that,

$$|\mathbb{E}[e_{\mathfrak{a}}(\lambda)] - \mathbb{E}[\exp(i\lambda(\operatorname{tr} f(H) - \mathbb{E}[\operatorname{tr} f(H)]))]| < \frac{|\lambda| N^{\mathfrak{a}} ||f''||_1}{N}.$$
(2.30)

Since  $\chi(y)$  is symmetric, the integral in the definition of  $e_{\mathfrak{a}}$  is real and so  $|e_{\mathfrak{a}}(\lambda)| \leq 1$ . We will use this estimate throughout the paper without further comment.

# 2.8 Estimates for derivatives of characteristic function

**Lemma 2.11.** Let f be admissible. We have that for  $i \neq j$ ,

$$\partial_{ij}e_{\mathfrak{a}} = -2\frac{\mathrm{i}\lambda e_{\mathfrak{a}}}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{z}}\tilde{f}(z))\partial_{z}G_{ji}(z)\mathrm{d}z\mathrm{d}\bar{z} = \mathcal{O}_{\prec}(|\lambda|\Phi), \qquad (2.31) \quad \{\mathtt{eqn:deleaij}\}$$

and

$$\partial_{ij}^2 e_{\mathfrak{a}} = \mathcal{O}_{\langle}(|\lambda|\Phi + |\lambda|^2 \Phi^2) + \lambda c_f e_{\mathfrak{a}}$$
(2.32) {eqn:deleaij2}

{lem:delea}

for a constant  $c_f$  satisfying  $c_f = \mathcal{O}_{\prec}(1)$ .

**Proof.** The equality in (2.31) follows from direct calculation using (B.5) and the fact that

$$\sum_{k} G_{ik}(z) G_{kj}(z) = \partial_z G_{ij}(z).$$
(2.33)

The inequality follows from the fact that  $\partial_z G_{ij}(z) = \mathcal{O}_{<}(N^{-1/2}|\text{Im}[z]|^{-3/2})$  (due to (2.9) and (2.27)) and (2.26). For the second estimate we have by direct calculation and an argument similar to the proof of (2.31),

$$\partial_{ij}^2 e_{\mathfrak{a}} = \mathcal{O}_{<}(|\lambda|^2 \Phi^2) + 2\frac{i\lambda e_{\mathfrak{a}}}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{z}}\tilde{f}(z))\partial_z (G_{ii}G_{jj} + G_{ij}^2) \mathrm{d}\bar{z}\mathrm{d}z.$$
(2.34) {eqn:delea2}

We now conclude (2.32) using that  $G_{ii}G_{jj} + G_{ij}^2 = m_{sc}(z)^2 + \mathcal{O}_{\prec}(\Psi(z))$ , and (2.26).

{lem:del-ea-ge

**Lemma 2.12.** Let f be admissible. Fix  $n \in \mathbb{N}$ . Uniformly in  $i \neq j$  we have,

$$|\partial_{ij}^{n} e_{\mathfrak{a}}(\lambda)| < ((1+|\lambda|)^{\lfloor n/2 \rfloor} (1+\Phi(1+|\lambda|)) + (1+|\lambda|)^{n} \Phi^{n}) < (1+|\lambda|)^{\lfloor n/2 \rfloor}$$
(2.35) {eqn:del-gen-

with the second inequality holding if  $|\lambda| \Phi \leq 1$ . If i = j then,

$$|\partial_{ii}^n e_{\mathfrak{a}}(\lambda)| < (1+|\lambda|)^n. \tag{2.36} \quad \{\texttt{eqn:del-gen-2}\}$$

If H is replaced by an  $\varepsilon$ -regular matrix, then for any  $\delta > 0$ , the above estimates hold with an additional factor of  $N^{\delta}$  on the right hand sides, as long as  $\varepsilon > 0$  is sufficiently small, depending on  $\delta$  and n.

**Proof.** We begin with (2.35). The derivative  $\partial_{ij}^n e_{\mathfrak{a}}(\lambda)$  is a sum of terms of the form,

$$e_{\mathfrak{a}}(\lambda) \prod_{r=1}^{m} \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{z}} \tilde{f}(z)) \partial_{ij}^{n_{r}} N m_{N}(z) \mathrm{d}z \mathrm{d}\bar{z}, \qquad (2.37) \quad \{\mathtt{eqn:del-integration}\}$$

where  $n_i \in \mathbb{N}$  s.t.  $\sum_{i=1}^{m} n_i = n$ . Similarly to the proof of Lemma 2.11 we have that,

$$\int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{z}} \tilde{f}(z)) \partial_{ij}^{n_r} N m_N(z) \mathrm{d}z \mathrm{d}\bar{z} = \mathcal{O}_{<} \left( \Phi + \mathbf{1}_{\{n_r > 1\}} \right).$$
(2.38) {eqn:del-integ

If  $m \leq \lfloor n/2 \rfloor$  then using (2.38) we can simply bound each integral in (2.37) by  $\mathcal{O}_{<}(1)$  yielding an overall estimate of  $\mathcal{O}_{<}((1 + |\lambda|)^{\lfloor n/2 \rfloor})$ .

On the other hand, if  $m > \lfloor n/2 \rfloor$ , then  $|\{r : n_r = 1\}| \ge 2m - n$ , and so we can bound (2.37) by  $\mathcal{O}_{<}((1 + |\lambda|)^m \Phi^{2m-n})$ . We then maximize this upper bound over choices of  $m \in [\lceil n/2 \rceil + 1, n \rceil]$ , yielding (2.35).

The estimate (2.36) simply follows by bounding the integral on the LHS of (2.38) by  $\mathcal{O}_{<}(1)$ . Finally, the statements for  $\varepsilon$ -regular matrices follows simply from the fact that the above estimates depended only on estimates for the resolvent entries.

# 2.9 Graphical notation

In order to develop estimates for quantities like  $\mathbb{E}[e_{\mathfrak{a}}G_{ij}(z)]$  we will be forced to consider more complicated quantities such as

$$\frac{(i\lambda)^2}{\pi^2} \int_{\Omega_a^2} (\partial_{u_1} \partial_{u_2} \mathbb{E}[G_{12}(u_1) G_{23}(u_1) G_{34}(w_1) G_{41}(u_2)]) \left( \prod_{i=1}^2 (\partial_{\bar{u}_i} \tilde{f}(u_i)) du_i d\bar{u}_i \right).$$
(2.39) {eqn:example-1}

In order to do so, we develop a graphical notation to assist with the evaluation of monomials in resolvent entries. Note that in (2.39), there are two variables of integration  $u_1$  and  $u_2$ , and a fixed variable  $w_1$ . In order to estimate such terms we will further apply cumulant expansions, which will result in more variables of integration  $u_i$  arising.

Due to this, we will need to distinguish between the the fixed variable  $w_1$  and the two integration variables  $u_1$  and  $u_2$  in (2.39) (and similarly for other terms with more integration variables and fixed variables). The following definition of encoding monomials of resolvent entries by graphs is natural, except for the fact that the introduction of the two vectors  $\boldsymbol{u}$  and  $\boldsymbol{w}$  appears unmotivated at first. The motivation is due to the fact, illustrated by (2.39), that we need to keep track of which variables are integrated over (the  $\boldsymbol{u}$ ) and which are fixed (the  $\boldsymbol{w}$ ).

**Definition 2.13** (Hypergraphs). A hypergraph  $\mathcal{G}$  is a graph of vertices V and directed edges E (multiple edges and self-loops are allowed), together with the following additional data. First, an index set  $\underline{i} \in [\![1, N]\!]^V$ . Secondly, a set of formal variables  $\{u_i\}_{i=1}^m$  for some  $m \in \mathbb{N}$ . Finally, two vectors  $\boldsymbol{u}, \boldsymbol{w} \in \mathbb{C}^E$  s.t. for all  $e \in E$ , exactly one of  $u_e$  or  $w_e$  is non-zero. Moreover, we assume that each non-zero entry of  $\boldsymbol{u}$  equals one of the  $\{u_i\}_{i=1}^m$ , and furthermore that each  $u_i$  appears at least once in  $\boldsymbol{u}$ .

Define  $\mathbf{z} = \mathbf{u} + \mathbf{w}$ . Note that  $\mathbf{z} \in \mathbb{C}^E$  and each entry contains either a variable denoted  $w_e$  or  $u_e$ . Given a hypergraph  $\mathcal{G}$ , the monomial associated to  $\mathcal{G}$  is,

$$\mathcal{M}(\mathcal{G}) := \prod_{e=(a,b)\in E} \left( G_{i_a i_b}(z_e) - \delta_{i_a, i_b} m_{\mathrm{sc}}(z_e) \right).$$
(2.40)

Note that we can regard this, for fixed w, as a function over the variables  $\{u_i\}_{i=1}^m$ .

We associate to a hypergraph  $\mathcal{G}$  two control parameters,

$$\tilde{\Psi}(\mathcal{G}) := \prod_{e \in E} \Psi(z_e), \qquad \tilde{\Psi}_1(\mathcal{G}) := \prod_{e \in E} \Psi_1(z_e).$$
(2.41)

Note that  $\mathcal{M}(\mathcal{G}) = \mathcal{O}_{\prec}(\tilde{\Psi}(\mathcal{G}))$  when we regard all of the entries  $z_e$  of z as lying in the domain  $\Omega_{\mathfrak{a}}$ .

The evaluation of a hypergraph  $\mathcal{G}$  is,

$$T(\mathcal{G}) := \frac{(\mathrm{i}\lambda)^m}{\pi^m} \int_{\Omega^m_{\mathfrak{a}}} \left( \partial_{u_1} \partial_{u_2} \dots \partial_{u_m} \mathbb{E}[e_{\mathfrak{a}} \mathcal{M}(\mathcal{G})] \right) \left( \prod_{i=1}^m (\partial_{\bar{u}_i} \tilde{f}(u_i)) \mathrm{d}u_i \mathrm{d}\bar{u}_i \right)$$
(2.42) {eqn:T-def}

Here, we are viewing  $\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(\mathcal{G})]$  as a function of the *m* variables  $\{u_i\}_{i=1}^m$  in order to define the derivatives wrt the  $u_i$  and the integration over the  $u_i$ .

We remark also that we can regard  $T(\mathcal{G})$  as a function of the non-zero entries of the vector w.

**Example.** Consider the case that (V, E) is a cycle on four edges and vertices. Assume that  $\underline{i} = (1, 2, 3, 4), \ \boldsymbol{u} = (u_1, u_1, 0, u_2)$  and  $\boldsymbol{w} = (0, 0, w_1, 0)$  (so that implicitly, the formal variables are  $\{u_i\}_{i=1}^2$ ). Then,

$$\mathcal{M}(\mathcal{G}) = G_{12}(u_1)G_{23}(u_1)G_{34}(w_1)G_{41}(u_2) \tag{2.43}$$

and

$$T(\mathcal{G}) = \frac{(i\lambda)^2}{\pi^2} \int_{\Omega_a^2} (\partial_{u_1} \partial_{u_2} \mathbb{E}[G_{12}(u_1) G_{23}(u_1) G_{34}(w_1) G_{41}(u_2)]) \left(\prod_{i=1}^2 (\partial_{\bar{u}_i} \tilde{f}(u_i)) du_i d\bar{u}_i\right) \quad (2.44)$$

I.e., this is (2.39) above.

**Definition 2.14** (Loops and lines). A loop is a graph with vertices  $V = \{1, \ldots, K\}$  and edges  $E = \{(1, 2), (2, 3), \ldots, (K - 1, K), (K, 1)\}$ . In the case that K = 1, a loop has only the edges  $\{(1, 1)\}$ . A line is a graph with vertices  $V = \{1, \ldots, K\}$  (with  $K \ge 2$ ) and edges  $E = \{(1, 2), (2, 3), \ldots, (K - 1, K)\}$ .

For a multi-index  $\underline{i}$ , we will fix the notational convention,

$$\{\underline{i}\} := \{i_v : v \in V\},\tag{2.45}$$

i.e.,  $\{\underline{i}\}$  is the set comprised of the entries of  $\underline{i}$ .

#### 15

### 2.10 Real symmetric vs complex Hermitian matrices

We will write in detail our proofs only for the case of real symmetric generalized Wigner matrices. The proofs for the complex Hermitian case are almost the same. We give the details of how the arguments can be modified to treat the complex Hermitian case in Appendix A.

#### 2.11 Organization of remainder of paper

In Sections 3 and 4 we develop two general estimates for graphs involving disjoint loops with possibly one line. These estimates are the backbone for what follows. In Section 5 we develop an intermediate expansion for the  $\mathbb{E}[e_{\mathfrak{a}}G_{ii}(z)]$  in terms of higher order terms to be estimated. These terms are of the form  $\mathbb{E}[e_{\mathfrak{a}}G_{ij}(z)G_{ji}(w)]$  and  $\mathbb{E}[e_{\mathfrak{a}}G_{ii}(z)G_{jj}(w)]$ . These terms are then estimated in Sections 6 and 7. We finally prove our main results in Section 8, along with proving various properties of the deterministic coefficients that arise.

In Section 9 we extend some of the results in [11] to generalized Wigner matrices.

# 3 Loops estimate

Throughout this section we will assume that f is an admissible function. We will develop an estimate for the following class of hypergraphs.

**Definition 3.1.** Fix  $z \in \mathbb{C}$ . We will say that a hypergraph  $\mathcal{G}$  is z-loop admissible if

- (1) Every entry of  $\underline{i}$  is unique
- (2) The graph (V, E) underlying  $\mathcal{G}$  is a union of disjoint loops.
- (3) Exactly one edge  $e \in \mathcal{G}$  satisfies  $w_e = z$ , and the loop containing e is of length at least 2.

For any hypergraph  $\mathcal{G}$  let  $E_w := \{e \in E : w_e \neq 0, u_e = 0\}$ . For later use, define,

$$\Psi_1(\mathcal{G}) := \prod_{e \in E_w} \Psi_1(z_e).$$
(3.1)

The main result of this section is the following estimate. The proof appears in Section 3.2.1 below.

**Proposition 3.2.** Let  $\mathcal{G}$  be z-loop admissible, and let L be the length of the loop containing z. Assume that u = 0 and that all entries of w lie in  $\Omega_{\mathfrak{a}}$ , and that  $|\lambda \Phi| \leq N^{-c}$  for some c > 0. If  $L \geq 3$ , then,

$$|T(\mathcal{G})| < \Psi_1(z)\Psi_1(\mathcal{G}) \tag{3.2}$$

**Remark.** The point of the above estimate is that the naive size of  $T(\mathcal{G})$ , using (2.10), is  $\mathcal{O}_{<}(\tilde{\Psi}_{1}(\mathcal{G}))$ , and so we gain a small factor  $\Psi_{1}(z)$  over this estimate, due to cancellation when taking the expectation. The restriction  $L \geq 3$  is optimal, as  $G_{12}(z)G_{21}(\bar{z}) = |G_{12}(z)|^2$  and so there is no improvement.

{sec:loops}

{def:loop-admi

{prop:main-loo

### 3.1 Self-consistent equation for loop graphs

This section is devoted to proving a self-consistent equation for expectations of monomials of resolvent entries,  $\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(\mathcal{G})]$ . We will assume throughout this section that  $\mathcal{G}$  is a z-loop admissible hypergraph and that all  $z_e$  satisfy  $z_e \in \Omega_{\mathfrak{a}}$  (and recall that f is assumed to be an admissible function). By relabelling indices we may assume that,

$$\mathcal{M}(\mathcal{G}) = G_{12}(z)M_1G_{a1}(w)M_2 \tag{3.3}$$

where  $M_1 = G_{23}G_{34} \dots G_{a-1,a}$ , and  $M_2$  is a collection of loops whose indices are disjoint from the loop  $G_{12}M_1G_{a1}$ . We also relabel the graph so that the vertices of the loop containing zare  $\{1, 2, \dots, a\}$ ; i.e., the vertex labels are the same as the index labels. If the loop containing z is of length 2, then  $M_1 = 1$  and  $G_{a1}(w) = G_{21}(w)$ . Note that it is possible that  $M_2 = 1$ . When the context is clear we will denote  $\tilde{\Psi} = \tilde{\Psi}(\mathcal{G})$ . Let,

$$\mathbb{T} := \{\underline{i}\} \setminus \{1\}. \tag{3.4} \quad \{\texttt{eqn:tau-def}\}$$

We will eventually apply a cumulant expansion to  $\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(\mathcal{G})]$ . In preparation, let us derive the following estimates.

**Lemma 3.3.** For any  $k \ge 1$ , the quantity,

$$\partial_{1j}^k(G_{j2}(z)M_1G_{a1}(w)M_2)$$
 (3.5) {eqn:del-loop

obeys the following estimates:

(i) If 
$$j = 1$$
, it is  $\mathcal{O}_{\prec}(\tilde{\Psi})$ .  
{it:gen-der-2}

(ii) If 
$$j = 2$$
 and  $2 = a$ , then it is  $\mathcal{O}_{\prec}(\tilde{\Psi}\Psi^{-1}(z)\Psi^{-1}(w))$ .

- (iii) If j = a and  $2 \neq a$ , then it is  $\mathcal{O}_{<}(\tilde{\Psi}\Psi^{-1}(w))$ .
- (iv) If  $j \in \{\underline{i}\}$  but  $j \neq 1, 2, a$  then it is  $\mathcal{O}_{<}(\bar{\Psi})$  as long as j is not the index of a self-loop  $G_{jj} m_{sc}$ . If it is the index of a self-loop  $G_{jj}(v) m_{sc}(v)$ , then any derivative of order at least 2 is  $\mathcal{O}_{<}(\bar{\Psi}\Psi(v)^{-1})$ , but the first derivative when k = 1 is still  $\mathcal{O}_{<}(\bar{\Psi})$ .
- (v) If  $j \notin \{\underline{i}\}$ , then it is  $\mathcal{O}_{\prec}(\tilde{\Psi})$

These estimates also hold for any  $\varepsilon$ -regular matrix with an additional factor of  $N^{\delta}$  on the RHS, for any  $\delta > 0$  as long as  $\varepsilon > 0$  is sufficiently small depending on  $\delta$  and k.

**Proof.** In general,  $\partial_{1j}^k(G_{ab}(v) - \delta_{ab}m_{sc}(v))$  is a sum of products of k + 1 resolvent entries, and the 2(k + 1) indices of these k + 1 resolvent entries are comprised of one copy each of a and b, and k copies each of 1 and j. Moreover, the indices a and b do not appear in the same resolvent entry. In particular, if either  $a \notin \{1, j\}$  or  $b \notin \{1, j\}$ , then this derivative is  $\mathcal{O}_{<}(\Psi(v))$ . From this observation, we see that the derivatives of  $M_2$  and  $M_1$  can be estimated by the same product of factors of  $\Psi(z_e)$  as can the original quantities  $M_2$  and  $M_1$ , except for the following case. If there is a diagonal entry  $G_{jj}(u) - m_{sc}(u)$  present in  $M_2$ , then the second and higher derivatives  $\partial_{1j}^k(G_{jj}(v) - m_{sc}(v))$  may contain a term that is a product of diagonal resolvent entries and so is no longer  $\mathcal{O}_{<}(\Psi(v))$ . These observations quickly prove items (iv) and (v). The remaining estimates of the lemma then just lists the various cases in

16

 $\{\texttt{sec:loop-self}$ 

 $\{\texttt{lem:del-loops}$ 

{it:gen-der-1}

{it:gen-der-3}

{it:gen-der-4}

which either the entry  $G_{j2}$  or  $G_{a1}$  has the indices (1, j) or when j = 2 and  $G_{j2}(z) = G_{22}(z)$  is no longer  $\mathcal{O}(\Psi(z))$ .

The statement about  $\varepsilon$ -regular matrices is simply due to the fact that all of the estimates follow from estimates on the resolvent entries, which the  $\varepsilon$ -regular matrices are assumed to obey (with the fixed  $N^{\varepsilon}$  loss).

In preparation for applying the cumulant expansion we derive the following estimate for the error terms.

**Lemma 3.4.** Let  $\delta > 0$  and assume  $|\lambda \Phi| \leq 1$ . For any  $\varepsilon$ -regular matrix and sufficiently small  $\varepsilon > 0$  we have that the quantity,

$$N^{-2}|\partial_{1j}^{3}(e_{\mathfrak{a}}G_{12}(z)M_{1}G_{a2}(w)M_{2})|$$
(3.6) {eqn:loops-fou

is bounded by  $N^{\delta-1}(1+|\lambda|)\tilde{\Psi}$  if  $j \in \{\underline{i}\}$  and  $N^{-2+\delta}(1+|\lambda|)\tilde{\Psi}$  if  $j \notin \{\underline{i}\}$ .

**Proof.** For the case j = 1, we see that from (2.36) the derivatives hitting  $e_{\mathfrak{a}}$  contribute at most a factor of  $(1 + |\lambda|)^3$  and from Lemma 3.3(i) that the resolvent entries and their derivatives contribute a factor of  $\tilde{\Psi}$ , yielding an upper bound of  $N^{-2+\delta}(1 + |\lambda|)^3 \tilde{\Psi}$  for this term. Similarly, the case j = 2 is bounded by  $N^{\delta-2}(1+|\lambda|)\tilde{\Psi}\Psi(z)^{-1}\Psi(w)^{-1} \leq N^{\delta-1}(1+|\lambda|)\tilde{\Psi}$ by applying (2.35) and Lemma 3.3(ii).

If  $j \in \{\underline{i}\}$  but  $j \neq 1, 2$ , then derivatives hitting resolvent entries can remove at most one factor of  $\Psi$  in the control parameter  $\tilde{\Psi}$ , by Lemma 3.3(iii),(iv). Combining this with (2.35) we see that these terms can be bounded by  $(1 + |\lambda|)N^{-3/2+\delta}\tilde{\Psi}$ .

Finally, if  $j \notin \{\underline{i}\}$ , then the resolvent entries and their derivatives always contribute a factor of  $\Psi$  by Lemma 3.3(v), and so we conclude by applying again (2.35).

By applying the above with the cumulant expansion, Lemma 2.6, we derive the following.

**Lemma 3.5.** We have, for  $|\lambda \Phi| \leq 1$ ,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)M_{1}G_{a1}(w)M_{2}] = \sum_{j} \frac{s_{1j}}{N}\mathbb{E}[\partial_{j1}(e_{\mathfrak{a}}G_{j2}(z)M_{1}G_{a1}(w)M_{2})] + \sum_{j} \frac{s_{1j}^{(3)}}{2N^{3/2}}\mathbb{E}[\partial_{1j}^{2}(e_{\mathfrak{a}}G_{j2}(z)M_{1}G_{a1}(w)M_{2})] + \mathcal{O}_{\prec}(N^{-1}(1+|\lambda|)\tilde{\Psi})$$
(3.7) {eqn:loops-exp

**Proof.** From the fact that (H - z)G = 1 we have,

$$z\mathbb{E}[G_{12}(z)M_1G_{a1}(w)M_2] = \sum_j \mathbb{E}[H_{1j}G_{j2}(z)M_1G_{a1}(w)M_2].$$
(3.8)

The result now follows from Lemma 2.6, and using Lemma 3.4 to estimate the terms at fourth order, and the fact that  $|H_{ij}| < N^{-1/2}$ .

We now estimate the third order terms.

**Lemma 3.6.** Assume  $|\lambda \Phi| \leq 1$ . We have for  $j \in \{\underline{i}\}$  that,

$$\left|\frac{1}{N^{3/2}}\mathbb{E}[\partial_{1j}^2(e_{\mathfrak{a}}G_{j2}M_1G_{a1}M_2)]\right| < (1+|\lambda|)\tilde{\Psi}N^{-1} + \delta_{a,2}\delta_{j,2}N^{-1/2}\tilde{\Psi}$$
(3.9) {eqn:loops-th:

 $\{\texttt{lem:loops-exp}$ 

{lem:loops-fou

{lem:loops-thi

For other terms we have that,

$$N^{-3/2} \sum_{j \notin \{\underline{i}\}} \left( |\mathbb{E}[(\partial_{1j} e_{\mathfrak{a}}) \partial_{1j} (G_{j2} M_1 G_{a1} M_2)]| + |\mathbb{E}[(e_{\mathfrak{a}}) \partial_{1j}^2 (G_{j2} M_1 G_{a1} M_2)]| \right) = \mathcal{O}_{\prec}(\tilde{\Psi} N^{-1/2}).$$
(3.10) {eqn:loops-thi

and

$$\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\hat{\sigma}_{1j}^2 e_{\mathfrak{a}}) G_{j2} M_1 G_{a1} M_2] = \mathcal{O}_{<}(N^{-1/2} \tilde{\Psi}).$$
(3.11) {eqn:loops-theorem (3.11)

**Proof.** We begin with (3.9), which will follow in a similar fashion to the first part of the proof of Lemma 3.4. For j = 1, we see from Lemma 3.3(i) and (2.36) that this term is  $\mathcal{O}_{<}((1 + |\lambda|)^2 \tilde{\Psi} N^{-3/2}) = \mathcal{O}_{<}(N^{-1}(1 + |\lambda|)\tilde{\Psi})$ . For  $j \in \{\underline{i}\}$  but  $j \neq 1$ , we consider first the case that both derivatives hit  $e_{\mathfrak{a}}$ . Then the resolvent entries contribute  $\mathcal{O}_{<}(\tilde{\Psi}\Psi(z)^{-1})$  (the factor  $\Psi(z)^{-1}$  coming in the case that j = 2) and so this term can be bounded by  $\mathcal{O}_{<}(N^{-3/2}(1 + |\lambda|)\tilde{\Psi}\Psi(z)^{-1}) = \mathcal{O}_{<}(N^{-1}(1 + |\lambda|)\tilde{\Psi})$ , using also (2.35). Here and below we use also that  $\Psi(z) \geq N^{-1/2}$ .

In the case that  $j \in \{\underline{i}\}, j \neq 1$  and either no or one derivatives hit  $e_{\mathfrak{a}}$ , we have the estimate  $\partial_{1j}e_{\mathfrak{a}} = \mathcal{O}_{\prec}(|\lambda|\Phi) = \mathcal{O}_{\prec}(1)$  from (2.31), as well as the facts that derivatives hitting the resolvent entries can remove at most two factors of  $\Psi(v)$  from  $\tilde{\Psi}$  if j = 2 and a = 2 and one factor otherwise, by items (ii), (iii) and (iv) of Lemma 3.3. These estimates combine to yield (3.9).

We now turn to (3.10). For both types of terms on the LHS, the resolvent entries and their derivatives are  $\mathcal{O}_{\prec}(\tilde{\Psi})$  by Lemma 3.3(v). We also have  $e_{\mathfrak{a}} = \mathcal{O}_{\prec}(1)$  and  $\partial_{1j}e_{\mathfrak{a}} = \mathcal{O}_{\prec}(1)$  by (2.31) and so (3.10) follows.

For (3.11) we have from (2.32) that,

$$\sum_{j \notin \underline{i}} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\hat{o}_{1j}^2 e_{\mathfrak{a}}) G_{j2}(z) M_1 G_{a1}(w) M_2]$$
  
= $i\lambda c_f \sum_{j \notin \underline{i}} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}} G_{j2}(z) M_1 G_{a1}(w) M_2] + \mathcal{O}_{<}(N^{-1/2} \tilde{\Psi}).$  (3.12)

Now from (2.9) we have that,

$$\sum_{j \notin \underline{i}} \frac{s_{1j}^{(3)}}{N^{3/2}} G_{j2}(z) = \mathcal{O}_{\prec}(N^{-1}\Psi(z))$$
(3.13)

and so (3.11) follows.

We now turn to the second order terms. We start with,

Lemma 3.7. We have,

$$\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}(\lambda) M_1 G_{a1}(w) M_2 \partial_{j1} G_{j2}(z)]$$
  
=  $-m_{\mathrm{sc}}(z) \mathbb{E}[e_{\mathfrak{a}} M_1 G_{a1}(w) M_2 G_{12}(z)] + \mathcal{O}_{\prec}(\Psi(z)\tilde{\Psi})$  (3.14)

{lem:loops-sec

**Proof.** By direct computation of the derivative we have,

$$\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} M_{1} G_{a1} M_{2} \partial_{j1} G_{j2}(z)]$$

$$= -\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} M_{1} G_{a1} M_{2} G_{jj}(z) G_{12}(z)] - \sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} M_{1} G_{a1} M_{2} G_{1j}(z) G_{j2}(z)]$$

$$= -m_{\rm sc}(z) \mathbb{E}[e_{\mathfrak{a}} M_{1} G_{a1} M_{2} G_{12}(z)] + \mathcal{O}_{\prec}(\Psi(z) \tilde{\Psi})$$
(3.15)

where in the last line we applied the entry-wise law (2.10).

We now define two operators on hypergraphs  $\mathcal{G}$ :

**Definition 3.8.** Let  $\mathcal{G} = (V, E, \underline{i}, \boldsymbol{u}, \boldsymbol{w})$  be a hypergraph with integration variables  $\{u_i\}_{i=1}^m$ . Given a vertex  $v \in V$ , an edge  $e_1 \in E$  and index j, we define a new hypergraph  $D_1(\mathcal{G}, v, j, e_1) = (V', E', \underline{i}', \boldsymbol{u}', \boldsymbol{w}')$  with the same integration variables  $\{u_i\}_{i=1}^m$  as follows. We add an additional vertex v' to its vertex set  $V' := V \cup \{v'\}$ . We add a new self-edge,  $E' := E \cup \{(v', v')\}$ . The new indices  $\underline{i}'$  are  $i'_l = i_l$  for  $l \in V \setminus \{v\}$ , and  $i'_v = j$  and  $i'_{v'} = i_v$ . The new spectral parameters are  $\boldsymbol{w}'_e = \boldsymbol{w}_e$  and  $\boldsymbol{u}'_e = \boldsymbol{u}_e$  for  $e \in E$  and  $\boldsymbol{w}'_{(v',v')} = \boldsymbol{w}_{e_1}$  and  $\boldsymbol{u}'_{(v',v')} = \boldsymbol{u}_{e_1}$ .

**Definition 3.9.** Let  $\mathcal{G} = (V, E, \underline{i}, u, w)$  be a hypergraph with integration variables  $\{u_i\}_{i=1}^m$ . Given an edge  $e \in E$ , an index j and another edge  $e_1 \in E$ , we define a new hypergraph  $D_2(\mathcal{G}, e, j, e_1) = (V', E', \underline{i}', u', w')$  with the same integration variables  $\{u_i\}_{i=1}^m$  as follows. Let us write e = (a, b). We add an additional vertex v' to its vertex set  $V' := V \cup \{v'\}$ . The new edges are the same as the old edges, except we remove e = (a, b) and add edges  $e_2 = (a, v')$  and  $e_3 = (v', b)$ . We set  $i'_v = i_v$  for  $v \in V$  and  $i'_{v'} = j$ . For all edges  $e' \in E \setminus \{e\}$  we set  $w'_{e'} = w_{e'}$  and  $u'_{e'} = u_{e'}$ . Finally, we set  $w'_{(a,v')} = w_{e_1}$ ,  $u'_{(a,v')} = u_{e_1}$  and  $w'_{(v',b)} = w_e$ ,  $u'_{(v',b)} = u_e$ .

In our setting, the above two operators arise when  $\partial_{j1}$  hits  $G_{a1}(w)$  in Lemma 3.10 below. The operator  $D_1$  arises from the  $G_{aj}(w)G_{11}(w)$  term in  $\partial_{j1}G_{a1} = -G_{aj}G_{11} - G_{a1}G_{1j}$ . This results in the addition of a self loop. The operator  $D_2$  arises from the  $G_{a1}G_{1j}$  term. This results in the loop containing z increasing in length by 1.

Lemma 3.10. We have,

$$\begin{split} &\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} M_{1} M_{2} G_{j2}(z) \partial_{j1}(G_{a1}(w))] \\ &= -m_{\mathrm{sc}}(w) \sum_{j \notin \mathbb{T}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{j2}(z) M_{1} G_{aj}(w) M_{2}] - \mathbf{1}_{\{a=2\}} \frac{s_{12} m_{\mathrm{sc}}(z) m_{\mathrm{sc}}(w)^{2}}{N} \mathbb{E}[e_{\mathfrak{a}} M_{2}] \\ &- \sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{1}(\mathcal{G}, 1, j, (a, 1))) + \mathcal{M}(D_{2}(\mathcal{G}, (1, 2), j, (a, 1))))] + \mathcal{O}_{\prec}(N^{-1/2} \tilde{\Psi}). \quad (3.16) \quad \{\mathsf{eqn:loops-sec}(w) \in \mathbb{C} \} \\ &= -\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{1}(\mathcal{G}, 1, j, (a, 1))) + \mathcal{M}(D_{2}(\mathcal{G}, (1, 2), j, (a, 1))))] + \mathcal{O}_{\prec}(N^{-1/2} \tilde{\Psi}). \quad (3.16) \quad \{\mathsf{eqn:loops-sec}(w) \in \mathbb{C} \} \\ &= -\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{1}(\mathcal{G}, 1, j, (a, 1))) + \mathcal{M}(D_{2}(\mathcal{G}, (1, 2), j, (a, 1))))] + \mathcal{O}_{\prec}(N^{-1/2} \tilde{\Psi}). \quad (3.16) \quad \{\mathsf{eqn:loops-sec}(w) \in \mathbb{C} \} \\ &= -\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{1}(\mathcal{G}, 1, j, (a, 1))) + \mathcal{M}(D_{2}(\mathcal{G}, (1, 2), j, (a, 1))))] + \mathcal{O}_{\prec}(N^{-1/2} \tilde{\Psi}). \quad (3.16) \quad \{\mathsf{eqn:loops-sec}(w) \in \mathbb{C} \} \\ &= -\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{1}(\mathcal{G}, 1, j, (a, 1))) + \mathcal{M}(D_{2}(\mathcal{G}, (1, 2), j, (a, 1))))] + \mathcal{O}_{\prec}(N^{-1/2} \tilde{\Psi}). \quad (3.16) \quad \{\mathsf{eqn:loops-sec}(w) \in \mathbb{C} \} \\ &= -\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{1}(\mathcal{G}, 1, j, (a, 1))) + \mathcal{M}(D_{2}(\mathcal{G}, (1, 2), j, (a, 1))))] + \mathcal{O}_{\bigstar}(N^{-1/2} \tilde{\Psi}). \quad (3.16) \quad \{\mathsf{eqn:loops-sec}(w) \in \mathbb{C} \}$$

**Proof.** By direct computation we have,

$$\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} M_1 M_2 G_{j2}(z) \partial_{j1} G_{a1}(w)] = -\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} M_1 M_2 G_{j2}(z) G_{11}(w) G_{aj}(w)] - \sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[M_1 M_2 G_{j2}(z) G_{a1}(w) G_{j1}(w)] \quad (3.17) \quad \{\text{eqn:loops-seeded}\}$$

{lem:loops-sec

{def:D1}

 $\{def:D2\}$ 

We now write,

$$\begin{split} \sum_{j} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} M_{1} M_{2} G_{j2}(z) G_{11}(w) G_{aj}(w)] &= \sum_{j \neq 2} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} M_{1} M_{2} G_{j2}(z) G_{11}(w) G_{aj}(w)] \\ &+ \mathbf{1}_{\{a=2\}} \frac{s_{12}}{N} \mathbb{E}[e_{\mathfrak{a}} M_{2} G_{22}(z) G_{11}(w) G_{22}(w)] + \mathcal{O}_{\prec} (N^{-1/2} \tilde{\Psi}) \\ &= m_{\mathrm{sc}}(w) \sum_{j \notin \mathbb{T}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{j2}(z) M_{1} G_{aj}(w) M_{2}] \\ &+ \sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{j2}(z) M_{1} G_{aj}(w) M_{2}(G_{11}(w) - m_{\mathrm{sc}}(w))] \\ &+ \mathbf{1}_{\{a=2\}} \frac{s_{12}}{N} \mathbb{E}[e_{\mathfrak{a}} M_{2} G_{22}(z) G_{11}(w) G_{22}(w)] + \mathcal{O}_{\prec} (N^{-1/2} \tilde{\Psi}) \end{split}$$
(3.18)

In the first estimate, we estimated the j = 2 term by  $\mathcal{O}_{<}(N^{-1}\tilde{\Psi}\Psi(z)^{-1}) = \mathcal{O}_{<}(N^{-1/2}\tilde{\Psi})$  if  $a \neq 2$ . In the second estimate we wrote  $G_{11}(w) = m_{\rm sc}(w) + (G_{11}(w) - m_{\rm sc}(w))$ , and also estimated the contribution of the terms  $j \in \{\underline{i}\}$  and  $j \neq 1, 2$  by  $\mathcal{O}_{<}(N^{-1/2}\tilde{\Psi})$ . The remaining terms on the second last line are of the form  $\mathcal{M}(D_1(\mathcal{G}, 1, j, (a, 1)))$ . Moreover, the terms on the third last line of (3.18) contribute the first term on the RHS of (3.16).

When a = 2 we can write the last term in (3.18) as,

$$\mathbf{1}_{\{a=2\}} \frac{s_{12}}{N} \mathbb{E}[e_{\mathfrak{a}} M_2 G_{22}(z) G_{11}(w) G_{22}(w)] = \mathbf{1}_{\{a=2\}} \frac{s_{12}}{N} m_{\mathrm{sc}}(z) m_{\mathrm{sc}}(w)^2 \mathbb{E}[e_{\mathfrak{a}} M_2] + \mathcal{O}(N^{-1/2} \tilde{\Psi}).$$
(3.19)

We also have,

$$\sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[M_1 M_2 G_{j2}(z) G_{a1}(w) G_{j1}(w)]$$
  
= 
$$\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[G_{1j}(w) G_{j2}(z) M_1 G_{a1}(w) M_2] + \mathcal{O}_{<}(N^{-1/2} \tilde{\Psi})$$
(3.20)

which yields the claim, as these terms are of the form  $\mathcal{M}(D_2(\mathcal{G}, (1, 2), j, (a, 1)))$ .

We introduce two more operations on hypergraphs  $D_3$  and  $D_4$ .

**Definition 3.11.** Let  $\mathcal{G} = (V, E, \underline{i}, u, w)$  be a hypergraph with integration variables  $\{u_i\}_{i=1}^m$ . Given two edges  $e_1 = (a, b)$  and  $e_2 = (c, d)$  in E, and an index j we define new hypergraphs  $D_3(\mathcal{G}, e_1, e_2, j)$  and  $D_4(\mathcal{G}, e_1, e_2, j)$  as follows. Both have the same integration variables  $\{u_i\}_{i=1}^m$  as before. In both cases we add a new vertex v' so that  $V' = V \cup \{v'\}$  and delete edges  $e_1$  and  $e_2$ . In both cases we add edge (v', b) and set  $i'_{v'} = j$ , and set  $u'_{(v', b)} = u_{e_1}$  and  $w'_{(v', b)} = w_{e_1}$ .

In the case of  $D_3$ , we add edges (c, v'), (a, d) and set  $u'_{(c,v')} = u'_{(a,d)} = u_{e_2}$ ,  $w'_{(c,v')} = w'_{(a,d)} = w_{e_2}$ . In the case of  $D_4$  we add edges (c, a) and (d, v') and set  $u'_{(c,a)} = u'_{(d,v')} = u_{e_2}$ ,  $w'_{(c,a)} = w'_{(d,v')} = w_{e_2}$ 

Lemma 3.12. We have,

$$\begin{split} &\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{j2}(z) G_{a1}(w) \partial_{j1}(M_{1}M_{2})] + \mathcal{O}_{<}(N^{-1/2} \tilde{\Psi}) \\ &= -\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \sum_{e \in E \setminus \{(1,2),(a,1)\}} \mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{3}(\mathcal{G},(1,2),e,j)) + \mathcal{M}(D_{4}(\mathcal{G},(1,2),e,j)))]. \quad (3.21) \quad \{\texttt{eqn:loops-sec}(M_{1}) \in \mathbb{E} \setminus \{(1,2),(a,1)\} \} \end{split}$$

 $\{def:D3D4\}$ 

{lem:loops-sec

**Proof.** First, since no resolvent entry in either  $M_1$  or  $M_2$  contains the index 1, we can estimate the terms for  $j \in \{\underline{i}\}$  by  $\mathcal{O}_{\prec}(N^{-1/2}\tilde{\Psi})$ . By inspection, the other derivatives generate the terms with  $D_3$  and  $D_4$ .

We introduce a final operation on hypergraphs.

**Definition 3.13.** Let  $\mathcal{G} = (V, E, \underline{i}, u, w)$  be a hypergraph with integration variables  $\{u_i\}_{i=1}^m$ . Given an edge  $e \in E$ , an index j and a spectral parameter u' we define a new hypergraph  $D_5(\mathcal{G}, e, j, u') = (V', E', \underline{i}', u', w')$  as follows. We add an additional integration variable  $u_{m+1} = u'$  so that the integration variables are  $\{u_i\}_{i=1}^{m+1}$ . Let us write e = (a, b). We add an additional vertex v' to its vertex set  $V' := V \cup \{v'\}$ . The new edges are the same as the old edges, except we remove e = (a, b) and add edges  $e_1 = (a, v')$  and  $e_2 = (v', b)$ . We set  $i'_v = i_v$  for  $v \in V$  and  $i'_{v'} = j$ . For all edges  $e' \in E \setminus \{e\}$  we set  $w'_{e'} = w_{e'}$  and  $u'_{e'} = u_{e'}$ . Finally, we set  $w'_{(a,v')} = 0$ ,  $u'_{(a,v')} = u_{m+1}$  and  $w'_{(v',b)} = w_e$ ,  $u'_{(v',b)} = u_e$ .

**Lemma 3.14.** We have for  $|\lambda \Phi| \leq 1$  that,

$$\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[(\partial_{j1}e_{\mathfrak{a}})G_{j2}(z)G_{a1}(w)M_{1}M_{2}]$$

$$= -\sum_{j\notin\{\underline{i}\}} \frac{2\mathrm{i}\lambda}{\pi} \frac{s_{1j}}{N} \int_{\Omega_{\mathfrak{a}}} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(D_{5}(\mathcal{G},(1,2),j,u))]\mathrm{d}u\mathrm{d}\bar{u} + \mathcal{O}_{<}(N^{-1/2}\tilde{\Psi}). \quad (3.22) \quad \{\mathrm{eqn:loops-sec}(N^{-1/2}\tilde{\Psi})\}$$

**Proof.** The j = 1 term can be estimated by  $\mathcal{O}_{<}(N^{-1}\tilde{\Psi}(1 + |\lambda|))$  by applying (2.36). The terms for  $j \in \{\underline{i}\}$  with  $j \neq 1$  can be estimated by  $\mathcal{O}_{<}(N^{-1/2}|\lambda|\Phi\tilde{\Psi})$  using (2.31). Since for  $j \neq 1$ ,

$$\partial_{1j}e_{\mathfrak{a}} = -\frac{2i\lambda}{\pi}e_{\mathfrak{a}}\int_{\Omega_{\mathfrak{a}}}\partial_{\bar{u}}\tilde{f}(u)\partial_{u}G_{1j}(u)\mathrm{d}u$$
(3.23)

the claim follows.

Define the vector,

$$v_j := \mathbb{E}[G_{j2}(z)M_1 G_{aj}(w)M_2] \mathbf{1}_{\{j \notin \mathbb{T}\}}$$
(3.24)

and the hypergraph  $\mathcal{G}^{(j)}$  to be the same as  $\mathcal{G}$  except we replace the vertex 1 with a vertex j with  $\underline{i}_j = j$ . The above expansions allow us to deduce the following self-consistent equation for the vector  $v_j$ . Recall the definition of  $\mathbb{T}$  in (3.4).

**Proposition 3.15.** We have for  $j \notin \mathbb{T}$ ,

$$\begin{aligned} v_{j} - m_{\rm sc}(z)m_{\rm sc}(w)\sum_{k}\frac{s_{jk}}{N}v_{k} &= \mathbf{1}_{\{a=2\}}\frac{m_{\rm sc}(z)^{2}m_{\rm sc}(w)^{2}s_{12}}{N}\mathbb{E}[e_{\mathfrak{a}}M_{2}] \\ &+ m_{\rm sc}(z)\sum_{k\notin\mathbb{T}\cup\{j\}}\frac{s_{jk}}{N}\mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{1}(\mathcal{G}^{(j)},j,k,(a,j))) + \mathcal{M}(D_{2}(\mathcal{G}^{(j)},(j,2),k,(a,j))))] \\ &+ m_{\rm sc}(z)\sum_{k\notin\mathbb{T}\cup\{j\}}\frac{s_{jk}}{N}\sum_{e\in E\setminus\{(j,2),(a,j)\}}\mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{3}(\mathcal{G}^{(j)},(j,2),e,k)) + \mathcal{M}(D_{4}(\mathcal{G}^{(j)},(j,2),e,k)))] \\ &+ m_{\rm sc}(z)\sum_{k\notin\mathbb{T}\cup\{j\}}\frac{2i\lambda}{\pi}\frac{s_{jk}}{N}\int_{\Omega_{\mathfrak{a}}}\partial_{\bar{u}}\tilde{f}(u)\partial_{u}\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(D_{5}(\mathcal{G}^{(j)},(j,2),k,u))]dud\bar{u} + \mathcal{O}_{<}(\Psi(z)\tilde{\Psi}). \end{aligned}$$

$$(3.25) \quad \{\text{eqn:loops-example}\}$$

{prop:loops-ex

 $\{def:D5\}$ 

{lem:loops-sec

**Proof.** For notational simplicity we do only the case j = 1. This then follows from Lemma 3.5, Lemma 3.6 to estimate the third order terms on the RHS of (3.7), and then Lemmas 3.7, 3.10, 3.12 and 3.14 to compute the second order terms. Note that we also use the identity  $m_{\rm sc}(z) + z = -1/m_{\rm sc}(z)$ .

Define the vector  $Y_j$  by,

$$Y_j := \mathbf{1}_{\{a=2\}} \mathbf{1}_{\{j \notin \mathbb{T}\}} \frac{m_{\rm sc}(z)^2 m_{\rm sc}(w)^2 s_{12}}{N} \mathbb{E}[e_{\mathfrak{a}} M_2].$$
(3.26)

For  $j \notin \mathbb{T}$ , define  $X_j$  to be the sum of the three summations over k on the last three lines of (3.25), and  $X_j = 0$  for  $j \in \mathbb{T}$ . Let **e** be the vector of all ones and define the matrices A and B by,

$$S = N^{-1} \mathbf{e} \mathbf{e}^T + A, \qquad B_{ij} = S_{ij} \mathbf{1}_{\{i \in \mathbb{T}\}}.$$
(3.27) {eqn:loops-AB

We can rewrite (3.25) as the vector equation,

$$(1 - m_{\rm sc}(z)m_{\rm sc}(w)(A - B))v = m_{\rm sc}(z)m_{\rm sc}(w)N^{-1}\mathbf{e}\mathbf{e}^Tv + X + Y + \varepsilon \qquad (3.28) \quad \{\mathsf{eqn:loops-exp}\}$$

where  $\varepsilon$  is a vector satisfying  $\|\varepsilon\|_{\infty} = \mathcal{O}_{\prec}(\Psi(z)\tilde{\Psi})$ . Note that the case when  $j \in \mathbb{T}$  is trivial because  $[(S - B)v]_j = 0$  for such j. With the above equation as input we can prove the following.

**Lemma 3.16.** For any  $j \notin \mathbb{T}$  we have that,

$$v_{j} = \sum_{k} C_{jk}(Y_{k} + X_{k} + Z_{k}) + \mathcal{O}_{<}(\Psi(z)\tilde{\Psi}) + \mathcal{O}_{<}\left(\frac{\Psi(z) + \Psi(w)}{N(|\mathrm{Im}[z]| + |\mathrm{Im}[w]|)}\tilde{\Psi}\Psi(z)^{-1}\Psi(w)^{-1}\right).$$
(3.29) {eqn:loops-exp

where  $C_{jk}$  are coefficients satisfying  $|C_{jk}| \leq C(\delta_{jk} + N^{-1})$ , and  $Z_k$  is the constant vector with entries,

$$Z_{k} = m_{\rm sc}(z)m_{\rm sc}(w)\frac{\delta_{a2}}{N}\mathbb{E}[e_{\mathfrak{a}}M_{2}]\left(\frac{m_{\rm sc}(z) - m_{\rm sc}(w)}{z - w} - m_{\rm sc}(w)m_{\rm sc}(w)\right).$$
(3.30)

Moreover, the  $C_{ik}$  are holomorphic in the variables  $z_e$  as they vary over  $\mathbb{C}\backslash\mathbb{R}$ .

**Proof.** We invert the vector equation (3.28) by moving  $(1 - m_{\rm sc}(z)m_{\rm sc}(w)(A - B))$  to the RHS. It follows that the coefficients will be  $C_{jk} := (1 - m_{\rm sc}(z)m_{\rm sc}(w)(A - B))_{jk}^{-1}$  which satisfy the desired estimates due to Lemma B.1 (the hypotheses of which are satisfied due to Lemma 2.8 and (2.17)). We compute,

$$\frac{1}{N}\mathbf{e}^{T}v = \frac{1}{N}\sum_{j\notin\mathbb{T}}\mathbb{E}[G_{aj}(w)G_{j2}(z)M_{1}M_{2}e_{\mathfrak{a}}]$$

$$= \frac{1}{N}\sum_{j}\mathbb{E}[G_{aj}(w)G_{j2}(z)M_{1}M_{2}e_{\mathfrak{a}}] - \mathbf{1}_{\{a=2\}}\frac{m_{\mathrm{sc}}(z)m_{\mathrm{sc}}(w)}{N}\mathbb{E}[M_{2}e_{\mathfrak{a}}] + \mathcal{O}_{<}(N^{-1/2}\tilde{\Psi})$$

$$= \frac{\delta_{a2}}{N}\mathbb{E}[e_{\mathfrak{a}}M_{2}]\left(\frac{m_{\mathrm{sc}}(z) - m_{\mathrm{sc}}(w)}{z - w} - m_{\mathrm{sc}}(w)m_{\mathrm{sc}}(w)\right) + \mathcal{O}_{<}(N^{-1/2}\tilde{\Psi})$$

$$+ \mathcal{O}_{<}\left(\frac{\Psi(z) + \Psi(w)}{N(|\mathrm{Im}[z]| + |\mathrm{Im}[w]|)}\tilde{\Psi}\Psi(z)^{-1}\Psi(w)^{-1}\right).$$
(3.31)

22

{lem:loops-exp

where in the last line we applied Lemma C.1. In the second line we used (2.10) in the case that a = 2 to replace the j = 2 term with the second term on the RHS. The claim now follows from this and (3.28).

# 3.2 Iterative expansion

The goal of this section is to prove Proposition 3.2 via an iterative expansion. Our iterative expansion will repeatedly use Lemma 3.16 to estimate  $T(\mathcal{G})$  in terms of contributions from a family of hypergraphs. We now define this family in Definition 3.17. We first require some notation.

Let  $\mathcal{G}$  be a z-loop admissible hypergraph. Denote the edge whose spectral parameter is z by  $e_z = (v_1, v_2)$  and assume  $i_{v_1} = a$ . Let  $e_1 = (v_3, v_1)$  be the edge leading to  $e_z$ . Let  $\mathcal{G}^{(j)}$  be the same hypergraph as  $\mathcal{G}$  but with  $i_v = j$ . Let  $\mathbb{T} := \{\underline{i}\} \setminus \{a\}$ .

**Definition 3.17.** For a z-loop admissible hypergraph  $\mathcal{G}$  as above, we let  $\mathcal{G}_z$  be the hypergraph obtained by removing the loop containing z from  $\mathcal{G}$  and let  $L_z(\mathcal{G})$  be the length of the loop containing z. We furthermore define a set of hypergraphs  $F_1(\mathcal{G})$  as follows. It is the union of the following sets.

- (i) The hypergraphs  $D_1(\mathcal{G}^{(j)}, v_1, k, e_1)$  for all j, k such that  $j \notin \mathbb{T}$  and  $k \notin \mathbb{T} \cup \{j\}$ .
- (ii) The hypergraphs  $D_2(\mathcal{G}^{(j)}, e_z, k, e_1)$  for all j, k such that  $j \notin \mathbb{T}$  and  $k \notin \mathbb{T} \cup \{j\}$ .
- (iii) The hypergraphs  $D_3(\mathcal{G}^{(j)}, e_z, e, k)$  and  $D_4(\mathcal{G}^{(j)}, e_z, e, k)$  where e varies over  $e \in E \setminus \{e_z, e_1\}$ and the indices j, k vary over  $j \notin \mathbb{T}$  and  $k \notin \mathbb{T} \cup \{j\}$ .
- (iv) The hypergraphs  $D_5(\mathcal{G}^{(j)}, e_z, k, u)$  where  $j \notin \mathbb{T}$  and  $k \notin \mathbb{T} \cup \{j\}$ .

Let now  $\mathcal{G}_0$  be a z-loop admissible hypergraph s.t.  $L_z(\mathcal{G}_0) \geq 3$  and u = 0. Assume that all entries of w line in  $\Omega_{\mathfrak{a}}$  for some  $\mathfrak{a} > 0$ . We define the following sets inductively. Let  $\mathcal{F}_0 := \{\mathcal{G}_0\}$  and for  $n \geq 1$  set,

$$\mathcal{F}_n := \bigcup_{\mathcal{H} \in \mathcal{F}_{n-1}} F_1(\mathcal{H}).$$
(3.32)

By inspection, one finds the following:

**Lemma 3.18.** (i) For any  $n \ge 1$ , all hypergraphs in  $\mathcal{F}_n$  contain n more edges than  $\mathcal{G}_0$ .

- (ii) Every graph in any  $\mathcal{F}_n$  is z-loop admissible.
- (iii) For every  $\mathcal{H} \in \mathcal{F}_n$  we have  $\hat{\Psi}_1(\mathcal{H}) \leq \tilde{\Psi}_1(\mathcal{G}_0) = \hat{\Psi}_1(\mathcal{G}_0)$ .
- (iv) If  $\mathcal{H}_1 \in F_1(\mathcal{H})$  for some  $\mathcal{H} \in \mathcal{F}_n$ , then

$$\tilde{\Psi}_1(\mathcal{H}_1) = \Psi_1(v)\tilde{\Psi}_1(\mathcal{H}) \tag{3.33}$$

for some v that equals some spectral parameter of  $\mathcal{H}$ .

The next lemma translates Lemma 3.16 into a form useable for our iteration argument.

{sec:loop-iter

{def:loops-fam

{lem:iterative

{lem:loops-exp

**Lemma 3.19.** Let  $n \ge 0$ . Uniformly in  $\mathcal{H} \in \mathcal{F}_n$  we have,

$$T(\mathcal{H}) = \mathbf{1}_{\{L_z(\mathcal{H})=2\}} a_1 T(\hat{\mathcal{H}}_z) + \sum_{\mathcal{H}_1 \in F_1(\mathcal{H})} a_{\mathcal{H}_1} T(\mathcal{H}_1) \\ + \mathcal{O}_{\prec}(\Psi_1(z)\hat{\Psi}_1(\mathcal{H}))$$
(3.34) {eqn:loops-exp

for some coefficients obeying

$$\sum_{\mathcal{H}_1 \in F_1(\mathcal{H})} |a_{\mathcal{H}_1}| \le C_n, \qquad |a_1| \le \frac{C_n}{N |\mathrm{Im}[z]|}.$$
(3.35)

**Proof.** This follows almost directly from applying Lemma 3.16 to the expectation in the integrand of the definition of  $T(\mathcal{H})$  in (2.42). In particular, the terms with  $Y_k$  and  $Z_k$  on the RHS of (3.29) correspond to the first term on the RHS of (3.34), and  $X_k$  terms correspond to to the sum on the RHS. The only thing to verify is that the error term on the RHS of (3.29) can be integrated in the  $u_i$  to obtain the error  $\mathcal{O}(\Psi_1(z)\hat{\Psi}_1(\mathcal{H}))$ . First, note that the error terms on the RHS of (3.29) are  $\mathcal{O}_{<}(\Psi_1(z)\tilde{\Psi}_1(\mathcal{H}))$  and are holomorphic functions in the spectral parameters of  $\mathcal{H}$  (as the error term is a difference of holomorphic functions). Let us assume that the hypergraph  $\mathcal{H}$  contains K independent integration variables  $u_1, \ldots, u_K$  (note that any single  $u_i$  may be repeated amongst the non-zero entries of  $\boldsymbol{u}$ ). We have that,

$$\tilde{\Psi}_1(\mathcal{H}) \le \hat{\Psi}_1(\mathcal{H}) \prod_{j=1}^K \Psi_1(u_j).$$
(3.36)

Therefore, the error on the RHS of (3.29) integrates to at most,

$$\mathcal{O}_{\prec}\left((|\lambda|\Phi)^{K}\Psi_{1}(z)\hat{\Psi}_{1}(\mathcal{H})\right)$$
(3.37)

by repeatedly applying (2.26).

We single out the case where  $L_z(\mathcal{H}) = 2$  for special treatment, as a naive estimate of the size of these terms would be too large.

**Lemma 3.20.** Let  $\mathcal{H} \in \mathcal{F}_n$  for some  $n \ge 0$ . Assume that  $L_z(\mathcal{H}) = 2$ , and let  $\mathcal{J} = \hat{\mathcal{H}}_z$ . Then, with  $a_1$  as in (3.34),

$$|a_1 T(\mathcal{J})| = \mathcal{O}_{\prec}(\Psi_1(z)\tilde{\Psi}_1(\mathcal{G}_0)).$$
(3.38)

**Proof.** First note that  $n \ge 1$  because by assumption  $L_z(\mathcal{G}_0) \ge 3$ . Let  $\hat{z}$  be the spectral parameter of the other edge in this loop. It either equals  $w_i$  for some  $w_i$  in the vector  $\boldsymbol{w}$  of  $\mathcal{H}$  or  $u_i$  for some integration variable. In either case, we claim that there is an edge  $e \in \mathcal{J}$  s.t.  $z_e = \hat{z}$ .

Consider the sequence of parents of  $\mathcal{H}$ , i.e., the sequence of graphs  $\hat{J}_i$  s.t.  $\hat{J}_0 = \mathcal{G}$ ,  $\hat{J}_i \in F_1(\hat{J}_{i-1})$  and  $\hat{J}_n = \mathcal{H}$ . Each of these graphs arise from one of the actions  $D_1, D_2, D_3, D_4$ or  $D_5$  as described above. Let k be the smallest integer s.t. the graphs  $\{\hat{J}_l\}_{k \leq l \leq n}$  all have the property that the loop containing z is of size 2 and the other spectral parameter is  $\hat{z}$ . Then  $1 \leq k \leq n$ . Consider how  $\hat{J}_k$  was constructed from  $\hat{J}_{k-1}$ . Since actions  $D_1, D_2$  and  $D_5$  either leave the z-loop unchanged or increase its length by 1, it cannot have arisen from either of  $\{lem:loop-2\}$ 

these operations. So it must have come from either  $D_3$  or  $D_4$ . But this action creates an additional edge elsewhere in the graph with the same spectral parameter of the edge that gets connected to the edge containing z. Therefore our claim that there is an edge  $e \in \mathcal{J}$  with  $z_e = \hat{z}$  is proven.

Now, using this and Lemma 3.18 we see that whether or not  $\hat{z}$  is an integration variable, the parameter  $\tilde{\Psi}_1(\mathcal{J})$  has at least one copy of  $\Psi_1(u_i)$  for each integration variable and at least one copy of  $\Psi(v)$  for all v that are not equal to z in the original graph  $\mathcal{G}_0$  (recall that we assume that  $\mathcal{G}_0$  has no integration variables). Therefore, if there are K independent variables of integration in  $\mathcal{H}$  we have that,

$$|a_1 T(\mathcal{J})| = \mathcal{O}_{\prec} \left( \frac{1}{N |\mathrm{Im}[z]|} (|\lambda| \Phi)^K \tilde{\Psi}_1(\mathcal{G}) \Psi_1(z)^{-1} \right) = \mathcal{O}_{\prec}(\Psi_1(z) \tilde{\Psi}_1(\mathcal{G})).$$
(3.39)

This completes the proof.

The following allows us to truncate our iterative expansion at large n.

**Lemma 3.21.** Let  $\mathcal{H} \in \mathcal{F}_n$ . Let  $\min_{z \in \mathcal{G}} N \operatorname{Im}[z] \ge N^{4\alpha}$  and  $|\lambda| \Phi \le N^{-4\alpha}$  for some  $\alpha > 0$  satisfying  $\alpha < \mathfrak{a}/10$ . Then,

$$|T(\mathcal{H})| \le N^{-n\alpha}.\tag{3.40}$$

for all N sufficiently large, depending only on n.

**Proof.** Suppose that there are K independent variables of integration in  $\mathcal{H}$ . Note that  $n \geq K$ . Then we have,

$$\mathcal{M}(\mathcal{H}) = \mathcal{O}_{\prec}(N^{-(n-K)2\alpha} \prod_{i=1}^{K} \Psi_1(u_i))$$
(3.41)

and so the claim follows after repeatedly applying (2.26).

#### 3.2.1 Proof of Proposition 3.2

For any n, we have by iterating (3.34) and applying Lemma 3.20 that,

$$|T(\mathcal{G})| < \Psi_1(z)\tilde{\Psi}_1(\mathcal{G}) + \max_{\mathcal{H}\in\mathcal{F}_n} |T(\mathcal{H})|.$$
(3.42)

For large enough n, the second term on the RHS is less than  $\Psi_1(z)\tilde{\Psi}_1(\mathcal{G})$  by Lemma 3.21 and so the claim follows.

#### **3.3** Better estimate in L = 2 case

The estimate of Proposition 3.2 applied in the case when the loop containing the spectral parameter z was of length at least 3. In this section we establish an estimate when the loop is of length 2 and the monomial  $M_2$  is of a specific form.

Specifically, we consider terms of the type,  $\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)X_n]$ , where

$$X_n := \prod_{k=3}^{2+n} (G_{kk}(v_k) - m_{\rm sc}(v_k)).$$
(3.43)

 $\{\texttt{sec:short-loo}$ 

{sec:main-loop

{lem:loops-ex]

and  $n \ge 1$  with  $X_0 = 1$ . Define now,

$$\tilde{\Psi} := \Psi(z)\Psi(w) \prod_{k=3}^{2+m} \Psi(v_k), \qquad \tilde{\Psi}_1 := \Psi_1(z)\Psi_1(w) \prod_{k=3}^{2+m} \Psi_1(v_k), \qquad (3.44)$$

We let  $\{\underline{i}\}$  be the set of indices appearing in  $X_n$  and  $G_{12}(z)G_{21}(w)$  and once again let  $\mathbb{T} := \{\underline{i}\}\setminus\{1\}$ .

**Lemma 3.22.** We have for  $|\lambda \Phi| \leq 1$ ,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)X_n] = \sum_j \frac{s_{1j}}{N}\mathbb{E}[\partial_{j1}(G_{j2}(z)G_{21}(w)X_ne_{\mathfrak{a}})] + \mathcal{O}_{\prec}(N^{-1/2}\tilde{\Psi})$$
(3.45)

**Proof.** This follows immediately from (3.7) and Lemma 3.6.

The following estimates the second order terms in the above expansion.

**Lemma 3.23.** We have for  $j \in \{\underline{i}\}$  and  $|\lambda \Phi| \leq 1$ ,

$$N^{-1}\mathbb{E}[\partial_{j1}(e_{\mathfrak{a}}G_{j2}(z)G_{21}(w)X_{n})] = -\frac{m_{\rm sc}(z)m_{\rm sc}(w)^{2}}{N}\mathbb{E}[X_{n}]\delta_{j=2} + \mathcal{O}_{\prec}(N^{-1/2}\tilde{\Psi})$$
(3.46) {eqn:weakgij-s

We have for  $j \notin \{\underline{i}\}$ 

$$|\mathbb{E}[G_{j2}(z)G_{21}(w)X_n\partial_{j1}e_{\mathfrak{a}}]| < |\lambda\Phi|\tilde{\Psi}_1\min\{\Psi_1(z),\Psi_1(w),\Phi\}$$
(3.47) {eqn:weakgij-second second se

and

$$|\mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{21}(w)\partial_{j1}X_n]| < \tilde{\Psi}_1\min\{\Psi_1(w),\Psi_1(z)\}.$$

$$(3.48) \quad \{\texttt{eqn:weakgij-set}\}$$

Finally,

$$\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} X_n \partial_{j1}(G_{j2}(z) G_{21}(w))] = -m_{\mathrm{sc}}(z) \mathbb{E}[G_{12}(z) G_{21}(w) X_n e_{\mathfrak{a}}] - m_{\mathrm{sc}}(w) \sum_{j \notin \mathbb{T}} \frac{s_{1j}}{N} \mathbb{E}[G_{j2}(z) G_{2j}(w) X_n e_{\mathfrak{a}}] - \sum_{j \notin \underline{i}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{j2}(z) G_{2j}(w) X_n (G_{11}(w) - m_{\mathrm{sc}}(w)) e_{\mathfrak{a}}] - \sum_{j \notin \{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[G_{12}(z) G_{21}(w) (G_{jj}(z) - m_{\mathrm{sc}}(z)) X_n e_{\mathfrak{a}}] + \mathcal{O}_{<}(\tilde{\Psi}_1 \min\{\Psi_1(z), \Psi_1(w)\})$$
(3.49) {eqn:weakgij-s

**Proof.** The estimate (3.46) follows easily using (2.35), (2.36) and (2.10) and direct computation. For (3.47) we have

$$\mathbb{E}[G_{j2}(z)G_{21}(w)X_n\partial_{1j}e_{\mathfrak{a}}] = \frac{-2\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u \mathbb{E}[G_{j1}(u)G_{j2}(z)G_{21}(w)X_3]\mathrm{d}u \qquad (3.50)$$

The expectation in the integral is  $\mathcal{O}_{\prec}(\tilde{\Psi}_1\Psi_1(u)\min\{\Psi_1(z),\Psi_1(w),\Psi_1(u)\})$  by Proposition 3.2, and so (3.47) follows by applying (2.26).

Since  $\partial_{j1}(G_{kk}(v_k) - m_{sc}(v_k)) = -2G_{k1}(v_k)G_{kj}(v_k)$ , the estimate (3.48) follows from Proposition 3.2. The estimate (3.49) follows in a similar manner, by applying Proposition 3.2. Note that we added back the term with j = 1 in the first sum on the RHS of (3.49) at a cost of  $\mathcal{O}_{<}(N^{-1}\tilde{\Psi}_1)$ .

{lem:weakgij-e

{lem:weakgij-s

Recalling the definition of  $\mathbb{T}$  above, define

$$v_j := \mathbb{E}[G_{j2}(z)G_{2j}(w)e_{\mathfrak{a}}X_n]\mathbf{1}_{\{j\notin\mathbb{T}\}}.$$
(3.51)

Then, Lemmas 3.22 and 3.23 imply, (recall A and B defined in (3.27)),

$$(1 - m_{\rm sc}(z)m_{\rm sc}(w)(A - B))v = m_{\rm sc}(z)m_{\rm sc}(w)N^{-1}\mathbf{e}\mathbf{e}^{T}v + U + V + \varepsilon \qquad (3.52) \quad \{\mathtt{eqn:weakgij-s} \}$$

where  $\|\varepsilon\|_{\infty} \leq \Psi_1 \min\{\Psi_1(z), \Psi_1(w)\}$  and

$$U_j = \mathbf{1}_{\{j \notin \mathbb{T}\}} \frac{m_{\rm sc}(z)^2 m_{\rm sc}(w)^2 s_{j2}}{N} \mathbb{E}[e_{\mathfrak{a}} X_n]$$
(3.53)

and

$$V_{j} = \mathbf{1}_{\{j \notin \mathbb{T}\}} m_{\rm sc}(z) \sum_{k \notin \mathbb{T} \cup \{j\}} \frac{s_{jk}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{k2}(z)G_{2k}(w)X_{n}(G_{kk}(w) - m_{\rm sc}(w))] + \mathbf{1}_{\{j \notin \mathbb{T}\}} m_{\rm sc}(z) \sum_{k \notin \mathbb{T} \cup \{j\}} \frac{s_{jk}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{j2}(z)G_{2j}(w)X_{n}(G_{kk}(z) - m_{\rm sc}(z))]$$
(3.54)

We therefore may obtain the following.

**Lemma 3.24.** For  $|\lambda \Phi| \leq 1$  we have,

$$\|v\|_{\infty} < \left( \left| \frac{m_{\rm sc}(z) - m_{\rm sc}(w)}{N(z - w)} \right| + N^{-1} \right) |\mathbb{E}[e_{\mathfrak{a}}X_n]| + \|V\|_{\infty} + \tilde{\Psi}_1 \min\{\Psi_1(z), \Psi_1(w)\}.$$
(3.55) {eqn:weakgij-i

**Proof.** By applying Lemma C.1 (similarly to the proof of Lemma 3.16) one sees that

$$|N^{-1}\mathbf{e}^{T}v| < \left(\left|\frac{m_{\rm sc}(z) - m_{\rm sc}(w)}{N(z - w)}\right| + N^{-1}\right) |\mathbb{E}[e_{\mathfrak{a}}X_{n}]| + \min\{\Psi_{1}(z), \Psi_{1}(w)\}\tilde{\Psi}_{1}.$$
 (3.56)

The claim now follows from (3.52) and Lemma B.1 (again, similar to the proof of Lemma 3.16).

Iterating the above, we can quickly conclude the following.

**Lemma 3.25.** We have for  $|\lambda \Phi| \leq 1$ , and  $u, v, w \in \Omega_{\mathfrak{a}}$ ,

**Proof.** Consider a vector v consisting of expectations of the form  $\mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{2j}(w)X_n]$ where  $X_n$  is a product of n terms of the form  $G_{kk}(v_k) - m_{sc}(v_k)$ . Then the vector V on the RHS of (3.55) is also of this form, except it has an additional term of the form  $G_{kk}(v) - m_{sc}(v)$ where v is either z or w. We therefore can iterate (3.55) repeatedly to the term  $\|V\|_{\infty}$  that appears on the RHS. The first iteration starts with the term on the LHS of (3.57). For this first iteration, the first term on the RHS of (3.55) is bounded by the first term on the RHS of (3.57). For the subsequent iterations, the first term on the RHS of (3.55) can be absorbed into the last term on the RHS of (3.57). At all steps of the iteration the last error term on the RHS of (3.55) can be absorbed into the last term on the RHS of (3.57). The iteration may be terminated in a fashion similar to the proof of Proposition 3.2. {lem:weak-g12g

# 4 Line estimate

In Section 3 we obtained estimates for expectations of resolvent entries whose associated graphs was a product of loops. In this section we obtain an estimate for hypergraphs containing a line. We again assume throughout that f is an admissible function. The overall strategy is roughly similar in that we will iterate the cumulant expansion and inspect which terms arise.

**Definition 4.1.** We will say that a hypergraph is line admissible if

- (i) Every entry of *i* is unique
- (ii) It is a disjoint union of loops and exactly one line.

We will assume  $\mathcal{M}(\mathcal{G}) = G_{12}(z)M_1M_2$  where  $M_1 = G_{23}G_{34}\ldots G_{K-1,K}$  and  $M_2$  is a disjoint union of loops. The parameter z will not play a particularly special role here, unlike the argument in Section 3. Our goal is to prove the following. The proof appears in Section 4.1.1.

**Proposition 4.2.** Let  $\mathcal{G}$  be a line admissible hypergraph with  $\boldsymbol{u} = 0$  and all  $z_e \in \Omega_{\mathfrak{a}}$ , and assume  $|\lambda \Phi| \leq N^{-c}$  for some c > 0. Then we have,

$$\left|\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(\mathcal{G})]\right| < N^{-1/2}\tilde{\Psi}.$$
(4.1)

We first develop some estimates in preparation for applying the cumulant expansion.

Lemma 4.3. The term,

obeys the following estimates:

(i) If 
$$j = 1$$
 it is  $\mathcal{O}_{<}(\Psi)$ . {it:line-der-

- (ii) If j = 2 it is  $\mathcal{O}_{\prec}(\tilde{\Psi}\Psi(z)^{-1})$ . {it:line-der-2
- (iii) If  $j \in \{i\}$  but  $j \neq 1, 2$ , then it is  $\mathcal{O}_{<}(\tilde{\Psi})$  as long as j is not the index of a self-loop  $(G_{jj}(v) m_{sc}(v))$ . If j is the index of a self-loop  $(G_{jj}(v) m_{sc}(v))$  then any derivative of order at least 2 is  $\mathcal{O}_{<}(\tilde{\Psi}\Psi(v)^{-1})$  but the first derivative is still  $\mathcal{O}_{<}(\tilde{\Psi})$  {it:line-der-3
- (iv) If  $j \notin \{\underline{i}\}$ , then it is  $\mathcal{O}_{\prec}(\tilde{\Psi})$

If  $\delta > 0$  and M is  $\varepsilon$ -regular, than the above estimates hold with an  $N^{\delta}$  factor as long as  $\varepsilon$  is sufficiently small.

**Proof.** The proof is almost identical to Lemma 3.3 and so is omitted.

**Lemma 4.4.** We have that for  $|\lambda \Phi| \leq 1$  that,

$$N^{-2}|\partial_{j1}^3(e_{\mathfrak{a}}G_{j2}M_1M_2e_{\mathfrak{a}})|$$

$$(4.3) \quad \{\texttt{eqn:line-foul}\}$$

is  $\mathcal{O}_{\prec}((1+|\lambda|)^2 \tilde{\Psi}/N^{3/2})$  for  $j \in \{\underline{i}\}$  and  $\mathcal{O}_{\prec}((1+|\lambda|)N^{-2}\tilde{\Psi})$  otherwise. The same estimates hold with an additional factor of  $N^{\delta}$  if M is  $\varepsilon$ -regular, for  $\varepsilon > 0$  sufficiently small.

28

( . . .

{prop:main-lin

{lem:line-four

{lem:line-der}

{def:line-admi

 $\{\texttt{sec:line}\}$ 

{it:line-der-4

**Proof.** When j = 1, the derivatives hitting  $e_{\mathfrak{a}}$  contribute at most  $(1 + |\lambda|)^3$  by (2.36), and the derivatives hitting resolvent entries can be estimated using Lemma 4.3(i), finding  $\mathcal{O}_{<}((1 + |\lambda|)^3 \tilde{\Psi} N^{-2})$ . For j = 2, we use (2.35) and Lemma 4.3(ii) to find an estimate of  $\mathcal{O}_{<}((1 + |\lambda|)\tilde{\Psi}/N^{3/2})$ . For  $j \in \{\underline{i}\}$  and  $j \neq 1, 2$  we use (2.35) and Lemma 4.3(iii) to find an estimate of estimate of  $\mathcal{O}_{<}((1 + |\lambda|)\tilde{\Psi}/N^{3/2})$ .

Finally, the terms when  $j \notin \{\underline{i}\}$  are estimated using (2.35) and Lemma 4.3(iv).

From this and the cumulant expansion, Lemma 2.6 we find the following. The proof is the same as Lemma 3.5 and so omitted.

**Lemma 4.5.** We have for  $|\lambda \Phi| \leq 1$ ,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)M_{1}M_{2}] = \sum_{j} \frac{s_{1j}}{N}\mathbb{E}[\partial_{j1}(G_{j2}(z)M_{1}M_{2}e_{\mathfrak{a}})] + \sum_{j} \frac{s_{1j}^{(3)}}{2N^{3/2}}\mathbb{E}[\partial_{1j}^{2}(G_{j2}(z)M_{1}M_{2}e_{\mathfrak{a}})] + \mathcal{O}_{<}((1+|\lambda|)N^{-1}\tilde{\Psi}).$$
(4.4) {eqn:line-expansion}

We now begin estimating the third order terms on the RHS of (4.4).

**Lemma 4.6.** The following hold for  $|\lambda \Phi| \leq 1$ . We have for  $j \in \{\underline{i}\}$  that,

$$\left| N^{-3/2} \mathbb{E}[\partial_{j1}^2 (e_{\mathfrak{a}} G_{j2} M_1 M_2)] \right| < (1 + |\lambda|) \tilde{\Psi} N^{-1}.$$
(4.5) {eqn:line-thir

For the other terms we have,

Ξ

$$\sum_{j \notin \{\underline{i}\}} \frac{1}{N^{3/2}} |\mathbb{E}[(\partial_{j1} e_{\mathfrak{a}}) \partial_{j1} (G_{j2} M_1 M_2)]| < |\lambda \Phi| N^{-1/2} \tilde{\Psi}$$

$$(4.6) \quad \{\texttt{eqn:line-thing} \in \mathbb{C} \}$$

$$\sum_{j \notin \{\underline{i}\}} \frac{1}{N^{3/2}} |\mathbb{E}[e_{\mathfrak{a}} \partial_{j1}^2 (G_{j2} M_1 M_2)]| < N^{-1/2} \tilde{\Psi}$$

$$(4.7) \quad \{\texttt{eqn:line-this}$$

$$\sum_{j \notin \{\underline{i}\}} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{j1}^2 e_{\mathfrak{a}}) G_{j2} M_1 M_2] = \mathcal{O}_{\prec}(|\lambda| \Phi N^{-1/2} \tilde{\Psi})$$
(4.8) {eqn:line-this

**Proof.** We start by proving (4.5). For j = 1 we use (2.36) and Lemma 4.3(i) to find an estimate of  $\mathcal{O}_{<}((1 + |\lambda|)^2 N^{-3/2} \tilde{\Psi})$ . When j = 2 we use (2.35) and Lemma 4.3(ii) to find an estimate of  $\mathcal{O}_{<}((1 + |\lambda|)\tilde{\Psi}N^{-1})$ . For other  $j \in \{\underline{i}\}$  we use (2.35) and Lemma 4.3(iii) to find an estimate of  $\mathcal{O}_{<}((1 + |\lambda|)\tilde{\Psi}N^{-1})$ . This completes the proof of (4.5).

The estimates (4.6) and (4.7) follow immediately from (2.32) and Lemma 4.3(iv). For (4.8) we use (2.32) to conclude,

$$\sum_{\substack{j \notin \{i\}\\}} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}^2 e_{\mathfrak{a}})G_{j2}M_1M_2] = \mathcal{O}_{\prec}(\tilde{\Psi}|\lambda\Phi|N^{-1/2}) + i\lambda c_f \sum_{\substack{j \notin \{i\}\\\\}} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}G_{j2}M_1M_2]$$
$$= \mathcal{O}_{\prec}(|\lambda\Phi|N^{-1/2}\tilde{\Psi}) + \mathcal{O}_{\prec}(N^{-1}|\lambda|\tilde{\Psi})$$
(4.9)

with the second line following from the isotropic local law (2.9). This completes the proof.  $\Box$ 

We now turn to computing the second order terms on the RHS of (4.4). The following is fairly straightforward.

29

 $\{\texttt{lem:line-expa}\}$ 

{lem:line-thir

{lem:line-seco

{lem:line-seco

{lem:line-seco

**Lemma 4.7.** We have for  $j \in \{\underline{i}\}$  and  $j \neq 2$ ,

$$\frac{1}{N} \left| \mathbb{E} \left[ \partial_{j1} (e_{\mathfrak{a}} G_{j2} M_1 M_2) \right] \right| < (1 + |\lambda|) N^{-1} \tilde{\Psi}$$

$$(4.10) \quad \{ \texttt{eqn:line-second} \}$$

and for j = 2,

$$\frac{1}{N} |\mathbb{E}[\partial_{j1}(e_{\mathfrak{a}}G_{j2}M_1M_2)]| < N^{-1/2}\tilde{\Psi}$$
(4.11)

**Proof.** These estimates follow from using (2.35), (2.36) and items (i), (ii) and (iii) of Lemma 4.3 as appropriate.

We recall here Definition 3.9 of the operation  $D_2$  on hypergraphs. We define here also another operation  $D_6$  on hypergraphs:

**Definition 4.8.** Given a hypergraph  $\mathcal{G} = (V, E, \underline{i}, u, w)$  with integration variables  $\{u_i\}_{i=1}^m$ , an index j and an edge  $e \in E$  we define a new hypergraph  $D_6(\mathcal{G}, j, e)$  with the same integration variables  $\{u_i\}_{i=1}^m$  as follows. We add a new vertex v' and a self-edge (v', v'). We set  $i'_{v'} = j$  and set  $u'_{(v',v')} = u_e$ ,  $w'_{(v',v')} = w_e$ . The other variables are all the same as the original graph. Lemma 4.9. We have,

$$\sum_{\substack{j \notin \{i\}}} \frac{s_{1j}}{N} \mathbb{E}[(\partial_{j1}G_{j2})M_1M_2e_{\mathfrak{a}}] = -m_{\mathrm{sc}}(z)\mathbb{E}[G_{12}(z)M_1M_2e_{\mathfrak{a}}] + \mathcal{O}_{\prec}(N^{-1}\tilde{\Psi})$$

$$-\sum_{\substack{j \notin \{i\}}} \frac{s_{1j}}{N}\mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_6(\mathcal{G}, j, (1, 2))) + \mathcal{M}(D_2(\mathcal{G}, (1, 2), j, (1, 2))))].$$
(4.12) {eqn:line-second

**Proof.** Since  $\partial_{j1}G_{j2} = -G_{jj}G_{12} - G_{1j}G_{j2}$  we have,

$$\sum_{\substack{j \notin \{\underline{i}\}\\N}} \frac{s_{1j}}{N} \mathbb{E}[(\partial_{j1}G_{j2})M_1M_2e_{\mathfrak{a}}] = -\sum_{\substack{j \notin \{\underline{i}\}\\N}} \frac{s_{1j}}{N} m_{\mathrm{sc}}(z) \mathbb{E}[G_{12}(z)M_1M_2e_{\mathfrak{a}}] - \sum_{\substack{j \notin \{\underline{i}\}\\N}} \frac{s_{1j}}{N} \mathbb{E}[G_{12}M_1M_2(G_{jj}(z) - m_{\mathrm{sc}}(z))e_{\mathfrak{a}}] - \sum_{\substack{j \notin \{\underline{i}\}\\N}} \frac{s_{1j}}{N} \mathbb{E}[G_{1j}(z)G_{j2}(z)M_1M_2e_{\mathfrak{a}}]$$
(4.13) {eqn:line-second

The first sum on the last line has terms of the form  $\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(D_6(\mathcal{G}, j, (1, 2))]$ . The second sum on the last line has terms of the form  $\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(D_2(\mathcal{G}, (1, 2), j, (1, 2))]$ . The claim follows from this and the fact that  $\sum_j \frac{s_{1j}}{N} = 1 + \mathcal{O}(N^{-1})$ .

We recall the definitions of the operations on hypergraphs  $D_3, D_4, D_5$  given in Definitions 3.11 and 3.13.

Lemma 4.10.

$$-\sum_{j\notin\{\underline{i}\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}G_{j2}\partial_{j1}(M_{1}M_{2})]$$

$$=\sum_{j\notin\{\underline{i}\}} \sum_{e\in E\setminus\{(1,2)\}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{3}(\mathcal{G},(1,2),e,j)) + \mathcal{M}(D_{4}(\mathcal{G},(1,2),e,j)))]$$
(4.14) {eqn:line-second

and

$$\sum_{\substack{j \notin \{\underline{i}\}\\ \pi}} \frac{s_{1j}}{N} \mathbb{E}[G_{j2}M_1M_2\partial_{j1}e_{\mathfrak{a}}]$$

$$= -\frac{2\mathrm{i}\lambda}{\pi} \sum_{\substack{j \notin \{\underline{i}\}\\ N}} \frac{s_{1j}}{N} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u(\partial_{\bar{u}}\tilde{f}(u))\partial_u \mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(D_5(\mathcal{G},(1,2),j,u))] \mathrm{d}u\mathrm{d}\bar{u} \qquad (4.15) \quad \{\text{eqn:line-second}\}$$

**Proof.** This is direct computation and so omitted.

We collect the above estimates into the following expansion for line admissible hypergraphs.

**Proposition 4.11.** Let  $\mathcal{G}$  be a line admissible hypergraph as above, and assume  $|\lambda \Phi| \leq 1$ . We have,

$$\begin{split} \mathbb{E}[e_{\mathfrak{a}}G_{12}(z)M_{1}M_{2}] &= m_{\mathrm{sc}}(z)\sum_{j\notin\{\underline{i}\}}\frac{s_{1j}}{N}\mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{6}(\mathcal{G},j,(1,2))) + \mathcal{M}(D_{2}(\mathcal{G},(1,2),j,(1,2))))] \\ &+ m_{\mathrm{sc}}(z)\sum_{j\notin\{\underline{i}\}}\sum_{e\in E\setminus\{(1,2)\}}\frac{s_{1j}}{N}\mathbb{E}[e_{\mathfrak{a}}(\mathcal{M}(D_{3}(\mathcal{G},(1,2),e,j)) + \mathcal{M}(D_{4}(\mathcal{G},(1,2),e,j)))] \\ &+ \frac{2\mathrm{i}\lambda m_{\mathrm{sc}}(z)}{\pi}\sum_{j\notin\{\underline{i}\}}\frac{s_{1j}}{N}\int_{\Omega_{\mathfrak{a}}}\mathrm{d}u(\partial_{\bar{u}}\tilde{f}(u))\partial_{u}\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(D_{5}(\mathcal{G},(1,2),j,u))]\mathrm{d}u\mathrm{d}\bar{u} + \mathcal{O}_{<}(N^{-1/2}\tilde{\Psi}) \end{split}$$

$$(4.16) \quad \{\mathrm{eqn:line-exx}\}$$

**Proof.** This follows from (4.4) and Lemmas 4.6, 4.7, 4.9 and 4.10.

# 4.1 Iterative expansion

We now carry out our iterative expansion for line-admissible hypergraphs. We begin by fixing some notation. Let  $\mathcal{G}$  be a line-admissible hypergraph, and let  $e = (v_1, v_2)$  be the first edge in this line. The following families of hypergraphs will appear in our iterative expansion.

**Definition 4.12.** Let  $\mathcal{G}$  and  $e = (v_1, v_2)$  be as above. We define a set of hypergraphs  $F_2(\mathcal{G})$  to be the union of the following sets of hypergraphs.

- (i) The graphs  $D_6(\mathcal{G}, j, e)$  and  $D_2(\mathcal{G}, e, j, e)$  for  $j \notin \{\underline{i}\}$
- (ii) The graphs  $D_3(\mathcal{G}, e, e_1, j)$  and  $D_4(\mathcal{G}, e, e_1, j)$  for  $j \notin \{\underline{i}\}$  and  $e_1 \in E \setminus \{e\}$ .
- (iii) The graphs  $D_5(\mathcal{G}, e, j, u)$  for  $j \notin \{\underline{i}\}$ .

For any hypergraph  $\mathcal{G}$  we recall that  $E_w$  is the support of the vector w and define,

$$\hat{\Psi}(\mathcal{G}) := \prod_{e \in E_w} \Psi(z_e).$$
(4.17)

Let now  $\mathcal{G}_0$  be a line-admissible hypergraph s.t.  $\boldsymbol{u} = 0$  and let  $\boldsymbol{e} = (v_1, v_2)$  be the first edge in this line. Set  $\mathcal{F}_0 := {\mathcal{G}_0}$  and inductively set

$$\mathcal{F}_n := \bigcup_{\mathcal{H} \in \mathcal{F}_{n-1}} F_2(\mathcal{H}).$$
(4.18)

By inspection, we find the following.

**Lemma 4.13.** (i) For any  $n \ge 1$ , all hypergraphs in  $\mathcal{F}_n$  contain n more edges than  $\mathcal{G}_0$ 

- (ii) Every graph in  $\mathcal{F}_n$  is line admissible
- (iii) For every  $H \in \mathcal{F}_n$  we have  $\hat{\Psi}(\mathcal{H}) \leq \tilde{\Psi}(\mathcal{G}_0) = \hat{\Psi}(\mathcal{G}_0)$

{sec:line-iter

{prop:line-exp

(iv) If  $\mathcal{H}_1 \in F_2(\mathcal{H})$  for some  $\mathcal{H} \in \mathcal{F}_n$ , then

$$\tilde{\Psi}(\mathcal{H}_1) = \Psi(v)\tilde{\Psi}(\mathcal{H}) \tag{4.19}$$

for some v that is a spectral parameter of  $\mathcal{H}$ .

**Lemma 4.14.** Let  $n \ge 0$ . Uniformly in  $\mathcal{H} \in \mathcal{F}_n$  we have,

$$T(\mathcal{H}) = \sum_{\mathcal{H}_1 \in \mathcal{F}_2(\mathcal{H})} b_{\mathcal{H}_1} T(\mathcal{H}_1) + \mathcal{O}_{<}(\tilde{\Psi}(\mathcal{G}_0) N^{-1/2})$$
(4.20) {eqn:line-expansion}

for some coefficients obeying,

$$\sum_{L_1 \in F_2(\mathcal{H})} |b_{\mathcal{H}_1}| \le C_n.$$
(4.21)

Let  $\alpha > 0$  satisfy  $\alpha < \frac{\mathfrak{a}}{10}$  as well as  $\min_{z \in \mathcal{G}} N |\operatorname{Im}[z]| \ge N^{4\alpha}$ , and  $|\lambda \Phi| \le N^{-4\alpha}$ . Then for all  $\mathcal{H} \in \mathcal{F}_n$  we have,

$$|T(\mathcal{H})| \le N^{-n\alpha}$$
. (4.22) {eqn:line-expa}

**Proof.** The expansion (4.20) follows from Proposition 4.11 in the exact same way as Lemma 3.19 follows from Lemma 3.16, and so we omit the detailed proof. Similarly, (4.22) follows in the same way as Lemma 3.21.

### 4.1.1 Proof of Proposition 4.2

By iterating (4.20) we see that,

$$|\mathbb{E}[e_{\mathfrak{a}}\mathcal{M}(\mathcal{G})]| < \max_{\mathcal{H}\in\mathcal{F}_n} |T(\mathcal{H})| + N^{-1/2}\tilde{\Psi}(\mathcal{G}).$$
(4.23)

The claim now follows from taking n sufficiently large and using (4.22).

# 4.2 A better estimate for $G_{ij}$

In this section we derive an even better estimate for  $\mathbb{E}[e_{\mathfrak{a}}G_{ij}(z)]$  with  $i \neq j$  than was derived in Proposition 4.2. We start first with the following. We set  $\mathbb{T} = \{1, 2\}$ .

**Lemma 4.15.** For  $|\lambda \Phi| \leq 1$  we have,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)] = \sum_{j\neq 1} \frac{s_{1j}}{N} \mathbb{E}[\partial_{j1}(G_{j2}(z)e_{\mathfrak{a}})] + \mathcal{O}_{\prec}(|\lambda|\Phi N^{-1/2}\Psi(z) + N^{-1/2}\Psi(z)^{2}).$$
(4.24) {eqn:gij-expansion}

**Proof.** First, by combining Lemma 4.5, (4.5), (4.6), (4.8) and (4.10) we see that,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)] = \sum_{j\neq 1} \frac{s_{1j}}{N} \mathbb{E}[\partial_{1j}(G_{j2}(z)e_{\mathfrak{a}})] + \sum_{j\notin \mathbb{T}} \frac{s_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}\partial_{1j}^{2}(G_{j2})] + \mathcal{O}_{\prec}((1+|\lambda|)N^{-1}\Psi(z) + |\lambda|\Phi N^{-1/2}\Psi(z)).$$
(4.25) {eqn:gij-expansion}

 $\{\texttt{sec:main-line}$ 

{lem:gij-expan

We now compute that for  $j \notin \mathbb{T}$  we have  $\partial_{1j}^2 G_{j2} = 2m_{\rm sc}(z)^2 G_{j2}(z) + \mathcal{O}_{\prec}(\Psi(z)^2)$  to see that,

$$\sum_{j \notin \mathbb{T}} \frac{s_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[e_{\mathfrak{a}} \partial_{1j}^2(G_{j2})] = m_{\mathrm{sc}}(z)^2 \sum_{j \notin \mathbb{T}} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}(G_{j2})] + \mathcal{O}_{\prec}(N^{-1/2}\Psi(z)^2)$$
(4.26) {eqn:gij-expansion}

and the first term on the RHS is  $\mathcal{O}_{<}(N^{-1}\Psi(z))$  by the istropic local law (2.9).

We now turn to computing the second order terms.

**Lemma 4.16.** Assume  $|\lambda \Phi| \leq 1$ . We have,

$$N^{-1}|\mathbb{E}[\partial_{21}(e_{\mathfrak{a}}G_{22})]| < N^{-1}|\lambda\Phi| + N^{-1}\Psi(z).$$
(4.27) {eqn:g12-second

{lem:gij-secon

{prop:gij-est}

and

$$\sum_{j \notin \mathbb{T}} \frac{s_{1j}}{N} \mathbb{E}[G_{j2}(z)\partial_{j1}e_{\mathfrak{a}}] = \mathcal{O}_{\prec}(|\lambda \Phi| N^{-1/2} \Psi(z))$$
(4.28) {eqn:g12-second

and

$$\sum_{j \notin \mathbb{T}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} \partial_{j1} G_{j2}(z)] = -m_{\mathrm{sc}}(z) \mathbb{E}[G_{12}(z) e_{\mathfrak{a}}] + \mathcal{O}_{\prec}(N^{-1/2} \Psi(z)^2).$$
(4.29) {eqn:gl2-second}

**Proof.** For (4.27) we have  $\partial_{21}G_{22} = \mathcal{O}_{\langle}(\Psi(z))$  and  $\partial_{12}e_{\mathfrak{a}} = \mathcal{O}(|\lambda|\Phi)$  by (2.35) and so the estimate follows. For (4.28) we have that,

$$\mathbb{E}[G_{j2}(z)\partial_{j1}e_{\mathfrak{a}}] = -\frac{2\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u}\mathbb{E}[e_{\mathfrak{a}}G_{1j}(u)G_{j2}(z)]\mathrm{d}u\mathrm{d}\bar{u}.$$
(4.30)

From Proposition 4.2 we have  $\mathbb{E}[e_{\mathfrak{a}}G_{1j}(u)G_{j2}(z)] = \mathcal{O}_{<}(N^{-1/2}\Psi(z)\Psi(u))$  and so (4.28) follows by applying (2.26). For (4.29) we have,

$$-\sum_{j\notin\mathbb{T}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}\partial_{j1}G_{j2}(z)] = \sum_{j\notin\mathbb{T}} \frac{s_{1j}}{N} m_{\mathrm{sc}}(z) \mathbb{E}[G_{12}(z)]$$

$$+\sum_{j\notin\mathbb{T}} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}G_{12}(z)(G_{jj}(z) - m_{\mathrm{sc}}(z)] + \sum_{j\notin\mathbb{T}} \frac{s_{1j}}{N} \mathbb{E}[G_{1j}(z)G_{j2}(z)e_{\mathfrak{a}}]$$

$$= m_{\mathrm{sc}}(z) \mathbb{E}[e_{\mathfrak{a}}G_{12}(z)] + \mathcal{O}_{\prec}(\Psi(z)^{2}N^{-1/2})$$

$$(4.31) \quad \{\mathrm{eqn}: \mathtt{gij-second}\}$$

with the last line following from applying Proposition 4.2 twice, as well as that  $\sum_{j\notin\mathbb{T}} \frac{s_{1j}}{N} = 1 + \mathcal{O}(N^{-1})$ .

The above two lemmas easily imply the following, which improves upon the estimate Proposition 4.2 in the special case of a single resolvent entry.

**Proposition 4.17.** If there is a c > 0 so that  $|\lambda \Phi| \leq N^{-c}$  then for  $i \neq j$  we have,

$$|\mathbb{E}[G_{ij}(z)e_{\mathfrak{a}}]| < |\lambda\Phi|N^{-1/2}\Psi(z) + N^{-1/2}\Psi(z)^{2}.$$
(4.32)

# 5 Preliminary expansion for $\mathbb{E}[e_{\mathfrak{a}}G_{11}]$

In this section we derive an expansion for  $\mathbb{E}[e_{\mathfrak{a}}G_{ii}(z)]$ . We will use it to derive an estimate for the expression  $\mathbb{E}[e_{\mathfrak{a}}(G_{ii}(z) - m_{sc}(z)]$  that is better than the naive estimate  $\mathcal{O}_{<}(\Psi(z))$ . Secondly, the first part of our expansion will be used to derive our main estimates on the characteristic function. Throughout this section we will use the notation  $z = E + i\eta$ , i.e.,  $\eta = |\mathrm{Im}[z]|$ . Throughout this section we will always assume that,

$$|\lambda \Phi| \le N^{-c} \tag{5.1}$$

for some c > 0 and that  $z \in \Omega_{\mathfrak{a}}$ .

We start with the following.

Lemma 5.1. We have that,

$$N^{-5/2} |\partial_{11}^4(e_{\mathfrak{a}} G_{11}(z))| < (1+|\lambda|)^4 N^{-5/2}.$$
(5.2)

and that for  $j \neq 1$ ,

$$N^{-5/2}|\partial_{j1}^4(e_{\mathfrak{a}}G_{1j}(z))| < N^{-5/2}\Psi(z)(1+|\lambda|)^2 + N^{-5/2}|\lambda\Phi|(1+|\lambda|)$$
(5.3) {eqn:G11-fifth

For any  $\delta > 0$ , the same estimates hold for any  $\varepsilon$ -regular matrix with an additional factor of  $N^{\delta}$  on the RHS, if  $\varepsilon$  is sufficiently small depending on  $\delta$ .

**Proof.** The estimate for j = 1 follows directly from (2.36). We turn to the estimate for  $j \neq 1$ . Note that for k = 0, 2, 4 we have that,

$$\partial_{1j}^k G_{1j} = \mathcal{O}_{\prec}(\Psi(z)) \tag{5.4}$$

by direct computation. The LHS is  $\mathcal{O}_{\prec}(1)$  otherwise. Furthermore, for k = 1, 3 we have

$$\partial_{1j}^{k} e_{\mathfrak{a}} = \mathcal{O}_{\prec}(|\lambda \Phi|(1+|\lambda|), \qquad (5.5) \quad \{\texttt{eqn:delea3}\}$$

and for k = 2, 4 it is  $\mathcal{O}_{<}((1 + |\lambda|)^2)$ . Here, the k = 1 case follows from (2.31), and the k = 2, 4 cases from (2.35). The k = 3 case follows from a direct computation similar to (2.32). The estimate (5.3) now follows. The statement about  $\varepsilon$ -regular matrices is now straightforward.

We now apply the cumulant expansion to arrive at our starting point for estimating  $\mathbb{E}[e_{\mathfrak{a}}G_{11}(z)].$ 

#### Lemma 5.2. We have,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{11}(z)] + \mathbb{E}[e_{\mathfrak{a}}] = \sum_{j} \frac{s_{1j}}{N} \mathbb{E}[\partial_{1j}(e_{\mathfrak{a}}G_{j1})] + \sum_{j} \frac{s_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[\partial_{1j}^{2}(e_{\mathfrak{a}}G_{j1})] + \sum_{j} \frac{s_{1j}^{(4)}}{6N^{2}} \mathbb{E}[\partial_{1j}^{3}(e_{\mathfrak{a}}G_{j1})] + \mathcal{O}_{\prec}((1+|\lambda|)^{4}N^{-5/2} + \Psi(z)N^{-3/2}(1+|\lambda|)^{2} + N^{-3/2}|\lambda\Phi|(1+|\lambda|))$$

$$(5.6) \quad \{\texttt{eqn:prelim-ex}\}$$

(2)

{sec:prelim}

{lem:G11-fifth

{lem:prelim-ex

**Proof.** This follows from the cumulant expansion of Lemma 2.6 to fifth order applied to,

$$z\mathbb{E}[G_{11}e_{\mathfrak{a}}] + \mathbb{E}[e_{\mathfrak{a}}] = \sum_{j} \mathbb{E}[H_{1j}G_{j1}e_{\mathfrak{a}}], \qquad (5.7)$$

using the estimates of Lemma 5.1.

In Sections 5.1 and 5.2 below, we will compute the fourth and third order terms, respectively, on the RHS of (5.6). The results will then be summarized in Section 5.3.

### 5.1 Fourth order terms

We first prove the following.

Lemma 5.3. We have,

$$\begin{split} &\sum_{j} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[\partial_{1j}^3(e_{\mathfrak{a}}G_{j1})] = -6 \sum_{j \neq 1} \frac{s_{1j}^{(4)}}{N^2} m_{\mathrm{sc}}(z)^4 \mathbb{E}[e_{\mathfrak{a}}] \\ &+3 \sum_{j \neq 1} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[(\partial_{1j}^2 e_{\mathfrak{a}})(\partial_{1j}G_{j1})] - 12 m_{\mathrm{sc}}(z)^3 \sum_{j \neq 1} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[e_{\mathfrak{a}}(G_{11}(z) - m_{\mathrm{sc}}(z))] \\ &+ \mathcal{O}_{\prec}(N^{-2}\eta^{-1} + (1 + |\lambda|)^3 N^{-2} + |\lambda \Phi| \Psi(z)(1 + |\lambda|) N^{-1}) \end{split}$$
(5.8) {eqn:prelim-formula}

**Proof.** The term when j = 1 can be estimated by  $\mathcal{O}_{<}(N^{-2}(1 + |\lambda|)^3)$  using (2.36). When  $j \neq 1$  we have that by direct computation and (2.10)

$$\partial_{j1}^3 G_{j1} = -6G_{11}^2 G_{jj}^2 + \mathcal{O}(\Psi(z)^2)$$
(5.9) {eqn:prelim-fo

and so,

$$\begin{aligned} & \frac{-1}{6} \sum_{j \neq 1} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[e_{\mathfrak{a}} \partial_{1j}^3(G_{j1})] = \sum_{j \neq 1} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[e_{\mathfrak{a}} G_{11}(z)^2 G_{jj}(z)^2] + \mathcal{O}_{<}(N^{-1}\Psi(z)^2) \\ & = \sum_{j \neq 1} \frac{s_{1j}^{(4)}}{N^2} \left\{ m_{\mathrm{sc}}(z)^4 \mathbb{E}[e_{\mathfrak{a}}] + 2m_{\mathrm{sc}}(z)^3 \mathbb{E}[e_{\mathfrak{a}}(G_{11}(z) + G_{jj}(z) - 2m_{\mathrm{sc}}(z))] \right\} + \mathcal{O}_{<}(N^{-1}\Psi(z)^2) \\ & = \sum_{j \neq 1} \frac{s_{1j}^{(4)}}{N^2} m_{\mathrm{sc}}(z)^4 \mathbb{E}[e_{\mathfrak{a}}] + 2m_{\mathrm{sc}}(z)^3 \sum_{j \neq 1} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[e_{\mathfrak{a}}(G_{11}(z) - m_{\mathrm{sc}}(z))] + \mathcal{O}_{<}(N^{-2}\eta^{-1}) \quad (5.10) \quad \{\mathrm{eqn:prelim-fo}\} \end{aligned}$$

The second line follows from writing  $G_{ii} = (G_{ii} - m_{sc}) + m_{sc}$  and expanding, and dropping all contributions quadratic and higher in  $(G_{ii} - m_{sc})$  (and similarly for the  $G_{jj}$  term). The last line follows from fluctuation averaging (2.7). This takes care of the terms when all of the  $\partial_{ij}$  hit  $G_{j1}$ .

We estimate the other contributions using,

$$(\partial_{1j}e_{\mathfrak{a}})(\partial_{1j}^{2}G_{1j}) = \mathcal{O}_{\prec}(|\lambda|\Phi\Psi(z)), \qquad \partial_{1j}^{3}e_{\mathfrak{a}} = \mathcal{O}_{\prec}(|\lambda\Phi|(1+|\lambda|))$$
(5.11)

which hold for  $j \neq 1$ . The first estimate follows from (2.31) and (2.10), and the second from (5.5).

For the remaining fourth order terms we compute,

35

 $\{\texttt{sec:prelim-fo}$ 

{lem:prelim-fo

{lem:prelim-fo

Lemma 5.4. We have,

$$\begin{split} \sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[(\partial_{1j}^2 e_{\mathfrak{a}})(\partial_{1j}G_{j1})] &= \mathcal{O}_{<}(|\lambda|N^{-2}\eta^{-1} + |\lambda|(1+|\lambda|)\Phi^2 N^{-1}) \\ &- \mathbb{E}[e_{\mathfrak{a}}]m_{\mathrm{sc}}(z)^2 \sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \frac{2\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u m_{\mathrm{sc}}(u)^2 \mathrm{d}u \mathrm{d}\bar{u} \\ &- \sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \frac{2\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u (m_{\mathrm{sc}}(z)^2 m_{\mathrm{sc}}(u)\mathbb{E}[e_{\mathfrak{a}}(G_{11}(u) - m_{\mathrm{sc}}(u))]) \mathrm{d}u \mathrm{d}\bar{u} \\ &- \sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \frac{2\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u (m_{\mathrm{sc}}(u)^2 m_{\mathrm{sc}}(z)\mathbb{E}[e_{\mathfrak{a}}(G_{11}(z) - m_{\mathrm{sc}}(z))]) \mathrm{d}u \mathrm{d}\bar{u} \end{split}$$
(5.12) {equiv.

**Proof.** By using that

$$\partial_{1j}G_{j1} = -G_{jj}G_{11} + \mathcal{O}_{\prec}(\Psi(z)^2)$$
(5.13) {eqn:prelim-fo

and

$$\partial_{1j}^2 e_{\mathfrak{a}} = \frac{2\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u (G_{11}(u)G_{jj}(u))\mathrm{d}u\mathrm{d}\bar{u} + \mathcal{O}_{<}(|\lambda\Phi^2|(1+|\lambda|)), \qquad (5.14) \quad \{\texttt{eqn:prelim-formula}\}$$

(the above following from (2.34)) we see that,

$$-\sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[(\partial_{1j}^2 e_{\mathfrak{a}})(\partial_{1j}G_{j1})] = \mathcal{O}_{<}(|\lambda|\Psi(z)^2 N^{-1} + |\lambda|(1+|\lambda|)\Phi^2 N^{-1}) \\ + \sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \frac{2i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u \mathbb{E}[e_{\mathfrak{a}}G_{11}(u)G_{jj}(u)G_{11}(z)G_{jj}(z)] dud\bar{u} \\ = \mathbb{E}[e_{\mathfrak{a}}]m_{\mathrm{sc}}(z)^2 \sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \frac{2i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u m_{\mathrm{sc}}(u)^2 dud\bar{u} \\ + \sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \frac{2i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u (m_{\mathrm{sc}}(z)^2 m_{\mathrm{sc}}(u)\mathbb{E}[e_{\mathfrak{a}}(G_{11}(u) + G_{jj}(u) - 2m_{\mathrm{sc}}(u))]) dud\bar{u} \\ \sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \frac{2i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u (m_{\mathrm{sc}}(u)^2 m_{\mathrm{sc}}(z)\mathbb{E}[e_{\mathfrak{a}}(G_{11}(z) + G_{jj}(z) - 2m_{\mathrm{sc}}(z))]) dud\bar{u} \\ + \mathcal{O}_{<}(|\lambda|\Psi(z)^2 N^{-1} + |\lambda|(1+|\lambda|)\Phi^2 N^{-1})$$
(5.15)

with the second estimate following from linearizing the  $G_{ii}$  and  $G_{jj}$  around  $m_{\rm sc}$ , similar to (5.10). The terms with  $G_{jj}$  on the third last and second last lines can now be estimated by  $\mathcal{O}_{<}(|\lambda|\Phi^2N^{-1})$  and  $\mathcal{O}_{<}(N^{-1}|\lambda|\eta^{-1})$ , respectively, using fluctuation averaging (2.7) and (2.26). This yields the claim.
## 5.2 Third order terms

We split up the estimation of the third order terms on the RHS of (5.6) into the diagonal and off-diagonal terms.

## 5.2.1 Diagonal terms

Starting with the diagonal term we have,

Lemma 5.5. We have,

$$\frac{s_{11}^{(3)}}{N^{3/2}} \mathbb{E}[\partial_{11}^{2}(e_{\mathfrak{a}}G_{11}(z))] = \frac{s_{11}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}] \left\{ m_{\mathrm{sc}}(z)^{3} + 2m_{\mathrm{sc}}(z)^{2} \left( \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}} \tilde{f}(u) \partial_{u} m_{\mathrm{sc}}(u) \right) + m_{\mathrm{sc}}(z) \left( -\frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}} \tilde{f}(u) \partial_{u} m_{\mathrm{sc}}(u) \right)^{2} + m_{\mathrm{sc}}(z) \left( \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}} \tilde{f}(u) \partial_{u} m_{\mathrm{sc}}(u)^{2} \right) \right\} \\
+ \mathcal{O}_{<}(N^{-3/2}\Psi(z)(1+|\lambda|)^{2} + |\lambda|(1+|\lambda|)\Phi N^{-3/2}) \tag{5.16} \quad \{ \text{eqn:prelim-there} \}$$

**Proof.** Using that  $\partial_{11}^2 G_{11} = m_{\rm sc}(z)^3 + \mathcal{O}(\Psi)$  we see that,

$$\frac{s_{11}^{(3)}}{N^{3/2}}\mathbb{E}[e_{\mathfrak{a}}\partial_{11}^{2}G_{11}(z)] = m_{\mathrm{sc}}(z)^{3}\frac{s_{11}^{(3)}}{N^{3/2}}\mathbb{E}[e_{\mathfrak{a}}] + \mathcal{O}_{\prec}(N^{-3/2}\Psi(z)).$$
(5.17)

Using the fact that  $\partial_{11}^k G_{11} = (-1)^k (G_{11})^{k+1} = (-1)^k m_{\rm sc}(z)^{k+1} + \mathcal{O}(\Psi)$  and that,

$$\partial_{11}e_{\mathfrak{a}} = -\frac{\mathrm{i}\lambda}{\pi}e_{\mathfrak{a}}\int_{\Omega_{\mathfrak{a}}}\mathrm{d}u\mathrm{d}\bar{u}\partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u)\mathrm{d}u\mathrm{d}\bar{u} + \mathcal{O}(|\lambda|\Phi)$$
  

$$\partial_{11}^{2}e_{\mathfrak{a}} = e_{\mathfrak{a}}\left(-\frac{\mathrm{i}\lambda}{\pi}\int_{\Omega_{\mathfrak{a}}}\mathrm{d}u\mathrm{d}\bar{u}\partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u)\mathrm{d}u\mathrm{d}\bar{u}\right)^{2}$$
  

$$+ e_{\mathfrak{a}}\left(\frac{\mathrm{i}\lambda}{\pi}\int_{\Omega_{\mathfrak{a}}}\mathrm{d}u\mathrm{d}\bar{u}\partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u)^{2}\mathrm{d}u\mathrm{d}\bar{u}\right) + \mathcal{O}(\Phi|\lambda|(1+|\lambda|))$$
(5.18)

we conclude the proof. We note that the above estimates follow from computing the derivatives explicitly (similar to the first equality in (2.35)), and then replacing all the appearances of  $G_{ii}(u)$  by  $m_{sc}(u)$  using (2.10) and (2.26).

## 5.2.2 Off-diagonal terms

The next few lemmas deal with the off-diagonal third order terms on the RHS of (5.6).

Lemma 5.6. We have,

$$\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}\partial_{j1}^{2}G_{j1}(z)] = 6m_{\mathrm{sc}}(z) \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}(G_{jj} - m_{\mathrm{sc}}(z))G_{1j}(z)] + \mathcal{O}_{\prec}(N^{-1/2}\Psi(z)^{3} + |\lambda\Phi|N^{-1}\Psi(z))$$
(5.19) (5.19) (5.19)

**Proof.** Since

$$\partial_{j1}^2 G_{j1} = 6G_{11}G_{jj}G_{1j} + \mathcal{O}_{\prec}(\Psi(z)^3)$$
(5.20) {eqn:prelim-th}

{lem:prelim-th

 $\{\texttt{sec:prelim-th}$ 

{lem:prelim-th

we have first that,

$$\sum_{j \neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}} \partial_{j1}^2 G_{j1}] = 6 \sum_{j \neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}} G_{11} G_{jj} G_{j1}] + \mathcal{O}_{<}(N^{-1/2} \Psi(z)^3).$$
(5.21)

Now we write,

$$\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} G_{11} G_{jj} G_{1j} = m_{\rm sc}(z)^2 \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} G_{1j} + m_{\rm sc}(z) \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} (G_{11} - m_{\rm sc}(z)) G_{1j} + m_{\rm sc}(z) \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} (G_{jj} - m_{\rm sc}(z)) G_{1j} + \mathcal{O}_{<}(N^{-1/2}\Psi(z)^3) = m_{\rm sc}(z)^2 \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} G_{1j} + m_{\rm sc}(z) \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} (G_{jj} - m_{\rm sc}(z)) G_{1j} + \mathcal{O}_{<}(N^{-1/2}\Psi(z)^3)$$
(5.22)

where we used the isotropic local law (2.9) in the second estimate. Now by Proposition 4.17 we have,

$$\sum_{j \neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}} G_{1j}(z)] = \mathcal{O}_{<}(N^{-1} \Psi(z)^2 + |\lambda \Phi| N^{-1} \Psi(z))$$
(5.23)

which completes the proof.

Lemma 5.7. We have,

$$\sum_{j \neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}e_{\mathfrak{a}})(\partial_{1j}G_{1j}(z))] = -\sum_{j \neq 1} \frac{m_{\mathrm{sc}}(z)s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}e_{\mathfrak{a}})(G_{jj}(z) - m_{\mathrm{sc}}(z))] \\ + \mathcal{O}_{\prec}(N^{-1/2}|\lambda\Phi|\Psi(z)^{2} + N^{-1}|\lambda|(1+|\lambda|)\Phi^{2})$$
(5.24) {eqn:prelim-th}

**Proof.** Since  $\partial_{1j}G_{1j} = -G_{jj}G_{11} + \mathcal{O}_{<}(\Psi(z)^2)$  and  $\partial_{1j}e_{\mathfrak{a}} = \mathcal{O}_{<}(|\lambda\Phi|)$  by (2.31) we have,

$$\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}e_{\mathfrak{a}})(\partial_{1j}G_{1j})] = -\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}e_{\mathfrak{a}})G_{11}G_{jj}] + \mathcal{O}_{<}(N^{-1/2}|\lambda\Phi|\Psi(z)^{2})$$

$$= -m_{\rm sc}(z)^{2} \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}e_{\mathfrak{a}})] - \sum_{j\neq 1} \frac{m_{\rm sc}(z)s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}e_{\mathfrak{a}})(G_{11} + G_{jj} - 2m_{\rm sc}(z))]$$

$$+ \mathcal{O}_{<}(N^{-1/2}|\lambda\Phi|\Psi(z)^{2}). \tag{5.25}$$

In the second estimate we linearized the  $G_{ii}$  and  $G_{jj}$  around  $m_{\rm sc}$ . We have,

$$\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}e_{\mathfrak{a}})(G_{11} - m_{sc}(z))]$$
  
=  $-\frac{2i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u} \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}G_{1j}(u)(G_{11}(z) - m_{sc}(z))] du d\bar{u} = \mathcal{O}_{\prec}(|\lambda\Phi|N^{-1}\Psi(z))$   
(5.26)

{lem:prelim-th

by the isotropic local law (2.9) and (2.26). Secondly, we have for  $j \neq 1$ ,

$$\mathbb{E}[\partial_{1j}e_{\mathfrak{a}}] = \mathcal{O}_{<}(N^{-1/2}|\lambda|(1+|\lambda|)\Phi^2)$$
(5.27)

by applying Proposition 4.17 and (2.26) to the formula in the first equality of (2.31). This completes the proof.

Lemma 5.8. We have,

$$\begin{split} &\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[G_{1j}(z)\partial_{1j}^{2}e_{\mathfrak{a}}] \\ &= \frac{2i\lambda}{\pi} \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \int_{\Omega_{\mathfrak{a}}} \mathrm{d} u \mathrm{d} \bar{u} (\partial_{\bar{u}} \tilde{f}(u)) \partial_{u} (m_{\mathrm{sc}}(u) \mathbb{E}[e_{\mathfrak{a}} G_{1j}(z) (G_{jj}(u) - m_{\mathrm{sc}}(u))]) \\ &+ \mathcal{O}_{<} (N^{-1/2} |\lambda \Phi^{2}| (1 + |\lambda|) \Psi(z) + |\lambda| N^{-1} \Psi(z)^{2}) \end{split}$$
(5.28) {eqn:prelim-th

**Proof.** We have by (5.14),

$$\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[G_{1j}(z)\partial_{1j}^{2}e_{\mathfrak{a}}] = \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}G_{1j}(z)] \frac{2\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u} m_{\mathrm{sc}}(u)^{2} \\ + \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \frac{2\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u} (m_{\mathrm{sc}}(u)\mathbb{E}[e_{\mathfrak{a}}G_{1j}(z)(G_{11}(u) + G_{jj}(u) - 2m_{\mathrm{sc}}(u))]) \\ + \mathcal{O}_{<}(N^{-1/2}\Phi^{2}|\lambda|(1+|\lambda|)\Psi(z))$$
(5.29)

The term with  $G_{11}(u) - m_{\rm sc}(u)$  on the second line contributes  $\mathcal{O}_{<}(N^{-1}|\lambda|\Phi\Psi(z))$  using the isotropic law (2.9). We use Proposition 4.17 to bound the term on the first line by  $\mathcal{O}_{<}(|\lambda|N^{-1}|\lambda\Phi|\Psi(z) + |\lambda|N^{-1}\Psi(z)^2)$ . The claim follows.

# 5.3 Intermediate expansion

Proposition 5.9. We have that,

$$\begin{split} z \sum_{a} \mathbb{E}[e_{\mathfrak{a}}G_{aa}(z)] + N\mathbb{E}[e_{\mathfrak{a}}] &= \sum_{ja} \frac{s_{1j}}{N} \mathbb{E}[\partial_{aj}(e_{\mathfrak{a}}G_{ja})] \\ &- \sum_{j \neq a} \frac{s_{aj}^{(4)}}{N^{2}} m_{\mathrm{sc}}(z)^{4} \mathbb{E}[e_{\mathfrak{a}}] - \mathbb{E}[e_{\mathfrak{a}}] m_{\mathrm{sc}}(z)^{2} \sum_{j \neq a} \frac{s_{aj}^{(4)}}{N^{2}} \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u}m_{\mathrm{sc}}(u)^{2} \mathrm{d}u \mathrm{d}\bar{u} \\ &+ \sum_{a} \frac{s_{aa}^{(3)}}{2N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}] \Big\{ m_{\mathrm{sc}}(z)^{3} + 2m_{\mathrm{sc}}(z)^{2} \left( \frac{\mathrm{i}\lambda}{\pi} e_{\mathfrak{a}} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right) \\ &+ m_{\mathrm{sc}}(z) \left( -\frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right)^{2} + m_{\mathrm{sc}}(z) \left( \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u)^{2} \right) \Big\} \\ &+ \mathcal{O}_{<}(N^{-1}\eta^{-1}(1+|\lambda|) + (1+|\lambda|)^{3}N^{-1} + N^{1/2}\Psi(z)^{3} + N^{1/2}|\lambda\Phi|\Psi(z)^{2}) \\ &+ \mathcal{O}_{<}(N^{1/2}|\lambda\Phi^{2}|(1+|\lambda|)\Psi(z)) \end{split}$$
(5.30) [eqn:intermedi

{peopinteenmedd

{lem:prelim-th

**Proof.** The proof starts by applying Lemma 5.2 to each term on the LHS of (5.30) but with the index 1 in (5.6) replaced by a. Then, one uses Lemmas 5.3, 5.4, 5.5, 5.6, 5.7 and 5.8 to deal with the terms that arise on the RHS of (5.6) (but again with the index 1 replaced by a). The only issue is that there are some extra terms in (5.8), (5.12), (5.19), (5.24) and (5.28) that we must estimate (they are simpler to estimate after taking the sum over a) in order to arrive at (5.30). Therefore, the remainder of the proof is devoted to estimating these terms.

The first extra term is the second term on the second line of (5.8). We estimate it by,

$$\sum_{j} \sum_{a \neq j} \frac{s_{aj}^{(4)}}{N^2} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - m_{\rm sc}(z))] = \mathcal{O}_{\prec}(N^{-1}\eta^{-1})$$
(5.31)

using fluctuation averaging (2.7). The next extra terms are those on the third and fourth lines of (5.12). By applying (2.7), one sees that the term on the third line is  $\mathcal{O}_{<}(|\lambda|\Phi^2)$  and the term on the fourth line is  $\mathcal{O}_{<}(|\lambda|N^{-1}\eta^{-1})$  (when we take the sum over *a*).

The next extra term is the term on the RHS of the first line of (5.19). We estimate it by,

$$\sum_{j} \sum_{a \neq j} \frac{s_{aj}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}(G_{jj}(z) - m_{\mathrm{sc}}(z))G_{aj}(z)]$$
  
= 
$$\sum_{j} \mathbb{E}\left[e_{\mathfrak{a}}(G_{jj} - m_{\mathrm{sc}}(z))\sum_{j \neq a} \frac{s_{aj}^{(3)}}{N^{3/2}}G_{aj}(z)\right] = \mathcal{O}_{<}(\Psi(z)^{2})$$
(5.32)

using the isotropic local law (2.9) for the sum over the index *a*. By almost the same argument one finds an estimate of  $\mathcal{O}_{<}(|\lambda\Phi|\Psi(z))$  for the terms on the RHS of the first lines of (5.24) and (5.28). This completes the proof.

As a simple corollary we find,

#### Corollary 5.10. We have,

$$\begin{split} z \sum_{a} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])] &= \sum_{ja} \frac{s_{aj}}{N} \left( \mathbb{E}[\partial_{aj}(e_{\mathfrak{a}}G_{ja}(z))] - \mathbb{E}[e_{\mathfrak{a}}]\mathbb{E}[\partial_{aj}G_{ja}(z)] \right) \\ &- \mathbb{E}[e_{\mathfrak{a}}]m_{\mathrm{sc}}(z)^{2} \sum_{j \neq a} \frac{s_{aj}^{(4)}}{N^{2}} \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u}m_{\mathrm{sc}}(u)^{2} \mathrm{d}u \mathrm{d}\bar{u} \\ &+ \sum_{a} \frac{s_{aa}^{(3)}}{2N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}] \left\{ 2m_{\mathrm{sc}}(z)^{2} \left( \frac{\mathrm{i}\lambda}{\pi} e_{\mathfrak{a}} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u}\partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right) \right. \\ &+ m_{\mathrm{sc}}(z) \left( -\frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u}\partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right)^{2} + m_{\mathrm{sc}}(z) \left( \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u}\partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right)^{2} \\ &+ \mathcal{O}_{<}(N^{-1}\eta^{-1}(1+|\lambda|) + (1+|\lambda|)^{3}N^{-1} + N^{1/2}\Psi(z)^{3} + N^{1/2}|\lambda\Phi|\Psi(z)^{2}) \\ &+ \mathcal{O}_{<}(N^{1/2}|\lambda\Phi^{2}|(1+|\lambda|)\Psi(z)) \end{split}$$
(5.33) {eqn:intermedian

as well as,

$$z \sum_{a} \mathbb{E}[G_{aa}(z)] + N = \sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[\partial_{aj}G_{ja}(z)] - \sum_{j \neq a} \frac{s_{aj}^{(4)}}{N^2} m_{\rm sc}(z)^4 + \sum_{a} \frac{s_{aa}^{(3)}}{2N^{3/2}} m_{\rm sc}(z)^3 + \mathcal{O}_{\prec}(N^{-1}\eta^{-1} + N^{1/2}\Psi(z)^3).$$
(5.34) {eqn:intermediation of the second second

**Proof.** The first estimate follows from applying (5.30) twice (once with  $\lambda = 0$ ) and subtracting the two results. The second just follows from taking  $\lambda = 0$  in (5.30).

## **5.4** Better estimate for $G_{11}$

In this section we derive an estimate for  $\mathbb{E}[e_{\mathfrak{a}}(G_{11}(z) - m_{sc}(z))]$  one order better than the naive estimate  $\Psi(z)$ . We start with the following.

**Proposition 5.11.** We have that,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{11}(z)] + \mathbb{E}[e_{\mathfrak{a}}] = \sum_{j} \frac{s_{1j}}{N} \mathbb{E}[\partial_{j1}(e_{\mathfrak{a}}G_{j1}(z))] + \mathcal{O}_{\prec}(N^{-1}(1+|\lambda|) + N^{-1/2}\Psi(z)^{2} + N^{-1/2}|\lambda\Phi|\Psi(z))$$
(5.35) {eqn:gll-expansion}

**Proof.** This follows from Lemma 5.2 and then using Lemmas 5.3, 5.5, 5.6, 5.7 and 5.8 to estimate the terms on the RHS of (5.6). Specifically, the errors in the  $\mathcal{O}_{<}$  on the last line of (5.6) are estimated by  $\mathcal{O}_{<}((1 + |\lambda|)N^{-1})$ . Then, all terms on the RHS of (5.8) and (5.16) can be estimated by  $\mathcal{O}_{<}((1 + |\lambda|)N^{-1})$ . All terms on the RHS of (5.19) can be estimated by  $\mathcal{O}_{<}(N^{-1/2}\Psi(z)^2)$ . All terms on the RHS of (5.24) and (5.28) can be estimated by  $\mathcal{O}_{<}(|\lambda\Phi|N^{-1/2}\Psi(z))$ .

We now compute the second order terms on the RHS of (5.35) in the following lemma.

Lemma 5.12. We have,

$$\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[G_{j1}\partial_{j1}e_{\mathfrak{a}}] = |\lambda|\mathcal{O}_{<}(\Phi\Psi(z)(\Psi(z) + \Phi(1+|\lambda|)) + \min\{\Psi_{1}(z)^{2}, \Phi^{2}\})$$
(5.36) {eqn:gl1-expansion}

and

$$\sum_{j} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} \partial_{j1} G_{j1}(z)] = -m_{\rm sc}(z) \mathbb{E}[e_{\mathfrak{a}} G_{11}(z)] + \mathcal{O}_{\prec}(N^{-1} \eta^{-1})$$
(5.37) {eqn:gll-expansion}

**Proof.** We start with (5.36). The j = 1 term contributes  $\mathcal{O}_{\prec}(|\lambda|N^{-1})$  using (2.36). The other terms we may write as,

$$\mathbb{E}[G_{1j}(z)\partial_{1j}e_{\mathfrak{a}}] = -\frac{2i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u}\mathbb{E}[e_{\mathfrak{a}}G_{1j}(z)G_{1j}(u)]\mathrm{d}u\mathrm{d}\bar{u}.$$
(5.38)

We now claim that the estimate (3.29), applied in the case that  $v_j = \mathbb{E}[e_{\mathfrak{a}}G_{1j}(z)G_{j1}(u)]$ , implies,

$$|\mathbb{E}[e_{\mathfrak{a}}G_{1j}(z)G_{1j}(u)]| < \Psi(z)\Psi(u)(\Psi(z) + \Psi(u) + |\lambda\Phi|) + \frac{1}{N(|\mathrm{Im}[z]| + |\mathrm{Im}[u]|)}.$$
 (5.39) {eqn:gll-expansion}

Indeed, in this case the terms involving  $X_k$  on the RHS of (3.29) are  $\mathcal{O}_{\langle}(\Psi(z)\Psi(u)(\Psi(u) + |\lambda\Phi|))$  (i.e., they involve hypergraphs with three edges, with the additional edge either having spectral parameter u or a new integration variable), the first error term on the RHS of (3.29) is  $\mathcal{O}_{\langle}(\Psi(z)^2\Psi(u))$ , and the error term on the second line of (3.29) is  $\mathcal{O}_{\langle}(N^{-1}(|\mathrm{Im}[z]| + |\mathrm{Im}[u]|)^{-1})$ . Furthermore, the terms involving  $Y_k$  and  $Z_k$  are  $\mathcal{O}_{\langle}(N^{-1}(|\mathrm{Im}[z]| + |\mathrm{Im}[u]|)^{-1})$  by their definition and (2.18).

{sec:g11-bette

{prop:g11-expa

{lem:g11-secon

Now, the the estimate (5.36) follows from (5.39) by applying (2.26). We now turn to (5.37). Since  $\partial_{1j}G_{1j} = -G_{11}G_{jj} + \mathcal{O}_{<}(\Psi(z)^2)$  we have,

$$\sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} \partial_{1j} G_{j1}(z)] = -m_{\rm sc}(z) \mathbb{E}[e_{\mathfrak{a}} G_{11}(z)] -\sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{11}(z) (G_{jj}(z) - m_{\rm sc}(z))] + \mathcal{O}_{<}(\Psi(z)^{2}).$$
(5.40)

The first term on the second line is  $\mathcal{O}_{<}(N^{-1}\eta^{-1})$  by fluctuation averaging (2.7), which completes the proof.

From (5.35), (5.36) and (5.37) we immediately conclude the following.

Proposition 5.13. We have,

$$\mathbb{E}[e_{\mathfrak{a}}(G_{11}(z) - m_{\mathrm{sc}}(z))] = \mathcal{O}_{\prec}(N^{-1}\eta^{-1} + |\lambda|[\Phi^2(1+|\lambda|)\Psi(z) + \min\{(N\eta)^{-1}, \Phi^2\}]) \quad (5.41)$$

Finally we have the following.

**Corollary 5.14.** Let  $z, w, v \in \Omega_{\mathfrak{a}}$ . We have,

$$\begin{aligned} |\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)(G_{33}(v) - m_{\mathrm{sc}}(v))]| &< \Psi_{1}(z)\Psi_{1}(w)\Psi_{1}(v)\min\{\Psi_{1}(z),\Psi_{1}(w)\}\\ \frac{1}{N(|\mathrm{Im}[z]| + |\mathrm{Im}[w]|)} \left(N^{-1}|\mathrm{Im}[v]|^{-1} + |\lambda|[\Phi^{2}(1 + |\lambda|)\Psi(v) + \min\{(N|\mathrm{Im}[v]|)^{-1},\Phi^{2}\}]\right) \end{aligned}$$

$$(5.42)$$

**Proof.** This follows immediately from using Proposition 5.13 to bound the first term on the second line of (3.57) of Lemma 3.25.

# **6** Estimates for $G_{12}G_{21}$

In this section we find a better an estimate for  $\mathbb{E}[e_{\mathfrak{a}}(\lambda)G_{12}(z)G_{21}(w)]$ . Throughout this section we assume that f is admissible and define  $z = E \pm i\eta$  and  $w = x \pm iy$ , for  $\eta, y > 0$ . We assume that  $z, w \in \Omega_{\mathfrak{a}}$  and that  $|\lambda \Phi| \leq N^{-c}$  for some c > 0. We begin with a cumulant expansion.

Lemma 6.1. We have,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)] = \sum_{j} \frac{s_{1j}}{N}\mathbb{E}[\partial_{j1}(G_{j2}(z)G_{21}(w)e_{\mathfrak{a}})] + \mathcal{O}_{\prec}(N^{-1/2}\Psi(z)\Psi(w)(\Psi(z) + \Psi(w)) + |\lambda\Phi|N^{-1/2}\Psi(z)\Psi(w))$$
(6.1) {eqn:g12g21-ex

**Proof.** From (3.7) and (3.9) we have,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)] = \sum_{j} \frac{s_{1j}}{N}\mathbb{E}[\partial_{1j}(G_{j2}(z)G_{21}(w)e_{\mathfrak{a}})] + \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{2N^{3/2}}\mathbb{E}[\partial_{1j}^{2}(e_{\mathfrak{a}}G_{j2}(z)G_{21}(w))] + \mathcal{O}_{<}(N^{-1}(1+|\lambda|)\Psi(z)\Psi(w))$$
(6.2) {eqn:g12g21-th

{cor:weak-g12g

{prop:g11-est}

 $\{\texttt{sec:g12g21}\}$ 

{lem:g12g21-ex

It therefore remains to estimate the sum on the second line of (6.2). For the j = 2 term in the sum we have,

$$|N^{-3/2}\mathbb{E}[\partial_{12}^2(e_{\mathfrak{a}}G_{22}(z)G_{12}(w))]| < (1+|\lambda|)N^{-3/2}\Psi(w) + N^{-3/2}|\lambda\Phi| + N^{-3/2}(\Psi(z) + \Psi(w))$$

$$(6.3) \quad \{\texttt{eqn:g12g21-th}\}$$

by direct computation, (2.31), (2.35) and (2.10).

By direct calculation and (2.10) we have for  $j \neq 1, 2$ ,

$$\begin{aligned} \partial_{1j}^2 G_{j2}(z) &= 2m_{\rm sc}(z)^2 G_{j2}(z) + \mathcal{O}_{\prec}(\Psi(z)^2), \quad \partial_{1j}^2 G_{21}(w) = 2m_{\rm sc}(w)^2 G_{12}(w) + \mathcal{O}_{\prec}(\Psi(w)^2) \\ \partial_{1j}G_{j2}(z) &= -G_{12}(z)m_{\rm sc}(z) + \mathcal{O}_{\prec}(\Psi(z)^2), \quad \partial_{1j}G_{21}(w) = -m_{\rm sc}(w)G_{j2}(w) + \mathcal{O}_{\prec}(\Psi(w)^2). \end{aligned}$$

$$(6.4) \quad \{\text{eqn:g12g21-th}\}$$

From the above we find,

$$\sum_{j \neq 1,2} \frac{s_{1j}^{(3)}}{N^{3/2}} \partial_{1j}^2 (G_{j2}(z)G_{21}(w)) = \mathcal{O}_{\prec}(N^{-1/2}\Psi(z)\Psi(w)(\Psi(z) + \Psi(w))$$
  
+ 
$$\sum_{j \neq 1,2} \frac{2s_{1j}^{(3)}}{N^{3/2}} ((m_{\rm sc}(z)^2 + m_{\rm sc}(w)^2)G_{j2}(z)G_{12}(w) + m_{\rm sc}(z)m_{\rm sc}(w)G_{12}(z)G_{j2}(w))$$
  
=  $\mathcal{O}_{\prec}(N^{-1/2}\Psi(z)\Psi(w)(\Psi(z) + \Psi(w))$  (6.5)

with the last line following from the isotropic local law (2.9).

Using the estimates on the second line of (6.4) and (2.31) we find,

$$\sum_{j \neq 1,2} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j} e_{\mathfrak{a}})(\partial_{1j}(G_{j2}(z)G_{21}(w)))] = \mathcal{O}_{\prec}(|\lambda \Phi| N^{-1/2} \Psi(z)\Psi(w)).$$
(6.6)

Using (2.32) we find,

$$\sum_{j \neq 1,2} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}^2 e_{\mathfrak{a}})(G_{j2}(z)G_{21}(w))] = \mathcal{O}_{\prec}(N^{-1/2}|\lambda\Phi|\Psi(z)\Psi(w))$$
(6.7)

using the isotropic local law (2.9). This completes the estimation of the sum on the second line of (6.2) and the proof.

We now estimate terms appearing on the RHS of the expansion (6.1).

**Proposition 6.2.** We have the estimates,

$$\frac{1}{N} |\mathbb{E}[\partial_{11}(e_{\mathfrak{a}}G_{12}(z)G_{21}(w))]| < (1+|\lambda|)N^{-1}\Psi(z)\Psi(w), \tag{6.8} \quad \{\texttt{eqn:g12g21-set}\}$$

and

$$\frac{s_{12}}{N} \mathbb{E}[\partial_{21}(e_{\mathfrak{a}}G_{22}(z)G_{21}(w))] = -\frac{s_{12}m_{\mathrm{sc}}(z)m_{\mathrm{sc}}(w)^{2}}{N} + \mathcal{O}_{<}(N^{-2}(\eta^{-1}+y^{-1})) + \mathcal{O}_{<}(|\lambda\Phi|N^{-1}\Psi(w)) + |\lambda|N^{-1}\mathcal{O}_{<}(\Phi^{2}(1+|\lambda|)\Psi(z) + \min\{N^{-1}y^{-1}+N^{-1}\eta^{-1},\Phi^{2}\})$$
(6.9) {eqn:g12g21-set}

{prop:g12g21-s

and for  $j \neq 1, 2$ ,

$$|\mathbb{E}[G_{j2}(z)G_{21}(w)\partial_{1j}e_{\mathfrak{a}}]| < |\lambda|\Psi_{1}(z)\Psi_{1}(w)\Phi\min\{\Psi_{1}(z),\Psi_{1}(w),\Phi\}.$$
(6.10) (eqn:g12g21-se

Finally,

$$\sum_{j \neq 1,2} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} \partial_{j1}(G_{j2}(z)G_{21}(w))] = -m_{\mathrm{sc}}(z)\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)]$$
  
$$-\sum_{j \neq 2} \frac{s_{1j}}{N}m_{\mathrm{sc}}(w)\mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{2j}(w)] + \mathcal{O}_{<}((\eta + y)^{-1}N^{-2}(y^{-1} + \eta^{-1}))$$
  
$$+\frac{|\lambda|}{N(\eta + y)}\mathcal{O}_{<}(\Phi^{2}(1 + |\lambda|)(\Psi(z) + \Psi(w)) + \min\{N^{-1}(y^{-1} + \eta^{-1}), \Phi^{2}\})$$
(6.11) {eqn:g12g21-set

**Proof.** The estimate (6.8) follows easily from (2.36) and (2.10). For the estimate (6.9), the term where the derivative hits  $e_{\mathfrak{a}}$  contributes  $\mathcal{O}_{<}(|\lambda\Phi|N^{-1}\Psi(w))$  by (2.31). The term where the derivative hits  $G_{22}(z)$  contributes  $\mathcal{O}_{<}(N^{-1}\Psi(z)\Psi(w))$ . Finally,

$$\frac{s_{12}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{22}(z) \partial_{21}(G_{21}(w))] = -\frac{s_{12}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{22}(z) G_{11}(w) G_{22}(w)] + \mathcal{O}_{<}(N^{-1} \Psi(w)^{2})$$

$$= -\frac{s_{12} m_{\rm sc}(z) m_{\rm sc}(w)^{2}}{N} + \mathcal{O}_{<}(N^{-2}(\eta^{-1} + y^{-1}))$$

$$+ N^{-1} |\lambda| \mathcal{O}_{<}(\Phi^{2}(1 + |\lambda|)(\Psi(z) + \Psi(w)) + \min\{N^{-1}y^{-1} + N^{-1}\eta^{-1}, \Phi^{2}\})$$
(6.12)

The second estimate follows from Proposition 5.13 after linearizing each of the  $G_{ii}$  around  $m_{\rm sc}$ .

Turning now to (6.10) we have by the equality in (2.31),

$$\mathbb{E}[G_{j2}(z)G_{21}(w)\partial_{j1}e_{\mathfrak{a}}] = -\frac{2i\lambda}{\pi}\int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u}\mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{21}(w)G_{1j}(u)]\mathrm{d}u\mathrm{d}\bar{u}.$$
(6.13)

From Proposition 3.2 we have,

$$|\mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{21}(w)G_{1j}(u)]| \le \Psi_1(z)\Psi_1(w)\Psi_1(u)\min\{\Psi_1(z),\Psi_1(w),\Psi_1(u)\}$$
(6.14)

and so (6.10) follows, using (2.26).

Turning now to (6.11) we have by direct computation,

$$\sum_{j \neq 1,2} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}}\partial_{j1}(G_{j2}(z)G_{21}(w))] = -m_{sc}(z)\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)] + \mathcal{O}_{<}(N^{-1}\Psi(w)\Psi(z))$$

$$-\sum_{j \neq 2} \frac{s_{1j}}{N}m_{sc}(w)\mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{2j}(w)] - \sum_{j \neq 1,2} \frac{s_{1j}}{N}\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)(G_{jj}(z) - m_{sc}(z))]$$

$$-\sum_{j \neq 1,2} \frac{s_{1j}}{N}\mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{2j}(w)(G_{11}(w) - m_{sc}(w))]$$

$$-\sum_{j \neq 1,2} \frac{s_{1j}}{N}\mathbb{E}[e_{\mathfrak{a}}(G_{1j}(z)G_{j2}(z)G_{21}(w) + G_{j2}(z)G_{21}(w)G_{j1}(w))]$$
(6.15) {eqn:g12g21-se}

The claimed estimates now follow from Corollary 5.14 applied to the term on the third line and second term on the second line of (6.15), and Proposition 3.2 applied to the terms on the last line.

Lemma 6.1 and Proposition 6.2 easily imply the following and so the proof is omitted.

{cor:g12g21-se

Corollary 6.3. We have that,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)] = -m_{\mathrm{sc}}(z)\mathbb{E}[e_{\mathfrak{a}}G_{12}(z)G_{21}(w)] - \sum_{j\neq 2}\frac{s_{1j}}{N}\mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{j2}(w)] - \frac{s_{12}m_{\mathrm{sc}}(z)m_{\mathrm{sc}}(w)^{2}}{N} + \mathcal{O}_{<}((Ny)^{-1}(N\eta)^{-1}) + \frac{|\lambda|}{N(y+\eta)}\mathcal{O}_{<}(\Phi^{2}(1+|\lambda|)(\Psi(z)+\Psi(w)) + \min\{N^{-1}(y^{-1}+\eta^{-1}),\Phi^{2}\}) + \mathcal{O}_{<}(|\lambda|\Psi_{1}(z)\Psi_{1}(w)\Phi\min\{\Psi_{1}(z),\Psi_{1}(w),\Phi\})$$
(6.16)

We define now,

$$v_j := \mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{2j}(w)]\mathbf{1}_{\{j\neq 2\}}.$$
(6.17)

Recall the definition of A in Lemma 2.8 and let B be the matrix  $B_{ij} = \frac{S_{ij}}{N} \mathbf{1}_{\{i=2\}}$ . Corollary 6.3 implies that we have the equation,

$$(1 - m_{\rm sc}(z)m_{\rm sc}(w)(A - B))v = m_{\rm sc}(z)m_{\rm sc}(w)N^{-1}ee^*v + X + \varepsilon$$
(6.18) {eqn:g12g21-se

where  $X_j := s_{j2} \mathbf{1}_{\{j \neq 2\}} \frac{m_{\rm sc}(z)^2 m_{\rm sc}(w)^2}{N}$  and the vector  $\varepsilon$  obeys,

$$\|\varepsilon\|_{\infty} < (Ny)^{-1} (N\eta)^{-1} ) + N^{-1} (y+\eta)^{-1} (|\lambda| [(\Phi^2(1+|\lambda|)(\Psi(z)+\Psi(w)) + \min\{N^{-1}(y^{-1}+\eta^{-1}), \Phi^2\}) + (|\lambda|\Psi_1(z)\Psi_1(w)\Phi\min\{\Psi_1(z), \Psi_1(w), \Phi\})$$
(6.19)

We now compute the inner product on the RHS of (6.18).

Lemma 6.4. We have,

$$N^{-1}\mathbf{e}\mathbf{e}^{*}v = \mathbb{E}[e_{\mathfrak{a}}]\left(\frac{m_{\mathrm{sc}}(z) - m_{\mathrm{sc}}(w)}{N(z-w)} - \frac{m_{\mathrm{sc}}(z)m_{\mathrm{sc}}(w)}{N}\right) + N^{-1}(y+\eta)^{-1}\mathcal{O}_{\prec}(N^{-1}(y^{-1}+\eta^{-1})) + N^{-1}(y+\eta)^{-1}\mathcal{O}_{\prec}(|\lambda|[(\Phi^{2}(1+|\lambda|)(\Psi(z)+\Psi(w)) + \min\{N^{-1}(y^{-1}+\eta^{-1}),\Phi^{2}\}])$$
(6.20)

**Proof.** We write,

$$N^{-1}\mathbf{e}\mathbf{e}^*v = \frac{1}{N}\sum_j \mathbb{E}[e_{\mathfrak{a}}G_{j2}(z)G_{2j}(w)] - \frac{1}{N}\mathbb{E}[e_{\mathfrak{a}}G_{22}(z)G_{22}(w)].$$
(6.21)

For the second term we write,

$$\frac{1}{N} \mathbb{E}[e_{\mathfrak{a}} G_{22}(z) G_{22}(w)] = \frac{m_{\rm sc}(z) m_{\rm sc}(w)}{N} \mathbb{E}[e_{\mathfrak{a}}] + N^{-1} \mathcal{O}_{\prec} (N^{-1}(y^{-1} + \eta^{-1}) + |\lambda| [\Phi^2(1 + |\lambda|)(\Psi(z) + \Psi(w)) + \min\{N^{-1}(y^{-1} + \eta^{-1}), \Phi^2\}])$$
(6.22)

using Proposition 5.13. For the first term we write,

$$\frac{1}{N} \sum_{j} \mathbb{E}[e_{\mathfrak{a}} G_{j2}(z) G_{2j}(w)] = \frac{m_{\rm sc}(z) - m_{\rm sc}(w)}{N(z - w)} \mathbb{E}[e_{\mathfrak{a}}] + \frac{1}{N(z - w)} \mathbb{E}[e_{\mathfrak{a}}(G_{22}(z) - m_{\rm sc}(z) + m_{\rm sc}(w) - G_{22}(w))].$$
(6.23)

{lem:g12g21-di

We now claim that,

$$\left| \frac{1}{N(z-w)} \mathbb{E}[e_{\mathfrak{a}}(G_{22}(z) - m_{\mathrm{sc}}(z) + m_{\mathrm{sc}}(w) - G_{22}(w))] \right|$$

$$< \frac{1}{N(y+\eta)} \left( N^{-1}(y^{-1} + \eta^{-1}) + |\lambda| [\Phi^{2}(1+|\lambda|)(\Psi(z) + \Psi(w)) + \min\{N^{-1}(y^{-1} + \eta^{-1}), \Phi^{2}\}] \right)$$

$$(6.24) \quad \{ \operatorname{eqn:g12g21-di} \}$$

which will complete the proof. In order to prove (6.24), WLOG assume  $\eta > y$ . If  $|z-w| > \eta/2$  then we can estimate the prefactor  $|N(z-w)|^{-1} < (N\eta)^{-1}$  and the expectation in (6.24) directly by Proposition 5.13, yielding the claim. Otherwise, assume that  $|z-w| < \eta/2$ . Then  $y \simeq \eta$  and  $\Psi(z) \simeq \Psi(w)$ . We then write,

$$\frac{1}{N(z-w)} \mathbb{E}[e_{\mathfrak{a}}(G_{22}(z) - m_{\rm sc}(z) + m_{\rm sc}(w) - G_{22}(w))]$$
  
=  $N^{-1} \int_{0}^{1} \partial_{s} \mathbb{E}[e_{\mathfrak{a}}(G_{22}(w + s(z-w)) - m_{\rm sc}(w + s(z-w)))] \mathrm{d}s.$  (6.25)

We conclude (6.24) by estimating the integrand using Proposition 5.13 and the Cauchy Integral formula.

Now for every k let us define the vector

$$v_j^{(k)} := \mathbb{E}[e_{\mathfrak{a}}G_{jk}(z)G_{kj}(w)]\mathbf{1}_{\{j\neq k\}}.$$
(6.26)

Define  $M = m_{\rm sc}(z)m_{\rm sc}(w)$  as well as,

$$F := M \frac{m_{\rm sc}(z) - m_{\rm sc}(w)}{N(z - w)} - \frac{M^2}{N}.$$
(6.27)

Define the matrix  $B^{(k)}$  by  $B^{(k)}_{ab} = \frac{s_{ab}}{N} \mathbf{1}_{\{a=k\}}$  and the vector,

$$Y_j^{(k)} := \frac{s_{jk}}{N} M^2 \mathbf{1}_{\{j \neq k\}}$$
(6.28)

We have the following equation for the  $v_j^{(k)}$ .

Corollary 6.5. We have that,

$$\begin{aligned} v_{j}^{(k)} &= \mathbb{E}[e_{\mathfrak{a}}] \sum_{i} (1 - m_{\mathrm{sc}}(z)m_{\mathrm{sc}}(w)(A - B^{(k)}))_{ji}^{-1} \left\{ Y_{i}^{(k)} + F \right\} \\ &+ \mathcal{O}_{<}((Ny)^{-1}(N\eta)^{-1}) \\ &+ \frac{|\lambda|}{N(y+\eta)} \mathcal{O}_{<}(\Phi^{2}(1+|\lambda|)(\Psi(z) + \Psi(w)) + \min\{N^{-1}(y^{-1}+\eta^{-1}), \Phi^{2}\}) \\ &+ \mathcal{O}_{<}(|\lambda|\Psi_{1}(z)\Psi_{1}(w)\Phi\min\{\Psi_{1}(z), \Psi_{1}(w), \Phi\}) \end{aligned}$$
(6.29) {eqn:g12g21-set}

**Proof.** This follows by replacing the index 2 by k in (6.18) and Lemma 6.4, after noticing that Lemma B.1 applies to bound the matrix elements of  $(1 - M(A - B^{(k)}))^{-1}$ .

For future use, we now compute the deterministic term that arises when we apply (6.29) to compute  $\sum_{ij} \frac{s_{ij}}{N} v_j^{(i)}$ .

Lemma 6.6. We have that,

$$\frac{1}{N} \sum_{ij} s_{ij} [(1 - M(A - B^{(i)}))^{-1} (Y^{(i)} + \mathbf{e}F)]_j = -M \mathrm{tr}S + M \mathrm{tr}(S(1 - MS)^{-1}) + \mathcal{O}_{\prec} (N^{-1}(y + \eta)^{-1})$$
(6.30) {eqn:g12g21-de

**Proof.** We first show that we can remove the operator  $B^{(i)}$  up to a negligible error. We have,

$$\sum_{j} s_{ij} [(1 - M(A - B^{(i)})^{-1}(Y^{(i)} + \mathbf{e}F)]_j - \sum_{j} s_{ij} [(1 - MA)^{-1}(Y^{(i)} + \mathbf{e}F)]_j$$
  
=  $((1 - M(A - B^{(i)}))^{-1}s^{(i)}) \cdot (B^{(i)}(1 - MA)^{-1}(Y^{(i)} + \mathbf{e}F))$  (6.31)

where  $s^{(i)}$  is the vector with entries  $s_{ij}$ . Now, since  $(1 - M(A - B^{(i)}))^{-1}$  and  $(1 - MA)^{-1}$  are bounded on  $\ell^{\infty}$ , the entries of the vector  $(1 - M(A - B^{(i)}))^{-1}s^{(i)}$  are bounded by a constant, and the entries  $(1 - MA)^{-1}(Y^{(i)} + \mathbf{e}F)$  are  $\mathcal{O}(N^{-1}(\eta + y)^{-1})$  (see (2.18)), and so

$$\left| \left( (1 - M(A - B^{(i)}))^{-1} s^{(i)} \right) \cdot \left( B^{(i)} (1 - MA)^{-1} (Y^{(i)} + \mathbf{e}F) \right) \right| < \frac{1}{N(y + \eta)} \sum_{ab} |B_{ab}^{(i)}| < \frac{1}{N^2(y + \eta)} \sum_{ab} \mathbf{1}_{\{a=i\}} < \frac{1}{N(y + \eta)}.$$
(6.32)

Therefore, up to an error of  $\mathcal{O}_{<}(N^{-1}(y+\eta)^{-1})$  it suffices to compute,

$$\frac{1}{N} \sum_{ij} s_{ij} [(1 - MA)^{-1} (Y^{(i)} + \mathbf{e}F)]_j.$$
(6.33)

The difference between the vector  $Y^{(i)}$  and the vector  $M^2 s^{(i)} N^{-1}$  is the vector with a single non-zero entry  $\mathbf{1}_{\{j=i\}} M^2 s_{ii}/N$ . Therefore,

$$\left| \frac{1}{N} \sum_{ij} s_{ij} [(1 - MA)^{-1} (Y^{(i)} - M^2 s^{(i)} N^{-1})]_j \right| < \frac{1}{N^2} \sum_{ijk} |(1 - MA)^{-1}_{jk}| \mathbf{1}_{\{k=i\}} = \frac{1}{N^2} \sum_{j,k} |(1 - MA)^{-1}_{jk}| < \frac{1}{N}$$
(6.34)

with the last estimate following from the fact that  $(1 - MA)^{-1}$  is bounded from  $\ell^{\infty}$  to  $\ell^{\infty}$ . We have therefore reduced the proof to computing,

$$\frac{1}{N}\sum_{ij}s_{ij}(1-MA)_{jk}^{-1}s_{ik}M^2N^{-1} + \frac{1}{N}\sum_{ij}s_{ij}((1-MA)^{-1}\mathbf{e}F)_j$$
(6.35) {eqn:determini

Since  $(1 - MA)^{-1}\mathbf{e} = \mathbf{e}$  by the definition of A we have,

$$\frac{1}{N}\sum_{ij}s_{ij}((1-MA)^{-1}\mathbf{e}F)_j = \frac{F}{N}\sum_{ij}s_{ij} = M\frac{m_{\rm sc}(z) - m_{\rm sc}(w)}{z-w} - M^2 = M\frac{M}{1-M} - M^2 \quad (6.36) \quad \{\texttt{eqn:determining} \in M^2 : M^2 = M\frac{M}{1-M} - M^2 \quad (6.36) \quad \{\texttt{eqn:determining} \in M^2 : M^2 = M\frac{M}{1-M} - M^2 \quad (6.36) \quad \{\texttt{eqn:determining} \in M^2 : M^2 = M\frac{M}{1-M} - M^2 \quad (6.36) \quad \{\texttt{eqn:determining} \in M^2 : M^2 : M^2 = M\frac{M}{1-M} - M^2 \quad (6.36) \quad \{\texttt{eqn:determining} \in M^2 : M$$

with the last equality using the identity (2.19). For the other term in (6.35) we have,

$$\frac{1}{N}\sum_{ij}s_{ij}(1-MA)_{jk}^{-1}s_{ik}M^2N^{-1} = M^2\operatorname{tr}(S(1-MA)^{-1}S).$$
(6.37)

On the other hand via the Sherman Morrison formula,

$$\frac{1}{1 - MS} = \frac{1}{1 - MA} + MN^{-1} \frac{\mathbf{e}\mathbf{e}^*}{1 - m_{\rm sc}(z)m_{\rm sc}(w)}$$
(6.38)

 $\mathbf{SO}$ 

$$\operatorname{tr}(S(1 - MA)^{-1}M^{2}S) = \operatorname{tr}(S(1 - MS)^{-1}M^{2}S) - M^{3}\frac{1}{1 - M}$$
$$= -M\operatorname{tr}S + M\operatorname{tr}(S(1 - MS)^{-1}) - M^{3}/(1 - M)$$
(6.39)

Therefore, using the above and (6.36) we find that the expression in (6.35) equals,

$$\frac{1}{N} \sum_{ij} s_{ij} (1 - MA)_{jk}^{-1} s_{ik} M^2 N^{-1} + \frac{1}{N} \sum_{ij} s_{ij} ((1 - MA)^{-1} \mathbf{e}F)_j$$
  
=  $-M \mathrm{tr}S + M \mathrm{tr} (S(1 - MS)^{-1}) - M^3 / (1 - M) + M \frac{M}{1 - M} - M^2$   
=  $-M \mathrm{tr}S + M \mathrm{tr} (S(1 - MS)^{-1})$  (6.40)

which yields the claim.

**Corollary 6.7.** For  $|\lambda \Phi| \leq 1$  we have,

$$\frac{1}{N} \sum_{i \neq j} s_{ij} \mathbb{E}[e_{\mathfrak{a}} G_{ij}(z) G_{ji}(w)] = \mathbb{E}[e_{\mathfrak{a}}] \left(-M \operatorname{tr} S + M \operatorname{tr}(S(1-MS)^{-1})\right) 
+ \mathcal{O}_{<}((Ny)^{-1}(\eta)^{-1}) 
+ \frac{|\lambda|}{y+\eta} \mathcal{O}_{<}(\Phi^{2}(1+|\lambda|)(\Psi(z)+\Psi(w)) + \min\{N^{-1}(y^{-1}+\eta^{-1}),\Phi^{2}\}) 
+ N \mathcal{O}_{<}(|\lambda|\Psi_{1}(z)\Psi_{1}(w)\Phi\min\{\Psi_{1}(z),\Psi_{1}(w),\Phi\})$$
(6.41)

**Proof.** We have,

$$\frac{1}{N}\sum_{i\neq j}s_{ij}\mathbb{E}[e_{\mathfrak{a}}G_{ij}(z)G_{ji}(w)] = \frac{1}{N}\sum_{i\neq j}s_{ij}v_{j}^{(i)} = \frac{1}{N}\sum_{i,j}s_{ij}v_{j}^{(i)}.$$
(6.42)

We then apply (6.29) and (6.30).

# 7 Estimates for $G_{11}G_{22}$

Let  $b_a \in \mathbb{C}$  for  $1 \le a \le N$  satisfy  $|b_a| \le C$  some fixed C > 0 and define,

$$X_i := \sum_{a \neq i} \frac{b_a}{N} (G_{aa}(w) - m_{\rm sc}(w)) = \mathcal{O}_{\prec}((Ny)^{-1})$$
(7.1) {eqn:Xi-est}

where the estimate follows by fluctuation averaging (2.7), where we denote  $w = x \pm iy$ , with y > 0. In this section we are going to find a good estimate for the quantity  $\mathbb{E}[e_{\mathfrak{a}}G_{ii}(z)X_i]$ . We again denote  $z = E \pm i\eta$  with  $\eta > 0$ . We will assume throughout this section that  $z, w \in \Omega_{\mathfrak{a}}$ . The following performs a cumulant expansion for this quantity.

{cor:gij-main-

{sec:g11g22}

{lem:giigjj-fo

{lem:giigjj-th

**Lemma 7.1.** For any j and  $k \ge 0$ ,

$$\partial_{j1}^{k} X_{1} = \mathcal{O}_{\prec}(\Psi(w)^{2} + \mathbf{1}_{\{k=0\}}(Ny)^{-1})$$
(7.2) {eqn:giigjj-fo

For  $|\lambda \Phi| \leq 1$  we have,

$$N^{-2}\partial_{11}^{3}(G_{11}(z)e_{\mathfrak{a}}X_{1}) = \mathcal{O}_{<}((1+|\lambda|)^{3}N^{-2}(Ny)^{-1})$$
(7.3) {eqn:giigjj-fo

and for  $j \neq 1$ ,

$$N^{-2}|\partial_{j1}^{3}(e_{\mathfrak{a}}G_{j1}(z)X_{1})| < N^{-2}(1+|\lambda|)(Ny)^{-1}).$$
(7.4) {eqn:giigjj-fo

For any  $\delta > 0$ , the above estimates also hold with an additional  $N^{\delta}$  factor on the RHS for any  $\varepsilon$ -regular matrix, if  $\varepsilon > 0$  is sufficiently small depending on  $\delta > 0$ .

We have for  $|\lambda \Phi| \leq 1$ ,

$$z\mathbb{E}[e_{\mathfrak{a}}G_{11}(z)X_{1}] + \mathbb{E}[e_{\mathfrak{a}}X_{1}] = \sum_{j} \frac{s_{1j}}{N} \mathbb{E}[\partial_{j1}(G_{j1}(z)e_{\mathfrak{a}}X_{1})] + \sum_{j} \frac{s_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[\partial_{j1}^{2}(G_{j1}(z)e_{\mathfrak{a}}X_{1})] + \mathcal{O}_{\prec}(N^{-1}(1+|\lambda|)(Ny)^{-1})$$
(7.5) {eqn:giigjj-fo

**Proof.** The estimate (7.2) follows from direct computation and (2.10). For (7.3), if all derivatives hit  $e_{\mathfrak{a}}$ , we can estimate this by  $(1 + |\lambda|)^3 N^{-2} (Ny)^{-1}$  using (7.1) and (2.36). Otherwise using (7.2) or (7.1) we find an estimate of  $\mathcal{O}_{<}((1 + |\lambda|)^2 N^{-2} (Ny)^{-1})$ , completing the proof of (7.3). For (7.4) we find an estimate of  $\mathcal{O}_{<}((1 + |\lambda|)N^{-2}\Psi(z)(Ny)^{-1})$  if no derivatives hit  $G_{1j}$  using (2.36), (7.1) and (2.10). Otherwise, if a derivative hits  $G_{1j}$  we find an estimate of  $\mathcal{O}_{<}(N^{-2}(1 + |\lambda|)(Ny)^{-1})$ . The statement about  $\varepsilon$ -regular matrices is clear. The expansion (7.5) now follows from applying the cumulant expansion Lemma 2.6, similar to, e.g., Lemma 3.5.

The following lemma takes care of the third order terms in the cumulant expansion above.

**Lemma 7.2.** Assume  $|\lambda \Phi| \leq 1$ . We have,

$$N^{-3/2}|\mathbb{E}[\partial_{11}^2(e_{\mathfrak{a}}G_{11}(z)X_1)]| < N^{-3/2}(1+|\lambda|)^2(Ny)^{-1}$$
(7.6) {eqn:giigjj-th}

and

$$\sum_{j \neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[G_{1j}(z)X_1 \partial_{j1}^2 e_{\mathfrak{a}}] = \mathcal{O}_{<}(|\lambda \Phi| N^{-1/2} \Psi(z)(Ny)^{-1})$$
(7.7) {eqn:giigjj-th}

and

$$\sum_{j \neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{j1} e_{\mathfrak{a}}) \partial_{j1}(G_{j1} X_1)] = \mathcal{O}_{\prec}(\Psi(z)(Ny)^{-1} N^{-1/2} |\lambda| \Phi)$$
(7.8) {eqn:giigjj-th}

and

$$\sum_{j \neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}} \partial_{j1}^2 (G_{j1} X_1)] = \mathcal{O}_{\prec}(N^{-1/2} (Ny)^{-1} \Psi(z))$$
(7.9) {eqn:giigjj-th}

**Proof.** The estimate (7.6) is straightforward from (7.2) and (2.36). The estimate (7.7) follows from the fact that  $\partial_{1j}^2 e_{\mathfrak{a}} = i\lambda c_f + \mathcal{O}_{<}(|\lambda\Phi|)$  (see (2.32)), the isotropic local law (2.9) and (7.1). For (7.8), the term where  $\partial_{j1}$  hits  $X_1$  is easily bounded by  $\mathcal{O}_{<}(N^{-1/2}|\lambda\Phi|\Psi(z)(Ny)^{-1})$  using (7.2) and (2.31). For the other term we have since  $\partial_{j1}G_{j1}(z) = -m_{\rm sc}(z)^2 + \mathcal{O}_{<}(\Psi(z))$ ,

$$\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[(\partial_{1j}e_{\mathfrak{a}})(\partial_{j1}G_{j1}(z))X_{1}] = -m_{\mathrm{sc}}(z)^{2} \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[X_{1}(\partial_{j1}e_{\mathfrak{a}})] + \mathcal{O}_{<}(\Psi(z)(Ny)^{-1}N^{-1/2}|\lambda|\Phi) = \mathcal{O}_{<}(\Psi(z)(Ny)^{-1}N^{-1/2}|\lambda|\Phi)$$
(7.10)

with the second estimate following from the isotropic local law (2.9) applied to the equality in (2.31). This completes (7.8). For (7.9), since  $\partial_{1j}^2 X_1 = \mathcal{O}_{\prec}((Ny)^{-1})$  and  $\partial_{j1}^2 G_{1j} = \mathcal{O}_{\prec}(\Psi(z))$  and  $\partial_{j1}G_{1j} = -m_{\rm sc}(z)^2 + \mathcal{O}_{\prec}(\Psi(z))$  we have,

$$\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}\partial_{j1}^{2}(G_{j1}X_{1})] = -2m_{\mathrm{sc}}(z)^{2} \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}(\partial_{j1}X_{1})] + \mathcal{O}_{<}(N^{-1/2}(Ny)^{-1}\Psi(z)).$$

$$(7.11) \quad \{\mathsf{eqn:giigjj-th}\}$$

Then,

$$-\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \partial_{j1} X_1 = 2 \sum_{j\neq 1} \sum_{a\neq 1} \frac{b_a s_{1j}^{(3)}}{N^{5/2}} G_{aj}(w) G_{a1}(w) = 2 \sum_{a\neq 1} G_{a1} \frac{b_a}{N^2} \sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{1/2}} G_{aj}$$
$$= 2 \sum_{a\neq 1} G_{a1} \frac{b_a}{N^{5/2}} s_{1a}^{(3)} m_{\rm sc}(w) + \mathcal{O}_{\prec}(N^{-1}\Psi(w)^2) = \mathcal{O}_{\prec}(N^{-1}\Psi(w)^2).$$
(7.12) {eqn:gigj-th}

Here, the third equality follows from the isotropic law (2.9). This completes the proof of (7.9).

We now compute the second order terms on the RHS of (7.5).

**Lemma 7.3.** Assume that there is a c > 0 so that  $|\lambda \Phi| \leq N^{-c}$ . Then we have

$$N^{-1}|\mathbb{E}[\partial_{11}(e_{\mathfrak{a}}G_{11}(z)X_{1})]| < N^{-1}(1+|\lambda|)(Ny)^{-1}$$
(7.13) {eqn:giigjj-set}

{lem:giigjj-se

and

$$\sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[G_{j1}(z)X_1 \partial_{j1} e_{\mathfrak{a}}] = \mathcal{O}_{\prec}(\Psi(z)|\lambda \Phi|(Ny)^{-1})$$
(7.14) {eqn:giigjj-se

and

$$\sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{j1}(z) \partial_{j1} X_1] = \mathcal{O}_{\prec}((Ny)^{-1} (N\eta)^{-1})$$
(7.15) {eqn:giigjj-se

and

$$\sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} X_1 \partial_{j1} G_{j1}] = -m_{\mathrm{sc}}(z) \mathbb{E}[G_{11} X_1 e_{\mathfrak{a}}] + \mathcal{O}_{\prec}((Ny)^{-1} (N\eta)^{-1})$$
(7.16) {eqn:giigjj-setal setal of a setal o

**Proof.** The estimates (7.13) and (7.14) are straightforward. For (7.15) we write,

$$-\sum_{j\neq 1} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} G_{j1}(z)\partial_{j1} X_{1}] = 2\sum_{j\neq 1} \sum_{a\neq 1} \frac{s_{1j}b_{a}}{N^{2}} \mathbb{E}[e_{\mathfrak{a}} G_{j1}(z)G_{1a}(w)G_{aj}(w)]$$
$$= 2\sum_{j\neq 1} \sum_{a\neq 1,j} \frac{s_{1j}b_{a}}{N^{2}} \mathbb{E}[e_{\mathfrak{a}} G_{j1}(z)G_{1a}(w)G_{aj}(w)] + \mathcal{O}_{\prec}(N^{-1}\Psi(z)\Psi(w))$$
(7.17)

Now (7.15) follows since  $\mathbb{E}[e_{\mathfrak{a}}G_{j1}(z)G_{1a}(w)G_{aj}(w)] = \mathcal{O}_{<}((Ny)^{-1}(N\eta)^{-1})$  for indices 1, a, j all distinct, by Proposition 3.2.

Finally, for (7.16) we have,

$$\sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} X_1 \partial_{j1} G_{j1}(z)] = -m_{\rm sc}(z) \sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} X_1 G_{11}(z)] - \sum_{j \neq 1} \frac{s_{1j}}{N} \mathbb{E}[e_{\mathfrak{a}} X_1 G_{11}(G_{jj}(z) - m_{\rm sc}(z))] + \mathcal{O}_{\prec}(\Psi(z)^2 (Ny)^{-1})$$
(7.18)

Now the first term on the second line is  $\mathcal{O}_{<}((N\eta)^{-1}(Ny)^{-1})$  by fluctuation averaging (2.7).  $\Box$ 

**Proposition 7.4.** Assume that  $|\lambda \Phi| \leq N^{-c}$  for some c > 0. For any *i* we have,

$$\sum_{a} \frac{b_{a}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{ii}(z) - m_{\mathrm{sc}}(z))(G_{aa}(w) - m_{\mathrm{sc}}(w))] = \mathcal{O}_{<}((Ny)^{-1}(N\eta)^{-1} + |\lambda\Phi|\Psi(z)(Ny)^{-1})$$
(7.19)

**Proof.** WLOG we can assume i = 1. Then the term in the sum with a = 1 contributes  $\mathcal{O}_{<}(N^{-1}\Psi(z)\Psi(w))$  and so can be ignored. The equation (7.5) combined with Lemmas 7.2 and 7.3 imply,

$$(z + m_{\rm sc}(z))\mathbb{E}[(G_{11}(z) - m_{\rm sc}(z))X_1] = \mathcal{O}_{<}((Ny)^{-1}(N\eta)^{-1} + |\lambda\Phi|\Psi(z)(Ny)^{-1})$$
(7.20)

and so the claim follows since  $|z + m_{\rm sc}(z)| \approx 1$ .

# 7.1 Expansion for expectation

In this section we derive an expansion for the expectation,

$$\frac{1}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{z}} \tilde{f}(z)) \sum_{a} \mathbb{E}[G_{aa}(z)] \mathrm{d}z \mathrm{d}\bar{z}.$$
(7.21)

We will use the expansion (5.34), and begin by computing the second order terms on its RHS.

Lemma 7.5. We have

$$\sum_{ja} \frac{s_{ja}}{N} \mathbb{E}[\partial_{ja} G_{ja}(z)] = -Nm_{\rm sc}(z)^2 - 2m_{\rm sc}(z) \sum_a \mathbb{E}[G_{aa}(z) - m_{\rm sc}(z)] + m_{\rm sc}(z)^2 {\rm tr}S - m_{\rm sc}(z)^2 {\rm tr}(S(1 - m_{\rm sc}(z)^2 S)^{-1}) + \mathcal{O}_{\prec}(N^{-1}\eta^{-2}).$$
(7.22) {eqn:exp-second

and so

7

$$\sum_{a} \mathbb{E}[G_{aa}(z) - m_{\rm sc}(z)] = \sum_{j \neq a} \frac{s_{aj}^{(4)}}{N^2} m_{\rm sc}(z)^3 m_{\rm sc}'(z) - \sum_{a} \frac{s_{aa}^{(3)}}{2N^{3/2}} m_{\rm sc}(z)^2 m_{\rm sc}'(z) - m_{\rm sc}(z) m_{\rm sc}'(z) + m_{\rm sc}'(z) m_{\rm$$

{prop:giigaa-e

{sec:expectati

{lem:exp-secon

**Proof.** We first have,

$$\sum_{ja} \frac{s_{ja}}{N} \mathbb{E}[\partial_{ja} G_{ja}(z)] = -\sum_{ja} \frac{s_{ja}}{N} \mathbb{E}[G_{aa}(z) G_{jj}(z)] - \sum_{j \neq a} \frac{s_{ja}}{N} \mathbb{E}[G_{ja}^2(z)]$$
(7.24) {eqn:exp-second

We then write,

$$\sum_{ja} \frac{s_{ja}}{N} \mathbb{E}[G_{aa}G_{jj}(z)] = Nm_{\rm sc}(z)^2 + 2m_{\rm sc}\sum_{a} \mathbb{E}[G_{aa}(z) - m_{\rm sc}(z)] + \sum_{ja} \frac{s_{ja}}{N} \mathbb{E}[(G_{aa}(z) - m_{\rm sc}(z))(G_{jj}(z) - m_{\rm sc}(z))]$$
(7.25)

Now by Proposition 7.4 the second line is  $\mathcal{O}_{<}(N^{-1}\eta^{-2})$ . By Corollary 6.7 we see that

$$\sum_{j \neq a} \frac{s_{ja}}{N} \mathbb{E}[G_{ja}^2(z)] = -m_{\rm sc}(z)^2 {\rm tr}S + m_{\rm sc}(z)^2 {\rm tr}(S(1 - m_{\rm sc}(z)^2 S)^{-1}) + \mathcal{O}_{\prec}(N^{-1}\eta^{-2}).$$
(7.26)

This completes the proof of (7.22). The equation (7.23) follows from substituting the result of (7.22) into (5.34) and using that  $m'_{\rm sc}(z) = -\frac{m_{\rm sc}(z)}{2m_{\rm sc}(z)+z}$ .

Lemma 7.6. Define,

We have,

$$\int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{z}}\tilde{f}(z))E(z)\mathrm{d}z\mathrm{d}\bar{z} = \int_{\mathbb{C}} (\partial_{\bar{z}}\tilde{f}(z))E(z)\mathrm{d}z\mathrm{d}\bar{z} + \mathcal{O}_{<}(N^{\mathfrak{a}-1}\|f''\|_{1,w})$$
(7.28) {eqn:exp-determinants} (7.28)

and

$$\frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{z}} \tilde{f}(z)) E(z) dz d\bar{z} = \frac{f(2) + f(-2)}{4} + \left( \sum_{j \neq a} \frac{s_{aj}^{(4)}}{N^2} \right) \frac{1}{2\pi} \int_{-2}^2 f(x) \frac{x^4 - 4x^2 + 2}{\sqrt{4 - x^2}} dx + \operatorname{tr}(S) \frac{1}{2\pi} \int_{-2}^2 f(x) \frac{2 - x^2}{\sqrt{4 - x^2}} dx + \left( \sum_a \frac{s_{aa}^{(3)}}{N^{3/2}} \right) \frac{1}{8\pi} \int_{-2}^2 f(x) \frac{x^3 - x^2 - 2x + 4}{\sqrt{4 - x^2}} dx + \frac{1}{\pi} \int_{-2}^2 \frac{f(x)}{\sqrt{4 - x^2}} \left( \operatorname{Re}[m_{\mathrm{sc}}(x + \mathrm{i}0)^2 \operatorname{tr}(S(1 - m_{\mathrm{sc}}(x + \mathrm{i}0)^2 S)^{-1})] \right) dx =: E_S(f).$$
(7.29) {eqn:exp-determinants}

Furthermore,

$$\left|\operatorname{Re}[m_{\mathrm{sc}}(x+\mathrm{i}0)^{2}\operatorname{tr}(S(1-m_{\mathrm{sc}}(x+\mathrm{i}0)^{2}S)^{-1})]\right| \lesssim 1$$
 (7.30) {eqn:exp-deter

for  $x \in (-2, 2)$ .

**Proof.** By the Sherman-Morrison formula we have that,

$$\operatorname{tr}(S(1 - m_{\operatorname{sc}}^2(z)S)^{-1}) = \operatorname{tr}(S(1 - m_{\operatorname{sc}}^2(z)A)^{-1}) + \frac{m_{\operatorname{sc}}(z)^2}{1 - m_{\operatorname{sc}}(z)^2}, \quad (7.31) \quad \{\operatorname{eqn:sherm-1}\}$$

with A defined as in Lemma 2.8. By Lemma B.1 the first term on the RHS is bounded. Since  $|m_{\rm sc}(z)| \leq 1 - c |\text{Im}[z]|$  for some c > 0 we have that,

$$|E(z)| \lesssim \frac{|m'_{\rm sc}(z)|}{|{\rm Im}[z]|}.$$
 (7.32)

From this, the estimate  $|m'_{\rm sc}(z)| \leq |4 - \operatorname{Re}[z]^2|^{-1/2}$ , and the explicit form of  $\partial_{\bar{z}}\tilde{f}(z)$  (see (2.6)), the estimate (7.28) follows.

We now turn to the computation of the LHS of (7.29). By [26, (4.69)] we have,

$$\frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{z}} \tilde{f}(z)) \sum_{j \neq a} \frac{s_{aj}^{(4)}}{N^2} m_{\rm sc}(z)^3 m_{\rm sc}'(z) \mathrm{d}z \mathrm{d}\bar{z} = \left(\sum_{j \neq a} \frac{s_{aj}^{(4)}}{N^2}\right) \frac{1}{2\pi} \int_{-2}^2 f(x) \frac{x^4 - 4x^2 + 2}{\sqrt{4 - x^2}} \mathrm{d}x \quad (7.33)$$

and by [26, (4.68)]

$$\frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{z}} \tilde{f}(z)) (-m_{\rm sc}(z) m_{\rm sc}'(z) {\rm tr} S) {\rm d}z {\rm d}\bar{z} = {\rm tr}(S) \frac{1}{2\pi} \int_{-2}^{2} f(x) \frac{2-x^2}{\sqrt{4-x^2}} {\rm d}x$$
(7.34)

and then by similar arguments as to the proof the above equalities,

$$\left(-\sum_{a} \frac{s_{aa}^{(3)}}{2N^{3/2}}\right) \frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{z}} \tilde{f}(z)) m_{\rm sc}(z)^2 m_{\rm sc}'(z) \mathrm{d}z \mathrm{d}\bar{z}$$
$$= \left(\sum_{a} \frac{s_{aa}^{(3)}}{N^{3/2}}\right) \frac{1}{8\pi} \int_{-2}^2 f(x) \frac{x^3 - x^2 - 2x + 4}{\sqrt{4 - x^2}} \mathrm{d}x.$$
(7.35)

We have only left to compute the contribution of the  $tr(S(1 - m_{sc}(z)^2 S)^{-1})$  term of E(z). We write,

$$m_{\rm sc}(z)m_{\rm sc}'(z){\rm tr}(S(1-m_{\rm sc}(z)^2S)^{-1}) = m_{\rm sc}(z)m_{\rm sc}'(z)\left({\rm tr}(A(1-m_{\rm sc}(z)^2A)^{-1}) + \frac{1}{1-m_{\rm sc}(z)^2}\right).$$
(7.36)

By [26, (4.70)],

$$\frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{z}} \tilde{f}(z)) \frac{m_{\rm sc}(z)m_{\rm sc}'(z)}{1 - m_{\rm sc}(z)^2} \mathrm{d}z \mathrm{d}\bar{z} = -\frac{1}{2\pi} \int_{-2}^{2} \frac{f(x)}{\sqrt{4 - x^2}} \mathrm{d}x + \frac{f(2) - f(-2)}{4}.$$
(7.37)

By Green's theorem (2.28),

$$\frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{z}} \tilde{f}(z)) m_{\rm sc}(z) m_{\rm sc}'(z) \operatorname{tr}(A(1 - m_{\rm sc}(z)^2 A)^{-1}) \mathrm{d}z \mathrm{d}\bar{z}$$
  
$$= \frac{1}{\pi} \int_{-2}^{2} \frac{f(x)}{\sqrt{4 - x^2}} \operatorname{Re} \left[ m_{\rm sc}(x + \mathrm{i}0)^2 \operatorname{tr}(A(1 - m_{\rm sc}(x + \mathrm{i}0)^2 A)^{-1}) \right] \mathrm{d}x.$$
(7.38)

Therefore,

$$\frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{z}} \tilde{f}(z)) m_{\rm sc}(z) m'_{\rm sc}(z) \operatorname{tr}(S(1 - m_{\rm sc}(z)^2 S)^{-1}) \mathrm{d}z \mathrm{d}\bar{z} \\
= \frac{f(2) - f(-2)}{4} + \frac{1}{\pi} \int_{-2}^{2} \frac{f(x)}{\sqrt{4 - x^2}} \left( \operatorname{Re}[m_{\rm sc}(x + \mathrm{i}0)^2 \operatorname{tr}(A(1 - m_{\rm sc}(x + \mathrm{i}0)^2 A)^{-1})] - \frac{1}{2} \right) \mathrm{d}x \\
= \frac{f(2) - f(-2)}{4} + \frac{1}{\pi} \int_{-2}^{2} \frac{f(x)}{\sqrt{4 - x^2}} \left( \operatorname{Re}[m_{\rm sc}(x + \mathrm{i}0)^2 \operatorname{tr}(S(1 - m_{\rm sc}(x + \mathrm{i}0)^2 S)^{-1})] \right) \mathrm{d}x. \quad (7.39)$$

The second equality follows from,

$$m_{\rm sc}(z)^2 {\rm tr}(S(1 - m_{\rm sc}(z)^2 S)^{-1}) = m_{\rm sc}(z)^2 {\rm tr}(A(1 - m_{\rm sc}(z)^2 A)^{-1}) + \frac{m_{\rm sc}(z)^2}{1 - m_{\rm sc}(z)^2}$$
(7.40) {eqn:sherm-3}

and  $m'_{\rm sc}(z) = \frac{m_{\rm sc}(z)^2}{1-m_{\rm sc}(z)^2}$  and  $\operatorname{Re}[m'_{\rm sc}(x+i0)] = -\frac{1}{2}$ . This completes the proof of (7.29). By taking the real part on both sides of (7.40) we conclude the estimate (7.30).

{prop:exp-corr

Proposition 7.7. We have that,

$$\mathbb{E}[\operatorname{tr} f(H)] = N \int f(x)\rho_{\mathrm{sc}}(x) \mathrm{d}x + E_S(f) + \mathcal{O}_{\prec}(N^{\mathfrak{a}-1} \| f'' \|_{1,w})$$
(7.41) {eqn:exp-correction}

**Proof.** From (7.28), (7.29) and (2.22) it suffices to show that

$$\int_{\Omega_{a}} (\partial_{\bar{z}} \tilde{f}(z)) (N\mathbb{E}[m_{N}(z) - m_{\rm sc}(z)] - E(z)) dz d\bar{z} = \mathcal{O}_{\prec}(N^{-1} \| f'' \|_{1,w}).$$
(7.42) {eqn:exp-1a}

From (7.23) we have that,

$$|N\mathbb{E}[m_N(z) - m_{\rm sc}(z)] - E(z)| < \frac{1}{N\eta^2 \sqrt{\eta + |4 - E^2|}}.$$
(7.43)

It is then straightforward to modify the proof of (2.26) to derive (7.42).

# 8 Characteristic function

In this section we complete our expansion for the characteristic function,

$$\mathbb{E}[e_{\mathfrak{a}}] = \mathbb{E}\left[\exp\left(\mathrm{i}\lambda \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))(\sum_{a} G_{aa}(u) - \mathbb{E}[G_{aa}(u)])\mathrm{d}u\mathrm{d}\bar{u}\right)\right].$$
(8.1)

Throughout this section we will assume that  $|\lambda \Phi| \leq N^{-c}$  for some c > 0. We will apply Stein's method which involves showing that,

$$\frac{\mathrm{d}}{\mathrm{d}\lambda}\mathbb{E}[e_{\mathfrak{a}}(\lambda)] \approx -\lambda V(f)\mathbb{E}[e_{\mathfrak{a}}(\lambda)]$$
(8.2)

for a certain function V(f) to be computed later. Differentiating  $\mathbb{E}[e_{\mathfrak{a}}(\lambda)]$  we see that we need to investigate the quantity,

$$\sum_{a} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])].$$
(8.3)

In order to compute this quantity, we will apply the expansion (5.33), on which the above appears on the LHS. In order to use (5.33), we must compute the first term on its RHS. We

 $\{\texttt{sec:char}\}$ 

write this term as,

$$\sum_{ja} \frac{s_{aj}}{N} \left( \mathbb{E}[\partial_{ja}(e_{\mathfrak{a}}G_{ja})] - \mathbb{E}[e_{\mathfrak{a}}]\mathbb{E}[\partial_{ja}G_{ja}] \right)$$

$$= -\frac{i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{w}}\tilde{f}(w))\partial_{w} \sum_{ja} \frac{s_{aj}}{N} (2 - \delta_{ja})\mathbb{E}[e_{\mathfrak{a}}G_{aj}(z)G_{ja}(w)] dw d\bar{w}$$

$$-\sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z)G_{jj}(z) - \mathbb{E}[G_{aa}(z)G_{jj}(z)])]$$

$$-\sum_{j\neq a} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aj}(z)^{2} - \mathbb{E}[G_{aj}(z)^{2}])]$$

$$(8.4) \quad \{\text{eqn:char-second}\}$$

{lem:char-seco

and begin computing the various terms on the RHS of the above. The terms on the last two lines are computed in the following. We will assume that  $z = E \pm i\eta$  with  $\eta > 0$  and that  $z \in \Omega_{\mathfrak{a}}$ .

Lemma 8.1. We have,

$$\sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z)G_{jj}(z) - \mathbb{E}[G_{aa}(z)G_{jj}(z)])] = 2m_{\mathrm{sc}}(z)\sum_{a} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])] + \mathcal{O}_{<}(N^{-1}\eta^{-2} + |\lambda|\Phi\Psi(z)\eta^{-1})$$
(8.5) {eqn:char-second

and

$$\begin{aligned} \left| \sum_{j \neq a} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aj}(z)^{2} - \mathbb{E}[G_{aj}(z)^{2}])] \right| &< N^{-1}\eta^{-2} \\ + (\eta)^{-1} |\lambda| [(\Phi^{2}(1+|\lambda|)\Psi(z) + \min\{N^{-1}\eta^{-1}, \Phi^{2}\}] + N|\lambda|\Psi_{1}(z)^{2}\Phi\min\{\Psi_{1}(z), \Phi\} \end{aligned}$$
(8.6) {eqn:char-second

**Proof.** The estimate (8.6) follows immediately from Corollary 6.7. We can write the LHS of (8.5) as,

$$\sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z)G_{jj}(z) - \mathbb{E}[G_{aa}(z)G_{jj}(z)])]$$

$$= 2\sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[G_{jj}(z)] \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])]$$

$$+ \sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])(G_{jj}(z) - \mathbb{E}[G_{jj}(z)])]$$

$$- \sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}] \mathbb{E}[(G_{aa}(z) - \mathbb{E}[G_{aa}(z))(G_{jj}(z) - \mathbb{E}[G_{jj}(z)])]$$
(8.7) {eqn:char-seco

Due to Proposition 5.13 and (2.7) we have,

$$2\sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[G_{jj}] \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])]$$
  
=2m<sub>sc</sub>(z)  $\sum_{a} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])] + \mathcal{O}_{\prec}(N^{-1}\eta^{-2})$  (8.8)

We now turn to the term on the second last line of (8.7). We have,

$$\sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])(G_{jj}(z) - \mathbb{E}[G_{jj}(z)])]$$

$$= \sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - m_{\mathrm{sc}}(z))(G_{jj}(z) - m_{\mathrm{sc}}(z))]$$

$$-2\sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - m_{\mathrm{sc}}(z))]\mathbb{E}[G_{jj}(z) - m_{\mathrm{sc}}(z)]$$

$$+ \sum_{ja} \mathbb{E}[e_{\mathfrak{a}}] \frac{s_{aj}}{N} \mathbb{E}[G_{aa} - m_{\mathrm{sc}}(z)]\mathbb{E}[G_{jj} - m_{\mathrm{sc}}(z)]$$
(8.9) {eqn:char-seco

By Proposition 7.4, the term on the second line is  $\mathcal{O}_{<}(N^{-1}\eta^{-2}+|\lambda|\Phi\Psi(z)\eta^{-1})$ . By Proposition 5.13 and (2.7), the terms on the third and fourth lines are  $\mathcal{O}_{<}(N^{-1}\eta^{-2})$ . Therefore,

$$\sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])(G_{jj}(z) - \mathbb{E}[G_{jj}(z)])] = \mathcal{O}_{\prec}(N^{-1}\eta^{-2} + |\lambda|\Phi\Psi(z)\eta^{-1}).$$
(8.10)

By almost the same argument, the same estimate (with  $\lambda = 0$ ) holds for the term on the fourth line of (8.7). This completes the proof of (8.5).

Recall the notation  $M := m_{\rm sc}(z)m_{\rm sc}(w)$ . The following lemma computes the first term on the RHS of (8.4).

Lemma 8.2. We have,

$$\frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{w}}\tilde{f}(w))\partial_{w} \sum_{j\neq a} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}G_{aj}(z)G_{ja}(w)] \mathrm{d}w \mathrm{d}\bar{w}$$

$$= \mathbb{E}[e_{\mathfrak{a}}] \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{w}}\tilde{f}(w))\partial_{w} \left(-M\mathrm{tr}S + M\mathrm{tr}(S(1-MS)^{-1})\right) \mathrm{d}w \mathrm{d}\bar{w} + |\lambda|\mathcal{O}_{<}(\eta^{-1}\Phi^{2}(1+|\lambda|))$$

$$(8.11) \quad \{\mathrm{eqn:char-var}\}$$

and

$$\frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{w}}\tilde{f}(w))\partial_{w} \sum_{a} \frac{s_{aa}}{N} \mathbb{E}[e_{\mathfrak{a}}G_{aa}(z)G_{aa}(w)] \mathrm{d}w \mathrm{d}\bar{w}$$
$$= \mathbb{E}[e_{\mathfrak{a}}] \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{w}}\tilde{f}(w))\partial_{w} M \mathrm{tr}S \mathrm{d}w \mathrm{d}\bar{w} + \mathcal{O}_{\prec}(\eta^{-1}\Phi^{2}|\lambda|). \tag{8.12}$$

**Proof.** The estimate (8.11) is an immediate application of Corollary 6.7, and using (2.26) to integrate the error terms in w. The estimate (8.12) follows from,

$$\sum_{a} \frac{s_{aa}}{N} G_{aa}(z) G_{aa}(w) = m_{\rm sc}(z) m_{\rm sc}(w) \operatorname{tr}(S) + \mathcal{O}_{<}(N^{-1}y^{-1} + N^{-1}\eta^{-1})$$
(8.13)

which follows from (2.10) and (2.7).

In the following we summarize our expansion for the characteristic function. In preparation, define

$$\tilde{s}_4 := \sum_{a \neq j} \frac{s_{aj}^{(4)}}{N^2}, \qquad \tilde{s}_3 := \sum_a \frac{s_{aa}^{(3)}}{N}$$
(8.14)

{lem:char-seco

and

$$a_{k,\mathfrak{a}} := \frac{1}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}} \tilde{f}(u)) \partial_{u}(m_{\mathrm{sc}}(u)^{k}) \mathrm{d}u \mathrm{d}\bar{u}.$$

$$(8.15) \quad \{\texttt{eqn:ak-def}\}$$

$$\{\texttt{prop:char-ste}\}$$

Proposition 8.3. We have,

$$\begin{aligned} &(z+2m_{\rm sc}(z))\sum_{a}\mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z)-\mathbb{E}[G_{aa}(z)])]\\ =&\mathbb{E}[e_{\mathfrak{a}}]\frac{\mathrm{i}\lambda}{\pi}\int_{\Omega_{\mathfrak{a}}}(\partial_{\bar{w}}\tilde{f}(w))\partial_{w}(-2M\mathrm{tr}(S(1-MS)^{-1}+M\mathrm{tr}S)\mathrm{d}w\mathrm{d}\bar{w}\\ &+\mathbb{E}[e_{\mathfrak{a}}]\bigg\{-\tilde{s}_{4}\mathrm{i}\lambda a_{2,\mathfrak{a}}m_{\rm sc}(z)^{2}+\frac{\tilde{s}_{3}\mathrm{i}\lambda}{2N^{1/2}}m_{\rm sc}(z)\left[2m_{\rm sc}(z)a_{1,\mathfrak{a}}+\mathrm{i}\lambda a_{1,\mathfrak{a}}^{2}+a_{2,\mathfrak{a}}\right]\bigg\}\\ &+\mathcal{O}_{\prec}\left(N^{-1}(1+|\lambda|)^{3}+N^{-1}\eta^{-2}+|\lambda\Phi\Psi(z)|\eta^{-1}+|\lambda|\eta^{-1}\Phi^{2}(1+|\lambda|)\right) \end{aligned} \tag{8.16}$$

Therefore,

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}\lambda} \mathbb{E}[e_{\mathfrak{a}}(\lambda)] &= -\lambda V_{\mathfrak{a}}(f) \mathbb{E}[e_{\mathfrak{a}}(\lambda)] + \mathrm{i}\lambda^2 B_{\mathfrak{a}}(f) \mathbb{E}[e_{\mathfrak{a}}(\lambda)] \\ &+ \mathcal{O}_{\prec}((1+|\lambda|)^3 N^{-1} + \Phi_w^2 (1+|\lambda|)^2) \end{aligned}$$

$$(8.17) \quad \{\texttt{eqn:stein-1}\}$$

where,

$$V_{\mathfrak{a}}(f) := \frac{1}{\pi^2} \int_{\Omega_{\mathfrak{a}}^2} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)) \partial_w \left( m_{\mathrm{sc}}'(z) m_{\mathrm{sc}}(w) (2\mathrm{tr}(S(1-MS)^{-1}) - \mathrm{tr}S) \right) \mathrm{d}w \mathrm{d}\bar{w} \mathrm{d}z \mathrm{d}\bar{z} + \frac{1}{2} \tilde{s}_4 a_{2,\mathfrak{a}}^2 - \frac{\tilde{s}_3}{N^{1/2}} a_{2,\mathfrak{a}} a_{1,\mathfrak{a}}$$
(8.18) {eqn:Va-def}

and

$$B_{\mathfrak{a}}(f) := \frac{\tilde{s}_3}{2N^{1/2}} (a_{1,\mathfrak{a}})^3.$$
(8.19)

**Proof.** The equation (8.16) follows from (5.33), (8.4) (which expands the first term on the RHS of (5.33)) and Lemmas 8.1 and 8.2 (which compute the terms on the RHS of (8.4)). The equation (8.17) then follows from using (8.16) to solve for the expression  $\sum_{a} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])]$  in the equality,

$$\frac{\mathrm{d}}{\mathrm{d}\lambda}\mathbb{E}[e_{\mathfrak{a}}(\lambda)] = \frac{\mathrm{i}}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{z}}\tilde{f}(z)) \sum_{a} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])] \mathrm{d}z \mathrm{d}\bar{z},$$
(8.20)

and using a slight modification of Lemma 2.10 to integrate the error term (i.e., we use the estimate  $|z+2m_{\rm sc}(z)|^{-1} \leq C/\sqrt{|E-2|+\eta}$  and arrive at errors with  $||f''||_{1,w}$  instead of  $||f''||_1$ .

# 8.1 Computation of deterministic coefficients

#### 8.1.1 Chebyshev polynomials

We recall now the Chebyshev polynomials of the first kind  $T_n(x)$  and coefficients  $t_n(f)$  as defined in Definition 1.3. We note the the orthogonality relation,

$$\int_{-2}^{2} T_n(x) T_n(x) \frac{\mathrm{d}x}{\sqrt{4-x^2}} = \begin{cases} 0, & \text{if } n \neq m \\ \pi, & \text{if } n = m = 0 \\ \frac{\pi}{2}, & \text{if } n = m \neq 0 \end{cases}$$
(8.21)

 $\{\texttt{sec:cheby-1}\}$ 

so that we have the expansion,

$$f = T_0(x)\frac{t_0(f)}{2} + \sum_{n \ge 1} T_n(x)t_n(f).$$
(8.22)

as  $T_n(x)$  is a complete orthogonal family of  $L^2((-2,2), \frac{\mathrm{d}x}{\sqrt{4-x^2}})$ . We have for  $x \in (-2,2)$ ,

$$T_n(x) = \frac{(-1)^n}{2} \left( m_{\rm sc}(x+{\rm i}0)^n + m_{\rm sc}(x-{\rm i}0)^n \right) = (-1)^n \operatorname{Re}\left[ m_{\rm sc}(x+{\rm i}0)^n \right]$$
(8.23) {eqn:Chebyshev

which is easily derived from (1.12). In particular,

$$t_n(f) = \frac{2(-1)^n}{\pi} \int_{-2}^2 f(x) \operatorname{Re}[m_{\mathrm{sc}}(x+\mathrm{i}0)^n] \frac{\mathrm{d}x}{\sqrt{4-x^2}}.$$
(8.24) {eqn:tn-msc}

## 8.1.2 Easy coefficients

In this section we compute the coefficients  $a_{k,\mathfrak{a}}$  defined in (8.15) in terms of the  $t_n(f)$ .

**Lemma 8.4.** For each  $k \ge 1$ ,

$$a_{k,\mathfrak{a}} = \frac{1}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u}(m_{\mathrm{sc}}^{k}(u))\mathrm{d}u\mathrm{d}\bar{u} = \frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u}(m_{\mathrm{sc}}^{k}(u))\mathrm{d}u\mathrm{d}\bar{u} + \mathcal{O}_{<}((N^{\mathfrak{a}-1})^{3/2} \|f''\|_{1})$$

$$(8.25) \quad \{\mathrm{eqn:easy-coeff}\}$$

and for  $k \geq 1$ ,

$$\frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{u}} \tilde{f}(u)) \partial_u(m_{\mathrm{sc}}^k(u)) \mathrm{d}u \mathrm{d}\bar{u} = (-1)^k \frac{k}{2} t_k(f).$$
(8.26) {eqn:easy-coef

**Proof.** The first equality in (8.25) is by definition. The estimate in (8.25) follows from the fact that  $|\partial_z m_{\rm sc}(z)| \leq |{\rm Im}[z]|^{-1/2}$  and direct computation.

For (8.26) we first note that Green's theorem (2.28) implies that,

$$\int_{\mathbb{C}} (\partial_{\bar{u}} \tilde{f}(u)) \frac{m_{\rm sc}(u)^k}{2m_{\rm sc}(u) + u} \mathrm{d}u \mathrm{d}\bar{u} = \int_{\mathbb{R}} f(x) \mathrm{Im} \left[ \frac{m_{\rm sc}(x + \mathrm{i}0)^k}{x + 2m_{\rm sc}(x + \mathrm{i}0)} \right] \mathrm{d}x.$$
(8.27)

The equality (8.26) follows from this, as well as the identities,  $\partial_u(m_{\rm sc}(u)^k) = -km_{\rm sc}(u)^k/(u+2m_{\rm sc}(u))$  and

$$\operatorname{Im}\left[\frac{m_{\rm sc}(x+{\rm i}0)^k}{x+2m_{\rm sc}(x+{\rm i}0)}\right] = -\frac{\operatorname{Re}[m_{\rm sc}(x+{\rm i}0)^k]}{\sqrt{4-x^2}},\tag{8.28}$$

and (8.24).

## 8.1.3 Variance functional

In this section we compute the variance function  $V_{\mathfrak{a}}(f)$  as defined in (8.18). Let us define,

$$F(z,w) := \partial_w \left( m'_{\rm sc}(z) m_{\rm sc}(w) (2 {\rm tr}(S(1-m_{\rm sc}(z)m_{\rm sc}(w)S)^{-1}) - {\rm tr}S) \right)$$
(8.29) {eqn:var-F-def

{sec:var-funct

{sec:easy}
{lem:easy}

Lemma 8.5. We have,

$$\int_{\Omega_{\mathfrak{a}}^{2}} (\partial_{\bar{z}}\tilde{f})(\partial_{\bar{w}}\tilde{f}(w))F(z,w)dzd\bar{z}dwd\bar{w}$$

$$= \int_{\mathbb{C}^{2}} (\partial_{\bar{z}}\tilde{f})(\partial_{\bar{w}}\tilde{f}(w))F(z,w)dzd\bar{z}dwd\bar{w} + \mathcal{O}_{\prec}((N^{\mathfrak{a}-1}(\|f''\|_{1,w} + \|f'\|_{1,w} + \|f\|_{1,w}))^{2}) \quad (8.30) \quad \{\texttt{eqn:var-aa1}\}$$

**Proof.** By the Sherman Morrison formula,

$$\operatorname{tr}(S(1-MS)^{-1}) = \operatorname{tr}(S(1-MA)^{-1}) + \frac{m_{\mathrm{sc}}(z)m_{\mathrm{sc}}(w)}{1-m_{\mathrm{sc}}(z)m_{\mathrm{sc}}(w)}.$$
(8.31) {eqn:sherman-1}

Since  $||MA||_{\ell^2 \to \ell^2} \le 1 - c$  for some c > 0 we can conclude, using Lemma B.1 that

$$\left|\partial_{w}[m'_{\rm sc}(z)m_{\rm sc}(w){\rm tr}(S(1-MA)^{-1})]\right| \leq |m'_{\rm sc}(z)m'_{\rm sc}(w)|.$$
 (8.32)

Since  $|1 - m_{\rm sc}(z)m_{\rm sc}(w)| \ge c(y + \eta)$  we conclude that,

$$|F(z,w)| \lesssim \frac{|m'_{\rm sc}(z)m'_{\rm sc}(w)|}{(y+\eta)^2}.$$
(8.33) {eqn:var-aa2}

From this estimate we easily conclude, by direct integration,

$$\left| \int_{\Omega^c_{\mathfrak{a}} \times \Omega^c_{\mathfrak{a}}} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)(w)) F(z,w) \mathrm{d}z \mathrm{d}\bar{z} \mathrm{d}w \mathrm{d}\bar{w} \right| < (N^{\mathfrak{a}-1} \| f'' \|_{1,w})^2.$$
(8.34)

as well as,

$$\left| \int_{\Omega_{\mathfrak{a}}^{c} \times \Omega_{\mathfrak{a}}} (\partial_{\bar{z}} \tilde{f}) (\partial_{\bar{w}} \tilde{f}(w)) F(z, w) \mathrm{d}z \mathrm{d}\bar{z} \mathrm{d}w \mathrm{d}\bar{w} \right| < (N^{\mathfrak{a}-1} (\|f''\|_{1,w} + \|f'\|_{1,w} + \|f\|_{1,w}))^{2}, \quad (8.35)$$

which yields the claim.

**Proposition 8.6.** We have that,

$$\frac{1}{\pi^2} \int_{\mathbb{C}^2} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)) \partial_w (m'_{\rm sc}(z) m_{\rm sc}(w) \operatorname{tr}(S(1 - m_{\rm sc}(z) m_{\rm sc}(w)S)^{-1}) \mathrm{d}z \mathrm{d}\bar{z} \mathrm{d}w \mathrm{d}\bar{w}$$

$$= \frac{1}{4} \sum_{j=1}^{\infty} j t_j(f)^2 \operatorname{tr}(S^j) \tag{8.36} \quad \{\operatorname{eqn:determ-value}$$

as well as,

$$\frac{1}{\pi^2} \int_{\mathbb{C}^2} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)) m'_{\rm sc}(z) m'_{\rm sc}(w) \mathrm{tr}S = \mathrm{tr}S \frac{1}{4} t_1(f)^2 \tag{8.37} \quad \{\mathrm{eqn:determ-value}\}$$

**Proof.** For  $0 < \delta < 1$  we have  $|1 - \delta m_{\rm sc}(z)m_{\rm sc}(w)| \ge 1 - \delta$  and so,

$$\left|\frac{1}{1-\delta m_{\rm sc}(z)m_{\rm sc}(w)} - \frac{1}{1-m_{\rm sc}(z)m_{\rm sc}(w)}\right| \le \frac{1-\delta}{|1-\delta m_{\rm sc}(z)m_{\rm sc}(w)|} \frac{1}{|1-m_{\rm sc}(z)m_{\rm sc}(w)|} \le \frac{1}{|1-m_{\rm sc}(w)|} \le \frac{1}{|1$$

{lem:var-1}

{prop:var-1}

Therefore using (8.31) and dominated convergence we find,

$$\int_{\mathbb{C}^2} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)) \partial_w (m'_{\rm sc}(z) m_{\rm sc}(w) \operatorname{tr}(S(1 - m_{\rm sc}(z) m_{\rm sc}(w)S)^{-1}) \mathrm{d}z \mathrm{d}\bar{z} \mathrm{d}w \mathrm{d}\bar{w}$$
$$= \lim_{\delta \to 1^-} \int_{\mathbb{C}^2} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)) \partial_w (m'_{\rm sc}(z) m_{\rm sc}(w) \operatorname{tr}(S(1 - \delta m_{\rm sc}(z) m_{\rm sc}(w)S)^{-1}) \mathrm{d}z \mathrm{d}\bar{z} \mathrm{d}w \mathrm{d}\bar{w}.$$
(8.39)

Now,

$$\partial_{w}(m_{\rm sc}'(z)m_{\rm sc}(w){\rm tr}(S(1-\delta m_{\rm sc}(z)m_{\rm sc}(w)S)^{-1}) = \partial_{w}\sum_{j=0}^{\infty}m_{\rm sc}'(z)m_{\rm sc}(w)^{j+1}m_{\rm sc}(z)^{j}\delta^{j}{\rm tr}S^{j+1}$$
$$=\sum_{j=0}^{\infty}(j+1)m_{\rm sc}'(z)m_{\rm sc}'(w)(\delta m_{\rm sc}(z)m_{\rm sc}(w))^{j}{\rm tr}S^{j+1}.$$
(8.40)

The integration of summation and differentiation is justified because  $|m_{\rm sc}(z)| \le 1$  and

$$\left|\operatorname{tr} S^{j}\right| \leq N \|S\|_{\ell^{2} \to \ell^{2}}^{j} \leq N.$$
(8.41) {eqn:trS-bd}

Therefore,

$$\frac{1}{\pi^2} \int_{\mathbb{C}^2} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)) \partial_w (m'_{\rm sc}(z) m_{\rm sc}(w) \operatorname{tr}(S(1 - \delta m_{\rm sc}(z) m_{\rm sc}(w)S)^{-1}) \mathrm{d}z \mathrm{d}\bar{z} \mathrm{d}w \mathrm{d}\bar{w}$$

$$= \sum_{j=1}^\infty \delta^{j-1} \frac{j t_j(f)^2}{4} \operatorname{tr}(S^j), \qquad (8.42)$$

using (8.24) as in the proof of Lemma 8.4. Again, interchanging the order of integration and summation is justified using that  $|m_{\rm sc}(z)| \leq 1$  and (8.41). Finally, by monotone convergence,

$$\lim_{\delta \to 1^{-}} \sum_{j=1}^{\infty} \delta^{j-1} \frac{jt_j(f)^2}{4} \operatorname{tr}(S^j) = \frac{1}{4} \sum_{j=1}^{\infty} jt_j(f)^2 \operatorname{tr}(S^j)$$
(8.43)

which completes the proof of (8.36). The equation (8.37) follows from (8.26).

{lem:var-2}

# Lemma 8.7. We have that,

$$V_{\mathfrak{a}}(f) = V_{1}(f) + \mathcal{O}_{\prec}(N^{\mathfrak{a}-1}(\|f''\|_{1,w} + \|f'\|_{1,w} + \|f\|_{1,w})$$
(8.44) {eqn: Va-V}

where  $V_1(f)$  is defined in (1.14). Furthermore,

$$V_1(f) \ge 0. \tag{8.45} \quad \{\texttt{eqn:V-pos}\}$$

We have also,

$$B_{\mathfrak{a}}(f) = \frac{\tilde{s}_3}{8N^{1/2}} t_1(f)^3 + \mathcal{O}_{\prec}(N^{\mathfrak{a}-3/2} \| f'' \|_1)$$
(8.46) {eqn:Ba-s}

**Proof.** First of all since  $\hat{s}_{4,1} = \tilde{s}_4 + \mathcal{O}(N^{-1})$  and  $|t_2(f)| \leq ||f||_{1,w}$ , we have that (8.44) follows from Lemmas 8.4 and 8.5 and Proposition 8.6. Similarly, (8.46) follows from Lemma 8.4. It remains to prove (8.45).

Let  $a = t_1(f)$  and  $b = t_2(f)$  for notational simplicity. We have,

$$\begin{split} V_1(f) \geq &\frac{a^2}{4} \operatorname{tr}(S) + b^2 \left( \operatorname{tr}(S^2) + \frac{\hat{s}_{4,1}}{2} \right) + \frac{\tilde{s}_3}{2N^{1/2}} ab \\ &= b^2 \left( \frac{1}{N^2} \sum_{i \neq j} s_{ij}^2 + \frac{s_{ij}^{(4)}}{2} \right) \\ &+ \frac{a^2}{4} \operatorname{tr}(S) + b^2 \left( \frac{1}{N^2} \sum_i s_{ii}^2 + \frac{s_{ii}^{(4)}}{2} \right) + ab \frac{\tilde{s}_3}{2N^{1/2}} \end{split}$$
(8.47) {eqn:var-nonnet

We recall that the third and fourth cumulants of a centered random variable X are given by,

$$\kappa_3(X) = \mathbb{E}[X^3], \qquad \kappa_4(X) = \mathbb{E}[X^4] - 3\mathbb{E}[X^2]^2.$$
(8.48)

The nonnegativity of the RHS of (8.47) follows from the following two inequalities. First,

$$\frac{1}{N^2} \sum_{i \neq j} 2s_{ij}^2 + s_{ij}^{(4)} = \sum_{i \neq j} \mathbb{E}[H_{ij}^4] - \mathbb{E}[H_{ij}^2]^2 = \sum_{i \neq j} \mathbb{E}[(H_{ij}^2 - \mathbb{E}[H_{ij}^2])^2] \ge 0$$
(8.49) {eqn:var-nonne}

Second,

$$\begin{aligned} \left| \frac{\tilde{s}_3}{2N^{1/2}} ab \right| &= \left| \frac{1}{2} \operatorname{Cov} \left( a \sum_i H_{ii}, b \sum_i H_{ii}^2 \right) \right| \le \frac{1}{4} \operatorname{Var} \left( a \sum_i H_{ii} \right) + \frac{1}{4} \operatorname{Var} \left( b \sum_i H_{ii}^2 \right) \\ &= \frac{a^2 \operatorname{tr}(S)}{4} + \frac{b^2}{4N^2} \left( \sum_i 2s_{ii}^2 + s_{ii}^{(4)} \right) \\ &\le \frac{a^2 \operatorname{tr}(S)}{4} + \frac{b^2}{2N^2} \left( \sum_i 2s_{ii}^2 + s_{ii}^{(4)} \right). \end{aligned}$$

$$(8.50) \quad \{ \operatorname{eqn:var-nonne} x = \frac{1}{2} \operatorname{var} x = \frac{1$$

The last inequality follows because  $\kappa_4(X) + 2\kappa_2(X)^2 \ge 0$  for any random variable X. Define,

$$\hat{V}_p(f) := \frac{1}{2} \sum_{j=1}^{\infty} j t_j(f)^2 \mathrm{tr} S^j.$$
(8.51)

{lem:var-3}

Lemma 8.8. We have that,

$$\hat{V}_{p}(f) = \frac{1}{2\pi^{2}} \int_{-2}^{2} \int_{-2}^{2} \left(\frac{f(x) - f(y)}{x - y}\right)^{2} \frac{4 - xy}{\sqrt{4 - x^{2}}\sqrt{4 - y^{2}}} \mathrm{d}x\mathrm{d}y + \frac{1}{\pi^{2}} \int_{-2}^{2} \int_{-2}^{2} f(x)f(y) \frac{g(x, y)}{\sqrt{4 - x^{2}}\sqrt{4 - y^{2}}} \mathrm{d}x\mathrm{d}y$$
(8.52)

where,

$$g(x,y) := \operatorname{Re}\left[\operatorname{tr}\frac{m_{\rm sc}(x+{\rm i}0)m_{\rm sc}(y+{\rm i}0)A}{(1-m_{\rm sc}(x+{\rm i}0)m_{\rm sc}(y+{\rm i}0)A)^2} + \operatorname{tr}\frac{m_{\rm sc}(x+{\rm i}0)m_{\rm sc}(y-{\rm i}0)A}{(1-m_{\rm sc}(x+{\rm i}0)m_{\rm sc}(y-{\rm i}0)A)^2}\right]$$
(8.53)

The function g obeys  $|g(x,y)| \leq 1$  for all  $x, y \in (-2,2)$ .

**Proof.** We need to re-compute the integral on the LHS of (8.36). By (8.31) we have that

$$\partial_w (m'_{\rm sc}(z)m_{\rm sc}(w){\rm tr}(S(1-m_{\rm sc}(z)m_{\rm sc}(w)S)^{-1})) = \frac{m'_{\rm sc}(z)m'_{\rm sc}(w)}{(1-m_{\rm sc}(z)m_{\rm sc}(w))^2} + m'_{\rm sc}(z)m'_{\rm sc}(w){\rm tr}\frac{A}{(1-MA)^2}.$$
(8.54)

By [26, (4.60)] and (2.19) we have,

$$\frac{2}{\pi^2} \int_{\mathbb{C}^2} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)) \frac{m_{\rm sc}'(z) m_{\rm sc}'(w)}{(1 - m_{\rm sc}(z) m_{\rm sc}(w))^2} dz d\bar{z} dw d\bar{w}$$
$$= \frac{1}{2\pi^2} \int_{-2}^2 \int_{-2}^2 \left( \frac{f(x) - f(y)}{x - y} \right)^2 \frac{4 - xy}{\sqrt{4 - x^2} \sqrt{4 - y^2}} dx dy \tag{8.55}$$

On the other hand, by applying Green's theorem (2.28) we see that,

$$\frac{2}{\pi^2} \int_{\mathbb{C}^2} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)) m'_{\rm sc}(z) m'_{\rm sc}(w) \operatorname{tr} \frac{A}{(1 - MA)^2} \mathrm{d}z \mathrm{d}\bar{z} \mathrm{d}w \mathrm{d}\bar{w} 
= \frac{1}{2\pi^2} \int_{-2}^2 \int_{-2}^2 \frac{f(x) f(y)}{\sqrt{4 - x^2}\sqrt{4 - y^2}} \bigg\{ h(x + \mathrm{i}0, y + \mathrm{i}0) + h(x - \mathrm{i}0, y - \mathrm{i}0) 
+ h(x - \mathrm{i}0, y - \mathrm{i}0) + h(x + \mathrm{i}0, y - \mathrm{i}0) \bigg\} \mathrm{d}x \mathrm{d}y$$
(8.56)

where  $h(z, w) = m_{\rm sc}(z)m_{\rm sc}(w)\operatorname{tr}(A(1 - MA)^{-2})$ . The claim now follows, with the bound on g following Lemma B.1.

# 8.2 Stein's method

We finally apply Stein's method to compute the characteristic function  $\mathbb{E}[e_{\mathfrak{a}}(\lambda)]$ .

**Lemma 8.9.** Suppose there is a c > 0 so that  $|\lambda \Phi| \leq N^{-c}$ . We have,

$$\begin{aligned} &\frac{\mathrm{d}}{\mathrm{d}\lambda} \mathbb{E}[e_{\mathfrak{a}}(\lambda)] = -\lambda V_{1}(f) \mathbb{E}[e_{\mathfrak{a}}(\lambda)] + \mathrm{i}\lambda^{2}B(f) \mathbb{E}[e_{\mathfrak{a}}(\lambda)] \\ &+ \mathcal{O}_{<}((1+|\lambda|)^{3}N^{-1} + \Phi_{w}^{2}(1+|\lambda|)^{2}) + \mathcal{O}_{<}(|\lambda|N^{-1+\mathfrak{a}}(\|f''\|_{1,w} + \|f'\|_{1,w} + \|f\|_{1,w}) \quad (8.57) \quad \{\texttt{eqn:stein-2}\} \end{aligned}$$

Consequently, if  $V_1(f) \ge c_1$  for some  $c_1 > 0$  then, for  $|\lambda \Phi| \le N^{-c}$  we have,

$$\mathbb{E}[e_{\mathfrak{a}}(\lambda)] = e^{-\lambda^2 V_1(f)/2 + i\lambda^3 B(f)/3} + \mathcal{O}_{<}(\Phi_w^2(1+|\lambda|) + N^{-1}(1+|\lambda|)^2) \\ + \mathcal{O}_{<}(N^{-1+\mathfrak{a}}(\|f''\|_{1,w} + \|f'\|_{1,w} + \|f\|_{1,w}))$$
(8.58) {eqn:result-1}

{lem:stein}

Otherwise for  $|\lambda \Phi| \leq N^{-c}$  we have,

$$\mathbb{E}[e_{\mathfrak{a}}(\lambda)] = e^{-\lambda^2 V_1(f)/2 + i\lambda^3 B(f)/3} + \mathcal{O}_{\prec}((1+|\lambda|)^4 N^{-1} + \Phi_w^2(1+|\lambda|)^3) + \mathcal{O}_{\prec}(|\lambda|^2 N^{-1+\mathfrak{a}}(\|f''\|_{1,w} + \|f'\|_{1,w} + \|f\|_{1,w})$$
(8.59) {eqn:result-2}

**Proof.** The estimate (8.57) follows from (8.17) and (8.44) and (8.46). Letting now  $\psi(\lambda) := \mathbb{E}[e_{\mathfrak{a}}(\lambda)]$  and  $\Phi(\lambda) := e^{-\lambda^2 V_1(f)/2 + i\lambda^3 B(f)/3}$  we have by differentiating  $\psi(\lambda)\Phi(\lambda)^{-1}$  that,

$$|\psi(\lambda) - \Phi(\lambda)| < \int_0^{|\lambda|} e^{(s^2 - \lambda^2)V_1(f)/2} \varepsilon(s) ds$$
(8.60)

with

$$\varepsilon(s) := (1+|s|)^3 N^{-1} + \Phi_w^2 (1+|s|)^2 + |s| N^{-1+\mathfrak{a}} (\|f''\|_{1,w} + \|f'\|_{1,w} + \|f\|_{1,w}).$$
(8.61)

The estimate (8.59) follows from the fact that  $|e^{(s^2-\lambda^2)V_1(f)/2}| \le 1$  for  $|s| \le |\lambda|$  since  $V_1(f) \ge 0$  from (8.47). The estimate (8.58) follows from using that

$$\int_{0}^{|\lambda|} e^{s^2 V_1(f)/2} |s|^k ds \lesssim (1+|\lambda|)^{k-1} e^{\lambda^2 V(f)/2}.$$
(8.62)

for  $k \ge 0$  if  $V_1(f) \ge c_1$ . This completes the proof.

#### 8.3 Proof of main theorems in the real symmetric case

In this section we prove Theorem 1.6 and Proposition 1.5 in the real symmetric case.

# 8.3.1 Proof of Theorem 1.6

The estimate for the expectation follows from Proposition 7.7. The two estimates for the characteristic function follow from Lemma 8.9, (2.22) and the just proved estimate for the expectation.

#### 8.3.2 Proof of Proposition 1.5

The lower bound for  $V_1(f)$  follows from (8.45). The upper bound for the part of  $V_1(f)$  that we denoted by  $\hat{V}_p(f)$  can be deduced from bounding the integral on the LHS of (8.36) using (8.33) and Lemma 2.10. The remaining components can be bounded by (8.37) and the fact that  $|t_n(f)| \leq C_n ||f||_{1,w}$  for any n > 0. The estimate for B(f) also follows from this. Finally, the estimate for  $E_1(f)$  follows from its definition as well as (7.30).

# 9 Application: max of log-characteristic polynomial

In this section we show how to extend the results of [11] to generalized Wigner matrices. We will discuss only the real symmetric case, the complex Hermitian case being similar. In this section we define the complex logarithm by  $\log(re^{i\theta}) := \log(r) + i\theta$  for  $\theta \in (-\pi, \pi]$ . Introduce,

$$L_N(z) := \sum_{j=1}^N \log(z - \lambda_j) - N \int_{\mathbb{R}} \log(z - x) \rho_{\rm sc}(x) \mathrm{d}x \tag{9.1}$$

where  $\lambda_j$  are the eigenvalues of a generalized Wigner matrix H. We will denote  $z = E + i\eta$ and assume  $\eta \ge 0$ . We will make the following assumption on the tail of the distribution of entries of H. {sec:real-proo

{sec:main-proo

 $\{\texttt{sec:funct-pro}$ 

{sec:max}

{ass:tail}

{thm:max}

{prop:max-lss}

**Assumption 9.1.** We assume that there is a c > 0 so that

$$\mathbb{P}\left[|H_{ij}| > \sqrt{N}x\right] \le c^{-1} \mathrm{e}^{-x^c} \tag{9.2}$$

for all x > 0, uniformly in i and j.

We will give a proof of the following. It extends Theorem 1.2 and Theorem 1.8(i) from the Wigner to generalized Wigner setting.

**Theorem 9.2.** Let H be a generalized Wigner matrix satisfying Assumption 9.1. Then, for any  $\varepsilon$  and  $\kappa > 0$ ,

$$\mathbb{P}\left[\sup_{|E|<2-\kappa}\frac{\operatorname{Re}[L_N(E)]}{\sqrt{2}\log N}\in(1-\varepsilon,1+\varepsilon)\right] = 1+o(1),\tag{9.3} \quad \{\operatorname{eqn:max-main}\right]$$

as well as, (for any choice of  $\pm$ )

and

for any choice of  $\pm$ .

The main input from our work will be the following.

**Proposition 9.3.** Fix  $K > 0, \frac{1}{2} > \varepsilon > 0, \kappa > 0$ . We have that,

$$\mathbb{P}\left[\sup_{z:|E|\leq 2-\kappa, \mathrm{e}^{-K(\log\log N)^{2}}\leq\eta\leq 1}|L_{N}(z)|>(\log N)^{\varepsilon}\right]\leq C\mathrm{e}^{-(\log N)^{\varepsilon}}$$
(9.6)

for some C > 0.

**Proof.** We show how to bound the real part of  $L_N(z)$ , with the imaginary part being similar. We start by proving the bound for fixed z. Define,

$$X_{z} := \frac{1}{2} \sum_{i} \log((\lambda_{i} - E)^{2} + \eta^{2}) - \frac{N}{2} \int \log((x - E)^{2} + \eta^{2}) \rho_{\rm sc}(x) dx = \operatorname{Re}[L_{N}(z)]$$
(9.7)

Let us denote by  $V_z$ ,  $\mathcal{E}_z$  and  $B_z$  the coefficients in Theorem 1.6 with the function  $\operatorname{Re}[\log(z-\cdot)]$ . By (9.20) below we have, for z as in the statement of the proposition, that  $V_z = -\log(\eta) + \mathcal{O}(1)$ , and that  $|B_z| \leq CN^{-1/2}$  and  $|\mathcal{E}_z| \leq C$ . It therefore follows from Theorem 1.6 that

$$\mathbb{E}[\mathrm{e}^{\mathrm{i}\lambda X_z}] = \mathrm{e}^{-\lambda^2 V_z/2 + \mathrm{i}\lambda \mathcal{E}_z} + \mathcal{O}_{\prec}(N^{-1/2}(1+|\lambda|)^4), \qquad (9.8) \quad \{\mathrm{eqn:max-char}\}$$

for all  $|\lambda| \leq N^{1/2-1/10}$ . Let  $u \geq 2$  and  $0.01 > \delta > 0$ . A straightforward argument using the rigidity estimates [20, Theorem 2.2] shows that,

$$\mathbb{P}\left[X_z - \mathcal{E}_z > u\right] \le \mathbb{P}\left[u < X_z - \mathcal{E}_z < N^{\delta}\right] + N^{-D}.$$
(9.9)

for any D > 0. Let  $\chi$  be a smooth function s.t.  $\chi(x) = 1$  for  $x \in (u, N^{\delta})$  and  $\chi(x) = 0$  for  $x \notin (u - 1, N^{\delta} + 1)$ . We have that for any M > 0,

$$|\hat{\chi}(\lambda)| \le N^{\delta} C_M (1+|\lambda|)^{-M}.$$
(9.10) {eqn:max-chi}

Then,

$$\mathbb{P}\left[u < X_{z} - \mathcal{E}_{z} < N^{\delta}\right] \leq \mathbb{E}[\chi(X_{z} - \mathcal{E}_{z})] \\
= \int_{\mathbb{R}} \hat{\chi}(\lambda) \mathbb{E}[e^{i\lambda X_{z}}] d\lambda \\
= \int_{|\lambda| \leq N^{\delta}} \hat{\chi}(\lambda) \mathbb{E}[e^{i\lambda(X_{E} - \mathcal{E}_{z})}] d\lambda + \mathcal{O}(N^{-1}) \\
= \int_{|\lambda| \leq N^{\delta}} \chi(\hat{\lambda}) e^{-\lambda^{2} V_{z}/2} d\lambda + \mathcal{O}(N^{6\delta} N^{-1/2}). \\
= \int_{\mathbb{R}} \chi(\hat{\lambda}) e^{-\lambda^{2} V_{z}/2} d\lambda + \mathcal{O}(N^{6\delta - 1/2})$$
(9.11)

Above, the third and fifth lines follow from (9.10) and the fact that  $V_z \ge 0$ . In the fourth line we used (9.8). Now let Z be a standard normal random variable. Then,

$$\int_{\mathbb{R}} \chi(\lambda) e^{-\lambda^2 V_z/2} d\lambda = \mathbb{E}[\chi(V_z^{1/2} Z)] \le \mathbb{P}\left[V_z^{1/2} Z > (u-1)\right] \le C e^{-(u-1)^2/(2V_z)}$$
(9.12)

From all of this we conclude that for all  $0 < \varepsilon < 1/2$  that

$$\mathbb{P}\left[|X_z| > (\log N)^{\varepsilon}\right] \le C e^{-(\log N)^{\varepsilon}}$$
(9.13) {eqn:max-est-1}

for some C > 0.

We have that

$$\partial_E X_z = N \operatorname{Re}[m_N(z) - m_{\rm sc}(z)], \qquad \partial_\eta X_z = N \operatorname{Im}[m_N(z) - m_{\rm sc}(z)]. \tag{9.14}$$

Let  $D_r(z)$  be the disc of radius r centered at  $z = E + i\eta$ . Assume that  $r < \eta/2$ . From the above estimates and [20, Theorem 2.1] we have that for some  $C_1 > 0$ ,

$$\mathbb{P}\left[\sup_{w\in D_r} |X_z - X_w| > e^{C_1(\log\log N)^2} \frac{r}{\eta}\right] \le N^{-D},$$
(9.15) {eqn:max-est-2}

We conclude the proof by taking a union bound of the estimate (9.13) over  $e^{C_2(\log \log N)^2}$  points, where  $C_2$  is chosen to depend on  $C_1$  and K. The claim follows.

#### 9.1 Proof of Theorem 9.2

In this section we detail how the proof of [11] can be modified to obtain the result for generalized Wigner matrices. The main point is that Proposition 9.3 is a sufficient substitute for [11, Corollary A.2] in order to obtain the main results. We first note that the main inputs of [11] from other works are: the rigidity estimates, from [20], listed as [11, Theorem 2.2]; the relaxation of eigenvalues under DBM, listed as [11, Proposition 4.1], using [10] as input; and the comparison techniques of [25], developed further in [11, Section 5]. All of these three input works prove their results for generalized Wigner matrices, and so these aspects of [11] hold for the generalized Wigner class without change.

We now give further details, beginning with the proof of (9.3). This theorem is proven in Section 6 of [11], building on the treatment of Gaussian divisible ensembles in Section 4, and the comparison arguments of Section 5. As indicated above, Section 5 goes through for generalized Wigner matrices without change, as it is based on [25] where the arguments were developed for generalized Wigner matrices. Section 4 requires some changes. Reading through line-by-line, one finds that the first place requiring changes is the proof of Proposition 4.3, which must be proven with the  $\lambda_k(t)$  (defined near [11, (4.1)]) starting from a generalized Wigner matrix instead of only a Wigner matrix. Reading through line-by-line, one can use Proposition 9.3 in place of [11, Corollary A.2] to obtain [11, (4.16)] and the estimate  $\mathbb{P}[\mathcal{G}_{\eta'}] =$ 1-o(1), with  $\mathcal{G}_{\eta'}$  defined in [11, (4.23)], in the same manner as in the proof of [11, Proposition 4.3]. Everything else is identical. The remainder of Section 4 is not needed for the proof of (9.3).

With [11, Proposition 4.3] extended to the case of generalized Wigner matrices, the proof of (9.3) given in [11, Section 6] goes through without change. We only remark that first: [11, Lemma 6.1] holds without change for generalized Wigner matrices. Second, the proof of the upper bound given in [11, Section 6.1] does not appear to be complete as of the writing of this paper, but the gap is extremely minor. In particular, we could not find the proof of the deterministic upper bound [11, (6.3)] within the paper. Instead, one can easily check that such an estimate holds with overwhelming probability, as long as one allows the set J to have cardinality  $Ne^{C(\log \log N)^2}$  for some large C > 0. Indeed, the estimate [11, (3.1)] allows one to add i/N to E. Then, estimates similar to (9.15) allow one to replace the supremum over a line with a supremum over the discrete set J. The arguments in [11, Section 5] are unchanged even though the size of J has increased slightly.

With the upper bound proven, we turn to the proof of the lower bound in Section 6.2 of [11]. The proof goes through without change, except that we could not quite follow all of the arguments in [11]. In particular, the proof of [11, (6.7)] relies on the proof of [11, (3.21)] the latter, as written, relies on [11, Lemma 2.5] which is proven only for  $\beta$ -ensembles. However one can see from the proof of [11, (3.21)] that the rigidity estimates of [11, Theorem 2.2] are sufficient to obtain [11, (6.7)].

For the remaining two results of Theorem 9.2 we first note that (9.4) and (9.5) are equivalent due to the fact that

$$|\{i : \lambda_i \le E\}| \ge k \iff \{\lambda_k \le E\},\tag{9.16}$$

and so it suffices to prove (9.5). For this, we follow directly the proof given in [11, Section 6.3]. Again, [11, Section 5] carries over without change to the generalized Wigner setting.

The proof of [11, (6.14)] follows from [11, Proposition 4.1] and Proposition 9.3 above. The rest of the proof is identical.

## 9.2 Estimates for the CLT functionals for the logarithm

Define,

$$V_{G\beta E}(f) := \frac{1}{2\pi^2 \beta} \int_{-2}^{2} \int_{-2}^{2} \left(\frac{f(x) - f(y)}{x - y}\right)^2 \frac{4 - xy}{\sqrt{4 - x^2}\sqrt{4 - y^2}} \mathrm{d}x \mathrm{d}y.$$
(9.17)

**Proposition 9.4.** Let  $\kappa > 0$  and let  $|E| \le 2 - \kappa$  and  $0 < \eta < 1$ . Define  $f_{\text{Re}}(x)$  and  $f_{\text{Im}}(x)$  by,

$$f_{\rm Re}(x) := {\rm Re}[\log(E + i\eta - x)], \qquad f_{\rm Im}(x) := {\rm Im}[\log(E + i\eta - x)]$$
(9.18)

Then,

$$V_{G\beta E}(f_{\rm Re}) = \beta^{-1} |\log(\eta)| + \mathcal{O}(1) = V_{G\beta E}(f_{\rm Im})$$

$$(9.19) \quad \{\texttt{eqn:max-var-1}\}$$

For any generalized Wigner matrix we have that for  $g = f_{Re}$  or  $g = f_{Im}$  that

 $V_{\beta}(g) = \beta^{-1} |\log(\eta)| + \mathcal{O}(1), \qquad \mathcal{E}_{\beta}(g) = \mathcal{O}(1), \qquad |B(g)| \le CN^{-1/2}. \tag{9.20} \quad \{\texttt{eqn:max-var-2}, a \le 0\}$ 

**Proof.** By [7, Lemma A.1] if we set,

$$L(z,w) := \int_{-2}^{2} \int_{-2}^{2} (\log(z-x) - \log(z-y))(\log(w-x) - \log(w-y))Q(x,y)dxdy \quad (9.21)$$

where,

$$Q(x,y) := \frac{4 - xy}{(x - y)^2 \sqrt{4 - x^2} \sqrt{4 - y^2}}$$
(9.22)

then

$$L(z,w) = 2\pi^2 \log\left[\frac{(z+R(z))(w+R(w))}{2(zw-4+R(z)R(w))}\right], \qquad R(z) := \sqrt{z^2-4}$$
(9.23)

for  $z, w \in \mathbb{C} \setminus (-\infty, 2]$  and R(z) defined with branch cut in [-2, 2]. Note that,

$$2\beta\pi^2 V_{G\beta E}(\text{Re}[\log(z-\cdot)]) = \frac{1}{4}(L(z,z) + 2L(z,\bar{z}) + L(\bar{z},\bar{z}))$$
(9.24)

and

$$2\beta\pi^2 V_{G\beta E}(\operatorname{Im}[\log(z-\cdot))) = \frac{1}{4}(2L(z,\bar{z}) - L(z,z) - L(\bar{z},\bar{z})).$$
(9.25)

It is clear that  $L(z, z) = \mathcal{O}(1) = L(\overline{z}, \overline{z})$  for  $|E| \le 2 - \kappa$  and  $0 < \eta < 10$ . Note that,

$$|z|^{2} - 4 + R(z)R(\bar{z}) = |z|^{2} - 4 + |R(z)|^{2} = E^{2} + \eta^{2} - 4 + |E^{2} - 4 - \eta^{2} + 2iE\eta|$$
  
$$= \frac{8\eta^{2}}{4 - E^{2}} + \mathcal{O}(\eta^{4}).$$
(9.26)

Hence,

$$\frac{1}{2}L(z,\bar{z}) = 2\pi^2 |\log \eta| + \mathcal{O}(1).$$
(9.27)

The claim (9.19) follows. The first estimate of (9.20) follows from the definition of V in (1.14) and Lemma 8.8. The estimate for  $\mathcal{E}_{\beta}(f)$  follows from its definition as well as (7.30). The estimate for B(f) follows directly from its definition.

#### Changes for complex Hermitian Wigner matrices Α

In the case of complex Hermitian Wigner matrices, we introduce the Wirtinger derivatives, for  $i \neq j$  by

$$\partial_{H_{ij}} = \frac{1}{2} \left( \partial_{\operatorname{Re}[H_{ij}]} - \mathrm{i} \partial_{\operatorname{Im}[H_{ij}]} \right), \qquad \partial_{\bar{H}_{ij}} = \frac{1}{2} \left( \partial_{\operatorname{Re}[H_{ij}]} + \mathrm{i} \partial_{\operatorname{Im}[H_{ij}]} \right)$$
(A.1)

We will denote  $\partial_{ij} = \partial_{H_{ij}}$  so that  $\partial_{ji} = \partial_{\bar{H}_{ij}}$ . We still denote  $\partial_{ii} = \partial_{H_{ii}}$ . In this section let us denote by  $\partial_{ij}^{(\mathbb{R})}$  the derivative of a function on the space of real symmetric matrices with respect to the (i, j)th element. The main algebraic difference between the complex Hermitian and real symmetric cases is that,

$$\partial_{ij}G_{ab}(z) = -G_{ai}(z)G_{jb}(z) \tag{A.2} \quad \{\texttt{eqn:derivative}\}$$

instead of

$$(1 + \delta_{ij})\partial_{ij}^{(\mathbb{R})}G_{ab}(z) = -G_{ai}(z)G_{jb}(z) - G_{aj}(z)G_{ib}(z).$$
(A.3) {eqn:derivative

Some general observations are then that any kind of monomial of Green's function elements that arises in the complex Hermitian case already has arisen in the real symmetric case. For example, in estimating third order contributions to cumulant expansions in the real case, we had to account for every term in an expression like  $(\partial_{ij}^{(\mathbb{R})})^2 G_{ab}(z)$ . In the complex case, the third derivatives will be combinations of  $\partial_{ij}\partial_{ji}G_{ab}$ ,  $\partial^2_{ij}G_{ab}$  and  $\partial^2_{ji}G_{ab}$ . These three expressions all contribute monomials in Green's function elements that already arose in the real case  $(\partial_{ij}^{(\mathbb{R})})^2 G_{ab}(z)$ , and so the work we have already done will apply without much change to the complex Hermitian case.

The main changes to the complex Hermitian case will then be that not every second order term that arose in the real symmetric case will arise in the complex Hermitian case. For example, when computing  $\mathbb{E}[G_{11}(z)]$  via the cumulant expansion, the second order terms in the real symmetric and complex Hermitian cases will be,

$$\partial_{1j}^{(\mathbb{R})}G_{j1}(z) = -G_{jj}(z)G_{11}(z) - G_{1j}(z)^2, \qquad \partial_{j1}G_{j1}(z) = -G_{jj}(z)G_{11}(z), \qquad (A.4) \quad \{\texttt{eqn:derivat} \} = -G_{jj}(z)G_{11}(z), \qquad (A.4)$$

respectively. So some second order terms present in the real symmetric case will not be there in the complex Hermitian case, simplifying the argument somewhat.

The only thing that one needs to be careful about is that in the real symmetric case one has  $G_{ij}(z) = G_{ji}(z)$  but in the complex Hermitian case,  $G_{ij}(\bar{z}) = G_{ji}(z)$ . We have already accounted for this in our notion of hypergraphs, in that we assume that the graphs are directed. In this way there is no ambiguity in the resolvent entry associated to an edge in our hypergraph (i.e., if e = (a, b) then the associated resolvent entry is  $G_{i_a i_b}(z)$  and not  $G_{i_b i_a}(z)$ ). Of course the estimates (2.10) and (2.9) are insensitive to whether we consider  $G_{ij}(z)$  or  $G_{ji}(z)$ .

However, there is one crucial aspect in a few of our proofs (in particular, Proposition 3.2) for which the order of indices matters. We sometimes use the fact that  $\sum_k G_{ik}(z)G_{ka}(w) =$  $\langle \delta_a, (H-z)^{-1}(H-w)^{-1}\delta_b \rangle$  crucially. We need to check that the terms that arise in our expansions are of the form  $G_{ik}(z)G_{ka}(w)$  and not  $G_{ik}(z)G_{ak}(w)$  as the same trick would not work for the latter monomials.

Let us now explain the underlying reason why the monomials we consider always have their indices appearing in the correct order. These terms all arise from second order terms in our

{a:cplx}

iv

cumulant expansions. The operation that generates the second order terms, differentiation by  $\partial_{ij}$ , has good algebraic properties that will not change the order that the indices appear in, and so our method will nonetheless work. I.e., if F is some monomial in resolvent entries, then the resolvent and cumulant expansion is schematically (keeping only the dominant second order terms)

$$G_{ab}F \to H_{aj}G_{jb}F \to \partial_{ja}(G_{jb}F).$$
 (A.5)

Then, due to (A.2), if  $\partial_{ja}$  hits any resolvent entry, then it adds j as a right index and a as a left index. Therefore, this expansion does not change whether a or b appear as right or left indices and introduces a new index j that appears once as a left index and once as a right index. Due to (A.8) below, a similar observation applies to when the derivative hits  $e_{\mathfrak{a}}$ .

In what follows we will go through the arguments of the main body of the paper, detailing the necessary changes. As is clear from the above discussion, for the most part the arguments needed to handle the complex Hermitian case are the same as the real symmetric cases and so little comment will be necessary. We mainly include this appendix for completeness, as there is little to no novelty introduced.

In the complex Hermitian case we fix the following notation,

$$\frac{s_{ij}}{N} := \mathbb{E}[|H_{ij}|^2], \qquad s_{ij}^{(k)} := \kappa_k(\sqrt{N} \mathrm{Re}[H_{ij}]), \qquad t_{ij}^{(k)} := \kappa_k(\sqrt{N} \mathrm{Im}[H_{ij}]), \qquad (A.6)$$

for  $k \ge 3$ . With this notation, by appling the cumulant expansion of Lemma 2.6 twice (once to the real part of  $H_{ij}$  and once to the imaginary part) we find that,

$$\mathbb{E}[H_{ij}F(H)] = \frac{s_{ij}}{N} \mathbb{E}[\partial_{ji}F(H)] + \sum_{k=3}^{K} \frac{s_{ij}^{(k)}}{(k-1)!N^{k/2}(1+\delta_{ij})^{k-1}} \mathbb{E}[(\partial_{ij}+\partial_{ji})^{k-1}F(H)] + \mathrm{i}\sum_{k=3}^{K} \frac{t_{ij}^{(k)}}{(k-1)!N^{k/2}} \mathbb{E}[(\mathrm{i}(\partial_{ij}-\partial_{ji}))^{k-1}F(H)] + \mathrm{error}$$
(A.7) {eqn:cumu-comp

with an error term having a similar form as in Lemma 2.6.

We first note that in Lemma 2.11, the equation (2.31) is replaced by,

$$\partial_{ij}e_{\mathfrak{a}} = -\frac{\mathrm{i}\lambda e_{\mathfrak{a}}}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{z}}\tilde{f}(z))\partial_{z}G_{ji}(z)\mathrm{d}z\mathrm{d}\bar{z} = \mathcal{O}_{<}(|\lambda\Phi|) \tag{A.8} \quad \{\texttt{eqn:delea-cpl}$$

and (2.32) is replaced by,

$$\partial_{ij}^{2-n}\partial_{ji}^{n}e_{\mathfrak{a}} = \mathcal{O}_{\prec}(|\lambda\Phi| + |\lambda|^{2}\Phi^{2}) + \mathbf{1}_{\{n=1\}}\lambda c_{f}.$$
(A.9) {eqn:delea-cpl

The proof is very similar. Moreover, the estimate in (2.35) holds with the LHS replaced by  $|\partial_{ij}^{n-m}\partial_{ji}^m e_{\mathfrak{a}}|$  for any  $0 \le m \le n$ . Again, the proof is similar.

# A.1 Loops estimate

We now detail the necessary changes in the complex Hermitian case in the arguments of Section 3. Definition 3.1 and the main result, Proposition 3.2 are unchanged. The statement

of Lemma 3.3 changes slightly as the estimates for the quantity (3.5) hold for any term of the form,

$$\partial_{1j}^{k-n} \partial_{j1}^n (G_{j2}(z) M_1 G_{a1}(w) M_2)$$
 (A.10) {eqn:del-loops

for any  $0 \le n \le k$ . The proof is almost identical (the order of derivative referred to in (iv) is in this case the power k in the expression (A.10)). Similarly, the statement of Lemma 3.4 is unchanged except that the  $\partial_{1j}^3$  in (3.6) is replaced by  $\partial_{1j}^{3-n} \partial_{j1}^n$  for  $0 \le n \le 3$  and the estimates hold for any such n.

In (3.7) of Lemma 3.5, the third order term on the second line is replaced by,

$$\sum_{j} \frac{s_{1j}^{(3)}}{2N^{3/2}(1+\delta_{1j})^2} \mathbb{E}[(\partial_{1j}+\partial_{j1})^2 (e_{\mathfrak{a}}G_{j2}(z)M_1G_{a1}(w)M_2)] -\mathrm{i} \sum_{j} \frac{t_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[(\partial_{1j}-\partial_{j1})^2 (e_{\mathfrak{a}}G_{j2}(z)M_1G_{a1}(w)M_2)], \qquad (A.11) \quad \{\mathrm{eqn:loops-exp}\}$$

due to (A.7).

The estimates in Lemma 3.6 for the third order terms must then be modified as follows. First, the estimates (3.9) and (3.11) hold with the  $\partial_{1j}^2$  on the LHS replaced by any of  $\partial_{1j}^2$ ,  $\partial_{1j}\partial_{j1}$  or  $\partial_{j1}^2$ . Similarly, the estimate (3.10) holds with any number of the  $\partial_{1j}$  appearing on the LHS replaced by  $\partial_{j1}$ . Again, the proof is almost identical.

After having addressed the third order terms, we now turn to the computation of the second order terms in the complex analog of (3.7). First, Lemma 3.7 is unchanged. For Lemma 3.10, we have that the term with  $D_2$  on the last line of (3.16) is not present, but the estimate (3.16) is otherwise unchanged. The difference is due to the fact that the second term on the RHS of (3.17) does not appear in the complex Hermitian case. In Lemma 3.12, we have that the term with  $D_4$  is not present in (3.21). Again, this is due to the action of operator  $\partial_{j1}$  in the complex case versus the real case (i.e., (A.2) vs (A.3)). Finally, the estimate (3.22) of Lemma 3.14 is unchanged except the prefactor  $\frac{2i\lambda}{\pi}$  is replaced by  $\frac{i\lambda}{\pi}$  due to (A.8). This completes the discussion of the second order terms in (3.7).

The outcome of all the above discussion is then that in (3.25) of Proposition 3.15, the terms with  $D_2$  and  $D_4$  do not appear, and that the  $2i\lambda$  is replaced by  $i\lambda$  in the last line. The rest of Section 3.1 is then unchanged, after accounting for this difference. The arguments of Section 3.2 and the proof of Proposition 3.2 given in Section 3.2.1 then go through without change.

We now turn to detailing how Section 3.3 changes in the complex case. First, the statement of Lemma 3.22 holds without change, given the changes to (3.7) and Lemma 3.6 we have outlined above. Next, the estimates (3.46), (3.47) and (3.48) hold without change, via similar proofs. The estimate (3.49) also holds without change except that in the proof, some terms estimated using Proposition 3.2 in the real symmetric case are simply not present in the complex case (and so the proof is easier). The remainder of the arguments of Section 3.3 go through without change and in particular the estimate (3.57) holds without change in the complex case.

#### A.2 Line estimates

We now detail the changes to Section 4 in the complex case. First, Definition 4.1 and the main result of this section, Proposition 4.2 are unchanged. For Lemma 4.3, the estimates of the term (4.2) hold with  $\partial_{j1}^k$  replaced by  $\partial_{1j}^{k-n}\partial_{j1}^n$  for any  $0 \le n \le k$ . The proof is similar. The estimates of Lemma 4.4 hold for the quantity (4.3) with the  $\partial_{j1}^3$  on the LHS replaced by  $\partial_{1j}^{3-n}\partial_{j1}^n$  for any  $0 \le n \le 3$ , and the proof is similar.

Now, in the expansion (4.4) of Lemma 4.5, the third order terms on the second line are replaced by,

$$\sum_{j} \frac{s_{1j}^{(3)}}{2N^{3/2}(1+\delta_{1j})^2} \mathbb{E}[(\partial_{1j}+\partial_{j1})^2 (e_{\mathfrak{a}}G_{j2}(z)M_1M_2)] -\mathrm{i} \sum_{j} \frac{t_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[(\partial_{1j}-\partial_{j1})^2 (e_{\mathfrak{a}}G_{j2}(z)M_1M_2)].$$
(A.12)

For the third order terms, the estimates (4.5), (4.6) (4.7), and (4.8) of Lemma 4.6 all hold, with any of the  $\partial_{j1}$  replaced by  $\partial_{1j}$ . The proofs are almost identical.

For the second order terms, the estimates of Lemma 4.7 hold without change. The estimate (4.12) of Lemma 4.9 holds except that the term with  $D_2$  on the second line is not present in the complex case. This is due to the fact that the second term on the last line of (4.13) is not present in the complex case. In Lemma 4.10, the equation (4.14) changes as the term with  $D_4$  in the second line is not present. In (4.15), the prefactor  $\frac{2i\lambda}{\pi}$  is replaced by  $\frac{i\lambda}{\pi}$ . Finally, the estimate (4.16) of Proposition 4.11 is unchanged except that the terms with  $D_2$  and  $D_4$  are not present, and the  $2i\lambda$  in the last line is replaced by  $i\lambda$ .

With the above, the arguments of Section 4.1 and the proof of Proposition 4.2 given in Section 4.1.1 are then unchanged.

We now turn to outlining how Section 4.2 changes. First, the result of Lemma 4.15 is unchanged, but the handling of the third order terms (i.e., the analog of the first term on the second line of (4.25)) changes somewhat. Specifically, one uses  $\partial_{1j}^{2-n} \partial_{j1}^n G_{j2}(z) = \mathbf{1}_{\{n=1\}} m_{\mathrm{sc}}(z)^2 G_{j2}(z) + \mathcal{O}_{\prec}(\Psi(z)^2)$  for  $0 \leq n \leq 2$ , and then an argument similar to (4.26) yields (4.24).

All of the three estimates of Lemma 4.16 are unchanged. Note that in the proof of (4.29), the second term on the second line of (4.31) is not present in the complex case. Finally, Proposition 4.17 follows in the complex case as in the real case.

## A.3 Preliminary expansion for $\mathbb{E}[e_{\mathfrak{a}}G_{11}]$

We now turn to the necessary changes to the arguments of Section 5 in the complex case. First, the estimates of Lemma 5.1 hold, with (5.3) also holding when  $\partial_{j1}^4$  on the LHS is replaced by  $\partial_{j1}^{4-n} \partial_{1j}^n$  for any  $0 \le n \le 4$ .

For the estimate (5.6), the third order terms are replaced with,

$$\sum_{j} \frac{s_{1j}^{(3)}}{2N^{3/2}(1+\delta_{1j})^2} \mathbb{E}[(\partial_{1j} + \partial_{j1})^2 (e_{\mathfrak{a}}G_{j1}(z))] - i\sum_{j} \frac{t_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[(\partial_{1j} - \partial_{j1})^2 (e_{\mathfrak{a}}G_{j1}(z))] \quad (A.13) \quad \{\texttt{eqn:cplx-c2}\}$$

and the fourth order terms are replaced by,

$$\sum_{j} \frac{s_{1j}^{(4)}}{6N^2(1+\delta_{1j})^3} \mathbb{E}[(\partial_{1j}+\partial_{j1})^3(e_{\mathfrak{a}}G_{j1}(z))] + \sum_{j} \frac{t_{1j}^{(4)}}{6N^2} \mathbb{E}[(\partial_{1j}-\partial_{j1})^3(e_{\mathfrak{a}}G_{j1}(z))]. \quad (A.14) \quad \{\texttt{eqn:cplx-c1}\}$$

We now turn to Section 5.1, which deals with the fourth order terms in (A.14). First, the estimate (5.8) is replaced by,

$$\sum_{j} \frac{s_{1j}^{(4)}}{N^2 (1+\delta_{1j})^3} \mathbb{E}[(\partial_{1j}+\partial_{j1})^3 (e_{\mathfrak{a}}G_{j1})] + \sum_{j} \frac{t_{1j}^{(4)}}{N^2} \mathbb{E}[(\partial_{1j}-\partial_{j1})^3 (e_{\mathfrak{a}}G_{j1}(z))]$$

$$= -m_{\rm sc}(z)^4 \sum_{j\neq 1} \frac{s_{1j}^{(4)}+t_{1j}^{(4)}}{N^2} 6\mathbb{E}[e_{\mathfrak{a}}] - 12m_{\rm sc}(z)^3 \sum_{j\neq 1} \frac{s_{1j}^{(4)}+t_{1j}^{(4)}}{N^2} \mathbb{E}[e_{\mathfrak{a}}(G_{11}(z)-m_{\rm sc}(z))]$$

$$+ 3\sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[((\partial_{1j}+\partial_{j1})^2 e_{\mathfrak{a}})(\partial_{1j}+\partial_{j1})G_{j1}] + 3\sum_{j\neq 1} \frac{t_{1j}^{(4)}}{N^2} \mathbb{E}[((\partial_{1j}-\partial_{j1})^2 e_{\mathfrak{a}})(\partial_{1j}-\partial_{j1})G_{j1}]$$

$$+ \mathcal{O}_{<}(N^{-2}\eta^{-1}+(1+|\lambda|)^3N^{-2}+|\lambda\Phi|\Psi(z)(1+|\lambda|). \qquad (A.15)$$

The proof is the same, except the identity

$$\partial_{j1}^{3-n} \partial_{1j}^{n} G_{j1}(z) = -2G_{jj}(z)^2 G_{11}(z)^2 \mathbf{1}_{\{n=1\}} + \mathcal{O}_{\prec}(\Psi(z)^2)$$
(A.16)

replaces (5.9). Instead of (5.12) of Lemma 5.4 we derive the estimate,

$$\sum_{j\neq 1} \frac{s_{1j}^{(4)}}{N^2} \mathbb{E}[((\partial_{1j} + \partial_{j1})^2 e_{\mathfrak{a}})(\partial_{1j} + \partial_{j1})G_{j1}] + \sum_{j\neq 1} \frac{t_{1j}^{(4)}}{N^2} \mathbb{E}[((\partial_{1j} - \partial_{j1})^2 e_{\mathfrak{a}})(\partial_{1j} - \partial_{j1})G_{j1}]$$

$$= -\mathbb{E}[e_{\mathfrak{a}}]m_{\mathrm{sc}}(z)^2 \sum_{j\neq 1} \frac{s_{1j}^{(4)} - t_{1j}^{(4)}}{N^2} \frac{2i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u m_{\mathrm{sc}}(u)^2 \mathrm{d}u \mathrm{d}\bar{u}$$

$$- \sum_{j\neq 1} \frac{s_{1j}^{(4)} - t_{ij}^{(4)}}{N^2} \frac{2i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u m_{\mathrm{sc}}(z)^2 m_{\mathrm{sc}}(u)\mathbb{E}[e_{\mathfrak{a}}(G_{11}(u) - m_{\mathrm{sc}}(u))]$$

$$- \sum_{j\neq 1} \frac{s_{1j}^{(4)} - t_{ij}^{(4)}}{N^2} \frac{2i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_u m_{\mathrm{sc}}(u)^2 m_{\mathrm{sc}}(z)\mathbb{E}[e_{\mathfrak{a}}(G_{11}(z) - m_{\mathrm{sc}}(z))]$$

$$+ \mathcal{O}_{<}(|\lambda|N^{-2}\eta^{-1} + |\lambda|(1 + |\lambda|)\Phi^2 N^{-1}). \tag{A.17}$$

The proof is the same, except that we use the estimates for n = 0, 1,

$$\hat{c}_{j1}^{1-n}\hat{c}_{1j}^{n}G_{j1}(z) = -\mathbf{1}_{\{n=1\}}G_{jj}G_{11} + \mathcal{O}_{\prec}(\Psi(z)^{2})$$
(A.18) {eqn:prelim-cp

and for m = 0, 1, 2,

$$\hat{c}_{j1}^{2-m}\hat{c}_{1j}^{m}e_{\mathfrak{a}} = \mathbf{1}_{\{m=1\}}\frac{\mathrm{i}\lambda}{\pi}\int_{\Omega_{\mathfrak{a}}}(\hat{c}_{\bar{u}}\tilde{f}(u))\hat{c}_{u}(G_{11}(u)G_{jj}(u)) + \mathcal{O}_{\prec}(|\lambda\Phi^{2}|(1+|\lambda|))$$
(A.19) (A.19) (A.19)

instead of (5.13) and (5.14). This completes our outline of the changes to the fourth order terms in (A.14)
We now turn to outlining the changes to the handling of the third order terms (A.13) in Section 5.2. First, estimate (5.16) of Lemma 5.5 holds without change. Next, estimate (5.19) of Lemma 5.6 changes to,

$$\sum_{j\neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}(\partial_{1j} + \partial_{j1})^2 G_{j1}(z)] - i \sum_{j\neq 1} \frac{t_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}(\partial_{1j} - \partial_{j1})^2 G_{j1}(z)]$$
  
= $m_{\mathrm{sc}}(z) \sum_{j\neq 1} \frac{1}{N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}(G_{jj} - m_{\mathrm{sc}}(z))(2(s_{ij}^{(3)} - \mathrm{i}t_{ij}^{(3)})G_{1j} + 4(s_{ij}^{(3)} + \mathrm{i}t_{ij}^{(3)})G_{j1})]$   
+ $\mathcal{O}_{<}(N^{-1/2}\Psi(z)^3 + |\lambda\Phi|N^{-1}\Psi(z)).$  (A.20)

The proof is the same except that we use

$$\partial_{1j}^{2-n} \partial_{j1}^n G_{j1}(z) = \mathbf{1}_{\{n=2\}} 2G_{jj} G_{11} G_{1j} + \mathbf{1}_{\{n=1\}} 2G_{jj} G_{11} G_{j1} + \mathcal{O}_{<}(\Psi(z)^3)$$
(A.21)

instead of (5.20) and we also use Proposition 4.17 to estimate both  $\mathbb{E}[e_{\mathfrak{a}}G_{1j}(z)]$  and  $\mathbb{E}[e_{\mathfrak{a}}G_{j1}(z)]$ . Instead of (5.24) of Lemma 5.7 we have,

$$\sum_{j \neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[((\partial_{1j} + \partial_{j1})e_{\mathfrak{a}})(\partial_{1j} + \partial_{j1})G_{j1}(z)] - i\sum_{j \neq 1} \frac{t_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[((\partial_{1j} - \partial_{j1})e_{\mathfrak{a}})(\partial_{1j} - \partial_{j1})G_{j1}(z)] \\ = \sum_{j \neq 1} \frac{m_{\mathrm{sc}}(z)}{N^{3/2}} \mathbb{E}[((s_{1j}^{(3)}(\partial_{1j} + \partial_{j1}) + it_{1j}^{(3)}(\partial_{1j} - \partial_{j1}))e_{\mathfrak{a}})(G_{jj}(z) - m_{\mathrm{sc}}(z))] \\ + \mathcal{O}_{<}(N^{-1/2}|\lambda\Phi|\Psi(z)^{2} + N^{-1}|\lambda|(1 + |\lambda|)\Phi^{2})$$
(A.22)

via almost the same proof, using the identity (A.18).

Instead of (5.28) of Lemma 5.8 we have,

$$\sum_{j \neq 1} \frac{s_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[((\partial_{1j} + \partial_{j1})^2 e_{\mathfrak{a}})G_{j1}(z)] - i \sum_{j \neq 1} \frac{t_{1j}^{(3)}}{N^{3/2}} \mathbb{E}[((\partial_{1j} - \partial_{j1})^2 e_{\mathfrak{a}})G_{j1}(z)]$$

$$= \frac{2i\lambda}{\pi} \sum_{j \neq 1} \frac{s_{1j}^{(3)} + it_{1j}^{(3)}}{N^{3/2}} \int_{\Omega_{\mathfrak{a}}} du d\bar{u} (\partial_{\bar{u}}\tilde{f}(u)) \partial_{u} m_{\mathrm{sc}}(u) \mathbb{E}[e_{\mathfrak{a}}G_{j1}(z)(G_{jj}(u) - m_{\mathrm{sc}}(u))]$$

$$+ \mathcal{O}_{<}(N^{-1/2} |\lambda \Phi^2| (1 + |\lambda|) \Psi(z) + |\lambda| N^{-1} \Psi(z)^2)$$
(A.23)

where we use (A.19) instead of (5.14). This completes the changes to the third order terms in (A.14).

We now turn to detailing the changes in Section 5.3. First, the estimate (5.30) of Propo-

sition 5.9 becomes,

$$\begin{aligned} z \sum_{a} \mathbb{E}[e_{\mathfrak{a}}G_{aa}(z) + N\mathbb{E}[e_{\mathfrak{a}}] &= \sum_{ja} \frac{s_{1j}}{N} \mathbb{E}[\partial_{ja}(e_{\mathfrak{a}}G_{ja})] \\ &- \sum_{j \neq a} \frac{s_{aj}^{(4)} + t_{aj}^{(4)}}{N^{2}} m_{\mathrm{sc}}(z)^{4} \mathbb{E}[e_{\mathfrak{a}}] - \mathbb{E}[e_{\mathfrak{a}}] m_{\mathrm{sc}}(z)^{2} \sum_{j \neq a} \frac{s_{aj}^{(4)} - t_{aj}^{(4)}}{N^{2}} \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u}m_{\mathrm{sc}}(u)^{2} \mathrm{d}u \mathrm{d}\bar{u} \\ &+ \sum_{a} \frac{s_{aa}^{(3)}}{2N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}] \Big\{ m_{\mathrm{sc}}(z)^{3} + 2m_{\mathrm{sc}}(z)^{2} \left( \frac{\mathrm{i}\lambda}{\pi} e_{\mathfrak{a}} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right) \\ &+ m_{\mathrm{sc}}(z) \left( -\frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right)^{2} + m_{\mathrm{sc}}(z) \left( \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u)^{2} \right) \Big\} \\ &+ \mathcal{O}_{<}(N^{-1}\eta^{-1}(1+|\lambda|) + (1+|\lambda|)^{3}N^{-1} + N^{1/2}\Psi(z)^{3} + N^{1/2}|\lambda\Phi|\Psi(z)^{2}) \\ &+ \mathcal{O}_{<}(N^{1/2}|\lambda\Phi^{2}|(1+|\lambda|)\Psi(z)) \end{aligned} \tag{A.24}$$

by using the above changes to the various lemmas of Sections 5.1 and 5.2. As in the proof of Proposition 5.9 in the real case, there are some extra terms that we can handle by (2.7) in the exact same manner.

We then find that (5.33) is replaced by,

$$\begin{split} z \sum_{a} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])] &= \sum_{ja} \frac{s_{aj}}{N} \left( \mathbb{E}[\partial_{ja}(e_{\mathfrak{a}}G_{ja}(z))] - \mathbb{E}[e_{\mathfrak{a}}]\mathbb{E}[\partial_{ja}G_{ja}(z)] \right) \\ &- \mathbb{E}[e_{\mathfrak{a}}]m_{\mathrm{sc}}(z)^{2} \sum_{j \neq a} \frac{s_{aj}^{(4)} + t_{aj}^{(4)}}{N^{2}} \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{u}}\tilde{f}(u))\partial_{u}m_{\mathrm{sc}}(u)^{2} \mathrm{d}u \mathrm{d}\bar{u} \\ &+ \sum_{a} \frac{s_{aa}^{(3)}}{2N^{3/2}} \mathbb{E}[e_{\mathfrak{a}}] \left\{ 2m_{\mathrm{sc}}(z)^{2} \left( \frac{\mathrm{i}\lambda}{\pi} e_{\mathfrak{a}} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right) \\ &+ m_{\mathrm{sc}}(z) \left( -\frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right)^{2} + m_{\mathrm{sc}}(z) \left( \frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} \mathrm{d}u \mathrm{d}\bar{u} \partial_{\bar{u}}\tilde{f}(u)\partial_{u}m_{\mathrm{sc}}(u) \right)^{2} \\ &+ \mathcal{O}_{<}(N^{-1}\eta^{-1}(1+|\lambda|) + (1+|\lambda|)^{3}N^{-1} + N^{1/2}\Psi(z)^{3} + N^{1/2}|\lambda\Phi|\Psi(z)^{2}) \\ &+ \mathcal{O}_{<}(N^{1/2}|\lambda\Phi^{2}|(1+|\lambda|)\Psi(z)) \end{split}$$
(A.25) {equ: intermedial

and that (5.34) is replaced by,

$$z \sum_{a} \mathbb{E}[G_{aa}(z)] + N = \sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[\partial_{ja}G_{ja}(z)] - \sum_{j \neq a} \frac{s_{aj}^{(4)} + t_{aj}^{(4)}}{N^2} m_{\rm sc}(z)^2 + \sum_{a} \frac{s_{aa}^{(3)}}{2N^{3/2}} m_{\rm sc}(z)^3 + \mathcal{O}_{\prec}(N^{-1}\eta^{-1} + N^{1/2}\Psi(z)^3).$$
(A.26) {eqn:intermediation of the second second

We now turn to outlining how Section 5.4 changes in the complex case. First, the estimate (5.35) of Proposition 5.11 holds without change. Indeed, the third and fourth order terms in the cumulant expansion are estimated using the estimates listed above, and the remainder of the terms are estimated in the same fashion as the proof of Proposition 5.11. The estimates (5.36) and (5.37) of Lemma 5.12 hold without change. Given these inputs, the rest of Section 5.4 then holds without change.

#### A.4 Estimates for $G_{12}G_{21}$

We now outline the changes to Section 6 in the complex case. First, the statement of Lemma 6.1 holds without change, but the proof changes somewhat. First, the third order term on the second line of (6.2) becomes

$$\sum_{j \neq 1} \frac{s_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[(\partial_{1j} + \partial_{j1})^2 (e_{\mathfrak{a}} G_{j2}(z) G_{21}(w))] - i \sum_{j \neq 1} \frac{t_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[(\partial_{1j} - \partial_{j1})^2 (e_{\mathfrak{a}} G_{j2}(z) G_{21}(w))]$$
(A.27)

In order to estimate this term, we then use, instead of (6.4), the estimate

$$\partial_{1j}^{2-n} \partial_{j1}^{n} G_{j2}(z) = \mathbf{1}_{\{n=1\}} m_{\rm sc}(z)^{2} G_{j2}(z) + \mathcal{O}_{\prec}(\Psi(z)^{2})$$
  

$$\partial_{1j}^{2-n} \partial_{j1}^{n} G_{21}(w) = \mathbf{1}_{\{n=1\}} m_{\rm sc}(w)^{2} G_{21}(w) + \mathcal{O}_{\prec}(\Psi(w)^{2})$$
  

$$\partial_{1j}^{1-m} \partial_{j1}^{m} G_{j2}(z) = -\mathbf{1}_{\{m=1\}} m_{\rm sc}(z) G_{12}(z) + \mathcal{O}_{\prec}(\Psi(z)^{2})$$
  

$$\partial_{1j}^{1-m} \partial_{j1}^{m} G_{21}(w) = -\mathbf{1}_{\{m=1\}} m_{\rm sc}(w) G_{2j}(w) + \mathcal{O}_{\prec}(\Psi(w)^{2})$$
(A.28)

The rest of the proof is similar (using at one point (A.9) instead of (2.32)). Next, the estimates of Proposition 6.2 all hold without change. In fact, the proofs are slightly easier as some terms we estimated in the real case are not present in the complex case (e.g., compare the two derivatives in (A.4)). After this, the remainder of the arguments of Section 6 do not involve any differentiation with respect to matrix entries and so the arguments are identical.

### A.5 Estimates for $G_{11}G_{22}$

We now outline the changes to Section 7 in the complex case. First, for Lemma 7.1, the estimate (7.3) is unchanged. The estimates (7.2) and (7.4) hold for any of the powers of  $\partial_{j1}$  on the LHS replaced by  $\partial_{1j}$ ; the proofs are similar. In (7.5), the third order term (the last term on the RHS of the first line) is replaced by,

$$\sum_{j} \frac{s_{1j}^{(3)}}{2N^{3/2}(1+\delta_{1j})^2} \mathbb{E}[(\partial_{1j}+\partial_{j1})^2 (e_{\mathfrak{a}}G_{j1}(z)X_1)] - \mathrm{i} \sum_{j} \frac{t_{1j}^{(3)}}{2N^{3/2}} \mathbb{E}[(\partial_{1j}-\partial_{j1})^2 (e_{\mathfrak{a}}G_{j1}(z)X_1)].$$
(A.29)

For the estimates of the third order terms carried out in Lemma 7.2, we have first that (7.6) is unchanged. The three remaining estimates hold also for any of the  $\partial_{j1}$  on the LHS replaced by  $\partial_{1j}$ , and the proofs are modified as follows. For (7.7), the proof uses (A.9). The proof of (7.8) uses

$$\partial_{j1}^{1-n} \partial_{1j}^{n} G_{j1}(z) = -\mathbf{1}_{\{n=1\}} m_{\rm sc}(z) + \mathcal{O}_{\prec}(\Psi(z)) \tag{A.30}$$

for n = 0, 1. The proof of (7.9) also uses this identity to reduce the estimate to estimating the first term on the RHS of (7.11) (note that this term is only present if a factor of  $\partial_{j1}$ hits  $G_{j1}(z)$ ). In this case, we need to estimate the first term on the RHS of (7.11) also with  $\partial_{j1}X_1$  replaced by  $\partial_{1j}X_1$ . The argument in (7.12) works for both of these cases, yielding the estimate.

All of the estimates of Lemma 7.3 are unchanged. The proofs are similar, except that some terms that were estimated in the real case are simply not present in the complex case. Finally, Proposition 7.4 and its proof are unchanged.

We now outline the changes to Section 7.1, in which we compute the correction to the expectation of a linear spectral statistic. First, recall that in the complex case, the expansion (5.34) is replaced by (A.26) above. Next, the estimates of Lemma 7.5 change somewhat. First, the two terms on the second line of (7.22) involving S are not present in the complex case. This is due to the fact that the second term on the RHS of (7.24) are not present in the complex case; the rest of the proof is the same. Consequently, (7.23) is replaced by,

$$\sum_{a} \mathbb{E}[G_{aa}(z) - m_{\rm sc}(z)] = \sum_{j \neq a} \frac{s_{aj}^{(4)} + t_{aj}^{(4)}}{N^2} m_{\rm sc}(z)^3 m_{\rm sc}'(z) - \sum_{a} \frac{s_{aa}^{(3)}}{2N^{3/2}} m_{\rm sc}(z)^2 m_{\rm sc}'(z) + \mathcal{O}_{\prec}(|m_{\rm sc}'(z)|N^{-1}\eta^{-2}),$$
(A.31)

where we note that the fourth order term also changed because (5.34) changed to (A.26). Consequently, the equations of Lemma 7.6 change. First, one should define E(z) in (7.27) as

$$E_{\mathbb{C}}(z) := \sum_{j \neq a} \frac{s_{aj}^{(4)} + t_{aj}^{(4)}}{N^2} m_{\rm sc}(z)^3 m_{\rm sc}'(z) - \sum_a \frac{s_{aa}^{(3)}}{2N^{3/2}} m_{\rm sc}(z)^2 m_{\rm sc}'(z).$$
(A.32)

Then, (7.28) holds with E(z) replaced by  $E_{\mathbb{C}}(z)$ . Equation (7.29) should be replaced with,

$$\frac{1}{\pi} \int_{\mathbb{C}} (\partial_{\bar{z}} \tilde{f}(z)) E(z) dz d\bar{z} = \left( \sum_{j \neq a} \frac{s_{aj}^{(4)} + t_{aj}^{(4)}}{N^2} \right) \frac{1}{2\pi} \int_{-2}^{2} f(x) \frac{x^4 - 4x^2 + 2}{\sqrt{4 - x^2}} dx + \left( \sum_{a} \frac{s_{aa}^{(3)}}{N^{3/2}} \right) \frac{1}{8\pi} \int_{-2}^{2} f(x) \frac{x^3 - x^2 - 2x + 4}{\sqrt{4 - x^2}} dx =: E_{S,\mathbb{C}}(f)$$
(A.33)

The estimate (7.30) is not needed in the complex case. Note that the majority of the proof of Lemma 7.6 is devoted to dealing with the tr $(S(1 - m_{\rm sc}(z)^2 S)^{-1})$  term which is not present in the complex case. Finally, in (7.41), the term  $E_S(f)$  should be replaced with  $E_{S,\mathbb{C}}(f)$ , but Proposition 7.7 and its proof is otherwise unchanged.

#### A.6 Characteristic function

We now outline how the arguments of Section 8 change in the complex case. First note that in the real symmetric case we rely on (5.33) whereas in the complex case we rely on (A.25). As in the real case, we must compute the second order terms in (A.25). In this case, we replace (8.4) with,

$$\sum_{ja} \frac{s_{aj}}{N} \left( \mathbb{E}[\partial_{ja}(e_{\mathfrak{a}}G_{ja}(z))] - \mathbb{E}[e_{\mathfrak{a}}]\mathbb{E}[\partial_{ja}G_{ja}(z)] \right)$$

$$= -\frac{i\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{w}}\tilde{f}(w))\partial_{w} \sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}G_{ja}(z)G_{aj}(w)] dw d\bar{w}$$

$$-\sum_{ja} \frac{s_{aj}}{N} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z)G_{jj}(z) - \mathbb{E}[G_{aa}(z)G_{jj}(z)])].$$
(A.34) {eqn:char-seco

For Lemma 8.1, the estimate (8.5) is unchanged and (8.6) is unneeded. The proof is identical. The estimates and proofs of Lemma 8.2 are also unchanged. As a consequence of replacing (5.33) by (A.25) and replacing (8.4) by (A.34), we have that (8.16) of Proposition 8.3 is replaced by,

$$(z + 2m_{\rm sc}(z))\sum_{a} \mathbb{E}[e_{\mathfrak{a}}(G_{aa}(z) - \mathbb{E}[G_{aa}(z)])]$$

$$= \mathbb{E}[e_{\mathfrak{a}}]\frac{\mathrm{i}\lambda}{\pi} \int_{\Omega_{\mathfrak{a}}} (\partial_{\bar{w}}\tilde{f}(w))\partial_{w}(-M\mathrm{tr}(S(1-MS)^{-1})\mathrm{d}w\mathrm{d}\bar{w}$$

$$+ \mathbb{E}[e_{\mathfrak{a}}]\left\{-\tilde{s}_{4,\mathbb{C}}\mathrm{i}\lambda a_{2,\mathfrak{a}}m_{\rm sc}(z)^{2} + \frac{\tilde{s}_{3}\mathrm{i}\lambda}{2N^{1/2}}m_{\rm sc}(z)\left[2m_{\rm sc}(z)a_{1,\mathfrak{a}} + \mathrm{i}\lambda a_{1,\mathfrak{a}}^{2} + a_{2,\mathfrak{a}}\right]\right\}$$

$$+ \mathcal{O}_{<}\left(N^{-1}(1+|\lambda|)^{3} + N^{-1}\eta^{-2} + |\lambda\Phi\Psi(z)|\eta^{-1} + |\lambda|\eta^{-1}\Phi^{2}(1+|\lambda|)\right) \qquad (A.35)$$

where,

$$\tilde{s}_{4,\mathbb{C}} := \sum_{a \neq j} \frac{s_{aj}^{(4)} + t_{aj}^{(4)}}{N^2}.$$
(A.36)

Consequently, (8.17) still holds after we replace  $V_{\mathfrak{a}}(f)$  by,

$$V_{\mathfrak{a},\mathbb{C}}(f) := \frac{1}{\pi^2} \int_{\Omega_{\mathfrak{a}}^2} (\partial_{\bar{z}} \tilde{f}(z)) (\partial_{\bar{w}} \tilde{f}(w)) \partial_w (m_{\mathrm{sc}}'(z) m_{\mathrm{sc}}(w) \mathrm{tr}(S(1-MS)^{-1})) \mathrm{d}w \mathrm{d}\bar{w} \mathrm{d}z \mathrm{d}\bar{z} + \frac{1}{2} \tilde{s}_{4,\mathbb{C}} a_{2,\mathfrak{a}}^2 - \frac{\tilde{s}_3}{N^{1/2}} a_{2,\mathfrak{a}} a_{1,\mathfrak{a}}.$$
(A.37)

We can leave  $B_{\mathfrak{a}}(f)$  unchanged.

Next, Sections 8.1.1 and 8.1.2 are unchanged. Some parts of Section 8.1.3 change. First, the relevant function replacing F in (8.29) is,

$$F_{\mathbb{C}}(z,w) := \partial_w (m'_{\rm sc}(z)m_{\rm sc}(w){\rm tr}(S(1-m_{\rm sc}(z)m_{\rm sc}(w)S)^{-1})).$$
(A.38)

With this replacement, the statement and proof of Lemma 8.5 are otherwise unchanged. Proposition 8.6 is unchanged. For Lemma 8.7, we replace  $V_1(f)$  with  $V_2(f)$  as defined in (1.14). With this definition the analog of (8.44) holds. For the remainder of Lemma 8.7 we need only verify that  $V_{\mathbb{C}}(f) \ge 0$ . For this we write with  $a = t_1(f)$  and  $b = t_2(f)$ ,

$$V_{\mathbb{C}}(f) \geq \frac{b^2}{2} \left( \sum_{i \neq j} \frac{s_{ij}^{(4)} + t_{ij}^{(4)}}{N^2} + \sum_{i \neq j} \frac{s_{ij}^2}{N^2} \right) + \frac{1}{4} a^2 \sum_i \frac{s_{ii}}{N} + \frac{b^2}{4} \left( \sum_i \frac{s_{ii}^{(4)}}{N^2} + 2 \sum_i \frac{s_{ii}^2}{N^2} \right) + \frac{ab}{2N^{1/2}} \left( \sum_i \frac{s_{ii}^{(3)}}{N^{3/2}} \right)$$
(A.39)

Non-negativity of the second line follows from the first two lines of (8.50). The non-negativity of the first line is proven in the same manner as (8.49) once one realizes that  $s_{ij}^2/N^2 = 2\mathbb{E}[\text{Re}[H_{ij}]^2]^2 + 2\mathbb{E}[\text{Im}[H_{ij}]^2]^2$ . Finally, Lemma 8.8 is unchanged.

The three estimates of Lemma 8.9 are unchanged after replacing  $V_1(f)$  with  $V_2(f)$ . Finally, the proofs in Section 8.3 are carried out in the same manner.

# **B** Linear algebra results

The following allows us to estimate matrix elements of operators similar to  $(1-m_{\rm sc}(z)m_{\rm sc}(w)S)^{-1}$ . **Lemma B.1.** Let M be an  $N \times N$  matrix with complex entries such that,

$$|M_{ij}| \le \frac{C}{N}, \qquad ||M||_{\ell^2 \to \ell^2} \le 1 - c$$
 (B.1)

for some C, c > 0. Then,

$$|(1-M)_{ij}^{-1}| \le C'(\delta_{ij}+N^{-1}).$$
 (B.2)

**Proof.** We have the identity,

$$\frac{1}{1-M} = 1 + M + M \frac{1}{1-M}M.$$
(B.3)

To prove the estimate it suffices to show that the matrix entries  $M(1-M)^{-1}M$  are bounded by C'/N. But this follows since,

$$|(M(1-M)^{-1}M)_{ab}| \le \frac{C^2}{N^2} \sum_{ij} |(1-M)_{ij}^{-1}| \le C'/N$$
(B.4)

where the last inequality uses the fact that  $\|(1-M)^{-1}\|_{\ell^2 \to \ell^2} \le C''$  for some C'' > 0.

### B.1 Derivatives of the resolvent wrt matrix entries

If we view the resolvent G = (H - z) as a function on the space of real symmetric  $N \times N$  matrices, then we have,

$$(1 + \delta_{ab})\partial_{ab}G_{ij} = -G_{ia}G_{bj} - G_{ib}G_{aj}.$$
(B.5) {eqn:delabgij}

# C Various proofs

Lemma C.1. We have,

$$\frac{1}{N}\sum_{j}G_{aj}(z)G_{j2}(w) = \frac{1}{N}\frac{G_{a2}(z) - G_{a2}(w)}{z - w} = \frac{\delta_{a2}}{N}\frac{m_{\rm sc}(z) - m_{\rm sc}(w)}{z - w} + \mathcal{O}_{\prec}\left(\frac{\Psi(z) + \Psi(w)}{N(|{\rm Im}[z]| + |{\rm Im}[w]|)}\right) \tag{C.1}$$

**Proof.** The first equality follows from  $(H-z)^{-1}(H-w)^{-1}(z-w) = (H-z)^{-1} - (H-w)^{-1}$ . For the estimate, let us assume that  $|\text{Im}[z]| \ge |\text{Im}[w]|$ , and for simplicity that Im[z] > 0, other cases being similar. If |z-w| > Im[z]/2, then the claim follows immediately from (2.10), as  $|z-w| \ge c \text{Im}[z]$ . On the other hand, if |z-w| < Im[z]/2, then the estimate follows from,

$$\frac{G_{a2}(z) - G_{a2}(w)}{z - w} = \int_0^1 (\partial_u G_{a2})(w + s(z - w)) ds$$
$$= \delta_{a2} \int_0^1 m'_{sc}(w + s(z - w)) ds + \mathcal{O}_{<}(\Psi(z)/\text{Im}[z])$$
(C.2)

{lem:G-dif}

#### C.1 Proof of Lemma 2.5

*H* and  $M^{(\theta)}$  be as in the statement of the lemma. Let  $\Delta_{ab}$  be the matrix that is 1 only for entries (a, b) and (b, a) and 0 otherwise. Assuming that  $\delta < \frac{1}{4}$  and using that  $||(M^{(\theta)} - z)^{-1}||_{\ell^2 \to \ell^2} \leq N$  we have by the resolvent expansion and (2.10) for *H*,

$$(M^{(\theta)} - z)_{ij}^{-1} = (H - z)_{ij}^{-1} + \sum_{m=1}^{20} \theta^m [(H - z)^{-1} (\Delta_{ab} (H - z)^{-1})^m]_{ij} + \mathcal{O}_{\prec} (N^{-2})$$
(C.3)

As long as, say,  $\delta < \varepsilon/2$  then every term in the sum is  $\mathcal{O}_{<}(N^{-1/2+\varepsilon})$  by applying the entrywise local law (2.10), and so (2.12) follows for  $M^{(\theta)}$ . Moreover, the terms for  $m \ge 2$  are  $\mathcal{O}_{<}(N^{-1+\varepsilon})$  so to establish (2.11) it suffices to to estimate the normalized trace of the term with m = 1. This term equals,

$$\frac{\theta}{N}\sum_{a}G_{ia}(z)G_{bi}(z) \prec \frac{\theta}{N\eta}(\operatorname{Im}[G_{aa}(z)] + \operatorname{Im}[G_{bb}(z)]) \prec \frac{1}{N\eta}$$
(C.4)

by the Ward identity. This completes the proof.

#### C.2 Proof of (2.22)

The difference between the LHS and the first term on the RHS of (2.22) is bounded by,

$$\int_{0 < y < N^{\mathfrak{a}^{-1}}} y |f''(x)| N |\operatorname{Im}[m_N(z)] - \mathbb{E}[\operatorname{Im}[m_N(z)]] | \mathrm{d}x \mathrm{d}y \tag{C.5}$$

Fixing an  $\varepsilon > 0$  we bound the integral over  $N^{\varepsilon-1} < y < N^{\mathfrak{a}-1}$  by  $\mathcal{O}_{<}(N^{\mathfrak{a}-1} || f'' ||_1)$  using (2.8). For the integral over  $y < N^{\varepsilon-1}$  we can use the fact that  $y \to y \operatorname{Im}[m_N(x + \mathrm{i}y)]$  is increasing to bound,

$$y \mathrm{Im}[m_N(x+\mathrm{i}y)] \le N^{\varepsilon-1} \mathrm{Im}[m_N(x+\mathrm{i}N^{\varepsilon-1})] < N^{\varepsilon-1}$$
(C.6)

with the second estimate following from (2.8). The claim now follows.

### C.3 Proof of Lemma 2.10

From the Cauchy integral formula,

$$\partial_z H(x + iy) = \frac{1}{2\pi i} \int_{|x + iy - w| = ry} \frac{H(w)}{(w - (x + iy))^2} dw$$
(C.7)

for any 0 < r < 1, and so

$$|\partial_z H(x+\mathrm{i}y)| \le \frac{C}{ry} \sup_{|w-(x+\mathrm{i}y)|=ry} |H(w)|. \tag{C.8} \quad \{\mathtt{eqn:a-H-est}\}$$

The estimate (2.27) follows by this and induction. The estimate (2.26) is a special case of (2.24) and so it remains to prove the latter.

a:eps-regular

{a:HS-est}

{a:H-est}

The terms on the LHS of (2.24) involving  $i(f(x) + iyf'(x))\chi'(y)$  are clearly bounded by the first term on the RHS of (2.24), and so it suffices to bound the integral involving only  $f''(x)y\chi(y)$ . Let  $T = ||f''||_1^{-1}$ . We first estimate,

$$\left| \int_{N^{\mathfrak{a}-1} < |y| < T} y\chi(y) f''(x) H(x + \mathrm{i}y) \mathrm{d}x \mathrm{d}y \right| \le C \|f''\|_1 \sup_{|x| \le 10} \int_{N^{\mathfrak{a}-1} < |y| < T} |yH(x + \mathrm{i}y)| \mathrm{d}y \quad (C.9)$$

and so this is bounded by the second term on the RHS of (2.24). By integration by parts and the fact that  $\partial_x H = \partial_z H$  by the Cauchy-Riemann equations we have,

$$\left| \int_{T < |y| < 2} \chi(y) y f''(x) H(x) dx dy \right| = \left| \int_{T < |y| < 2} \chi(y) y f'(x) \partial_z H(x) dx dy \right|$$
  
$$\leq \|f'\|_1 \sup_{|x| \le 10} \int_{T < |y| < 2} |S(x + iy)|$$
(C.10)

where we used (C.8) in the last inequality (with r = 1/2). This completes the proof.

#### C.4 Semicircle calculations

## C.4.1 Proof of Lemma 2.9

The estimate (2.17) follows from [14, (4.2)]. The estimate (2.18) is a consequence of (2.19) and [14, (4.2)]. The identity (2.19) follows from [27, Lemma B.1].

# References

- [1] K. Adhikari, I. Jana, and K. Saha. Linear eigenvalue statistics of random matrices with a variance profile. *Random Matrices: Theory and Applications*, 10(03):2250004, 2021.
- [2] O. H. Ajanki, L. Erdős, and T. Krüger. Universality for general Wigner-type matrices. Probability Theory and Related Fields, 169:667–727, 2017.
- [3] J. Alt, L. Erdős, T. Krüger, and D. Schröder. Correlated random matrices: band rigidity and edge universality. 2020.
- [4] G. W. Anderson and O. Zeitouni. A CLT for a band matrix model. Probab. Theory Related Fields, 134(2):283–338, 2006.
- [5] Z. Bai, X. Wang, and W. Zhou. CLT for linear spectral statistics of Wigner matrices. *Electron. J. Probab.*, 14:2391–2417, 2009.
- [6] Z. Bai and J. Yao. On the convergence of the spectral empirical process of Wigner matrices. *Bernoulli*, 11(6):1059–1092, 2005.
- [7] J. Baik and J. O. Lee. Fluctuations of the free energy of the spherical Sherrington-Kirkpatrick model. *Journal of Statistical Physics*, 165:185–224, 2016.
- [8] Z. Bao and Y. He. Quantitative CLT for linear eigenvalue statistics of Wigner matrices. The Annals of Applied Probability, 33(6B):5171–5207, 2023.

{a:semi-calc}

- [9] A. Bloemendal, L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Isotropic local laws for sample covariance and generalized Wigner matrices. 2014.
- [10] P. Bourgade. Extreme gaps between eigenvalues of Wigner matrices. J. Eur. Math. Soc., to appear, 2021.
- [11] P. Bourgade, P. Lopatto, and O. Zeitouni. Optimal rigidity and maximum of the characteristic polynomial of Wigner matrices. arXiv preprint arXiv:2312.13335, 2023.
- [12] S. Chatterjee. Fluctuations of eigenvalues and second order Poincaré inequalities. Probability Theory and Related Fields, 143(1):1–40, 2009.
- [13] M. Diaz, J. A. Mingo, and S. T. Belinschi. On the global fluctuations of block Gaussian matrices. Probability Theory and Related Fields, 176(1):599–648, 2020.
- [14] L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. The local semicircle law for a general class of random matrices. 2013.
- [15] L. Erdős. The matrix Dyson equation and its applications for random matrices. arXiv preprint arXiv:1903.10060, 2019.
- [16] L. Erdős and A. Knowles. The Altshuler–Shklovskii formulas for random band matrices I: the unimodular case. *Communications in Mathematical Physics*, 333(3):1365–1416, 2015.
- [17] L. Erdős and A. Knowles. The Altshuler–Shklovskii formulas for random band matrices II: the general case. In Annales Henri Poincaré, volume 16, pages 709–799. Springer, 2015.
- [18] L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Delocalization and diffusion profile for random band matrices. *Communications in Mathematical Physics*, 323:367–416, 2013.
- [19] L. Erdős and H.-T. Yau. A dynamical approach to random matrix theory, volume 28. American Mathematical Soc., 2017.
- [20] L. Erdős, H.-T. Yau, and J. Yin. Rigidity of eigenvalues of generalized Wigner matrices. Advances in Mathematics, 229(3):1435–1515, 2012.
- [21] A. Guionnet. Large deviations upper bounds and central limit theorems for noncommutative functionals of Gaussian large random matrices. Ann. Inst. Henri Poincaré Probab. Stat., 38(3):341–384, 2002.
- [22] I. Jana, K. Saha, and A. Soshnikov. Fluctuations of linear eigenvalue statistics of random band matrices. Theory of Probability & Its Applications, 60(3):407–443, 2016.
- [23] A. Khorunzhy, B. Khoruzhenko, and L. Pastur. On the 1/n corrections to the green functions of random matrices with independent entries. *Journal of Physics A: Mathematical* and General, 28(1):L31, 1995.
- [24] A. M. Khorunzhy, B. A. Khoruzhenko, and L. A. Pastur. Asymptotic properties of large random matrices with independent entries. *Journal of Mathematical Physics*, 37(10):5033–5060, 1996.

- [25] B. Landon, P. Lopatto, and J. Marcinek. Comparison theorem for some extremal eigenvalue statistics. *The Annals of Probability*, 48(6):2894–2919, 2020.
- [26] B. Landon and P. Sosoe. Applications of mesoscopic CLTs in random matrix theory. Ann. Appl. Probab., 30(6):2769–2795, 2020.
- [27] B. Landon and P. Sosoe. Almost-optimal bulk regularity conditions in the CLT for Wigner matrices. arXiv preprint arXiv:2204.03419, 2022.
- [28] J. O. Lee and K. Schnelli. Local law and Tracy–Widom limit for sparse random matrices. Probab. Theory Related Fields, 171(1):543–616, 2018.
- [29] L. Li and A. Soshnikov. Central limit theorem for linear statistics of eigenvalues of band random matrices. arXiv preprint arXiv:1304.6744, 2013.
- [30] Y. Li and Y. Xu. On fluctuations of global and mesoscopic linear statistics of generalized Wigner matrices. *Bernoulli*, 2(27):1057–1076, 2021.
- [31] A. Lodhia and N. Simm. Mesoscopic linear statistics of Wigner matrices. arXiv preprint arXiv:1503.03533, 2015.
- [32] A. Lytova and L. Pastur. Central limit theorem for linear eigenvalue statistics of random matrices with independent entries. Ann. Probab., 37(5):1778–1840, 2009.
- [33] M. Shcherbina. Central limit theorem for linear eigenvalue statistics of the Wigner and sample covariance random matrices. J. Math. Phys., Analysis, Geometry, (7):176–192, 2011.
- [34] M. Shcherbina. On fluctuations of eigenvalues of random band matrices. Journal of Statistical Physics, 161:73–90, 2015.