# THE POSITIVE ORTHOGONAL GRASSMANNIAN

YASSINE EL MAAZOUZ - YELENA MANDELSHTAM

The Plücker positive region  $OGr_+(k, 2k)$  of the orthogonal Grassmannian emerged as the positive geometry behind the ABJM scattering amplitudes. In this paper we initiate the study of the positive orthogonal Grassmannian  $OGr_+(k, n)$  for general values of k, n. We determine the boundary structure of the quadric  $OGr_+(1, n)$  in  $\mathbb{P}^{n-1}_+$  and show that it is a positive geometry. We show that  $OGr_+(k, 2k + 1)$  is isomorphic to  $OGr_+(k+1, 2k+2)$  and connect its combinatorial structure to matchings on [2k+2]. Finally, we show that in the case n > 2k + 1, the *positroid cells* of  $Gr_+(k, n)$  do not induce a CW cell decomposition of  $OGr_+(k, n)$ .

#### 1. Introduction

Let  $n \ge k$  be positive integers and denote by Gr(k,n) the *Grassmannian* of *k*dimensional subspaces of  $\mathbb{C}^n$ . The *positive Grassmannian*  $Gr_+(k,n)$  is the semialgebraic set in Gr(k,n) where all Plücker coordinates are real and nonnegative. Following the seminal work of Lusztig in [14–16], Postnikov introduced and studied  $Gr_+(k,n)$  in [17] from a combinatorial point of view. The matroid stratification of the Grassmannian [10] induces a natural decomposition of  $Gr_+(k,n)$ into the so-called *positroid cells*. These cells can be indexed by combinatorial objects like *Grassmann necklaces, decorated permutations, plabic graphs and Le diagrams*.

After Postnikov's landmark paper [17], the positive Grassmannian became a rich object of research in Algebraic Combinatorics [8, 18, 20]. Its study accelerated, in recent years, largely due to its unexpected and profound connection to Physics, in particular shallow water waves [1, 13] and scattering amplitudes in quantum field theory [2–4].

The object of study in the present article is the *positive orthogonal Grass*mannian  $OGr^{\omega}_{+}(k,n)$  which we define as follows.

**Definition 1.1.** Let  $\omega : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$  be a non-degenerate symmetric bilinear form, then  $OGr^{\omega}_+(k,n)$  is the semi-algebraic set of *k*-dimensional subspaces *V* of  $\mathbb{R}^n$  such that:

AMS 2010 Subject Classification: 14M15, 15B48, 05E14.

Keywords: Orthogonal Grassmannian, Flag variety, Positive geometry, Gröbner basis, Straightening laws

- *V* is *isotropic* with respect to  $\omega$  i.e.  $\omega(x, y) = 0$  for any  $x, y \in V$ ,
- the Plücker coordinates of V all have the same sign.

We denote the algebraic variety of isotropic k-dimensional subspaces of  $\mathbb{C}^n$  by  $OGr^{\omega}(k,n)$ . The semi-algebraic set  $OGr^{\omega}_+(k,n)$  is the intersection of the algebraic variety  $OGr^{\omega}(k,n)$  with the positive Grassmannian  $Gr_+(k,n)$ .

In the special case n = 2k and  $\omega(x, y) = \sum_{i=1}^{2k} (-1)^{i-1} x_i y_i$ , this object was first studied in the context of ABJM scattering amplitudes [12] and later connected to the Ising model in statistical physics [9]. In this paper we initiate the study of OGr<sup> $\omega$ </sup><sub>+</sub>(*k*,*n*) for general values of *k*,*n* with respect to the quadratic form

$$\omega_0(x,y) = x_1 y_1 - x_2 y_2 + \dots + (-1)^{n-1} x_n y_n.$$
(1)

We note that the choice of the quadratic form  $\omega$  is extremely important. For certain quadratic forms the variety  $OGr^{\omega}(k,n)$  has no real points. Moreover, even if one chooses  $\omega$  such that  $OGr^{\omega}(k,n)$  has real points, the positive region  $OGr^{\omega}_{+}(k,n)$  may not be full-dimensional and sometimes can even be empty.

**Example 1.2.** Let  $\omega(x,y) = x_1y_1 - x_2y_2 - x_3y_3 - x_4y_4$  be the Lorentzian inner product on  $\mathbb{R}^4$ . The variety OGr<sup> $\omega$ </sup>(2,4) has no real points. To see why suppose that *V* is a real point in OGr<sup> $\omega$ </sup>(2,4) and let  $x, y \in \mathbb{R}^4$  be a basis of *V*. We have  $\omega(x,y) = \omega(x,x) = \omega(y,y) = 0$ , so  $x_1, y_1 \neq 0$  and we may assume that  $x_1 = y_1 = 1$ . Then z = x - y satisfies  $z_1 = 0$  and  $\omega(z,z) = -(z_2^2 + z_3^2 + z_4^2) = 0$ . So we deduce that z = 0 and hence x = y which is a contradiction.

Let  $\omega(x,y) = x_1y_1 + x_2y_2 - x_3y_3 - x_4y_4$ . The variety OGr(2,4)<sup> $\omega$ </sup> has real points, for example the rowspan of the matrix:

$$M(\theta) = \begin{bmatrix} 1 & 0 & \cos(\theta) & -\sin(\theta) \\ 0 & 1 & \sin(\theta) & \cos(\theta) \end{bmatrix} \text{ for } \theta \in [0, 2\pi].$$
(2)

However, the semialgebraic set  $OGr^{\omega}_{+}(2,4)$  is zero-dimensional. To see why, if V is a point in  $OGr^{\omega}_{+}(2,4)$  then the first Plücker coordinate  $p_{12}$  of V does not vanish, otherwise V would contain a non-zero vector of the form (0,0,a,b) and such a vector cannot be orthogonal to itself. Now since  $p_{12} \neq 0$ , the space V is then the row span of some matrix of the form (2) and one can check that the only such matrix with nonnegative minors is the matrix  $M(\pi/2)$ .

This article is organized as follows. In Section 2 we collect some facts on the geometry of the orthogonal Grassmannian  $OGr^{\omega}(k,n)$ . In particular we exhibit a set of equations that cut out OGr(k,n) in  $\mathbb{P}(\wedge^k \mathbb{C}^n)$ , a Gröbner basis for the quadratic ideal they generate, and discuss the effect of different choices for the quadratic form  $\omega$ . In Section 3 we investigate  $OGr^{\omega_0}_+(1,n)$  with respect to the alternating form (1), and show that it is a positive geometry. Section 4 is devoted to  $OGr^{\omega_0}_+(k, 2k+1)$ . In Section 5 we initiate the study of  $OGr^{\omega_0}_+(k, n)$  starting with the case k = 2. Already in this specific case, we show that the positroid decomposition is no longer sufficient to induce a cell decomposition of  $OGr^{\omega_0}_+(2, n)$  and pose questions for future work.

## **2.** Commutative algebra and geometry of OGr(k, n)

In this section we collect some facts on the algebraic variety  $OGr^{\omega}(k,n)$  over  $\mathbb{C}$ . Since all non-degenerate symmetric bilinear forms over  $\mathbb{C}$  are isomorphic to

$$(x,y) := x_1y_1 + \dots + x_ny_n,$$

up to a linear change of variables, the varieties  $OGr^{\omega}(k,n)$  for different  $\omega$  are isomorphic. So, in this section we may assume that  $\omega$  is the standard inner product  $(\cdot, \cdot)$ , and we suppress  $\omega$  and write OGr(k,n). We recall that the Grassmannian Gr(k,n) can be embedded in Plücker space  $\mathbb{P}(\wedge^k \mathbb{C}^n)$ . The  $\binom{n}{k}$  coordinates of this projective space are called Plücker coordinates which we denote by  $p_I$  for any subset  $I = \{i_1 < i_2 < \cdots < i_k\}$  of [n]. In Physics, the Plücker coordinates are often denoted by  $p_I = \langle i_1 i_2 \dots i_k \rangle$ .

**Theorem 2.1.** In addition to the Plücker relations, the orthogonal Grassmannian OGr(k,n) is cut out in  $\mathbb{P}(\wedge^k \mathbb{C}^n)$  by the following  $\frac{1}{2}\binom{n}{k-1}\binom{n}{\binom{n}{k-1}+1}$  equations:

$$\sum_{\ell=1}^{n} \varepsilon(I\ell) \varepsilon(J\ell) \ p_{I\ell} p_{J\ell} = 0, \quad \text{for ordered sets } I, J \in \binom{[n]}{k-1}.$$
(3)

where  $\varepsilon(I\ell)$  denotes the sign of the permutation that sorts the string  $I\ell$ .

*Proof.* Let  $p = (p_I)_{I \in \binom{[n]}{k}}$  be a point in OGr(k, n) and let V be its corresponding isotropic space in  $\mathbb{C}^n$ . The space V is the rowspan of a rank k matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & \dots & a_{2n} \\ \vdots & \vdots & \dots & \ddots & \vdots \\ a_{k1} & a_{k2} & \dots & \dots & a_{kn} \end{bmatrix}$$

For  $I = \{i_1 < \cdots < i_k\}$ , a Plücker coordinate  $p_I$  can then be written as follows:

$$p_I = \sum_{\sigma \in S_k} \varepsilon(\sigma) \prod_{s=1}^k a_{\ell j_{\sigma(s)}},$$

where  $S_k$  is the symmetric group on [k]. Now let  $I = (i_1 < \cdots < i_{k-1})$  and  $J = (j_1 < \cdots < j_{k-1})$  be ordered subsets of [n], then:

$$\sum_{\ell=1}^{n} \varepsilon(I\ell) \varepsilon(J\ell) p_{I\ell} p_{J\ell} = \sum_{\ell=1}^{n} \sum_{\sigma, \tau \in S_k} \operatorname{sign}(\sigma\tau) \prod_{s=1}^{k-1} a_{\sigma(s)i_s} a_{\tau(s)j_s} a_{\sigma(k)\ell} a_{\tau(k)\ell}$$
$$= \sum_{\sigma, \tau \in S_k} \operatorname{sign}(\sigma\tau) \prod_{s=1}^{k-1} a_{\sigma(s)i_s} a_{\tau(s)j_s} \underbrace{\left(\sum_{\ell=1}^{n} a_{\sigma(k)\ell} a_{\tau(k)\ell}\right)}_{=0}.$$
(4)

Since V is isotropic, the last sum in the right-hand-side is 0 so we deduce that:

$$\sum_{\ell=1}^n \varepsilon(I\ell)\varepsilon(J\ell)p_{I\ell}p_{J\ell}=0.$$

Conversely, let *p* be a point in Gr(k,n) such that:

$$\sum_{\ell=1}^{n} \varepsilon(I\ell) \varepsilon(J\ell) p_{I\ell} p_{J\ell} = 0 \quad \text{for all } I, J \subset \binom{[n]}{k}.$$

Since the quadratic form  $(\cdot, \cdot)$  is invariant under the action of the symmetric group  $S_n$ , we can use the action of this group on Gr(k, n) and, without loss of generality, assume that  $p_{1,2,...,k} \neq 0$ . So we can write the vector space in  $\mathbb{C}^n$  represented by p as the rowspan of the matrix:

$$A = \begin{bmatrix} p_{1,2,\dots,k} & 0 & \dots & 0 & (-1)^{k-1} p_{2,3,\dots,k,k+1} & (-1)^{k-1} p_{2,3,\dots,k,k+2} & \dots & (-1)^{k-1} p_{2,3,\dots,k,n} \\ 0 & \ddots & \dots & \vdots & (-1)^{k-2} p_{1,3,\dots,k,k+1} & (-1)^{k-2} p_{1,3,\dots,k,k+2} & \dots & (-1)^{k-2} p_{1,3,\dots,k,n} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \\ 0 & \dots & 0 & p_{1,2,\dots,k} & p_{1,2,\dots,k-1,k+1} & p_{1,2,\dots,k-1,k+2} & \dots & p_{1,2,\dots,k-1,n} \end{bmatrix}.$$

From the equations (3), we can see that the rows of this matrix are orthogonal to themselves and to one another so  $p \in OGr(k, n)$  and this finishes the proof.  $\Box$ 

**Example 2.2.** The orthogonal Grassmannian OGr(2,5) is cut out by a system of 5 Plücker relations and 15 relations of the form (3):

$$p_{12}^2 + p_{13}^2 + p_{14}^2 + p_{15}^2, \ p_{12}^2 + p_{23}^2 + p_{24}^2 + p_{25}^2, \ \dots, \ p_{15}^2 + p_{25}^2 + p_{35}^2 + p_{45}^2,$$
  
$$p_{13}p_{23} + p_{14}p_{24} + p_{15}p_{25}, \ -p_{12}p_{23} + p_{14}p_{34} + p_{15}p_{35}, \ \dots, \ p_{14}p_{15} + p_{24}p_{25} + p_{34}p_{35}.$$

The ideal generated by these relations is the prime ideal of OGr(2,5).

**Remark 2.3.** In Physics notation the equations (3) become

$$\sum_{\ell=1}^n \langle i_1 i_2 \dots i_{k-1} \ell \rangle \langle j_1 j_2 \dots j_{k-1} \ell \rangle = 0$$

This is because  $\varepsilon(I\ell)p_{I\ell} = \langle i_1 i_2 \dots i_{k-1}\ell \rangle$ . The relations (3) can also be obtained from the so-called *co-circuit matrices* in [7, Section 3]. We denote by *P* the  $\binom{n}{k-1} \times n$  matrix whose rows are indexed by size k-1 subsets *I* in [*n*] and its entries are given by:

$$P_{I\ell} = \begin{cases} \varepsilon(I\ell) \ p_{I\ell}, & \text{if } \ell \notin I \\ 0, & \text{otherwise } \ell \in I \end{cases}, \text{ for } I \in {[n] \choose k-1} \text{ and } 1 \le \ell \le n.$$

The equations (3) are equivalent to  $PP^T = 0$ . For a different quadratic form  $\omega$  whose matrix in the standard basis of  $\mathbb{C}^n$  is  $\Omega$ , the equations for  $OGr^{\omega}(k,n)$  are obtained by setting  $P\Omega P^T = 0$  instead of (3). For example for the alternating bilinear form (1), the equations (3) become:

$$(P\Omega P^T)_{IJ} = \sum_{\ell=1}^n (-1)^{\ell-1} \varepsilon(I\ell) \varepsilon(J\ell) p_{I\ell} p_{J\ell} = 0, \quad \text{for} \quad I, J \in \binom{[n]}{k-1}.$$
(5)

**Proposition 2.4.** The variety OGr(k, n) is empty if n < 2k. When n = 2k it splits into two irreducible connected components, and it is irreducible when n > 2k. Moreover we have:

dim(OGr(k,n)) = 
$$k(n-k) - {\binom{k+1}{2}}$$
 for  $n \ge 2k$ .

*Proof.* The proof for the first part of the statement can be found in [11, Proposition on page 735]. For the dimension count it is more convenient to work with the quadratic form:

$$\omega(x,y) = x_1y_n + x_2y_{n-1} + \dots + x_ny_1.$$

Let  $\Omega$  be its corresponding matrix,  $X = (x_{i,j})$  be a  $k \times (n-k)$  matrix of indeterminates, and consider the  $k \times n$  matrix  $A = [Id_k|X] = 0$ . The equations  $A\Omega A^T = 0$  are equivalent to an expression of each entry in the lower right corner of X:

X =	$x_{1,1}$	•••			$x_{1,n-k-1}$	$\boldsymbol{x}_{1,n-k}$	
	$x_{2,1}$	•••		•••	$x_{2,n-k-1}$	$x_{2,n-k}$	
	:			. · ·		÷	,
	$x_{k,1}$		$\boldsymbol{x}_{k,n-2k+1}$			$\boldsymbol{x}_{k,n-k}$	

in terms of all the other entries of the matrix *A* outside that corner. So we deduce that  $\dim(OGr(k,n)) = \dim(OGr^{\omega}(k,n)) = k(n-k) - \binom{k+1}{2}$ .

From now on, we fix positive integers k, n such that  $n \ge 2k$ . Following [7], let  $Y_{k,n}$  denote *Young's lattice*. This is a poset whose elements are subsets of size k in [n] and the order relation in  $Y_{k,n}$  is:

$$\langle i_1 < \dots < i_k \rangle \le \langle j_1 < \dots < j_k \rangle$$
 if  $i_1 \le j_1, i_2 \le j_2, \dots, i_{k-1} \le j_{k-1}$  and  $i_k \le j_k$ .

We denote by  $\widetilde{Y}_{k,n}$  another copy of Young's lattice with elements  $[i'_1 < \cdots < i'_{n-k}]$  and order relation:

$$[i'_1 < \dots < i'_{n-k}] \le [j'_1 < \dots < j'_{n-k}]$$
 if  $i'_1 \ge j'_1, \dots, i'_{n-k} \ge j'_{n-k}$ .

Finally we denote by  $\mathcal{P}_{k,n}$  the poset which, as a set, is the disjoint union of  $Y_{k,n}$  and  $\widetilde{Y}_{k,n}$ . All order relations in  $Y_{k,n}$  and  $\widetilde{Y}_{k,n}$  remain order relations in  $\mathcal{P}_{k,n}$  and in addition to these relations we have  $\binom{2k}{k}$  covering relations:

$$[j'_1 < \cdots < j'_{n-k}] < \langle i_1 < \cdots < i_k \rangle$$

whenever  $\{1, 2, 3, \dots, 2k\} = \{i_1, \dots, i_k\} \sqcup \{j_1, \dots, j_k\}$  is a partition where the set  $\{j_1, \dots, j_k\}$  is the complement  $[n] \setminus \{j'_1, \dots, j'_{n-k}\}$ .




Figure 1: The poset  $\mathcal{P}_{2,6}$  is created from  $Y_{2,6}$  and  $\widetilde{Y}_{2,6}$  by adding the six covering relations in red.

An incomparable pair  $(\langle i_1, \ldots, i_k \rangle, \langle j_1, \ldots, j_k \rangle)$  or  $(\langle i_1, \ldots, i_k \rangle, [j'_1, \ldots, j'_{n-k}])$ in the poset  $\mathcal{P}_{k,n}$  yields a non-semistandard Young tableau  $\mu$  of shape (k,k) or  $\lambda$  of shape (n-k,k):

$$\mu = \begin{bmatrix} j_1 & \cdots & j_{\ell-1} & \mathbf{j}_{\ell} & \mathbf{j}_{\ell+1} & \cdots & \mathbf{j}_k \\ \mathbf{i}_1 & \cdots & \mathbf{i}_{\ell-1} & \mathbf{i}_{\ell} & \mathbf{i}_{\ell+1} & \cdots & \mathbf{i}_k \end{bmatrix},$$

$$\lambda = \begin{bmatrix} j'_1 & \cdots & j'_{\ell-1} & \mathbf{j}'_{\ell} & \mathbf{j}'_{\ell+1} & \cdots & \mathbf{j}'_k & \cdots & \mathbf{j}'_{n-k} \\ \mathbf{i}_1 & \cdots & \mathbf{i}_{\ell-1} & \mathbf{i}_{\ell} & \mathbf{i}_{\ell+1} & \cdots & \mathbf{i}_k \end{bmatrix}.$$

$$(6)$$

The tableau  $\mu$  or  $\lambda$  being non-semistandard means that there exists an index  $\ell$  in [k] such that:

$$i_1 < \dots < i_\ell < j_\ell < \dots < j_k$$
 or  $i_1 < \dots < i_\ell < j'_\ell < \dots < j'_{n-k}$ . (7)

We pick  $\ell$  to be the smallest index with this property. The strictly increasing sequences of integers in (7) are highlighted in bold in (6). Now for the tableau  $\mu$  of type (k,k), consider the  $\pi$  permutations of the sequence  $i_1 < \cdots < i_\ell < j_\ell < \cdots < j_k$  which make the first  $\ell$  entries and the last  $k - \ell + 1$  entries separately increasing. Summing over these permutations, each such tableaux  $\mu$  of type (k,k) yields a quadric

$$f_{\mu} := \sum_{\pi} \operatorname{sign}(\pi) \langle \pi(i_1), \dots, \pi(i_{\ell}), i_{\ell+1}, \dots i_k \rangle \langle j_1, \dots, j_{\ell-1}, \pi(j_{\ell}), \dots, \pi(j_k) \rangle.$$
(8)

Similarly, for the tableau  $\lambda$  of type (n - k, k), summing over the permutations  $\pi$  of the sequence  $i_1 < \cdots < i_{\ell} < j'_{\ell} < \cdots < j'_{n-k}$  which make the first  $\ell$  entries and the last  $n - k - \ell + 1$  entries separately increasing, we get a quadric

$$f_{\lambda} := \sum_{\pi} \operatorname{sign}(\pi) \ \langle \pi(i_1), \dots, \pi(i_{\ell}), i_{\ell+1}, \dots i_k \rangle \ [j'_1, \dots, j'_{\ell-1}, \pi(j'_{\ell}), \dots, \pi(j'_k)].$$
(9)

Here, whenever  $J' = \{j'_1 < \cdots < j'_{n-k}\}$  and  $[n] \setminus J' = \{\overline{j}_1 < \cdots < \overline{j}_k\}$  we set

$$[j'_1, \dots, j'_{n-k}] := (-1)^{\sum_{r=1}^{n-k} j'_r} \langle \bar{j}_1, \dots, \bar{j}_k \rangle.$$

These quadrics are called *straightening laws* in standard monomial theory. We note that, under any reverse lexicographic term order  $\prec$  in  $\mathbb{C}[p_I]$  given by a linear extension of the poset  $Y_{k,n}$ , the leading monomial of  $f_{\mu}$  in (8) is the binomial  $p_{i_1...i_k} p_{j_1...j_k}$  and the leading monomial of  $f_{\lambda}$  in (9) is the binomial  $p_{i_1...i_k} p_{\bar{j}_1...i_k} p_{\bar{j}_1...i_k}$ .

**Example 2.5.** The pair  $\langle 12 \rangle$  and [1356] are incomparable in  $\mathcal{P}_{2,6}$ , see Figure 1. This pair corresponds to the following non-semistandard Young tableau:



There are 15 permutations  $\pi$  that make the two first entries and the remaining 3 entries of  $\{1,2,3,5,6\}$  separately increasing. Among these 15 permutations, only 4 leave 1 in the bottom row of  $\lambda$ . The quadric  $f_{\lambda}$  in this case has the following expression:

$$f_{\lambda} = -\underline{p_{12}p_{24}} - p_{13}p_{24} + p_{15}p_{45} + p_{16}p_{46}.$$

**Theorem 2.6.** The quadrics in (8) and (9) form a Gröbner basis for the ideal  $I_{k,n}$  in  $\mathbb{C}[p_I]$  generated by the Plücker relations and the quadratic equations in (3) with respect to any monomial ordering given by a linear extension of the poset  $\mathcal{P}_{k,n}$ .

*Proof.* The ideal  $I_{k,n}$  is the image of the ideal  $I_{k,n,0}$  of the spinor-helicity variety SH(k, n, 0) in [7] under the map:

$$\boldsymbol{\varphi}: \mathbb{C}[p_I, q_{J'}] \mapsto \mathbb{C}[p_I], \quad p_I \mapsto p_I, \ q_{J'} \mapsto (-1)^{\sum_{j' \in J'} j'} p_{[n] \setminus J'}.$$

The quadrics (8) and (9) are the images of a Gröbner basis of the ideal  $I_{k,n,0}$ [7, Theorem 2.7] under the map  $\varphi$ . So we deduce that  $f_{\mu}$  and  $f_{\lambda}$  in (8) and (9) are members of the ideal  $I_{k,n}$  for any non-semistandard Young tableaux  $\mu$  of type (k,k) and  $\lambda$  of type (n-k,k) and the initial monomials  $p_{i_1...i_k} p_{j_1...j_k}$  and  $p_{i_1...i_k} p_{\overline{j_1}...\overline{j_k}}$  are in the initial ideal in $\prec$  (I<sub>k,n</sub>). Now let  $g \in I_{k,n}$  and write  $g = \varphi(h)$ for some  $h \in I_{k,n,0}$ . After scaling h with a suitable nonzero scalar in  $\mathbb{C}$ , we can write:

$$h = \underline{\boldsymbol{p}^{\boldsymbol{a}}\boldsymbol{q}^{\boldsymbol{b}}} + \sum_{\ell=1}^{N} c_{\ell} \boldsymbol{p}^{\boldsymbol{a}_{\ell}} \boldsymbol{q}^{\boldsymbol{b}_{\ell}},$$

where  $\boldsymbol{a}, \boldsymbol{a}_{\ell}$  are exponent vectors for the variables  $(p_I, I \in {\binom{[n]}{k}})$  and  $\boldsymbol{b}, \boldsymbol{b}_{\ell}$  are exponent vectors for the variables  $(q_{J'}, J' \in {\binom{[n]}{k}})$ . The underlined monomial in h is the leading monomial with respect to  $\prec$ , so it is divisible by some binomial  $p_I q_{J'}$  where I', J' is an incomparable pair in  $\mathcal{P}_{k,n}$ . We claim that the leading term of  $g = \varphi(h)$  is  $\varphi(\boldsymbol{p}^a \boldsymbol{q}^b)$ . To see why let  $1 \leq \ell \leq N$ , we then have  $\boldsymbol{p}^{a_\ell} \boldsymbol{q}^{b_\ell} \prec \boldsymbol{p}^a \boldsymbol{q}^b$  and after applying the map  $\varphi$  this inequality continues to hold.

Hence  $in_{\prec}(I_{k,n})$  is generated by the binomials  $p_I p_J$  where  $I, J' = [n] \setminus J$  is an incomparable pair in  $\mathcal{P}_{k,n}$ . These are exactly the initial monomials of the quadrics (8) and (9).

Our next result gives a formula for the degree of OGr(k, n) for n > 2k.

**Proposition 2.7.** Let n > 2k,  $m := \lfloor n/2 \rfloor$ , and set  $D := k(n-k) - \binom{k+1}{2}$ . The degree of OGr(k, n) in the Plücker embedding is

$$D! \cdot \left(\prod_{\substack{1 \le i \le k \\ k < j \le m}} \frac{1}{(2m - i - j)(j - i)}\right) \left(\prod_{1 \le i < j \le k} \frac{2}{2m - i - j}\right), \qquad \text{if } n = 2m,$$

$$D! \cdot \left(\prod_{j \le m} \frac{2}{(2m - i - j)}\right) \left(\prod_{j \le m} \frac{1}{(2m - i - j)}\right) \left(\prod_{j \le m} \frac{2}{(2m - i - j)}\right) \qquad \text{if } n = 2m + 1$$

$$(10)$$

$$D! \cdot \left(\prod_{1 \le i \le k} \frac{2}{2m - 2i + 1}\right) \left(\prod_{\substack{1 \le i \le k \\ k < j \le m}} \frac{1}{(2m - i - j)(j - i)}\right) \left(\prod_{1 \le i < j \le k} \frac{2}{2m - i - j + 1}\right), \quad \text{if } n = 2m + 1.$$

*Proof.* Fix n > 2k and recall that OGr(k,n) is irreducible in this case. Let R be homogeneous coordinate ring of OGr(k,n) endowed with its natural  $\mathbb{Z}$ -grading. By the Borel-Weil-Bott theorem, each degree  $\ell$  piece  $R_{\ell}$  of R is an irreducible representation  $V_{\ell\lambda}$  of SO(n) corresponding to the highest weight vector  $\ell\lambda$  where  $\lambda := e_1 + ... + e_k \in \mathbb{R}^m$  and  $m := \lfloor n/2 \rfloor$ . Furthermore, we can compute the dimensions of these representations using the Weyl dimension formula as follows. Recall that the positive roots of SO(n) are

$$\Phi^{+} = \begin{cases} \{e_{i} \pm e_{j}\}_{1 \le i < j \le m}, & \text{if } n = 2m\\ \{e_{i} \pm e_{j}\}_{1 \le i < j \le m} \cup \{e_{i}\}_{0 < i \le m}, & \text{if } n = 2m+1. \end{cases}$$

The Weyl dimension formula states that for any integer  $\ell \ge 0$  we have

$$\dim V_{\ell\lambda} = \frac{\prod_{\alpha \in \Phi^+} \langle \rho + \ell\lambda, \alpha \rangle}{\prod_{\alpha \in \Phi^+} \langle \rho, \alpha \rangle} = \prod_{\alpha \in \Phi^+} \left( 1 + \ell \frac{\langle \lambda, \alpha \rangle}{\langle \rho, \alpha \rangle} \right), \tag{11}$$

where  $2\rho$  be the sum of the positive roots  $\Phi^+$ , and  $\langle \cdot, \cdot \rangle$  is the standard inner product on the root space  $\mathbb{R}^m$ . This is equal to

$$2\rho = \begin{cases} (2m-2)e_1 + (2m-4)e_2 + \dots + e_{m-1} & n = 2m \\ (2m-1)e_1 + (2m-3)e_2 + \dots + e_m & n = 2m+1. \end{cases}$$

Plugging this into the dimension formula above for n = 2m yields

$$\dim V_{\ell\lambda} = \prod_{\substack{1 \le i \le k \\ k < j \le m}} \left( \left( 1 + \frac{1}{2m - i - j} \ell \right) \left( 1 + \frac{1}{j - i} \ell \right) \right) \prod_{1 \le i < j \le k} \left( 1 + \frac{2}{2m - i - j} \ell \right).$$
(12)

When n = 2m + 1 we obtain

$$\dim V_{\ell\lambda} = \prod_{1 \le i \le k} \left( 1 + \frac{2}{2m - 2i + 1} \ell \right) \prod_{\substack{1 \le i \le k \\ k < j \le m}} \left( \left( 1 + \frac{1}{2m - i - j + 1} \ell \right) \left( 1 + \frac{1}{j - i} \ell \right) \right) \cdot \prod_{1 \le i < j \le k} \left( 1 + \frac{2}{2m - i - j + 1} \ell \right).$$

This determines the Hilbert polynomial  $H(\ell) := \dim(V_{\ell\lambda})$  of the coordinate ring R, which is a polynomial in  $\ell$  of degree D. The degree of OGr(k, n) is then the leading coefficient of the polynomial D! H(X).

Next we prove that  $I_{k,n}$  is a prime ideal for n > 2k.

**Lemma 2.1.** The incomparable pairs (I,J') in the poset  $P_{k,n}$  with  $I \in {\binom{[n]}{k}}$  and  $J' \in {\binom{[n]}{n-k}}$  such that the pair  $(I,[n] \setminus J')$  is comparable in  $Y_{k,n}$  are in bijection with semistandard Young Tableaux of shape (k-1,k-1) and fillings in [n]. Their number is

$$\frac{1}{k-1}\binom{n+1}{k}\binom{n}{k-2}.$$

*Proof.* Note that the incomparable pairs (I, J') in question here are exactly the non-semistandard Young tableaux  $\lambda$  of shape (n - k, k) with fillings in [n] such that the Young tableau  $\lambda^c$  of shape (k, k) obtained by taking the complement of the first row of  $\lambda$  in [n] is semistandard. To show the result, it then suffices to exhibit a bijection between the following two sets

$$\left\{(S_1,S_2):S_1,S_2\in\binom{[n]}{k-1},S_1\leq S_2\right\}$$

and

$$\left\{ (T_1, T_2) : T_1, T_2 \in \binom{[n]}{k}, \ T_1 \leq T_2 \text{ and } T_1^c \not\leq T_2 \right\}.$$

Here if  $A = \{a_1 < ..., a_k\}, B = \{b_1 < ..., b_k\} \in {[n] \choose k}$ , by  $A \le B$  we mean  $a_1 \le b_1, a_2 \le b_2, ..., a_k \le b_k$ . Similarly, if  $C := \{c_1 < ... < c_{n-k}\}$  we write  $C \le B$  if  $c_1 \le b_1, ..., c_k \le b_k$ .

Before we present the bijection, the following notation and observations will be helpful. For any  $h \in [n]$  and set  $S \subseteq [n]$  we denote by  $S^h$  the set  $S \cap [h]$  of elements in *S* that are at most *h*. Now it is not hard to verify that  $S \leq T$  in Young's lattice if and only if  $S^h \geq T^h$  for all  $h \in [n]$ .

The bijection is as follows. Fix  $S_1 \leq S_2 \in {\binom{[n]}{k-1}}$  and let  $L = S_1 \cap S_2$  and  $R = S_1^c \cap S_2^c$ . Let *h* be minimal such that  $|L^h| < |R^h|$ . Note that *h* is guaranteed to exist since  $|R^n| = n - 2(k-1) + |L^n| > |L^n|$ , and further note that we have  $|L^h| + 1 = |R^h|$ . Then we define

$$T_1 = S_1 \setminus L^h \cup R^h, \quad T_2 = S_2 \setminus L^h \cup R^h.$$

It is easy to see that  $T_1, T_2 \in {\binom{[n]}{k}}$  with  $T_1 \leq T_2$ . Now we show that  $T_1^c \leq T_2$ . Consider the set  $(T_1^c)^h = \{t \in T_1 : t \leq h\} = S_2^h$ , while  $T_2^h = S_2^h \setminus L^h \cup R^h$ . Thus  $|(T_1^c)^h| = |S_2^h|$  but  $|T_2^h| = |S_2^h| - |L^h| + |R^h| = |S_2^h| + 1 > |(T_1^h)^c|$ , and therefore  $T_1^c \leq S_2$ .

The inverse map, starting from  $T_1 \leq T_2 \in \binom{n}{k}$  with  $T_1^c \leq T_2$  is given by

$$S_1 = T_1 \setminus \mathbb{R}^h \cup \mathbb{L}^h, \quad S_2 = T_2 \setminus \mathbb{R}^h \cup \mathbb{L}^h,$$

where  $L = T_1 \cap T_2$ ,  $R = T_1^c \cap T_2^c$ , and *h* is minimal such that  $|L^h| > |R^h|$ . We show that such a *h* exists. Observe that  $T_1^c \not\leq T_2$  implies that there is some *h* for which  $|(T_1^c)^h| < |T_2^h|$ , while  $T_1 \leq T_2$  implies  $|T_2^h| \leq |T_1^h|$ . We have

$$|T_2^h| = |T_2^h \cap T_1^h| + |T_2^h \cap (T_1^c)^h|$$
 and  $|(T_1^c)^h| = |(T_1^c)^h \cap T_2^h| + |(T_1^c)^h \cap (T_2^c)^h|.$ 

Together these give  $|L^h| = |T_1^h \cap T_2^h| > |(T_1^c)^h \cap (T_2^c)^h| = |R^h|$ , as desired. The defined map yields  $S_1 \le S_2 \in {\binom{[n]}{k-1}}$ , so our provided map is a bijection.

**Remark 2.8.** We remark that the proof of Lemma 2.1 may be more intuitive to the reader if translated into the non-crossing lattice path formulation of semistandard Young tableaux, with a row of a Young tableau defining a path by its left steps [19, Chapter 7]. Under this formulation, taking the complement of a row of a Young tableau corresponds to reflecting the corresponding path. In the proof, the sets L and R correspond respectively to shared left and right steps of the two paths, and the superscript h denotes a cutoff at height h. The bijective map reflects the picture of the two lattice paths up to the point at which the number of shared right steps exceeds shared left steps by one (see Figure 2).




Figure 2: A lattice path depiction of the bijection in Lemma 2.1. The red path crosses the reflection of the blue path, pictured as a dotted line.

**Theorem 2.9.** When n > 2k, the ideal  $I_{k,n}$  in  $\mathbb{C}[p_I]$  generated by the Plücker relations and the quadratic equations in (3) is the prime ideal of OGr(k,n). In particular, the degree of  $I_{k,n}$  is given by (10).

*Proof.* By Kostant's theorem, the prime ideal of OGr(k,n) in  $\mathbb{C}[p_I]$  is generated in degree 2. To show that  $I_{k,n}$  is exactly the prime ideal of OGr(k,n) it is then enough to prove that the dimension of the degree 2 piece of the  $\mathbb{Z}$ -graded ring  $\mathbb{C}[p_I]/I_{k,n}$  is equal to dim $V_{2\lambda}$  from the proof of Proposition 2.7. To do so we need to count the dimension of the degree 2 part of  $I_{k,n}$ .

This dimension is equal to the sum of the number of incomparable pairs in the poset  $Y_{k,n}$  together with the mixed incomparable pairs in Lemma 2.1. So the dimension of the degree 2 part of  $\mathbb{C}[p_I]/I_{k,n}$  is

$$\frac{1}{k}\binom{n+1}{k+1}\binom{n}{k-1} - \frac{1}{k-1}\binom{n+1}{k}\binom{n}{k-2}.$$
(13)

Using (12), we compute dim( $V_{2\lambda}$ ) and we find that it is equal to (13).

We now shift our attention to the special case n = 2k and consider quadratic forms  $\omega$  of the following shape for some subset  $S \subset [n]$ :

$$\boldsymbol{\omega}(x,y) = \sum_{i \notin S} x_i y_i - \sum_{i \in S} x_i y_i.$$

**Proposition 2.10.** The ratio of any two complementary Plücker coordinates in the function field  $\mathbb{C}(\operatorname{OGr}(k, 2k))$  is either  $\pm 1$  if k + |S| is even or  $\pm \sqrt{-1}$  otherwise. Moreover, for any point in  $\operatorname{OGr}(k, 2k)$ , the ratio of one pair of complementary Plücker coordinates uniquely determines the ratio of all the other pairs.

*Proof.* Suppose  $\tilde{I}, \tilde{J} \in {[2k] \choose k}$  are disjoint, and consider elements  $i \in \tilde{I}$  and  $j \in \tilde{J}$ . By setting  $I = \tilde{I} \setminus \{i\}$  and  $J = \tilde{J} \setminus \{j\}$  in (3), we obtain the following relation:

$$p_{\tilde{I}}p_{Ji}\pm p_{Ij}p_{\tilde{J}}=0,$$

where the sign is determined by the form and I, J, i, j. This relation tells us that

$$\frac{p_{\tilde{I}}}{p_{\tilde{J}}} = \pm \frac{p_{Ij}}{p_{Ji}}.$$

Thus, for complementary sets  $\tilde{I}, \tilde{J}$  the ratio of their Plücker coordinates is equal, up to sign, to the ratio of the Plücker coordinates of the pair of complementary sets obtained by exchanging one element of  $\tilde{I}$  with one element of  $\tilde{J}$ . By interchanging all elements of  $\tilde{I}$  with those of  $\tilde{J}$  we obtain an equation of the form:

$$\frac{p_{\tilde{I}}}{p_{\tilde{J}}} = \pm \frac{p_{\tilde{J}}}{p_{\tilde{I}}}.$$

Note that the sign in the equation in not ambiguous. If the sign is positive, then the ratio is  $\pm 1$ , while if it is negative, the ratio is  $\pm \sqrt{-1}$ . Moreover, note that any pair of complementary Plücker coordinates can be obtained by exchanging elements of  $\tilde{I}$  and  $\tilde{J}$ . Therefore, once the ratio  $p_{\tilde{I}}/p_{\tilde{J}}$  for some complementary I, J is known, all the other complementary ratios are uniquely determined.

We now show that whether the complementary ratios are  $\pm 1$  or  $\pm \sqrt{-1}$  is determined by the parity of k + |S|. As noted above, since any pair of complementary sets can be obtained by exchanging elements from a different pair, it is sufficient to consider  $p_{\bar{I}}/p_{\bar{J}}$  for one complementary pair  $\tilde{I}, \tilde{J}$ . We fix  $\tilde{I} =$  $\{1, 3, ..., 2k - 1\}$  and  $\tilde{J} = \{2, 4, ..., 2k\}$ . We then proceed by iteratively exchanging the smallest odd element of  $\tilde{I}$  with the smallest even element of  $\tilde{J}$ . Observe that, as a result of our choice of sets and exchanged elements, we have  $\varepsilon(I'\ell)\varepsilon(J'\ell) = 1$  for any I', J' and  $\ell$  obtained as a result of partially completing the exchange process so  $(-1)^{1(\ell_1 \in S)} p_{I'\ell_1} p_{J'\ell_1} + (-1)^{1(\ell_2 \in S)} p_{I'\ell_2} p_{J'\ell_2} = 0$  at each step. Putting all steps together to swap I and J, we find:

$$(-1)^{|S|+k} p_{1,3,\dots,2k-1} p_{1,3,\dots,2k-1} = p_{2,4,\dots,2k} p_{2,4,\dots,2k}.$$
  
So we deduce that  $(p_{1,3,\dots,2k-1}/p_{2,4,\dots,2k})^2 = (-1)^{|S|+k}.$ 

By virtue of of Proposition 2.10, for any point  $p \in OGr^{\omega_0}(k, 2k)$  we have

$$p_I = p_{I^c}$$
 for any  $I \in {[2k] \choose k}$  or  $p_I = -p_{I^c}$  for any  $I \in {[2k] \choose k}$ . (14)

**Corollary 2.11.** The ideal  $I_{k,2k}$  is not prime.

*Proof.* The ideal  $I_{k,2k}$  is generated by quadrics. Equation 14 exhibits products of linear polynomials that vanish on OGr<sup> $\omega_0$ </sup>(k, 2k). So  $I_{k,2k}$  is not a prime ideal.  $\Box$ 

We have then shown that  $OGr^{\omega_0}(k, 2k)$  has two connected components. In this article, we define the *standard* component of  $OGr^{\omega_0}(k, 2k)$  to be the connected component where  $p_I = p_{I^c}$  for all  $I \in \binom{[2k]}{k}$ . We will denote the semi-algebraic set in the standard component where all Plücker coordinates are real and have the same sign by  $OGr^{\omega_0}_+(k, 2k)$ .

## **3.** The positive orthogonal Grassmannian $OGr_+(1,n)$

In this section we switch gears to study the positive geometry of  $OGr(1,n) = OGr^{\omega_0}(1,n)$  where  $\omega_0$  is the sign alternating quadratic form (1). For the remainder of this article, unless specifically mentioned, we always work with this form. We denote by (p,q) the signature of  $\omega_0$  where  $p = \lceil \frac{n}{2} \rceil$  and  $q = \lfloor \frac{n}{2} \rfloor$ .

We think of the elements of [n] as vertices of a regular *n*-gon ordered clockwise from 1 to *n*. For each pair of non-empty subsets  $A \subset [n] \cap (2\mathbb{Z} + 1)$  and  $B \subset [n] \cap 2\mathbb{Z}$ , there exists a unique cycle  $\sigma(A, B)$  in the symmetric group  $S_n$  such that  $\sigma(A, B)$  has exactly one excedance and the support of  $\sigma(A, B)$  is  $A \sqcup B$ . The set of such permutations<sup>1</sup>  $\sigma(A, B)$  is denoted  $\mathfrak{S}_{1,n}$ . The set  $\mathfrak{S}_{1,n}$  is endowed with a partial order given by:

 $\sigma(C,D) \preceq \sigma(A,B) \iff C \subseteq A \text{ and } D \subseteq B.$ 

For  $\sigma(A,B) \in \mathfrak{S}_{1,n}$ , we denote by  $\Pi_{\sigma}$  the subset of  $\mathbb{P}^{n-1}_+$  where  $x_i = 0$  if and only if *i* is a fixed point of  $\sigma(A,B)$  i.e.  $i \notin A \sqcup B$ .

**Theorem 3.1.** The positive orthogonal Grassmannian  $OGr_+(1,n)$  is combinatorially isomorphic to the product of simplices  $\Delta_{p-1} \times \Delta_{q-1}$ . More precisely, the following holds:

1. 
$$\operatorname{OGr}_+(1,n) = \bigsqcup_{\sigma \in \mathfrak{S}_{1,n}} \operatorname{OGr}_+(1,n) \cap \Pi_{\sigma}.$$

2.  $\overline{\mathrm{OGr}_+(1,n)\cap\Pi_{\sigma}} = \bigsqcup_{\tau\preceq\sigma}\mathrm{OGr}_+(1,n)\cap\Pi_{\tau}.$ 

<sup>&</sup>lt;sup>1</sup>These are decorated permutations with all fixed points having a "+" decoration.

3. If  $A = \{i_1 < \cdots < i_r\}$  and  $B = \{j_1 < \cdots < j_m\}$  and  $\sigma = \sigma(A, B)$  the cell OGr<sub>+</sub>(1,*n*)  $\cap \prod_{\sigma(A,B)}$  can be parameterized as follows. For each  $t_1, \ldots, t_{r-1}$  and  $s_1, \ldots, s_{m-1}$  in  $\mathbb{R}_{>0}$  we get a point  $x \in OGr_+(1,n) \cap \prod_{\sigma(A,B)}$  by setting  $x_i = 0$  whenever *i* is a fixed point of  $\sigma(A, B)$  and:

$$x_{i_{1}} = \frac{e^{t_{1}} - e^{-t_{1}}}{e^{t_{1}} + e^{-t_{1}}}, \quad x_{i_{2}} = \frac{2}{e^{t_{1}} + e^{-t_{1}}} \frac{e^{t_{2}} - e^{-t_{2}}}{e^{t_{2}} + e^{-t_{2}}}, \dots, \quad x_{i_{r-1}} = \frac{2}{e^{t_{r-1}} + e^{-t_{r-1}}} \prod_{\ell=1}^{r-1} \frac{e^{t_{\ell}} - e^{t_{\ell}}}{e^{t_{\ell}} + e^{-t_{\ell}}},$$

$$x_{j_{1}} = \frac{e^{s_{1}} - e^{-s_{1}}}{e^{s_{1}} + e^{-s_{1}}}, \quad x_{j_{2}} = \frac{2}{e^{s_{1}} + e^{-s_{1}}} \frac{e^{s_{2}} - e^{-s_{2}}}{e^{s_{2}} + e^{-s_{2}}}, \dots, \quad x_{j_{m-1}} = \frac{2}{e^{s_{m-1}} + e^{-s_{m-1}}} \prod_{\ell=1}^{m-1} \frac{e^{s_{\ell}} - e^{s_{\ell}}}{e^{s_{\ell}} + e^{-s_{\ell}}}.$$
(15)

*Proof.* The semi-algebraic set  $OGr_+(1,n)$  in  $\mathbb{P}^{n-1}$  is cut out by:

$$\sum_{i \in [n] \cap (2\mathbb{Z}+1)} x_i^2 = \sum_{j \in [n] \cap 2\mathbb{Z}} x_j^2 \quad \text{and} \quad x \in \mathbb{P}^{n-1}_+.$$

The cells  $OGr_+(1,n) \cap \Pi_{\sigma(A,B)}$  correspond to the boundaries of  $OGr_+(1,n)$  where the point  $x \in \mathbb{P}^{n-1}_+$  satisfies:

$$\sum_{i \in A} x_i^2 = \sum_{j \in B} x_j^2 \quad \text{and} \quad x_i \neq 0 \iff i \in A \sqcup B.$$
(16)

The closure of such a cell is obtained by driving some of the coordinates indexed by *A* or *B* to 0. So the first and second statement follow. The face structure described in here is the same face structure as the product  $\Delta_{p-1} \times \Delta_{q-1}$  of simplices of dimensions p-1 and q-1.

For the third statement, to parametrize the cell  $OGr_+(1,n) \cap \Pi_{\sigma(A,B)}$  it is enough to parametrize the set of points  $(x_i)_{i \in A} \in \mathbb{R}^{|A|}_{>0}$  and  $(x_j)_{j \in B} \in \mathbb{R}^{|B|}_{>0}$  on the unit spheres  $S^{|A|-1}$  and  $S^{|B|-1}$ . This is exactly what is done in (15).  $\Box$ 

**Example 3.2.** The orthogonal Grassmannian  $OGr_+(1,5)$  has the same combinatorial structure as  $\Delta_1 \times \Delta_2$ . The poset of the boundaries of  $OGr_+(1,5)$  is depicted in Figure 3.

The next theorem shows that  $OGr_+(1,n)$  is a positive geometry. To give its canonical form, it is convenient to permute<sup>2</sup> the coordinates of  $\mathbb{P}^{n-1}$  and write  $OGr_+(1,n)$  as follows:

OGr<sub>+</sub>(1,n) = { (x<sub>1</sub>:...:x<sub>n</sub>) 
$$\in \mathbb{P}^{n-1}_+$$
:  $x_1^2$  + ... +  $x_p^2$  -  $x_{p+1}^2$  - ... -  $x_n^2$  = 0 }.

**Theorem 3.3.** The semi-algebraic set  $OGr_+(1,n) \subset \mathbb{P}^{n-1}_+$  is a positive geometry in the sense of [2] with canonical form:

$$\Omega = (1 + u_{2,1}^2 + u_{3,1}^2 + \dots + u_{p,1}^2) \frac{du_{2,1} \wedge du_{3,1} \wedge \dots \wedge du_{n-1,1}}{u_{2,1} u_{3,1} \cdots u_{n-1,1} u_{n,1}^2}$$

Here,  $u_{i,j} := x_i/x_1$  in the projective coordinates  $(x_1 : \cdots : x_n)$  of  $\mathbb{P}^{n-1}$ .

<sup>&</sup>lt;sup>2</sup>Here, since k = 1, permuting the coordinates does not change the signs of the "minors".




Figure 3: The Hasse diagram of the poset structure on  $\mathfrak{S}_{1.5}$ .

*Proof.* For ease of notation we write  $u_i$  for  $u_{i,1}$ . In the way the form  $\Omega$  is defined, it is clear that for  $p + 1 \le i \le n - 1$ :

$$\operatorname{Res}_{u_i=0}(\Omega) = (-1)^{i+1} (1 + u_2^2 + \dots + u_p^2) \frac{du_2 \wedge \dots \wedge du_i \wedge \dots \wedge du_{n-1}}{u_2 \dots \widehat{u_i} \dots u_{n-1} u_n^2}$$

For  $2 \le i \le p$  we can rewrite  $\Omega$  as:  $\Omega = -(u_{p+1}^2 + \dots + u_n^2) \frac{du_2 \land du_3 \land \dots \land du_{n-1}}{u_2 \ u_3 \ \dots \ u_{n-1} \ u_n^2}$ , and then:

$$\operatorname{Res}_{u_i=0}(\Omega) = (-1)^{i+1} (-u_{p+1}^2 - \dots - u_n^2) \frac{du_2 \wedge \dots \wedge du_i \wedge \dots \wedge du_{n-1}}{u_1 \dots \hat{u_i} \dots u_{n-1} u_n^2}$$

For the boundary corresponding to  $u_n = 0$  note that we have:

$$u_2du_2+\cdots+u_pdu_p-u_{p+1}du_{p+1}-\cdots-u_ndu_n=0$$

Taking the wedge on the right with  $du_3 \wedge \cdots \wedge du_{n-1}$  we get:

$$u_2 du_2 \wedge \cdots \wedge du_{n-1} - u_n du_n \wedge du_3 \wedge \cdots \wedge du_{n-1} = 0.$$

So we can rewrite  $\Omega$  as follows:

$$\Omega = (1 + u_2^2 + \dots + u_p^2) \frac{du_2 \wedge du_3 \wedge \dots \wedge du_{n-1}}{u_2 \ u_3 \ \dots \ u_{n-1} \ u_n^2} = (1 + u_2^2 + \dots + u_p^2) \frac{du_n \wedge du_3 \wedge \dots \wedge du_{n-1}}{u_2^2 u_3 \dots u_{n-1} u_n}$$

We then get the residue at  $u_n = 0$ :

$$\operatorname{Res}_{u_n=0}(\Omega) = (1+u_2^2+\cdots+u_p^2) \frac{du_3\wedge\cdots\wedge du_{n-1}}{u_2^2u_3\ldots u_{n-1}}.$$

The residue at the boundary where  $x_1 = 0$  can be computed similarly by switching to a different affine chart (eg. set  $x_2 = 1$ ). This covers all the boundaries of OGr<sub>+</sub>(1,*n*) and since the residue at each boundary gives the same form for a lower dimensional positive orthogonal Grassmannian OGr<sub>+</sub>(1,*n*), we can carry on taking residues for lower dimensional boundaries in the same way.

**Example 3.4** (OGr<sub>+</sub>(1,4)). The points  $(x_1 : x_2 : x_3 : x_4)$  in the positive orthogonal Grassmannian OGr<sub>+</sub>(1,4) in  $\mathbb{P}^3$  are those that satisfy:

$$x_1^2 - x_2^2 + x_3^2 - x_4^2 = 0$$
 and  $x_1, x_2, x_3, x_4 \ge 0$ .

We can see that  $OGr_+(1,4)$  is a curvy quadrilateral inside the 3-simplex in  $\mathbb{P}^3_+$  as depicted in Figure 4. It is a positive geometry and its canonical form is:

$$\Omega = (1 + u_{3,1}^2) \frac{du_{2,1} \wedge du_{3,1}}{u_{3,1} u_{2,1} u_{4,1}^2}.$$

where  $u_{i,j} = x_i/x_j$ .




Figure 4: The positive Grassmannian  $OGr_+(1,4)$  is the red region in the tetrahedron  $\mathbb{P}^3_+$ . The boundaries of  $OGr_+(1,4)$  lie on the facets of  $\mathbb{P}^3_+$ .

#### 4. The positive Orthogonal Grassmannian $OGr_+(k, 2k+1)$

We recall that we are working with the sign alternating form (1). The positroid cells of  $Gr_+(k, 2k)$  induce a cell decomposition on the positive orthogonal Grassmannian  $OGr_+(k, 2k)$ , and the cells of this decomposition are indexed by fixed-point-free involutions of [2k]. The face structure of  $OGr_+(k, 2k)$  and the parametrization of its cells are studied in detail in [9, Section 5].

One of the reasons positroid cells induce a cell decomposition of the positive orthogonal Grassmannian  $OGr_+(k, 2k)$  is that the latter can be obtained by slicing  $Gr_+(k, 2k)$  by a linear space, see (14). In general, one can obtain  $OGr_+(k, n)$  by slicing the positive flag variety with a linear space as we shall now explain.

For a subspace V in  $\mathbb{C}^n$  of dimension k, we denote by  $V^{\perp}$  its orthogonal complement with respect to the form (1).

**Lemma 4.1.** The Hodge star map  $Gr(k,n) \rightarrow Gr(n-k,n), V \mapsto V^{\perp}$  is given in *Plücker coordinates by:* 

$$q_J = p_{J^c}, \quad \text{for any } J \in \binom{[n]}{n-k},$$

where the  $p_I$  and  $q_J$ 's are the Plücker coordinates in Gr(k,n) and Gr(n-k,n) respectively. In particular it restricts to an isomorphism of positive geometries between  $Gr_+(k,n)$  and  $Gr_+(n-k,n)$ .

*Proof.* If  $(p_I)_I$  are the Plücker coordinates of *V*, the Plücker coordinates  $q_J$  of  $V^{\perp}$  are obtained as follows:

$$q_J = s_J t_J p_{J^c}$$
 for any  $J \in \binom{[n]}{n-k}$ ,

where  $t = \prod_{j \in J} \omega_0(e_j, e_j)$  and  $s_J$  is the sign of the permutation that sorts the string  $JJ^c$ . We compute the signs  $s_J$  and  $t_J$  and get the following:

$$s_J = (-1)^{\sum_{j \notin J} j - k(k+1)/2}$$
 and  $t_J = (-1)^{|J \cap 2\mathbb{Z}|}$ 

From this it is not so hard to see that the sign  $s_J t_J$  does not depend on J. Since we work with projective coordinates, we deduce that  $q_J = p_{J^c}$ , hence the Hodge star map is indeed an isomorphism between  $Gr_+(k,n)$  to  $Gr_+(n-k,n)$ .

Let  $\mathcal{F}(k,n)$  be the 2-step flag variety of partial flags  $V \subset W \subset \mathbb{C}^n$  where  $\dim(V) = k$  and  $\dim(W) = n - k$ . The positive part  $\mathcal{F}_+(k,n)$  of  $\mathcal{F}(k,n)$  is the semi-algebraic set of points  $(V,W) \in \operatorname{Gr}_+(k,n) \times \operatorname{Gr}_+(n-k,n)$  such that  $(V,W) \in \mathcal{F}(k,n)$ . We denote by  $\mathcal{D}$  the *diagonal* subset of  $\mathbb{P}^{\binom{n}{k}} \times \mathbb{P}^{\binom{n}{n-k}}$  i.e.:

$$\mathcal{D} := \left\{ (p,q) \colon p_I = q_{I^c} \quad \text{for any } I \in {[n] \choose k} \right\}.$$

**Proposition 4.1.** The positive orthogonal Grassmannian  $OGr_+(k,n)$  is the intersection of the positive flag variety  $\mathcal{F}_+(k,n)$  with  $\mathcal{D}$  i.e.:

$$OGr_{+}(k,n) = \mathcal{F}_{+}(k,n) \cap \mathcal{D}.$$
(17)

*Proof.* This follows immediately from Lemma 4.1.

This motivates the choice of the sign alternating form (1) in [9, 12]. However, unlike  $\mathcal{F}_+(k, 2k) \cong \text{Gr}_+(k, 2k)$ , the positive region  $\mathcal{F}_+(k, n)$  is not well understood<sup>3</sup> for general k. This motivates the following question:

**Problem 4.2.** Study the face structure of  $\mathcal{F}_+(k,n)$  and find a parametrization of its cells for general (k,n).

The following result is known to the experts, see [7, Proposition 5.1] and [6, Equation (1.7)]. We include a proof here for completeness.

**Proposition 4.3.** The homogeneous coordinate rings of the 2-step flag variety  $\mathcal{F}(k, 2k+1)$  and the Grassmannian Gr(k+1, 2k+2) are isomorphic.

*Proof.* Note that the cone over  $\mathcal{F}(k, 2k+1) \subset \mathbb{P}^{\binom{2k+1}{k}-1} \times \mathbb{P}^{\binom{2k+1}{k+1}-1}$  is an affine variety  $\widehat{\mathcal{F}}(k, 2k+1) \subset \mathbb{A}^{\binom{2k+1}{k}} \times \mathbb{A}^{\binom{2k+1}{k+1}} = \mathbb{A}^{\binom{2k+2}{k+1}}$  and the cone over the Grassmannian  $\operatorname{Gr}(k+1, 2k+2)$  is the affine variety  $\widehat{\operatorname{Gr}}(k+1, 2k+2) \subset \mathbb{A}^{\binom{2k+2}{k+1}}$ . To prove the above statement, we show that  $\widehat{\mathcal{F}}(k, 2k+1) = \widehat{\operatorname{Gr}}(k+1, 2k+2)$  in  $\mathbb{A}^{\binom{2k+2}{k+1}}$ . To that end, it suffices exhibit a quasi affine variety U that is open and dense in both  $\widehat{\operatorname{Gr}}(k+1, 2k+2), \widehat{\mathcal{F}}(k, 2k+1) \subset \mathbb{A}^{\binom{2k+2}{k+1}}$ .

The points of the variety  $U \subset \mathbb{A}^{\binom{2k+2}{k+1}}$  are obtained by taking taking the  $(k+1) \times (k+1)$ -minors of the  $(k+1) \times (2k+2)$  matrices of the following form:

$$\begin{bmatrix} 1 & \cdots & 0 & * & \cdots & * & 0 \\ \vdots & \ddots & \vdots & \vdots & \cdots & \vdots & \vdots \\ 0 & \cdots & 1 & * & \cdots & * & 0 \\ 0 & \cdots & 0 & t & * \cdots & * & z \end{bmatrix},$$
(18)

with  $* \in \mathbb{A}$  and  $t, z \in \mathbb{C}^{\times}$ . It is clear how U sits inside  $\widehat{\operatorname{Gr}}(k+1, 2k+2)$ . We now describe how to obtain points  $(\widehat{V}, \widehat{W}) \in \widehat{\mathcal{F}}(k, 2k+1)$  from the matrices (18). We obtain  $\widehat{W}$  by taking  $(k+1) \times (k+1)$  minors excluding the last column of (18), while  $\widehat{V}$  is obtained by taking the  $k \times k$  minors of the upper left  $k \times (2k+1)$  part of (18), all multiplied by z. This shows that  $\widehat{\mathcal{F}}(k, 2k+1) = \widehat{\operatorname{Gr}}(k+1, 2k+2)$  is the Zariski closure of U in  $\mathbb{A}^{\binom{2k+2}{k+1}}$  which finishes the proof.  $\Box$ 

**Theorem 4.4.** The positive orthogonal Grassmannians  $OGr_+(k, 2k + 1)$  and  $OGr_+(k+1, 2k+2)$  can be identified through a linear isomorphism.

<sup>&</sup>lt;sup>3</sup>The Lusztig positive part of  $\mathcal{F}(k,n)$  is well understood but it can be shown that the Plücker positive region  $\mathcal{F}_+(k,n)$  strictly contains the Lusztig positive region when n > 2k + 1, see [5].

*Proof.* We denote the Plücker coordinates on OGr(k, 2k + 1) by  $p_I$  and those of OGr(k + 1, 2k + 2) by  $q_J$ . The orthogonal Grassmannian OGr(k, 2k + 1) is irreducible and the map:

$$\Phi_k: \quad \operatorname{OGr}(k+1,2k+2) \quad \to \quad \operatorname{OGr}(k,2k+1) \\
(q_J) \qquad \mapsto \quad p_I = q_{I \cup \{2k+2\}}$$
(19)

is an isomorphism between the OGr(k, 2k + 1) (which is irreducible) and the standard component in OGr(k + 1, 2k + 2). It is not so difficult to see that this isomorphism restricts to an isomorphism between the positive orthogonal Grassmannians OGr $_+(k, 2k + 1)$  and OGr $_+(k + 1, 2k + 2)$ .

**Remark 4.5.** The equations that cut out OGr(k, 2k + 1) in Gr(k, 2k + 1) are all quadrics. So it is remarkable that we can still describe the face structure of  $OGr_+(k, 2k + 1)$  from our understanding of the face structure of  $OGr_+(k + 1, 2k + 2)$  which is obtained by taking a linear slice of  $Gr_+(k + 1, 2k + 2)!$ 

**Example 4.6** (OGr<sub>+</sub>(2,5)). The orthogonal Grassmannian OGr<sub>+</sub>(2,5) is isomorphic to OGr<sub>+</sub>(3,6). The Hasse diagram of the face poset of the latter is given in [9, Figure 7]. Figure 5 gives the same Hasse diagram but in the realizable permutations of [5]. Each of the cells can be parameterized by combining the isomorphism in (19) and [9, Theorem 5.17 (i)].




Figure 5: The face poset of  $OGr_+(2,5)$  matches that of  $OGr_+(3,6)$ . See Figure 7 in [9].

We finish this section by explaining how one goes from matchings  $\tau$  on [2k+2] to the admissible permutations in [2k+1] i.e. permutations  $\sigma$  of [2k+1] with corresponding positroid cell  $\Pi_{\sigma}$  such that  $\Pi_{\sigma} \cap OGr_{+}(k, 2k+1)$  is nonempty.

Let *c* denote the chord in  $\tau$  attached to the vertex 2k + 2 and, starting from the vertex 2k + 2, consider the largest sequence  $c = c_1, c_2, \ldots, c_r$  of pairwise intersecting chords of  $\tau$ . Denote the 2r vertices of these chords by  $i_1 < \cdots < i_{2r-1} < 2k + 2$ . Then the cell  $\Pi_{\tau} \cap \text{OGr}_+(k+1, 2k+2)$  is isomorphic to the cell  $\Pi_{\sigma} \cap \text{OGr}_+(k, 2k+1)$  where  $\sigma$  is the permutation of [2k+1] obtained by replacing the chords  $c_1, \ldots, c_r$  with the unique cycle with support  $\{i_1, \ldots, i_{2r-1}\}$ and *r* excedances. See Figure 6 for an example.




Figure 6: A matching  $\tau$  of [2k+2] and the corresponding permutation  $\sigma$  of [2k+1] for k = 7. In the left figure, starting vertex 16 (in blue), the chords in red are longest sequence of chords  $c_1, \ldots, c_r$  that intersect each other pairwise. In the figure on the right, the blue vertex is deleted and the chords in red are turned into the unique cycle with support  $\{1, 2, 4, 8, 11, 13, 15\}$  and 4 excedances.

## 5. What goes wrong for $OGr^+(k, n)$ when n > 2k + 1 and k > 1?

In this section we show why positroid cells fail to induce a cell decomposition of  $OGr_+(k,n)$  as soon as n > 2k + 1 and k > 1. Let us start with the following:

**Definition 5.1.** For any positroid  $\mathcal{M}$  of type (k, n) and for any pair of subsets I, J of [n] of size k - 1 we define the following two subsets of [n]:

$$A_{IJ}^{\pm}(\mathscr{M}) = \left\{ \ell \in [n] \colon I\ell, J\ell \in \mathscr{M} \text{ and } (-1)^{\ell-1} \varepsilon_{I\ell} \varepsilon_{J\ell} = \pm 1 \right\}.$$

We say that  $\mathcal{M}$  is an orthopositroid if for any  $I, J \in {[n] \choose k-1}$  we have:

$$A^+_{IJ}(\mathscr{M}) = \emptyset \quad \iff \quad A^-_{IJ}(\mathscr{M}) = \emptyset.$$

**Example 5.2.** Let n = 5 and consider the two following positroids:

$$\mathcal{M}_1 = \{\{1,2\},\{1,4\},\{2,5\},\{4,5\}\} \text{ and } \mathcal{M}_2 = \{\{1,2\},\{1,3\},\{2,4\},\{3,4\}\}.$$

We then have:

$$A_{24}^+(\mathscr{M}_1) = \emptyset$$
 and  $A_{24}^-(\mathscr{M}_1) = \{2\}.$ 

So  $\mathcal{M}_1$  is **not** an orthopositroid. One can check that  $\mathcal{M}_2$  is an orthopositroid.

The motivation behind this definition is that if X is a point in  $OGr_+(k,n)$  and  $\mathcal{M}_X$  is its associated positroid then  $\mathcal{M}_X$  is necessarily an orthopositroid in the sense of Definition 5.1. This is because the Plücker coordinates of X satisfy the equations (5).

Since we will show that positroid cells dot not induce a CW cell decomposition of  $OGr_+(k,n)$ , we refrain from elaborating more on the realizability of orthopositroids for general k, leaving that discussion for upcoming work, and state the following

**Conjecture 5.3.** The orthopositroids in the sense of Definition 5.1 are realizable i.e. for any orthopositroid  $\mathcal{M}$  of type (k,n), there exists  $X \in OGr_+(k,n)$  such that  $\mathcal{M} = \mathcal{M}_X$ .

Let us start with OGr<sub>+</sub>(2,6). A exhaustive computation shows that, out of all the positroids  $\mathcal{M}$  (or decorated permutations  $\sigma$ ) of type (2,6), there are exactly 99 orthopositroids (or admissible permutations). All of these orthopositroids are realizable in OGr<sub>+</sub>(2,6) and are listed in Table 1. Let us now focus on the following orthopositroid cells  $\sigma$  and  $\tau$  in OGr<sub>+</sub>(2,6):



We shall show that  $C_{\sigma} = \Pi_{\sigma} \cap OGr_{+}(2, 6)$  and  $C_{\tau} = \Pi_{\tau} \cap OGr_{+}(2, 6)$ , which are cells of dimension 2, have the combinatorial type of a triangle and a square respectively. We start by giving generic matrices  $M_{\sigma}, M_{\tau}$  that describe the points of the cells  $C_{\sigma}, C_{\tau}$  respectively:

$$M_{\sigma} = \begin{bmatrix} 1 & 1 & 0 & 0 & -x & -x \\ 0 & 0 & 1 & 1 & y & y \end{bmatrix}, \quad \text{where } x, y > 0,$$
$$M_{\tau} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & a & b & c \end{bmatrix}, \quad \text{where } \begin{cases} a, b, c > 0 \\ 1 + b^2 = a^2 + c^2 \end{cases}$$

Note that on the one hand, the closure of the cell  $C_{\tau}$  in OGr<sub>+</sub>(2,6) is isomorphic to OGr<sub>+</sub>(1,4) so Theorem 3.1 implies that  $\overline{C_{\tau}}$  has the same combinatorial type

as a square. On the other hand, the closure of the cell  $C_{\sigma}$  has the combinatorial type of a triangle. Its edges are given by:

 $e_1 = \begin{bmatrix} 1 & 1 & 0 & 0 & b & b \\ 0 & 0 & 1 & 1 & 0 & 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 1 & 1 & b & b & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix}, \quad e_3 = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & b & b \end{bmatrix} \quad b \ge 0.$ 

The edge  $e_1$  is one of the diagonals of the "square"  $C_{\tau}$ . So the cell  $C_{\sigma}$  glues with the cell  $C_{\tau}$  as in Figure 7.



Figure 7: A cartoon of the cell  $C_{\sigma}$  (in green) glued to the cell  $C_{\tau}$  (in red).

This shows that, to get a cell decomposition, the positroid cell  $C_{\tau}$  needs to be subdivided into two cells and its two halves together with the diagonal that separate them need to be added to the 99 orthopositroids. This shows that the positroid cells are not enough to induce a cell decomposition on OGr<sub>+</sub>(2,6).

In general this problem arises as soon as n > 2k + 1. This is because whenever n > 2k + 1 we have  $n - 6 \ge 2(k - 2)$ , so by extending a  $2 \times 6$  matrix in OGr<sub>+</sub>(2,6) by a  $(k-2) \times (n-6)$  as follows:

			 											1	
												( <b>0</b> )			
Ι.												(0)			
			1 												
				0	1	1	0	0	*	*	*	*	*	*	
L			(0)									*			

We can then realize each positroid cell in  $OGr_+(2,6)$  as some positroid cell  $\Pi_{\sigma} \cap OGr_+(k,n)$  of  $OGr_+(k,n)$  and the same argument as above applies again. This highlights the need for new combinatorics to give a cell decomposition of  $OGr_+(k,n)$  for n > 2k+1.

**Problem 5.4.** Find a cell decomposition for  $OGr_+(k, n)$  when n > 2k + 1 and describe the combinatorics behind its face poset.




Table 1: The 99 orthopositroids of type (2,6) (in permutation form) in the middle column. The dimensions of their corresponding cells in OGr<sub>+</sub>(2,6) are given in the leftmost column. The number of cells in each dimension is given in the rightmost column.

Acknowlegements: We are grateful to Elizabeth Pratt for pointing us to the formula (11) used in the proof of Proposition 2.7 and Theorem 2.9. We also thank Lara Bossinger and Sebastian Seemann for helpful comments. This project started during our visit to the Max Planck Institute for Mathematics in the Sciences. We thank MPI MiS for its generous hospitality. YEM was partially supported by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) SFB-TRR 195 "Symbolic Tools in Mathematics and their Application." YM is supported by the National Science Foundation under Award DMS-2402069 and the Bob Moses Fund of the Institute for Advanced Study.

#### REFERENCES

- ABENDA, S., AND GRINEVICH, P. G. Rational degenerations of m-curves, totally positive Grassmannians and KP2-solitons. *Comm. Math. Phys.* 361, 3 (2018), 1029–1081.
- [2] ARKANI-HAMED, N., BAI, Y., AND LAM, T. Positive geometries and canonical forms. *Journal of High Energy Physics 2017*, 11 (2017), 1–124.
- [3] ARKANI-HAMED, N., BOURJAILY, J., CACHAZO, F., GONCHAROV, A., POST-NIKOV, A., AND TRNKA, J. Grassmannian geometry of scattering amplitudes. Cambridge University Press, Cambridge, 2016.
- [4] ARKANI-HAMED, N., AND TRNKA, J. The amplituhedron. *Journal of High Energy Physics 2014*, 10 (2014), 30.
- [5] BARKLEY, G., BORETSKY, J., EUR, C., AND GAO, J. On two notions of total positivity for generalized partial flag varieties of classical lie types. *arXiv:2410.11804* (2024).
- [6] BOSSINGER, L., AND LI, J.-R. Cluster structures on spinor helicity and momentum twistor varieties, 2024.
- [7] EL MAAZOUZ, Y., PFISTER, A., AND STURMFELS, B. Spinor-helicity varieties. *arXiv:2406.17331* (2024).
- [8] GALASHIN, P., KARP, S. N., AND LAM, T. The totally nonnegative Grassmannian is a ball. *Adv. Math.* 397 (2022), Paper No. 108123, 23.
- [9] GALASHIN, P., AND PYLYAVSKYY, P. Ising model and the positive orthogonal Grassmannian. *Duke Math. J. 169*, 10 (2020), 1877–1942.
- [10] GELFAND, I., GORESKY, M., MACPHERSON, R., AND SERGANOVA, V. Combinatorial geometries, convex polyhedra, and schubert cells. *Advances in Mathematics* 63, 3 (1987), 301–316.
- [11] GRIFFITHS, P., AND HARRIS, J. Principles of algebraic geometry. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1994. Reprint of the 1978 original.
- [12] HUANG, Y.-T., WEN, C., AND XIE, D. The positive orthogonal Grassmannian and loop amplitudes of ABJM. J. Phys. A 47, 47 (2014), 474008, 48.
- [13] KODAMA, Y., AND WILLIAMS, L. Kp solitons and total positivity for the grassmannian. *Inventiones mathematicae 198* (2014), 637–699.
- [14] LUSZTIG, G. Total positivity in reductive groups. In *Lie theory and geometry*, vol. 123 of *Progr. Math.* Birkhäuser Boston, Boston, MA, 1994, pp. 531–568.
- [15] LUSZTIG, G. Introduction to total positivity. In *Positivity in Lie theory: open problems*, vol. 26 of *De Gruyter Exp. Math.* de Gruyter, Berlin, 1998, pp. 133–145.
- [16] LUSZTIG, G. Total positivity in partial flag manifolds. *Represent. Theory 2* (1998), 70–78.
- [17] POSTNIKOV, A. Total positivity, grassmannians, and networks. *arXiv:0609764* (2006).

- [18] POSTNIKOV, A. Positive Grassmannian and polyhedral subdivisions. In Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. IV. Invited lectures (2018), World Sci. Publ., Hackensack, NJ, pp. 3181–3211.
- [19] SAGAN, B. E. *Combinatorics: The art of counting*, vol. 210. American Mathematical Soc., 2020.
- [20] WILLIAMS, L. K. The positive Grassmannian, the amplituhedron, and cluster algebras. In *Proceedings of the International Congress of Mathematicians*. Vol. 6. Sections 12–14. EMS Press, Berlin, 2023, pp. 4710–4737.

YASSINE EL MAAZOUZ California Institute of Technology e-mail: maazouz@caltech.edu

> YELENA MANDELSHTAM Institute for Advanced Study *e-mail:* yelenam@ias.edu