EXAMPLES OF CYCLIC POLYNOMIALLY BOUNDED OPERATORS THAT ARE NOT SIMILAR TO CONTRACTIONS

MARIA F. GAMAL'

ABSTRACT. A question if a polynomially bounded operator is similar to a contraction was posed by Halmos and was answered in the negative by Pisier. His counterexample is an operator of infinite multiplicity, while all its restrictions on invariant subspaces of finite multiplicity are similar to contractions. In the paper, cyclic polynomially bounded operators which are not similar to contractions and are quasisimilar to C_0 -contractions or to isometries are constructed. The construction is based on a perturbation of the sequence of finite dimensional operators which is uniformly polynomially bounded, but is not uniformly completely polynomially bounded, constructed by Pisier.

1. INTRODUCTION

Let \mathcal{H} be a Hilbert space, and let $T: \mathcal{H} \to \mathcal{H}$ be an operator on \mathcal{H} . T is called *polynomially bounded*, if there exists a constant M such that

(1.1)
$$||p(T)|| \le M ||p||_{\infty}$$
 for every polynomial p .

The smallest constant M which satisfies (1.1) is denoted by $M_{pb}(T)$ and is called the *polynomial bound* of T. If T is not polynomially bounded, then $M_{pb}(T) = \infty$.

For a natural number $n \ a \ n \times n$ matrix can be regarded as an operator on ℓ_n^2 , its norm is denoted by the symbol $\|\cdot\|_{\mathcal{L}(\ell_n^2)}$. For a family of polynomials $[p_{ij}]_{i,j=1}^n$ put

$$[p_{ij}]_{i,j=1}^n \|_{H^{\infty}(\ell_n^2)} = \sup\{ \| [p_{ij}(z)]_{i,j=1}^n \|_{\mathcal{L}(\ell_n^2)}, \ z \in \mathbb{D} \}.$$

For an operator T on \mathcal{H} and a family of polynomials $[p_{ij}]_{i,j=1}^n$ the operator $[p_{ij}(T)]_{i,j=1}^n$ acting on the space $\bigoplus_{j=1}^n \mathcal{H}$ is defined. T is called *completely polynomially bounded*, if there exists a constant M such that

(1.2)
$$\|[p_{ij}(T)]_{i,j=1}^n\| \le M \|[p_{ij}]_{i,j=1}^n\|_{H^{\infty}(\ell_n^2)}$$
for every family of polynomials $[p_{ij}]_{i,j=1}^n$.

The smallest constant M which satisfies (1.2) is denoted by $M_{cpb}(T)$ and is called the *complete polynomial bound* of T. If T is not complete polynomially bounded, then $M_{cpb}(T) = \infty$.

An operator T is called a *contraction* if $||T|| \leq 1$. The following criterion for an operator to be similar to a contraction is proved in [Pa]:

²⁰¹⁰ Mathematics Subject Classification. Primary 47A60; Secondary 47A65, 47A16, 47A20.

Key words and phrases. Polynomially bounded operator, similarity, contraction, unilateral shift, isometry, C_0 -contraction, C_0 -operator.

An operator T is similar to a contraction if and only if $M_{cpb}(T) < \infty$, and

$$M_{cpb}(T) = \inf\{\|X\| \|X^{-1}\| : \|XTX^{-1}\| \le 1\},\$$

where the operators X are invertible.

The question if a polynomially bounded operator is similar to a contraction was posed by Halmos [H] and was answered in the negative by Pisier [Pi]. The counterexample is an appropriate Foguel–Hankel operator. Recall that Γ is a (vector-valued) Hankel operator if and only if $S_k^*\Gamma = \Gamma S_n$, where $1 \leq n, k \leq \infty$, S_n is the unilateral shift of multiplicity n. A Foguel–Hankel operator is an operator of the form

$$\begin{pmatrix} S_k^* & \Gamma \\ \mathbb{O} & S_n \end{pmatrix},$$

where $1 \le n, k \le \infty$, Γ is a Hankel operator. There exists a Hankel operator Γ such that the operator

$$T = \begin{pmatrix} S_{\infty}^* & \Gamma \\ \mathbb{O} & S_{\infty} \end{pmatrix}$$

is polynomially bounded and is not similar to a contraction [Pi]. Since $\mu_{S_{\infty}} = \infty$, $\mu_T = \infty$, where μ_T is the multiplicity of the operator T.

On the other hand, if $1 \leq n < \infty$ or $1 \leq k < \infty$, and Γ is a Hankel operator, then the Foguel–Hankel operator $\begin{pmatrix} S_k^* & \Gamma \\ \mathbb{O} & S_n \end{pmatrix}$ is polynomially bounded if and only if it is similar to a contraction [DP, Theorem 4.4 and Corollary 4.5], [Pe, Theorem 15.1.2], see references in [DP] and in [Pe] to the history of the question. Therefore, if a Foguel–Hankel operator T is polynomially bounded, and if \mathcal{M} is an invariant subspace of T such that $\mu_{T|\mathcal{M}} < \infty$, then $T|_{\mathcal{M}}$ is similar to a contraction, even if T is not similar to a contraction, see Lemma 1.3 below for the proof.

The purpose of this paper is to construct a polynomially bounded operator T such that T is not similar to a contraction and $\mu_T = 1$, that is, T is cyclic. The main result of the paper is Theorem 7.1, where a polynomially bounded operator T is constructed such that T is not similar to a contraction and there exist quasiaffinities X and Y, an outer function $g \in H^{\infty}$, and a cyclic C_0 -contraction T_0 such that $XT = T_0X$, $YT_0 = TY$, and $XY = g(T_0)$. This result allows us to construct polynomially bounded operators that are not similar to contractions and are quasisimilar to cyclic absolutely continuous isometries: to the unilateral shift, to the bilateral shift, to absolutely continuous cyclic reductive unitary operators (Section 2). In particular, we provide a negative answer to Question 18 from [K]: "if the polynomially bounded operator T is a quasisfinite transform of a singular unitary operator, then T is similar to this unitary operator, see [M], [AT], or [K, Theorem 17].)

The main construction is based on a perturbation of the sequence $\{R_N\}_N$ of finite dimensional operator R_N such that

 $\sup_{N} M_{pb}(R_N) < \infty \quad \text{and} \quad \sup_{N} M_{cpb}(R_N) = \infty,$

from [DP] and [Pi], see (5.8) below.

We need the following notation and definition.

 $\mathcal{A}(\mathbb{D})$ is the *disk algebra*, and H^{∞} is the Banach algebra of all analytic bounded functions in \mathbb{D} . The uniform norm on \mathbb{D} is denoted by the symbol $\|\cdot\|_{\infty}$.

For $w \in \mathbb{D}$ put

(1.3)
$$\beta_w(z) = \frac{w-z}{1-\overline{w}z}, \quad z \in \mathbb{D}.$$

Clearly, $\beta_w \circ \beta_w(z) = z$ for every $z \in \mathbb{D}$, and for every $\varphi \in H^{\infty}$ we have $\varphi \circ \beta_w \in H^{\infty}$, $\|\varphi \circ \beta_w\|_{\infty} = \|\varphi\|_{\infty}$, and if $\varphi \in \mathcal{A}(\mathbb{D})$, then $\varphi \circ \beta_w \in \mathcal{A}(\mathbb{D})$.

Since polynomials are dense in $\mathcal{A}(\mathbb{D})$ in the uniform norm, polynomials in (1.1) and (1.2) can be replaced by functions from $\mathcal{A}(\mathbb{D})$. Let $\sigma(T) \subset \overline{\mathbb{D}}$, and let $w \in \mathbb{D}$. Then the operator $\beta_w(T) = (w - T)(I - \overline{w}T)^{-1}$ is well defined, $p(\beta_w(T)) = (p \circ \beta_w)(T)$, and it is easy to see that

(1.4)
$$M_{pb}(\beta_w(T)) = M_{pb}(T)$$
 and $M_{cpb}(\beta_w(T)) = M_{cpb}(T)$.

The following proposition is simple, but important in our construction.

Proposition 1.1. Let $\{T_N\}_N$ be a family of operators such that $\sup_N ||T_N|| < \infty$. Put $T = \bigoplus_N T_N$. Then $M_{pb}(T) = \sup_N M_{pb}(T_N)$ and $M_{cpb}(T) = \sup_N M_{cpb}(T_N)$.

Proof. Since $||T|| = \sup_N ||T_N||$ and $p(T) = \bigoplus_N p(T_N)$ for every polynomial p, we have that $||p(T)|| = \sup_N ||p(T_N)||$. Therefore, the equality for M_{pb} is fulfilled. Denote by \mathcal{H}_N the spaces on which T_N act. Let $[p_{ij}]_{i,j=1}^n$ be a family of polynomials. The operator $R = [p_{ij}(T)]_{i,j=1}^n$ acts on the space $\mathcal{K} = \bigoplus_{j=1}^n \bigoplus_N \mathcal{H}_N$. Rewriting this space as $\mathcal{K} = \bigoplus_N \bigoplus_{j=1}^n \mathcal{H}_N$ we obtain that $R = \bigoplus_N [p_{ij}(T_N)]_{i,j=1}^n$. Therefore, $||[p_{ij}(T)]_{i,j=1}^n|| = \sup_N ||[p_{ij}(T_N)]_{i,j=1}^n||$, and we conclude that the equality for M_{cpb} is fulfilled. \Box

Corollary 1.2. Let $\{T_N\}_N$ be a family of operators such that $\sigma(T_N) \subset \overline{\mathbb{D}}$ for every index N, and let $\{w_N\}_N \subset \mathbb{D}$. Suppose that $\sup_N \|\beta_{w_N}(T_N)\| < \infty$, where β_{w_N} are defined in (1.3). Put $T = \bigoplus_N \beta_{w_N}(T_N)$. Then $M_{pb}(T) =$ $\sup_N M_{pb}(T_N)$ and $M_{cpb}(T) = \sup_N M_{cpb}(T_N)$.

Proof. The corollary follows from Proposition 1.1 and the equalities (1.4).

The paper is organized as follows. In Section 2, it is shown how to construct operators quasisimilar to isometries and having given compressions on their semi-invariant subspaces (not for arbitrary given compressions). Sections 3 and 4 contain preliminary results concerning functions and operators, respectively. Most of them, excepting maybe Theorem 3.8, are simple or known, and are given to achieve a complete exposition. In Section 5 the sequence of finite dimensional operators from [DP], [Pi] is described, because the knowledge of the structure of these operators is needed to construct an appropriate perturbation of them. In Section 6 the perturbation of the sequence of finite dimensional operators described in Section 5 is constructed. Then this perturbation is used to construct a cyclic polynomially bounded operator which is not similar to a contraction (Corollary 6.4). This operator is quasisimilar to a C_0 -contraction, but the author doesn't know if the product of intertwining quasiaffinities is an outer function of this C_0 -contraction. In Section 7 an additional construction is given that guarantees the existence of such a function. It allows to apply the results of Section 2 to a C_0 -operator constructed in Section 7.

In the rest part of Introduction, the needed notation and definitions are given, and Lemma 1.3 is formulated and proved.

Let \mathcal{H} be a (complex, separable) Hilbert space, and let \mathcal{M} be a (linear, closed) subspace. By $I_{\mathcal{H}}$ and $P_{\mathcal{M}}$ the identical operator on \mathcal{H} and the orthogonal projection from \mathcal{H} onto \mathcal{M} are denoted, respectively.

Let T and R be operators on Hilbert spaces \mathcal{H} and \mathcal{K} , respectively, and let $X : \mathcal{H} \to \mathcal{K}$ be a linear bounded transformation such that X intertwines T and R, that is, XT = RX. If X is unitary, then T and R are called unitarily equivalent, in notation: $T \cong R$. If X is invertible, that is, X has a bounded inverse X^{-1} , then T and R are called similar, in notation: $T \approx R$. If X is a quasiaffinity, that is, ker $X = \{0\}$ and clos $X\mathcal{H} = \mathcal{K}$, then T is called a quasiaffine transform of R, in notation: $T \prec R$. If $T \prec R$ and $R \prec T$, then T and R are called quasisimilar, in notation: $T \sim R$.

The multiplicity μ_T of an operator T acting on a space \mathcal{H} is the minimum dimension of its reproducing subspaces:

$$\mu_T = \min\{\dim E : E \subset \mathcal{H}, \quad \bigvee_{n=0}^{\infty} T^n E = \mathcal{H}\}.$$

An operator T is called *cyclic*, if $\mu_T = 1$. Let \mathcal{M} be an invariant subspace of T, that is, \mathcal{M} is a subspace of \mathcal{H} such that $T\mathcal{M} \subset \mathcal{M}$. It is well known and easy to see that

(1.5)
$$\mu_{P_{\mathcal{M}^{\perp}}T|_{\mathcal{M}^{\perp}}} \le \mu_T \le \mu_{T|_{\mathcal{M}}} + \mu_{P_{\mathcal{M}^{\perp}}T|_{\mathcal{M}^{\perp}}}$$

and

(1.6)
$$\mu_T \ge \dim \ker T^*$$

(see, for example, [Ni, II.D.2.3.1]). It follows immediately from the definition of quasiaffine transform that if $T \prec R$, then $\mu_R \leq \mu_T$.

 \mathbb{D} is the open unit disk, \mathbb{T} is the unit circle, H^2 is the Hardy space on \mathbb{D} , H^{∞} is the Banach algebra of all bounded analytic functions in \mathbb{D} . The unilateral shift S of multiplicity 1 is the operator of multiplication by the independent variable on H^2 . For an inner function $\theta \in H^{\infty}$ the subspace θH^2 is invariant for S, put

(1.7)
$$\mathcal{K}_{\theta} = H^2 \ominus \theta H^2 \quad \text{and} \quad T_{\theta} = P_{\mathcal{K}_{\theta}} S|_{\mathcal{K}_{\theta}}.$$

If T is a polynomially bounded operator, then $T = T_a + T_s$, where T_a is an absolutely continuous polynomially bounded operator, that is, a H^{∞} functional calculus is well-defined for T_a , and T_s is similar to a singular unitary operator, see [M] or [K]. In this paper, absolutely continuous polynomially bounded operators are regarded. An absolutely continuous polynomially bounded operator T is called a C_0 -operator, if there exists $\varphi \in H^{\infty}$ such that $\varphi(T) = \mathbb{O}$, see [BP]; if a C_0 -operator is a contraction, it is called a C_0 -contraction, see [SFBK]. For an inner function θ , T_{θ} is a C_0 -contraction, because $\theta(T_{\theta}) = \mathbb{O}$.

For a cardinal number $n, 0 \leq n \leq \infty, H_n^2$ is the orthogonal sum of n copies of H^2 , and the unilateral shift S_n is the operator of multiplication

by the independent variable on H_n^2 . It is well known and easy to see that $S_n \cong \bigoplus_{l=1}^n S$ and $\mu_{S_n} = n$.

Lemma 1.3. Let $1 \le n, k \le \infty$ be cardinal numbers, and let $\Gamma: H_n^2 \to H_k^2$ be an operator such that $S_k^* \Gamma = \Gamma S_n$ (that is, Γ is a Hankel operator). Put

$$T = \begin{pmatrix} S_k^* & \Gamma \\ \mathbb{O} & S_n \end{pmatrix},$$

T acts on $H_k^2 \oplus H_n^2$. Suppose T is polynomially bounded, and \mathcal{M} is an invariant subspace for T such that $\mu_{T|_{\mathcal{M}}} < \infty$. Then $T|_{\mathcal{M}}$ is similar to a contraction.

Proof. Put $\mathcal{M}_1 = (H_k^2 \oplus \{0\}) \vee \mathcal{M}$ and $T_1 = T|_{\mathcal{M}_1}$. It follows from the definition of the multiplicity that $\mu_{T_1} \leq \mu_{T|_{H_k^2 \oplus \{0\}}} + \mu_{T|_{\mathcal{M}}}$. Since $T|_{H_k^2 \oplus \{0\}} = S_k^*$ and $\mu_{S_k^*} = 1$ for $1 \leq k \leq \infty$ (see, for example, [Ni, Lemma II.D.2.4.20]), we obtain that $\mu_{T_1} < \infty$. Put $\mathcal{N} = \mathcal{M}_1 \ominus (H_k^2 \oplus \{0\})$. Then $\mathcal{N} \subset \{0\} \oplus H_n^2$, $S_n \mathcal{N} \subset \mathcal{N}$, and $P_{\mathcal{N}} T_1|_{\mathcal{N}} = S_n|_{\mathcal{N}}$. Put $\Gamma_1 = \Gamma|_{\mathcal{N}}$. Then

$$T_1 = \begin{pmatrix} S_k^* & \Gamma_1 \\ \mathbb{O} & S_n |_{\mathcal{N}} \end{pmatrix}$$

and $S_k^* \Gamma_1 = \Gamma_1 S_n |_{\mathcal{N}}$.

By (1.5), $\mu_{P_N T_1|_N} \leq \mu_{T_1} < \infty$. Therefore, there exists $n_1, 1 \leq n_1 < \infty$, such that $S_n|_N \cong S_{n_1}$. Thus,

$$T_1 \cong \begin{pmatrix} S_k^* & \Gamma_2 \\ \mathbb{O} & S_{n_1} \end{pmatrix},$$

where Γ_2 is such that $S_k^*\Gamma_2 = \Gamma_2 S_{n_1}$. Since $n_1 < \infty$, T_1 is similar to a contraction by [DP, Theorem 4.4]. Since $T|_{\mathcal{M}} = T_1|_{\mathcal{M}}$, $T|_{\mathcal{M}}$ is similar to a contraction, too.

2. Construction of operators quasisimilar to isometries

The following definition and notation will be used in this section. An operator T is called *power bounded*, if $\sup_{n\geq 1} ||T^n|| < \infty$. For an operator T, let $\{T\}'$ denote the commutant of T, that is, the algebra of all operators that commute with T.

The following lemma is very simple, therefore, its proof is omitted.

Lemma 2.1. Suppose \mathcal{H}_0 , \mathcal{K}_0 , \mathcal{H}_1 , \mathcal{K}_1 are Hilbert spaces, $Y_0: \mathcal{H}_0 \to \mathcal{K}_0$, $Y_1: \mathcal{H}_1 \to \mathcal{K}_1$, $Z: \mathcal{H}_1 \to \mathcal{K}_0$ are operators. Put

$$Y = \begin{pmatrix} Y_0 & Z \\ \mathbb{O} & Y_1 \end{pmatrix}.$$

Then

(i) if Y_0 and Y_1 are quasiaffinities, then Y is a quasiaffinity;

- (ii) if Y is a quasiaffinity, then $\operatorname{clos} Y_1 \mathcal{H}_1 = \mathcal{K}_1$;
- (iii) if Y_0 is invertible and Y is a quasiaffinity, then Y_1 is a quasiaffinity.

The following proposition will be applied to construct an example of a polynomially bounded operator, which is not similar to a contraction and is a quasiaffine transform of S.

Proposition 2.2. Suppose \mathcal{H} , \mathcal{K} , \mathcal{M} are Hilbert spaces, $T_0: \mathcal{H} \to \mathcal{H}$, $R_0: \mathcal{K} \to \mathcal{K}$, $V: \mathcal{M} \to \mathcal{M}$, $Y_0: \mathcal{H} \to \mathcal{K}$, $K: \mathcal{K} \to \mathcal{M}$, $Z: \mathcal{H} \to \mathcal{M}$ are operators. Moreover, suppose $Y_0T_0 = R_0Y_0$. Put $A = VZ - ZT_0 + KY_0$,

$$T = \begin{pmatrix} V & A \\ \mathbb{O} & T_0 \end{pmatrix}, \quad R = \begin{pmatrix} V & K \\ \mathbb{O} & R_0 \end{pmatrix}, \quad and \quad Y = \begin{pmatrix} I_{\mathcal{M}} & Z \\ \mathbb{O} & Y_0 \end{pmatrix}.$$

Then YT = RY. If R and T_0 are power bounded, then T is power bounded. If R and T_0 are polynomially bounded, then T is polynomially bounded.

Proof. The equality YT = RY is a straightforward consequence of the definition of T, R, and Y and the equality $Y_0T_0 = R_0Y_0$. For a polynomial p set $A_{(p)} = P_{\mathcal{M}}p(T)|_{\mathcal{H}}$ and $K_{(p)} = P_{\mathcal{M}}p(R)|_{\mathcal{K}}$. From the equality Yp(T) = p(R)Y writing in the matrix form it is easy to see that

$$p(T) = \begin{pmatrix} p(V) & A_{(p)} \\ \mathbb{O} & p(T_0) \end{pmatrix}, \text{ and } A_{(p)} = p(V)Z - Zp(T_0) + K_{(p)}Y_0.$$

Now suppose that R and T_0 are polynomially bounded. Since $p(V) = p(R)|_{\mathcal{M}}$, we have $||p(V)|| \leq M_{pb}(R)||p||_{\infty}$, and the estimate $||K_{(p)}|| \leq M_{pb}(R)||p||_{\infty}$ follows from the definition of $K_{(p)}$. Also, $||p(T_0)|| \leq M_{pb}(T_0)||p||_{\infty}$. Therefore,

$$||A_{(p)}|| \le M_{pb}(R) ||p||_{\infty} ||Z|| + ||Z||M_{pb}(T_0)||p||_{\infty} + M_{pb}(R)||p||_{\infty} ||Y_0||$$

= $((M_{pb}(R) + M_{pb}(T_0))||Z|| + M_{pb}(R)||Y_0||) ||p||_{\infty}.$

Since $||p(T)|| \leq \sqrt{3} \max(||p(V)||, ||A_{(p)}||, ||p(T_0)||)$, we obtain that

$$\|p(T)\| \le \sqrt{3} \max\left(M_{pb}(R), (M_{pb}(R) + M_{pb}(T_0))\|Z\| + M_{pb}(R)\|Y_0\|, M_{pb}(T_0)\right)\|p\|_{\infty}.$$

Thus, if R and T_0 are polynomially bounded, then T is polynomially bounded.

If we suppose only that R and T_0 are power bounded, then the proof of the power boundedness of T is the same.

Corollary 2.3. Suppose $\theta \in H^{\infty}$ is an inner function, $\mathcal{K}_{\theta} = H^2 \ominus \theta H^2$, $T_{\theta} = P_{\mathcal{K}_{\theta}}S|_{\mathcal{K}_{\theta}}, \ \mathcal{K}_{\theta} = P_{\theta H^2}S|_{\mathcal{K}_{\theta}}, \ \mathcal{H} \text{ is a Hilbert space, } T_0: \mathcal{H} \to \mathcal{H}, \ Y_0: \mathcal{H} \to \mathcal{K}_{\theta}, \ Z: \mathcal{H} \to \theta H^2 \text{ are operators. Moreover, suppose } Y_0T_0 = T_{\theta}Y_0.$ Put $A = S|_{\theta H^2}Z - ZT_0 + K_{\theta}Y_0,$

$$T = \begin{pmatrix} S|_{\theta H^2} & A \\ \mathbb{O} & T_0 \end{pmatrix} \quad and \quad Y = \begin{pmatrix} I_{\theta H^2} & Z \\ \mathbb{O} & Y_0 \end{pmatrix}$$

Then YT = SY. If Y_0 is a quasiaffinity, then Y is a quasiaffinity. If T_0 is polynomially bounded, then T is polynomially bounded. If T_0 is not similar to a contraction, then T is not similar to a contraction.

Proof. Put $\mathcal{M} = \theta H^2$, $V = S|_{\theta H^2}$, $\mathcal{K} = \mathcal{K}_{\theta}$, $K = K_{\theta}$, $R_0 = T_{\theta}$, and apply Proposition 2.2. Then R = S, therefore, YT = SY. By Proposition 2.2, the polynomially boundedness of T_0 implies the polynomially boundedness of T. If Y_0 is a quasiaffinity, then Y is a quasiaffinity by Lemma 2.1 (i). If Tis similar to a contraction, the same holds for the compression of T to any semi-invariant subspace, in particular, for T_0 . Therefore, if T_0 is not similar to a contraction, then T is not similar to a contraction. **Proposition 2.4.** Suppose $\theta \in H^{\infty}$ is an inner function, T is the operator from Corollary 2.3, and Y_0 from Corollary 2.3 is a quasiaffinity. Then $T \sim S$ if and only if there exist a quasiaffinity $X_0: \mathcal{K}_{\theta} \to \mathcal{H}$ and an outer function $g \in H^{\infty}$ such that $T_0X_0 = X_0T_{\theta}$ and $Y_0X_0 = g(T_{\theta})$.

Proof. The "if" part. Put $K_{(g)} = P_{\theta H^2}g(S)|_{\mathcal{K}_{\theta}}, W = K_{(g)} - ZX_0$, and

$$X = \begin{pmatrix} g(S)|_{\theta H^2} & W \\ \mathbb{O} & X_0 \end{pmatrix}.$$

Since g is outer, $g(S)|_{\theta H^2}$ is a quasiaffinity, and X is a quasiaffinity by Lemma 2.1 (i). It remains to prove that XS = TX. Writing this equality in matrix form, it is easy to see that it is sufficient to prove the equality

(2.1)
$$g(S)|_{\theta H^2} K_{\theta} + W T_{\theta} = S|_{\theta H^2} W + A X_0.$$

Using the definition of W and A we obtain that

$$g(S)|_{\theta H^2} K_{\theta} + WT_{\theta} = g(S)|_{\theta H^2} K_{\theta} + K_{(q)}T_{\theta} - ZX_0T_{\theta},$$

and

$$S|_{\theta H^2}W + AX_0 = S|_{\theta H^2}K_{(g)} - S|_{\theta H^2}ZX_0 + S|_{\theta H^2}ZX_0 - ZT_0X_0 + K_\theta Y_0X_0$$
$$= S|_{\theta H^2}K_{(g)} - ZT_0X_0 + K_\theta g(T_\theta).$$

Using the equality $T_0X_0 = X_0T_\theta$, we infer that (2.1) is equivalent to the equality

$$g(S)|_{\theta H^2} K_{\theta} + K_{(g)} T_{\theta} = S|_{\theta H^2} K_{(g)} + K_{\theta} g(T_{\theta}).$$

But this equality follows from the equality g(S)S = Sg(S) written in the matrix form.

The "only if" part. Suppose that a quasiaffinity Y is from Corollary 2.3, and $X: H^2 \to \theta H^2 \oplus \mathcal{H}$ is a quasiaffinity such that XS = TX. Since $YX \in \{S\}'$, there exists a function $g \in H^{\infty}$ such that YX = g(S), and, since YX is a quasiaffinity, g is outer. Writing $X = \begin{pmatrix} X_1 & W \\ X_2 & X_0 \end{pmatrix}$ with respect to the decompositions of the spaces $H^2 = \theta H^2 \oplus \mathcal{K}_{\theta}$ and $\theta H^2 \oplus \mathcal{H}$ and regarding the equality YX = g(S) with respect to these decompositions, we obtain that $Y_0X_0 = g(T_{\theta})$ and $Y_0X_2 = \mathbb{O}$. Since ker $Y_0 = \{0\}$, we conclude that $X_2 = \mathbb{O}$. Thus, $X = \begin{pmatrix} X_1 & W \\ \mathbb{O} & X_0 \end{pmatrix}$. By Lemma 2.1 (ii), $\operatorname{clos} X_0\mathcal{K}_{\theta} = \mathcal{H}$. Since ker $g(T_{\theta}) = \{0\}$, from the equality $Y_0X_0 = g(T_{\theta})$ we conclude that ker $X_0 =$ $\{0\}$. From the equality XS = TX we conclude that $X_0T_{\theta} = T_0X_0$.

Remark 2.5. The conditions $T_0X_0 = X_0T_\theta$ and $Y_0T_0 = T_\theta Y_0$ guarantee that $Y_0X_0 \in \{T_\theta\}'$, consequently, there exists a function $\varphi \in H^\infty$ such that $Y_0X_0 = \varphi(T_\theta)$. The condition that X_0 and Y_0 are quasiaffinities guarantees that the inner factor of φ is coprime with θ . For every $f \in H^\infty$ such that the inner factor of f is coprime with θ one can regard $f(T_\theta)Y_0$ instead of Y_0 , because $f(T_\theta)Y_0$ is a quasiaffinity which intertwines T_θ with T_0 . Also, for every $h \in H^\infty$ the equality $(f\varphi)(T_\theta) = (\varphi f + \theta h)(T_\theta)$ holds. But there exist functions φ and θ such that φ is coprime with θ and the function $\varphi f + \theta h$ is not outer for every $f, h \in H^\infty$, see [No] for (it seems the first) example of such functions φ and θ . Now we see that to construct a polynomially bounded operator T such that $T \prec S$ and T is not similar to a contraction, it is sufficient to construct a polynomially bounded operator T_0 such that $T_0 \sim T_\theta$ for some inner function θ , and T_0 is not similar to a contraction. This is done in Corollary 6.4. To construct a polynomially bounded operator T such that $T \sim S$ and T is not similar to a contraction, it is sufficient to construct a polynomially bounded operator T such that $T \sim S$ and T is not similar to a contraction, it is sufficient to construct a polynomially bounded operator T_0 such that $T_0 \sim T_\theta$ for some inner function θ , T_0 is not similar to a contraction, and there exist quasiaffinities X_0 and Y_0 and an outer function $g \in H^\infty$ such that $T_0 X_0 = X_0 T_\theta$, $Y_0 T_0 = T_\theta Y_0$, and $Y_0 X_0 = g(T_\theta)$. This is done in Theorem 7.1. The function g is from Lemma 3.4. The operator Z from Proposition 2.2 can be zero in both cases. The operator Z was considered to show that the choice of Z does not allow to avoid the condition on the existence of an *outer* function g such that $Y_0 X_0 = g(T_\theta)$ in this construction.

Put $\chi(z) = z, z \in \mathbb{T}$. Let U be the bilateral shift of multiplicity 1, that is, the operator of multiplication by χ on $L^2(\mathbb{T})$. U has the following form with respect to the decomposition $L^2(\mathbb{T}) = H^2 \oplus H^2_-$:

(2.2)
$$U = \begin{pmatrix} S & K \\ \mathbb{O} & S_* \end{pmatrix},$$

where S is the unilateral shift of multiplicity 1, $K: H_{-}^2 \to H^2$ acts by the formula $K\chi^{-n} = 0, n \ge 2, K\chi^{-1} = \chi^0, S_*: H_{-}^2 \to H_{-}^2$ acts by the formula $S_*\chi^{-n} = \chi^{-n+1}, n \ge 2, S_*\chi^{-1} = 0.$

The following propositions will be applied to construct an example of a polynomially bounded operator, which is not similar to a contraction and is quasisimilar to U.

Proposition 2.6. Suppose \mathcal{H} is a Hilbert space, and $T_1: \mathcal{H} \to \mathcal{H}$ and $Y_1: \mathcal{H}^2 \to \mathcal{H}$ are operators such that $Y_1S = T_1Y_1$. Put $x_0 = Y_1\chi^0$, $A: \mathcal{H}^2 \to \mathcal{H}$, $A\chi^{-1} = x_0$, $A\chi^{-n} = 0$, $n \geq 2$,

$$T = \begin{pmatrix} T_1 & A \\ \mathbb{O} & S_* \end{pmatrix} \quad and \quad Y = \begin{pmatrix} Y_1 & \mathbb{O} \\ \mathbb{O} & I_{H^2_-} \end{pmatrix}.$$

Then YU = TY.

If T_1 is power bounded, then T is power bounded. If T_1 is polynomially bounded, then T is polynomially bounded.

Proof. Clearly, $Y_1K = A$. From this equality and (2.2) we conclude that YU = TY. Note that

$$T^* = \begin{pmatrix} (S_*)^* & A^* \\ \mathbb{O} & T_1^* \end{pmatrix}$$

with respect to the decomposition $H^2_- \oplus \mathcal{H}$. Applying Proposition 2.2 with $T_0 = T_1^*$, $R_0 = S^*$, $V = (S_*)^*$, $Y_0 = Y_1^*$, $Z = \mathbb{O}$, K^* instead of K and A^* instead of A, and taking into account that $R = U^*$, we obtain the conclusion of Proposition 2.2 for T^* , and, consequently, for T.

Proposition 2.7. Suppose $g \in H^{\infty}$ is an outer function, \mathcal{H} is a Hilbert space, $T_1: \mathcal{H} \to \mathcal{H}$ is an operator, $Y_1: H^2 \to \mathcal{H}$ and $X_1: \mathcal{H} \to H^2$ are

quasiaffinities such that $Y_1S = T_1Y_1$, $X_1T_1 = SX_1$, and $X_1Y_1 = g(S)$. Let T and Y be defined as in Proposition 2.6. Put $K_{(q)} = P_{H^2}g(U)|_{H^2}$ and

$$X = \begin{pmatrix} X_1 & K_{(g)} \\ \mathbb{O} & g(S_*) \end{pmatrix}.$$

Then X and Y are quasiaffinities such that YU = TY, XT = UX, and XY = g(U).

Proof. Since g is outer, $g(S_*)$ is a quasiaffinity. By Lemma 2.1 (i), X and Y are quasiaffinities. The equality YU = TY is proved in Proposition 2.6. Using the matrix forms of U, T, and X, and the equality $X_1T_1 = SX_1$, it is easy to see that the equality XT = UX follows from the equality

(2.3)
$$X_1A + K_{(g)}S_* = SK_{(g)} + Kg(S_*)$$

We show that $X_1A = g(S)K$. Indeed, for $n \ge 2$ we have that $A\chi^{-n} = 0$ and $K\chi^{-n} = 0$. Furthermore, $X_1A\chi^{-1} = X_1x_0 = X_1Y_1\chi^0 = g$, and $g(S)K\chi^{-1} = g(S)\chi^0 = g$. To prove (2.3) it remains to note that $SK_{(g)} + Kg(S_*) = g(S)K + K_{(g)}S_*$, and this equality follows from the equality g(U)U = Ug(U) written in the matrix form. The equality XY = g(U)can be easily obtained from the definitions of X and Y and the matrix form of g(U).

Corollary 2.8. Suppose T_1 satisfies the conditions of Proposition 2.7, and T is defined in Proposition 2.7. If T_1 is polynomially bounded, then T is polynomially bounded. If T_1 is not similar to a contraction, then T is not similar to a contraction.

Proof. The assertion about polynomial boundedness is proved in Proposition 2.6. The assertion about similarity to a contraction follows from the same reasons as at the end of the proof of Corollary 2.3. \Box

The following lemma shows that there exist polynomially bounded operators that are quasisimilar to cyclic reductive unitaries and are not similar to contractions. A function g in the example will be $g(z) = \exp(-(\frac{1+z}{1-z})^{\alpha})$, $z \in \operatorname{clos} \mathbb{D} \setminus \{1\}, 0 < \alpha < 1$, see Lemma 3.4. The interested readers can find sets σ satisfying the condition of the lemma themselves.

Lemma 2.9. Suppose T is an operator, and X and Y are quasiaffinities such that YU = TY, and XT = UX. Since $XY \in \{U\}'$, there exists a function $g \in L^{\infty}$ such that XY = g(U). Let $\sigma \subset \mathbb{T}$ be a measurable set and let $\delta > 0$ be such that $|g| \ge \delta$ a.e. on σ . Put $\mathcal{M} = \operatorname{clos} YL^2(\sigma)$, where $L^2(\sigma) = \{h \in L^2 : h = 0 \text{ a.e. on } \mathbb{T} \setminus \sigma\}$. Then

(2.4)
$$T|_{\mathcal{M}} \approx U|_{L^2(\sigma)}$$

and $P_{\mathcal{M}^{\perp}}T|_{\mathcal{M}^{\perp}} \sim U|_{L^{2}(\mathbb{T}\setminus\sigma)}$. If T is power bounded and is not similar to a contraction, then $P_{\mathcal{M}^{\perp}}T|_{\mathcal{M}^{\perp}}$ is not similar to a contraction.

Proof. Clearly, $T\mathcal{M} \subset \mathcal{M}$, and

$$\operatorname{clos} X\mathcal{M} = \operatorname{clos} XYL^2(\sigma) = \operatorname{clos} g(U)L^2(\sigma) = L^2(\sigma).$$

Put

$$X_{\sigma} = X|_{\mathcal{M}} \colon \mathcal{M} \to L^2(\sigma) \quad \text{and} \quad Y_{\sigma} = Y|_{L^2(\sigma)} \colon L^2(\sigma) \to \mathcal{M}.$$

Then X_{σ} and Y_{σ} are quasiaffinities. Since $X_{\sigma}Y_{\sigma} = g(U)|_{L^{2}(\sigma)}$ is invertible, we conclude that X_{σ} and Y_{σ} are invertible. Since $X_{\sigma}T|_{\mathcal{M}} = U|_{L^{2}(\sigma)}X_{\sigma}$, (2.4) is proved.

Put $T_1 = P_{\mathcal{M}^{\perp}}T|_{\mathcal{M}^{\perp}}$, $X_1 = P_{L^2(\mathbb{T}\setminus\sigma)}X|_{\mathcal{M}^{\perp}}$, and $Y_1 = P_{\mathcal{M}^{\perp}}Y|_{L^2(\mathbb{T}\setminus\sigma)}$. It is easy to see that $X_1T_1 = U|_{L^2(\mathbb{T}\setminus\sigma)}X_1$ and $Y_1U|_{L^2(\mathbb{T}\setminus\sigma)} = T_1Y_1$. By Lemma 2.1 (iii), X_1 and Y_1 are quasiaffinities. By [B, Corollary 2.2] applied to T^* we have that $T \approx U|_{L^2(\sigma)} \oplus T_1$. If T_1 is similar to a contraction, we conclude that T is similar to a contraction. Also, to show that T is similar to a contraction, if T_1 is similar to a contraction, [C, Corollary 4.2] can be applied. \Box

3. Preliminaries: function theory

For $\lambda \in \mathbb{D}$ denote by b_{λ} a Blaschke factor: $b_{\lambda}(z) = \frac{|\lambda|}{\lambda} \frac{\lambda - z}{1 - \overline{\lambda} z}, z \in \mathbb{D}$. Note that

(3.1)
$$b_{\lambda} \circ \beta_w = \zeta_{w,\lambda} b_{\beta_w(\lambda)}, \text{ where } \zeta_{w,\lambda} \in \mathbb{T}.$$

Recall that β_w is defined in (1.3).

The following lemma is very simple, but useful.

Lemma 3.1. Let $\{B_N\}_N$ be a sequence of finite Blaschke products. Then there exists a sequence $\{r_N\}_N$ such that $0 < r_N < 1$ and for every sequence $\{w_N\}_N \subset \mathbb{D}$ such that $|w_N| \ge r_N$ there exists a sequence $\{\zeta_N\}_N \subset \mathbb{T}$ such that the product $\prod_N \zeta_N B_N \circ \beta_{w_N}$ converges.

Proof. Denote by Λ_N the set of zeros of B_N , then $B_N = \prod_{\lambda \in \Lambda_N} b_{\lambda}^{k_{\lambda}}$, where $1 \leq k_{\lambda} < \infty$ is the multiplicity of λ . For $w_N \in \mathbb{D}$ put $\zeta_N = \prod_{\lambda \in \Lambda_N} \overline{\zeta}_{w_N,\lambda}^{k_{\lambda}}$, where $\zeta_{w_N,\lambda}$ are from (3.1), then $\zeta_N B_N \circ \beta_{w_N} = \prod_{\lambda \in \Lambda_N} b_{\beta_{w_N}(\lambda)}^{k_{\lambda}}$. Let $\{\eta_N\}_N$ be a sequence such that $\eta_N > 0$ and $\sum_N \eta_N < \infty$. Since $|\beta_w(\lambda)| \to 1$ when $|w| \to 1$ and Λ_N is finite, there exists $0 < r_N < 1$ such that $\sum_{\lambda \in \Lambda_N} k_{\lambda}(1 - |\beta_w(\lambda)|) \leq \eta_N$ for $w \in \mathbb{D}$, $|w| \geq r_N$. Let $w_N \in \mathbb{D}$, and let $|w_N| \geq r_N$. Then

$$\sum_{N}\sum_{\lambda\in\Lambda_{N}}k_{\lambda}(1-|\beta_{w_{N}}(\lambda)|)\leq\sum_{N}\eta_{N}<\infty,$$

that is, the product $\prod_{N} \prod_{\lambda \in \Lambda_N} b_{\beta_{w_N}(\lambda)}^{k_\lambda} = \prod_{N} \zeta_N B_N \circ \beta_{w_N}$ converges. \Box

The following lemma is a particular case of [SFBK, VI.1.6]. Also, it can be proved in a straightforward manner.

Lemma 3.2. Let $w \in \mathbb{D}$. Put

$$U_w \colon H^2 \to H^2$$
, $(U_w h)(z) = \frac{(1 - |w|^2)^{1/2}}{1 - \overline{w}z} (h \circ \beta_w)(z)$, $z \in \mathbb{D}$, $h \in H^2$,

where β_w is defined in (1.3). Then U_w is unitary, $U_w = U_w^{-1}$, $U_w \mathcal{K}_{\theta \circ \beta_w} = \mathcal{K}_{\theta}$ and $U_w|_{\mathcal{K}_{\theta \circ \beta_w}} T_{\theta \circ \beta_w} = \beta_w(T_{\theta})U_w|_{\mathcal{K}_{\theta \circ \beta_w}}$ for every inner function $\theta \in H^{\infty}$, where the space \mathcal{K}_{θ} and the operator T_{θ} are defined in (1.7).

The following lemma can be easily proved by induction. Therefore, its proof is omitted.

Lemma 3.3. There exist functions $c_{nk} \colon \mathbb{D} \to \mathbb{C}$ such that $\sup_{\mathbb{D}} |c_{nk}| < \infty$ for every $n \ge 1$, $0 \le k \le n-1$, and for every analytic function $g \colon \mathbb{D} \to \mathbb{C}$ and every $w \in \mathbb{D}$

(3.2)
$$(g \circ \beta_w)^{(n)}(z) = \sum_{k=0}^{n-1} g^{(n-k)}(\beta_w(z)) \frac{c_{nk}(w)}{(1-\overline{w}z)^{2n-k}}.$$

Namely, $c_{10}(w) = |w|^2 - 1$, $c_{n+1,0}(w) = (|w|^2 - 1)c_{n0}(w)$,

$$c_{n+1,k}(w) = (|w|^2 - 1)c_{nk}(w) + (2n - k + 1)\overline{w}c_{n,k-1}(w), \quad 1 \le k \le n - 1,$$

 $c_{n+1,n}(w) = (n+1)\overline{w}c_{n,n-1}(w).$

We need the following lemmas.

Lemma 3.4. There exists an outer function $g \in H^{\infty}$ such that $g^{(n)}(r) \to 0$ when $r \in (0,1), r \to 1$, for every $n \ge 0$.

Proof. Let $0 < \alpha < 1$, put $g(z) = \exp(-(\frac{1+z}{1-z})^{\alpha})$, $z \in \mathbb{D}$. It is easy to prove by induction that

$$g^{(n)}(z) = g(z) \sum_{l=1}^{\kappa(n)} a_{nl} \left(\frac{1+z}{1-z}\right)^{\gamma_{nl}} (1-z)^{\eta_{nl}}, \quad z \in \mathbb{D},$$

where $\kappa(n) < \infty$, a_{nl} , γ_{nl} , $\eta_{nl} \in \mathbb{R}$. Therefore, $g^{(n)}(r) \to 0$ when $r \in (0, 1)$, $r \to 1$, for every $n \ge 0$. In [Ni, Example I.A.4.3.7, p.71] it is proved that $g \in H^{\infty}$ and g is outer.

Lemma 3.5. Suppose $\Lambda \subset \mathbb{D}$ is finite, $1 \leq k_{\lambda} < \infty$ for every $\lambda \in \Lambda$, and $B = \prod_{\lambda \in \Lambda} b_{\lambda}^{k_{\lambda}}$. Then there exists C > 0 which depends on B such that

dist
$$(\varphi, BH^{\infty}) \leq C \max_{\lambda \in \Lambda, 0 \leq k \leq k_{\lambda} - 1} |\varphi^{(k)}(\lambda)|$$
 for every $\varphi \in H^{\infty}$.

Proof. Recall that for the Blaschke product B the space \mathcal{K}_B and the operator T_B on \mathcal{K}_B are defined by (1.7). Put $u_{\lambda k}(z) = \frac{B(z)}{(z-\lambda)^k}$, $z \in \mathbb{D}$, $\lambda \in \Lambda$, $1 \leq k \leq k_{\lambda}$. Then $\{u_{\lambda k}\}_{\lambda \in \Lambda, 1 \leq k \leq k_{\lambda}}$ is a basis of \mathcal{K}_B , and, since dim $\mathcal{K}_B < \infty$, there exist an invertible operator X on \mathcal{K}_B and an orthonormal basis $\{e_{\lambda k}\}_{\lambda \in \Lambda, 1 \leq k \leq k_{\lambda}}$ of \mathcal{K}_B such that $Xu_{\lambda k} = e_{\lambda k}$, $\lambda \in \Lambda$, $1 \leq k \leq k_{\lambda}$. Let $\varphi \in H^{\infty}$. Then

$$X\varphi(T_B)X^{-1}e_{\lambda k} = \sum_{l=0}^{k-1} \frac{\varphi^{(l)}(\lambda)}{l!} e_{\lambda,k-l}, \quad \lambda \in \Lambda, \quad 1 \le k \le k_{\lambda},$$

therefore, there exists $C_1 > 0$ which depends on $\sum_{\lambda \in \Lambda} k_{\lambda}$ only such that $||X\varphi(T_B)X^{-1}|| \leq C_1 \max_{\lambda \in \Lambda, 0 \leq k \leq k_{\lambda} - 1} |\varphi^{(k)}(\lambda)|$. Thus,

$$\|\varphi(T_B)\| \le \|X\| \|X^{-1}\| C_1 \max_{\lambda \in \Lambda, 0 \le k \le k_\lambda - 1} |\varphi^{(k)}(\lambda)|.$$

By Nehari's theorem, $dist(\varphi, BH^{\infty}) = \|\varphi(T_B)\|$.

Lemma 3.6. Suppose B is a finite Blaschke product, and $a, \gamma \in (0, 1)$. Then there exists $r \in (0, 1)$ such that $|(B \circ \beta_w)(z)| \ge \gamma$ for z, $w \in \mathbb{D}$, $|z| \le a$, $r \le |w|$, where β_w is defined in (1.3).

Proof. Set $B = \prod_{\lambda \in \Lambda} b_{\lambda}^{k_{\lambda}}$, where $\Lambda \subset \mathbb{D}$ is a finite set, and $1 \leq k_{\lambda} < \infty$ for every $\lambda \in \Lambda$. Put $\kappa = \sum_{\lambda \in \Lambda} k_{\lambda}$.

It is easy to see from (3.1) that

$$1 - |(b_{\lambda} \circ \beta_w)(z)|^2 = \frac{(1 - |\beta_w(\lambda)|^2)(1 - |z|^2)}{|1 - \overline{\beta}_w(\lambda)z|^2}$$

From this equality and the relations $|\beta_w(\lambda)| \to 1$ when $w \in \mathbb{D}$, $|w| \to 1$, we have that $|(b_{\lambda} \circ \beta_w)(z)| \to 1$ when $w \in \mathbb{D}, |w| \to 1$ uniformly on $\{z : |z| \le a\}$ for every $\lambda \in \mathbb{D}$. Therefore, for every $\lambda \in \mathbb{D}$ there exists $r_{\lambda} \in (0, 1)$ such that $|(b_{\lambda} \circ \beta_w)(z)| \geq \gamma^{1/\kappa}$ for $z, w \in \mathbb{D}, |z| \leq a, r_{\lambda} \leq |w|$. Put $r = \max_{\lambda \in \Lambda} r_{\lambda}$. Then

$$|(B \circ \beta_w)(z)| = \prod_{\lambda \in \Lambda} |(b_\lambda \circ \beta_w)(z)|^{k_\lambda} \ge \prod_{\lambda \in \Lambda} (\gamma^{1/\kappa})^{k_\lambda} = \gamma$$

 $\in \mathbb{D}, |z| \le a, r \le |w|.$

Proposition 3.7. Suppose C > 0, $\Lambda \subset (0,1)$ is finite, $1 \leq k_{\lambda} < \infty$ for every $\lambda \in \Lambda$, $B = \prod_{\lambda \in \Lambda} b_{\lambda}^{k_{\lambda}}$, φ , $g \in H^{\infty}$, $\varphi(\lambda) \neq 0$ for every $\lambda \in \Lambda$, and $g^{(k)}(r) \to 0$ when $r \in (0,1), r \to 1$, for every $k \ge 0$. Then there exists r_0 , $0 < r_0 < 1$, such that for every $w \in [r_0, 1)$ there exists $f \in H^{\infty}$ such that $g \circ \beta_w - f \varphi \in BH^{\infty}$ and $\operatorname{dist}(f, BH^{\infty}) \leq C$, where β_w is defined in (1.3).

Proof. We have $g \circ \beta_w - f\varphi \in BH^\infty$ if and only if

(3.3)
$$(g \circ \beta_w)^{(n)}(\lambda) = (f\varphi)^{(n)}(\lambda)$$
 for every $\lambda \in \Lambda$, $0 \le n \le k_\lambda - 1$.

Put

for z, w

$$\varphi_{\lambda nk} = \frac{n!}{k!(n-k)!} \varphi^{(n-k)}(\lambda), \quad 0 \le k \le n, \quad 0 \le k \le k_{\lambda} - 1,$$
$$\varphi_{\lambda nk} = 0, \quad 0 \le n \le k - 1 \le k_{\lambda} - 1.$$

Then (3.3) is equivalent to

(3.4)
$$\{\varphi_{\lambda nk}\}_{n,k=0}^{k_{\lambda}-1} \cdot \{f^{(k)}(\lambda)\}_{k=0}^{k_{\lambda}-1} = \{(g \circ \beta_w)^{(n)}(\lambda)\}_{n=0}^{k_{\lambda}-1}.$$

Since $\{\varphi_{\lambda nk}\}_{n,k=0}^{k_{\lambda}-1}$ is a lower triangular matrix, and $\varphi_{\lambda nn} = \varphi(\lambda), 0 \leq n \leq \infty$ $k_{\lambda} - 1$, det $\{\varphi_{\lambda nk}\}_{n,k=0}^{k_{\lambda}-1} = \varphi(\lambda)^{k_{\lambda}} \neq 0$ for $\lambda \in \Lambda$, therefore, the matrix $\{\varphi_{\lambda nk}\}_{n,k=0}^{k_{\lambda}-1}$ is invertible. Put $\Phi_{\lambda} = (\{\varphi_{\lambda nk}\}_{n,k=0}^{k_{\lambda}-1})^{-1}$, then (3.4) is equivalent to

(3.5)
$$\{f^{(k)}(\lambda)\}_{k=0}^{k_{\lambda}-1} = \Phi_{\lambda} \cdot \{(g \circ \beta_w)^{(n)}(\lambda)\}_{n=0}^{k_{\lambda}-1}.$$

Since Λ is a finite set and $k_{\lambda} < \infty$, for every $w \in \mathbb{D}$ there exists a function $f \in H^{\infty}$ which satisfies (3.5). By Lemma 3.5, there exists C_1 which depends on B only such that $\operatorname{dist}(f, BH^{\infty}) \leq C_1 \max_{\lambda \in \Lambda, 0 \leq k \leq k_{\lambda}-1} |f^{(k)}(\lambda)|$. Since $\Lambda \subset (0,1), \ \beta_w(\lambda) \in (-1,1), \text{ if } w \in (0,1), \text{ and } \beta_w(\lambda) \to 1 \text{ when } w \in (0,1),$ $w \to 1$. From (3.2) and the condition on g we conclude that

$$\|\{(g \circ \beta_w)^{(n)}(\lambda)\}_{n=0}^{k_{\lambda}-1}\| \to 0 \text{ when } w \to 1, w \in (0,1),$$

for every $\lambda \in \Lambda$. Therefore, we infer from (3.5) that for every $\lambda \in \Lambda$ there exists $r_{\lambda} \in (0,1)$ such that $\|\{f^{(k)}(\lambda)\}_{k=0}^{k_{\lambda}-1}\| \leq C/C_1$, if $w \in [r_{\lambda}, 1)$. Now $r_0 = \max_{\lambda \in \Lambda} r_{\lambda} < 1$ satisfies to the conclusion of the proposition. \Box

The following theorem is the main result of this section. The condition (3.6) from the theorem is the *generalized Carleson condition*, see [Ni, Theorem II.C.3.2.14, p.164].

Recall that β_w is defined in (1.3).

Theorem 3.8. Suppose C > 0, $g \in H^{\infty}$, and $g^{(k)}(r) \to 0$ when $r \in (0,1)$, $r \to 1$, for every $k \geq 0$. Furthermore, suppose B_N are finite Blaschke products with zeros from (0,1), $\varphi_N \in H^{\infty}$, $\varphi_N(\lambda) \neq 0$ for every $\lambda \in \mathbb{D}$ such that $B_N(\lambda) = 0$, for every index N. Then there exist $\delta > 0$ and sequences of $w_N \in (0,1)$, of $\zeta_N \in \mathbb{T}$, and of $\psi_N \in H^{\infty}$ such that the product $\prod_N \zeta_N B_N \circ \beta_{w_N}$ converges,

(3.6)
$$\left|\prod_{N} \zeta_{N}(B_{N} \circ \beta_{w_{N}})(z)\right| \geq \delta \inf_{N} \left|(B_{N} \circ \beta_{w_{N}})(z)\right| \text{ for every } z \in \mathbb{D},$$

$$g - \psi_N \cdot \varphi_N \circ \beta_{w_N} \in (B_N \circ \beta_{w_N}) H^{\infty}$$
, and $\operatorname{dist}(\psi_N, (B_N \circ \beta_{w_N}) H^{\infty}) \leq C$.

Proof. Denote by $\{r_{1N}\}_N$ a sequence from Lemma 3.1 applied to $\{B_N\}_N$, and by r_{2N} a quantity from Proposition 3.7 applied to C, B_N , φ_N , and g. Put $r_N = \max(r_{1N}, r_{2N})$.

Let $\gamma_n \in (0,1)$ be such that $\delta = \prod_n \gamma_n$ converges, put $\delta_N = \prod_{n=1}^N \gamma_n$. We construct the sequence $\{w_N\}_N$ such that $w_N \in [r_N, 1)$ by induction. Let $w_1 \in [r_1, 1)$ be arbitrary. Clearly, $|(B_1 \circ \beta_{w_1})(z)| \ge \delta_1 |(B_1 \circ \beta_{w_1})(z)|$ for every $z \in \mathbb{D}$. Suppose that $w_n \in [r_n, 1), 1 \le n \le N$, are such that (3.7)

$$|\theta_N(z)| \ge \delta_N \inf_{1 \le n \le N} |(B_n \circ \beta_{w_n})(z)| \quad \text{for every } z \in \mathbb{D}, \text{ where } \theta_N = \prod_{n=1}^N B_n \circ \beta_{w_n}$$

Since θ_N is a finite Blaschke product, there exists $a \in (0, 1)$ such that

(3.8)
$$|\theta_N(z)| \ge \gamma_{N+1}$$
 for every $z \in \mathbb{D}, |z| \ge a$

We have from (3.8) that

(3.9)
$$\inf_{1 \le n \le N} |(B_n \circ \beta_{w_n})(z)| \ge \gamma_{N+1} \quad \text{for every} \quad z \in \mathbb{D}, \quad |z| \ge a.$$

By Lemma 3.6, there exists $w_{N+1} \in [r_{N+1}, 1)$ such that

$$(3.10) \qquad |(B_{N+1} \circ \beta_{w_{N+1}})(z)| \ge \gamma_{N+1} \quad \text{for every } z \in \mathbb{D}, \quad |z| \le a.$$

We show that (2,11)

$$|(\partial_N \cdot B_{N+1} \circ \beta_{w_{N+1}})(z)| \ge \gamma_{N+1} \delta_N \inf_{1 \le n \le N+1} |(B_n \circ \beta_{w_n})(z)| \quad \text{for every } z$$

We consider four cases.

First case: $a \leq |z| < 1$, $\inf_{1 \leq n \leq N+1} |(B_n \circ \beta_{w_n})(z)| = \inf_{1 \leq n \leq N} |(B_n \circ \beta_{w_n})(z)|$. By (3.7) and (3.9),

$$\begin{aligned} |(\theta_N \cdot B_{N+1} \circ \beta_{w_{N+1}})(z)| &\geq \gamma_{N+1} \delta_N \inf_{1 \leq n \leq N} |(B_n \circ \beta_{w_n})(z)| \\ &= \gamma_{N+1} \delta_N \inf_{1 < n \leq N+1} |(B_n \circ \beta_{w_n})(z)|. \end{aligned}$$

Second case: $a \leq |z| < 1$, $\inf_{1 \leq n \leq N+1} |(B_n \circ \beta_{w_n})(z)| = |(B_{N+1} \circ \beta_{w_{N+1}})(z)|$. By (3.8),

$$|(\theta_N \cdot B_{N+1} \circ \beta_{w_{N+1}})(z)| \ge \gamma_{N+1} |(B_{N+1} \circ \beta_{w_{N+1}})(z)| = \gamma_{N+1} \inf_{1 \le n \le N+1} |(B_n \circ \beta_{w_n})(z)| \le \gamma_{N+1} |(B_N \circ \beta_{w_{N+1}})(z)| \le \gamma_{N+1} |(B_N \circ \beta_{w_{N+1}})(z)$$

N

 $\in \mathbb{D}.$

Third case: $|z| \leq a$, $\inf_{1 \leq n \leq N+1} |(B_n \circ \beta_{w_n})(z)| = \inf_{1 \leq n \leq N} |(B_n \circ \beta_{w_n})(z)|$. By (3.7) and (3.10),

$$|(\theta_N \cdot B_{N+1} \circ \beta_{w_{N+1}})(z)| \ge |(B_{N+1} \circ \beta_{w_{N+1}})(z)|\delta_N \inf_{1 \le n \le N} |(B_n \circ \beta_{w_n})(z)|$$

$$\geq \gamma_{N+1}\delta_N \inf_{1\leq n\leq N} |(B_n \circ \beta_{w_n})(z)| = \gamma_{N+1}\delta_N \inf_{1\leq n\leq N+1} |(B_n \circ \beta_{w_n})(z)|$$

Fourth case: $|z| \leq a$, $\inf_{1 \leq n \leq N+1} |(B_n \circ \beta_{w_n})(z)| = |(B_{N+1} \circ \beta_{w_{N+1}})(z)|$. By (3.7) and (3.10),

$$|(\theta_N \cdot B_{N+1} \circ \beta_{w_{N+1}})(z)| \ge |(B_{N+1} \circ \beta_{w_{N+1}})(z)|\delta_N \inf_{1 \le n \le N} |(B_n \circ \beta_{w_n})(z)|$$

$$\ge |(B_{N+1} \circ \beta_{w_{N+1}})(z)|\delta_N \gamma_{N+1} = \gamma_{N+1}\delta_N \inf_{1 \le n \le N+1} |(B_n \circ \beta_{w_n})(z)|.$$

Since $\delta_{N+1} = \gamma_{N+1}\delta_N$ and $0 < \delta_N < 1$, the relation (3.11) is proved. Thus, (3.7) is proved by induction for all N. Let $N \to \infty$ in both parts of (3.7), then (3.6) follows. By Lemma 3.1, there exists a sequence $\zeta_N \in \mathbb{T}$ such that the product $\prod_N \zeta_N B_N \circ \beta_{w_N}$ converges.

Let f_N be functions from Proposition 3.7 applied to C, B_N , φ_N , and g, with $w = w_N$. Put $\psi_N = f_N \circ \beta_{w_N}$. Then $\psi_N \in H^{\infty}$, $g - \psi_N \cdot \varphi_N \circ \beta_{w_N} \in (B_N \circ \beta_{w_N}) H^{\infty}$ and $\operatorname{dist}(\psi_N, (B_N \circ \beta_{w_N}) H^{\infty}) = \operatorname{dist}(f_N, B_N H^{\infty}) \leq C$. \Box

4. Preliminaries: Jordan Operators

For $N \geq 1$ and $\lambda \in \mathbb{D}$ define the operator $T_{N,\lambda} \colon \mathbb{C}^{N+1} \to \mathbb{C}^{N+1}$ acting by the formula $T_{N,\lambda}e_0 = \lambda e_0$, $T_{N,\lambda}e_n = \lambda e_n + e_{n-1}$, $1 \leq n \leq N$, where $\{e_n\}_{n=0}^N$ is an orthonormal basis of \mathbb{C}^{N+1} . It is well known that $T_{N,\lambda} \approx T_{b_{\lambda}^{N+1}}$, but $\|T_{N,\lambda}\| \leq 1$ if and only if $\lambda = 0$.

Lemma 4.1. Let $N \geq 1$, and let $\varepsilon > 0$. Then there exists $\delta > 0$ such that for every $\lambda \in \mathbb{D}$, $|\lambda| < \delta$, there exists an operator $X : \mathbb{C}^{N+1} \to \mathcal{K}_{b_{\lambda}^{N+1}}$ such that $XT_{N,\lambda} = T_{b_{\lambda}^{N+1}}X$, $||X|| \leq 1 + \varepsilon$, and $||X^{-1}|| \leq 1 + \varepsilon$, where the space $\mathcal{K}_{b_{\lambda}^{N+1}}$ and the operator $T_{b_{\lambda}^{N+1}}$ are defined in (1.7).

Proof. Put

$$h_n(z) = b_{\lambda}^{N+1}(z) \frac{(1-|\lambda|^2)^{1/2}(1-\overline{\lambda}z)^n}{(z-\lambda)^{n+1}}, \ z \in \mathbb{D}, \text{ and } c_{nk} = \frac{n!}{k!(n-k)!} \frac{\overline{\lambda}^{n-k}}{(1-|\lambda|^2)^n},$$

 $0 \le k \le n, \ 0 \le n \le N$. Then $\{h_n\}_{n=0}^N$ is an orthonormal basis of $\mathcal{K}_{b_{\lambda}^{N+1}}$. Put

$$X \colon \mathbb{C}^{N+1} \to \mathcal{K}_{b_{\lambda}^{N+1}}, \quad Xe_n = \sum_{k=0}^n c_{nk}h_k.$$

Then X has the upper triangular form in the orthonormal bases $\{e_n\}_{n=0}^N$ and $\{h_n\}_{n=0}^N$. Also, $XT_{N,\lambda} = T_{b_{\lambda}^{N+1}}X$. To see that, put $u_n = Xe_n$, then $u_n(z) = b_{\lambda}^{N+1}(z)\frac{(1-|\lambda|^2)^{1/2}}{(z-\lambda)^{n+1}}, z \in \mathbb{D}$, therefore, $T_{b_{\lambda}^{N+1}}u_0 = \lambda u_0, T_{b_{\lambda}^{N+1}}u_n = \lambda u_n + u_{n-1}, 1 \leq n \leq N$.

Let D be a diagonal matrix with the elements c_{nn} , $0 \le n \le N$, put A = X - D, then $(D^{-1}A)^{N+1} = \mathbb{O}$, therefore, $X^{-1} = \sum_{n=0}^{N} (-1)^n (D^{-1}A)^n D^{-1}$. Also, $\|D\| = \frac{1}{(1-|\lambda|^2)^N}$, and $\|D^{-1}\| = 1$. If $0 \le k \le n-1$, then $c_{nk} \to 0$ when $|\lambda| \to 0$, and, since A is a finite matrix with elements c_{nk} and 0, then
$$\begin{split} \|A\| &\to 0 \text{ when } |\lambda| \to 0. \text{ Since } \|X\| \leq \|D\| + \|A\| \text{ and } \|X^{-1}\| \leq \sum_{n=0}^{N} \|A\|^{n}, \\ \|X\| \to 1 \text{ and } \|X^{-1}\| \to 1 \text{ when } |\lambda| \to 0. \text{ Therefore, there exists } \delta > 0 \text{ such that if } |\lambda| < \delta \text{ then } \|X\| \leq 1 + \varepsilon, \text{ and } \|X^{-1}\| \leq 1 + \varepsilon. \end{split}$$

Suppose \mathcal{H} is a Hilbert space, and $N \geq 1$. Define the shift operator $\mathbf{S}_{\mathcal{H}N}$ on $\bigoplus_{n=0}^{N} \mathcal{H}$ as following:

(4.1)
$$\mathbf{S}_{\mathcal{H}N} = \begin{pmatrix} \mathbb{O} & I_{\mathcal{H}} & \dots & \mathbb{O} & \mathbb{O} \\ \dots & \dots & \dots & \dots & \dots \\ \mathbb{O} & \mathbb{O} & \dots & \mathbb{O} & I_{\mathcal{H}} \\ \mathbb{O} & \mathbb{O} & \dots & \mathbb{O} & \mathbb{O} \end{pmatrix}.$$

Note that $\mathbf{S}_{\mathcal{H}N}^{N+1} = \mathbb{O}$, and $\mathbf{S}_{\mathcal{H}N}^N \neq \mathbb{O}$. Also, $\mu_{\mathbf{S}_{\mathcal{H}N}} = \dim \mathcal{H}$.

Corollary 4.2. Suppose \mathcal{H} is a Hilbert space, $\mathbf{d} = \dim \mathcal{H} < \infty$, $\{e_j\}_{j=1}^{\mathbf{d}}$ is an orthonormal basis of \mathcal{H} , $\{\lambda_j\}_{j=1}^{\mathbf{d}} \subset \mathbb{D}$, the operator $D: \mathcal{H} \to \mathcal{H}$ acts by the formula $De_j = \lambda_j e_j$, $1 \leq j \leq \mathbf{d}$. Furthermore, suppose $N \geq 1$, and put $T = \bigoplus_{n=0}^{N} D + \mathbf{S}_{\mathcal{H}N}$. Then $T \approx \bigoplus_{j=1}^{\mathbf{d}} T_{b_{\lambda_j}^{N+1}}$. Moreover, let $\varepsilon > 0$, and let δ be from Lemma 4.1 applied to N and ε . If $|\lambda_j| < \delta$ for every $1 \leq j \leq \mathbf{d}$, then there exists an operator $X: \bigoplus_{n=0}^{N} \mathcal{H} \to \bigoplus_{j=1}^{\mathbf{d}} \mathcal{K}_{b_{\lambda_j}^{N+1}}$ such that $XT = (\bigoplus_{j=1}^{\mathbf{d}} T_{b_{\lambda_j}^{N+1}})X$, $||X|| \leq 1 + \varepsilon$, and $||X^{-1}|| \leq 1 + \varepsilon$.

Proof. We have $\mathcal{H} = \bigoplus_{j=1}^{\mathbf{d}} \mathbb{C}e_j$, therefore, $\bigoplus_{n=0}^{N} \mathcal{H} = \bigoplus_{j=1}^{\mathbf{d}} \bigoplus_{n=0}^{N} \mathbb{C}e_j$. The spaces $\bigoplus_{n=0}^{N} \mathbb{C}e_j$ are invariant for T, and $T|_{\bigoplus_{n=0}^{N} \mathbb{C}e_j} = T_{N,\lambda_j}$. Thus, $T = \bigoplus_{j=1}^{\mathbf{d}} T_{N,\lambda_j}$, and it remains to apply Lemma 4.1.

Lemma 4.3. Suppose \mathcal{H} is a Hilbert space, $\mathbf{d} = \dim \mathcal{H} < \infty$, $\{e_j\}_{j=1}^{\mathbf{d}}$ is an orthonormal basis of \mathcal{H} , $\{\lambda_j\}_{j=1}^{\mathbf{d}}$, $\{\nu_j\}_{j=1}^{\mathbf{d}} \subset \mathbb{D}$, the operators $D: \mathcal{H} \to \mathcal{H}$, $D_{\star}: \mathcal{H} \to \mathcal{H}$ act by the formulas $De_j = \lambda_j e_j$, $1 \leq j \leq \mathbf{d}$, $D_{\star}e_j = \nu_j e_j$, $1 \leq j \leq \mathbf{d}$. Let $N \geq 1$, and let $A: \bigoplus_{n=0}^{N} \mathcal{H} \to \bigoplus_{n=0}^{N} \mathcal{H}$ be an arbitrary operator. Put $B = \prod_{j=1}^{\mathbf{d}} b_{\lambda_j}^{N+1} b_{\nu_j}^{N+1}$. Put $T_0 = \bigoplus_{n=0}^{N} \mathcal{D} + \mathbf{S}_{\mathcal{H}N}$ and $T_1 = \bigoplus_{n=0}^{N} D_{\star} + \mathbf{S}_{\mathcal{H}N}^{*}$. Define the operator T on the space $\bigoplus_{n=0}^{N} \mathcal{H} \oplus \bigoplus_{n=0}^{N} \mathcal{H}$ as follows:

$$T = \begin{pmatrix} T_1 & A \\ \mathbb{O} & T_0 \end{pmatrix}.$$

If $\lambda_j \neq \lambda_k$, $\nu_j \neq \nu_k$ for $j \neq k$, and $\lambda_j \neq \nu_k$ for every $1 \leq j, k \leq \mathbf{d}$, then $T \approx T_B$, where T_B is defined in (1.7).

Proof. Put $p(z) = \prod_{j=1}^{\mathbf{d}} (z - \lambda_j)^{N+1}$. Then $p(T_0) = \mathbb{O}$, $p(T_1)$ has bounded inverse, because $\lambda_j \neq \nu_k$ for every $1 \leq j, k \leq \mathbf{d}$ and T_1 acts on a finite dimensional space, and

$$p(T) = \begin{pmatrix} p(T_1) & A_0 \\ \mathbb{O} & \mathbb{O} \end{pmatrix},$$

where A_0 is an appropriate operator. Put $Y_0 = p(T_1)^{-1}A_0$ and $Y = \begin{pmatrix} I & Y_0 \\ \mathbb{O} & I \end{pmatrix}$. The equality $YT = (T_1 \oplus T_0)Y$ follows from the definition of Y and the equality p(T)T = Tp(T) writing in matrix form. Thus, $T \approx T_1 \oplus T_0$.

By Corollary 4.2, $T_0 \approx \bigoplus_{j=1}^{\mathbf{d}} T_{b_{\lambda_j}^{N+1}}$ and $T_1 \approx \bigoplus_{j=1}^{\mathbf{d}} T_{b_{\nu_j}^{N+1}}$. Since *B* is a finite Blaschke product and all λ_j , ν_k are pairwise distinct, we conclude that $T \approx T_B$.

5. Preliminaries: Foguel–Hankel operators with truncated shifts

We introduce the following notation (see [DP], Sec. 1.3]): $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ $\begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix}$

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{C} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{I}_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{D}(a,c) = \begin{pmatrix} a & 0 \\ 0 & c \end{pmatrix}, \quad a,c \in \mathbb{C}.$$

For $N > 1$ put

(5.1)
$$\mathbf{C}_{Nj} = \mathbf{V}^{\otimes j} \otimes \mathbf{C} \otimes \mathbf{I}_2^{\otimes N-j}, \quad 0 \le j \le N,$$

and for the families $\{a_l\}_{l=0}^N$, $\{c_l\}_{l=0}^N \subset \mathbb{C}$ put

(5.2)
$$\mathbf{D}(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N) = \mathbf{D}(a_0, c_0) \otimes \ldots \otimes \mathbf{D}(a_N, c_N)$$

and

Here $A^{\otimes j}$ denotes the tensor product of j copies of A. \mathbf{C}_{Nj} , $\mathbf{D}(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)$ and $\mathbf{D}_j(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)$ are $2^{N+1} \times 2^{N+1}$ matrices, that is, operators on $\mathbb{C}^{2^{N+1}}$, and $\mathbf{D}(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)$ and $\mathbf{D}_j(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)$, $0 \leq j \leq N$, are diagonal with respect to the standard basis in $\mathbb{C}^{2^{N+1}}$. From the equalities

 $\mathbf{VD}(a,c) = \mathbf{D}(a,c)\mathbf{V}, \quad \mathbf{I}_2\mathbf{D}(a,c) = \mathbf{D}(a,c)\mathbf{I}_2 \text{ and } \mathbf{CD}(a,c) = \mathbf{D}(c,a)\mathbf{C}$ we conclude that

(5.4)
$$\mathbf{C}_{Nj}\mathbf{D}(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N) = \mathbf{D}_j(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)\mathbf{C}_{Nj}, \quad 0 \le j \le N.$$

Lemma 5.1. For every $0 < \delta < 1$ there exist families $\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N \subset (0, \delta^{1/(N+1)})$ such that the elements of a diagonal matrix $\mathbf{D}(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)$ are from $(0, \delta)$ and are pairwise distinct.

Proof. The elements of a diagonal matrix $\mathbf{D}(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)$ are the products of N+1 factors, each of which is equal to a_l or c_l for some $l, 0 \leq l \leq N$. Therefore, if $0 < a_l < \delta^{1/(N+1)}$ and $0 < c_l < \delta^{1/(N+1)}$ for $0 \leq l \leq N$, then the elements of a diagonal matrix $\mathbf{D}(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)$ are from $(0, \delta)$. The choice of $\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N \subset (0, \delta^{1/(N+1)})$ is such that the elements of a diagonal matrix $\mathbf{D}(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)$ are pairwise distinct by induction. Page of inductions $a_l \leq a_l$

The choice of $\{a_l\}_{l=0}^N$, $\{c_l\}_{l=0}^N \subset (0, \delta^{1/(N+1)})$ is such that the elements of a diagonal matrix $\mathbf{D}(\{a_l\}_{l=0}^N, \{c_l\}_{l=0}^N)$ are pairwise distinct by induction. Base of induction: $a_0 \neq c_0$. Suppose $1 \leq n \leq N$, and suppose that $\{a_l\}_{l=0}^n$, $\{c_l\}_{=0}^n \subset (0, \delta^{1/(N+1)})$ are such that the elements $\{d_k\}_{k=1}^{2^{n+1}}$ of the matrix $\mathbf{D}(a_0, c_0) \otimes \ldots \otimes \mathbf{D}(a_n, c_n)$ are pairwise distinct. Let a_{n+1} , $c_{n+1} \subset (0, \delta^{1/(N+1)})$ be such that $c_{n+1} < a_{n+1}d_k$ for all $k, 1 \leq k \leq 2^{n+1}$. The elements of $\mathbf{D}(a_0, c_0) \otimes \ldots \otimes \mathbf{D}(a_{n+1}, c_{n+1})$ are $a_{n+1}d_k$ and $c_{n+1}d_k$, $1 \leq k \leq 2^{n+1}$. It is easy to see that $\{a_l\}_{l=0}^{n+1}$ and $\{c_l\}_{l=0}^{n+1}$ satisfy the inductional assumption again. \Box

Suppose \mathcal{H} is a Hilbert space, $N \geq 1$, and $A_j: \mathcal{H} \to \mathcal{H}, 0 \leq j \leq N$, are operators. Define a Hankel operator $\Gamma(\{A_j\}_{j=0}^N)$ on $\bigoplus_{j=0}^N \mathcal{H}$ as follows:

(5.5)
$$\Gamma(\{A_j\}_{j=0}^N) = \begin{pmatrix} \mathbb{O} & \mathbb{O} & \dots & \mathbb{O} & A_N \\ \mathbb{O} & \mathbb{O} & \dots & A_N & A_{N-1} \\ \dots & \dots & \dots & \dots & \dots \\ \mathbb{O} & A_N & \dots & A_{N-1} & A_1 \\ A_N & A_{N-1} & \dots & A_1 & A_0 \end{pmatrix}.$$

Put $\Gamma(k, \{A_j\}_{j=0}^N) = \Gamma(A_k, \dots, A_N, \mathbb{O}, \dots, \mathbb{O}), 0 \le k \le N$. Then $\mathbf{S}_{\mathcal{H}N}^* \Gamma(\{A_j\}_{j=0}^N) = \Gamma(\{A_j\}_{j=0}^N) \mathbf{S}_{\mathcal{H}N} \text{ and } \Gamma(\{A_j\}_{j=0}^N) \mathbf{S}_{\mathcal{H}N}^k = \Gamma(k, \{A_j\}_{j=0}^N),$ where $\mathbf{S}_{\mathcal{H}N}$ is defined in (4.1).

Define the operator $Q_N(\{A_j\}_{j=0}^N)$ on $\bigoplus_{i=0}^N \mathcal{H} \oplus \bigoplus_{i=0}^N \mathcal{H}$ as follows:

(5.7)
$$Q_N(\{A_j\}_{j=0}^N) = \begin{pmatrix} \mathbf{S}_{\mathcal{H}N}^* & \Gamma(\{A_j\}_{j=0}^N) \\ \mathbb{O} & \mathbf{S}_{\mathcal{H}N} \end{pmatrix}.$$

Operators from (5.7) are analogs of Foguel-Hankel operators (truncated shifts are used in the construction instead of the forward and backward shifts). We will call such operators "truncated" Foguel-Hankel operators in this paper. It is easy to see that

$$(Q_N(\{A_j\}_{j=0}^N))^{N+2} = \mathbb{O}$$
 and $(Q_N(\{A_j\}_{j=0}^N))^N \neq \mathbb{O}$

(Note that $(Q_N(\{A_j\}_{j=0}^N))^{N+1} = \mathbb{O}$ if and only if $A_N = \mathbb{O}$). Also,

$$\mu_{Q_N(\{A_j\}_{j=0}^N)} \ge \mu_{\mathbf{S}_{\mathcal{H}N}} = \dim \mathcal{H}.$$

The following is proved in [DP, Theorem 3.1 and Remark 3.6], see also [Pi], [Pe, Ch.15.3]. Let $\{\alpha_{Nj}\}_{j=0}^N \subset \mathbb{C}$. Put

(5.8)
$$R_N(\{\alpha_{Nj}\}_{j=0}^N) = Q_N(\{\alpha_{Nj}\mathbf{C}_{Nj}\}_{j=0}^N),$$

where Q_N is defined in (5.7), and \mathbf{C}_{Nj} are defined in (5.1). The operator R_N acts on $\bigoplus_{j=0}^N \mathbb{C}^{2^{N+1}} \oplus \bigoplus_{j=0}^N \mathbb{C}^{2^{N+1}}$. Then

$$M_{pb}(R_N)^2 \le 3(4 \sup_{1 \le n \le N+1} n^2 \sum_{j=n-1}^N |\alpha_{Nj}|^2 + 1)$$

and

$$M_{cpb}(R_N)^2 \ge \frac{1}{4} \sum_{j=1}^{N+1} j^2 |\alpha_{N,j-1}|^2.$$

Thus, if $\{\alpha_{Nj}\}_{j=0}^N$, $N \ge 1$, are such that

(5.9)
$$\sup_{N} \sup_{1 \le n \le N+1} n^2 \sum_{j=n-1}^{N} |\alpha_{Nj}|^2 < \infty \quad \text{and} \quad \sup_{N} \sum_{j=1}^{N+1} j^2 |\alpha_{N,j-1}|^2 = \infty,$$

then $\sup_N M_{pb}(R_N) < \infty$ and $\sup_N M_{cpb}(R_N) = \infty$. For example, one can take $\alpha_{Nj} = \alpha_j = (j+1)^{-3/2}, \ 0 \le j \le N$, for every N. Let $\{\alpha_{Nj}\}_{j=0}^N \subset \mathbb{C}, \ N \ge 1$, satisfy (5.9), and let $R_N = R_N(\{\alpha_{Nj}\}_{j=0}^N)$ be

defined in (5.8). By Proposition 1.1, $R = \bigoplus_N R_N$ is polynomially bounded

and is not similar to a contraction. The minimal function of R is the least common multiple of the minimal functions of R_N , that is, of χ^{N+2} or χ^{N+1} , where $\chi(z) = z, z \in \mathbb{D}$, therefore, the zero function. Thus, R is not a C_0 operator. Let $\{w_N\}_N \subset \mathbb{D}$ be such that $\sum_N (N+2)(1-|w_N|) < \infty$. Set $B = \prod_N b_{w_N}^{N+2}$ and $R = \bigoplus_N \beta_{w_N}(R_N)$, where β_w is defined in (1.3) and b_w is a Blaschke factor. By Corollary 1.2, R is polynomially bounded and is not similar to a contraction. Also, $B(R) = \mathbb{O}$. But $\mu_R \ge \mu_{\beta_{w_N}(R_N)} = \mu_{R_N} \ge 2^{N+1}$ for every N (see (1.5) and (1.6)), therefore, $\mu_R = \infty$.

6. Perturbation of "truncated" Foguel-Hankel operators

Before formulating the main result of this section, we introduce the following notation. Set $q_n(\lambda, \nu) = \sum_{k=0}^{n-1} \lambda^{n-1-k} \nu^k$, $n \ge 1$, $\lambda, \nu \in \mathbb{C}$, $q_0 = 0$. If $\lambda \ne \nu$, then $q_n(\lambda, \nu) = \frac{\lambda^n - \nu^n}{\lambda - \nu}$. Furthermore,

$$\lambda^n + q_n(\lambda,\nu)\nu = q_{n+1}(\lambda,\nu),$$

(6.1)
$$\begin{array}{l} \frac{n!}{k!(n-k)!}(\lambda^{n-k}+q_{n-k}(\lambda,\nu)\nu) = \frac{n!}{k!(n-k)!}q_{n+1-k}(\lambda,\nu) \\ = \frac{(n+1)!}{k!(n+1-k)!}q_{n+1-k}(\lambda,\nu) - \frac{n!}{(k-1)!(n+1-k)!}q_{n+1-k}(\lambda,\nu), \quad 1 \le k \le n. \end{array}$$

Lemma 6.1. Suppose $N \ge 1$, $0 < \delta < 1$, p is (an analytic) polynomial, and $\hat{p}(n) = 0$ for $0 \le n \le N + 1$. Then

$$\left|\sum_{n \ge N+2} \widehat{p}(n) \frac{n!}{k!(n-k)!} q_{n-k}(\lambda,\nu)\right| \le \frac{(k+1)(k+2)}{(1-\delta)^{(k+3)}} \delta \|p\|_{\infty}$$

for $\lambda, \nu \in \mathbb{C}, |\lambda| \leq \delta, |\nu| \leq \delta, 0 \leq k \leq N.$

Proof. We have

$$\sum_{n \ge N+2} \widehat{p}(n) \frac{n!}{k!(n-k)!} q_{n-k}(\lambda,\nu) = \frac{1}{k!} \frac{p^{(k)}(\lambda) - p^{(k)}(\nu)}{\lambda - \nu}, \quad \text{if } \lambda \neq \nu,$$

and

$$\sum_{n\geq N+2}\widehat{p}(n)\frac{n!}{k!(n-k)!}q_{n-k}(\lambda,\lambda) = \frac{1}{k!}p^{(k+1)}(\lambda).$$

If $0 < \delta < 1$ and $|\lambda| < \delta$, $|\nu| < \delta$, $\lambda \neq \nu$, then

$$\left|\frac{p^{(k)}(\lambda) - p^{(k)}(\nu)}{\lambda - \nu}\right| \le \sup_{|z| \le \delta} |p^{(k+1)}(z)| \le \sup_{|z| \le \delta} |p^{(k+1)}(z) - p^{(k+1)}(0)|$$
$$\le \sup_{|z| \le \delta} |p^{(k+2)}(z)| \delta \le \frac{(k+2)!\delta}{(1-\delta)^{(k+3)}} \|p\|_{\infty} \quad \text{for } 0 \le k \le N,$$

because $p^{(k+1)}(0) = 0$ for $0 \le k \le N$.

If $\lambda = \nu$, the conclusion of the lemma follows from the estimate of $p^{(k+1)}$ above.

The following theorem is the main result of this section.

Theorem 6.2. Suppose $N \ge 1$, \mathcal{H} is a Hilbert space, D, D_{\star} , D_j , A_j are operators on \mathcal{H} such that $D_jA_j = A_jD$, and D_{\star} , D_j for $0 \le j \le N$ are simultaneously diagonalizable with respect to some orthonormal basis. Define the operators R and T on the space $\bigoplus_{n=0}^{N} \mathcal{H} \oplus \bigoplus_{n=0}^{N} \mathcal{H}$ as follows:

$$R = Q_N(\{A_j\}_{j=0}^N) = \begin{pmatrix} \mathbf{S}_{\mathcal{H}N}^* & \Gamma(\{A_j\}_{j=0}^N) \\ \mathbb{O} & \mathbf{S}_{\mathcal{H}N} \end{pmatrix}$$

and

$$T = R + \bigoplus_{n=0}^{N} D_{\star} \oplus \bigoplus_{n=0}^{N} D = \begin{pmatrix} \bigoplus_{n=0}^{N} D_{\star} + \mathbf{S}_{\mathcal{H}N}^{*} & \Gamma(\{A_{j}\}_{j=0}^{N}) \\ \mathbb{O} & \bigoplus_{n=0}^{N} D + \mathbf{S}_{\mathcal{H}N} \end{pmatrix},$$

where $\Gamma(\{A_j\}_{j=0}^N)$ is defined in (5.5) and $\mathbf{S}_{\mathcal{H}N}$ is defined in (4.1). For a polynomial p put

$$R_{(p)} = P_{\bigoplus_{n=0}^{N} \mathcal{H} \oplus \{0\}} p(R)|_{\{0\} \oplus \bigoplus_{n=0}^{N} \mathcal{H}} \text{ and } T_{(p)} = P_{\bigoplus_{n=0}^{N} \mathcal{H} \oplus \{0\}} p(T)|_{\{0\} \oplus \bigoplus_{n=0}^{N} \mathcal{H}}$$

that is, $R_{(p)}$ and $T_{(p)}$ are the operators from the right upper corner in the

that is, $R_{(p)}$ and $I_{(p)}$ are the operators from the right upper corner in the matrix forms of p(R) and p(T), respectively.

Let $\varepsilon > 0$. Then there exists $\delta > 0$ which depends on N, $\{A_j\}_{j=0}^N$, and ε , such that if $\|D_{\star}\| < \delta$ and $\|D_j\| < \delta$, $0 \le j \le N$, then

 $||T_{(p)} - R_{(p)}|| \le \varepsilon ||p||_{\infty}$ for every polynomial p.

Proof. Recall that $\chi(z) = z, z \in \mathbb{D}$. To simplify notation set $\mathbf{S} = \mathbf{S}_{\mathcal{H}N}$, $K_{\star} = \bigoplus_{n=0}^{N} D_{\star}, K = \bigoplus_{n=0}^{N} D, T_{(n)} = T_{(\chi^n)}, R_{(n)} = R_{(\chi^n)}$. We have $T_{(0)} = R_{(0)} = \mathbb{O}$ and $T_{(1)} = R_{(1)} = \Gamma(\{A_j\}_{j=0}^N)$.

Since $\mathbf{S}^* K_{\star} = K_{\star} \mathbf{S}^*$ and $\mathbf{S}^{N+1} = \mathbb{O}$,

(6.2)
$$(\mathbf{S}^* + K_\star)^n = \sum_{k=0}^{\min(n,N)} \frac{n!}{k!(n-k)!} K_\star^{n-k} (\mathbf{S}^*)^k.$$

If the operators L_j on \mathcal{H} are such that $D_j L_j = L_j D$, $0 \leq j \leq N$, then

(6.3)
$$\Gamma(n, \{L_j\}_{j=0}^N) K = \Gamma(n, \{D_j L_j\}_{j=0}^N) \text{ for } n \ge 0$$

Also, for arbitrary operators L_j on \mathcal{H} , $0 \leq j \leq N$,

(6.4)
$$K^k_{\star}\Gamma(n, \{L_j\}_{j=0}^N) = \Gamma(n, \{D^k_{\star}L_j\}_{j=0}^N) \text{ for } n \ge 0, \ k \ge 0.$$

From the equality $R_{(n)} = \sum_{k=0}^{n-1} (\mathbf{S}^*)^{n-1-k} \Gamma(\{A_j\}_{j=0}^N) \mathbf{S}^k$ and (5.6), taking into account that $\mathbf{S}^{N+1} = \mathbb{O}$, we obtain that (6.5)

$$R_{(n)} = \Gamma(n-1, \{nA_j\}_{j=0}^N), \quad 1 \le n \le N+1, \text{ and } R_{(n)} = \mathbb{O}, \quad n \ge N+2.$$

Since D_{\star} and D_j for $0 \leq j \leq N$ are simultaneously diagonalizable with respect to some orthonormal basis, we have that $D_{\star}D_j = D_jD_{\star}, 0 \leq j \leq N$, therefore, $q_n(D_{\star}, D_j)$ is well defined and (6.1) is fulfilled for D_{\star} and D_j instead of λ and ν . We prove by induction that

(6.6)
$$T_{(n)} = \sum_{k=0}^{\min(n-1,N)} \Gamma(k, \{\frac{n!}{k!(n-k)!}q_{n-k}(D_{\star}, D_{j})A_{j}\}_{j=0}^{N}), \quad n \ge 1.$$

Base of induction: if n = 1, then (6.6) is clearly fulfilled. Assume that (6.6) is fulfilled for some $n, 1 \le n \le N$. We have from (5.6), (6.1), (6.2), (6.3), and (6.4) that

$$\begin{split} T_{(n+1)} &= (\mathbf{S}^* + K_\star)^n \Gamma(\{A_j\}_{j=0}^N) + T_{(n)}(\mathbf{S} + K) \\ &= \left(\sum_{k=0}^n \frac{n!}{k!(n-k)!} K_\star^{n-k}(\mathbf{S}^*)^k\right) \Gamma(\{A_j\}_{j=0}^N) \\ &+ \sum_{k=0}^{n-1} \Gamma(k, \{\frac{n!}{k!(n-k)!} q_{n-k}(D_\star, D_j) A_j\}_{j=0}^N) (\mathbf{S} + K) \\ &= \sum_{k=0}^n \Gamma(k, \{\frac{n!}{k!(n-k)!} D_\star^{n-k} A_j\}_{j=0}^N) + \sum_{k=0}^{n-1} \Gamma(k+1, \{\frac{n!}{k!(n-k)!} q_{n-k}(D_\star, D_j) A_j\}_{j=0}^N) \\ &+ \sum_{k=0}^{n-1} \Gamma(k, \{\frac{n!}{k!(n-k)!} q_{n-k}(D_\star, D_j) D_j A_j\}_{j=0}^N) \\ &= \Gamma(\{D_\star^n A_j\}_{j=0}^N) + \Gamma(\{q_n(D_\star, D_j) D_j A_j\}_{j=0}^N) + \Gamma(n, \{A_j\}_{j=0}^N) + \Gamma(n, n\{A_j\}_{j=0}^N) \\ &+ \sum_{k=1}^{n-1} \Gamma\left(k, \{\frac{n!}{k!(n-k)!} q_{n-k}(D_\star, D_j) D_j A_j\}_{j=0}^N\right) \\ &= \Gamma(\{q_{n+1}(D_\star, D_j) + \frac{n!}{k!(n-k)!} q_{n-k}(D_\star, D_j) D_j A_j\}_{j=0}^N) \\ &= \Gamma(\{q_{n+1}(D_\star, D_j) A_j\}_{j=0}^N) + \Gamma(n, \{(n+1)A_j\}_{j=0}^N) \\ &+ \sum_{k=1}^{n-1} \Gamma(k, \{\frac{(n+1)!}{k!(n+1-k)!} q_{n+1-k}(D_\star, D_j) A_j\}_{j=0}^N) \\ &= \sum_{k=0}^n \Gamma(k, \{\frac{(n+1)!}{k!(n+1-k)!} q_{n+1-k}(D_\star, D_j) A_j\}_{j=0}^N). \end{split}$$

Thus, (6.6) for $1 \le n \le N+1$ is proved. Now assume that (6.6) is proved for some $n, n \ge N+1$. Acting as in the case $1 \le n \le N+1$, and taking into account that $\Gamma(N, \{L_j\}_{j=0}^N) \mathbf{S} = \mathbb{O}$ for arbitrary operators L_j on \mathcal{H} , we obtain that

$$T_{(n+1)} = \sum_{k=0}^{N} \Gamma(k, \{\frac{n!}{k!(n-k)!} D_{\star}^{n-k} A_j\}_{j=0}^{N}) + \sum_{k=0}^{N-1} \Gamma(k+1, \{\frac{n!}{k!(n-k)!} q_{n-k}(D_{\star}, D_j) A_j\}_{j=0}^{N}) + \sum_{k=0}^{N} \Gamma(k, \{\frac{n!}{k!(n-k)!} q_{n-k}(D_{\star}, D_j) D_j A_j\}_{j=0}^{N}) = \Gamma(\{D_{\star}^n A_j\}_{j=0}^{N}) + \Gamma(q_n(D_{\star}, D_j) D_j A_j\}_{j=0}^{N}) + \sum_{k=1}^{N} \Gamma\left(k, \{\frac{n!}{k!(n-k)!} D_{\star}^{n-k} + \frac{n!}{(k-1)!(n-k+1)!} q_{n-k+1}(D_{\star}, D_j) + \frac{n!}{k!(n-k)!} q_{n-k}(D_{\star}, D_j) D_j A_j\}_{j=0}^{N}\right)$$

$$= \sum_{k=0}^{N} \Gamma(k, \{\frac{(n+1)!}{k!(n+1-k)!}q_{n+1-k}(D_{\star}, D_{j})A_{j}\}_{j=0}^{N})$$

(we apply (6.1)).

We infer from (6.5) and (6.6) that

$$T_{(n)} = R_{(n)} + \sum_{k=0}^{n-2} \Gamma(k, \{\frac{n!}{k!(n-k)!}q_{n-k}(D_{\star}, D_j)A_j\}_{j=0}^N), \quad 2 \le n \le N+1.$$

Since $\|\Gamma\{L_j\}_{j=0}^N\| \leq \sum_{j=0}^N \|L_j\|$ for arbitrary operators L_j , $0 \leq j \leq N$, we have that

(6.7)
$$\|T_{(n)} - R_{(n)}\| \leq \sum_{k=0}^{n-2} \|\Gamma(k, \{\frac{n!}{k!(n-k)!}q_{n-k}(D_{\star}, D_{j})A_{j}\}_{j=0}^{N})\|$$
$$\leq \sum_{k=0}^{n-2} \sum_{j=k}^{N} \frac{n!}{k!(n-k)!} \|q_{n-k}(D_{\star}, D_{j})\| \|A_{j}\|, \quad 2 \leq n \leq N+1.$$

If $0 < \delta < 1$ and $||D_{\star}|| < \delta$, $||D_j|| < \delta$, $0 \le j \le N$, then $||q_{n-k}(D_{\star}, D_j)|| < (N+1)\delta$ for $0 \le k \le n-2$, $2 \le n \le N+1$. We infer from (6.7) then there exists a constant C which depends on N and $||A_j||$, $0 \le j \le N$, such that

(6.8)
$$||T_{(n)} - R_{(n)}|| \le C\delta$$
 for every $n, 2 \le n \le N + 1$,

if $0 < \delta < 1$ and $||D_{\star}|| < \delta$, $||D_j|| < \delta$, $0 \le j \le N$. For a polynomial p put

 $\frac{N+1}{N+1}$

$$p_{1N} = \sum_{n=0} \widehat{p}(n)\chi^n$$
 and $p_{2N} = \sum_{n\geq N+2} \widehat{p}(n)\chi^n$.

We infer from (6.8) that

$$\|T_{(p_{1N})} - R_{(p_{1N})}\| = \left\|\sum_{n=2}^{N+1} \widehat{p}(n)(T_{(n)} - R_{(n)})\right\|$$

$$\leq \sum_{n=2}^{N+1} |\widehat{p}(n)| \|T_{(n)} - R_{(n)}\| \leq \sum_{n=2}^{N+1} \|p\|_{\infty} C\delta = NC\delta \|p\|_{\infty},$$

that is, there exists a constant C_1 which depends on N and $||A_j||, 0 \le j \le N$, such that

(6.9)
$$||T_{(p_{1N})} - R_{(p_{1N})}|| \le C_1 \delta ||p||_{\infty}$$

for every polynomial p, if $0 < \delta < 1$ and $||D_*|| < \delta$, $||D_j|| < \delta$, $0 \le j \le N$. We infer from (6.5) that $R_{(p_{2N})} = \mathbb{O}$ for every polynomial p. From (6.6) we have

$$T_{(p_{2N})} = \sum_{n \ge N+2} \widehat{p}(n) T_{(n)} = \sum_{n \ge N+2} \widehat{p}(n) \sum_{k=0}^{N} \Gamma(k, \{\frac{n!}{k!(n-k)!} q_{n-k}(D_{\star}, D_{j}) A_{j}\}_{j=0}^{N})$$
$$= \sum_{k=0}^{N} \Gamma\left(k, \{\sum_{n \ge N+2} \widehat{p}(n) \frac{n!}{k!(n-k)!} q_{n-k}(D_{\star}, D_{j}) A_{j}\}_{j=0}^{N}\right),$$

therefore,

(6.10)
$$\|T_{(p_{2N})}\| \leq \sum_{k=0}^{N} \left\| \Gamma\left(k, \left\{ \sum_{n \geq N+2} \widehat{p}(n) \frac{n!}{k!(n-k)!} q_{n-k}(D_{\star}, D_{j}) A_{j} \right\}_{j=0}^{N} \right) \right\|$$
$$\leq \sum_{k=0}^{N} \sum_{j=k}^{N} \left\| \sum_{n \geq N+2} \widehat{p}(n) \frac{n!}{k!(n-k)!} q_{n-k}(D_{\star}, D_{j}) \right\| \|A_{j}\|.$$

Denote by λ_l and ν_{jl} the eigenvectors of D_{\star} and D_j , $0 \leq j \leq N$ (recall that D_{\star} and D_j for $0 \leq j \leq N$ are simultaneously diagonalizable with respect to some orthonormal basis). We have (6.11)

$$\left\|\sum_{n\geq N+2}^{n} \widehat{p}(n) \frac{n!}{k!(n-k)!} q_{n-k}(D_{\star}, D_{j})\right\| = \sup_{l} \left|\sum_{n\geq N+2}^{n} \widehat{p}(n) \frac{n!}{k!(n-k)!} q_{n-k}(\lambda_{l}, \nu_{jl})\right|,$$

where $0 \le k \le N$. From (6.10), (6.11), and Lemma 6.1 we conclude that there exists a constant C which depends on N and $||A_j||$, $0 \le j \le N$, such that

(6.12)
$$||T_{(p_{2N})}|| \le C\delta ||p_{2N}||_{\infty}$$

for every polynomial p, if $0 < \delta \leq 1/2$, $||D_{\star}|| < \delta$ and $||D_j|| < \delta$, $0 \leq j \leq N$. There exists a constant c (which does not depend on N) such that $||p_{2N}||_{\infty} \leq c \log N ||p||_{\infty}$ for every polynomial $p, N \geq 2$ [T, 13.4.3]. We conclude from (6.12) that there exists a constant C_2 which depends on N and $||A_j||, 0 \leq j \leq N$, such that

(6.13)
$$||T_{(p_{2N})}|| \le C_2 \delta ||p||_{\infty}$$

<

for every polynomial p, if $0 < \delta \le 1/2$ and $||D_*|| < \delta$, $||D_j|| < \delta$, $0 \le j \le N$. Finally, if $0 < \delta \le 1/2$ and $||D_*|| < \delta$, $||D_j|| < \delta$, $0 \le j \le N$, then from

Finally, if $0 < \delta \le 1/2$ and $||D_{\star}|| < \delta$, $||D_j|| < \delta$, $0 \le j \le N$, then from (6.9) and (6.13)

$$||T_{(p)} - R_{(p)}|| = ||T_{(p_{1N})} - R_{(p_{1N})} + T_{(p_{2N})}||$$
$$||T_{(p_{1N})} - R_{(p_{1N})}|| + ||T_{(p_{2N})}|| \le (C_1 + C_2)\delta||p||_{\infty}$$

for every polynomial p. Put $\delta = \varepsilon/(C_1 + C_2)$. The theorem is proved. \Box

Theorem 6.3. There exist sequences of operators $\{T_N\}_N$ and of finite Blaschke products $\{B_N\}_N$ with zeros from (0,1) such that $\sup_N M_{pb}(T_N) < \infty$, $\sup_N M_{cpb}(T_N) = \infty$, and $T_N \approx T_{B_N}$.

Proof. Let $\{\alpha_{Nj}\}_{j=0}^N$, $N \ge 1$, satisfy (5.9), and let $R_N = R_N(\{\alpha_{Nj}\}_{j=0}^N)$ be defined in (5.8). Then $\sup_N M_{pb}(R_N) < \infty$, and there exists a sequence of families of polynomials $[p_{Nij}]_{i,j=1}^{2^{N+1}}$, $N \ge 1$, such that

(6.14)
$$\|[R_{N,(p_{Nij})}]_{i,j=1}^{2^{N+1}}\| \ge C_N \|[p_{Nij}]_{i,j=1}^{2^{N+1}}\|_{H^{\infty}(\ell_{2^{N+1}}^2)}, \text{ where } C_N \to \infty,$$

here $R_{N,(p)}$ are defined for R_N as in Theorem 6.2 (see [DP, Theorem 3.1]). Let a sequence $\{\varepsilon_N\}_N$ be such that $\varepsilon_N > 0$ and

(6.15)
$$C = \sup_{N} 2^{N+1} \varepsilon_N < \infty.$$

Applying Lemma 4.1 to N and ε_N and Theorem 6.2 to N, R_N , and ε_N we obtain a sequence $\{\delta_N\}_N$ such that δ_N satisfies the conclusion of Lemma 4.1 and Theorem 6.2 for every N. Let $\{a_{Nl}\}_{l=0}^N$ and $\{c_{Nl}\}_{l=0}^N$ be from Lemma 5.1 applied to δ_N . Put $D_N = \mathbf{D}(\{a_{Nl}\}_{l=0}^N, \{c_{Nl}\}_{l=0}^N)$, and $D_{Nj} = \mathbf{D}_j(\{a_{Nl}\}_{l=0}^N, \{c_{Nl}\}_{l=0}^N), 0 \leq j \leq N$, where **D** and **D**_j are defined in (5.2) and (5.3), respectively. Denote the eigenvalues of D_N by λ_{Nl} , $1 \leq l \leq 2^{N+1}$. By Lemma 5.1, $\lambda_{Nl} \in (0, \delta_N)$ and $\lambda_{Nl} \neq \lambda_{Nk}$, $l \neq k$. Also, D_{Nj} are diagonal with respect to the standard basis in $\mathbb{C}^{2^{N+1}}$, and the elements of D_{Nj} are from $(0, \delta_N)$. Let $\nu_{Nl} \in (0, \delta_N)$ be such that $\nu_{Nl} \neq \nu_{Nk}$ for $l \neq k$, and $\nu_{Nl} \neq \lambda_{Nk}$ for all $l, k, 1 \leq k, l \leq 2^{N+1}$. Denote by $D_{N\star}$ the diagonal operator with respect to the standard basis on $\mathbb{C}^{2^{N+1}}$, with the eigenvalues ν_{Nl} . Put $B_N = \prod_{l=1}^{2^{N+1}} b_{\lambda_{Nl}}^{N+1} b_{\nu_{Nl}}^{N+1}$ and

$$T_N = R_N + \bigoplus_{n=0}^N D_{N\star} \oplus \bigoplus_{n=0}^N D_N.$$

By Lemma 4.3, $T_N \approx T_{B_N}$.

Put $T_{N0} = \bigoplus_{n=0}^{N} D_N + \mathbf{S}_{\mathbb{C}^{2^{N+1}}N}$ and $T_{N1} = \bigoplus_{n=0}^{N} D_{N\star} + \mathbf{S}_{\mathbb{C}^{2^{N+1}}N}^*$. By Corollary 4.2 applied to T_{N0} and T_{N1} there exist operators X_{N0} and X_{N1} such that $X_{N0}T_{N0} = (\bigoplus_{l=1}^{2^{N+1}} T_{b_{\lambda_{Nl}}^{N+1}}) X_{N0}, X_{N1}T_{N1} = (\bigoplus_{l=1}^{2^{N+1}} T_{b_{\nu_{Nl}}^{N+1}}) X_{N1}, ||X_{N0}|| \le 1 + 1$

 $\varepsilon_N, \|X_{N0}^{-1}\| \leq 1 + \varepsilon_N, \|X_{N1}\| \leq 1 + \varepsilon_N, \text{ and } \|X_{N1}^{-1}\| \leq 1 + \varepsilon_N.$ Therefore,

(6.16)
$$M_{pb}(T_{N0}) \le (1+\varepsilon_N)^2$$
 and $M_{pb}(T_{N1}) \le (1+\varepsilon_N)^2$.

By Theorem 6.2,

(6.17)
$$||T_{N,(p)} - R_{N,(p)}|| \le \varepsilon_N ||p||_{\infty}$$
 for every polynomial p .

From (6.16) and (6.17) we conclude that $\sup_N M_{pb}(T_N) < \infty$. From (6.15) and (6.17) we have that

(6.18)

$$\begin{split} \left\| \left[T_{N,(p_{Nij})} - R_{N,(p_{Nij})} \right]_{i,j=1}^{2^{N+1}} \right\| &\leq 2^{N+1} \sup_{1 \leq i,j \leq 2^{N+1}} \| T_{N,(p_{Nij})} - R_{N,(p_{Nij})} \| \\ &\leq 2^{N+1} \varepsilon_N \sup_{1 \leq i,j \leq 2^{N+1}} \| p_{Nij} \|_{\infty} \leq C \left\| \left[p_{Nij} \right]_{i,j=1}^{2^{N+1}} \right\|_{H^{\infty}(\ell_{2^{N+1}}^2)}. \end{split}$$

If $C_N > C$, then from (6.14) and (6.18)

$$\|[T_{N,(p_{Nij})}]_{i,j=1}^{2^{N+1}}\| \ge \|[R_{N,(p_{Nij})}]_{i,j=1}^{2^{N+1}}\| - \|[T_{N,(p_{Nij})} - R_{N,(p_{Nij})}]_{i,j=1}^{2^{N+1}}\| \\ \ge (C_N - C)\|[p_{Nij}]_{i,j=1}^{2^{N+1}}\|_{H^{\infty}(\ell_{2^{N+1}}^2)},$$

therefore, $\sup_N M_{cpb}(T_N) = \infty$.

Corollary 6.4. There exist an operator T and a Blashcke product B with zeros from (0,1) such that T is polynomially bounded, T is not similar to a contraction, and $T \sim T_B$, where T_B is defined in (1.7).

Proof. Let $\{T_N\}_N$ and $\{B_N\}_N$ be the sequences of operators and of finite Blaschke products with zeros from (0, 1) from Theorem 6.3. By Lemma 3.1, there exists a sequence $\{w_N\}_N \subset (0, 1)$ such that $B = \prod_N \zeta_N B_N \circ \beta_{w_N}$ converges, and $B_N \circ \beta_{w_N}$ are pairwise coprime.

Put $T = \bigoplus_N \beta_{w_N}(T_N)$. By Lemma 3.2, $\beta_{w_N}(T_N) \approx \beta_{w_N}(T_{B_N}) \cong T_{B_N \circ \beta_{w_N}}$, therefore, $T \sim T_B$. By Corollary 1.2, T is polynomially bounded, and T is not similar to a contraction.

Applying Corollary 2.3 to the operator T from Corollary 6.4, one can obtain a polynomially bounded operator \mathbf{T} such that $\mathbf{T} \prec S$, and \mathbf{T} is not similar to a contraction. Denote by X and Y quasiaffinities such that $YT = T_BY$ and $XT_B = TX$. By [SFBK, Theorem X.2.10], there exists a function $\varphi \in H^{\infty}$ such that $YX = \varphi(T_B)$. The constructions of the operators R_N from [DP] and of the operator T from Corollary 6.4 are explicit, therefore, the quasiaffinities X and Y and a function φ can be computed, but the author cannot do it. Also, since \mathbf{T} is polynomially bounded and $\mathbf{T} \prec S$, it follows from [BP] that $\mathbf{T} \sim S$ if and only if $\mu_{\mathbf{T}} = 1$. But the author cannot compute $\mu_{\mathbf{T}}$.

7. The construction of quasisimilar operators such that the product of intertwining quasiaffinities is an outer function of operators

Theorem 7.1. There exist an operator $T: \mathcal{H} \to \mathcal{H}$, a Blashcke product *B* with zeros from (0,1), an outer function $g \in H^{\infty}$, and quasiaffinities $X: \mathcal{H} \to \mathcal{K}_B, Y: \mathcal{K}_B \to \mathcal{H}$, such that *T* is polynomially bounded, *T* is not similar to a contraction, $XT = T_BX$, $YT_B = TY$, and $XY = g(T_B)$, where T_B is defined in (1.7).

Proof. Let $\{T_N\}_N$ and $\{B_N\}_N$ be the sequences of operators and of finite Blaschke products with zeros from (0,1) from Theorem 6.3, respectively. Let C > 0 be fixed. Denote by \mathcal{H}_N the finite dimensional spaces on which T_N acts. There exists invertible operators $X_N \colon \mathcal{H}_N \to \mathcal{K}_{B_N}, Y_N \colon \mathcal{K}_{B_N} \to \mathcal{H}_N$ such that $X_N T_N = T_{B_N} X_N, Y_N T_{B_N} = T_N Y_N, ||X_N|| \leq C, ||Y_N|| \leq C$. By [SFBK, Theorem X.2.10], there exist functions $\varphi_N \in H^{\infty}$ such that $X_N Y_N = \varphi_N(T_{B_N})$. Note that $\varphi_N(\lambda) \neq 0$ for every $\lambda \in \mathbb{D}$ such that $B_N(\lambda) = 0$, for every index N.

Let g be from Lemma 3.4. Applying Theorem 3.8 to C, g, sequences of B_N and of φ_N we obtain $\delta > 0$ and sequences of $w_N \in (0, 1)$, of $\zeta_N \in \mathbb{T}$ and of $\psi_N \in H^{\infty}$ which satisfy the conclusion of Theorem 3.8. Put

$$B = \prod_{N} \zeta_N B_N \circ \beta_{w_N}$$
 and $T = \bigoplus_N \beta_{w_N}(T_N).$

By Theorem 6.3 and Corollary 1.2, T is polynomially bounded, and T is not similar to a contraction.

Put $U_N = U_{w_N}|_{\mathcal{K}_{B_N \circ \beta_{w_N}}} : \mathcal{K}_{B_N \circ \beta_{w_N}} \to \mathcal{K}_{B_N}$, where U_w is defined in Lemma 3.2. Then U_N is unitary, and $U_N T_{B_N \circ \beta_{w_N}} = \beta_{w_N}(T_{B_N})U_N$. Put

$$X_{1N} = U_N^{-1}(\psi_N \circ \beta_{w_N})(T_{B_N})X_N \quad \text{and} \quad Y_{1N} = Y_N U_N.$$

Then

$$X_{1N}\beta_{w_N}(T_N) = T_{B_N \circ \beta_{w_N}} X_{1N}, \quad Y_{1N}T_{B_N \circ \beta_{w_N}} = \beta_{w_N}(T_N)Y_{1N}$$
$$X_{1N}Y_{1N} = g(T_{B_N \circ \beta_{w_N}}), \quad ||X_{1N}|| \le C^2, \quad ||Y_{1N}|| \le C,$$

and X_{1N} and Y_{1N} are quasiaffinities. Indeed, $X_N\beta_{w_N}(T_N) = \beta_{w_N}(T_{B_N})X_N$ and $Y_N\beta_{w_N}(T_{B_N}) = \beta_{w_N}(T_N)Y_N$. Therefore,

$$X_{1N}\beta_{w_N}(T_N) = U_N^{-1}\psi_N(\beta_{w_N}(T_{B_N}))\beta_{w_N}(T_{B_N})X_N$$

= $U_N^{-1}\beta_{w_N}(T_{B_N})\psi_N(\beta_{w_N}(T_{B_N}))X_N = T_{B_N\circ\beta_{w_N}}U_N^{-1}\psi_N(\beta_{w_N}(T_{B_N}))X_N$
= $T_{B_N\circ\beta_{w_N}}X_{1N}.$

Also,

$$Y_{1N}T_{B_N\circ\beta_{w_N}} = Y_N U_N T_{B_N\circ\beta_{w_N}} = Y_N\beta_{w_N}(T_{B_N})U_N = \beta_{w_N}(T_N)Y_{1N}.$$

Furthermore,

 $(\psi_N \circ \beta_{w_N})(T_{B_N}) X_N Y_N = (\psi_N \circ \beta_{w_N})(T_{B_N}) \varphi_N(T_{B_N}) = (\varphi_N \cdot \psi_N \circ \beta_{w_N})(T_{B_N}).$ Since $g - \psi_N \cdot \varphi_N \circ \beta_{w_N} \in (B_N \circ \beta_{w_N}) H^{\infty}$, we have that $g \circ \beta_{w_N} - \varphi_N \cdot \psi_N \circ \beta_{w_N} \in B_N H^{\infty}$, therefore, $(\varphi_N \cdot \psi_N \circ \beta_{w_N})(T_{B_N}) = (g \circ \beta_{w_N})(T_{B_N}) = g(\beta_{w_N}(T_{B_N})).$ Thus, $X_{1N}Y_{1N} = U_N^{-1}g(\beta_{w_N}(T_{B_N}))U_N = g(T_{B_N \circ \beta_{w_N}}).$ By Nehari's theorem,

$$||X_{1N}|| \le ||(\psi_N \circ \beta_{w_N})(T_{B_N})|| ||X_N|| \le C \operatorname{dist}(\psi_N \circ \beta_{w_N}, B_N H^{\infty})$$
$$= C \operatorname{dist}(\psi_N, (B_N \circ \beta_{w_N}) H^{\infty}) \le C^2.$$

Clearly, $||Y_{1N}|| = ||Y_N|| \leq C$. Since $X_{1N}Y_{1N} = g(T_{B_N \circ \beta_{w_N}})$ and g is outer, $X_{1N}Y_{1N}$ is a quasiaffinity, and, since Y_{1N} is invertible, we conclude that X_{1N} is a quasiaffinity.

We have $\mathcal{K}_B = \bigvee_N \mathcal{K}_{B_N \circ \beta_{w_N}}$. Define a linear mapping $J \colon \mathcal{K}_B \to \bigoplus_N \mathcal{K}_{B_N \circ \beta_{w_N}}$ by the formula $J \sum_N x_N = \bigoplus_N x_N$, where $x_N \in \mathcal{K}_{B_N \circ \beta_{w_N}}$, the cardinality of the set of N such that $x_N \neq 0$ is finite. Since $\{B_N \circ \beta_{w_N}\}_N$ satisfy the condition (3.6), J is expanded on \mathcal{K}_B , is bounded and invertible (its inverse is bounded) [Ni, Theorems II.C.3.1.4 and II.C.3.2.14]. Since the spaces $\mathcal{K}_{B_N \circ \beta_{w_N}}$ are invariant for T_B^* , it is easy to see that $JT_B^* = (\bigoplus_N T_{B_N \circ \beta_{w_N}})^*J$.

Define the following operators:

$$Z: \oplus_N \mathcal{H}_N \to \mathcal{K}_B, \quad Z = J^{-1}(\oplus_N Y_{1N}^*),$$
$$W: \mathcal{K}_B \to \oplus_N \mathcal{H}_N, \quad W = (\oplus_N X_{1N}^*)J.$$

It is easy to see that Z and W are quasiaffinities. Also,

(7.1)
$$ZT^* = T_B^*Z, WT_B^* = T^*W, \text{ and } ZW = g_*(T_B^*),$$

where $g_*(z) = \overline{g(\overline{z})}, z \in \mathbb{D}$. Indeed,

$$ZT^* = J^{-1}(\bigoplus_N Y_{1N}^*) \left(\bigoplus_N (\beta_{w_N}(T_N)) \right)^* = J^{-1} \left(\bigoplus_N Y_{1N}^* (\beta_{w_N}(T_N))^* \right)$$
$$= J^{-1} \left(\bigoplus_N T_{B_N \circ \beta_{w_N}} \right)^* \left(\bigoplus_N Y_{1N}^* \right) = T_B^* J^{-1} \left(\bigoplus_N Y_{1N}^* \right) = T_B^* Z.$$

Also,

$$WT_B^* = (\bigoplus_N X_{1N}^*) JT_B^* = (\bigoplus_N X_{1N}^*) (\bigoplus_N T_{B_N \circ \beta_{w_N}})^* J$$

= $(\bigoplus_N (\beta_{w_N}(T_N))^*) (\bigoplus_N X_{1N}^*) J = T^* W.$

Furthermore,

$$ZW = J^{-1}(\bigoplus_N Y_{1N}^*)(\bigoplus_N X_{1N}^*)J = J^{-1}(\bigoplus_N (X_{1N}Y_{1N})^*)J$$
$$= J^{-1}g_*(\bigoplus_N (T_{B_N \circ \beta_{w_N}})^*)J = g_*(T_B^*).$$

The conclusion of the theorem follows from (7.1).

References

[AT] T. Ando and K. Takahashi, On operators with unitary ρ -dilations, Ann. Polon. Math., **66** (1997), 11–14.

[B] C. Badea, Perturbations of operators similar to contractions and the commutator equation. *Stud. Math.*, **150** (2002), No.3, 273-293.

[BP] H. Bercovici and B. Prunaru, Quasiaffine transforms of polynomially bounded operators, *Arch. Math. (Basel)*, **71** (1998), 384–387.

[C] G. Cassier, Generalized Toeplitz operators, restriction to invariant subspaces and similarity problems. *J. Oper. Theory*, **53** (2005), No. 1, 49-89.

[DP] K. R. Davidson and V. I. Paulsen, Polynomially bounded operators, J. reine angew. Math, 487 (1997), 153-170.

[H] P. R. Halmos, Ten problems in Hilbert space, *Bull. Am. Math. Soc.*, **76** (1970), 887-933.

[K] L. Kérchy, Quasianalytic polynomially bounded operators, *Operator Theory: the State of the Art*, Theta, Bucharest, 2016, 75–101.

[M] W. Mlak, Algebraic polynomially bounded operators, Ann. Polon. Math., **29** (1974), 133–139.

[Ni] N. K. Nikolski, Operators, functions, and systems: an easy reading. Volume I: Hardy, Hankel, and Toeplitz, Volume II: Model operators and systems, Math. Surveys and Monographs **92**, AMS, 2002.

[No] E. A. Nordgren, The ring N^+ is not adequate, Acta Sci. Math. (Szeged), **36** (1974), 203–204.

[Pa] V. I. Paulsen, Every completely polynomially bounded operator is similar to a contraction, J. Funct. Anal., 55 (1984), 1-17.

[Pe] V. V. Peller, *Hankel operators and their applications*, Springer Monographs in Math. New York, NY, Springer, 2003.

[Pi] G. Pisier, A polynomially bounded operator on Hilbert space which is not similar to a contraction, J. Am. Math. Soc., **10** (1997), No.2, 351-369.

[SFBK] B. Sz.-Nagy, C. Foias, H. Bercovici and L. Kérchy, *Harmonic analysis of operators on Hilbert space*, Springer, New York, 2010.

[T] E. C. Titchmarsh, *The theory of functions*. 2nd ed. London, Oxford University Press, 1975.

St. Petersburg Branch, V. A. Steklov Institute of Mathematics, Russian Academy of Sciences, Fontanka 27, St. Petersburg, 191023, Russia

Email address: gamal@pdmi.ras.ru