PLANAR ROOTED LINE ARRANGEMENTS AND AN OPERAD FOR FACTORIZED SCATTERING

DENIS BASHKIROV

ABSTRACT. We introduce two topological non- Σ operad structures on planar line arrangements subject to a certain geometric order condition, ensuring a well-defined notion of particle ordering on a distinguished line. This is interpreted in terms of scattering diagrams in purely elastic (1+1)dimensional theories. We discuss a possible approach to factorized scattering in operadic terms.

1. INTRODUCTION

Consider a system of n identical classical non-relativistic particles of equal mass m = 1 on a line, governed by the Hamiltonian $H = \frac{1}{2} \sum_{i=1}^{n} p_i^2 + g^2 \sum_{i < j} V(q_i - q_j)$, where q_i and p_i denote the position and momentum of the i-th particle respectively, V is a pairwise interaction potential and g is a real coupling constant. The potential V = V(q) is assumed to be repulsive, symmetric about q = 0, falling off sufficiently rapidly as $|q| \to +\infty$ and impenetrable, meaning $V(q) \to +\infty$ as $|q| \rightarrow 0$. The key examples of such V's include the rational $V(q) = \frac{1}{q^2}$ and the hyperbolic $V(q) = \frac{a^2}{\sinh^2(aq)}$ Calogero-Moser potentials. In the latter case the parameter a > 0 is the inverse of the characteristic interaction range, and the rational case is restored in the limit $a \rightarrow 0$. Both cases are hallmark examples of integrable many-body systems, where integrability can be established by explicitly constructing a corresponding Lax pair [Mos75][Kul76][Per90, Chapter 3]. This had received further development in Olshanetsky–Perelomov's projection method [OP81] and Kazhdan– Kostant–Sternberg's Hamiltonian reduction [KKS78] [Eti14]. A feature that makes these systems particularly amenable to algebraic and combinatorial analysis is existence of asymptotically-free zones for the particles and a simple asymptotic behavior of solutions. Namely, for $1 \le i \le n$, the latter are characterized by the asymptotic prescriptions $q_i(t) = p_i^- t + q_i^- + o(1)$ for $t \to -\infty$, and $q_i(t) = p_i^+ t + q_i^+ + o(1)$ for $t \to +\infty$ for a set of constants $\{p_i^{\pm}, q_i^{\pm}\}$. Furthermore, by the assumptions on V, the particles cannot overtake each other and thus can be indexed in such a way that $q_1(t) < q_2(t) < \cdots < q_n(t)$ for all t. This implies $p_1^- > p_2^- > \cdots > p_n^-$ and $p_1^+ < p_2^+ < \cdots < p_n^+$. If, as in the Calogero-Moser case, the Lax matrix can be chosen to admit diagonal asymptotic limits for $t \to \pm \infty$, then the sets of the incoming $\{p_i^-\}$ and outgoing $\{p_i^+\}$ momenta can be shown to be equal; in fact, $p_i^- = p_{n-i+1}^+$ for $1 \le i \le n$. The *n* independent integrals of motion can be chosen to be asymptotically symmetric functions of momenta $\sum_{i=1}^{n} p_i^k$ for $1 \le k \le n$. One says that such a system supports *purely elastic scattering*. This property is not exclusive to the above model and is known to emerge in other settings as well, including (relativistic) field-theoretic ones, such as soliton scattering in the sine-Gordon model [Zam77][Tor22], and other quantum integrable systems [ZZ79],[Dor07] and references therein.

Adhering to the classical Galilean set-up in our exposition, consider an initial (t = 0) configuration of n particles on a line at positions $q_1 < q_2 < \cdots < q_n$ with incoming asymptotic momenta $p_1 > p_2 > \cdots > p_n$. The conditions imply that no interaction took place before t = 0, and each particle will eventually catch up and interact (strictly once) with any other. By equality of the asymptotic momenta sets, every single interaction of two, or more, particles results in a mere exchange of asymptotic momenta between them. The evolution of such a system can be geometrically encoded by an arrangement of n lines in a qt-plane – a scattering diagram, where the *i*-th line has the slope p_i and passes through the point t = 0, $q = q_i$.



FIGURE 1. A scattering diagram.

Note that while such a line arrangement does contain all the data that determines the corresponding scattering scenario, it does not depict the particles worldlines as such. The lines are the asymptotes, and by our assumptions on the interaction potential, in the vicinity of any point of their intersection, the trajectories of pairwise-interacting particles are disjoint. As an exercise, the reader may trace a possible set of trajectories on the figure above.



This is a reflection presentation of a scattering process pertaining to our classical picture. A complimentary point of view on combinatorics of particle dynamics is that of the transmission presentation. Namely, we may assume that each particle always retains its asymptotic momentum p_i , while an interaction amounts to changing the relative order of the particles involved, as if they pass through each other, accompanied by phase shifts. The latter are discrepancies δ_i in the asymptotic positions that show up upon identifying the *i*-th incoming particle with the (n-i+1)-th outgoing particle in the transmission picture. Specifically, $\delta_i = q_{n-i+1}^+ - q_i^-$. Due to $p_{n-i+1}^+ = p_i^-(=p_i)$, $q_i(t) = p_i^-t + q_i^- + o(1)$ for $t \to -\infty$ and $q_{n-i+1}(t) = p_{n-i+1}^+t + q_{n-i+1}^+ + o(1)$ for $t \to +\infty$, we have $\delta_i = \lim_{t\to +\infty} [q_{n-i+1}(t) - q_i(-t) - 2p_it]$. The values $\delta_1, \ldots, \delta_n$ characterize the scattering process and can be thought of as a classical counterpart of a quantum-mechanical S-matrix. For instance, in the rational Calogero-Moser case all δ_i 's are known to vanish [Aru19, Chapter 4].



FIGURE 2. The reflection and the transmission presentations of an elastic two-body scattering process.

Integrability of both classical and quantum pure elastic scattering processes is associated with a particular feature that the classical scattering data or the *S*-matrix possess. This is usually stated as the *factorizability* property thereof. Namely, in presence of sufficiently many independent conserved quantities, the complete *S*-matrix accounting for multiparticle collisions can be shown

3

to be representable as a composition of the two-body S-matrices. A way to think about it in terms of scattering diagrams is that any such diagram can be perturbed into a line arrangement in a general position, where only pairwise line intersections appear, $\frac{n(n-1)}{2}$ of them in total for n lines. The invariance of a quantum S-matrix under such a perturbation is expressed as the Yang-Baxter relation.

$$\sum_{a,b,c} S_{ij}^{ba}(\theta_1 - \theta_2) S_{bk}^{nc}(\theta_1 - \theta_3) S_{ac}^{ml}(\theta_2 - \theta_3) = \sum_{a,b,c} S_{jk}^{bc}(\theta_2 - \theta_3) S_{ic}^{al}(\theta_1 - \theta_3) S_{ab}^{nm}(\theta_1 - \theta_2),$$

where the spectral parameter θ_i is identified with the rapidity or momentum of the *i*-th particle; cf. figure 16 below. Note that in the quantum case the particles may have internal degrees of freedom, thus the *S*-matrix indices above, and come in different flavors. The invariance of the particles count in this setting is not immediate.

While, effectively, the data of a scattering diagram is that of the initial positions and asymptotic momenta of classical particles, or that of the relative order and asymptotic momenta of quantum ones, the line arrangement interpretation proves itself useful in the following way.

1.1. An operadic symmetry. We would like to take an look at the integrability in (1 + 1)dimensional scattering through the lens of a certain *operadic* symmetry, meaning a consistency condition with respect to an operation of fusing, or composing, multiple initial particle configurations together. As before, consider the setting of the classical scattering on a line with initial configurations characterized uniquely by the ordered positions $q_1 < \cdots < q_n$ and momenta $p_1 > \cdots > p_n$ of the particles. A mere set-theoretic union of two initial configurations does not in general result in a well-defined initial configuration due to a possible violation of the order condition on the momenta or possible particle overlaps. As a more refined approach, a composition of two particle configurations a and b can be defined by replacing an interval between two adjacent particles of aby an appropriately rescaled, both in terms of positions and momenta, copy of b. Note that such a rescaling procedure can be nonlinear and is not unique, since ordered positions and momenta can always be perturbed. The substitution operation $(-\circ -)$ is manifestly non-commutative and requires an additional parameter i – the insertion position, or the index of the gap between two adjacent particles of a, where b is going to be mapped to.



FIGURE 3. Composing two initial particle configurations.

For initial configurations a and b consisting of m and n particles respectively, such an operation is expected to return a valid initial configuration $a \circ_i b$ consisting of m + n - 2 particles. As noted above, this alone does not determine $a \circ_i b$ uniquely, unless b is a 2-particle configuration. As a possible bootstrap, among all possible ways of defining $(-\circ_i -)$ -compositions, we can single out the ones that collectively exhibit a certain "chronological" symmetry¹. Namely, a natural condition to impose is independence of an iteratively constructed initial particle configuration on the order in which pairwise compositions of three, or more, configurations a_1, \ldots, a_k are performed, as long as

 $^{^{1}}$ One may think of an additional time axis orthogonal to the original spacetime – the time axis of an agent preparing an initial configuration for us.

a chosen relative order of the operands in such a compound composition remains the same. Here, by a relative order on a_1, \ldots, a_k we mean a partial order defined by the cover relation $a_i < a_j$ iff a_j gets (eventually) substituted into a_i . The Hasse diagram of such a poset is a rooted tree, which, upon adding some extra leaves, encodes the corresponding compositional pattern.



FIGURE 4. An iterated composition of three particle configurations and the corresponding tree. Another way to assemble the same configuration is $(a \circ_2 b) \circ_2 c$.

For three configurations a, b and c the discrete "chronological" invariance condition takes the form

$$(a \circ_i b) \circ_j c = a \circ_i (b \circ_{j-i+1} c)$$

or

$$(a \circ_i b) \circ_{i+l-1} c = (a \circ_i c) \circ_i b,$$

depending on the ranges, where the insertion position indices i and j fall in. In the latter identity, l + 1 is the total number of particles in b. The invariance conditions for higher compositions can be deduced from here inductively. The equations above are a form of generalized associativity formalized by the notion of an operad. The affine geometry of line arrangements gets handy when we construct an explicit example of such an operad of particle configurations.

Theorem. There is a structure of a non- Σ operad $\widehat{\mathcal{L}}$ on scattering diagrams (equivalently, on the initial particle configurations) with binary generators parametrized by rooted line arrangements of rank 3.

The presence of an operad structure on scattering diagrams prompts to bring up its counterpart at the level of S-matrices. Indeed, as we show, the S-matrix data of

$$\left.\begin{array}{c} \text{Initial/final} \\ \text{states} \end{array}\right\} \rightarrow \left\{\begin{array}{c} \text{Transition amplitudes} \\ \text{(classical phase shifts)} \end{array}\right.$$

of purely elastic scattering can be recast as a morphism of operads. Namely one may think of a C-valued (1+1)-dimensional purely elastic scattering theory as of an operad morphism $S: \widehat{\mathcal{L}} \to \mathcal{P}$, where $\widehat{\mathcal{L}}$ is an operad of scattering diagrams and \mathcal{P} is a non- Σ operad in a symmetric monoidal category C. In particular, the S-matrix factorization property translates into S being entirely determined by its restriction onto the monoid $\widehat{\mathcal{L}}(1)$, that is, on two crossing lines arrangements.

2. Preliminaries

2.1. **Operads.** We recall the terminology and the basic concepts of operadic algebra. A point of view that will be convenient for us to maintain throughout the paper is that an *operad* \mathcal{P} is a container of combinatorial objects defined inductively by means of iteratively putting together, or *composing*, some atomic pieces – the *generators* of \mathcal{P} . Any generator as well as any composite object a of \mathcal{P} is characterized by a positive integer, the *arity*, that measures receptivity of a to forming new compositions with any given element of \mathcal{P} . That is, the arity m of $a \in \mathcal{P}$ counts the

5

number of possible composites $a \circ_i b$ for i = 1, ..., m that a can form for any fixed $b \in \mathcal{P}$. In many natural cases, the process of building combinatorial objects via iterated compositions exhibits a certain symmetry – a consistency condition of associativity that lies at the core of the definition of an operad. Furthermore, to enhance a merely combinatorial setting, we would like the elements of \mathcal{P} to have an internal structure (for instance, topological) respected by the composition operations. Moreover, it is desired to have a well-defined notion of a congruence on an operad that would enable one to impose *relations* among the composites. A common approach to implement all this is to encapsulate all the combinatorial data of \mathcal{P} into a collection of objects and morphisms of a chosen category C that would host all the generators and relations.

A particular implementation of such a structure, which we find most suitable for our purposes, is that of a non- Σ , or a non-symmetric, operad [Gir18], [LV12, Section 5.9], [MSS02, Chapter 1]. Descriptively, in a non- Σ operad \mathcal{P} , the elements are modeled on planar² rooted trees, while the operation of forming a composite $a \circ_i b$ corresponds to grafting two planar rooted trees a, b together. We recall that the grafting operation is defined in the following way. First, by invoking planarity, we enumerate the leaves of all the rooted trees involved. The indexing scheme that we adhere to is defined by traversing any given planar rooted tree in the clockwise caterpillar order, starting at the root, and enumerating all the (non-root) leaves in their order of appearance. Next, upon having all the leaves indexed, the result of grafting b onto the *i*-th leaf of a is defined as the tree $a \circ_i b$, formed by attaching the root of b onto the -th leaf of a, while preserving its planar structure, and re-indexing all the leaves of the resulting tree accordingly.



FIGURE 5. Planar rooted trees grafting.

In a sense, a non- Σ operad is a structure that we get upon allowing all the tree vertices in the above picture to carry some extra data and allowing trees decorated in this manner to be subject to relations expressible in terms of the vertices data and the grafting operation. This is formalized in the following way. Let C be a category that we assume to be closed symmetric monoidal, complete and cocomplete. The usual category **Sets** of sets, the category **Top** of topological spaces and the category \Bbbk -(gr)Vect of (graded) vector spaces over a field \Bbbk are the main use-cases for us. Note that each of these categories is concrete, meaning it comes equipped with a (forgetful) faithful functor $C \rightarrow$ **Sets** of *elements*. Accordingly, in what follows, we will assume, at times, superfluously, that C is concrete.

Definition 1. We say that a collection of objects $\mathcal{P} = \{\mathcal{P}(n)\}_{n \geq 1}$ and morphisms

$$(-\circ_i -): \mathcal{P}(m) \otimes \mathcal{P}(n) \to \mathcal{P}(m+n-1)$$

in C, defined for all $m, n \ge 1$ and i = 1, ..., m is a non- Σ operad in C, provided that these morphisms, called *partial compositions*, satisfy the following conditions. For any $k, l, m \ge 1$, $a \in \mathcal{P}(k)$, $b \in \mathcal{P}(l)$, $c \in \mathcal{P}(m)$, we have

(1)
$$(a \circ_i b) \circ_j c = a \circ_i (b \circ_{j-i+1} c)$$

²In the literature, the term *plane tree*, referring to a tree equipped with an embedding into an oriented Euclidean plane considered up to an orientation-preserving isotopy, is used interchangeably. Combinatorially, this additional data amounts to defining a cyclic order on the set F_v of all half-edges going out of a vertex v for each vertex v of a given tree.

for all $1 \le i \le k$, $i \le j \le i+l-1$, and (2) $(a \circ_i b) \circ_{j+l-1} c = (a \circ_j c) \circ_i b$

for all $1 \le i < j \le k$.

The parameter n in $\mathcal{P} = \{\mathcal{P}(n)\}_{n\geq 1}$ is referred to as the *arity*. Conditions (1) and (2) hold for the operation of planar rooted trees grafting, and are called the sequential and the parallel associativity, respectively. A morphism $f : \mathcal{P} \to \mathcal{Q}$ of two non- Σ operads in C is a collection of morphisms $f_n : \mathcal{P}(n) \to \mathcal{Q}(n)$ in C, defined for all $n \geq 1$, such that $f_{m+n-1}(a \circ_i b) = f_m(a) \circ_i f_n(b)$ for any $a \in \mathcal{P}(m), b \in P(n), m, n \geq 1$ and $1 \leq i \leq m$. The non- Σ operads in C and their morphisms form a category nsOp(C). There are natural notions of a *suboperad*, one- and two-sided *ideal*, *free operad* and the *quotient operad* extending the corresponding concepts from the realm of monoids and associative algebras. The reader may consult either of the references cited above for a detailed exposition on these notions, if needed. The term $non-\Sigma$, or non-symmetric, is meant to indicated the absence of any equivariance conditions imposed on the partial compositions. A comprehensive account of common equivariance relations emerging for operads can be found in [Yau21].

Example 1. Consider planar rooted trees with the simplest topology, the oriented paths. On such trees, grafting reduces to oriented paths concatenation $(-\circ_1 -)$, which is a monoidal product thereon, and there are no other partial compositions.

More generally, any monoid A in C gives rise to a non- Σ operad \mathcal{P} in C concentrated in arity 1. Namely, we set $\mathcal{P}(1) \coloneqq A$ and $\mathcal{P}(n) \coloneqq \mathbf{0}$ for n > 1, where $\mathbf{0}$ is an initial object in C. The partial composition $\circ_1 : \mathcal{P}(1) \otimes \mathcal{P}(1) \to \mathcal{P}(1)$ is the monoidal product on A, while in all other instances, a partial composition $\circ_i : \mathcal{P}(m) \otimes \mathcal{P}(n) \to \mathcal{P}(m+n-1)$ is, necessarily, the identity morphism $id_{\mathbf{0}}: \mathbf{0} \to \mathbf{0}$. Conversely, given any operad \mathcal{P} in C, by (1), $\mathcal{P}(1)$ is a semi-group in C.

Example 2. The simplest non-trivial example of a non- Σ operad is that of the associative operad As in C. Namely, for $n \ge 1$, we let $As(n) \coloneqq \mathbf{1}$, where $\mathbf{1}$ is a (chosen) monoidal unit of C. Any partial composition $\circ_i : As(m) \otimes As(n) \to As(m + n - 1)$ in this operad is the unit multiplication $\mathbf{1} \otimes \mathbf{1} \to \mathbf{1}$ of C. If a symmetric monoidal category C is pointed, meaning $\mathbf{1}$ is a terminal object of C, this operad is a terminal object in the category nsOp(C). Thus the qualification the simplest given above. The categories Sets and Top with their standard symmetric monoidal structures are pointed, while the category \Bbbk -Vect is not.

In terms of trees, As can be described as the operad, where in each arity $n \ge 1$, the component As(n) contains a single tree with n leaves and no decorations. We take such a unique tree representative to be a planar rooted n-leaf corolla and define partial compositions as grafting followed by an edge contraction. Identification of two trees upon an edge contraction is an example of a *relation*. As a minimal set of *generators* for this operad we can take a single 2-leaf corolla, that can be used to produce all higher-arity corollas by iterated self-compositions, and a 1-leaf corolla that behaves as a neutral element with respect to the $(-\circ_i -)$'s.



FIGURE 6. Evaluating a partial composition in As.

Example 3. As a minimal enhancement, the associative operad As defined above can be given a twist. Let C be cartesian, and let A be a monoid in C. The *semidirect product* $As \rtimes A$ of the associative operad by a monoid A is known as the *word operad* WA. By definition [SW03], the elements of $As \rtimes A$ can be identified with planar corollas, whose leaves are decorated by elements of A, while partial compositions of As are promoted to be homogeneous with respect to these coefficients.



By reading the coefficients off the leaves of a decorated corolla in their canonical order, an element of WA(m) can be identified with a vector, or a *word*, $(a_1, \ldots, a_m) \in A^m$ of elements of A, that can be thought of as a decoration that we assign to the unique vertex of a *m*-leaf planar rooted corolla. In terms of words, the partial compositions take the form

$$(a_1, \ldots, a_m) \circ_i (b_1, \ldots, b_n) = (a_1, \ldots, a_{i-1}, a_i \cdot b_1, a_i \cdot b_2, \ldots, a_i \cdot b_n, a_{i+1}, \ldots, a_m)$$

By virtue of various bijections, multiple classical families of combinatorial objects are know to admit a description in terms of the word operad for different choices of A [Gir15; Bas24].

Example 4. For any $V \in Ob(C)$, the corresponding *endomorphism* operad End_V in C is defined by setting $End_V(n) := \underline{Hom}_C(V^{\otimes n}, V)$ for each $n \ge 1$, where $\underline{Hom}_C(-, -)$ denotes the internal hom in C. The partial compositions are given by the substitution

 $(f \circ_i g)(x_1, \ldots, x_{m+n-1}) = f(x_1, \ldots, x_{i-1}, g(x_i, \ldots, x_{i+n-1}), x_{i+n}, \ldots, x_n).$

Given a non- Σ operad \mathcal{P} in C and $V \in \operatorname{Ob}(C)$, a representation of \mathcal{P} , or a \mathcal{P} -algebra structure on V, is an operad morphism $\mathcal{P} \to End_V$ in nsOp(C). For example, upon taking $C = \Bbbk$ -Vect, a representation of the operad As of example 2 on a \Bbbk -vector space V is equivalent to the data of an associative \Bbbk -algebra on V. Indeed, by definition, for any $n \ge 1$, As(n) is a 1-dimensional vector space. Hence, a non- Σ operad morphism $As \to End_V$ amounts to picking a single \Bbbk -linear map $\mu_n : V^{\otimes n} \to V$ for each $n \ge 1$ in such a way that a system of defining relations $\mu_m \circ_i \mu_n = \mu_{m+n-1}$ of As holds as an equality of \Bbbk -linear maps on $V^{\otimes m+n-1}$ for all $m, n \ge 1$ and $1 \le i \le m$. In particular, $\mu := \mu_2 : V \otimes V \to V$ is a bilinear product subject to the edge-contraction relation

$$\mu \circ_1 \mu = \mu_3 = \mu \circ_2 \mu$$

expressing the usual associativity condition.

Example 5. For $n \ge 1$, let PRT(n) be the set of all (isomorphism classes of) planar rooted trees with n leaves. Tautologically, the tree grafting operation turns the collection $PRT := \{PRT(n)\}_{n\ge 1}$ into a non- Σ operad. For any $k \ge 1$, the subcollection $PRT_k := \{PRT_k(n)\}_{n\ge 1}$ consisting of k-ary planar rooted trees forms a suboperad of PRT. Each of these suboperads PRT_k is generated by a single element, the k-leaf corolla.

Example 6. For $n \ge 1$, let $\mathcal{D}_1(n)$ be the set of all collections of n closed non-trivial subintervals, with pairwise disjoint interiors, embedded into the unit interval I = [0, 1]. Such a set can be given a natural topology as a subspace of $[0, 1]^{2n}$, and subintervals within any configuration $a \in \mathcal{D}_1(n)$ can be enumerated naturally in their order of appearance as I is traced from 0 to 1. For any two subinterval configurations $a \in \mathcal{D}_1(m)$, $b \in \mathcal{D}_1(n)$ and $1 \le i \le m$, the partial composition $a \circ_i b$ is defined by mapping the unit interval containing configuration b onto the *i*-th subinterval of a via an orientation-preserving affine map. This results in a configuration of m + n - 1 subintervals in [0, 1] and, in fact, turns $\mathcal{D}_1 = \{\mathcal{D}_1(n)\}_{n\ge 1}$ into an operad in Top, known as the $(\text{non-}\Sigma)$ operad of *little* 1-*disks*. This operad can be regarded as the topological version of the associative operad As. Indeed,

for each $n \ge 1$, any two subinterval configurations in $\mathcal{D}_1(n)$ can be homotopically transformed to each other, and the subintervals enumeration in any partial composition $a \circ_i b$ matches the one from the definition of As in example 2. This equivalence identifies $\pi_0(\mathcal{D}_1)$ with As viewed as non- Σ operads in Sets.



FIGURE 7. An element of $\mathcal{D}_1(3)$.

To familiarize oneself with partial composition of little disks and the theme of recursion frequently emerging in the context of operads, an interested reader might consider the suboperad of \mathcal{D}_1 generated by a certain element of arity 2 – the configuration $[0, 1/3] \cup [2/3, 1]$. This is a non- Σ operad of pre-Cantor sets, isomorphic to PRT_2 . The Cantor set itself can be used as a left module generator over this non- Σ operad in Top.

Another suboperad of \mathcal{D}_1 that will be particularly useful for us by virtue of encoding static particle configurations on a line is a special case of a tiling operad that we discuss below.

3. Operads of tilings and configuration spaces.

Self-similar tilings of polyhedra (*reptiles* and *irreptiles* [Rei03][Gol66]), provide natural examples of combinatorial operads. For instance, consider tilings of a L-shaped tromino by homothetically rescaled smaller copies thereof.



FIGURE 8. Two self-similar L-shaped tilings.

The arity of a tiling is defined as the number of elementary tiles contained within. A partial composition $a \circ_i b$ of a tiling a of arity m and a tiling b of arity n amounts to substituting a homothetically rescaled, translated and, possibly, rotated copy of b for the *i*-th tile of a. This predicates on the existence of a certain tile-enumerating scheme that allows one to address each tile by its unique index i. We define such a scheme inductively in the following way. For any tile configuration that is indecomposable in the sense of the \circ_i -products, the enumeration of the tiles is a choice to be made (in general, this could be subject to a consistency condition across the indecomposables if there are relations between them). Next, for tilings a, b of arities m and n respectively, and any $1 \le i \le m$, all the tiles of a with indicies j < i retain their respective indicies in $a \circ_i b$. Any tile of a of index j > i changes it to j' := j + n - 1. All the tiles of b, originally indexed $1, 2, \ldots, n$, acquire indicies $i, i + 1, \ldots, i + n - 1$ respectively in $a \circ_i b$; cf. leaves enumeration in $a \circ_i b$ in example 2.

A tiling pattern shown on the left of figure 8 (Golomb's configuration), can be used as a single generator of arity four for a family of self-similar tromino tilings forming a non- Σ operad (in Sets). The enumeration of four tiles within the generator is arbitrary. The enumeration of tiles in all composite configurations follows the scheme described above. The non- Σ operad generated in this manner is isomorphic to the operad of 4-ary planar rooted trees PRT_4 . A concrete isomorphism depends on the choice of the tiles enumeration in the generator. The figure below depicts such a correspondence based on the enumeration of the generator tiles as shown on the right.



FIGURE 9. A tiling generated by Golomb's configuration and the corresponding tree. On the right: the choice of the generator that induces the corresponding operadic bijection.

Note that this is a proper suboperad of the operad of all possible self-similar tromino tilings. For instance, the tiling of arity 6 shown on the right of figure 8[Kho10] is not a composite of Golomb's configurations, and is, in fact, \circ_i -indecomposable.

As a geometric model, a self-similar tiling operad can be presented as a suboperad of the word operad WG, where $G \leq Aut(X)$ is the group of the admissible tile transformations and X is the corresponding ambient space. For the above example of trominos embedded in a Euclidean plane, G can be taken to be the orientation-preserving affine similarity group – the subgroup of the affine group $Aff(\mathbb{R}^2)$ restricted to scaling by s > 0, translations and rotations. To this end, we fix a reference copy of a basic tile T in X, such as a single tromino in a plane as in the above example. We will assume that T has no non-trivial automorphisms in G, by putting some framing on T, if needed. Then any tiling of T consisting of pieces T_1, T_2, \ldots, T_n can be uniquely encoded by the word $(g_1, g_2, \ldots, g_n) \in G^n$, where each g_i satisfies $g_iT = T_i$. These are subject to $\bigcup_{i=1}^n T_i = T$ and $int(T_i) \cap int(T_j) = \emptyset$ for $i \neq j$. Note that the tiles are enumerated, all g_i 's are distinct, and $|\mathbb{S}_n|$ ways of relabeling the tiles correspond to all the letter permutations within the word.

Lemma 1. The family $\mathcal{T} = \{T(n)\}_{n\geq 1}$, where T(n) consists of all tiling words $(g_1, g_2, \ldots, g_n) \in G^n$ as above, is a non- Σ suboperad $\mathcal{W}G$. of It is (non-canonically) isomorphic to the operad of labeled self-similar tilings of T.

Proof. For any
$$(g_1, \ldots, g_m) \in \mathcal{T}(m)$$
, $(h_1, \ldots, h_n) \in \mathcal{T}(n)$ and $1 \le i \le m$, we have

(3)
$$(g_1, \ldots, g_m) \circ_i (h_1, \ldots, h_n) = (g_1, \ldots, g_{i-1}, g_i h_1, \ldots, g_i h_n, g_{i+1}, \ldots, g_m).$$

Multiplying $\bigcup_{i=1}^{n} h_i T = T$ on the left by g_i we deduce that the subword $g_{i-1}, g_i h_1, \ldots, g_i h_n$ yields a valid tiling of $g_i T$. Thus, (3) encodes a valid tiling of T. All the topological conditions are met due to G acting homeomorphically on X.

Non-canonicity is due to making a choice of a template tile $T \subset X$. For any other choice gT, the tiling operad is $gTg^{-1} = \{gT(n)g^{-1}\}_{n\geq 1}$, where G acts on each T(n) diagonally.

3.1. Operads of static particle configurations. Consider a particular one-dimensional instance of the self-similar tiling operad. Namely, let C be the operad of tilings of the closed unit interval I = [0,1] by subintervals. Equivalently, C is the suboperad of the little 1-disks operad \mathcal{D}_1 of example 6 that consists of all maximally stretched-out subinterval configurations in [0,1]. These are collections of closed non-degenerate subintervals $I_1, \ldots, I_n \subseteq I$ with pairwise disjoint interiors,

such that $\bigcup_{i=1}^{n} I_i = I$. As a topological space, C(n) can be identified with the configuration space $UConf_{n-1}(0,1)$ of n-1 unlabeled points in (0,1), where pairwise-adjacent intervals I_i, I_{i+1} meet.



FIGURE 10. An element of $UConf_2(0,1) \simeq C(3)$.

The family of configuration spaces $UConf(0,1) = \{UConf_{n-1}(0,1)\}_{n\geq 1}$ acquires the structure of a non- Σ operad as a suboperad of the little 1-disks operad through a bijective correspondence with C.



FIGURE 11. Evaluating a partial composition in C.

Note that a configuration of n-1 points in (0,1) is an element of arity n of this operad. Furthermore, despite being unlabeled, the points come with a canonical enumeration induced by the usual order on [0,1]. The lengths $\lambda_1, \ldots, \lambda_n$ of the consecutive tiling intervals I_1, \ldots, I_n define a coordinate system on $UConf_{n-1}(0,1)$, which identifies the latter with the interior $int(\Delta^{n-1})$ of the standard geometric (n-1)-simplex. In fact, it identifies $UConf_{n-1}(0,1)$ with a suboperad the word operad $\mathcal{W}[0,1]$, where [0,1] with its multiplicative structure is treated as a topological monoid. The partially-defined addition on [0,1] allows one to cut $UConf_{n-1}(0,1)$ out of the cube $\mathcal{W}[0,1](n) :=$ $[0,1]^n$ by means of an affine equation $\lambda_1 + \cdots + \lambda_n = 1$ with $\lambda_i > 0$ for all $1 \le i \le n$. Non- Σ operad \mathcal{C} admits a family of binary generators $P_{\lambda}, 0 < \lambda < 1$. In terms of interval tilings, P_{λ} 's are two-interval configurations of the form $[0,\lambda] \cup [\lambda,1]$, or equivalently, a point $0 < \lambda < 1$ in $UConf_1(0,1)$. These generators are subject to the barycentric relations

(4)
$$P_{\alpha+\beta} \circ_1 P_{\frac{\alpha}{\alpha+\beta}} = P_{\alpha} \circ_2 P_{\frac{\beta}{1-\alpha}}$$

for any $\alpha, \beta > 0$, $\alpha + \beta < 1$. Any convex set $X \subset \mathbb{R}^N$ is a *C*-algebra in Top.

For our purposes, we need an operad structure for particle configurations on the entire line \mathbb{R} . One way to get it is to transfer an operad structure from UConf(0,1) to $UConf(\mathbb{R})$ along a homeomorphism $(0,1) \simeq \mathbb{R}$. On the other hand, a more natural approach that circumvents having to choose such a homeomorphism is to use the specifics of the line geometry. Namely, since any finite point configuration Q in \mathbb{R} is contained in an interval $[Q_1, Q_n]$, where Q_1 and Q_n are the well-defined leftmost and rightmost points of Q respectively, there is a unique orientation-preserving affine map sending $[Q_1, Q_n]$ to any interval $[P_1, P_m]$ by mapping Q_1 to P_1 and Q_n to P_m ; cf. figure 3.

Proposition 1 provides a way to formalize it. This model serves as a precursor to the main construction of section 4. To this end, let G be the orientation-preserving affine group $Aff_+(\mathbb{R}^1)$. Then a tiling of I = [0, 1] by subintervals I_1, \ldots, I_n of lengths $\lambda_1, \ldots, \lambda_n$ can be identified with the tuple

(5)
$$\left(\begin{bmatrix} \lambda_1 & b_1 \\ 0 & 1 \end{bmatrix}, \dots, \begin{bmatrix} \lambda_n & b_n \\ 0 & 1 \end{bmatrix} \right) \in G^n$$

subject to the boundary matching and the normalization conditions that read

(6)
$$\begin{bmatrix} \lambda_i & b_i \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \lambda_{i+1} & b_{i+1} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad 1 \le i \le n-1$$

and

(7)
$$\begin{bmatrix} \lambda_1 & b_1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} \lambda_n & b_n \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

respectively. Lemma 1 identifies \mathcal{C} with the suboperad of $\mathcal{W}G$ consisting of all such tuples.

Upon omitting the normalization condition (7), a tuple (5) can be identified with a tiling of an arbitrary interval $[b_1, b_n + \lambda_n] \subset \mathbb{R}$ by *n* subintervals $[b_1, b_2], \ldots, [b_{n-1}, b_n], [b_n, b_n + \lambda_n]$. Equivalently, this is a configuration of n+1 points $b_1, b_2, \ldots, b_n, b_n+\lambda_n$ in \mathbb{R} . At the same time, the partial compositions $(-\circ_i -)$ of the word operad $\mathcal{W}G$ no longer descent onto the set of such tuples. Indeed, as an example, both $a = \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \right)$ and $b = \left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \right)$ satisfy the matching condition,

but do not satisfy the normalization one. Upon computing $a \circ_1 b = \left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \right)$ we find out that it is not a valid tiling word. Indeed, the matching condition for the 2nd and 3rd entries does not hold (also, can be seen from having two equal entries).

The partial compositions are to be modified, and this can be done by employing a variation of the moving frame method [Olv03].

Proposition 1. Let $\mathcal{Q} = {\mathcal{Q}(n)}_{n \geq 1}$ be a collection of objects of a category C with a group G acting on each component $\mathcal{Q}(n)$, $\rho = \{\rho_n : \mathcal{Q}(n) \to G\}$ be a family of equivariant maps, $\rho_n(g \cdot z) = g \cdot \rho_n(z)$ for any $n \ge 1$, $z \in \mathcal{Q}(n)$, $g \in G$. For $n \ge 1$, let $\mathcal{P}(n) \coloneqq \{z \in \mathcal{Q}(n) | \rho(z) = e\}$ If $\mathcal{P} = \{\mathcal{P}(n)\}_{n \ge 1}$ has the structure of a non- Σ operad, this structure can be lifted to one on Q. Namely, for $a \in Q(m)$, $b \in \mathcal{Q}(n), 1 \leq i \leq m, we set$

(8)
$$a *_i b \coloneqq \rho(a) [\rho(a)^{-1} a \circ_i \rho(b)^{-1} b]$$

In words, we evaluate a composition within a distinguished reference frame associated with \mathcal{P} and then state the result in the reference frame of a in \mathcal{Q} .

Proof. First, note that $a *_i b$ is well-defined. Indeed, for any $z \in \mathcal{Q}$, we have $\rho(\rho(z)^{-1}z) = \rho(z)^{-1}\rho(z) = \rho(z)^{-1}\rho(z)$ e. Thus, $\rho(z)^{-1}z \in \mathcal{P}$, and the partial composition in the brackets is well-defined. Next, observe that

$$\rho(a *_i b) = \rho(\rho(a)[\rho(a)^{-1}a \circ_i \rho(b)^{-1}b]) = \rho(a)\rho(\underbrace{\rho(a)^{-1}a \circ_i \rho(b)^{-1}b}_{\in \mathcal{P}}) = \rho(a)$$

The last equality is due to $\rho(z) = e$ for any $z \in \mathcal{P}$.

Now, let $a, b, c \in \mathcal{Q}$ and i, j be as in the hypothesis of the sequential associativity axiom (1). Then

$$(a *_{i} b) *_{j} c = \rho(a *_{i} b) [\rho(a *_{i} b)^{-1}(a *_{i} b) \circ_{j} \rho(c)^{-1} c] = \rho(a) [\rho(a)^{-1}(a *_{i} b) \circ_{j} \rho(c)^{-1} c]$$

= $\rho(a) [(\rho(a)^{-1}a \circ_{i} \rho(b)^{-1}b) \circ_{j} \rho(c)^{-1}c] \stackrel{(1)}{=} \rho(a) [\rho(a)^{-1}a \circ_{i} (\rho(b)^{-1}b \circ_{j-i+1} \rho(c)^{-1}c)]$
= $\rho(a) [\rho(a)^{-1}a \circ_{i} \rho(b)^{-1}(b *_{j-i+1} c)] = \rho(a) [\rho(a)^{-1}a \circ_{i} \rho(b *_{j-i+1} c)^{-1}(b *_{j-i+1} c)] = a *_{i} (b *_{j-i+1} c).$

The proof of parallel associativity is analogous and differs only by applying (2) instead of (1) in the second line above.

In our case, we take $\mathcal{Q}(n)$ to be the set of all tuples (5) subject to the boundary matching condition (6) for all $n \ge 1$. The set is taken with the diagonal left action by $G = Aff_+(\mathbb{R}^1)$. The

11

normalization map $z \mapsto \rho(z)^{-1}z$ is defined by taking $\rho_n(z) \coloneqq \begin{bmatrix} \lambda_1 + \dots + \lambda_n & b_1 \\ 0 & 1 \end{bmatrix}$, where z is as in (5). The corresponding cross-section operad \mathcal{P} is the operad \mathcal{C} from above. Note that points 0 and 1 are no longer distinguished in \mathcal{Q} . For this reason, any elements of $\mathcal{Q}(n)$ is identified with a configuration of n + 1 points in \mathbb{R} . In what follows, we will denote this non- Σ operad \mathcal{Q} by $\widehat{\mathcal{C}}$.



FIGURE 12. Evaluating a partial composition in \mathcal{C} .

As an upshot, starting with an non- Σ operad \mathcal{P} and a group G, there is a way to extend \mathcal{P} to a larger G-equivariant operad \mathcal{Q} . Such an extension is parametrized by a choice of a moving frame $\rho = \{\rho_n : \mathcal{Q}(n) \to G\}_{n \ge 1}$. This will be used again in the next section.

4. An operad of planar rooted line arrangements

Let L be a collection of n + 1 lines $\ell_0, \ell_1, \ldots, \ell_n$ embedded into an oriented Euclidean plane \mathbb{R}^2 . Assume that line ℓ_0 is given a direction. The open half-plane that lies to the left of ℓ_0 with respect to this direction will be referred to as the *upper* half-plane of L.

Definition 2. We say that a collection of lines $L = \{\ell_0, \ell_1, \ldots, \ell_n\}$ in \mathbb{R}^2 is a *planar rooted line* arrangement of rank n if

- (1) no two lines among $\ell_0, \ell_1, \ldots, \ell_n$ are parallel or overlapping;
- (2) for any two lines ℓ_i , ℓ_j , such that $i, j \neq 0$ and $i \neq j$, the point of intersection $\ell_i \cap \ell_j$ lies in the upper half-plane;
- (3) the order in which the points of intersection $L_i := \ell_0 \cap \ell_i$, for $i \ge 1$, appear on the oriented line ℓ_0 as we trace it in the positive direction agrees with the natural order on the line indices $1, \ldots, n$.

The distinguished line ℓ_0 will be called the *root* of an arrangement L.

Note that the lines $\ell_0, \ell_1, \ldots, \ell_n$ of L are not assumed to be in a general position. A planar rooted line arrangement may contain a subarrangement of three, or more, lines intersecting at a single point. As an immediate observation, note that by condition (2) of the above definition, an orientation of the root line ℓ_0 of L induces a canonical orientation on each line ℓ_i for $i \ge 1$. Specifically, for $i \ge 1$, a directional vector of ℓ_i stemming out of L_i is chosen to lie in the upper-half plane with respect to ℓ_0 .



FIGURE 13. A planar rooted line arrangement of rank 4.

In terms of scattering diagrams, the root line ℓ_0 represents the one-dimensional space at time t = 0. The remaining n lines represent the worldline asymptotes of n particles undergoing a scattering process in transition presentation. For $n \ge 1$, the set of all rank-n planar rooted line arrangements is a real 2n-dimensional manifold. As a possible set of global coordinates one can take positions of L_i on ℓ_0 and the oriented angles θ_i from ℓ_0 to ℓ_i for $1 \le i \le n$, or momenta $p_i = \operatorname{ctg}(\theta_i)$ identifying this 2n-dimensional manifold with with $UConf_n(\mathbb{R}) \times UConf_n((0,\pi))$, or $UConf_n(\mathbb{R}) \times UConf_n(\mathbb{R})$ respectively. Here, as before, $UConf_n(X)$ is the configuration space of n unlabeled, but naturally ordered in our case, points in X. Note that in the copy of $UConf(\mathbb{R})$ for the momenta the order of point enumeration is reversed – the slower particles are further right. Equivalently, we may think that to each particle we assign its negative momentum.

In what follows, we will consider two examples of a non- Σ operad structure on planar rooted line arrangements. We start off with a simpler one, but it is the second one – affine-geometric, that will be of particular interest to us.

4.1. Construction \mathbf{I} – decoupling positions and momenta. As we have seen in the previous section, configuration spaces $UConf_n(\mathbb{R})$ assemble into an operad \widehat{C} , where $UConf_n(\mathbb{R}) = \widehat{C}(n-1)$ for $n \geq 2$. Then the coordinate presentation $UConf_n(\mathbb{R}) \times UConf_n(\mathbb{R})$ of rank-*n* arrangements in terms of positions and negative momenta yields a non- Σ operad structure on the space of all planar rooted line arrangements isomorphic to $\widehat{C} \times \widehat{C}$. Again, note that a rank-*n* arrangement is identified with an element of $\widehat{C}(n-1) \times \widehat{C}(n-1)$. Another model thereof is an operad of polygonal chains.

For a rank-(n + 1) planar rooted line arrangement, or equivalently, for an initial configuration of n + 1 particles, consider the polygonal chain $P_1P_2 \dots P_{n+1}$ in \mathbb{R}^2 , where $P_i = (q_i, -p_i)$, the initial position and the negative momentum of the *i*-th particle. Note that both the q and the $-p_i$ coordinates are strictly monotonically increasing for $1 \le i \le n$. For all $n \ge 2$, let us define $\mathcal{P}(n)$ to be the set of all polygonal chains in the qp-plane with this property with n + 1 vertices. For any two polygonal chains $P \in \mathcal{P}(n), Q \in \mathcal{P}(m)$ and $1 \le i \le n$, we define their partial composition $P \circ_i Q$ as follows. The operation amounts to replacing the *i*-th line segment P_iP_{i+1} with a rescaled and translated copy of Q so that Q_1 maps to P_i and Q_{m+1} maps to P_{i+1} . Here, rescaling stands for the transformation $(q, p) \mapsto \left(\frac{q(P_{i+1})-q(P_i)}{q(Q_{m+1})-q(Q_1)} \cdot q, \frac{p(P_{i+1})-p(P_i)}{p(Q_{m+1})-p(Q_1)} \cdot p\right)$, where q(K) and p(K) denote the q,p-coordinates of a point K respectively. This path substitution operation turns \mathcal{P} into a non- Σ operad isomorphic to $\widehat{C} \times \widehat{C}$. Correspondingly, this operad is generated by two topological families of arity-two generators subject to quadratic relations. The latter are translated and rescaled barycentric relations (4).





FIGURE 14. Evaluating $P \circ_3 Q \in \mathcal{P}(6)$.

4.2. Construction II – using affine symmetries. For $n \ge 1$, let $\widehat{\mathbb{L}}(n)$ be the space of all rooted line arrangements of rank n+1. Let $m, n \ge 1$, $P \in \widehat{\mathbb{L}}(m)$, $Q \in \widehat{\mathbb{L}}(n)$ and $1 \le i \le m$. Consider the points of intersection of the following pairs of lines of P: (p_0, p_i) , (p_i, p_{i+1}) , (p_0, p_{i+1}) , and denote them by P_i , $P_{i,i+1}$, P_{i+1} respectively. Similarly, let Q_1 , $Q_{1,m+1}$, Q_{m+1} denote, in their respective order, the points of intersection of the following pairs of lines: (q_0, q_1) , (q_1, q_{m+1}) , (q_0, q_{m+1}) . Here, as per our usual notation, p_0 and q_0 denote the root lines of P and Q respectively. Note that neither of these two triples of points is collinear. Then, by the fundamental theorem of affine geometry, there exists a unique orientation-preserving affine transformation $T_i \in Aff(\mathbb{R}^2)$ that sends the ordered triple $(Q_1, Q_{1,m+1}, Q_{m+1})$ to $(P_i, P_{i,i+1}, P_{i+1})$. Such an affine transformation sends the lines q_0, q_1 and q_{m+1} to p_0 , p_i and p_{i+1} respectively.



FIGURE 15. Evaluating $P *_2 Q \in \widehat{L}(4)$ for $P \in \widehat{L}(3)$, $Q \in \widehat{L}(2)$.

Lemma 2. Let $P \cup T_i(Q)$ be the union of the line arrangement P and the lines in the image of the arrangement Q under T_i . Then, upon appropriate enumeration of the lines, $P *_i Q \coloneqq P \cup T_i(Q)$ is a well-defined planar rooted line arrangement of rank n + m - 1.

Proof. First, since an affine transformation T_i is bijective and preserves the property of lines being parallel, no two lines in the image $T_i(Q)$ are overlapping or parallel to each other. Furthermore, the angles that the lines in the image $T_i(Q)$ make with the root line of arrangement P are strictly in between the angle θ_i between the line p_i and p_0 and the angle θ_{i+1} between the line p_{i+1} and p_0 in arrangement P. Hence, no line in the image $T_i(Q)$ can be parallel to any of the lines of P. This verifies condition (1) of Definition 2.

By the same set of inequalities for the angles, any line in the image $T_i(Q)$ intersects any line of P in the upper half-plane. Since T_i respects the orientations of the root lines p_0 and q_0 , all the intersections between the lines in $T_i(Q)$ lie in the upper-half plane as well. This verifies condition (2) of Definition 2.

To make condition (3) hold, we enumerate the lines in $P \cup T_i(Q) = \{r_1, r_2, \dots, r_{n+m-1}\}$ by setting

$$r_{j} \coloneqq \begin{cases} p_{j}, & j \leq i \\ T(q_{j-i+1}), & i < j < i+m . \\ p_{j-m+1}, & i+m \leq j \end{cases}$$

We will verify that the partial compositions $(-*_i -)$ satisfy the generalized associativity conditions.

Theorem 1. The collection $\widehat{L} := {\widehat{L}(n)}_{n \ge 1}$ with partial compositions $P *_i Q$ defined above is a non- Σ operad in Top.

We will prove the statement for *normalized* planar rooted line arrangements L first, and will extend it to $\widehat{\mathbf{L}}$ by means of proposition 1. The operations $(-*_i -)$ defined on $\widehat{\mathbf{L}}$ above will be identified as the partial compositions of this extended operad. To this end, for $n \ge 1$, let $\mathbf{L}(n)$ be the set of all planar rooted line arrangements P of rank n + 1 bounded by the lines t = 0, q = 0 and q + t = 1in the qt-plane. These are the root line p_0 , the first line p_1 and the last line p_{n+1} of any $P \in \mathbf{L}(n)$ respectively. That is, L consists of the scattering diagrams drawn in the rest frame of the first particle, with units chosen in such a way that the last particle approaches the first one at the unit speed from the right.

Lemma 3. Let $L := \{L(n)\}_{n \ge 1}$ and $(-\circ_i -)$ denote the restriction of the operation $(-*_i -)$ defined above onto L. Then L with these partial compositions is a non- Σ operad.

Proof. We will identify L as a suboperad of a certain word operad WG, similarly to how we did it for static particle configurations. The relevant group for us is the four-dimensional abelian subgroup

 $G ext{ of } Aff(\mathbb{R}^2) ext{ that can be identified with the group of matrices of the form } T = \begin{bmatrix} * & * & * \\ 0 & * & 0 \\ 0 & 0 & 1 \end{bmatrix}, ext{ where }$

all the entries, except possibly $T_{1,2}$, are non-negative, and $T_{1,1} > 0$.

Consider a fixed template – the planar rooted line arrangement of rank 2 consisting of the lines t = 0, q = 0, q + t = 1 that we denote by τ_0, τ_L and τ_R respectively. Then any planar rooted line arrangement $P \in L(n)$ can be uniquely encoded by the tuple (T_1, \ldots, T_n) , where T_i is the affine map sending points (0,0), (1,0), (0,1) to P_i, P_{i+1} and $P_{i,i+1}$ respectively; we use the notation as in lemma 2. That is, T_i keeps track on the point on intersection of the lines p_i and p_{i+1} .



Concretely, $T_i = \begin{bmatrix} a_{i+1} - a_i & b_i - a_i & a_i \\ 0 & c_i & 0 \\ 0 & 0 & 1 \end{bmatrix}$ for $P_i = (a_i, 0), P_{i+1} = (a_{i+1}, 0), P_{i,i+1} = (b_i, c_i)$. Note that

the T_i 's are dependent. Namely, any tuple (T_1, \ldots, T_n) encoding a planar rooted line arrangement is subject to the matching and normalization conditions akin to (6), (7). The matching condition reads

(9)
$$T_i \tau_R = T_{i+1} \tau_L, \quad 1 \le i < n.$$

Equivalently, it can be restated as collinearity of points $(a_{i+1}, 0), (b_i, c_i), (b_{i+1}, c_{i+1})$, where the last two points can be the same. The normalization condition amounts to

(10)
$$T_1\tau_L = \tau_L, T_n\tau_R = \tau_R.$$

Preservation of the root line, $T_i \tau_0 = \tau_0$, is automatic by the form of T_i . Such a tuple (T_1, \ldots, T_n) is an element of the word operad $\mathcal{W}G$, where G is as defined above. The geometric definition of a composition $P *_i Q$ of planar rooted line arrangements translates to

(11)
$$(T_1, \dots, T_n) \circ_i (S_1, \dots, S_m) = (T_1, \dots, T_{i-1}, T_i S_1, \dots, T_i S_m, T_{i+1}, \dots, T_n)$$

in terms of tuples in $\mathcal{W}G$. It remains to show that the set of all tuples subject to the matching and normalization conditions stated above is closed in $\mathcal{W}G$.

The matching conditions for the subranges T_1, \ldots, T_{i-1} and T_{i+1}, \ldots, T_n on the right-hand side of (11) follow immediately from the matching conditions for (T_1, \ldots, T_n) . The matching condition for T_iS_1, \ldots, T_iS_m follows upon multiplying sides of $S_j\tau_R = S_{j+1}\tau_L$ for $1 \le j < m$ by T_i on the left. In the remaining two edge cases, by matching of T_i 's and the normalization condition for S_1, \ldots, S_n we have

$$T_{i-1}\tau_R = T_i\tau_L = T_i(S_1\tau_L) = (T_iS_1)\tau_L$$

and similarly for the adjacent entries $T_i S_m, T_{i+1}$. The normalization condition on the right-hand side of (11) follows directly from the one holding for T_1, T_n and S_1, S_m .

Proof (of theorem 1). The argument is analogous to the one used for non- Σ operad \widehat{C} of static particle configurations discussed in the preceding section. First, note that any planar rooted line arrangement $P \in \widehat{L}(n)$ is determined by a tuple $(T_1, \ldots, T_n) \in WG(n)$ subject to the matching condition (9), but with no regard for the normalization (10). The tuples of this form are not closed under the partial compositions of the word operad WG. A correction is provided by proposition 1 upon relating \widehat{L} to non- Σ operad L of normalized planar rooted line arrangements defined above.

Indeed, since (9) remains valid upon multiplying both sides by any $g \in G$ on the left, $\widehat{L}(n)$ is naturally acted on by G for any $n \geq 1$. The moving frame $\rho_n : \widehat{L}(n) \to G$ that normalizes \widehat{L} onto L is defined by assigning to $P \in \widehat{L}(n)$ the unique affine transform that sends points (0,0), (1,0), (0,1) to P_1, P_{n+1} and $P_{1,n+1}$ respectively. Note that it always exists and can be worked out explicitly as a 3by-3 matrix, even though the resulting formula is cumbersome. Now, an application of proposition 1 extends the non- Σ operad structure provided by lemma 3 from L to \widehat{L} . The geometric description of $P *_i Q$ stated before matches (8) provided by proposition 1.

Corollary 1. Let F be any planar rooted line arrangement. Then all planar rooted line arrangements containing F as a subarrangement form a suboperad of $\widehat{\mathcal{L}}$. Indeed, since a partial composition $P *_i Q$ amounts to only drawing new lines on P, then $F \subset P$ persists in $P *_i Q$.

By identifying a rank-*n* planar rooted line arrangement with the data of a purely elastic scattering process of *n* particles, the construction of the $\widehat{\mathcal{L}}$ yields an example of an operad structure on initial particle configurations that we have been after. By theorem 1, to compute a composition of initial particle configurations $a \circ_i b$, we switch to the normalized rest frame of the first particle a_1 , do the work there according to the affine-geometric rules prescribed by operad L, and return the answer

back covariantly in the reference frame of a. This is what $\widehat{\mathcal{L}}$ does. Naturally, thanks to the natural G-action on $\widehat{\mathcal{L}}$, the result can be returned in the reference frame of b, or elsewhere.

Lemma 4. Non- Σ operad \widehat{L} is generated by the symmetry group $\widehat{L}(1) = G$ and three topological families of binary operations corresponding to three possible combinatorial types of rank-3 arrangements, shown below.



FIGURE 16. Combinatorial types of rank-3 arrangements.

Proof. The proof amounts to observing that evaluating a partial composition $Q *_i M$ with either one of the line arrangements M on figure 16 amounts to drawing a new line in between the lines q_i and q_{i+1} of Q. Any diagram with n > 2 lines can be produced in this way by starting with a planar root arrangement of rank 2 and consecutively adding more lines one by one. More formally, let $P \in \widehat{L}(n)$ for n > 2 and $P' = \{p_0, p_1, \ldots, p_{n-1}, p_{n+1}\}$ be the arrangement obtained by removing the penultimate line p_n of P. Then $P' \in \widehat{\mathcal{L}}(n-1)$, and $P = P' *_{n-1} M$, where $M = \{p_0, p_{n-1}, p_n, p_{n+1}\}$. The latter is necessarily of one of the above forms, and induction applied to P' yields the result. \Box

5. Symmetries of $\widehat{\mathcal{L}}$ and factorized scattering

Let us address the question of what symmetries can non- Σ operad of scattering diagrams $\widehat{\mathcal{L}}$ support. In what follows, G denotes the subgroup of $Aff(\mathbb{R})$ described in lemma 3. First, there is a global gauge action by adjunction $g\widehat{\mathcal{L}}g^{-1}$, where $g \in Aff(\mathbb{R}^2)$ acts on each $\widehat{\mathcal{L}}(n) \in G^n \leq Aff(\mathbb{R}^2)$ diagonally; cf. lemma 1. Also, partial compositions commute with translations along the q-axis, or for that matter, with multiplication action by $g \in G$. In our non-relativistic setting the crossing symmetry does not arise. As far as the CPT goes, no charge conjugation is defined, since the lines of a planar rooted arrangements as such are not assumed to have any internal degrees of freedom. That said, one may consider a semidirect product $\widehat{\mathcal{L}} \rtimes H$, where H is the symmetry group of a particle. Note that in this setting the charges would be defined up to a global phase. Indeed, an element of $(\widehat{\mathcal{L}} \rtimes H)(n)$ is a rank-(n + 1) line arrangement with n values of H distributed among n + 1 lines (particles).

The *P*-symmetry amounts to switching the direction of the root line, or equivalently, to flipping a diagram about the *t*-axis. The equivariance with respect to this involution π reads $\pi(P *_i Q) = \pi(P) *_{m-i+1} \pi(Q)$ for any $P \in \widehat{\mathcal{L}}(m)$, $Q \in \widehat{\mathcal{L}}(n)$, $1 \le i \le m$. In fact, such a symmetry exists in any word operad $\mathcal{W}M$, where π reverses a word.

The *T*-symmetry can be introduced upon extending our definition of a planar rooted line arrangements to configurations where line intersections may appear in both the upper and the lower half-planes (but not on the root line). The symmetry group *G* should be modified appropriately; the construction of $\hat{\mathcal{L}}$ remains the same. The time reversal τ amounts, in such a setting, to flipping a diagram vertically about the *q*-axis. The equivariance reads $\tau(P \circ_i Q) = \tau(P) \circ_{m-i+1} \tau(Q)$ for P,Q as above.

A crucial property for integrability of both classical and quantum pure elastic scattering processes are good covariance properties of scattering data with respect to *Baxter's Z-symmetry*. By

definition, Z-symmetry on $P \in \widehat{\mathcal{L}}(n)$ amounts to shifting any line of P parallel to itself without violating the ordered momenta condition. For instance, different combinatorial types of rank-3 arrangements, as shown on figure 16, can be related to each other by such a transformation. More generally, using Z-symmetry, any rank-(n + 1) arrangement $P \in \widehat{\mathcal{L}}(n)$ can be transformed to an arrangement of n+1 lines intersecting at a single point. We can describe all of them in the following way. Let ℓ_0 be a root line and A be a point in the upper half-plane. Since affine maps preserve intersections, the space $\widehat{\mathcal{L}}_A$ of all planar rooted line arrangements consisting of lines intersecting at A is a suboperad of $\widehat{\mathcal{L}}$. Moreover, since affine maps preserve distance ratios along a single line, $\widehat{\mathcal{L}}_A$ is isomorphic to the operad of static particle configurations $\widehat{\mathcal{C}}$. Indeed, all the momenta can be recovered from the particle positions and A.



FIGURE 17. Evaluating $P *_2 Q \in \widehat{L}_A(4)$ for $P \in \widehat{L}_A(3)$, $Q \in \widehat{L}_A(2)$.

Since parallel translations commute with affine maps, there is a surjective non- Σ operad morphism $p: \widehat{\mathcal{L}} \to \widehat{\mathcal{L}}_A \simeq \widehat{\mathcal{C}}$ for any A in the upper half-plane. This is the Z-symmetry transformation of P into a configuration of lines intersecting at A described above. These symmetries are useful if we would like to study invariants of planar rooted line arrangements. In terms of our physical model, a quest for invariants can be formalized as follows.

Definition 3. Let C be a symmetric monoidal category subject to our usual technical constraints, and \mathcal{P} be a non- Σ operad in C. A *C*-valued (1+1 dimensional purely elastic) scattering theory is a non- Σ operad morphism $s: \widehat{\mathcal{L}} \to \mathcal{P}$ over Sets.

By lemma 4, defining s amounts to specifying s on all planar rooted line arrangements of ranks 2 and 3. Line arrangements of rank 2 is the monoid $\widehat{\mathcal{L}}(1)$, which is identified with our affine symmetry group $G \leq Aff(\mathbb{R}^2)$. So $\mathcal{P}(1)$ contains a homomorphic image thereof. To handle rank-3 arrangements, we can impose, for the sake of simplicity, the Z-symmetry and consider scattering theories that do not distinguish between Z-symmetric scattering diagrams of different combinatorial types shown on figure 16. These are scattering theories where multiparticle collisions can be reduced to pairwise interactions. Specifically, we will say that s has the Yang-Baxter property if s factorizes through p:



By the global gauge action on $\widehat{\mathcal{L}}$ this is independent from the choice of A. As a concrete implementation, a k-Vect-valued scattering theory in the above sense can be constructed from the data of an R-matrix, or equivalently, by recasting Zamolodchikov-Faddeev's algebras in operadic terms.

6. Outlook

A merit of the operadic approach to the subject is in its functoriality. This becomes relevant if we are interested in generalizations and deformations of scattering theories. In particular, it could be interesting to look for scattering theories $\widehat{\mathcal{L}} \to \mathcal{P}$, where \mathcal{P} is valued in a category with good homotopy properties, as well as to modify the source operad $\widehat{\mathcal{L}}$, for instance, by considering pseudoline arrangements with some additional data that would yield a reasonable operad structure thereon. One may look for scattering theories associated to various combinatorial and algebraic invariants of line arrangements or configuration spaces of points on a line. We briefly elaborate on these points below.

6.1. Related polytopes. A construction that we may naturally associate to a planar rooted line arrangement is a polytopal chain in a permutahedron. Indeed, in transmission presentation, a scattering process of n particles can be thought of as a path in \mathbb{R}^n with the braid hyperplane arrangement. A permutahedron is a dual object thereof.

More specifically, for any t > 0, the cross section of a scattering diagram by a horizontal line t = const is, generically, a configuration of n distinct points in \mathbb{R} , but there may (and will, for n > 1) be up to $\frac{n(n-1)}{2}$ special values of t corresponding to the time moments when a lines intersection, or a multiple lines intersections, take place for some t. Any collision pattern at moment t is described by a partition $I(t) = \{I_1(t), I_2(t), \ldots, I_k(t)\}$ of the set $\{1, 2, \ldots, n\}$, where a block $I_j(t)$ of cardinality $|I_j(t)| > 2$ registers the fact of a simultaneous collision of particles $\alpha \in I_j(t)$ at moment t, or equivalently, represents a point of common intersection of the lines ℓ_{α} for $\alpha \in I_j(t)$. In particular, generically, in absence of collisions, I(t) is the finest partition $\{\{1\}, \{2\}, \ldots, \{n\}\}$. The simultaneous collision of all n particles is represented by the trivial single block partition. In a general position, any planar rooted lines arrangement has $\frac{n(n-1)}{2}$ singular values of t, when I(t) is a partition with n-2 blocks of size 1 and one extra block $\{i, j\}$.

Furthermore, for any t > 0, the partition I(t) carries an extra structure, and in fact can be equipped with a natural total order on the set of blocks. Indeed, invoking the orientation on the ambient \mathbb{R}^2 , we can define I' < I'' for $I', I'' \in I(t)$ iff on the horizontal cross section of a scattering diagram for a given t the intersection of lines indexed by I' appears strictly to the left of the intersection of the lines indexed by I''. Now, the set of all ordered partitions of $\{1, 2, \ldots, n\}$ is a lattice that admits a polytopal realization as the face lattice of the (n-1)-dimensional permutahedron P_{n-1} . Indeed, this is one of the known compactifications of the configuration space of points in a closed interval [LTV10]. Thus a scattering diagram gives rise to a polytopal chain in P_{n-1} that starts at the vertex ($\{1\}, \{2\}, \ldots, \{n\}$) and arrives at the vertex ($\{n\}, \{n-1\}, \ldots, \{1\}$). For a rank n planar rooted line arrangement in a general position, such a chain will traverse only the edges and the vertices of P_{n-1} . This corresponds to having strictly pairwise collisions. As an example, the three combinatorially distinct rank 3 planar rooted line arrangements shown on figure 16 give rise to three paths on a hexagon P_2 as shown on figure 18.



FIGURE 18. Three chains in P_2 .

The two sides of the Yang-Baxter identity $S_{12}S_{13}S_{23} = S_{23}S_{13}S_{12}$ are encoded by two paths, consisting of three edges each, connecting the vertices $\{1\}\{2\}\{3\}$ and $\{3\}\{2\}\{1\}$. The scattering scenarios with only pairwise interactions are parametrized by reduced decompositions of the longest element $n(n-1)\ldots 1$ of \mathbb{S}_n .

6.2. Relation to the Swiss-cheese operad. Given a line arrangement $P \in \widehat{\mathcal{L}}(n)$, we define its upper envelope U_P as the unique polygonal chain made of the segments lying on the lines of P such that all points of intersection of the lines of P lie in the closed region bounded by U_P and the root line. That is, constructing U_P amounts to going along the lines of P, starting at point P_1 and finishing at P_{n+1} , in such a way that all points of intersection of the lines of the lines of P remain on the right hand side with respect to the direction of motion.

For a given $P \in \widehat{\mathcal{L}}(n)$, there exists a sufficiently small $\epsilon > 0$ such that thickening each line segments of U_P to a rectangular band of width ϵ results in a subset P_{ϵ} of \mathbb{R}^2 homeomorphic [this needs to be refined] to a half disk with n half-disks at the bottom cut out and with m holes in the bulk, where m is the number of the connected bounded subsets of $\mathbb{R}^2 \setminus P$. In particular, $m = \frac{n(n-1)}{2}$ if P is in a general position.



FIGURE 19. Morphing a planar rooted line arrangement of rank 4 to an element of SC(3,3).

REFERENCES

That is, $P \in \widehat{\mathcal{L}}(n)$ gives rise (non-uniquely) to an element $P_{\epsilon} \in \mathcal{SC}(n,m)$ of the Swiss-cheese operad $\mathcal{SC}[Vor99]$.

Acknowledgments. The author is grateful to Branislav Jurčo, Tomáš Procházka, Dmitry Roytenberg and Alexander Voronov for stimulating communications. The work is supported by RVO:67985840 of the Institute of Mathematics of the Czech Academy of Sciences and the Praemium Academiae grant of M. Markl.

References

- [Aru19] G. Arutyunov. Elements of Classical and Quantum Integrable Systems. 2019. DOI: 10. 1007/978-3-030-24198-8.
- [Bas24] D. Bashkirov. "Lattice operads and operad filtrations". Version v3 (2024). arXiv:2212.14833 [math.RA]. arXiv: 2212.14833 [math.RA]. URL: https://arxiv.org/abs/2212.14833.
- [Dor07] P. Dorey. "Exact S-matrices". Conformal Field Theories and Integrable Models: Lectures Held at the Eötvös Graduate Course, Budapest, Hungary, 13–18 August 1996. Springer, 2007, pp. 85–125.
- [Eti14] P. I. Etingof. Calogero-Moser systems and representation theory. Zurich Lectures in Advanced Mathematics. European Mathematical Society, 2014.
- [Gir15] S. Giraudo. "Combinatorial operads from monoids". Journal of Algebraic Combinatorics 41 (2015), pp. 493–538.
- [Gir18] S. Giraudo. *Nonsymmetric operads in combinatorics*. Vol. 193. Lecture Notes in Mathematics. Springer International Publishing, 2018. ISBN: 978-3-030-02074-3.
- [Gol66] S. W. Golomb. "Tiling with polyominoes". Journal of Combinatorial Theory 1.2 (1966), pp. 280–296.
- [KKS78] D. Kazhdan, B. Kostant, and S. Sternberg. "Hamiltonian group actions and dynamical systems of Calogero type". Communications on Pure and Applied Mathematics 31.4 (1978), pp. 481–507.
- [Kho10] T. Khovanova. L-Reptiles. https://blog.tanyakhovanova.com/2010/04/1-reptiles/. Accessed: 2023-12-19. 2010.
- [Kul76] P. P. Kulish. "Factorization of the classical and the quantum S matrix and conservation laws". *Theoretical and Mathematical Physics* 26 (1976), pp. 132–137.
- [LTV10] P. Lambrechts, V. Turchin, and I. Volić. "Associahedron, cyclohedron and permutohedron as compactifications of configuration spaces". Bulletin of the Belgian Mathematical Society-Simon Stevin 17.2 (2010), pp. 303–332.
- [LV12] J.-L. Loday and B. Vallette. *Algebraic Operads*. Vol. 346. Grundlehren der mathematischen Wissenschaften. Berlin, Heidelberg: Springer, 2012. ISBN: 978-3-642-30361-6.
- [MSS02] M. Markl, S. Shnider, and J. Stasheff. *Operads in algebra, topology and physics*. Vol. 96. Mathematical Surveys and Monographs. American Mathematical Society, 2002.
- [Mos75] J. Moser. "Three integrable Hamiltonian systems connected with isospectral deformations". Advances in Mathematics 16.2 (1975), pp. 197–220.
- [OP81] M. A. Olshanetsky and A. M. Perelomov. "Classical integrable finite-dimensional systems related to Lie algebras". *Physics Reports* 71.5 (1981), pp. 313–400.
- [Olv03] P. J. Olver. "Moving frames". Journal of Symbolic Computation 36.3-4 (2003), pp. 501–512.
- [Per90] A. M. Perelomov. Integrable systems of classical mechanics and Lie algebras. Vol. 1. Birkhäuser texts in physics. Birkhäuser, 1990.
- [Rei03] M. Reid. "Tiling with similar polyominoes". Journal of Recreational Mathematics 31.1 (2003), pp. 15–24.
- [SW03] P. Salvatore and N. Wahl. "Framed discs operads and Batalin–Vilkovisky algebras". *Quarterly Journal of Mathematics* 54.2 (2003), pp. 213–231.

REFERENCES

| [Tor 22] | A. Torrie | lli. $Lon T$ | <i>Lectures</i> | on | Sine- $Gordon$ | and | Thirring. | 2022. | arXiv: | 2211 | . 01186 |
|----------|-----------|--------------|-----------------|----|----------------|-----|-----------|-------|--------|------|---------|
| | [hep-th]. | | | | | | | | | | |

- [Vor99] A. A. Voronov. "The Swiss-cheese operad". *Contemporary Mathematics* 239 (1999), pp. 365–374.
- [Yau21] D. Yau. Infinity operads and monoidal categories with group equivariance. World Scientific, 2021.
- [Zam77] A. B. Zamolodchikov. "Exact two-particle S-matrix of quantum sine-Gordon solitons". Communications in Mathematical Physics 55 (1977), pp. 183–186.
- [ZZ79] A. B. Zamolodchikov and A. B. Zamolodchikov. "Factorized S-matrices in two dimensions as the exact solutions of certain relativistic quantum field theory models". Annals of physics 120.2 (1979), pp. 253–291.

CZECH ACADEMY OF SCIENCES, INSTITUTE OF MATHEMATICS, ŽITNÁ ,25, 115 67 PRAGUE 1, CZECH REPUBLIC