A Peculiar Proof of the Martingale Convergence Theorem

P.J. Fitzsimmons

Department of Mathematics
University of California, San Diego
9500 Gilman Drive
La Jolla, CA 92093-0112
pfitzsimmons@ucsd.edu

ABSTRACT

We prove the Martingale Convergence Theorem by using the work of L. Dubins and I. Monroe about embedding a given discrete-time martingale in the sample paths of a Brownian motion.

We provide a new (if ridiculous) proof of the Martingale Convergence Theorem.

Let $M = (M_n)_{n=0}^{\infty}$ be a discrete time martingale. With no real loss of generality we assume $M_0 = 0$. We consider first the case of L^1 -bounded martingales.

(1) Theorem. (J.L. Doob) Suppose that

(2)
$$K := \sup_{n} \mathbf{E}|M_n| < \infty.$$

Then

$$M_{\infty} := \lim_{n \to \infty} M_n$$

exists in **R** a.s., and M_{∞} is integrable.

Proof. Because $n \mapsto |M_n|$ is a submartingale, we deduce from (2) that

$$\mathbf{E}|M_T| \le K$$

for each a.s. finite stopping time T. We know from the work of Monroe [M] and Dubins [D] (see also [C; Thm. 10.3]) that the martingale M can be embedded in Brownian motion as follows. Let (B_t) be a 1-dimensional Brownian motion with $B_0 = 0$, defined on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbf{P})$ satisfying the usual conditions. Here (\mathcal{F}_t) denotes the filtration generated by B, suitably completed. Then there is a sequence $0 = T_0 \leq T_1 \leq T_2 \leq \cdots$ of standard (\mathcal{F}_t) -stopping times such that the process M' defined by $M'_n := B_{T_n}$, $n = 0, 1, 2, \ldots$ has the same law as the original martingale M. (In this context, we say that a stopping time T is "standard" provided the stopped process $t \mapsto B_{t \wedge T}$ is uniformly integrable. In particular, such a stopping time is finite a.s.) To show that $\lim_n M_n$ exists a.s., it therefore suffices to show that M' has the same property. To take advantage of (3) we use Tanaka's formula (see [RY; p. 222]):

(4)
$$|B_t| = \int_0^t \operatorname{sgn}(B_s) dB_s + L_t, \qquad t \ge 0,$$

where L is the local time of B at 0. Consequently, if T is an (\mathcal{F}_t) -stopping time, then

(5)
$$\mathbf{E}[L_{T \wedge m}] = \mathbf{E}|B_{T \wedge m}|, \quad \forall m > 0.$$

In particular,

(6)
$$\mathbf{E}[L_{T_n \wedge m}] = \mathbf{E}|B_{T_n \wedge m}|, \qquad n = 0, 1, 2, \dots,$$

and by standardness of T_n and monotone convergence, after a passage to the limit in (6) as $m \to \infty$ we obtain

(7)
$$\mathbf{E}[L_{T_n}] = \mathbf{E}|B_{T_n}| = \mathbf{E}|M_n| \le K, \qquad n = 0, 1, 2, \dots$$

Let $T_{\infty} := \lim_n T_n$. Because $t \mapsto L_t$ is monotone:

(8)
$$\mathbf{E}[L_{T_{\infty}}] \le K < \infty.$$

But $L_{\infty} = \infty$ a.s. because of the recurrence of one-dimensional Brownian motion. It follows from (8) that

$$\mathbf{P}[T_{\infty} < \infty] = 1.$$

Therefore the limit

$$M_{\infty}' := \lim_{n} M_{n}' = \lim_{n} B_{T_{n}} = B_{T_{\infty}}$$

exists in **R**, a.s. The integrability of M'_{∞} follows from Fatou's lemma. \square

The same sort of proof can be given for the convergence theorem for non-negative supermartingales. The proof is more elementary because Tanaka's formula is not needed. Indeed, let $X = (X_n)_{n=0}^{\infty}$ be a non-negative supermartingale with $X_0 = 1$ (without loss of generality). Let the Brownian motion (B_t) be as before. We know from Gushchin and Urusov [GU] that X can be embedded in the geometric Brownian motion $Z_t = \exp(B_t - t/2), t \geq 0$. That is, there is an increasing sequence $0 = T_0 \leq T_1 \leq \cdots \leq T_n \leq T_{n+1} \leq \infty$ of (\mathcal{F}_t) -stopping times such that $X'_n := Z_{T_n}, n = 0, 1, 2, \ldots$ has the same distribution as X. (N.B.: The stopping times T_n are now allowed to take the value ∞ . By the law of large numbers for Brownian motion, $\lim_{t\to\infty} Z_t = 0$ a.s., so we take $Z_{\infty} := 0$.) By monotonicity we have $T_{\infty} := \lim_n T_n \in [0, \infty]$, and then by path continuity $\lim_n X'_n = \lim_n Z_{T_n} = Z_{T_{\infty}}$ a.s. It follows that $\lim_n X_n$ exists in $[0, \infty[$ a.s. \square

It may be noted that because the geometric Brownian motion is a time change of Brownian motion on $[0, \infty[$ absorbed in state 0, the embedding result for non-negative supermartingales can be restated as follows: As before, let $B = (B_t)_{t\geq 0}$, be a Brownian motion, but now started at 1; let $\tau := \inf\{t > 0 : B_t = 0\}$. If $X = (X_n)_{n=0}^{\infty}$ is a non-negative supermartingale with $X_0 = 1$ then there is a sequence (T_n) of stopping times of B with $0 = T_0 \leq T_1 \leq T_2 \leq \cdots \leq T_n \leq T_{n+1} \leq \cdots \leq \tau$ such that $(B_{T_n})_{n=0}^{\infty}$ has the same law as X.

The embedding of Gushchin and Urusov uses randomized stopping times, but this can be avoided when embedding a supermartingale in absorbing Brownian motion by employing Falkner's refinement $[\mathbf{F}; \text{Thm. 1}]$ of Rost's theorem $[\mathbf{R}]$ for transient Markov processes.

References

- [C] Chacon, R.V.: Potental processes, Trans. Amer. Math. Soc. 226 (1977) 39–58.
- [D] Dubins, L.E.: On a theorem of Skorohod, Ann. Math. Statist. 39 (1968) 2094–2097.
- [F] Falkner, N.: Stopped distributions for Markov processes in duality, Z. Wahrsch. Verw. Gebiete 62 (1983) 43–51.
- [GU] Gushchin, A.A., Urusov, M.A.: Processes that can be embedded in a geometric Brownian motion, *Theory Probab. Appl.* **60** (2016) 246–262.
- [M] Monroe, I.: On embedding right continuous martingales in Brownian motion, *Ann. Probab.* **43** (1972) 1293–1311.
- [RY] Revuz, D., Yor, M.: Continuous martingales and Brownian motion, (3rd ed.). Springer-Verlag, Berlin, 1999.
 - [R] Rost, H.: The stopping distributions of a Markov process, *Invent. Math.* 14 (1971), 1–16.