# Upper Bounded Current Fluctuation in One-Dimensional Driven Diffusive Systems

Jiayin Gu<sup>®</sup><sup>∗1</sup> and Fan Zhang<sup>®</sup><sup>†2</sup>

<sup>1</sup>School of Physics and Technology, Nanjing Normal University, Nanjing 210023, China <sup>2</sup>School of Physics, Peking University, Beijing 100871, China

#### Abstract

The current fluctuation in one-dimensional driven diffusive systems is reported to have a upper bound that is set by its mean value together with the driving force. The resulting inequality arises from interactions between the diffusive particles, and the bound is approached in near-equilibrium systems or far-from-equilibrium systems with weak interactions between the diffusive particles. Three systems of increasing physical complexity are used to illustrate how the inequality is arrived at. The first system consists of only two reservoirs with particles randomly exchanging between them at constant rates. For this system, the bound is shown to be saturated. The second one is the symmetric simple exclusion process, where the inequality is rigorously proven. The third one is the transport of charged particles driven by an external electric field, and numerical studies for this system also support the existence of the inequality.

### 1 Introduction

Understanding the transport behavior of interacting particles has always been one of the central questions in nonequilibrium statistical physics [1–3]. In the last few decades, progress were made from extensive investigations of stochastic lattice systems, such as exclusion process [4, 5], zero-range process [5, 6], and contact process [7]. Moreover, general statements can also be made on systems in far-from-equilibrium regimes. On the one hand, the microscopic reversibility was revealed to constrain the current fluctuations in nonequilibrium steady-state systems, leading to the establishment of fluctuation theorems [8–16]. Close to equilibrium, these theorems reduce to the well-known fluctuation-dissipation relations such as the Green-Kubo formulae [17]. They also imply the Onsager reciprocal relations [18] as well as the generalized nonlinear ones up to arbitrary orders [16, 19–25]. On the other hand, great advances were initiated by the discovery of the so-called thermodynamic uncertainty relations (TURs) [26–28], which provide general lower bounds on the current fluctuation in terms of its mean value and the overall entropy production. These relations are interpreted as the thermodynamic cost of precision and find applications in deriving upper bounds on the efficiency of molecular motors [29], inferring entropy production [30], and even understanding sperm dynamics [31].

Despite the achievements so far, open questions about transport behavior in nonequilibrium systems can still be raised. For example, how do the interactions between the diffusive particles influence the current fluctuation? In this Letter, we try to answer this question, and accidentally, this leads us to discover a upper bound on the current fluctuation. For a one-dimensional nonequilibrium driven diffusive system whose the mean current J and its diffusivity D are defined in the following main text, we report a new inequality as follows:

$$2D \le J \coth(A/2),\tag{1}$$

where A stands for the affinity driving the system out of equilibrium [32–34]. If the system is isothermal at the inverse temperature  $\beta \equiv 1/(k_{\rm B}T)$ , the affinity is given in terms of the difference of chemical

<sup>\*</sup>gujiayin@njnu.edu.cn

<sup>&</sup>lt;sup>†</sup>van314159@pku.edu.cn

potentials between the left and right reservoirs,  $A = \beta(\mu_L - \mu_R)$ . The upper bound is saturated in the case where the system is close to equilibrium. This can be proven by invoking the fluctuation-dissipation relation J = DA that the inequality should reduce to it in the limit  $A \to 0$ . In addition, the original TUR in Ref. [26] can be developed to yield an equivalent form  $D \ge J/A$ , where the implied lower bound tends to coincide with the our reported upper bound as  $A \to 0$ . This also proves that the equality sign asymptotically holds when the system is near equilibrium. Moreover, when the system is driven far form equilibrium, the upper bound is also saturated in the case where the interactions between the diffusive particles are absent.

The suppressed current fluctuation can be understood intuitively by recalling the subdiffusion of tagged particles in a medium [35–37]. In normal diffusion, the particles undergo Brownian motion independently. In this case, the mean square displacement (MSD) of tracers in one dimension grows linearly with the time. However, in the exclusion process where the tracers can not bypass neighboring particles and consequently undergo subdiffusion, the growth of MSD scales with square root of time,  $\langle x_t^2 \rangle \sim \sqrt{t}$ , which is comparatively much slower. The case of interacting particles diffusing in driven systems follows the same physics. In the following, three systems of increasing physical complexity are employed to elucidate this essential physics step by step. The inequality (1) is expected to hold in general. It directly implies that the current fluctuation can be estimated from the mean current and the driving force or affinity. Moreover, if the three quantities are accurately measured, the inequality can also be used to infer whether the interactions between diffusive particles are present or not.

#### 2 Particle Exchanges between Two Reservoirs

We start from the simplest driven diffusive system where particles exchange stochastically between two reservoirs at constant rates  $W_+$  and  $W_-$ , as shown in the top panel of Figure 1. This system can be seen as a coarse-grained model at the highest level of description for the particle transport in a general nonequilibrium system such that the intermediate subsystem is integrated out. The physics of particle exchanges is simplified as a superposition of two independent Poisson processes. Let Z denote the cumulative particle transfers from the left to the right reservoir during the time interval [0, t] and  $\mathcal{P}(Z, t)$ be its probability distribution, then the cumulant generating function in terms of the counting parameter  $\lambda$  can be defined as

$$Q(\lambda) = \lim_{t \to \infty} -\frac{1}{t} \ln \sum_{Z = -\infty}^{+\infty} e^{-\lambda Z} \mathcal{P}(Z, t),$$
(2)

whose analytical expression for this system can be explicitly given by (see Appendix A)

$$Q(\lambda) = W_+ \left(1 - e^{-\lambda}\right) + W_- \left(1 - e^{+\lambda}\right).$$
(3)

It can be shown that this cumulant generating function satisfies the Gallavotti-Cohen symmetry  $Q(\lambda) = Q(A - \lambda)$  [10, 11, 13], where

$$A \equiv \ln \frac{W_+}{W_-} \tag{4}$$

is defined as the affinity. This symmetry is the direct consequence of the fluctuation theorem. The mean current and its diffusivity are calculated as follows:

$$J \equiv \lim_{t \to \infty} \frac{\langle Z \rangle}{t} = \left. \frac{\partial Q}{\partial \lambda} \right|_{\lambda=0} = W_{+} - W_{-},\tag{5}$$

$$D \equiv \lim_{t \to \infty} \frac{\langle (Z - Jt)^2 \rangle}{2t} = -\frac{1}{2} \frac{\partial^2 Q}{\partial \lambda^2} \Big|_{\lambda=0} = \frac{W_+ + W_-}{2}.$$
 (6)

It can be easily seen from Eqs. (4)-(6) that A, J and D are not mutually independent such that an equality

$$2D = J \coth(A/2) \tag{7}$$

exists between them. The immediate implication is that if this simple system for particle exchanges can truly be regarded as an accurate coarse-grained model for general nonequilibrium systems then this



Figure 1: Schematic representations of the three one-dimensional driven diffusive systems. Top panel: The simplest system consisting of only two reservoirs with particles stochastically exchanging between them. Middle panel: The symmetric simple exclusion process. The black (white) circles denote occupied (vacant) sites. Bottom panel: Charged particles transporting in a conductive channel.

equality can be employed to test fluctuation theorem indirectly. Since the fluctuation theorem is the property of large deviation [13, 38–40], it is generally very difficult to access rare events that are essential to directly test the fluctuation theorem. The equality (7) provides an alternative way to bypass this conundrum as it is relatively easy to access the mean current J and its diffusivity D. Actually, this equality has already been used to test the fluctuation theorem for the charge transport in diodes [41] and transistors [20]. It will be shown in the following that, for general nonequilibrium systems where the interactions between diffusive particles are present, this equality is violated. Nevertheless, this equality serves as the starting point to arrive at the inequality (1).

## 3 Symmetric Simple Exclusion Process

We now consider a paradigmatic model in nonequilibrium statistical physics, the symmetric simple exclusion process (SSEP) [4, 5, 42–45]. This process describes the hopping of hard-core particles in a one-dimensional chain, only driven at the boundries by the inflow and outflow of particles. As shown in the middle panel of Figure 1, the particles stochastically hop to a vacant neighboring site at a rate of 1 in the bulk and enter (exit) at the left and right boundries at rates  $\alpha$  ( $\gamma$ ) and  $\delta$  ( $\beta$ ). This system was extensively investigated during the last few decades, and therefore a wealth of analytical knowledge has accumulated [42, 44, 46–48]. Suppose this system consists of L sites that are indexed 1, 2,  $\cdots L$  from the left to right, then the average occupation number at *i*-th site is [42–44]

$$\langle n_i \rangle = \frac{\rho_{\rm L}(L+b-i) + \rho_{\rm R}(i-1+a)}{L+a+b-1},$$
(8)

where

$$\rho_{\rm L} = \frac{\alpha}{\alpha + \gamma}, \qquad \qquad a = \frac{1}{\alpha + \gamma},$$
(9)

$$\rho_{\rm R} = \frac{\delta}{\beta + \delta}, \qquad \qquad b = \frac{1}{\beta + \delta}. \tag{10}$$

Here,  $\rho_{\rm L}$  and  $\rho_{\rm R}$  are identified as the densities of the left and right reservoirs. If these two densities are unequal, the mean current is induced, given by

$$J = \langle n_i - n_{i+1} \rangle = \frac{\rho_{\rm L} - \rho_{\rm R}}{\tilde{L}},\tag{11}$$

where  $\tilde{L} \equiv L + a + b - 1$ . Moreover, the diffusivity of the current can also be given by [42–44]

$$D = \frac{1}{2\tilde{L}} \left(\rho_{\rm L} + \rho_{\rm R} - 2\rho_{\rm L}\rho_{\rm R}\right) + \frac{a(a-1)(2a-1) + b(b-1)(2b-1) - \tilde{L}(\tilde{L}-1)(2\tilde{L}-1)}{6\tilde{L}^3(\tilde{L}-1)} \left(\rho_{\rm L} - \rho_{\rm R}\right)^2, \quad (12)$$

The affinity can also be evaluated [42-44],

$$A = \ln \frac{\rho_{\rm L} \bar{\rho}_{\rm R}}{\bar{\rho}_{\rm L} \rho_{\rm R}} = \ln \frac{\alpha \beta}{\gamma \delta},\tag{13}$$

where  $\bar{\rho}_{\rm L} \equiv 1 - \rho_{\rm L}$  and  $\bar{\rho}_{\rm R} \equiv 1 - \rho_{\rm R}$  are densities quantifying vacancy. This affinity can be identified through Schnakenberg's graph analysis of the master equation for the jump process [49]. It is given as the ratio of product of transition rates along the cyclic path over the product of transition rates along the reversed cyclic path<sup>1</sup>. The implication of affinities identified from cycles for the graph leads to the establishment of the fluctuation theorem in the early form [50].

At this point, we are ready to calculate 2D/J with the expressions (11)-(12), reading

$$\frac{2D}{J} = \frac{\rho_{\rm L} + \rho_{\rm R} - 2\rho_{\rm L}\rho_{\rm R}}{\rho_{\rm L} - \rho_{\rm R}} + \frac{a(a-1)(2a-1) + b(b-1)(2b-1) - \tilde{L}(\tilde{L}-1)(2\tilde{L}-1)}{3\tilde{L}^2(\tilde{L}-1)}(\rho_{\rm L} - \rho_{\rm R}).$$
(14)

Remarkably, the first term on the right-hand side can be further simplifed as

$$\frac{\rho_{\rm L} + \rho_{\rm R} - 2\rho_{\rm L}\rho_{\rm R}}{\rho_{\rm L} - \rho_{\rm R}} = \frac{\rho_{\rm L}\bar{\rho}_{\rm R} + \rho_{\rm R}\bar{\rho}_{\rm L}}{\rho_{\rm L}\bar{\rho}_{\rm R} - \rho_{\rm R}\bar{\rho}_{\rm L}} = \frac{\alpha\beta + \delta\gamma}{\alpha\beta - \delta\gamma} = \frac{e^A + 1}{e^A - 1} = \frac{e^{A/2} + e^{-A/2}}{e^{A/2} - e^{-A/2}} = \coth(A/2), \quad (15)$$

giving what we expect. To prove that the inequality (1) holds in this system, we only need to show that the second term is non-positive. Taking L = 1, we have

second term = 
$$-\frac{2ab}{(a+b)^2} \left(\rho_{\rm L} - \rho_{\rm R}\right),$$
 (16)

which is non-positive (see Appendix B). Here,  $\rho_{\rm L} \ge \rho_{\rm R}$  is implicitly assumed. When  $L \ge 2$ , the denominator is always positive and the numerator decreases as L increases. The maximum of the numerator at L = 2 is

$$-6\left(a^{2}b + a^{2} + ab + ab^{2} + b^{2}\right)\left(\rho_{\rm L} - \rho_{\rm R}\right),\tag{17}$$

indicating that the second term is also non-positive, as expected (see Appendix B). It is easy to notice that the density difference,  $\rho_{\rm L} - \rho_{\rm R}$ , determines whether the equality sign in Eq. (1) holds or not. When the two densities are equal,  $\rho_{\rm L} = \rho_{\rm R}$ , the equality sign holds. This corresponds to the case where the system is in equilibrium. The equality sign can also asymptotically hold when  $1 \gg \rho_{\rm L} > \rho_{\rm R} > 0$ , i.e., the two densities are both very low. This translates to the situation where the exclusion between diffusive particles is negligible. Graphical presentations are shown in the left panel of Figure 2, where the defined quantity  $\chi \equiv 2DA/J$  is both bounded from below by 2 (TUR) and bounded form upper by  $A \coth(A/2)$ . It is clearly that the upper bound is saturated in the low-density limit. What's unexpected, we find that the upper bound seems to be saturated also in the high-density limit.

## 4 Transport of Charged Particles

Next, we consider the transport of charged particles in a conductive channel at the mesoscopic level. As shown in the bottom panel of Figure 1, the channel can be thought of as a three-dimensional rod extending in x-axis from 0 to l. In the transverse y- and z- directions, the section area is denoted by  $\Sigma$ . Two kinds

 $<sup>^{1}</sup>$ The cyclic path can be constructed here as a particle entering the empty chain from the left, then jumping to the right and eventually exiting the chain.



Figure 2: (Color online) Graphical representations of the bounded current fluctuation. Left panel: The behavior of  $\chi \equiv 2DA/J$  as a function of A with different particle densities (controlled by a parameter  $x \equiv \gamma/\alpha$  with small value for high density and large value for low density) for SSEP. The parameter values are  $\gamma = \beta = 1$  and L = 10. Moreover, the condition  $y = e^A x$  with  $y \equiv \beta/\delta$  is also imposed. The black solid line is the affinity-dependent upper bound  $A \coth(A/2)$ , and the gray solid line is the lower bound 2 from the TUR. The dash lines are depicted from Eqs. (11)-(13). Right panel: The comparison between numerical affinities (24) and theoretical affinities (23) for the transport of charged particles. It is set that  $\bar{N}_{\rm L} = \bar{N}_{\rm R} = N_- = n_-\Omega$ , and takes different values in three cases, as shown in the legend. The asterisks are numerical points with dashed lines joining them. The dot dash line indicates the equality between both kinds of affinities. The parameter values used in simulation are  $\beta = e = 1.0$ ,  $D = \epsilon = 0.01$ ,  $\Omega = 10000$ ,  $\Delta x = 0.1$ , L = 10. The numerical affinities are computed with J and D evaluated over time interval [0, 5000] with 20000 data.

of charged particles are supposed to be distributed in this channel: mobile positive-charged particles with the density n(x), and anchored negative-charged particles with uniform density  $n_-(x) = n_-$ . The charge density is therefore given by  $\rho(x) = e[n(x) - n_-]$ , where e = |e| represents the elementary electric charge. Both terminals are in contact with reservoirs fixing the densities of mobile positive-charged particles with  $n_{\rm L}$ ,  $n_{\rm R}$ , and electric potentials with  $\Phi_{\rm L}$ ,  $\Phi_{\rm R}$ . The electric potential  $\Phi(x)$  across the channel fluctuates and is determined through Poisson equation by the fluctuating charge distributions as well as the boundary conditions [51].

The basic idea for simulating the system is to first discretize the system in space and then establish master equation to describe the stochastic evolution of the system state [52]. For this purpose, the channel is discretized into cells, each with width  $\Delta x$  and volume  $\Omega = \Sigma \Delta x$ . There are a total of  $L = l/\Delta x$  of such cells. The reservoirs are modeled as two cells containing fixed number of positive-charged particles  $\bar{N}_{\rm L} = n_{\rm L}\Omega$ ,  $\bar{N}_{\rm R} = n_{\rm R}\Omega$ . The system state is specified by the numbers  $\{N_1, N_2, \dots, N_L\}$  of the mobile charged particles in the intermediate cells. It evolves in time according to the network

$$\bar{N}_{\rm L} \xrightarrow[W_0^{(+)}]{W_0^{(-)}} N_1 \xrightarrow[W_1^{(+)}]{W_1^{(-)}} N_2 \xrightarrow[W_2^{(+)}]{W_2^{(-)}} \cdots \xrightarrow[W_{i-1}^{(+)}]{W_{i-1}^{(-)}} N_i \xrightarrow[W_{i-1}^{(+)}]{W_{i-1}^{(-)}} \cdots \xrightarrow[W_{L-2}^{(+)}]{W_{L-2}^{(+)}} N_{L-1} \xrightarrow[W_{L-1}^{(+)}]{W_{L-1}^{(+)}} N_L \xrightarrow[W_{L-1}^{(+)}]{W_{L-1}^{(+)}} \bar{N}_{\rm R},$$
(18)

with the transition rates given by

$$W_i^{(+)} = \frac{D}{\Delta x^2} \psi(\Delta U_{i,i+1}) N_i, \qquad (19)$$

$$W_i^{(-)} = \frac{D}{\Delta x^2} \psi(\Delta U_{i+1,i}) N_{i+1}.$$
 (20)

Here, D denotes the diffusion coefficient of mobile particles, not to be confused with diffusivity of the current.  $\Delta U_{i,i+1}$  is the intrinsic energy change of the system associated with one particle jumping from the *i*-th to (i + 1)-th cell and is given by [52]

$$\Delta U_{i,i+1} = e(\Phi_{i+1} - \Phi_i) + \frac{e^2 L \Delta x^2}{2(L+1)\epsilon\Omega},$$
(21)

where  $\epsilon$  is the permittivity. The function  $\psi(\Delta U)$  is defined as

$$\psi(\Delta U) = \frac{\beta \Delta U}{\exp(\beta \Delta U) - 1},\tag{22}$$

which guarantees the detailed balance in equilibrium,  $\psi(\Delta U) = \psi(-\Delta U) \exp(-\beta \Delta U)$ . The master equation can be readily written down from the network (18) and random trajectories can be generated with the Gillespie algorithm [53]. A much faster algorithm based on Langevin-type stochastic process is actually implemented in simulation (see Appendix C).

In numerical simulation, we impose the boundary condition  $\bar{N}_{\rm L} = \bar{N}_{\rm R}$ . So, the affinity in theory is

$$A^{(t)} = \beta e (\Phi_{\rm L} - \Phi_{\rm R}). \tag{23}$$

The integrated current Z crossing a section over a large time interval [0, t] is counted. The mean current J and its diffusivity D can be extracted from the counting statistics according to their definition. Inspired by the first particle-exchange system, we numerically estimate the the affinity from these two cumulants

$$A^{(n)} = \ln \frac{2D + J}{2D - J}.$$
(24)

The results are displayed in the right panel of Figure 2, where we compared  $A^{(n)}$  and  $A^{(t)}$  in three cases of different particle densities. We notice that  $A^{(n)}$  does not always agree with  $A^{(t)}$ , the former is generally greater than or equal to the latter,  $A^{(n)} \ge A^{(t)}$ , leading to the inequality (1). When the system is near equilibrium, they are approximately equal to each other. However, when system is driven far from equilibrium, the approximate equality is only found in the case of very low particle densities. It can be argued that the charged particles with low densities play a negligible role in determining the potential  $\Phi(x)$  by Poisson equation, the fluctuating electric field in this case can be approximated as a static background field. It can be also exactly proven that, in the case of low-density limit, the cumulant generating function for the current has the same form as Eq. (3) with the equivalent global transition rates given by (see Appendix C)

$$W_{+} = \frac{DN_{\rm L}}{\Delta x^2(L+1)} \frac{\beta e(\Phi_{\rm R} - \Phi_{\rm L})}{\exp\left[\beta e(\Phi_{\rm R} - \Phi_{\rm L})\right] - 1},\tag{25}$$

$$W_{-} = \frac{DN_{\rm R}}{\Delta x^2(L+1)} \frac{\beta e(\Phi_{\rm L} - \Phi_{\rm R})}{\exp\left[\beta e(\Phi_{\rm L} - \Phi_{\rm R})\right] - 1}.$$
(26)

This precisely shows that the equality sign in the Eq. (1) holds in the low-density limit. The physical interpretation is that in this case the electrostatic repulsive interactions between charged particles play no role, and the transport of charged particles is reduced to normal driven Brownian motion. On the contrary, when the electrostatic repulsive interactions are present, the current fluctuation is suppressed. In addition, the inset of the right panel of Figure 2 shows that the mean current is not affected by the repulsive interactions, following the Ohm law.

#### 5 Conclusion

In this work, we have reported an inequality that sets an upper bound for the current fluctuation in one-dimensional driven diffusive systems. Three concrete systems are used to demonstrate how this inequality is established. The first particle-exchange system is very simple and illuminating; it enables the quantification of the current fluctuation in terms of the mean current and the affinity that are relatively easy to measure. The second and third systems show that the interactions between diffusive particles generally suppress the current fluctuation. Although the inequality is obtained through a case-by-case study, it captures the essential physics of driven transport of interacting particles. The current fluctuation in more complex systems, such as quantum ballistic transport system and the diffusion-reaction system containing more than one particle species, can also be shown to be limited by our bound. We hope to report this in future publications. In addition, an upper bound on the fluctuations of flux observables of trajectories has been recently derived in Ref. [54]. However, this bound is proved rather mathematically, and relies on an input quantity which is not easy to access. By contrast, our upper bound is more specific and physical. It would interesting to investigate the connection between this bound and our bound. To conclude, the newly found inequality represents a important result in nonequilibrium statistical physics.

## A Particle Exchanges between Two Reservoirs

Let Z denote the cumulative particle transfers from the left to the right reservoir during the time interval [0, t], then the probability distribution of Z at time t obeys the master equation

$$\frac{\mathrm{d}\mathcal{P}(Z,t)}{\mathrm{d}t} = \left[W_{+}\left(\mathrm{e}^{-\partial_{Z}}-1\right) + W_{-}\left(\mathrm{e}^{+\partial_{Z}}-1\right)\right]\mathcal{P}(Z,t)$$
(27)

We now define the moment generating function of signed cumulated flux,

$$G(s,t) \equiv \sum_{Z=-\infty}^{+\infty} s^{Z} \mathcal{P}(Z,t), \qquad (28)$$

whose time derivative is as follows,

$$\partial_t G(s,t) = W_+ \sum_{Z=-\infty}^{+\infty} s^Z \mathcal{P}(Z-1,t) + W_- \sum_{Z=-\infty}^{+\infty} s^Z \mathcal{P}(Z+1,t) - (W_+ + W_-) \sum_{Z=-\infty}^{+\infty} s^Z \mathcal{P}(Z,t) \\ = \left( W_+ s + \frac{W_-}{s} - W_+ - W_- \right) G(s,t).$$
(29)

For convenience, the initial probability distribution is taken to be  $\mathcal{P}(Z,0) = \delta_{Z,0}$ , so

$$G(s,0) = \sum_{Z=-\infty}^{\infty} s^Z \delta_{Z,0} = 1,$$
(30)

and

$$G(s,t) = \exp\left[\left(W_{+}s + \frac{W_{-}}{s} - W_{+} - W_{-}\right)t\right].$$
(31)

The cumulant generating function can thus be obtained,

$$Q(\lambda) \equiv \lim_{t \to \infty} -\frac{1}{t} \ln G\left(e^{-\lambda}, t\right) = W_+ \left(1 - e^{-\lambda}\right) + W_- \left(1 - e^{+\lambda}\right).$$
(32)

## **B** Symmetric Simple Exclusion Process

From the analytical expression of J and D, we can readily evaluate 2D/J, reading

$$\frac{2D}{J} = \frac{\rho_{\rm L} + \rho_{\rm R} - 2\rho_{\rm L}\rho_{\rm R}}{\rho_{\rm L} - \rho_{\rm R}} + \frac{a(a-1)(2a-1) + b(b-1)(2b-1) - L(L-1)(2L-1)}{3\tilde{L}^{2}(\tilde{L}-1)}(\rho_{\rm L}-\rho_{\rm R}) 
= \frac{\rho_{\rm L}\bar{\rho}_{\rm R} + \rho_{\rm R}\bar{\rho}_{\rm L}}{\rho_{\rm L}\bar{\rho}_{\rm R} - \rho_{\rm R}\bar{\rho}_{\rm L}} + \frac{a(a-1)(2a-1) + b(b-1)(2b-1) - \tilde{L}(\tilde{L}-1)(2\tilde{L}-1)}{3\tilde{L}^{2}(\tilde{L}-1)}(\rho_{\rm L}-\rho_{\rm R}) 
= \coth(A/2) + \frac{a(a-1)(2a-1) + b(b-1)(2b-1) - \tilde{L}(\tilde{L}-1)(2\tilde{L}-1)}{3\tilde{L}^{2}(\tilde{L}-1)}(\rho_{\rm L}-\rho_{\rm R}).$$
(33)

The second term can be proved to be non-positive. Here, we implicitly assume that  $\rho_{\rm L} - \rho_{\rm R}$ . So  $\rho_{\rm L} - \rho_{\rm R}$  is irrelevant in determing the sign of the second term. When L = 1, the numerator of the second term  $(\rho_{\rm L} - \rho_{\rm R} \text{ ignored})$  is

$$\begin{aligned} \text{numerator} &= a(a-1)(2a-1) + b(b-1)(2b-1) - \tilde{L}(\tilde{L}-1)(2\tilde{L}-1) \\ &= a(a-1)(2a-1) + b(b-1)(2b-1) - (a+b)(a+b-1)(2a+2b-1) \\ &= a(2a^2 - 3a+1) + b(2b^2 - 3b+1) - (a+b)(a+b-1)(2a+2b-1) \\ &= 2a^3 + 2b^3 - 3(a^2 + b^2) + a + b - (a+b)(a+b-1)(2a+2b-1) \\ &= 2(a+b)(a^2 - ab+b^2) - 3(a+b)^2 + 6ab + a + b - (a+b)(a+b-1)(2a+2b-1) \\ &= (a+b)(2a^2 - 2ab + 2b^2 - 3a - 3b+1) + 6ab - (a+b)[2(a+b)^2 - 3(a+b)+1] \\ &= (a+b)(2a^2 - 2ab + 2b^2 - 2(a+b)^2] + 6ab \\ &= (a+b)(-6ab) + 6ab \\ &= -6ab(a+b-1). \end{aligned}$$

The denominator is

denominator = 
$$3(a+b)^2(a+b-1)$$
. (34)

So, the second term is

$$2\mathrm{nd} \ \mathrm{term} = -\frac{2ab}{(a+b)^2},\tag{35}$$

which is negative. When  $L \ge 2$ , the denominator is positive, and the numerator takes the maximum value at L = 2. So

maximum of numerator 
$$=a(a-1)(2a-1) + b(b-1)(2b-1) - \tilde{L}(\tilde{L}-1)(2\tilde{L}-1)$$
$$=a(a-1)(2a-1) + b(b-1)(2b-1) - (a+b+1)(a+b)(2a+2b+1)$$
$$=(a+b)(2a^2 - 2ab + 2b^2 - 3a - 3b+1) + 6ab - (a+b+1)(a+b)(2a+2b+1)$$
$$=(a+b)(2a^2 - 2ab + 2b^2 - 3a - 3b+1) + 6ab - (a+b)[2(a+b)^2 + 3(a+b) + 1]$$
$$=(a+b)(-6ab - 6a - 6b) + 6ab$$
$$= -6(a^2b + a^2 + ab + ab^2 + ab + b^2 - ab)$$
$$= -6(a^2b + a^2 + ab + ab^2 + b^2),$$
(36)

which is negative.

## C Transport of Charged Particles

## C.1 Langevin-Type Stochastic Process

The probability  $\mathcal{P}(\mathbf{N}, t)$  that the cells contain the particle numbers  $\mathbf{N}$  for time t obeys the master equation

$$\frac{\mathrm{d}\mathcal{P}}{\mathrm{d}t} = \sum_{i=0}^{L} \left[ \left( \mathrm{e}^{+\partial_{N_i}} \mathrm{e}^{-\partial_{N_{i+1}}} - 1 \right) W_i^{(+)} + \left( \mathrm{e}^{-\partial_{N_i}} \mathrm{e}^{+\partial_{N_{i+1}}} - 1 \right) W_i^{(-)} \right] \mathcal{P}.$$
(37)

When  $N_i \gg 1$ , the operators  $\exp(\pm \partial_{N_i})$  can be expanded up to second order in the partial derivatives  $\partial_{N_i}$  in Eq. (37). In this way, we get the Fokker-Planck equation

$$\partial_{t}\mathscr{P} = \sum_{i=0}^{L} \left\{ -\partial_{N_{i}} \left[ \left( W_{i-1}^{(+)} - W_{i-1}^{(-)} - W_{i}^{(+)} + W_{i}^{(-)} \right) \mathscr{P} \right] + \partial_{N_{i}} \partial_{N_{i+1}} \left[ - \left( W_{i}^{(+)} + W_{i}^{(-)} \right) \mathscr{P} \right] \right\} + \partial_{N_{i}} \partial_{N_{i+1}} \left[ - \left( W_{i}^{(+)} + W_{i}^{(-)} \right) \mathscr{P} \right] \right\}.$$
(38)

for the time evolution of the probability density  $\mathscr{P}$ . This shows that the variables  $N_i$  obeys the following stochastic differential equations of Langevin type,

$$\frac{\mathrm{d}N_i}{\mathrm{d}t} = F_{i-1} - F_i,\tag{39}$$

expressed in terms of the fluxes

$$F_i = W_i^{(+)} - W_i^{(-)} + \sqrt{W_i^{(+)} + W_i^{(-)}} \xi_i(t),$$
(40)

and the Gaussian white noises  $\xi_i(t)$  satisfying the properties:

$$\langle \xi_i(t) \rangle = 0, \tag{41}$$

$$\langle \xi_i(t)\xi_j(t')\rangle = \delta_{ij}\delta(t-t'). \tag{42}$$

#### C.2 The Case of Low-Density Limit

In the low-density limit, the fluctuating electric field is approximated as a static background field. In this case, the energy difference associated with particle transitions between discretized cells are given by

$$\Delta U_{i,i+1} = e(\Phi_{i+1} - \Phi_i) = -\frac{e(\Phi_{\rm L} - \Phi_{\rm R})}{L+1},\tag{43}$$

$$\Delta U_{i+1,i} = e(\Phi_i - \Phi_{i+1}) = + \frac{e(\Phi_{\rm L} - \Phi_{\rm R})}{L+1},\tag{44}$$

which can be uniformly expressed as

$$\Delta U^{(\pm)} = \mp \frac{e(\Phi_{\rm L} - \Phi_{\rm R})}{L+1}.$$
(45)

Correspondingly,  $\psi(\Delta U)$  is denoted as

$$\psi^{(\pm)} = \psi(\Delta U^{(\pm)}) = \frac{\beta \Delta U^{(\pm)}}{\exp\left(\beta \Delta U^{(\pm)}\right) - 1}.$$
(46)

The local transition rates are thus expressed as

$$W_i^{(+)} = \frac{D}{\Delta x^2} \psi^{(+)} N_i = k_+ N_i, \tag{47}$$

$$W_i^{(-)} = \frac{D}{\Delta x^2} \psi^{(-)} N_{i+1} = k_- N_{i+1}, \tag{48}$$

with the rate constants  $k_+$  and  $k_-$  defined as obvious. The rates corresponding to the transitions from the reservoirs to the system are

$$W_0^{(+)} = k_+ \bar{N}_{\rm L}, \qquad W_L^{(-)} = k_- \bar{N}_{\rm R}.$$
 (49)

Clearly, all these transition rates are determined locally, and as such the transport system in the lowdensity limit is linear. We consider the time evolution of the probability

$$\mathcal{P}(Z, N_1, \cdots, N_L, t) \tag{50}$$

that the cells contain given particle numbers and that the signed cumulated number Z of particles is transferred from the I-th to the (I + 1)-th cells during time interval [0, t]. This probability is ruled by the following master equation,

$$\frac{d\mathcal{P}}{dt} = k_{+}\bar{N}_{L} \left(e^{-\partial_{N_{1}}} - 1\right) \mathcal{P} + k_{-} \left(e^{+\partial_{N_{1}}} - 1\right) N_{1}\mathcal{P} + k_{+} \left(e^{+\partial_{N_{1}}} e^{-\partial_{N_{2}}} - 1\right) N_{1}\mathcal{P} \\
+ \sum_{i=2}^{I-1} \left[k_{-} \left(e^{+\partial_{N_{i}}} e^{-\partial_{N_{i-1}}} - 1\right) N_{i}\mathcal{P} + k_{+} \left(e^{+\partial_{N_{i}}} e^{-\partial_{N_{i+1}}} - 1\right) N_{i}\mathcal{P}\right] \\
+ k_{-} \left(e^{+\partial_{N_{I}}} e^{-\partial_{N_{I-1}}} - 1\right) N_{I}\mathcal{P} + k_{+} \left(e^{+\partial_{N_{I}}} e^{-\partial_{N_{I+1}}} e^{-\partial_{2}} - 1\right) N_{I}\mathcal{P} \\
+ k_{-} \left(e^{+\partial_{N_{I+1}}} e^{-\partial_{N_{I}}} e^{+\partial_{2}} - 1\right) N_{I+1}\mathcal{P} + k_{+} \left(e^{+\partial_{N_{I}}} e^{-\partial_{N_{I+2}}} - 1\right) N_{I+1}\mathcal{P} \\
+ \sum_{i=I+2}^{L-1} \left[k_{-} \left(e^{+\partial_{N_{i}}} e^{-\partial_{N_{i-1}}} - 1\right) N_{i}\mathcal{P} + k_{+} \left(e^{+\partial_{N_{i}}} e^{-\partial_{N_{i+1}}} - 1\right) N_{i}\mathcal{P}\right] \\
+ k_{-}\bar{N}_{R} \left(e^{-\partial_{N_{L}}} - 1\right) \mathcal{P} + k_{-} \left(e^{+\partial_{N_{L}}} e^{-\partial_{N_{L-1}}} - 1\right) N_{L}\mathcal{P} + k_{+} \left(e^{+\partial_{N_{L}}} - 1\right) N_{L}\mathcal{P}.$$
(51)

Using the method of Ref. [55], this master equation is solved by introducing the moment generating function

$$G(\eta, s_1, \cdots, s_L, t) = \sum_{Z, N_1, \cdots, N_L} \eta^Z \prod_i s_i^{N_i} \mathcal{P}(Z, N_1, \cdots, N_L, t).$$
(52)

where

$$\eta = e^{-\lambda},\tag{53}$$

and  $\lambda$  is the counting parameter for the particle transfers Z. This moment generating function is ruled by the following first-order partial differential equation,

$$\begin{aligned} \partial_t G + [k_-(s_1 - 1) + k_+(s_1 - s_2)] \,\partial_{s_1} G \\ &+ \sum_{i=2}^{I-1} [k_-(s_i - s_{i-1}) + k_+(s_i - s_{i+1})] \,\partial_{s_i} G \\ &+ [k_-(s_I - s_{I-1}) + k_+(s_I - \eta s_{I+1})] \,\partial_{s_I} G \\ &+ [k_-(s_{I+1} - \eta^{-1} s_I) + k_+(s_{I+1} - s_{I+2})] \,\partial_{s_{I+1}} G \\ &+ \sum_{i=I+1}^{L-1} [k_-(s_i - s_{i-1}) + k_+(s_i - s_{i+1})] \,\partial_{s_i} G \\ &+ [k_-(s_L - s_{L-1}) + k_+(s_L - 1)] \,\partial_{s_L} G \\ &= [k_+ \bar{N}_{\rm L}(s_1 - 1) + k_- \bar{N}_{\rm R}(s_L - 1)] \,G, \end{aligned}$$
(54)

which, in vectoral notations, can be written in the following form,

$$\partial_t G + (\mathbf{L} \cdot \mathbf{s} + \mathbf{f}) \cdot \partial_\mathbf{s} G = (\mathbf{g} \cdot \mathbf{s} + h)G \tag{55}$$

where

$$\mathbf{L} \equiv \begin{pmatrix} k_{-} + k_{+} & -k_{+} & & & \\ -k_{-} & k_{-} + k_{+} & -k_{+} & & & \\ & \ddots & \ddots & \ddots & & \\ & & -k_{-} & k_{-} + k_{+} & -\eta k_{+} & & \\ & & -\eta^{-1} k_{-} & k_{-} + k_{+} & -k_{+} & \\ & & & \ddots & \ddots & \ddots & \\ & & & & -k_{-} & k_{-} + k_{+} & -k_{+} \\ & & & & -k_{-} & k_{-} + k_{+} & -k_{+} \\ & & & & -k_{-} & k_{-} + k_{+} & -k_{+} \\ \end{pmatrix},$$
(56)

$$\mathbf{s} \equiv \begin{pmatrix} s_1 \\ s_2 \\ \vdots \\ s_{L-1} \\ s_L \end{pmatrix}, \qquad \mathbf{f} \equiv - \begin{pmatrix} k_- \\ 0 \\ \vdots \\ 0 \\ k_+ \end{pmatrix}, \qquad \mathbf{g} \equiv \begin{pmatrix} k_+ \bar{N}_L \\ 0 \\ \vdots \\ 0 \\ k_- \bar{N}_R \end{pmatrix}, \tag{57}$$

and

$$h \equiv -k_+ \bar{N}_{\rm L} - k_- \bar{N}_{\rm R}.\tag{58}$$

The parameter  $\eta$  in matrix **L** appears in the *I*-th and the (I + 1)-th rows. From the matrix **L**, we can define  $\mathbf{L}_0$  by setting  $\eta = 1$  and thus  $\lambda = 0$ . So, we have the relations

$$\mathbf{f} = -\mathbf{L}_0 \cdot \mathbf{1},\tag{59}$$

$$h = -\mathbf{g} \cdot \mathbf{1},\tag{60}$$

where  $\mathbf{1}$  denotes the vector with all entries equal to one. Besides, the stationary values of particle numbers are given by

$$\boldsymbol{\Gamma}_0 = \boldsymbol{\mathsf{L}}_0^{-1\mathrm{T}} \cdot \boldsymbol{\mathrm{g}}.$$
 (61)

The first-order partial differential equation (55) can be solved by the method of characteristics. The equations for the characteristics are given by

$$\frac{\mathrm{d}\mathbf{s}}{\mathrm{d}t} = \mathbf{L} \cdot \mathbf{s} + \mathbf{f},\tag{62}$$

$$\frac{\mathrm{d}G}{\mathrm{d}t} = (\mathbf{g} \cdot \mathbf{s} + h)G. \tag{63}$$

The solution of Eq. (62) gives the characteristics

$$\mathbf{s} = e^{\mathbf{L}t} \cdot \left[ \mathbf{s}_0 + \mathbf{L}^{-1} \cdot \left( \mathbf{I} - e^{-\mathbf{L}t} \right) \cdot \mathbf{f} \right].$$
(64)

Replacing in Eq. (63), we obtain after integration that

$$G = G_0 \exp\left[\mathbf{g} \cdot \mathbf{L}^{-1} \cdot \left(\mathbf{I} - e^{-\mathbf{L}t}\right) \cdot \left(\mathbf{s} + \mathbf{L}^{-1} \cdot \mathbf{f}\right) + \left(h - \mathbf{g} \cdot \mathbf{L}^{-1} \cdot \mathbf{f}\right) t\right],\tag{65}$$

where I denotes the identity matrix. The initial condition being the Poisson distribution describing the steady state and the counter reset to zero Z = 0, we have that

$$G_0(\eta, \mathbf{s}_0) = e^{\Gamma_0 \cdot (\mathbf{s}_0 - \mathbf{1})}.$$
(66)

The solution of Eq. (55) is thus given by

$$G(\eta, \mathbf{s}, t) = \exp\left[\mathbf{g} \cdot \mathbf{L}^{-1} \cdot \left(\mathbf{I} - e^{-\mathbf{L}t}\right) \cdot \left(\mathbf{s} + \mathbf{L}^{-1} \cdot \mathbf{f}\right) + \left(h - \mathbf{g} \cdot \mathbf{L}^{-1} \cdot \mathbf{f}\right) t\right] \\ \times \exp\left\{\Gamma_0 \cdot \left[e^{-\mathbf{L}t} \cdot \mathbf{s} - \mathbf{L}^{-1} \cdot \left(\mathbf{I} - e^{-\mathbf{L}t}\right) \cdot \mathbf{f} - 1\right]\right\}.$$
(67)

The cumulant generating function of the signed cumulated transfers of particles from the I-th to the (I + 1)-th cell is defined as

$$Q(\lambda) \equiv \lim_{t \to \infty} -\frac{1}{t} \ln \left[ G(\eta = e^{-\lambda}, \mathbf{1}, t) \right] = \mathbf{g} \cdot \left( \mathbf{1} + \mathbf{L}^{-1} \cdot \mathbf{f} \right), \tag{68}$$

where the positivity of  $L_0$  has been used to obtain the explicit expression. We observe that

$$\mathbf{L} = \mathbf{M} \cdot \mathbf{L}_0 \cdot \mathbf{M}^{-1},\tag{69}$$

where

$$\mathbf{M} = \eta \mathbf{P}_{\mathrm{L}} + \mathbf{P}_{\mathrm{R}} \tag{70}$$

with the projection matrices

$$\mathbf{P}_{\mathrm{L}} = \begin{pmatrix} 1 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 \end{pmatrix} \qquad \text{and} \qquad \mathbf{P}_{\mathrm{R}} = \begin{pmatrix} 0 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 1 \end{pmatrix}.$$
(71)

The identity matrix in  $\mathbf{P}_{\mathrm{L}}$  is of dimension  $I \times I$ , while the identity matrix in  $\mathbf{P}_{\mathrm{R}}$  is  $(L - I) \times (L - I)$ . Since the projection matrices satisfy the condition  $\mathbf{P}_{\mathrm{L}} + \mathbf{P}_{\mathrm{R}} = \mathbf{I}$ , we thus have

$$\mathbf{M} = \mathbf{I} + (\eta - 1)\mathbf{P}_{\mathrm{L}},\tag{72}$$

$$\mathbf{M}^{-1} = \mathbf{I} + (\eta^{-1} - 1)\mathbf{P}_{\mathrm{L}}.$$
(73)

From the above related expressions, the cumulant generating function (68) can be written in the following form,

$$Q(\lambda) = \mathbf{g} \cdot \left[ \mathbf{I} - \mathbf{M} \cdot \mathbf{L}_0^{-1} \cdot \mathbf{M}^{-1} \cdot \mathbf{L}_0 \right] \cdot \mathbf{1}.$$
(74)

Because of Eqs. (72)-(73), we find that

$$Q(\lambda, t) = \mathbf{g} \cdot \left[ (1 - \eta) \mathbf{P}_{\rm L} + (1 - \eta^{-1}) \mathbf{L}_0^{-1} \cdot \mathbf{P}_{\rm L} \cdot \mathbf{L}_0 - (2 - \eta - \eta^{-1}) \mathbf{P}_{\rm L} \cdot \mathbf{L}_0^{-1} \cdot \mathbf{P}_{\rm L} \cdot \mathbf{L}_0 \right] \cdot \mathbf{1}.$$
 (75)

Using Eq. (61) and  $\mathbf{P}_{\mathrm{R}} = \mathbf{I} - \mathbf{P}_{\mathrm{L}}$ , the cumulant generating function becomes

$$Q(\lambda) = W_{+} \left( 1 - e^{-\lambda} \right) + W_{-} \left( 1 - e^{+\lambda} \right),$$
(76)

with the global transition rates given by

$$W_{+} = \boldsymbol{\Gamma}_{0} \cdot \boldsymbol{\mathsf{L}}_{0} \cdot \boldsymbol{\mathsf{P}}_{\mathrm{L}} \cdot \boldsymbol{\mathsf{L}}_{0}^{-1} \cdot \boldsymbol{\mathsf{P}}_{\mathrm{R}} \cdot \boldsymbol{\mathsf{L}}_{0} \cdot \boldsymbol{1},$$
(77)

$$W_{-} = \boldsymbol{\Gamma}_{0} \cdot \boldsymbol{\mathsf{L}}_{0} \cdot \boldsymbol{\mathsf{P}}_{\mathrm{R}} \cdot \boldsymbol{\mathsf{L}}_{0}^{-1} \cdot \boldsymbol{\mathsf{P}}_{\mathrm{L}} \cdot \boldsymbol{\mathsf{L}}_{0} \cdot \boldsymbol{1}.$$
(78)

The global transition rates (77)-(78) can be developed as

$$W_{+} = k_{+}^{2} \bar{N}_{\rm L} \left( \mathbf{L}_{0}^{-1} \right)_{1L} \qquad \text{and} \qquad W_{-} = k_{-}^{2} \bar{N}_{\rm R} \left( \mathbf{L}_{0}^{-1} \right)_{L1}.$$
(79)

Inverting the matrix  $\mathbf{L}_0$ , we get

$$\left( \mathbf{L}_{0}^{-1} \right)_{ij} = \begin{cases} \frac{k_{+}^{i-i} \left( k_{+}^{i} - k_{-}^{i} \right) \left( k_{+}^{L+1-j} - k_{-}^{L+1-j} \right)}{(k_{+} - k_{-}) \left( k_{+}^{L+1-} - k_{-}^{L+1} \right)} & \text{if } i \leq j, \\ \frac{k_{-}^{i-j} \left( k_{+}^{j} - k_{-}^{j} \right) \left( k_{+}^{L+1-i} - k_{-}^{L+1-i} \right)}{(k_{+} - k_{-}) \left( k_{+}^{L+1-} - k_{-}^{L+1} \right)} & \text{if } i > j. \end{cases}$$

$$(80)$$

So, the two global transition rates are calculated as

$$W_{+} = \bar{N}_{\rm L} \frac{k_{+}^{L+1}(k_{+} - k_{-})}{k_{+}^{L+1} - k_{-}^{L+1}} = \frac{D\bar{N}_{\rm L}}{\Delta x^{2}(L+1)} \frac{\beta e(\Phi_{\rm L} - \Phi_{\rm R})}{1 - \exp\left[-\beta e(\Phi_{\rm L} - \Phi_{\rm R})\right]},\tag{81}$$

$$W_{-} = \bar{N}_{\rm R} \frac{k_{-}^{L+1}(k_{+} - k_{-})}{k_{+}^{L+1} - k_{-}^{L+1}} = \frac{D\bar{N}_{\rm R}}{\Delta x^{2}(L+1)} \frac{\beta e(\Phi_{\rm R} - \Phi_{\rm L})}{1 - \exp\left[-\beta e(\Phi_{\rm R} - \Phi_{\rm L})\right]}.$$
(82)

#### References

- [1] Herbert Spohn. Large Scale Dynamics of Interacting Particles. Springer, 1991.
- [2] C. Kipnis and C. Landim. Scaling Limits of Interacting Particle Systems. Springer, 1999.
- [3] Thomas M. Liggett. Stochastic Interacting Systems: Contact, Voter and Exclusion Processes. Springer-Verlag, 1999.
- T. E. Harris. "Diffusion with "Collisions" between Particles". In: Journal of Applied Probability 2 (1965), pp. 323–338.
- [5] Frank Spitzer. "Interaction of Markov Processes". In: Advances in Mathematics 5 (1970), pp. 246–290.
- [6] E. Levine, D. Mukamel, and G. M. Schütz. "Zero-Range Process with Open Boundaries". In: Journal of Statistical Physics 120 (2005), pp. 759–778.
- [7] T. E. Harris. "Contact Interactions on a Lattice". In: The Annals of Probability 2 (1974), pp. 969– 988.
- [8] Denis J. Evans, E. G. D. Cohen, and G. P. Morriss. "Probability of Second Law Violations in Shearing Steady States". In: *Physical Review Letters* 71 (1993), pp. 2401–2404.
- [9] Denis J. Evans and Debra J. Searles. "The Fluctuation Theorem". In: Advances in Physics 51 (2002), pp. 1529–1585.
- G. Gallavotti and E. G. D. Cohen. "Dynamical Ensembles in Nonequilibrium Statistical Mechanics". In: *Physical Review Letters* 74 (1995), pp. 2694–2697.
- [11] G. Gallavotti and E. G. D. Cohen. "Dynamical Ensembles in Stationary States". In: Journal of Statistical Physics 80 (1995), pp. 931–970.
- [12] Jorge Kurchan. "Fluctuation Theorem for Stochastic Dynamics". In: Journal of Physics A: Mathematical and General 31 (1998), pp. 3719–3729.

- [13] Joel L. Lebowitz and Herbert Spohn. "A Gallavotti-Cohen-Type Symmetry in the Large Deviation Functional for Stochastic Dynamics". In: *Journal of Statistical Physics* 95 (1999), pp. 333–365.
- [14] Christian Maes. "The Fluctuation Theorem as a Gibbs Property". In: Journal of Statistical Physics 95 (1999), pp. 367–392.
- [15] R. J. Harris and G. M. Schütz. "Fluctuation Theorems for Stochastic Dynamics". In: Journal of Statistical Mechanics: Theory and Experiment, P07020 (2007).
- [16] Pierre Gaspard. "Multivariate Fluctuation Relations for Currents". In: New Journal of Physics 15, 115014 (2013).
- [17] Robert Zwanzig. "Time-Correlation Functions and Transport Coefficients in Statistical Mechanics". In: Annual Review of Physical Chemistry (1965), pp. 67–102.
- [18] H. B. G. Casimir. "On Onsager's Principle of Microscopic Reversibility". In: Reviews of Modern Physics 17 (1945), pp. 343–350.
- [19] David Andrieux and Pierre Gaspard. "Fluctuation Theorem and Onsager Reciprocity Relations". In: The Journal of Chemical Physics 121 (2004), pp. 6167–6174.
- [20] Jiayin Gu and Pierre Gaspard. "Microreversibility, Fluctuations and Nonlinear Transport in Transistors". In: *Physical Review E* 99, 012137 (2019).
- [21] M. Barbier and P. Gaspard. "Microreversibility, Nonequilibrium Current Fluctuations, and Response Theory". In: Journal of Physics A: Mathematical and Theoretical 51, 355001 (2018).
- [22] M. Barbier and P. Gaspard. "Microreversibility and Nonequilibrium Response Theory in Magnetic Fields". In: Journal of Physics A: Mathematical and Theoretical 52, 025003 (2019).
- [23] Jiayin Gu and Pierre Gaspard. "Counting Statistics and Microreversibility in Stochastic Models of Transistors". In: Journal of Statistical Mechanics: Theory and Experiment, 103206 (2020).
- [24] Yu-Xin Wu, Jiayin Gu, and H. T. Quan. "Full Counting Statistics and Fluctuation Theorem for the Currents in the Discrete Model of Feynman's Ratchet". In: *Physical Review E* 106, 014154 (2022).
- [25] Fan Zhang, Jiayin Gu, and H. T. Quan. "Full Counting Statistics, Fluctuation Relations, and Linear Response Properties in a One-Dimensional Kitaev Chain". In: *Physical Review E* 108, 024110 (2023).
- [26] Andre C. Barato and Udo Seifert. "Thermodynamic Uncertainty Relation for Biomolecular Processes". In: *Physical Review Letters* 114, 158101 (2015).
- [27] Todd R. Gingrich, Jordan M. Horowitz, Nikolay Perunov, et al. "Dissipation Bounds All Steady-State Current Fluctuations". In: *Physical Review Letters* 116, 120601 (2016).
- [28] Jordan Horowitz and Todd R. Gingrich. "Thermodynamic Uncertainty Relations Constrain Non-Equilibrium Fluctuations". In: Nature Physics 16 (2020), pp. 15–20.
- [29] Wonseok Hwang and Changbong Hyeon. "Energetic Costs, Precision, and Transport Efficiency of Molecular Motors". In: The Journal of Physics Chemistry Letters 9 (2018), pp. 513–520.
- [30] Sreekanth K. Manikandan, Deepak Gupta, and Supriya Krishnamurthy. "Inferring Entropy Production from Short Experiments". In: *Physical Review Letters*, 120603 (2020).
- [31] C. Maggi, F. Saglimbeni, V. Carmona Sosa, et al. "Thermodynamic Limits of Sperm Swimming Precision". In: *PRX Life* 1, 013003 (2023).
- [32] T. De Donder and P. Van Rysselberghe. Thermodynamic Theory of Affinity. Stanford University Press, 1936.
- [33] I. Prigogine. Introduction to Thermodynamics of Irreversible Processes. Wiley, 1967.
- [34] Dilip Kondepudi and Ilya Prigogine. Modern Thermodynamics: From Heat Engines to Dissipative Structures. John Wiley & Sons, 1998.
- [35] Richard Arratia. "The Motion of a Tagged Particle in the Simple Symmetric Exclusion System on Z". In: The Annals of Probability 11 (1983), pp. 362–373.
- [36] Herbert Spohn. "Tracer Diffusion in Lattice Gases". In: Journal of Statistical Physics 59 (1990), pp. 1227–1239.

- [37] Takashi Imamura, Kirone Mallick, and Tomohiro Sasamoto. "Large Deviations of a Tracer in the Symmetric Exclusion Process". In: *Physical Review Letters*, 160601 (2017).
- [38] Richard S. Ellis. Entropy, Large Deviations, and Statistical Mechanics. Springer, 2006.
- [39] Hugo Touchette. "The Large Deviation Approach to Statistical Mechanics". In: Physics Reports 478 (2009), pp. 1–69.
- [40] Amir Dembo and Ofer Zeitouni. Large Deviations Techniques and Applications. Second Edition. Springer, 2010.
- [41] Jiayin Gu and Pierre Gaspard. "Stochastic Approach and Fluctuation Theorem for Charge Transport in Diodes". In: *Physical Review E* 97, 052138 (2018).
- [42] Bernard Derrida. "Non-Equilibrium Steady States: Fluctuations and Large Deviations of the Density and of the Current". In: Journal of Statistical Mechanics: Theory and Experiment, P07023 (2007).
- [43] B. Derrida, B. Douçot, and P.-E. Roche. "Current Fluctuations in the One-Dimensional Symmetric Exclusion Process with Open Boundaries". In: *Journal of Statistical Physics* 115 (2004), pp. 717– 748.
- [44] Bernard Derrida. "Microscopic Versus Macroscopic Approaches to Non-Equilibrium Systems". In: Journal of Statistical Mechanics: Theory and Experiment, P01030 (2007).
- [45] Kirone Mallick. "The Exclusion Process: A Paradigm for Non-Equilibrium Behavior". In: Physica A: Statistical Mechanics and its Applications 418 (2015), pp. 17–48.
- [46] Pavel L. Krapivsky, Sidney Redner, and Eli Ben-Naim. A Kinetic View of Statistical Physics. Cambridge University Press, 2010.
- [47] Sheldon Katz, Joel L. Lebowitz, and Herbert Spohn. "Nonequilibrium Steady States of Stochastic Lattice Gas Models of Fast Ionic Conductors". In: *Journal of Statistical Physics* 34 (1983), pp. 497– 537.
- [48] Joel L. Lebowitz, Errico Presutti, and Herbert Spohn. "Microscopic Models of Hydrodynamic Behavior". In: Journal of Statistical Physics 51 (1988), pp. 841–862.
- [49] J. Schnakenberg. "Network Theory of Microscopic and Macroscopic Behavior of Master Equation Systems". In: *Reviews of Modern Physics* 48 (1976), pp. 571–585.
- [50] D.-Q. Jiang, M. Qian, and M.-P. Qian. Mathematical Theory of Nonequilibrium Steady States: On the Frontier of Probability and Dynamical Systems. Springer, 2004.
- [51] John David Jackson. Classical Electrodynamics. Third Edition. Wiley, 1999.
- [52] David Andrieux and Pierre Gaspard. "Stochastic Approach and Fluctuation Theorem for Ion Transport". In: Journal of Statistical Mechanics: Theory and Experiment, P02057 (2009).
- [53] Daniel T. Gillespie. "A General Method for Numerically Simulating the Stochastic Time Evolution of Coupled Chemical Reactions". In: *Journal of Computational Physics* 22 (1976), pp. 403–434.
- [54] George Bakewell-Smith, Federico Girotti, Mădălin Guţă, et al. "General Upper Bounds on Fluctuations of Trajectory Observables". In: *Physical Review Letters* 131, 197101 (2023).
- [55] Pierre Gaspard and Raymond Kapral. "Finite-Time Fluctuation Theorem for Diffusion-Influenced Surface Reactions". In: Journal of Statistical Mechanics: Theory and Experiment 8, 083206 (2018).