## RATIONAL TORSION OF GENERALISED DRINFELD MODULAR JACOBIANS OF PRIME POWER LEVEL

### MAR CURCÓ IRANZO

ABSTRACT. For a prime  $\mathfrak{p} \subseteq \mathbb{F}_q[T]$  and a positive integer r, we consider the generalised Jacobian  $J_0(\mathfrak{n})_{\mathfrak{m}}$  of the Drinfeld modular curve  $X_0(\mathfrak{n})$  of level  $\mathfrak{n} = \mathfrak{p}^r$ , with respect to the modulus  $\mathfrak{m}$  consisting of all cusps on the modular curve. We show that the  $\ell$ -primary part of the group  $J_0(\mathfrak{n})_{\mathfrak{m}}(\mathbb{F}_q(T))_{tor}[\ell^{\infty}]$  is trivial for all primes  $\ell$  not dividing  $q(q^2-1)$ . Our results establish a function field analogue to those of Yamazaki–Yang [27] for the classical case.

### 1. Introduction

1.1. **Torsion of modular (semi-)abelian varieties.** Given a positive integer N, let  $X_0(N)$  denote the modular curve of level N for the congruence subgroup  $\Gamma_0(N) = \{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \} \in \operatorname{SL}_2(\mathbb{Z}) : c \equiv 0 \pmod{N} \}$ . This is a projective nonsingular curve defined over  $\mathbb{Q}$ . For N prime, Ogg [18, Conjecture 2] considered the Jacobian variety  $J_0(N)$  of  $X_0(N)$  as a natural family of candidates of higher-dimensional abelian varieties whose torsion could be determined. His predictions on  $J_0(N)$ , based on computational evidence, are now known as Ogg's Conjecture. To state it, write  $C_N$  for the image of the degree-zero divisors with support at the cusps of  $X_0(N)$  in  $J_0(N)$  and define  $C_N(\mathbb{Q}) := C_N \cap J_0(N)(\mathbb{Q})$ ; Ogg proved that, for  $N \geq 5$  prime, the cyclic group  $C_N(\mathbb{Q})$  has order  $n = \frac{p-1}{\gcd(p-1,12)}$  and made the following conjecture: the cyclic group  $C_N(\mathbb{Q})$  of order n = 1 is the full torsion subgroup  $J_0(N)(\mathbb{Q})_{\text{tors}}$  of  $J_0(N)(\mathbb{Q})$ . The conjecture was proved later by Mazur [17, Theorem I] using the Eisenstein ideal machinery developed in his work. A natural generalisation of this conjecture is the following.

**Conjecture A** (Generalised Ogg's conjecture). Let N be a positive integer. Then

$$C_N(\mathbb{Q}) = J_0(N)(\mathbb{Q})_{tors}$$
.

Notice that by theorems of Manin [16, Corollary 3.6] and Drinfeld [5] the group  $C_N$  is finite, thus, the inclusion  $C_N(\mathbb{Q}) \subseteq J_0(N)(\mathbb{Q})_{\text{tors}}$  always holds. Moreover, if C(N) is the subgroup formed by the equivalence classes of  $\mathbb{Q}$ -rational divisors in  $C_N(\mathbb{Q})$ , i.e., the divisor classes fixed under the action of the absolute Galois group of  $\mathbb{Q}$ , a further conjecture states that  $C_N(\mathbb{Q})$  is equal to C(N).

**Conjecture B** (Rational cuspidal groups conjecture). Let N be a positive integer. Then

$$C_N(\mathbb{Q}) = C(N)$$
.

A strategy for proving Conjectures A and B consists in decomposing the groups C(N),  $C_N(Q)$  and  $J_0(\mathbb{Q})_{\text{tors}}$  into the direct sum of their  $\ell$ -primary parts for all primes  $\ell$  and showing that each respective  $\ell$ -primary part equals that of the other groups. In this spirit, the strongest result to date says that  $C(N)[\ell^{\infty}]$  and  $J_0(\mathbb{Q})_{\text{tors}}[\ell^{\infty}]$ 

<sup>2020</sup> Mathematics Subject Classification. 11G09 (11G18, 11F03, 14G05, 14H40, 14G35).

Key words and phrases. Generalized Jacobian, Drinfeld modular curves, Cuspidal divisor group, Rational torsion.

agree for all primes  $\ell$  with  $\ell \geq 5$ , or  $\ell = 3$  and either  $3^2 \nmid N$  or N has a prime divisor congruent to -1 modulo 3. This result is due to Yoo [29] and is based on previous work of Ohta [19] on the Eistenstein ideal. See Section 1 in [29] for a summary of more (partial) results on Conjectures A and B. On the other hand, the elaborate task of determining the structure of C(N) has recently been accomplished in further work by Yoo [28]. A meticulous study of modular units, in particular of  $\eta$ -quotients, allows him to determine the structure and generators of  $C(N)[\ell^{\infty}]$  for all  $\ell$ . Hence, as of today we can say that the structure of  $J_0(\mathbb{Q})_{tors}[\ell^{\infty}]$  is understood for the primes  $\ell$  in Yoo's result [29, Theorem 1.4].

In a parallel direction, a number of authors have moved on to determining the structure of  $J_0(N)_{\mathbf{m}}(\mathbb{Q})_{\mathrm{tors}}$ , where  $J_0(N)_{\mathbf{m}}(\mathbb{Q})_{\mathrm{tors}}$  is the generalised Jacobian of  $X_0(N)$  with respect to a cuspidal modulus  $\mathbf{m}$ . That is, if we let  $\mathrm{Cusp}(X_0(N))$  be the set of cusps of  $X_0(N)$  and  $\mathbf{m} = \sum_{\omega \in \mathrm{Cusp}(X_0(N))} \omega$ , the generalised Jacobian is the group of invertible line bundles on the singular curve that results from  $X_0(N)$  after the identification of all the points in  $\mathbf{m}$ . The general construction of this group variety can be found in [?] and [25]. In our case,  $J_0(N)_{\mathbf{m}}(\mathbb{Q})_{\mathrm{tors}}$  is what is called a semi-abelian variety; that is, the extension of  $J_0(N)$  – an abelian variety – by a commutative affine linear algebraic group  $L_{\mathbf{m}}$  isomorphic to a number of copies of  $\mathbb{G}_m$ . The group  $J_0(N)_{\mathbf{m}}(\mathbb{Q})$  is not not necessarily finitely generated anymore, but the subgroup  $J_0(N)_{\mathbf{m}}(\mathbb{Q})_{\mathrm{tors}}$  is finite. Just like the Jacobian  $J_0(N)$ , the generalised Jacobian  $J_0(N)_{\mathbf{m}}$  is an object of high (arithmetic) relevance when it come to the study of the curve  $X_0(N)$ . With this underlying motivation, Yamazaki-Yang [27], Wei-Yamazaki [26], Jordan-Ribet-Scholl [14] and the author [4] study the structure of  $J_0(N)_{\mathbf{m}}(\mathbb{Q})_{\mathrm{tors}}$  for different N and with different approaches. The following result of Yamazaki-Yang is relevant for us in this paper.

**Theorem 1.1.** [27, Theorem 1.1.3] Let p be a prime number and n be a positive integer. For  $\ell \nmid 2p$ , the group  $J_0(p^n)_{\mathfrak{m}}(\mathbb{Q})[\ell^{\infty}]$  is trivial. Moreover, if Conjectures A and B hold, we have

$$J_0(p^n)_{\mathbf{m}}(\mathbb{Q})_{\mathrm{tors}} \simeq \prod_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \mathbb{Z}/(2p^i\mathbb{Z}) \times \prod_{i=1}^{\lfloor \frac{n}{2} \rfloor} \mathbb{Z}/(2p^i\mathbb{Z}).$$

1.2. **Torsion of Drinfeld modular (semi-)abelian varieties.** In the current paper, we study an analogue of the results of Yamazaki-Yang in the function field setting.

**Notation 1.2.** Let p be a prime and fix  $q = p^t$ . We fix the following notation for the rest of the paper.

```
= finite field of characteristic p with q elements
              = \mathbb{F}_a[T] polynomial ring in T over \mathbb{F}_a
A
K
              = \mathbb{F}_{q}(T) rational function field
              =\mathbb{F}_q((\pi)) the completion of K at the infinite place; \pi:=T^{-1}
              = |\cdot|_{\infty} = normalised absolute value on K_{\infty}; that is, |T|_{\infty} := q
|\cdot|
0_{\infty}
              = \mathbb{F}_q[[\pi]] ring of integers in K_{\infty}
              = the completion of an algebraic closure of K_{\infty}
\mathbb{C}_{\infty}
G
              = group scheme GL(2) over \mathbb{F}_a
              = scalar matrices in G
Z
```

Before introducing our main result, we present the analogue of Conjectures A and B for Drinfeld modular curves. Given an monic element  $\mathfrak{n} \in \mathcal{A}$ , let  $X_0(\mathfrak{n})$  denote the Drinfeld modular curve of level  $\mathfrak{n}$  for the congruence subgroup  $\Gamma_0(\mathfrak{n})$  consisting of matrices in  $\Gamma = G(\mathcal{A})$  that are upper triangular modulo  $\mathfrak{n}$  – see Section 2.

We regard  $X_0(\mathfrak{n})$  as a smooth projective curve defined over K – see [10]. Let  $J_0(\mathfrak{n})$  be the Jacobian variety of  $X_0(\mathfrak{n})$  and  $J_0(\mathfrak{n})(K)_{\text{tors}}$  its rational torsion subgroup. The Lang-Néron theorem [15], which generalises the Mordell-Weil theorem, asserts that  $J_0(\mathfrak{n})(K)$  is finitely generated, and thus that  $J_0(\mathfrak{n})(K)_{\text{tors}}$  is finite. Similar as before, let  $\mathcal{C}_\mathfrak{n}$  be the cuspidal subgroup of  $J_0(\mathfrak{n})$ , that is, the image of the degree-zero divisors with support at the cusps of  $X_0(\mathfrak{n})$  in  $J_0(\mathfrak{n})$ , and let  $\mathcal{C}_\mathfrak{n}(K) := \mathcal{C}_\mathfrak{n} \cap J_0(\mathfrak{n})(K)$  be the rational cuspidal subgroup of  $J_0(\mathfrak{n})$ . Moreover, let  $\mathcal{C}(\mathfrak{n})$  be the rational cuspidal divisor class group of  $X_0(\mathfrak{n})$ , which is the subgroup of  $\mathcal{C}_\mathfrak{n}$  formed by the equivalence classes of K-rational divisors in  $C_N(\mathbb{Q})$ , i.e., the divisor classes fixed under the action of the absolute Galois group of K. Since  $\mathcal{C}_\mathfrak{n}$  is a finite group – it is generated by a finite set of torsion elements [8] –, we have the tower of inclusions

(1) 
$$\mathcal{C}(\mathfrak{n}) \subseteq \mathcal{C}_{\mathfrak{n}}(K) \subseteq J_0(\mathfrak{n})(K)_{\text{tors}}.$$

Analogues of Conjectures A and B in this setting predict the equality of two the inclusions.

**Conjecture C.** Let  $n \in A$  be monic. Then we have

$$\mathcal{C}(\mathfrak{n}) = \mathcal{C}_{\mathfrak{n}}(K) = J_0(\mathfrak{n})(K)_{\text{tors}}.$$

As analogous results to those of Ogg and Mazur in this direction, Gekeler [7] computed the order of the cyclic group  $\mathcal{C}_p$  and Pál [20] proved Conjecture C for  $\mathfrak{n}=\mathfrak{p}$  prime. Their techniques resembled those of the classical modular setting, that is, they employed a study of the discriminant function and the development of the appropriate Eisenstein ideal machinery in this setting. Further progress in determining the structure of  $\mathcal{C}(\mathfrak{n})$  and proving Conjecture C in full generality can be found in works of e.g. Papikian-Wei [21], Papikian [22], [23] and Ho [12], [13]—see [1] for a thorough survey on the topic.

In the Drinfeld case, one can also consider the generalised Jacobian  $J_0(\mathfrak{n})_{\mathbf{m}}$  of  $X_0(\mathfrak{n})$  with cuspidal modulus  $\mathbf{m} = \sum_{\omega \in \operatorname{Cusp}(X_0(\mathfrak{n}))} \omega$ . In [26], Wei-Yamazaki compute the structure of  $J_0(\mathfrak{n})_{\mathbf{m}}(K)_{\text{tors}}$  up to  $q(q^2-1)$ -torsion. The goal of this paper is two-fold: compute the structure of the group  $J_0(\mathfrak{n})$  for  $\mathfrak{n} = \mathfrak{p}^r$  and compare the results for this level to the ones in the classical case when  $N = p^r$  given in 1.1. Our main result is the following.

**Theorem 1.3.** Let  $\mathfrak{p} \in \mathcal{A}$  monic irreducible and r an integer. For all odd primes  $\ell$  with  $\ell \nmid q(q^2-1)$  the subgroup  $J_0(\mathfrak{p}^r)_{\mathbf{m}}(\mathbb{Q})[\ell^{\infty}]$  is trivial.

The results in Theorem 1.3 show that the situation in the Drinfeld case is analogous to the one in Theorem 1.1.

The structure of the paper is the following. In Section 2 we present some background on the Drinfeld modular curves  $X_0(\mathfrak{n})$  and their cusps. In particular, in Theorem 2.2 we describe the residue fields of the cuspidal closed points of  $X_0(\mathfrak{n})$ . We also review known results on  $J_0(\mathfrak{n})$  and  $\mathfrak{C}(\mathfrak{n})$  in the direction of Conjecture C. In Section 3 we recall some results on the Drinfeld generalised Jacobian  $J_0(\mathfrak{n})_{\mathfrak{m}}$ . We introduce a group homomorphism denoted by  $\delta$  and reduce the proof of Theorem 1.3 to determining the injectivity of a related homomorphism  $\overline{\delta}$  on  $\mathfrak{C}(\mathfrak{n})$ . Finally, in the Sections 4 and 5 we prove the main result. The strategy of the proof then goes as follows. In Section 4 we develop the tools to compute the image of certain cuspidal divisors appearing in Theorem 2.8. In Section 5 we use that computation to construct the matrix that represents the map  $\overline{\delta}|_{\mathfrak{C}(\mathfrak{p}^r)}$ . Finally, that representation allows us to see that  $\overline{\delta}|_{\mathfrak{C}(\mathfrak{p}^r)}$  is injective.

### 2. BACKGROUND

Let  $\Omega = \mathbb{C}_{\infty} - K_{\infty}$  be the Drinfeld upper half plane. The group  $\Gamma_0(\mathfrak{n})$  acts on  $\Omega$  by linear fractional transformations. The Drinfeld modular curve  $Y_0(\mathfrak{n})$  can be described analytically as the quotient  $\Gamma_0(\mathfrak{n}) \setminus \Omega$ . The points of  $Y_0(\mathfrak{n})$  parametrise rank 2 Drinfeld modules equipped with a cyclic subgroup of order  $\mathfrak{n}$ . Then,  $X_0(\mathfrak{n})$  is the unique smooth projective curve over K containing  $Y_0(\mathfrak{n})$ , that is, the canonical non-singular compactification of  $Y_0(\mathfrak{n})$ . The set of cusps of  $X_0(\mathfrak{n})$  is  $\mathrm{Cusp}(X_0(\mathfrak{n})) = X_0(\mathfrak{n}) \setminus Y_0(\mathfrak{n})$ . after base change to  $\mathbb{C}_{\infty}$ , there is a canonical 1-to-1 correspondence between cusps of  $X_0(\mathfrak{n})_{\mathbb{C}_{\infty}}$  and  $\Gamma_0(\mathfrak{n}) \setminus \mathbb{P}_1(K)$ . A divisor D of  $X_0(\mathfrak{n})_{\mathbb{C}_{\infty}}$  with  $\mathrm{supp}(D) \subseteq \mathrm{Cusps}(X_0(\mathfrak{n})_{\mathbb{C}_{\infty}})$  is called a cuspidal divisor. Then the group  $\mathfrak{C}_{\mathfrak{n}}$  (resp.  $\mathfrak{C}(\mathfrak{n})$ ) defined in the introduction is the subgroup of  $J_0(\mathfrak{n})$  of classes of cuspidal divisors (resp. K-rational cuspidal divisors). A recent remarkable result on Conjecture  $\mathbb{C}$  is the following.

**Theorem 2.1.** [13, Theorem 1.7] Let  $\mathfrak{n} = \mathfrak{p}^r \in \mathcal{A}$  be a prime power ideal with  $r \geq 2$ . For any prime  $\ell \nmid q(q-1)$ , we have

$$\mathfrak{C}(\mathfrak{p}^r)[\ell^{\infty}] = J_0(\mathfrak{p}^r)(K)_{\text{tors}}[\ell^{\infty}].$$

Motivated by Conjecture  $\mathbb{C}$ , a number of authors have studied the structure of the group  $\mathbb{C}(\mathfrak{n})$  in order to understand the torsion of  $J_0(\mathfrak{n})$  better. For that, understanding the points in  $\mathrm{Cusps}(X_0(\mathfrak{n}))$  is crucial. We proceed by giving some background knowledge about cusps.

**Theorem 2.2.** ([7, pg. 196], [21, Lemma 3.1(1)]) Let  $\mathfrak{n} \in \mathcal{A}$  be monic. The cusps of  $X_0(\mathfrak{n})_{\mathbb{C}_{\infty}}$  are in bijection with  $\Gamma_0(\mathfrak{n})/\mathbb{P}^1(K)$ . Every cusp in  $\mathrm{Cusp}(X_0(\mathfrak{n}))_{\mathbb{C}_{\infty}}$  has a representative  $\begin{bmatrix}\mathfrak{a}\\\mathfrak{m}\end{bmatrix}$  for some elements  $\mathfrak{a},\mathfrak{m} \in \mathcal{A}$ , with  $\mathfrak{m} \mid \mathfrak{n}$  and  $\mathrm{gcd}(\mathfrak{a},\mathfrak{n}) = 1$ . In particular, we can describe them as the set

$$\operatorname{Cusp}(X_0(\mathfrak{n})_{\mathbb{C}_\infty}) \simeq \left\{ \left[ \begin{smallmatrix} \mathfrak{a} \\ \mathfrak{m} \end{smallmatrix} \right] \in \Gamma_0(\mathfrak{n}) \setminus \mathbb{P}^1(K) : \mathfrak{a}, \mathfrak{m} \in \mathcal{A} \text{ monic, } \mathfrak{m} \mid \mathfrak{n}, \text{ and } \gcd(\mathfrak{a}, \mathfrak{n}) = 1 \right\}.$$

For two divisors m and m' of n, we have

$$\left[\begin{smallmatrix}\mathfrak{a}\\\mathfrak{m}\end{smallmatrix}\right] = \left[\begin{smallmatrix}\mathfrak{a}'\\\mathfrak{m}'\end{smallmatrix}\right] \iff \mathfrak{m} = \mathfrak{m}' \text{ and } k\mathfrak{a} \equiv \mathfrak{a}' \pmod{\mathfrak{d}},$$

where  $\mathfrak{d} = \mathfrak{d}(\mathfrak{m}) := \gcd(\mathfrak{m}, \mathfrak{n}/\mathfrak{m})$  and  $k \in \mathbb{F}_a^{\times}$ .

A cusp  $\omega$  of  $X_0(\mathfrak{n})_{\mathbb{C}_{\infty}}$  with a representative of the form  $[\mathfrak{m}]$  as in Theorem 2.2 is said to be of height  $\mathfrak{m}$  [11, (2.4)]. Let  $(P_{\mathfrak{m}})$  be the divisor given by the sum of all the cusps of  $X_0(\mathfrak{n})_{\mathbb{C}_{\infty}}$  of fixed height  $\mathfrak{m} \mid \mathfrak{n}$ . The cuspidal divisors  $(P_{\mathfrak{m}})$  are K-rational, that is, they are invariant under the action  $\operatorname{Gal}(\overline{K}/K)$ , cf. [9, Prop 6.3]. Indeed, the cusps of  $X_0(\mathfrak{n})_{\mathbb{C}_{\infty}}$  of the same height form an orbit under  $\operatorname{Gal}(\overline{K}/K)$ . For  $\mathfrak{m} = \mathfrak{p}^i$  with  $i \in \{0, \ldots, r\}$ , we let  $P_i$  be the closed K-point of  $X_0(\mathfrak{n})/K$  given by  $P_i := (P_{\mathfrak{m}})$ . The following results describe the residue fields of the closed points  $P_i$ .

**Proposition 2.3.** Let  $\mathfrak{n} = \mathfrak{p}^r$  and let  $\mathfrak{m} \mid \mathfrak{n}$ . Then the orbit of the cusp  $\begin{bmatrix} 1 \\ \mathfrak{m} \end{bmatrix}$  under the action of  $GL_2(\mathcal{A})$  is of size  $\varphi(\mathfrak{m}, \mathfrak{n}/\mathfrak{m})/(q-1)$ .

*Proof.* This follows from Theorem 2.2.

**Definition 2.4.** For a polynomial  $M \in \mathbb{F}_p[T]$ , let K(M) denote the cyclotomic function field obtained after adjoining the M-torsion of the Carlitz module in  $\overline{\mathbb{F}_p(T)}$  to  $K = \mathbb{F}_p(T)$ . See [3] for a survey on cyclotomic function fields.

**Theorem 2.5.** Let  $\mathfrak{m} = \mathfrak{p}^i$ ,  $\mathfrak{d} = \mathfrak{d}(\mathfrak{m}) = \gcd(\mathfrak{m}, \mathfrak{n}/\mathfrak{m}) = \mathfrak{p}^{\min\{i, r-i\}}$ , and let  $P_i$  be the closed point defined by the cuspidal divisor  $(P_m)$ . The residue field  $K(P_i)$  of the point  $P_i$  is the field  $K^+(\mathfrak{d})$ , that is, the maximal real subfield of the  $\mathfrak{d}$ -th cyclotomic field  $K(\mathfrak{d})$  of K.

*Proof.* By classical results of Gekeler [9, Rmk 4.7], we know that the cusps of  $X_0(\mathfrak{n})(\mathbb{C}_{\infty})$  are  $K^+(\mathfrak{n})$ -rational points. Hence, we know that  $K(P_i)$  is a subgroup of the cyclic group  $K^+(\mathfrak{n})$ . Consider now the cusp  $\begin{bmatrix} 1 \\ \mathfrak{m} \end{bmatrix}$ . By Proposition 2.3, we know that the orbit of  $\begin{bmatrix} 1 \\ \mathfrak{m} \end{bmatrix}$  under the action of  $GL_2(\mathcal{A})$  is of size  $\varphi(\mathfrak{m},\mathfrak{n}/\mathfrak{m})/(q-1)$ . It follows that the field  $K(P_i)$  is the unique degree  $\varphi(\mathfrak{m},\mathfrak{n}/\mathfrak{m})/(q-1)$  subfield of the cyclic extension  $K^+(\mathfrak{n})/K$ . That is, we get that  $K(P_i) = K(\mathfrak{d})^+$ .

In particular, there are two points in  $\operatorname{Cusp}(X_0(\mathfrak{n}))$  defined over K: the cusp at 0 (given by the class  $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ ) and the cusp at  $\infty$  (given by the class  $\begin{bmatrix} 1 \\ n \end{bmatrix}$ ). We denote them by  $\omega_0$  and  $\omega_\infty$  respectively.

When  $\mathfrak{n} = \mathfrak{p}$  is a monic irreducible polynomial  $\mathfrak{p} \in \mathcal{A}$ ,  $\omega_0$  and  $\omega_{\infty}$  are the only cusp points of  $X_0(\mathfrak{n})$ . Hence, the group  $\mathcal{C}(\mathfrak{n})$  is the cyclic group generated by the class of the divisor  $\omega_0 - \omega_\infty$ . Gekeler shows in [7] that the latter is an element of order  $N(\mathfrak{p})$ , see Definition 2.6. When the level  $\mathfrak{n}$  is not prime, determining the structure of C(n) is more involved. In [12], Ho studies the structure of the group C(n) for  $n = p^r$  a power of a monic irreducible polynomial  $\mathfrak{p} \in \mathcal{A}$ . In particular, he constructs a basis of  $\mathcal{C}(\mathfrak{p}^r)$  for any  $r \geq 1$ . The constructed basis will be very useful to prove our results. For convenience of the reader, we reproduce the results of Ho below.

**Definition 2.6.** Fix a prime  $\mathfrak{p} \in \mathcal{A}$ . We define the integers  $M(\mathfrak{p})$  and  $N(\mathfrak{p})$  as follows:

$$M(\mathfrak{p}) := \frac{|\mathfrak{p}|^2 - 1}{q^2 - 1}$$
 and  $N(\mathfrak{p}) := \begin{cases} \frac{|\mathfrak{p}| - 1}{q^2 - 1}, & \text{if deg}(\mathfrak{p}) \text{ is even;} \\ \frac{|\mathfrak{p}| - 1}{q - 1}, & \text{otherwise.} \end{cases}$ 

**Definition 2.7.** [12, Section 3.2] For each  $i \in \{1, ..., r-2\}$  we define  $C_i := P_i - \deg(P_i)[\infty] \in \operatorname{Div}_{\text{cusp}}^0(X_0(\mathfrak{n})(K))$ . We further define  $D_0$  and  $D_{r-1} \in \text{Div}^0_{\text{cusp}}(X_0(\mathfrak{n})(K))$  as

$$D_0 := C_0 + (q-1) \left( \sum_{1 \le i \le \lfloor \frac{r}{2} \rfloor} C_i + \sum_{\lfloor \frac{r}{2} \rfloor + 1 \le i \le r-1} |\mathfrak{p}|^{2i-r} C_i \right)$$

and  $D_{r-1}$  through the case-by-case definition given in the Appendix – see Definition 6.1 – for convenience of the reader.

**Theorem 2.8.** [12, Theorem 3.5] Let  $\mathfrak{p} \in A$  be a prime and  $r \geq 2$ . The elements  $C_i$ ,  $D_{r-1}$ , and  $D_0$  are defined in Definition 2.7. Then

$$\mathcal{C}(\mathfrak{p}^r) = \left(\bigoplus_{i=1}^m \langle \overline{C_i} \rangle\right) \oplus \left(\bigoplus_{i=m+1}^{r-2} \langle \overline{C_i} - |\mathfrak{p}|C_{i+1} \rangle\right) \oplus \langle \overline{D_{r-1}} \rangle \oplus \langle \overline{D_0} \rangle,$$

where  $m := \left| \frac{r-1}{2} \right|$  and

- (1)  $\operatorname{ord}(\overline{C_i}) = |\mathfrak{p}|^{r-i} M(\mathfrak{p}) \text{ for } 1 \le i \le m;$
- (2) ord $(\underline{C_i |\mathfrak{p}|C_{i+1}}) = |\mathfrak{p}|^i M(\mathfrak{p})$  for  $m+1 \le i \le r-2$ ;
- (3)  $\operatorname{ord}(\overline{D_{r-1}}) = M(\mathfrak{p});$ (4)  $\operatorname{ord}(\overline{D_0}) = N(\mathfrak{p}).$

Theorem 2.8 is obtained after taking a close look at the modular units of  $X_0(\mathfrak{n})$  for  $\mathfrak{n}=\mathfrak{p}^r$ . In particular, in [12] the author constructs a map  $g: \mathrm{Div}^0(\mathrm{Cusp}(X_0(\mathfrak{n})) \longrightarrow \mathcal{E}_{\mathfrak{n}} \otimes_{\mathbb{Z}} \mathbb{Q}$  which, in analogy to the classical case, to each degree zero cuspidal divisor D attaches a modular unit that has D as a divisor. This is done through so-called  $\Delta$ -quotients.

**Definition 2.9.** Let  $\mathfrak{n} \in \mathcal{A}$  and consider the Drinfeld discriminant function  $\Delta(z): \Omega \to \mathbb{C}_{\infty}$  defined by Gekeler in [7]. For  $\mathfrak{m} \in \mathcal{A}$  monic, let  $\Delta_{\mathfrak{m}} := \Delta(\mathfrak{m}z)$ . All functions  $\Delta_{\mathfrak{m}}$  with  $\mathfrak{m} \mid \mathfrak{n}$  are modular forms on  $\Omega$  of weight  $q^2-1$  and type 0 for  $\Gamma_0(\mathfrak{n})$ , see [7]. A  $\Delta$ -quotient on  $\Gamma_0(\mathfrak{n})$  is a function in  $\mathcal{E}_{\mathfrak{n}} \otimes_{\mathbb{Z}} \mathbb{Q}$  for  $\Gamma_0(\mathfrak{n})$  of the form  $\prod_{\mathfrak{m} \mid \mathfrak{n}} \Delta_{\mathfrak{m}}(z)^{r_{\mathfrak{m}}}$ , with  $r_{\mathfrak{m}} \in \mathbb{Q}$  and  $\sum_{\mathfrak{m} \mid \mathfrak{n}} r_{\mathfrak{m}} = 0$ .

**Remark 2.10.** Notice that since  $\frac{\Delta}{\Delta_{\mathfrak{m}}} \in \mathcal{E}_{\mathfrak{n}}$  for all  $\mathfrak{m} \mid \mathfrak{n}$ , a  $\Delta$ -quotient is indeed an element of  $\mathcal{E}_{\mathfrak{n}} \otimes_{\mathbb{Z}} \mathbb{Q}$ .

The map g can then be constructed by understanding the order of the zeroes of the functions  $\Delta_{\mathfrak{m}}$  at all the cusps of  $X_0(\mathfrak{n})$ . In [12], the author uses the orders of such functions for  $\mathfrak{n} = \mathfrak{p}^r$  described in [7] and constructs

$$g: \operatorname{Div}^{0}(\operatorname{Cusp}(X_{0}(\mathfrak{n}))(K) \longrightarrow \mathcal{E}_{\mathfrak{n}} \otimes_{\mathbb{Z}} \mathbb{Q}$$

$$C:= \sum a_{\mathfrak{m}} P_{\mathfrak{m}} \longmapsto \prod \Delta_{\mathfrak{m}}(z)^{r_{\mathfrak{n}}}$$

$$C := \sum_{\mathfrak{m} \mid \mathfrak{n}} a_{\mathfrak{m}} P_{\mathfrak{m}} \qquad \longmapsto \qquad \prod_{\mathfrak{m} \mid \mathfrak{n}} \Delta_{\mathfrak{m}}(z)^{r_{\mathfrak{m}}}$$

where the linear map g is described by

(2) 
$$\begin{pmatrix} r_1 \\ r_{\mathfrak{p}} \\ \vdots \\ r_{\mathfrak{p}^r} \end{pmatrix} = \frac{1}{(q-1)|\mathfrak{p}|^{r-1}(|\mathfrak{p}|^2-1)} \Upsilon(\mathfrak{p}^r) \cdot \begin{pmatrix} a_1 \\ a_{\mathfrak{p}} \\ \vdots \\ a_{\mathfrak{p}^r} \end{pmatrix},$$

and the matrix  $\Upsilon(\mathfrak{p}^r) = (\upsilon[i][j])_{i,j}$  is described by

$$\upsilon[i][j] = \begin{cases} (|\mathfrak{p}|^2 + 1)|\mathfrak{p}|^{m(j) - 1}, & \text{if } 1 \le i = j \le r - 1; \\ -|\mathfrak{p}|^{m(j)}, & \text{if } |i - j| = 1 \text{ and } j \ne 0, r; \\ (q - 1)|\mathfrak{p}|, & \text{if } (i, j) = (0, 0) \text{ or } (r, r); \\ 1 - q, & \text{if } (i, j) = (1, 0) \text{ or } (r - 1, r); \\ 0, & \text{otherwise.} \end{cases}$$

Then, the strategy to prove Theorem 2.8 is to use an appropriate root of the  $\Delta$ -quotient  $\frac{\Delta(z)}{\Delta(\mathfrak{n}z)}$  to to find an optimal way of computing the order of a cuspidal divisor D in  $\mathcal{C}(\mathfrak{n})$ . This is also used to determine whether two  $\Delta$ -quotients have associated cuspidal divisors that generate groups in  $J_0(\mathfrak{n})$  with trivial intersection. This allows Ho to determine a cyclic decomposition of  $\mathcal{C}(\mathfrak{n})$ .

The functions  $g(D) \in \mathcal{E}_{\mathfrak{n}} \otimes_{\mathbb{Z}} \mathbb{Q}$  for  $D \in \{C_i : 1 \leq i \leq \lfloor \frac{r-1}{2} \rfloor\} \cup \{C_i - |\mathfrak{p}|C_{i+1} : \lfloor \frac{r+1}{2} \rfloor \leq i \leq r-2\} \cup \{D_{r-1}\}$  will play an important role in proving our results. In particular, we will need a way of computing their leading Fourier coefficients at the different cusps of  $X_0(\mathfrak{n})$ . The following result provides such a tool and will be used later in Proposition 3.2.

**Lemma 2.11.** Let  $\begin{bmatrix} \mathfrak{a}' \\ \mathfrak{m}' \end{bmatrix} \in \operatorname{Cusp}(X_0(\mathfrak{n})(\mathbb{C}_{\infty}))$  be a cusp with  $\mathfrak{m}' = \mathfrak{p}^k$  and  $\gcd(\mathfrak{a}', \mathfrak{m}') = 1$ . Let the matrix  $A = \begin{pmatrix} a & b \\ \mathfrak{p}^k & \mathfrak{a}' \end{pmatrix} \in \operatorname{SL}_2(\mathcal{A})$  be such that  $A^{-1} \cdot \omega_{\infty} = \begin{bmatrix} \mathfrak{a}' \\ \mathfrak{m}' \end{bmatrix}$ . Then, for a monic divisor  $\mathfrak{m} \mid \mathfrak{n}$  we have the identity

$$\Delta_{\mathfrak{m}}(A^{-1} \cdot z) = \left(\frac{d}{\mathfrak{m}}(-\mathfrak{p}^k z + a)\right)^{q^2 - 1} \Delta\left(\frac{d^2}{\mathfrak{m}}z + \frac{\eta d}{\mathfrak{m}}\right),$$

where  $d = \gcd(\mathfrak{m}, \mathfrak{p}^k)$  and  $\eta \in A$ .

Proof. By definition we have

$$\Delta_{\mathfrak{m}}(A^{-1}z) = \Delta\left(\frac{\mathfrak{m}\alpha'z - \mathfrak{m}b}{\mathfrak{p}^kz + a}\right) = \Delta(\alpha z),$$

where  $\alpha = \binom{\mathfrak{m}\alpha' - \mathfrak{m}b}{\mathfrak{p}^k}$ . Take  $L = \binom{x}{u} \binom{y}{v} \in \operatorname{SL}_2(\mathcal{A})$  with  $x = \frac{\mathfrak{m}\alpha'}{\gcd(\mathfrak{p}^k,\mathfrak{m}\alpha')} = \frac{\mathfrak{m}\alpha'}{d}$  and  $u = \frac{-\mathfrak{p}^k}{d}$ . Then we have  $L^{-1}\alpha = \binom{*}{0} \binom{*}{\mathfrak{m}/d}$ . And since  $\det(L^{-1}\alpha) = \det(L^{-1})\det(\alpha) = 1 \cdot \mathfrak{m} = \mathfrak{m}$ , we infer that  $L^{-1}\alpha = \binom{d}{0} \binom{\eta}{\mathfrak{m}/d}$ , where  $\eta$  is some element in A. Since  $\Delta(z)$  is a modular form of weight  $q^2 - 1$  under elements of  $\operatorname{SL}_2(A)$ , we get the following:

$$\Delta_{\mathfrak{m}}(A^{-1}z) = \Delta(\alpha z) = \Delta(LL^{-1}\alpha z)$$

$$= \left(u\left(\frac{dz+\eta}{\mathfrak{m}/d}+v\right)z+v\right)^{q^2-1}\Delta\left(\frac{dz+\eta}{\mathfrak{m}/d}\right) = \left(\frac{d}{\mathfrak{m}}(-\mathfrak{p}^kz+a)\right)^{q^2-1}\Delta\left(\frac{d^2}{\mathfrak{m}}z+\frac{\eta d}{\mathfrak{m}}\right).$$

### 3. The generalised Jacobian

Let  $J_0(\mathfrak{n})_{\mathbf{m}}$  be the generalised Jacobian of  $X_0(\mathfrak{n})$  with respect to the modulus  $\mathbf{m} := \sum_{P \in C} P$ , where  $C := \text{Cusps}(X_0(\mathfrak{n}))$ . That is, we have

$$J_0(\mathfrak{n})_{\mathbf{m}}(K) = \mathrm{Div}^0(X_0(\mathfrak{n}) \setminus C) / \{\mathrm{div}(f) \mid f \in K(X_0(\mathfrak{n}))^{\times}, f \equiv 1 \pmod{\mathbf{m}} \}.$$

Hence,  $J_0(\mathfrak{n})_{\mathbf{m}}$  fits into the exact sequence

$$(3) 0 \to L_{\mathbf{m}} \to J_0(\mathfrak{n})_{\mathbf{m}} \to J_0(\mathfrak{n}),$$

where  $L_{\mathbf{m}}$  is defined by the quotient

$$0 \to \mathbb{G}_m \to \prod_{i=0}^r \operatorname{Res}_{K(P_i)/K} \mathbb{G}_m \to L_{\mathbf{m}} \to 0.$$

After applying the exact functor  $H^1(-,K)$ , and the left-exact functor  $\text{Hom}(\mathbb{Q}/\mathbb{Z},-)$  to the sequence in Equation (3), we obtain the exact sequence

$$(4) 0 \to \bigoplus_{i=0}^{r-1} K(P_i)_{\text{tors}}^{\times} \to J_0(\mathfrak{n})_{\mathbf{m}}(K)_{\text{tors}} \to J_0(\mathfrak{n})(K)_{\text{tors}} \xrightarrow{\delta} \bigoplus_{i=0}^{r-1} K(P_i)_{\text{tors}} \otimes \mathbb{Q}/\mathbb{Z}.$$

Hence, to understand  $J_0(\mathfrak{n})_{\mathbf{m}}(K)_{\text{tors}}$ , we need to understand the kernel of the map  $\delta$  appearing in Equation (4). The key lemma to do so is Lemma 2.3.1 in [27]. We present their result here in the form that will be useful to us.

**Lemma 3.1** (cf. [27, Lemma 2.3.1]). Fix  $\mathfrak{n} = \mathfrak{p}^r$  with  $\mathfrak{p} \subseteq \mathcal{A}$  monic irreducible and r a positive integer. Let  $D = \sum_{i=0}^r a_i \cdot P_i \in \operatorname{Div}^0(X_0(\mathfrak{n}))$  be a degree-zero divisor supported on m such that  $[D] \in J_0(\mathfrak{n})(K)_{\operatorname{tors}}$ . Let  $n \in \mathbb{Z}_{>0}$  so that  $n \cdot [D] = 0$ , i.e., so that there exists  $f \in K(X_0(\mathfrak{n}))$  such that  $\operatorname{div}(f) = n \cdot D$ . We have

$$\delta([D]) = \left(\left(\frac{f}{t_{P_r}^{na_r}}\right)(P_r)\left(\frac{t_{P_i}^{na_i}}{f}\right)(P_i)\right)_{i=0}^{r-1} \otimes \frac{1}{n} \in \bigoplus_{i=0}^{r-1} K(P_i)^{\times} \otimes \mathbb{Q}/\mathbb{Z},$$

where  $t_{P_i}$  is a uniformiser at the closed point  $P_i \in X_0(\mathfrak{n})$ . Notice that this description does not depend on the choice of f nor on that of  $t_{P_i}$ .

Since the map  $\delta$  is a group homomorphism, we have the decomposition

$$(5) \qquad \ker(\delta|_{J_0(\mathfrak{n})(\mathbb{Q})_{\mathrm{tors}}}) = \bigoplus_{\ell \text{ prime}} \ker(\delta|_{J_0(\mathfrak{n})(\mathbb{Q})_{\mathrm{tors}}}) [\ell^{\infty}] = \bigoplus_{\ell \text{ prime}} \ker(\delta|_{J_0(\mathfrak{n})(\mathbb{Q})_{\mathrm{tors}}[\ell^{\infty}]}).$$

By Theorem 2.1, for all  $\ell \nmid q(q-1)$  we have

(6) 
$$\ker(\delta|_{J_0(\mathfrak{n})(\mathbb{Q})_{\text{tors}}[\ell^{\infty}]}) = \ker(\delta|_{C(\mathfrak{n})[\ell^{\infty}]}) = \ker(\delta|_{C(\mathfrak{n})})[\ell^{\infty}].$$

In other words, to compute  $\ker(\delta|_{J_0(N)(\mathbb{Q})_{\text{tors}}[\ell^{\infty}]})$ , it is enough to understand the map  $\delta$  restricted to the subgroup  $C(\mathfrak{n})$ . The next proposition takes a closer look at  $\delta$  when applied to elements of  $C(\mathfrak{n})$  when  $\mathfrak{n} = \mathfrak{p}^r$  is a prime power.

**Proposition 3.2.** Fix  $\mathfrak{n} = \mathfrak{p}^r$  with  $\mathfrak{p} \subseteq \mathcal{A}$  monic irreducible and r a positive integer. Let  $D \in \mathfrak{C}(\mathfrak{n})$  be a cuspidal divisor such that for some integer  $n(D) \in \mathbb{Z}_{>0}$  we have  $g(n(D) \cdot D) = \prod_{\mathfrak{m} \mid \mathfrak{n}} \Delta_{\mathfrak{m}}(z)^{k_{\mathfrak{m}}} \in K(X_0(\mathfrak{n}))$  for some  $k_{\mathfrak{m}} \in \mathbb{Z}$ . We have

$$\delta(D) = \left(\prod_{\mathfrak{m} \mid \mathfrak{n}} \left(\frac{\gcd(\mathfrak{p}^k, \mathfrak{m})}{\mathfrak{m}}\right)^{-(q^2-1)k_{\mathfrak{m}}} \prod_{\mathfrak{m} \mid \mathfrak{n}} e_L \left(\frac{\eta_{\mathfrak{m}} \gcd(\mathfrak{p}^k, \mathfrak{m})}{\mathfrak{m}}\right)^{-k_{\mathfrak{m}}}\right)^{r-1} \otimes \frac{1}{n(D)},$$

where  $\eta_{\mathfrak{m}} \in \mathcal{A}$  and  $e_L$  is the Carlitz exponential associated to a lattice L as defined in [6].

*Proof.* From the explicit expression in Lemma 3.1 we see that to prove the result we want to find a formula for  $\left(\frac{g(\mathfrak{n}(D)\cdot D)(z)}{t_{P_i}^{\mathfrak{n}(D)a_i}}\right)(P_i)$  for all closed points  $P_i$ . Since  $P_i$  is a Galois orbit, it is enough to consider one representative  $\omega\in \operatorname{Cusp}(X_0(\mathfrak{n})(\mathbb{C}_\infty))$  for each i. Take a cusp representative  $\omega=\left[\begin{smallmatrix}\mathfrak{a}'\\\mathfrak{m}'\end{smallmatrix}\right]\in\operatorname{Cusp}(X_0(\mathfrak{n})(\mathbb{C}_\infty))$  and a matrix  $A\in\operatorname{SL}_2(A)$  such that  $A^{-1}\cdot\omega_\infty=\left[\begin{smallmatrix}\mathfrak{a}'\\\mathfrak{m}'\end{smallmatrix}\right]$ . The Fourier expansion of  $\Delta_\mathfrak{m}(z)$  at  $\omega$  is the expansion of  $\Delta_\mathfrak{m}(A^{-1}z)$  at  $\omega_\infty$ . Take  $t_{\omega_\infty}\in K(X_0(\mathfrak{n}))^\times$  to be a uniformiser at  $\omega_\infty$  which has 1 as the leading term of its Fourier expansion. Let  $h(z):=\prod_{\mathfrak{m}\mid\mathfrak{n}}\Delta_\mathfrak{m}(z)^{k_\mathfrak{m}}\in\mathcal{E}_\mathfrak{n}\otimes_\mathbb{Z}\mathbb{Q}$  be any  $\Delta$ -quotient. Then we have

$$\left(\frac{h(z)}{t_{\omega_{\infty}}^{\operatorname{ord}_{\infty}(g)}}\right)(\omega_{\infty}) = 1.$$

Now take the uniformiser at  $[\omega]$  given by

$$t_{\omega} := t_{\omega_{\infty}} \circ A \in K(X_0(\mathfrak{n})).$$

The identity in Lemma 2.11 yields

$$\begin{split} &\left(\frac{h(z)}{t_{\omega}^{\operatorname{ord}_{\omega}(g)}}\right)(\omega) = \left(\frac{h(z) \circ A^{-1}}{t_{\omega_{\infty}}^{\operatorname{ord}_{\omega}(g)}}\right)(\omega_{\infty}) = \\ &\prod_{\mathfrak{m} \mid \mathfrak{n}} \left(\frac{\gcd(\mathfrak{p}^{k}, \mathfrak{m})}{\mathfrak{m}}\right)^{(q^{2}-1)k_{\mathfrak{m}}} \frac{\prod_{\mathfrak{m} \mid \mathfrak{n}} \Delta\left(\frac{\gcd(\mathfrak{p}^{k}, \mathfrak{m})^{2}}{\mathfrak{m}}z + \frac{\eta_{(\mathfrak{m}, \mathfrak{p}^{k})}\gcd(\mathfrak{p}^{k}, \mathfrak{m})}{\mathfrak{m}}\right)^{k_{\mathfrak{m}}}}{t_{\omega_{\infty}}^{\operatorname{ord}_{\omega}(g)}}(\omega_{\infty}); \end{split}$$


where  $\eta_{(\mathfrak{m},\mathfrak{p}^k)}$  is some element in  $\mathcal{A}$ . The product formula for the function  $\Delta$  found by Gekeler in [6, Theorem pg.137] finishes the proof.

**Remark 3.3.** Take the notation as in the statement of Proposition 3.2. Note that the Fourier expansion of  $\Delta$  at a cusp of height  $\mathfrak{m}=p^i$  has  $K(P_i)$ -rational coefficients, see [6]. For a fixed  $\mathfrak{m} \mid \mathfrak{n}$ , recall that  $\mathfrak{d}=\mathfrak{d}(\mathfrak{m})=\gcd(\mathfrak{m},\mathfrak{n}/\mathfrak{m})$ . Let  $\mathfrak{O}_{\mathfrak{d}}:=\langle \mathfrak{p}\rangle\subseteq (K(\mathfrak{d})^+)^\times$  be the subgroup of  $(K(\mathfrak{d})^+)^\times$  generated by  $\mathfrak{p}$ . The previous proposition shows that  $\operatorname{im}(\delta|_{\mathfrak{C}(\mathfrak{n})})\subseteq\bigoplus_{i=0}^{r-1}(\mathfrak{O}_{\mathfrak{d}}\cdot\mu(K(P_i)))\otimes\mathbb{Q}/\mathbb{Z}$ .

For the rest of the paper we fix  $\mathfrak{n} = \mathfrak{p}^r$  with  $\mathfrak{p} \subseteq \mathcal{A}$  monic irreducible and r a positive integer.

**Definition 3.4.** Consider the subgroup  $\Lambda_{\mathfrak{d}}$  given by the image of  $\mathfrak{O}_{\mathfrak{d}}$  in the quotient  $K(P_i)^{\times}/\mu(K(P_i))$ . We denote  $\Lambda := \bigoplus_{\mathfrak{m} \mid \mathfrak{n}} \Lambda_{\mathfrak{d}(\mathfrak{m})}$ .

By Remark 3.3 and since all torsion units  $x \in \mu(K(P_i))$  with  $n \mid \operatorname{ord}(x)$  satisfy  $x \otimes \frac{a}{b} = 1 \otimes \frac{a}{nb}$  and the latter is in turn the identity element in  $K(P_i) \otimes \mathbb{Q}/\mathbb{Z}$ , the map  $\delta$  factorises as



**Lemma 3.5.** The map  $\phi$  in the diagram of (7) is an injection.

*Proof.* For each  $\mathfrak{m}=p^i$  with  $\mathfrak{m}\mid \mathfrak{n}$ , let  $\mathfrak{d}=\mathfrak{d}(\mathfrak{m})=\gcd(\mathfrak{m},\mathfrak{n}/\mathfrak{m})$  as before. Consider the following tensoring map on  $\mathbb{Z}$ -modules:

$$\phi_{\mathfrak{d}(\mathfrak{m})}: \mathbb{Q}/\mathbb{Z} \longrightarrow K(P_i)^{\times} \otimes \mathbb{Q}/\mathbb{Z}, \quad x \mapsto \mathfrak{p} \otimes x.$$

If we prove that for each  $\mathfrak{d}$  the map  $\phi_{\mathfrak{d}}$  is a split injection, then the injectivity of  $\phi$  follows since  $\Lambda \otimes \mathbb{Q}/\mathbb{Z} \simeq \bigoplus_{\mathfrak{d}} (\Lambda_{\mathfrak{d}} \otimes \mathbb{Q}/\mathbb{Z})$  and  $(\bigoplus_i K(P_i^{\times})) \otimes \mathbb{Q}/\mathbb{Z} \simeq \bigoplus_i (K(P_i)^{\times} \otimes \mathbb{Q}/\mathbb{Z})$ . Since  $\mathbb{Q}/\mathbb{Z}$  is an injective module, splitting of  $\phi_{\mathfrak{d}}$  follows automatically if we prove its injectivity. Now the prime  $\mathfrak{p} \in K(P_i)$  is not torsion, thus its image  $\mathfrak{O}_{\mathfrak{d}}$  is a free  $\mathbb{Z}$ -module (note that  $\mathbb{F}_q$  is the only torsion part in  $K(P_i)^{\times}$ ). Hence, the  $\phi_{\mathfrak{d}}$  is injective for all  $\mathfrak{m}$ .

Remark 3.6. The factorisation given in Equation (7) together with Lemma 3.5 show that we have the equality  $\ker(\delta|_{C(\mathfrak{n})}) = \ker(\overline{\delta}|_{C(\mathfrak{n})})$ . Since the map  $\overline{\delta}$  no longer takes the torsion units into account, it is relatively easier to compute. In the next section we will explicitly compute the images of the generators in Theorem 2.8 under  $\overline{\delta}$ .

As a result of a technical study of  $\overline{\delta}$  we obtain the main result which we recall below.

**Theorem 3.7.** Fix  $\mathfrak{n} = \mathfrak{p}^r$  with  $\mathfrak{p} \subseteq \mathcal{A}$  monic irreducible and r a positive integer. The map  $\overline{\delta}|_{\mathfrak{C}(\mathfrak{n})}$  is an injection. In particular, for all odd primes  $\ell$  with  $\ell \nmid q(q^2-1)$  the subgroup  $J_0(\mathfrak{p}^r)_{\mathfrak{m}}(\mathbb{Q})[\ell^{\infty}]$  is trivial.

*Proof.* This is a consequence of Equations (6), Lemma 3.5 and Theorem 5.3 proven in Section 5, which states that  $\overline{\delta}|_{\mathcal{C}(\mathfrak{n})}$  is injective.

The goal of the coming two sections is to prove Theorem 5.3. In Section 4 we compute the image of the cuspidal divisors appearing in Theorem 2.8. This will allow us to find a matrix representation of the map  $\overline{\delta}|_{\mathfrak{C}(\mathfrak{p}^r)}$ . In Section 5 we will use this representation to determine the kernel of  $\overline{\delta}|_{\mathfrak{C}(\mathfrak{n})}$ .

# 4. The image of $\overline{\delta}$ .

Now we proceed with the computations of the map  $\overline{\delta}$ . The main goal of this section is to compute  $\operatorname{im}(\overline{\delta}|_{\mathcal{C}(\mathfrak{p}^r)})$  in Proposition 4.3.

Remark 4.1. Recall from Proposition 3.2 that the product  $\prod_{\mathfrak{m}\mid\mathfrak{n}}\left(\frac{\gcd(\mathfrak{p}^k,\mathfrak{m})}{\mathfrak{m}}\right)^{(q^2-1)k_{\mathfrak{m}}}\prod_{\mathfrak{m}\mid\mathfrak{n}}e_L\left(\frac{\eta_{\mathfrak{m}}\gcd(\mathfrak{p}^k,\mathfrak{m})}{\mathfrak{m}}\right)^{k_{\mathfrak{m}}}$  appears in the expression of  $\delta(\overline{D})$  for  $D\in \mathcal{C}(\mathfrak{n})$ . From [3, Theorem 8.4] we have that the factor of the product  $\prod_{\mathfrak{m}\mid\mathfrak{n}}e_L\left(\frac{\eta_{\mathfrak{m}}\gcd(\mathfrak{p}^k,\mathfrak{m})}{\mathfrak{m}}\right)^{k_{\mathfrak{m}}}\in K(P_i)$  is a torsion unit. Hence, by Lemma 3.5, in order to show that the map  $\overline{\delta}$  in Equation (7) is injective we will only need to control the product  $\prod_{\mathfrak{m}\mid\mathfrak{n}}\left(\frac{\gcd(\mathfrak{p}^k,\mathfrak{m})}{\mathfrak{m}}\right)^{(q^2-1)k_{\mathfrak{m}}}$  in the product expression.

The product expression appearing Proposition 3.2 together with Remark 4.1 is used in the following for the computation of the images of the divisors  $\{C_i\}_{i=1}^{\left\lfloor\frac{r-1}{r}\right\rfloor}$ ,  $\{C_i-C_{i-1}\}_{i=\left\lfloor\frac{r+1}{2}\right\rfloor}^{r-2}$  and  $\{D_i\}_{i=0,r-1}$  under the map  $\overline{\delta}$ .

Letting  $r \ge 2$ , by Theorem 2.8 we have

$$\mathcal{C}(\mathfrak{p}^r) = \left(\bigoplus_{i=1}^m \langle \overline{C_i} \rangle\right) \oplus \left(\bigoplus_{i=m+1}^{r-2} \langle \overline{C_i - |\mathfrak{p}|C_{i+1}} \rangle\right) \oplus \langle \overline{D_{r-1}} \rangle \oplus \langle \overline{D_0} \rangle,$$

where  $m := \lfloor \frac{r-1}{2} \rfloor$ .

**Notation 4.2.** Let  $\sigma \in \Lambda$  a tuple of the form

$$\sigma = (\mathfrak{p}^{\sigma_k})_{k=0}^{r-1};$$

where each for each k, the element  $\mathfrak{p}^{\sigma_k}$  lies in  $\Lambda_{\mathfrak{d}}$  with  $\mathfrak{d} = \mathfrak{d}(\mathfrak{p}^k)$ .

**Proposition 4.3.** The images of  $\overline{C_i}$  for  $i \in \{1, ..., \lfloor \frac{r-1}{2} \rfloor\}$  under the map  $\overline{\delta}$  are given by

(8) 
$$\overline{\delta}(\overline{C_i}) = (\mathfrak{p}^{-\sigma(i)_k})_{k=0}^{r-1} \otimes \frac{q+1}{|\mathfrak{p}|^{r-i}(|\mathfrak{p}|^2-1)},$$

where

(9) 
$$\sigma(i)_{k} = \begin{cases} (r-i)(|\mathfrak{p}|^{2}-2|\mathfrak{p}|+1)+|\mathfrak{p}|-1 & \text{if } 0 \leq k \leq i-1; \\ (r-i)(|\mathfrak{p}|^{2}-2|\mathfrak{p}|+1)+2|\mathfrak{p}|-1 & \text{if } k=i; \\ (r-k)(|\mathfrak{p}|^{2}-2|\mathfrak{p}|+1)+|\mathfrak{p}|-1 & \text{if } i+1 \leq k \leq r-1. \end{cases}$$

Similarly, for  $i \in \{\lfloor \frac{r+1}{2} \rfloor, \dots, r-2 \}$  we have

(10) 
$$\overline{\delta}(\overline{C_i - |\mathfrak{p}|C_{i+1}}) = (\mathfrak{p}^{-\sigma(i)_k})_{k=0}^{r-1} \otimes \frac{q+1}{|\mathfrak{p}|^i(|\mathfrak{p}|^2-1)},$$

where

(11) 
$$\sigma(i)_{k} = \begin{cases} (|\mathfrak{p}|-1)^{2} & \text{if } 0 \leq k \leq i-1; \\ |\mathfrak{p}|^{2}-|\mathfrak{p}|+1 & \text{if } k=i; \\ -|\mathfrak{p}| & \text{if } k=i+1; \\ 0 & \text{if } i+2 \leq k \leq r-1. \end{cases}$$

Finally, we have

(12) 
$$\overline{\delta}(\overline{D_0}) = (\mathfrak{p}^{-(q-1)\sigma(0)_k})_{k=0}^{r-1} \otimes \frac{q+1}{|\mathfrak{p}|-1}, \quad \text{where} \quad \sigma(0)_k = 1$$

for all  $0 \le k \le r - 1$ , and

(13) 
$$\overline{\delta}(\overline{D_{r-1}}) = (\mathfrak{p}^{-\sigma(r-1)_k})_{k=1}^{r-1} \otimes \frac{q+1}{|\mathfrak{p}|^2 - 1},$$

where the exponents  $\sigma(r-1)_k$  can be found in Appendix 6 – see Remark 6.2.

*Proof.* We first determine  $\overline{\delta}(\overline{C_i})$  for  $i \in \{1, \dots, \lfloor \frac{r-1}{2} \rfloor\}$ . Let  $m = \min\{i, r-i\}$ . Then [12, Theorem 3.6] states that

$$g(C_i) = \left(\frac{\Delta_{\mathfrak{p}^i}^{|\mathfrak{p}|^2+1} \Delta_{\mathfrak{p}^{r-1}}^{|\mathfrak{p}|-1} \Delta_{\mathfrak{p}^r}^{|\mathfrak{p}|}}{\Delta_{\mathfrak{p}^{i-1}}^{|\mathfrak{p}|} \Delta_{\mathfrak{p}^{i+1}}^{|\mathfrak{p}|} \Delta_{\mathfrak{p}^r}^{|\mathfrak{p}|}}\right)^{\frac{1}{|\mathfrak{p}|^r - m(|\mathfrak{p}|^2 - 1)(q - 1)}}.$$

Since by Theorem 2.8 we have  $\operatorname{ord}(\overline{C_i}) = |\mathfrak{p}|^{r-i} M(\mathfrak{p})$ , using Proposition 3.2 we have

$$\overline{\delta}(\overline{C_i}) = (\mathfrak{p}^{-\frac{\sigma(i)_k}{q-1}})_{k=0}^r \otimes \frac{q^2 - 1}{|\mathfrak{p}|^{r-i}(|\mathfrak{p}|^2 - 1)} = (\mathfrak{p}^{-\sigma(i)_k})_{k=0}^r \otimes \frac{q+1}{|\mathfrak{p}|^{r-i}(|\mathfrak{p}|^2 - 1)}$$

with  $\sigma(i)_k$  as in Equation (9).

On the other hand, for  $i = \lfloor \frac{r+1}{2} \rfloor, \dots, r-2$  and  $m = \min\{i, r-i\}$ , [12, Lemma 3.8] computes

$$g(C_i - |\mathfrak{p}|C_{i+1}) = \left(\frac{\Delta_{\mathfrak{p}^i}^{|\mathfrak{p}|^2 + |\mathfrak{p}| + 1} \Delta_{\mathfrak{p}^{i+2}}^{|\mathfrak{p}|}}{\Delta_{\mathfrak{p}^{i-1}}^{|\mathfrak{p}|} \Delta_{\mathfrak{p}^{i+1}}^{|\mathfrak{p}|^2 + |\mathfrak{p}| + 1}}\right)^{\frac{1}{|\mathfrak{p}|^i(|\mathfrak{p}|^2 - 1)(q - 1)}}.$$

Since by Theorem 2.8 we have  $\operatorname{ord}(\overline{C_i - |\mathfrak{p}|C_{i+1}}) = |\mathfrak{p}|^i M(\mathfrak{p})$ , using Proposition 3.2 we obtain

$$\overline{\delta}(\overline{C_i - |\mathfrak{p}|C_{i+1}}) = (\mathfrak{p}^{-\frac{\sigma(i)_k}{q-1}})_{k=0}^r \otimes \frac{q^2-1}{|\mathfrak{p}|^i(|\mathfrak{p}|^2-1)} = (\mathfrak{p}^{-\sigma(i)_k})_{k=0}^r \otimes \frac{q+1}{|\mathfrak{p}|^i(|\mathfrak{p}|^2-1)}$$

with  $\sigma(i)_k$  as in Equation (11).

We are left with computing  $\overline{\delta}(\overline{D_i})$  for i=0,r-1. We start with i=0. By [12, Lemma 3.9], we have

$$g(D_0) = \left(\frac{\Delta_{\mathfrak{p}^{r-1}}}{\Delta_{\mathfrak{n}^r}}\right)^{\frac{1}{|\mathfrak{p}|-1}},$$

and Theorem 2.8 gives  $\operatorname{ord}(\overline{D_0}) = N(\mathfrak{p})$ . Hence, Proposition 3.2 yields Equation (12). Finally for i = r - 1 we have a case distinction for the definition of  $D_{r-1}$  – see [12, Section 3.2]. The modular units  $g(D_{r-1})$  are given in [12, Lemma 3.8] for all different cases. Using this lemma, if  $\sigma(r-1)_k$  denotes the exponent at  $\mathfrak{p}$  of the leading coefficient of the q-expansion of the modular unit  $g(M(\mathfrak{p}) \cdot D_{r-1})$  at the cusp  $\frac{1}{\mathfrak{p}^k}$  for  $0 \le k \le r-1$ , then we have

$$\overline{\delta}(\overline{D_{r-1}}) = (\mathfrak{p}^{-\sigma(r-1)_k})_{k=0}^{r-1} \otimes \frac{q+1}{|\mathfrak{p}|^2-1}.$$

For convenience of the reader, the computation of the entries  $\sigma(r-1)_k$  for  $0 \le k \le r-1$  can be found in the Appendix 6.

# 5. Injectivity of the map $\delta$ .

In this section, we show that the map  $\overline{\delta}$  is injective to complete the proof of Theorem 1.3. Using Proposition 4.3, we will first find a matrix representing the map  $\overline{\delta}$ . To do so, we introduce the following notation.

**Notation 5.1.** Recall that for a divisor  $D \in \{C_i : 1 \le i \le \lfloor \frac{r-1}{2} \rfloor\} \cup \{C_i - |\mathfrak{p}|C_{i+1} : \lfloor \frac{r+1}{2} \rfloor \le i \le r-2\} \cup \{D_{r-1}\},$  we denote by  $\sigma(i)_k$  the exponent of  $\mathfrak{p}$  at the k-th coordinate of the tuple of element in the tensor product  $\bigoplus_{k=0}^{r-1} K(P_k)^{\times} \otimes \mathbb{Q}/\mathbb{Z}$  of the representatives  $\overline{\delta}(\overline{D})$  given in Equations (9), (11) and (13).

Notice that, since we can take  $\frac{q+1}{|\mathfrak{p}|^r(|\mathfrak{p}|^2-1)} \in \mathbb{Q}/\mathbb{Z}$  as common factor of the images of the divisors  $\{C_i\}_{i=1}^{\lfloor \frac{r-1}{2} \rfloor}$ ,  $\{C_i-|\mathfrak{p}|C_{i+1}\}_{i=\lfloor \frac{r+1}{2} \rfloor}^{r-2}$  and  $D_{r-1}$  computed in Proposition 4.3, the  $r \times r$  matrix  $(M_{\delta}(i,j))_{i,j}$  given by

$$\mathbf{M}_{\delta}(i,j) := -1 \cdot \begin{cases} |\mathfrak{p}|^r (q-1)\sigma(0)_j & \text{for } i=0, \ j=0,\dots,r-1; \\ |\mathfrak{p}|^i \sigma(i)_j & \text{for } i=1,\dots,\lfloor\frac{r-1}{2}\rfloor, \ j=0,\dots,r-1; \\ \sigma(i)_j & \text{for } i=\lfloor\frac{r+1}{2}\rfloor,\dots,r-1, \ j=0,\dots,r-1; \\ |\mathfrak{p}|^r \sigma(i)_j & \text{for } i=r-1, \ j=0,\dots,r-1; \end{cases}$$

represents the map  $\overline{\delta}$  defined in Equation (7). Notice that the entries of  $\mathbf{M}_{\delta}$  are labelled from 0 to r-1 instead of the ususal 1 to r. To finish the proof of Theorem 1.3, we need to check the injectivity of  $\overline{\delta}$ . That is, we want to show that  $\det(\mathbf{M}_{\delta}) \neq 0$ .

**Notation 5.2.** For a matrix  $M = (m_{i,j})_{0 \le i \le n-1, 0 \le j \le m-1}$  of dimension  $n \times m$ , we use the following notation throughout the section.

· For  $0 \le i \le n-1$ , we denote by M[i] the i-th row of M.

- · For  $0 \le j \le m-1$ , we denote by M[[j]] the j-th column of M.
- · For  $0 \le i \le n-1$  and  $0 \le j \le m-1$ , we denote by M(i,j) the (i,j)-th entry of M.
- · For a given i, we represent the row M[i] of M as  $M[i] = (*_0, ..., *_{m-1})$ , where  $*_j$  is the entry M(i, j). We will write the subindex j only when necessary to clarify the position of the entries of a given tuple.

# **Theorem 5.3.** The matrix $\mathbf{M}_{\delta}$ representing the map $\delta$ has maximal rank.

We will take the rest of the section to prove Theorem 5.3. To do so, ill use the following auxiliary matrix. Let

$$M_{\delta}(i,j) := \begin{cases} \sigma(0)_{j} & \text{for } i = 0, \ j = 0, \dots, r-1; \\ \sigma(i)_{j} & \text{for } i = 1, \dots, \lfloor \frac{r-1}{2} \rfloor, \ j = 0, \dots, r-1; \\ \sigma(i)_{j} & \text{for } i = \lfloor \frac{r+1}{2} \rfloor, \dots, r-1, \ j = 0, \dots, r-1; \\ \sigma(r-1)_{j} & \text{for } i = r-1, \ j = 0, \dots, r-1. \end{cases}$$

The matrices  $\mathbf{M}_{\delta}$  and  $M_{\delta}$  differ only from each other by rescaling the rows. Hence, to check that  $\mathbf{M}_{\delta}$  has non-zero determinant, it is enough to check that  $\det(M_{\delta}) \neq 0$ . If  $r \leq 6$ , showing  $\det(M_{\delta}) \neq 0$  can be done by a straightforward computation. However, to prove the result in more generality for  $r \geq 7$ , we will first transform the matrix  $M_{\delta}$  into a matrix  $M_{\delta}^h$  with the same determinant that is easier to manipulate. We will do that in two steps: we will first do a row reduction to produce many zeroes above the main diagonal of the matrix, and then we will do a column reduction to create many zeroes in the first column. The combination of these steps will be enough to allow us to prove that  $\det(M_{\delta}) \neq 0$ .

**Step 1:** Let  $r \ge 7$ . Let  $M_{\delta}^H$  be the matrix obtained from  $M_{\delta}$  after applying the following <u>row</u> transformations:

```
 \begin{array}{l} \cdot \ M_{\delta}^{H}[1] = M_{\delta}[1] - M_{\delta}[2], \ \text{for} \ i = 1; \\ \cdot \ M_{\delta}^{H}[i] = 2M_{\delta}[i] - M_{\delta}[i-1] - M_{\delta}[i+1], \ \text{for} \ 1 < i \leq \lfloor \frac{r-1}{2} \rfloor - 1; \\ \cdot \ M_{\delta}^{H}[\lfloor \frac{r-1}{2} \rfloor] = 2M_{\delta}[\lfloor \frac{r-1}{2} \rfloor] + \sum_{k=1}^{\lfloor r-3/2 \rfloor} (p^{k}-1)M_{\delta}[r-1-k] - M_{\delta}[(r-3)/2]; \\ \cdot \ M_{\delta}^{H}[\lfloor \frac{r+1}{2} \rfloor] = M_{\delta}[\lfloor \frac{r+1}{2} \rfloor] - \Big(M_{\delta}[\lfloor \frac{r-1}{2} \rfloor] + \sum_{k=1}^{\lfloor r-3/2 \rfloor} (p^{k}-1)M_{\delta}[r-1-k] \Big); \\ \cdot \ M_{\delta}^{H}[i] = M_{\delta}[i] - M_{\delta}[i-1], \ \text{for} \ \lfloor \frac{r+1}{2} \rfloor < i \leq r-1. \end{array}
```

**Claim 5.4.** The matrix  $M_{\delta}^{H}$  is a matrix of the form

where  $\alpha = |\mathfrak{p}|^{\lfloor \frac{r+1}{2} \rfloor} - |\mathfrak{p}|^{\lfloor \frac{r-1}{2} \rfloor}$  and  $\beta = |\mathfrak{p}|^{\lfloor \frac{r+1}{2} \rfloor} - |\mathfrak{p}|^{\lfloor \frac{r-1}{2} \rfloor} + |\mathfrak{p}|^2 - 2|\mathfrak{p}| + 1$ . More formally, we have

$$MAR \ CURCO \ IRANZO$$
 
$$if \ i = 0;$$
 
$$((|\mathfrak{p}|-1)^2, |\mathfrak{p}|^2 - |\mathfrak{p}|+1, -|\mathfrak{p}|, 0, \dots, 0) \qquad if \ i = 1;$$
 
$$(0, \dots, 0, -|\mathfrak{p}|_{i-1}, (|\mathfrak{p}|^2 + 1)_i, -|\mathfrak{p}|_{i+1}, 0, \dots, 0) \qquad if \ 1 < i < \lfloor \frac{r-1}{2} \rfloor;$$
 
$$(\alpha, \dots, \alpha, (\alpha + |\mathfrak{p}|)_{\lfloor \frac{r-3}{2} \rfloor}, \beta_{\lfloor \frac{r-1}{2} \rfloor}, |\mathfrak{p}| \left( |\mathfrak{p}|^{\lfloor \frac{r-1}{2} \rfloor} - 1 \right)_{\lfloor \frac{r+1}{2} \rfloor}, 0, \dots, 0) \qquad if \ i = \lfloor \frac{r-1}{2} \rfloor;$$
 
$$(-\alpha, \dots, -\alpha, (-\alpha - |\mathfrak{p}|)_{\lfloor \frac{r-1}{2} \rfloor}, (-|\mathfrak{p}|^{\lfloor \frac{r+1}{2} \rfloor} + |\mathfrak{p}|^2 + 1)_{\lfloor \frac{r+1}{2} \rfloor}, -|\mathfrak{p}|_{\lfloor \frac{r+3}{2} \rfloor}, 0, \dots, 0) \qquad if \ i = \lfloor \frac{r+1}{2} \rfloor;$$
 
$$(0, \dots, 0, -|\mathfrak{p}|_{i-1}, (|\mathfrak{p}|^2 + 1)_i, -|\mathfrak{p}|_{i+1}, 0, \dots, 0) \qquad if \ i = r-1.$$

*Proof.* Notice that, for  $1 < i \le \lfloor \frac{r-1}{2} \rfloor - 1$ , we can rewrite the transformations as follows:

$$M_{\delta}^{H}[i] = (M_{\delta}[i] - M_{\delta}[i+1]) - (M_{\delta}[i-1] - M[i]).$$

Hence, it is enough to look at the differences  $M_{\delta}[i] - M_{\delta}[i+1]$  for  $1 \le i \le \lfloor \frac{r-1}{2} - 1 \rfloor$ . From Equation (9) it follows that

$$M_{\delta}[i] - M_{\delta}[i+1] = ((|\mathfrak{p}|^2 - 2|\mathfrak{p}| + 1), \dots, (|\mathfrak{p}|^2 - 2|\mathfrak{p}| + 1)_{i-1}, (|\mathfrak{p}|^2 - |\mathfrak{p}| + 1)_i, -|\mathfrak{p}|_{i+1}, 0, \dots, 0),$$

for  $1 \le i \le \lfloor \frac{r-3}{2} \rfloor$ . Hence, we have

$$M_{\delta}^{H}[1] = ((|\mathfrak{p}|-1)^{2}, |\mathfrak{p}|^{2} - |\mathfrak{p}|+1, -|\mathfrak{p}|, 0, \dots, 0)$$

for i = 1 and

$$(M_{\delta}[i] - M_{\delta}[i+1]) - (M_{\delta}[i-1] - M[i]) = (0, \dots, 0, -|\mathfrak{p}|_{i-1}, (|\mathfrak{p}|^2 + 1)_i, -|\mathfrak{p}|_{i+1}, 0, \dots, 0)$$

for  $1 < i < \lfloor \frac{r-3}{2} \rfloor$ . Similarly, from Equation (11) it follows that

$$M_{\delta}^{H}[i] = M_{\delta}[i] - M_{\delta}[i-1] = (0, \dots, 0, -|\mathfrak{p}|_{i-1}, (|\mathfrak{p}|^{2}+1)_{i}, -|\mathfrak{p}|_{i+1}, 0, \dots, 0),$$

for  $\lfloor \frac{r+1}{2} \rfloor < i \le r-1$ . Hence, we are left with computing  $M_{\delta}^H[\lfloor \frac{r-1}{2} \rfloor]$  and  $M_{\delta}^H[\lfloor \frac{r+1}{2} \rfloor]$ . Recall from **Step 1** that

$$M_{\delta}^{H}[\lfloor \frac{r-1}{2} \rfloor] = 2M_{\delta}[\lfloor \frac{r-1}{2} \rfloor] + \sum_{k=1}^{\lfloor r-3/2 \rfloor} (p^{k}-1)M_{\delta}[r-1-k] - M_{\delta}[(r-3)/2] = 2M_{\delta}[\lfloor \frac{r-1}{2} \rfloor]$$

$$\left(M_{\delta}[\lfloor\frac{r-1}{2}\rfloor] + \sum_{k=1}^{\lfloor r-3/2\rfloor}(p^k-1)M_{\delta}[r-1-k]\right) - \left(M_{\delta}[(r-3)/2] - M_{\delta}[\lfloor\frac{r-1}{2}\rfloor]\right)$$

and

$$M_{\delta}^{H}[\lfloor \frac{r+1}{2} \rfloor] = M_{\delta}[\lfloor \frac{r+1}{2} \rfloor] - \left(M_{\delta}[\lfloor \frac{r-1}{2} \rfloor] + \sum_{k=1}^{\lfloor r-3/2 \rfloor} (p^{k}-1)M_{\delta}[r-1-k]\right).$$

Hence if we show that

$$(14) M_{\delta}[\lfloor \frac{r-1}{2} \rfloor] + \sum_{k=1}^{\lfloor r-3/2 \rfloor} (p^{k}-1) M_{\delta}[r-1-k] = (\beta, \dots, \beta, (\beta+|\mathfrak{p}|)_{\lfloor \frac{r-1}{2} \rfloor}, |\mathfrak{p}| (|\mathfrak{p}|^{\lfloor \frac{r-1}{2} \rfloor}-1)_{\lfloor \frac{r+1}{2} \rfloor}, 0, \dots, 0)$$

then the equalities

$$M_{\delta}^{H}[\lfloor \frac{r-1}{2} \rfloor] = (\alpha, \dots, \alpha, (\alpha + |\mathfrak{p}|)_{\lfloor \frac{r-3}{2} \rfloor}, (\beta + 2|\mathfrak{p}|)_{\lfloor \frac{r-1}{2} \rfloor}, |\mathfrak{p}| (|\mathfrak{p}|^{\lfloor \frac{r-1}{2} \rfloor} - 1)_{\lfloor \frac{r+1}{2} \rfloor}, 0, \dots, 0)$$

and

$$M_{\delta}^{H}\left[\left\lfloor \frac{r+1}{2}\right\rfloor\right] = (-\alpha, \ldots, -\alpha, (-\alpha - |\mathfrak{p}|)_{\lfloor \frac{r-1}{2}\rfloor}, (-|\mathfrak{p}|^{\lfloor \frac{r+1}{2}\rfloor} + |\mathfrak{p}|^{2} + 1)_{\lfloor \frac{r+1}{2}\rfloor}, -|\mathfrak{p}|_{\lfloor \frac{r+3}{2}\rfloor}, 0, \ldots, 0)$$

follow and finish the proof of the claim. Thus, to complete the proof we only need to verify Equation (14). Fix  $j \geq \lfloor \frac{r+3}{2} \rfloor$ . For all  $\lfloor \frac{r+1}{2} \rfloor \leq i \leq r-2$  we have  $M_{\delta}[i][j] = (|\mathfrak{p}|-1)^2$  if  $i \geq j+1$ ,  $M_{\delta}[i][j] = |\mathfrak{p}|^2 - |\mathfrak{p}|+1$  if i=j,  $M_{\delta}[i][j] = -|\mathfrak{p}|$  if i = j-1 and  $M_{\delta}[i][j] = 0$  if  $i \leq j-2$ . Hence,

$$\left( M_{\delta} \left[ \left\lfloor \frac{r-1}{2} \right\rfloor \right] + \sum_{k=1}^{\lfloor r-3/2 \rfloor} (p^{k}-1) M_{\delta} \left[ r-1-k \right] \right)_{j} =$$

$$(r-j)(|\mathfrak{p}|^{2}-2|\mathfrak{p}|+1) + |\mathfrak{p}|-1 - |\mathfrak{p}|(|\mathfrak{p}|^{r-j}-1) + (|\mathfrak{p}|^{2}-|\mathfrak{p}|+1)(|\mathfrak{p}|^{r-j-1}-1) +$$

$$(|\mathfrak{p}|-1)^{2} \sum_{k=1}^{r-2-j} (|\mathfrak{p}|^{r-1-j-k}-1) =$$

$$(r-j)(|\mathfrak{p}|^{2}-2|\mathfrak{p}|+1) + |\mathfrak{p}|-1 - |\mathfrak{p}|(|\mathfrak{p}|^{r-j}-1) + (|\mathfrak{p}|^{2}-|\mathfrak{p}|+1)(|\mathfrak{p}|^{r-j-1}-1) +$$

$$(|\mathfrak{p}|-1)^{2}|\mathfrak{p}|^{r-1-j} \sum_{k=1}^{r-2-j} |\mathfrak{p}|^{-k} - (|\mathfrak{p}|-1)^{2}(r-2-j) =$$

$$(r-j)(|\mathfrak{p}|^{2}-2|\mathfrak{p}|+1) + |\mathfrak{p}|-1 - |\mathfrak{p}|(|\mathfrak{p}|^{r-j}-1) + (|\mathfrak{p}|^{2}-|\mathfrak{p}|+1)(|\mathfrak{p}|^{r-j-1}-1) +$$

$$(|\mathfrak{p}|-1)(|\mathfrak{p}|^{r-1-j}-|\mathfrak{p}|) - (|\mathfrak{p}|-1)^{2}(r-2-j) = 0.$$

Moreover, for  $j \in \{\lfloor \frac{r+1}{2} \rfloor, \lfloor \frac{r-1}{2} \rfloor\}$  we have

$$\left( M_{\delta} \left[ \left\lfloor \frac{r-1}{2} \right\rfloor \right] + \sum_{k=1}^{\lfloor r-3/2 \rfloor} (p^{k}-1) M_{\delta} \left[ r-1-k \right] \right)_{\left\lfloor \frac{r+1}{2} \right\rfloor} =$$

$$(r-\lfloor \frac{r+1}{2} \rfloor) (|\mathfrak{p}|^{2}-2|\mathfrak{p}|+1) + |\mathfrak{p}|-1 + (|\mathfrak{p}|^{2}-|\mathfrak{p}|+1) (|\mathfrak{p}|^{r-\lfloor \frac{r+1}{2} \rfloor-1}-1) +$$

$$(|\mathfrak{p}|-1)^{2} \sum_{k=1}^{r-1-j} (|\mathfrak{p}|^{r-1-j-k}-1) =$$

$$(r-\lfloor \frac{r+1}{2} \rfloor) (|\mathfrak{p}|^{2}-2|\mathfrak{p}|+1) + |\mathfrak{p}|-1 + (|\mathfrak{p}|^{2}-|\mathfrak{p}|+1) (|\mathfrak{p}|^{r-\lfloor \frac{r+1}{2} \rfloor-1}-1) +$$

$$(|\mathfrak{p}|-1) (|\mathfrak{p}|^{r-\lfloor \frac{r+1}{2} \rfloor-1}-1) - (p-1)^{2} (r-1-j) =$$

$$p(p^{r-\lfloor \frac{r+1}{2} \rfloor}-1) = p(p^{\lfloor \frac{r-1}{2} \rfloor}-1);$$

and

$$\left(M_{\delta}\left[\lfloor\frac{r-1}{2}\rfloor\right] + \sum_{k=1}^{\lfloor r-3/2\rfloor} (p^k-1)M_{\delta}[r-1-k]\right)_{\lfloor\frac{r-1}{2}\rfloor} =$$

$$\lfloor \frac{r+1}{2} \rfloor (|\mathfrak{p}|^2 - |\mathfrak{p}| + 1) + |\mathfrak{p}| - 1 + (|\mathfrak{p}| - 1)^2 \sum_{k=1}^{\lfloor \frac{r-3}{2} \rfloor} (p^k - 1) = 1 - |\mathfrak{p}| + |\mathfrak{p}|^2 - |\mathfrak{p}|^{\lfloor \frac{r-1}{2} \rfloor} + |\mathfrak{p}|^{\lfloor \frac{r+1}{2} \rfloor}.$$

Finally, for  $0 \le j < \lfloor \frac{r-1}{2} \rfloor$  we have

$$\left(M_{\delta}[\lfloor\frac{r-1}{2}\rfloor] + \sum_{k=1}^{\lfloor r-3/2\rfloor}(p^k-1)M_{\delta}[r-1-k]\right)_{\lfloor\frac{r+1}{2}\rfloor} = \\ \lfloor\frac{r+1}{2}\rfloor(|\mathfrak{p}|^2-2|\mathfrak{p}|+1) + |\mathfrak{p}|-1 + (|\mathfrak{p}|-1)^2\sum_{k=1}^{\lfloor\frac{r-3}{2}\rfloor}(|\mathfrak{p}|^k-1) = 1 - 2|\mathfrak{p}| + |\mathfrak{p}|^2 - |\mathfrak{p}|^{\lfloor\frac{r-1}{2}\rfloor} + |\mathfrak{p}|^{\lfloor\frac{r+1}{2}\rfloor}.$$

This completes the proof of the claim.

**Step 2:** Let  $r \ge 7$ . Let  $M_{\delta}^h$  be the matrix obtained after performing the <u>column</u> transformation to  $M_{\delta}^H$ :  $\cdot\ M^h_{\delta}[[1]] = M^H_{\delta}[[1]] - M^H_{\delta}[[2]].$ 

**Claim 5.5.** The matrix  $M_{\delta}^{h}$  is a matrix of the form

 $\textit{where } \alpha = |\mathfrak{p}|^{\lfloor \frac{r+1}{2} \rfloor} - |\mathfrak{p}|^{\lfloor \frac{r-1}{2} \rfloor}, \ \beta = |\mathfrak{p}|^{\lfloor \frac{r+1}{2} \rfloor} - |\mathfrak{p}|^{\lfloor \frac{r-1}{2} \rfloor} + |\mathfrak{p}|^2 - 2|\mathfrak{p}| + 1 \ \textit{and} \ \sigma(r-1)_0 - \sigma(r-1)_1 = \pm 1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p}|) \ \textit{for all } \beta = -1 + |\mathfrak{p}| g_1(|\mathfrak{p$ some  $g_1 \in \mathbb{Z}[x]$ . More formally, we have

$$M_{\delta}^{H}[i] = \begin{cases} (0,1,\ldots,1) & \text{if } i=0;\\ (-|\mathfrak{p}|,|\mathfrak{p}|^{2}-|\mathfrak{p}|+1,-|\mathfrak{p}|,0,\ldots,0) & \text{if } i=1;\\ (|\mathfrak{p}|,-|\mathfrak{p}|,|\mathfrak{p}|^{2}+1,-|\mathfrak{p}|,0,\ldots,0) & \text{if } i=2;\\ (0,\alpha,\ldots,\alpha,(\alpha+|\mathfrak{p}|)_{\lfloor\frac{r-3}{2}\rfloor},(\beta+2|\mathfrak{p}|)_{\lfloor\frac{r-1}{2}\rfloor},|\mathfrak{p}|\left(|\mathfrak{p}|^{\lfloor\frac{r-1}{2}\rfloor}-1\right)_{\lfloor\frac{r+1}{2}\rfloor},0,\ldots,0) & \text{if } i=\lfloor\frac{r-1}{2}\rfloor;\\ (0,\alpha,\ldots,\alpha,(\alpha+|\mathfrak{p}|)_{\lfloor\frac{r-2}{2}\rfloor},(\beta+2|\mathfrak{p}|)_{\lfloor\frac{r-1}{2}\rfloor},|\mathfrak{p}|\left(|\mathfrak{p}|^{\lfloor\frac{r-1}{2}\rfloor}-1\right)_{\lfloor\frac{r+1}{2}\rfloor},0,\ldots,0) & \text{if } i=\lfloor\frac{r-1}{2}\rfloor;\\ (0,-\alpha,\ldots,-\alpha,(-\alpha-|\mathfrak{p}|)_{\lfloor\frac{r-1}{2}\rfloor},(-|\mathfrak{p}|^{\lfloor\frac{r+1}{2}\rfloor}+|\mathfrak{p}|^{2}+1)_{\lfloor\frac{r+1}{2}\rfloor},-|\mathfrak{p}|_{\lfloor\frac{r+3}{2}\rfloor},0,\ldots,0) & \text{if } i=\lfloor\frac{r+1}{2}\rfloor;\\ (0,\ldots,0,-|\mathfrak{p}|_{i-1},(|\mathfrak{p}|^{2}+1)_{i},-|\mathfrak{p}|_{i+1},0,\ldots,0) & \text{if } i=r-1. \end{cases}$$
 Proof. The proof follows from Claim 5.4 after a direct computation.

*Proof.* The proof follows from Claim 5.4 after a direct computation.

Remark 5.6. Notice that since the determinant of a matrix is invariant under row and column transformations, we have  $\det(M_{\delta}) = \det(M_{\delta}^H) = \det(M_{\delta}^h)$ . Hence, verifying  $\det(M_{\delta}) \neq 0$  is equivalent to proving  $\det(M_{\delta}^h) \neq 0$ .

**Example 5.7.** Take r = 7. The matrix  $M_{\delta}$  is given by

$$M_{\delta} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 6|\mathfrak{p}|^2 - 11|\mathfrak{p}| + 5 & 6|\mathfrak{p}|^2 - 10|\mathfrak{p}| + 5 & 5|\mathfrak{p}|^2 - 9|\mathfrak{p}| + 4 & 4|\mathfrak{p}|^2 - 7|\mathfrak{p}| + 3 & 3|\mathfrak{p}|^2 - 5|\mathfrak{p}| + 2 & 2|\mathfrak{p}|^2 - 3|\mathfrak{p}| + 1 & (|\mathfrak{p}| - 1)|\mathfrak{p}| \\ 5|\mathfrak{p}|^2 - 9|\mathfrak{p}| + 4 & 5|\mathfrak{p}|^2 - 9|\mathfrak{p}| + 4 & 5|\mathfrak{p}|^2 - 8|\mathfrak{p}| + 4 & 4|\mathfrak{p}|^2 - 7|\mathfrak{p}| + 3 & 3|\mathfrak{p}|^2 - 5|\mathfrak{p}| + 2 & 2|\mathfrak{p}|^2 - 3|\mathfrak{p}| + 1 & (|\mathfrak{p}| - 1)|\mathfrak{p}| \\ 4|\mathfrak{p}|^2 - 7|\mathfrak{p}| + 3 & 4|\mathfrak{p}|^2 - 7|\mathfrak{p}| + 3 & 4|\mathfrak{p}|^2 - 6|\mathfrak{p}| + 3 & 3|\mathfrak{p}|^2 - 5|\mathfrak{p}| + 2 & 2|\mathfrak{p}|^2 - 3|\mathfrak{p}| + 1 & (|\mathfrak{p}| - 1)|\mathfrak{p}| \\ (|\mathfrak{p}| - 1)^2 & (|\mathfrak{p}| - 1)^2 &$$

where

$$\begin{split} \sigma(6) = & (4|\mathfrak{p}|^4 - 12|\mathfrak{p}|^3 + 5|\mathfrak{p}|^2 - |\mathfrak{p}| + 4, 4|\mathfrak{p}|^4 - 12|\mathfrak{p}|^3 + 4|\mathfrak{p}|^2 + |\mathfrak{p}| + 3, 4|\mathfrak{p}|^4 - 11|\mathfrak{p}|^3 + 4|\mathfrak{p}|^2 + 2|\mathfrak{p}| + 1, \\ & |\mathfrak{p}| \Big( 4|\mathfrak{p}|^3 - 10|\mathfrak{p}|^2 + 3|\mathfrak{p}| + 3 \Big), |\mathfrak{p}| \Big( 3|\mathfrak{p}|^3 - 8|\mathfrak{p}|^2 + 2|\mathfrak{p}| + 3 \Big), |\mathfrak{p}| \Big( 2|\mathfrak{p}|^3 - 5|\mathfrak{p}|^2 + |\mathfrak{p}| + 2 \Big), |\mathfrak{p}| \Big( |\mathfrak{p}|^3 - 2|\mathfrak{p}|^2 + 2 \Big). \end{split}$$

After Step 1 we obtain

$$M^H_{\delta} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ (|\mathfrak{p}|-1)^2 & |\mathfrak{p}|^2 - |\mathfrak{p}| + 1 & -|\mathfrak{p}| & 0 & 0 & 0 & 0 & 0 \\ 0 & -|\mathfrak{p}| & |\mathfrak{p}|^2 + 1 & -|\mathfrak{p}| & 0 & 0 & 0 & 0 \\ (|\mathfrak{p}|-1)|\mathfrak{p}|^3 & (|\mathfrak{p}|-1)|\mathfrak{p}|^3 & |\mathfrak{p}| \big( |\mathfrak{p}|^3 - |\mathfrak{p}|^2 - 1 \big) & |\mathfrak{p}|^4 - |\mathfrak{p}|^3 + |\mathfrak{p}|^2 + 1 & |\mathfrak{p}| \big( |\mathfrak{p}|^3 - 1 \big) & 0 & 0 \\ -(|\mathfrak{p}|-1)|\mathfrak{p}|^3 & -(|\mathfrak{p}|-1)|\mathfrak{p}|^3 & -(|\mathfrak{p}|-1)|\mathfrak{p}|^3 & -|\mathfrak{p}|^4 + |\mathfrak{p}|^3 - |\mathfrak{p}| & -|\mathfrak{p}|^4 + |\mathfrak{p}|^2 + 1 & -|\mathfrak{p}| & 0 \\ 0 & 0 & 0 & 0 & -|\mathfrak{p}| & |\mathfrak{p}|^2 + 1 & -|\mathfrak{p}| \\ \sigma(6)_1 & \sigma(6)_2 & \sigma(6)_3 & \sigma(6)_4 & \sigma(6)_5 & \sigma(6)_6 & \sigma(6)_7 \end{pmatrix}.$$

After Step 2 we obtain

$$M_{\delta}^{h} = \begin{pmatrix} 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ -|\mathfrak{p}| & |\mathfrak{p}|^{2} - |\mathfrak{p}| + 1 & -|\mathfrak{p}| & 0 & 0 & 0 & 0 \\ |\mathfrak{p}| & -|\mathfrak{p}| & |\mathfrak{p}|^{2} + 1 & -|\mathfrak{p}| & 0 & 0 & 0 & 0 \\ 0 & (|\mathfrak{p}| - 1)|\mathfrak{p}|^{3} & |\mathfrak{p}| (|\mathfrak{p}|^{3} - |\mathfrak{p}|^{2} - 1) & |\mathfrak{p}|^{4} - |\mathfrak{p}|^{3} + |\mathfrak{p}|^{2} + 1 & |\mathfrak{p}| (|\mathfrak{p}|^{3} - 1) & 0 & 0 \\ 0 & -(|\mathfrak{p}| - 1)|\mathfrak{p}|^{3} & -(|\mathfrak{p}| - 1)|\mathfrak{p}|^{3} & -|\mathfrak{p}|^{4} + |\mathfrak{p}|^{3} - |\mathfrak{p}| & -|\mathfrak{p}|^{4} + |\mathfrak{p}|^{2} + 1 & -|\mathfrak{p}| & 0 \\ 0 & 0 & 0 & 0 & -|\mathfrak{p}| & |\mathfrak{p}|^{2} + 1 & -|\mathfrak{p}| \\ (|\mathfrak{p}| - 1)^{2} & \sigma(6)_{2} & \sigma(6)_{3} & \sigma(6)_{4} & \sigma(6)_{5} & \sigma(6)_{6} & \sigma(6)_{7} \end{pmatrix}.$$

Notice that, from Claim 5.5, to compute the determinant of the matrix  $M_{\delta}^h$  we need to take a look at the determinant of the  $(r-1)\times(r-1)$  submatrix of  $M^h_{\delta}$  obtained by deleting the (r-1)th row and the 1st column; call this submatrix M. In particular, M is a Hessenberg matrix (after moving the first row to the last one), that is, it satisfies M[i][j] = 0 for all j > i+1. We will use the following theorem about determinants of Hessenberg matrices to prove Theorem 5.3.

**Theorem 5.8.** [2] Let  $M = (m_{i,j})_{1 \le i,j \le m}$  be an  $m \times m$  Hessenberg matrix. Let M(k) denote the submatrix of Mgiven by

$$M(k) := \begin{pmatrix} m_{1,1} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ m_{1,k} & \cdots & m_{k,k} \end{pmatrix}.$$

Set  $\det(M(0)) = 1$ . Then the determinants  $\{\det(M(k))\}_{k \ge 0}$  satisfy  $\det(M(0)) = 1$ ,  $\det(M(1)) = m_{1,1}$  and

$$\det(M(n)) = m_{n,n} \det(M(n-1)) + \sum_{i=1}^{n-1} \left( (-1)^{n-i} m_{n,i} \left( \prod_{j=i}^{n-1} m_{j,j+1} \det(M(i-1)) \right) \right).$$

Now we are ready to show that  $\det(M_{\delta}^h) \neq 0$ .

**Proposition 5.9.** We have  $det(M_{\delta}^h) = \pm 1 + |\mathfrak{p}|f(|\mathfrak{p}|)$  for some  $f(x) \in \mathbb{Z}[x]$ .

*Proof.* For  $r \le 6$ , a quick computation shows:

- Case 1: r=2. We have  $\det(M_{\delta}^h)=1$ .
- Case 2: r = 3. We have  $\det(M_{\delta}^{\tilde{h}}) = 1$ .
- Case 3: r = 4. We have  $\det(M_s^h) = -1$ .
- Case 4: r = 5. We have  $\det(M_{\delta}^{\bar{h}}) = 1$ .
- Case 5: r = 6. We have  $\det(M_{\delta}^{h}) = -1 |\mathfrak{p}|^2 + |\mathfrak{p}|^3 + |\mathfrak{p}|^4 |\mathfrak{p}|^5$ .

Hence, the statement is true for r < 7. Assume  $r \ge 7$ . Then  $M_{\delta}^h$  has the form stated in Claim 5.5. Hence, if we develop the determinant of  $M_{\delta}^h$  from the first column, from Claim 5.5 we have

$$\det(M_{\delta}^h) = |\mathfrak{p}|(M_{\delta}^h)_{2,0} + |\mathfrak{p}|(M_{\delta}^h)_{3,0} \pm (\sigma(r-1)_0 - \sigma(r-1)_1)(M_{\delta}^h)_{r-1,0},$$

where  $(M_{\delta}^h)_{i,j}$  denotes the minor of the matrix  $M_{\delta}^h$  formed after deleting the *i*-th row and *j*-th column. Hence, to prove the result, it is enough to check that

$$(\sigma(r-1)_0 - \sigma(r-1)_1)(M_{\delta}^h)_{r-1,0} = \pm 1 + |\mathfrak{p}|g(|\mathfrak{p}|)$$

for some  $g(x) \in \mathbb{Z}[x]$ . By the expression of  $\sigma(r-1)_k$  computed in the Appendix – see Remark 6.2 – we have that  $(\sigma(r-1)_0 - \sigma(r-1)_1) = \pm 1 + |\mathfrak{p}|g_1(|\mathfrak{p}|)$  for all  $r \geq 2$ . We will now use Theorem 5.8 to show that  $(M_{\delta}^h)_{r-1,1} = 1 + |\mathfrak{p}|g_2(|\mathfrak{p}|)$  for some  $g_2 \in \mathbb{Z}[x]$ . Let  $M := M_{\delta}^h$  and M(n) be the upper left  $n \times n$  matrix block of M. We proceed by induction on n. We claim that for all  $n \in \{1, \ldots, r-1\}$  there exists  $f_n(x) \in \mathbb{Z}[x]$  such that

$$\det(M(n)) = 1 + |\mathfrak{p}|f_n(|\mathfrak{p}|).$$

**Base case:** n = 1. We have  $M(n) = 1 - |\mathfrak{p}| + |\mathfrak{p}|^2$ . Hence, the claim is satisfied.

**Induction step:** Assume the statement is true for all M(i) with i < n. Notice the following two facts of the matrix M:

- $\cdot M[i][i] = 1 + |\mathfrak{p}|h_i(|\mathfrak{p}|);$
- $M[i][i+1] = |\mathfrak{p}|l_i(|\mathfrak{p}|);$

for all  $0 \le i \le n$ , where  $h_i(x), l_i(x) \in \mathbb{Z}[x]$ . Hence, by Theorem 5.8 we have

$$\det(M(n)) = m_{n,n}\det(M(n-1)) + \sum_{i=1}^{n-1} \left( (-1)^{n-i} m_{n,i} \left( \prod_{j=i}^{n-1} m_{j,j+1} \det(M(i-1)) \right) \right) =$$

$$(1 + |\mathfrak{p}|h_n(|\mathfrak{p}|))(1 + |\mathfrak{p}|f_{n-1}(|\mathfrak{p}|)) + \sum_{i=1}^{n-1} \left( (-1)^{n-i} m_{n,i} \left( \prod_{j=i}^{n-1} (|\mathfrak{p}|l_j(|\mathfrak{p}|))(1 + |\mathfrak{p}|f_{i-1}(|\mathfrak{p}|)) \right) \right) =$$

$$(1 + |\mathfrak{p}|h_n(|\mathfrak{p}|))(1 + |\mathfrak{p}|f_{n-1}(|\mathfrak{p}|)) + |\mathfrak{p}|t_n(|\mathfrak{p}|) = 1 + |\mathfrak{p}|g_n(|\mathfrak{p}|),$$

for some  $t_n(x) \in \mathbb{Z}[x]$ . Finally, we conclude that  $\det(M(n)) = 1 + |\mathfrak{p}|f_n(|\mathfrak{p}|)$  for all n.

Taking  $g_2(x) = f_{r-1}(x)$  we get

$$(\sigma(r-1)_0 - \sigma(r-1)_1)(M_\delta^h)_{r-1,0} = (\pm 1 + |\mathfrak{p}|g_1(x))(1 + |\mathfrak{p}|g_2(x)) = \pm 1 + |\mathfrak{p}|g(x),$$

for  $g(x) \in \mathbb{Z}[x]$ , which proves the proposition.

As a corollary, we obtain the following.

**Corollary 5.10.** The matrix  $M_{\delta}^h$  satisfies  $det(M_{\delta}^h) \neq 0$ .

*Proof.* By Proposition 5.9 we have  $\det(M^h_{\delta}) = \pm 1 + |\mathfrak{p}|f(|\mathfrak{p}|)$  for some  $f(x) \in \mathbb{Z}[x]$ . If  $\det(M^h_{\delta}) = 0$ , then we have  $\pm 1 = |\mathfrak{p}|f(|\mathfrak{p}|)$ . But then we would get that  $|\mathfrak{p}|$  divides either 1 or -1, which is not possible because  $|\mathfrak{p}| = q^{\deg(\mathfrak{p})} \geq 3$ . Hence,  $\det(M^h_{\delta}) \neq 0$  as we wanted.

Now we can prove Theorem 5.3.

*Proof of Theorem 5.3.* This follows from Corollary 5.10 together with Remark 5.6.

## 6. APPENDIX

**Definition 6.1.** ([12, 3.2]) The following case distinction defines the cuspidal divisor  $D_{r-1}$  for all  $r \ge 2$  in Theorem 2.8.

- (1) If r = 2, let  $D_{r-1} := C_1$ .
- (2) If  $r \ge 3$  and  $r \equiv 3 \mod 4$ , let

$$\begin{split} D_{r-1} &:= C_{r-1} - (|\mathfrak{p}|^r - |\mathfrak{p}|^{r-2})C_1 \\ &+ \sum_{2 \leq i \leq \frac{r-1}{2}} (|\mathfrak{p}|^{r-1} - |\mathfrak{p}|^{r-2} - |\mathfrak{p}|^{r-2i+1} + |\mathfrak{p}|^{r-2i})C_i \\ &- \sum_{\lfloor \frac{r+1}{2} \rfloor \leq i \leq r-2} (|\mathfrak{p}|^i - |\mathfrak{p}|^{\frac{r-1}{2}} + |\mathfrak{p}|^{i-\frac{r-1}{2}} - 1)(C_i - |\mathfrak{p}|C_{i+1}). \end{split}$$

(3) If  $r \ge 4$  and  $r \equiv 0 \mod 4$ , let

$$\begin{split} D_{r-1} &:= C_{r-1} - (|\mathfrak{p}|^r - |\mathfrak{p}|^{r-2})C_1 \\ &+ \sum_{2 \leq i \leq \frac{r}{2} - 1} (|\mathfrak{p}|^{r-1} - |\mathfrak{p}|^{r-2} - |\mathfrak{p}|^{r-2i+1} + |\mathfrak{p}|^{r-2i})C_i \\ &+ \sum_{\frac{r}{2} \leq i \leq r-2} (|\mathfrak{p}|^{i+1} - 2|\mathfrak{p}|^i + |\mathfrak{p}|^{\frac{r}{2}} - |\mathfrak{p}|^{i-\frac{r}{2}+1} + 1)(C_i - |\mathfrak{p}|C_{i+1}) \\ &- \sum_{\frac{r}{2} + 1 \leq i \leq r-3} (|\mathfrak{p}|^{i+1} - |\mathfrak{p}|^{\frac{r}{2}} + |\mathfrak{p}|^{i-\frac{r}{2}+1} - 1)(C_i - |\mathfrak{p}|C_{i+1}). \end{split}$$

(4) If  $r \ge 5$  and  $r \equiv 1 \mod 4$ , let

$$\begin{split} D_{r-1} &:= C_{r-1} - (|\mathfrak{p}|^r - |\mathfrak{p}|^{r-2})C_1 \\ &+ \sum_{2 \leq i \leq \frac{r-1}{2}} (|\mathfrak{p}|^{r-1} - |\mathfrak{p}|^{r-2} - |\mathfrak{p}|^{r-2i+1} + |\mathfrak{p}|^{r-2i})C_i \\ &- \sum_{\substack{\lfloor \frac{r+1}{2} \rfloor \leq i \leq r-2 \\ i : \text{ odd}}} (2|\mathfrak{p}|^{i+1} - |\mathfrak{p}|^i - |\mathfrak{p}|^{\frac{r-1}{2}} + |\mathfrak{p}|^{i-\frac{r-1}{2}} - 1)(C_i - |\mathfrak{p}|C_{i+1}) \\ &- \sum_{\substack{\frac{r+3}{2} \leq i \leq r-3 \\ i : \text{ even}}} (|\mathfrak{p}|^i - |\mathfrak{p}|^{\frac{r-1}{2}} + |\mathfrak{p}|^{i-\frac{r-1}{2}} - 1)(C_i - |\mathfrak{p}|C_{i+1}). \end{split}$$

(5) If  $r \ge 6$  and  $r \equiv 2 \mod 4$ , let

$$\begin{split} D_{r-1} &:= C_{r-1} - (|\mathfrak{p}|^r - |\mathfrak{p}|^{r-2}) C_1 \\ &+ \sum_{2 \leq i \leq \frac{r}{2} - 1} (|\mathfrak{p}|^{r-1} - |\mathfrak{p}|^{r-2} - |\mathfrak{p}|^{r-2i+1} + |\mathfrak{p}|^{r-2i}) C_i \\ &- \sum_{\frac{r}{2} \leq i \leq r-2} (|\mathfrak{p}|^{i+1} - |\mathfrak{p}|^{\frac{r}{2}} + |\mathfrak{p}|^{i-\frac{r}{2}+1} - 1) (C_i - |\mathfrak{p}|C_{i+1}). \end{split}$$

**Proposition 6.2.** The following table contains the computations of  $\sigma(r-1)_k$  for all possible  $r \ge 2$  and  $0 \le k \le r-1$ .

• *Case 1:* r = 2. *Then* 

$$\sigma(r-1)_k = \begin{cases} |\mathfrak{p}| - 1 & \text{if } k = 0; \\ |\mathfrak{p}| & \text{if } k = 1. \end{cases}$$

• *Case 2:* r = 3. *Then* 

$$\sigma(r-1)_k = \begin{cases} -2|\mathfrak{p}|^3 + 3|\mathfrak{p}|^2 + |\mathfrak{p}| - 2 & \text{if } k = 0; \\ -2|\mathfrak{p}|^3 + 2|\mathfrak{p}|^2 + |\mathfrak{p}| - 1 & \text{if } k = 1; \\ -|\mathfrak{p}|^3 + |\mathfrak{p}|^2 + |\mathfrak{p}| & \text{if } k = 2. \end{cases}$$

• *Case 3*: r = 4.

$$\sigma(r-1)_k = \begin{cases} -2|\mathfrak{p}|^3 + 3|\mathfrak{p}| - 3 & \text{if } k = 0; \\ -2|\mathfrak{p}|^3 + |\mathfrak{p}|^2 + 3|\mathfrak{p}| - 2 & \text{if } k = 1; \\ -2 + 2|\mathfrak{p}| + |\mathfrak{p}|^2 - |\mathfrak{p}|^3 & \text{if } k = 2; \\ -|\mathfrak{p}| \left( |\mathfrak{p}|^2 - 2 \right) & \text{if } k = 3. \end{cases}$$

• *Case 4*: r = 5.

$$\sigma(r-1)_{k} = \begin{cases} -3|\mathfrak{p}|^{3} + 4|\mathfrak{p}|^{2} + 2|\mathfrak{p}| - 3 & \text{if } k = 0; \\ -3|\mathfrak{p}|^{3} + 3|\mathfrak{p}|^{2} + 2|\mathfrak{p}| - 2 & \text{if } k = 1; \\ -2|\mathfrak{p}|^{3} + 3|\mathfrak{p}|^{2} + |\mathfrak{p}| - 2 & \text{if } k = 2; \\ -2|\mathfrak{p}|^{3} + |\mathfrak{p}|^{2} + |\mathfrak{p}| (|\mathfrak{p}|^{2} - 2) & \text{if } k = 3; \\ |\mathfrak{p}|^{2} & \text{if } k = 4. \end{cases}$$

$$\begin{cases} -3|\mathfrak{p}|^{3} + 3|\mathfrak{p}|^{2} + 4|\mathfrak{p}| - 4 & \text{if } k = 0; \\ -3|\mathfrak{p}|^{3} + 3|\mathfrak{p}|^{2} + 4|\mathfrak{p}| - 4 & \text{if } k = 0; \end{cases}$$

• *Case 5:* r = 6.

$$\sigma(r-1)_k = \begin{cases} -3|\mathfrak{p}|^3 + 3|\mathfrak{p}|^2 + 4|\mathfrak{p}| - 4 & \text{if } k = 0; \\ -3|\mathfrak{p}|^3 + 2|\mathfrak{p}|^2 + 4|\mathfrak{p}| - 3 & \text{if } k = 1; \\ -2|\mathfrak{p}|^3 + 2|\mathfrak{p}|^2 + 3|\mathfrak{p}| - 3 & \text{if } k = 2; \\ |\mathfrak{p}|^2 + 3|\mathfrak{p}| - 2|\mathfrak{p}| - 2 & \text{if } k = 3; \\ -|\mathfrak{p}|^3 + 2|\mathfrak{p}| - 1 & \text{if } k = 4; \\ |\mathfrak{p}| & \text{if } k = 5. \end{cases}$$

*Case 6:*  $r \ge 7$  *and*  $r \equiv 3 \pmod{4}$ .

$$\sigma(r-1)_k = \begin{cases} -\frac{(|\mathfrak{p}|-1)}{2} \left( (5-3r) + (r-1)|\mathfrak{p}| + 2(r-1)|\mathfrak{p}|^2 + (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} - (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = 0; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( (7-3r) + (r+1)|\mathfrak{p}| + 2(r-1)|\mathfrak{p}|^2 + (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} - (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = 1; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( (7-3r) + (r-1)|\mathfrak{p}| + 2(r-2)|\mathfrak{p}|^2 + (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} - (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = 2; \\ \frac{(|\mathfrak{p}|-1)}{2} \left( (-2k+3r-3) - (r-1)|\mathfrak{p}| + 2(k-r)|\mathfrak{p}|^2 - (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} + (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = \frac{r-1}{2}; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -2(r-1) + (r-1)|\mathfrak{p}| + (r+1)|\mathfrak{p}|^2 + (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} - (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = \frac{r-1}{2}; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -2(r-3) + (r-1)|\mathfrak{p}| + (r-1)|\mathfrak{p}|^2 + (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} - (r-1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = \lfloor \frac{r+1}{2} \rfloor; \\ (|\mathfrak{p}|-1) \left( -2(k-r+1) + (k-r)|\mathfrak{p}| + (k-r)|\mathfrak{p}|^2 + (k-r+1)|\mathfrak{p}|^{\frac{r-3}{2}} + (r-k)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = r-2; \\ -2 + 4|\mathfrak{p}|-2|\mathfrak{p}|^3 + |\mathfrak{p}|^{\frac{r-3}{2}} - 3|\mathfrak{p}|^{\frac{r-1}{2}} + 2|\mathfrak{p}|^{\frac{r+1}{2}} & \text{if } k = r-1. \end{cases}$$

Case 7:  $r \ge 8$  and  $r \equiv 0 \pmod{4}$ .

$$\sigma(r-1)_k = \begin{cases} -\frac{(|\mathfrak{p}|-1)}{2} \left( (4-3r) + (r+2)|\mathfrak{p}| + 2(r-2)|\mathfrak{p}|^2 + (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} - (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = 0; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -3(r-2) + (r+4)|\mathfrak{p}| + 2(r-2)|\mathfrak{p}|^2 + (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} - (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = 1; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -3(r-2) + (r+2)|\mathfrak{p}| + 2(r-3)|\mathfrak{p}|^2 + (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} - (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = 2; \\ \frac{(|\mathfrak{p}|-1)}{2} \left( -(2k+3r-2) - (r+2)|\mathfrak{p}| + 2(k-r+1)|\mathfrak{p}|^2 - (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} + (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = \frac{r}{2}-1; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -2r + (r+2)|\mathfrak{p}| + r|\mathfrak{p}|^2 + (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} - (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = \frac{r}{2}-1; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -2(r-1) + r|\mathfrak{p}| + (r-2)|\mathfrak{p}|^2 + (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} - r|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = \frac{r}{2}; \\ (|\mathfrak{p}|-1) \left( -2(k-r+1) + |\mathfrak{p}|(k-r-2) + |\mathfrak{p}|^2 (k-r) + (k-r+1)|\mathfrak{p}|^{\frac{r}{2}-2} + (r-k)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } \frac{r}{2} < k < r-2, \\ k = \frac{r}{2} + j \text{ and } j \text{ even}; \\ (|\mathfrak{p}|-1) \left( -2k+2r-1 \right) + (k-r)|\mathfrak{p}| + (k-r+1)|\mathfrak{p}|^2 + (k-r+1)|\mathfrak{p}|^{\frac{r}{2}-2} + (r-k)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = r-2; \\ (|\mathfrak{p}|-1) \left( -3 + 2|\mathfrak{p}| + |\mathfrak{p}|^2 + |\mathfrak{p}|^{\frac{r}{2}-2} - 2|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = r-2; \\ 3|\mathfrak{p}|-|\mathfrak{p}|^2 - |\mathfrak{p}|^3 - |\mathfrak{p}|^{\frac{r}{2}-1} + |\mathfrak{p}|^{\frac{r}{2}} & \text{if } k = r-1. \end{cases}$$

23

Case 8:  $r \ge 9$  and  $r \equiv 1 \pmod{4}$ .

$$\sigma(r-1)_k = \begin{cases} \frac{(|\mathfrak{p}|-1)}{2} \left( 2(r-2) + (r-1)|\mathfrak{p}|-3(r-1)|\mathfrak{p}|^2 - (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} + (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = 0; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -2(r-3) - (r-3)|\mathfrak{p}|+3(r-1)|\mathfrak{p}|^2 + (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} - (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = 1; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -2(r-3) - (r-1)|\mathfrak{p}|+(3r-5)|\mathfrak{p}|^2 + (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} - (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = 2; \\ \frac{(|\mathfrak{p}|-1)}{2} \left( -2(k-r+1) + (r-1)|\mathfrak{p}|+2(k-3r+1)|\mathfrak{p}|^2 - (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} + (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = \frac{r-1}{2}; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -(r-1) - (r-1)|\mathfrak{p}|+2r|\mathfrak{p}|^2 + (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} - (r+1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = \frac{r-1}{2}; \\ -\frac{(|\mathfrak{p}|-1)}{2} \left( -(r-5) - (r-5)|\mathfrak{p}|+2(r-1)|\mathfrak{p}|^2 + (r-3)|\mathfrak{p}|^{\frac{r-3}{2}} - (r-1)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = \frac{r+1}{2}; \\ (|\mathfrak{p}|-1) \left( (-k+r-2) + (-k+r-2)|\mathfrak{p}|+2(k-r)|\mathfrak{p}|^2 + (k-r+1)|\mathfrak{p}|^{\frac{r-3}{2}} + (r-k)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } \frac{r+1}{2} < k < r-2, \\ k = \frac{r+3}{2} + j \text{ and } j \text{ odd}; \\ (|\mathfrak{p}|-1) \left( (-k+r-1) + (r-k)|\mathfrak{p}|+(2k-2r+1)|\mathfrak{p}|^2 + (k-r+1)|\mathfrak{p}|^{\frac{r-3}{2}} + (r-k)|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = r-2; \\ -(|\mathfrak{p}|-1) \left( 4|\mathfrak{p}|^2 + |\mathfrak{p}|^{\frac{r-3}{2}} - 2|\mathfrak{p}|^{\frac{r-1}{2}} \right) & \text{if } k = r-1. \end{cases}$$

*Case 9:*  $r \ge 10$  *and*  $r \equiv 2 \pmod{4}$ .

$$\sigma(r-1)_k = \begin{cases} \frac{(|\mathfrak{p}|-1)}{2} \left( 2(r-2) + (r-2)|\mathfrak{p}| + (4-3r)|\mathfrak{p}|^2 - (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} + (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = 0; \\ \frac{(|\mathfrak{p}|-1)}{2} \left( 2(r-3) + (r-4)|\mathfrak{p}| - 3(r-2)|\mathfrak{p}|^2 + (4-3r)|\mathfrak{p}|^2 - (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} + (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = 1; \\ \frac{(|\mathfrak{p}|-1)}{2} \left( 2(r-3) + (r-2)|\mathfrak{p}| - (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} + (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = 2; \\ \frac{(|\mathfrak{p}|-1)}{2} \left( -2(k-r+1) + (r-2)|\mathfrak{p}| + (2k-3r+2)|\mathfrak{p}|^2 - (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} + (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = \frac{r}{2} - 1; \\ \frac{(|\mathfrak{p}|-1)}{2} \left( (r+(r-2)|\mathfrak{p}| - 2r|\mathfrak{p}|^2 - (r-2)|\mathfrak{p}|^{\frac{r}{2}-2} + (r+2)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = \frac{r}{2}; \\ (|\mathfrak{p}|-1) \left( (-k+r-1) + (-k+r-2)|\mathfrak{p}| + (2k-2r+1)|\mathfrak{p}|^2 + (k-r+1)|\mathfrak{p}|^{\frac{r}{2}-2} + (r-k)|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = r-2. \\ -(|\mathfrak{p}|-1) \left( -1 + 3|\mathfrak{p}|^2 |\mathfrak{p}|^{\frac{r}{2}-2} - 2|\mathfrak{p}|^{\frac{r}{2}-1} \right) & \text{if } k = r-1. \end{cases}$$

#### REFERENCES

- [1] C. Armana, S.-Y. K. Ho and M. Papikian, Ogg's conjectures over function fields, https://arxiv.org/abs/2410.05502, 2024.
- [2] N. D. CAHILL, J. R. D'ERRICO, D. A. NARAYAN AND J. Y. NARAYAN, Fibonacci determinants, The College Mathematics Journal, vol. 33 (2002), no. 3, pp. 221–225.
- [3] K. CONRAD., Carlitz extensions, Available at https://kconrad.math.uconn.edu/blurbs/gradnumthy/carlitz.pdf.
- [4] M. Curcó-Iranzo, Rational torsion of generalised modular Jacobians of odd level, To appear in Research in the Mathematical Sciences.
- [5] V. G. Drinfeld, Two theorems on modular curves, Funktsional'nyi Analiz i ego Prilozheniya, vol. 7 (1973), no. 2, pp. 83-84.
- [6] E.-U. GEKELER, A product expansion for the discriminant function of Drinfel'd modules of rank two, J. Number Theory, vol. 21 (1985), no. 2, pp. 135–140.
- [7] ——, On the Drinfeld discriminant function, Compositio Math., vol. 106 (1997), no. 2, pp. 181–202.
- [8] —, A note on the finiteness of certain cuspidal divisor class groups, Israel J. Math., vol. 118 (2000), pp. 357-368.
- [9] \_\_\_\_\_, Invariants of some algebraic curves related to Drinfeld modular curves, J. Number Theory, vol. 90 (2001), no. 1, pp. 166–183.
- [10] \_\_\_\_\_\_, Drinfel'd modular curves, Lecture Notes in Mathematics, vol. 1231, Springer-Verlag, Berlin, 1986.
- [11] E.-U. GEKELER AND U. NONNENGARDT, Fundamental domains of some arithmetic groups over function fields, Internat. J. Math., vol. 6 (1995), no. 5, pp. 689–708.
- [12] S.-Y. K. Ho., On the rational cuspidal divisor class groups of Drinfeld modular curves  $X_0(p^r)$ , Int. J. Number Theory, vol. **20** (2024), no. 8, pp. 2099–2123. *Int. J. Number Theory*, 20(8):2099–2123, 2024.
- [13] —, The rational torsion subgroup of  $J_0(\mathfrak{p}^r)$ , Research in the Mathematical Sciences, vol. 11 (2024), no. 4.
- [14] B. W. JORDAN AND K. A. RIBET AND A. J. SCHOLL, *Modular curves and Néron models of generalized Jacobians*, Compositio Mathematica, vol. **160** (2024), no. 5, pp. 945–981.
- [15] S. LANG AND A. NÉRON, Rational points of abelian varieties over function fields, Amer. J. Math., vol. 81 (1959), pp. 95-118.
- [16] J. I. Manin, Parabolic points and zeta-functions of modular curves, Izv. Akad. Nauk SSSR Ser. Mat., vol. 6 (1972), no. 1, pp. 19-66.
- [17] B. MAZUR, *Modular curves and the Eisenstein ideal (with an appendix by B. Mazur and M. Rapoport)*, Publications Mathématiques de l'Institut des Hautes Études Scientifiques, vol. 47 (1977), no. 1 pp. 33–186.
- [18] ——, Diophantine equations and modular forms, Bulletin of the American Mathematical Society, vol. 81 (1975), no. 1, pp. 14–27.
- [19] M. Ohta, Eisenstein ideals and the rational torsion subgroups of modular Jacobian varieties, Journal of the Mathematical Society of Japan, vol. **65** (2013), no. 3, pp. 733–772.
- [20] A. PÁL, On the torsion of the Mordell-Weil group of the Jacobian of Drinfeld modular curves, Doc. Math., vol. 10 (2005), pp. 131–198.
- [21] M. PAPIKIAN AND F.-T. WEI, On the Eisenstein ideal over function fields, J. Number Theory, vol. 161 (2016), pp. 384-434.
- [22] ——, The rational torsion subgroups of Drinfeld modular Jacobians and Eisenstein pseudo-harmonic cochains, Mathematische Zeitschrift, vol. **287** (2017), no. 1-2, pp. 521–546.
- [23] ——, *Drinfeld discriminant function and Fourier expansion of harmonic cochains*, Mathematische Annalen, vol. **388** (2024), no. 2, pp. 1379–1435.
- [24] M. ROSENLICHT, Generalised Jacobian Varieties, Annals of Mathematics, vol. 56 (1954), no. 3, pp. 505-530.
- [25] J.P. SERRE, *Groupes algébriques et corps de classes*, Publications de l'Institut de Mathématique de l'Université de Nancago VII, Hermann, Paris (1959).
- [26] F.-T. WEI AND T. YAMAZAKI, Rational torsion of generalized Jacobians of modular and Drinfeld modular curves, in Forum Mathematicum, vol. 31 (2019), no. 3, pp. 647–659.
- [27] T. YAMAZAKI AND Y. YANG, Rational torsion on the generalised Jacobian of a modular curve with cuspidal modulus, Documenta Mathematica, vol. 21 (2016), pp. 1669-1690.
- [28] ——, The rational cuspidal divisor class group of  $X_0(N)$ , Journal of Number Theory, vol. **242** (2023), pp. 278–401.
- [29] ——, The rational torsion subgroup of  $J_0(N)$ , Advances in Mathematics, vol. **426** (2023), no. 109111, pp. 30.

Email address: m.curcoiranzo@uu.nl

MATHEMATICAL INSTITUTE, UTRECHT UNIVERSITY, PO BOX 80010, 3508 TA, UTRECHT, THE NETHERLANDS