Strong Gaussian approximations with random multipliers^{*}

Fabian Mies

Delft University of Technology[†]

December 20, 2024

Abstract

One reason why standard formulations of the central limit theorems are not applicable in high-dimensional and non-stationary regimes is the lack of a suitable limit object. Instead, suitable distributional approximations can be used, where the approximating object is not constant, but a sequence as well. We extend Gaussian approximation results for the partial sum process by allowing each summand to be multiplied by a data-dependent matrix. The results allow for serial dependence of the data, and for high-dimensionality of both the data and the multipliers. In the finite-dimensional and locally-stationary setting, we obtain a functional central limit theorem as a direct consequence. An application to sequential testing in non-stationary environments is described.

1 Introduction

Donsker's theorem states that for iid, centered, d-variate random vectors X_t , the partial sum process $S_n(u) = \frac{1}{\sqrt{n}} \sum_{t=1}^{\lfloor un \rfloor} X_t$, $u \in [0, 1]$, converges weakly to $\Sigma^{\frac{1}{2}}B(u)$ in the Skorokhod space D[0, 1], where B(u) is a standard d-variate Brownian motion B(u), and $\Sigma = \text{Cov}(X_t)$ is the covariance matrix of the random vectors. If the X_t are not iid but stationary and satisfy suitable ergodicity properties, e.g. strong mixing, then the same functional central limit theorem (FCLT) holds, but with $\Sigma = \sum_{h=-\infty}^{\infty} \text{Cov}(X_t, X_{t+h})$ the long-run variance (Billingsley, 1999). Another deviation from the iid assumption is a potential non-stationarity of the X_t . In the latter case, it is difficult to formulate a FCLT because it is in general unclear how to specify a limit random element or

^{*}Funded by the Federal Ministry of Education and Research (BMBF) and the Ministry of Culture and Science of the German State of North Rhine-Westphalia (MKW) under the Excellence Strategy of the Federal Government and the Länder.

[†]This work was done while the author was employed at RWTH Aachen University, Germany.

distribution. One mathematical trick to solve this challenge is the asymptotic framework of locally stationary time series introduced by Dahlhaus (1997). Here, the nonstationarity is reparametrized from the absolute time $t \in \{1, \ldots, n\}$ to the relative time $\frac{t}{n} \in [0, 1]$, such that for $\frac{t}{n} \approx u$, the time series X_t may be well approximated by a stationary time series $X_t(u)$. A rather general model formalizing this idea is introduced in Dahlhaus et al. (2019), see also Section 2 below. Under suitable conditions, the locallystationary version of Donsker's Theorem is of the form $S_n(u) \Rightarrow \int_0^u \Sigma^{\frac{1}{2}}(u) dB(u)$, where $\Sigma(u) = \sum_{h=-\infty}^{\infty} \text{Cov}(X_t(u), X_{t+h}(u))$ is the long-run variance of the local approximating time series.

Moving beyond the locally stationary framework, the FCLT breaks down because there is no well-defined limit object. In the same way, the FCLT breaks down in a highdimensional asymptotic setting, where $d = d_n \to \infty$ as $n \to \infty$. What can be salvaged in these situations is a good approximation of $S_n(u)$ in terms of a Gaussian process $B_n(u)$, say, such that $\sup_{u \in [0,1]} ||S_n(u) - B_n(u)|| \to 0$ in probability as $n \to \infty$. That is, we approximate the sequence $S_n(u)$ by a sequence of Gaussian processes, instead of a single limiting Gaussian process. Approximations of this kind have been known for iid data as strong approximations, or Hungarian couplings (Komlós et al., 1975, 1976), where optimal rates of approximation have been achieved; see Zaitsev (2013) for an overview. More recent contributions for multivariate and / or dependent data are due to Liu and Lin (2009); Wu and Zhou (2011); Berkes et al. (2014); Karmakar and Wu (2020); Bonnerjee et al. (2024). For non-stationary and high-dimensional time series, a corresponding result has been derived in Mies and Steland (2023); see also Mies and Steland (2024). In this article, we extend the latter approximation in two regards:

- (i) We exploit a stronger covariance bound to improve the approximation rates of Mies and Steland (2023).
- (ii) We extend the results to partial sums with data-dependent multipliers, i.e. $\widetilde{S}_n(u) = \frac{1}{\sqrt{n}} \sum_{t=1}^{\lfloor un \rfloor} \widehat{g}_t X_t$ for random matrices $\widehat{g}_t \in \mathbb{R}^{m \times d}$, with potentially growing dimension $d \to \infty, m \to \infty$.

Notation For a vector $x \in \mathbb{R}^d$, we denote by $||x|| = (\sum_{i=1}^d x_i^2)^{1/2}$ the Euclidean norm, and for a matrix $A \in \mathbb{R}^{d \times d}$, A^T denotes its transpose, $\operatorname{tr}(A)$ its trace, $||A||_{\operatorname{tr}} = \operatorname{tr}((A^T A)^{\frac{1}{2}})$ its trace norm. For any matrix $A \in \mathbb{R}^{m \times d}$, we denote by $||A|| = ||A||_F = \sqrt{\operatorname{tr}(A^T A)}$ the Frobenius norm. We use \Rightarrow to denote weak convergence of probability measures resp. random elements. The \mathcal{L}_q norm of a random vector X is denoted as $||X||_{\mathcal{L}_q} = (\mathbb{E}||X||^q)^{1/q}$, i.e. we always use the Euclidean base-norm. For two symmetric matrices A, B, the notation $A \succeq B$ means that A - B is positive semidefinite.

2 Sequential approximations for time series

A general class of non-stationary and potentially nonlinear time series may be formulated as follows. For iid random seeds $\epsilon_i \sim U(0, 1)$, and non-random functions $G_{t,n} : \mathbb{R}^\infty \to \mathbb{R}^d$, $t = 1, \ldots, n$, define the sequence of non-stationary time series $\{X_{t,n}, t = 1, \ldots, n\}$ with $X_{t,n} \in \mathbb{R}^d$ as

$$X_{t,n} = G_{t,n}(\boldsymbol{\epsilon}_t), \quad t = 1, \dots, n,$$

$$\boldsymbol{\epsilon}_t = (\boldsymbol{\epsilon}_t, \boldsymbol{\epsilon}_{t-1}, \dots) \in \mathbb{R}^{\infty}.$$

Throughout this section, we assume that $\mathbb{E}(X_{t,n}) = 0$. For an independent copy $\tilde{\epsilon}_i \sim U(0, 1)$, define furthermore

$$\tilde{\boldsymbol{\epsilon}}_{t,h} = (\epsilon_t, \dots, \epsilon_{t-h+1}, \tilde{\epsilon}_{t-h}, \epsilon_{t-h-1} \dots) \in \mathbb{R}^{\infty}, \\ \boldsymbol{\epsilon}_{t,h} = (\epsilon_t, \dots, \epsilon_{t-h+1}, \tilde{\epsilon}_{t-h}, \tilde{\epsilon}_{t-h-1} \dots) \in \mathbb{R}^{\infty}.$$

To express the mixing-type behavior of $X_{t,n}$, we employ the physical dependence measure introduced by Wu (2005), and defined as

$$\delta_n(h) := \max_{t=1,\dots,n} \|G_{t,n}(\boldsymbol{\epsilon}_0) - G_{t,n}(\tilde{\boldsymbol{\epsilon}}_{0,h})\|_{\mathcal{L}_q}.$$

for some $q \ge 2$. The dependence of δ_n on q will be implicit in the remainder of this article.

Remark 1. For a nonstationary linear process of the form $X_{t,n} = \sum_{i=0}^{\infty} a_{t,n,i} \eta_{t-i}$ with iid random vectors $\eta_t \in \mathbb{R}^d$ and matrices $a_{t,n,i} \in \mathbb{R}^{d \times d}$, we may write $\eta_t = h(\epsilon_t)$ for a measurable mapping $h : \mathbb{R} \to \mathbb{R}^d$. The physical dependence measure may then be computed as $\delta_n(h) = \|\eta_1\|_{\mathcal{L}_q} \sup_t \|\alpha_{t,n,h}\|_{\text{op}}$, where $\|A\|_{\text{op}}$ denotes the Euclidean operator norm of a matrix $A \in \mathbb{R}^{d \times d}$. Upper bounds on the physical dependence measure may also be established for a broader class of time-varying ARMA processes as in Subba Rao (1970); Grenier (1983); Moulines et al. (2005); Dahlhaus and Polonik (2009); Giraud et al. (2015). Here, $\delta_n(h) \to 0$ polynomially as $h \to \infty$ if the roots of the autoregressive poolynomial are bounded away from one (Mies, 2023, Example 1).

We first impose the following set of assumptions, for some $\Gamma_n \ge 1$ and $\beta > 1$:

$$\|G_{1,n}(\boldsymbol{\epsilon}_0)\|_{\mathcal{L}_2} + \sum_{t=2}^n \|G_{t,n}(\boldsymbol{\epsilon}_0) - G_{t-1,n}(\boldsymbol{\epsilon}_0)\|_{\mathcal{L}_2} \le \Theta_n \Gamma_n, \tag{A.1}$$
$$\delta_n(h) \le \Theta_n(h+1)^{-\beta}, \tag{A.2}$$

As a first result, we find that the Gaussian approximation result presented in Mies and Steland (2023) can be slightly improved in terms of the dependence on the exponent β . In particular, the following property has not been fully exploited in the proofs therein.

Proposition 1. Assumption (A.2) implies that $\|\operatorname{Cov}(X_{t,n}, X_{s,n})\|_{\mathrm{tr}} \leq \Theta_n^2 C_\beta (|s-t|+1)^{-\beta}$.

To state the Gaussian approximation result, define

$$\xi(q,\beta) = \begin{cases} \frac{q-2}{6q-4}, & \beta \ge \frac{3}{2}, \beta > \frac{2q}{q+2}, \\ \frac{\beta-1}{4\beta-2}, & \beta \ge \frac{3}{2}, \beta \le \frac{2q}{q+2}, \\ \frac{(\beta-1)(q-2)}{4q\beta-3q-2}, & \beta < \frac{3}{2}, \beta > \frac{2q}{q+2}, \\ \frac{(\beta-1)^2}{2\beta^2-1-\beta}, & \beta < \frac{3}{2}, \beta \le \frac{2q}{q+2}, \end{cases} \qquad q > 2, \beta > 1$$

Theorem 2. Let $X_{t,n} = G_{t,n}(\boldsymbol{\epsilon}_t)$ with $\mathbb{E}(X_{t,n}) = 0$ satisfy (A.1) and (A.2), for some q > 2, and suppose $d \leq cn$ for some c > 0. Suppose furthermore that $\beta > 1$, which implies that the local long-run covariance matrix $\Sigma_{t,n} = \sum_{h=-\infty}^{\infty} \operatorname{Cov}(G_{t,n}(\boldsymbol{\epsilon}_0), G_{t,n}(\boldsymbol{\epsilon}_h))$ is well-defined. Then, on a potentially different probability space, there exist random vectors $(X'_{t,n})_{t=1}^n \stackrel{d}{=} (X_{t,n})_{t=1}^n$ and independent, mean zero, Gaussian random vectors $Y_t^* \sim \mathcal{N}(0, \Sigma_{t,n})$ such that

$$\left(\mathbb{E}\max_{k\leq n}\left\|\frac{1}{\sqrt{n}}\sum_{t=1}^{k}(X'_{t,n}-Y^*_{t,n})\right\|^2\right)^{\frac{1}{2}}\leq C\Theta_n\Gamma_n^{\frac{\beta-1}{2\beta}}\sqrt{\log(n)}\left(\frac{d}{n}\right)^{\xi(q,\beta)}.$$
(1)

Remark 2. The essential improvement of Theorem 2 over Theorem 3.1(ii) of Mies and Steland (2023) is that the new formulation only requires $\beta > 1$, instead of $\beta > 2$. This is possible by exploiting Proposition 1 throughout the proof.

3 Strong approximation with random multipliers

In this section, we extend the Gaussian approximation result to also allow for datadependent multipliers. That is, we consider the partial sums $\frac{1}{\sqrt{n}} \sum_{t=1}^{n} \hat{g}_{t,n} X_{t,n}$ for random matrices $\hat{g}_{t,n} \in \mathbb{R}^{m \times d}$, with potentially growing dimensions $d = d_n \to \infty$, $m = m_n \to \infty$. The matrices $\hat{g}_{t,n}$ are assumed to converge to some non-random matrix $g_{t,n}$ in the sense that

$$\Lambda_n := \sqrt{\sum_{t=1}^n \|\hat{g}_{t,n} - g_{t,n}\|_{\mathcal{L}_2}^2} = o(\sqrt{n})$$

Moreover, the latter sequences of matrices should admit some regularity in time, which we express in terms of the quantities

$$\Psi_n := \sum_{t=2}^n \|g_{t,n} - g_{t-1,n}\|, \qquad \Phi_n = \max_{t=1,\dots,n} \|g_{t,n}\|.$$

In all the terms Λ_n , Ψ_n , and Φ_n , the dependence on the dimension d_n and m_n is implicit in the norms. Besides these technical restrictions, we impose the important qualitative restriction that, for some sequence $L_n \in \mathbb{N}$ which is not necessarily diverging,

$$\widehat{g}_{t,n}$$
 is ϵ_{t-L_n} -measurable. (A.M)

Thus, if $\hat{g}_{t,n}$ is based on an estimator, then it may only use the data up to time $t - L_n$. A similar measurability assumption was imposed in Mies (2023). To express the relation between the delay L_n and the serial dependence of the time series, define the term

$$\Xi_n := \sum_{h=L_n}^{\infty} \delta_n(h).$$

Remark 3. Note that (A.2) implies that $\Xi_n \leq \Phi_n L_n^{1-\beta}$, but in some situations, it might be possible to derive sharper bounds on Ξ_n . In particular, if the $X_{t,n}$ are indeed independent, then $\Xi_n = 0$ for any sequence $L_n \geq 1$, thus greatly reducing the measurability conditions on $\widehat{g}_{t,n}$. In the latter case, (A.2) also holds for all $\beta > 0$.

The local long run covariance matrix of the process $g_{t,n}X_{t,n}$ is given by

$$\Sigma_{t,n} = \sum_{h=-\infty}^{\infty} g_{t,n} \operatorname{Cov}[G_{t,n}(\boldsymbol{\epsilon}_0), G_{t,n}(\boldsymbol{\epsilon}_h)] g_{t,n}^T.$$

Note that $\Sigma_{t,n}$ is finite for $\beta > 1$ by virtue of Proposition 1.

Theorem 3. Let Assumptions (A.1) and (A.2) hold, and suppose that $\hat{g}_{t,n}$ is ϵ_{t-L_n} measurable for some $L_n \geq 1$. Then, upon potentially enlarging the probability space, there exist independent Gaussian random vectors $Y_{t,n} \sim \mathcal{N}(0, \Sigma_{t,n})$, such that

$$\begin{aligned} & \left| \max_{k=1,\dots,n} \frac{1}{\sqrt{n}} \sum_{t=1}^{k} [\hat{g}_{t,n} X_{t,n} - Y_{t,n}] \right| \\ &= \mathcal{O}_P \left(n^{-\frac{1}{2}} \Lambda_n \Theta_n + \Lambda_n \Xi_n + n^{\frac{1}{q} - \frac{1}{2}} L_n \Phi_n \right) \\ &+ \mathcal{O}_P \left(\Phi_n \Theta_n \left(\Phi_n \Theta_n \Gamma_n + \Psi_n \Theta_n \right)^{\frac{\beta - 1}{2\beta}} \sqrt{\log(n)} \left(\frac{m}{n} \right)^{\xi(q,\beta)} \right) \end{aligned}$$

This result is also applicable if $m = m_n \to \infty$.

Multiplier central limit theorems are often used in the literature to study the behavior of bootstrap methods, see e.g. (van der Vaart and Wellner, 1996, Sec. 2.9). There, the process $X_{t,n}$ and the multipliers are independent, which is not the case in our setting. Instead, Assumption (A.4) implies that the $\hat{g}_{t,n}$ are almost deterministic, up to some estimation error. The contribution of Theorem 3 is to take this estimation error explicitly into account. Compared to Theorem 2, it allows for greater flexibility in the decay of the physical dependence measure $\delta_n(h)$ as expressed via the quantity Ξ_n , rather than assuming a polynomial decay.

4 Functional central limit theorem

In the classical finite-dimensional setting where d and m are constant, the previous results also allow for the formulation of a functional central limit theorem. We embed the idea of local stationarity in our framework by considering a local asymptotic mapping $G_u: \mathbb{R}^{\infty} \to \mathbb{R}^d, u \in [0, 1]$, such that, as $n \to \infty$,

$$\int_0^1 \|G_{\lfloor vn \rfloor, n}(\boldsymbol{\epsilon}_0) - G_v(\boldsymbol{\epsilon}_0)\|_{\mathcal{L}_2} \, dv \to 0.$$
(A.3)

Then $X_t(u) = G_u(\boldsymbol{\epsilon}_t)$ is the local stationary approximation in the sense of Dahlhaus (1997). Analogously, we suppose that there exists another class of matrices $g_u \in \mathbb{R}^{m \times d}$, $u \in [0, 1]$, such that $\sup_u ||g_u|| < \infty$ and, as $n \to \infty$,

$$\int_0^1 \|g_{\lfloor vn\rfloor,n} - g_v\| \, dv \to 0. \tag{A.4}$$

Since we are now working with fixed dimensions d and m, the choice of the matrix norm is irrelevant. To formulate the functional central limit theorem, introduce the asymptotic local long-run variance matrix

$$\Sigma_u = \sum_{h=-\infty}^{\infty} g_u \operatorname{Cov}[G_u(\boldsymbol{\epsilon}_0), G_u(\boldsymbol{\epsilon}_h)] g_u^T.$$

Lemma 4. Suppose that (A.2), (A.3), (A.4) hold, and $\Theta_n, \Phi_n = \mathcal{O}(1)$. For independent Gaussian random vectors $Y_{t,n} \sim \mathcal{N}(0, \Sigma_{t,n})$, we have the following weak convergence in the Skorokhod space D[0, 1]:

$$\frac{1}{\sqrt{n}}\sum_{t=1}^{\lfloor un \rfloor} Y_{t,n} \Rightarrow \int_0^u \Sigma_v^{\frac{1}{2}} dW_v.$$

Combining Theorem 3 and Lemma 4, we immediately obtain the main result of this paper.

Theorem 5. Suppose that $\hat{g}_{t,n}$ is $\boldsymbol{\epsilon}_{t-L}$ -measurable for $L = L_n$, and let (A.1), (A.2), (A.3), (A.4) hold with $\Theta_n, \Phi_n = \mathcal{O}(1)$ such that

$$\Lambda_n \left(n^{-\frac{1}{2}} + \Xi_n \right) \to 0, \qquad L_n = o \left(n^{\frac{1}{2} - \frac{1}{q}} \right),$$
$$\left(\Gamma_n + \Psi_n \right)^{\frac{\beta - 1}{2\beta}} \sqrt{\log(n)} n^{-\xi(q,\beta)} \to 0.$$

Then

$$\frac{1}{\sqrt{n}} \sum_{t=1}^{\lfloor un \rfloor} \hat{g}_{t,n} [X_{t,n} - \mathbb{E}(X_{t,n})] \Rightarrow \int_0^u \Sigma_v^{\frac{1}{2}} dW_v,$$

where weak convergence holds in the Skorokhod space D[0, 1].

5 Example: studentized partial sums

In the sequel, we outline a statistical application of the FCLT with data-dependent multipliers. Consider a heteroskedastic sequence of independent *d*-variate random vectors $X_{t,n} = \Sigma(t/n)^{\frac{1}{2}}\eta_t$, for iid random vectors η_t with $\operatorname{Cov}(\eta_t) = I_{d\times d}$, $\mathbb{E}(\eta_t) = 0$, $\mathbb{E}\|\eta_t\|^q < \infty$ for some q > 2, and a Lipschitz-continuous function $\Sigma : [0, 1] \to \mathbb{R}^{d\times d}$. Suppose that each $\Sigma(u)$ is symmetric positive definite, and that $\Sigma(u)$ is uniformly elliptic: there exists some small c > 0 such that $\Sigma(u) \succeq cI_{d \times d}$, i.e. $\Sigma(u) - cI_{d \times d}$ is positive semidefinite for all $u \in [0, 1]$. The FCLT for locally-stationary data, e.g. Theorem 5 with $\hat{g}_{t,n} = I_{d \times d}$, yields that

$$S_n(u) = \frac{1}{\sqrt{n}} \sum_{t=1}^{\lfloor un \rfloor} X_{t,n} \Rightarrow \int_0^u \Sigma(v)^{\frac{1}{2}} dW_v.$$
(2)

Although technically a special case of the previous theorem, the latter convergence is rather standard in the literature and does not exploit our new results. Instead, to showcase the application of Theorem 5, we estimate $\Sigma(u)$ locally by the window average

$$\widehat{\Sigma}(t/n) = \frac{1}{k_n} \sum_{j=1}^{k_n} X_{t-j} X_{t-j}^T, \qquad t = k_n + 1, \dots, n.$$

For $t \leq k_n$, let $\widehat{\Sigma}(t/n) = 0$. To enforce the uniform ellipticity, we define

$$\widetilde{\Sigma}(t/n) = \begin{cases} \widehat{\Sigma}(t/n), & \text{if } \widehat{\Sigma}(t/n) \succeq cI_{d \times d}, \\ cI_{d \times d}, & \text{otherwise.} \end{cases}$$

We consider the studentized partial sum process

$$S_n^*(u) = \frac{1}{\sqrt{n}} \sum_{t=k_n+1}^{\lfloor un \rfloor} \widetilde{\Sigma}(t/n)^{-\frac{1}{2}} X_{t,n}$$

Theorem 6. For $k_n = \lfloor n^{\frac{2}{3}} \rfloor$, the studentized partial sum process converges weakly in the Skorokhod space to a standard Brownian motion,

$$S_n^*(u) \Rightarrow \int_0^u dW_v. \tag{3}$$

Both limits (2) and (3) can be used to test the null hypothesis $H_0: \mathbb{E}(X_{t,n}) = 0$ via the test statistic

$$T_{n}^{*} = \sup_{u \in [0,1]} \|S_{n}^{*}(u)\| \Rightarrow \sup_{u \in [0,1]} \left\| \int_{0}^{u} dW_{v} \right\|,$$
$$T_{n} = \sup_{u \in [0,1]} \|S_{n}(u)\| \Rightarrow \sup_{u \in [0,1]} \left\| \int_{0}^{u} \Sigma(v)^{\frac{1}{2}} dW_{v} \right\|$$

The supremum may be evaluated exactly as S_n and S_n^* are step functions. Because T_n^* weights the observations according to their inverse variance, we expect this statistic to be more powerful. This is supported by simulation results reported in Table 1, for sample size n = 10000. Here, the data is generated as $X_{t,n} = \sigma(\frac{t}{n})(Z_t - 1) + \mu$ with $Z_t \sim \text{Exp}(1)$, and $\sigma(u) = 1.2 + \sin(6\pi u)$. It is found that both the size distortion of T_n^* is slightly bigger, but its power is much higher than for the test statistic T_n . However, for

μ	0	.002	.004	.006	.008	.010	.012	.014	.016	.018	.020
Power of T_n	.048	.053	.054	.060	.078	.094	.119	.151	.184	.213	.254
Power of T_n^*	.066	.079	.097	.134	.194	.261	.362	.456	.559	.656	.743

Table 1: Power of the test statistics T_n and T_n^* for sample size n = 10000. Reported results are based on 10^4 simulations each.

smaller sample sizes (n = 1000), we observed that T_n^* suffers from a more pronounced upward size distortion (test size 0.097).

A second important distinction is that T_n^* enables a sequential testing procedure. Since the critical value c_{α}^* of $\sup_u ||W_u||$ does not depend on the unknown function Σ , we can reject H_0 as soon as $||S_n^*(u)|| > c_{\alpha}^*$. Although the non-studentized test statistic $||S_n(u)||$ may also be computed sequentially, the corresponding critical value $c_{\alpha}(\Sigma)$ will depend on the whole function $\Sigma(u)$. However, at time t, the best we can achieve is to estimate $\Sigma(u)$ for $u \in [0, \frac{t}{n}]$.

The idea of local studentization can also be extended to non-stationary time series. In this situation, we need to replace $\hat{\Sigma}(t/n)$ by a suitable estimator of the long-run variance matrix. The technical details are beyond the scope of this article.

Proofs

Proof of Proposition 1

Observe that the random vectors $Y_{t,n,i} = \mathbb{E}(X_{t,n}|\boldsymbol{\epsilon}_i) - \mathbb{E}(X_{t,n}|\boldsymbol{\epsilon}_{i-1}), i \in \mathbb{Z}$, are martingale differences, with

$$\|Y_{t,n,i}\|_{\mathcal{L}_q} \leq \begin{cases} \|G_{t,n}(\boldsymbol{\epsilon}_t) - G_{t,n}(\boldsymbol{\epsilon}_{t,t-i})\|_{\mathcal{L}_q} \leq \Theta_n(t-i+1)^{-\beta}, & \text{if } i \leq t, \\ 0, & \text{if } i > t. \end{cases}$$

The analogous bound holds for $Y_{s,n,i} = \mathbb{E}(X_{s,n}|\boldsymbol{\epsilon}_i) - \mathbb{E}(X_{s,n}|\boldsymbol{\epsilon}_{i-1})$. Thus,

$$\|\operatorname{Cov}(X_{t,n}, X_{s,n})\|_{\operatorname{tr}} = \left\|\sum_{i \in \mathbb{Z}} \operatorname{Cov}(Y_{t,n,i}, Y_{s,n,i})\right\|_{\operatorname{tr}}$$

$$\leq \sum_{i \leq (s \wedge t)} \mathbb{E} \|Y_{t,n,i}Y_{s,n,i}^{T}\|_{\operatorname{tr}}$$

$$\stackrel{(\dagger)}{=} \sum_{i \leq (s \wedge t)} \mathbb{E} \left(\|Y_{t,n,i}\| \cdot \|Y_{s,n,i}\|\right)$$

$$\leq \sum_{i \leq (s \wedge t)} \|Y_{t,n,i}\|_{\mathcal{L}_{2}} \cdot \|Y_{s,n,i}\|_{\mathcal{L}_{2}}$$

$$\leq \Theta_{n}^{2} \sum_{k=0}^{\infty} (k+1)^{-\beta} (k+|s-t|+1)^{-\beta}$$

$$\leq \Theta_n^2 C_\beta (|s-t|+1)^{-\beta}.$$

At the step (†), we use that $||xy^T||_{tr} = ||x|| \cdot ||y||$, see for example (Mies and Steland, 2023, Lemma 6.3).

Proof of Theorem 2

We show how to improve various steps in the proof of Theorem 3.1 of Mies and Steland (2023). For consistency with the latter reference, in this proof, L denotes a block length parameter distinct from the lag sequence L_n of the main text.

By using the improved inequality $\|\operatorname{Cov}(G_{t,n}(\boldsymbol{\epsilon}_r), G_{t,n}(\boldsymbol{\epsilon}_s))\|_F \leq C\Theta^2(|r-s|+1)^{-\beta}$ from Proposition 1, we may improve (22) therein to

$$\left\|\Sigma_{t,n} - \Sigma_{t,n}^{l}\right\|_{\mathrm{tr}} \le C\Theta_{n}^{2} L^{(1-\beta)\vee(-1)},$$

and (23) therein to

$$\|\Sigma_{t,n}^{l} - \operatorname{Cov}(Y_{t,n}')\|_{\mathrm{tr}} \le C\Theta_{n}^{2} \left[\frac{1}{\Theta_{n}} \sum_{s=t_{l-1}+1}^{t_{l}} \|G_{s,n}(\boldsymbol{\epsilon}_{0}) - G_{s-1,n}(\boldsymbol{\epsilon}_{0})\|_{\mathcal{L}_{2}}\right]^{1-\frac{1}{\beta}}$$

Hence, we can improve (26) therein to

$$\left(\mathbb{E}\max_{k\leq n} \left\|\sum_{t=1}^{k} Y_{t,n}^{*} - Y_{t,n}^{\prime}\right\|^{2}\right)^{\frac{1}{2}} \leq C\Theta_{n} n^{\frac{1}{2}} L^{\frac{1-\beta}{2}\vee(-\frac{1}{2})} + C\Theta_{n} L^{\frac{1}{2}} \Gamma_{n}^{\frac{\beta-1}{2\beta}} M^{\frac{1}{2\beta}}.$$

1

That is, all occurrences of β in (26) therein are replaced by $\beta + 1$. As in (27) therein, we obtain for any $L \leq c\frac{n}{d}$,

$$\begin{split} &\left(\mathbb{E}\max_{k\leq n}\left\|\frac{1}{\sqrt{n}}\sum_{t=1}^{k}(X_{t,n}'-Y_{t,n}^{*})\right\|^{2}\right)^{\frac{1}{2}} \\ &\leq C\Theta_{n}\sqrt{\log(n)}\left(\frac{dL}{n}\right)^{\frac{1}{4}-\frac{1}{2q}}+C\Theta\left(L^{1-\beta}+L^{-\frac{1}{2}}\right) \\ &\quad +C\Theta_{n}L^{\frac{1-\beta}{2}\vee(-\frac{1}{2})}+C\Theta_{n}n^{-\frac{1}{2}}L^{\frac{1}{2}}\Gamma_{n}^{\frac{\beta-1}{2\beta}}M^{\frac{1}{2\beta}} \\ &\leq C\Theta_{n}\Gamma_{n}^{\frac{\beta-1}{2\beta}}\sqrt{\log(n)}\left\{\left(\frac{dL}{n}\right)^{\frac{q-2}{4q}}+L^{1-\beta}+L^{-\frac{1}{2}}+\left(\frac{L}{n}\right)^{\frac{\beta-1}{2\beta}}\right\} \\ &\leq \begin{cases} C\Theta_{n}\Gamma_{n}^{\frac{\beta-1}{2\beta}}\sqrt{\log(n)}\left\{\left(\frac{dL}{n}\right)^{\frac{q-2}{4q}}+L^{-\frac{1}{2}}\right\}, \quad \beta\geq\frac{3}{2}, \ \beta>\frac{2q}{q+2}, \\ C\Theta_{n}\Gamma_{n}^{\frac{\beta-1}{2\beta}}\sqrt{\log(n)}\left\{\left(\frac{dL}{n}\right)^{\frac{\beta-1}{2\beta}}+L^{-\frac{1}{2}}\right\}, \quad \beta\geq\frac{3}{2}, \ \beta>\frac{2q}{q+2}, \\ C\Theta_{n}\Gamma_{n}^{\frac{\beta-1}{2\beta}}\sqrt{\log(n)}\left\{\left(\frac{dL}{n}\right)^{\frac{q-2}{4q}}+L^{1-\beta}\right\}, \quad \beta<\frac{3}{2}, \ \beta>\frac{2q}{q+2}, \\ C\Theta_{n}\Gamma_{n}^{\frac{\beta-1}{2\beta}}\sqrt{\log(n)}\left\{\left(\frac{dL}{n}\right)^{\frac{q-2}{2\beta}}+L^{1-\beta}\right\}, \quad \beta<\frac{3}{2}, \ \beta>\frac{2q}{q+2}. \end{split}$$

The value of L has been left unspecified and may now be chosen. In the three cases, the rate-optimal choice L^* of L is, respectively,

$$L^* \simeq \begin{cases} \left(\frac{n}{d}\right)^{\frac{q-2}{3q-2}}, & \beta \ge \frac{3}{2}, \ \beta > \frac{2q}{q+2}; \\ \left(\frac{n}{d}\right)^{\frac{\beta-1}{2\beta-1}}, & \beta \ge \frac{3}{2}, \ \beta \le \frac{2q}{q+2}; \\ \left(\frac{n}{d}\right)^{\frac{q-2}{4q\beta-3q-2}}, & \beta < \frac{3}{2}, \ \beta > \frac{2q}{q+2}; \\ \left(\frac{n}{d}\right)^{\frac{\beta-1}{2\beta^2-1-\beta}}, & \beta < \frac{3}{2}, \ \beta \le \frac{2q}{q+2}; \end{cases}$$

These choices achieve the rates (1).

Proof of Theorem 3

Denote $L = L_n$.

De-randomizing the multiplier: Assume w.l.o.g. that $X_{t,n}$ is centered, and denote $e_t = \hat{g}_{t,n} - g_{t,n}$. Decompose

$$\sum_{t=1}^{k} e_{t} X_{t,n} = \sum_{j=0}^{\lceil \frac{n}{L} \rceil - 1} \sum_{t=jL+1}^{(j+1)L \wedge k} e_{t} X_{t,n}$$

$$= \sum_{j=0}^{\lceil \frac{n}{L} \rceil - 1} \sum_{t=jL+1}^{(j+1)L \wedge k} e_{t} \widetilde{X}_{t,n} + \sum_{j=0}^{\lceil \frac{n}{L} \rceil - 1} \sum_{t=jL+1}^{(j+1)L \wedge k} e_{t} [X_{t,n} - \widetilde{X}_{t,n}], \qquad (4)$$
for $\widetilde{X}_{t,n} = G_{t,n}(\widetilde{\epsilon}_{t,t-jL}^{j}), \qquad t = jL + 1, \dots, (j+1)L.$

Here, $\boldsymbol{\epsilon}_{t,h} = (\epsilon_t, \dots, \epsilon_{t-h+1}, \tilde{\epsilon}_{t-h}^j, \tilde{\epsilon}_{t-h-1}^j, \dots)$ where $\tilde{\epsilon}_t^j \sim U(0, 1)$ are independent for all t and all j. The decomposition is chosen such that the terms $\sum_{t=jL+1}^{(j+1)L} e_t \widetilde{X}_{t,n}$ are martingale differences, and e_t and $\widetilde{X}_{t,n}$ are independent. Hence

$$\mathbb{E} \max_{r=1,\dots,\lfloor\frac{n}{L}\rfloor-1} \left\| \sum_{j=0}^{r} \sum_{t=jL+1}^{(j+1)L} e_t \widetilde{X}_{t,n} \right\|^2 = \sum_{j=0}^{\lfloor\frac{n}{L}\rfloor-1} \mathbb{E} \left\| \sum_{t=jL+1}^{(j+1)L} e_t \widetilde{X}_{t,n} \right\|^2$$
$$= \sum_{j=0}^{\lfloor\frac{n}{L}\rfloor-1} \sum_{s,t=jL+1}^{(j+1)L} \operatorname{tr} \left[\mathbb{E} \left(e_t \widetilde{X}_{t,n} \widetilde{X}_{s,n}^T e_s^T \right) \right]$$
$$= \sum_{j=0}^{\lfloor\frac{n}{L}\rfloor-1} \sum_{s,t=jL+1}^{(j+1)L} \mathbb{E} \left(\operatorname{tr} \left[e_t \widetilde{X}_{t,n} \widetilde{X}_{s,n}^T e_s^T \right] \right)$$
$$= \sum_{j=0}^{\lfloor\frac{n}{L}\rfloor-1} \sum_{s,t=jL+1}^{(j+1)L} \mathbb{E} \left(\operatorname{tr} \left[e_s^T e_t \widetilde{X}_{t,n} \widetilde{X}_{s,n}^T \right] \right)$$
$$= \sum_{j=0}^{\lfloor\frac{n}{L}\rfloor-1} \sum_{s,t=jL+1}^{(j+1)L} \operatorname{tr} \left[\mathbb{E} (e_s^T e_t) \mathbb{E} \left(\widetilde{X}_{t,n} \widetilde{X}_{s,n}^T \right) \right]$$

$$\leq \sum_{j=0}^{\lfloor \frac{n}{L} \rfloor - 1} \sum_{s,t=jL+1}^{(j+1)L} \mathbb{E} \| e_s^T e_t \|_F \cdot \left\| \mathbb{E} \left(\widetilde{X}_{t,n} \widetilde{X}_{s,n}^T \right) \right\|_F$$

$$\leq \Theta_n^2 \sum_{j=0}^{\lfloor \frac{n}{L} \rfloor - 1} \sum_{s,t=jL+1}^{(j+1)L} \| e_s \|_{\mathcal{L}_2} \| e_t \|_{\mathcal{L}_2} (|s-t|+1)^{-\beta}$$

$$\leq \Theta_n^2 \sum_{j=0}^{\lfloor \frac{n}{L} \rfloor - 1} \sum_{s,t=jL+1}^{(j+1)L} (\| e_s \|_{\mathcal{L}_2}^2 + \| e_t \|_{\mathcal{L}_2}^2) (|s-t|+1)^{-\beta}$$

$$= \Theta_n^2 \sum_{j=0}^{\lfloor \frac{n}{L} \rfloor - 1} \sum_{s,t=jL+1}^{(j+1)L} 2 \| e_s \|_{\mathcal{L}_2}^2 (|s-t|+1)^{-\beta}$$

$$\leq C_\beta \Theta_n^2 \sum_{j=0}^{\lfloor \frac{n}{L} \rfloor - 1} \sum_{s,t=jL+1}^{(j+1)L} \| e_s \|_{\mathcal{L}_2}^2 \quad (\text{because } \beta > 1)$$

$$= C_\beta \Theta_n^2 \Lambda_n^2.$$

Moreover, the block discretization error may be bounded uniformly in k as

$$\max_{k=1,\dots,n} \max_{j=0,\dots,\lceil \frac{n}{L}\rceil-1} \sum_{t=(jL+1)\lor k}^{(j+1)L\land k} e_t \widetilde{X}_{t,n} \le \Phi_n L_n \max_{t=1,\dots,n} |\widetilde{X}_{t,n}| = \mathcal{O}_P\left(\Phi_n L_n n^{\frac{1}{q}}\right).$$

Regarding the second term in (4), we find that

$$\max_{k=1,\dots,n} \left| \sum_{j=0}^{\lceil \frac{n}{B} \rceil - 1} \sum_{t=jB+1}^{(j+1)B \land k} e_t [X_{t,n} - \widetilde{X}_{t,n}] \right| \le \left[\sum_{t=1}^n e_t^2 \right]^{\frac{1}{2}} \left[\sum_{j=0}^{\lceil \frac{n}{B} \rceil - 1} \sum_{t=jB+1}^{(j+1)B \land n} [X_{t,n} - \widetilde{X}_{t,n}]^2 \right]^{\frac{1}{2}} = \mathcal{O}_P \left(\Lambda_n \right) \mathcal{O}_P \left(\sqrt{n} \Xi_n \right)$$

We have thus established that

$$\frac{1}{\sqrt{n}}\sum_{t=1}^{k}\hat{g}_{t,n}X_{t,n} = \frac{1}{\sqrt{n}}\sum_{t=1}^{k}g_{t,n}X_{t,n} + \mathcal{O}_P\left(n^{-\frac{1}{2}}\Lambda_n\Theta_n + \Lambda_n\Xi_n + n^{\frac{1}{q}-\frac{1}{2}}L_n\Phi_n\right),$$

uniformly in $k = 1, \ldots, n$.

Sequential Gaussian approximation: Denote $Z_{t,n} = H_{t,n} = g_{t,n}G_{t,n}$, such that $g_{t,n}X_{t,n} = H_{t,n}(\boldsymbol{\epsilon}_t)$, and long-run covariance matrix $\Sigma_{t,n} = \sum_{h=\infty}^{\infty} \operatorname{Cov}(H_{t,n}(\boldsymbol{\epsilon}_0), H_{t,n}(\boldsymbol{\epsilon}_h))$. Because $g_{t,n}$ is bounded, this time series satisfies $\|H_{t,n}(\boldsymbol{\epsilon}_0) - H_{t,n}(\tilde{\boldsymbol{\epsilon}}_{0,h})\|_{\mathcal{L}_q} \leq \Phi_n \Theta_n (h+1)^{-\beta}$. Moreover,

$$\sum_{t=2}^{n} \|H_{t,n}(\boldsymbol{\epsilon}_{0}) - H_{t-1,n}(\boldsymbol{\epsilon}_{0})\|_{\mathcal{L}_{2}} \leq \sum_{t=2}^{n} |g_{t,n}| \|G_{t,n} - G_{t-1,n}\|_{\mathcal{L}_{2}} + \|G_{t-1,n}\|_{\mathcal{L}_{2}} |g_{t,n} - g_{t-1,n}| \leq \Phi_{n}\Theta_{n}\Gamma_{n} + \Psi_{n}\Theta_{n}.$$

We now apply Theorem 2: Upon potentially enlarging the probability space, there exist independent Gaussian random vectors $Y_{t,n} \sim \mathcal{N}(0, \Sigma_{t,n})$ such that

$$\left\|\max_{k=1,\dots,n}\frac{1}{\sqrt{n}}\sum_{t=1}^{k}Z_{t,n}-Y_{t,n}\right\|_{\mathcal{L}_{2}} \leq C_{\beta,q}\Phi_{n}\Theta_{n}\left(\Phi_{n}\Theta_{n}\Gamma_{n}+\Psi_{n}\Theta_{n}\right)^{\frac{\beta-1}{2\beta}}\sqrt{\log(n)}\left(\frac{m}{n}\right)^{\xi(q,\beta)}$$

Proof of Lemma 4

Define the processes

$$I_n(u) = \int_0^u \Sigma_{\lfloor vn \rfloor, n}^{\frac{1}{2}} dW_v, \qquad J_n(u) = \frac{1}{\sqrt{n}} \sum_{t=1}^{\lfloor un \rfloor} Y_{t, n},$$

such that $(J_n(u))_{u \in [0,1]} \stackrel{d}{=} (I_n(\lfloor un \rfloor/n))_{u \in [0,1]}$. Because $\Sigma_{t,n}$ is bounded, we find that $\sup_{u \in [0,1]} |I_n(\lfloor un \rfloor/n) - I_n(u)| \stackrel{\mathbb{P}}{\to} 0$. The process $I_n(u)$ is fully determined via its quadratic variation $[I_n]_u = \int_0^u \Sigma_{\lfloor vn \rfloor,n} dv$. To establish the weak convergence, it suffices to show that $[I_n]_u \to \int_0^u \Sigma_v dv$ for each u. Note that this convergence will then be uniform by monotonicity of the limit, and because the $\Sigma_{t,n}$ are bounded.

To show convergence of $[I_n]_u$, we denote $\gamma_{t,n}(h) = \text{Cov}[G_{t,n}(\boldsymbol{\epsilon}_0), G_{t,n}(\boldsymbol{\epsilon}_h)]$ and $\gamma_u(h) = \text{Cov}[G_u(\boldsymbol{\epsilon}_0), G_u(\boldsymbol{\epsilon}_h)]$, and observe that

$$\begin{split} \|\Sigma_{t,n} - \Sigma_v\| &\leq \sum_{h=-\infty}^{\infty} \|g_{t,n} - g_v\| \|\gamma_{t,n}(h)\| \|g_{t,n}\| \\ &+ \sum_{h=-\infty}^{\infty} \|g_v\| \|\gamma_{t,n}(h) - \gamma_v(h)\| \|g_{t,n}\| \\ &+ \sum_{h=-\infty}^{\infty} \|g_v\| \|\gamma_{t,n}(h)\| \|g_{t,n} - g_v\| \\ &\leq C \|g_{t,n} - g_v\| \sum_{h=-\infty}^{\infty} (|h|+1)^{-\beta} + C \sum_{h=-\infty}^{\infty} \|G_{t,n} - G_v\|_{\mathcal{L}_2} \wedge (|h|+1)^{-\beta}. \end{split}$$

Thus,

$$\begin{split} &\int_0^1 \left\| \Sigma_{\lfloor vn \rfloor, n} - \Sigma_v \right\| dv \\ &\leq C \int_0^1 \left\| g_{\lfloor vn \rfloor, n} - g_v \right\| dv + \sum_{h = -\infty}^\infty \left[\int_0^1 \left\| G_{\lfloor vn \rfloor, n} - G_v \right\|_{\mathcal{L}_2} dv \wedge (|h| + 1)^{-\beta} \right]. \end{split}$$

By virtue of assumptions (A.3) and (A.4), and by dominated convergence, the latter term tends to zero as $n \to \infty$, thus completing the proof.

Proof of Theorem 6

Because of the Lipschitz-continuity of Σ , $X_{t,n}$ satisfies (A.1) and (A.2) for any $\beta > 1$, with $\Gamma_n = 1$ and $\Theta_n = \Theta(\beta) = \mathcal{O}(1)$. Assumption (A.3) also holds, with $G_u(\epsilon_t) = \Sigma(u)^{\frac{1}{2}} \eta_t$.

The multipliers are given by $\widehat{g}_{t,n} = \widehat{\Sigma}(t/n)^{-\frac{1}{2}}$, and we define $g_u = \Sigma(u)^{-\frac{1}{2}}$ and

$$\begin{cases} \Sigma(t/n)^{-\frac{1}{2}}, & t = k_n + 1, \dots, n, \\ cI_{d \times d}, & t = 1, \dots, k_n, \end{cases}$$

Then $\Psi_n, \Phi_n = \mathcal{O}(1)$, (A.4) holds, and (A.M) with $L_n = 1$ and $\Theta_n = 0$, due to independence of the $X_{t,n}$.

It remains to compute Λ_n . A standard bias-variance decomposition exploiting the Lipschitz continuity of $\Sigma(u)$ yields, for $t = k_n + 1, \ldots, n$,

$$\mathbb{E}\|\widetilde{\Sigma}(t/n) - \Sigma(t/n)\|^2 \leq C\left(\left(\frac{k_n}{n}\right)^2 + \frac{1}{k_n}\right) \leq Cn^{-\frac{2}{3}},$$

for $k_n \simeq n^{-\frac{1}{3}}$ and a constant C changing from line to line, but depending on neither n nor t. Because the mapping $A \mapsto A^{-\frac{1}{2}}$ is Lipschitz-continuous on the set $\{A \succeq cI_{d \times d}\}$, we readily find that $\mathbb{E}\|\widehat{g}_{t,n} - g_{t,n}\|^2 \leq Cn^{-\frac{2}{3}}$ for all $t = 1, \ldots, n$. Thus, $\Lambda_n = \mathcal{O}(n^{\frac{1}{6}})$, which shows that 5 is applicable.

References

- Berkes, I., Liu, W., and Wu, W. B. (2014). Komlós-Major-Tusnády approximation under dependence. Annals of Probability, 42(2):794–817.
- Billingsley, P. (1999). Convergence of Probability Measures. John Wiley & Sons.
- Bonnerjee, S., Karmakar, S., and Wu, W. B. (2024). Gaussian Approximation For Non-stationary Time Series with Optimal Rate and Explicit Construction. <u>Annals of</u> <u>Statistics</u>, 52(5):2293–2317.
- Dahlhaus, R. (1997). Fitting time series models to nonstationary processes. <u>Annals of</u> Statistics, 25(1):1–37.
- Dahlhaus, R. and Polonik, W. (2009). Empirical spectral processes for locally stationary time series. Bernoulli, 15(1):1–39.
- Dahlhaus, R., Richter, S., and Wu, W. B. (2019). Towards a general theory for nonlinear locally stationary processes. Bernoulli, 25(2):1013–1044.
- Giraud, C., Roueff, F., and Sanchez-Perez, A. (2015). Aggregation of predictors for nonstationary sub-linear processes and online adaptive forecasting of time varying autoregressive processes. <u>The Annals of Statistics</u>, 43(6):2412–2450.

- Grenier, Y. (1983). Time-dependent ARMA modeling of nonstationary signals. <u>IEEE</u> Transactions on Acoustics, Speech, and Signal Processing, 31(4):899–911.
- Karmakar, S. and Wu, W. B. (2020). Optimal Gaussian Approximation For Multiple Time Series. Statistica Sinica, 30:1399–1417.
- Komlós, J., Major, P., and Tusnády, G. (1975). An approximation of partial sums of independent RV'-s, and the sample DF. I. <u>Zeitschrift für Wahrscheinlichkeitstheorie</u> und Verwandte Gebiete, 32(1-2):111–131.
- Komlós, J., Major, P., and Tusnády, G. (1976). An approximation of partial sums of independent R.V.'s and the sample DF. II. <u>Zeitschrift für Wahrscheinlichkeitstheorie</u> und Verwandte Gebiete, 34:33–58.
- Liu, W. and Lin, Z. (2009). Strong approximation for a class of stationary processes. Stochastic Processes and their Applications, 119:249–280.
- Mies, F. (2023). Functional estimation and change detection for nonstationary time series. Journal of the American Statistical Association, 118(542):1011–1022.
- Mies, F. and Steland, A. (2023). Sequential Gaussian approximation for nonstationary time series in high dimensions. Bernoulli, 29(4):3114–3140.
- Mies, F. and Steland, A. (2024). Projection inference for high-dimensional covariance matrices with structured shrinkage targets. <u>Electronic Journal of Statistics</u>, 18(1):1643–1676.
- Moulines, E., Priouret, P., and Roueff, F. (2005). On recursive estimation for time varying autoregressive processes. The Annals of Statistics, 33(6).
- Subba Rao, T. (1970). The Fitting of Non-stationary Time-series Models with Timedependent Parameters. Journal of the Royal Statistical Society. Series B (Statistical Methodology), 32(2):312–322.
- van der Vaart, A. W. and Wellner, J. A. (1996). Weak Convergence and Empirical Processes With Applications to Statistics.
- Wu, W. B. (2005). Nonlinear system theory: Another look at dependence. <u>Proceedings</u> of the National Academy of Sciences, 102(40):14150–14154.
- Wu, W. B. and Zhou, Z. (2011). Gaussian approximations for non-stationary multiple time series. Statistica Sinica, 21(3):1397–1413.
- Zaitsev, A. Y. (2013). The accuracy of strong Gaussian approximation for sums of independent random vectors. Russian Mathematical Surveys, 68(4):721–761.