# CHARACTERIZATION OF MINIMAL TRIPOTENTS VIA ANNIHILATORS AND ITS APPLICATION TO THE STUDY OF ADDITIVE PRESERVERS OF TRUNCATIONS

LEI LI, SIYU LIU, AND ANTONIO M. PERALTA

Abstract. The contributions in this note begin with a new characterization of (positive) scalar multiples of minimal tripotents in a general  $JB^*$ -triple  $E$ , proving that a non-zero element  $a \in E$  is a positive scalar multiple of a minimal tripotent in  $E$  if, and only if, its inner quadratic annihilator (that is, the set  $\pm q$  {a} = {b ∈ E : {a, b, a} = 0}) is maximal among all inner quadratic annihilators of single elements in  $E$ . We subsequently apply this characterization to the study of surjective additive maps between atomic JBW<sup>∗</sup> -triples preserving truncations in both directions. Let  $A : E \to F$  be a surjective additive mapping between atomic JBW<sup>\*</sup>-triples, where  $E$  contains no one-dimensional Cartan factors as direct summands. We show that A preserves truncations in both directions if, and only if, there exists a bijection  $\sigma : \Gamma_1 \to \Gamma_2$ , a bounded family  $(\gamma_k)_{k \in \Gamma_1} \subseteq \mathbb{R}^+$ , and a family  $(\Phi_k)_{k \in \Gamma_1}$ , where each  $\Phi_k$  is a (complex) linear or a conjugate-linear (isometric) triple isomorphism from  $C_k$  onto  $C_{\sigma(k)}$ satisfying  $\inf_k{\gamma_k} > 0$ , and

$$
A(x) = \left(\gamma_k \Phi_k \left(\pi_k(x)\right)\right)_{k \in \Gamma_1}, \text{ for all } x \in E,
$$

where  $\pi_k$  denotes the canonical projection of E onto  $C_k$ .

#### 1. INTRODUCTION

Preservers of truncations are currently studied in different settings, from preservers on the space  $B(H)$ , of all bounded linear operators on a Hilbert space H (cf.  $[31, 40, 45]$  $[31, 40, 45]$  $[31, 40, 45]$  $[31, 40, 45]$ ), to the wider setting of Cartan factors and general JB<sup>\*</sup>-triples (see [\[27\]](#page-30-1)). We recall that, for  $a, b \in B(H)$ , we say that a is a truncation of b if  $aa^*a = ab^*a$ . It is widely known that every C<sup>\*</sup>-algebra, A, is a JB<sup>\*</sup>-triple with respect to the triple product

<span id="page-0-0"></span>
$$
\{a, b, c\} := \frac{1}{2}(ab^*c + cb^*a) \quad (a, b, c \in A).
$$
 (1)

Furthermore, in terms of the just defined triple product,  $a$  is a truncation of  $b$  if and only if  $\{a, a, a\} = \{a, b, a\}$ . So, the relation "being a truncation of" fully makes sense in the wider setting of JB<sup>\*</sup>-triples (see Section [2](#page-3-0) for the detailed definitions).

A very recent study shows that if  $\Delta: A \to B$  is a (non-necessarily linear nor continuous) bijection between atomic JBW<sup>∗</sup> -triples preserving the truncation of triple products in both directions, that is,

a is a truncation of 
$$
\{b, c, b\}
$$
  $\Leftrightarrow$   $\Delta(a)$  is a truncation of  $\{\Delta(b), \Delta(c), \Delta(b)\}$ ,

<sup>2010</sup> Mathematics Subject Classification. Primary 47B49 Secondary 46C99, 17C65, 47B48, 47C15, 46H40.

Key words and phrases. Cartan factor, JB∗-triple, truncation, tripotents, triple isomorphisms, preservers.

and satisfying its restriction to each rank-one Cartan factor in  $A$ , if any, is a continuous mapping, then  $\Delta$  is an isometric real-linear triple isomorphism [\[27\]](#page-30-1). A related precedent in the particular case that  $A = B = B(H)$  was considered by X. Jia, W. Shi, and G. Ji in [\[31\]](#page-30-0).

Just like the studies on (non-necessarily linear) bijections preserving  $\lambda$ -Aluthge transforms on products in  $[11, 12, 18]$  $[11, 12, 18]$  $[11, 12, 18]$  $[11, 12, 18]$  was preceded by the description of the bijective linear transformations between von Neumann factors that commute with the  $\lambda$ -Aluthge transform in [\[5\]](#page-29-0), it seems natural to pose the challenge of describing all linear surjection between (atomic) JB<sup>∗</sup> -triples preserving truncations in both directions. Observe that this problem is independent from the commented result on bijections preserving truncations of triple products in both directions. Moreover, it could be also asked whether the linearity of the mapping is absolutely necessary to obtain the description. Concerning this question, in [\[45,](#page-31-1) Theorem 2.3], J. Yao and G. Ji. prove that for each complex Hilbert space H with  $\dim(H) \geq 2$ , the following statements are equivalent for every additive and surjective map  $A: B(H) \to B(H)$ :

- (a) A preserves truncations of operators in both directions;
- (b) There exist a nonzero scalar  $\alpha \in \mathbb{C}$  and maps  $u, v : H \to H$  which are both unitary operators or both anti-unitary operators such that  $A(x) = \alpha u x v$  for all  $x \in B(H)$  or  $A(x) = \alpha u x^* v$  for all  $x \in B(H)$ .

Observe that statement  $(b)$  above is equivalent to say that  $A$  is a positive multiple of a (complex) linear or a conjugate-linear triple automorphism on  $B(H)$ .

As commented before  $B(H)$  is an example of C<sup>\*</sup>-algebra, and a JB<sup>\*</sup>-triple. It can be also regarded inside the collection of complex Banach spaces called Cartan factors. We recall their definition with the aim to provide some basic background on Cartan factors for the readers. Let  $H$  and  $K$  be complex Hilbert spaces. The spaces  $B(H, K)$  of all bounded linear operators between H and K form the socalled *Cartan factors of type* 1. Suppose next that  $j : H \to H$  is a conjugation on H (i.e., a conjugate-linear isometry of period 2). The subspaces of  $B(H)$  given by  $C_2 = \{a \in B(H) : a = -ja * j\}$  and  $C_3 = \{a \in B(H) : a = ja * j\}$  determine the classes of *Cartan factors of type* 2 and 3, respectively. All these spaces are equipped with the triple product given in [\(1\)](#page-0-0). A Banach space V is called a *Cartan factor of type* 4 or a *spin factor* if it admits a complete inner product  $\langle \cdot | \cdot \rangle$  and a conjugation  $x \mapsto \bar{x}$ , for which the norm of V is given by

$$
||x||^2 = \langle x|x\rangle + \sqrt{\langle x|x\rangle^2 - |\langle x|\bar{x}\rangle|^2}
$$

and the triple product of V is defined by

$$
\{x, y, z\} = \langle x|y\rangle z + \langle z|z\rangle x - \langle x|\overline{z}\rangle \overline{y}, \quad (x, y, z \in V).
$$

The *Cartan factor of type* 5 *and* 6 (also called *exceptional Cartan factors*) are the finite dimensional Banach spaces  $M_{1,2}(\mathbb{O})$  and  $H_3(\mathbb{O})$ , of all  $1 \times 2$  matrices with entries in the (complex) octonions  $\mathbb{O}$ , and all  $3 \times 3$  hermitian matrices with entries in  $\mathbb{O}$ , respectively (see [\[28\]](#page-30-5) for more details).

In our main result in this note we study surjective additive mappings preserving truncations in both directions between JB<sup>∗</sup>-triples which can be written as  $\ell_{\infty}$ -sums of families of Cartan factors. Our main conclusion, established in Theorem [5.15,](#page-28-0) reads as follows: Let  $E = \bigoplus^{\ell_{\infty}}$  $k \in \Gamma_1$  $C_k$  and  $F = \bigoplus^{\ell_{\infty}}$  $j \in \Gamma_2$  $C_j$  be atomic JBW<sup>\*</sup>-triples, where

 ${C_k}_{k \in \Gamma_1}$  and  ${\tilde{C}_j}_{j \in \Gamma_2}$  are two families of Cartan factors. Suppose, additionally, that  $E$  contains no one-dimensional Cartan factors as direct summands (i.e.,  $\dim(C_k) \geq 2$  for all k). Let  $A : E \to F$  be a surjective additive mapping. Then the following statements are equivalent:

- (a) A preserves truncations in both directions.
- (b) There exists a bijection  $\sigma : \Gamma_1 \to \Gamma_2$ , a bounded family  $(\gamma_k)_{k \in \Gamma_1} \subseteq \mathbb{R}^+$ , and a family  $(\Phi_k)_{k \in \Gamma_1}$ , where each  $\Phi_k$  is a (complex) linear or a conjugate-linear (isometric) triple isomorphism from  $C_k$  onto  $C_{\sigma(k)}$  satisfying  $\inf_k \{ \gamma_k \} > 0$ , and

$$
A(x) = \left(\gamma_k \Phi_k \left(\pi_k(x)\right)\right)_{k \in \Gamma_1}, \text{ for all } x \in E,
$$

where  $\pi_k$  denotes the canonical projection of E onto  $C_k$ .

In particular, every surjective additive mapping  $A: E \to F$  preserving truncations in both directions is a real-linear bijection, and the  $JB^*$ -triples  $E$  and  $F$  are isometrically triple isomorphic. Moreover, if  $A$  satisfies  $(a)$  or  $(b)$  and there exists an element  $x_0$  in E satisfying  $||A\pi_k(x_0)|| = ||\pi_k(x_0)|| \neq 0$ , for all  $k \in \Gamma_1$ , the mapping A is a real-linear (isometric) triple isomorphism. Note that the implication  $(b) \Rightarrow (a)$  is clear. We shall see in Example [3.2](#page-8-0) that the hypothesis affirming that  $\dim(C_k) \geq 2$  for all k cannot be relaxed.

It is known that a JB<sup>∗</sup> -triple has rank-one if and only if it is a complex Hilbert space regarded as a type 1 Cartan factor (cf. Section [3\)](#page-7-0). Therefore, every additive mapping A between two rank-one JB<sup>∗</sup> -triples is nothing but an additive mapping between two Hilbert spaces. We shall see below that, in such a case, A preserves truncations if, and only if, it preserves Euclidean orthogonality between elements in the corresponding Hilbert spaces (cf. Remark [3.1\)](#page-7-1). We are thus in a position to apply a recent result on additive preservers of Euclidean orthogonality between complex Hilbert spaces, obtained in  $[37]$ , to deduce that if A is a surjective additive mapping between rank-one JB<sup>\*</sup>-triples of dimension  $\geq 2$ , and A preserves truncations (in one direction), then  $A$  is a positive scalar multiple of a (complex) linear or a conjugate-linear isometry (see Proposition [3.3\)](#page-8-1).

The proof of the main result in this note has also required the development of a tool of independent interest. In Theorem [2.4](#page-5-0) we establish that a non-zero element a in a JB<sup>\*</sup>-triple  $E$  is a (positive) scalar multiple of a minimal tripotent in  $E$ , if and only if, its inner quadratic annihilator  $\perp^q \{a\} = \{b \in E : \{a, b, a\} = 0\}$  is maximal among all inner quadratic annihilators of single elements in E. This result generalises a previous characterization of rank-one operators in  $B(H)$  obtained by J. Yao and G. Ji in [\[45,](#page-31-1) Lemma 2.2].

We have essentially described the main results in sections 2 and 3. Section [4](#page-9-0) is devoted to analyse the basic properties of the surjective additive maps preserving truncations in both directions between two general  $JB^*$ -triples  $E$  and  $F$ . It is shown that any such a mapping must be automatically injective and maps scalar multiples of minimal tripotents in  $E$  (if any) to scalar multiples of minimal tripotents in  $F$ . Moreover, for each minimal tripotent e in E we have  $A(\mathbb{C}e) = \mathbb{C}A(e) = \mathbb{C}r(A(e)),$ where  $r(A(e))$  denotes the range tripotent of  $A(e)$ , which in this case lies in F (see Lemma [4.1\)](#page-9-1). Further technical conclusions assure that if  $w_1$  and  $w_2$  are two collinear (respectively, orthogonal) minimal tripotents in  $E$ , the corresponding range tripotents  $r(A(w_1))$  and  $r(A(w_2))$  are minimal and collinear (respectively, orthogonal) in  $F$  (cf. Propositions [4.5](#page-14-0) and [4.7\)](#page-14-1).

The concluding section is devoted to establish our main result. We prove that if  $A: E \to F$  is a surjective additive mapping preserving truncations in both directions between two atomic JBW<sup>\*</sup>-triples, where  $E$  contains no one-dimensional Cartan factors as direct summands, the mapping  $A_r : e \mapsto r(A(r))$ , sending each minimal tripotent in E to the range tripotent of  $A(e)$  in F is a bijective map preserving collinearity and orthogonality in both directions from the set  $\mathcal{U}_{min}(E)$  of all minimal tripotents in E onto the corresponding set  $\mathcal{U}_{min}(F)$  (see Proposition [5.8\)](#page-22-0). Furthermore, when restricted to each Cartan factor  $C_k$  in the decomposition of E, the mapping  $A_r$  either preserves or reverses triple transition pseudo-probabilities between all minimal tripotents in  $C_k$ , actually  $A|_{C_k}$  must be a positive scalar multiple of a linear or conjugate-linear triple isomorphism from  $C_k$  onto a Cartan factor in the decomposition of  $F$  (see Proposition [5.14\)](#page-26-0). We can finally apply the description of all bijective maps preserving triple transition pseudo-probabilities between the sets of minimal tripotents in two atomic JBW<sup>∗</sup> -triples borrowed from [\[44\]](#page-31-2) to complete the proof of our main result.

### 2. Quadratic annihilators and minimal tripotents

<span id="page-3-0"></span>It is perhaps worth to begin this section by recalling the definition of JB<sup>∗</sup> -triples. A *JB*<sup>∗</sup> *-triple* (see [\[34\]](#page-30-7)) is a Banach space together with a continuous triple product

$$
\{\cdot, \cdot, \cdot\} : E \times E \times E \to E : (x, y, z) \mapsto \{x, y, z\}
$$

for all  $x, y, z \in E$ , which is linear in the first and third variables, conjugate-linear in the middle one, and satisfies the following axioms:

(i) For each a, b in E, the operators  $L(a, b) : E \to E$ ,  $L(a, b)(c) := \{a, b, c\}$  satisfy the identity

$$
L(w, v){x, y, z} = {L(w, v)x, y, z} - {x, L(v, w)y, z} + {x, y, L(w, v)z},
$$

for all  $x, y, z, w, v \in E$ ; (*Jordan identity*)

(ii) For each  $x \in E$ , the operator  $L(x, x)$  is hermitian with non-negative spectrum;  $(iii)$   $\|\{x, x, x\}\| = \|x\|^3$ for all x ∈ E. (*Gelfand-Naimark axiom*) All Cartan factors are JB<sup>∗</sup> -triples.

A JBW<sup>∗</sup> -triple is a JB<sup>∗</sup> -triple whose underlying Banach space is a dual Banach space. Every JBW<sup>\*</sup>-triple admits an unique (isometric) predual and its triple product is separately continuous [\[2\]](#page-29-1). It is worth to note that all von Neumann algebras are examples of JBW<sup>\*</sup>-triples, and the bidual,  $E^{**}$ , of a JB<sup>\*</sup>-triple E is a JBW<sup>\*</sup>triple [\[16\]](#page-30-8). JBW<sup>∗</sup> -triples satisfy extra geometric algebraic properties, for example, they contain an abundant collection of tripotents; since tripotents in a JB<sup>∗</sup> -triple  $E$  coincide with the extreme points of the closed unit ball of  $E$  (see, for example, [\[17,](#page-30-9) Corollary 4.8] or [\[10,](#page-30-10) Theorem 4.2.34]). Note that the set of all tripotents in a general JB<sup>∗</sup> -triple might be empty.

The relation "being a truncation of" between elements in a JB<sup>∗</sup> -triple can be better understood via inner quadratic annihilators. The *inner quadratic annihilator* of a subset S of a JB<sup>\*</sup>-triple E is defined as the set

$$
{}^{\perp_q}S = \{b \in E: Q(s)(b) = 0, \ \forall \, s \in S\} = \bigcap_{s \in S} \ker(Q(s)).
$$

We note that, given an element s in a JB<sup>\*</sup>-triple E, we denote by  $Q(s)$  the conjugatelinear mapping on E given by  $Q(s)(x) = \{s, x, s\}$ . Observe that <sup>⊥</sup><sup>q</sup>S is a closed linear subspace of E since  $Q(s)$  is a bounded conjugate-linear operator on E for every s. It is known that for each element  $a$  in  $E$  we have

<span id="page-4-0"></span>
$$
{}^{\perp_q}\{a\} = E \cap \left(E_0^{**}(r(a)) \oplus E_1^{**}(r(a))\right) = \{x \in E : P_2(r(a))(x) = 0\},\qquad(2)
$$

where  $r(a)$  denotes the range tripotent of a in  $E^{**}$ , and  $E_j(e)$  stands for the Peirce-j subspace associated to e, and  $P_i(e)$  stands for the projection of E onto  $E_i(e)$  (cf. [ $27$ , Lemma 2.2]). In case that E is a JBW<sup>\*</sup>-triple, we can further conclude that

$$
{}^{\perp_q}{a} = E_0(r(a)) \oplus E_1(r(a)) = \{x \in E : P_2(r(a))(x) = 0\},\
$$

where  $r(a)$  denotes the range tripotent of a in E.

Some of the notions employed in the previous paragraph are perhaps less known for non-experts. The missing details will be fully explained now. For example, an element e in a JB<sup>\*</sup>-triple E is called a *tripotent* if  $\{e, e, e\} = e$ . In such a case, we can always E according to the so-called *Peirce decomposition* induced by e in the form

$$
E = E_0(e) \oplus E_1(e) \oplus E_2(e), \tag{3}
$$

where each  $E_j(e) := \{a \in E : L(e,e)(a) = \frac{j}{2}a\}$  is a subtriple of E called the *Peirce* j-subspace  $(j = 0, 1, 2)$ . Triple products among elements in Peirce subspaces follow certain patterns known as *Peirce rules* or *Peirce arithmetics*, namely,  ${E_i(e), E_j(e), E_k(e)} \subseteq E_{i-j+k}(e)$  for all  $i, j, k = \{0, 1, 2\}$  and

$$
{E_0(e), E_2(e), E} = {E_2(e), E_0(e), E} = 0,
$$

where  $E_{i-j+k}(e) = \{0\}$  if  $i-j+k \neq \{0,1,2\}$ . The Peirce 2-subspace  $E_2(e)$  is actually a unital JB<sup>∗</sup> -algebra with identity e, where the Jordan product and involution operations are given by  $a \circ_e b := \{a, e, b\}$  and  $a^{*_e} := \{e, a, e\}$ , respectively (cf. [\[29\]](#page-30-11) or  $[10,$  Fact 4.2.14, Proposition 4.2.22, and Corollary 4.2.30]). An illustrative example is given by  $C^*$ -algebras (and hence by  $B(H)$  spaces), regarded as JB<sup>\*</sup>-triples with respect to the triple product in  $(1)$ , where tripotents correspond to partial isometries.

We can classify tripotents in terms of the summands in the corresponding Peirce decomposition. For example, a tripotent e is called (*algebraically*) *minimal* (respectively, *complete* or *maximal*) if  $E_2(e) = \mathbb{C}e \neq \{0\}$  (respectively,  $E_0(e) = \{0\}$ ). We shall write  $\mathcal{U}(E)$ ,  $\mathcal{U}_{min}(E)$  and  $\mathcal{U}_{max}(E)$  for the sets of all tripotents, all minimal tripotents, and maximal tripotents in  $E$ , respectively. We employed the word "algebraic" because there is also a notion of minimality associated with the partial ordering on tripotents.

<span id="page-4-1"></span>The characterization of the inner quadratic annihilator in [\(2\)](#page-4-0) can be now applied to deduce the following result.

Lemma 2.1. *Let* e, v *be two minimal tripotents in a JB*<sup>∗</sup> *-triple* E*, and let us pick two non-zero elements*  $a \in \mathbb{C}e$  *and*  $b \in \mathbb{C}v$ *. Then* a *and* b *are linearly dependent if,* and only if,  $\perp_q \{a\} = \perp_q \{b\}.$ 

Elements  $x, y$  in a JB<sup>\*</sup>-triple E are said to be *orthogonal*  $(x \perp y$  in short) if  $L(x, y) = 0$  (see, for example, Lemma 1 and the comments around it in [\[8\]](#page-29-2) for additional properties). When the relation of orthogonality is restricted to  $\mathcal{U}(E)$ , it is known that  $w, v \in \mathcal{U}(E)$ , are orthogonal if, and only if,  $w \in E_0(v)$ . It is known that  $a \perp b$  in E implies that they are M-orthogonal, that is,  $||a+b|| = \max{||a||, ||b||}$ (see [\[25,](#page-30-12) Lemma  $1.3(a)$ ]).

Let us recall that two tripotents  $w_1, w_2$  in a JB<sup>\*</sup>-triple E are called *collinear*  $(w_1\top w_2$  in short) if  $w_i \in E_1(w_j)$  for all  $i \neq j$  in  $\{1,2\}$ . We say that  $w_2$  governs  $w_1$  $(w_2 \vdash w_1 \text{ in short})$  if  $w_1 \in E_2(w_2)$  and  $w_2 \in E_1(w_1)$ .

The natural partial ordering on  $\mathcal{U}(E)$  is defined as follows: for  $e, u \in \mathcal{U}(E)$  the symbol  $e \leq u$  means that  $u-e \in \mathcal{U}(E)$  and  $u-e \perp e$ , or equivalently, e is a projection in the unital JB<sup>\*</sup>-algebra  $E_2(u)$ . We note that these relations of orthogonality and partial ordering among elements in  $\mathcal{U}(E)$  agree with the original notions of orthogonality and order for C<sup>∗</sup> -algebras in that particular setting. Clearly, every (algebraic) minimal tripotent is order minimal, however the reciprocal statement is not necessarily true as shown by the unit element in  $C[0, 1]$ . Under the stronger assumption that  $E$  is a JBW<sup>\*</sup>-triple minimal and order minimal tripotents agree (cf.  $[17, Corollary 4.8]$  and  $[3, Lemma 4.7]$ ).

For each element  $a$  in a JBW<sup>\*</sup>-triple  $E$ , there exists a smallest tripotent  $e$  in E (called the *range tripotent* of a) satisfying that a is a positive element in the JBW<sup>\*</sup>-algebra  $E_2(e)$ . We write  $r(a)$  for the range tripotent of a. It is known that  $r(a)$  also coincides the range projection of a in  $E_2(r(a))$ .

<span id="page-5-1"></span>**Remark 2.2.** It is known from [\[27,](#page-30-1) Lemma 2.4] that an element  $a$  in a JB<sup>\*</sup>-triple  $E$  is a truncation of another element b in E if, and only if, one of the following equivalent statements holds:

(a)  $b = a + z$  for some  $z \in E$  with  $\{a, z, a\} = 0$  (i.e.  $z \in {}^{\perp_q}(a)$ );

(b)  $a = P_2(r(a))(b)$  where  $r(a)$  is the range tripotent of a in  $E^{**}$ .

<span id="page-5-2"></span>The following lemma is an obvious consequence of the characterization given in Remark [2.2.](#page-5-1)

**Lemma 2.3.** Let  $A: E \to F$  be an additive mapping between  $JB^*$ -triples. Then A *preserves truncations if, and only if, it preserves inner quadratic annihilators.*

Preservers of rank-one operators on subalgebras of the space  $B(X)$ , of all bounded linear operators on a nontrivial real or complex Banach space  $X$ , have been intensively studied, and we have (see, for example, [\[42\]](#page-31-3) and the historical review therein).

We recall that a JBW<sup>\*</sup>-triple is called *atomic* if it coincides with the weak<sup>\*</sup>closure of the linear span of all its minimal tripotents  $(cf.[25, 26])$  $(cf.[25, 26])$  $(cf.[25, 26])$  $(cf.[25, 26])$  $(cf.[25, 26])$ . Atomic JB<sup>\*</sup>triple can be concretely represented as  $\ell_{\infty}$ -sums of families of Cartan factors [\[26,](#page-30-13) Proposition 2 and Theorem E].

In  $B(X)$  the notion of rank-one operator is quite natural and needs no extra explanation. But, do we have an equivalent notion in more general structures? In [\[45,](#page-31-1) Lemma 2.2] J. Yao and G. Ji established that a non-zero operator  $a$  in  $B(H)$ is a rank-one operator if, and only if,  $\perp^q \{a\} = \{x \in B(H) : ax^*a = 0\}$  is maximal among all inner quadratic annihilators of elements in  $B(H)$ . It is natural to ask whether a similar conclusion holds when  $B(H)$  is replaced with a von Neumann algebra, a C<sup>∗</sup> -algebra or a general JB<sup>∗</sup> -triple. Our first main result presents an argument to solve this question.

<span id="page-5-0"></span>Theorem 2.4. *Let* a *be a non-zero element in a JB*<sup>∗</sup> *-triple* E*. Then the following statements are equivalent.*

- (a) a *is a* (*positive*) *scalar multiple of a minimal tripotent in* E*.*
- (b) The inner quadratic annihilator of a,  $\perp^q \{a\}$ , is a maximal element in the set *of all inner quadratic annihilators of single elements in* E*.*

*Proof.* (a)  $\Rightarrow$  (b) Suppose  $a = \lambda e$ , where  $\lambda \in \mathbb{C} \setminus \{0\}$  and e is a minimal tripotent. It is easy to see that  $\perp^q \{a\} = \perp^q \{e\} = E_1(e) \oplus E_0(e)$  (cf. [\(2\)](#page-4-0), for latter equality). Since  $E = E_2(e) \oplus E_1(e) \oplus E_0(e)$  with  $E_2(e) = \mathbb{C}e$ , it follows that  $\perp_q \{a\}$  is a onecodimensional closed subspace of  $E$ , and hence maximal among subspaces. Having in mind that the inner quadratic annihilator associated to each subset of  $E$  is a closed linear subspace, we deduce the desired property for  $\perp_q \{a\}$ .

 $(b) \Rightarrow (a)$  As we shall see now, it is relatively easy to prove that a is a positive scalar multiple of a tripotent, but showing that this tripotent is in fact minimal in  $E$ is more difficult. By local theory, the JB<sup>\*</sup>-subtriple of  $E$  generated by the element a, denoted by  $E_a$ , is (isometrically) JB<sup>\*</sup>-triple isomorphic to a commutative C<sup>\*</sup>algebra of the form  $C_0(\Omega)$  for some locally compact Hausdorff space  $\Omega \subseteq (0, ||a||)$ , such that  $\Omega \cup \{0\}$  is compact, and under the corresponding identification a is identified with the inclusion mapping of  $\Omega$  into C (cf. [\[33,](#page-30-14) Corollary 4.8], [\[34,](#page-30-7) Corollary 1.15] and [\[23,](#page-30-15) [24\]](#page-30-16)). It is also known that the range tripotent of a in  $E^{**}$  is a unit element in  $E_a^{**}$ . If  $\Omega$  contains more than one point, we can take two orthogonal non-zero elements  $c, b$  in  $E_a \cong C_0(\Omega)$  (positive in the local order) whose range tripotents satisfy  $r(b), r(c) \leq r(a)$  in  $E_a^{**}$ , and also as tripotents in  $E^{**}$ . It follows from the inequality  $r(b) \leq r(a)$  and [\(2\)](#page-4-0) that  $\perp^q a$   $\subseteq \perp^q b$ . Since  $c \perp b$  we have  $c \in {}^{\perp_q}{\{b\}},$  and  $\{a, c, a\} \neq 0$  gives  $c \notin {}^{\perp_q}{\{a\}}$ . Therefore  ${}^{\perp_q}{\{a\}} \subsetneqq {}^{\perp_q}{\{b\}},$  which contradicts the maximality of  $\perp^q \{a\}$ . Therefore  $\Omega$  reduces to a single element, and hence a is a positive scalar multiple of a tripotent  $e$  in  $E_a$ , and clearly a tripotent in E. We have thus proved that  $a = \lambda e$ , for  $\lambda \in \mathbb{R}^+$  and a tripotent e in E.

Our next goal will consist in proving that  $e$  is a minimal tripotent in  $E$ . We recall that we can decompose  $E^{**}$  as the direct sum of two weak<sup>\*</sup>-closed ideal A and N, where A, called the atomic part of  $E^{**}$ , is generated by all minimal tripotent in  $E^{**}$ , while N contains no minimal tripotents (cf.  $[25,$  Theorem 2]). It is also known that if  $\pi : E^{**} \to A$  denotes the canonical projection of  $E^{**}$  onto A, and  $\iota_E : E \hookrightarrow E^{**}$  stands for the canonical inclusion, the mapping  $\Phi = \pi \circ \iota_E : E \to A$ is an isometric triple monomorphism with weak<sup>∗</sup> -dense range (see [\[26,](#page-30-13) Proposition 1 and its proof]). We further know from the just quoted reference that A is an atomic JBW<sup>\*</sup>-triple, that is, it can be written as an  $\ell_{\infty}$ -sum of Cartan factors (cf.  $[26,$  Proposition 2 and Theorem E]). It is clear that in order to prove that e is minimal in E it suffices to show that  $\Phi(e)$  is minimal in A.

Suppose that  $\Phi(e)$  is not minimal in A. Since the latter is an atomic JBW<sup>\*</sup>-triple, we can find two orthogonal minimal tripotents  $v_1, v_2$  in A such that  $\Phi(e) \geq v_1, v_2$ in A. Since  $e \in E$ , it is easy to check, from the weak<sup>\*</sup>-density of  $\Phi(E)$  in E and the weak<sup>\*</sup>-continuity of Peirce projections, that  $\Phi(E)_2(\Phi(e)) = A_2(e) \cap \Phi(E)$ is weak<sup>\*</sup>-dense in  $A_2(e)$ . Now, Kadison's transitivity theorem for JB<sup>\*</sup>-triples ([\[7,](#page-29-4) Theorem 3.3] or [\[19,](#page-30-17) Corollary 1.3] and the explanation in [\[20,](#page-30-18) §3] for completeness), applied to A and  $\Phi(E)$ , implies the existence of two norm-one pairwise orthogonal elements  $\Phi(b), \Phi(c)$  in  $\Phi(E)_2(\Phi(e))$  such that  $\Phi(b) = v_1 + P_0(v_1)(\Phi(b))$  and  $\Phi(c) =$  $v_2 + P_0(v_2)(\Phi(c))$ . Since  $\Phi(b) \in \Phi(E)_2(\Phi(e))$  it can be easily deduced from Peirce arithmetic that  $\perp^q {\{\Phi(e)\}} = \perp^q {\{\Phi(a)\}} \subseteq \perp^q {\{\Phi(b)\}}$ . It follows from  $\Phi(c) \perp \Phi(b)$ that  $\Phi(c) \in {}^{\perp_q} {\Phi(b)}$ , while  $\Phi(c) \notin {}^{\perp_q} {\Phi(a)}$  because  $\Phi(c) \in \Phi(E)_2(\Phi(e))$ . This contradicts the assumed maximality of <sup> $\perp_q \{a\}$ </sup>. Therefore  $\Phi(e)$  must be minimal in A (equivalently, e must be minimal in E), as desired.  $\square$  The next corollary is an interesting characterization of minimal tripotents.

Corollary 2.5. *Let* a *be a norm-one element in a JB*<sup>∗</sup> *-triple* E*. Then the following statements are equivalent.*

- (a) a *is a minimal tripotent in* E*.*
- (b) The inner quadratic annihilator of a,  $\perp^q \{a\}$ , is maximal among the set of all *inner quadratic annihilators of elements in* E*.*

From Theorem [2.4](#page-5-0) one can derive a characterisation of positive scalar multiples of minimal partial isometries in C<sup>∗</sup> -algebras, which seems to be also new.

Corollary 2.6. *Let* a *be a non-zero element in a C*<sup>∗</sup> *-algebra* A*. Then the following statements are equivalent.*

- (a) a *is a* (*positive*) *scalar multiple of a minimal partial isometry in* A*.*
- <span id="page-7-0"></span>(b) The inner quadratic annihilator of a,  $\perp^q \{a\}$ , is maximal among the set of all *inner quadratic annihilators of elements in* A*.*

## 3. THE CASE OF RANK-ONE JB<sup>\*</sup>-TRIPLES

A general JB<sup>∗</sup> -triple need not contain a single non-zero tripotent. However, every Cartan factor coincides with the weak<sup>\*</sup>-closure of the linear of its minimal tripotents, and the same occurs to any arbitrary orthogonal or direct sum of a family of Cartan factors. Actually a JBW<sup>\*</sup>-triple coincides with the weak<sup>\*</sup>-closure of the linear span of its minimal tripotents (also called atomic) if, and only if, it is a direct sum of a family of Cartan factors  $[26,$  Proposition 2. It is perhaps worth to review the notion of rank. A tripotent  $e$  in a JB<sup>\*</sup>-triple  $E$  has rank–n with  $n \in \mathbb{N}$ (denoted by  $r(e) = n$ ), if it can be written as the sum of n mutually orthogonal minimal tripotents in E. A subset S of E is called *orthogonal* if  $0 \notin S$  and  $x \perp y$ for every  $x \neq y$  in S. The minimal cardinal number r satisfying card(S)  $\leq r$  for every orthogonal subset  $S \subseteq E$  is called the *rank* of E (cf. [\[36\]](#page-30-19), [\[6\]](#page-29-5) and [\[4\]](#page-29-6) for basic results on the rank of a Cartan factor and a JBW<sup>∗</sup> -triple, and its connection with the reflexivity of the underlying Banach space).

Let us illustrate the previous notions with some examples. Let  $H$  be a complex Hilbert space regarded as a type 1 Cartan factor of the form  $H \cong B(H, \mathbb{C})$ , that is we consider its Hilbert norm and the triple product given by

$$
\{a, b, c\} = \frac{1}{2} (\langle a|b \rangle c + \langle c|b \rangle a), \quad (a, b, c \in H).
$$

It is easy to see that every norm-one element in  $H$  is a minimal and complete tripotent in  $H$ , and the latter has rank–one. Actually, a JB<sup>\*</sup>-triple has finite rank if, and only if, it is reflexive, and it has rank-one precisely when it is isometrically isomorphic to a complex Hilbert space regarded as a type 1 Cartan factor (see the discussion in  $[4, §3]$ ,  $[36, page 210]$ ,  $[15, Corollary in page 308]$ ). In case that H and K are complex Hilbert spaces with dimensions  $m \leq n$ , respectively, the JB<sup>\*</sup>-triple  $B(H, K)$  has rank–m.

<span id="page-7-1"></span>The relationship "being a trunction of" admits a particular nice characterization in the case of rank-one JB<sup>∗</sup> -triples.

**Remark 3.1.** Let  $H$  be a Hilbert space regarded as a type 1 Cartan factor, and let x, y be non-zero elements in H. It is easy to see that x is a truncation of y (i.e.  ${x, x, x} = {x, y, x}$  if, and only if,  $\langle y|x \rangle = \langle x|x \rangle$ , equivalently,  $y = x + z$  with  $\langle x|z \rangle = 0$ . We can alternatively say that the inner quadratic annihilator of x is precisely the (Euclidean) orthogonal complement of  $x$ , that is,

$$
{}^{\perp_q}{x} = {x}^{\perp_2} = {z \in H : \langle z|x \rangle = 0}.
$$

In this setting we write  $x \perp_2 y$  if  $\langle x|y \rangle = 0$ . So, the study of mappings between Hilbert spaces preserving (Euclidean) orthogonality is equivalent to the study of mappings preserving truncations.

Every non-zero element x in H can be written in the form  $x = ||x|| \frac{x}{||x||}$ , where  $\frac{x}{\|x\|}$  is a minimal tripotent in H. It is easy to check that  $x, y \in H \setminus \{0\}$  satisfy  $x \perp_2 y$ if, and only if,  $\frac{x}{\|x\|}$  and  $\frac{y}{\|y\|}$  are collinear.

Motivated by the previous remark, we recall a well-known definition. Let  $H$ and K be two real or complex Hilbert spaces. A mapping  $\Delta: H \to K$  preserves (Euclidean) orthogonality if

$$
\forall x, y \in H, \ x \perp_2 y \Rightarrow \Delta(x) \perp_2 \Delta(y).
$$

<span id="page-8-0"></span>The mapping  $\Delta$  preserves (Euclidean) orthogonality in both directions if the equivalence  $x \perp_2 y \Leftrightarrow \Delta(x) \perp_2 \Delta(y)$  holds for all  $x, y \in H$ .

Example 3.2. In case that our JB<sup>\*</sup>-triple  $E$  reduces to the complex field, the characterization of additive surjective maps preserving truncations in both directions on E is simply hopeless, since every additive surjective mapping on  $\mathbb C$  preserves truncations (equivalently, (Euclidean) orthogonality) in both directions. We can easily find an additive bijection on C preserving orthogonality in both directions (cf. the introduction of [\[37\]](#page-30-6)).

<span id="page-8-1"></span>We shall see next that the counterexample above can only occur when the Hilbert space one–dimensional.

**Proposition 3.3.** Let E and F be rank-one  $JB^*$ -triples with  $dim(E) \geq 2$ . Suppose *that*  $A: E \to F$  *is a surjective additive mapping preserving truncations, then* A *is a positive scalar multiple of a* (*complex*) *linear or a conjugate-linear isometry.*

*Proof.* As we commented above, under our hypotheses, E and F must be complex Hilbert spaces regarded as type 1 Cartan factors. By Lemma [2.3](#page-5-2) and Remark [3.1,](#page-7-1) A preserves (Euclidean) orthogonality of the underlying Hilbert spaces. Since A is surjective and additive by hypotheses, the desired conclusion is now a consequence of Theorem 2.1 in [\[37\]](#page-30-6).

<span id="page-8-2"></span>**Remark 3.4.** Let  $E$  be a JB<sup>\*</sup>-triple. Then

 $E = \mathbb{C} \Leftrightarrow \perp^{\perp_q} \{a\} = \{0\}, \text{ for all } a \in E \setminus \{0\}$ 

The "only if" implication is clear. For the "if" implication, observe that if  $a$  and  $b$  are two orthogonal non-zero elements in E, we have  $0 \neq b \in \perp^q \{a\}$ , which contradicts the assumption. So,  $E$  has rank-one, and thus  $E$  is a complex Hilbert space. If  $\dim(E) \geq 2$ , we can take two non-zero vectors  $a, b \in E$  with  $\langle a|b \rangle = 0$ . In this case  $0 \neq b \in \perp^q \{a\}$ , which is also a contradiction.

#### 4. Additive preservers of truncations

<span id="page-9-0"></span>We have already observed in Lemma [2.3](#page-5-2) that an additive mapping between JB<sup>\*</sup>triples preserves truncations if, and only if, it preserves inner quadratic annihilators. In this section we study the first properties of surjective additive preservers of truncations. It is well-known that a mapping between real or complex normed spaces is additive if, and only if, it is  $\mathbb Q$ -linear. We shall use this fact without any further mentioning. From now on, the symbol T will stand for the unit sphere of  $\mathbb{C}.$ 

<span id="page-9-1"></span>**Lemma 4.1.** Let  $A: E \to F$  be a surjective additive mapping between JB<sup>\*</sup>-triples. *Suppose additionally that* A *preserves truncations of elements in both directions, that is,*

 ${a, b, a} = {a, a, a} \text{ in } E \Leftrightarrow {A(a), A(b), A(a)} = {A(a), A(a), A(a)}.$ 

*Then the following statements hold:*

- (a) A *is injective.*
- (b)  $dim(E) > 2$  *if, and only if,*  $dim(F) > 2$ *.*
- (c) A *maps minimal tripotents in* E *to positive scalar multiples of minimal tripotents in* F*, Moreover, if* e *is a minimal tripotent in* E *and* λ *is a non-zero complex number, then* A(λe) *is a positive scalar multiple of a minimal tripotent in* F, that is,  $A(\mathbb{C} \mathcal{U}_{min}(E)) = \mathbb{C} \mathcal{U}_{min}(F)$ .
- (d) For each minimal tripotent  $e$  in  $E$  we have  $A(\mathbb{C}e) = \mathbb{C}A(e) = \mathbb{C}r(A(e)).$
- (e) For each minimal tripotent  $e \in E$ , we have  $r(A^{-1}(r(A(e)))) \in \mathbb{T}e$ .

*Proof.* (a) By the additivity of A it suffices to prove that  $A(a) = 0$  implies  $a = 0$ . Let us take  $a \in E$  with  $A(a) = 0$ . It then follows that  $\{A(a), A(b), A(a)\} = 0$  ${A(a), A(a), A(a)}$  for all  $b \in E$ , and thus  ${a, b, a} = {a, a, a}$  for all  $b \in E$ . Taking  $b = 0$  we get  $||a||^3 = ||\{a, a, a\}|| = 0.$ 

(b) Since A is a bijection and, by Lemma [2.3,](#page-5-2)  $A(\perp^q \{a\}) = \perp^q \{A(a)\}\$ for all  $a \in E$ , the desired equivalence is a consequence of the characterization of one-dimensional JB<sup>∗</sup> -triples given in Remark [3.4.](#page-8-2)

(c) This statement follows from Lemma [2.3](#page-5-2) and Theorem [2.4.](#page-5-0)

(d) Since  $A(0) = 0$ , it suffices to prove that for each  $\lambda \in \mathbb{C}\backslash\{0\}$  the element  $A(\lambda e)$  lies in  $\mathbb{C}A(e)$ . Lemma [2.3](#page-5-2) implies that

$$
\perp_q \{A(\lambda e)\} = A(\perp_q \{ \lambda e\}) = A(\perp_q \{ e\}) = \perp_q \{A(e)\}.
$$

Since, by (b) and (a), the element  $A(\lambda e)$  is a positive multiple of a minimal tripotent in F, Lemma [2.1](#page-4-1) assures that  $A(\lambda e) \in \mathbb{C}A(e)$ . The rest follows from the bijectivity of A.

 $(e)$  We know from  $(d)$  that

$$
A(e) = \gamma r(A(e))
$$
 and  $A^{-1}(r(A(e))) = \delta r(A^{-1}(r(A(e))))$ ,

<span id="page-9-2"></span>for some positive real numbers  $\gamma$  and  $\delta$ , where  $r(A(e))$  and  $r(A^{-1}(r(A(e))))$  are minimal tripotents in F and E, respectively. It follows from  $(c)$ , applied to  $A^{-1}$ , that  $e = A^{-1}A(e) = A^{-1}(\gamma r(A(e))) \in \mathbb{C}A^{-1}(r(A(e))),$  which implies that  $A^{-1}(r(A(e)))$ =  $\lambda e$  for some non-zero complex number  $\lambda$ , and hence  $r(A^{-1}(r(A(e)))) \in \mathbb{T}e$ , as desired.  $\Box$ 

**Remark 4.2.** Let  $E$  and  $F$  be JB<sup>\*</sup>-triples. Thanks to the previous lemma, we can conclude that every surjective additive mapping  $A: E \to F$  preserving truncations in both directions is a bijection (and  $A^{-1}$  enjoys the same property). Moreover, for each minimal tripotent  $e \in E$ , the range tripotent of  $A(e)$  (in  $F^{**}$ ) is a minimal tripotent and lies in F, and there exists a unique positive number  $\gamma(e, A)$  (depending on e and A) such that  $A(e) = \gamma(e, A)r(A(e))$ . We further know the existence of a unique  $\gamma_1(e, A) \in \mathbb{T}$  satisfying  $A^{-1}(r(A(e))) = \gamma_1(e, A)r(A^{-1}(r(A(e))))$ . We keep this notation henceforth.

A *quadrangle* in E is an ordered quadruple  $(u_1, u_2, u_3, u_4)$  formed by tripotents in E satisfying  $u_1 \perp u_3$ ,  $u_2 \perp u_4$ ,  $u_1 \perp u_2 \perp u_3 \perp u_4 \perp u_5$  and  $u_4 = 2{u_1, u_2, u_3}$  (the latter equality also holds if the indices are permutated cyclically –e.g.  $u_2 = 2{u_3, u_4, u_1}.$ An ordered triplet  $(w_1, u, w_2)$  of tripotents in E is called a *trangle* if  $w_1 \perp w_2$ ,  $u \vdash w_1$ ,  $u \vdash w_2$  and  $w_1 = Q(u)(w_2)$  (see, the references [\[15,](#page-30-20) [41\]](#page-31-4) for additional details).

Let  $(e_1, e_2, e_3, e_4)$  (respectively,  $(w_1, u, w_2)$ ) be a quadrangle (respectively, a trangle) of tripotents in a JB<sup>\*</sup>-triple  $E$ . It is part of the folklore in JB<sup>\*</sup>-triple theory that in case that  $E$  is a Cartan factor and  $e_1, e_2, e_3$ , and  $e_2$  are minimal tripotents (respectively, u is a rank-2 tripotent and  $w_1$  and  $w_2$  are minimal tripotents), then the element  $\alpha e_1 + \beta e_2 + \gamma e_4 + \delta e_3$  (respectively,  $\alpha w_1 + \beta u + \delta w_2$ ) is a minimal tripotent in E for every  $\alpha, \beta, \gamma, \delta \in \mathbb{C}$  with  $|\alpha|^2 + |\beta|^2 + |\gamma|^2 + |\delta|^2 = 1$ , and  $\alpha\delta - \beta\gamma = 0$ (respectively, for every  $\alpha, \beta, \delta \in \mathbb{C}$  with  $|\alpha|^2 + 2|\beta|^2 + |\delta|^2 = 1$ , and  $\alpha\delta - \beta^2 = 0$ ) (cf. [\[21,](#page-30-21) Lemma 3.10]). We can now prove a stronger property via an alternative and self-contained argument.

## <span id="page-10-0"></span>Lemma 4.3. *Let* E *be a JB*<sup>∗</sup> *-triple. The the following statements hold:*

- (a) Let  $(e_1, e_2, e_3, e_4)$  *be a quadrangle of tripotents in* E. Then for every  $\alpha, \beta, \gamma, \delta \in$  $\mathbb{C}$  with  $|\alpha|^2 + |\beta|^2 + |\gamma|^2 + |\delta|^2 = 1$ , and  $\alpha\delta - \beta\gamma = 0$ , the element  $v = \alpha e_1 + \beta e_2$  $\beta e_2 + \gamma e_4 + \delta e_3$  *is a tripotent in* E. Furthermore, if  $e_1$  and  $e_3$  (respectively,  $e_2$ ) and  $e_4$ ) are minimal tripotents, the tripotents  $v, e_2$ , and  $e_4$  (respectively,  $v, e_1$ , *and* e3) *are minimal.*
- (b) Let  $(w_1, u, w_2)$  be a trangle of tripotents in E. Then for every  $\alpha, \beta, \delta \in \mathbb{C}$  with  $|\alpha|^2 + 2|\beta|^2 + |\delta|^2 = 1$ , and  $\alpha\delta - \beta^2 = 0$ , the element  $v = \alpha w_1 + \beta u + \delta w_2$  is a *tripotent in* E. Furthermore, if  $w_1$  and  $w_2$  are minimal tripotents, the tripotent v *is minimal.*

*Proof.* (a) Let  $(e_1, e_2, e_3, e_4)$  be a quadrangle of tripotents in E, and let  $\alpha, \beta, \gamma, \delta$  be complex numbers satisfying the hypotheses above. Having in mind the definition of quadrangle and Peirce arithmetic, and a bit of patience, we compute the cube of the element  $v$ ,

$$
\{v, v, v\} = \{\alpha e_1 + \beta e_2 + \gamma e_4 + \delta e_3, \alpha e_1 + \beta e_2 + \gamma e_4 + \delta e_3, \alpha e_1 + \beta e_2 + \gamma e_4 + \delta e_3\}
$$
  
\n
$$
= |\alpha|^2 \Big( \alpha \{e_1, e_1, e_1\} + \beta \{e_1, e_1, e_2\} + \gamma \{e_1, e_1, e_4\} + \delta \{\underline{e_1, e_1, e_3}\} + \alpha \overline{\beta} \Big( \alpha \{\underline{e_1, e_2, e_1}\} + \beta \{e_1, e_2, e_2\} + \gamma \{\underline{e_1, e_2, e_4}\} + \delta \{e_1, e_2, e_3\} \Big)
$$
  
\n
$$
+ \alpha \overline{\gamma} \Big( \alpha \{\underline{e_1, e_4, e_1}\} + \beta \{\underline{e_1, e_2, e_2}\} + \gamma \{\underline{e_1, e_2, e_4}\} + \delta \{\underline{e_1, e_2, e_3}\} \Big)
$$
  
\n
$$
+ \alpha \overline{\delta} \Big( \alpha \{\underline{e_1, e_4, e_1}\} + \beta \{\underline{e_1, e_4, e_2}\} + \gamma \{\underline{e_1, e_4, e_4}\} + \delta \{\underline{e_1, e_4, e_3}\} \Big)
$$
  
\n
$$
+ \alpha \overline{\delta} \Big( \alpha \{\underline{e_1, e_3, e_1}\} + \beta \{\underline{e_1, e_3, e_2}\} + \gamma \{\underline{e_1, e_3, e_4}\} + \delta \{\underline{e_1, e_3, e_3}\} \Big)
$$

+ 
$$
\beta \overline{\alpha} \left( \alpha \{e_2, e_1, e_1\} + \beta \{e_2, e_1, e_2\} + \gamma \{e_2, e_1, e_4\} + \delta \{e_2, e_2, e_3\} \right)
$$
  
+  $\beta \overline{\beta} \left( \alpha \{e_2, e_2, e_1\} + \beta \{e_2, e_2, e_2\} + \gamma \{e_2, e_2, e_1, e_4\} + \delta \{e_2, e_2, e_3\} \right)$   
+  $\beta \overline{\gamma} \left( \alpha \{e_2, e_1, e_1\} + \beta \{e_2, e_3, e_2\} + \gamma \{e_2, e_1, e_4\} + \delta \{e_2, e_2, e_3\} \right)$   
+  $\beta \overline{\delta} \left( \alpha \{e_2, e_3, e_1\} + \beta \{e_2, e_3, e_2\} + \gamma \{e_2, e_3, e_4\} + \delta \{e_2, e_3, e_3\} \right)$   
+  $\gamma \overline{\alpha} \left( \alpha \{e_4, e_1, e_1\} + \beta \{e_4, e_1, e_2\} + \gamma \{e_4, e_1, e_4\} + \delta \{e_4, e_1, e_3\} \right)$   
+  $\gamma \overline{\beta} \left( \alpha \{e_4, e_2, e_3\} + \beta \{e_4, e_1, e_2\} + \gamma \{e_4, e_3, e_4\} + \delta \{e_4, e_2, e_3, e_3\} \right)$   
+  $\gamma \overline{\gamma} \left( \alpha \{e_4, e_3, e_1\} + \beta \{e_4, e_2, e_2\} + \gamma \{e_4, e_3, e_4\} + \delta \{e_4, e_2, e_3\} \right)$   
+  $\gamma \overline{\gamma} \left( \alpha \{e_4, e_3, e_1\} + \beta \{e_4, e_3, e_2\} + \gamma \{e_4, e_3, e_4\} + \delta \{e_4, e_3, e_3\} \right)$   
+  $\delta \overline{\alpha} \left( \alpha \{e_3, e_2, e_1$ 

Since, clearly,  $e_1, e_2, e_4, e_3 \in E_2(e_1 + e_3)$ , all the tripotents listed above lie in  $E_2(e_1 + e_3)$ . If  $e_1$  and  $e_3$  are minimal tripotents, the Peirce-2 subspace  $E_2(e_1 + e_3)$ is a spin factor (cf. [\[32,](#page-30-22) Lemma 3.6]). Since spin factors have rank-2 (see, for example, [\[36,](#page-30-19) Table 1 in page 210]),  $e_2 \perp e_4$ , and  $e_2 + e_4 \in E_2(e_1 + e_3)$ , it follows that  $e_2$  and  $e_4$  must be minimal in  $E_2(e_1 + e_2)$ , and hence in E by [\[9,](#page-29-7) Lemma 3.3]. Consequently,  $e_2$  and  $e_4$  are minimal tripotents too. It is part of the folklore in  $JB<sup>*</sup>$ -triple theory that under these conditions  $v$  is a minimal tripotent.

We can alternatively consider the element  $\tilde{v} = \overline{\delta}e_1 - \overline{\gamma}e_2 - \overline{\beta}e_4 + \overline{\alpha}e_3$ . The arguments above also prove that  $\tilde{v}$  is a tripotent in E. By following similar computations to those at the beginning of this proof we deduce that

$$
\{v, v, \tilde{v}\} = \{\alpha e_1 + \beta e_2 + \gamma e_4 + \delta e_3, \alpha e_1 + \beta e_2 + \gamma e_4 + \delta e_3, \overline{\delta} e_1 - \overline{\gamma} e_2 - \overline{\beta} e_4 + \overline{\alpha} e_3\}
$$
  
\n
$$
= |\alpha|^2 \overline{\delta} e_1 - \frac{1}{2} |\alpha|^2 \overline{\gamma} e_2 - \frac{1}{2} |\alpha|^2 \overline{\beta} e_4 - \frac{1}{2} \alpha \overline{\beta} \overline{\gamma} e_1 + \frac{1}{2} \alpha \overline{\beta} \overline{\alpha} e_4 - \frac{1}{2} \alpha \overline{\gamma} \overline{\beta} e_1 + \frac{1}{2} \alpha \overline{\gamma} \overline{\alpha} e_2
$$
  
\n
$$
+ \frac{1}{2} \beta \overline{\alpha} \overline{\delta} e_2 - \frac{1}{2} \beta \overline{\alpha} \overline{\beta} e_3 + \frac{1}{2} \beta \overline{\beta} \overline{\delta} e_1 - \beta \overline{\beta} \overline{\gamma} e_2 + \frac{1}{2} \beta \overline{\beta} \overline{\alpha} e_3 - \frac{1}{2} \beta \overline{\delta} \overline{\beta} e_1 + \frac{1}{2} \beta \overline{\delta} \overline{\alpha} e_2
$$
  
\n
$$
+ \frac{1}{2} \gamma \overline{\alpha} \overline{\delta} e_4 - \frac{1}{2} \gamma \overline{\alpha} \gamma e_3 + \frac{1}{2} \gamma \overline{\gamma} \overline{\delta} e_1 - \gamma \overline{\gamma} \overline{\beta} e_4 + \frac{1}{2} \gamma \overline{\gamma} \overline{\alpha} e_3 - \frac{1}{2} \gamma \overline{\delta} \gamma e_1 + \frac{1}{2} \gamma \overline{\delta} \overline{\alpha} e_4
$$
  
\n
$$
+ \frac{1}{2} \delta \overline{\beta} \delta e_4 - \frac{1}{2} \delta \overline{\beta} \gamma e_3 + \frac{1}{2} \delta \overline{\gamma} \overline{\delta} e_2 - \frac{1}{2} \delta \overline{\gamma} \overline{\beta} e_3 - \frac{1}{2} \delta \overline{\delta} \gamma e_2 - \frac{1}{2} \delta \over
$$

and thus  $v \perp \tilde{v}$ . Since  $v, \tilde{v}$  are both non-zero and lie in  $E_2(e_1 + e_3)$ , we conclude that v and  $\tilde{v}$  are minimal tripotents in  $E$ .

Clearly, the roles of  $e_1, e_3$  and  $e_2, e_4$  can be exchanged in this final argument by just having in mind that  $(e_2, e_1, e_4, e_3)$  is a quadrangle too.

(b) The proof follows similar lines to those in the proof of  $(a)$  by just observing the following identities. First, since by assumptions  $\{u, w_1, u\} = w_2$ , the element  $w_1 + w_2$  is a self-adjoint tripotent in the JB<sup>\*</sup>-algebra  $E_2(u)$ . Moreover, since  $\{w_1 + w_2, w_1 + w_2, u\} = \{w_1, w_1, u\} + \{w_2, w_2, u\} = u$ , we can deduce, via Peirce arithmetic, that  $u = \{w_1 + w_2, u, w_1 + w_2\} = \{w_1, u, w_1\} + \{w_2, u, w_2\} + \{w_3, w_4, w_5\}$  $2\{w_1, u, w_2\} = 2\{w_1, u, w_2\}.$  For the sake of brevity, the remaining computations are left to the reader.

The relationship "being in the inner quadratic annihilator" is not, in general, reflexive. For example, the tripotents  $e =$  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$  and  $w =$  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  satisfy that  $w \in {}^{\perp_q} \{e\}$  but  $e \notin {}^{\perp_q} \{w\}.$ 

It is shown in [\[15,](#page-30-20) Lemma in page 306] that each family  $\{w_j\}_j$  of mutually collinear minimal tripotents in a JB<sup>∗</sup> -triple generates a norm-closed subspace isometrically isomorphic to a complex Hilbert space in which the family  $\{w_j\}_j$  is an orthonormal basis. We shall see a kind of reciprocal to this property in the next key proposition.

<span id="page-12-0"></span>Proposition 4.4. *Let* w1, w<sup>2</sup> *be two minimal tripotents in a JB*<sup>∗</sup> *-triple* E*. Then the following statements hold:*

- (a)  $w_1 \in {}^{\perp_q}(w_2)$  if, and only if,  $w_2 \in {}^{\perp_q}(w_1)$ .
- (b) Suppose that  $w_2 \in {}^{\perp_q}(w_1)$  and there exist  $\gamma_1, \gamma_2 \in \mathbb{R}^+$ ,  $q_1, q_2 \in \mathbb{Q}$  with  $q_1^2 + q_2^2 =$ 1, and  $q_1q_2 \neq 0$ , such that the element  $q_1\gamma_1w_1 + q_2\gamma_2w_2$  *is a positive multiple of a minimal tripotent in*  $E$ *. Then*  $w_1$  *and*  $w_2$  *are collinear.*

*Proof.* As in the proof of Theorem [2.4,](#page-5-0) we can embed E inside the atomic part,  $\mathcal{A}_{E^{**}}$ , of its bidual space,  $E^{**}$ , via an isometric triple isomorphism  $\Psi_E : E \hookrightarrow \mathcal{A}_{E^{**}}$ with weak<sup>∗</sup>-dense image, and  $A_{E^{**}}$  can be represented as an  $\ell_{\infty}$ -sum of a family of Cartan factors  $\{C_i\}_i$  (cf. [\[25,](#page-30-12) Theorem 2] and [\[26,](#page-30-13) Proposition 1 and its proof]). Clearly,  $\Psi_E(w_1)$  and  $\Psi_E(w_2)$  are minimal tripotents in  $\mathcal{A}_{E^{**}} = \bigoplus^{\infty} C_i$ , and hence each one of them belongs to a unique Cartan factor in the  $\ell_{\infty}\text{-sum.}$  Let us assume that  $\Psi_E(w_j) \in C_{i_j}$   $(j = 1, 2)$ .

(a) Assume that  $w_2 \in {}^{\perp_q}(w_1)$  (equivalently,  $\Psi_E(w_2) \in {}^{\perp_q}(\Psi_E(w_1))$ ). If  $i_1 \neq i_2$ , the tripotents  $\Psi_E(w_1)$  and  $\Psi_E(w_2)$  (equivalently,  $w_1, w_2$ ) are orthogonal in  $\mathcal{A}_{E^{**}}$ (respectively, in E), which implies that  $w_1 \in {}^{\perp_q}(w_2)$  and  $w_2 \in {}^{\perp_q}(w_1)$ . We can therefore assume that  $\Psi_E(w_1)$  and  $\Psi_E(w_2)$  belong to the same Cartan factor  $C_{i_1}$ (i.e.  $i_1 = i_2$ ). We are thus, in a position to apply [\[21,](#page-30-21) Lemma 3.10] to deduce that one of the following statements holds:

(i) There exist minimal tripotents  $e_2, e_3, e_4$  in  $C_{i_1}$  such that  $(\Psi_E(w_1), e_2, e_3, e_4)$ is a quadrangle and  $\Psi_E(w_2) = \alpha \Psi_E(w_1) + \beta e_2 + \gamma e_4 + \delta e_3$  with  $\alpha, \beta, \gamma, \delta \in \mathbb{C}$ ,  $|\alpha|^2 + |\beta|^2 + |\gamma|^2 + |\delta|^2 = 1$ , and  $\alpha\delta = \gamma\beta$ ;

(*ii*) There exist a minimal tripotent  $\tilde{e} \in C_{i_1}$  and a rank-two tripotent  $u \in C_{i_1}$  such that  $(\Psi_E(w_1), u, \tilde{e})$  is a trangle and  $\Psi_E(w_2) = \alpha \Psi_E(w_1) + \beta u + \delta \tilde{e}$  with  $\alpha, \beta, \delta \in \mathbb{C}$ and  $\alpha\delta = \beta^2$ .

Since  $\Psi_E(w_2) \in {}^{\perp_q} {\{\Psi_E(w_1)\}}$ , it follows that in each of the previous cases we have  $\alpha = 0$ . In case (ii) the element  $\beta$  must be also zero, and hence  $\Psi_E(w_2) = \delta \tilde{e}$ is orthogonal to  $\Psi_E(w_1)$ , and consequently,  $\Psi_E(w_1) \in {}^{\perp_q} {\{\Psi_E(w_2)\}}$ . In case (*i*),  $\beta\gamma = 0$ , implies that  $\Psi_E(w_2) = \gamma e_4 + \delta e_3$  or  $\Psi_E(w_2) = \beta e_2 + \delta e_3$ . In both cases, by Peirce arithmetic, we have  ${\Psi_E(w_2), \Psi_E(w_1), \Psi_E(w_2)} = {\gamma e_4 + \delta e_3, \Psi_E(w_1), \gamma e_4 + \delta e_4}$  $\{\delta e_3\} = \gamma \{e_4, \Psi_E(w_1), e_4\} = 0$ , and similarly  $\{\beta e_2 + \delta e_3, \Psi_E(w_1), \beta e_2 + \delta e_3\} = 0$  $\beta^2 \{e_2, \Psi_E(w_1), e_2\} = 0.$  This shows that  $\Psi_E(w_1) \in {}^{\perp_q} {\{\Psi_E(w_2)\}}.$ 

(b) Let us assume the hypothesis in the statement and in the opening paragraph of this proof. If  $i_1 \neq i_2$ , the tripotents  $w_1$  and  $w_2$  (equivalently,  $\Psi_E(w_1) = w_1$  and  $\Psi_E(w_2) = w_1$  are orthogonal in  $\mathcal{A}_{E^{**}}$  (respectively, in E). By assumptions, the element  $q_1\gamma_1w_1+q_2\gamma_2w_2$  is a minimal tripotent in E, which is impossible. Therefore the elements  $\Psi_E(w_1) = w_1$  and  $\Psi_E(w_2) = w_2$  belong to the same Cartan factor  $C_{i_1}$ . Lemma 3.10 in [\[21\]](#page-30-21) assures that one of the following statements holds:

(i) There exist minimal tripotents  $e_2, e_3, e_4$  in  $C_{i_1}$  such that the elements  $w_1, e_2,$  $e_3$ , and  $e_4$  form a quadrangle, and

$$
w_2 = \alpha w_1 + \beta e_2 + \gamma e_4 + \delta e_3
$$

with  $\alpha, \beta, \gamma, \delta \in \mathbb{C}, |\alpha|^2 + |\beta|^2 + |\gamma|^2 + |\delta|^2 = 1$ , and  $\alpha\delta = \gamma\beta$ . In such a case we have

 $q_1\gamma_1w_1 + q_2\gamma_2w_2 = q_1\gamma_1w_1 + q_2\gamma_2\alpha w_1 + q_2\gamma_2\beta e_2 + q_2\gamma_2\gamma e_4 + q_2\gamma_2\delta e_3.$ 

Now, by the quadrangle's properties, we get  $(q_1\gamma_1 + q_2\gamma_2\alpha) q_2\gamma_2\delta = q_2^2\gamma_2^2\beta\gamma$ , and thus  $\gamma_1\gamma_2q_1q_2\delta=0$ , equivalently,  $\delta=0$ . The condition  $w_2 \in {}^{\perp_q}(w_1)$  implies that  $\alpha = 0$ . Therefore, either  $w_2 = \beta e_2$ ,  $(\gamma = 0)$  or  $w_2 = \gamma e_4$  ( $\beta = 0$ ), and hence  $w_1 \top w_2$ . (*ii*) There exist a minimal tripotent  $\tilde{e} \in C_{i_1}$  and a rank-two tripotent  $u \in C_{i_1}$  such that  $(\Psi_E(w_1), u, \tilde{e})$  is a trangle and  $\Psi_E(w_2) = \alpha \Psi_E(w_1) + \beta u + \delta \tilde{e}$  with  $\alpha, \beta, \delta \in \mathbb{C}$ and  $\alpha\delta = \beta^2$ . The condition  $w_2 \in {}^{\perp_q}(w_1)$  gives  $\alpha = 0 = \beta$ , and thus  $w_2 = \delta\tilde{e}$ is orthogonal to  $w_1$ , which contradicts that  $q_1\gamma_1w_1 + q_2\gamma_2w_2$  is a positive scalar multiple of a minimal tripotent in  $E$ . This shows that this second case is impossible.  $\Box$ 

As a consequence of the previous proposition, we can conclude next that a surjective additive mapping between JB<sup>∗</sup> -triples preserving truncations in both directions "somehow preserves" collinear minimal tripotents.

<span id="page-14-0"></span>**Proposition 4.5.** Let  $A : E \to F$  be a surjective additive mapping between  $JB^*$ . *triples. Suppose additionally that* A *preserves truncations of elements in both directions.* Let  $w_1, w_2$  be two collinear minimal tripotents in E. Then the range *tripotents*  $r(A(w_1))$  *and*  $r(A(w_2))$  *are minimal and collinear in* F.

*Proof.* Since  $w_1$  and  $w_2$  are collinear, it is not hard to see that for every  $q_1, q_2 \in \mathbb{Q}$ with  $q_1^2 + q_2^2 = 1$ , the element  $q_1w_1 + q_2w_2$  is a minimal tripotent in E, and thus, Remark [4.2](#page-9-2) assures that

$$
A(q_1w_1 + q_2w_2) = q_1\gamma(w_1, A)r(A(w_1)) + q_2\gamma(w_2, A)r(A(w_2))
$$

is a positive scalar multiple of a minimal tripotent in  $F$ . It follows from Proposi-tion [4.4](#page-12-0)(b) that  $r(A(w_1))\top r(A(w_2))$ .

We say that two tripotents e and v in a JB<sup>\*</sup>-triple E are *compatible* if  $P_j(e)P_k(v) =$  $P_k(v)P_j(e)$  for all  $i, k \in \{0, 1, 2\}$ . If  $v \in E_j(e)$  for some  $j \in \{0, 1, 2\}$ , then e and v are compatible  $[29, (1.10)].$ 

<span id="page-14-2"></span>**Lemma 4.6.** Let  $w_1, w_2$  be orthogonal tripotents in a JB<sup>\*</sup>-triple E. Then

$$
E_1(w_1) \subseteq {}^{\perp_q} \{w_2\}.
$$

*Proof.* Let us take  $x \in E_1(w_1)$ . Since  $w_1$  and  $w_2$  are compatible, we conclude that  $E_1(w_1) = (E_1(w_1) \cap E_0(w_2)) \oplus (E_1(w_1) \cap E_1(w_2)) \oplus (E_1(w_1) \cap E_2(w_2))$ . Observe that  $w_2 \perp w_1$  implies that  $E_2(w_2) \subseteq E_0(w_1)$ , and thus  $E_1(w_1) \cap E_2(w_2) = \{0\}$ . We can therefore write  $x = x_0 + x_1$  with  $x_j \in E_1(w_1) \cap E_j(w_2)$ . Therefore, we deduce from Peirce arithmetic that

$$
\{w_2, x, w_2\} = \{w_2, x_0, w_2\} + \{w_2, x_1, w_2\} = \{w_2, x_1, w_2\} \in E_{4-1}(w_2) = \{0\},\
$$

which shows that  $x \in \perp^q$  $\{w_2\}.$ 

<span id="page-14-1"></span>We can now describe the images of two orthogonal minimal tripotents under a surjective additive mapping preserving truncations.

**Proposition 4.7.** Let E and F be  $JB^*$ -triples, and let  $A : E \to F$  be a surjective *additive mapping preserving truncations in both directions. Suppose additionally that* F is atomic. Let  $w_1, w_2$  be two orthogonal minimal tripotents in E. Then the *range tripotents*  $r(A(w_1))$  *and*  $r(A(w_2))$  *are orthogonal in* F.

*Proof.* Since F is atomic, it can be expressed in the form  $F = \bigoplus^{\ell_{\infty}} C_i$ , where each  $C_i$  is a Cartan factor. If  $A(w_1)$  and  $A(w_2)$  lie in two different Cartan factors in the decomposition of  $F$ , then they are clearly orthogonal and the same occurs to  $r(A(w_1))$  and  $r(A(w_2))$ , which gives the desired statement. We can therefore assume that  $A(w_1)$  and  $A(w_2)$  belong to the same Cartan factor  $C_{i_1}$ . We are again in a position to apply  $\left|21\right|$ , Lemma 3.10 to reduce our study to one of the following cases:

(*i*) There exist minimal tripotents  $e_2, e_3, e_4$  in  $C_{i_1}$  such that the ordered quadruple  ${r(A(w_1)), e_2, e_3, e_4}$  is a quadrangle, and

$$
r(A(w_2)) = \alpha r(A(w_1)) + \beta e_2 + \gamma e_4 + \delta e_3
$$

with  $\alpha, \beta, \gamma, \delta \in \mathbb{C}, |\alpha|^2 + |\beta|^2 + |\gamma|^2 + |\delta|^2 = 1$ , and  $\alpha\delta = \gamma\beta$ . By hypotheses,  $A(w_2) =$  $\gamma(w_2, A)r(A(w_2)) \in {}^{\perp_q}\{A(w_1)\}\$ , and thus  $\alpha = 0$ , which also implies that  $\beta = 0$ or  $\gamma = 0$ . We shall only consider the first case, since the other one can be treated via similar arguments. Assume then that  $r(A(w_2)) = \gamma e_4 + \delta e_3$ . Suppose  $\gamma \neq 0$ . The element  $e_4$  in  $C_{i_1} \subset F$  satisfies  $e_4 \text{Tr}(A(w_1))$  and  $\{r(A(w_2)), e_4, r(A(w_2))\}$  $\gamma^2 e_4 + \delta \gamma e_3 \neq 0.$ 

Proposition [4.5](#page-14-0) implies that  $r(A^{-1}(e_4)) \top r(A^{-1}(r(A(w_1))))$ . Observe that by Lemma [4.1](#page-9-1)(*d*) we have  $r(A^{-1}(r(A(w_1)))) = e^{i\theta} w_1$  for some real  $\theta$ . Now, it follows from  $r(A^{-1}(e_4)) \top r(A^{-1}(r(A(w_1))))$  that  $w_1 \top r(A^{-1}(e_4))$ .

On the other hand, since  $w_1 \perp w_2$  and  $A^{-1}(e_4) = \gamma(e_4, A^{-1})r(A^{-1}(e_4)) \in$  $E_1(w_1)$ , we deduce from Lemma [4.6](#page-14-2) that  $A^{-1}(e_4) \in {}^{\perp_q}(w_2)$ . Therefore the hypotheses on the mapping A lead to  $e_4 \in {}^{\perp_q} \{A(w_2)\}\,$ , which contradicts that

$$
\{r(A(w_2)), v_4, r(A(w_2))\} \neq 0.
$$

This implies that  $\gamma = 0$ , and hence

$$
A(w_2) = \gamma(w_2, A)\delta e_3 \perp A(w_1) = \gamma(w_1, A)r(A(w_1)),
$$

as desired.

(ii) There exist a minimal tripotent  $\tilde{e} \in C_{i_1}$  and a rank-two tripotent  $u \in C_{i_1}$ such that  $(r(A(w_1)), u, \tilde{e})$  is a trangle and  $r(A(w_2)) = \alpha r(A(w_1)) + \beta u + \delta \tilde{e}$  with  $\alpha, \beta, \delta \in \mathbb{C}$  and  $\alpha\delta = \beta^2$ . The condition  $w_2 \in {}^{\perp_q}(w_1)$  (equivalently,  $A(w_2) \in$  $\perp^q \{A(w_1)\}\$  gives  $\alpha = 0 = \beta$ , and thus  $A(w_2) = \gamma(w_2, A)\delta\tilde{e}$  is orthogonal to  $A(w_1) =$  $\gamma(w_1, A)r(A(w_1)).$ 

## 5. ATOMIC JBW<sup>\*</sup>-TRIPLES

Let E be a JBW<sup>\*</sup>-triple with predual  $E_*$ . The extreme points of the closed unite ball of  $E_*$  are called *atoms* or *pure atoms* of E. The symbol  $\partial_e(\mathcal{B}_{E_*})$  will denote the set of all atoms of E. A celebrated result by Y. Friedman and B. Russo shows that for each  $\varphi \in \partial_e(\mathcal{B}_{E_*})$  there exists a unique minimal tripotent e in E satisfying  $P_2(e)(x) = \varphi(x)e$  for all  $x \in E$  (cf. [\[25,](#page-30-12) Proposition 4]). In case that E is an atomic JBW<sup>\*</sup>-triple,  $E_*$  is precisely the norm closure of the linear span of the atoms of E [\[25,](#page-30-12) Theorem 1]. In particular,  $\partial_e(\mathcal{B}_{E_*})$ , equivalently  ${P_2(e) : e \text{ minimal tripotent in } E},$  is a norming set for E.

A (closed) subtriple I of a JB<sup>∗</sup> -triple E is said to be an *ideal* (respectively, an *inner ideal*) of E if  $\{E, E, I\}$  +  $\{E, I, E\}$   $\subseteq$  I (respectively,  $\{I, E, I\}$   $\subseteq$  I). If e is a tripotent of E, it follows from Peirce arithmetic that  $E_0(e)$  and  $E_2(e)$  are inner ideals of  $E$ . A JBW<sup>\*</sup>-triple  $E$  is called a factor if there does not exist a decomposition of E as a direct sum of two non-zero ideals  $I, J$ , or equivalently, if {0} and E are the only weak<sup>∗</sup> -closed ideals of E. The Cartan factors are precisely the JBW<sup>\*</sup>-triple factors  $E$  containing a minimal tripotent (cf.  $[30,$  Corollary 1.8]). It is known that every tripotent  $e$  in a Cartan factor  $C$  admits a representation  $e = \sum_i e_i$ , where the series converges with respect to the weak<sup>\*</sup>-topology of C and  ${e_i}_i$  is a family of mutually orthogonal minimal tripotents in C (cf. [\[36,](#page-30-19) page 200]).

Let F be a JBW<sup>\*</sup>-subtriple of a JBW<sup>\*</sup>-triple E, and let e be a tripotent in F. Clearly, e being minimal in  $E$  implies that e is minimal in  $F$ . We cannot, in general, conclude that  $e$  being minimal in  $F$  is equivalent to  $e$  being minimal in  $E$ . However, if F is an inner ideal of E, Lemma 3.3 in  $[9]$  assures that every minimal tripotent in  $F$  is minimal in  $E$ .

<span id="page-16-0"></span>For later purposes we need to revisit some structure results and characterizations of atomic JBW<sup>∗</sup> -triples, which are not explicit in the literature.

## Proposition 5.1. *Let* E *be a JBW*<sup>∗</sup> *-triple. Then the following statements hold:*

- (a) E *is atomic if, and only if, for every non-zero tripotent* v *in* E *there exists a minimal tripotent*  $e \in E$  *satisfying*  $e \leq v$ .
- (b) If E is atomic and e is a tripotent in E, the Peirce subspaces  $E_i(e)$  are atomic  $JBW^*$ -triples for all  $j \in \{0, 1, 2\}$ .

*Proof.* (a) The "only if" implication is clear by structure theory, as we have seen above, every non-zero tripotent  $v$  in a Cartan factor, and hence in an atomic JBW<sup>\*</sup>triple E, can be written in the form  $v = \text{weak}^*$ - $\sum_i e_i$ , where  $\{e_i\}_i$  is a family of mutually orthogonal minimal tripotents in E, and hence  $e_i \leq v$  for all i.

For the "if" implication, recall that every element  $a$  in a JBW\*-triple  $E$  can be approximated in norm by a finite linear combination of mutually orthogonal non-zero tripotents  $e_1, \ldots, e_k$  in E (cf. [\[29,](#page-30-11) Lemma 3.11]). Let us fix a non-zero tripotent  $e_j$ . By hypothesis, there exists a minimal tripotent v satisfying  $v \leq e_j$ . We can find, via Zorn's lemma, a maximal family  $\{v_k^j\}_k$  of mutually orthogonal minimal tripotents in E with  $v_k^j \le e_j$ . The series  $\sum_k v_k^j$  is summable in the weak<sup>\*</sup>topology of E and  $w^*$ - $\sum_k v_k^j \le e_j$  [\[29,](#page-30-11) Corollary 3.13]. If  $e_j - \sum_k v_k^j \ne 0$ , then by the assumptions on E, there exists another minimal tripotent w with  $w \leq$  $e_j - \sum_k v_k^j$ , which implies that  $\{v_k^j\}_k \cup \{w\}$  is another family of mutually orthogonal minimal tripotents in E bounded by  $e_j$ , which is impossible. Therefore,  $e_j$  can be approximated in the weak<sup>∗</sup> -topology by a finite sum of mutually orthogonal minimal tripotents in  $E$ . Consequently,  $a$  can be approximated in the weak<sup>\*</sup>-topology of  $E$ by a finite linear combination of mutually orthogonal minimal tripotents in E.

(b) By Peirce arithmetic, the Peirce subspaces  $E_0(e)$  and  $E_2(e)$  are inner ideals of E. By applying that E is atomic and  $(a)$ , we deduce that for each non-zero tripotent v in  $E_j(e)$ , there exists a minimal tripotent  $w \in E$  with  $w \leq v$ . Observe that in this case  $w = \{v, w, v\} \in \{E_j(e), E, E_j(e)\}$ , and having in mind that  $E_j(e)$ is an inner ideal for all  $j = 0, 2$ , we conclude that w is a minimal tripotent in  $E_j(e)$  $(j = 0, 2)$ . It follows from  $(a)$  that  $E_j(e)$  is an atomic JBW<sup>\*</sup>-triple for all  $j = 0, 2$ .

We deal next with  $E_1(e)$ . By assumptions, E is an atomic JBW<sup>\*</sup>-triple, and hence,  $E = \bigoplus^{\ell_{\infty}}$ 

k  $C_k$  for a certain family of Cartan factors  $\{C_k\}$ . Let  $e = (e_k)$  be

a tripotent in E (where each  $e_k$  is a tripotent in  $C_k$ ). Since  $E_j(e)$  coincides with the  $\ell_{\infty}$ -sum of the family  $\{(C_k)_i(e_k)\}_k$ , we can clearly assume that E is a Cartan factor. If E has finite rank, the Peirce-j subspace  $E_i(e)$  also has finite rank, and it is known that in such a case  $E_j(e)$  must be a reflexive atomic JB<sup>\*</sup>-triple ([\[6,](#page-29-5) Proposition 4.5,  $[4,$  Proof of Theorem 2.3 and  $[14]$ . We can therefore assume that E is a Cartan factor with infinite rank.

We shall distinguish the three remaining cases for  $j = 1$ .

 $E = B(H, K)$  is a type 1 Cartan factor, where H and K are infinite dimensional complex Hilbert spaces. Each tripotent  $e$  in  $E$  is a partial isometry (equivalently,  $ee^*$  is a projection in  $B(K)$  and  $e^*e$  is a projection in  $B(H)$ ), and  $E_1(e)$  $ee^*B(H,K)(1-e^*e) \oplus (1-ee^*)B(H,K)e^*e$ . It is easy to check that the summands  $ee^*B(H,K)(1-e^*e) = B(ee^*(H), (1-e^*e)(K))$  are  $B((1-ee^*)(H), e^*e(K))$  are orthogonal type 1 Cartan factors, which proves that  $E_1(e)$  is an atomic JBW<sup>\*</sup>-triple. Observe that  $E_1(e)$  is not, in general, a factor.

Let j be a conjugation on a complex Hilbert space H, and set  $a^t = ja^*j$  for all  $a \in B(H)$ . Pick a tripotent e in the type 2 Cartan factor

$$
C = B(H)_{skew} = \{a \in B(H) : a^t = -a\}.
$$

Clearly, e is a partial isometry in  $B(H)$  satisfying  $e^t = -e$ ,  $e^*j = -je$ , and  $je^* =$  $-ej$ . For each  $a \in C_1(e)$ , the elements  $a_1 = ee^*a(1-e^*e)$  and  $a_2 = (1-e^*)ae^*e$  are orthogonal with  $a_2 = -a_1^t$  and  $a = a_1 + a_2$ . The mapping  $a \mapsto a_1 = ee^*a(1 - e^*e)$ is an isometric triple isomorphism from  $C_1(e)$  onto  $B((1-e^*e)(H), ee^*(H))$ , which proves that  $C_1(e)$  is a Cartan factor. The case of the type 3 Cartan factor  $C =$  $B(H)_{symm} = \{a \in B(H) : a^t = a\}$  follows via similar arguments.

<span id="page-17-1"></span>The following identity principle is almost explicit in [\[27,](#page-30-1) Theorem 3.7], we include here a simplified version for completeness reasons.

**Proposition 5.2.** *Let*  $\Delta : E \to E$  *be a (non-necessarily additive) bijection preserving truncations and inner quadratic annihilators of elements, where* E *is an atomic JBW*<sup>∗</sup> *-triple. Suppose additionally that for each minimal tripotent* e *in* E *and each complex number*  $\lambda$  *we have*  $\Delta(\lambda e) = \lambda e$ *. Then*  $\Delta$  *is the identity mapping on* E.

*Proof.* Let us take an element  $a \in E$ , and a minimal tripotent  $e \in E$  supported at a pure atom  $\varphi_e$  in  $E_*$ . We reduce to the following two cases:

*Case 1.* If  $\varphi_e(a) \neq 0$  then  $\varphi_e(a)e$  is a truncation of a since

$$
\{\varphi_e(a)e, \varphi_e(a)e, \varphi_e(a)e\} = \varphi_e(a)^2 \overline{\varphi_e(a)}e = \varphi_e(a)^2 Q(e) P_2(e)(a)
$$
  
= 
$$
\varphi_e(a)^2 Q(e)(a) = \{\varphi_e(a)e, a, \varphi_e(a)e\}.
$$

The hypotheses on  $\Delta$  imply that  $\Delta(\varphi_e(a)e) = \varphi_e(a)e$  is a truncation of  $\Delta(a)$ , and hence

$$
\varphi_e(a)^2 \overline{\varphi_e(a)} e = \{ \Delta(\varphi_e(a)e), \Delta(\varphi_e(a)e), \Delta(\varphi_e(a)e) \}
$$
  
= 
$$
\{ \Delta(\varphi_e(a)e), \Delta(a), \Delta(\varphi_e(a)e) \} = \varphi_e(a)^2 \{ e, \Delta(a), e \}
$$
  
= 
$$
\varphi_e(a)^2 \overline{\varphi_e(\Delta(a))} e,
$$

which gives

 $\varphi_e(a) = \varphi_e(\Delta(a))$ , equivalently,  $\varphi_e(a - \Delta(a)) = 0$ .

*Case 2.* If  $\varphi_e(a) = 0$ , equivalently,  $P_2(e)(a) = 0 \Leftrightarrow a \in {\perp_q} \{e\}$ , and thus, the assumptions on  $\Delta$  assure that  $\Delta(a) \in {}^{\perp_q} {\{\Delta(e)\}} = {}^{\perp_q} {e}$ , therefore  $\varphi_e(\Delta(a) - a) =$  $\varphi_e(\Delta(a)) - \varphi_e(a) = 0.$ 

We have proved that  $\varphi_e(a - \Delta(a)) = 0$ , for all  $e \in \mathcal{U}_{min}(E)$ ,  $a \in E$ . The desired conclusion follows from the fact that pure atoms of E separate the points in E.  $\Box$ 

<span id="page-17-0"></span>In our next result we describe the Peirce-1 subspace associated with a minimal tripotent in an atomic JB<sup>∗</sup> -triple in terms of inner quadratic annihilators.

Lemma 5.3. *Let* e *be a minimal tripotent in an atomic JBW*<sup>∗</sup> *-triple* E*. Then we have*

$$
E_1(e) = {}^{\perp_q} \{e\} \bigcap \left( \bigcap_{\substack{v \in \mathcal{U}_{min}(E) \\ v \perp e}} {}^{\perp_q} \{v\} \right) = {}^{\perp_q} \{ \mathbb{C}e\} \bigcap \left( \bigcap_{\substack{v \in \mathcal{U}_{min}(E) \\ v \perp e}} {}^{\perp_q} \{ \mathbb{C}v\} \right).
$$

*Proof.* The second equality is clear, we shall only prove the first one.

 $(2)$  Take an element a in the intersection given in the right-hand-side above. Since  $a \in {}^{\perp_q}{e}$  it follows that  $a \in E_1(e) \oplus E_0(e)$ . Similarly  $a \in E_1(v) \oplus E_0(v)$ for all  $v \in \mathcal{U}_{min}(E)$  with  $v \perp e$ , equivalently, for all  $v \in \mathcal{U}_{min}(E_0(e))$ . Therefore  $P_2(v)(P_0(e)(a)) = 0$  for all  $v \in \mathcal{U}_{min}(E_0(e))$ . Having in mind that  $E_0(e)$  is an atomic JBW<sup>\*</sup>-triple (see Proposition [5.1](#page-16-0)(b)), and the fact that the set  $\{P_2(v)$ :  $v \in \mathcal{U}_{min}(E_0(e))$  separates the points of  $E_0(e)$ , we get  $P_0(e)(a) = 0$ , and hence  $a = P_1(e)(a) \in E_1(e)$ .

(⊆) Take now  $a \in E_1(e)$ . Clearly  $a \in \perp^q \{e\} = E_0(e) \oplus E_1(e)$ . If  $v \in \mathcal{U}_{min}(E)$ with  $v \perp e$ , it follows from Lemma [4.6](#page-14-2) that  $E_1(e) \subseteq {}^{\perp_q}{v}$ .

We can now show that every surjective additive mapping preserving truncations in both directions between atomic JBW<sup>∗</sup> -triples preserves Peirce-1 subspaces associated to minimal tripotents.

<span id="page-18-1"></span>**Proposition 5.4.** *Let* E and F *be atomic JBW<sup>\*</sup>-triples, and let*  $A : E \to F$  *be a surjective additive mapping preserving truncations in both directions. Then for each minimal tripotent e in* E *we have*  $A(E_1(e)) = F_1(r(A(e)))$ .

*Proof.* Lemma [4.1](#page-9-1)(c),  $A(\mathbb{C} \mathcal{U}_{min}(E)) = \mathbb{C} \mathcal{U}_{min}(F)$ . Take any  $v \in \mathcal{U}_{min}(E)$  with  $v \perp e$ . Lemma [4.1](#page-9-1)(d) assures that  $A(\mathbb{C}v) = \mathbb{C}A(v) = \mathbb{C}r(A(v))$ , while Proposi-tion [4.7](#page-14-1) proves that  $r(A(v)) \perp r(A(e))$ . This proves that

$$
A(\mathbb{C} \ \{v \in \mathcal{U}_{min}(E) : v \perp e\}) \subseteq \mathbb{C} \ \{w \in \mathcal{U}_{min}(F) : w \perp r(A(e))\}.
$$

Similarly, since  $r(A^{-1}(r(A(e)))) \in \mathbb{T}e$  (see Lemma [4.1](#page-9-1)(*e*)),

$$
A^{-1}(\mathbb{C} \{w \in \mathcal{U}_{min}(F): w \perp r(A(e))\}) \subseteq \mathbb{C} \{v \in \mathcal{U}_{min}(E): v \perp e\},\
$$

and thus

<span id="page-18-0"></span>
$$
A\left(\mathbb{C}\ \left\{v\in\mathcal{U}_{min}(E):v\perp e\right\}\right)=\mathbb{C}\ \left\{w\in\mathcal{U}_{min}(F):w\perp r(A(e))\right\}.\tag{4}
$$

Finally, by Lemma [5.3,](#page-17-0) [\(4\)](#page-18-0), and the fact that A preserves inner quadratic annihilators we derive that

$$
A(E_1(e)) = A\left(\bigcup_{q \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} A\left(\bigcup_{q \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} A\left(\bigcup_{q \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} A\left(\bigcup_{q \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} A\left(\bigcup_{q \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} A\left(\bigcup_{q \in \mathcal{U}_{min}(E)} A\left(\bigcup_{q \in \mathcal{U}_{min}(E)} \bigcap_{v \in \mathcal{U}_{min}(E)} A\left(\bigcup_{q \in \mathcal{U}_{min
$$

$$
= {}^{\perp_q} \{ \mathbb{C}r(A(e)) \} \bigcap \left( \bigcap_{\substack{v \in \mathcal{U}_{min}(E) \\ v \perp e}} {}^{\perp_q} \{ \mathbb{C}r(A(v)) \} \right)
$$
  

$$
= {}^{\perp_q} \{ \mathbb{C}r(A(e)) \} \bigcap \left( \bigcap_{\substack{w \in \mathcal{U}_{min}(F) \\ w \perp r(A(e))}} {}^{\perp_q} \{ \mathbb{C}w \} \right) = F_1(r(A(e))).
$$

We have therefore shown that  $A(E_1(e)) \subseteq F_1(r(A(e)))$ . Similarly,

$$
A^{-1}\Big(F_1(r(A(e)))\Big) \subseteq E_1\Big(r\big(A^{-1}(r(A(e)))\big)\Big) = E_1(e),
$$

<span id="page-19-1"></span>which gives  $A(E_1(e)) = F_1(r(A(e)))$ .

Corollary 5.5. Let E and F be atomic JBW<sup>\*</sup>-triples, and let  $A : E \to F$  be a *surjective additive mapping preserving truncations in both directions. Then for each rank-2 tripotent e in E we have*  $A(E_2(e)) = F_2(r(A(e)))$ .

*Proof.* Since  $e$  has rank-2, we can find two orthogonal minimal tripotents  $e_1, e_2$  in E such that  $e = e_1 + e_2$ . It is well-known that

$$
E_2(e) = E_2(e_1) \oplus E_1(e_1) \cap E_1(e_2) \oplus E_2(e_2) = \mathbb{C}e_1 \oplus E_1(e_1) \cap E_1(e_2) \oplus \mathbb{C}e_2.
$$

Observe that  $A(e) = A(e_1) + A(e_2) = \gamma(e_1, A)r(A(e_1)) + \gamma(e_2, A)r(A(e_2))$ , where  $r(A(e_1))$  and  $r(A(e_2))$  are two orthogonal minimal tripotents in F, and hence  $r(A(e)) = r(A(e_1)) + r(A(e_2)).$  Furthermore,  $A^{-1}(r(A(e))) = A^{-1}(r(A(e_1))) +$  $A^{-1}(r(A(e_2)))$ , and hence

$$
r\Big(A^{-1}\Big(r(A(e))\Big)\Big) = \gamma_1 e_1 + \gamma_2 e_2,
$$

for suitable  $\gamma_1, \gamma_2 \in \mathbb{T}$ .

The additivity of A, Lemma [4.1,](#page-9-1) Proposition [4.7,](#page-14-1) and Proposition [5.4](#page-18-1) lead to  $A(E_2(e)) = \mathbb{C}r(A(e_1)) \oplus A(E_1(e_1) \cap E_1(e_2)) \oplus \mathbb{C}r(A(e_2))$ 

 $\subseteq \mathbb{C}r(A(e_1)) \oplus A(E_1(e_1)) \cap A(E_1(e_2)) \oplus \mathbb{C}r(A(e_2))$ 

 $=\mathbb{C}r(A(e_1)) \oplus F_1(r(A(e_1))) \cap F_1(r(A(e_2))) \oplus \mathbb{C}r(A(e_2)) = F_2(r(A(e))).$ 

That is,  $A(E_2(e)) \subseteq F_2(r(A(e)))$ . Similarly

$$
A^{-1}\Big(F_2\big(r(A(e))\big)\Big) \subseteq E_2\Big(r\big(A^{-1}(r(A(e))\big)\Big) = E_2(\gamma_1e_1 + \gamma_2e_2) = E_2(e),
$$
  
which concludes the proof.

<span id="page-19-0"></span>We can now extend some useful consequences of our previous results.

**Proposition 5.6.** *Let* E and F be atomic JBW<sup>\*</sup>-triples, that is,  $E = \bigoplus_{k=1}^{\ell_{\infty}}$  $k \in \Gamma_1$ C<sup>k</sup> *and*

 $F = \bigoplus^{\ell_{\infty}}$  $j \in \Gamma_2$  $C_j$ , where  $\{C_k\}_{k\in\Gamma_1}$  and  $\{C_j\}_{j\in\Gamma_2}$  are two families of Cartan factors.

*Suppose, additionally, that* E *contains no one-dimensional Cartan factors as direct summands* (*i.e.,*  $dim(C_k) \geq 2$  *for all* k). Let  $A : E \to F$  be a surjective additive

*mapping preserving truncations in both directions. Then the following statements hold:*

- (a) F *contains no one-dimensional Cartan factors as direct summands.*
- (b) For each  $k \in \Gamma_1$  there exists a unique  $\sigma(k) \in \Gamma_2$  such that  $A(C_k) = \tilde{C}_{\sigma(k)}$ , and *the mapping*  $\sigma : \Gamma_1 \to \Gamma_2$  *is a bijection. Furthermore, the ranks of*  $C_k$  *and*  $\tilde{C}_{\sigma(k)}$  *coincide.*

*Proof.* (a) Arguing by contradiction we assume that  $C_{j_0} = \mathbb{C}$  for some  $j_0$ . Let  $e_{j_0}$ be any norm-one element in  $C_{j_0}$ . Observe that  $e_{j_0}$  is a minimal tripotent in F and  $F_1(e_{j_0}) = \{0\}$ . Let us write  $A^{-1}(e_{j_0}) = \gamma(e_{j_0}, A^{-1})r(A^{-1}(e_{j_0}))$ , where  $r(A^{-1}(e_{j_0}))$ is a minimal tripotent in E and  $\gamma(e_{j_0}, A^{-1}) > 0$  (cf. Lemma [4.1](#page-9-1) and Remark [4.2\)](#page-9-2). It follows that  $r(A^{-1}(e_{j_0}))$  belongs to a unique Cartan factor  $C_{k_0}$  in the decomposition of E. We know from the hypotheses that dim  $(C_{k_0}) \geq 2$ . Therefore,  $E_1(r(A^{-1}(e_{j_0}))\neq \{0\}$ . Lemma [4.1](#page-9-1)(e) implies that  $r(A(r(A^{-1}(e_{j_0})))) \in \mathbb{T}e_{j_0}$ , and hence Proposition [5.4](#page-18-1) implies that

$$
\{0\} \neq A\Big(E_1\big(r(A^{-1}(e_{j_0}))\big)\Big) = F_1\Big(r\Big(A\big(r(A^{-1}(e_{j_0}))\big)\Big)\Big) = F_1(e_{j_0}) = \{0\},\
$$

which is impossible.

(b) Fix  $k \in \Gamma_1$  and a minimal tripotent  $e_k \in C_k$ . Lemma [4.1](#page-9-1)(c) (see also Remark [4.2\)](#page-9-2) assures that  $r(A(e_k))$  is a minimal tripotent in F, and hence it belongs to a unique Cartan factor in the decomposition of  $F$ , we denote this latter factor by  $C_{\sigma(k)}$ .

Given a minimal tripotent  $w \in C_k$  with  $w \top e_k$ , Proposition [4.5](#page-14-0) assures that  $r(A(w))\top r(A(e_k))$ , and thus  $r(A(w)), A(w) \in C_{\sigma(k)}$ .

Let v be any other minimal tripotent in  $C_k$ .

If  $C_k$  has rank-one, it is a complex Hilbert space with dimension  $\geq 2$ , it follows form the discussion in Remark [3.1](#page-7-1) that we can find a minimal tripotent  $w \in C_k$ such that  $w\top e_k$  and v is a linear combination of w and  $e_k$ . It follows that  $A(v) \in$  $A(\mathbb{C}w) + A(\mathbb{C}e_k) = \mathbb{C}r(A(w)) + \mathbb{C}r(A(e_k)) \in C_{\sigma(k)}$  (cf. Lemma [4.1](#page-9-1)(*e*)).

Assuming that  $C_k$  has rank  $\geq 2$ , it follows from [\[21,](#page-30-21) Lemma 3.10] that one of the next statements holds:

(1) There exists a quadrangle of minimal tripotents  $(e_k, e_2, e_3, e_4)$  such that v is a linear combination of  $e_k, e_2, e_3$ , and  $e_4$ , and thus

$$
A(v) \in \mathbb{C}r(A(e_k)) + \mathbb{C}r(A(e_2)) + \mathbb{C}r(A(e_3)) + \mathbb{C}r(A(e_4)) \in \widetilde{C}_{\sigma(k)}.
$$

(2) There exists a trangle of the form  $(e_k, u, \tilde{e}_k)$ , such that  $\tilde{e}_k$  is minimal, u is a rank-2 tripotent, and v is a linear combination of  $e_k$ ,  $u$ , and  $\tilde{e}_k$ . Observe that  $\frac{e_k \pm u + \tilde{e}_k}{2}$  are minimal tripotents with  $\frac{e_k \pm u + \tilde{e}_k}{2} \notin {}^{\perp_q}$  {Ce<sub>k</sub>} (cf. Lemma [4.3\)](#page-10-0). It follows that  $A(\frac{e_k \pm u + \tilde{e}_k}{2}) \notin {}^{\perp_q} {\{ \mathbb{C}A(e_k) \}} = {}^{\perp_q} {\{ r(A(e_k)) \}},$  and thus  $A(\frac{e_k \pm u + \tilde{e}_k}{2}) \in$  $C_{\sigma(k)}$ . We therefore have

$$
A(v) \in A(\mathbb{C}e_k) + A(\mathbb{C}u) + A(\mathbb{C}\tilde{e}_k)
$$
  
\n
$$
\subseteq \mathbb{C}A(e_k) + \mathbb{C}A\left(\frac{e_k + u + \tilde{e}_k}{2}\right) + \mathbb{C}A\left(\frac{e_k - u + \tilde{e}_k}{2}\right) + \mathbb{C}A(\tilde{e}_k) \in \widetilde{C}_{\sigma(k)}.
$$

The previous arguments show that

$$
A\left(\mathbb{C}\,\mathcal{U}_{min}(C_k)\right)\subseteq\mathbb{C}\,\mathcal{U}_{min}(\widetilde{C}_{\sigma(k)})\subset\widetilde{C}_{\sigma(k)}.
$$

By applying a similar reasoning to  $A^{-1}$  we deduce that

<span id="page-21-0"></span>
$$
A\left(\mathbb{C}\,\mathcal{U}_{min}(C_k)\right) = \mathbb{C}\,\mathcal{U}_{min}(\widetilde{C}_{\sigma(k)}).
$$
\n<sup>(5)</sup>

The  $\sigma : \Gamma_1 \to \Gamma_2$ ,  $k \mapsto \sigma(k)$  is well-defined and a bijection because A is bijective.

Fix 
$$
k_0 \in \Gamma_1
$$
, and set  $C_{k_0}^{\perp} := \bigoplus_{k \in \Gamma_1, k \neq k_0}^{\ell_{\infty}} C_k$  and  $\widetilde{C}_{\sigma(k_0)}^{\perp} = \bigoplus_{j \in \Gamma_2, j \neq \sigma(k_0)}^{\ell_{\infty}} \widetilde{C}_j$ . Clearly,  
\n $C_{k_0}^{\perp}$  and  $\widetilde{C}_{\sigma(k_0)}^{\perp}$  are atomic JBW\*-triples with  $\mathcal{U}_{min}(C_{k_0}) = \bigcup_{k \in \Gamma_1, k \neq k_0}^{\ell_{\infty}} \mathcal{U}_{min}(C_k)$  and

 $\mathcal{U}_{min}\left(\tilde{C}_{k_0}^{\perp}\right) = \qquad \bigcup$  $j\in\Gamma_2,j\neq\sigma(k_0)$  $\mathcal{U}_{min}(\tilde{C}_j)$ . It follows from [\(5\)](#page-21-0) that

$$
A\left(\mathbb{C}\,\mathcal{U}_{min}\left(C_{k_0}^{\perp}\right)\right)=\mathbb{C}\,\mathcal{U}_{min}\left(\tilde{C}_{\sigma(k_0)}^{\perp}\right).
$$

For each  $x \in C_{k_0}$ , we have  $x \perp \mathbb{C}$   $\mathcal{U}_{min}(C_{k_0}^{\perp})$ , which assures that x lies in  $\perp^{\perp_q}\Big\{\mathbb{C}\ \mathcal{U}_{min}\left(C_{k_0}^{\perp}\right)\Big\},$  and hence

$$
A(x) \in {}^{\perp_q}\Big\{A\left(\mathbb{C}\,\mathcal{U}_{min}\left(C_{k_0}^{\perp}\right)\right)\Big\} = {}^{\perp_q}\Big\{ \mathbb{C}\,\mathcal{U}_{min}\left(\tilde{C}_{\sigma(k_0)}^{\perp}\right)\Big\}.
$$

This implies that  $P_2(v)(A(x)) = 0$ , for all  $v \in \mathcal{U}_{min}(\tilde{C}_{\sigma(k_0)}^{\perp})$  . Now, by applying that the set  $\left\{P_2(v) : v \in \mathcal{U}_{min}\left(\tilde{C}^{\perp}_{\sigma(k_0)}\right) \right\}$  separates the points in  $\tilde{C}^{\perp}_{\sigma(k_0)}$ , we arrive to  $A(x) \in \tilde{C}_{\sigma(k_0)}$ . We have therefore shown that  $A(C_{k_0}) \subseteq \tilde{C}_{\sigma(k_0)}$ , and consequently  $A(C_{k_0}) = \tilde{C}_{\sigma(k_0)}.$ 

The final statement is a consequence of Proposition [4.7](#page-14-1) and the fact that in an atomic JBW<sup>∗</sup> -triple the rank coincides with the cardinality of a maximal set of mutually orthogonal minimal tripotents.

The next proposition relies on the main result in [\[37\]](#page-30-6) combined with the presence of JB<sup>∗</sup> -subtriples isometrically isomorphic to Hilbert spaces of dimension bigger than or equal to 2.

<span id="page-21-1"></span>**Proposition 5.7.** Let E and F be JBW<sup>\*</sup>-triples, and let  $A: E \to F$  be a surjective *additive mapping preserving truncations in both directions. Suppose that*  $e_1, \ldots, e_n$ *are mutually collinear minimal tripotents in* E *with*  $n \geq 2$ . Then the restriction of A *to the JB<sup>\*</sup>-subtriple of* E generated by  $e_1, \ldots, e_n$  *is a positive scalar multiple of a* (*complex) linear or a conjugate-linear isometry.*

*Proof.* By Lemma [4.1](#page-9-1) and Proposition [4.5,](#page-14-0) the elements  $r(A(e_1)), \ldots, r(A(e_n))$  are mutually collinear minimal tripotents in  $F$ . Observe that the JB<sup>\*</sup>-subtriple generated by  $e_1, \ldots, e_n$  (respectively, by  $r(A(e_1)), \ldots, r(A(e_n)))$ ) is isometrically isomorphic to an *n*-dimensional complex Hilbert space in which  $e_1, \ldots, e_n$  (respectively,  $r(A(e_1)), \ldots, r(A(e_n)))$  is an orthonormal basis (cf. [\[15,](#page-30-20) Lemma in page 306]). Let  $E_0$  and  $F_0$  denote the JB<sup>\*</sup>-subtriples of E and F generated by  $e_1, \ldots, e_n$  and

 $r(A(e_1)), \ldots, r(A(e_n))$ , respectively. A new application of Lemma [4.1](#page-9-1) and the additivity of A implies that

$$
A(E_0) = A(\mathbb{C}e_1 \oplus \ldots \oplus \mathbb{C}e_n) = \mathbb{C}r(A(e_1)) \oplus \ldots \oplus \mathbb{C}r(A(e_n)) = F_0,
$$

that is,  $A|_{E_0}$  is a bijective additive mapping preserving truncations in both directions. Proposition [3.3](#page-8-1) gives the desired conclusion.

<span id="page-22-0"></span>Back to the setting of atomic JBW<sup>\*</sup>-triples set in Proposition [5.6](#page-19-0) we can improve our previous conclusions.

Proposition 5.8. *Under the hypotheses of Proposition [5.6,](#page-19-0) the following statements hold:*

- (a) For each minimal tripotent  $e \in E$  one of the next statements holds: (a.1)  $A(\lambda e) = \gamma(e, A)\lambda r(A(e)) = \lambda A(e)$ , *for all*  $\lambda \in \mathbb{C}$ *.*  $(a.2)$   $A(\lambda e) = \gamma(e, A)\overline{\lambda}r(A(e)) = \overline{\lambda}A(e)$ , *for all*  $\lambda \in \mathbb{C}$ *.*
- (b) *If*  $e, v \in \mathcal{U}_{min}(E)$  *with*  $e \top v$ *, we have*  $\gamma(e, A) = \gamma(v, A)$ *.*
- (c) The mapping  $A_r : \mathcal{U}_{min}(E) \to \mathcal{U}_{min}(F)$ ,  $e \mapsto A_r(e) := r(A(e))$  is a bijective *mapping preserving collinearity and orthogonality in both directions.*

*Proof.* (a) Let us fix an arbitrary minimal tripotent  $e \in E$ . Suppose v is another minimal tripotent in E such that  $v\tau e$  (we can always assume the existence of such minimal tripotent v since  $\dim(C_k) \geq 2$  for all  $k \in \Gamma_1$ , compare, for example Section [3](#page-7-0) and  $[21, \text{ Lemma } 3.10]$ . Let  $E_0$  denote the JB<sup>\*</sup>-subtriple of E generated by e and v. By Proposition [5.7](#page-21-1) the restriction of A to  $E_0$  is a positive scalar multiple of a (complex) linear or a conjugate-linear isometry, that is, there exists  $\gamma_{E_0} > 0$  such that  $\gamma_{E_0}^{-1} A|_{E_0}$  is a (complex) linear or a conjugate-linear isometry. Consequently, by Remark [4.2,](#page-9-2) one of the next statements holds:

- $(a.1) A(\lambda e) = \gamma(e, A)\lambda r(A(e)) = \lambda A(e)$  for all  $\lambda \in \mathbb{C}$ , and the same identity holds when  $e$  is replaced by any norm-one element in  $E_0$ .
- $(a.2)$   $A(\lambda e) = \gamma(e, A)\overline{\lambda}r(A(e)) = \overline{\lambda}A(e)$  for all  $\lambda \in \mathbb{C}$ , and the same identity holds when  $e$  is replaced by any norm-one element in  $E_0$ .

It also follows from the above that

$$
\gamma(e, A) = \|A(e)\| = \gamma_{E_0} \|e\| = \gamma_{E_0} \|v\| = \|A(v)\| = \gamma(v, A),
$$

which proves the statement in  $(b)$ .

(c) It follows from Lemma [4.1](#page-9-1)(a) and (b) that the mapping  $A_r$  is well-defined and injective. The surjectivity of  $A_r$  is a straightforward consequence of (a) and the commented Lemma [4.1.](#page-9-1) The remaining properties of  $A<sub>r</sub>$  follow from Proposition [4.5](#page-14-0) and Proposition [4.7.](#page-14-1)

As we shall see later, three dimensional spin factors deserve to be treated independently. It is well-known that each three dimensional Hilbert space can be isometrically identified with the complex Banach space  $S_2$  of all  $2 \times 2$  symmetric complex matrices of the form  $\begin{pmatrix} \alpha & \beta \\ \beta & \delta \end{pmatrix}$  with  $\alpha$ ,  $\beta$  and  $\delta$  in  $\mathbb C$  (see, for example, [\[32,](#page-30-22) proof of Lemma 3.4]).

<span id="page-22-1"></span>**Proposition 5.9.** Let F be an atomic JBW<sup>\*</sup>-triple expressed in the form  $F =$  $\overset{\ell_{\infty}}{\bigtriangleup}$  $j \in \Gamma_2$  $C_j$ , where  $\{C_j\}_{j\in\Gamma_2}$  is a family of Cartan factors. Let  $A : S_2 \to F$  be a *surjective additive mapping preserving truncations in both directions. Then the following statements hold:*

- $(a) F = S_2.$
- (b) A *is a positive scalar multiple of a surjective (complex) linear or a conjugatelinear isometry.*

*Proof.* (a) It follows from Proposition [5.6](#page-19-0)(b) that F must be a Cartan factor of rank-two. Let  $e_1 =$  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ ,  $e_2 =$  $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ , and  $u =$  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ . By combining that A is surjective with Proposition [5.8](#page-22-0) we get

$$
F = A(\mathbb{C}e_1 \oplus \mathbb{C}u \oplus \mathbb{C}e_2) = \mathbb{C}r(A(e_1)) \oplus \mathbb{C}A(u) \oplus \mathbb{C}r(A(e_2)),
$$

where  $r(A(e_1))$  and  $r(A(e_2))$  are mutually orthogonal minimal tripotents in F, and hence  $2 \leq \dim(F) \leq 3$ . Comparing [\[36,](#page-30-19) Table in page 210] or [\[35,](#page-30-25) Table in page 475]. the unique Cartan factor satisfying these properties is  $S_2$ , that is, the 3-dimensional spin factor.

(b) We keep the notation above. Note that  $(S_2)_1(e_1) = (S_2)_1(e_2) = \mathbb{C}u$ , and thus Proposition [5.4,](#page-18-1) implies that  $\mathbb{C}A(u) = A(\mathbb{C}u) = F_1(r(A(e_1))) = F_1(r(A(e_2))).$ It is known that we can find a linear triple automorphism  $\Phi: S_2 \to S_2$  satisfying  $\Phi(r(A(e_1))) = e_1, \Phi(r(A(e_2))) = e_2$  and  $\Phi(r(A(u))) = \mu u$  for some  $\mu \in \mathbb{T}$  (see, for example, [\[36,](#page-30-19) Proposition 5.8]). The mapping  $\Phi A : S_2 \to S_2$  satisfies the same hypotheses of A. We also know that  $\mathbb{C}r(A(u)) = F_1(r(A(u))) \ni A(u) = \alpha r(A(u))$  for some positive  $\alpha$ . Observe that the elements  $v_{\pm} := \frac{e_1 \pm u + e_2}{2}$  are mutually orthogonal minimal tripotents (cf. Lemma [4.3\)](#page-10-0), and thus by Lemma [4.1](#page-9-1) and Proposition [4.7,](#page-14-1) their images under  $\Phi A$  must be positive scalar multiples of two mutually orthogonal minimal tripotents, that is,

$$
\Phi A \left( \frac{e_1 \pm u + e_2}{2} \right) = \frac{\gamma(e_1, A)e_1 \pm \alpha \mu u + \gamma(e_2, A)e_2}{2} = \begin{pmatrix} \gamma(e_1, A) & \pm \alpha \mu \\ \pm \alpha \mu & \gamma(e_2, A) \end{pmatrix}
$$

are positive scalar multiples of mutually orthogonal minimal tripotents, which assures that

$$
\gamma(e_1, A)\gamma(e_2, A) = \alpha^2 \mu^2 \Rightarrow \gamma(e_1, A)\gamma(e_2, A) = \alpha^2 \text{ and } \mu^2 = 1,
$$
  

$$
\gamma(e_1, A)^2 = \alpha^2 |\mu|^2, \text{ and } \gamma(e_2, A)^2 = \alpha^2 |\mu|^2.
$$

We have therefore shown that  $\gamma(e_1, A) = \alpha = \gamma(e_2, A), \mu = \pm 1.$ 

Having in mind Proposition  $5.8(a)$  $5.8(a)$  one of the next cases holds:

*Case* (I)  $A(\lambda e_1) = \lambda A(e_1)$ , for all  $\lambda \in \mathbb{C}$ . Again, by the dichotomy given by Proposition  $5.8(a)$  $5.8(a)$  one of the next statements is satisfied:

*Case*  $(I.a) A(\lambda v_+) = \lambda A(v_+)$  for all  $\lambda \in \mathbb{C}$ . The identity

$$
\lambda \frac{A(e_1) + A(u) + A(e_2)}{2} = \lambda A(v_+) = A(\lambda v_+) = \frac{A(\lambda e_1) + A(\lambda u) + A(\lambda e_2)}{2},
$$

combined with the fact that  $A(e_1) \perp A(e_2)$  and  $A(u) \in F_1(r(A(e_1))) = F_1(r(A(e_2))),$ implies that  $A(\lambda u) = \lambda A(u)$  and  $A(\lambda e_2) = \lambda A(e_2)$  for all  $\lambda \in \mathbb{C}$ . Therefore, since every  $x \in S_2$  writes uniquely in the form  $x = \lambda_1 e_1 + \lambda_2 u + \lambda_3 e_2$  for  $\lambda_j \in \mathbb{C}$ , we arrive to

$$
A(x) = \lambda_1 A(e_1) + \lambda_2 A(u) + \lambda_3 A(e_2),
$$

which clearly assures that  $A$  is (complex) linear. We further know that

$$
\Phi A(x) = \Phi A \left( \begin{pmatrix} \lambda_1 & \lambda_2 \\ \lambda_2 & \lambda_3 \end{pmatrix} \right) = \gamma(e_1, A) \begin{pmatrix} \lambda_1 & \mu \lambda_2 \\ \mu \lambda_2 & \lambda_3 \end{pmatrix} \quad (\lambda_1, \lambda_2, \lambda_3 \in \mathbb{C}),
$$

and consequently  $A$  is a positive scalar multiple of a surjective isometry on  $S_2$ . *Case* (*I.b*)  $A(\lambda v_+) = \overline{\lambda} A(v_+)$  for all  $\lambda \in \mathbb{C}$ . In this case, the identity

$$
\overline{\lambda} \frac{A(e_1) + A(u) + A(e_2)}{2} = \overline{\lambda} A(v_+) = A(\lambda v_+) = \frac{A(\lambda e_1) + A(\lambda u) + A(\lambda e_2)}{2},
$$

is incompatible with the assumptions we made at the beginning of this *Case* (I) since  $A(e_1) \perp A(e_2)$  and  $A(u) \in F_1(r(A(e_1))) = F_1(r(A(e_2))).$ 

*Case* (II)  $A(\lambda e_1) = \overline{\lambda}A(e_1)$  for all  $\lambda \in \mathbb{C}$ , we deduce, via similar arguments to those given in *Case*  $(I)$  above that A is positive scalar multiple of a conjugate-linear isometry.  $\Box$ 

In the next lemma we shall analyse when two minimal tripotents  $e$  and  $v$  in a Cartan factor C with rank  $\geq 2$  can be connected via a linear combination of the elements in a trangle but not in a quadrangle. Recall that the weak<sup>∗</sup> -closed ideal generated by any minimal tripotent in a Cartan factor  $C$  is the whole  $C$ .

<span id="page-24-1"></span>Lemma 5.10. *Let* e *and* v *be two minimal tripotents in a Cartan factor* C*. Assume that* C *has rank* $\geq$  2, *and we cannot find a quadrangle of minimal tripotents*  $(e, e_2, e_3, e_4)$  in C such that v is a linear combination of the elements in this quad*rangle. Then* C *is a Cartan factor of type* 3*, and there exists a trangle of the form*  $(e, u, \tilde{e})$  *with*  $\tilde{e}$  *minimal and u having rank-2 such that* v *is a linear combination of*  $e, u, and \tilde{e}, and the JBW<sup>∗</sup> -subtriple of C generated by this triangle coincides with$  $C_2(e + \tilde{e}) = C_2(u)$  *and is isometrically isomorphic to*  $S_2$ *.* 

*Proof.* Lemma 3.10 in [\[21\]](#page-30-21) combined with the hypotheses assure the existence of a trangle of the form  $(e, u, \tilde{e})$  with  $\tilde{e}$  minimal and u having rank-2, such that v is a linear combination of  $e, u$ , and  $\tilde{e}$ .

The tripotent  $u \in C_1(e)$  governs e, and hence  $e \not\top u$ . By combining this observation with the assumptions on  $C$ , the Classification Scheme in [\[15,](#page-30-20) page 305] (see also Proposition in page 308 in the same reference) assures that C is a Cartan factor of type 3. The rest is clear from [\[21,](#page-30-21) Lemma 3.10] and the general form of minimal and rank-2 tripotents in a type 3 Cartan factor.  $\Box$ 

<span id="page-24-0"></span>The behaviour of a surjective additive mapping preserving truncations in both directions on quadrangles of minimal tripotents is even easier to determine.

**Proposition 5.11.** Let E and F be atomic JBW<sup>\*</sup>-triples, that is,  $E = \bigoplus_{k=1}^{\ell_{\infty}}$  $k \in \Gamma_1$  $C_k$ 

and  $F = \bigoplus^{\ell_{\infty}}$  $j \in \Gamma_2$  $C_j$ , where  $\{C_k\}_{k\in\Gamma_1}$  and  $\{C_j\}_{j\in\Gamma_2}$  are two families of Cartan factors.

*Suppose, additionally, that* E *contains no one-dimensional Cartan factors as direct summands* (*i.e., dim*( $C_k$ )  $\geq$  2 *for all* k)*. Let*  $A : E \rightarrow F$  *be a surjective additive mapping preserving truncations in both directions. Suppose*  $(e_1, e_2, e_3, e_4)$  *is a quadrangle of minimal tripotents in* E*. Then the restriction of* A *to the JBW*<sup>∗</sup> *-subtriple* E<sup>0</sup> *of* E *generated by the elements in the quadrangle is a positive scalar multiple of a* (*complex*) *linear or a conjugate-linear isometry.*

*Proof.* Observe first that since  $e_1 \top e_2 \top e_3 \top e_4$ , Proposition [5.8](#page-22-0)(b) implies that

$$
\boldsymbol{\gamma}(e_1, A) = \boldsymbol{\gamma}(e_2, A) = \boldsymbol{\gamma}(e_3, A) = \boldsymbol{\gamma}(e_4, A).
$$

Observe that  $w = \frac{e_1 + e_2 + e_3 + e_4}{2}$  is a minimal tripotent in E (see Lemma [4.3\)](#page-10-0), and thus by Proposition  $5.8(\tilde{a})$  $5.8(\tilde{a})$  one of the next cases holds:

*Case* (I)  $A(\lambda w) = \lambda A(w)$  for all  $\lambda \in \mathbb{C}$ . The identity

$$
\lambda \frac{A(e_1) + A(e_2) + A(e_3) + A(e_4)}{2} = \lambda A(w)
$$
  
=  $A(\lambda w) = \frac{A(\lambda e_1) + A(\lambda e_2) + A(\lambda e_3) + A(\lambda e_4)}{2}$ 

,

combined with the fact that  $r(A(e_1))\top r(A(e_2))\top r(A(e_3))\top r(A(e_1)), r(A(e_1))$  ⊥  $r(A(e_3))$ , and  $r(A(e_2)) \perp r(A(e_4))$  proves that  $A(\lambda e_i) = \lambda A(e_i)$  for all  $j \in$  ${1, 2, 3, 4}, \lambda \in \mathbb{C}$ . Therefore

$$
A(\lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3 + \lambda_4 e_4) = \lambda_1 A(e_1) + \lambda_2 A(e_2) + \lambda_3 A(e_3) + \lambda_4 A(e_4) \quad (\lambda_j \in \mathbb{C}),
$$
  
assume that the restriction of A to the linear span of  $e_1$ ,  $e_2$ ,  $e_3$ ,  $e_4$ , is (complex) linear

assures that the restriction of A to the linear span of  $e_1, e_2, e_3, e_4$  is (complex) linear, while,

$$
\lambda_1 A(e_1) + \lambda_2 A(e_2) + \lambda_3 A(e_3) + \lambda_4 A(e_4)
$$
  
=  $\gamma(e_1, A) (\lambda_1 r(A(e_1)) + \lambda_2 r(A(e_2)) + \lambda_3 r(A(e_3)) + \lambda_4 r(A(e_4))) (\lambda_j \in \mathbb{C}),$ 

implies that the restriction of A to  $E_0 = \text{span}\{e_1, e_2, e_3, e_4\}$  is a positive scalar multiple of an isometry. It can be easily checked that the minimal tripotents  $r(A(e_1)), r(A(e_2)), r(A(e_3)), r(A(e_4))$  form a quadrangle in  $C_{\sigma(k)} \subseteq F$ , and  $A(E_0)$  $=$  span $\{r(A(e_1)), r(A(e_2)), r(A(e_3)), r(A(e_4))\}.$ 

*Case* (II)  $A(\lambda w) = \overline{\lambda}A(w)$  for all  $\lambda \in \mathbb{C}$ . Similar arguments to those given above prove that the restriction of A to  $E_0 = \text{span}\{e_1, e_2, e_3, e_4\}$  is a positive scalar multiple of a conjugate-linear isometry, and  $A|_{E_0}: E_0 \to A(E_0)$  is a positive multiple of a conjugate-linear triple isomorphism.  $\Box$ 

### 5.1. Connections with preservers of triple transition pseudo-probabilities.

As we have seen in Proposition [5.8](#page-22-0) Under the hypotheses of Proposition [5.6,](#page-19-0) given two atomic JBW<sup>\*</sup>-triples  $E$  and  $F$ , such that  $E$  contains no one-dimensional Cartan factor summands, and a surjective additive mapping  $A: E \to F$  preserving truncations in both directions, we can associate A with a bijection  $A_r : \mathcal{U}_{min}(E) \rightarrow$  $\mathcal{U}_{min}(F)$  preserving collinearity and orthogonality in both directions which is given by  $A_r(e) := r(A(e))$ . However, we lack of a result saying that A maps tripotents to positive scalar multiples of tripotents. So, we must restrict ourself to the sets of minimal tripotents. The remaining part of our arguments will rely on the connections between surjective additive mappings preserving truncations and preservers of triple transition pseudo-probabilities. Let  $e$  and  $v$  be two minimal tripotents in a JBW<sup>∗</sup> -triple E. We recall that the *triple transition pseudo-probability* (of transitioning) from  $e$  to  $v$  is the complex number given by

$$
TTP(e, v) = \varphi_v(e),\tag{6}
$$

where  $\varphi_v$  is the unique pure atom in  $E_*$  supported at v (cf. [\[43,](#page-31-5) [44\]](#page-31-2)). The triple transition pseudo-probability between minimal tripotents is a symmetric mapping, that is,  $TTP(e, v) = TTP(v, e)$ , for all  $e, v \in \mathcal{U}_{min}(E)$ . In the case that e and v are minimal projections in  $B(H)$ , the triple transition pseudo-probability from e to v is precisely the usual transition probability from  $e$  to  $v$  in the celebrated Wigner's theorem. It is known that every (linear) triple isomorphism  $T$  between JBW<sup>\*</sup>-triples is automatically weak<sup>\*</sup>-continuous (cf.  $[29,$  Corollary 3.22]), and hence T preserves triple transition pseudo-probabilities between minimal tripotents. Reciprocally, the main result in [\[44\]](#page-31-2) shows that every bijective mapping preserving triple transition pseudo-probabilities between the sets of minimal tripotents of two atomic JBW<sup>∗</sup> triples is precisely the restriction of a (complex-)linear triple isomorphism between the corresponding JBW<sup>∗</sup> -triples.

<span id="page-26-1"></span>**Remark 5.12.** Let  $S : E \to F$  be a conjugate-linear triple isomorphism between two JBW<sup>\*</sup>-triples. It is also known that in this case S must be also weak<sup>\*</sup>continuous (see, for example, [\[39,](#page-31-6) Proposition 2.3]). Let us take two minimal tripotents  $e, v \in E$  and the corresponding pure atoms  $\varphi_e, \varphi_v$  supported at e and v, respectively. Clearly  $S(e)$  and  $S(v)$  are minimal tripotents in F whose supporting functional are denoted by  $\varphi_{S(e)}, \varphi_{S(v)}$ , respectively. By definition we have

$$
\varphi_{S(v)}(S(e))S(v) = P_2(S(v))(S(e)) = \{S(v), \{S(v), S(e), S(v)\}, S(v)\}
$$
  
=  $\{S(v), S(\{v, e, v\}), S(v)\} = S(P_2(v)(e)) = S(\varphi_v(e)v)$   
=  $\overline{\varphi_v}(e)S(v)$ ,

which assures that

$$
TTP(S(e), S(v)) = \overline{TTP(e, v)} = TTP(v, e).
$$

<span id="page-26-2"></span>**Remark 5.13.** Let  $F$  be a JBW<sup>\*</sup>-subtriple of a JBW<sup>\*</sup>-triple  $E$ . Let us consider two tripotents  $e, v \in F$  which are minimal tripotents in E. Then the triple transition pseudo-probability from e to v does not change when computed in  $E$  or in  $F$ . Namely, for each minimal tripotent  $e \in E$  belonging also to F we have  $E_2(e)$  =  $\mathbb{C}e = F_2(e)$ . If  $\phi$  is any functional in  $E_*$  such that  $\|\phi\| = 1 = \phi(e) = 1$ , we have  $\phi = \phi P_2(e) = \phi|_{E_2(e)} P_2(e)$  (cf. [\[25,](#page-30-12) Proposition 1]). So, if  $\varphi_v^E \in \partial_e(\mathcal{B}_{E_*})$ and  $\varphi_v^F \in \partial_e(\mathcal{B}_{F_*})$  denote the unique pure atoms of E and F supported at v, respectively, it follows that  $\varphi_v^E|_F = \varphi_v^F$  –in other words, there is a unique normpreserving extension of  $\varphi_v^F$  to an element in  $E_*$ .

<span id="page-26-0"></span>We are now in a position to culminate our technical arguments.

Proposition 5.14. *Under the hypotheses of Proposition [5.6,](#page-19-0) the following statements hold:*

- (a) For each  $k \in \Gamma_1$  there exists a (unique) positive constant  $\gamma_k$  such that  $\gamma_k =$  $\gamma(w, A)$  *for all minimal tripotent*  $w \in C_k$ *, that is,*  $\gamma(w, A)$  *does not change* when w runs in the set,  $\mathcal{U}_{min}(C_k)$ , of all minimal tripotents of each Cartan *factor*  $C_k$ .
- (b) For each  $k \in \Gamma_1$  one of the next statements holds:
	- (b.1)  $A(\lambda e) = \gamma_k \lambda r(A(e)) = \lambda A(e)$ , *for all*  $e \in \mathcal{U}_{min}(C_k)$  *and*  $\lambda \in \mathbb{C}$ *.*
	- (b.2)  $A(\lambda e) = \gamma_k \overline{\lambda} r(A(e)) = \overline{\lambda} A(e)$ , *for all*  $e \in \mathcal{U}_{min}(C_k)$  *and*  $\lambda \in \mathbb{C}$ *.*
- (c) For each  $k \in \Gamma_1$  one of the next statements holds:
	- (c.1) *If*  $A(\lambda w) = \lambda A(w)$ , *for all*  $w \in \mathcal{U}_{min}(C_k)$  *and*  $\lambda \in \mathbb{C}$ *, the mapping*  $A_r$  *preserves triple transition pseudo-probabilities between elements in*  $\mathcal{U}_{min}(C_k)$ *, that is,*  $TTP(A_r(e), A_r(v)) = TTP(e, v)$ *, for all*  $e, v \in \mathcal{U}_{min}(C_k)$ *.*
	- (c.2) If  $A(\lambda w) = \lambda A(w)$ , for all  $w \in \mathcal{U}_{min}(C_k)$  and  $\lambda \in \mathbb{C}$ , the mapping  $A_r$  re*verses triple transition pseudo-probabilities between elements in*  $\mathcal{U}_{min}(C_k)$ ,

that is, 
$$
TTP(A_r(e), A_r(v)) = \overline{TTP(e, v)} = TTP(v, e)
$$
, for all  $e, v \in U_{min}(C_k)$ .

(d) The families  $(\gamma_k)_{k \in \Gamma_1}$  and  $(\gamma_k^{-1})_{k \in \Gamma_1}$  are bounded.

*Proof.* Pick an arbitrary  $k \in \Gamma_1$  and two minimal tripotents  $e, v \in C_k$ . We shall distinguish three main cases.

*Case* (1):  $C_k$  has rank-one. By Proposition [5.6](#page-19-0)(b), the restricted mapping  $A|_{C_k} : C_k \to \tilde{C}_{\sigma(k)}$  is a surjective mapping preserving truncations in both directions between two rank-one Cartan factors with  $\dim(C_k) \geq 2$ . Proposition [3.3](#page-8-1) proves that  $A|_{C_k}$  is a positive scalar multiple of a (complex) linear or a conjugatelinear isometry, and hence the desired conclusion in  $(a)$  and  $(b)$  are trivially true.

*Case* (2):  $C_k$  has rank  $\geq 2$  and there exists a quadrangle of minimal tripotents of the form  $(e, e_2, e_3, e_4)$  in  $C_k$  such that v is a linear combination of the elements in the quadrangle. We deduce from Proposition [5.11](#page-24-0) that the restriction of A to the JBW<sup>\*</sup>-subtriple  $E_0 \subseteq C_k$  spanned by the set  $\{e, e_2, e_3, e_4\}$  is a positive scalar multiple of a(complex) linear or a conjugate-linear isometry. Consequently, the conclusions in  $(a)$  and  $(b)$  trivially hold for e and v, and the arbitrariness of thee two elements concludes the proof.

*Case* (3):  $C_k$  has rank  $\geq 2$ , but v cannot be written as a linear combination of the elements in a quadrangle of minimal tripotents of the form  $(e, e_2, e_3, e_4)$  in  $C_k$ . Lemma [5.10](#page-24-1) implies that  $C_k$  is a type 3 Cartan factor and there exists a trangle of the form  $(e, u, \tilde{e})$  with  $\tilde{e}$  minimal and u having rank-2 such that v is a linear combination of  $e, u$ , and  $\tilde{e}$ , and the JBW<sup>\*</sup>-subtriple of C generated by this trangle coincides with  $C_2(e+\tilde{e}) = C_2(u)$  and is isometrically isomorphic to  $S_2$ . Corollary [5.5](#page-19-1) shows that  $A(C_2(e+\tilde{e})) = F_2(r(A(e+\tilde{e})))$ . We also know that  $r(A(e+\tilde{e}))$  is a rank-2 tripotent in  $F$  (cf. Lemma [4.1](#page-9-1) and Proposition [4.7](#page-14-1) combined with the additivity of A), and  $F_2(r(A(e + \tilde{e})))$  is an atomic JBW<sup>\*</sup>-triple (cf. Proposition [5.1\)](#page-16-0). We can now conclude from Proposition [5.9](#page-22-1) the  $F_2(r(A(e + \tilde{e}))) = S_2$  and the restriction of A to  $E_0 = C_2(e + \tilde{e})$  is a positive scalar multiple of a surjective (complex) linear or a conjugate-linear isometry. As before, this shows that  $(a)$  and  $(b)$  hold for e and  $\upsilon$ .

(c) We have shown in the proof of (a) and (b) that for each  $k \in \Gamma_1$ , and any two minimal tripotents  $e, v \in C_k$ , we can find a JBW<sup>\*</sup>-subtriple  $E_0 \subseteq C_k$  containing e and v such that  $\gamma_k^{-1}A|_{E_0}: E_0 \to A(E_0)$  is a linear or a conjugate-linear isometry, equivalently, a linear or a conjugate-linear triple isomorphism between JBW<sup>\*</sup>-triples (see  $[35,$  Proposition 5.5]). It is also clear from  $(a)$  and Lemma [4.1](#page-9-1) that  $A_r|_{\mathcal{U}_{min}(E_0)} = \gamma_k^{-1} A|_{\mathcal{U}_{min}(E_0)}$ . It follows that  $TTP(A_r(e), A_r(v)) = TTP(e, v)$ if  $A(\lambda w) = \gamma_k \lambda r(A(w)) = \lambda A(w)$ , for all  $w \in \mathcal{U}_{min}(C_k)$  and all  $\lambda \in \mathbb{C}$ , and  $TTP(A_r(e), A_r(v)) = \overline{TTP(e, v)} = TTP(v, e)$  if  $A(\lambda w) = \gamma_k \overline{\lambda}r(A(w)) = \overline{\lambda}A(w)$ , for all  $w \in \mathcal{U}_{min}(C_k)$  and  $\lambda \in \mathbb{C}$  (cf. Remark [5.12,](#page-26-1) the preceding comments to this result, and Remark [5.13\)](#page-26-2).

(d) For each  $k \in \Gamma_1$  pick a minimal tripotent  $e_k \in C_k$ . The element  $e = (e_k)_{k \in \Gamma_1}$ is a tripotent in  $E$  satisfying

$$
\gamma_{k_0} = ||A(e_{k_0})|| = ||A(e_{k_0}) + A(e - e_{k_0})|| = ||A(e)||.
$$

If in this argument we replace A with  $A^{-1}$  we obtain the boundedness of the family  $(\gamma_k^{-1})_{k \in \Gamma_1}$ .

<span id="page-28-0"></span>We can now state our main result on surjective additive mappings between atomic JBW<sup>∗</sup> -triples preserving truncations in both directions.

**Theorem 5.15.** Let E and F be atomic JBW<sup>\*</sup>-triples, that is,  $E = \bigoplus_{k=1}^{\ell_{\infty}}$  $k \in \Gamma_1$ C<sup>k</sup> *and*  $\ell_{\infty}$ 

$$
F = \bigoplus_{j \in \Gamma_2} \widetilde{C}_j, \text{ where } \{C_k\}_{k \in \Gamma_1} \text{ and } \{\widetilde{C}_j\}_{j \in \Gamma_2} \text{ are two families of Cartan factors.}
$$

*Suppose, additionally, that* E *contains no one-dimensional Cartan factors as direct summands* (*i.e.,*  $dim(C_k) \geq 2$  *for all* k). Let  $A : E \rightarrow F$  be a surjective additive *mapping. Then the following statements are equivalent:*

- (a) A *preserves truncations in both directions.*
- (b) *There exists a bijection*  $\sigma : \Gamma_1 \to \Gamma_2$ , a bounded family  $(\gamma_k)_{k \in \Gamma_1} \subseteq \mathbb{R}^+$ , and *a family*  $(\Phi_k)_{k \in \Gamma_1}$ , *where each*  $\Phi_k$  *is a (complex) linear or a conjugate-linear* (*isometric*) *triple isomorphism from*  $C_k$  *onto*  $C_{\sigma(k)}$  *satisfying*  $\inf_k \{ \gamma_k \} > 0$ , *and*

$$
A(x) = \left(\gamma_k \Phi_k \left(\pi_k(x)\right)\right)_{k \in \Gamma_1}, \text{ for all } x \in E,
$$

*where*  $\pi_k$  *denotes the canonical projection of* E *onto*  $C_k$ .

*Moreover, if* A *preserves truncations in both directions and there exists an element*  $x_0$  *in* E satisfying  $||A\pi_k(x_0)|| = ||\pi_k(x_0)|| \neq 0$ , for all  $k \in \Gamma_1$ , then the mapping A *is a real-linear* (*isometric*) *triple isomorphism.*

*Proof.* We shall only prove that  $(a) \Rightarrow (b)$ . Let  $\sigma : \Gamma_1 \rightarrow \Gamma_2$  denote the bijec-tion given by Proposition [5.6](#page-19-0)(b), and let  $(\gamma_k)_{k \in \Gamma_1}$  stand for the bounded family of positive numbers whose existence is assured by Proposition [5.14.](#page-26-0)

Fix now  $k \in \Gamma_1$ . By Proposition [5.6](#page-19-0) and Lemma [4.1,](#page-9-1) the restricted mapping  $A|_{C_k}: C_k \to C_{\sigma(k)}$  is a bijection preserving truncations in both directions. Proposition [5.14](#page-26-0) implies that one (and just one) of the next statements holds:

- (1)  $A(\lambda e) = \gamma_k \lambda r(A(e)) = \lambda A(e) = \gamma_k \lambda A_r(e)$ , for all  $e \in \mathcal{U}_{min}(C_k)$  and  $\lambda \in \mathbb{C}$ , and  $TTP(A_r(e), A_r(v)) = TTP(e, v)$ , for all  $e, v \in \mathcal{U}_{min}(C_k)$ ..
- (2)  $A(\lambda e) = \gamma_k \lambda r(A(e)) = \lambda A(e) = \gamma_k \lambda A_r(e)$ , for all  $e \in \mathcal{U}_{min}(C_k)$  and  $\lambda \in \mathbb{C}$ , and  $TTP(A_r(e), A_r(v)) = TTP(e, v) = TTP(v, e)$ , for all  $e, v \in \mathcal{U}_{min}(C_k)$ .

In case (1), the mapping  $A_r^{-1}: \mathcal{U}_{min}(\tilde{C}_{\sigma(k)}) \to \mathcal{U}_{min}(C_k)$  is a bijection preserving triple transition pseudo-probabilities (see also Proposition  $5.8(c)$  $5.8(c)$ ). Corollary 3.3 in [\[44\]](#page-31-2) assures the existence of a linear (isometric) triple isomorphism  $\Psi_k : C_{\sigma(k)} \to C_k$ whose restriction to  $\mathcal{U}_{min}(\tilde{C}_{\sigma(k)})$  is  $A_r^{-1}$ . We also know from Proposition [5.14](#page-26-0) and the assumptions in this case (1), that  $A|_{\mathcal{U}_{min}(C_k)} = \gamma_k A_r$ , and the mapping  $T_k = \gamma_k^{-1} \Psi_k A|_{C_k} : C_k \to C_k$  is an additive bijection preserving truncations in both directions, which also satisfies  $T_k(\lambda e) = \lambda e$  for all  $e \in \mathcal{U}_{min}(C_k)$  and  $\lambda \in \mathbb{C}$ . The identity principle in Proposition [5.2](#page-17-1) now implies that  $T_k$  is the identity mapping on  $C_k$ , and thus  $A|_{C_k} = \gamma_k \Psi_k^{-1}$  is a positive scalar multiple of an isometric linear triple isomorphism  $\Phi_k = \Psi_k^{-1}$  from  $C_k$  onto  $\widetilde{C}_{\sigma(k)}$ .

Suppose now that  $A|_{C_k}: C_k \to C_{\sigma(k)}$  satisfies the properties in case (2). It is known that we can always find a conjugation (i.e., a conjugate-linear triple automorphism of period-2)  $\Psi_k$  :  $C_{\sigma(k)} \to C_{\sigma(k)}$  (cf. [\[36,](#page-30-19) Theorem 4.1] and [\[38\]](#page-30-26) for additional details in the case of exceptional Cartan factors). The mapping

 $\Psi_k A|_{C_k}: C_k \to C_{\sigma(k)}$  satisfies the properties in case (1), and hence, by the previous conclusion  $\Psi_k A|_{C_k} = \alpha_k \widetilde{\Phi}_k$ , where  $\alpha_k \in \mathbb{R}^+$  and  $\widetilde{\Phi}_k : C_k \to C_{\sigma(k)}$  is a linear and isometric triple isomorphism. Necessarily  $\alpha_k = \gamma_k$ . Finally,  $A|_{C_k} = \gamma_k \Psi_k \widetilde{\Phi}_k = \gamma_k \Phi_k$ , where  $\Phi_k = \Psi_k \Phi_k : C_k \to C_{\sigma(k)}$  is a conjugate-linear triple isomorphism.

The final statement is clear since the existence of such element  $x_0 \in E$  implies that  $\gamma_k = 1$  for all k.

Our last corollary, which is interesting by itself, is a straightforward consequence of our main theorem.

**Corollary 5.16.** Let C and  $\tilde{C}$  be Cartan factors with  $dim(C) > 2$ . Then every *surjective additive mapping*  $A: C \to \tilde{C}$  *preserving truncations in both directions is a positive scalar multiple of a linear or a conjugate-linear (isometric) triple isomorphism. Moreover, if we also assume that*  $||A(x_0)|| = ||x_0||$  *for some non-zero*  $x_0$  *in* C*, the mapping* A *is a linear or a conjugate-linear* (*isometric*) *triple isomorphism.*

Acknowledgements L. Li supported by National Natural Science Foundation of China (Grant No. 12171251). A.M. Peralta supported by grant PID2021- 122126NB-C31 funded by MICIU/AEI/10.13039/501100011033 and by ERDF/EU, by Junta de Andalucía grant FQM375, IMAG–María de Maeztu grant CEX2020- 001105-M/AEI/10.13039/501100011033 and (MOST) Ministry of Science and Technology of China grant G2023125007L.

Part of this work was completed during a visit of A.M. Peralta to Nankai University and the Chern Institute of Mathematics, which he thanks for the hospitality.

Statements and Declarations. All authors declare that they have no conflicts of interest to disclose.

Data availability. There is no data associate for this submission.

#### **REFERENCES**

- [1] A. Blanco, A. Turnšek, On maps that preserve orthogonality in normed spaces, Proc. Roy. Soc. Edinburgh Sect. A 136 (2006), 709–716.
- <span id="page-29-1"></span>[2] T. Barton, R. M. Timoney, Weak∗-continuity of Jordan triple products and its applications, Math. Scand., 59(2) (1986), 177–191.
- <span id="page-29-3"></span>[3] M. Battaglia, Order theoretic type decomposition of JBW∗-triples, Q. J. Math., Oxf. II. Ser., 42(166) (1991), 129–147.
- <span id="page-29-6"></span>[4] J. Becerra Guerrero, G. López Pérez, A. M. Peralta, A. Rodríguez-Palacios, Relatively weakly open sets in closed balls of Banach spaces, and real JB∗-triples of finite rank, Math. Ann. 330 (2004), no. 1, 45–58.
- <span id="page-29-0"></span>[5] F. Botelho, L. Molnár, G. Nagy, Linear bijections on von Neumann factors commuting with λ-Aluthge transform, Bull. Lond. Math. Soc. 48 (2016), No. 1, 74–84.
- <span id="page-29-5"></span>[6] L.J. Bunce, Ch.-H. Chu, Compact operations, multipliers and Radon-Nikodym property in JB∗-triples, Pacific J. Math. 153 (1992), 249–265.
- <span id="page-29-4"></span>[7] L.J. Bunce, F.J. Fernández-Polo, J.M. Moreno, A.M. Peralta, A Saitô–Tomita–Lusin theorem for JB<sup>\*</sup>-triples and applications, *Quart. J. Math. Oxford Ser.* 57 (2006), 37–48.
- <span id="page-29-2"></span>[8] M. Burgos, F.J. Fernández-Polo, J.J. Garcés, J.M. Martínez, A.M. Peralta, Orthogonality preservers in C<sup>∗</sup>-algebras, JB<sup>∗</sup>-algebras and JB<sup>∗</sup>-triples. J. Math. Anal. Appl., 348(1) (2008), 220–233.
- <span id="page-29-7"></span>[9] D. Cabezas, M. Martín, A.M. Peralta, The Daugavet equation for polynomials on C<sup>∗</sup>-algebras and JB<sup>\*</sup>-triples, Adv. Math. 439 (2024), Article ID 109479.

- <span id="page-30-10"></span>[10] M. Cabrera García, A. Rodríguez Palacios, Non-associative normed algebras. Volume 1. The Vidav-Palmer and Gelfand-Naimark theorems, volume 154 of Encycl. Math. Appl.. Cambridge: Cambridge University Press, 2014.
- <span id="page-30-3"></span><span id="page-30-2"></span>[11] F. Chabbabi, Product commuting maps with the λ-Aluthge transform, J. Math. Anal. Appl.,  $449(1)$  (2017), 589-600.
- [12] F. Chabbabi, M. Mbekhta, Jordan product maps commuting with the λ-Aluthge transform, J. Math. Anal. Appl., 450(1) (2017), 293–313.
- [13] J. Chmieliński, Linear mappings approximately preserving orthogonality, J. Math. Anal. Appl. 304 (2005), No. 1, 158–169.
- <span id="page-30-24"></span>[14] C.-H. Chu, B. Iochum, On the Radon-Nikodým property in Jordan triples, Proc. Am. Math. Soc. 99 (1987), 462-464.
- <span id="page-30-20"></span>[15] T. Dang, Y. Friedman, Classification of JBW∗-triple factors and applications, Math. Scand., 61 (1987), (2), 292–330.
- <span id="page-30-8"></span>[16] S. Dineen, The second dual of a JB<sup>∗</sup> triple system. In Complex analysis, functional analysis and approximation theory (Campinas, 1984), volume 125 of North-Holland Math. Stud., pages 67–69. North-Holland, Amsterdam, 1986.
- <span id="page-30-9"></span>[17] C.M. Edwards, G.T. Rüttimann, On the facial structure of the unit balls in a JBW∗-triple and its predual, J. London Math. Soc. (2), 38(2) (1988), 317–332.
- <span id="page-30-4"></span>[18] A.B.A. Essaleh, A.M. Peralta, Preservers of λ-Aluthge transforms, Linear Algebra Appl., 554 (2018), 86–119.
- <span id="page-30-17"></span>[19] F.J. Fernández-Polo, A.M. Peralta, Compact tripotents and the Stone–Weierstrass Theorem for  $C^*$ -algebras and JB<sup>\*</sup>-triples, *J. Operator Theor.* **58**(1) (2007), 157–173.
- <span id="page-30-18"></span>[20] F.J. Fernández-Polo, A.M. Peralta, Non-commutative generalisations of Urysohn's lemma and hereditary inner ideals, Journal of Functional Analysis 259 (2010), 343–358
- <span id="page-30-21"></span>[21] F.J. Fernández-Polo, A.M. Peralta, Low rank compact operators and Tingley's problem, Adv. Math., 338 (2018), 1–40.
- [22] Y. Friedman, A.M. Peralta, Representation of symmetry transformations on the sets of tripotents of spin and Cartan factors, Anal. Math. Phys.,  $12(1)$  (2022), Id/No 37.
- <span id="page-30-15"></span>[23] Y. Friedman, B. Russo, Contractive projections on  $C_0(K)$ , Trans. Am. Math. Soc., 273 (1982), 57-73.
- <span id="page-30-16"></span>[24] Y. Friedman, B. Russo, Function representation of commutative operator triple systems, J. Lond. Math. Soc., II. Ser., 27 (1983), 513-524.
- <span id="page-30-12"></span>[25] Y. Friedman, B. Russo, Structure of the predual of a JBW<sup>\*</sup>-triple, *J. Reine Angew. Math.*, 356 (1985), 67–89.
- <span id="page-30-13"></span>[26] Y. Friedman, B. Russo, The Gelfand-Naimark theorem for JB<sup>\*</sup>-triples, Duke Math. J., 53 (1986), 139–148.
- <span id="page-30-1"></span>[27] J.J. Garcés, L. Li, A.M. Peralta, S. Su, Maps preserving the truncation of triple products on Cartan factors, Results Math. 79 (2024), 287. https://doi.org/10.1007/s00025-024-02305-1
- <span id="page-30-5"></span>[28] J. Hamhalter, O.F.K. Kalenda, A.M. Peralta, Determinants in Jordan matrix algebras, Linear Multilinear Algebra, 71 (6) (2023), 961–1002.
- <span id="page-30-11"></span>[29] G. Horn, Characterization of the predual and ideal structure of a JBW∗-triple, Math. Scand., 61(1) (1987), 117–133.
- <span id="page-30-23"></span><span id="page-30-0"></span>[30] G. Horn, Classification of JBW<sup>∗</sup>-triples of type I, Math. Z. 196 (1987), 271–291.
- [31] X. Jia, W. Shi, G. Ji, Maps preserving the truncation of products of operators, Ann. Funct. Anal.,  $13(3)$  (2022), Id/No 40.
- <span id="page-30-22"></span>[32] O.F.K. Kalenda, A.M. Peralta, Extension of isometries from the unit sphere of a rank-2 Cartan factor, Anal. Math. Phys., 11 (2021) (1), Id/No 15.
- <span id="page-30-14"></span>[33] W. Kaup, Algebraic characterization of symmetric complex Banach manifolds, Math. Ann., 228 (1977), 39–64.
- <span id="page-30-7"></span>[34] W. Kaup, A Riemann mapping theorem for bounded symmetric domains, Math Z., 183 (1983), 503–530.
- <span id="page-30-25"></span>[35] W. Kaup, Üher die Klassifikation der symmetrlschen Hermiteschen Mannigfaltigkeiten unendlicher Dimension I, Math. Ann. 257 (1981), 463–486.
- <span id="page-30-19"></span><span id="page-30-6"></span>[36] W. Kaup, On real Cartan factors, *Manuscr. Math.*, **92**(2) (1997), 191–222.
- [37] L. Li, S. Liu, A.M. Peralta, Additive orthogonality preserving mappings between complex inner product spaces, preprint 2024. arXiv: 2410.08101
- <span id="page-30-26"></span>[38] O. Loos, *Bounded symmetric domains and Jordan pairs*, Mathematical Lectures. Irvine: University of California at Irvine 1977.
- [39] J. Martínez, A.M. Peralta, Separate weak∗-continuity of the triple product in dual real JB∗ triples, Math. Z. 234 (2000), No. 4, 635–646.
- <span id="page-31-0"></span>[40] Y. Mao and G. Ji, Truncations of operators in  $\mathcal{B}(H)$  and their preservers, Adv. Operator Theory, 9 (2024), Id/No 32.
- <span id="page-31-4"></span>[41] E. Neher, Jordan triple systems by the grid approach. Lecture Notes in Mathematics, 1280. Springer-Verlag, Berlin, 1987.
- <span id="page-31-3"></span>[42] M. Omladič, P. Šemrl, Additive mappings preserving operators of rank one, Linear Algebra Appl., 182 (1993), 239–256.
- <span id="page-31-5"></span>[43] A.M. Peralta, Maps preserving triple transition pseudo-probabilities, RIMS Kôkyûroku Bessatsu, B93 (2023), 1–28,.
- <span id="page-31-2"></span>[44] A.M. Peralta, Preservers of triple transition pseudo-probabilities in connection with orthogonality preservers and surjective isometries, Result. Math., 78(2) (2023), Id/No 51.
- <span id="page-31-1"></span>[45] J. Yao and G. Ji, Additive maps preserving the truncation of operators, J. Math. Res. Appl., 42(1) (2022), 89–94.
- (L. Li) School of Mathematical Sciences and LPMC, Nankai University, 300071 Tianjin, China.

Email address: leilee@nankai.edu.cn

(S. Liu) School of Mathematical Sciences and LPMC, Nankai University, 300071 Tianjin, China.

Email address: 760659676@qq.com

(A.M. Peralta) Instituto de Matemáticas de la Universidad de Granada (IMAG), Departamento de Análisis Matemático, Facultad de Ciencias, Universidad de Granada, 18071 Granada, Spain.

Email address: aperalta@ugr.es

<span id="page-31-6"></span>