AN INVERSE THEOREM FOR GENERALIZED ARITHMETIC PROGRESSION WITH MILD MULTIPLICATIVE PROPERTY

ERNIE CROOT AND JUNZHE MAO

ABSTRACT. In this paper we prove a structural theorem for generalized arithmetic progressions in \mathbb{F}_p which contain a large product set of two other progressions.

1. INTRODUCTION

Let p be a prime number, \mathbb{F}_p be the finite field with p elements. Recall that a generalized arithmetic progression (GAP) of rank d in \mathbb{F}_p is a set of the form

$$B = x_0 + \{\sum_{i=1}^d a_i x_i : a_i \in [0, N_i - 1]\}$$

for some elements $x_0, x_1, \ldots, x_d \in \mathbb{F}_p$ and positive integers $2 \leq N_1, \ldots, N_d \leq p$. We say *B* is proper if $|B| = N_1 N_2 \ldots N_d$. GAPs play an important role in additive combinatorics, as they exhibit an almost periodic behavior under addition. Another example of additively structured sets is Bohr sets. Given $\Gamma \subset \mathbb{F}_p$ and $\epsilon > 0$, the Bohr set $B(\Gamma, \epsilon)$ is defined as

$$B(\Gamma, \epsilon) = \left\{ x \in \mathbb{F}_p : \left\| \frac{xr}{p} \right\| < \epsilon, \forall r \in \Gamma \right\}.$$

Here $\|\cdot\|$ denotes the distance to the nearest integer. One can think of Bohr sets as approximate subspaces or (additive) subgroups in \mathbb{F}_p . It turns out that GAPs and Bohr sets are closely related. Using the geometry of numbers, one can show that any Bohr set $B(\Gamma, \epsilon)$ must contain a large symmetric proper GAP with bounded rank. This fact has been used to prove a foundational result in additive combinatorics, namely Freiman's theorem: given a finite set $A \subset \mathbb{F}_p$ and some C > 0, if $|A + A| \leq C|A|$, where

$$A + A = \{a_1 + a_2 : a_1, a_2 \in A\},\$$

then there exists a GAP B such that $A \subset B$ and $|B| \ll_C |A|$. Here we use the Vinogradov notation \ll_C meaning that there is some constant K > 0 depending on C so that $|B| \leq K|A|$. Freiman's theorem tells us that GAPs are the "standard model" for sets with rich additive structures. For more properties and applications of Bohr sets and GAPs, we refer the reader to [9].

One of the reasons people are concerned about these additively structured sets is the sum-product phenomenon, which basically says no set can have rich additive structure and multiplicative structure simultaneously. For example, in [3] Erdös and Szemerédi conjectured that if a set $A \subset \mathbb{Z}$ satisfies

$$|A + A| \le K|A$$

for some constant K > 0, then for any $\epsilon > 0$, $|AA| \gg |A|^{2-\epsilon}$, where

$$AA = \{a_1a_2 : a_1, a_2 \in A\}$$

This has been confirmed for $A \subset \mathbb{R}$ by a result of Elekes and Ruzsa [2], and recently verified for thin sets $A \subset \mathbb{F}_p$ by a result of Kerr, Mello and Shparlinski [6]. The case when $A \subset \mathbb{F}_p$ and $|A| \sim p^{1/2}$ remains open. We refer the reader to [7, 8] for some relevant results. To prove such an estimate, a

common method is to bound the multiplicative energy E(A) of A, defined as the number of solutions to the equation

$$a_1a_2 = a_3a_4, \ a_1, a_2, a_3, a_4 \in A.$$

By Freiman's theorem, this reduces to bounding the multiplicative energy of GAPs in \mathbb{F}_p . Better bounds on the multiplicative energy also have many applications in number theory. For example, Chang [1] used a nontrivial estimate for the multiplicative energy to give a Burgess-type bound for character sums over proper GAPs. Following a similar approach Hanson [4] generalized this to character sums over regular Bohr sets. In fact, if one can prove the optimal bound

$$E(B) \ll |B|^{2+o(1)}$$

for any proper GAP $B \subset \mathbb{F}_p$, then for any nonprincipal Dirichlet character χ of \mathbb{F}_p , Burgess's bound can be fully generalized to

$$\sum_{n \in B} \chi(n) = O(p^{-\delta}|B|)$$

whenever $|B| \ge p^{1/4+\epsilon}$. With current techniques one can prove near-optimal bounds on the multiplicative energy of GAPs with some extra structure. For example, Kerr [5] proved the following result:

Theorem 1 (Kerr). Let $P \subset \mathbb{F}_p$ be a GAP given by

$$P = x_0 + \{\sum_{i=1}^d a_i x_i : a_i \in [1, N]\}$$

and suppose

$$P' = \{\sum_{i=1}^{d} a_i x_i : |a_i| \le N^2\}$$

is proper. Then $E(P) \ll |P|^2 (\log N)^{2d+1}$.

The above discussion suggests that we can only hope for some mild multiplicative structure in a GAP or Bohr set. Let's consider

$$B = \{a + bt : |a|, |b| \le N\},\$$

which is a symmetric GAP. Assume t satisfies a quadratic polynomial equation mod p

$$c_0 + c_1 t + t^2 = 0$$

with $c_i \in \mathbb{Z}$ bounded by some constant C independent of p. Let $A = \{a' + b't : |a'|, |b'| \leq \frac{1}{2C}N^{1/2}\}$, then obviously A satisfies the condition of Kerr's theorem. As a result E(A) is small and A does not possess a rich multiplicative structure. On the other hand, it's easy to check that $AA \subset B$. This set B gives us an example of a GAP with some mild multiplicative properties, i.e. containing a large product set. It raises the following question: if a GAP B contains a product set AA with $|B|^{1/2} \ll |A| \ll |B|^{1/2}$ and A also a GAP, what can we say about B? In this paper we show that under some natural assumptions on A and B, the generators of B must be given by the roots of some low-height, low-degree polynomials.

Notations. For any element $x \in \mathbb{F}_p$, we define its integer height

$$|x| = \min\{|a| : a \in \mathbb{Z}, a \equiv x \pmod{p}\}.$$

For a polynomial $f \in \mathbb{Z}[x]$, let $\overline{f}(x)$ denote the projection of f(x) into $\mathbb{F}_p[x]$, let H(f) denote the height of f, which is the maximum of the absolute value of coefficients of f. We use $M_n(\mathbb{F}_p)$ to denote the ring of $n \times n$ matrices over \mathbb{F}_p .

For any x > 0, let $[x] = \{1, 2, \dots, \lfloor x \rfloor\}$, where $\lfloor x \rfloor$ denote the integer part of x.

Given $x \in \mathbb{F}_p$, $A \subset \mathbb{F}_p$, we write the dilation of A by x as $x \cdot A = \{xa : a \in A\}$.

For any $n \in \mathbb{N}, A \subset \mathbb{F}_p$, we write the iterated sumset as

$$nA = \underbrace{A + A + \dots + A}_{n}$$

2. Main Results

We first consider a simple case, namely a proper GAP B of rank 2 and size at most $o(p^{2/3})$. We show that if B contains the product set of a sub-GAP A of size $|A| \approx |B|^{1/2}$, then the generators of B must be "algebraic" in the sense that they are the roots of some low-height quadratic polynomials.

Theorem 2. Let 0 < c < 1, $N = N(p) \in \mathbb{N}$ with $\lim_{p\to\infty} N(p) = \infty$, $N = o(p^{1/3})$, $B = \{a + bt : |a|, |b| \le N\} \subset \mathbb{F}_p$ be proper and $|B| = o(p^{2/3})$. Suppose $A = \{a' + b't : |a'|, |b'| \le cN^{1/2}\}$ satisfies $AA \subset B$. Then there is a constant C depending only on c so that when p is sufficiently large, there exists some quadratic polynomial $f \in \mathbb{Z}[x] \setminus \{0\}$ with $H(f) \le C$ and $\overline{f}(t) = 0$ in \mathbb{F}_p .

Proof. Since $t^2 \in AA \subset B$, we have

(1)
$$t^2 = a_0 + b_0 t$$

for some $|a_0|, |b_0| \leq N$. We claim that for any integer m, n with $|m|, |n| \leq 5N$, $m - nt \neq 0$ in \mathbb{F}_p . Suppose for contradiction there exists $|m_0|, |n_0| \leq 5N$ so that $m_0 - n_0 t = 0$. Without loss of generality we may assume $gcd(m_0, n_0) = 1$. Then (1) implies

$$(\frac{m_0}{n_0})^2 = a_0 + \frac{b_0 m_0}{n_0}$$

and hence

(2)
$$m_0^2 = a_0 n_0^2 + b_0 m_0 n_0 \text{ in } \mathbb{F}_p.$$

By assumption $|B| = (2N + 1)^2 = o(p^{2/3})$, when p is sufficiently large we get

$$m_0^2 < \frac{p}{2}, \ |an_0^2 + bm_0n_0| < \frac{p}{2}$$

Thus (2) holds not only in \mathbb{F}_p but also in \mathbb{Z} . This implies $n_0 \mid m_0^2$ and hence $|n_0| = 1, m_0 \equiv \pm t \pmod{p}$. From the fact that B is proper we also know $|m_0| \ge |t| \ge N + 1$, therefore

$$m_0^2 \pm b_0 m_0 \ge N + 1 > a_0$$

which is a contradiction to (2). This finishes the proof of the claim.

Now since $\lambda t^2 \in B$ for any $\lambda \in \Pi := \{uv : |u|, |v| \le cN^{1/2}\}$, consider

$$\{(\lambda + \mu)t^2 : \lambda, \mu \in \Pi\}$$

This set will include wt^2 for any integer $|w| < c^2 N/2$. (To see this, take $b = \lfloor cN^{1/2} \rfloor$ and then consider the base-*b* expansion of all positive integers $\leq c^2 N/2$, and note that such integers can be written as $ab + 1 \cdot r$, where $0 \leq a \leq cN^{1/2}$ and $0 \leq r \leq cN^{1/2}$.) Suppose $\max(|a_0|, |b_0|) \geq \frac{8}{c^2}$, let

$$w = \left\lfloor \frac{2N+1}{\max(|a_0|, |b_0|)} \right\rfloor + 1,$$

it is easy to check that $w < c^2 N/2$ when p is large, hence $wt^2 \in 2B$. On the other hand, $2N < \max(|wa_0|, |wb_0|) \leq 3N$. From the claim we know

$$wt^2 = wa_0 + wb_0t \notin 2B_2$$

a contradiction. Hence $\max(|a_0|, |b_0|) \leq \frac{8}{c^2}$ and equation (1) gives the polynomial we want.

The proof strongly depends on the property that enlarging B by some constant will still give us a proper GAP. We do not know if it remains true for $|B| \ge p^{2/3}$. This motivates us to define a special class of GAPs.

Definition 1. A generalized arithmetic progression $B = x_0 + \{\sum_{i=1}^d a_i x_i : a_i \in [0, N_i - 1]\}$ is called κ -isolated if there is no nonzero $(a_1, ..., a_d) \in \mathbb{Z}^d$ with $|a_i| \leq \kappa N_i$, for all i = 1, ..., d, such that $\sum_i a_i x_i = 0$.

Working with isolated GAPs we are able to generalize Theorem 2 to arbitrary rank and product set of two different GAPs. Our main result is the following:

Theorem 3. Let $0 < c, c', \epsilon < 1, d \in \mathbb{N}, N = N(p) \in \mathbb{N}$ with $\lim_{p \to \infty} N(p) = \infty$,

$$B = \{\sum_{i=1}^{d} a_i x_i : |a_i| \le N\} \subset \mathbb{F}_p$$

be 6-isolated with $x_1 = 1$. Suppose there exist proper GAPs

$$A = \{\sum_{i=1}^{d} a_i y_i : |a_i| \le cN^{1-\epsilon}\}, \ A' = \{\sum_{i=1}^{d} a_i y'_i : |a_i| \le c'N^{\epsilon}\}$$

such that $AA' \subset B$. Then there is a constant C = C(c, c', d) so that when p is sufficiently large, for any $i \in [d]$, there exists some polynomial $f_i \in \mathbb{Z}[x] \setminus \{0\}$ of degree at most d with $H(f_i) \leq C$ and $\overline{f_i}(x_i) = 0$ in \mathbb{F}_p .

To prove Theorem 3, we need the following lemma from linear algebra.

Lemma 4. Let $T = (t_{ij}) \in M_d(\mathbb{F}_p)$. Suppose there exists $C \ge 1$ so that $|t_{ij}| \le C$ and T is not invertible. Then there exists a nonzero vector $(a_1, a_2, \ldots, a_d) \in \mathbb{F}_p^d$ with $|a_i| \le d!C^d$ such that

$$(a_1, a_2, \cdots, a_d)T = 0$$

Proof. Let

$$\vec{v_i} = (t_{i1}, t_{i2}, \dots, t_{id}) \in \mathbb{F}_p^d$$

be the *i*th row vector of T. If $\vec{v_i} = 0$ for some i, we can take $a_i = 1$ and $a_j = 0$ for $j \neq i$. Hence we may assume $\vec{v_i} \neq 0$ for all i. Without loss of generality, assume $\{\vec{v_1}, \ldots, \vec{v_k}\}$ forms a maximal linearly independent set, since T is not invertible we must have k < d. Let $T' \in M_k(\mathbb{F}_p)$ be a submatrix of T using the first k rows so that $\operatorname{rank}(T') = k$. Suppose the columns of T' correspond to the $j_1 < j_2 < \cdots < j_k$ th columns of T, let

$$\vec{u}_i = (t_{ij_1}, t_{ij_2}, \dots, t_{ij_k}) \in \mathbb{F}_p^k,$$

We claim that $\vec{u}_{(k+1)} \neq 0$. Indeed, by our choice of $\{v_i\}$, there exists a linear combination

$$\vec{v}_{(k+1)} = \sum_{i=1}^k b_i \vec{v}_i,$$

where the b_i 's are not all zero since $\vec{v}_{k+1} \neq 0$. This implies

$$\vec{u}_{(k+1)} = \sum_{i=1}^k b_i \vec{u}_i$$

and hence $\vec{u}_{k+1} \neq 0$.

Now by adding the (k + 1)th row and another column to T', we obtain a matrix $T'' \in M_{k+1}(\mathbb{F}_p)$ with rank(T'') = k and all the row vectors of T'' are nonzero. It follows that the adjugate matrix $\operatorname{adj}(T'')^1$ of T'' has the (k + 1)th row $(a_1, a_2, \ldots, a_{k+1})$ with $a_{k+1} \neq 0$. Since $\operatorname{det}(T'') = 0$, we get

$$\sum_{i=1}^{k+1} a_i \vec{u}_i = 0.$$

¹Which is the $(k+1) \times (k+1)$ matrix such that $\operatorname{adj}(T'') \cdot T'' = \operatorname{det}(T'')I_{k+1}$, where I_{k+1} is the $(k+1) \times (k+1)$ identity matrix.

From the fact that $\{\vec{v}_1, \vec{v}_2, \ldots, \vec{v}_k\}$ forms a maximal linearly independent set we can deduce

$$\sum_{i=1}^{k+1} a_i \vec{v}_i = 0.$$

Set $a_i = 0$ for all i > k + 1, since each entry of $\operatorname{adj}(T'')$ is the determinant of some submatrix of T'', it follows that $|a_i| \le d!C^d$. This gives us a nonzero vector $(a_1, \ldots, a_d) \in \mathbb{F}_p^d$ so that

$$(a_1, a_2, \cdots, a_d)T = \sum_{i=1}^d a_i \vec{v}_i = 0.$$

Proof of Theorem 3. For any $i, j \in [d]$,

(3)
$$y_i y'_j = \sum_k a^{(k)}_{ij} x_k$$
, where $|a^{(k)}_{i,j}| \le N$

Now since $\lambda y_i y'_j \in B$ for any $\lambda \in \Pi := \{uv : |u| \le cN^{1-\epsilon}, |v| \le c'N^{\epsilon}\}$, consider

$$\{(\lambda+\mu)y_iy_j' : \lambda, \mu \in \Pi\},\$$

As in the proof of Theorem 2, this set will include $wy_iy'_j$ for any |w| < c''N, where 0 < c'' < 1 is some constant depending on c and c'. This implies that

$$wy_iy'_i \in AA' + AA' \subset 2B$$

for any |w| < c'' N. Therefore we have

$$w\sum_{k}a_{ij}^{(k)}x_k = \sum_{k}b_w^{(k)}x_k$$

where $|b_w^{(k)}| \le 2N$. If

$$\max\{|a_{ij}^{(k)}|: k \in [d]\} \ge \frac{3}{c''},$$

then for some |w| < c''N we would have

$$2N < \max\{|wa_{ij}^{(k)}| : k \in [d]\} \le 4N,$$

contradicts with B being 6-isolated. Hence

(4)
$$\max\{|a_{ij}^{(k)}|:k\in[d]\}<\frac{3}{c''}$$

Now fix some $i \in [d]$, and let $\vec{v} = (x_1, x_2, \dots, x_d)^T$, $\vec{u'} = (y'_1, y'_2, \dots, y'_d)^T \in \mathbb{F}_p^d$. Also, let $T \in M_d(\mathbb{F}_p)$ be the matrix whose entry in the *j*th row and *k*th column is

$$T_{j,k} = a_{ij}^{(k)}$$

so that (3) becomes

(5)
$$T\vec{v} = y_i \vec{u'}$$

From (4) it follows that all the entries of the adjugate matrix adj(T) are bounded by

$$C = (d-1)!3^{d-1}/(c'')^{d-1}.$$

Moreover, T must be invertible since otherwise from Lemma 4, there exists a nonzero vector $(a_1, a_2, \ldots, a_d) \in \mathbb{F}_p^d$ with $|a_i|$ bounded by some constant depending on c'' and

$$(a_1, a_2, \ldots, a_d)T = 0.$$

Combining this with (5) gives us

$$y_i \cdot \sum_{j=1}^a a_j y'_j = (a_1, a_2, \dots, a_d) T \vec{v} = 0,$$

which would imply that A' is not proper, a contradiction.

Multiplying (5) by adj(T) on the left on both sides we get

$$\det(T)\vec{v} = y_i \cdot \operatorname{adj}(T)\vec{u'}.$$

Taking linear combinations of coordinates on both sides (using coefficients b_j), we can deduce

$$A_1 := \{\det(T) \sum_j b_j x_j : |b_j| < \frac{c'}{C} N^{\epsilon}\} \subset y_i \cdot A'.$$

Let $\vec{u} = (y_1, y_2, \dots, y_d)^T$. As before, we define a matrix $T' \in M_d(\mathbb{F}_p)$ where the entry in the *j*th row, *k*th column is

$$T'_{j,k} = a^{(k)}_{ji}$$

Again (3) becomes

(6) $T'\vec{v} = \vec{u}y'_i,$

and from (4), all the entries of $\operatorname{adj}(T')$ are bounded by C. Using the same argument that we did earlier for the matrix T, we can deduce that T' is also invertible since A is proper. Multiplying both sides of (6) on the left by $\operatorname{adj}(T')$ we get

$$A_2 := \{\det(T')\sum_j b_j x_j : |b_j| < \frac{c}{C}N^{1-\epsilon}\} \subset A \cdot y'_i.$$

Hence

$$A_1A_2 \subset y_i \cdot (A'A) \cdot y_i' \subset y_i By_i'.$$

Now for any $j, k \in [d]$, we have

$$\det(T)\det(T')x_jx_k = y_i\left(\sum_{l=1}^d s_{jk}^{(l)}x_l\right)y_i'$$

and the set

$$\{\det(T)\det(T')wx_jx_k: |w| < \frac{cc'}{C^2}N\}$$

will be a subset of $y_i(2B)y'_i$ from the same argument as the one at the beginning of our proof. Again from the fact that B is 6-isolated we would have

$$\max\{|s_{ik}^{(l)}| : l \in [d]\} < C'$$

for some C' = C'(c, c', d). Thus there exists $T_j \in M_d(\mathbb{F}_p)$ whose entries are bounded by C' so that

$$\det(T)\det(T')x_j\vec{v} = y_i(T_j\vec{v})y'_i.$$

This equality implies that for any $j \in [d]$, $\det(T)\det(T')x_j/(y_iy'_i)$ is an eigenvalue of T_j , hence is the root of the characteristic polynomial of T_j , which has degree d and height bounded by some constant depending on C'. From Lemma 4 and the fact that B is proper, T_1 must be invertible. Notice that all these eigenvalues share the same eigenvector \vec{v} , thus

$$(\operatorname{adj}(T_1)T_j)\vec{v} = \det(T_1)T_1^{-1}T_j\vec{v} = \det(T_1)\frac{x_j}{x_1}\vec{v} = \det(T_1)x_j\vec{v}.$$

This shows that $\det(T_1)x_j$ is the root of the characteristic polynomial of $\operatorname{adj}(T_1)T_j$, hence x_j is the root of some polynomial of degree at most d with bounded height.

Remark 1. The same proof works for non-symmetric GAP B under slightly stronger conditions. Consider B - B, which is a symmetric GAP, and assume the two symmetric GAPs A - A and A' - A' are proper, then $AA' \subset B$ implies that $(A - A)(A' - A') \subset 2(B - B)$. If B is 24-isolated, then 2(B - B) is 6-isolated and we can repeat the proof for 2(B - B), A - A and A' - A'.

Remark 2. If we remove the condition that $x_1 = 1$, we can only prove that the ratio x_i/x_j of any two generators x_i, x_j satisfies a low-height, low-degree polynomial equation. This is what we expect since the condition $AA' \subset B$ is not invariant under dilation in general.

Since the conclusion of Theorem 3 is about roots of polynomials, which also make sense in the non-commutative ring $M_n(\mathbb{F}_p)$, a natural question is if we consider GAPs in $M_n(\mathbb{F}_p)$, can we get a result similar to Theorem 3? It turns out that under some extra conditions, we can modify the proof of Theorem 3 to get the following:

Corollary 5. Let $0 < c, c', \epsilon < 1, n, d \in \mathbb{N}, N = N(p) \in \mathbb{N}$ with $\lim_{p \to \infty} N(p) = \infty$,

$$B = \{\sum_{i=1}^{d} a_i X_i : |a_i| \le N\} \subset M_n(\mathbb{F}_p)$$

be 6-isolated with $X_1 = I_n$. Suppose there exist proper GAPs

$$A = \{\sum_{i=1}^{d} a_i Y_i : |a_i| \le cN^{1-\epsilon}\}, \ A' = \{\sum_{i=1}^{d} a_i Y'_i : |a_i| \le c'N^{\epsilon}\}$$

such that $AA' \subset B$, $A'A \subset B$. Moreover, assume Y_i, Y'_j are invertible for some $i, j \in [d]$. Then there is a constant C = C(c, c', d) so that when p is sufficiently large, for any $k \in [d]$, there exists some polynomial $f_k \in \mathbb{Z}[x] \setminus \{0\}$ of degree at most d with $H(f_k) \leq C$ and $\overline{f}_k(X_k) = 0$ in $M_n(\mathbb{F}_p)$.

Proof. Replacing x_i, y_i, y'_i by X_i, Y_i, Y'_i in the proof of Theorem 3, in the same way we obtain

$$A_1 \subset Y_i \cdot A', \ A_2 \subset A \cdot Y'_i$$

By assumption $A'A \subset B$, it follows that

$$A_1 A_2 \subset Y_i \cdot B \cdot Y'_i.$$

Now for any $k \in [d]$, there exists $T_k \in M_d(\mathbb{F}_p)$ with bounded entries so that

$$\det(T)\det(T')X_k\vec{v} = Y_i(T_k\vec{v})Y'_i$$

Since $X_1 = I_n$, Y_i, Y'_j are invertible, by Lemma 4 and the fact that B is proper, we conclude that T_1 must be invertible. Moreover,

$$T_k \vec{v} = \det(T) \det(T') Y_i^{-1} X_k \vec{v} (Y_j')^{-1}, \ \det(T) \det(T') T_1^{-1} \vec{v} = Y_i \vec{v} Y_j'.$$

Therefore

$$(\operatorname{adj}(T_1)T_k)\vec{v} = \det(T_1)T_1^{-1}T_k\vec{v} = \det(T_1)Y_i^{-1}X_kY_i\vec{v}.$$

Let $f_k(x) \in \mathbb{Z}[x]$ be the characteristic polynomial of $\operatorname{adj}(T_1)T_k$, then by Cayley-Hamilton's theorem

$$\overline{f}_k(\det(T_1)Y_i^{-1}X_kY_i)\vec{v} = \overline{f}_k(\operatorname{adj}(T_1)T_k)\vec{v} = 0.$$

In particular, using the fact $X_1 = I_n$ we can deduce

$$Y_i^{-1}\overline{f}_k(\det(T_1)X_k)Y_i = \overline{f}_k(\det(T_1)Y_i^{-1}X_kY_i) = 0,$$

which implies

$$\overline{f}_k(\det(T_1)X_k) = 0$$

3. Open questions

A natural question is can we weaken the condition on B or A to get the same result as in Theorem 3. A simple example shows that if we remove the isolated condition on B and allow A to be some arbitrary set with $|A| \approx |B|^{1/2}$, then the conclusion of Theorem 3 is no longer true. Let $B = \{a + bN : a, b \in [0, N - 1]\} = [0, N^2 - 1]$, which is a degenerate rank-2 GAP. We can take A = [0, N - 1] so that $AA \subset B$ and $|A| \approx |B|^{1/2}$. By choosing N to be about $p^{1/4}$ so that it's not the root of a low-height, low-degree polynomial, we've got a counterexample. From the argument of Theorem 2 we guess the isolated condition could probably be replaced by some restriction on the size of B.

One can also consider unbalanced GAPs. Suppose

$$B = \{\sum_{i=1}^{d} a_i x_i : |a_i| \le N_i\} \subset \mathbb{F}_p$$

and

$$A = \{\sum_{i=1}^{d} a_i y_i : |a_i| \le cM_i\}, \ A' = \{\sum_{i=1}^{d} a_i y'_i : |a_i| \le c'M'_i\}$$

with $|A||A'| \approx |B|$. If we make no restriction on the N_i 's, then there is no hope to obtain the same result. For example, B can be constructed with $N_i = 1$ for some i, and the corresponding x_i is not the root of some low-height, low-degree polynomial. The question is whether the conclusion of Theorem 3 remains true if we assume

$$\lim_{p \to \infty} N_i = \lim_{p \to \infty} M_i = \lim_{p \to \infty} M'_i = \infty$$

for each $i \in [d]$ to rule out the extreme case.

References

- M.-C. Chang. On a question of Davenport and Lewis and new character sum bounds in finite fields. Duke Math. J., 145(3):409-442, 2008.
- [2] G. Elekes and I. Z. Ruzsa. Few sums, many products. Studia Sci. Math. Hungar., 40(3):301-308, 2003.
- [3] P. Erdős and E. Szemerédi. On sums and products of integers. In Studies in pure mathematics, pages 213–218. Birkhäuser, Basel, 1983.
- [4] B. Hanson. Character sums over Bohr sets. Canad. Math. Bull., 58(4):774-786, 2015.
- [5] B. Kerr. Some multiplicative equations in finite fields. Finite Fields Appl., 75:Paper No. 101883, 28, 2021.
- [6] B. Kerr, J. Mello, and I. E. Shparlinski. An effective local-global principle and additive combinatorics in finite fields. J. Anal. Math., 152(1):109–135, 2024.
- [7] A. Mohammadi and S. Stevens. Attaining the exponent 5/4 for the sum-product problem in finite fields. Int. Math. Res. Not. IMRN, (4):3516–3532, 2023.
- [8] M. Rudnev, G. Shakan, and I. D. Shkredov. Stronger sum-product inequalities for small sets. Proc. Amer. Math. Soc., 148(4):1467–1479, 2020.
- [9] T. Tao and V. Vu. Additive combinatorics, volume 105 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2006.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160 USA *Email address*: ecroot@math.gatech.edu

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160 USA *Email address:* jmao87@gatech.edu