# Delayed Feedback in Online Non-Convex Optimization: A Non-Stationary Approach with Applications

Felipe Lara\*

Cristian Vega\*

December 20, 2024

#### **Abstract**

We study non-convex delayed-noise online optimization problems by evaluating dynamic regret in the non-stationary setting when the loss functions are quasar-convex. In particular, we consider scenarios involving quasar-convex functions either with a Lipschitz gradient or weakly smooth and, for each case, we ensure bounded dynamic regret in terms of cumulative path variation achieving sub-linear regret rates. Furthermore, we illustrate the flexibility of our framework by applying it to both theoretical settings such as zeroth-order (bandit) and also to practical applications with quadratic fractional functions. Moreover, we provide new examples of non-convex functions that are quasar-convex by proving that the class of differentiable strongly quasiconvex functions (Polyak 1966) are strongly quasar-convex on convex compact sets. Finally, several numerical experiments validate our theoretical findings, illustrating the effectiveness of our approach.

*Keywords*: Non-convex online optimization; Delayed algorithms; Quasar-convexity; Bandit; Quadratic fractional programming.

# 1 Introduction

Online optimization [16, 48, 40, 17] is an outstanding mathematical model for sequential decision-making which involves continuous optimization, machine learning and data sciences among others. It is used to provide automated decisions based on data, and its goal is to minimize the cumulative loss (or regret) of those decisions. This model has attracted the attention of the scientific community from several research fields in virtue of its applications in a variety of technological real-world problems, such as portfolio selection [2, 6], dynamic pricing [9], decision trees [19], machine learning [5], shortest paths [26], network resource allocation [45, 44] and generalized linear models [37] among others.

Each decision involves a loss determined by a function, which is revealed only after it is made. It can be viewed as a structured, repeated game between a learner and an adversary. At each round t, a player selects a decision  $x_t$  from a convex feasible set  $\mathcal{X} \subset \mathbb{R}^p$ . Then, an adversary gives a loss or cost function  $f_t \colon \mathcal{X} \to \mathbb{R}$ , and the player incurs a loss  $f_t(x_t)$ . The aim is to minimize the (stationary) regret  $R_T$ , defined as:

$$R_T := \sum_{t=1}^{T} f_t(x_t) - \min_{x \in \mathcal{X}} \sum_{t=1}^{T} f_t(x),$$

which is the difference between the cumulative loss of the algorithm and the cumulative loss of the best fixed decision in hindsight. Since the loss function is only revealed after the decision is made, we cannot expect to beat the opponent. Significant regret is required to be sub-linear in T, ensuring that as T becomes sufficiently large, the player performs as well to the best fixed decision.

A classical assumption in the literature is that the loss functions are assumed to be convex. This assumption is not only in virtue of the several applications of convex functions but also because of theoretical limitations

<sup>\*</sup>Instituto de Alta investigación (IAI), Universidad de Tarapacá, Arica, Chile. E-mail: felipelaraobreque@gmail.com; flarao@academicos.uta.cl. Web: felipelara.cl, ORCID-ID: 0000-0002-9965-0921

<sup>†</sup>Instituto de Alta investigación (IAI), Universidad de Tarapacá, Arica, Chile. E-mail: cristianvegacereno6@gmail.com, ORCID-ID: 0000-0001-7792-0137

since when we are dealing with non-convex loss functions, several useful properties are no longer valid as, for instance, local information is no longer global. In the very recent years, several efforts for dealing with non-convex loss functions have been developed, mainly motivated by several real-world applications in which classes of non-convex functions are more natural than convex ones. An interesting case occurs when dealing with (strongly) quasar-convex functions introduced in [7, 13, 21, 15] (with different names). This class of functions provides us several useful properties which are sufficient for ensuring good convergence rates for the gradient method, even when quasar-convex functions may have several local minimizers which are not necessarily global (see Example 3.7).

Another classical assumption is to assumed that the strategy is unchanged throughout the periods. However, in several real-world scenarios, regret is not adequate for evaluating performance. For instance, when aiming to maximize profit after several trades, the profit depends on the best investment at each round, rather than on the best investment for the sum of all loss functions at the final round. In such cases, the so-called non-stationary regret [4], which compares the cumulative loss with the minimum of each loss, is more suitable:

$$R_T^{NS} = \sum_{t=1}^T f_t(x_t) - \sum_{t=1}^T f_t(x_t^*), \tag{1.1}$$

where  $x_t^*$  represents the solution that minimizes  $f_t(x)$  over  $\mathcal{X}$ . Since  $f_t(x_t^*) \leq f_t(x^*)$ , for every  $t \in \{1, \dots, T\}$ , it follows that non-stationary regret is always greater or equal to the stationary regret, i.e.,  $R_T \leq R_T^{NS}$ . Moreover, dynamic regret has a theoretical advantage since the sum of non-convex functions may not have an unique minimizer even when each individual function has its own unique minimizer, for instance, strongly quasiconvex and strongly quasar-convex functions has unique global minimizers, but the sum of even two of these functions may have more than one minimizer. This makes dynamic regret particularly useful for addressing non-convexity.

Observe that the previous approach does not consider the potential delay between making a decision and receiving feedback, supposing that the loss function  $f_t$  is revealed immediately after the player makes their decision  $x_t$ . Delays like this have been extensively studied in [42, 38, 23, 43] and is common in real-world applications such as in economic analysis or online advertising [18]. For example, when assessing the impact of new economic or financial data, or political decisions with economic consequences, feedback is often delayed. Similarly, in online advertising, the output of the loss function may depend on whether a user clicks an ad. Users might take additional time before providing feedback, and non-feedback may only be inferred if the user does not click the ad after a sufficiently long period, which can make this situation uncertain for some time after the ad has been viewed.

Another classical extension of online optimization is the bandit setting, where the gradient of  $f_t$  is approximated by a zeroth-order approximation of it, making it suitable for problems [47] L. ZHANG, T. YANG, J. YI, R. JIN, Z.H. ZHOU, Improved dynamic regret for non-degenerate functions. where the gradient is not directly available or is computationally expensive to compute, even when the function itself is known explicitly. This extension proves useful in several applications in fields such as economics, robotics, statistics, machine learning and its online version has been extensively studied in [10, 1, 30, 39].

In this paper, we study online optimization in a very general setting: First, we deal with the non-convex case by assuming that the loss functions are not necessarily convex; in particular, we assume them to be quasar-convex (see [7, 13, 21, 15]), a class of functions which includes convex functions but also several classes of non-convex ones; second, we incorporate a delay between the decision and receiving feedback, which is a very realistic assumption for applied models in which the data needs extra time for being analyzed/computed; third, we assume a noise on the collecting data for obtaining more flexibility and applicability in our online optimization model. Under this general setting, we ensure bounded dynamic regret with sub-linear rate. Furthermore, we provide new examples of non-convex functions which belong to quasar-convexity as the class of strongly quasiconvex functions (Polyak [35]), which has been proved to be useful in several applications in mathematical models and algorithms (see [12, 22, 28, 29]). Finally, we apply our results in classes of (non-convex) quasar-convex functions as generalized linear models (GLM henceforth) and quadratic fractional functions, which are (strongly) quasiconvex. Several non-convex numerical experiments illustrate our findings.

The structure of the paper is as follows:In Section 2, we analyze the state of the art of the proposed problem under convexity/nonconvexity assumptions and with and without delay. In Section 3, we introduce the notation along with some preliminary concepts. Section 4 presents the main algorithm and provides a regret bound. Additionally, we demonstrate the flexibility of our approach by extending the algorithm and its theoretical guarantees to the bandit setting. In Section 5, we present new examples of quasar-convex functions that will be used in the subsequent section. Section 6 evaluates the algorithm's performance on three numerical applications: highly non-convex functions, generalized linear models, and quadratic fractional functions. Finally, Section 7 offers conclusions and future research directions.

## 2 State of the Art and Related Work

The standard algorithm for online optimization is the online gradient descent (OGD) algorithm. At each step, the algorithm updates its decision based on the gradient of the loss function at the current decision:

$$x_{t+1} = P_{\mathcal{X}}(x_t - \eta_t \nabla f_t(x_t)),$$

where  $P_{\mathcal{X}}$  denotes the projection onto the convex set  $\mathcal{X}$ , and  $\eta_t$  is the step-size (learning rate). Several papers have established sub-linear regret bounds for different classes of functions. For convex functions without delay, [48] proved a stationary regret bound of  $\mathcal{O}(T^{\frac{1}{2}})$ , while [16] improved this result to a regret rate of  $\mathcal{O}(\ln(T))$  for strongly convex functions. In [10] and [1] is extended for the bandit setting for convex and strongly convex, respectively.

In [38] is proposed the delayed version of OGD (DOGD) algorithm, and it is proved that DOGD with a constant learning rate  $\eta_t = \frac{1}{L(T+D)^{\frac{1}{2}}}$  attains a regret bound of  $\mathcal{O}(D^{\frac{1}{2}})$ , where D is sum of the delay. The extension of DOGD for the bandit setting was studied in [30]. Observe that, the constant learning rate used in DOGD does not take advantage of the strong convexity of loss functions. In standard OGD, [16] established a  $\ln(T)$  regret bound for  $\beta$ -strongly convex functions by setting  $\eta_t = \frac{1}{\beta t}$ . This learning rate leverages the increasing nature of the inverse of  $\frac{1}{\eta_t}$  by the modulus of strong convexity of  $f_t$ , where  $\frac{1}{\eta_t} - \frac{1}{\eta_{t-1}} = \beta$ . Inspired by this, [42] proposes an initialization where  $\frac{1}{\eta_0} = 0$ . Otherwise,  $\frac{1}{\eta_t} - \frac{1}{\eta_{t-1}} = a\beta$ , where a is the number of gradient that arrives at round t. DOGD for strongly convex functions achieves a regret bound of  $\mathcal{O}(d \ln(T))$ , where d is the maximum delay. The results for the stationary setting are summarized in Table 1.

In [4], it is shown that without any additional assumptions regarding how the loss functions evolve over time, the non-stationary regret can grow linearly with time T. To achieve a sub-linear non-stationary regret, it is essential to impose additional assumptions on the variation of the loss functions over time, often quantified by the path variation. Specifically, [4] establishes non-stationary regret bounds of  $\mathcal{O}(\hat{V}_T^{\frac{1}{3}}T^{\frac{2}{3}})$  for online gradient descent with noisy gradient feedback for convex functions,  $\mathcal{O}(\hat{V}_T^{\frac{1}{2}}T^{\frac{1}{2}})$  for strongly convex functions, and  $\mathcal{O}(\hat{V}_T^{\frac{1}{3}}T^{\frac{2}{3}})$  for convex functions with noisy gradient feedback and strongly convex functions with noisy function values. Where  $\hat{V}_T$  denotes the cumulative variation of the functions.

In [4] is achieved a bound of  $\mathcal{O}(T^{\frac{1}{3}}V_T^{\frac{2}{3}})$  for convex functions and  $\mathcal{O}(T^{\frac{1}{2}}V_T^{\frac{1}{2}})$  for strongly convex functions, where  $V_T$  is the cumulative path defined as follows:

$$V_T = \sum_{t=1}^{T-1} \|x_t^* - x_{t+1}^*\|.$$

In the delayed setting, [38] obtained a stationary regret bound of  $\mathcal{O}(T^{\frac{1}{2}})$  for convex functions, while [42] showed that strongly convex functions yield a logarithmic regret bound of  $\mathcal{O}(\ln(T))$ . In the context of non-stationary regret, [4] achieved a bound of  $\mathcal{O}(T^{\frac{1}{3}}V_T^{\frac{2}{3}})$  for convex functions and  $\mathcal{O}(T^{\frac{1}{2}}V_T^{\frac{1}{2}})$  for strongly convex functions, both without delay. In [31] the bound for strongly convex was improved to  $\mathcal{O}(T^{\frac{1}{2}}V_T^{\frac{1}{2}})$ . For the delayed setting in [43] provided a non-stationary regret bound of  $\mathcal{O}((dTV_T)^{\frac{1}{2}})$  for convex functions.

In [27] is studied OGD for functions satisfying Polyak-Łojasiewicz condition, while in [47] is studied for functions that have quadratic growth. In [46] composition of increasing functions with linear terms are studied. In [11] weakly pseudo-convex functions are considered achieving a stationary regret bound of  $\mathcal{O}(V_T)$  without delay. Lastly, for quasar-convex functions, [37] provided several significant results: a stationary regret bound of  $\mathcal{O}(T^{\frac{1}{2}}V_T^{\frac{1}{2}})$  under delay, a bound of  $\mathcal{O}(V_T)$  for weakly smooth and quasar-convex functions without delay, and another bound of  $\mathcal{O}(V_T)$  for strongly quasar-convex functions without delay. The results for the non-stationary setting are summarized in Table 2.

Work	Function	Delay	Regret Bound
[48]	Convex	No	$\mathcal{O}(T^{\frac{1}{2}})$
[16]	Strongly Convex	No	$\mathcal{O}(\ln(T))$
[38]	Convex	Yes	$\mathcal{O}(D^{rac{1}{2}})$
[42]	Strongly Convex	Yes	$\mathcal{O}(d\ln(T))$

Table 1: Regret bounds for stationary regret

Work	Function	Delay	Regret Bound
[4]	Convex	No	$\mathcal{O}(T^{\frac{1}{3}}V_T^{\frac{2}{3}})$
[31]	Strongly Convex	No	$\mathcal{O}(V_T)$
[43]	Convex	Yes	$\mathcal{O}((dTV_T))$
[37]	Quasar-Convex	No	$\mathcal{O}(T^{\frac{1}{2}}V_T^{\frac{1}{2}})$
[37]	Weakly Smooth & Quasar-Convex	No	$\mathcal{O}(V_T)$
[37]	Strongly Quasar-Convex	No	$\mathcal{O}(V_T)$
Our work	Quasar-Convex	Yes	$\mathcal{O}(dT^{\frac{1}{2}}V_T^{\frac{1}{2}})$
Our work	Weakly Smooth & Quasar-Convex	Yes	$\mathcal{O}(dV_T)$

Table 2: Contribution and Comparison of non-stationary regret with related works

## 3 Preliminaries

Let  $f: \mathbb{R}^p \to \mathbb{R}$  be a real-valued function. It is indicated by  $\operatorname{epi} f := \{(x,t) \in \mathbb{R}^p \times \mathbb{R} : f(x) \leq t\}$  the epigraph of f, by  $S_{\lambda}(f) := \{x \in \mathbb{R}^p \mid f(x) \leq \lambda\}$  the sub-level set of f at the height  $\lambda \in \mathbb{R}$  and by  $\operatorname{argmin} f$  the set of all minimal points of f.

A function f is said to be:

(a) convex if, given any  $x, y \in \mathbb{R}^p$ , then

$$f(\lambda x + (1 - \lambda)y) \le \lambda f(x) + (1 - \lambda)f(y), \ \forall \ \lambda \in [0, 1], \tag{3.1}$$

(b) strongly convex with modulus  $\gamma \in ]0, +\infty[$  if for all  $x, y \in \mathbb{R}^p$  and all  $\lambda \in [0, 1]$ , we have

$$f(\lambda y + (1 - \lambda)x) \le \lambda f(y) + (1 - \lambda)f(x) - \lambda(1 - \lambda)\frac{\gamma}{2} ||x - y||^2, \tag{3.2}$$

(c) quasiconvex if, given any  $x, y \in \mathbb{R}^p$ , then

$$f(\lambda x + (1 - \lambda)y) \le \max\{f(x), f(y)\}, \ \forall \ \lambda \in [0, 1],$$
 (3.3)

(d) strongly quasiconvex [35] with modulus  $\gamma \in ]0, +\infty[$  if for all  $x, y \in \mathbb{R}^p$  and all  $\lambda \in [0, 1]$ , we have

$$f(\lambda y + (1 - \lambda)x) \le \max\{f(y), f(x)\} - \lambda(1 - \lambda)\frac{\gamma}{2}||x - y||^2.$$
 (3.4)

It is said that f is strictly convex (resp. strictly quasiconvex) if the inequality in (3.1) (resp. (3.3)) is strict whenever  $x \neq y$  and  $\lambda \in ]0,1[$ .

The relationship between all these notions is summarized below (we denote quasiconvex by qcx):

$$\begin{array}{cccc} \text{strongly convex} & \Longrightarrow & \text{strictly convex} & \Longrightarrow & \text{convex} \\ & & & \downarrow & & \downarrow \\ \text{strongly qcx} & \Longrightarrow & \text{strictly qcx} & \Longrightarrow & \text{qcx} \end{array}$$

All the reverse statements does not hold in general (see [8, 14, 28]). Note that lsc strongly quasiconvex functions have an unique minimizer on closed convex sets, as we recall next.

**Lemma 3.1.** ([28, Corollary 3]) Let  $K \subseteq \mathbb{R}^p$  be a closed and convex set and  $f: K \to \mathbb{R}$  be a lsc and strongly quasiconvex function on K with modulus  $\gamma > 0$ . Then argmin f is a singleton.

Furthermore, the unique minimizer  $x^* \in K$  of the strongly quasiconvex function f satisfies a quadratic growth condition (see [25]):

$$f(x^*) + \frac{\gamma}{8} ||x^* - y||^2 \le f(y), \ \forall \ y \in K.$$
 (3.5)

For differentiable strongly quasiconvex functions, we have the following characterization [41].

**Lemma 3.2.** ([41, Theorems 1 and 6]) Let  $K \subseteq \mathbb{R}^p$  be a convex and open set and  $f: K \to \mathbb{R}$  be a differentiable function. Then f is strongly quasiconvex with modulus  $\gamma \geq 0$  if and only if for every  $x, y \in K$ , we have

$$f(x) \le f(y) \implies \langle \nabla f(y) \mid x - y \rangle \le -\frac{\gamma}{2} ||x - y||^2.$$

The previous result extends the well-known characterization for differentiable quasiconvex functions given by [3]  $(\gamma = 0)$ .

Before continuing, we present some examples of strongly quasiconvex functions that are not (necessarily) convex.

- **Remark 3.3.** (i) Let  $f: \mathbb{R}^p \to \mathbb{R}$  be given by  $f(x) = ||x||^{\alpha}$ , with  $\alpha \in ]0,2]$ . Clearly, f is non-convex, but it is strongly quasiconvex on any  $\mathbb{B}(0,r)$ , r > 0, by [32, Corollary 3.9]  $(0 < \alpha < 1)$ , [24, Theorem 2]  $(\alpha = 1)$  and [41, page 27]  $(1 < \alpha \le 2)$ .
  - (ii) Let  $A, B \in \mathbb{R}^{p \times p}$ ,  $a, b \in \mathbb{R}^p$ ,  $\alpha, \beta \in \mathbb{R}$ , and  $f : \mathbb{R}^p \to \mathbb{R}$  be the functions given by:

$$f(x) = \frac{g(x)}{q(x)} = \frac{\frac{1}{2}\langle Ax \mid x \rangle + \langle a \mid x \rangle + \alpha}{\frac{1}{2}\langle Bx \mid x \rangle + \langle b \mid x \rangle + \beta}.$$
 (3.6)

Take 0 < m < M and define:

$$K := \{ x \in \mathbb{R}^p \mid m \le q(x) \le M \}.$$

If *A* is a positive definite matrix and at least one of the following conditions holds:

- (a) B = 0 (the null matrix),
- (b) g is non-negative on K and B is negative semi-definite,
- $(c) \ \ g$  is non-positive on K and B is positive semi-definite,

then f is strongly quasiconvex on K with modulus  $\gamma = \frac{\sigma_{\min}(A)}{M}$  by [22, Proposition 4.1], where  $\sigma_{\min}(A)$  is the minimum eigenvalue of A.

- (iii) Let  $f_1, f_2 \colon \mathbb{R}^p \to \mathbb{R}$  be two strongly quasiconvex functions with modulus  $\gamma_1, \gamma_2 > 0$ , respectively. Then  $f := \max\{f_1, f_2\}$  is strongly quasiconvex with modulus  $\gamma := \min\{\gamma_1, \gamma_2\} > 0$  (see [41]).
- (iv) Let  $\alpha > 0$  and  $f : \mathbb{R}^p \to \mathbb{R}$  be a strongly quasiconvex function with modulus  $\gamma > 0$ . Then  $\alpha f$  is strongly quasiconvex with modulus  $\gamma \alpha > 0$  (straightforward).
- (v) If  $A: \mathbb{R}^p \to \mathbb{R}^p$  is a linear operator and f is a strongly quasiconvex function with modulus  $\gamma \geq 0$ , then  $g:=f\circ A$  is strongly quasiconvex with modulus  $\gamma\sigma_{\min}(A)\geq 0$  [12] (where  $\sigma_{\min}(A)$  is the minimum positive eigenvalue of A).

**Definition 3.4.** (See, for instance,[21, Definition 1]) Let  $\kappa \in (0,1]$  and  $f: \mathbb{R}^p \to \mathbb{R}$  be a differentiable function. Then f is said to be  $\kappa$ -quasar-convex with respect to  $x^* \in \underset{\mathbb{R}^p}{\operatorname{argmin}} f$  if

$$f(x^*) \ge f(x) + \frac{1}{\kappa} \langle \nabla f(x) \mid x^* - x \rangle, \ \forall \ x \in \mathbb{R}^p.$$

Further, for  $\gamma \geq 0$ , the function f is  $(\kappa, \gamma)$ -strongly quasar-convex if for all  $x \in \mathbb{R}^p$ ,

$$f(x^*) \ge f(x) + \frac{1}{\kappa} \langle \nabla f(x) \mid x^* - x \rangle + \frac{\gamma}{2} ||x^* - x||^2.$$

If  $\kappa=1$ , then a function is 1-quasar-convex if and only if it is star-convex (see [34]). Similarly, a function is  $(1,\gamma)$ -quasar-convex if and only if it is quasi-strongly convex (see [33]). These notions are variants of convexity and strong convexity, respectively, where the conditions are required to hold only at a minimizer  $x^*$ , rather than for all points in  $\mathbb{R}^p$ . Furthermore, note that as  $\kappa$  decreases, the function becomes "more non-convex," indicating that the term  $\frac{1}{\kappa}\langle \nabla f(x) \mid x^* - x \rangle$  can become more negative. We now introduce a characterization of quasar-convexity.

**Lemma 3.5.** [21, Lemma 11] Let  $f: \mathcal{X} \to \mathbb{R}$  be differentiable with  $x^* \in \underset{\mathbb{R}^p}{\operatorname{argmin}} f$ , where the domain  $\mathcal{X} \subseteq \mathbb{R}^p$  is open and convex. Then, f is quasar-convex with modulus  $\kappa \in (0,1]$  if and only if

$$f(\lambda x^* + (1 - \lambda)x) \le \kappa \lambda f(x^*) + (1 - \kappa \lambda)f(x), \ \forall \ x \in \mathcal{X}, \ \forall \ \lambda \in [0, 1].$$

**Proposition 3.6.** Let  $f_1: \mathbb{R} \to \mathbb{R}$  be  $\kappa$ -quasar-convex with respect to the point 0, such that  $\min_{x \in \mathbb{R}} f_1(0) = f_1(0) = 0$ , a  $C_1$ , and let  $f_2: S^{d-1} \to \mathbb{R}$  be an arbitrary non-negative  $C_1$  function defined on the unit sphere of  $\mathbb{R}^p$ , satisfying  $f_2 \geq 0$ . We define the function f as  $f(x) = f_1(\|x\|)f_2\left(\frac{x}{\|x\|}\right)$  with f(0) = 0. Then, the function f is non-negative,  $C_1$ , and  $\kappa$ -quasar-convex on  $\mathbb{R}^p$  with respect to  $x^* = 0$ . Moreover, the gradient of f(x) is given by:

$$\nabla f(x) = \frac{x}{\|x\|} f_1'(\|x\|) f_2\left(\frac{x}{\|x\|}\right) + \frac{1}{\|x\|} \left( \text{Id } -\frac{xx^\top}{\|x\|^2} \right) f_1(\|x\|) \nabla f_2\left(\frac{x}{\|x\|}\right).$$

*Proof.* Since both  $f_1$  and  $f_2$  are  $C_1$ , it follows that f(x) is also  $C_1$ . Applying the product rule component-wise, we obtain:

$$\nabla f(x) = \nabla_x f_1(\|x\|) f_2\left(\frac{x}{\|x\|}\right) + f_1(\|x\|) \nabla_x f_2\left(\frac{x}{\|x\|}\right)$$
$$= \frac{x}{\|x\|} f_1'(\|x\|) f_2\left(\frac{x}{\|x\|}\right) + \frac{1}{\|x\|} \left(\operatorname{Id} - \frac{xx^{\top}}{\|x\|^2}\right) f_1(\|x\|) \nabla f_2\left(\frac{x}{\|x\|}\right).$$

The non-negativity of  $f_1$  and  $f_2$  implies that f is non-negative as well. To prove the quasar-convexity of f, we show that for all  $x \in \mathbb{R}^p$  and  $\lambda \in [0, 1]$ , the following inequality holds:

$$f((1-\lambda)x) = f(\lambda x^* + (1-\lambda)x) < (1-\kappa\lambda)f(x) + \lambda\kappa f(x^*) = (1-\kappa\lambda)f(x).$$

Using the quasar-convexity of f at  $x^*$ , we have:

$$f((1-\lambda)x) = f_1((1-\lambda)||x||)f_2\left(\frac{x}{||x||}\right) \le (1-\kappa\lambda)f_1(||x||)f_2\left(\frac{x}{||x||}\right) = (1-\kappa\lambda)f(x).$$

This concludes the proof.

**Example 3.7.** Consider the function  $f: \mathbb{R}^2 \to \mathbb{R}$ , defined as:

$$f(x) = f_1(||x||) f_2\left(\frac{x}{||x||}\right),$$

where  $f_1(t) = \frac{t^2}{1+t^2}$  and

$$f_2(x_1, x_2) = \frac{1}{4N} \sum_{i=1}^{N} (a_i \sin(b_i x_1)^2 + c_i \cos(d_i x_2)^2),$$

with N=10. Here,  $a_i$  and  $c_i$  are independently and uniformly distributed on [0,20], while  $b_i$  and  $d_i$  are independently and uniformly distributed on [-25,25]. The function is illustrated on the left side of Figure 3.7.

By Lemma 3.6, this function is  $\frac{1}{1+R^2}$ -quasar-convex on the ball centered at the origin with radius R, since  $f_1$  is  $\frac{1}{1+R^2}$ -quasar-convex on [-R,R], independent of the specific form of  $f_2$ . Note that this function is similar to the one introduced in [20], with the main difference being that our function  $f_2$  remains positive and not lower-bounded by 1, whereas  $f_1$  is not strongly quasar-convex. Moreover, for any fixed direction, it behaves like  $cf_1(\|x\|)$ , where c is a constant. However, since  $f_2$  is a linear combination of high-frequency trigonometric functions, it can exhibit highly non-convex behavior. This is evident when examining the segment between two arbitrary points u and v that does not contain the minimizer, as shown on the left side of Figure 3.7.




Figure 1: An illustration of the non-convex function f(x) described in Example 3.7. On the left-hand side, a 3D plot of the function f(x) is displayed, while on the right-hand side, an arbitrary segment that does not contain the minimizer is shown, highlighting the non-convexity of f.

A differentiable function  $f: \mathbb{R}^p \to \mathbb{R}$  is said to be ([21, Definition 1])  $\Gamma$ -weakly smooth with respect to  $x^* \in \underset{x \in \mathbb{R}^p}{\operatorname{argmin}} f(x)$ , if for any  $x \in \mathbb{R}^p$ , we have

$$\|\nabla f(x)\|^2 \le \Gamma \left( f(x) - f(x^*) \right).$$

The following lemma provides a method to interchange the limits of a double sum, and will be applied in the proof of Theorem 4.2.

**Lemma 3.8.** Let  $\{a_k\}_{k=1}^T$  be a positive sequence. Then, the following holds:

$$\sum_{t=1}^{T-1} \sum_{k=t+1}^{\min(T,t+d-1)} a_k = \sum_{k=2}^{T} \sum_{t=\max(2,k-d+1)}^{k-1} a_k \le (d-1) \sum_{k=1}^{T} a_k.$$
(3.7)

*Proof.* For each t, the index k ranges from t+1 to  $\min(T, t+d-1)$ , which can be rewritten as a sum over k. For each  $k \in [2, T]$ , t ranges from  $\max(2, k-d+1)$  to k-1. Thus,

$$\sum_{t=1}^{T-1} \sum_{k=t+1}^{\min(T,t+d-1)} a_k = \sum_{k=2}^{T} \sum_{t=\max(2,k-d+1)}^{k-1} a_k. \le (d-1) \sum_{t=1}^{T} a_k.$$

where the last inequality is since the inner sum has at most d-1 terms, giving the bound and completing the proof.

## 4 Main Results

In this section, inspired by the techniques developed in [42], we derive non-stationary regret bounds for the iterates generated by DOGD when applied to quasar-convex functions. In each round t, DOGD queries the gradient  $\nabla f_t(x_t)$ , but due to the delay, this gradient is received at the end of round  $t' = t + d_t - 1$  and can be utilized in round  $t + d_t$ , where  $d_t \in \mathbb{Z}^+$  represents a non-negative integer delay. DOGD updates the decision  $x_t$  with the sum of the gradients received by the end of round t. The set of rounds whose gradients are received at each round t is denoted by  $\mathcal{F}_t = \{k \in [T] \mid k + d_k - 1 = t\}$ , where  $[T] := \{1, \ldots, T\}$ . In the setting without delays, for all t, D = T,  $d = d_t = 1$ , t' = t, and  $\mathcal{F}_t = \{t\}$ . The detailed procedure for DOGD is summarized in Algorithm 1. Specifically, we address cases where the functions are L-Lipschitz and  $\Gamma$ -weakly smooth. Unlike previous works, we tackle the more challenging scenario where only a gradient estimate is available rather than full access to the gradient. This will allow us to explore the bandit setting of this algorithm, where only a zeroth-order gradient estimate is available, and analyze the convergence of the offline version of this algorithm.

## Algorithm 1 Delayed projected Online Gradient Descent for quasar-convex functions

- 1: **Initialization:** Choose an arbitrary vector  $x_1 \in \mathcal{X}$  and a sequence of step-sizes  $\{\eta_t\}_{t=1}^T$
- 2: **for** t = 1, 2, ..., T **do**
- 3: Query  $r_t \approx \nabla f_t(x_t)$ .

4: 
$$x_{t+1} = \begin{cases} P_{\mathcal{X}} \left( x_t - \eta_t \sum_{k \in \mathcal{F}_t} r_k \right) & \text{if } |\mathcal{F}_t| > 0 \\ x_t & \text{otherwise} \end{cases}$$

5: end for

# 4.1 Delayed Online Gradient Descent for Quasar-Convex Functions with Non-Stationary Regret

In the quasar-convex case, the following assumptions will be needed:

#### Assumption 4.1.

- (B1) The radius of the convex decision set  $\mathcal{X}$  is bounded by R, i.e.,  $||x|| \leq R$  for all  $x \in \mathcal{X}$ .
- (B2) Each loss function  $f_t$  is  $\kappa$ -quasar-convex for some  $x_t^* \in \operatorname{argmin} f_t(x)$ .
- (B3) Each loss function  $f_t \colon \mathcal{X} \to \mathbb{R}$  is L-Lipschitz.
- (B4) Each loss function  $f_t \colon \mathcal{X} \to \mathbb{R}$  is  $\Gamma$ -weakly smooth.

To simplify the notation, for every  $t \in [T]$ , we define  $r_t = \nabla f_t(x_t) + m_t$ , where  $m_t \in \mathbb{R}^p$  represents the deterministic gradient error. We assume that  $m_t$  is bounded for some constant  $\delta_t > 0$ . The cumulative gradient error and the cumulative squared gradient over the horizon T are defined as:

$$\Delta_T = \sum_{t=1}^T \delta_t$$
 and  $\Lambda_T = \sum_{t=1}^T \delta_t^2$ ,

respectively. Additionally, we define

$$x'_{t+1} = \begin{cases} x_t - \eta_t \sum_{k \in \mathcal{F}_t} r_k & \text{if } |\mathcal{F}_t| > 0\\ x_t & \text{otherwise} \end{cases}$$

which the term inside of the projection. Then, by definition, for every  $t \in [T + d - 1]$ , it holds that

$$\eta_t \sum_{k \in \mathcal{F}_t} r_k = (x_t - x'_{t+1}). \tag{4.1}$$

According to Algorithm 1, there could exist some feedback that arrives after the round T and is not used to update the decision. However, it is useful for the analysis. Therefore, for  $t \in [T+1, T+d-1]$ , we also define  $\mathcal{F}_t = \{k \in [T] \mid k+d_k-1=t\}$ , and the virtual update as:

$$x_{t+1} = \begin{cases} P_{\mathcal{X}} \left( x_t - \eta_t \sum_{k \in \mathcal{F}_t} r_k \right) & \text{if } |\mathcal{F}_t| > 0, \\ x_t & \text{otherwise.} \end{cases}$$

Analogously, for any  $t \in [T+1, T+d-1]$ , we also define

$$x'_{t+1} = \begin{cases} x_t - \eta_t \sum_{k \in \mathcal{F}_t} r_k & \text{if } |\mathcal{F}_t| > 0, \\ x_t & \text{otherwise.} \end{cases}$$

Let us also define  $s = \min \left\{ t \in [T+d-1] \mid |\mathcal{F}_t| > 0 \right\}$  . Observe that

$$\bigcup_{t=s}^{T+d-1} \mathcal{F}_t = \bigcup_{t=1}^{T+d-1} \mathcal{F}_t = [T],$$

and  $\mathcal{F}_i \cap \mathcal{F}_j = \emptyset$  for any  $i \neq j$ . Combining these results, we obtain that, for every finite sequence  $\{a_k\}_{k=1}^T$  of real numbers, we have that:

$$\sum_{t=1}^{T} a_t = \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_t} a_k. \tag{4.2}$$

Our main result is given below.

**Theorem 4.2.** *Under assumptions (B1)-(B2), the following assertions hold:* 

(i) If (B3) holds, the regret of Algorithm 1 with a constant step-size  $0 < \eta_t = \eta$ , can be upper-bounded by:

$$R_T^{NS} \le \frac{2R^2}{\eta \kappa} + \frac{3R + \eta L(d-1)}{\eta \kappa} V_T + \frac{L^2d + 4(d-1)L^2}{2\kappa} \eta T + \frac{\eta d}{2\kappa} \Lambda_T + \frac{\eta dL + 2R + 2\eta(d-1)L}{\kappa} \Delta_T \quad \textbf{(4.3)}$$

(ii) If (B4) holds, the regret of Algorithm 1 with a constant step-size  $0 < \eta_t = \eta < \frac{1}{\bar{\alpha}} = \frac{2\kappa}{\left(d+4d^{\frac{1}{2}}(d-1)\right)\Gamma}$ , can be upper-bounded by:

$$R_T^{NS} \le \frac{b^2 + 2ac + b\left(b^2 + 4ac\right)^{\frac{1}{2}}}{2a^2},$$
 (4.4)

where  $a = 1 - \alpha \eta > 0$ ,

$$b = \frac{(2R(d-1)\Gamma V_T)^{\frac{1}{2}}}{\kappa} + \frac{(2d)^{\frac{1}{2}}(d-1)\eta + \eta d}{\kappa} (\Gamma \Lambda_T)^{\frac{1}{2}} \quad \text{and} \quad c = \frac{2R^2}{\kappa} + \frac{3RV_T}{\kappa} + \frac{\eta d}{2\kappa} \Lambda_T + \frac{2R}{\kappa} \Delta_T.$$

*Proof.* By assumption (B2), we have for every  $t \in [T]$  that:

$$f_{t}(x_{t}) - f_{t}(x_{t}^{*}) \leq \frac{1}{\kappa} \langle \nabla f_{t}(x_{t}) \mid x_{t} - x_{t}^{*} \rangle = \frac{1}{\kappa} \left( \langle \nabla f_{t}(x_{t}) \mid x_{t} - x_{t'} \rangle + \left\langle \nabla f_{t}(x_{t}) \mid x_{t'} - x_{\bar{t'}}^{*} \right\rangle + \left\langle \nabla f_{t}(x_{t}) \mid x_{\bar{t'}} - x_{\bar{t'}}^{*} \right\rangle \right)$$

$$= \frac{1}{\kappa} \langle \nabla f_{t}(x_{t}) \mid x_{t} - x_{t'} \rangle + \frac{1}{\kappa} \langle r_{t} \mid x_{t'} - x_{\bar{t'}}^{*} \rangle - \frac{1}{\kappa} \langle m_{t} \mid x_{t'} - x_{\bar{t'}}^{*} \rangle + \frac{1}{\kappa} \langle \nabla f_{t}(x_{t}) \mid x_{\bar{t'}} - x_{\bar{t'}}^{*} \rangle. \tag{4.5}$$

Summing (4.5) from 1 to T and multiplying by  $\kappa$ , we obtain:

$$\kappa R_{T}^{NS} \leq \sum_{t=1}^{T} \left\langle \nabla f_{t}(x_{t}) \mid x_{\bar{t}'}^{*} - x_{t}^{*} \right\rangle + \sum_{t=1}^{T} \left( \left\langle r_{t} \mid x_{t'} - x_{\bar{t}'}^{*} \right\rangle - \left\langle m_{t} \mid x_{t'} - x_{\bar{t}'}^{*} \right\rangle \right) + \sum_{t=1}^{T} \left\langle \nabla f_{t}(x_{t}) \mid x_{t} - x_{t'} \right\rangle \\
\leq \sum_{t=1}^{T} \left\langle \nabla f_{t}(x_{t}) \mid x_{\bar{t}'}^{*} - x_{t}^{*} \right\rangle + \sum_{t=1}^{T} \left( \left\langle r_{t} \mid x_{t'} - x_{\bar{t}'}^{*} \right\rangle + \|m_{t}\| \|x_{t'} - x_{\bar{t}'}^{*}\| \right) + \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_{t}} \left\langle \nabla f_{k}(x_{k}) \mid x_{k} - x_{t} \right\rangle \\
\leq \sum_{t=1}^{T} \left\langle \nabla f_{t}(x_{t}) \mid x_{t} - x_{t'} \right\rangle + \sum_{t=1}^{T} \left\langle r_{t} \mid x_{t'} - x_{\bar{t}'}^{*} \right\rangle + 2R \frac{p^{\frac{1}{2}}Gh}{2} T + \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_{t}} \|\nabla f_{k}(x_{k})\| \|x_{\bar{t}}^{*} - x_{k}^{*}\|. \tag{4.6}$$

where the second and third inequalities follow from the Cauchy-Schwarz.

Now, we now establish an upper bound for the second term on the right side of (4.6):

$$\sum_{t=1}^{T} \left\langle r_{t} \mid x_{t'} - x_{\bar{t}'}^{*} \right\rangle = \sum_{t=s}^{T+d-1} \left\langle \sum_{k \in \mathcal{F}_{t}} r_{k} \mid x_{t} - x_{\bar{t}}^{*} \right\rangle = \sum_{t=s}^{T+d-1} \frac{1}{\eta} \left\langle x_{t} - x_{t+1}' \mid x_{t} - x_{\bar{t}}^{*} \right\rangle$$

$$= \sum_{t=s}^{T+d-1} \frac{1}{2\eta} \left( \|x_{t} - x_{\bar{t}}^{*}\|^{2} - \|x_{t+1}' - x_{\bar{t}}^{*}\|^{2} + \|x_{t} - x_{t+1}'\|^{2} \right)$$

$$= \sum_{t=s}^{T+d-1} \frac{1}{2\eta} \left( \|x_{t} - x_{\bar{t}}^{*}\|^{2} - \|x_{t+1}' - x_{\bar{t}}^{*}\|^{2} + \eta^{2} \left\| \sum_{k \in \mathcal{F}_{t}} r_{k} \right\|^{2} \right)$$

$$\leq \sum_{t=s}^{T+d-1} \frac{1}{2\eta} \left( \|x_{t} - x_{\bar{t}}^{*}\|^{2} - \|x_{t+1} - x_{\bar{t}}^{*}\|^{2} + \eta^{2} \left\| \sum_{k \in \mathcal{F}_{t}} r_{k} \right\|^{2} \right)$$

$$\leq \sum_{t=s}^{T+d-1} \frac{1}{2\eta} \left( \|x_{t} - x_{\bar{t}}^{*}\|^{2} - \|x_{t+1} - x_{\bar{t}}^{*}\|^{2} + \eta^{2} \sum_{k \in \mathcal{F}_{t}} |\mathcal{F}_{t}| \|r_{k}\|^{2} \right)$$

$$= \frac{1}{2\eta} \sum_{t=s}^{T-1} \left( \|x_{t} - x_{\bar{t}}^{*}\|^{2} - \|x_{t+1} - x_{\bar{t}}^{*}\|^{2} \right) + \frac{1}{2\eta} \sum_{t=T}^{T+d-1} \left( \|x_{t} - x_{\bar{t}}^{*}\|^{2} - \|x_{t+1} - x_{\bar{t}}^{*}\|^{2} \right) + \frac{\eta}{2} \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_{t}} |\mathcal{F}_{t}| \|r_{k}\|^{2}$$

$$= \frac{1}{2\eta} \sum_{t=s}^{T-1} \left( \|x_{t} - x_{\bar{t}}^{*}\|^{2} - \|x_{t+1} - x_{\bar{t}}^{*}\|^{2} \right) + \frac{1}{2\eta} \left( \|x_{T} - x_{T}^{*}\|^{2} - \|x_{T+d} - x_{T}^{*}\|^{2} \right) + \frac{\eta}{2} \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_{t}} |\mathcal{F}_{t}| \|r_{k}\|^{2}$$

$$\leq \frac{1}{2\eta} \sum_{t=s}^{T-1} \left( \|x_{t} - x_{\bar{t}}^{*}\|^{2} - \|x_{t+1} - x_{\bar{t}}^{*}\|^{2} \right) + \frac{1}{2\eta} \left( \|x_{T} - x_{T}^{*}\|^{2} - \|x_{T+d} - x_{T}^{*}\|^{2} \right) + \frac{\eta d}{2} \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_{t}} |\mathcal{F}_{t}| \|r_{k}\|^{2}$$

$$\leq \frac{1}{2\eta} \sum_{t=s}^{T-1} \left( \|x_{t} - x_{\bar{t}}^{*}\|^{2} - \|x_{t+1} - x_{\bar{t}}^{*}\|^{2} \right) + \frac{1}{2\eta} \left( \|x_{T} - x_{T}^{*}\|^{2} - \|x_{T+d} - x_{T}^{*}\|^{2} \right) + \frac{\eta d}{2} \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_{t}} |\mathcal{F}_{t}| \|r_{k}\|^{2}$$

$$\leq \frac{1}{2\eta} \sum_{t=s}^{T-1} \left( \|x_{t} - x_{t}^{*}\|^{2} - \|x_{t+1} - x_{t}^{*}\|^{2} \right) + \frac{1}{2\eta} \left( \|x_{T} - x_{T}^{*}\|^{2} - \|x_{T+d} - x_{T}^{*}\|^{2} \right) + \frac{\eta d}{2} \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_{t}} |\mathcal{F}_{t}| \|r_{k}\|^{2}$$

$$\leq \frac{1}{2\eta} \sum_{t=s}^{T-1} \left( \|x_{t} - x_{t}^{*}\|^{2} - \|x_{t+1} - x_{t}^{*}\|^{2} \right) + \frac{1}{2\eta} \left( \|x_{T} - x_{T}^{*}\|^{2} - \|x_{T+d} - x_{T}^{*}\|^{2}$$

where the first equality follows from (4.2), the third and fifth equalities follow from (4.1). The first inequality is due to the non-expansiveness of the projection operator, while the second inequality results from the convexity of the norm. The final inequality holds because, for every  $k \in [T+d]$ ,  $|\mathcal{F}_k| \leq d$ .

We now proceed to develop the last sum in 4.7. It follows from (4.2) and (4.1) that

$$\sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_t} \|r_k\|^2 = \sum_{t=1}^T \|\nabla f_t(x_t) + m_t\|^2 \le \sum_{t=1}^T \left( \|\nabla f_t(x_t)\|^2 + 2\|\nabla f_t(x_t)\|\delta_t + \delta_t^2 \right)$$

$$\le \sum_{t=1}^T \left( \|\nabla f_t(x_t)\|^2 + 2\|\nabla f_t(x_t)\|\delta_t \right) + \Lambda_T, \tag{4.8}$$

where the last inequality follows from the boundedness of  $m_t$ . On the other hand, for every  $t \in [T-1]$ , we have that:

$$-\|x_{t+1} - x_t^*\|^2 \le -\|x_{t+1} - x_{t+1}^*\|^2 + (2\|x_{t+1} - x_t^*\| + \|x_{t+1}^* - x_t^*\|)\|x_{t+1}^* - x_t^*\|$$

$$\le -\|x_{t+1} - x_{t+1}^*\|^2 + 6R\|x_{t+1}^* - x_t^*\|.$$
(4.9)

Plugging (4.8) and (4.9) in (4.7) yields:

$$\sum_{t=1}^{T} \langle r_{k} \mid x_{t'} - x_{t'}^{*} \rangle \leq \frac{1}{2\eta} \sum_{t=s}^{T-1} \left( \|x_{t} - x_{t}^{*}\|^{2} - \|x_{t+1} - x_{t}^{*}\|^{2} \right) + \frac{1}{2\eta} \left( \|x_{T} - x_{T}^{*}\|^{2} - \|x_{T+d} - x_{T}^{*}\|^{2} \right) \\
+ \frac{\eta d}{2} \sum_{t=1}^{T} \left( \|\nabla f_{t}(x_{t})\|^{2} + 2\|\nabla f_{t}(x_{t})\|\delta_{t} \right) + \frac{\eta d}{2} \Lambda_{T} \\
\leq \frac{1}{2\eta} \sum_{t=s}^{T-1} \left( \|x_{t} - x_{t}^{*}\|^{2} - \|x_{t+1} - x_{t+1}^{*}\|^{2} + 6R\|x_{t+1}^{*} - x_{t}^{*}\| \right) \\
+ \frac{1}{2\eta} \left( \|x_{T} - x_{T}^{*}\|^{2} - \|x_{T+d} - x_{T}^{*}\|^{2} \right) + \frac{\eta d}{2} \sum_{t=1}^{T} \left( \|\nabla f_{t}(x_{t})\|^{2} + 2\|\nabla f_{t}(x_{t})\|\delta_{t} \right) + \frac{\eta d}{2} \Lambda_{T} \\
\leq \frac{1}{2\eta} \|x_{s} - x_{s}^{*}\|^{2} + \frac{3R}{\eta} \sum_{t=s}^{T-1} \|x_{t+1}^{*} - x_{t}^{*}\| + \frac{\eta d}{2} \sum_{t=1}^{T} \left( \|\nabla f_{t}(x_{t})\|^{2} + 2\|\nabla f_{t}(x_{t})\|\delta_{t} \right) + \frac{\eta d}{2} \Lambda_{T} \\
\leq \frac{2R^{2}}{\eta} + \frac{3R}{\eta} V_{T} + \frac{\eta d}{2} \sum_{t=1}^{T} \left( \|\nabla f_{t}(x_{t})\|^{2} + 2\|\nabla f_{t}(x_{t})\|\delta_{t} \right) + \frac{\eta d}{2} \Lambda_{T}, \tag{4.10}$$

where the last inequality follows from assumption (B1) and from the definition of  $V_T$ .

(i) Case  $f_t$  is L-Lipschitz: We now proceed to upper bound the last term in the right-hand side of (4.6):

$$\sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_t} \|\nabla f_k(x_k)\| \|x_{\overline{t}}^* - x_k^*\| \le L \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_t} \|x_{\overline{t}}^* - x_k^*\| = L \sum_{t=1}^{T} \|x_{\overline{t'}}^* - x_t^*\| \\
\le L \sum_{t=1}^{T-1} \sum_{k=t+1}^{t+d_t-1} \|x_{\overline{k}}^* - x_{\overline{k-1}}^*\| = L \sum_{t=1}^{T-1} \sum_{k=t+1}^{\overline{t+d_t-1}} \|x_k^* - x_{k-1}^*\| \le L \sum_{t=1}^{T-1} \sum_{k=t+1}^{\overline{t+d-1}} \|x_k^* - x_{k-1}^*\| \\
\le L(d-1) \sum_{t=1}^{T-1} \|x_{t+1}^* - x_t^*\| = L(d-1)V_T, \tag{4.11}$$

where the first inequality follows from assumption (B3), the first equality is due to (4.2), and the second inequality comes from the triangle inequality. The second equality holds because  $\|x_k^* - x_{k-1}^*\| = 0$  when  $k \ge T+1$ , the third inequality is given by  $d = \max_{t \in [T]} d_t$ , while the fourth follows from (3.7).

Substituting (4.10) and (4.11) into (4.6) and rearranging terms, we obtain:

$$\kappa R_T^{NS} \leq \frac{2R^2}{\eta} + \frac{3R}{\eta} V_T + \frac{\eta d}{2} \sum_{t=1}^T \left( \|\nabla f_t(x_t)\|^2 + 2\|\nabla f_t(x_t)\| \delta_t \right) + \frac{\eta d}{2} \Lambda_T + L(d-1)V_T + 2R\Delta_T \\
+ \sum_{t=1}^T \|\nabla f_t(x_t)\| \|x_t - x_{t'}\| \\
\leq \frac{2R^2}{\eta} + \frac{3R}{\eta} V_T + \frac{\eta d}{2} \sum_{t=1}^T \left( L^2 + 2L\delta_t \right) + \frac{\eta d}{2} \Lambda_T + L(d-1)V_T + 2R\Delta_T + \sum_{t=1}^T L \|x_t - x_{t'}\| \\
\leq \frac{2R^2}{\eta} + \frac{3R}{\eta} V_T + \frac{\eta dTL^2}{2} + (\eta dL + 2R)\Delta_T + \frac{\eta d}{2} \Lambda_T + L(d-1)V_T + \sum_{t=1}^T L \|x_t - x_{t'}\| \tag{4.12}$$

Then, it remains to bound  $\sum_{t=1}^{T} L ||x_t - x_{t'}||$ . To do so, we use the triangle inequality to obtain:

$$||x_{t'} - x_t|| \leq \sum_{i=t}^{t'-1} ||x_{i+1} - x_i|| \leq \sum_{i=t}^{t'-1} ||x'_{i+1} - x_i|| = \sum_{i=\max(t,s)}^{t'-1} ||x'_{i+1} - x_i|| = \eta \sum_{i=\max(t,s)}^{t'-1} ||\sum_{k \in \mathcal{F}_i} r_k||$$

$$= \eta \sum_{i=\max(t,s)}^{t'-1} ||\sum_{k \in \mathcal{F}_i} (\nabla f_k(x_k) + m_k)|| \leq \eta \sum_{i=\max(t,s)}^{t'-1} ||\sum_{k \in \mathcal{F}_i} \nabla f_k(x_k)|| + \eta \sum_{i=\max(t,s)}^{t'-1} ||\sum_{k \in \mathcal{F}_i} m_k||$$

$$\leq \eta \sum_{i=\max(t,s)}^{t'-1} ||\mathcal{F}_i|L + \eta \sum_{i=\max(t,s)}^{t'-1} \sum_{k \in \mathcal{F}_i} ||m_k||$$

$$(4.13)$$

where the second inequality is due to the non-expansiveness of the projection, the second equality is due to (4.1). Summing (4.13) from 1 to T we have that

$$\sum_{t=1}^{T} \|x_{t'} - x_t\| \le \eta \sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t'-1} |\mathcal{F}_i| L + \eta \sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t'-1} \sum_{k \in \mathcal{F}_i} \delta_k$$

$$\le \eta \sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t'-1} |\mathcal{F}_i| L + \eta \sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t'-1} \sum_{k \in \mathcal{F}_i} \delta_k$$
(4.14)

To bound these terms, first note that  $s \le d$ . Indeed, if s > d, then for every  $k \in \mathcal{F}_s$ ,  $k + d_k - 1 = s > d$ , implying  $1 \notin \mathcal{F}_s$ . By the definition of s,  $d_1 = 1 + d_1 - 1 > s > d$ , contradicting the maximality of d. We then consider two cases:

(a) Case s > T: In this case, the algorithm does not update, and we have:

$$\sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t'-1} |\mathcal{F}_i| = \sum_{t=1}^{T} \sum_{i=s}^{t'-1} |\mathcal{F}_i| = \sum_{t=1}^{T} \sum_{i=s}^{t+d_t-2} |\mathcal{F}_i| \le \sum_{t=1}^{T} \sum_{i=s}^{T+d-2} |\mathcal{F}_i| \le (d-1) \sum_{t=s}^{T+d-1} |\mathcal{F}_i| \le (d-1)T,$$

where the first equality holds because  $s \ge T$ , ensuring  $\max(t,s) \ge t$  for all  $t \in [T]$ . The second equality and first inequality follow from  $t' = t + d_t - 1 \le T + d - 1$ , while the second inequality holds since  $d \ge s > T$ . Similarly,

$$\sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t'-1} \sum_{k \in \mathcal{F}_i} ||m_k|| = \sum_{t=1}^{T} \sum_{i=s}^{t'-1} \sum_{k \in \mathcal{F}_i} ||m_k|| = \sum_{t=1}^{T} \sum_{i=s}^{t+d_t-2} \sum_{k \in \mathcal{F}_i} ||m_k|| \le \sum_{t=1}^{T} \sum_{i=s}^{T+d-2} \sum_{k \in \mathcal{F}_i} \delta_k$$

$$\le (d-1) \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_i} \delta_k \le (d-1) \Delta_T,$$

(b) Case  $s \le T$ : In this case, we have:

$$\begin{split} \sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t'-1} |\mathcal{F}_i| &= \sum_{t=1}^{s-1} \sum_{i=s}^{t'-1} |\mathcal{F}_i| + \sum_{t=s}^{T} \sum_{i=t}^{t'-1} |\mathcal{F}_i| \leq \sum_{t=1}^{s-1} \sum_{i=s}^{T+d-2} |\mathcal{F}_i| + \sum_{t=s}^{T} \sum_{i=t}^{t+d-2} |\mathcal{F}_i| \\ &= \sum_{t=1}^{s-1} \sum_{i=s}^{T+d-2} |\mathcal{F}_i| + \sum_{t=s}^{T+d-2} \sum_{i=\max(s,t-d+2)}^{t} |\mathcal{F}_t| \\ &\leq (s-1) \sum_{i=s}^{T+d-2} |\mathcal{F}_i| + (d-1) \sum_{t=s}^{T+d-2} |\mathcal{F}_t| \leq 2(d-1) \sum_{t=s}^{T+d-2} |\mathcal{F}_t| \leq 2(d-1)T, \end{split}$$

where the first inequality follows from  $t' = t + d_t - 1 \le t + d - 1 \le T + d - 1$ , the second equality by reordering the sum, and the third inequality since  $d \ge s$ . Similarly,

$$\sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t'-1} \sum_{k \in \mathcal{F}_i} ||m_k|| = \sum_{t=1}^{s-1} \sum_{i=s}^{t'-1} \sum_{k \in \mathcal{F}_i} ||m_k|| + \sum_{t=s}^{T} \sum_{i=t}^{t'-1} \sum_{k \in \mathcal{F}_i} ||m_k||$$

$$\leq \sum_{t=1}^{s-1} \sum_{i=s}^{T+d-2} \sum_{k \in \mathcal{F}_i} \delta_k + \sum_{t=s}^{T} \sum_{i=t}^{t+d-2} \sum_{k \in \mathcal{F}_i} \delta_k$$

$$= \sum_{t=1}^{s-1} \sum_{i=s}^{T+d-2} \sum_{k \in \mathcal{F}_i} \delta_k + \sum_{t=s}^{T+d-2} \sum_{i=\max(s,t-d+2)} \sum_{k \in \mathcal{F}_t} \delta_k$$

$$\leq (s-1) \sum_{i=s}^{T+d-2} \sum_{k \in \mathcal{F}_i} \delta_k + (d-1) \sum_{t=s}^{T+d-2} |\mathcal{F}_t|$$

$$\leq 2(d-1) \sum_{t=s}^{T+d-2} \sum_{k \in \mathcal{F}_t} \delta_k \leq 2(d-1)\Delta_T.$$

Then, combining the last inequality with (4.12) and (4.14), we get:

$$R_T^{NS} \leq \frac{2R^2}{\eta\kappa} + \frac{3R + \eta L(d-1)}{\eta\kappa} V_T + \frac{L^2d + 4(d-1)L^2}{2\kappa} \eta T + \frac{\eta d}{2\kappa} \Lambda_T + \frac{\eta dL + 2R + 2\eta(d-1)L}{\kappa} \Delta_T.$$

and the proof of this bound is complete.

(ii) Case  $f_t$  is weakly smooth: We now proceed to upper bound the second term in (4.6):

$$\left(\sum_{t=1}^{T} \left\langle \nabla f_{t}(x_{t}) \mid x_{\overline{t'}}^{*} - x_{t}^{*} \right\rangle \right)^{2} \leq \left(\sum_{t=1}^{T} \|x_{\overline{t'}}^{*} - x_{t}^{*}\|^{2}\right) \left(\sum_{t=1}^{T} \|\nabla f_{t}(x_{t})\|^{2}\right) 
\leq 2R \left(\sum_{t=1}^{T} \|x_{\overline{t'}}^{*} - x_{t}^{*}\|\right) \left(\Gamma \sum_{t=1}^{T} (f_{t}(x_{t}) - f_{t}(x_{t}^{*}))\right) 
\leq 2R\Gamma \left(\sum_{t=1}^{T} \sum_{k=t+1}^{t+d_{t}-1} \|x_{\overline{k}}^{*} - x_{\overline{k}-1}^{*}\|\right) R_{T}^{NS} 
= 2R\Gamma \left(\sum_{t=1}^{T-1} \sum_{k=t+1}^{t+d_{t}-1} \|x_{k}^{*} - x_{k-1}^{*}\|\right) R_{T}^{NS} 
\leq 2R\Gamma \left(\sum_{t=1}^{T-1} \sum_{k=t+1}^{t+d_{t}-1} \|x_{k}^{*} - x_{k-1}^{*}\|\right) R_{T}^{NS} 
\leq 2R(d-1)\Gamma \left(\sum_{t=2}^{T} \|x_{k}^{*} - x_{k-1}^{*}\|\right) R_{T}^{NS} 
= 2R(d-1)\Gamma V_{T} R_{T}^{NS},$$
(4.15)

where the first inequality follows from applying Cauchy-Schwarz two times, the second from assumption (B4), and the third from the triangle inequality and the definition of  $R_T^{NS}$ . The first equality holds since  $\|x_k^* - x_{k-1}^*\| = 0$  for  $k \ge T+1$ . The fourth inequality holds as  $d = \max_{t \in [T]} d_t$  and the last from Lemma 3.8.

Now, we control the term  $\sum_{t=1}^{T} \|\nabla f_t(x_t)\| \delta_t$ . Applying the Cauchy-Schwarz inequality, we obtain:

$$\left(\sum_{t=1}^{T} \|\nabla f_t(x_t)\| \delta_t\right)^2 \le \left(\sum_{t=1}^{T} \|\nabla f_t(x_t)\|^2\right) \Lambda_T \le \left(\Gamma \sum_{t=1}^{T} \left(f_t(x_t) - f_t(x_t^*)\right)\right) \Lambda_T = \Gamma R_T^{NS} \Lambda_T. \tag{4.16}$$

Plugging (4.15), (4.16), and (4.10) into (4.6) we get:

$$R_{T}^{NS} \leq \frac{2R^{2}}{\eta \kappa} + \frac{3R}{\eta \kappa} V_{T} + \frac{\eta d}{2\kappa} \sum_{t=1}^{T} \|\nabla f_{t}(x_{t})\|^{2} + \frac{\eta d}{2\kappa} \Lambda_{T} + \frac{(2R(d-1)\Gamma)^{\frac{1}{2}}}{\kappa} \left(V_{T} R_{T}^{NS}\right)^{\frac{1}{2}} + \frac{\eta d}{\kappa} \left(\Gamma R_{T}^{NS} \Lambda_{T}\right)^{\frac{1}{2}} + \frac{2R}{\kappa} \Delta_{T} + \frac{1}{\kappa} \sum_{t=1}^{T} \|\nabla f_{t}(x_{t})\| \|x_{t} - x_{t'}\|,$$

$$(4.17)$$

where the second inequality follows from assumption (B4). To bound the regret, we must control the term  $\sum_{t=1}^{T} \|\nabla f_t(x_t)\| \|x_t - x_{t'}\|$ . Applying the Cauchy-Schwarz inequality, we obtain:

$$\left(\sum_{t=1}^{T} \|\nabla f_t(x_t)\| \|x_t - x_{t'}\|\right)^2 \le \left(\sum_{t=1}^{T} \|\nabla f_t(x_t)\|^2\right) \left(\sum_{t=1}^{T} \|x_t - x_{t'}\|^2\right) \le \Gamma R_T^{NS} \left(\sum_{t=1}^{T} \|x_t - x_{t'}\|^2\right). \tag{4.18}$$

Then, it remains to bound the second term on the right side of (4.18):

$$||x_{t'} - x_t||^2 \le (d - 1) \sum_{i=t}^{t'-1} ||x_{i+1} - x_i||^2 \le (d - 1) \sum_{i=t}^{t'-1} ||x'_{i+1} - x_i||^2 = (d - 1) \sum_{i=\max(t,s)}^{t'-1} ||x'_{i+1} - x_i||^2$$

$$= (d - 1)\eta^2 \sum_{i=\max(t,s)}^{t'-1} \left\| \sum_{k \in \mathcal{F}_i} r_k \right\|^2 \le 2(d - 1)\eta^2 \sum_{i=\max(t,s)}^{t'-1} \sum_{k \in \mathcal{F}_i} |\mathcal{F}_i| \left( ||\nabla f_k(x_k)||^2 + ||m_k||^2 \right)$$

$$\le 2d(d - 1)\eta^2 \sum_{i=\max(t,s)}^{t'-1} \sum_{k \in \mathcal{F}_i} \left( ||\nabla f_k(x_k)||^2 + \delta_k^2 \right), \tag{4.19}$$

where the first and third inequalities are due to the convexity of the norm. The first equality holds as  $||x_{k+1} - x_k|| = 0$  for  $k \in [s-1]$ , the second from (4.1), and the final inequality since  $|\mathcal{F}_k| \leq d$  for all  $k \in [t]$ .

Summing (4.19) from 1 to T we have that

$$\sum_{t=1}^{T} \|x_{t'} - x_{t}\|^{2} \leq 2d(d-1)\eta^{2} \sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t'-1} \sum_{k \in \mathcal{F}_{i}} \left( \|\nabla f_{k}(x_{k})\|^{2} + \delta_{k}^{2} \right) \\
= 2d(d-1)\eta^{2} \sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t+d_{t}-2} \sum_{k \in \mathcal{F}_{i}} \left( \|\nabla f_{k}(x_{k})\|^{2} + \delta_{k}^{2} \right) \\
\leq 2d(d-1)\eta^{2} \sum_{t=1}^{T} \sum_{i=\max(t,s)}^{t+d-2} \sum_{k \in \mathcal{F}_{i}} \left( \|\nabla f_{k}(x_{k})\|^{2} + \delta_{k}^{2} \right) \\
\leq 2d(d-1)^{2}\eta^{2} \sum_{t=s}^{T+d-1} \sum_{k \in \mathcal{F}_{t}} \left( \|\nabla f_{k}(x_{k})\|^{2} + \delta_{k}^{2} \right) \\
\leq 2d(d-1)^{2}\eta^{2} \sum_{t=1}^{T} \|\nabla f_{t}(x_{t})\|^{2} + 2d(d-1)^{2}\eta^{2}\Lambda_{T} \\
\leq 2\Gamma d(d-1)^{2}\eta^{2} R_{T}^{NS} + 2d(d-1)^{2}\eta^{2}\Lambda_{T} \tag{4.20}$$

where the second inequality follows from the definition of d, the third by Lemma 3.8, and the fourth by (4.2). The fifth equality is due to assumption (B4). Combining (4.18) and (4.20), we obtain:

$$\sum_{t=1}^{T} \|\nabla f_t(x_t)\| \|x_t - x_{t'}\| \le (2d)^{\frac{1}{2}} \Gamma(d-1) \eta R_T^{NS} + (2d\Gamma R_T^{NS})^{\frac{1}{2}} (d-1) \eta (\Lambda_T)^{\frac{1}{2}}.$$
 (4.21)

Substituting (4.21) into (4.17), we obtain

$$\begin{split} R_T^{NS} \leq & \frac{2R^2}{\eta \kappa} + \frac{3R}{\eta \kappa} V_T + \frac{2R}{\kappa} \Delta_T + \frac{\eta d}{2\kappa} \Lambda_T + \frac{\eta d\Gamma}{2\kappa} R_T^{NS} + \frac{(2d)^{\frac{1}{2}} \Gamma(d-1)\eta}{\kappa} R_T^{NS} \\ & + \frac{(2R(d-1)\Gamma)^{\frac{1}{2}}}{\kappa} \left( V_T R_T^{NS} \right)^{\frac{1}{2}} + \frac{(2d\Gamma)^{\frac{1}{2}} (d-1)\eta}{\kappa} \left( \Lambda_T R_T^{NS} \right)^{\frac{1}{2}} + \frac{\eta d}{\kappa} \left( \Gamma R_T^{NS} \Lambda_T \right)^{\frac{1}{2}}. \end{split}$$

Rearranging terms, we get that  $aR_T^{NS}-b\left(R_T^{NS}\right)^{\frac{1}{2}}\leq c$ . Since  $R_T^{NS}\geq 0$ , by solving the second-order inequality, we get,

$$R_T^{NS} \le \left(\frac{b + \sqrt{b^2 + 4ac}}{2a}\right)^2 \le \frac{b^2 + 2ac + b\left(b^2 + 4ac\right)^{\frac{1}{2}}}{2a^2},$$

which completes the proof.

As an immediate consequence of Theorem 4.2, we obtain the following corollary when  $m_k \equiv 0$ .

**Corollary 4.3.** Under assumptions (B1)-(B2) and considering that the full gradient is available, i.e., for all  $t \in [T]$ ,  $r_t = \nabla f_t(x_t)$ , the following assertions hold:

(i) If (B3) holds, the regret of Algorithm 1 with a constant step-size  $0 < \eta_t = \eta$ , can be upper-bounded by:

$$R_T^{NS} \le \frac{2R^2}{\eta \kappa} + \frac{3RV_T}{\eta \kappa} + \frac{\eta L^2 T d}{2\kappa} + \frac{L(d-1)V_T}{\kappa} + \frac{2\eta(d-1)L^2 T}{\kappa}$$

(ii) If (B4) holds, the regret of Algorithm 1 with a constant step-size  $0 < \eta_t = \eta < \frac{1}{\bar{\alpha}} = \frac{2\kappa}{\left(d+2d^{\frac{1}{2}}(d-1)\right)\Gamma}$ , can be upper-bounded by:

$$R_T^{NS} \leq \frac{\bar{b}^2 + 2\bar{a}\bar{c} + \bar{b}\left(\bar{b}^2 + 4\bar{a}\bar{c}\right)^{\frac{1}{2}}}{2\bar{a}^2},$$

where  $\bar{a}=1-\bar{\alpha}\eta>0$ ,  $\bar{b}=\frac{(2R(d-1)\Gamma V_T)^{\frac{1}{2}}}{\kappa}$ , and  $\bar{c}=\frac{2R^2+3RV_T}{\kappa}$ . Proof. If  $m_t=0$ , then  $\delta_t=0$ , leading to  $\Lambda_T=\Delta_T=0$ . Moreover, in (4.19), it is unnecessary to impose an upper bound of 2, which will impact the computation of  $\alpha$  and subsequently affect the value of  $\bar{a}$ .

Remark 4.4. In this remark, we only focus on the implications of Corollary 4.3 under the condition that the error is zero. In the next subsection, we will consider the case where the error is not necessarily zero.

- (i) If  $f_t$  is L-Lipschitz for every  $t \in [T]$ , then the regret bound (4.3) can be interpreted as follows: The term  $\frac{2R^2}{n\kappa}$  represents the initialization error, which decreases as the step size  $\eta$  increases. The second term,  $\frac{\eta \kappa}{\eta \kappa}$ , incorporates the cumulative path variation  $V_T$  and also diminishes with a larger  $\eta$ . In contrast, the terms  $\frac{\eta L^2 T d}{2\kappa}$  and  $\frac{2\eta (d-1)L^2 T}{\kappa}$  grow with  $\eta$ , suggesting a trade-off between these terms and the first two. The term  $\frac{L(d-1)V_T}{\kappa}$  highlights that, in the presence of delay (d>1), there is an influence of both the cumulative path variation and the delay, independent of the choice of the step size, which is different
- (ii) If  $V_T$  grows sub-linearly, i.e.,  $V_T=\mathcal{O}(T^\alpha)$  with  $\alpha<1$ , then selecting  $\eta=\mathcal{O}(T^{\frac{\alpha-1}{2}})$  ensures a sub-linear regret bound of  $\mathcal{O}(dT^{\frac{1+\alpha}{2}})$ , as stated in Theorem 4.2. Specifically, the optimal regret bound is obtained by setting

$$\eta = \left(\frac{2R(2R + 3V_T)}{TL^2(5d - 4)}\right)^{\frac{1}{2}}.$$

With this choice of  $\eta$ , Theorem 4.2 yields a regret bound of

$$\frac{\left(2RTL^{2}(5d-4)(2R+3V_{T})\right)^{\frac{1}{2}}}{\kappa} + \frac{L(d-1)V_{T}}{\kappa}.$$

However, this choice of  $\eta$  requires prior knowledge of  $V_T$ , which may not always be available.

(iii) If d = 1, meaning that there is not delay, then (4.4) becomes,

$$R_T^{NS} \le \frac{2R^2}{\eta \kappa} + \frac{3RV_T}{\eta \kappa} + \frac{\eta L^2 T d}{2\kappa},$$

which is the same regret obtained in [37, Theorem 1 (ii)]. When  $d \geq 1$  we extend the results for quasar-convex functions.

(iv) If  $f_t$  is  $\Gamma$ -weakly smooth for every  $t \in [T]$ , then the regret bound (4.3) can be viewed as the square of the sum of two terms arising from solving a second-order inequality. The first term, b, represents the square root of the cumulative path variation and is not null if there exists a delay (d > 1). The second term,  $(b^2 + 4ac)^{\frac{1}{2}}$ , includes both b and the initialization error along with  $V_T$ . Both terms are dominated by a factor of order  $V_T^{\frac{1}{2}}$ , making the overall regret grow as  $\mathcal{O}(dV_T)$ . Then, if  $V_T$  grows sub-linearly, the algorithm ensures a sub-linear regret bound. Since the feasible set is bounded  $V_T$  is at most linear. Then, similar to [37, Theorem 1], the order of the regret for  $\Gamma$ -weakly smooth function is better than the one of L-Lipschitz. Furthermore, the choice of the step size  $\eta$  is crucial: a smaller  $a=1-\alpha\eta$  can impact the regret bound significantly, as it appears in the denominator, thereby magnifying the overall regret. But, in this setting, the choice of  $\eta$  does not require prior knowledge of  $V_T$ .

(v) If d = 1, indicating no delay, then (4.4) simplifies to

$$R_T^{NS} \le \frac{c}{a} = \frac{2R^2 + 3RV_T}{(1 - \alpha n)\kappa},$$

which matches the regret bound given in [37, Theorem 1 (i)]. Furthermore, the step size  $\eta$  must satisfy the same condition, i.e.,  $\eta \in \left]0, \frac{2\kappa}{\Gamma}\right[$ .

- (vi) Note that for  $\kappa=1$  and under the assumption of differentiability, our results apply to star-convex functions with delay. To the best of our knowledge, this is the first work to establish a non-stationary regret bound for this class of functions in the presence of delays. A similar result holds for convex functions; however, in the case where d=1, we do not recover the bounds proposed by [4], which are of the order  $\mathcal{O}\left(V_T^{\frac{2}{3}}T^{\frac{1}{3}}\right)$ . In the delayed setting, we extend the work of [43] by providing non-stationary regret bounds for quasar-convex functions but assuming that  $f_t$  is  $\Gamma$ -weakly smooth.
- (vii) For both regret bounds, a decrease in the constant  $\kappa$  indicates that the functions become "more non-convex" resulting in larger regret bounds. This highlights the impact of the level of non-convexity on the performance of the algorithm.

## 4.2 Extension to the Bandit Setting with Delays

In this section, we study the zeroth-order version of Algorithm 1, which allows the method to address more complex scenarios where decision-making relies on partial gradient information and delayed feedback. To handle the bandit setting, following previous studies [1, 39, 42], we introduce two changes to Algorithm 1.

First, instead of querying the full gradient  $\nabla f_t(x_t)$ , we estimate it by querying the function  $f_t$  at (p+1) points:  $x_t$  and  $\{x_t+h_te_i\}_{i=1}^p$ , where  $\{e_i\}_{i=1}^p$  represents the canonical basis and  $h_t>0$  is the discretization parameter. The gradient approximation is given by:

$$r_t = \sum_{i=1}^p \frac{f_t(x_t + h_t e_i) - f_t(x_t)}{h_t} e_i,$$
(4.22)

and arrives at the end of round t'.

Second, we ensure the feasibility of these points, as  $x_t$  is feasible, but  $\{x_t + h_t e_i\}_{i=1}^p$  may not be. To maintain feasibility, we restrict the feasible set to a subset of the original decision set  $\mathcal{X}$ , defined as:

$$\mathcal{X}_h = \left\{ \left(1 - \frac{h}{R}\right) x \mid x \in \mathcal{X} \right\},\,$$

where  $h = \max_{t \in [T]} h_t \in ]0, R[$ . This ensures that for any  $x_t \in \mathcal{X}_h$  and for all  $i \in [p]$ , the point  $x_t + h_t e_i$  remains feasible.

**Theorem 4.5.** Let assumptions (B1) and (B2) hold, and assume that each loss function  $f_t: \mathcal{X} \to \mathbb{R}$  is G-smooth. Then, the regret of Algorithm 1, using a constant step-size  $0 < \eta_t = \eta < \frac{1}{\alpha_1} = \frac{\kappa}{\left(d + 4d^{\frac{1}{2}}(d-1)\right)G}$ , projecting over  $\mathcal{X}_h$ , and  $r_t$  as in (4.22), can be upper-bounded by:

$$R_T^{NS} \le \frac{b_1^2 + 2a_1c_1 + b_1 \left(b_1^2 + 4a_1c_1\right)^{\frac{1}{2}}}{2a_1^2},$$

where  $a_1 = 1 - \alpha_1 \eta > 0$ ,

$$b_1 = \frac{\left(2R(d-1)\Gamma V_T\right)^{\frac{1}{2}}}{\kappa} + \frac{\left(2d\right)^{\frac{1}{2}}(d-1)\eta + \eta d}{\kappa} \left(\Gamma\bar{\Lambda}_T\right)^{\frac{1}{2}} \quad \text{and} \quad c_1 = \frac{2R^2}{\kappa} + \frac{3RV_T}{\kappa} + \frac{\eta d}{2\kappa}\bar{\Lambda}_T + \frac{2R}{\kappa}\bar{\Delta}_T.$$

with  $\bar{\Lambda}_T=rac{pG^2}{4}\sum_{t=1}^T h_t^2$  and  $\bar{\Delta}_T=rac{p^{\frac{1}{2}}G}{2}\sum_{t=1}^T h_t.$ 

*Proof.* Since  $\nabla f_t$  is G-Lipschitz, it follows from [36, Subsection 1.1.2], that for every  $x \in \mathbb{R}^p$ ,  $t \in [T]$ , and  $i \in [n]$ 

$$|f_t(x_t + h_t e_i) - f_t(x_t) - h_t \langle \nabla f_t(x_t) \mid e_i \rangle| \le \frac{G h_t^2}{2},$$

which implies that

$$\left\| \sum_{i=1}^{p} \left[ \frac{f_t(x_t + h_t e_i) - f_t(x_t)}{h_t} e_i \right] - \nabla f_t(x_t) \right\| = \sqrt{\sum_{i=1}^{p} \left( \frac{f_t(x_t + h_t e_i) - f_t(x_t)}{h_t} - \langle \nabla f_t(x_t) \mid e_i \rangle \right)^2} \le \frac{p^{\frac{1}{2}} G h_t}{2}.$$

Then, we can set  $\delta_t \equiv \frac{p^{\frac{1}{2}}Gh_t}{2}$ , which gives that  $\bar{\Lambda}_T = \frac{pG^2}{4}\sum_{t=1}^T h_t^2$  and  $\bar{\Delta}_T = \frac{p^{\frac{1}{2}}G}{2}\sum_{t=1}^T h_t$ . On the other hand, from [37, Proposition 1] G-smooth implies 2G-weakly smooth. We conclude the proof applying Theorem 4.2 (ii).  $\Box$  **Remark 4.6.** In this remark, we explore the implications of our formulation.

- (i) Since *G*-smooth implies 2*G*-weakly smooth, we used Theorem 4.2 (ii), which requires assumption (B3) to hold, making Assumption (B1) unnecessary.
- (ii) By Theorem 4.5, for every  $x \in \mathbb{R}^p$  and sufficiently small  $h_t > 0$ , we have:

$$\sum_{i=1}^{p} \left[ \frac{f_t(x_t + h_t e_i) - f_t(x_t)}{h_t} e_i \right] - \nabla f(x) = \mathcal{O}(h_t^2).$$

Therefore, when  $h_t > 0$ , the gradient approximation is biased, so our method is more flexible than the stochastic methods that require unbiased estimators, such as [37].

(iii) Theorem 4.2 also applies to a 2p-point gradient estimator:

$$\sum_{i=1}^{p} \left[ \frac{f(x_t + h_t e_i) - f_t(x_t - h_t e_i)}{2h_t} e_i \right].$$

From [36, Subsection 1.1.2], for every  $x \in \mathbb{R}^p$  and  $i \in [n]$ :

$$|f(x + he_i) - f(x - he_i) - 2h\langle \nabla f(x) \mid e_i \rangle| \le Gh^2.$$

This implies:

$$\left\| \sum_{i=1}^{p} \left[ \frac{f(x+he_i) - f_t(x_t - he_i)}{2h} e_i \right] - \nabla f(x) \right\| \le \frac{p^{\frac{1}{2}}Gh}{2},$$

and we proceed analogously to Theorem 4.5. However, the main drawback is the increased number of function queries required.

(iv) From the definitions of  $a_1$ ,  $b_1$ , and  $c_1$ , we deduce that  $a_1$  is constant,  $b_1$  scales with the maximum of  $\bar{\Lambda}_T$  and  $V_T^{\frac{1}{2}}$ , and  $c_1$  scales with the maximum of  $\bar{\Delta}_t$ ,  $\bar{\Lambda}_T$ , and  $V_T$ . Consequently, the overall regret grows as  $\mathcal{O}(\max\{V_T,\bar{\Delta}_T,\bar{\Lambda}_T\})$ , making the choice of  $\{h_t\}_{t\in[T]}$  crucial. If  $V_T$  grows sub-linearly, i.e.,  $V_T=\mathcal{O}(T^\alpha)$  with  $\alpha<1$ , selecting  $h_t\equiv\mathcal{O}(T^{\alpha-1})$  ensures a sub-linear regret of  $\mathcal{O}(T^\alpha)=\mathcal{O}(V_T)$ . However, this choice of  $h_t$  requires prior knowledge of  $V_T$ , which may not always be available.

(v) Since the sum of quasar-convex functions is not necessarily quasar-convex, we cannot directly apply the techniques from [42, Theorem 2]. We extend the results presented in [43] for quasar-convex functions in the bandit setting. We also extend the results of [42, Theorem 2] to quasar-convex functions with dynamic regret, requiring weaker assumptions at the cost of losing the logarithmic regret rate. Observe that, we do not assume the existence of a constant r such that  $\mathbb{B}(0,r) \subset \mathcal{X}$ .

Now, we study the convergence of the delayed zeroth-order method for strongly quasar-convex functions in an offline setting, which will be applied to find the minimizer for applications in quadratic fractional functions.

Corollary 4.7. Let  $f: \mathcal{X} \to \mathbb{R}$  be  $(\kappa, \mu)$ -strongly quasar-convex with respect to  $x^* \in \operatorname{argmin}_{x \in \mathcal{X}} f(x)$ , and assume that f is G-smooth. Let  $\{r_t\}_{t \in \mathbb{N}}$  be defined as in (4.22) by the sequence  $\{h_t\}_{t \in \mathbb{N}}$ , where  $h_t$  is bounded by a constant h. Let the convex decision set  $\mathcal{X}$  be bounded with radius R. Consider the iterates  $\{x_t\}_{t \in \mathbb{N}}$  generated by Algorithm 1 using a constant step-size  $0 < \eta_t = \eta < \frac{1}{\alpha_1} = \frac{\kappa}{\left(d + 4d^{\frac{1}{2}}(d-1)\right)G}$ , where  $f_t \equiv f$ , with projection onto  $\mathcal{X}_h$ . Assume

further that  $\bar{\Delta}_t$  and  $\bar{\Lambda}_t$  are sub-linear terms. Then,  $X_t \to x^*$  as  $t \to +\infty$ , where  $X_T = \frac{1}{T} \sum_{k=1}^T x_t$  for all T > 0.

*Proof.* By the convexity of the norm, the  $(\kappa, \mu)$ -strong quasar-convexity of f with respect to  $x^*$ , and [37, Lemma 1 (ii)], for any T > 0, we obtain:

$$||X_T - x^*||^2 \le \frac{\sum_{t=1}^T ||x_t - x^*||^2}{T} \le \frac{8G\sum_{t=1}^T (f(x_t) - f(x^*))}{T\kappa^2 \mu^2} \le \frac{8GR_T}{T\kappa^2 \mu^2}.$$
 (4.23)

Since  $f_t \equiv f$  we obtain that  $V_T = 0$ . Moreover, since  $\bar{\Lambda}_T$  and  $\bar{\Delta}_T$  are sub-linear terms, the right-hand side of (4.23) tends to zero, thereby ensuring the convergence of the average iterate.

**Remark 4.8.** As far as we know, this is the first theorem that proved the convergence of the iterates of a delayed zeroth-order version of gradient descent for quasar-convex functions. However, the proof relies on making the regret tends to zero over time, which does not fully exploit the advantages of  $(\kappa, \mu)$ -strong quasar-convexity. For example, in [37, Proposition 2], it is shown that under similar assumptions, an exponential convergence rate can be achieved when the full gradient is available and there is no delay.

# 5 Examples and Applications

### 5.1 New Classes of Quasar-Convex Functions

In this section, we prove that strongly quasiconvex functions are (strongly) quasar-convex on convex and compact sets. To do this, we first present the following result:

**Proposition 5.1.** Let f be a differentiable function such that there exists  $\gamma, \kappa > 0$  such that

$$f(x) \le f(y) \implies \langle \nabla f(y) \mid x - y \rangle \le -\frac{\gamma}{2} ||x - y||^2, \tag{5.1}$$

$$f(x) \le f(y) \implies \langle \nabla f(y) \mid x - y \rangle \le \kappa (f(x) - f(y)). \tag{5.2}$$

Then

$$f(x) \le f(y) \implies f(x) \ge f(y) + \frac{2}{\kappa} \langle \nabla f(y) \mid x - y \rangle + \frac{\gamma}{2\kappa} ||x - y||^2.$$

*Proof.* It follows from (5.1) and (5.2) that

$$\kappa(f(y) - f(x)) + 2\langle \nabla f(y) \mid x - y \rangle + \frac{\gamma}{2} ||x - y||^2 \le 0, \ \forall \ x \in S_{f(y)}(f),$$

and the result follows.

In the next theorem, we prove that differentiable strongly quasiconvex functions are quasar-convex:

**Theorem 5.2.** Let  $K \subseteq \mathbb{R}^p$  be a convex and compact set,  $f: \mathbb{R}^p \to \mathbb{R}$  be a differentiable strongly quasiconvex with modulus  $\gamma > 0$  and  $\underset{K}{\operatorname{argmin}} f = \{x^*\}$ . Then there exists  $n_1 \in \mathbb{N}$  such that

$$\langle \nabla f(y) \mid x^* - y \rangle \leq \frac{4}{n} \left( f(x^*) - f(y) \right), \ \forall \ n \geq n_1.$$

As a consequence, f is quasar-convex with modulus  $\kappa = \frac{4}{n_1+4} \in ]0,1[$  and  $\gamma > 0$  on K. Proof. By Lemma 3.1, argmin  $f = \{x^*\}$ . Then, by equation (3.5), we have

$$f(x^*) + \frac{\gamma}{8} ||x^* - y||^2 \le f(y) \iff 0 < \frac{\gamma}{2} ||x^* - y||^2 \le 4(f(y) - f(x^*)), \ \forall \ y \in K.$$

Since K is compact and f is continuous, there exists M > 0 such that  $f(y) \leq M$  for all  $y \in K$ , thus,

$$0 < \frac{\gamma}{2} ||x^* - y||^2 \le 4(f(y) - f(x^*)) \le 4(M - f(x^*)), \ \forall \ y \in K.$$

By Archimedean axiom, there exists  $n_1 \in \mathbb{N}$  such that,

$$0 < \frac{4}{n}(M - f(x^*)) \le \frac{\gamma}{2} ||x^* - y||^2, \ \forall \ n \ge n_1.$$

Using now relation (5.1) (which holds in virtue of Lemma 3.2), we have

$$0 < \frac{4}{n}(M - f(x^*)) \le \frac{\gamma}{2} ||x^* - y||^2 \le \langle \nabla f(y) \mid y - x^* \rangle, \ \forall \ n \ge n_1.$$

Finally, since  $f(y) \leq M$  for all  $y \in K$ , we have for all  $n \geq n_1$  that

$$\langle \nabla f(y) \mid x^* - y \rangle \le \frac{4}{n} (f(x^*) - M) \le \frac{4}{n} (f(x^*) - f(y)), \ \forall \ y \in K,$$

i.e., f is quasar-convex with modulus  $\kappa = \frac{4}{n_1 + 4} \in \,]0,1[$  on K.

As a consequence, we have the desired result.

**Corollary 5.3.** Let  $K \subseteq \mathbb{R}^p$  be a convex and compact set,  $f: \mathbb{R}^p \to \mathbb{R}$  be a differentiable strongly quasiconvex with modulus  $\gamma > 0$  and argmin  $f = \{x^*\}$ . Then there exists  $n_1 \in \mathbb{N}$  such that f is strongly quasar-convex with modulus  $\kappa = \frac{2}{n_1 + 4} \in ]0, \frac{1}{2}[$  and  $\frac{\gamma}{\kappa} > 0$  on K.

*Proof.* Since f is a differentiable strongly quasiconvex function with modulus  $\gamma > 0$ , it satisfies relation (5.1) by Lemma 3.2, while in virtue of Theorem 5.2, there exists  $n_1 \in \mathbb{N}$  for which f satisfies relation (5.2) when  $x = x^*$  with  $\kappa = \frac{4}{n_1 + 4} \in ]0,1[$ . Therefore, it follows from Proposition 5.1 (applied to  $x = x^*$ ) that

$$f(x^*) \ge f(y) + \frac{2}{\kappa} \langle \nabla f(y) \mid x^* - y \rangle + \frac{\gamma}{2\kappa} ||x^* - y||^2, \ \forall \ y \in K,$$

i.e., f is strongly quasar-convex with modulus  $\frac{\gamma}{\kappa} > 0$  and  $\kappa = \frac{2}{n_1 + 4} \in ]0, \frac{1}{2}[$  on K.

- **Remark 5.4.** (i) The reverse statement in the previous results does not hold in general. Indeed, the function in Example 3.7 is strongly quasar-convex and it is not strongly quasiconvex (it is not even quasiconvex) because there are sublevel sets that are not convex, while the function  $f(x) = (x^2 + \frac{1}{8})^{1/6}$  is quasar-convex on  $\mathbb{R}$  with  $\kappa = 1/2$ , and it is not strongly quasiconvex as a consequence of [28, Theorem 1].
- (ii) On the whole space  $\mathbb{R}^p$ , a sufficient condition for differentiable strongly quasiconvex functions to be quasar-convex is that their gradient should be Lipschitz continuous (see [29, page 15]). The extension of Corollary 5.3 and/or Theorem 5.2 to the whole space remains as open problems.

As noted in Corollary 5.3, even when differentiable strongly quasiconvex functions are strongly quasar-convex, it could be hard to determine the parameters  $\kappa \in ]0,1]$  and  $\gamma>0$  of strong quasar-convexity. However, in the next proposition, we show how to compute these parameters for the case of quadratic fractional functions.

**Proposition 5.5.** Let  $A, B \in \mathbb{R}^{p \times p}$ ,  $a, b \in \mathbb{R}^p$ ,  $\alpha, \beta \in \mathbb{R}$ , and  $f : \mathbb{R}^p \to \mathbb{R}$  be the functions given by:

$$f(x) = \frac{g(x)}{q(x)} = \frac{\frac{1}{2}\langle Ax \mid x \rangle + \langle a \mid x \rangle + \alpha}{\frac{1}{2}\langle Bx \mid x \rangle + \langle b \mid x \rangle + \beta}.$$

Take 0 < m < M and define:

$$K := \{ x \in \mathbb{R}^p \mid m \le q(x) \le M \}.$$

*If A is a positive definite matrix and at least one of the following conditions holds:* 

- (a) B = 0 (the null matrix),
- (b) g is non-negative on K and B is negative semi-definite,
- (c) q is non-positive on K and B is positive semi-definite,

then the function f is  $\kappa$ -quasar-convex on K with  $\kappa = \frac{m}{M}$  and is  $(\frac{\kappa}{2}, \gamma)$ -strongly quasar-convex on K, where  $\gamma = \frac{\sigma_{\min}(A)}{m}$ . Moreover, for every  $\lambda \in (0,1)$ , f is  $(\lambda \kappa, (\lambda^{-1}-1)\frac{\sigma_{\min}(A)}{4M})$ -strongly quasar-convex on K. Proof. Since q(x) > 0 for all  $x \in \text{dom}(g) \cap K$ , we have  $\text{dom}(f) = \text{dom}(g) \cap K = K$ . By Remark 3.3, f is

strongly quasiconvex on K, ensuring a unique minimizer  $x^* \in \operatorname{argmin}_K f(x)$ . For every  $\lambda \in [0,1]$ , it holds that

$$f(\lambda x^* + (1 - \lambda)x) \le \frac{\lambda g(x^*) + (1 - \lambda)g(x)}{\lambda q(x^*) + (1 - \lambda)q(x)}$$
  
$$\iff f(\lambda x^* + (1 - \lambda)x) \le f(x) + \frac{\lambda q(x^*)}{\lambda q(x^*) + (1 - \lambda)q(x)} \left(f(x^*) - f(x)\right).$$

Using that  $q(x^*) \ge m$  and  $q(x) \le M$ , it follows that

$$f(\lambda x^* + (1 - \lambda)x) \le f(x) + \lambda \frac{m}{M} \left( f(x^*) - f(x) \right).$$

By Lemma 3.5, f is  $\kappa$ -quasar-convex on K with  $\kappa = \frac{m}{M}$ . Moreover, applying Lemma 3.2, we obtain

$$\langle \nabla f(x) \mid x^* - x \rangle < \kappa \left( f(x^*) - f(x) \right).$$

Furthermore, using Lemma 3.5,

$$\langle \nabla f(x) \mid x^* - x \rangle \le -\frac{\sigma_{\min}(A)}{2M} \|x - x^*\|^2.$$

Combining the above, we get

$$\frac{\sigma_{\min}(A)}{2M\kappa} \|x - x^*\|^2 + \frac{2}{\kappa} \langle \nabla f(x) \mid x^* - x \rangle \le \lambda \kappa \left( f(x^*) - f(x) \right)$$

which implies that f is  $(\frac{\kappa}{2}, \gamma)$ - strongly quasar-convex.

Finally, for fixed  $\lambda \in (0,1)$ , we have

$$\left\langle \nabla f(x) \mid x - x^* \right\rangle \geq \kappa \left( f(x) - f(x^*) \right) = \kappa \lambda \left( f(x) - f(x^*) \right) + \left( 1 - \lambda \right) \kappa \left( f(x) - f(x^*) \right),$$

where the last term satisfies

$$(1 - \lambda) \kappa \left( f(x) - f(x^*) \right) \ge \frac{(1 - \lambda) \kappa \sigma_{\min}(A)}{8M} \|x - x^*\|^2.$$

Thus, f is  $\left(\lambda\kappa, \frac{(1-\lambda)\kappa\sigma_{\min}(A)}{4M\lambda}\right)$ -strongly quasar-convex on K.

**Remark 5.6.** The parameter  $\lambda$  introduces a trade-off between  $\kappa$  and  $\gamma$ . By setting  $\lambda = 1$ , we obtain the largest constant for quasar-convexity, but losing strong quasar-convexity. In contrast, setting  $\lambda \leq \frac{m}{4M+m}$ , we achieve a larger constant for strong convexity, greater than  $\gamma$ , but this significantly reduces the constant for quasarconvexity. By selecting the constants as  $(\frac{\kappa}{2}, \gamma)$ , the quasar-convexity constant becomes larger than the previous choice. However, the strong quasar-convexity parameter cannot be arbitrarily chosen or adjusted in this case.

In the next example, we check that quadratic fractional functions satisfies the assumptions of our algorithm.

**Example 5.7.** Let  $A, B \in \mathbb{R}^{p \times p}$ ,  $a, b \in \mathbb{R}^p$ ,  $\alpha, \beta \in \mathbb{R}$ ,  $f : \mathbb{R}^p \to \mathbb{R}$ , 0 < m < M and K defined as in Proposition 5.5. Then the gradient is given by:

$$\nabla f(x) = \frac{q(x)(Ax+a) - g(x)(Bx+b)}{(q(x))^2}.$$

Additionally, if K is bounded by R, then f is L-smooth, where

$$L = \frac{\|A\|^2}{m} + 2\frac{(\|B\|R + \|b\|)(\|A\|R + \|a\|)}{m^2} + \frac{(\|A\|R^2 + 2\|a\|R + 2\alpha)(\|B\|R + \|b\|)^2}{m^3} + \frac{(\frac{1}{2}\|A\|R^2 + \|a\|R + \alpha)\|B\|^2}{m^2}.$$

Indeed, the Hessian f(x) is given by:

$$\nabla^2 f(x) = \frac{A^\top A}{q(x)} - \frac{(Bx+b)(Ax+a)^\top}{(q(x))^2} - \frac{(Ax+a)(Bx+b)^\top}{(q(x))^2} - g(x)\frac{B^\top B}{(q(x))^2} + 2g(x)\frac{(Bx+b)(Bx+b)^\top}{(q(x))^3}.$$

We now proceed to bound the Hessian, assuming that K is bounded. Since  $q(x) \ge m$ , we can bound the terms as follows:

$$\|\nabla^{2} f(x)\| \leq \frac{\|A\|^{2}}{m} + 2\frac{(\|B\|\|x\| + \|b\|)(\|A\|\|x\| + \|a\|)}{m^{2}} + 2\frac{(\frac{1}{2}\|A\|\|x\|^{2} + \|a\|\|x\| + \alpha)(\|B\|\|x\| + \|b\|)^{2}}{m^{3}} + \frac{(\frac{1}{2}\|A\|\|x\|^{2} + \|a\|\|x\| + \alpha)\|B\|^{2}}{m^{2}} \leq L.$$

Thus, f is L-smooth.

# 5.2 Other Non-Convex Examples

In this subsection, we check two non-convex examples of quasar-convex functions that satisfy the assumptions of our algorithm.

**Example 5.8.** We set  $\mathcal{X} = \{x \in \mathbb{R}^p \mid ||x|| \leq R\}$  and set the function  $f \colon \mathcal{X} \to \mathbb{R}$ . The function is defined as:

$$f(x) = g(\|x\|)q\left(\frac{x}{\|x\|}\right),\tag{5.3}$$

where  $g(t) = \frac{t^2}{1+t^2}$  and  $q(x) = \sum_{i=1}^p a_i \sin^2(b_i x_i)$  with  $\{a_i\}_{i=1}^p \subseteq [0, m_1]^p$  and  $\{b_i\}_{i=1}^p \subseteq [-m_2, m_2]^p$ . This function is  $\frac{2}{R^2+1}$ -quasar-convex on  $\mathcal{X}$  with respect to  $x^* = 0$  (see Proposition 3.5), and  $V_T = 0$ . The gradient of f is given by:

$$\nabla f(x) = \frac{x}{\|x\|} g'(\|x\|) q\left(\frac{x}{\|x\|}\right) + \frac{1}{\|x\|} \left( \text{Id} - \frac{xx^{\top}}{\|x\|^2} \right) g(\|x\|) \nabla q\left(\frac{x}{\|x\|}\right)$$

where, for every  $u \in \mathbb{R}^p$  and  $\alpha > 0$ ,  $\nabla q(u)$  and  $g'(\alpha)$  are given by:

$$\nabla q(u) = (a_i b_i \sin(2b_i u_i))_{i=1}^p \text{ and } g'(\alpha) = \frac{2\alpha}{(\alpha^2 + 1)^2}.$$

Hence, f is  $m_1p + m_1m_2p^{\frac{1}{2}}$  Lipchitz continuous. Indeed,

$$\|\nabla f(x)\| \le \left\| \frac{x}{\|x\|} \right\| \|g'(\|x\|)\| \left\| q\left(\frac{x}{\|x\|}\right) \right\| + \left\| \frac{g(\|x\|)}{\|x\|} \right\| \left\| \operatorname{Id} - \frac{xx^{\top}}{\|x\|^{2}} \right\| \left\| \nabla q\left(\frac{x}{\|x\|}\right) \right\|$$

$$\le \left\| q\left(\frac{x}{|x|}\right) \right\| + \left\| \nabla q\left(\frac{x}{|x|}\right) \right\| \le m_{1}p + m_{1}m_{2}p^{\frac{1}{2}}.$$

**Example 5.9.** In this example, we focus on GLM using the logistic function, which is an example of a quasarconvex and weakly-smooth function, and provides an instance where we can control the path variation  $V_T$ . We set the feasible set as  $\mathcal{X} = \{x \in \mathbb{R}^p \mid ||x|| \leq R\}$ . The function  $f \colon \mathcal{X} \to \mathbb{R}$  is defined as follows:

$$f(x) = \frac{1}{2m} \sum_{i=1}^{m} \left( \sigma(\langle a_i \mid x \rangle) - b_i \right)^2, \tag{5.4}$$

where, for every  $i \in [m]$ ,  $a^i \in \mathbb{R}^p$  and  $b^i = \sigma(\langle a^i_t \mid x^* \rangle)$ , with  $\sigma$  denoting the logistic function defined as follows  $\sigma(\alpha) = \frac{e^{\alpha}}{1+e^{\alpha}}$ . Note that

$$\sigma'(\alpha) = \frac{e^{\alpha}}{(1 + e^{\alpha})^2}$$
 and  $\sigma''(\alpha) = \frac{e^{\alpha}(1 - e^{\alpha})}{(1 + e^{\alpha})^3}$ ,

which implies that  $\sigma'$  is bounded by  $\frac{1}{4}$ . The gradient of f is given by:

$$\nabla f(x) = \frac{1}{m} \sum_{i=1}^{m} \sigma'(\langle a_i \mid x \rangle) \left( \sigma(\langle a_i \mid x \rangle) - b_i \right) a_i^{\top},$$

Hence f is  $\Gamma$ -weakly smooth with  $\Gamma = \frac{\max\limits_{i \in [m]} \|a_i\|^2}{8}$ . Indeed, since  $4\sigma' \leq 1$ , and  $f(x^*) = 0$ , we have, for every  $x \in \mathcal{X}$ , that

$$\|\nabla f(x)\|^{2} \leq \frac{1}{m} \sum_{i=1}^{m} \|\sigma'(\langle a_{i} \mid x \rangle) (\sigma(\langle a_{i} \mid x \rangle) - b_{i}) a_{i}^{\top}\|^{2} \leq \frac{1}{m} \sum_{i=1}^{m} \sigma'(\langle a_{i} \mid x \rangle)^{2} (\sigma(\langle a_{i} \mid x \rangle) - b_{i}))^{2} \|a_{i}^{\top}\|^{2}$$

$$\leq \frac{\max_{i \in [m]} \|a_{i}\|^{2}}{16m} \sum_{i=1}^{m} (\sigma(\langle a_{i} \mid x \rangle) - b_{i}))^{2} = \frac{\max_{i \in [m]} \|a_{i}\|^{2}}{8} (f(x) - f(x^{*})),$$

Since  $f(x^*) = 0$  and the positiveness of f, for  $\kappa = \min(1, 8\sigma'(R))$ , we have that

$$f(x^*) \ge \left(1 - \frac{8\sigma'(R)}{\kappa}\right) f(x)$$

$$\ge -\frac{2\sigma'(R)}{\kappa} \frac{1}{2m} \sum_{i=1}^{m} \left(\sigma(\langle a_i \mid x \rangle) - \sigma(\langle a_i \mid x^* \rangle)\right) (\langle a_i \mid x \rangle - \langle a_i \mid x^* \rangle) + f(x)$$

$$\ge -\frac{1}{\kappa} \frac{1}{2m} \sum_{i=1}^{m} 2\sigma'(\langle a_i \mid x \rangle)\right) (\sigma(\langle a_i \mid x \rangle) - \sigma(\langle a_i \mid x^* \rangle)) (\langle a_i \mid x \rangle - \langle a_i \mid x^* \rangle) + f(x)$$

$$\ge \frac{1}{\kappa} \langle \nabla f(x) \mid x^* - x \rangle + f(x),$$

which implies that f is  $\kappa$ -quasar-convex.

# 6 Numerical Experiments

In this section, we present numerical experiments to evaluate the performance of Delayed Online Gradient Descent on quasar-convex functions. All numerical examples are implemented in Python on a laptop equipped with an AMD Ryzen 5 3550Hz processor, Radeon Vega Mobile Gfx, and 32 GB of RAM. The corresponding code can be downloaded at <a href="https://github.com/cristianvega1995/Delayed-Feedback-in-Online-Quasar-Convex-Optimization-A-Non-Stationary-Approach">https://github.com/cristianvega1995/Delayed-Feedback-in-Online-Quasar-Convex-Optimization-A-Non-Stationary-Approach</a>.

We conduct experiments in three different settings: highly non-convex functions, generalized linear models, and quadratic fractional functions. In all experiments, unless otherwise specified, we set T=20000 and run the algorithm for  $d \in \{1, 5, 10, 20\}$ . At each iteration, we set  $r_t = \nabla f_t(x_t)$  and the delay  $d_t$  is drawn from a

discrete uniform distribution over the range [d]. Each experiment is repeated 20 times with random realizations of the sequence  $\{f_t\}_{t\in[T]}$ , and we report both the average performance and its standard deviation.

For the first experiment, we consider the Lipschitz continuous function presented in (5.3). We set R=100, p=100. For every  $t\in [T]$ ,  $\{a_i^t\}_{i=1}^p$  are independently and uniformly distributed on  $[0,1]^p$ , and  $\{b_i^T\}_{i=1}^p$  are independently and uniformly distributed on  $[-2.5,2.5]^p$ . Since  $x_t^*=0$ , we have  $f_t(x_t^*)=0$  for every  $t\in [T]$ , and  $V_T=0$ . The initialization is given such that  $x_0\sim U(0.2,0.4)^p$ . The step-size is given by  $\eta=\frac{2R}{L}\left(T(5d-4)\right)^{-\frac{1}{2}}$ , which according to Remark 4.4(ii) is the optimal step-size for minimize the upper bound of the regret.




Figure 2: Average regret of Algorithm 1 on the Lipschitz continuous function defined in (5.3) withh  $d \in \{1, 5, 10, 20\}$ .

Algorithm	Iter	std
d = 1	1476	67.18
d = 5	6707	315.02
d = 10	10015	540.10
d=20	14469	749.89

Table 3: First iteration when the error is less than 0.1 (iter), along with the standard deviation of the final iterate (std), as shown in Figure 2.

Figure 2 illustrates the performance of delayed online gradient descent on highly non-convex functions. As observed, the regret curves grow sub-linearly with the number of iterations and stabilize and flat after a certain number of iterations. Additionally, the regret increases with the delay, meaning that larger delays result in higher regret and require more iterations to stabilize, as shown in Table 3. These results support our

Algorithm	Iter	std
d = 20	45694	2375.71
d = 50	71466	3746.63
d = 100	103445	6024.52
d = 150	127445	5842.88
d = 200	142963	5784.98

Table 4: First iteration when the error is less than 0.1 (iter), along with the standard deviation of the final iterate (std), as shown in Figure 3.




Figure 3: Average regret of Algorithm 1 on the Lipschitz continuous function defined in (5.3) in a high-delay setting with  $d \in \{20, 50, 100, 150, 200\}$ .

theoretical predictions that larger delays lead to greater regret. A similar trend is observed for the standard deviation, which also increases as the delay grows.

We also conduct experiments in a high-delay setting, where  $d \in \{20, 50, 100, 150, 200\}$  and T = 200000, while keeping the other parameters as in the previous experiment. As seen in Figure 3 and Table 4, the regret curves grow sub-linearly with the number of iterations and eventually stabilize. The delay causes the regret to increase, and the number of iterations needed to stabilize the curve also grows, supporting our earlier findings. However, in this case the standard deviation tend to stabilize around 6000 and is not increases with delay. Note that although the same class of functions and parameters are used, this experiment is not directly comparable to the previous one because the step-size decreases as T increases. For instance, when d = 20, the regret curve stabilizes at iteration 45.203, compared to 14.469 in the previous experiment, due to the smaller step-size (reduced by a factor of  $\sqrt{10}$ ), highlighting the importance of the step-size choice.

For the second experiment, we consider the function presented in (5.4), which is weakly smooth. We set R=1 and p=100. The sequence of optimal solutions is initialized as  $x_1^* \sim \mathcal{N}(0, \operatorname{Id}_p)$ , and evolves over time steps  $t \in [T-1]$  according to the update rule:

$$x_{t+1}^* = x_t^* + \frac{0.1}{t^{\frac{1}{2}}} v_t,$$

where  $v_1,\ldots,v_{T-1}\sim\mathcal{N}(0,\operatorname{Id}_p)$  are independent and identically distributed standard Gaussian vectors. This implies that  $f_t(x_t^*)=0$  for every  $t\in[T]$ , and that  $V_T=\mathcal{O}(T^{\frac{1}{2}})$ . The initial iterate is sampled as  $x_0\sim\mathcal{N}(0,\operatorname{Id}_p)$  and subsequently normalized. At each round t, we collect m=1000 i.i.d. samples  $\{(a_t^i,b_t^i)\}_{i=1}^m$ , where the inputs  $a_t^i\sim\mathcal{N}(0,\operatorname{Id}_p)$ , and the outputs are defined as  $b_t^i=\sigma(\langle a_t^i\mid x_t^*\rangle)$ . The step-size is set as

$$\eta_t = 0.99 \frac{2\kappa}{(d+2d^{\frac{1}{2}}(d-1))\Gamma}.$$
(6.1)

Figure 4 illustrates the performance of delayed online gradient descent on GLM. As observed in the previous experiment, the regret and the standard deviation increase with the delay, and all the regret curves grow sublinearly. Note that in this example, when there is no delay, the cumulative regret is significantly lower than the other regret curves and stabilizes much faster. This highlights the substantial impact that delay has on cumulative regret, as larger delays result in both higher regret and slower stabilization.

To explore the impact of path variation  $V_T$ , we keep d,  $\Gamma$ ,  $\kappa$ ,  $\theta$ , and R constant while varying  $V_T$ . Consequently, the upper bound on the regret will depend only on  $V_T$ . To achieve this, we modify the equation for  $x_t^*$  by




Figure 4: Average regret of Algorithm 1 on the weakly smooth function defined in (5.4).

Algorithm	Iter	std
d = 1	78	0.029
d = 5	1862	0.053
d = 10	5328	0.127
d = 20	15728	0.366

Table 5: First iteration when the error is less than  $10^{-4}$  (iter), along with the standard deviation of the final iterate (std), as shown in Figure 4.

introducing a parameter a as follows:

$$x_{t+1}^* = x_t^* + \frac{0.1}{t^2} v_t,$$

where  $\mathbf{a} \in \{0.0625, 0.125, 0.25, 0.5, 1\}$ , and  $\{v_t\}_{t \in [T]}$  is sampled as in the previous experiment. This implies that  $V_T = \mathcal{O}(T^{1-\mathbf{a}})$  for every  $t \in [T]$ . The initial iterate  $x_0$  is sampled similarly, and for each time step  $t \in [T]$ ,  $\eta_t$ ,  $b_t$ , and  $A_t$  are sampled as in the previous example. Even though we are comparing different functions, normalizing,  $A_t$  and  $b_t$  we ensure that all parameters except  $V_T$  remain the same, allowing us to study the effect of  $V_T$  on the upper bound of the regret.

As seen in Figure 5, the regret decreases as the value a increases. All regret curves, except for a=0.0625, exhibit sub-linear growth, indicating that the curves may require more iterations to flatten, rather than suggesting linear growth. In Table 6, we observe that as a increases, the error decreases more quickly. However, the standard deviation remains consistent across different values of a. These results reinforce our findings regarding the impact of cumulative path variation on regret.

For the third experiment, we consider the function defined in (3.6). We set  $B_t \equiv 0$ , R = 10, and p = 50. The

Algorithm	Iter	std
a = 0.0625	2939	0.084
a = 0.125	2135	0.070
a = 0.25	2021	0.064
a = 0.5	1945	0.068
a = 1	1867	0.073

Table 6: First iteration when the error is less than  $10^{-4}$  (iter), along with the standard deviation of the final iterate (std), as shown in Figure 5.




Figure 5: Average regret of Algorithm 1 on the weakly smooth function defined in (5.4) for differents  $V_T$ .

parameters of the function  $f_t$  are initialized as follows:  $A_0$  is a positive definite matrix with norm 1, and  $a_0$  and  $b_0$  are independently sampled from a standard normal distribution, then normalized to have a norm of 0.1. For each  $t \in [T]$ , the parameters update according to the following recursions:

$$A_t = \frac{A_{t-1} + 0.01t^{-\frac{1}{2}}v_t^1}{\|A_{t-1} + 0.01t^{-\frac{1}{2}}v_t^1\|}, \quad a_t = \frac{a_{t-1} + 0.01t^{-\frac{1}{2}}v_t^2}{10\|a_{t-1} + 0.01t^{-\frac{1}{2}}v_t^2\|}, \quad \text{and} \quad b_t = \frac{b_{t-1} + 0.01t^{-\frac{1}{2}}v_t^3}{10\|b_{t-1} + 0.01t^{-\frac{1}{2}}v_t^3\|},$$

where  $v_t^1$  is a positive definite matrix and  $v_t^2$ ,  $v_t^3$  are independent, identically distributed standard Gaussian vectors. We also set  $\beta_t \equiv \|b_t\|R + 100$  and  $\alpha_t = 10$ , which ensures that for all  $x \in \mathcal{X}$ ,  $x \in K$  with m = 100 and  $M = \|b_t\|R + \beta$ . We set the step-size  $\eta$  as in (6.1). Since the sequence of optimal solutions is not available a priori (and consequently  $V_T$ ), we implement a gradient descent subroutine at each iteration  $t \in [T]$ , which, as guaranteed by Corollary 4.7, converges to the unique solution. The stopping criterion for this subroutine is based on the relative error, with a tolerance level of  $10^{-6}$ . To improve execution time, we use a warm-start strategy, initializing the subroutine at iteration t with the solution from iteration t = 1.




Figure 6: Average regret of Algorithm 1 on the weakly smooth function defined in (3.6) for differents  $V_T$ .

Figure 6 illustrates the performance of delayed online gradient descent on quadratic fractional functions. As observed in all the previous examples, the regret curves grow sub-linearly with the number of iterations and stabilize and flat after a certain number of iterations. Observe that the regret increases with the delay. In this

Algorithm	Iter	std
d = 1	162	$3.1 \cdot 10^{-4}$
d = 5	4126	$5.9 \cdot 10^{-3}$
d = 10	12069	$1.2 \cdot 10^{-2}$
d=20	19985	$1.7 \cdot 10^{-2}$

Table 7: First iteration when the error is less than  $10^{-5}$  (iter), along with the standard deviation of the final iterate (std), as shown in Figure 6.

Algorithm	Iter	std	Time [s]
Full gradient	5394	$7.6 \cdot 10^{-3}$	1.0
a = 1	5394	$7.8 \cdot 10^{-3}$	19.9
a = 0.8	5495	$8.5 \cdot 10^{-3}$	19.7
a = 0.6	6853	$1.5 \cdot 10^{-2}$	19.5
a = 0.4	-	$5.4 \cdot 10^{-2}$	19.6

Table 8: First iteration where the error falls below  $10^{-5}$  (iter) and the standard deviation of the final iterate (std), as illustrated in Figure 7. The average execution time for each experiment is also included.

example, the regret increases with the delay. In Table 7 shows that larger delays result in more iterations to stabilize the curve. The same results are obtained for the standard deviation.

To investigate the impact of error when the full gradient is unavailable, and consequently how  $\Lambda_T$  and  $\Delta_T$  impact in the regret, we set the parameters as in the previous experiment, but fixing d=5 since the curve stabilizes faster than d=10, which will allow a wider range to select the parameter a. However, in this case, we define  $r_k$  as in (4.22) with  $h_t=\frac{1}{t^a}$ , where  $a\in\{0.4,0.6,0.8,1\}$ . As a result, the upper bound on the regret depends only on  $\Lambda_T$  and  $\Delta_T$ , allowing us to analyze their effects on the regret's upper bound. We also compare the results with the case when the full gradient is available ( $h_t\equiv 0$ ). This leads to  $\Delta_T\geq \mathcal{O}(T^{1-a})$  for  $a\in\{0.4,0.6,0.8\}$ , and  $\Delta_T\geq \mathcal{O}(\ln(T))$  when a=1. As illustrated in Figure 7, the regret decreases




Figure 7: Average regret of Algorithm 1 with d=5 on the weakly smooth function defined in (3.6) for differents  $\Delta_T$  and  $\Lambda_T$ .

as a increases. For a=1 and a=0.8, the regret curve closely follows the full gradient curve, indicating a good approximation. A similar behaviour is seen when a=0.6, though the gradient approximation is less accurate compared to the previous cases. When a=0.4, the regret increases, and the error does not drop below  $10^{-5}$  within 20.000 iterations. Table 8 shows that as a increases, fewer iterations are required to reach the given tolerance level. A similar trend is observed with the standard deviation. Furthermore, the standard deviation for the full gradient and for a=1 are comparable, though the gradient computation in the

bandit setting incurs significantly higher execution time. This raises the question of exploring cheaper gradient approximations, such as the two-point estimator proposed in [1, 42].

## 7 Conclusions and Future Work

In this work, we introduced a projected gradient descent algorithm to address online quasar-convex optimization in a delayed feedback setting for both weakly smooth and Lipschitz gradient functions. We established a sub-linear dynamic regret rate for the proposed algorithm, which depends on the cumulative path variation, the delay, the radius of the set, and the time horizon. Furthermore, we extend our algorithm to the (zerothorder) bandit setting and the convergence of the delayed zeroth-order projected gradient descent method for strongly quasar-convex functions. We also provide new examples of quasar-convex functions that improve the applicability of online optimization to quasar-convex functions, such as strongly quasiconvex functions. Experimental results validated our theoretical findings, highlighting the influence of path variation and delay on algorithmic behavior.

Future research will extend this work for more general settings such as mirror descent, inertial schemes, second-order methods, and non-smooth functions. Another research direction is to investigate stochastic error to include more complex zeroth-order schemes, especially a two point stimator as in [10, 42]. We also want to study how to weaken conditions to consider unbounded feasible sets and thus consider linear constraints.

## 8 Declarations

## 8.1 Competing Interests

There are no conflicts of interest or competing interests related to this manuscript.

# 8.2 Funding

This research was partially supported by ANID—Chile through Fondecyt Regular 1241040 (Lara)

## References

- [1] A. AGARWAL, O. DEKEL, L. XIAO, Optimal algorithms for online convex optimization with multi-point bandit feedback. In: Colt, pp. 28–40. Citeseer (2010).
- [2] A. AGARWAL, E. HAZAN, S. KALE, R. SHAPIRE, Algorithms for portfolio management based on the newton method. In Proc. 23rd ICML, (2006).
- [3] K. Arrow, A. Enthoven, Quasiconcave programming. Econometrica 29, 779–800, (1961).
- [4] O. Besbes, Y. Gur, A. Zeevi, Non-stationary stochastic optimization. Oper. Res. **63**(5), 1227–1244, (2015).
- [5] A. Blum, On-line algorithms in machine learning. Online algorithms: the state of the art pp. 306–325, (2005).
- [6] A. Blum, A. Kalai, Universal portfolios with and without transaction costs. Mach. Learn. **35**, 193–205, (1999).
- [7] A. CABOT, H. ENGLER, S. GADAT, On the long time behavior of second order differential equations with asymptotically small dissipation. Trans. Amer. Math. Soc. **361**(0), 5983–6017, (2009).
- [8] A. CAMBINI, L. MARTEIN, Generalized convexity and optimization: Theory and applications. Springer, (2009).
- [9] X. CHEN, Y. LIU, G. HONG, An online learning approach to dynamic pricing and capacity sizing in service systems. Oper. Res., (2023).

- [10] A.D. FLAXMAN, A.T. KALAI, H.B. MCMAHAN, Online convex optimization in the bandit setting: gradient descent without a gradient. arXiv preprint cs/0408007, (2004).
- [11] X. GAO, X. LI, S. ZHANG, Online learning with non-convex losses and non-stationary regret. In: Inter. Conf. Artif. Intel. Stati., pp. 235–243. PMLR, (2018).
- [12] S.-M. GRAD, F. LARA, R.T. MARCAVILLACA, Strongly quasiconvex functions: what we know (so far). arXiv:2410.23055, (2024).
- [13] S. GUMINOV, A. GASNIKOV, I. KURUZOV, Accelerated methods for weakly quasi convex optimization problems. Comp. Manag. Sci. 20, 36, (2023).
- [14] N. HADJISAVVAS, S. KOMLOSI, S. SCHAIBLE, Handbook of Generalized Convexity and Generalized Monotonicity. Springer, (2005).
- [15] M. HARDT, T. MA, B. RECHT, Gradient descent learns linear dynamical systems. JMLR 19, 1–44, (2018).
- [16] E. HAZAN, A. AGARWAL, S. KALE, Logarithmic regret algorithms for online convex optimization. Mach. Learn. **69**(2), 169–192, (2007).
- [17] E. HAZAN ET AL.: Introduction to online convex optimization. Found. Trends Optim. **2**(3-4), 157–325, (2016).
- [18] X. HE ET AL.: Practical lessons from predicting clicks on ads at facebook. In: Proc. Eighth Inter. AdKDD, pp. 1–9, (2014).
- [19] D.P. HELMBOLD, R.E. SCHAPIRE, Predicting nearly as well as the best pruning of a decision tree. In: Proc. Eighth Annual Conf. Comp. Learning Theory, pp. 61–68, (1995).
- [20] J. HERMANT, J.F. AUJOL, C. DOSSAL, A. RONDEPIERRE, Study of the behaviour of Nesterov accelerated gradient in a non convex setting: the strongly quasar convex case. arXiv:2405.19809, (2024).
- [21] O. HINDER, A. SIDFORD, N. SOHONI, Near-optimal methods for minimizing star-convex functions and beyond. In: COLT, pp. 1894–1938. PMLR (2020).
- [22] A. IUSEM, F. LARA, R.T. MARCAVILLACA, L.H. YEN, A two-step proximal point algorithm for nonconvex equilibrium problems with applications to fractional programming. J. Glob. Optim. **90**, 755–779, (2024).
- [23] P. JOULANI, A. GYORGY, C. SZEPESVÁRI, Online learning under delayed feedback. In: ICML, pp. 1453–1461. PMLR (2013).
- [24] M. JOVANOVIC, A note on strongly convex and quasiconvex functions. Math. Notes, 60, 584–585, (1996).
- [25] A. KABGANI, F. LARA, Strong subdifferentials: theory and applications in nonconvex optimization. J. Glob. Optim. 84, 349–368, (2022).
- [26] A. KALAI, S. VEMPALA, Efficient algorithms for online decision problems. J. Comput. Syst. Sci. **71**(3), 291–307, (2005).
- [27] S. KIM, L. MADDEN, E. DALL'ANESE, Online stochastic gradient methods under sub-weibull noise and the polyak-Łojasiewicz condition. In: 2022 IEEE 61st Conf. Decis. Control, pp. 3499–3506. (2022).
- [28] F. LARA, On strongly quasiconvex functions: existence results and proximal point algorithms. J. Optim. Theory Appl. **192**, 891–911, (2022).
- [29] F. LARA, R.T. MARCAVILLACA, P.T. VUONG, Characterizations, dynamical systems and gradient methods for strongly quasiconvex functions, arXiv:2410.03534, (2024).
- [30] B. Li, T. Chen, G.B. Giannakis, Bandit online learning with unknown delays. In: The 22nd Inter. Conf. Artif. Intel. Stat., pp. 993–1002. PMLR, (2019).
- [31] A. MOKHTARI, S. SHAHRAMPOUR, A. JADBABAIE, A. RIBEIRO, Online optimization in dynamic environments: Improved regret rates for strongly convex problems. Proc. IEEE Conf. Decis. Control pp. 7195–7201, (2016).
- [32] N.M. NAM, J. SHARKANSKY, On strong quasiconvexity of functions in infinite dimensions, arXiv:2409.17450, (2024).
- [33] I. NECOARA, Y. NESTEROV, F. GLINEUR, Linear convergence of first order methods for non-strongly convex optimization. Math. Programm. **175**, 69–107, (2019).
- [34] Y. NESTEROV, B.T. POLYAK, Cubic regularization of newton method and its global performance. Math. Programm. **108**(1), 177–205, (2006).
- [35] B.T. Polyak, Existence theorems and convergence of minimizing sequences in extremum problems with restrictions. Soviet Math. 7, 72–75, (1966).

- [36] B.T. POLYAK, Introduction to optimization. Optimization Softwares. New York. (1987).
- [37] Y.M. Pun, I. Shames, Online non-stationary stochastic quasar-convex optimization. arXiv:2407.03601, (2024).
- [38] K. QUANRUD, D. KHASHABI, Online learning with adversarial delays. Adv. Neural Inf. Process. Syst. 28, (2015).
- [39] A. SAHA, A. TEWARI, Improved regret guarantees for online smooth convex optimization with bandit feedback. In: Proc. 14th Inter. Conf. Art. Intel. Stat., pp. 636–642. JMLR W&CP., (2011).
- [40] S. SHALEV-SHWARTZ ET AL., Online learning and online convex optimization. Found. Trends Mach. Learn. 4(2), 107–194, (2012).
- [41] A. VLADIMIROV, J. NESTEROV, J. CHEKANOV, O ravnomerno kvazivypuklyh funkcionalah [on uniformly quasiconvex functionals]. Vestn. Mosk. un-ta, vycis. mat. i kibern., 4, 18–27, (1978).
- [42] Y. WAN, W.W. Tu, L. ZHANG, Online strongly convex optimization with unknown delays. Mach. Learn., 111(3), 871–893, (2022).
- [43] Y. WAN, C. YAO, M. SONG, L. ZHANG, Non-stationary online convex optimization with arbitrary delays. arXiv:2305.12131, (2023).
- [44] J. Wang, M. Dong, B. Liang, G. Boudreau, H. Abou-Zeid, Delay-tolerant oco with long-term constraints: Algorithm and its application to network resource allocation. IEEE/ACM Trans. Netw., **31**(1), 147–163, (2022).
- [45] J. Wang, B. Liang, M. Dong, G. Boudreau, H. Abou-Zeid, Delay-tolerant constrained oco with application to network resource allocation. In: IEEE INFOCOM 2021-IEEE Conf. Comp. Comm., pp. 1–10. (2021).
- [46] L. Zhang, T. Yang, R. Jin, Z.H. Zhou, Online bandit learning for a special class of non-convex losses. In: Proc. AAAI Conf. Artif. Intel., 29, (2015).
- [47] L. Zhang, T. Yang, J. Yi, R. Jin, Z.H. Zhou, Improved dynamic regret for non-degenerate functions. Adv. Neural Inf. Process. Syst., **30**, (2017).
- [48] M. ZINKEVICH, Online convex programming and generalized infinitesimal gradient ascent. In: Proc. 20th ICML, (ICML-03), pp. 928—936, (2003).