

SEVERAL NEW WITTEN RIGIDITY THEOREM FOR ELLIPEIC GENUS

JIANYUN GUAN; KEFENG LIU; AND YONG WANG*

ABSTRACT. Using the Liu's method, we prove a new Witten rigidity theorem of elliptic genus of twisted Dirac operators in even dimensional spin manifolds under the circle action. Combined with the Han-Yu's method, we prove the Witten rigidity theorems of elliptic genus of twisted Toplitz operators of odd-dimensional spin manifolds under the circle action. Moreover, we have obtained several similar Witten rigidity theorems of elliptic genus.

1. INTRODUCTION

The study of the rigidity of the signature operator, the Dolbeault operator and the Dirac operator on different manifolds is an interesting topic in geometry and mathematical physics. Around 1982, motivated by physics, Witten proved the rigidity of the twisted Dirac operator $D \otimes TX$ for compact homogeneous spin manifolds. In 1988, Witten derived a series of twisted Dirac operators on the free loop space LM of a spin manifold M [3]. In this work, Witten was surprised to discover that the elliptic genus, constructed topologically by Landweber-Stone[5] and Ochanine[4], is the index of one of the operators. Inspired by physics, Witten conjectured that these elliptic operators should be rigid. Further, in 1989, the Witten conjecture was first proved by Taubes[8] and Bott-Taubes[7]. Hirzebruch [6] and Krichever[9] proved Witten conjecture for almost complex manifold case. In 1996, Liu used modular invariance to give a simple and uniform proof, as well as various extensive generalizations of Witten conjecture[12, 13]. And Liu also established several new vanishing theorems in this work. To further generalize, in 2000, Dessai established the rigidity and vanishing theorems for spinc case. Liu-Ma[14, 15] and Liu-Ma-Zhang[16, 17] generalized the rigidity and vanishing theorems to the family case on the levels of equivariant Chern character and of equivariant K-theory.

On the other hand, in 2009, Liu and Wang established the rigidity for twisted Toeplitz operators associated to the Witten bundles by approach[12, 13] under the assumption that the fixed point sets of the group action are 1-dimensional. This work is the first to study the Witten-type rigidity of Toplitz operators[18]. In 2015, Han and Yu extend the rigidity and vanishing properties for twisted Toeplitz operators to the cases of fixed

2010 Mathematics Subject Classification: 58C20; 57R20; 53C80.

Key words and phrases: Twisted Dirac operators; Twisted Toeplitz operators; Equivariant index theorem; Witten rigidity theorem

* *Corresponding author.*

points of general dimensions[20]. In [23, 24], we construct some new $SL(2, \mathbb{Z})$ modular forms, $\Gamma^0(2)$ modular forms and $\Gamma_0(2)$ modular forms in spin manifolds by the modular invariance of characteristic forms. Moreover, we get some divisibility results of index of the twisted Dirac operators and the twisted Toeplitz operators on spin manifolds. The purpose of this paper is to study the rigidity properties of these elliptic genus associated with twisted Dirac operators and twisted Toplitz operators in [23, 24] in the case of fixed points of general dimension by using Liu's method and Han-Yu's method.

The structure of this paper is briefly described below: In Section 2, we have introduce some definitions and basic concepts that we will use in the paper. In Section 3, we prove the Witten-type rigidity of twisted Dirac operators in even dimensions. Finally, in section 4, we prove the Witten-type rigidity of twisted Toplitz operators in odd dimensions.

2. CHARACTERISTIC FORMS AND MODULAR FORMS

The purpose of this section is to review the necessary knowledge on characteristic forms and modular forms that we are going to use.

2.1 characteristic forms

Let M be a Riemannian manifold. Let ∇^{TM} be the associated Levi-Civita connection on TM and $R^{TM} = (\nabla^{TM})^2$ be the curvature of ∇^{TM} . Let $\widehat{A}(TM, \nabla^{TM})$ and $\widehat{L}(TM, \nabla^{TM})$ be the Hirzebruch characteristic forms defined respectively by (cf. [19])

$$(2.1) \quad \widehat{A}(TM, \nabla^{TM}) = \det^{\frac{1}{2}} \left(\frac{\frac{\sqrt{-1}}{4\pi} R^{TM}}{\sinh(\frac{\sqrt{-1}}{4\pi} R^{TM})} \right),$$

$$(2.2) \quad \widehat{L}(TM, \nabla^{TM}) = \det^{\frac{1}{2}} \left(\frac{\frac{\sqrt{-1}}{2\pi} R^{TM}}{\tanh(\frac{\sqrt{-1}}{4\pi} R^{TM})} \right).$$

Let E, F be two Hermitian vector bundles over M carrying Hermitian connection ∇^E, ∇^F respectively. Let $R^E = (\nabla^E)^2$ (resp. $R^F = (\nabla^F)^2$) be the curvature of ∇^E (resp. ∇^F). If we set the formal difference $G = E - F$, then G carries an induced Hermitian connection ∇^G in an obvious sense. We define the associated Chern character form as

$$(2.3) \quad \text{ch}(G, \nabla^G) = \text{tr} \left[\exp\left(\frac{\sqrt{-1}}{2\pi} R^E\right) \right] - \text{tr} \left[\exp\left(\frac{\sqrt{-1}}{2\pi} R^F\right) \right].$$

For any complex number t , let

$$\wedge_t(E) = \mathbf{C}|_M + tE + t^2 \wedge^2(E) + \cdots, \quad S_t(E) = \mathbf{C}|_M + tE + t^2 S^2(E) + \cdots$$

denote respectively the total exterior and symmetric powers of E , which live in $K(M)[[t]]$. The following relations between these operations hold,

$$(2.4) \quad S_t(E) = \frac{1}{\wedge_{-t}(E)}, \quad \wedge_t(E - F) = \frac{\wedge_t(E)}{\wedge_t(F)}.$$

Moreover, if $\{\omega_i\}, \{\omega'_j\}$ are formal Chern roots for Hermitian vector bundles E, F respectively, then

$$(2.5) \quad \text{ch}(\wedge_t(E)) = \prod_i (1 + e^{\omega_i t})$$

Then we have the following formulas for Chern character forms,

$$(2.6) \quad \text{ch}(S_t(E)) = \frac{1}{\prod_i (1 - e^{\omega_i t})}, \quad \text{ch}(\wedge_t(E - F)) = \frac{\prod_i (1 + e^{\omega_i t})}{\prod_j (1 + e^{\omega'_j t})}.$$

If W is a real Euclidean vector bundle over M carrying a Euclidean connection ∇^W , then its complexification $W_{\mathbf{C}} = W \otimes \mathbf{C}$ is a complex vector bundle over M carrying a canonical induced Hermitian metric from that of W , as well as a Hermitian connection $\nabla^{W_{\mathbf{C}}}$ induced from ∇^W . If E is a vector bundle (complex or real) over M , set $\tilde{E} = E - \dim E$ in $K(M)$ or $KO(M)$.

2.2 Some properties about the Jacobi theta functions and modular forms

We first recall the four Jacobi theta functions are defined as follows (cf. [1]):

$$(2.7) \quad \theta(v, \tau) = 2q^{\frac{1}{8}} \sin(\pi v) \prod_{j=1}^{\infty} [(1 - q^j)(1 - e^{2\pi\sqrt{-1}v} q^j)(1 - e^{-2\pi\sqrt{-1}v} q^j)],$$

$$(2.8) \quad \theta_1(v, \tau) = 2q^{\frac{1}{8}} \cos(\pi v) \prod_{j=1}^{\infty} [(1 - q^j)(1 + e^{2\pi\sqrt{-1}v} q^j)(1 + e^{-2\pi\sqrt{-1}v} q^j)],$$

$$(2.9) \quad \theta_2(v, \tau) = \prod_{j=1}^{\infty} [(1 - q^j)(1 - e^{2\pi\sqrt{-1}v} q^{j-\frac{1}{2}})(1 - e^{-2\pi\sqrt{-1}v} q^{j-\frac{1}{2}})],$$

$$(2.10) \quad \theta_3(v, \tau) = \prod_{j=1}^{\infty} [(1 - q^j)(1 + e^{2\pi\sqrt{-1}v} q^{j-\frac{1}{2}})(1 + e^{-2\pi\sqrt{-1}v} q^{j-\frac{1}{2}})],$$

where $q = e^{2\pi\sqrt{-1}\tau}$ with $\tau \in \mathbf{H}$, the upper half complex plane. Let

$$(2.11) \quad \theta'(0, \tau) = \frac{\partial \theta(v, \tau)}{\partial v} \Big|_{v=0}.$$

Then the following Jacobi identity (cf. [1]) holds,

$$(2.12) \quad \theta'(0, \tau) = \pi \theta_1(0, \tau) \theta_2(0, \tau) \theta_3(0, \tau).$$

Denote

$$SL_2(\mathbf{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbf{Z}, ad - bc = 1 \right\}$$

the modular group. Let $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ be the two generators of $SL_2(\mathbf{Z})$. They act on \mathbf{H} by $S\tau = -\frac{1}{\tau}$, $T\tau = \tau + 1$.

Definition 2.1 A modular form over Γ , a subgroup of $SL_2(\mathbf{Z})$, is a holomorphic function $f(\tau)$ on \mathbf{H} such that

$$(2.13) \quad f(g\tau) := f\left(\frac{a\tau + b}{c\tau + d}\right) = \chi(g)(c\tau + d)^k f(\tau), \quad \forall g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma,$$

where $\chi : \Gamma \rightarrow \mathbf{C}^*$ is a character of Γ . k is called the weight of f .

Let

$$\Gamma_0(2) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbf{Z}) \mid c \equiv 0 \pmod{2} \right\},$$

$$\Gamma^0(2) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbf{Z}) \mid b \equiv 0 \pmod{2} \right\},$$

be the two modular subgroups of $SL_2(\mathbf{Z})$. It is known that the generators of $\Gamma_0(2)$ are T , ST^2ST , the generators of $\Gamma^0(2)$ are STS , T^2STS (cf. [1]).

3. TWISTED DIRAC OPERATOR AND WITTEN RIGIDITY THEOREM IN ODD DIMENSIONS

Let M be a $4k$ -dimensional spin manifold and $\Delta(M)$ be the spinor bundle. Let $\widetilde{T_{\mathbf{C}}M} = T_{\mathbf{C}}M - \dim M$. Set

$$(3.1) \quad \Theta_1(T_{\mathbf{C}}M) = \bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_{\mathbf{C}}M}) \otimes \bigotimes_{m=1}^{\infty} \Lambda_{q^m}(\widetilde{T_{\mathbf{C}}M}),$$

$$(3.2) \quad \Theta_2(T_{\mathbf{C}}M) = \bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_{\mathbf{C}}M}) \otimes \bigotimes_{m=1}^{\infty} \Lambda_{-q^{m-\frac{1}{2}}}(\widetilde{T_{\mathbf{C}}M}),$$

$$(3.3) \quad \Theta_3(T_{\mathbf{C}}M) = \bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_{\mathbf{C}}M}) \otimes \bigotimes_{m=1}^{\infty} \Lambda_{q^{m-\frac{1}{2}}}(\widetilde{T_{\mathbf{C}}M}).$$

Let V be a rank $2l$ real vector bundle on M . Moreover, $V_{\mathbf{C}} = V \otimes \mathbf{C}$. Set

$$(3.4) \quad Q_1(V_{\mathbf{C}}) = \Delta(V) \otimes \bigotimes_{n=1}^{\infty} \Lambda_{q^n}(\widetilde{V_{\mathbf{C}}}),$$

$$(3.5) \quad Q_2(V_{\mathbf{C}}) = \bigotimes_{n=1}^{\infty} \Lambda_{-q^{n-\frac{1}{2}}}(\widetilde{V_{\mathbf{C}}}),$$

$$(3.6) \quad Q_3(V_{\mathbf{C}}) = \bigotimes_{n=1}^{\infty} \Lambda_{q^{n-\frac{1}{2}}}(\widetilde{V_{\mathbf{C}}}).$$

Let

$$(3.7) \quad \Phi_0(M) = \Delta(M) \otimes \Theta_1(T_{\mathbf{C}}M) + 2^{2k}\Theta_2(T_{\mathbf{C}}M) + 2^{2k}\Theta_3(T_{\mathbf{C}}M),$$

$$(3.8) \quad \Phi(M, V) = \Phi_0 \otimes (Q_1(V)) \otimes (Q_2(V)) \otimes (Q_3(V)).$$

Let M be a closed smooth spin Riemannian manifold which admits a circle action. Without loss of generality, we may assume that S^1 acts on M isometrically and preserves the spin structure of M .

Let V be an oriented real rank- $2l$ vector bundle on a manifold M , and V is equipped with an S^1 action which restricts on each V -fiber over M to a linear action preserving that fiber. Then the associated twisted Dirac operator is S^1 -equivariant, which implies that the corresponding orthogonal projection P_+ is also S^1 -equivariant.

Next we assume $g = e^{2\pi it} \in S^1$ be a generator of the action group and let F_α denote the fixed submanifold of the circle action on M . In general, F_α is not connected. Let \mathbf{N} denote the normal bundle to F_α in M , which can be identified as the orthogonal complement of TF_α in $TM|_{F_\alpha}$. Then we have the following S^1 -equivariant decomposition when restricted upon F_α ,

$$(3.9) \quad TM|_{F_\alpha} = \mathbf{N}_1 \oplus \cdots \oplus \mathbf{N}_{2r} \oplus TF_\alpha,$$

where each $\mathbf{N}_\beta, \beta = 1, \dots, 2r$ is a complex vector bundle, and that g acts on \mathbf{N}_β by $e^{2\pi i m_\beta t}$. We assume $\{\pm 2\pi i x_i, 1 \leq i \leq 2r\}$ be the Chern roots of $\mathbf{N}_\beta \otimes \mathbf{C}$. Let $\{\pm 2\pi i y_j, 1 \leq j \leq 2s\}$ be the Chern roots of $TF_\alpha \otimes \mathbf{C}$.

Similarly, let

$$(3.10) \quad V|_{F_\alpha} = V_1 \oplus \cdots \oplus V_l \oplus V_0^{\mathbb{R}}$$

be the S^1 -equivariant decomposition of the restrictions of V over F_α , where V_ν is a complex vector bundle and $V_0^{\mathbb{R}}$ is the real subbundle of $V|_{F_\alpha}$. Assume that g acts on V_ν by $e^{2\pi i n_\nu t}$. And let $\{\pm 2\pi i z_\nu, 1 \leq \nu \leq l\}$ be the Chern roots of $V_\nu \otimes \mathbf{C}$. Let $\{\pm 2\pi i z_\nu^0\}$ be the Chern roots of $V_0^{\mathbb{R}} \otimes \mathbf{C}$.

We suppose $p_1(\cdot)_{S^1}$ denote the first S^1 -equivariant pontrjagin class, then we have the following rigidity theorems.

Theorem 3.1. *For an even dimensional connected spin manifold with non-trivial S^1 -action, if $3p_1(V)_{S^1} = 0$, then the twisted Dirac operators $\mathcal{D} \otimes \Phi$ are rigid.*

First, we calculate the corresponding Lefschetz numbers.

Proposition 3.2. *The Lefschetz numbers $\mathcal{D} \otimes \Phi$ are*

$$(3.11) \quad \begin{aligned} \mathcal{L}(g; \tau) &= 2^{2s+2r+l} \left(\frac{1}{\pi}\right)^{2r} \sum_{\alpha=1}^{2s} \int_{F_\alpha} \prod_{j=1}^{2s} \left(y_j \frac{\theta'(0, \tau)}{\theta(y_j, \tau)} \right) \\ &\cdot \prod_{i=1}^{2r} \left(\frac{\theta'(0, \tau)}{\theta(x_i + m_i t, \tau)} \right) \sum_{\omega=1}^3 \left(\prod_{j=1}^{2s} \frac{\theta_\omega(y_j, \tau)}{\theta_\omega(0, \tau)} \prod_{i=1}^{2r} \frac{\theta_\omega(x_i + m_i t, \tau)}{\theta_\omega(0, \tau)} \right) \\ &\cdot \prod_{\nu=1}^l \frac{\theta_1(z_\nu + \eta_\nu t, \tau) \theta_2(z_\nu + \eta_\nu t, \tau) \theta_3(z_\nu + \eta_\nu t, \tau)}{\theta_1(0, \tau) \theta_2(0, \tau) \theta_3(0, \tau)}, \end{aligned}$$

Proof. By Lefschetz fixed point formula

$$(3.12) \quad \mathcal{L}(g) = \sum_i \int_{F_i} \hat{A}(TF_i) \left[Pf \left(2 \sinh \left(\frac{\Omega^\perp}{4\pi} + \sqrt{-1} \frac{\Theta_j}{2} \right) \right) \right]^{-1} \text{ch}_g(E).$$

We can calculate it directly

$$(3.13) \quad \hat{A}(TF_\alpha) \left[Pf \left(2 \sinh \left(\frac{\Omega^\perp}{4\pi} + \sqrt{-1} \frac{\Theta_\beta}{2} \right) \right) \right]^{-1} = \prod_{j=1}^{2s} \frac{\pi y_j}{\sinh \pi y_j} \prod_{i=1}^{2r} \frac{1}{\sinh \pi(x_i + m_i t)},$$

(3.14)

$$\begin{aligned} \text{ch}_g(\Phi_0) &= \left(\frac{2}{\pi} \right)^{2s+2r} \prod_{j=1}^{2s} \sin(\pi y_j) \left(\frac{\theta'(0, \tau)}{\theta(y_j, \tau)} \right) \prod_{i=1}^{2r} \sin(\pi(x_i + m_i t)) \left(\frac{\theta'(0, \tau)}{\theta(x_i + m_i t, \tau)} \right) \\ &\quad \cdot \sum_{\omega=1}^3 \left(\prod_{j=1}^{2s} \frac{\theta_\omega(y_j, \tau)}{\theta_\omega(0, \tau)} \prod_{i=1}^{2r} \frac{\theta_\omega(x_i + m_i t, \tau)}{\theta_\omega(0, \tau)} \right), \end{aligned}$$

$$(3.15) \quad \text{ch}_g \left(\Delta(V) \otimes \bigotimes_{n=1}^{\infty} \Lambda_{q^n}(\widetilde{V}_{\mathbf{C}}) \right) = 2^l \prod_{\nu=1}^l \frac{\theta_1(z_\nu + \eta_\nu t, \tau)}{\theta_1(0, \tau)},$$

$$(3.16) \quad \text{ch}_g \left(\bigotimes_{n=1}^{\infty} \Lambda_{-q^{n-\frac{1}{2}}}(\widetilde{V}_{\mathbf{C}}) \right) = \prod_{\nu=1}^l \frac{\theta_2(z_\nu + \eta_\nu t, \tau)}{\theta_2(0, \tau)},$$

$$(3.17) \quad \text{ch}_g \left(\bigotimes_{n=1}^{\infty} \Lambda_{q^{n-\frac{1}{2}}}(\widetilde{V}_{\mathbf{C}}) \right) = \prod_{\nu=1}^l \frac{\theta_3(z_\nu + \eta_\nu t, \tau)}{\theta_3(0, \tau)}.$$

So, it's true that we get Proposition 3.2 by (3.12)-(3.17). \square

In the follows, let us view the above expression as defining a function $\mathcal{L}'(t, \tau)$, such that $\mathcal{L}'(t, \tau) = \mathcal{L}(g; \tau)$. The expression for $\mathcal{L}'(t, \tau)$ involves only complex-differentiable functions, so we can extend its domain to every complex number t and choice of τ in the open upper half-plane, i.e. $\mathcal{L}'(t, \tau)$ where the expression exists in $\mathbf{C} \times \mathbf{H}$. The Witten rigidity theorems are equivalent to that these $\mathcal{L}'(t, \tau)$ are independent of t . Then, we have the following lemma.

Lemma 3.3. *Let $(t, \tau) \in \mathbf{C} \times \mathbf{H}$ be in the domain of $\mathcal{L}'(t, \tau)$.*

(1) *Then $\mathcal{L}'(t, \tau) = \mathcal{L}'(t + a, \tau)$ for any $a \in 2\mathbb{Z}$.*

(2) *If $3p_1(V)_{S^1} = 0$, then $\mathcal{L}'(t, \tau) = \mathcal{L}'(t + a\tau, \tau)$ for any $a \in 2\mathbb{Z}$.*

Proof. We have the following transformation formulas of theta-functions[7]:

$$\begin{aligned} \theta(t+1, \tau) &= -\theta(t, \tau), & \theta(t+\tau, \tau) &= -q^{-1/2} e^{-2\pi i t} \theta(t, \tau), \\ \theta_1(t+1, \tau) &= -\theta_1(t, \tau), & \theta_1(t+\tau, \tau) &= q^{-1/2} e^{-2\pi i t} \theta_1(t, \tau), \\ \theta_2(t+1, \tau) &= \theta_2(t, \tau), & \theta_2(t+\tau, \tau) &= -q^{-1/2} e^{-2\pi i t} \theta_2(t, \tau), \end{aligned}$$

$$\theta_3(t+1, \tau) = \theta_3(t, \tau), \quad \theta_3(t+\tau, \tau) = -q^{-1/2} e^{-2\pi i t} \theta_3(t, \tau),$$

where q here is equal to $e^{\pi i \tau}$.

By reusing the formula in the first column, for $a \in 2\mathbb{Z}$, we can easily verify that $\theta, \theta_1, \theta_2, \theta_3$ terms is unchanged in \mathcal{L}' . So (1) is true.

For (2), under the replacement $t \rightarrow t + a\tau$ for a an even integer, we apply the second column formula to get the following formula:

$$\begin{aligned} \theta(t+a\tau) &= (-1)^a e^{-2\pi i(at+a^2\tau/2)} \theta(t, \tau), \\ \theta_1(t+a\tau) &= e^{-2\pi i(at+a^2\tau/2)} \theta_1(t, \tau), \\ \theta_2(t+a\tau) &= (-1)^a e^{-2\pi i(at+a^2\tau/2)} \theta_2(t, \tau), \\ \theta_3(t+a\tau) &= e^{-2\pi i(at+a^2\tau/2)} \theta_3(t, \tau). \end{aligned}$$

Then, under the replacement $t \rightarrow t + a\tau$ for a an even integer, the following transformation property holds:

$$(3.18) \quad \theta_\mu(z_\nu + n_\nu(t+a\tau), \tau) = e^{-2\pi i a(n_\nu z_\nu + n_\nu^2(t+a^2\tau/2))} \theta_\mu(z_\nu + n_\nu t, \tau),$$

where $\theta_\mu \in \{\theta, \theta_1, \theta_2, \theta_3\}$.

By the condition $3p_1(V)_{S^1} = 0$ implies that

$$3p_1(V)_{S^1} = p_1(\otimes_3 V)_{S^1} = 0,$$

then, we have

$$\sum_{\nu=1}^l n_\nu z_\nu = 0, \quad \sum_{\nu=1}^l n_\nu^2 = 0.$$

Thus, we can see that for each connected component F_α of a fixed locus of M , \mathcal{L}' is completely unchanged. (2) certification. \square

This lemma tells us that for fixed τ these $\mathcal{L}'(t, \tau)$ are meromorphic functions on the torus $\mathbf{C}/2\mathbb{Z} \times 2\mathbb{Z}\tau$. Therefore, to get the rigidity we only need to prove that they are holomorphic in t . We will actually prove that they are holomorphic in (t, τ) on $\mathbf{C} \times \mathbf{H}$.

Given

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$$

define its modular transformation on $\mathbf{C} \times \mathbf{H}$ by

$$g(t, \tau) = \left(\frac{\tau}{c\tau + d}, \frac{a\tau + b}{c\tau + d} \right).$$

This defines a group action. Obviously two generators of $SL_2(\mathbb{Z})$,

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

act by

$$S(t, \tau) = \left(\frac{t}{\tau}, -\frac{1}{\tau} \right), \quad T(t, \tau) = (t, \tau + 1).$$

Then, we have the following transformation formulas

Lemma 3.4. (1) If $3p_1(V)_{S^1} = 0$, then the action of S ,

$$(3.19) \quad \mathcal{L}'\left(\frac{t}{\tau}, -\frac{1}{\tau}\right) = \tau^{2k} \mathcal{L}'(t, \tau).$$

(2) The action of T , that

$$(3.20) \quad \mathcal{L}'(t, \tau + 1) = \mathcal{L}'(t, \tau).$$

Proof. We have the following transformation laws of Jacobi theta-functions under the actions of S and T (cf. [1]):

$$\begin{aligned} \theta\left(\frac{t}{\tau}, -\frac{1}{\tau}\right) &= \frac{1}{\mathbf{i}} \sqrt{\frac{\tau}{\mathbf{i}}} e^{\pi i t^2 / \tau} \theta(t, \tau), \quad \theta(t, \tau + 1) = e^{\pi i / 4} \theta(t, \tau); \\ \theta_1\left(\frac{t}{\tau}, -\frac{1}{\tau}\right) &= \frac{1}{\mathbf{i}} \sqrt{\frac{\tau}{\mathbf{i}}} e^{\pi i t^2 / \tau} \theta_2(t, \tau), \quad \theta_1(t, \tau + 1) = e^{\pi i / 4} \theta_1(t, \tau); \\ \theta_2\left(\frac{t}{\tau}, -\frac{1}{\tau}\right) &= \frac{1}{\mathbf{i}} \sqrt{\frac{\tau}{\mathbf{i}}} e^{\pi i t^2 / \tau} \theta_1(t, \tau), \quad \theta_2(t, \tau + 1) = \theta_3(t, \tau); \\ \theta_3\left(\frac{t}{\tau}, -\frac{1}{\tau}\right) &= \frac{1}{\mathbf{i}} \sqrt{\frac{\tau}{\mathbf{i}}} e^{\pi i t^2 / \tau} \theta_3(t, \tau), \quad \theta_3(t, \tau + 1) = e^{\pi i / 4} \theta_2(t, \tau); \\ \theta'\left(0, -\frac{1}{\tau}\right) &= \left(\frac{\tau}{\mathbf{i}}\right)^{3/2} \theta'(0, \tau), \quad \theta'(0, \tau + 1) = e^{\pi i / 4} \theta'(0, \tau). \end{aligned}$$

Then, we obtain the following transformation formulas,

$$(3.21) \quad y_j \frac{\theta'(0, -1/\tau)}{\theta(y_j, -1/\tau)} = e^{-\pi i \tau y_j^2} \tau y_j \frac{\theta'(0, \tau)}{\theta(\tau y_j, \tau)},$$

$$(3.22) \quad \frac{\theta'(0, -1/\tau)}{\theta(x_i + m_i t / \tau, -1/\tau)} = e^{-\pi i \tau (x_i + m_i t / \tau)^2} \tau \frac{\theta'(0, \tau)}{\theta(\tau x_i + m_i t, \tau)},$$

$$(3.23) \quad \frac{\theta_1(y_j, -1/\tau)}{\theta_1(0, -1/\tau)} = e^{\pi i \tau y_j^2} \frac{\theta_2(\tau y_j, \tau)}{\theta_2(0, \tau)},$$

$$(3.24) \quad \frac{\theta_1(x_i + m_i t / \tau, -1/\tau)}{\theta_1(0, -1/\tau)} = e^{\pi i \tau (x_i + m_i t / \tau)^2} \frac{\theta_2(\tau x_i + m_i t, \tau)}{\theta_2(0, \tau)},$$

$$(3.25) \quad \frac{\theta_2(y_j, -1/\tau)}{\theta_2(0, -1/\tau)} = e^{\pi i \tau y_j^2} \frac{\theta_1(\tau y_j, \tau)}{\theta_1(0, \tau)},$$

$$(3.26) \quad \frac{\theta_2(x_i + m_i t / \tau, -1/\tau)}{\theta_2(0, -1/\tau)} = e^{\pi i \tau (x_i + m_i t / \tau)^2} \frac{\theta_1(\tau x_i + m_i t, \tau)}{\theta_1(0, \tau)},$$

$$(3.27) \quad \frac{\theta_3(y_j, -1/\tau)}{\theta_3(0, -1/\tau)} = e^{\pi i \tau y_j^2} \frac{\theta_3(\tau y_j, \tau)}{\theta_3(0, \tau)},$$

$$(3.28) \quad \frac{\theta_3(x_i + m_i t/\tau, -1/\tau)}{\theta_3(0, -1/\tau)} = e^{\pi i \tau (x_i + m_i t/\tau)^2} \frac{\theta_3(\tau x_i + m_i t, \tau)}{\theta_3(0, \tau)},$$

$$(3.29) \quad \frac{\theta_1(z_\nu + n_\nu t/\tau, -1/\tau)}{\theta_1(0, -1/\tau)} = e^{\pi i \tau (z_\nu + n_\nu t/\tau)^2} \frac{\theta_2(\tau z_\nu + n_\nu t, \tau)}{\theta_2(0, \tau)}.$$

Combining the condition $3p_1(V)_{S^1} = 0$, and (3.23)-(3.31), we obtain (3.19).

For (2), we use the transformation laws of Jacobi theta-functions under the action of T , which we can easily verify to get (3.20). \square

Lemma 3.5. *For any function*

$$\mathcal{L}'(t, \tau)$$

its modular transformation is holomorphic in $(t, \tau) \in \mathbf{R} \times \mathbf{H}$.

Proof. The proof is almost the same as the proof of [12, Lemma 2.3] except that we use Proposition 3.2 instead of the corresponding Lefschetz fixed point formulas. \square

Proof of Theorem 3.1. We prove that these \mathcal{F}_λ are holomorphic on $\mathbf{C} \times \mathbf{H}$ which implies the rigidity of Theorem 3.1. By (3.11), we see that the possible poles of $\mathcal{L}'(t, \tau)$ can be written in the form $t = \frac{k}{l}(c\tau + d)$ for integers k, l, c, d with $(c, d) = 1$.

We can always find integers a, b such that $ad - bc = 1$. Then the matrix $g_0 = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \in SL_2(\mathbb{Z})$ induces an action

$$\mathcal{L}'(g_0(t, \tau)) = \mathcal{F}\left(\frac{t}{-c\tau + a}, \frac{d\tau - b}{-c\tau + a}\right).$$

Now, if $t = \frac{k}{l}(c\tau + d)$ is a polar divisor of $\mathcal{F}(t, \tau)$, then one polar divisor of $\mathcal{L}'(g_0(t, \tau))$ is given by

$$\frac{t}{-c\tau + a} = \frac{k}{l} \left(c \frac{d\tau - b}{-c\tau + a} + d \right)$$

which gives $t = \frac{k}{l}$.

By Lemma 3.4, we know that up to some constant, $\mathcal{L}'(g_0(t, \tau))$ is still one of $\{\mathcal{L}'(t, \tau)\}$. This contradicts Lemma 3.5, therefore, this completes the proof of Theorem 3.1.

4. TWISTED TOPLITZ OPERATOR AND WITTEN RIGIDITY THEOREM IN ODD DIMENSIONS

We recall the odd Chern character of a smooth map g from M to the general linear group $GL(N, \mathbf{C})$ with N a positive integer (see [20]). Let d denote a trivial connection on $\mathbf{C}^N|_M$. We will denote by $c_g(M, [g])$ the cohomology class associated to the closed n -form

$$(4.1) \quad c_n(\mathbf{C}^N|_M, g, d) = \left(\frac{1}{2\pi\sqrt{-1}} \right)^{\frac{(n+1)}{2}} \text{Tr}[(g^{-1}dg)^n].$$

The odd Chern character form $\text{ch}(\mathbf{C}^N|_M, g, d)$ associated to g and d by definition is

$$(4.2) \quad \text{ch}(\mathbf{C}^N|_M, g, d) = \sum_{n=1}^{\infty} \frac{n!}{(2n+1)!} c_{2n+1}((\mathbf{C}^N|_M, g, d)).$$

Let the connection ∇_u on the trivial bundle $\mathbf{C}^N|_M$ defined by

$$(4.3) \quad \nabla_u = (1-u)d + ug^{-1} \cdot d \cdot g, \quad u \in [0, 1].$$

Then we have

$$(4.4) \quad d\text{ch}(\mathbf{C}^N|_M, g, d) = \text{ch}(\mathbf{C}^N|_M, d) - \text{ch}(\mathbf{C}^N|_M, g^{-1} \cdot d \cdot g).$$

Now let $g : M \rightarrow SO(N)$ and we assume that N is even and large enough. Let E denote the trivial real vector bundle of rank N over M . We equip E with the canonical trivial metric and trivial connection d . Set

$$\nabla_u = d + ug^{-1}dg, \quad u \in [0, 1].$$

Let R_u be the curvature of ∇_u , then

$$(4.5) \quad R_u = (u^2 - u)(g^{-1}dg)^2.$$

We also consider the complexification of E and g extends to a unitary automorphism of $E_{\mathbb{C}}$. The connection ∇_u extends to a Hermitian connection on $E_{\mathbb{C}}$ with curvature still given by (3.6). Let $\Delta(E)$ be the spinor bundle of E , which is a trivial Hermitian bundle of rank $2^{\frac{N}{2}}$. We assume that g has a lift to the Spin group $\text{Spin}(N) : g^{\Delta} : M \rightarrow \text{Spin}(N)$. So g^{Δ} can be viewed as an automorphism of $\Delta(E)$ preserving the Hermitian metric. We lift d on E to be a trivial Hermitian connection d^{Δ} on $\Delta(E)$, then

$$(4.6) \quad \nabla_u^{\Delta} = (1-u)d^{\Delta} + u(g^{\Delta})^{-1} \cdot d^{\Delta} \cdot g^{\Delta}, \quad u \in [0, 1]$$

lifts ∇_u on E to $\Delta(E)$. Let $Q_j(E), j = 1, 2, 3$ be the virtual bundles defined as follows:

$$(4.7) \quad Q_1(E) = \Delta(E) \otimes \bigotimes_{n=1}^{\infty} \wedge_{q^n}(\widetilde{E}_{\mathbb{C}});$$

$$(4.8) \quad Q_2(E) = \bigotimes_{n=1}^{\infty} \wedge_{-q^{n-\frac{1}{2}}}(\widetilde{E}_{\mathbb{C}});$$

$$(4.9) \quad Q_3(E) = \bigotimes_{n=1}^{\infty} \wedge_{q^{n-\frac{1}{2}}}(\widetilde{E}_{\mathbb{C}}).$$

Let g on E have a lift $g^{Q(E)}$ on $Q(E)$ and ∇_u have a lift $\nabla_u^{Q(E)}$ on $Q(E)$. Following [20], we defined $\text{ch}(Q_j(E), g^{Q_j(E)}, d, \tau)$ for $j = 1, 2, 3$ as following

$$(4.10) \quad \text{ch}(Q_j(E), \nabla_0^{Q_j(E)}, \tau) - \text{ch}(Q_j(E), \nabla_1^{Q_j(E)}, \tau) = d\text{ch}(Q_j(E), g^{Q_j(E)}, d, \tau),$$

where

$$(4.11) \quad \text{ch}(Q_1(E), g^{Q_1(E)}, d, \tau) = -\frac{2^{N/2}}{8\pi^2} \int_0^1 \text{Tr} \left[g^{-1} dg \frac{\theta'_1(R_u/(4\pi^2), \tau)}{\theta_1(R_u/(4\pi^2), \tau)} \right] du,$$

and for $j = 2, 3$

$$(4.12) \quad \text{ch}(Q_j(E), g^{Q_j(E)}, d, \tau) = -\frac{1}{8\pi^2} \int_0^1 \text{Tr} \left[g^{-1} dg \frac{\theta'_j(R_u/(4\pi^2), \tau)}{\theta_j(R_u/(4\pi^2), \tau)} \right] du.$$

Let M be a $(4k - 1)$ -dimensional spin manifold and $\Delta(M)$ be the spinor bundle. Let $\widetilde{T_{\mathbf{C}}M} = T_{\mathbf{C}}M - \dim M$. Set

$$(4.13) \quad \Psi_1(M, V, E) = \Phi_0 \otimes \left(\Delta(V) \otimes \bigotimes_{n=1}^{\infty} \Lambda_{q^n}(\widetilde{V_{\mathbf{C}}}) \right) \otimes (Q_1(E), g^{Q_1(E)}),$$

$$(4.14) \quad \Psi_2(M, V, E) = \Phi_0 \otimes \left(\bigotimes_{n=1}^{\infty} \Lambda_{-q^{n-\frac{1}{2}}}(\widetilde{V_{\mathbf{C}}}) \right) \otimes (Q_2(E), g^{Q_2(E)}),$$

$$(4.15) \quad \Psi_3(M, V, E) = \Phi_0 \otimes \left(\bigotimes_{n=1}^{\infty} \Lambda_{q^{n-\frac{1}{2}}}(\widetilde{V_{\mathbf{C}}}) \right) \otimes (Q_3(E), g^{Q_3(E)}).$$

Let M be an odd dimensional closed smooth spin Riemannian manifold which admits a circle action. Let V be an S^1 -equivariant complex vector bundle over M carrying an S^1 -invariant Hermitian connection. In addition, we assume $g : M \rightarrow GL(N, \mathbf{C})$ is S^1 -invariant, i.e.,

$$(4.16) \quad g(hx) = g(x), \text{ for any } h \in S^1 \text{ and } x \in M.$$

Thus the twisted Toeplitz operator $\mathcal{T} \otimes \Psi_i$, $i = 1, 2, 3$ is S^1 -equivariant.

We consider the fixed point case. Similarly, we have the following S^1 -equivariant decomposition when restricted upon F_{α} ,

$$(4.17) \quad TM|_{F_{\alpha}} = \mathbf{N}_1 \oplus \cdots \oplus \mathbf{N}_{2r} \oplus TF_{\alpha},$$

where each \mathbf{N}_{β} , $\beta = 1, \dots, 2r$ is a complex vector bundle such that $h \in S^1$ acts on it by $e^{2\pi i m_i t}$. We assume $\{\pm 2\pi i x_i, 1 \leq i \leq 2r\}$ be the Chern roots of $\mathbf{N}_{\beta} \otimes \mathbf{C}$. Let $\{\pm 2\pi i y_j, 1 \leq j \leq 2s\}$ be the Chern roots of $TF_{\alpha} \otimes \mathbf{C}$. And let

$$(4.18) \quad V|_{F_{\alpha}} = V_1 \oplus \cdots \oplus V_l \oplus V_0^{\mathbb{R}}$$

be the S^1 -equivariant decomposition of the restrictions of V over F_{α} , where V_{ν} is a complex vector bundle and $V_0^{\mathbb{R}}$ is the real subbundle of $V|_{F_{\alpha}}$. Assume that $h \in S^1$ acts on V_{ν} by $e^{2\pi i n_{\nu} t}$. And let $\{\pm 2\pi i z_{\nu}, 1 \leq \nu \leq l\}$ be the Chern roots of $V_{\nu} \otimes \mathbf{C}$. Let $\{\pm 2\pi i z_{\nu}^0\}$ be the Chern roots of $V_0^{\mathbb{R}} \otimes \mathbf{C}$.

We suppose $p_1(\cdot)_{S^1}$ denote the first S^1 -equivariant pontrjagin class, then we have the following rigidity theorems.

Theorem 4.1. *For an odd dimensional connected spin manifold with non-trivial S^1 -action, if $p_1(V)_{S^1} = 0$ and $c_3(E, g, d) = 0$, then the Toeplitz operators $\mathcal{T} \otimes \Psi_i$, $i = 1, 2, 3$ are rigid.*

First, we calculate the corresponding Lefschetz numbers.

Proposition 4.2. *The Lefschetz numbers $\mathcal{T} \otimes \Psi_i$, $i = 1, 2, 3$ are*

$$(4.19) \quad \begin{aligned} \mathcal{L}_1(g; \tau) &= 2^{2s+2r+al} \left(\frac{1}{\pi}\right)^{2r} \sum_{\alpha=1}^{2s} \int_{F_\alpha} \prod_{j=1}^{2s} \left(y_j \frac{\theta'(0, \tau)}{\theta(y_j, \tau)}\right) \\ &\cdot \prod_{i=1}^{2r} \left(\frac{\theta'(0, \tau)}{\theta(x_i + m_i t, \tau)}\right) \sum_{\omega=1}^3 \left(\prod_{j=1}^{2s} \frac{\theta_\omega(y_j, \tau)}{\theta_\omega(0, \tau)} \prod_{i=1}^{2r} \frac{\theta_\omega(x_i + m_i t, \tau)}{\theta_\omega(0, \tau)}\right) \\ &\cdot \prod_{\nu=1}^l \frac{\theta_1^a(z_\nu + \eta_\nu t, \tau)}{\theta_1^a(0, \tau)} \text{ch}(Q_1(E), g^{Q_1(E)}, d, \tau), \end{aligned}$$

$$(4.20) \quad \begin{aligned} \mathcal{L}_2(g; \tau) &= 2^{2s+2r} \left(\frac{1}{\pi}\right)^{2r} \sum_{\alpha=1}^{2s} \int_{F_\alpha} \prod_{j=1}^{2s} \left(y_j \frac{\theta'(0, \tau)}{\theta(y_j, \tau)}\right) \\ &\cdot \prod_{i=1}^{2r} \left(\frac{\theta'(0, \tau)}{\theta(x_i + m_i t, \tau)}\right) \sum_{\omega=1}^3 \left(\prod_{j=1}^{2s} \frac{\theta_\omega(y_j, \tau)}{\theta_\omega(0, \tau)} \prod_{i=1}^{2r} \frac{\theta_\omega(x_i + m_i t, \tau)}{\theta_\omega(0, \tau)}\right) \\ &\cdot \prod_{\nu=1}^l \frac{\theta_2^b(z_\nu + \eta_\nu t, \tau)}{\theta_2^b(0, \tau)} \text{ch}(Q_2(E), g^{Q_2(E)}, d, \tau), \end{aligned}$$

and

$$(4.21) \quad \begin{aligned} \mathcal{L}_3(g; \tau) &= 2^{2s+2r} \left(\frac{1}{\pi}\right)^{2r} \sum_{\alpha=1}^{2s} \int_{F_\alpha} \prod_{j=1}^{2s} \left(y_j \frac{\theta'(0, \tau)}{\theta(y_j, \tau)}\right) \\ &\cdot \prod_{i=1}^{2r} \left(\frac{\theta'(0, \tau)}{\theta(x_i + m_i t, \tau)}\right) \sum_{\omega=1}^3 \left(\prod_{j=1}^{2s} \frac{\theta_\omega(y_j, \tau)}{\theta_\omega(0, \tau)} \prod_{i=1}^{2r} \frac{\theta_\omega(x_i + m_i t, \tau)}{\theta_\omega(0, \tau)}\right) \\ &\cdot \prod_{\nu=1}^l \frac{\theta_3^c(z_\nu + \eta_\nu t, \tau)}{\theta_3^c(0, \tau)} \text{ch}(Q_3(E), g^{Q_3(E)}, d, \tau). \end{aligned}$$

Proof. Similar to Proposition 3.2, we can calculate equations (4.19), (4.20) and (4.21) by using Lefschetz fixed point formula, (4.11) and (4.12). \square

Next, let us view the above expression as defining a function $\mathcal{F}_\lambda(t, \tau)$, for each $\lambda \in \{1, 2, 3\}$, i.e. $\mathcal{F}_\lambda(t, \tau) = \mathcal{L}_\lambda(g; \tau)$. The expression for $\mathcal{F}_\lambda(t, \tau)$ involves only complex-differentiable functions, so we can extend its domain to every complex number t and choice of τ in the open upper half-plane, i.e. $\mathcal{F}_\lambda(t, \tau)$ where the expression exists in $\mathbf{C} \times \mathbf{H}$. The Witten rigidity theorems are equivalent to that these $\mathcal{F}_\lambda(t, \tau)$ are independent of t . Then, we have the following lemma.

Lemma 4.3. *Let $(t, \tau) \in \mathbf{C} \times \mathbf{H}$ be in the domain of $\mathcal{F}_\lambda(t, \tau)$, $\lambda \in \{1, 2, 3\}$.*

(1) Then $\mathcal{F}_\lambda(t, \tau) = \mathcal{F}_\lambda(t + a, \tau)$ for any $a \in 2\mathbb{Z}$.

(2) If $p_1(V)_{S^1} = 0$, then $\mathcal{F}_\lambda(t, \tau) = \mathcal{F}_\lambda(t + a\tau, \tau)$ for any $a \in 2\mathbb{Z}$.

Proof. Similar to Lemma 3.3. □

Next, we will actually prove that these \mathcal{F}_λ are holomorphic in (t, τ) on $\mathbf{C} \times \mathbf{H}$. Similar to Lemma 3.4, we have the following lemma

Lemma 4.4. *(1) If $p_1(V)_{S^1} = 0$, and $c_3(E, g, d) = 0$, then the action of S ,*

$$(4.22) \quad \mathcal{F}_1\left(\frac{t}{\tau}, -\frac{1}{\tau}\right) = 2^{al+N/2}\tau^{2k}\mathcal{F}_2(t, \tau),$$

$$(4.23) \quad \mathcal{F}_2\left(\frac{t}{\tau}, -\frac{1}{\tau}\right) = 2^{-(al+N/2)}\tau^{2k}\mathcal{F}_1(t, \tau),$$

$$(4.24) \quad \mathcal{F}_3\left(\frac{t}{\tau}, -\frac{1}{\tau}\right) = \tau^{2k}\mathcal{F}_3(t, \tau).$$

(2) The action of T , then

$$(4.25) \quad \mathcal{F}_1(t, \tau + 1) = \mathcal{F}_1(t, \tau),$$

$$(4.26) \quad \mathcal{F}_2(t, \tau + 1) = \mathcal{F}_3(t, \tau),$$

$$(4.27) \quad \mathcal{F}_3(t, \tau + 1) = \mathcal{F}_2(t, \tau).$$

Proof. By Proposition 2.2 in [20], we have if $c_3(E, g, d) = 0$, then for any integer $i \geq 1$,

$$(4.28) \quad \left\{ \text{ch} \left(Q_1(E), g^{Q_1(E)}, d, -\frac{1}{\tau} \right) \right\}^{4i-1} = 2^{N/2} \left\{ \tau^{2i} \text{ch} \left(Q_2(E), g^{Q_2(E)}, d, \tau \right) \right\}^{4i-1},$$

$$(4.29) \quad \left\{ \text{ch} \left(Q_2(E), g^{Q_2(E)}, d, -\frac{1}{\tau} \right) \right\}^{4i-1} = 2^{-N/2} \left\{ \tau^{2i} \text{ch} \left(Q_1(E), g^{Q_1(E)}, d, \tau \right) \right\}^{4i-1},$$

$$(4.30) \quad \left\{ \text{ch} \left(Q_3(E), g^{Q_3(E)}, d, -\frac{1}{\tau} \right) \right\}^{4i-1} = \left\{ \tau^{2i} \text{ch} \left(Q_3(E), g^{Q_3(E)}, d, \tau \right) \right\}^{4i-1}.$$

Combining the condition $p_1(V)_{S^1} = 0$ and (3.21)-(3.29), we obtain (4.22). The formulas (4.23) and (4.24) can be verified in a similar way.

For (2), we use the transformation laws of Jacobi theta-functions under the action of T and Proposition 2.2 in [20], which we can easily verify to get (4.25), (4.26) and (4.27). □

Lemma 4.5. *For any function*

$$\mathcal{F}_\lambda, \quad \lambda = \{1, 2, 3\}$$

its modular transformation is holomorphic in $(t, \tau) \in \mathbf{R} \times \mathbf{H}$.

Proof. The proof is almost the same as the proof of [12, Lemma 2.3] except that we use Proposition 4.2 instead of the corresponding Lefschetz fixed point formulas. \square

Proof of Theorem 4.1. We prove that these \mathcal{F}_λ are holomorphic on $\mathbf{C} \times \mathbf{H}$ which implies the rigidity of Theorem 4.1. We denote by \mathcal{F} one of the functions: $\{\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3\}$. By (4.19), (4.20) and (4.21), we see that the possible poles of $\mathcal{F}(t, \tau)$ can be written in the form $t = \frac{k}{l}(c\tau + d)$ for integers k, l, c, d with $(c, d) = 1$.

We can always find integers a, b such that $ad - bc = 1$. Then the matrix $g_1 = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \in SL_2(\mathbb{Z})$ induces an action

$$\mathcal{F}(g_1(t, \tau)) = \mathcal{F}\left(\frac{t}{-c\tau + a}, \frac{d\tau - b}{-c\tau + a}\right).$$

Now, if $t = \frac{k}{l}(c\tau + d)$ is a polar divisor of $\mathcal{F}(t, \tau)$, then one polar divisor of $\mathcal{F}(g_1(t, \tau))$ is given by

$$\frac{t}{-c\tau + a} = \frac{k}{l} \left(c \frac{d\tau - b}{-c\tau + a} + d \right)$$

which gives $t = \frac{k}{l}$.

By Lemma 4.4, we know that up to some constant, $\mathcal{F}(g_1(t, \tau))$ is still one of $\{\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3\}$. This contradicts Lemma 4.5, therefore, this completes the proof of Theorem 4.1.

Similarly, use Liu's method [12]. In the odd-dimensional case, we can get the following theorem

Theorem 4.6. *For an odd dimensional connected spin manifold with non-trivial S^1 -action.*

(1) *If $3p_1(V)_{S^1} = 0$ and $c_3(E, g, d) = 0$, then the Toeplitz operators $\mathcal{T} \otimes \Phi \otimes (Q_1(E), g^{Q_1(E)})$, $\mathcal{T} \otimes \Phi \otimes (Q_2(E), g^{Q_2(E)})$, $\mathcal{T} \otimes \Phi \otimes (Q_3(E), g^{Q_3(E)})$ and $\mathcal{T} \otimes \Phi \otimes (Q(E), g^{Q(E)})$ are rigid.*

(2) *If $p_1(V)_{S^1} = 0$ and $c_3(E, g, d) = 0$, then the Toeplitz operators $\mathcal{T} \otimes \Psi_i \otimes (Q(E), g^{Q(E)})$ are rigid. Where $Q(E) = Q_1(E) \otimes Q_2(E) \otimes Q_3(E)$.*

5. ACKNOWLEDGEMENTS

The second author was supported in part by NSFC No.11771070. The authors also thank the referee for his (or her) careful reading and helpful comments.

6. DATA AVAILABILITY

No data was gathered for this article.

7. CONFLICT OF INTEREST

The authors have no relevant financial or non-financial interests to disclose.

REFERENCES

- [1] K. Chandrasekharan, *Elliptic Functions*, Springer-Verlag, 1985.
- [2] E. Witten, Elliptic genera and quantum field theory. *Comm. Math. Phys.* 109(1987), 525-536.
- [3] E. Witten, The index of the Dirac operator in loop space. In: *Elliptic Curves and Modular Forms in Algebraic Topology* (Princeton, NJ, 1986), Lecture Notes in Math., vol. 1326, pp. 161-181. Springer, Berlin (1988).
- [4] S. Ochanine, Sur les genres multiplicatifs définis par des intégrales elliptiques. *Topology* 26(1987), 143-151.
- [5] P. S. Landweber, R. E. Stong, Circle actions on Spin manifolds and characteristic numbers. *Topology* 27(1988), 145-161.
- [6] F. Hirzebruch, Elliptic genera of level N for complex manifolds. In: *Differential Geometrical Methods in Theoretical Physics* (Como, 1987), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. 250, pp. 37-63. Kluwer Acad. Publ., Dordrecht (1988).
- [7] R. Bott, C. Taubes, On the rigidity theorems of Witten. *J. Amer. Math. Soc.* 2 (1989), 137-186.
- [8] C. H. Taubes, S^1 -actions and elliptic genera. *Comm. Math. Phys.* 122(1989), 455-526.
- [9] I. M. Krichever, Generalized elliptic genera and Baker-Akhiezer functions. *Mat. Zametki* 47(1990), 34-45.
- [10] A. Dessai, Rigidity theorems for Spinc-manifolds. *Topology* 39(2000), 239-258.
- [11] K. Liu, Modular invariance and characteristic numbers. *Comm. Math. Phys.* 174 (1995), 29-42.
- [12] K. Liu, On modular invariance and rigidity theorems. *J. Differential Geom.* 41(1995), 343-396.
- [13] K. Liu, On elliptic genera and theta-functions. *Topology* 35 (1996), 617-640.
- [14] K. Liu, X. Ma, On family rigidity theorems. I. *Duke Math. J.* 102(2000), 451-474.
- [15] K. Liu, X. Ma, On family rigidity theorems for Spinc manifolds. In: *Mirror Symmetry, IV* (Montreal, QC, 2000), AMS/IP Stud. Adv. Math., vol. 33, pp. 343-360. Amer. Math. Soc., Providence, RI (2002).
- [16] K. Liu, X. Ma, W. Zhang, Spinc manifolds and rigidity theorems in K-theory. *Asian J. Math.* 4(2000), 933-959.
- [17] K. Liu, X. Ma, W. Zhang, Rigidity and vanishing theorems in K-theory. *Comm. Anal. Geom.* 11(2003), 121-180.
- [18] K. Liu, Y. Wang, Rigidity theorems on odd dimensional manifolds. *Pure Appl. Math. Q.* 5(3, Special Issue: In honor of Friedrich Hirzebruch. Part 2), 1139-1159 (2009).
- [19] W. Zhang, *Lectures on Chern-weil Theory and Witten Deformations*. Nankai Tracks in Mathematics Vol. 4, World Scientific, Singapore, 2001.
- [20] F. Han, J. Yu, On the Witten rigidity theorem for odd dimensional manifolds, arXiv:1504.03007, 2015.
- [21] Y. Wang, $SL(2, \mathbf{Z})$ modular forms and anomaly cancellation formulas, arXiv preprint arXiv:2304.01458, 2023.
- [22] Y. Wang, J. Guan, $SL(2, \mathbf{Z})$ Modular Forms and Anomaly Cancellation Formulas, arXiv preprint arXiv:2304.01458, 2023.
- [23] J. Guan, Y. Wang, $SL(2, \mathbf{Z})$ Modular Forms and Anomaly Cancellation Formulas II, arXiv preprint arXiv:2309.11833, 2023.
- [24] J. Guan, Y. Wang, and H. Liu, $SL(2, \mathbf{Z})$ modular forms and Witten genus in odd dimensions, arXiv preprint arXiv: arXiv:2401.06944.

J. Guan

School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin, 130024, China

E-mail: *guanjy@nenu.edu.cn*

K. Liu

Mathematical Science Research Center, Chongqing University of Technology, Chongqing,
400054, China

E-mail: *kefeng@cqut.edu.cn*

Y. Wang

School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin,
130024, China

E-mail: *wangy581@nenu.edu.cn*