# Lie Symmetries for the Shallow Water Magnetohydrodynamics Equations in a Rotating Reference Frame

Andronikos Paliathanasis<sup>1,2,3</sup>\* and Amlan Halder<sup>3</sup>

<sup>1</sup> Institute of Systems Science, Durban University of Technology, PO Box 1334, Durban 4000, Republic of South Africa

 $2$  Departamento de Matemáticas, Universidad Católica del Norte,

Avda. Angamos 0610, Casilla 1280 Antofagasta, Chile

<sup>3</sup> School of Sciences, Woxsen University,

Hyderabad 502345, Telangana, India

December 20, 2024

#### Abstract

We perform a detailed Lie symmetry analysis for the hyperbolic system of partial differential equations that describe the one-dimensional Shallow Water magnetohydrodynamics equations within a rotating reference frame. We consider a relaxing condition  $\nabla (h\mathbf{B}) \neq 0$  for the one-dimensional problem, which has been used to overcome unphysical behaviors. The hyperbolic system of partial differential equations depends on two parameters: the constant gravitational potential  $g$  and the Coriolis term  $f_0$ , related to the constant rotation of the reference frame. For four different cases, namely  $g = 0$ ,  $f_0 = 0$ ;  $g \neq 0$ ,  $f_0 = 0$ ;  $g = 0$ ,  $f_0 \neq 0$ ; and  $g \neq 0$ ,  $f_0 \neq 0$  the admitted Lie symmetries for the hyperbolic system form different Lie algebras. Specifically the admitted Lie algebras are the  $L^{10} = \{A_{3,3} \rtimes A_{2,1}\} \otimes_s A_{5,34}^a$ ;  $L^8 = A_{2,1} \rtimes A_{6,22}$ ;  $L^7 = A_{3,5} \rtimes \{A_{2,1} \rtimes A_{2,1}\}$ ; and

<sup>∗</sup>Email: anpaliat@phys.uoa.gr

 $L^6 = A_{3,5} \rtimes A_{3,3}$  respectively, where we use the Morozov-Mubarakzyanov-Patera classification scheme. For the general case where  $f_0g \neq 0$ , we derive all the invariants for the Adjoint action of the Lie algebra  $L^6$  and its subalgebras, and we calculate all the elements of the one-dimensional optimal system. These elements are then considered to define similarity transformations and construct analytic solutions for the hyperbolic system.

Keywords: Lie symmetries; shallow water magnetohydrodynamics; one-dimensional optimal system; similarity transformations

### 1 Introduction

Saint-Venant in [\[1\]](#page-26-0) introduced a set of partial differential equations to describe the flow of a thin layer of inviscid fluid. These equations form a hyperbolic system known as the Shallow-Water (SW) equations. The SW equations can model water flow in rivers, lakes, and other physical systems where the flow creates a thin layer, allowing the vertical fluid movements to be neglected. For various applications of the SW equations, we refer the reader to [\[2–](#page-26-1)[7\]](#page-27-0) and the references therein. In an ideal physical system, the SW equations describe the conservation of mass and momentum. In a real physical system, gravitational force terms are introduced in the momentum equation, along with the Coriolis force, friction, and other effects [\[3\]](#page-26-2). On the other hand, in multi-fluid flows with nonzero interaction and energy transfer between the two fluids, the equation of motion for the mass is modified [\[8,](#page-27-1)9].

The Magnetohydrodynamic Shallow-Water system (SWMHD) was introduced to describe the solar tachocline [\[10\]](#page-27-3), a thin layer in the Sun that marks the transition in angular velocity behavior between the radiative zone and the convection zone [\[11\]](#page-27-4). The tachocline is believed to play an important role in the evolution of solar behavior and structure. Moreover, it has been proposed as the main mechanism for explaining the solar dynamo; for more details, see [\[12\]](#page-27-5). In [\[13\]](#page-27-6), it was shown that SW waves in the tachocline lead to a dispersion equation that can describe Alfven waves and magneto-gravity waves [\[13\]](#page-27-6). The Coriolis term in the rotating reference frame results in waves that do not change shape, while in the absence of the Coriolis force, singularities in the waves appear. In [\[14\]](#page-27-7), slow and fast magnetic Rossby waves were identified for the tachocline. The SWMHD system has been applied to various physical systems beyond the tachocline [\[15–](#page-27-8)[17\]](#page-28-0).

In this study, we focus on the algebraic properties of the SWMHD system within a rotating reference frame and under a constant gravitational force. Specifically, we apply Lie symmetry analysis [\[18](#page-28-1)[–21\]](#page-28-2) to determine all the point transformations that leave the SWMHD system invariant. Symmetry analysis is a crucial approach in the study of nonlinear differential equations [\[22–](#page-28-3)[27\]](#page-28-4) and has various applications in SW models [\[28,](#page-28-5) [29,](#page-28-6) [31, 31–](#page-29-0)[35\]](#page-29-1) as well as MHD equations [\[36–](#page-29-2)[42\]](#page-30-0). The Lie symmetry analysis for the one-dimensional SWMHD equations was recently presented in [\[43\]](#page-30-1). In [\[43\]](#page-30-1), a non-rotating reference frame was assumed, and Gauss' law for the magnetic field was imposed. However, as demonstrated, this requirement for the one-dimensional system can lead to unphysical results, and this constraint can be relaxed when studying the one-dimensional system [\[44–](#page-30-2)[47\]](#page-30-3). The inclusion of this term restores Galilean invariance for the SWMHD equations.

The structure of the paper is as follows:

In Section [2,](#page-2-0) we present the basic definitions and properties of Lie symmetry analysis. In particular, we introduce the constraint equation, where a given differential equation remains invariant under the action of an infinitesimal one-parameter point transformation. We discuss the main application of Lie symmetry analysis, which is the concept of similarity transformations, and how these transformations can be applied to reduce a differential equation. Furthermore, we outline the main steps for calculating the one-dimensional system for the admitted Lie symmetries of the differential equation.

In Section [3,](#page-5-0) we introduce the SWMHD system in the presence of a gravitational field and in a rotating reference frame, which leads to the appearance of a Coriolis force. Since we focus on the one-dimensional system, we consider a set of five hyperbolic partial differential equations. We also consider a system with a relaxed condition for Gauss' law for the magnetic field, as this leads to physically acceptable solutions for the one-dimensional system.

The Lie symmetry classification problem for the SWMHD equations is presented in Section [4.](#page-6-0) Additionally, in Sections [5](#page-15-0) and [6,](#page-17-0) we present the derivation of the one-dimensional optimal system and its application in constructing similarity solutions, respectively. Finally, in Section [7,](#page-24-0) we provide our conclusions.

### <span id="page-2-0"></span>2 Preliminaries

In this section, we introduce the fundamental concepts and properties of Lie symmetry analysis for differential equations. We explore the use of similarity transformations and their application in deriving solutions. Finally, we define the one-dimensional optimal system.

#### 2.1 Lie symmetries

Consider the independent variables  $y^i$  and the dependent variables  $\Psi^A$ , which are defined in the jet space  $B = B(y^i, \Psi^A, \Psi^A_{i,i}, ...)$ . We note that in this work, we use Einstein's summation convention.

Let the function  $\mathbf{H} = \mathbf{H}(y^i, \Psi^A, \Psi^A, \Psi^A)$  describe a system of first-order differential equations. In this study, we focus on the case of first-order differential equations, but the following analysis can also be extended to higher-dimensional differential equations.

Under the application of the one-parameter infinitesimal point transformation

<span id="page-3-1"></span><span id="page-3-0"></span>
$$
\bar{y}^i = y^i + \varepsilon \xi^i(y^k, \Psi^B) \tag{1}
$$

$$
\bar{\Psi}^A = \Psi^A + \varepsilon \eta^A (y^k, \Psi^B) \tag{2}
$$

with  $\varepsilon^2 \to 0$ , the set of differential equations **H**, becomes

$$
\bar{\mathbf{H}} = \bar{\mathbf{H}} \left( \bar{y}^i \left( \varepsilon, y^k, \Psi^B \right), \bar{\Psi}^A \left( \varepsilon, y^k, \Psi^B \right), \bar{\Psi}^A_{,i} \left( \varepsilon, y^k, \Psi^B \right) \right). \tag{3}
$$

Thus, the set of differential equations H remain invariant under the application of the infinitesimal transformation  $(1)$ ,  $(2)$  if and only if  $[19, 20]$  $[19, 20]$ 

$$
\bar{\mathbf{H}}\left(\bar{y}^{i}\left(\varepsilon,y^{k},\Psi^{B}\right),\bar{\Psi}^{A}\left(\varepsilon,y^{k},\Psi^{B}\right),\bar{\Psi}_{,i}^{A}\left(\varepsilon,y^{k},\Psi^{B}\right)\right) = \mathbf{H}(y^{i},\Psi^{A},\Psi_{,i}^{A}),\tag{4}
$$

that is [\[19,](#page-28-7) [20\]](#page-28-8)

<span id="page-3-2"></span>
$$
\lim_{\varepsilon \to 0} \frac{\bar{\mathbf{H}}\left(\bar{y}^i\left(\varepsilon, y^k, \Psi^B\right), \bar{\Psi}^A\left(\varepsilon, y^k, \Psi^B\right), \bar{\Psi}^A_{,i}\left(\varepsilon, y^k, \Psi^B\right)\right) - \mathbf{H}(y^i, \Psi^A, \Psi^A_{,i})}{\varepsilon} = 0.
$$
 (5)

When the latter condition is true we shall say that the generator

$$
\mathbf{X} = \frac{\partial \bar{y}^i}{\partial \varepsilon} \partial_i + \frac{\partial \bar{\Psi}^A}{\partial \varepsilon} \partial_A,\tag{6}
$$

of the infinitesimal transformation  $(1)-(2)$  $(1)-(2)$  $(1)-(2)$  is a Lie symmetry vector for the system of differential equations H.

The symmetry condition [\(5\)](#page-3-2) can be written in the equivalent form [\[19,](#page-28-7) [20\]](#page-28-8)

$$
\mathbf{X}^{[1]} \left( \mathbf{H} \right) = 0,\tag{7}
$$

or

<span id="page-3-3"></span>
$$
\mathbf{X}^{[1]}(\mathbf{H}) = \lambda \mathbf{H} , \quad \text{mod } \mathbf{H} = 0,
$$
\n(8)

in which  $X^{[1]}$  is the first extension/prolongation vector of X in the jet space, and  $\lambda$  is a function which should derived.

Condition [\(8\)](#page-3-3) is referred to as the Lie symmetry condition, which leads to a system of linear differential equations for the coefficient functions  $\xi^{i}(y^{k}, \Psi^{B})$  and  $\eta^{A}(y^{k}, \Psi^{B})$  of the generator vector X.

The prolongation vector  $X^{[1]}$  is given by the following expression

$$
\mathbf{X}^{[1]} = \mathbf{X} + \eta^{[1]A} \left( x^k, \Psi^B, \Psi^B_{,i} \right) \partial_{u^A_{,i}},\tag{9}
$$

with

$$
\eta^{[1]A}\left(x^{k}, \Psi^{B}, \Psi^{B}_{,i}\right) = D_{i}\left(\eta^{A}\right) - \Psi^{A}_{,j} D_{i} \xi^{j},\tag{10}
$$

in which  $D_i$  is the total derivative operator, that is,

$$
D_i = \frac{\partial}{\partial x^i} + \Psi^B_{,i} \frac{\partial}{\partial \Psi^B} + \dots \,. \tag{11}
$$

Therefore,  $\eta^{[1]}$  is expressed as

$$
\eta_{[i]}^{A} = \eta_{,i}^{A} + \Psi_{,i}^{B} \eta_{,B}^{A} - \xi_{,i}^{j} \Psi_{,j}^{A} - \Psi_{,i}^{A} \Psi_{,j}^{B} \xi_{,B}^{j} . \tag{12}
$$

#### 2.2 Similarity transformation

Assume now that  $X = \xi^i(y^k, \eta^B) \partial_i$  is a Lie symmetry vector for the system of differential equations  $H = H(y^i, \Psi^A, \Psi^A_i)$ . Then, there exist a set of variables  $y^i \to z^{\beta}$ , so that the Lie symmetry vector to read [\[19,](#page-28-7) [20\]](#page-28-8)

$$
\mathbf{X} = \partial_{\beta},\tag{13}
$$

which are known as normal coordinates. Hence the set of differential equations in the new variables are defined as

$$
\mathbf{H} = \mathbf{H}(z^{\beta}, \Psi^{A}, \Psi^{A}_{,\beta}).
$$
\n(14)

This change of variables is known as a similarity transformation, which, in the case of partial differential equations, is applied to reduce the number of independent variables.

In essence, the requirement for a similarity transformation implies that the solution of the differential equations remains invariant along the surface defined by the Lie symmetry vector.

If  $\Psi^A(y^k)$  is a solution for the differential equation, then  $D_i\Psi^A$ , is also a solution. Moreover, if  $X = \xi^i \partial_i + \eta^A \partial_A$  is a Lie symmetry for the set of differential equations H, then by definition if  $\Psi^A(y^k)$  is a solution, the  $X(\Psi^A(y^k))$  remain a solution, i.e.  $X(\Psi^A(y^k)) = \bar{\Psi}^A(y^k)$ , where  $\bar{\Psi}^A(y^k)$  is also a solution.

Therefore, the operator  $X' = X - f^i D_i$  is also a symmetry vector, where  $f^k$  is an arbitrary function. Thus, for  $f^k = \xi^k$  the vector field X' is simplified as follows [\[19,](#page-28-7)20]

$$
\mathbf{X}' = \left(\eta^A - \xi^k \Psi_{,k}\right) \partial_A. \tag{15}
$$

Therefore, we can define the operator

$$
\eta^A - \xi^k \Psi_{,k} = a_0 \Psi. \tag{16}
$$

This condition results in the derivation of the similarity transformation associated with the vector field  $X$ . The two approaches are equivalent, meaning that for a given symmetry vector, the corresponding similarity transformation is unique.

However, not all symmetry vectors lead to similarity transformations that describe independent solutions. In the following, we define the one-dimensional optimal system, which is essential for classifying all independent solutions that arise from the application of similarity transformations. For the derivation of the one-optimal system for a given algebra we refer the reader to [\[21\]](#page-28-2)

## <span id="page-5-0"></span>3 Shallow Water Magnetohydrodynamics

The equations of SWMHD for a hyperbolic system of partial differential equations, that is  $[10, 49, 50],$  $[10, 49, 50],$  $[10, 49, 50],$  $[10, 49, 50],$ 

<span id="page-5-1"></span>
$$
h_{,t} + \nabla (h\mathbf{u}) = 0, \qquad (17)
$$

$$
(h\mathbf{u})_{,t} + \nabla \left(h\mathbf{u}\mathbf{u} - h\mathbf{B}\mathbf{B} + \frac{g}{2}h^2\right) + 2\mathbf{\Omega} \times h\mathbf{u} = 0, \qquad (18)
$$

$$
(h\mathbf{B})_{,t} + \nabla (h\mathbf{B}\mathbf{u} - h\mathbf{u}\mathbf{B}) + \mathbf{u}\nabla (h\mathbf{B}) = 0, \qquad (19)
$$

where  $h$  is the layer depth of the incompressible, perfectly conducting fluid with an equation of state parameter  $p=\frac{1}{2}$  $\frac{1}{2}gh^2$ . The vector field **u** represents the horizontal component of the velocity, **B** is the magnetic field, q is the gravitational constant, and  $\Omega$ , originating from the rotating frame, provides the Coriolis force term.

From Maxwell's equations, we obtain the constraint  $\nabla (h\mathbf{B}) = 0$ . However, enforcing this constraint leads to degenerate numerical solutions for the Riemann problem with one-dimensional initial conditions [\[48,](#page-30-6)[51](#page-30-7)[–53\]](#page-30-8). To address this issue, a relaxed condition  $\nabla (h\mathbf{B}) \neq 0$ has been introduced in several studies for the one-dimensional problem. This modification helps restore the Galilean invariance of the equations [\[54\]](#page-31-0) and can lead to physically acceptable solutions for the one-dimensional Riemann problem [\[52\]](#page-30-9). For more details, we refer the reader to the discussions in [\[55,](#page-31-1) [56\]](#page-31-2).

Assuming that the dependent variables are functions of time  $t$  and the spatial coordinate x, the components of the horizontal velocity are  $\mathbf{u} = (u(t, x), v(t, x))^T$ , and the components of the magnetic field are  $\mathbf{B} = (a(t, x), b(t, x))^T$ .

From [\(17\)](#page-5-1), [\(18\)](#page-5-1), and [\(19\)](#page-5-1), we obtain the hyperbolic system of five partial differential equations [\[56\]](#page-31-2).

<span id="page-6-1"></span>
$$
h_{,t} + (hu)_{,x} = 0, \t\t(20)
$$

$$
(hu)_{,t} + \left(hu^2 + \frac{g}{2}h^2 - ha^2\right)_{,x} + f_0hv = 0,\tag{21}
$$

$$
(hv)_{,t} + (huv - hab)_{,x} - f_0hu = 0,
$$
\n(22)

$$
(ha)_{,t} + u (ha)_{,x} = 0,
$$
\n(23)

$$
(hb)_{,t} + (h (ub - va))_{,x} + v (ha)_{,x} = 0,
$$
\n(24)

where  $\Omega = -\frac{1}{2}$  $rac{1}{2}f_0$ .

In the following we proceed with the symmetry analysis for the hyperbolic system [\(20\)](#page-6-1),  $(21), (22), (23)$  $(21), (22), (23)$  $(21), (22), (23)$  $(21), (22), (23)$  $(21), (22), (23)$  and  $(24).$  $(24).$ 

We consider the infinitesimal one-parameter point transformation

$$
\bar{t} = t + \varepsilon \xi^{t} (t, x, h, \mathbf{u}, \mathbf{B}), \quad \bar{x} = x + \varepsilon \xi^{x} (t, x, h, \mathbf{u}, \mathbf{B}), \tag{25}
$$

$$
\bar{u} = u + \varepsilon \eta^{u} (t, x, h, \mathbf{u}, \mathbf{B}) , \quad \bar{v} = v + \varepsilon \eta^{v} (t, x, h, \mathbf{u}, \mathbf{B}) , \tag{26}
$$

$$
\bar{a} = a + \varepsilon \eta^{a} (t, x, h, \mathbf{u}, \mathbf{B}), \bar{b} = b + \varepsilon \eta^{b} (t, x, h, \mathbf{u}, \mathbf{B}), \qquad (27)
$$

$$
\bar{h} = h + \varepsilon \eta^h(t, x, h, \mathbf{u}, \mathbf{B}), \qquad (28)
$$

with generator the vector field

$$
X = \xi^t \partial_t + \xi^x \partial_x + \eta^h \partial_h + \eta^u \partial_u + \eta^v \partial_v + \eta^a \partial_a + \eta^b \partial_b.
$$
 (29)

The Lie symmetry condition results in a set of constraint equations for the coefficients of the vector field X. The constraint system and the admitted Lie symmetries depend on the presence of the gravitational field and the Coriolis force. We examine all possible cases separately.

# <span id="page-6-0"></span>4 Lie symmetry analysis for the SWMHD system

In the following lines we present the Lie symmetry conditions [\(8\)](#page-3-3) for the SWMHD system. The admitted Lie symmetries for the SWMHD system depend on the value of the free parameters  $f_0$ and g. It follows that there are four different cases for the values of the gravitational field and of the Coriolis term. Specifically, for the following four cases (a)  $g = 0$ ,  $f_0 = 0$ , (b) $g \neq 0$ ,  $f_0 = 0$ ,

(c)  $g = 0$ ,  $f_0 \neq 0$  and (d)  $g \neq 0$ ,  $f_0 \neq 0$ , it follows that the admitted Lie symmetries for the SWMHD system [\(20\)](#page-6-1), [\(21\)](#page-6-1), [\(22\)](#page-6-1), [\(23\)](#page-6-1) and [\(24\)](#page-6-1) form different dimensional Lie algebra. The fist two cases, describe nonrotating systems with or without gravitational field, while cases (c) and (d) describe rotating systems with or without gravitational field.

#### 4.1 Free system

Let us now assume that the SWMHD equations are free, that is, there is not a gravitational field, i.e.  $g = 0$ , and there is not any Coriolis force term, i.e.  $f_0 = 0$ . Therefore, the hyperbolic system describes the SWMHD equations is simplified as follows

$$
h_{,t} + (hu)_{,x} = 0, \t\t(30)
$$

$$
(hu)_{,t} + (hu^2 - ha^2)_{,x} = 0,\t\t(31)
$$

$$
(hv)_{,t} + (huv - hab)_{,x} = 0,
$$
\n
$$
(32)
$$

$$
(ha)_{,t} + u(ha)_{,x} = 0,
$$
\n(33)

$$
(hb)_{,t} + (h (ub - va))_{,x} + v (ha)_{,x} = 0,
$$
\n(34)

The Lie symmetry conditions for this system are

$$
\xi^{t} = \xi^{t}(t), \ \xi^{x} = \xi^{x}(t,x), \ \eta^{h} = \eta^{h}(h),
$$
  

$$
\eta^{u} = (\xi^{x}_{,x} - \xi^{t}_{,t}) u + \xi^{x}_{,t}, \ \eta^{v} = \eta^{v}(u,v), \tag{35}
$$

$$
\eta^{a} = -a \left( \xi_{,t}^{t} - \xi_{,x}^{x} \right), \ \eta^{b} = \eta^{b} \left( h, a, b \right), \tag{36}
$$

$$
\xi_{,tt}^t = 0, \ \xi_{,tt}^x = 0, \ \xi_{,xx}^x = 0, \ \xi_{,tx}^x = 0,\tag{37}
$$

$$
\eta_{,uu}^v = 0, \ \eta_{,vv}^v = 0, \ \eta_{,uv}^v = 0,\tag{38}
$$

$$
h\eta_{,h}^{h} - \eta_{,h}^{h} = 0 \ , a\eta_{,a}^{h} - \left(2a\eta_{,u}^{v} - \eta_{,h}^{h} + \eta_{,v}^{v}b\right) = 0, \tag{39}
$$

$$
\eta_{,b}^{b} - \eta_{,v}^{v} = 0, \ h\eta_{,h}^{b} - \left(a\eta_{,u}^{v} - \eta^{b} + \eta_{,v}^{v}b\right) = 0. \tag{40}
$$

The solution of this system provides the generic generator  $X$  for the infinitesimal transformations that leave the SWMHD equations invariant. The arbitrary constants in the generic vector field  $\bf{X}$  determine the specific admitted Lie symmetries for the hyperbolic system.

#### 4.1.1 Lie symmetries

The solution of the symmetry conditions for the free SWMHD hyperbolic equations provide a ten-dimensional Lie algebra  $L^{10}$  consisted by the vector fields

$$
X_1 = \partial_t, X_2 = \partial_x, X_3 = t\partial_t + x\partial_x, X_4 = h\partial_h, X_5 = t\partial_x + \partial_u, X_6 = \partial_v,
$$
  

$$
X_7 = u\partial_v + a\partial_b, X_8 = v\partial_v + b\partial_b, X_9 = t\partial_t - (u\partial_u + a\partial_a), X_{10} = \frac{1}{ah}\partial_b.
$$

where we see that the Galileon symmetries are preserved to the system.

For the admitted vector fields we can define the corresponding one-parameter point transformations, indeed we derive

$$
X_1 : \overline{t} = t + \varepsilon_1,
$$
  
\n
$$
X_2 : \overline{x} = x + \varepsilon_2,
$$
  
\n
$$
X_3 : \overline{t} = e^{\varepsilon_3}t, \overline{x} = e^{\varepsilon_3}x,
$$
  
\n
$$
X_4 : \overline{h} = e^{\varepsilon_4}h,
$$
  
\n
$$
X_5 : \overline{t} = e^{\varepsilon_5}t, \overline{u} = e^{\varepsilon_5}u,
$$
  
\n
$$
X_6 : \overline{v} = v + \varepsilon
$$
  
\n
$$
X_7 : \overline{v} = v + \varepsilon_7u, \overline{b} = b + \varepsilon_7a,
$$
  
\n
$$
X_8 : \overline{v} = e^{\varepsilon_8}v, \overline{b} = e^{\varepsilon_8}b,
$$
  
\n
$$
X_9 : \overline{t} = e^{\varepsilon_9}t, \overline{u} = e^{-\varepsilon_9}u, \overline{a} = e^{-\varepsilon_9}a
$$
  
\n
$$
X_{10} : \overline{b} = b + \varepsilon_{10}ah.
$$

In order to understand the algebraic properties of the admitted symmetries we derive the commutators and the adjoint representation  $Ad(e^{(\varepsilon \mathbf{X}_i)})$ . W remark that the Adjoint operator is defined by the Lie bracket [, ] as follows

$$
Ad\left(\exp\left(\epsilon X_A\right)\right)X_B = X_B - \epsilon \left[X_A, X_B\right] + \frac{1}{2}\epsilon^2 \left[X_A, \left[X_A, X_B\right]\right] + \dots \tag{41}
$$

The one-dimensional subalgebras of the admitted Lie symmetries which are not related through the adjoint representation form the one-dimensional optimal system. The determination of the one-dimensional system is essential in order to perform a complete classification of all the possible similarity transformations and solutions.

The commutators for the ten-dimensional Lie algebra  $L^{10}$  are presented in Table [1,](#page-9-0) while the adjoint presentation table is given in Tables [2](#page-9-1) and [3.](#page-10-0) Therefore from the commutators we infer that the admitted  $L^{10}$  Lie Algebra is the  $\{A_{3,3} \rtimes A_{2,1}\} \rtimes A_{5,34}^a$  in the Morozov-Mubarakzyanov-Patera classification scheme [\[57–](#page-31-3)[61\]](#page-31-4).

$[\mathbf{X}_i, \mathbf{X}_j]$	$\mathbf{X}_1$			$\mathbf{X}_2$ $\mathbf{X}_3$ $\mathbf{X}_4$	$\mathbf{X}_5$	$\mathbf{X}_6$	$\mathbf{X}_7$	$\mathbf{X}_8$	$\mathbf{X}_9$	$\mathbf{X}_{10}$
$\mathbf{X}_1$	$\theta$	$\vert 0 \vert$	$X_1$	$\overline{0}$	$X_2$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$X_1$	$\overline{0}$
$\mathbf{X}_2$	$\overline{0}$	$\vert 0 \vert$	$X_2$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\theta$	$\theta$	$\overline{0}$
$\mathbf{X}_3$		$-X_1$ $-X_2$ 0		$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\boldsymbol{0}$	$\theta$	$\theta$
$\mathbf{X}_4$	$\theta$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\theta$	$\begin{matrix} 0 \end{matrix}$	$-X_{10}$
$\mathbf{X}_5$	$-X_2$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$X_6$	$\vert 0 \vert$	$-X_5$	$\overline{0}$
$\mathbf{X}_6$	$\overline{0}$	$\overline{0}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\boldsymbol{0}$	$\begin{matrix} 0 \end{matrix}$	$X^6$	$\theta$	$\overline{0}$
$\mathbf{X}_7$	$\overline{0}$	$\overline{0}$	$\overline{0}$		0 $-X_6$	$\overline{0}$	$\overline{0}$	$X^7$	$X^7$	$\overline{0}$
$\mathbf{X}_8$	$\theta$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$-X^6$ $-X^7$		$\overline{0}$	$\overline{0}$	$-X_{10}$
$\mathbf{X}_9$	$-X_1$	$\overline{0}$	$\overline{0}$	$\overline{0}$		$X_5$ 0 $-X^7$		$\overline{0}$	$\theta$	$X_{10}$
$\mathbf{X}_{10}$	$\theta$	$\theta$	$\begin{matrix} 0 \end{matrix}$	$X_{10}$	$\overline{0}$	$\overline{0}$	$\begin{matrix} 0 \end{matrix}$		$X_{10}$ $-X_{10}$	$\overline{0}$

<span id="page-9-0"></span>Table 1: Commutators of the ten admitted Lie symmetries for the free SWMHD hyperbolic system.

Table 2: Adjoint representation of the ten admitted Lie symmetries for the free SWMHD hyperbolic <u>system  $(1/2)$ .</u>

<span id="page-9-1"></span>

$Ad\left(e^{(\varepsilon \mathbf{X}_i)}\right) \mathbf{X}_j$	$\mathbf{X}_1$	$\mathbf{X}_2$	$\mathbf{X}_3$	$\mathbf{X}_4$	$\mathbf{X}_5$
$\mathbf{X}_1$	$X_1$	$X_2$	$X_3-\varepsilon X_1$	$X_4$	$X_5 - 2X_2$
$\mathbf{X}_2$	$X_1$	$X_2$	$X_3-\varepsilon X_2$	$X_4$	$X_5$
$\mathbf{X}_3$	$e^{\varepsilon}X_1$	$e^{\varepsilon}X_2$	$X_3$	$X_4$	$X_5$
$\mathbf{X}_4$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$
$\mathbf{X}_5$	$X_1+\varepsilon X_2$	$X_2$	$X_3$	$X_4$	$X_5$
$\mathbf{X}_6$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$
$\mathbf{X}_7$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5 + \varepsilon X^6$
$\mathbf{X}_8$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$
$\mathbf{X}_9$	$e^{\varepsilon}X_1$	$X_2$	$X_3$	$X_4$	$e^{-\varepsilon}X_5$
$\mathbf{X}_{10}$	$X_1$	$X_2$	$X_3$	$X_4 - \varepsilon X_{10}$	$X_5$

<span id="page-10-0"></span>

$Ad\left(e^{(\varepsilon{\bf X}_i)}\right){\bf X}_j$	$\mathbf{X}_6$	$\mathbf{X}_7$	$\mathbf{X}_8$	$\mathbf{X}_9$	$\mathbf{X}_{10}$
$\mathbf{X}_1$	$X_6$	$X_7$	$X_8$	$X_9 - \varepsilon X_1$	$X_{10}$
$\mathbf{X}_2$	$X_6$	$X_7$	$X_8$	$X_9$	$X_{10}$
$\mathbf{X}_3$	$X_6$	$X_7$	$X_8$	$X_9$	$X_{10}$
$\mathbf{X}_4$	$X_6$	$X_7$	$X_8$	$X_9$	$e^{\varepsilon}X_{10}$
$\mathbf{X}_5$	$X_6$	$X_7 - \varepsilon X_6$	$X_8$	$X_9+\varepsilon X_5$	$X_{10}$
$\mathbf{X}_6$	$X_6$	$X_7$	$X_8 - \varepsilon X_6$	$X_9$	$X_{10}$
$\mathbf{X}_7$	$X_6$	$X_7$	$X_8 - \varepsilon X_7$	$X_9 - \varepsilon X_7$	$X_{10}$
$\mathbf{X}_8$	$e^{\varepsilon}X_6$	$e^{\varepsilon}X_7$	$X_8$	$X_9$	$e^{\varepsilon}X_{10}$
$\mathbf{X}_9$	$X_6$	$e^{\varepsilon}X_7$	$X_8$	$X_9$	$e^{-\varepsilon}X_{10}$
$\mathbf{X}_{10}$	$X_6$	$X_7$		$X_8 - \varepsilon X_{10}$ $X_9 + \varepsilon X_{10}$	$X_{10}$

Table 3: Adjoint representation of the ten admitted Lie symmetries for the free SWMHD hyperbolic system  $(2/2)$ .

### 4.2 Constant gravitational field

We continue our discussion by introducing a constant gravitational field in non-rotating frame, that is  $f_0 = 0$ . Hence, the SWMHD equations take the following form

$$
h_{,t} + (hu)_{,x} = 0, \t\t(42)
$$

$$
(hu)_{,t} + (hu^2 - ha^2)_{,x} + ghh_{,x} = 0,
$$
\n(43)

$$
(hv)_{,t} + (huv - hab)_{,x} = 0,
$$
\n
$$
(44)
$$

$$
(ha)_{,t} + u (ha)_{,x} = 0,
$$
\n(45)

$$
(hb)_{,t} + (h (ub - va))_{,x} + v (ha)_{,x} = 0.
$$
 (46)

Hence, the Lie symmetry condition leads to the following set of constraint equations

$$
\xi^{t} = \xi^{t}(t), \ \xi^{x} = \xi^{x}(t,x), \ \eta^{h} = 2h\left(\xi_{,x}^{x} - \xi_{,t}^{t}\right),\tag{47}
$$

$$
\xi_{,tt}^t = 0, \ \xi_{,tt}^x = 0, \ \xi_{,xx}^x = 0, \ \xi_{,tx}^x = 0,\tag{48}
$$

$$
\eta^{u} = u\left(\xi_{,x}^{x} - \xi_{,t}^{t}\right) + \xi_{,t}^{x}, \ \eta^{a} = a\left(\xi_{,x}^{x} - \xi_{,t}^{t}\right),\tag{49}
$$

$$
a\eta_{,a}^{b} - h\eta_{,h}^{b} = 0, \ b\eta_{,b}^{b} - h\eta_{,h}^{b} - \eta^{b} = 0, \ h\eta_{,hh}^{b} + 2\eta_{,h}^{b} = 0,
$$
\n
$$
(50)
$$

$$
\eta^v = \eta^v(v), \ b\eta^v_{,v} - h\eta^b_{,h} - \eta^b = 0. \tag{51}
$$

$[\mathbf{X}_i,\mathbf{X}_j] = \mathbf{X}_1 = \mathbf{X}_2 = \mathbf{X}_3 - \mathbf{X}_5 = \mathbf{X}_6$							$\mathbf{X}_8$ $\mathbf{X}_{10}$	Y
$\mathbf{X}_1$	$\boldsymbol{0}$				0 $X_1$ $X_2$ 0	$\overline{0}$	$\overline{0}$	$X_1$
$\mathbf{X}_2$	$\overline{0}$				$0 \t X_2 \t 0 \t 0$	$\overline{0}$	$\overline{0}$	$\Omega$
$\mathbf{X}_3$		$-X_1$ $-X_2$ 0 0			$\overline{0}$	$\overline{0}$	$\overline{0}$	$\Omega$
$\mathbf{X}_5$	$-X_2$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$	$-X_5$
$\mathbf{X}_6$	$\boldsymbol{0}$	$\begin{matrix} 0 \end{matrix}$	$\overline{0}$	$\overline{0}$		$0 \t X^6$	$\overline{0}$	$\Omega$
$\mathbf{X}_8$	$\overline{0}$	$\overline{0}$	$\overline{0}$		$0 \quad -X^6$		0 $-X_{10}$	$\theta$
$\mathbf{X}_{10}$	$\boldsymbol{0}$	$\overline{0}$	$\overline{0}$	$\overline{0}$				0 $X_{10}$ 0 $-3X_{10}$
Y	$-X_1$	$\overline{0}$			$0 \quad X_5 \quad 0$	$\overline{0}$	$3X_{10}$	$\overline{0}$

<span id="page-11-0"></span>Table 4: Commutators of the eight admitted Lie symmetries for the SWMHD hyperbolic system in a constant gravitational field.

#### 4.2.1 Lie symmetries

The solution of the Lie symmetry conditions for the SWMHD hyperbolic equations within a constant gravitational field leads to the derivation of eight Lie symmetries with form the  $L^8$ Lie algebra.

The elements of the Lie algebra  $L^8$  are

$$
X_1
$$
,  $X_2$ ,  $X_3$ ,  $X_5$ ,  $X_6$ ,  $X_8$ ,  $X_{10}$ ,  $Y = X^9 - 2X^4$ ,

where the Galileon symmetries are preserved. Moreover, for vector field Y we derive the oneparameter point transformation

$$
Y: \bar{t} = e^{\varepsilon_Y}t, \ \bar{u} = e^{-\varepsilon_Y}u, \ \bar{a} = e^{-\varepsilon_Y}a, \ \bar{h} = e^{-2\varepsilon_4}h.
$$

In Tables [4](#page-11-0) and [5](#page-12-0) we present the commutators and the adjoint representation of the Lie symmetries. Thus, the admitted eight Lie symmetries form the  $A_{2,1} \rtimes A_{6,22}$ , Lie algebra in the Morozov-Mubarakzyanov-Patera classification scheme [\[57–](#page-31-3)[61\]](#page-31-4).

$Ad\left(e^{(\varepsilon {\bf X}_i)}\right) {\bf X}_j$	$\mathbf{X}_1$	$\mathbf{X}_2$	$\mathbf{X}_3$	$\mathbf{X}_5$	$\mathbf{X}_6$	$\mathbf{X}_8$		$X_{10}$ Y
$\mathbf{X}_1$	$X_1$		$X_2$ $X_3 - \varepsilon X_1$ $X_5 - 2X_2$		$X_6$	$X_8$	$X_{10}$	$Y-\varepsilon X_1$
$\mathbf{X}_2$	$X_1$		$X_2$ $X_3 - \varepsilon X_2$	$X_5$	$X_6$	$X_8$	$X_{10}$	Y
$\mathbf{X}_3$			$e^{\varepsilon}X_1$ $e^{\varepsilon}X_2$ $X_3$	$X_5$	$X_6$	$X_8$	$X_{10}$	Y
$\mathbf{X}_5$	$X_1 + \varepsilon X_2$ $X_2$		$X_3$	$X_5$	$X_6$	$X_8$	$X_{10}$	$Y + \varepsilon X_5$
$\mathbf{X}_6$	$X_1$	$X_2$	$X_3$	$X_5$	$X_6$	$X_8 - \varepsilon X_6$ $X_{10}$		Y
$\mathbf{X}_8$	$X_1$	$X_2$	$X_3$	$X_5$	$e^{\varepsilon}X_6$	$X_8$	$e^{\varepsilon}X_{10}$	Y
$\mathbf{X}_{10}$	$X_1$	$X_2$	$X_3$	$X_5$		$X_6$ $X_8 - \varepsilon X_{10}$ $X_{10}$ $Y + 3\varepsilon X_8$		
Y	$e^{\varepsilon}X_1$	$X_2$	$X_3$	$e^{-\varepsilon}X_5$	$X_6$		$X_8$ $e^{-3\varepsilon}X_{10}$ Y	

<span id="page-12-0"></span>Table 5: Adjoint representation of the eight admitted Lie symmetries for the SWMHD hyperbolic system in a constant gravitational field.

### 4.3 Coriolis force without gravitational field

In the absence of the gravitational field, but in the presence of a constant rotational frame the one-dimensional SWMHD equations [\(20\)](#page-6-1), [\(21\)](#page-6-1), [\(22\)](#page-6-1), [\(23\)](#page-6-1) and [\(24\)](#page-6-1) read

$$
h_{,t} + (hu)_{,x} = 0, \t\t(52)
$$

$$
(hu)_{,t} + (hu^2 - ha^2)_{,x} + f_0hv = 0,
$$
\n(53)

$$
(hv)_{,t} + (huv - hab)_{,x} - f_0hu = 0,
$$
\n(54)

$$
(ha)_{,t} + u(ha)_{,x} = 0,
$$
\n(55)

$$
(hb)_{,t} + (h (ub - va))_{,x} + v (ha)_{,x} = 0,
$$
\n(56)

For this hyperbolic system the Lie symmetry condition provides the constraint equations

$$
\xi^{t} = \xi^{t}(t), \ \xi^{x} = \xi^{x}(t,x), \ \eta^{h} = \eta(h), \ \eta^{v} = \eta^{v}(v), \ \eta^{b} = \eta^{b}(a,b,h), \tag{57}
$$

$$
\eta^{a} = a\left(\xi_{,x}^{x} - \xi_{,t}^{t}\right), \ \eta^{h} = 2H\left(\xi_{,x}^{x} - \xi_{,t}^{t}\right), \ \eta^{u} = u\left(\xi_{,x}^{x} - \xi_{,t}^{t}\right) + \xi_{,t}^{x},\tag{58}
$$

$$
\xi_{,tt}^t = 0, \ \xi_{,tt}^x = 0, \ \xi_{,xx}^x = 0, \ \xi_{,tx}^x = 0,\tag{59}
$$

$$
a\eta^{b} - h\eta^{b}_{,h} = 0, \ b\eta^{b}_{,b} - h\eta^{b}_{,h} - \eta^{b} = 0,
$$
\n(60)

$$
h\eta_{,hh}^b + 2\eta_{,h}^b = 0, \ b\eta_{,v}^v - \eta_{,h}^b - \eta^b = 0.
$$
 (61)

<span id="page-13-0"></span>

-0							
$[\mathbf{X}_i,\mathbf{X}_j]$				$\mathbf{X}_1$ $\mathbf{X}_2$ $\mathbf{X}_4$ $\mathbf{X}_{10}$		$\mathbf{Z}_1$ $\mathbf{Z}_2$ $\mathbf{Z}_3$	
$\mathbf{X}_1$	$\theta$	$\begin{matrix} 0 \end{matrix}$	$\overline{0}$	$\begin{matrix} 0 \end{matrix}$			0 $f_0Z_3 - f_0Z_2$
$\mathbf{X}_2$	$\theta$	$\vert 0 \vert$	$\overline{0}$	$\begin{matrix} 0 \end{matrix}$	$X_2$	$\overline{0}$	$\theta$
$\mathbf{X}_4$	$\theta$	$\overline{0}$	$\overline{0}$	$-X_{10}$	$\vert 0 \vert$	$\theta$	$\theta$
$\mathbf{X}_{10}$	$\theta$			0 $X_{10}$ 0 $2X_{10}$		$\overline{0}$	$\theta$
$\mathbf{Z}_1$	$\theta$			$-X_2$ 0 $-2X_{10}$		0 $-Z_2$ $-Z_3$	
${\bf Z}_2$	$-f_0Z_3$ 0		$\overline{0}$	0 $Z_2$		$\theta$	$\theta$
$\mathbf{Z}_3$	$f_0Z_2$ 0		$\theta$	$\boldsymbol{0}$	$Z_3$	$\theta$	$\theta$

Table 6: Commutators of the seven admitted Lie symmetries for the SWMHD hyperbolic system in a rotating frame.

#### 4.3.1 Lie symmetries

The solution of the Lie symmetry conditions leads to the seven dimensional Lie algebra  $L^7$  with elements the vector fields

$$
X_1, X_2, X_4, X_{10}, Z_1 = X_3 + X_8 - X_9,
$$
  
\n
$$
Z_2 = \sin(f_0 t) \partial_x + f_0 (\cos(f_0 t) \partial_u + \sin(f_0 t) \partial_v),
$$
  
\n
$$
Z_3 = \cos(f_0 t) \partial_x + f_0 (-\sin(f_0 t) \partial_u + \cos(f_0 t) \partial_v).
$$

We remark that the Galileon symmetry is lost when the Coriolis term appears in the system.

Thus, for the generator vector fields  $Z_1$ ,  $Z_2$  and  $Z_3$  we calculate the one-parameter point transformation

$$
Z_1 : \bar{x} = e^{\varepsilon_1^Z} x, \ \bar{u} = e^{\varepsilon_1^Z} u, \ \bar{v} = e^{\varepsilon_1^Z} v, \ \bar{a} = e^{\varepsilon_1^Z} a, \ \bar{b} = e^{\varepsilon_1^Z} b,
$$
  
\n
$$
Z_2 : \bar{x} = x + \varepsilon_2^Z \sin(f_0 t), \ \bar{u} = u + \varepsilon_2^Z f_0 \cos(f_0 t), \ \bar{v} = v + \varepsilon_2^Z f_0 \sin(f_0 t),
$$
  
\n
$$
Z_3 : \ \bar{x} = x + \varepsilon_3^Z \cos(f_0 t), \ \bar{u} = u - \varepsilon_3^Z f_0 \sin(f_0 t), \ \bar{v} = v + \varepsilon_3^Z f_0 \cos(f_0 t),
$$

Hence, the commutators and the adjoint representation for the Lie algebra  $L^7$  are presented respectively in Tables [6,](#page-13-0) [7](#page-14-0) and [8.](#page-14-1) Hence, in the Morozov-Mubarakzyanov-Patera classification scheme [\[57–](#page-31-3)[61\]](#page-31-4) it follows that  $L^7$  is the Lie algebra  $A_{3,5} \rtimes \{A_{2,1} \rtimes A_{2,1}\}.$ 

$\sim$	$( - ) - )$			
$Ad\left(e^{\left(\varepsilon{\bf X}_i\right)}\right){\bf X}_j$	$\mathbf{X}_1$	$\mathbf{X}_2$	$\mathbf{X}_4$	$\mathbf{X}_{10}$
$\mathbf{X}_1$	$X_1$	$X_2$	$X_3$	$X_{10}$
$\mathbf{X}_2$	$X_1$	$X_2$	$X_3$	$X_{10}$
$\mathbf{X}_4$	$X_1$	$X_2$	$X_3$	$e^{\varepsilon}X_{10}$
$\mathbf{X}_{10}$	$X_1$	$X_2$	$X_3 - \varepsilon X_{10}$	$X_{10}$
$\mathbf{Z}_1$	$X_1$	$e^{\varepsilon}X_2$	$X_3$	$e^{2\varepsilon}X_{10}$
${\bf Z}_2$	$X_1 + \varepsilon f_0 Z_3$	$X_2$	$X_3$	$X_{10}$
$\mathbf{Z}_3$	$X_1 - \varepsilon f_0 Z_2$	$X_2$	$X_3$	$X_{10}$

<span id="page-14-0"></span>Table 7: Adjoint representation of the seven admitted Lie symmetries for the SWMHD hyperbolic system in a <u>rotating frame  $(1/2)$ .</u>

<span id="page-14-1"></span>Table 8: Adjoint representation of the seven admitted Lie symmetries for the SWMHD hyperbolic system in a rotating frame (2/2).

$Ad\left(e^{(\varepsilon \mathbf{X}_i)}\right) \mathbf{X}_j$	$\mathbf{Z}_1$	$\mathbf{Z}_2$	$\mathbf{Z}_3$
$\mathbf{X}_1$	$Z_1$	$\cos(f_0 \varepsilon) Z_2 - \sin(f_0 \varepsilon) Z_3 \quad \sin(f_0 \varepsilon) Z_2 + \sin(f_0 \varepsilon) Z_3$	
$\mathbf{X}_2$	$Z_1 - \varepsilon X_2$	$Z_2$	$Z_3$
$\mathbf{X}_4$	$Z_1$	$Z_2$	$Z_3$
$\mathbf{X}_{10}$	$Z_1-2\varepsilon X_{10}$	$Z_2$	$Z_3$
$\mathbf{Z}_1$	$Z_1$	$e^{\varepsilon}Z_2$	$e^{\varepsilon}Z_3$
$\mathbf{Z}_2$	$Z_1-\varepsilon Z_2$	$Z_2$	$Z_3$
$\mathbf{Z}_3$	$Z_1 - \varepsilon Z_3$	$Z_2$	$Z_3$

#### 4.4 Constant gravitational field and Coriolis force

For the general case of the one-dimensional hyperbolic system  $(20)$ ,  $(21)$ ,  $(22)$ ,  $(23)$  and  $(24)$ with  $gf_0 \neq 0$ , and g,  $f_0$  be constants, the Lie symmetry conditions are expressed as follows

$$
\xi^{t} = \xi_{0}^{t}, \ \xi^{x} = \xi^{x}(t), \ \eta^{b} = \eta^{b}(a, h),
$$
  

$$
\xi_{,ttt}^{x} + f_{0}^{2}\xi_{,t}^{x} = 0, \ \xi_{,xx}^{x} = 0, \ \xi_{,tx}^{x} = 0, \ a\eta_{,a}^{b} + \eta^{b} - b\xi_{,x}^{x} = 0,
$$
  

$$
\eta_{,h}^{b} - \xi_{,x}^{x}, \ h\eta_{,h}^{b} + \eta^{b} - b\xi_{,h}^{b} = 0, \ \eta^{a} = a\xi_{,x}^{x}, \ \eta^{h} = 2h\xi_{,x}^{x},
$$
  

$$
\eta^{u} = \xi_{,x}^{x}u + \xi_{,t}^{x}, \ \eta^{v} = \xi_{,x}^{x}v - \frac{1}{f_{0}}\xi_{,tt}^{x}.
$$

#### 4.4.1 Lie symmetries

The Lie symmetries for the hyperbolic system  $(20)$ ,  $(21)$ ,  $(22)$ ,  $(23)$  and  $(24)$  with g,  $f_0$  nonzero constants are

$$
X_1
$$
,  $X_2$ ,  $X_{10}$ ,  $Z_1$ ,  $Z_2$ ,  $Z_3$ .

which form the sixth-dimensional Lie algebra  $G^6$ . We remark that the Galileon symmetry is lost.

We observe that  $L^6 \subseteq L^7$  thus, the commutators are those presented in Table [6](#page-13-0) and the adjoint representation is that given in Table [7.](#page-14-0) Therefore,  $L^6$  is the  $A_{3,5} \rtimes A_{3,3}$  Lie algebra in the Morozov-Mubarakzyanov-Patera classification scheme [\[57–](#page-31-3)[61\]](#page-31-4).

# <span id="page-15-0"></span>5 One-dimensional optimal system for the SWMHD system

We proceed with deriving the one-dimensional optimal system for the six-dimensional Lie algebra admitted by the SWMHD system under a constant gravitational field and a nonzero Coriolis term.

The first step in deriving the one-dimensional optimal system is to find the invariants  $\phi(\mathbf{a}, \mathbf{z})$ of the adjoint representation. Based on the commutators of the Lie symmetries presented in Table [6,](#page-13-0) we define the following linear system of homogeneous partial differential equations

$$
z_1 \phi_{,a_2} = 0, \t\t(62)
$$

$$
2z_1\phi_{,a_{10}} = 0, \t\t(63)
$$

$$
z_2 \phi_{,z_3} - z_3 \phi_{,z_2} = 0, \tag{64}
$$

$$
f_0 a_1 \phi_{,z_3} - z_1 \phi_{,z_2} = 0, \tag{65}
$$

$$
f_0 a_1 \phi_{,z_2} + z_1 \phi_{,z_3} = 0, \tag{66}
$$

$$
a_2 \phi_{,z_2} + 2a_{10} \phi_{,a_{10}} + z_2 \phi_{,z_2} + z_3 \phi_{,z_3} = 0. \tag{67}
$$

Parameters  $a_1, a_2, a_{10}, z_1, z_2$  and  $z_3$  are the coefficient constants for the generic vector field

$$
\mathbf{X} = a_1 X_1 + a_2 X_2 + a_{10} X_{10} + z_1 Z_1 + z_2 Z_2 + z_3 Z_3. \tag{68}
$$

The solution of the latter system is  $\phi = \phi(a_1, z_1)$ , from where we infer that the invariants of the Adjoint representation are the  $a_1$  and  $z_1$ . This means that under the Action of the adjoint representation the vector filed  $X$  can be expressed as

$$
\mathbf{X}_{(I)} = a_1 X_1 + z_1 Z_1. \tag{69}
$$

On the hand, for  $z_1 = 0$  and  $a_1 \neq 0$ , we find  $\phi = \phi(a_1, a_{10})$  from where it follows that the generic vector field can be expressed as

$$
\mathbf{X}_{(II)} = a_1 X_1 + a_2 X_2 + a_{10} X_{10}.\tag{70}
$$

Furthermore, in the case  $z_1 \neq 0$  and  $a_1 = 0$ , we calculate  $\phi = \phi(z_1)$ , then it follows

$$
\mathbf{X}_{(III)} = z_1 Z_1. \tag{71}
$$

Moreover, when  $z_1 = 0$  and  $a_1 = 0$ , it follows  $\phi = \phi(a_2, a_{10}, z_2, z_3)$ , thus, the invariants are the  $a_2$ ,  $a_{10}$ ,  $z_2$  and  $z_3$  and the equivalent vector field reads

$$
\mathbf{X}_{(IV)} = a_2 X_2 + a_{10} X_{10} + z_2 Z_2 + z_3 Z_3. \tag{72}
$$

From the above we conclude that the one-dimensional optimal system for the Lie algebra

 $G<sup>6</sup>$  is consisted by the following one-dimensional Lie algebras

$$
\{X_1\}, \{X_2\}, \{X_3\}, \{X_{10}\}, \{Z_1\}, \{Z_2\}, \{Z_3\},
$$
  

$$
\{a_1X_1 + z_1Z_1\}, \{a_1X_1 + a_2X_2\}, \{a_1X_1 + a_{10}X_{10}\},
$$
  

$$
\{a_1X_1 + a_2X_2 + a_{10}X_{10}\}, \{a_2X_2 + a_{10}X_{10}\},
$$
  

$$
\{a_2X_2 + z_2Z_2\}, \{a_2X_2 + z_3Z_3\}, \{a_{10}X_{10} + z_2Z_2\},
$$
  

$$
\{a_{10}X_{10} + z_3Z_3\}, \{z_2Z_2 + z_3Z_3\}, \{a_2X_2 + a_{10}X_{10} + z_2Z_2\},
$$
  

$$
\{a_2X_2 + a_{10}X_{10} + z_3Z_3\}, \{a_{10}X_{10} + z_2Z_2 + z_3Z_3\},
$$
  

$$
\{a_2X_2 + a_{10}X_{10} + z_2Z_2 + z_3Z_3\}.
$$

### <span id="page-17-0"></span>6 Similarity solutions

We continue our analysis by constructing similarity solutions derived from the identified Lie symmetries. For each symmetry vector, we define a similarity transformation that can reduce the hyperbolic system  $(20)$ ,  $(21)$ ,  $(22)$ ,  $(23)$ , and  $(24)$  to a system of ordinary differential equations.

In the following, we use elements of the one-dimensional optimal system to define the similarity transformations, write the reduced system, and determine when a closed-form solution can be derived.

#### 6.1 Reduction with  $X_1$

The Lie symmetry vector  $X_1$  leads to the construction of the invariants  $x, h, u, v, a, b$ . Hence, we define the similarity transformation

$$
h = h(x), \ u = u(x), \ v = v(x), \ a = a(x), \ b = b(x), \tag{73}
$$

which describes static solution.

By replacing in  $(20)$ ,  $(21)$ ,  $(22)$ ,  $(23)$  and  $(24)$  we end with the following system or ordinary differential equations

$$
(hu)_{,x} = 0,\t(74)
$$

$$
\left(hu^2 + \frac{g}{2}h^2 - ha^2\right)_{,x} + f_0hv = 0,\t\t(75)
$$

$$
(huv - hab)_{,x} - f_0hu = 0,
$$
\n(76)

$$
u\left(ha\right)_{,x} = 0,\tag{77}
$$

$$
(h (ub - va))_{,x} + v (ha)_{,x} = 0,
$$
\n
$$
(78)
$$

For  $u \neq 0$ , it follows that

$$
u = \frac{u_0}{h}, \ v = -\frac{f_0 u_0 a_0}{a_0^2 - u_0^2} t + v_0,\tag{79}
$$

$$
a = \frac{a_0}{h}, \ b = -\frac{f_0 u_0^2}{a_0^2 - u_0^2} t + b_0,\tag{80}
$$

while  $h(t)$  satisfies the nonlinear differential equation

$$
\frac{(gh^3 + a_0^2 - u_0^2)}{h}h_{,t} - f_0^2 \frac{f_0 u_0^2}{a_0^2 - u_0^2}t + v_0 h = 0.
$$
\n(81)

The latter solution exist for initial conditions in which  $a_0^2 - u_0^2 \neq 0$  and  $u_0 \neq 0$ .

### 6.2 Reduction with  $X_2$

The Lie symmetry vector  $X_2$  leads to the similarity transformation

$$
h = h(t), u = u(t), v = v(t), a = a(t), b = b(t),
$$
\n(82)

which describes stationary solution.

Thus we derive the reduced system described by the following set of ordinary differential equations

 $h_{,t} = 0,$  (83)

$$
(hu)_{,t} + f_0 hv = 0,\t\t(84)
$$

$$
(hv)_{,t} - f_0 hu = 0,\t\t(85)
$$

$$
(ha)_{,t} = 0,\t\t(86)
$$

$$
(hb)_{,t} = 0,\t\t(87)
$$

from where it follows

$$
h = h_0, \ a = a_0, \ b = b_0,\tag{88}
$$

and

$$
u(t) = u_0 \cos(f_0 t) - v_0 \sin(f_0 t), \qquad (89)
$$

$$
v(t) = u_0 \sin(f_0 t) + v_0 \cos(f_0 t). \tag{90}
$$

### **6.3** Reduction with  $X_3$

The application of the Lie symmetry  $X_3$  gives the similarity transformation

$$
\sigma = \frac{x}{t}, \ h = h(\sigma), \ u = u(\sigma), \ v = v(\sigma), \ a = a(\sigma), \ b = b(\sigma), \tag{91}
$$

from where it follows the constant solution

$$
h = h_0, \ u = u_0, \ a = a_0, \ v = 0, \ b = 0. \tag{92}
$$

#### 6.4 Reduction with  $Z_1$

From the Lie symmetry vector  $Z_1$  we construct the similarity transformation

$$
h = x2 H(t), u = xU(t), v = xV(t), a = xA(t), b = xB(t).
$$
 (93)

Therefore, the hyperbolic system  $(20)$ ,  $(21)$ ,  $(22)$ ,  $(23)$  and  $(24)$  is reduced as follows

$$
H_{,t} + 3UH = 0,\t\t(94)
$$

$$
(HU)_{,t} + H\left(4\left(U^2 - A^2\right) + 2gH + f_0V\right) = 0,\t(95)
$$

$$
(HV)_{,x} + H(4(UV - AB) - f_0U) = 0,
$$
\n(96)

$$
(HA)_{,t} + 3UAH = 0,\t\t(97)
$$

$$
(HB)_{,t} + H(4UB - VA) = 0. \tag{98}
$$

Thus, by replacing  $H(t) = H_0 e^{-3 \int U(t) dt}$  the latter system is simplified as

$$
U_{,t} + U^2 - 4A + f_0 V + 2g e^{-3 \int U dt} = 0, \tag{99}
$$

$$
V_{,t} + UV - 4AB - f_0U = 0, \t(100)
$$

$$
A_{,t} = 0, \tag{101}
$$

$$
B_{,t} + (4UB - VA) = 0. \t(102)
$$

The analytic solution is expressed in elliptic integrals.

### 6.5 Reduction with  $Z_2$

The Lie symmetry vector  $Z_2$  provides the similarity transformation

$$
h = h(t), \ u = \cot(f_0 t) f_0 x + U(t), \ v = f_0 x + V(t), \ a = a(t), \ b = b(t).
$$
 (103)

Hence, the SWMHD equations provides the closed-form solution

$$
h(t) = \frac{h_0}{\sin(f_0 t)}, \ V(t) = V_0, \ a = a_0 \sin(f_0 t), \ b = -b_0 \cos(f_0 t) + b_1,
$$
 (104)

and

$$
U(t) = \frac{U_0}{\sin\left(f_0 t\right)} - h_0 \cot\left(f_0 t\right). \tag{105}
$$

The solution describes a shock wave with wall  $f_0t = n\pi$ ,  $n \in \mathbb{N}$ .

#### 6.6 Reduction with  $Z_3$

The Lie symmetry vector  $Z_3$  provides the similarity transformation

$$
h = h(t), \ u = -\tan(f_0 t) f_0 x + U(t), \ v = f_0 x + V(t), \ a = a(t), \ b = b(t).
$$
 (106)

Hence, the SWMHD equations provides the closed-form solution

$$
h(t) = \frac{h_0}{\cos\left(f_0 t\right)}, \ V(t) = V_0, \ a = a_0 \cos\left(f_0 t\right), \ b = -b_0 \sin\left(f_0 t\right) + b_1,\tag{107}
$$

and

$$
U(t) = \frac{U_0}{\cos(f_0 t)} - h_0 \tan(f_0 t).
$$
 (108)

The solution describes a shock wave with wall  $f_0 t = n \frac{\pi}{2}$  $\frac{\pi}{2}$ ,  $n \in \mathbb{N}$ . The similarity solutions  $Z_2$  and  $Z_3$  are related through the change of variables  $f_0 t \rightarrow f_0 t + \frac{\pi}{4}$  $\frac{\pi}{4}$ .

# **6.7** Reduction with  $X_1 + a_2X_2$

The Lie symmetry  $X_1 + a_2 X_2$  provides the similarity transformation

$$
\xi = x - a_2 t, \ h = h(\xi), \ u = u(\xi), \ v = v(\xi), \ a = a(\xi), \ b = b(\xi), \tag{109}
$$

where  $a_2$  is the speed of the travel-waves.

By replacing in the SWMHD equations we end with the system

$$
u = a_2 + \frac{u_0}{h(\xi)}, \ a = \frac{a_0}{h(\xi)}, \ b = \frac{a_0}{u_0}v(\xi) + b_0,
$$
\n(110)

<span id="page-20-0"></span>
$$
u_0 \left( 1 - \left( \frac{a_0}{u_0} \right)^2 \right) v_{,\xi} - f_0 \left( a_2 h + u_0 \right) = 0, \tag{111}
$$

$$
\frac{(a_0^2 - u_0^2)}{h^2} h_{,\xi} + \frac{1}{2} g \left( h^2 \right)_{,\xi} + f_0 v h = 0. \tag{112}
$$

The solution is expressed in elliptic integrals. The numerical solution of the system [\(111\)](#page-20-0), [\(112\)](#page-20-0) is given in Fig. [1.](#page-21-0)



<span id="page-21-0"></span>Figure 1: Qualitative evolution of the functions  $h(\xi)$  and  $v(\xi)$  as they are given by numerical simulation of the nonlinear system  $(111)$ ,  $(112)$ .

# **6.8** Reduction with  $X_2 + z_2Z_2$

From the vector field  $X_2 + z_2 Z_2$  we define the similarity transformation

$$
\zeta = x + \frac{z_2}{f_0} \cos(f_0 t), \quad h = h(\zeta), \quad a = a(\zeta), \quad b = b(\zeta), \tag{113}
$$

$$
u = z_2 \sin(f_0 t) + U(\zeta), \ v = -z_2 \cos(f_0 t) + V(\zeta).
$$
 (114)

Therefore the hyperbolic system  $(20)$ ,  $(21)$ ,  $(22)$ ,  $(23)$  and  $(24)$  provides

$$
a = \frac{a_0}{h}, \ U = \frac{u_0}{h}, \ b = -\frac{f_0 a_0 u_0}{a_0^2 - u_0^2} \zeta + b_0 \tag{115}
$$

$$
V = -\frac{f_0 u_0^2}{a_0^2 - u_0^2} \zeta + v_0,\tag{116}
$$

and

<span id="page-21-1"></span>
$$
\frac{1}{2} \left( \frac{1}{h^2} \right)_{,\zeta} \left( u_0^2 - a_0^2 - gh^3 \right) - \frac{f_0^2 u_0^2}{a_0^2 - u_0^2} \zeta + f_0 v_0 = 0. \tag{117}
$$

Numerical simulations of the latter nonlinear ordinary differential equations are presented in Fig. [2](#page-22-0)



<span id="page-22-0"></span>Figure 2: Qualitative evolution for the function  $h(\zeta)$  as it is given by numerical simulation of the nonlinear differential equation [\(117\)](#page-21-1).

# **6.9** Reduction with  $X_{10} + z_2Z_2$

From the Lie symmetry vector  $X_{10} + z_2 Z_2$  we find the similarity transformation

$$
h = h(t), \ u = \cot(f_0 t) f_0 x + U(t), \ v = f_0 x + V(t), \qquad (118)
$$

$$
a = a(t), b = \frac{x}{z_2 \sin(f_0 t) h(t) a(t)} + B(t).
$$
 (119)

Thus, from the SWMHD equations it follows

$$
h(t) = \frac{h_0}{\sin\left(f_0 t\right)}, \ V\left(t\right) = \frac{t}{h_0 z_2} + V_0,\tag{120}
$$

$$
a(t) = a_0 \sin\left(f_0 t\right),\tag{121}
$$

$$
U(t) = \frac{U_0}{\sin(f_0 t)} + \frac{1}{h_0 z_2} \left( \cot(f_0 t) (t + h_0 V_0 z_2) - \frac{1}{f_0} \right),
$$

and

$$
B(t) = -a_0 \cos\left(f_0 t\right) + \frac{t}{\left(h_0 z_2\right)^2 a_0 \sin\left(f_0 t\right)} + \frac{\left(V_0 + U_0 \cos\left(f_0 t\right)\right)}{h_0 z_2 a_0 \sin\left(f_0 t\right)} + B_0.
$$
 (122)

We observe that this solution describes shoch waves with wall at  $f_0t = n\pi$ .

### **6.10** Reduction with  $X_{10} + z_3 Z_3$

From the application of the Lie symmetry vector  $X_{10} + z_3 Z_3$  we derive the similarity transformation

$$
h = h(t), \ u = -\tan(f_0 t) f_0 x + U(t), \ v = f_0 x + V(t), \tag{123}
$$

$$
a = a(t), b = \frac{x}{z_3 \cos(f_0 t) h(t) a(t)} + B(t).
$$
 (124)

Thus, from the SWMHD equations it follows

$$
h(t) = \frac{h_0}{\cos\left(f_0 t\right)}, \ V\left(t\right) = \frac{t}{h_0 z_2} + V_0,\tag{125}
$$

$$
a(t) = a_0 \cos(f_0 t), \qquad (126)
$$

$$
U(t) = \frac{U_0}{\cos(f_0 t)} - \frac{1}{h_0 z_3} \left( \tan(f_0 t) (t + h_0 V_0 z_2) + \frac{1}{f_0} \right),
$$
\n(127)

and

$$
B(t) = a_0 \sin\left(f_0 t\right) + \frac{t}{\left(h_0 z_3\right)^2 a_0 \cos\left(f_0 t\right)} + \frac{\left(V_0 - U_0 \sin\left(f_0 t\right)\right)}{h_0 z_2 a_0 \cos\left(f_0 t\right)} + B_0.
$$
\n(128)

We observe that this solution describe solitons. We observe that this solution describes shoch waves with wall at  $f_0t = n\pi$ . The latter two solutions are related through the transformation  $f_0 t \rightarrow f_0 t + \frac{\pi}{4}$  $\frac{\pi}{4}$ .

# **6.11** Reduction with  $X_2 + a_{10}X_{10} + z_2Z_2$

The generic vector field  $X_2 + a_{10}X_{10} + z_2Z_2$  leads to the derivation of the similarity transformation

$$
h = h(t), \ u = \frac{z_2 \cos\left(f_0 t\right) f_0}{1 + z_2 \cos\left(f_0 t\right)} x + U(t), \ v = \frac{z_2 \sin\left(f_0 t\right)}{1 + z_2 \sin\left(f_0 t\right)} + V(t), \tag{129}
$$

$$
a = a(t), b = \frac{x}{h(t) a(t) (1 + z_2 \cos(f_0 t))} + B(t).
$$
 (130)

By replacing the later transformation in the hyperbolic SWMHD [\(20\)](#page-6-1), [\(21\)](#page-6-1), [\(22\)](#page-6-1), [\(23\)](#page-6-1) and [\(24\)](#page-6-1) system we determine the closed-form solution

$$
h(t) = \frac{h_0}{1 + z_2 \sin\left(f_0 t\right)},\tag{131}
$$

$$
a(t) = a_0 \left(1 + z_2 \sin\left(f_0 t\right)\right),
$$
  
\n
$$
V(t) = \frac{\sin\left(f_0 t\right) V_0 + \cos\left(f_0 t\right) U_0}{1 + z_2 \sin\left(f_0 t\right)} + \frac{a_{10} z_2}{h_0} \frac{t}{\left(1 + z_2 \sin\left(f_0 t\right)\right)},
$$
\n(132)

$$
U(t) = \frac{f_0 h_0 (U_0 \cos (f_0 t) + V_0 \sin (f_0 t)) + a_{10} z_2 \sin (f_0 t)}{h_0 f_0 (1 + z_2 \sin (f_0 t))},
$$
(133)

and the magnetic field  $B(t)$  is given by the first-order linear ordinary differential equation

$$
B_{,t} = -\frac{h_{,t}}{h}B - \frac{a_{10}U}{ha\left(1+z_2\sin\left(f_0t\right)\right)} + \frac{z_2^2 f_0 \left(V\sin\left(2f_0t\right) - 2a\sin\left(f_0t\right)^2\right) + z_2 f_0 \left(a\sin\left(f_0t\right) - V\cos\left(f_0t\right)\right)}{2\left(1-z_2\sin^2\left(f_0t\right)\right)}.
$$
\n(134)

When  $|z_2| \leq 1$ , then solitons exist.

Moreover, the vector field  $X_2 + a_{10}X_{10} + z_3Z_3$  leads to the derivation of a similar similarity solution which can derived after a phase transition.

In Fig. [3](#page-25-0) we present the qualitative evolution ofor the similarity solution derived before. The plots are for unitary values of the free parameters.

### <span id="page-24-0"></span>7 Conclusions

In this work, we investigated the Lie symmetries of a hyperbolic system consisting of five firstorder nonlinear partial differential equations that describe an SWMHD system. Hyperbolic systems are crucial for describing various physical phenomena. Symmetry analysis is a powerful tool for the analytical study of nonlinear differential equations, and Lie symmetries offer a systematic approach to better understand these systems.

In the context of the SW equations, Lie symmetry analysis has been employed to determine analytic solutions and conservation laws. Similarly, Lie theory has played a significant role in the study of MHD systems. We considered a one-dimensional SWMHD model with the relaxing condition  $\nabla (h\mathbf{B}) \neq 0$ , which is physically acceptable as long as the boundary condition  $\nabla(h\mathbf{B})=0$  is met.

The introduction of the term  $\nabla(h\mathbf{B})$  restores Galilean symmetries in the free SWMHD system without the Coriolis term. We found that the admitted symmetries number ten and form the Lie algebra  $A_{3,3} \rtimes A_{2,1} \rtimes A_{5,34}^a$ . When a constant gravitational field is present, the number of admitted Lie symmetries reduces to eight, and the Lie algebra is  $A_{2,1} \rtimes A_{6,22}$ . These results differ from those of the one-dimensional SWMHD system where the condition  $\nabla (h\mathbf{B}) = 0$  is imposed.

Furthermore, when a Coriolis force term appears, related to the rotating reference frame, the number of admitted Lie symmetries is further reduced to seven or six, depending on the presence of the gravitational field. The corresponding Lie algebras are  $A_{3,5} \rtimes A_{2,1} \rtimes A_{2,1}$  and  $A_{3,5} \rtimes A_{3,3}$ , respectively. For the most general case, where the Lie symmetries form the  $A_{3,5} \rtimes A_{3,3}$  Lie algebra, we calculated the one-dimensional optimal system and used similarity transformations to reduce the hyperbolic system to a system of ordinary differential equations, from which we



<span id="page-25-0"></span>Figure 3: Qualitative evolution for the similarity solution of  $h(t)$ ,  $a(t)$ ,  $V(t)$ ,  $U(t)$  as they are given by the application of the Lie symmetry vector  $X_2 + a_{10}X_{10} + z_2Z_2$ .

derived the analytic solution. The one-dimensional optimal system was obtained by determining all the invariants of the adjoint action for the  $A_{3,5} \rtimes A_{3,3}$  Lie algebra and its subalgebras.

In this study, we have used symmetries to construct invariants that are primarily applied to reduce the system of hyperbolic equations. Nevertheless, another important feature of symmetry analysis is that it can be used to identify conservation laws [\[62\]](#page-31-5). There are different approaches in the literature regarding how to use symmetries to identify conservation laws. For systems that follow from a variational principle, Noether's theorem is mainly applied. Another approach is Ibragimov's conservation law theorem [\[63\]](#page-31-6). For other studies, we refer the reader to [\[64](#page-31-7)[–66\]](#page-31-8) and the references therein.

In future work, we plan to further investigate the algebraic properties of this specific SWMHD model by applying the theory of higher-order symmetries and determining conservation laws related to the symmetry vectors. Additionally, the symmetry analysis of the SWMHD system in Lagrangian coordinates will be of special interest.

Data Availability Statements: Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

Code Availability Statements: Code sharing is not applicable to this article as no code was used in this study.

Acknowledgements: AP thanks the support of Vicerrectoria de Investigación y Desarrollo Tecnológico (Vridt) at Universidad Católica del Norte through Núcleo de Investigación Geometría Diferencial y Aplicaciones, Resolución Vridt No - 096/2022 and Resolución Vridt No -098/2022. Part of this work was supported by Proyecto Fondecyt Regular 2024, Folio 1240514, Etapa 2024. AP thanks AH and the Woxsen University for the hospitality provided while part of this work was performed.

### References

- <span id="page-26-0"></span>[1] A.J.C. Barre de Saint-Venant, Théorie du mouvement non permanent des eaux, avec application aux crues des rivières et a l'introduction de marées dans leurs lits, Comptes Rendus de l'Académie des Sciences 73, 147 (1871)
- <span id="page-26-1"></span>[2] O. Castro-Orgaz and W.H. Hager, Shallow Water Hydraulics, Springer Cham, New York (2019)
- <span id="page-26-2"></span>[3] N. Paldor, Shallow Water Waves on the Rotating Earth, Springer Cham, New York (2015)
- [4] B.K. Valery and P. James Lynch, Fundamentals of Shallow Water Acoustics, Springer New York, New York (2012)
- [5] P. García-Navarro, J. Murillo, J. Fernández-Pato, I. Echeverribar and M. Morales-Hernandez, The shallow water equations and their application to realistic cases, Enviromental Fluid Mechanics 19, 1235 (2019)
- [6] A. Kurganov, Finite-volume schemes for shallow-water equations, Acta Numerica 27, 289 (2018)
- <span id="page-27-0"></span>[7] I. Kinnmark, The Shallow Water Wave Equations: Formulation, Analysis and Application, Lecture Notes in Engineering, Springer Berlin, Heidelberg (1986)
- <span id="page-27-1"></span>[8] Y. Taitel and D. Barena, Encyclopedia of Two-Phase Heat Transfer and Flow I, Fundamentals and Methods Volume 1 : Modeling of Gas Liquid Flow in Pipes, edited by J.R. Thome, World Scientiffic Publishing Co. Pte. Ltd., Singapure, (2015)
- <span id="page-27-2"></span>[9] G.F. Hewitt, Two-Phase Flow and Its Applications: Past, Present, and Future, Heat Transfer Engineering, 4, 67 (1983)
- <span id="page-27-3"></span>[10] P.A. Gilman, Magnetohydrodynamic "shallow water" equations for the solar tachocline, Astroph. J. 544, L79 (2000)
- <span id="page-27-4"></span>[11] E.A Spiegel and J.-P. Zahn, The solar tachocline, A&A 265, 106 (1992).
- <span id="page-27-5"></span>[12] D.W. Hyghes, R. Rosner and N.O. Weiss, The Solar Tachocline, Cambridge University Press, Cambridger (2009)
- <span id="page-27-6"></span>[13] D.A. Schecter, J.F Boyd and P.A. Gilman, "Shallow-Water" Magnetohydrodynamic Waves in the Solar Tachocline, Ap. J. 551, L185 (2001)
- <span id="page-27-7"></span>[14] T. V. Zaqarashvili, R. Oliver, J.L. Ballester and B.M. Shergelashvili, Rossby waves in "shallow water" magnetohydrodynamics, A&A 470, 815 (2007)
- <span id="page-27-8"></span>[15] A.W. Hindle, P.J. Bunsby and T.M. Rogers, Observational Consequences of Shallow-water Magnetohydrodynamics on Hot Jupiters, Ap.J. Lett. 916, L8 (2021)
- [16] D. Alonso-Oran, Asymptotic Shallow Models Arising in Magnetohydrodynamics, Water Waves 3, 371 (2021)
- <span id="page-28-0"></span>[17] A. Petrosyan, D. Klimachkov, M. Fedotova and T. Zinyakov, Shallow Water Magnetohydrodynamics in Plasma Astrophysics. Waves, Turbulence, and Zonal Flows, Atmosphere 11, 314 (2020)
- <span id="page-28-1"></span>[18] N.H. Ibragimov, CRC Handbook of Lie Group Analysis of Differential Equations, Volume I: Symmetries, Exact Solutions, and Conservation Laws, CRS Press LLC, Florida (2000)
- <span id="page-28-7"></span>[19] G.W. Bluman and S. Kumei, Symmetries of Differential Equations, Springer-Verlag, New York, (1989)
- <span id="page-28-8"></span>[20] H. Stephani, Differential Equations: Their Solutions Using Symmetry, Cambridge University Press, New York, (1989)
- <span id="page-28-2"></span>[21] P.J. Olver, Applications of Lie Groups to Differential Equations, Springer-Verlag, New York, (1993)
- <span id="page-28-3"></span>[22] B. Gwaxa, S. Jamal and A.G. Johnpillai, On the Optimal System and Series Solutions of Fifth-Order Fujimoto-Watanabe Equations 17, 557 (2023)
- [23] S. Moyo and P.G.L. Leach, Symmetry Properties of Autonomous Integrating Factors, Sigma 1, 024 (2005)
- [24] T. Christodoulakis, A. Karagiorgos and A. Zampeli, Symmetries in Classical and Quantum Treatment of Einstein's Cosmological Equations and Mini-Superspace Actions, Symmetry 10, 70 (2018)
- [25] T. Christodoulakis, N. Dimakis and P.A. Terzis, Lie point and variational symmetries in minisuperspace Einstein gravity, J. Math. Phys.: Math. Theor. 47, 095202 (2014)
- [26] A. Paliathanasis, Geometric Linearization for Constraint Hamiltonian Systems, Symmetry 16, 988 (2024)
- <span id="page-28-4"></span>[27] A.I. Aliy, J.S. Yusuf, M.M. Nauman, D.U. Ozsahin, B.G. Agaie, J.H. Zaini and H. Umar, Lie Symmetry Analysis and Explicit Solutions to the Estevez–Mansfield–Clarkson Equation, Symmetry 16, 1194 (2024)
- <span id="page-28-5"></span>[28] A.A. Chesnokov, Symmetries and exact solutions of the shallow water equations for a two-dimensional shear flow, J. Appl. Mech. Techn. Phys. 49, 737 (2008)
- <span id="page-28-6"></span>[29] B. Bira, T.R. Sekhar and D. Zeidan, Application of Lie groups to compressible model of two-phase flows, Comput. Math. Appl. 71, 46 (2016)
- [30] A.A. Chesnokov, Symmetries and exact solutions of the rotating shallow-water equations, Eur. J. Appl. Math. 20, 461 (2009)
- <span id="page-29-0"></span>[31] A.V. Aksenov and K.P. Druzhkov, Conservation laws and symmetries of the shallow water system above rough bottom, J. Phys.: Conf. Ser. 722, 012001 (2016)
- [32] A. Paliathanasis, One-dimensional optimal system for 2D Rotating Ideal Gas, Symmetry 11, 1115 (2019)
- [33] A. Paliathanasis, Similarity transformations for modified shallow water equations with density dependence on the average temperature, Int. J. Nonl. Sci. Num. Sim. 24, 1095 (2023)
- [34] A.S. Pavlenko, Symmetries and solutions to equations of two-dimensional motions of polytropic gas, Sib. Elektron. Mat. Izv., 2, 291 (2005) `
- <span id="page-29-1"></span>[35] C. Huo and L. Li, Lie Symmetry Analysis, Particular Solutions and Conservation Laws of a New Extended (3+1)-Dimensional Shallow Water Wave Equation, Symmetry 14, 1855 (2022)
- <span id="page-29-2"></span>[36] V.A. Dorodnitsyn, E.I. Kaptsov, R.V. Kozlov and S. Meleshko, One-dimensional MHD flows with cylindrical symmetry: Lie symmetries and conservation laws, Int. J. Non-Linear Mechanics 148, 104290 (2023)
- [37] G.M. Webb and G.P. Zank, Fluid relabelling symmetries, Lie point symmetries and the Lagrangian map in magnetohydrodynamics and gas dynamics, J. Phys. A.: Math. Theor. 40, 545 (2007)
- [38] O. Bogoyavlenskij, Restricted Lie point symmetries and reductions for ideal magnetohydrodynamics equilibria, J. Engineering Mathematics 66, 141 (2010)
- [39] M. Liu and H. Dong, On the existence of solution, Lie symmetry analysis and conservation law of magnetohydrodynamic equations, CNSNS 87, 105277 (2020)
- [40] A. Paliathanasis, Group properties and solutions for the 1D Hall MHD system in the cold plasma approximation, Eur. Phys. J. Plus 136, 538 (2021)
- [41] G.M. Webb and R.L. Mace, Potential Vorticity in Magnetohydrodynamics, J. Plasma Phys. 81, 905810115 (2015)
- <span id="page-30-0"></span>[42] S.V. Meleshko, E.I. Kaptsov, S. Moyo and G.M. Webb, Group classification of the twodimensional magnetogasdynamics equations in Lagrangian coordinates, Math.Meth. Appl. Sci. 46, 15367 (2023)
- <span id="page-30-1"></span>[43] E.I. Kaptsov, S.V. Meleshko and V.A. Dorodnitsyn, Symmetries and conservation laws of the one-dimensional shallow water magnetohydrodynamics equations in Lagrangian coordinates, J. Phys. A.: Math. Theor. 55, 495202 (2022)
- <span id="page-30-2"></span>[44] F. Bouchut and X. Lhebrand, A 5-wave relaxation solver for the shallow water MHD system, J. Sci. Comput. 68, 92 (2015)
- [45] F. Bouchut and X. Lhebrand, A multi well-balanced scheme for the shallow water MHD system with topography, Numerische Mathematik 136, 875 (2017)
- [46] J.L. Ye, H.B. Guo and Y.B. Hu, An initial-boundary value problem for the one-dimensional rotating shallow water magnetohydrodynamic equations, J. Math. Anal. Appl. 527, 127422 (2023)
- <span id="page-30-3"></span>[47] J. Mak, S.D Griffiths and D.W. Hughes, Shear flow instabilities in shallow-water magnetohydrodynamics, J. Fluid Mechanics 788, 767 (2016)
- <span id="page-30-6"></span>[48] P.J. Dellar, Lattice Kinetic Schemes for Magnetohydrodynamics, J. Comput. Phys. 172, 392 (2001)
- <span id="page-30-4"></span>[49] H. De Sterck, Hyperbolic theory of the "shallow water" magnetohydrodynamics equation, Physics of Plasmas 8, 3293 (2001)
- <span id="page-30-5"></span>[50] P.J. Dellar, Dispersive shallow water magnetohydrodynamics, Physics of Plasmas 10, 581 (2003)
- <span id="page-30-7"></span>[51] K.G. Powell, An approximate Riemann solver for magnetohydrodynamics (that works in more than one dimension), NASA CR-194902 and ICASE Report No. 94-24, NASA Langley Research Center (1994)
- <span id="page-30-9"></span>[52] P. Janhunen, A Positive Conservative Method for Magnetohydrodynamics Based on HLL and Roe Methods, J. Comput. Phys. 160, 649 (2000)
- <span id="page-30-8"></span>[53] I.V. Sokolov, K.G. Powell, O. Cohen and T.I. Gombosi, Computational MagnetoHydrodynamics, Based on Solution of the Well-Posed Riemann Problem, Numerical Modeling of Space Plasma Flows: Astronum 2007, Edited by N.V. Pogorelov, E. Audit, and G.P. Zank, ASP Conference Series, Vol. 385 (2008)
- <span id="page-31-0"></span>[54] P. J. Morrison, in Mathematical Methods in Hydrodynamics and Integrability of Dynamical Systems, edited by M. Tabor and Y. M. Treve, AIP Conference Proceedings: American Institute of Physics, New York, Vol. 88 of (1982)
- <span id="page-31-1"></span>[55] P.J. Dellar, Hamiltonian and symmetric hyperbolic structures of shallow water magnetohydrodynamics, Physics of Plasmas 9, 1130 (2002)
- <span id="page-31-2"></span>[56] F. Bouchut and X. Lhébrard, A 5-Wave Relaxation Solver for the Shallow Water MHD System, J. Sci. Comput. 68, 98 (2016)
- <span id="page-31-3"></span>[57] V.V. Morozov, Classification of six-dimensional nilpotent Lie algebras, Izvestia Vysshikh Uchebn Zavendenii Matematika 5, 161 (1958)
- [58] G.M. Mubarakzyanov, On solvable Lie algebras, Izvestia Vysshikh Uchebn Zavendenii Matematika 32, 114 (1963)
- [59] G.M. Mubarakzyanov, Classification of real structures of five-dimensional Lie algebras, Izvestia Vysshikh Uchebn Zavendenii Matematika 34, 99 (1963)
- [60] G.M. Mubarakzyanov, Classification of solvable six-dimensional Lie algebras with one nilpotent base element, Izvestia Vysshikh Uchebn Zavendenii Matematika 35, 104 (1963)
- <span id="page-31-4"></span>[61] J. Patera, R. T. Sharp, P. Winternitz and H. Zassenhaus, Invariants of real low-dimensional Lie algebras, J. Math. Phys. 17, 986 (1976)
- <span id="page-31-5"></span>[62] E. Noether, Invariante Variationsprobleme, Nach. Konig. Gesell. Wissen. Gottingen, Math.-phys. Kl 235, (1918)
- <span id="page-31-6"></span>[63] N.H. Ibragimov, A new conservation theorem, J. Math. Anal. Appl. 333, 311 (2007)
- <span id="page-31-7"></span>[64] M.L. Gandarias, Weak self-adjoint differential equations, J. Phys. A: Math. Theor. 44, 262001 (2011)
- [65] Z. Yi, Symmetries and Two Types of Non-Noether Conservation Laws of Birkhoffian System with Unilateral Constraints, Commun. Theor. Phys. 45, 239 (2006)
- <span id="page-31-8"></span>[66] S.C. Anco, On the incompleteness of Ibragimov's conservation law theorem and its equivalence to a standard formula using symmetries and adjoint-symmetries,Symmetry 9, 33 (201)